On a t-exactness property of the Harish-Chandra transform

Roman Gonin, Andrei Ionov, Kostiantyn Tolmachov

Abstract

Using hyperbolic localization, we identify the nearby cycles along the Vinberg degeneration with the composition of Radon and Harish-Chandra functors, both considered for the category of character sheaves. This provides a new, simple proof of the exactness of this composition, extending previously known results to arbitrary monodromy and more general sheaf-theoretic set-ups.

1 Introduction.

1.1 Notations. For a stack \mathcal{X} defined over \mathbb{C} , let $D^b(\mathcal{X})$ be the bounded algebraically constructible derived category of sheaves on \mathcal{X} with analytic topology, with coefficients in a ring \mathbb{k} . All the stacks we encounter will be quotient stacks of the form X/H, where X is a scheme and H is an algebraic group over \mathbb{C} . The sheaf theory for such stacks is described in [BL06]. We write $pt = \operatorname{Spec}(\mathbb{C})$. We write $\underline{\mathbb{k}}_{\mathcal{X}}$ for the constant sheaf on \mathcal{X} .

For an algebraic group H acting on a scheme X and its subgroup H^\prime let

$$\operatorname{For}_{H'}^H \colon D^b(H \setminus X) \to D^b(H' \setminus X)$$

stand for the forgetful functor, that is the *-pullback along the projection $H' \setminus X \to H \setminus X$. Let

$$\operatorname{Av}_{H',!}^{H}, \operatorname{Av}_{H',*}^{H} \colon D^{b}(H' \setminus X) \to D^{b}(H \setminus X)$$

stand for the ! and *-pushforward from $H' \setminus X$ to $H \setminus X$, respectively. If H' is clear from the context, we will omit it from the notation, writing For^H, $Av_!^H$, Av_*^H , respectively.

We will adopt the same notation for the right action of the group, writing R For, R Av for the right forgetful and averaging functors, when the emphasis is needed.

1.2 Background and the main result. Let G be a reductive group, B a fixed Borel subgroup, $U \subset B$ its unipotent radical and $T \subset B$ a maximal torus. The Harish-Chandra transform \mathfrak{hc}_1 for constructible sheaves was introduced by Lusztig [Lus85] in his study of character sheaves. It is given by a pull-push functor along the horocycle correspondence

$$G/_{\mathrm{Ad}}G \leftarrow G/_{\mathrm{Ad}}B \rightarrow (U\backslash G/U)/_{\mathrm{Ad}}T$$

 $\mathfrak{hc}_{\mathfrak{l}} := \mathrm{Av}_{\mathfrak{l}}^U \circ \mathrm{For}^G$.

It was observed by Ginzburg [Gin89] that it produces objects in the categorical center of the finite Hecke category attached to a reductive group G and its Borel subgroup B. This observation has been elaborated upon in various contexts in (among other works) [BFO12], [BN09], [Lus], [Lus15], [HL23], [BITV23]. Another remarkable property of the Harish-Chandra transform on the character sheaves, playing a key role in [BFO12], is that it becomes *t*-exact and Verdier selfdual after composing it with the Radon transform functor R_* . The latter transform is given by the averaging with respect to the unipotent radical U^- of the opposite Borel subgroup B^- , or, equivalently, a pull-push functor along the following diagram:

$$(U \setminus G/U)/_{\mathrm{Ad}}T \leftarrow (G/U)/_{\mathrm{Ad}}T \rightarrow (U^- \setminus G/U)/_{\mathrm{Ad}}T,$$

 $R_* := \operatorname{Av}^{U^-}_* \circ \operatorname{For}^{U}[2\dim U] \colon D^b\left((U \setminus G/U)/_{\operatorname{Ad}}T\right) \to D^b\left((U^- \setminus G/U)/_{\operatorname{Ad}}T\right).$

For a local system \mathcal{L} of finite order, let $D^b_{\mathfrak{C}_{\mathcal{L}}}(G)$ be the derived category of character sheaves with the character \mathcal{L} (see Section 2.2 for the precise definition). Our main result is as follows:

Theorem 1.2.1 (Corollary 2.3.2 and Remark 2.3.3). The functor $R_* \circ \mathfrak{hc}_!$, when restricted to the category $D^b_{\mathfrak{C}_{\mathcal{L}}}(G)$, is t-exact with respect to the perverse t-structure and commutes with the Verdier duality.

We prove the Theorem 1.2.1 by identifying $R_* \circ \mathfrak{hc}_!$ with the nearby cycles functor along the Vinberg degeneration. The key idea is to use that the nearby cycles functor commutes with the hyperbolic localization, as shown in [Nak16], [Ric19].

All our methods are geometric and can be restated with the same proofs, up to standard modifications, for $D^b(\mathcal{X})$ for stacks \mathcal{X} over \mathbb{C} being replaced by either of the following:

- the bounded derived category of holonomic *D*-modules,
- the bounded constructible derived category of $\overline{\mathbb{Q}}_{\ell}$ -sheaves on a stack \mathcal{X} defined over an algebraically closed field of characteristic p > 0 for primes $p \neq \ell$ as in [LO08].

We expect that the methods of the present paper can be extended to handle the parabolic case. We intend to treat it in the subsequent publication.

1.3 Relation to the previous works. The idea to relate the nearby cycles functor along the Vinberg degeneration with the functor $R_* \circ \mathfrak{hc}_!$ appears already in [BFO12], where an analogue of Theorem 1.2.1 in the setting of D-modules is also proved. The same idea was used in [CY17] to prove the Theorem 1.2.1 for unipotent character sheaves (that is for a trivial local system \mathcal{L}), with methods applicable to either of the sheaftheoretic cases, and also in the more general setting associated to a symmetric pair. It is very plausible that the methods of [CY17] can be adapted to treat the general monodromy \mathcal{L} . However, we believe that the systematic use of hyperbolic localization presented in this paper makes for a particularly short and straightforward proof.

1.4 Acknowledgments. We would like to thank Roman Bezrukavnikov, Alexander Yom Din, and Yakov Varshavsky for helpful discussions. The work of R.G. was supported by the EPSRC grant EP/V053787/1. R.G. also acknowledges the hospitality of the Hebrew University of Jerusalem during his postdoctoral fellowship. While working on the project, K. T. was at different times funded by the EPSRC programme grant EP/R034826/1 and by the Deutsche Forschungsgemeinschaft SFB 1624 grant, Projektnummer 506632645.

2 Setup and the statement of the main results.

2.1 Limit functors. For a scheme \widetilde{X} and a map $f: \widetilde{X} \to \mathbb{A}^1$, let

$$\psi_f \colon D^b(\widetilde{X} \times_{\mathbb{A}^1} \mathbb{G}_m) \to D^b(\widetilde{X} \times_{\mathbb{A}^1} \mathrm{pt}).$$

stand for the nearby cycles functor. Here $\times_{\mathbb{A}^1}$ denotes the fiber product with respect to embeddings $\text{pt} \to \mathbb{A}^1 \leftarrow \mathbb{G}_m$: the embedding of 0 into \mathbb{A}^1 and the complementary open embedding, respectively.

Assume that \mathbb{G}_m acts on X and let $a: \mathbb{G}_m \times X \to X$ be the action map. Consider $\widetilde{X} = X \times \mathbb{A}^1$. Let $f: \widetilde{X} \to \mathbb{A}^1$ be the projection to the second factor. Following [CY17], define the limit functor

$$\mathbb{L}: D^b(X) \to D^b(X), \quad \mathbb{L} = \psi_f \circ a^*[1].$$

Let *H* be an algebraic group acting on *X*, and let us fix a cocharacter $\gamma \colon \mathbb{G}_m \to H$. This defines the \mathbb{G}_m -action on *X*, and we write

$$\mathbb{L}_{\gamma}: D^b(X) \to D^b(X)$$

for the corresponding limit functor.

Let G be a reductive group and T its maximal torus. Choice of a regular cocharacter $\lambda \colon \mathbb{G}_m \to T$ fixes a choice of a pair of opposite Borel subgroups $B, B^- \supset T$ for which λ is dominant and antidominant, respectively. Let U and U^- be the unipotent radicals of B and B^- , respectively.

Let $W = N_G(T)/T$ be the Weyl group of G, and let $w_0 \in W$ be its longest element. We have two projections $q_B \colon B \to T$ and $q_{B^-} \colon B^- \to T$.

Let $K \simeq G$ stand for the diagonal subgroup of $G \times G$, and define three subgroups $K_0, K_\infty, K' \subset G \times G$ as the following fiber products

(1)
$$K_0 \simeq B \times_T B^-, \quad K_\infty \simeq B^- \times_T B, \quad K' \simeq B \times_T B,$$

with respect to the homomorphisms $q_B \colon B \to T$ and $q_{B^-} \colon B^- \to T$.

Fix a cocharacter of $H=G\times G$ given by

$$\gamma: \mathbb{G}_m \to G \times G, \quad \gamma = (\lambda^{-1}, \lambda).$$

Let X be a scheme equipped with a left action of $G \times G$. As shown in [CY17, Lemma 4.12], the limit functor \mathbb{L}_{γ} associated with this case, can be lifted to either of the following functors:

(2)
$$\mathbb{L}_{\gamma} \colon D^{b}(K \backslash X) \to D^{b}(K_{0} \backslash X),$$

(3)
$$\mathbb{L}_{\gamma} \colon D^{b}(K' \backslash X) \to D^{b}(K' \backslash X).$$

Remark 2.1.1. Consider the constant group scheme $\widetilde{G} = G \times \mathbb{P}^1$ over \mathbb{P}^1 . One can define its subgroup scheme \widetilde{K} , with the fibers $K_t = \gamma(t)K\gamma(t)^{-1}$ for $t \neq 0, \infty$, and the fibers over 0 and ∞ given by K_0 and K_∞ , see (1). One can verify that \widetilde{K} is a well-defined smooth group scheme over \mathbb{P}^1 , see [DG14, Proposition 2.5.2].

Let $\widetilde{X} = X \times \mathbb{P}^1$ and $\widetilde{K}' = K' \times \mathbb{P}^1$ be the constant scheme/constant group scheme over \mathbb{P}^1 . Informally, the functors (2) and (3) can be thought of as the limit functor for the families $\widetilde{K} \setminus \widetilde{X}$, $\widetilde{K}' \setminus \widetilde{X} \to \mathbb{P}^1$. We refer to [CY17] for details. We collect the required properties of this functor in the following lemma. These follow from the standard properties of the nearby cycles functor, see [SGA06, XIII].

Lemma 2.1.2. Let X_1, X_2 be schemes equipped with a \mathbb{G}_m -action, and let $f : X_1 \to X_2$ be a \mathbb{G}_m -equivariant map.

- (a) There is a natural transformation $f_! \circ \mathbb{L} \to \mathbb{L} \circ f_!$ of functors $D^b(X_1) \to D^b(X_2)$. This natural transformation is an isomorphism if f is proper.
- (b) There is a natural transformation $f^* \circ \mathbb{L} \to \mathbb{L} \circ f^*$ of functors $D^b(X_2) \to D^b(X_1)$. This natural transformation is an isomorphism if f is smooth.

Consider a Cartesian diagram

$$\begin{array}{ccc} X_1 \times_S X_2 & \xrightarrow{p_1} & X_1 \\ p_2 & & & \downarrow \\ X_2 & \xrightarrow{q_2} & S \end{array}$$

of schemes with a \mathbb{G}_m action, where all the arrows are \mathbb{G}_m -equivariant.

(c) The following diagram, where the horizontal arrows are induced by the natural transformations from (a) and (b) and the vertical ones are base change isomorphisms, is commutative

(d) When restricted to a subcategory of C_m-monodromic sheaves on X₁, the functor
 L is isomorphic to the identity functor.

Remark 2.1.3. It may be convenient for further arguments to think about the commutativity of the diagram (4) as follows. Consider the cube diagram below. Its front and back faces are identical Cartesian squares, and its arrows coming out of the plane of the paper are limit functors \mathbb{L} .



The proposition says that two natural transformations between red and dashed blue paths $D^b(X_1) \to D^b(X_2)$, one given by composition of natural transformations on the top, bottom and right faces of the cube, and another along left, bottom and front faces, coincide.

2.2 Character sheaves. Let X be a scheme with an action of an algebraic torus T, and let $a: T \times X \to X$, $p: T \times X \to X$ be the action and projection maps, respectively. Let \mathcal{L} be a rank 1 local system on T. We say that $\mathcal{F} \in D^b(X)$ is weakly \mathcal{L} -equivariant if there is an isomorphism $a^*\mathcal{F} \simeq \mathcal{L} \boxtimes \mathcal{F}$. We denote by $D^b_{\mathcal{L}}(X)$ the full idempotent-complete triangulated subcategory of $D^b(X)$ generated by weakly \mathcal{L} -equivariant complexes. We call the complexes in $D^b_{\mathcal{L}}(X)$ \mathcal{L} -monodromic complexes or monodromic complexes with generalized monodromy \mathcal{L} .

If $T = \mathbb{G}_m$ and \mathcal{L} is trivial, we will write $D^b_{mon}(X)$ for $D^b_{\mathcal{L}}(X)$ and refer to this category as a category of \mathbb{G}_m -monodromic complexes.

For a scheme X with an action of an algebraic group H which is normalized by the action of T, we will write $D^b_{\mathcal{L}}(H \setminus X)$ for the full triangulated subcategory of $D^b(H \setminus X)$ consisting of objects \mathcal{F} with For^H $\mathcal{F} \in D^b_{\mathcal{L}}(X)$.

We call a local system \mathcal{L} on T rational if $\alpha^* \mathcal{L}$ is trivial for some surjective homomorphism $\alpha: T \to T$.

Let $X = G \times G$, and consider actions of $G \times G$ on X by left and right multiplication. Consider a functor

$$\mathfrak{hc}_{!}\colon D^{b}(K\backslash X/K)\to D^{b}(K\backslash X/K'), \quad \mathfrak{hc}_{!}={}^{R}\mathrm{Av}_{K\cap K',!}^{K'}\circ {}^{R}\mathrm{For}_{K\cap K'}^{K},$$

where the superscript R on averaging and forgetful functor is used to emphasized that these are with respect to the groups acting on the right. The notation \mathfrak{hc} stands for Harish-Chandra, and the functor thus defined is commonly called the Harish-Chandra functor. Let

$$\chi \colon D^b(K \setminus X/K') \to D^b(K \setminus X/K), \quad \chi = {^R}\operatorname{Av}_{K \cap K',!}^K \circ {^R}\operatorname{For}_{K \cap K'}^{K'}$$

The torus T acts on $K \setminus X/K'$ on the right. Let \mathcal{L} be a rational rank 1 local system on T. The derived category of character sheaves with character \mathcal{L} , denoted by $D^b_{\mathfrak{C}_{\mathcal{L}}}(G)$, is the full triangulated idempotent-complete subcategory of $D^b(K \setminus X/K)$ generated by the essential image of $D^b_{\mathcal{L}}(K \setminus X/K')$ under the functor χ .

2.3 Main results. Consider the diagram

(5)
$$D^{b}(K \setminus X/K) \xrightarrow{\mathbb{L}_{\gamma}} D^{b}(K_{0} \setminus X/K)$$
$$Av_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \downarrow \qquad \qquad \downarrow Av_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}}$$
$$D^{b}(K' \setminus X/K) \xrightarrow{\mathbb{L}_{\gamma}} D^{b}(K' \setminus X/K)$$

Here $\alpha \colon \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \to \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}$ is the natural transformation given by Lemma 2.1.2.

We will prove the following

Theorem 2.3.1. When restricted to the category of character sheaves $D^b_{\mathfrak{C}_{\mathcal{L}}}(G)$, the transformation α is an isomorphism.

Corollary 2.3.2. After the restriction to the category of character sheaves $D^b_{\mathfrak{C}_{\mathcal{L}}}(G)$, there is an isomorphism of functors

$$\mathbb{L}_{\gamma} \simeq \operatorname{Av}_{B,*}^{K_0} \circ \operatorname{For}_B^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^K.$$

Proof. By Lemma 2.1.2 (d), the lower horizontal arrow in the diagram (5) is isomorphic to the identity functor when restricted to the monodromic subcategory. We obtain an isomorphism

$$\operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_0} \circ \mathbb{L}_{\gamma} \simeq \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}.$$

It is well-known that, restricted to the monodromic subcategory, the functor $\operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^{K_0}$ is an equivalence, with inverse given by $\operatorname{Av}_{B,*}^{K_0} \circ \operatorname{For}_B^{K'}$, and the Corollary follows. \Box

Remark 2.3.3. To match the notations of the introduction, note that

$$K \setminus X/K = G \setminus G \times G/G \simeq G/_{\mathrm{Ad}}G,$$

$$K' \setminus X/K = (B \times_T B) \setminus G \times G/G \simeq (U \setminus G/U)/_{\mathrm{Ad}}T,$$

$$K_0 \setminus X/K = (B \times_T B^-) \setminus G \times G/G \simeq (U^- \setminus G/U)/_{\mathrm{Ad}}T.$$

We will reduce Theorem 2.3.1 to a similar statement of the form "nearby cycles functor commutes with the averaging functor" with X/K replaced first by X/K' and then by $\mathcal{B} \times \mathcal{B}$, where $\mathcal{B} = G/B$ is the flag variety of G. Consider the following diagram, which is similar to diagram (5), with X/K replaced by X/K'.

Here α' : $\operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \to \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}$ is again the natural transformation given by Lemma 2.1.2.

We have the following

Theorem 2.3.4. When restricted to the monodromic category $D^b_{\mathcal{L}}(X/K')$, the transformation α' is an isomorphism.

Let $\mathcal{B} = G/B$ be the flag variety of G. Consider the following diagram, which is similar to diagram (5), with X/K' replaced by $\mathcal{B} \times \mathcal{B}$.

Here $\alpha'': \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \to \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}$ is again the natural transformation given by Lemma 2.1.2.

Theorem 2.3.5. The transformation α'' is an isomorphism.

The implication Theorem $2.3.4 \Rightarrow$ Theorem 2.3.1 will be deduced from the diagram chase using the standard properties of the nearby cycles functor.

For unipotent monodromy, Theorem 2.3.5 implies the Theorem 2.3.4 in a similar way.

Proof of Theorem 2.3.4 and a simpler Theorem 2.3.5 will use properties of the hyperbolic localization functor, which we proceed to explain.

3 Hyperbolic localization.

3.1 Braden's theorem. We recall some notions from [DG14]. Let X be a scheme and $a: \mathbb{G}_m \times X \to X$ be a \mathbb{G}_m -action. Consider the scheme of fixed points $X^{\mathbb{G}_m}$. Let by X^+ and X^- the attracting and the repelling schemes respectively. Consider the diagram



where i_{\pm} is the inclusion and p_{\pm} is taking the limit. Let us define the functors of hyperbolic restriction

(8)
$$\mathbb{H}^+ = (p_+)_! \circ i_+^* \qquad \mathbb{H}^- = (p_-)_* \circ i_-^!$$

Theorem 3.1.1 ([Bra03], [DG14]). There is a natural transformation $\mathbb{H}^- \to \mathbb{H}^+$ which is an isomorphism on \mathbb{G}_m -monodromic complexes.

Denote by \mathbb{H} the restriction of \mathbb{H}^{\pm} to the \mathbb{G}_m -monodromic subcategory.

3.2 Hyperbolic localization commutes with nearby cycles. For a scheme \widetilde{X} and a map $f: \widetilde{X} \to \mathbb{A}^1$ one has the nearby cycles functor ψ_f as in Subsection 2.1. Let $a': \mathbb{G}_m \times \widetilde{X} \to \widetilde{X}$ be an action which commutes with the map f for the trivial action on \mathbb{A}^1 . Consider the fixed points of the action and the map $f_0: \widetilde{X}^{\mathbb{G}_m} \to \mathbb{A}^1$. The natural transformations from Lemma 2.1.2 yield natural transformations

(9)
$$\mathbb{H}^+ \circ \psi_f \to \psi_{f_0} \circ \mathbb{H}^+ \qquad \psi_{f_0} \circ \mathbb{H}^- \to \mathbb{H}^- \circ \psi_f$$

Theorem 3.2.1. ([Nak16, Proposition 5.4.1 (2)], [Ric19, Theorem 3.3] Restricted to the category of \mathbb{G}_m -monodromic sheaves $D^b_{mon}(\tilde{X}^{\mathbb{G}_m} \times_{\mathbb{A}^1} \mathbb{G}_m)$ the natural transformations (9) is an isomorphism $\mathbb{H} \circ \psi_f \simeq \psi_{f_0} \circ \mathbb{H}$.

3.3 Hyperbolic localization commutes with Radon transform. Case of the flag variety. Consider the \mathbb{G}_m -action on $\mathcal{B} \times \mathcal{B}$ given by a cocharacter (λ, λ) for a dominant λ . This gives us the functor

$$\mathbb{H}: D^b_{mon}(\mathcal{B} \times \mathcal{B}) \to D^b(\mathcal{B}^{\mathbb{G}_m} \times \mathcal{B}^{\mathbb{G}_m})$$

Moreover, consider U acting on the right factor of $\mathcal{B} \times \mathcal{B}$. This gives us functors

$$For^{U}: D^{b}_{mon}(\mathcal{B} \times U \backslash \mathcal{B}) \to D^{b}_{mon}(\mathcal{B} \times \mathcal{B})$$
$$Av^{U}_{!}: D^{b}_{mon}(\mathcal{B} \times \mathcal{B}) \to D^{b}_{mon}(\mathcal{B} \times U \backslash \mathcal{B})$$

The following lemma is straightforward, see [CGY19, Sect. 1.4.5].

Lemma 3.3.1. Counit map induces a natural isomorphism

 $\mathbb{H}\simeq\mathbb{H}\circ For^U\circ Av^U_!$

As a corollary we obtain the following proposition

Proposition 3.3.2. (a) The unit map $\operatorname{Id} \to \operatorname{For}_B^{K'} \circ \operatorname{Av}_{B,!}^{K'}$ induces an isomorphism of functors making the following diagram commutative:

$$D^{b}(K \backslash \mathcal{B} \times \mathcal{B}) \xrightarrow{\operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}} D^{b}(K' \backslash \mathcal{B} \times \mathcal{B})$$

$$\xrightarrow{\mathbb{H} \circ \operatorname{For}^{K}} D^{b}(\mathcal{B}^{\mathbb{G}_{m}} \times \mathcal{B}^{\mathbb{G}_{m}})$$

(b) The same holds for K replaced by K_0 .

Proof. Using basechange for the following Cartesian diagram

$$\begin{array}{ccc} \mathcal{B} \times \mathcal{B} & \longrightarrow & B \backslash \mathcal{B} \times \mathcal{B} \\ & & \downarrow & & \downarrow \\ \mathcal{B} \times U \backslash \mathcal{B} & \longrightarrow & K' \backslash \mathcal{B} \times \mathcal{B} \end{array}$$

we get

$$\begin{split} \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^K &\simeq \operatorname{For}^U \circ \operatorname{For}_U^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^K &\simeq \\ & \operatorname{For}^U \circ \operatorname{Av}_!^U \circ \operatorname{For}^B \circ \operatorname{For}_B^K &\cong \operatorname{For}^U \circ \operatorname{Av}_!^U \circ \operatorname{For}^K \end{split}$$

The sheaves in the essential image of the For^K are \mathbb{G}_m -monodromic, which allows us to use Lemma 3.3.1. This finishes the proof of (a). The case of K_0 is similar.

3.4 Hyperbolic localization commutes with Radon transform. Case of the base affine space. Consider the \mathbb{G}_m -action on X/K' given by a cocharacter

$$\begin{aligned} \mathbf{a}_{w_1,w_2} \colon \mathbb{G}_m \times X/K' \to X/K' \\ \mathbf{a}_{w_1,w_2}\left(t, (g_1, g_2)\right) &= \left(\lambda(t) \cdot g_1 \cdot w_1^{-1}\left(\lambda(t^{-1})\right), \lambda(t) \cdot g_2 \cdot w_2^{-1}\left(\lambda\left(t^{-1}\right)\right)\right)\end{aligned}$$

for a dominant λ . This gives us the functors of hyperbolic localization

$$\mathbb{H}_{w_1,w_2}\colon D^b_{mon}(X/K')\to D^b(X/K')^{\mathbb{G}_m}.$$

For the rest of this subsection we fix w_1, w_2 and shorten $\mathbb{H} := \mathbb{H}_{w_1, w_2}$.

Note that $(X/K')^{\mathbb{G}_m} = (w_1 B \times w_2 B)/K'$. Moreover, consider U acting on the right factor of $X/K' = G \times G/K'$. This gives us functors

$$For^{U}: D^{b}_{mon}(U \setminus X/K') \to D^{b}_{mon}(X/K')$$
$$Av^{U}_{!}: D^{b}_{mon}(X/K') \to D^{b}_{mon}(U \setminus X/K')$$

The following lemma is straightforward generalization of [CGY19, Sect. 1.4.5].

Lemma 3.4.1. The unit map induces a functor isomorphism

$$\mathbb{H} \simeq \mathbb{H} \circ For^U \circ Av_1^U$$

Proof. The map p_+ factors

$$(X/K')^+ \to U \setminus (X/K')^+ \xrightarrow{\tilde{p}} (X/K')^{\mathbb{G}_m}$$

Hence

$$(p_+)_! \circ For^U \circ Av^U_! = (\tilde{p}_+)_! \circ Av^U_! \circ For^U \circ Av^U_! \simeq (\tilde{p}_+)_! \circ Av^U_! = (p_+)_! \cdot Av^U_! =$$

Then we have the following chain of isomorphisms

(10)
$$\mathbb{H} \circ For^U \circ Av^U_! = (p_+)_! \circ i^*_+ \circ For^U \circ Av^U_! \simeq$$
$$(p_+)_! \circ For^U \circ Av^U_! \circ i^*_+ \simeq (p_+)_! \circ i^*_+ = \mathbb{H},$$

where the second isomorphism is the base change and the third isomorphism is explained above. $\hfill \Box$

As a corollary we obtain the following proposition

Proposition 3.4.2. (a) The unit map $\operatorname{Id} \to \operatorname{For}_B^{K'} \circ \operatorname{Av}_{B,!}^{K'}$ induces an isomorphism of functors for the following diagram

$$D^{b}(K\backslash X/K') \xrightarrow{\operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}} D^{b}(K'\backslash X/K')$$

$$\xrightarrow{\mathbb{H} \circ \operatorname{For}^{K}} D^{b}(X/K')^{\mathbb{G}_{m}} \xrightarrow{\mathbb{H} \circ \operatorname{For}^{K'}} D^{b}(X/K')$$

(b) The same holds for K replaced by K_0 .

Proof. Using basechange for the following Cartesian diagram

$$\begin{array}{ccc} X/K' & \longrightarrow & B \backslash X/K' \\ & & \downarrow \\ U \backslash X/K' & \longrightarrow & K' \backslash X/K' \end{array}$$

we get

$$\begin{split} \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \operatorname{For}^{U} \circ \operatorname{For}_{U}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \\ \operatorname{For}^{U} \circ \operatorname{Av}_{!}^{U} \circ \operatorname{For}^{B} \circ \operatorname{For}_{B}^{K} \cong \operatorname{For}^{U} \circ \operatorname{Av}_{!}^{U} \circ \operatorname{For}^{K} \end{split}$$

The sheaves in the essential image of the For^K are \mathbb{G}_m -monodromic on the attracting locus, which allows us to use Lemma 3.4.1. This finishes the proof of (a). The case of K_0 is identical.

Informally, one can refer to the proposition above as hyperbolic restriction commutes with Radon transform $R_! = \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^{K_0}$.

4 Proof of the main results.

4.1 Proof for the flag variety. In this subsection we give a proof of Theorem 2.3.5. The following Lemma is clear.

Lemma 4.1.1. The functor

$$\mathbb{H} \circ \operatorname{For}^{K'} \colon D^b(K' \setminus \mathcal{B} \times \mathcal{B}) \to D^b(\mathcal{B}^{\mathbb{G}_m} \times \mathcal{B}^{\mathbb{G}_m})$$

 $is \ conservative.$

Using the lemma above it is enough to check that

(11)
$$\mathbb{H} \circ \operatorname{For}^{K'}(\alpha'') \colon \mathbb{H} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \to \\ \mathbb{H} \circ \operatorname{For}^{K'} \circ \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}$$

is an isomorphism. We have the following chain of isomorphisms

(12)
$$\mathbb{H} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \simeq \mathbb{H} \circ \operatorname{For}^{K_{0}} \circ \mathbb{L}_{\gamma} \simeq$$
$$\mathbb{H} \circ \mathbb{L}_{\gamma} \circ \operatorname{For}^{K} \simeq \mathbb{L}_{\gamma} \circ \mathbb{H} \circ \operatorname{For}^{K} \simeq \mathbb{H} \circ \operatorname{For}^{K}$$

The first map is inverse of the unit transformation, see Proposition 3.3.2(b). The second isomorphism is given by Lemma 2.1.2. The third isomorphism is Theorem 3.2.1. The last isomorphism comes from the fact that family is $\mathcal{B}^{\mathbb{G}_m} \times \mathcal{B}^{\mathbb{G}_m}$ is trivial.

Analogously, we have the following chain of isomorphism

(13)
$$\mathbb{H} \circ \operatorname{For}^{K'} \circ \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \mathbb{H} \circ \mathbb{L}_{\gamma} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq$$
$$\mathbb{L}_{\gamma} \circ \mathbb{H} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \mathbb{L}_{\gamma} \circ \mathbb{H} \circ \operatorname{For}^{K} \simeq \mathbb{H} \circ \operatorname{For}^{K} .$$

The first isomorphism comes from Lemma 2.1.2. The second isomorphism is Theorem 3.2.1. The third map is inverse of the unit transformation, see Proposition 3.3.2(a). The last isomorphism comes from the fact that family is $\mathcal{B}^{\mathbb{G}_m} \times \mathcal{B}^{\mathbb{G}_m}$ is trivial.

A diagram chase along the base change transformations implies that the natural transformations (11), (12), and (13) form a commutative triangle.

Remark 4.1.2. Consider the diagonal embedding $\Delta : K \setminus \mathcal{B} \to K \setminus \mathcal{B} \times \mathcal{B}$. The localization of the object $\mathbb{L}_{\gamma}(\Delta_* \underline{\Bbbk}_{K \setminus \mathcal{B}})$ in the setting of *D*-modules is known as a semiregular module, see [Don08]. For $w \in W$ let $\mathcal{O}_w = UwB \subset \mathcal{B}, \mathcal{O}_w^- = U^-wB \subset \mathcal{B}$ be the Bruhat cells, and let $j_w : \mathcal{O}_w \to \mathcal{B}, j'_w : \mathcal{O}_w^- \to \mathcal{B}$ be the corresponding locally-closed embeddings. Let $\Delta_w = j_w!\underline{\Bbbk}_{\mathcal{O}_w}[\dim \mathcal{O}_w], \nabla_w^- = j'_{w*}\underline{\Bbbk}_{\mathcal{O}_w^-}[\dim \mathcal{O}_w^-]$. Using Lemma 3.2.1 it is easy to see that $\operatorname{For}^{K_0} \mathbb{L}_{\gamma}(\Delta_* \underline{\Bbbk}_{K \setminus \mathcal{B}})[\dim K_0]$ admits a filtration with associated graded pieces given by $\Delta_{w_0w^{-1}} \boxtimes \nabla_w, w \in W$. After localization, this recovers [Don08, Theorem 11.14]. See also [Che23] for a related discussion in the affine setting.

4.2 Proof for monodromic sheaves. In this subsection we give a proof of Theorem 2.3.4.

Denote $\mathbb{H}_{\bullet} = \bigoplus_{w_1, w_2} \mathbb{H}_{w_1, w_2}$.

Lemma 4.2.1. The functor

$$\mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \colon D^{b}(K' \setminus X/K') \to \bigoplus_{w_{1},w_{2}} D^{b}(w_{1}B \times w_{2}B/K')$$

is conservative.

Proof. Let $\mathcal{F}' \neq 0$ be a complex in $D^b(K' \setminus X/K')$, and let $\mathcal{F} = \operatorname{For}^{K'} \mathcal{F}'$. Assume that $\mathbb{H}_{\bullet} \mathcal{F} = 0$.

For $(w_1, w_2) \in W^2$, write $\mathcal{O}_{w_1, w_2} = w_1 B \times w_2 B/K'$, and let $j_{w_1, w_2} : \mathcal{O}_{w_1, w_2} \to X/K'$ be the embedding. Pick a pair $(w_1, w_2) \in W$ such that \mathcal{O}_{w_1, w_2} is open in the support of \mathcal{F} . Let

(14)
$$j_{w_1,w_2} j_{w_1,w_2}^* \mathcal{F} \to \mathcal{F} \to \mathcal{Z}$$

be the standard triangle associated to the natural transformation

$$j_{w_1,w_2!}j_{w_1,w_2}^* \to \mathrm{Id}_{D^b(X/K')}.$$

It is easy to see that $\mathbb{H}_{v_1,v_2}(j_{w_1,w_2!}\mathcal{G}) \simeq 0$ unless $(v_1,v_2) = (w_1,w_2)$ for any $\mathcal{G} \in D^b(\mathcal{O}_{w_1,w_2})$. Applying \mathbb{H}_{\bullet} to the triangle (14) and using this consideration, we get that

$$\mathbb{H}_{w_1, w_2} j_{w_1, w_2!} j_{w_1, w_2}^* \mathcal{F} = 0,$$

which is easily seen to imply $j_{w_1,w_2}^* \mathcal{F} \simeq 0$, since this complex is smooth along the left K'-orbits. This contradicts our assumptions on the support of \mathcal{F} .

Using the lemma above it is enough to check that

(15)
$$\mathbb{H}_{\bullet} \circ \operatorname{For}^{K'}(\alpha') \colon \mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \to$$
$$\mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \circ \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K}$$

is an isomorphism. We have the following chain of isomorphisms

(16)
$$\mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K_{0}} \circ \mathbb{L}_{\gamma} \simeq \mathbb{H}_{\bullet} \circ \operatorname{For}^{K_{0}} \circ \mathbb{L}_{\gamma} \simeq \mathbb{H}_{\bullet} \circ \operatorname{For}^{K} \simeq \mathbb{L}_{\gamma} \circ \mathbb{H}_{\bullet} \circ \operatorname{For}^{K}.$$

The first map is inverse of the unit transformation, see Proposition 3.4.2(b). The second isomorphism is given by Lemma 2.1.2. The third isomorphism is Theorem 3.2.1.

Analogously, we have the following chain of isomorphism

(17)
$$\mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \circ \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \mathbb{H}_{\bullet} \circ \mathbb{L}_{\gamma} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq$$
$$\mathbb{L}_{\gamma} \circ \mathbb{H}_{\bullet} \circ \operatorname{For}^{K'} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_{B}^{K} \simeq \mathbb{L}_{\gamma} \circ \mathbb{H}_{\bullet} \circ \operatorname{For}^{K}.$$

The first isomorphism comes from Lemma 2.1.2. The second isomorphism is Theorem 3.2.1. Note that the sheaves in the essential image of the For^{K'} are \mathbb{G}_m -monodromic on the attracting locus, which makes the theorem applicable. The third map is inverse of the unit transformation, see Proposition 3.4.2(a).

Diagram chase along the base change transformations implies that the natural transformations (15), (16), and (17) form a commutative triangle.

4.3 From monodromic sheaves to character sheaves. Here we prove the implication Theorem $2.3.4 \Rightarrow$ Theorem 2.3.1. The following Lemma is clear.

Lemma 4.3.1. Let $\Phi, \Psi: \mathcal{C} \to \mathcal{D}$ be two exact functors between triangulated categories \mathcal{C}, \mathcal{D} and let $\alpha: \Phi \to \Psi$ be a natural transformation. Assume that for any object $c \in \mathcal{C}$, there is an object c' containing c as a direct summand, such that $\alpha_{c'}: \Phi(c') \to \Psi(c')$ is an isomorphism. Then α is an isomorphism.

By Lemma 4.3.1 it is enough to show that the natural transformation

$$\alpha \circ \chi \colon \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^K \circ \mathbb{L}_{\gamma} \circ \chi \to \mathbb{L}_{\gamma} \circ \operatorname{Av}_{B,!}^{K'} \circ \operatorname{For}_B^K \circ \chi$$

is an isomorphism. To this end, consider the following cube diagram, where in each cardinal plane parallel arrows have the same name.



Iterating Lemma 2.1.2 (c) we get that the two natural transformations between red and dashed blue paths $D^b(K \setminus X/K') \to D^b(K' \setminus X/K)$, one given by composition of natural transformations on the top, back and right faces of the cube, and another along left, bottom and front faces, coincide. Note that the natural transformation corresponding to the right face is exactly the natural transformation $\alpha \circ \chi$. Thus, it is enough to show that natural transformations corresponding to the other five faces are isomorphisms.

Natural transformation corresponding to the left face is an isomorphism by Theorem 2.3.4. The back and front faces are compositions of base change isomorphisms. The top and bottom face are isomorphisms by Lemma 2.1.2 (a) and (b).

References

- [BFO12] Roman Bezrukavnikov, Michael Finkelberg, and Victor Ostrik. Character D-modules via Drinfeld center of Harish-Chandra bimodules. *Inventiones* mathematicae, 188(3):589–620, June 2012.
- [BITV23] Roman Bezrukavnikov, Andrei Ionov, Kostiantyn Tolmachov, and Yakov Varshavsky. Equivariant derived category of a reductive group as a categorical center. arXiv preprint arXiv:2305.02980, 2023.
- [BL06] Joseph Bernstein and Valery Lunts. *Equivariant Sheaves and Functors*. Springer, 2006.
- [BN09] David Ben-Zvi and David Nadler. The character theory of a complex group. arXiv preprint arXiv:0904.1247, 2009.

- [Bra03] Tom Braden. Hyperbolic localization of intersection cohomology. *Transform. Groups*, 8(3):209–216, 2003.
- [CGY19] T.-H. Chen, D. Gaitsgory, and A. Yom Din. On the Casselman-Jacquet functor. 101:73–112, 2019.
- [Che23] Lin Chen. Nearby cycles on Drinfeld-Gaitsgory-Vinberg interpolation grassmannian and long intertwining functor. *Duke Math. J.*, 172(3):447–543, 2023.
- [CY17] Tsao-Hsien Chen and Alexander Yom Din. A formula for the geometric Jacquet functor and its character sheaf analogue. *Geometric and Functional* Analysis, 27(4):772–797, 2017.
- [DG14] Vladimir Drinfeld and Denis Gaitsgory. On a theorem of Braden. Transformation Groups, 19(2):313–358, June 2014.
- [Don08] Dmitry Donin. Lie algebras of differential operators and D-modules. PhD thesis, University of Toronto, 2008.
- [Gin89] Victor Ginzburg. Admissible modules on a symmetric space. Astérisque, (173-74):199–255, 1989.
- [HL23] Quoc P. Ho and Penghui Li. Graded character sheaves, HOMFLY-PT homology, and Hilbert schemes of points on C². arXiv preprint arXiv:2305.01306, 2023.
- [LO08] Yves Laszlo and Martin Olsson. The six operations for sheaves on Artin stacks II: Adic coefficients. Publications Mathématiques de l'IHÉS, 107:169– 210, 2008.
- [Lus] George Lusztig. Truncated convolution of character sheaves. Bulletin of the Institute of Mathematics Academia Sinica, 10(1):1–72.
- [Lus85] George Lusztig. Character sheaves I. Advances in Mathematics, 56(3):193– 237, 1985.
- [Lus15] George Lusztig. Non-unipotent character sheaves as a categorical centre. arXiv preprint arXiv:1506.04598, 2015.
- [Nak16] Hiraku Nakajima. Lectures on perverse sheaves on instanton moduli spaces. arXiv preprint arXiv:1604.06316, 2016.
- [Ric19] Timo Richarz. Spaces with \mathbb{G}_m -action, hyperbolic localization and nearby cycles. Journal of Algebraic Geometry, 28:251–289, 2019.
- [SGA06] Groupes de Monodromie en Geometrie Algebrique: Seminaire de Geometrie Algebrique du Bois-Marie 1967-1969 (SGA 7 II), Pierre Deligne, Nicholas Katz, volume 340. Springer, 2006.

R. Gonin, School of Mathematics, University of Cardiff, Cardiff, United Kingdom

E-mail address: goninr@cardiff.ac.uk

A. Ionov, Department of Mathematics, Boston College, Chestnut Hill, Massachusetts, USA

E-mail address: ionov@bc.edu

K. Tolmachov, Department of Mathematics, University of Hamburg, Hamburg, Germany

E-mail address: tolmak@khtos.com