

# Orthogonal Oscillator Representations, Laplace Equations and Intersections of Determinantal Varieties <sup>1</sup>

Hengjia Zhang<sup>2</sup> and Xiaoping Xu<sup>3</sup>

HLM, Institute of Mathematics, Academy of Mathematics & System Sciences  
Chinese Academy of Sciences, Beijing 100190, P.R. China  
& School of Mathematics, University of Chinese Academy of Sciences,  
Beijing 100049, P.R. China

## Abstract

Associated varieties are geometric objects appearing in infinite-dimensional representations of semisimple Lie algebras (groups). By applying Fourier transformations to the natural orthogonal oscillator representations of special linear Lie algebras, Luo and the second author (2013) obtained a big family of infinite-dimensional irreducible representations of the algebras on certain spaces of homogeneous solutions of the Laplace equation. In this paper, we prove that the associated varieties of these irreducible representations are the intersections of explicitly given determinantal varieties. This provides an explicit connection among representation theory, partial differential equations and algebraic geometry.

*Keywords:* special linear Lie algebra; orthogonal oscillator representation; associated variety; irreducible module; filtrations; determinantal variety.

## 1 Introduction

In 1971, Bernstein [11] defined a variety associated with a filtrated module of the algebra of differential operators, in order to construct the fundamental solution of a linear partial differential equation with constant coefficients. A primitive ideal of the universal enveloping algebra of a semisimple Lie algebra is the kernel of an irreducible representation of the Lie algebra. Borho and Jantzen [13] (1977) found a translation principle among primitive ideals and used it to prove the Dixmier orbit map is well defined for  $sl(n)$ . Vogan [33] (1978) found particularly refined techniques in the description of the leading coefficient in the Hilbert-Samuel polynomial of a Harish-Chandra module for a semisimple Lie algebra and its maximal compact subalgebra, where the filtration of the module is naturally induced from the canonical filtration of the universal enveloping algebra of the Lie algebra. Moreover, he [34] (1980) derived a new order relation in primitive ideals and used it to prove a Joseph's conjecture on primitive ideals and multiplicities in the composition series of principal series representations.

Borho and Brylinsky [12] (1982) proved that the kernel of the representation of the universal enveloping algebra of a semisimple Lie algebra by global differential operators on a complete homogeneous space of the corresponding Lie group is the annihilator of a generalized Verma module, and the associated variety (the term firstly appeared) of

---

<sup>1</sup>2010 Mathematical Subject Classification. Primary 17B10, 22E47; Secondary 14M12, 35C05.

<sup>2</sup>Corresponding author.

<sup>3</sup>Research supported by National Key R&D Program of China 2020YFA0712600.

the ideal is irreducible. Joseph [22] (1983) gave an extensive review on the classification of primitive ideals in the universal enveloping algebra of a semisimple Lie algebra. He [23] (1984) continued his investigation on associated varieties aiming for a way to compute them and to relate it more naturally to the nilpotent orbits and the corresponding Springer representations of the Weyl group. Moreover, he [24] (1985) proved that the associated varieties are the Zariski closures of nilpotent orbits. Matumote [27] (1987) proved a necessary condition for the existence of certain Whittaker vectors in terms of associated varieties. Vogan [35] (1991) proved that the associated varieties of  $(G, K)$ -modules are the Zariski closures of union of  $K$ -nilpotent orbits. Melnikov [29] (1993) conditionally proved that the associated variety of an irreducible highest weight  $sl(n)$ -module is irreducible and Williamson [36] (1995) found a counter example. Barbasch and Bozicevic [6] (1999) derived a formula for the associated varieties of induced modules, which is analogous to the one for wave front set of a derived functor module obtained by Barbasch and Vogan in 1980. There are also many other interesting significant works related to associated varieties (e.g., cf. [7, 8, 24, 39, 40, 21, 41, 28, 19, 25, 30, 32, 31, 1, 5, 3, 4]). It seems desirable to know if there exist geometrically sophisticated explicit known algebraic varieties that are associated varieties.

Determinantal varieties are natural important geometric objects in algebraic geometry. Barshay [9] (1973) used them to prove the coordinate ring of the Veronese is arithmetically normal and Cohen-Macaulay. Catanese [14] (1981) used symmetric determinantal varieties to prove Babbage's conjecture. de Concini, Eisenbud and Procesi [15] (1980) found an important connection of Young diagrams with determinantal varieties. Griffiths [18] (1983) found that determinantal varieties play important roles in the infinitesimal variations of Hodge structure. Enright and Hunziker [16] (2004) gave explicit formulas for the numerator polynomials of the Hilbert series associated with the closures of the nilpotent orbits corresponding to Wallach representations through determinantal varieties. Moreover, Enright, Hunziker and Prueett [17] (2014) presented a natural generalization of Young diagrams for Hermitian symmetric spaces and used it to study syzygies of determinantal varieties. Bernardara, Bolognesi and Faenzi [10] (2016) derived a homological projective duality for determinantal varieties. Wong [37] (2017) gave an explicit calculation of the Euler characteristic of the spaces of classical vacua in terms of the Euler characteristics of suitably chosen determinantal subvarieties. Homma and Manabe [20] (2018) showed that determinantal Calabi-Yau varieties in Grassmannians are closely related to Givental I-functions.

By applying Fourier transformations to the natural orthogonal oscillator representations of special linear Lie algebras, Luo and the second author [26] (2013) obtained a big family of infinite-dimensional irreducible representations of the algebras on certain spaces of homogeneous solutions of the Laplace equation. Bai [2] (2015) found the Gel'fand-Kirillov dimensions of these representations. In this paper, we prove that the associated varieties of these irreducible representations are intersections of explicitly given determinantal varieties. This provides an explicit connection among representation theory, partial differential equations and algebraic geometry. It also partially meets Bernstein's original concern in [11]. Below we give a more detailed technical introduction to our work.

Let  $\mathbb{F}$  be a field with characteristic 0, and let  $n > 1$  be an integer. Denote by  $\mathcal{A} = \mathbb{F}[x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n]$  the polynomial algebra in  $2n$  variables. Let  $E_{r,s}$  be the square

matrix with 1 as its  $(r, s)$ -entry and 0 as the others. The special linear Lie algebra

$$sl(n) = \sum_{1 \leq i < j \leq n} (\mathbb{F}E_{i,j} + \mathbb{F}E_{j,i}) + \sum_{r=1}^{n-1} \mathbb{F}(E_{r,r} - E_{r+1,r+1}) \quad (1.1)$$

with the commutator as its Lie bracket. For any two integers  $p \leq q$ , we denote  $\overline{p, q} = \{p, p+1, \dots, q\}$ . The natural orthogonal oscillator representation  $\pi$  of  $sl(n)$  on  $\mathcal{A}$  is given as follows:

$$\pi(E_{i,j}) = x_i \partial_{x_j} - y_j \partial_{y_i} \quad \text{for } i, j \in \overline{1, n}. \quad (1.2)$$

With respect to the above representation, the Laplace operator

$$\Delta = \sum_{i=1}^n \partial_{x_i} \partial_{y_i}. \quad (1.3)$$

Denote by  $\mathbb{N}$  the set of nonnegative integers. For  $\ell_1, \ell_2 \in \mathbb{N}$ , we denote by  $\mathcal{A}_{\ell_1, \ell_2}$  the subspace of homogeneous polynomials with the degree  $\ell_1$  in  $\{x_1, x_2, \dots, x_n\}$  and the degree  $\ell_2$  in  $\{y_1, y_2, \dots, y_n\}$ . Then  $\mathcal{A} = \bigoplus_{\ell_1, \ell_2 \in \mathbb{N}} \mathcal{A}_{\ell_1, \ell_2}$  is an  $\mathbb{N}^2$ -graded algebra.

Set the subspace of homogeneous harmonic polynomials as follows:

$$\mathcal{H}_{\ell_1, \ell_2} = \{f \in \mathcal{B}_{\ell_1, \ell_2} \mid \Delta(f) = 0\}. \quad (1.4)$$

Then  $\mathcal{H}_{\ell_1, \ell_2}$  forms an irreducible  $sl(n)$ -module with highest weight  $\ell_1 \lambda_1 + \ell_2 \lambda_{n-1}$  (e.g., cf. [38]), where  $\lambda_i$  denotes the  $i$ th fundamental weight of  $sl(n)$ .

Denote by  $\mathcal{F}_z$  the Fourier transformation on  $z$ . Fix  $n_1, n_2 \in \overline{1, n}$  with  $n_1 \leq n_2$ . Letting (1.2) act on the space

$$\mathcal{A}' = \mathcal{F}_{x_1} \cdots \mathcal{F}_{x_{n_1}} \mathcal{F}_{x_{n_2+1}} \cdots \mathcal{F}_{x_n}(\mathcal{A}) \quad (1.5)$$

of distributions, one can obtain a big family of infinite-dimensional irreducible representations of  $sl(n)$  on certain spaces of homogeneous solutions of the Laplace equation. However, this picture is not convenient to be dealt with. Instead, we consider another equivalent representation as follows. In the representation formulas given in (1.2), applying the Fourier transformations on operators:

$$\partial_{x_r} \mapsto -x_r, \quad x_r \mapsto \partial_{x_r}, \quad \partial_{y_s} \mapsto -y_s, \quad y_s \mapsto \partial_{y_s} \quad \text{for } r \in \overline{1, n_1}, \quad s \in \overline{n_2+1, n}, \quad (1.6)$$

we obtain a new representation  $\tilde{\pi}$  on  $\mathcal{A}$ . The representations  $(\pi, \mathcal{A}')$  and  $(\tilde{\pi}, \mathcal{A})$  are equivalent (or isomorphic). Under the changes in (1.6), the Laplace operator becomes

$$\tilde{\Delta} = \sum_{i=1}^{n_1} x_i \partial_{y_i} - \sum_{r=n_1+1}^{n_2} \partial_{x_r} \partial_{y_r} + \sum_{s=n_2+1}^n y_s \partial_{x_s}. \quad (1.7)$$

Take the degree of  $\{x_1, \dots, x_{n_1}, y_{n_2+1}, \dots, y_n\}$  as  $-1$  and that of  $\{x_{n_1+1}, \dots, x_n, y_1, \dots, y_{n_2}\}$  as  $1$ . Denote by  $\mathbb{Z}$  the ring of integers. For  $\ell_1, \ell_2 \in \mathbb{Z}$ , we denote by  $\mathcal{A}_{(\ell_1, \ell_2)}$  the subspace of homogeneous polynomials with the degree  $\ell_1$  in  $\{x_1, x_2, \dots, x_n\}$  and the degree  $\ell_2$  in  $\{y_1, y_2, \dots, y_n\}$ . Then  $\mathcal{A} = \bigoplus_{\ell_1, \ell_2 \in \mathbb{Z}} \mathcal{A}_{(\ell_1, \ell_2)}$  is a  $\mathbb{Z}^2$ -graded algebra. Set

$$\mathcal{H}_{(\ell_1, \ell_2)} = \{f \in \mathcal{A}_{(\ell_1, \ell_2)} \mid \tilde{\Delta}(f) = 0\}. \quad (1.8)$$

The following is a result from [26]:

**Proposition 1** For  $\ell_1, \ell_2 \in \mathbb{Z}$ , the necessary and sufficient condition for  $\mathcal{H}_{(\ell_1, \ell_2)}$  to be an infinite-dimensional irreducible  $sl(n)$ -module is as follows:

(1) When  $n_1 + 1 < n_2$ , (a)  $\ell_1 + \ell_2 \leq n_1 - n_2 + 1$ ; (b)  $n_2 = n$ ,  $\ell_1 \in \mathbb{N}$  and  $\ell_2 = 0$ ; (c)  $n_2 = n$ ,  $\ell_2 \in \mathbb{N}$  and  $\ell_1 \geq n_1 - n + 2$ ;

(2) If  $n_1 + 1 = n_2$ ,  $\ell_1 + \ell_2 \leq 0$  or  $n_2 = n$  and  $0 \leq \ell_2 \leq \ell_1$ ;

(3) In the case  $n_1 = n_2$ ,  $\ell_1 + \ell_2 \leq 0$  and: (a)  $\ell_2 \leq 0$ ,  $n_1 < n - 1$  and  $n \geq 3$ ; (b)  $\ell_1 \leq 0$ ,  $1 < n_1 < n$  and  $n \geq 3$ ; (c)  $\ell_1, \ell_2 \leq 0$ ,  $n_1 = 1$  and  $n = 2$ .

The above irreducible modules are infinite-dimensional highest weight modules with distinct highest weights. For instance, if  $n_1 + 1 < n_2 < n$  and  $m_1, m_2 \in \mathbb{N}$  with  $m_1 + m_2 \geq n_2 - n_1 - 1$ ,  $\mathcal{H}_{(-m_1, -m_2)}$  has a highest-weight vector  $x_{n_1}^{m_1} y_{n_2+1}^{m_2}$  of weight  $m_1 \lambda_{n_1-1} - (m_1 + 1) \lambda_{n_1} - (m_2 + 1) \lambda_{n_2} + m_2 (1 - \delta_{n_2, n-1}) \lambda_{n_2+1}$ .

Let  $\mathfrak{g}$  be a finite-dimensional semisimple Lie algebra and let  $U(\mathfrak{g})$  be the universal enveloping algebra of  $\mathfrak{g}$ . Denote  $U_0(\mathfrak{g}) = \mathbb{F}$  and for  $0 < m \in \mathbb{Z}$ ,  $U_m(\mathfrak{g}) = \mathbb{F} + \sum_{r=1}^m \mathfrak{g}^r$ . Then  $\{U_k(\mathfrak{g}) \mid k \in \mathbb{N}\}$  forms a filtration of  $U(\mathfrak{g})$ . Suppose that  $M$  is a finitely generated  $U(\mathfrak{g})$ -module and  $M_0$  is a finite-dimensional subspace of  $M$  satisfying

$$(U(\mathfrak{g}))(M_0) = M. \quad (1.9)$$

Setting  $M_k = (U_k(\mathfrak{g}))(M_0)$ , we get a filtration  $\{M_k \mid k \in \mathbb{N}\}$  of  $M$ .

The associated graded module is given by

$$\overline{M} = \text{gr}(M; M_0) = \bigoplus_{k=0}^{\infty} \overline{M}_k \quad \text{with} \quad \overline{M}_k = M_k / M_{k-1}, \quad (1.10)$$

and it is a finitely generated graded  $S(\mathfrak{g})$ -module, since graded algebra  $\text{gr}(U(\mathfrak{g}))$  is canonically isomorphic to the symmetric tensor algebra  $S(\mathfrak{g})$  by the PBW theorem. Here we identify  $S(\mathfrak{g})$  with  $\mathbb{F}[\mathfrak{g}^*]$ , the polynomial ring over  $\mathfrak{g}^*$ , in the canonical way through the Killing form of  $\mathfrak{g}$ .

The annihilator of  $\overline{M}$  in  $S(\mathfrak{g})$  is defined as

$$\text{Ann}_{S(\mathfrak{g})}(\overline{M}) = \{\xi \in S(\mathfrak{g}) \mid \xi(v) = 0, \forall v \in \overline{M}\}, \quad (1.11)$$

which is a graded ideal of  $S(\mathfrak{g})$ . It defines the associated variety in the dual space  $\mathfrak{g}^*$ :

$$\mathcal{V}(M) = \{f \in \mathfrak{g}^* \mid u(f) = 0, \forall u \in \text{Ann}_{S(\mathfrak{g})}(\overline{M})\}. \quad (1.12)$$

The variety does not depend on the choice of the subspace  $M_0$  satisfying (1.9).

Let  $S_1$  and  $S_2$  be subsets of  $\overline{1, n}$ . Denote the determinantal ideal  $I_t(S_1, S_2)$  to be the ideal generated by all the  $t$ -minors in the matrix  $(z_{i,j})_{i \in S_1, j \in S_2}$ , and  $\mathcal{V}_t(S_1, S_2) = \mathcal{V}(I_t)$  is called a *determinantal variety*. For convenience, we denote  $I_t(S_1, S_2) = 0$  and  $\mathcal{V}_t(S_1, S_2) = \mathbb{A}^{|S_1||S_2|}$ , when  $t > |S_1|$  or  $t > |S_2|$ .

Recall  $n_1, n_2 \in \overline{1, n}$  with  $n_1 \leq n_2$ . Set

$$J_1 = \overline{1, n_1}, \quad J_2 = \overline{n_1 + 1, n_2}, \quad J_3 = \overline{n_2 + 1, n}. \quad (1.13)$$

The following is our main theorem in this paper:

**Main Theorem** If  $n_1 \leq n_2$ ,  $\ell_1 \leq 0$  or  $\ell_2 \leq 0$ ,

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_3(J_2 \cup J_3, J_1 \cup J_2) \cap \mathcal{V}_2(J_2, J_1) \cap \mathcal{V}_2(J_3, J_2) \cap \mathcal{V}_1(J_2, J_2). \quad (1.14)$$

If  $n_1 < n_2 = n$  and  $\ell_1, \ell_2 > 0$ ,

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_2(J_2, J_1). \quad (1.15)$$

The theorem covers all the irreducible  $sl(n)$ -modules  $\mathcal{H}_{(\ell_1, \ell_2)}$  listed in Proposition 1. In particular, when  $n_1 = n_2$  and  $|J_1|, |J_3| \geq 3$ ,

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_3(J_3, J_1) \quad (1.16)$$

is exactly a determinantal variety. When  $n_2 = n$ , the projective version of the associated variety

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_2(J_2, J_1) \quad (1.17)$$

is the well-known Serge variety in algebraic geometry.

The paper is organized as follows. In Section 2, we explicitly describe the filtration of  $\mathcal{H}_{(\ell_1, \ell_2)}$  with respect to the representation  $\tilde{\pi}$  and a properly selected finite-dimensional subspace  $M_0$  by an adjusted degree  $\mathfrak{d}$  (cf. (2.46)-(2.48)). In Section 3, we study certain linear independence on certain subsets in the spanning subsets of filtration subspaces by introducing the notion of “3-chain”, which plays a key role in next section. Finally in Section 4, we present a proof of the main theorem by using certain “order” on the products of quadratic alternating polynomials.

## 2 Explicit Description of the Filtration for $\mathcal{H}_{(\ell_1, \ell_2)}$

In this section, we select a special subspace  $M_0$  of  $M = \mathcal{H}_{(\ell_1, \ell_2)}$  and determine a spanning subset for each  $M_k = (U_k(\mathfrak{g}))(M_0)$ . Moreover, a new degree function  $\mathfrak{d}$  on  $M$  is introduced in order to determine the structure of  $M_k$ .

In the rest of this paper, we only consider the representation  $\tilde{\pi}$  of  $sl(n)$  obtained under the Fourier transformations (1.6), whose representation formulas are given by

$$\tilde{\pi}(E_{i,j}) = E_{i,j}^x - E_{j,i}^y \quad \text{for } i, j \in \overline{1, n} \quad (2.1)$$

with

$$E_{i,j}^x = \begin{cases} -x_j \partial_{x_i} - \delta_{i,j} & \text{if } i, j \in \overline{1, n_1}; \\ \partial_{x_i} \partial_{x_j} & \text{if } i \in \overline{1, n_1}, j \in \overline{n_1 + 1, n}; \\ -x_i x_j & \text{if } i \in \overline{n_1 + 1, n}, j \in \overline{1, n_1}; \\ x_i \partial_{x_j} & \text{if } i, j \in \overline{n_1 + 1, n} \end{cases} \quad (2.2)$$

and

$$E_{i,j}^y = \begin{cases} y_i \partial_{y_j} & \text{if } i, j \in \overline{1, n_2}; \\ -y_i y_j & \text{if } i \in \overline{1, n_2}, j \in \overline{n_2 + 1, n}; \\ \partial_{y_i} \partial_{y_j} & \text{if } i \in \overline{n_2 + 1, n}, j \in \overline{1, n_2}; \\ -y_j \partial_{y_i} - \delta_{i,j} & \text{if } i, j \in \overline{n_2 + 1, n}. \end{cases} \quad (2.3)$$

Moreover, we adopt the notion

$$\xi(f) = \tilde{\pi}(\xi)(f) \quad \text{for } \xi \in sl(n), f \in \mathcal{A}. \quad (2.4)$$

Denote

$$x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}, \quad y^\alpha = y_1^{\alpha_1} y_2^{\alpha_2} \cdots y_n^{\alpha_n} \quad \text{for } \alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}^n. \quad (2.5)$$

In the rest of this section, we always assume  $n_1 < n_2$ . Define the linear operator  $T$  on  $\mathcal{A}$  by

$$T(x^\alpha y^\beta) = \sum_{i=0}^{\infty} \frac{(x_{n_1+1} y_{n_1+1})^i \left( \tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}} \right)^i}{\prod_{r=1}^i (\alpha_{n_1+1} + r) (\beta_{n_1+1} + r)} (x^\alpha y^\beta) \quad (2.6)$$

for  $\alpha, \beta \in \mathbb{N}^n$ . Then we have

$$T\tilde{\pi}(E_{i,j}) = \tilde{\pi}(E_{i,j})T \quad \text{for } n_1 + 1 \neq i, j \in \overline{1, n}. \quad (2.7)$$

Moreover, the set

$$\{T(x^\alpha y^\beta) \mid \alpha, \beta \in \mathbb{N}^n, \alpha_{n_1+1} \beta_{n_1+1} = 0, x^\alpha y^\beta \in \mathcal{A}_{\langle \ell_1, \ell_2 \rangle}\} \quad (2.8)$$

forms a basis of the module  $\mathcal{H}_{\langle \ell_1, \ell_2 \rangle}$ , as shown in [26]. Furthermore, we introduce the notion

$$\begin{aligned} & N \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \\ &= \left\{ x^\alpha y^\beta \in \mathcal{A}_{\langle \ell_1, \ell_2 \rangle} \mid \sum_{r=1}^{n_1} \alpha_r = k_{11}, \sum_{r=n_1+1}^{n_2} \alpha_r = k_{12}, \sum_{r=n_2+1}^n \alpha_r = k_{13}, \right. \\ & \quad \left. \sum_{r=1}^{n_1} \beta_r = k_{21}, \sum_{r=n_1+1}^{n_2} \beta_r = k_{22}, \sum_{r=n_2+1}^n \beta_r = k_{23}, \alpha_{n_1+1} \beta_{n_1+1} = 0 \right\} \quad (2.9) \end{aligned}$$

and

$$TN \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} = \text{Span} \left\{ T(f) \mid f \in N \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right\} \quad (2.10)$$

where  $k_{11}, k_{12}, k_{13}, k_{21}, k_{22}, k_{23} \in \mathbb{N}$ . Now we take

$$M_0 = \begin{cases} TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & 0 & -\ell_2 \end{pmatrix} & \text{if } \ell_1 \leq 0, \ell_2 \leq 0, \\ TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 & 0 \end{pmatrix} & \text{if } \ell_1 \leq 0, \ell_2 \geq 0, \\ TN \begin{pmatrix} 0 & \ell_1 & 0 \\ 0 & 0 & -\ell_2 \end{pmatrix} & \text{if } \ell_1 \geq 0, \ell_2 \leq 0, \end{cases} \quad (2.11)$$

which is a finite-dimensional subspace of  $M = \mathcal{H}_{\langle \ell_1, \ell_2 \rangle}$ . Set

$$M_k = U_k(\mathfrak{sl}(n))(M_0) \quad \text{for } k \in \mathbb{N}. \quad (2.12)$$

For convenience, we denote

$$X_S = \{x_i \mid i \in S\}, \quad Y_S = \{y_i \mid i \in S\} \quad \text{for } S \subset \overline{1, n}. \quad (2.13)$$

In order to express  $M_k$  in terms of the basis given in (2.8), we need the following conclusions.

**Lemma 2.1.** Fix  $k, k_{13}, k_{21} \in \mathbb{N}$ . For each  $j, i_{1,j} \in J_1, i_{3,j} \in J_3, v_0 \in \mathbb{F} [X_{J_1}, X_{J_2 \setminus \{n_1+1\}}, Y_{J_3}] \cup \mathbb{F} [X_{J_1}, Y_{J_2 \setminus \{n_1+1\}}, Y_{J_3}]$ , we have

$$\begin{aligned} & (-1)^k \frac{k!}{(k - k_{13})!} T \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^{k_{13}} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{k-k_{13}} \right) \\ &= E_{i_{3,k_{13}}, n_1+1} \cdots E_{i_{3,1}, n_1+1} E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k}} (v_0) \end{aligned} \quad (2.14)$$

when  $k \geq k_{13}$ , and

$$\begin{aligned} & (-1)^k \frac{k!}{(k - k_{21})!} T \left( v_0 \cdot \left( \prod_{j=1}^{k_{21}} y_{i_{1,j}} \right) \left( \prod_{j=1}^k y_{i_{3,j}} \right) \cdot y_{n_1+1}^{k-k_{21}} \right) \\ &= E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k_{21}}} E_{i_{3,1}, n_1+1} \cdots E_{i_{3,k}, n_1+1} (v_0) \end{aligned} \quad (2.15)$$

if  $k \geq k_{21}$ .

*Proof.* For arbitrary  $k \geq 0$ , we prove (2.14) by induction on  $k_{13}$ . According to (2.1)-(2.3),

$$\tilde{\pi}(E_{i_2, i_1}) = -x_{i_1} x_{i_2} - y_{i_1} \partial_{y_{i_2}}, \quad \tilde{\pi}(E_{i_3, i_2}) = x_{i_3} \partial_{x_{i_2}} + y_{i_3} y_{i_2} \quad (2.16)$$

for  $i_1 \in J_1, i_2 \in J_2$  and  $i_3 \in J_3$ .

When  $k_{13} = 0$ , the first equation of the above expression gives

$$E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k}} (v_0) = (-1)^k \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \cdot x_{n_1+1}^k \right); \quad (2.17)$$

that is, (2.14) holds. Indeed, the numerator  $k!$  plays this role of normalization.

Assume that when  $k_{13} = k' - 1 \leq k - 1$ ,

$$\begin{aligned} & (-1)^k \frac{k!}{(k - k' + 1)!} T \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^{k'-1} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{k-k'+1} \right) \\ &= E_{i_{3,k'-1}, n_1+1} \cdots E_{i_{3,1}, n_1+1} E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k}} (v_0). \end{aligned} \quad (2.18)$$

When  $k_{13} = k'$ , we set

$$a_0 = (-1)^k \frac{k!}{(k - k' + 1)!} \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^{k'-1} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{k-k'+1} \right), \quad (2.19)$$

and

$$\begin{aligned} a_l &= (-1)^k \frac{k!}{(k - k' + 1)!} \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \cdot x_{n_1+1}^{k-k'+1} \right) \\ &\quad \times \frac{(x_{n_1+1} y_{n_1+1})^l (k - k' + 1)!}{(k - k' + l + 1)! l!} \tilde{\Delta}^l (f_0) \\ &= (-1)^k \frac{k!}{(k - k' + l + 1)! l!} \left( v_0 \left( \prod_{j=1}^k x_{i_{1,j}} \right) x_{n_1+1}^{k-k'+l+1} y_{n_1+1}^l \cdot \tilde{\Delta}^l (f_0) \right), \end{aligned} \quad (2.20)$$

where

$$f_0 = \prod_{j=1}^{k'-1} x_{i_3,j}. \quad (2.21)$$

According to (1.7),

$$\tilde{\Delta} x_{i_1,j} = x_{i_1,j} \tilde{\Delta}, \quad \tilde{\Delta} y_{i_3,l} = y_{i_3,l} \tilde{\Delta} \quad (2.22)$$

and

$$[\tilde{\Delta}, v_0] = - \sum_{n_1+1 \neq j \in J_2} (\partial_{x_j}(v_0) \partial_{y_j} + \partial_{y_j}(v_0) \partial_{x_j}). \quad (2.23)$$

Since

$$\partial_{x_{n_1+1}} \partial_{y_{n_1+1}} (x_{n_1+1}^{k-k'+1}) = 0, \quad (2.24)$$

we have

$$T(a_0) = \sum_{r=0}^{k'-1} a_r \quad (2.25)$$

by (2.6). Thus the second equation in (2.16) yields

$$\begin{aligned} & E_{i_3,k',n_1+1} \cdots E_{i_3,1,n_1+1} E_{n_1+1,i_1,1} \cdots E_{n_1+1,i_1,k} (v_0) \\ &= E_{i_3,k',n_1+1} \left( (-1)^k \frac{k!}{(k-k'+1)!} T \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_1,j} \right) \left( \prod_{j=1}^{k'-1} x_{i_3,j} \right) \cdot x_{n_1+1}^{k-k'+1} \right) \right) \\ &= E_{i_3,k',n_1+1} (T(a_0)) \\ &= \left( x_{i_3,k'} \partial_{x_{n_1+1}} \right) \left( \sum_{l=0}^{k'} a_l \right) + y_{i_3,k'} y_{n_1+1} \left( \sum_{l=0}^{k'} a_l \right) \\ &= x_{i_3,k'} \partial_{x_{n_1+1}} (a_0) + \sum_{l=1}^{k'} \left( y_{i_3,k'} y_{n_1+1} a_{l-1} + x_{i_3,k'} \partial_{x_{n_1+1}} (a_l) \right). \end{aligned} \quad (2.26)$$

We set

$$b_0 = x_{i_3,k'} \partial_{x_{n_1+1}} (a_0) = (-1)^k \frac{k!}{(k-k')!} \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_1,j} \right) \left( \prod_{j=1}^{k'} x_{i_3,j} \right) \cdot x_{n_1+1}^{k-k'} \right) \quad (2.27)$$

by (2.19) and

$$b_l = y_{i_3,k'} y_{n_1+1} a_{l-1} + x_{i_3,k'} \partial_{x_{n_1+1}} (a_l) \quad \text{for } l \geq 1. \quad (2.28)$$

We will show that (2.26) is equivalent to  $T(b_0)$ .

According to (1.7),

$$[\tilde{\Delta}, x_{i_3}] = y_{i_3} \quad \text{for } i_3 \in J_3. \quad (2.29)$$

Moreover, (2.29) and the second equation in (2.22) show

$$[\tilde{\Delta}^{s+1}, x_{i_3}] = (s+1) y_{i_3} \tilde{\Delta}^s \quad \text{for } s \in \mathbb{N}. \quad (2.30)$$

Thus for each  $l \in \mathbb{N}$ , we have

$$\tilde{\Delta}^l (f_0 \cdot x_{i_3}) = x_{i_3} \tilde{\Delta}^l (f_0) + l \tilde{\Delta}^{l-1} (f_0) y_{i_3}. \quad (2.31)$$



Therefore,

$$\begin{aligned}
b_l &= y_{i_3,k'} y_{n_1+1} a_{l-1} + x_{i_3,k'} \partial_{x_{n_1+1}}(a_l) \\
&= (-1)^k \frac{k!}{(k-k'+l)! (l-1)!} (y_{i_3,k'} y_{n_1+1} \cdot v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'+l} y_{n_1+1}^{l-1} \cdot \tilde{\Delta}^{l-1}(f_0)) + \\
&\quad x_{i_3,k'} \partial_{x_{n_1+1}} \left( (-1)^k \frac{k!}{(k-k'+l+1)! l!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'+l+1} y_{n_1+1}^l \cdot \tilde{\Delta}^l(f_0)) \right) \\
&= (-1)^k \frac{k!}{(k-k'+l)! (l-1)!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'} (x_{n_1+1} y_{n_1+1})^l) \cdot \tilde{\Delta}^{l-1}(f_0) y_{i_3,k'} \\
&\quad + (-1)^k \frac{k!}{(k-k'+l)! l!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'} (x_{n_1+1} y_{n_1+1})^l \cdot \tilde{\Delta}^l(f_0)) x_{i_3,k'} \\
&= \frac{(-1)^k k!}{(k-k'+l)! l!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'} (x_{n_1+1} y_{n_1+1})^l) (l \tilde{\Delta}^{l-1}(f_0) y_{i_3,k'} + x_{i_3,k'} \tilde{\Delta}^l(f_0)) \\
&= \frac{(-1)^k k!}{(k-k'+l)! l!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'} (x_{n_1+1} y_{n_1+1})^l) \cdot \tilde{\Delta}^l(f_0 x_{i_3,k'}) \\
&= \frac{(-1)^k k!}{(k-k'+l)! l!} (v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) x_{n_1+1}^{k-k'} (x_{n_1+1} y_{n_1+1})^l) \cdot (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^l (f_0 x_{i_3,k'}) \\
&= \frac{(x_{n_1+1} y_{n_1+1})^l}{(\prod_{r=1}^l (k-k'+r))! l!} (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^l \left( \frac{(-1)^k k!}{(k-k')!} v_0 \left( \prod_{j=1}^k x_{i_1,j} \right) \left( \prod_{j=1}^{k'} x_{i_3,j} \right) x_{n_1+1}^{k-k'} \right) \\
&= \frac{(x_{n_1+1} y_{n_1+1})^l}{(\prod_{r=1}^l (k-k'+r))! l!} (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^l (b_0) \tag{2.32}
\end{aligned}$$

by (2.21) and (2.27). Thus

$$\sum_{l=0}^{k'} b_l = T(b_0). \tag{2.33}$$

The proof of the second formula (2.15) is similar to the first one.  $\square$

**Corollary 2.1.** Fix  $k, \alpha_{n_1+1}, \beta_{n_1+1} \in \mathbb{N}$ . For  $j, i_{1,j} \in J_1, i_{3,j} \in J_3, v_1 \in \mathbb{F}[X_{J_1}, X_{J_2 \setminus \{n_1+1\}}, Y_{J_3}] \cup \mathbb{F}[X_{J_1}, Y_{J_2 \setminus \{n_1+1\}}, Y_{J_3}]$ , we have

$$\frac{k!}{\alpha_{n_1+1}!} T \left( v_1 \left( \prod_{j=1}^{k-\alpha_{n_1+1}} x_{i_3,j} \right) \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) = E_{i_3, k-\alpha_{n_1+1}, n_1+1} \cdots E_{i_3, 1, n_1+1} (v_1 x_{n_1+1}^k), \tag{2.34}$$

when  $k \geq \alpha_{n_1+1}$ , and

$$\frac{k!}{\beta_{n_1+1}!} T \left( v_1 \left( \prod_{j=1}^{k-\beta_{n_1+1}} y_{i_1,j} \right) \cdot y_{n_1+1}^{\beta_{n_1+1}} \right) = E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1, k-\beta_{n_1+1}}} (v_1 y_{n_1+1}^k), \tag{2.35}$$

where  $k \geq \beta_{n_1+1}$ .

*Proof.* By (2.14) and (2.17), we have

$$\begin{aligned} & \frac{k!}{(k-k_{13})!} T \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^{k_{13}} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{k-k_{13}} \right) \\ &= E_{i_3, k_{13}, n_1+1} \cdots E_{i_3, 1, n_1+1} E_{n_1+1, i_1, 1} \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \cdot x_{n_1+1}^k \right). \end{aligned} \quad (2.36)$$

Redenote

$$k_{13} = k - \alpha_{n_1+1}. \quad (2.37)$$

Then (2.36) becomes

$$\begin{aligned} & \frac{k!}{\alpha_{n_1+1}!} T \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^{k-\alpha_{n_1+1}} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{k-k_{13}} \right) \\ &= E_{i_3, k-\alpha_{n_1+1}, n_1+1} \cdots E_{i_3, 1, n_1+1} E_{n_1+1, i_1, 1} \left( v_0 \cdot \left( \prod_{j=1}^k x_{i_{1,j}} \right) \cdot x_{n_1+1}^k \right). \end{aligned} \quad (2.38)$$

Note that in the proof of Lemma 2.1, all the arguments still work if we replace the polynomial  $v_0$  by the rational function

$$v_0 = \frac{v_1}{\prod_{j=1}^k x_{i_{1,j}}}. \quad (2.39)$$

Substituting the above equation into (2.38), we obtain (2.34). Equation (2.35) can be obtained similarly.  $\square$

For  $m \in \mathbb{N}$  and  $r \in \{1, 2, 3\}$ , we denote

$$X_{J_r}^m = \{x_{j_1}^{\alpha_{j_1}} x_{j_2}^{\alpha_{j_2}} \cdots x_{j_k}^{\alpha_{j_k}} \mid \{j_1, \dots, j_k\} = J_r; \alpha_{j_1}, \dots, \alpha_{j_k} \in \mathbb{N}; \sum_{i=1}^k \alpha_{j_i} = m\} \quad (2.40)$$

and

$$Y_{J_r}^m = \{y_{j_1}^{\alpha_{j_1}} y_{j_2}^{\alpha_{j_2}} \cdots y_{j_k}^{\alpha_{j_k}} \mid \{j_1, \dots, j_k\} = J_r; \alpha_{j_1}, \dots, \alpha_{j_k} \in \mathbb{N}; \sum_{i=1}^k \alpha_{j_i} = m\}. \quad (2.41)$$

Next we want to introduce a degree function  $\mathfrak{d} : \mathcal{A}_{(\ell_1, \ell_2)} \rightarrow \mathbb{N}$  which can help us determine  $M_k/M_{k-1}$ . Our motivation comes from the facts as follow. Suppose  $\ell_1 = -m_1$  and  $\ell_2 = -m_2$  with  $m_1, m_2 \in \mathbb{N}$ . By (2.11), we have

$$M_0 = \text{Span}\{X_{J_1}^{m_1} Y_{J_3}^{m_2}\}. \quad (2.42)$$

Applying  $E_{i_2, i_1}$  and  $E_{i_3, i_2}$  to  $M_0$  with  $i_r \in J_r$ , we have

$$\{X_{J_1}^{m_1+1} x_{i_2} Y_{J_3}^{m_2}, X_{J_1}^{m_1} y_{i_2} Y_{J_3}^{m_2+1} \mid i_2 \in J_2\} \subset M_1 \setminus M_0 \quad (2.43)$$

by (2.1)-(2.3). Fix  $k, k_{13}, k_{21} \in \mathbb{N}$ . For each  $j, i_{1,j} \in J_1, i_{3,j} \in J_3, v_0 \in \mathbb{F} [X_{J_1}, X_{J_2 \setminus \{n_1+1\}}, Y_{J_3}] \cup \mathbb{F} [X_{J_1}, Y_{J_2 \setminus \{n_1+1\}}, Y_{J_3}]$ , we have

$$\begin{aligned} & (-1)^k k! T \left( X_{J_1}^{m_1} Y_{J_3}^{m_2} \left( \prod_{j=1}^k x_{i_{1,j}} \right) \left( \prod_{j=1}^k x_{i_{3,j}} \right) \right) \\ &= E_{i_{3,k}, n_1+1} \cdots E_{i_{3,1}, n_1+1} E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k}} (X_{J_1}^{m_1} Y_{J_3}^{m_2}) \subset M_{2k}. \end{aligned} \quad (2.44)$$

When  $k_{21} = k$  in (2.15), we find

$$\begin{aligned} & (-1)^k k! T \left( X_{J_1}^{m_1} Y_{J_3}^{m_2} \left( \prod_{j=1}^k y_{i_{1,j}} \right) \left( \prod_{j=1}^k y_{i_{3,j}} \right) \right) \\ &= E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1,k}} E_{i_{3,1}, n_1+1} \cdots E_{i_{3,k}, n_1+1} (X_{J_1}^{m_1} Y_{J_3}^{m_2}) \subset M_{2k}. \end{aligned} \quad (2.45)$$

The above three expressions motivate us to define  $\mathfrak{d} : \mathcal{A}_{(\ell_1, \ell_2)} \rightarrow \mathbb{N}$  by

$$\mathfrak{d}(x^\alpha y^\beta) = 2 \sum_{i \in J_3} \alpha_i + \sum_{i \in J_2} \alpha_i + 2 \sum_{i \in J_1} \beta_i + \sum_{i \in J_2} \beta_i - \frac{\ell_1 + |\ell_1| + \ell_2 + |\ell_2|}{2}, \quad (2.46)$$

$$\mathfrak{d}(\lambda f) = \mathfrak{d}(f), \quad \forall \lambda \in \mathbb{F} \setminus \{0\}, \quad (2.47)$$

$$\mathfrak{d}\left(\sum_{i=1}^m f_i\right) = \max\{\mathfrak{d}(f_1), \dots, \mathfrak{d}(f_m)\}, \quad \text{where } f_i \text{ are monomials.} \quad (2.48)$$

The last term in (2.46) is to ensure  $\mathfrak{d}(M_0) = 0$ .

For  $\alpha, \beta \in \mathbb{N}^n$  with  $\alpha_{n_1+1} \beta_{n_1+1} = 0$  and  $(x_{n_1+1} y_{n_1+1}) (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}}) (x^\alpha y^\beta) \neq 0$ , we have

$$\begin{aligned} & \mathfrak{d}((x_{n_1+1} y_{n_1+1}) (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}}) (x^\alpha y^\beta)) \\ &= \mathfrak{d}((x_{n_1+1} y_{n_1+1}) (\sum_{i=1}^{n_1} x_i \partial_{y_i} - \sum_{r=n_1+2}^{n_2} \partial_{x_r} \partial_{y_r} + \sum_{s=n_2+1}^n y_s \partial_{x_s}) (x^\alpha y^\beta)) \\ &= \max\{\mathfrak{d}((x_{n_1+1} y_{n_1+1} x_i \partial_{y_i}) (x^\alpha y^\beta)), \mathfrak{d}((x_{n_1+1} y_{n_1+1} \partial_{x_r} \partial_{y_r}) (x^\alpha y^\beta)), \\ & \quad \mathfrak{d}((x_{n_1+1} y_{n_1+1} y_s \partial_{x_s}) (x^\alpha y^\beta)) \mid i \in J_1, n_1+2 \leq r \leq n_2, s \in J_3\} \\ &= \mathfrak{d}(x^\alpha y^\beta). \end{aligned} \quad (2.49)$$

In general,

$$\mathfrak{d}((x_{n_1+1} y_{n_1+1})^i (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^i (x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta) \text{ or } 0, \quad (2.50)$$

where  $\alpha_{n_1+1} \beta_{n_1+1} = 0$ .

Recall that  $T$  is injective. Hence

$$\mathfrak{d}(T(x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta), \quad \text{where } \alpha_{n_1+1} \beta_{n_1+1} = 0. \quad (2.51)$$

For convenience, we denote

$$L_3 = J_3 \times J_1, \quad L_2 = J_3 \times J_2, \quad L_1 = J_2 \times J_1, \quad L = L_1 \cup L_2 \cup L_3. \quad (2.52)$$

From the above properties, we have

$$\mathfrak{d}(E_{i,j}(f)) \leq \begin{cases} \mathfrak{d}(f) + 2 & \text{if } (i,j) \in L_3, \\ \mathfrak{d}(f) + 1 & \text{if } (i,j) \in L_1 \cup L_2, \\ \mathfrak{d}(f) & \text{if } (i,j) \notin L. \end{cases} \quad (2.53)$$

We set

$$TN(k) = \text{Span}\{T(x^\alpha y^\beta) \mid \mathfrak{d}(x^\alpha y^\beta) = k, x^\alpha y^\beta \in \mathcal{A}_{(\ell_1, \ell_2)}, \alpha_{n_1+1} \beta_{n_1+1} = 0\}, \quad (2.54)$$

$$TN_k = \text{Span}\{T(x^\alpha y^\beta) \mid \mathfrak{d}(x^\alpha y^\beta) \leq k, x^\alpha y^\beta \in \mathcal{A}_{(\ell_1, \ell_2)}, \alpha_{n_1+1} \beta_{n_1+1} = 0\}. \quad (2.55)$$

According to (2.1)-(2.3), the alternating polynomials

$$x_{i_1} x_{i_3} - y_{i_1} y_{i_3} = -\tilde{\pi}(E_{i_3, i_1}) \quad \text{for } i_1 \in J_1, i_3 \in J_3 \quad (2.56)$$

(cf. (1.13)). By (2.7),

$$T(f)(x_{i_1} x_{i_3} - y_{i_1} y_{i_3}) = T(f(x_{i_1} x_{i_3} - y_{i_1} y_{i_3})) \quad \text{for } f \in \mathcal{A}. \quad (2.57)$$

Set

$$P = \{x_{i_1} x_{i_3} - y_{i_1} y_{i_3} \mid i_1 \in J_1, i_3 \in J_3\}. \quad (2.58)$$

Then for  $r \in \mathbb{N}$ ,

$$\begin{aligned} & TN \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} P^r \\ &= \text{Span}\{T(f) p_1 \cdot p_2 \cdots p_r \mid f \in N \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix}, p_i \in P, 1 \leq i \leq r\} \\ &= T \left( N \begin{pmatrix} k_{12} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} P^r \right) \subset \mathcal{H}_{(\ell_1, \ell_2)}. \end{aligned} \quad (2.59)$$

Moreover, we have

$$\mathfrak{d}(f \cdot p) = \mathfrak{d}(f) + 2 \quad \text{for } p \in P, 0 \neq f \in \mathcal{A}_{(\ell_1, \ell_2)}. \quad (2.60)$$

Set

$$\begin{aligned} V_k &= \text{Span}\{TN(i), TN(k-i)P^i \mid i = 0, 1, \dots, k\} \\ &= \text{Span}\{TN_k, TN(k-i)P^i \mid i = 1, \dots, k\}. \end{aligned} \quad (2.61)$$

Then  $M_0 = V_0 = TN_0 = TN(0)$ . Denote  $\mathfrak{g} = \mathfrak{sl}(n)$ .

**Proposition 2.1.** *For  $k \in \mathbb{N}$ , we have  $M_k = U_k(\mathfrak{g})(M_0) = V_k$ . Moreover,  $M = \mathcal{H}_{(\ell_1, \ell_2)} = U(\mathfrak{g})(M_0)$ .*

*Proof.* We prove  $M_i = V_i$  for any  $i \in \mathbb{N}$  by induction. It holds when  $i = 0$ . Suppose that  $M_i = V_i$  with  $i \leq k-1$ . Consider  $i = k$ . Note

$$M_k = U_k(\mathfrak{g})(M_0) = \mathfrak{g}(M_{k-1}) = \mathfrak{g}(V_{k-1}). \quad (2.62)$$

Let  $f \in TN_{k-1}$ . If  $(i, j) \notin L_3$ ,

$$\mathfrak{d}(E_{i,j}(f)) \leq \mathfrak{d}(f) + 1 \Rightarrow E_{i,j}(f) \in V_k. \quad (2.63)$$

When  $(i, j) \in L_3$ , (2.56) yields

$$E_{i,j}(f) \in TN_{k-1}P \subset V_k. \quad (2.64)$$

Fix  $g \in TN_{k-r-1}P^r$  with  $r \in \overline{1, k-1}$ . If  $(i, j) \in L_3$ , then

$$E_{i,j}(g) \in TN_{k-r-1}P^{r+1} \subset V_k. \quad (2.65)$$

According (2.56)-(2.58), we may assume

$$g = E_{i_1, j_1} \cdots E_{i_r, j_r}(g') \quad (2.66)$$

with  $g' \in TN_{k-r-1} \subset V_{k-r-1} = M_{k-r-1}$  and  $(i_1, j_1), \dots, (i_r, j_r) \in L_3$ . When  $(i, j) \notin L_3$ ,

$$\begin{aligned} E_{i,j}(g) &= \sum_{s=1}^r (E_{i_1, j_1} \cdots E_{i_{s-1}, j_{s-1}} [E_{i,j}, E_{i_s, j_s}] E_{i_{s+1}, j_{s+1}} \cdots E_{i_r, j_r}(g')) \\ &\quad + E_{i_1, j_1} \cdots E_{i_r, j_r} E_{i,j}(g') \subset M_{k-1} + P^r TN_{k-r} \subset V_k \end{aligned} \quad (2.67)$$

by (2.61). Thus we have proved  $M_k \subset V_k$ .

Next we want to prove  $V_k \subset M_k$ . Since the case of  $\ell_2 \leq 0 \leq \ell_1$  and the case of  $\ell_1 \leq 0 \leq \ell_2$  are symmetric, it is enough to prove it in the following two cases.

*Case 1.*  $\ell_1, \ell_2 \leq 0$ .

We write  $\ell_1 = -m_1$ ,  $\ell_2 = -m_2$ , where  $m_1, m_2 \in \mathbb{N}$ . For  $k_{13}, \alpha_{n_1+1} \in \mathbb{N}$ , Lemma 2.1 gives

$$\begin{aligned} (-1)^{k_{13} + \alpha_{n_1+1}} \frac{(k_{13} + \alpha_{n_1+1})!}{\alpha_{n_1+1}!} T \left( X_{J_1}^{m_1} X_{J_3}^{m_2} \cdot \left( \prod_{j=1}^{k_{13} + \alpha_{n_1+1}} x_{i_{1,j}} \right) \left( \prod_{j=1}^{k_{13}} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) = \\ E_{i_{3,1}, n_1+1} \cdots E_{i_{3, k_{13}}, n_1+1} E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1, k_{13} + \alpha_{n_1+1}}} (X_{J_1}^{m_1} Y_{J_3}^{m_2}), \end{aligned}$$

where  $i_{1,r} \in J_1, i_{3,s} \in J_3$ . This shows that

$$T \left( X_{J_1}^{m_1} Y_{J_3}^{m_2} \cdot \left( \prod_{j=1}^{k_{13} + \alpha_{n_1+1}} x_{i_{1,j}} \right) \left( \prod_{j=1}^{k_{13}} x_{i_{3,j}} \right) \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2k_{13} + \alpha_{n_1+1}}, \quad (2.68)$$

which implies

$$T \left( N \begin{pmatrix} m_1 + k_{13} + \alpha_{n_1+1} & 0 & k_{13} \\ 0 & 0 & m_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2k_{13} + \alpha_{n_1+1}}. \quad (2.69)$$

Next we want to prove

$$T \left( N \begin{pmatrix} m_1 + k_{13} + \alpha_{n_1+1} & 0 & k_{13} \\ k_{21} & 0 & m_2 + k_{21} \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2(k_{13} + k_{21}) + \alpha_{n_1+1}}. \quad (2.70)$$

by induction on  $k_{21}$ . It holds for  $k_{21} = 0$ . Suppose that it holds for  $k_{21} \leq r$ . Let

$$h \in N \begin{pmatrix} m_1 + k_{13} + \alpha_{n_1+1} & 0 & k_{13} \\ r & 0 & m_2 + r \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}}. \quad (2.71)$$

Note

$$T(h \cdot y_{i_1} y_{i_3}) = -T(h \cdot (x_{i_1} x_{i_3} - y_{i_1} y_{i_3})) + T(h \cdot x_{i_1} x_{i_3}). \quad (2.72)$$

Moreover,

$$\begin{aligned} T(h \cdot x_{i_1} x_{i_3}) &\in \left( N \begin{pmatrix} m_1 + k_{13} + \alpha_{n_1+1} + 1 & 0 & k_{13} + 1 \\ r & 0 & m_2 + r \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \\ &\subset M_{2(k_{13}+1)+2r+\alpha_{n_1+1}} \end{aligned} \quad (2.73)$$

by the inductual assumption, and

$$\begin{aligned} &-T(h \cdot (x_{i_1} x_{i_3} - y_{i_1} y_{i_3})) \\ &= -(x_{i_1} x_{i_3} - y_{i_1} y_{i_3}) T(h) = E_{i_3, i_1}(T(h)) \\ &\subset E_{i_3, i_1} \left[ T \left( N \begin{pmatrix} m_1 + k_{13} + \alpha_{n_1+1} & 0 & k_{13} \\ r & 0 & m_2 + r \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \right] \\ &\subset E_{i_3, i_1}(M_{2k_{13}+2r+\alpha_{n_1+1}}) \subset M_{2k_{13}+2r+1+\alpha_{n_1+1}} \subset M_{2k_{13}+2r+2+\alpha_{n_1+1}} \end{aligned} \quad (2.74)$$

by (2.56) and (2.57). So the above three expressions imply

$$T(h \cdot y_{i_1} y_{i_3}) \in M_{2k_{13}+2r+2+\alpha_{n_1+1}}. \quad (2.75)$$

This proves (2.70). Symmetrically, we have

$$T \left( N \begin{pmatrix} m_1 + k_{13} & 0 & k_{13} \\ k_{21} & 0 & m_2 + k_{21} + \beta_{n_1+1} \end{pmatrix} \cdot y_{n_1+1}^{\beta_{n_1+1}} \right) \in M_{2(k_{13}+k_{21})+\beta_{n_1+1}}. \quad (2.76)$$

Let  $k_{12}, k_{22} \in \mathbb{N}$ . For convenience, we redenote

$$k_1 = m_1 + k_{12} + k_{13} + \alpha_{n_1+1}, \quad k_2 = m_2 + k_{21} + k_{22} \quad (2.77)$$

and

$$N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} = \left\{ x^\alpha y^\beta \in N \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \mid \alpha_{n_1+1} = \beta_{n_1+1} = 0 \right\} \quad (2.78)$$

(cf. (2.9)). We want to prove

$$T \left( N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+\alpha_{n_1+1}} \quad (2.79)$$

by induction on  $k_{12} + k_{22}$ . When  $k_{12} + k_{22} = 0$ , it is exactly (2.70). Assume that it holds for  $k_{12} + k_{22} < k_0$ . Let  $i_1 \in J_1$  and  $i_2 \in J_2 \setminus \{n_1 + 1\}$ . According to (2.1)-(2.3) and (2.7),

$$\tilde{\pi}(E_{i_2, i_1}) = -x_{i_1} x_{i_2} - y_{i_1} \partial_{y_{i_2}}, \quad \tilde{\pi}(E_{i_2, i_1}) T = T \tilde{\pi}(E_{i_2, i_1}). \quad (2.80)$$

Let

$$h \in N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \quad \text{with } k_{12} + k_{22} = k_0 - 1. \quad (2.81)$$

We have

$$E_{i_2, i_1}(T(h)) = T(E_{i_2, i_1}(h)) = -T(x_{i_1} x_{i_2} h) - T(y_{i_1} \partial_{y_{i_2}}(h)). \quad (2.82)$$

Note

$$x_{i_1} x_{i_2} h \in N' \begin{pmatrix} k_1 + 1 & k_{12} + 1 & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \quad (2.83)$$

and

$$\begin{aligned} T(y_{i_1} \partial_{y_{i_2}}(h)) &\in T \left( N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} + 1 & k_{22} - 1 & k_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \\ &\subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+1+\alpha_{n_1+1}}. \end{aligned} \quad (2.84)$$

On the other hand,

$$E_{i_2, i_1}(T(h)) \in E_{i_2, i_1}(M_{2k_{21}+2k_{13}+k_{12}+k_{22}+\alpha_{n_1+1}}) \subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+1+\alpha_{n_1+1}}. \quad (2.85)$$

Thus

$$T(x_{i_1} x_{i_2} h) = -E_{i_2, i_1}(T(h)) - T(y_{i_1} \partial_{y_{i_2}}(h)) \in M_{2k_{21}+2k_{13}+k_{12}+k_{22}+1+\alpha_{n_1+1}}. \quad (2.86)$$

This proves

$$T \left( N' \begin{pmatrix} k_1 + 1 & k_{12} + 1 & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+1+\alpha_{n_1+1}}. \quad (2.87)$$

By applying  $E_{i_3, i_2}$  with  $i_2 \in J_2 \setminus \{n_1 + 1\}$  and  $i_3 \in J_3$ , we can similarly obtain

$$T \left( N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} + 1 & k_2 + 1 \end{pmatrix} \cdot x_{n_1+1}^{\alpha_{n_1+1}} \right) \subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+1+\alpha_{n_1+1}}. \quad (2.88)$$

Hence (2.79) holds for  $k_{12} + k_{22} = k_0$ . By induction, (2.79) holds for any  $k_{12}, k_{22} \in \mathbb{N}$ . Using the symmetry of  $x'_i$ 's and  $y'_j$ 's, we have

$$T \left( N' \begin{pmatrix} k_1 & k_{12} & k_{13} \\ k_{21} & k_{22} & k_2 \end{pmatrix} \cdot y_{n_1+1}^{\beta_{n_1+1}} \right) \subset M_{2k_{21}+2k_{13}+k_{12}+k_{22}+\beta_{n_1+1}}, \quad (2.89)$$

where

$$k_1 = m_1 + k_{12} + k_{13}, \quad k_2 = m_2 + k_{21} + k_{22} + \beta_{n_1+1}. \quad (2.90)$$

Expressions (2.54), (2.79) and (2.89) show

$$TN(\ell) \subset M_\ell \quad \text{for } \ell \in \mathbb{N}. \quad (2.91)$$

For  $0 < r \in \mathbb{N}$ , any element  $w \in TN(k - r) \cdot P^r$  can be written as

$$w = E_{i_1, j_1} \cdots E_{i_r, j_r}(h) \in E_{i_1, j_1} \cdots E_{i_r, j_r}(M_{k-r}) \subset M_k \quad (2.92)$$

by (2.56)-(2.58), where  $h \in TN(k - r) \subset M_{k-r}$ ,  $i_1, \dots, i_r \in J_1$  and  $j_1, \dots, j_r \in J_2$ . Now Expressions (2.55), (2.91) and (2.92) yield  $V_k \subset M_k$ . Recall we have already proved

$M_k \subset V_k$  before Case 1. Thus  $M_k = V_k$ . By induction, the first conclusion in the proposition holds in this case.

*Case 2.*  $\ell_1 \leq 0 \leq \ell_2$ .

In this case,

$$M_0 = TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 & 0 \end{pmatrix}. \quad (2.93)$$

For any  $r \in \mathbb{N}$  and

$$g \in N \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 + r & r \end{pmatrix}, \quad (2.94)$$

$$(\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})(g) = \left( \sum_{i=1}^{n_1} x_i \partial_{y_i} - \sum_{r=n_1+2}^{n_2} \partial_{x_r} \partial_{y_r} + \sum_{s=n_2+1}^n y_s \partial_{x_s} \right)(g) = 0 \quad (2.95)$$

by (1.7) and (2.9). Thus (2.6) and (2.10) give

$$TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 + r & r \end{pmatrix} = \text{Span} \left\{ N \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 + r & r \end{pmatrix} \right\}. \quad (2.96)$$

According to (2.1)-(2.3), we have

$$\tilde{\pi}(E_{i_3, i_2}) = x_{i_3} \partial_{x_{i_2}} + y_{i_3} y_{i_2} \quad \text{for } i_2 \in J_2, i_3 \in J_3. \quad (2.97)$$

For  $g$  in (2.94), we get

$$E_{i_3, i_2}(g) = y_{i_3} y_{i_2} g \in N \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & \ell_2 + r + 1 & r + 1 \end{pmatrix}. \quad (2.98)$$

By (2.96), (2.98) and induction, we can prove

$$TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & k_{23} + \ell_2 & k_{23} \end{pmatrix} \subset M_{k_{23}} \quad \text{for } k_{23} \in \mathbb{N}. \quad (2.99)$$

Let  $k_{21}, k_{22}, k_{23}, \beta_{n_1+1} \in \mathbb{N}$  such that  $k_{22} \geq \beta_{n_1+1}$  and

$$k_{21} + k_{22} - k_{23} = \ell_2. \quad (2.100)$$

Take

$$v_1 \in X_{J_1}^{-\ell_1} Y_{J_2 \setminus \{n_1+1\}}^{k_{22} - \beta_{n_1+1}} Y_{J_3}^{k_{23}}. \quad (2.101)$$

Then

$$v_1 y_{n_1+1}^{k_{21} + \beta_{n_1+1}} \in TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ 0 & k_{23} + \ell_2 & k_{23} \end{pmatrix} \subset M_{k_{23}} \quad (2.102)$$

and

$$v_1 \left( \prod_{j=1}^{k_{21}} y_{i_{1,j}} \right) \cdot y_{n_1+1}^{\beta_{n_1+1}} \in N \begin{pmatrix} -\ell_1 & 0 & 0 \\ k_{21} & k_{22} & k_{23} \end{pmatrix}. \quad (2.103)$$

By (2.35) with  $k = k_{21} + \beta_{n_1+1}$ , we have

$$\begin{aligned} & \frac{(k_{21} + \beta_{n_1+1})!}{\beta_{n_1+1}!} T \left( v_1 \left( \prod_{j=1}^{k_{21}} y_{i_{1,j}} \right) \cdot y_{n_1+1}^{\beta_{n_1+1}} \right) \\ &= E_{n_1+1, i_{1,1}} \cdots E_{n_1+1, i_{1, k_{21}}} \left( v_1 y_{n_1+1}^{k_{21} + \beta_{n_1+1}} \right) \subset M_{k_{21} + k_{23}}. \end{aligned} \quad (2.104)$$



This shows

$$TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \subset M_{k_{21}+k_{23}}. \quad (2.105)$$

Since

$$\begin{aligned} T(h \cdot x_{i_1} x_{i_3}) &= T(h \cdot y_{i_1} y_{i_3}) + T(h \cdot (x_{i_1} x_{i_3} - y_{i_1} y_{i_3})) \\ &= T(h \cdot y_{i_1} y_{i_3}) - E_{i_3, i_1}[T(h)] \end{aligned} \quad (2.106)$$

for  $i_1 \in J_1$  and  $i_3 \in J_3$ , we have

$$\begin{aligned} &TN \begin{pmatrix} -\ell_1 + 1 & 0 & 1 \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \\ &\subset \mathfrak{g} \left[ TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right] + TN \begin{pmatrix} -\ell_1 & 0 & 0 \\ k_{21} + 1 & k_{22} & k_{23} + 1 \end{pmatrix} \\ &\subset M_{k_{21}+k_{23}+1} + M_{k_{21}+k_{23}+2} \subset M_{k_{21}+k_{23}+2}. \end{aligned} \quad (2.107)$$

Repeatedly applying (2.106), we can prove

$$TN \begin{pmatrix} k_{11} & 0 & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \subset M_{k_{21}+k_{23}+2k_{13}}. \quad (2.108)$$

by induction.

For  $i_1 \in J_1$  and  $i_2 \in J_2 \setminus \{n_1 + 1\}$ , (2.1)-(2.3) and (2.7) show

$$T(x_{i_1} x_{i_2} h) = -E_{i_2, i_1}(T(h)) - T(y_{i_1} \partial_{y_{i_2}} h), \quad (2.109)$$

Denote

$$N^* \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} = \left\{ x^\alpha y^\beta \in N \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \mid \alpha_{n_1+1} = 0 \right\}. \quad (2.110)$$

Applying (2.109) to (2.108), we get

$$\begin{aligned} &T \left[ N^* \begin{pmatrix} k_{11} & 1 & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right] \\ &\subset \mathfrak{g} \left[ TN \begin{pmatrix} k_{11} - 1 & 0 & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right] + TN \begin{pmatrix} k_{11} - 1 & 0 & k_{13} \\ k_{21} + 1 & k_{22} - 1 & k_{23} \end{pmatrix} \\ &\subset M_{k_{21}+k_{23}+2k_{13}+1} + M_{k_{21}+1+k_{23}+2k_{13}} \subset M_{k_{21}+k_{23}+2k_{13}+1}. \end{aligned} \quad (2.111)$$

By (2.108), (2.111) and induction on  $k_{12}$ , we can deduce

$$T \left[ N^* \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right] \subset M_{k_{21}+k_{23}+2k_{13}+k_{12}} = M_{2k_{21}+k_{22}+2k_{13}+k_{12}-\ell_2}. \quad (2.112)$$

Denote

$$N^\# \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} = \left\{ x^\alpha y^\beta \in N \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \mid \beta_{n_1+1} = 0 \right\}. \quad (2.113)$$

By the symmetry of  $x'_i$ s and  $y'_j$ s, we have

$$T \left[ N^\# \begin{pmatrix} k_{11} & k_{12} & k_{13} \\ k_{21} & k_{22} & k_{23} \end{pmatrix} \right] \subset M_{k_{21}+k_{23}+2k_{13}+k_{12}} = M_{2k_{21}+k_{22}+2k_{13}+k_{12}-\ell_2}. \quad (2.114)$$

Expressions (2.112) and (2.114) yield  $TN_k \subset M_k$ . According to (2.92), we have  $V_k \subset M_k$ . Recall we have already proved  $M_k \subset V_k$  before Case 1. Thus  $M_k = V_k$ . By induction, the first conclusion in the proposition holds in this case.

By the symmetry of  $x'_i$ s and  $y'_j$ s, the first conclusion in the proposition holds in any case. Thanks to (2.55) and (2.61),

$$M = \mathcal{H}_{(\ell_1, \ell_2)} \subset \bigcup_{k=0}^{\infty} TN_k \subset \bigcup_{k=0}^{\infty} V_k = \bigcup_{k=0}^{\infty} M_k = M. \quad (2.115)$$

So (1.9) holds.  $\square$

According to (2.55) and (2.61),  $M_k = V_k$  is spanned by the set

$$\begin{aligned} \mathcal{S}_k = & \{T(x^\alpha y^\beta), T(x^\alpha y^\beta)P^{k-\mathfrak{d}(x^\alpha y^\beta)} \mid x^\alpha y^\beta \in \mathcal{A}_{(\ell_1, \ell_2)}, \\ & \alpha_{n_1+1}\beta_{n_1+1} = 0, \mathfrak{d}(x^\alpha y^\beta) \leq k\}. \end{aligned} \quad (2.116)$$

### 3 Products of Alternating Polynomials

In this section, we want to study the linear dependence among the elements in  $\mathcal{S}_k$ .

Denote

$$Z = \{z_{j,i} \mid i \in J_1, j \in J_3\} \quad (3.1)$$

as a set of  $n_1(n - n_2)$  variables. The notion is compatible with (2.56). Set

$$\mathcal{B} = \mathbb{F}[Z] \quad (3.2)$$

the polynomial algebra in  $Z$ . Denote by  $\mathcal{B}_m$  the subspace of its homogeneous polynomials with degree  $m$  and by  $Z^{(m)}$  the set of monomials with degree  $m$ .

Define associative algebra homomorphisms  $\varphi_x, \varphi_y : \mathcal{B} \rightarrow \mathcal{A}$  by

$$\varphi_x(z_{j,i}) = x_i x_j, \quad \varphi_y(z_{j,i}) = y_i y_j \quad \text{for } i \in J_1, j \in J_3. \quad (3.3)$$

Then  $\ker \varphi_x$  and  $\ker \varphi_y$  are graded ideals of  $\mathcal{B}$ . For subset  $S$  of a polynomial algebra, we use  $\langle S \rangle$  to denote the ideal generated by  $S$ . Let

$$\mathcal{R}_2 = \left\langle \begin{array}{c|c} z_{j_1, i_1} & z_{j_1, i_2} \\ \hline z_{j_2, i_1} & z_{j_2, i_2} \end{array} \mid i_1, i_2 \in J_1, j_1, j_2 \in J_3 \right\rangle. \quad (3.4)$$

**Lemma 3.1.** *We have*

$$\ker \varphi_x = \ker \varphi_y = \mathcal{R}_2. \quad (3.5)$$

*Proof.* First, we want to prove  $\ker \varphi_x = \mathcal{R}_2$ . Let  $S_r$  be the permutation group on  $\{1, 2, \dots, r\}$ . We define a linear action of the direct product group  $S_r \times S_r$  on  $\mathcal{B}_r$  by

$$(\sigma_1, \sigma_2) \left( \prod_{s=1}^r z_{j_s, i_s} \right) = \prod_{s=1}^r z_{j_{\sigma_1(s)}, i_{\sigma_2(s)}}, \quad (3.6)$$

where  $(j_1, i_1) \geq (j_2, i_2) \geq \dots \geq (j_r, i_r)$  in lexical order. Moreover,

$$\varphi_x(g) = \varphi_x[\sigma(g)] \quad (3.7)$$

for each  $\sigma \in S_r \times S_r$  and  $g \in \mathcal{B}_r$ .

Define

$$\mathcal{I}_1 \left( \prod_{s=1}^k z_{j_s, i_s} \right) = \{i_1, i_2, \dots, i_k\} \subset J_1, \quad \mathcal{I}_3 \left( \prod_{s=1}^k z_{j_s, i_s} \right) = \{j_1, j_2, \dots, j_k\} \subset J_3. \quad (3.8)$$

For example,  $\mathcal{I}_1(z_{n,1}z_{n-1,2}z_{n,2}) = \{1, 2, 2\}$ . Let  $f \in \ker \varphi_x \cap \mathcal{B}_r$ . Since the image under  $\varphi_x$  of the monomials in  $\mathcal{B}$  with different  $\mathcal{I}_1$  and  $\mathcal{I}_3$  are linearly independent, we may assume that  $f = \sum_{k=1}^m a_k f_k$  is a linear combination of the monomials with

$$\mathcal{I}_1(f_1) = \mathcal{I}_1(f_2) = \dots = \mathcal{I}_1(f_m), \quad \mathcal{I}_3(f_1) = \mathcal{I}_3(f_2) = \dots = \mathcal{I}_3(f_m). \quad (3.9)$$

Moreover,  $\sum_{k=1}^m a_k = 0$  by  $\varphi_x(f)=0$ . There exists  $\sigma_k \in S_r \times S_r$  such that  $\sigma_k(f_k) = f_1$ . Thus

$$f = \sum_{k=1}^m a_k f_k = \sum_{k=2}^m a_k \sigma_k(f_1) + a_1 f_1 = \sum_{k=2}^m a_k (\sigma_k(f_1) - f_1). \quad (3.10)$$

Next, we want to prove  $\sigma(f) - f \in \mathcal{R}_2$  for any  $\sigma \in S_r \times S_r$  and  $f \in Z^{(r)}$ . Firstly,  $\mathcal{R}_2$  is invariant under  $S_r \times S_r$  because

$$\begin{aligned} & \sigma((z_{i,j}z_{i',j'} - z_{i',j}z_{i,j'})g) \\ &= (z_{\sigma_1(i), \sigma_2(j)}z_{\sigma_1(i'), \sigma_2(j')} - z_{\sigma_1(i'), \sigma_2(j)}z_{\sigma_1(i), \sigma_2(j')}) \sigma(g) \in \mathcal{R}_2 \end{aligned} \quad (3.11)$$

for each  $g \in Z^{(r-2)}$ . We set

$$\sigma = (\sigma_1^1 \cdots \sigma_1^p, \sigma_2^1 \cdots \sigma_2^q), \quad (3.12)$$

where  $\sigma_i^j \in S_r$  are transpositions. Note that

$$\begin{aligned} & ((\iota_1 \iota_2), 1) \cdot \left( \prod_{s=1}^r z_{j_s, i_s} \right) - \prod_{s=1}^r z_{j_s, i_s} \\ &= \left( \prod_{s \neq \iota_1, \iota_2} z_{i_s, j_s} \right) \cdot (z_{j_{\iota_2}, i_{\iota_1}} z_{j_{\iota_1}, i_{\iota_2}} - z_{j_{\iota_1}, i_{\iota_1}} z_{j_{\iota_2}, i_{\iota_2}}) \in \mathcal{R}_2. \end{aligned} \quad (3.13)$$

Symmetrically,

$$(1, (\iota_1 \iota_2)) \left( \prod_{s=1}^r z_{j_s, i_s} \right) - \prod_{s=1}^r z_{j_s, i_s} \in \mathcal{R}_2. \quad (3.14)$$

Let  $h \in Z^{(r)}$ . We have

$$\begin{aligned}
\sigma(g) - g &= (\sigma_1^1 \cdots \sigma_1^p, \sigma_2^1 \cdots \sigma_2^q)(g) - g \\
&= \sum_{s=1}^p ((\sigma_1^s, 1)[(\sigma_1^{s+1} \cdots \sigma_1^p, \sigma_2^1 \cdots \sigma_2^q)(g)] - (\sigma_1^{s+1} \cdots \sigma_1^p, \sigma_2^1 \cdots \sigma_2^q)(g)) \\
&\quad + \sum_{t=1}^q ((1, \sigma_2^t)[(1, \sigma_2^{t+1} \cdots \sigma_2^q)(g)] - (1, \sigma_2^{t+1} \cdots \sigma_2^q)(g)) \in \mathcal{R}_2. \tag{3.15}
\end{aligned}$$

Hence

$$f = \sum_{k=2}^m a_k (\sigma_k(f_1) - f_1) \in \mathcal{R}_2 \tag{3.16}$$

by (3.13). So  $\ker \varphi_x \cap \mathcal{B}_r \subset \mathcal{R}_2$  for each  $r \geq 2$ . Symmetrically,  $\ker \varphi_y \cap \mathcal{B}_r \subset \mathcal{R}_2$ . Furthermore, (3.3) and (3.5) imply  $\varphi_x(\mathcal{R}_2) = \varphi_y(\mathcal{R}_2) = \{0\}$ . Thus  $\ker \varphi_x = \ker \varphi_y = \mathcal{R}_2$ .  $\square$

Set

$$z_{n+1,0} = 0, \quad z_{n+1,i} = x_i, \quad z_{j,0} = y_j \quad \text{for } i \in J_1, j \in J_3. \tag{3.17}$$

Denote

$$J'_1 = \{0\} \cup J_1, \quad J'_3 = J_3 \cup \{n+1\}. \tag{3.18}$$

Let

$$\mathcal{C} = \mathbb{F}[z_{j,i} \mid j \in J'_3, i \in J'_1, (j,i) \neq (n+1,0)] \tag{3.19}$$

be the algebra of polynomials in the variables  $\{z_{j,i} \mid j \in J'_3, i \in J'_1, (j,i) \neq (n+1,0)\}$ . Denote by  $\mathcal{C}_k$  the subspace of homogenous polynomials in  $\mathcal{C}$  with degree  $k$ . Define an associative algebra homomorphism  $\phi : \mathcal{C} \rightarrow \mathcal{A}$  by

$$\phi(z_{j,i}) = x_i x_j - y_i y_j, \quad \phi(x_i) = x_i, \quad \phi(y_j) = y_j \quad i \in J_1, j \in J_3. \tag{3.20}$$

For  $0 < m \in \mathbb{N}$ , we set

$$\widehat{J}_1^m = \{\{i_1, i_2, \dots, i_m\} \mid i_s \in J_1\}, \quad \widehat{J}_3^m = \{\{j_1, j_2, \dots, j_m\} \mid i_t \in J_1\}. \tag{3.21}$$

Let  $\{(j_1, i_1), (j_2, i_2), \dots, (j_r, i_r)\} \subset \mathbb{N}^2$ . If there exist three distinct  $1 \leq k_1, k_2, k_3 \leq r$  such that  $i_{k_1} < i_{k_2} < i_{k_3}$  and  $j_{k_1} < j_{k_2} < j_{k_3}$ , we say that  $\{(j_1, i_1), (j_2, i_2), \dots, (j_r, i_r)\}$  contains a 3-chain  $(j_{k_1}, i_{k_1}) \prec (j_{k_2}, i_{k_2}) \prec (j_{k_3}, i_{k_3})$ . For  $k_1, k_2, k_3 \in \mathbb{N}$ , we define

$$\mathcal{I}_1^x \left[ \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right] = \{s_{1,1}, \dots, s_{1,k_1}\}, \tag{3.22}$$

$$\mathcal{I}_1 \left[ \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right] = \{s_{1,1}, \dots, s_{1,k_1}, i_1, \dots, i_{k_3}\}, \tag{3.23}$$

$$\mathcal{I}_3^y \left[ \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right] = \{s_{2,1}, \dots, s_{2,k_2}\}, \tag{3.24}$$

$$\mathcal{I}_3 \left[ \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right] = \{s_{2,1}, \dots, s_{2,k_2}, j_1, \dots, j_{k_3}\}, \tag{3.25}$$

$$\begin{aligned} & \mathcal{I} \left[ \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right] \\ &= \{(n+1, s_{1,1}), \dots, (n+1, s_{1,k_1}), (j_1, i_1), \dots, (j_{k_3}, i_{k_3}), (s_{2,1}, 0), \dots, (s_{2,k_2}, 0)\}, \end{aligned} \quad (3.26)$$

where

$$\{s_{1,1}, \dots, s_{1,k_1}, i_1, \dots, i_{k_3}\} \in \widehat{J}_1^{k_1+k_3}, \quad \{s_{2,1}, \dots, s_{2,k_2}, j_1, \dots, j_{k_3}\} \in \widehat{J}_3^{k_2+k_3}. \quad (3.27)$$

In the following, we use the lexical order  $\leq$  on  $\mathbb{N}^2$ . Fix  $\mathcal{I}_1 \in \widehat{J}_1^{k_1+k_3}$  and  $\mathcal{I}_3 \in \widehat{J}_3^{k_2+k_3}$ . we define

$$\begin{aligned} G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} &= \left\{ g = \left( \prod_{t_1=1}^{k_1} x_{s_{1,t_1}} \right) \left( \prod_{t_2=1}^{k_2} y_{s_{2,t_2}} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \mid \mathcal{I}_1(g) = \mathcal{I}_1, \right. \\ & \quad \mathcal{I}_3(g) = \mathcal{I}_3, \text{ no 3-chains in } \mathcal{I}(g); s_{1,1} \leq \dots \leq s_{1,k_1}, \\ & \quad \left. s_{2,1} \leq \dots \leq s_{2,k_2}, (j_1, i_1) \leq \dots \leq (j_{k_3}, i_{k_3}) \right\}. \end{aligned} \quad (3.28)$$

Set

$$V_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} = \text{Span}\{G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3}\}. \quad (3.29)$$

**Lemma 3.2.** *The set  $\phi(G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3})$  is a linearly independent set in  $\mathcal{A}$ .*

*Proof.* For convenience, we make the following convention:

$$x_0 = 0, \quad y_{n+1} = 0. \quad (3.30)$$

For  $i_1, i_2 \in J'_1$  with  $i_1 \neq i_2$ , we define an associative algebra endomorphism  $\Gamma_{i_1, i_2}^y$  of  $\mathcal{C}$  by

$$\Gamma_{i_1, i_2}^y(z_{j, i_1}) = z_{j, i_2}, \quad \Gamma_{i_1, i_2}^y(z_{j, i'}) = z_{j, i'} \quad \text{for } j \in J'_3, i_1 \neq i' \in J'_1. \quad (3.31)$$

In particular,

$$\Gamma_{i_1, 0}^y(z_{j, i_1}) = y_j, \quad \Gamma_{i_1, 0}^y(x_{i_1}) = 0 \quad (3.32)$$

by (3.17) and (3.31).

For any  $\mathcal{I} \in (\widehat{J}_1^m \cup \widehat{J}_3^m)$ , we denote by  $\mathbf{c}(\mathcal{I})$  the number of distinct elements in  $\mathcal{I}$ . We want to prove that  $\phi(G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3})$  is linearly dependent by induction on

$$\ell_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} = k_1 + k_2 + k_3 + \mathbf{c}(\mathcal{I}_1) + \mathbf{c}(\mathcal{I}_3). \quad (3.33)$$

First  $\ell_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} \geq 2$  by definition. When  $\ell_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} = 2$ ,  $G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} = \{x_i\}$  for some  $i \in J_1$  or  $\{y_j\}$  for some  $j \in J_3$ . The lemma trivially holds. Assume the lemma holds for  $\ell_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} < l \in \mathbb{N}$ . Consider  $\ell_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} = l$ . Suppose that  $\phi(G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3})$  is linearly dependent. Then there exists  $0 \neq f \in V_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3}$  such that  $\phi(f) = 0$ . Write

$$f = \sum_{s=1}^m a_s f_s, \quad (3.34)$$

where  $0 \neq a_1, a_2, \dots, a_m \in \mathbb{F}$  and  $f_1, f_2, \dots, f_m \in G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3}$  are distinct. If  $\text{g.c.d}(f_1, \dots, f_m) = h \neq 1$ , then  $f_s = g_s h$  for  $s \in \overline{1, m}$ . We have  $f = h(\sum_s a_s g_s)$  and  $0 \neq g = \sum_s a_s g_s \in V_{k'_1, k'_2, k'_3}^{\mathcal{I}'_1, \mathcal{I}'_3}$  with  $\ell_{k'_1, k'_2, k'_3}^{\mathcal{I}'_1, \mathcal{I}'_3} < l$ . By inductual assumption,  $\phi(G_{k'_1, k'_2, k'_3}^{\mathcal{I}'_1, \mathcal{I}'_3})$  is linearly independent, which implies  $\phi(g) \neq 0$ . On the other hand,

$$0 = \phi(f) = \phi(h)\phi(g) \implies \phi(g) = 0 \quad (3.35)$$

because  $\phi(h) \neq 0$  due to  $h$  being a monomial. It is a contradiction. Thus

$$\text{g.c.d}(f_1, \dots, f_m) = 1. \quad (3.36)$$

If  $k_3 = 0$ , then  $f = a\xi$  with  $\xi \in \mathbb{F}[X_{J_1}, Y_{J_3}]$ . Then  $0 = \phi(f) = f$ , which leads to a contradiction. Hence  $k_3 > 0$ . Write


$$f = \sum_{s=1}^m h_s g_s \quad \text{with } h_s \in \mathbb{F}[X_{J_1}, Y_{J_3}], \quad g_s \in Z^{(k_3)} \quad (3.37)$$

(cf. the below of (3.2)). Denote

$$i_0 = \min\{\mathcal{S}_1(g_s) \mid s \in \overline{1, m}\}. \quad (3.38)$$

Due to (3.36), there exists some  $\Gamma_{i_0, 0}^y(f_s) \neq 0$ . For any  $\Gamma_{i_0, 0}^y(f_t) \neq 0$ , we have  $i_0 \notin \mathcal{S}_1(h_t)$ . Write

$$\Gamma_{i_0, 0}^y(f_t) = h'_t g'_t \quad \text{with } h'_t \in \mathbb{F}[X_{J_1}, Y_{J_3}], \quad g'_t \in Z^{(k_3)}. \quad (3.39)$$



**Fig. 1** The case  $i_1 = i'_1$

There exist  $\mathcal{I}_1^* \in \widehat{\mathcal{J}}_1^{k_1+k_3-\iota}$ ,  $\mathcal{I}_3^* \in \widehat{\mathcal{J}}_3^{k_2+k_3}$  and  $0 < \iota \in \mathbb{Z}$  such that

$$\mathcal{S}_1(\Gamma_{i_0, 0}^y(f_t)) = \mathcal{I}_1^*, \quad \mathcal{S}_3(\Gamma_{i_0, 0}^y(f_t)) = \mathcal{I}_3^*, \quad |\mathcal{S}_1(g'_t)| = k_3 - \iota. \quad (3.40)$$

If  $\mathcal{S}(h'_t g'_t)$  contains a 3-chain  $(j_1, i_1) \prec (j_2, i_2) \prec (j_3, i_3)$ , then  $i_1 = 0$ . So  $\mathcal{S}(f_t)$  contains 3-chain  $(j_1, i_0) \prec (j_2, i_2) \prec (j_3, i_3)$  (cf. Fig.1), which contradicts (3.28).

Thus  $\mathcal{S}(h'_t g'_t)$  does not contain 3-chains. Therefore

$$\Gamma_{i_0, 0}^y(f) \in V_{k_1, k_2+\iota, k_3-\iota}^{\mathcal{I}_1^*, \mathcal{I}_3^*}. \quad (3.41)$$

Moreover,

$$\mathbf{c}(\mathcal{I}_1^*) = \mathbf{c}(\mathcal{I}_1) - 1, \quad \mathbf{c}(\mathcal{I}_3^*) = \mathbf{c}(\mathcal{I}_3). \quad (3.42)$$

Hence

$$\ell_{k_1, k_2 + \iota, k_3 - \iota}^{\mathcal{I}_1^*, \mathcal{I}_3^*} = l - 1. \quad (3.43)$$

By inductual assumption,  $\phi(G_{k_1, k_2 + \iota, k_3 - \iota}^{\mathcal{I}_1^*, \mathcal{I}_3^*})$  is linearly independent. Note

$$\phi(\Gamma_{i_0, 0}^y(f)) = \phi(f)|_{x_{i_0}=0, y_{i_0}=1} = 0. \quad (3.44)$$

So

$$\Gamma_{i_0, 0}^y(f) = 0. \quad (3.45)$$

Take  $i_1, i_2 \in \mathcal{I}_1$  such that  $i_1 < i_2$  and there is no  $i' \in \mathcal{I}_1$  satisfying  $i_1 < i' < i_2$ . Then


$$\Gamma_{i_2, i_1}^y(f_s) \neq 0 \quad \text{for } s \in \overline{1, m}. \quad (3.46)$$

Write

$$\Gamma_{i_2, i_1}^y(f_s) = \xi_s \eta_s \quad \text{with } \xi_s \in \mathbb{F}[X_{J_1}, Y_{J_3}], \quad \eta_s \in Z^{k_3}. \quad (3.47)$$

If  $\mathcal{S}(\xi_s \eta_s)$  contains a 3-chain  $(j'_1, i'_1) \prec (j'_2, i'_2) \prec (j'_3, i'_3)$ , then  $i_1 \in \{i'_1, i'_2, i'_3\}$ . When  $i_1 = i'_1$ ,  $\mathcal{S}(f_s)$  contains the 3-chain  $(j'_1, i_2) \prec (j'_2, i'_2) \prec (j'_3, i'_3)$ . In Case  $i_1 = i'_2$ ,  $\mathcal{S}(f_s)$  contains the 3-chain  $(j'_1, i'_1) \prec (j'_2, i_2) \prec (j'_3, i'_3)$  (cf. Fig.2(a)). If  $i_1 = i'_3$ ,  $\mathcal{S}(f_s)$  contains the 3-chain  $(j'_1, i'_1) \prec (j'_2, i'_2) \prec (j'_3, i_2)$  (cf. Fig.2(b)). These contradict (3.28). Thus  $\mathcal{S}(\xi_s \eta_s)$  does not contain a 3-chain. There exist  $0 < \iota' \in \mathbb{Z}$  and  $\mathcal{I}_1^\# \in \widehat{\mathcal{J}}_1^{k_1 + k_3 - \iota'}$  such that

$$\Gamma_{i_2, i_1}^y(f) \in V_{k_1, k_2, k_3}^{\mathcal{I}_1^\#, \mathcal{I}_3}. \quad (3.48)$$



**Fig. 2** The cases  $i_1 = i'_2$  and  $i_1 = i'_3$

In this case,

$$\mathbf{c}(\mathcal{I}_1^\#) = \mathbf{c}(\mathcal{I}_1) - 1. \quad (3.49)$$

So

$$\ell_{k_1, k_2, k_3}^{\mathcal{I}_1^\#, \mathcal{I}_3} = l - 1. \quad (3.50)$$

By inductual assumption,  $\phi(G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3})$  is linearly independent. Note

$$\phi(\Gamma_{i_2, i_1}^y(f)) = \phi(f)|_{x_{i_2}=x_{i_1}, y_{i_2}=y_{i_1}} = 0. \quad (3.51)$$

This yields

$$\Gamma_{i_2, i_1}^y(f) = 0. \quad (3.52)$$

For  $j_1, j_2 \in J'_3$  with  $j_1 > j_2$ , we define an associative algebra endomorphism  $\Gamma_{j_2, j_1}^x$  of  $\mathcal{C}$  by

$$\Gamma_{j_2, j_1}^x(z_{j_2, i}) = z_{j_1, i}, \quad \Gamma_{j_2, j_1}^x(z_{j', i}) = z_{j', i} \quad \text{for } j_2 \neq j' \in J'_3, i \in J'_1. \quad (3.53)$$

In particular,

$$\Gamma_{j_2, n+1}^x(z_{j_2, i}) = x_i, \quad \Gamma_{j_2, n+1}^x(y_{j_2}) = 0 \quad (3.54)$$

by (3.17) and (3.53). Set

$$j_0 = \max\{\mathcal{S}_3(g_s) \mid s \in \overline{1, m}\}. \quad (3.55)$$

Take  $j_1, j_2 \in \mathcal{I}_3$  such that  $j_1 > j_2$  and there is no  $j' \in \mathcal{I}_3$  satisfying  $j_2 < j' < j_1$ . Symmetrically, we have

$$\Gamma_{j_0, n+1}^x(f) = 0 \quad (3.56)$$

and

$$\Gamma_{j_2, j_1}^x(f) = 0. \quad (3.57)$$

Suppose that the set of distinct elements of  $\mathcal{I}_1$  is

$$\Upsilon_1 = \{s_1, s_2, \dots, s_{\ell_1}\} \quad \text{with } s_1 < s_2 < \dots < s_{\ell_1}. \quad (3.58)$$

By (3.45) and (3.52),

$$\Gamma_{s_1, 0}^y(f) = \Gamma_{s_2, s_1}^y(f) = \dots = \Gamma_{s_{\ell_1}, s_{\ell_1-1}}^y(f) = 0. \quad (3.59)$$

Write the set of distinct elements of  $\mathcal{I}_3$  as

$$\Upsilon_3 = \{t_1, t_2, \dots, t_{\ell_3}\} \quad \text{with } t_1 < t_2 < \dots < t_{\ell_3}. \quad (3.60)$$

According to (3.56) and (3.57),

$$\Gamma_{t_1, t_2}^x(f) = \Gamma_{t_2, t_3}^x(f) = \dots = \Gamma_{t_{\ell_3-1}, t_{\ell_3}}^x(f) = \Gamma_{t_{\ell_3}, n+1}^x(f) = 0. \quad (3.61)$$

If  $k_1 = 0$  and  $\Gamma_{t_{\ell_3}, n+1}^x(f) = 0$ , this implies  $f = 0$ , which contradicts  $f \neq 0$ . Thus  $k_1 > 0$ . Symmetrically,  $k_2 > 0$ . Suppose that  $\mathcal{S}(f_\iota)$  does not contain any 2-chain  $(j, 0) \prec (j', s_1)$  with  $j, j' \in \mathcal{I}_3$ . Then

$$j \geq j' \quad \text{for } j \in \mathcal{S}_3^y(f_\iota), (j', s_1) \in \mathcal{S}(f_\iota). \quad (3.62)$$

If  $y_{t_{\ell_3}} \not\mid f_\iota$ , there exists  $(t_{\ell_3}, i) \in \mathcal{S}(f_\iota)$  for some  $s_1 < i \in \mathcal{I}_1$ ; that is  $z_{t_{\ell_3}, i} \mid f_\iota$ . Since  $\mathcal{S}(f_\iota)$  does not contain 3-chain  $(j_1, 0) \prec (t_{\ell_3}, i) \prec (n+1, i')$ , we have

$$i' \leq i \quad \text{for } i' \in \mathcal{S}_1^x(f_\iota). \quad (3.63)$$

By  $\Gamma_{t_{\ell_3}, n+1}^x(f) = 0$ , there exists another  $f_{\iota'}$  such that  $x_i z_{t_{\ell_3}, i'} \mid f_{\iota'}$  with  $i' < i$  and

$$\mathcal{S}_3^y(f_\iota) = \mathcal{S}_3^y(f_{\iota'}). \quad (3.64)$$



Now  $\mathcal{S}(f_{s'})$  contains a 3-chain  $(j_1, 0) \prec (t_{\ell_3}, i') \prec (n+1, i)$ , which contradicts (3.28). Thus  $y_{t_{\ell_3}} | f_{s'}$ .


Assume  $\mathcal{S}(f_{s'})$  contains a 2-chain  $(j, 0) \prec (j', s_1)$  with  $j, j' \in \mathcal{I}_3$ ,  $y_{t_{\ell_3}} \nmid f_{s'}$  and  $z_{t_{\ell_3}, i} | f_{s'}$  for some  $s_1 < i \in \mathcal{I}_1$ . Since  $\mathcal{S}(f_{s'})$  does not contain 3-chain  $(j_1, 0) \prec (t_{\ell_3}, i) \prec (n+1, i')$ , we have

$$i' \leq i \quad \text{for } i' \in \mathcal{S}_1^x(f_{s'}). \quad (3.65)$$

By  $\Gamma_{t_{\ell_3}, n+1}^x(f) = 0$ , there exists another  $f_{s''}$  such that  $x_i z_{t_{\ell_3}, i'} | f_{s''}$  with  $i' < i$  and

$$\mathcal{S}_3^y(f_{s'}) = \mathcal{S}_3^y(f_{s''}). \quad (3.66)$$

Now  $\mathcal{S}(f_{s''})$  contains a 3-chain  $(j_1, 0) \prec (t_{\ell_3}, i') \prec (n+1, i)$ , which contradicts (3.28) (cf. Fig.3). Thus  $y_{t_{\ell_3}} | f_{s''}$ .



**Fig. 3** If  $y_{t_{\ell_3}} \nmid f_{s'}$

Assume  $\mathcal{S}(f_{s'})$  contains a 2-chain  $(j, 0) \prec (j', s_1)$  with  $j, j' \in \mathcal{I}_3$ ,  $y_{t_{\ell_3}} \nmid f_{s'}$  and  $z_{t_{\ell_3}, i} | f_{s'}$  forces  $i = s_1$ . If there exists  $s_1 < i' \in \mathcal{S}_1^x(f_{s'})$ ,  $\mathcal{S}(f_{s'})$  contains a 3-chain  $(j, 0) \prec (j', s_1) \prec (n+1, i')$ , which contradicts (3.28). Therefore,  $\mathcal{S}_1^x(f_{s'}) = \{s_1, s_1, \dots, s_1\}$ , which yields  $\mathcal{S}_1^x(\Gamma_{t_{\ell_3}, n+1}^x(f_{s''})) = \{s_1, s_1, \dots, s_1\}$ . This leads  $\Gamma_{t_{\ell_3}, n+1}^x(f) \neq 0$ , and contradicts (3.61). So  $y_{t_{\ell_3}} | f_{s'}$ . This leads a contradiction to (3.36). Then there does not exist  $0 \neq f \in V_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3}$  such that  $\phi(f) = 0$ , that is,  $\phi(G_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3})$  is linearly independent.  $\square$

Let

$$\mathcal{R}_3 = \left\langle \left( \begin{array}{ccc} z_{j_1, i_1} & z_{j_1, i_2} & z_{j_1, i_3} \\ z_{j_2, i_1} & z_{j_2, i_2} & z_{j_2, i_3} \\ z_{j_3, i_1} & z_{j_3, i_2} & z_{j_3, i_3} \end{array} \middle| i_1, i_2, i_3 \in J'_1, j_1, j_2, j_3 \in J'_3 \right) \right\rangle \quad (3.67)$$

be an ideal of  $\mathcal{C}$ .

**Lemma 3.3.** For  $k \in \mathbb{N}$ , we have  $\ker \phi = \mathcal{R}_3$ .

*Proof.* We take  $\leq$  be the lexical order on  $J'_3 \times J'_1$  (cf. (1.13) and (3.18)). For

$$S_1 = \{(j_1, i_1), \dots, (j_k, i_k)\}, \quad S_2 = \{(j'_1, i'_1), \dots, (j'_k, i'_k)\} \quad (3.68)$$

with  $(j_s, i_s), (j'_s, i'_s) \in J'_3 \times J'_1$ . Since the notion of sets  $\{\dots\}$  is independent of the order of its elements, we may assume

$$(j_1, i_1) \leq \dots \leq (j_k, i_k), \quad (j'_1, i'_1) \leq \dots \leq (j'_k, i'_k). \quad (3.69)$$

Take the lexical order  $\leq$  on  $(J'_3 \times J'_1)^k$  and define  $S_1 \leq S_2$  if

$$((j_1, i_1), \dots, (j_k, i_k)) \leq ((j'_1, i'_1), \dots, (j'_k, i'_k)) \text{ in } (J'_3 \times J'_1)^k. \quad (3.70)$$

Assume the above  $S_1 \leq S_2$  and

$$T_1 = \{(m_1, \ell_1), \dots, (m_\nu, \ell_\nu)\}, \quad T_2 = \{(m'_1, \ell'_1), \dots, (m'_\nu, \ell'_\nu)\} \quad (3.71)$$

with  $(m_t, \ell_t), (m'_t, \ell'_t) \in J'_3 \times J'_1$  also satisfying  $T_1 \leq T_2$ . Then

$$\{S_1, T_1\} \leq \{S_2, T_2\}. \quad (3.72)$$

Set

$$\begin{aligned} \widehat{G}_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} &= \left\{ g = \left( \prod_{t_1=1}^{k_1} x_{s_1, t_1} \right) \left( \prod_{t_2=1}^{k_2} y_{s_2, t_2} \right) \left( \prod_{t_3=1}^{k_3} z_{j_{t_3}, i_{t_3}} \right) \right. \\ &\quad \left. \mid \mathcal{I}_1(g) = \mathcal{I}_1, \mathcal{I}_3(g) = \mathcal{I}_3; s_{1,1} \leq \dots \leq s_{1, k_1}, \right. \\ &\quad \left. s_{2,1} \leq \dots \leq s_{1, k_2}, (j_1, i_1) \leq \dots \leq (j_{k_3}, i_{k_3}) \right\}. \end{aligned} \quad (3.73)$$

Let  $f \in \ker \phi \cap \mathcal{C}_k$  for some  $k \in \mathbb{N}$ . We may assume

$$f = \sum_{s=1}^m a_s f_s \quad \text{with } a_s \in \mathbb{F}, f_s \in \widehat{G}_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3} \quad (3.74)$$

for some fixed  $k_1, k_2, k_3$  satisfying  $k_1 + k_2 + k_3 = k$ , and  $\mathcal{I}_1 \in \widehat{\mathcal{J}}_1^{k_1+k_3}$ ,  $\mathcal{I}_3 \in \widehat{\mathcal{J}}_3^{k_2+k_3}$ .

Suppose some  $\mathcal{I}(f_t)$  contains a 3-chain  $(j_1, i_1) \prec (j_2, i_2) \prec (j_3, i_3)$ . Write  $f_t = z_{j_1, i_1} z_{j_2, i_2} z_{j_3, i_3} g_t$  with  $g_t \in \mathcal{C}_{k-3}$ . Then

$$\begin{aligned} f_t &= z_{j_1, i_1} z_{j_2, i_2} z_{j_3, i_3} g_t \equiv (z_{j_1, i_2} z_{j_2, i_1} z_{j_3, i_3} + z_{j_1, i_3} z_{j_2, i_2} z_{j_3, i_1} \\ &\quad + z_{j_1, i_1} z_{j_2, i_3} z_{j_3, i_2} - z_{j_1, i_2} z_{j_2, i_3} z_{j_3, i_1} - z_{j_1, i_3} z_{j_2, i_1} z_{j_3, i_2}) g_t \pmod{\mathcal{R}_3}. \end{aligned} \quad (3.75)$$

Note

$$\begin{aligned} \mathcal{I}(f_t) &= \{(j_1, i_1), (j_2, i_2), (j_3, i_3), \mathcal{I}(g_t)\} \\ &< \{(j_1, i_2), (j_2, i_1), (j_3, i_3), \mathcal{I}(g_t)\} = \mathcal{I}(z_{j_1, i_2} z_{j_2, i_1} z_{j_3, i_3} g_t), \end{aligned} \quad (3.76)$$

$$\begin{aligned} \mathcal{I}(f_t) &= \{(j_1, i_1), (j_2, i_2), (j_3, i_3), \mathcal{I}(g_t)\} \\ &< \{(j_1, i_3), (j_2, i_2), (j_3, i_1), \mathcal{I}(g_t)\} = \mathcal{I}(z_{j_1, i_3} z_{j_2, i_2} z_{j_3, i_1} g_t), \end{aligned} \quad (3.77)$$

$$\begin{aligned} \mathcal{I}(f_t) &= \{(j_1, i_1), (j_2, i_2), (j_3, i_3), \mathcal{I}(g_t)\} \\ &< \{(j_1, i_1), (j_2, i_3), (j_3, i_2), \mathcal{I}(g_t)\} = \mathcal{I}(z_{j_1, i_1} z_{j_2, i_3} z_{j_3, i_2} g_t), \end{aligned} \quad (3.78)$$

$$\begin{aligned} \mathcal{I}(f_t) &= \{(j_1, i_1), (j_2, i_2), (j_3, i_3), \mathcal{I}(g_t)\} \\ &< \{(j_1, i_2), (j_2, i_3), (j_3, i_1), \mathcal{I}(g_t)\} = \mathcal{I}(z_{j_1, i_2} z_{j_2, i_3} z_{j_3, i_1} g_t), \end{aligned} \quad (3.79)$$

$$\begin{aligned} \mathcal{I}(f_t) &= \{(j_1, i_1), (j_2, i_2), (j_3, i_3), \mathcal{I}(g_t)\} \\ &< \{(j_1, i_3), (j_2, i_1), (j_3, i_2), \mathcal{I}(g_t)\} = \mathcal{I}(z_{j_1, i_3} z_{j_2, i_1} z_{j_3, i_2} g_t). \end{aligned} \quad (3.80)$$

Continue this process for each  $f_s$ , we can prove that there exists  $f' \in V_{k_1, k_2, k_3}^{\mathcal{I}_1, \mathcal{I}_3}$  (cf. (3.29)) such that

$$f \equiv f' \pmod{\mathcal{R}_3}. \quad (3.81)$$

It is easy to verify  $\phi(\mathcal{R}_3) = \{0\}$  (cf. (3.67)). Thus  $0 = \phi(f) = \phi(f')$ . By Lemma 3.2,  $f' = 0$ ; that is  $f \in \mathcal{R}_3$ . Since  $\ker \phi$  is a graded subspace of  $\mathcal{C}$ , we have  $\ker \phi = \mathcal{R}_3$ .  $\square$

## 4 Proof of the Main Theorem

In this section, we prove our main theorem. We define a notion *ord* to single out the powers of  $P$  in  $M_k$ , and find the ideals of the associated varieties by the linear relationship we found in the last section.

For  $f \in M_k = V_k$  (cf. (2.61)), we define the *P-order* of  $f$  in  $M_k$  as

$$\text{ord}_k(f) = \min \{s \in \mathbb{N} \mid f \in \text{Span}\{TN_k, TN(k-r)P^r \mid r = 0, 1, 2, \dots, s\}\}. \quad (4.1)$$

In particular,  $\text{ord}_k(f) = 0 \iff f \in TN_k$ .

There are some properties of *P-order*:

1. If  $f_i \in M_k$ , then

$$\text{ord}_k \left( \sum_{i=1}^m f_i \right) \leq \max \{ \text{ord}_k(f_i) \mid i \in \overline{1, m} \}. \quad (4.2)$$

2. If  $f \in TN_{k-r}P^r$ , then

$$\text{ord}_k(f) \leq r. \quad (4.3)$$

3. If  $f \in M_{k_1}$  and  $k_1 \leq k_2$ , then

$$\text{ord}_{k_1}(f) \geq \text{ord}_{k_2}(f). \quad (4.4)$$

**Lemma 4.1.** *For  $f \in M_k$ , we have*

$$\mathfrak{d}(f) \leq k + \text{ord}_k(f). \quad (4.5)$$

Moreover,  $\mathfrak{d}(f) = k + \text{ord}_k(f)$  if and only if  $f \notin TN_{k-1}$ .

*Proof.* For  $f \in V_k$  with  $\text{ord}_k(f) = r$ , we write

$$f = \sum_{s_r=1}^{\ell_r} g_{r,s_1} f_{r,s_r} + \sum_{s_{r-1}=1}^{\ell_{r-1}} g_{r-1,s_{r-1}} f_{r-1,s_{r-1}} + \dots + f_0, \quad (4.6)$$

where  $g_{i,s_i} \in P^{(i)}$ ,  $f_{i,s_i} \in TN(k-i)$  for  $s_i \in \overline{1, \ell_i}$ , and  $f_0 \in TN_k$ . Then

$$\begin{aligned} \mathfrak{d}(f) &= \max \{ \mathfrak{d}(g_{i,s_i} f_{i,s_i}), f_0 \mid 1 \leq i \leq r, s_i \in \overline{1, \ell_i} \} \\ &= \max \{ \mathfrak{d}(f_{i,s_i}) + 2i, \mathfrak{d}(f_0) \mid 1 \leq i \leq r, s_i \in \overline{1, \ell_i} \} \\ &\leq \max \{ k + i \mid i \in \overline{0, r} \} = k + r. \end{aligned} \quad (4.7)$$

In other words,

$$\mathfrak{d}(f) \leq k + \text{ord}_k(f). \quad (4.8)$$

Note that in Equation (4.7), equality holds if and only if  $\mathfrak{d}(f_0) = k$ . Thus  $\mathfrak{d}(f) = k + \text{ord}_k(f)$  if and only if  $f \notin TN_{k-1}$ .  $\square$

Fix  $1 \leq p \in \mathbb{N}$ , we recall  $\mathfrak{g}^p \subset U(\mathfrak{g})$  with  $\mathfrak{g} = sl(n)$  and  $M_k = U_k(\mathfrak{g})(M_0)$  (cf. (2.11)). Define

$$\mathfrak{k}_{(p)} = \{\xi \in \mathfrak{g}^p \mid \xi(M_k) \subset M_{k+p-1}, k \in \mathbb{N}\}. \quad (4.9)$$

Recall the associated graded  $\mathfrak{g}$ -module  $\overline{M}$  in (1.10) and the annihilator of  $\overline{M}$  in (1.11) is a graded ideal in  $S(\mathfrak{g})$ . Define  $S_{(p)}(\mathfrak{g}) = U_p(\mathfrak{g})/U_{p-1}(\mathfrak{g})$  and

$$I_{(p)} = \{\eta \in S_{(p)}(\mathfrak{g}) \mid \eta(v) = 0 \text{ for } v \in \overline{M}\}. \quad (4.10)$$

Then  $\text{Ann}_{S(\mathfrak{g})}(\overline{M}) = \sum_{p=1}^{\infty} I_{(p)}$  and we are going to compute  $I_{(p)}$  for different values of  $p$ . First, we assume  $|J_1|, |J_2|, |J_3| \geq 2$ . Denote

$$\begin{aligned} \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} &= E_{j_1, i_1} E_{j_2, i_2} E_{j_3, i_3} + E_{j_1, i_2} E_{j_2, i_3} E_{j_3, i_1} + E_{j_1, i_3} E_{j_2, i_1} E_{j_3, i_2} \\ &\quad - E_{j_1, i_2} E_{j_2, i_1} E_{j_3, i_3} - E_{j_1, i_3} E_{j_2, i_2} E_{j_3, i_1} - E_{j_1, i_1} E_{j_2, i_3} E_{j_3, i_2}, \end{aligned} \quad (4.11)$$

$$\Delta_{i_1, i_2}^{j_1, j_2} = E_{j_1, i_1} E_{j_2, i_2} - E_{j_1, i_2} E_{j_2, i_1} \quad (4.12)$$

for  $i_1, i_2, i_3, j_1, j_2, j_3 \in \overline{1, n}$ .

According to (2.1)-(2.3), we have

$$\tilde{\pi}(E_{j_1, i_1}) = -x_{i_1} x_{j_1} - y_{i_1} \partial_{y_{j_1}}, \quad \tilde{\pi}(E_{j_2, i_2}) = x_{j_2} \partial_{x_{i_2}} + y_{i_2} y_{j_2} \quad (4.13)$$

$$\tilde{\pi}(E_{j_3, i_3}) = -x_{i_3} x_{j_3} + y_{i_3} y_{j_3} \quad \text{for } (j_s, i_s) \in L_s. \quad (4.14)$$

(cf. (2.52)). Thus

$$E_{j, i}(M_0) \not\subset M_0, \quad E_{j', i'}(M_k) \subset M_k \quad \text{when } (j, i) \in L, (j', i') \notin L, k \in \mathbb{N} \quad (4.15)$$

and

$$I_{(1)} = \text{Span}\{E_{j, i} \mid (j, i) \notin L\}. \quad (4.16)$$

**Lemma 4.2.** *If  $\ell_1, \ell_2 \leq 0$ ,*

$$I_{(2)} \equiv \left\{ \Delta_{i_1, i'_1}^{j_1, j'_1}, \Delta_{i_2, i'_2}^{j_2, j'_2} \mid (j_1, i_1), (j'_1, i'_1) \in L_1, (j_2, i_2), (j'_2, i'_2) \in L_2 \right\} \pmod{\langle I_{(1)} \rangle}. \quad (4.17)$$

*If  $\ell_1 \leq 0$  and  $\ell_2 > 0$ ,*

$$I_{(2)} \equiv \left\{ \Delta_{i_2, i'_2}^{j_2, j'_2} \mid (j_2, i_2), (j'_2, i'_2) \in L_2 \right\} \pmod{\langle I_{(1)} \rangle}, \quad (4.18)$$

*and*

$$\left\{ \left( \Delta_{i_1, i'_1}^{j_1, j'_1} \right)^{\ell_2+1} \mid (j_1, i_1), (j'_1, i'_1) \in L_1 \right\} \in I_{(2\ell_2+2)}. \quad (4.19)$$

*In the case  $\ell_1 > 0$  and  $\ell_2 \leq 0$ ,*

$$I_{(2)} \equiv \left\{ \Delta_{i_1, i'_1}^{j_1, j'_1} \mid (j_1, i_1), (j'_1, i'_1) \in L_1 \right\} \pmod{\langle I_{(1)} \rangle}, \quad (4.20)$$

*and*

$$\left\{ \left( \Delta_{i_2, i'_2}^{j_2, j'_2} \right)^{\ell_1+1} \mid (j_2, i_2), (j'_2, i'_2) \in L_2 \right\} \in I_{(2\ell_1+2)}. \quad (4.21)$$

*Proof.* Suppose

$$\xi = \sum_{(j,i),(j',i') \in L} a_{j,i;j',i'} E_{j,i} E_{j',i'} \in \mathfrak{k}_{(2)} \quad (4.22)$$

and set

$$v_0 = x_1^{\max\{0,-\ell_1\}} y_n^{\max\{0,-\ell_2\}} x_{n+1}^{\max\{\ell_1,0\}} y_{n+1}^{\max\{\ell_2,0\}} \in M_0. \quad (4.23)$$

Then  $\xi(v_0) \in M_1$  by (4.9). Note that

$$\mathfrak{d}(x_{i_1} x_{j_1}) = \mathfrak{d}(y_{i_2} y_{j_2}) = 1, \quad \mathfrak{d}(x_{i_3} x_{j_3}) = \mathfrak{d}(y_{i_3} y_{j_3}) = 2 \quad (4.24)$$

as multiplication operators on  $M$  for  $(j_s, i_s) \in L_s$  (cf. (2.46)-(2.48)). Hence

$$\text{the coefficient of } x_{i_1} x_{j_1}' x_{j_1} x_{j_1}' v_0 = a_{j_1, i_1; j_1', i_1'} + a_{j_1', i_1'; j_1, i_2} + a_{j_1', i_1; j_1, i_1'} + a_{j_1, i_1'; j_1', i_1} = 0, \quad (4.25)$$

$$\text{the coefficient of } y_{i_2} y_{j_2}' y_{j_2} y_{j_2}' v_0 = a_{j_2, i_2; j_2', i_2'} + a_{j_2', i_2'; j_2, i_2} + a_{j_2', i_2; j_2, i_2'} + a_{j_2, i_2'; j_2', i_2} = 0, \quad (4.26)$$

$$\text{the coefficient of } x_{i_3} x_{j_3} y_{i_3}' y_{j_3}' v_0 = a_{j_3, i_3; j_3', i_3'} + a_{j_3', i_3'; j_3, i_3} = 0, \quad (4.27)$$

$$\text{the coefficient of } x_{i_1} x_{j_1} y_{i_3} y_{j_3} v_0 = a_{j_1, i_1; j_3, i_3} + a_{j_3, i_3; j_1, i_1} = 0, \quad (4.28)$$

$$\text{the coefficient of } x_{i_1} x_{j_1} y_{i_2} y_{j_2} v_0 = a_{j_1, i_1; j_2, i_2} + a_{j_2, i_2; j_1, i_1} = 0, \quad (4.29)$$

$$\text{the coefficient of } y_{i_2} y_{j_2} y_{i_3} y_{j_3} v_0 = a_{j_2, i_2; j_3, i_3} + a_{j_3, i_3; j_2, i_2} = 0 \quad (4.30)$$

for  $(j_s, i_s) \in L_s$  with  $s = 1, 2, 3$ . On the other hand,

$$E_{j,i} E_{j',i'} - E_{j',i'} E_{j,i} \in U_1(\mathfrak{g}) = \mathfrak{g}. \quad (4.31)$$

By Lemma 3.1,

$$\begin{aligned} \xi \in & \sum_{(j_1, i_1), (j_1', i_1') \in L_1; (j_1, i_1) \leq (j_1', i_1')} a_{j_1, i_1; j_1', i_1'} (E_{j_1, i_1} E_{j_1', i_1'} - E_{j_1', i_1'} E_{j_1, i_1}) \\ & + \sum_{(j_2, i_2), (j_2', i_2') \in L_2; (j_2, i_2) \leq (j_2', i_2')} a_{j_2, i_2; j_2', i_2'} (E_{j_2, i_2} E_{j_2', i_2'} - E_{j_2', i_2'} E_{j_2, i_2}) + U_1(\mathfrak{g}). \end{aligned} \quad (4.32)$$

Let  $v_i = T(x^\alpha y^\beta) \in M_i$  with  $\mathfrak{d}(T(x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta) = i$  (cf. (2.51)). If  $j_1 = j_1'$ , we naturally have

$$E_{j_1, i_1} E_{j_1', i_1'} - E_{j_1', i_1'} E_{j_1, i_1} = 0. \quad (4.33)$$

So we assume  $j_1 \neq j_1'$ . Suppose that there exists  $j_3 \in J_3$  such that  $\beta_{j_3} > 0$ . According to (1.7),

$$[\tilde{\Delta}, x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1}'}] = \partial_{y_{j_1}} \partial_{y_{j_1}'} - \partial_{y_{j_1}'} \partial_{y_{j_1}} = 0. \quad (4.34)$$

Assume  $\beta \neq 0$ . Since

$$x_{i_1} T = T x_{i_1}, \quad y_{j_3} T = T y_{j_3} \quad \text{for } i_1 \in J_1, j_3 \in J_3 \quad (4.35)$$

by (1.7) and (2.6), we have

$$\begin{aligned} \Delta_{i_1, i_1'}^{j_1, j_1'}(v_i) &= (E_{j_1, i_1} E_{j_1', i_1'} - E_{j_1', i_1'} E_{j_1, i_1})(v_i) \\ &= (x_{i_1} x_{j_1} + y_{i_1} \partial_{y_{j_1}})(x_{i_1'} x_{j_1'} + y_{i_1'} \partial_{y_{j_1}'}) - (x_{i_1'} x_{j_1} + y_{i_1'} \partial_{y_{j_1}})(x_{i_1} x_{j_1}' + y_{i_1} \partial_{y_{j_1}'}) \\ &= (x_{i_1'} y_{i_1} - x_{i_1} y_{i_1'})(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1}'}) (T(x^\alpha y^\beta)) \\ &= (x_{i_1'} y_{i_1} y_{j_3} - x_{i_1} y_{i_1'} y_{j_3})(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1}'}) (T(x^\alpha y^{\beta - \epsilon_{j_3}})) \\ &= E_{j_3, i_1} ((x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1}'}) T(x^{\alpha + \epsilon_{i_1'}} y^{\beta - \epsilon_{j_3}})) \\ &\quad - E_{j_3, i_1'} ((x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1}'}) T(x^{\alpha + \epsilon_{i_1}} y^{\beta - \epsilon_{j_3}})) \end{aligned} \quad (4.36)$$

Then

$$\tilde{\Delta}((x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})T(x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}})) = (x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})\tilde{\Delta}(T(x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}})) = 0. \quad (4.37)$$

By (1.4), we have  $(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})T(x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}}) \in M$ . Furthermore, according to (2.51),

$$\begin{aligned} & \mathfrak{d} \left( (x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})T(x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}}) \right) \\ &= \mathfrak{d} \left( (x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}} \right) \leq i \end{aligned} \quad (4.38)$$

So

$$(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})T(x^{\alpha+\epsilon_{i_1'}} y^{\beta-\epsilon_{j_3}}) \in M_i. \quad (4.39)$$

Similarly, we have

$$(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})T(x^{\alpha+\epsilon_{i_1}} y^{\beta-\epsilon_{j_3}}) \in M_i. \quad (4.40)$$

Hence

$$\Delta_{i_1, i_1'}^{j_1, j_1'}(v_i) \in M_{i+1}. \quad (4.41)$$

Next we consider  $\beta = 0$ . If  $j_1, j_1' \neq n_1 + 1$ , then

$$\begin{aligned} \Delta_{i_1, i_1'}^{j_1, j_1'}(v_i) &= (x_{i_1'} y_{i_1} - x_{i_1} y_{i_1'})(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})(T(x^\alpha)) \\ &= (x_{i_1'} y_{i_1} - x_{i_1} y_{i_1'})(T(x_{j_1'} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_1'}})(x^\alpha)) = 0 \end{aligned} \quad (4.42)$$

by (4.34). So we may assume  $j_1 = n_1 + 1$  and  $j_1' \neq n_1 + 1$ . Now

$$v_i = T(x^\alpha) = x^\alpha + \sum_{\iota=1}^{\infty} \frac{\alpha_{n_1+1}!(x_{n_1+1}y_{n_1+1})^\iota (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^{\iota-1}}{(\alpha_{n_1+1} + \iota)! \iota!} \sum_{j=n_2+1}^n \alpha_j y_j x^{\alpha-\epsilon_j}. \quad (4.43)$$

Now

$$\begin{aligned} \Delta_{i_1, i_1'}^{j_1, j_1'}(v_i) &= \sum_{\iota=1}^{\infty} \sum_{j=n_2+1}^n \frac{\alpha_{n_1+1}! \alpha_j}{(\alpha_{n_1+1} + \iota)! \iota!} y_j (x_{i_1'} y_{i_1} - x_{i_1} y_{i_1}') \\ &\quad \times (x_{j_1'} \partial_{y_{n_1+1}} - x_{n_1+1} \partial_{y_{j_1'}})(x_{n_1+1} y_{n_1+1})^\iota (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^{\iota-1} x^{\alpha-\epsilon_j} \\ &= \sum_{j=n_2+1}^n \frac{\alpha_j (x_{i_1'} y_j y_{i_1} - x_{i_1} y_j y_{i_1}')}{\alpha_{n_1+1} + 1} \sum_{\iota=1}^{\infty} \frac{(\alpha_{n_1+1} + 1)!}{(\alpha_{n_1+1} + 1 + \iota - 1)! (\iota - 1)!} \\ &\quad \times (x_{n_1+1} y_{n_1+1})^{\iota-1} (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^{\iota-1} x^{\alpha+\epsilon_{n_1+1}+\epsilon_{j_1'}-\epsilon_j} \\ &= \sum_{j=n_2+1}^n \frac{\alpha_j (x_{i_1'} (y_j y_{i_1} - x_j x_{i_1}) + x_{i_1} (x_j x_{i_1}' - y_j y_{i_1}'))}{\alpha_{n_1+1} + 1} T(x^{\alpha+\epsilon_{n_1+1}+\epsilon_{j_1'}-\epsilon_j}) \\ &= \sum_{j=n_2+1}^n \frac{\alpha_j (x_{i_1'} E_{j, i_1} - x_{i_1} E_{j, i_1}')}{\alpha_{n_1+1} + 1} T(x^{\alpha+\epsilon_{n_1+1}+\epsilon_{j_1'}-\epsilon_j}) \\ &= \sum_{j=n_2+1}^n \frac{\alpha_j}{\alpha_{n_1+1} + 1} [E_{j, i_1}(T(x^{\alpha+\epsilon_{i_1}'+\epsilon_{n_1+1}+\epsilon_{j_1'}-\epsilon_j})) \\ &\quad - E_{j, i_1'}(T(x^{\alpha+\epsilon_{i_1}+\epsilon_{n_1+1}+\epsilon_{j_1'}-\epsilon_j}))], \end{aligned} \quad (4.44)$$

where we have used

$$\begin{aligned} & [x_{j'_1}, (\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^{\iota-1}] (x^{\alpha+\epsilon_{n_1+1}-\epsilon_j}) \\ &= (\iota-1)(\tilde{\Delta} + \partial_{x_{n_1+1}} \partial_{y_{n_1+1}})^{\iota-2} \partial_{y_{j'_1}} (x^{\alpha+\epsilon_{n_1+1}-\epsilon_j}) = 0. \end{aligned} \quad (4.45)$$

Since

$$\mathfrak{d}(x_{n_1+1}) = \mathfrak{d}(x_{j'_1}) = 1, \quad \mathfrak{d}(x_j) = 2 \quad (4.46)$$

as operators on  $\mathcal{A}$ , we have

$$\mathfrak{d}(T(x^{\alpha+\epsilon_{i'_1}+\epsilon_{n_1+1}+\epsilon_{j'_1}-\epsilon_j})), \mathfrak{d}(T(x^{\alpha+\epsilon_{i_1}+\epsilon_{n_1+1}+\epsilon_{j'_1}-\epsilon_j})) \leq i. \quad (4.47)$$

Thus

$$\Delta_{i_1, i'_1}^{j_1, j'_1}(v_i) \in M_{i+1}. \quad (4.48)$$

Suppose  $\ell_1, \ell_2 < 0$ , then there always exist  $j_3 \in J_3$  and  $i_1 \in J_1$  such that  $\beta_{j_3} > 0$  and  $\alpha_{i_1} > 0$ . For  $t_1, \dots, t_s \in J_1$ ,  $t'_1, \dots, t'_s \in J_3$  and  $T(x^\alpha y^\beta) \in M_{i-s}$ ,

$$\begin{aligned} & \Delta_{i_1, i'_1}^{j_1, j'_1} [(y_{t_1} y_{t'_1} - x_{t_1} x_{t'_1}) \cdots (y_{t_s} y_{t'_s} - x_{t_s} x_{t'_s}) T(x^\alpha y^\beta)] \\ &= \Delta_{i_1, i'_1}^{j_1, j'_1} [E_{t'_1, t_1} \cdots E_{t'_s, t_s} (T(x^\alpha y^\beta))] \\ &\in E_{t'_1, t_1} \cdots E_{t'_s, t_s} [\Delta_{i_1, i'_1}^{j_1, j'_1} (T(x^\alpha y^\beta))] + U_{s+1}(\mathfrak{g})(T(x^\alpha y^\beta)) \\ &\subset U_s(\mathfrak{g})(M_{i-s+1}) + U_{s+1}(\mathfrak{g})(M_{i-s}) \subset M_{i+1}. \end{aligned} \quad (4.49)$$

By (2.61) and Proposition 2.1, we get

$$\Delta_{i_1, i'_1}^{j_1, j'_1}(M_i) \subset M_{i+1}. \quad (4.50)$$

Hence  $\Delta_{i_1, i'_1}^{j_1, j'_1} \in I_{(2)}$  for each  $(j_1, i_1), (j'_1, i'_1) \in L_1$  (cf. (2.52) and (4.10)). Symmetrically, when there exists  $i_1 \in J_1$  such that  $\alpha_{i_1} > 0$ , we also have  $\Delta_{i_2, i'_2}^{j_2, j'_2} \in I_{(2)}$  for each  $(j_2, i_2), (j'_2, i'_2) \in L_2$ . Note that there always exist  $j_3 \in J_3$  and  $i_1 \in J_1$  such that  $\beta_{j_3} > 0$  and  $\alpha_{i_1} > 0$ , when  $\ell_1, \ell_2 < 0$ . Thus (4.17) holds.

Suppose  $\ell_2 > 0$  and  $\ell_1 \leq 0$ . Take  $\xi$  in (4.22) and  $v_0$  in (4.23). Expressions (4.24)-(4.32) still hold. For each  $j_1 \in J_2$ , we set  $w_{j_1} = y_{j_1}^{\ell_2} x_1^{-\ell_1} \in M_0$ . Recall that  $\xi(w_{j_1}) \in M_1$  (cf. (4.22) and (4.9)). For  $(j_1, i_1), (j'_1, i'_1) \in L_1$ ,

$$\text{the coefficient of } x_{j'_1} x_{i'_1} y_{i_1} y_{j_1}^{\ell_2-1} x_1^{-\ell_1} \text{ in } \xi(w_{j_1}) = \ell_2(a_{j_1, i_1; j'_1, i'_1} + a_{j'_1, i'_1; j_1, i_1}) = 0. \quad (4.51)$$

By (4.22), (4.26)-(4.31) and (4.51),

$$\xi \in \sum_{(j_2, i_2), (j'_2, i'_2) \in L_2; (j_2, i_2) \leq (j'_2, i'_2)} a_{j_2, i_2; j'_2, i'_2} (E_{j_2, i_2} E_{j'_2, i'_2} - E_{j'_2, i'_2} E_{j_2, i_2}) + U_1(\mathfrak{g}). \quad (4.52)$$

Using the same arguments as in (4.34)-(4.50), we can prove (4.18).

Let  $v_i = T(x^\alpha y^\beta) \in M_i$  with  $\mathfrak{d}(T(x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta) = i$  (cf. (2.51)). When  $\sum_{j \in J_3} \beta_j > 0$ , we have shown that  $\Delta_{i_1, i'_1}^{j_1, j'_1}(v_i) \in M_{i+1}$  for  $(j_1, i_1), (j'_1, i'_1) \in L_1$ . Consider  $\sum_{j_3 \in J_3} \beta_{j_3} = 0$ , we have  $\sum_{j_2 \in J_2} \beta_{j_2} \leq \ell_2$ . Then

$$\left( \Delta_{i_1, i'_1}^{j_1, j'_1} \right)^{\ell_2+1} (v_i) = (x_{i'_1} y_{i_1} - x_{i_1} y_{i'_1})^{\ell_2+1} (x_{j'_1} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j'_1}})^{\ell_2+1} (T(x^\alpha y^\beta)) = 0. \quad (4.53)$$

Hence  $\left(\Delta_{i_1, i'_1}^{j_1, j'_1}\right)^{\ell_2+1} (v_i) \in M_{i+2\ell_2+1}$ . Using the same arguments as in (4.49)-(4.50), we can prove (4.19). Similarly, when  $\ell_1 > 0$  and  $\ell_2 \leq 0$ , (4.20) and (4.21) holds.  $\square$

Denote

$$\bar{I}_{(2)} = \left\{ \Delta_{i_1, i'_1}^{j_1, j'_1}, \Delta_{i_2, i'_2}^{j_2, j'_2} \mid (j_1, i_1), (j'_1, i'_1) \in L_1, (j_2, i_2), (j'_2, i'_2) \in L_2 \right\}. \quad (4.54)$$

**Lemma 4.3.** *For each  $p \geq 3$ ,  $\ell_1 \leq 0$  or  $\ell_2 < 0$ , we have*

$$I_{(p)} \equiv \langle \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \mid i_1, i_2, i_3 \in J_1 \cup J_2; j_1, j_2, j_3 \in J_2 \cup J_3 \rangle \pmod{\langle I_{(1)}, \bar{I}_{(2)} \rangle}. \quad (4.55)$$

*Proof.* Suppose  $p \geq 3$ , we set

$$\xi = \sum_{l=1}^m c_l \left( \prod_{s=1}^{k_{1,l}} E_{j_{1,l,s}, i_{1,l,s}} \prod_{t=1}^{k_{2,l}} E_{j_{2,l,t}, i_{2,l,t}} \prod_{r=1}^{k_{3,l}} E_{j_{3,l,r}, i_{3,l,r}} \right) \in \mathfrak{k}_{(p)} \quad (4.56)$$

(cf. (4.9)), where  $k_{1,l}, k_{2,l}, k_{3,l} \in \mathbb{N}$  satisfying  $k_{1,l} + k_{2,l} + k_{3,l} = p$ , and

$$(j_{1,l,s}, i_{1,l,s}) \in L_1, (j_{2,l,t}, i_{2,l,t}) \in L_2, (j_{3,l,r}, i_{3,l,r}) \in L_3, \quad (4.57)$$

$$c_l \in \mathbb{F} \quad \text{for each } l \in \overline{1, m}, \quad s \in \overline{1, k_{1,l}}, \quad t \in \overline{1, k_{2,l}}, \quad r \in \overline{1, k_{3,l}}. \quad (4.58)$$

For any  $v_0 \in M_0$ , we consider the partial sum  $h = \sum_{l=1}^m h_l$  of highest degree in  $\xi(v_0)$ , where

$$h_l = (-1)^{k_{1,l}} c_l \left( \prod_{s=1}^{k_{1,l}} x_{j_{1,l,s}} x_{i_{1,l,s}} \prod_{t=1}^{k_{2,l}} y_{j_{2,l,t}} y_{i_{2,l,t}} \prod_{r=1}^{k_{3,l}} (y_{j_{3,l,r}} y_{i_{3,l,r}} - x_{j_{3,l,r}} x_{i_{3,l,r}}) \right) v_0. \quad (4.59)$$

We will show that  $h = 0$  can be derived from  $\xi(v_0) \in M_{p-1}$ .

Suppose that the set of distinct elements of  $\{k_{3,l} \mid l \in \overline{1, m}\}$  is  $\{j_1, \dots, j_N\}$ , and  $j_1 < j_2 < \dots < j_N$ . For  $\iota \in \overline{1, N}$ , we set

$$\bar{h}_\iota = \sum_{l \in \overline{1, m}: k_{3,l} = j_\iota} h_l. \quad (4.60)$$

Suppose that  $\bar{h}_\iota \neq 0$  for some  $\iota \in \overline{1, N}$ . According to (2.48):

$$\mathfrak{d}(\xi(v_0)) \geq \mathfrak{d}(\bar{h}_\iota) = p + j_\iota. \quad (4.61)$$

Denote the degree in  $\{x_{i_3} \mid i_3 \in J_3\}$  as  $\deg_{x_{J_3}}(\xi(v_0))$  and the degree in  $\{y_{i_1} \mid i_1 \in J_1\}$  as  $\deg_{y_{J_1}}(\xi(v_0))$ . For  $\iota' \in \mathbb{N}$  and  $f = \sum_{i=1}^{m'} f_i \in \mathcal{A}$ , where  $f_i$  are distinct monomials in  $\mathcal{A}$ , we define

$$\mathfrak{r}(f, \iota') = \min\{\deg_{x_{J_3}}(f_i) + \deg_{y_{J_1}}(f_i) \mid \mathfrak{d}(f_i) = \iota', i \in \overline{1, m'}\}. \quad (4.62)$$

Then we have

$$\mathfrak{r}(\xi(v_0), p + j_\iota) \leq \mathfrak{r}(\bar{h}_\iota, \mathfrak{d}(\bar{h}_\iota)) = j_\iota. \quad (4.63)$$

Since  $\xi(v_0) \in M_{p-1}$  and  $\xi(v_0) \notin TN_{p-1}$ , according to Lemma 4.1, we have

$$\mathfrak{d}(\xi(v_0)) = \text{ord}_{p-1}(\xi(v_0)) + p - 1 \Rightarrow \text{ord}_{p-1}(\xi(v_0)) \geq 1 + j_\iota. \quad (4.64)$$



Recall (4.6) and (2.1). For  $r = \text{ord}_{p-1}(\xi(v_0)) \geq 1 + j_\ell$  in this case, we can write

$$\xi(v_0) = \sum_{s_r=1}^{\ell_r} g_{r,s_r} f_{r,s_r} + \sum_{s_{r-1}=1}^{\ell_{r-1}} g_{r-1,s_{r-1}} f_{r-1,s_{r-1}} + \cdots + f_0, \quad (4.65)$$

where  $g_{i,s_i} \in P^{(i)}$ ,  $f_{i,s_i} \in TN(p-1-i)$  for  $s_i \in \overline{1, \ell_i}$ , and  $f_0 \in TN_{p-1}$ . Note that for  $s_i \in \overline{1, \ell_i}$  and  $i \in \overline{1, r}$ ,

$$\mathfrak{d}(g_{i,s_i} f_{i,s_i}) = p-1-i+2i = p-1+i \quad (4.66)$$

by (2.60). Since  $\overline{h}_\ell \neq 0$  and  $\mathfrak{d}(\overline{h}_\ell) = p + j_\ell$ , we have

$$\sum_{s_{j_\ell+1}=1}^{\ell_{j_\ell+1}} g_{j_\ell+1,s_{j_\ell+1}} f_{j_\ell+1,s_{j_\ell+1}} \neq 0. \quad (4.67)$$

Therefore,

$$\mathfrak{r}(\xi(v_0), p + j_\ell) = \mathfrak{r} \left( \sum_{s_{j_\ell+1}=1}^{\ell_{j_\ell+1}} g_{j_\ell+1,s_{j_\ell+1}} f_{j_\ell+1,s_{j_\ell+1}}, p + j_\ell \right) \geq j_\ell + 1, \quad (4.68)$$

which contradicts (4.63). Hence,  $\overline{h}_\ell = 0$  for  $w \in \overline{1, N}$ . Therefore  $h = 0$ .

By considering different monomials in  $X_{J_2}$  and  $Y_{J_2}$ , we may assume that  $k_{1,l}, k_{2,l}, k_{3,l}$  in (4.59) are independent of  $l \in \overline{1, m}$ ; that is,  $k_{1,l} = k_1, k_{2,l} = k_2, k_{3,l} = k_3$  for  $l \in \overline{1, m}$ . Thus (4.59) yields

$$0 = \sum_{l=1}^m c_l \left( \prod_{s=1}^{k_1} x_{i_{1,l},s} \prod_{t=1}^{k_2} y_{j_{2,l},t} \prod_{r=1}^{k_3} (y_{j_{3,l},r} y_{i_{3,l},r} - x_{j_{3,l},r} x_{i_{3,l},r}) \right) = h'. \quad (4.69)$$

According to (3.17),

$$\phi \left( \sum_{l=1}^m c_l \left( \prod_{s=1}^{k_1} z_{n+1,i_{1,l},s} \prod_{t=1}^{k_2} z_{j_{2,l},t,0} \prod_{r=1}^{k_3} z_{j_{3,l},r,i_{3,l},r} \right) \right) = h' = 0. \quad (4.70)$$

By Lemma 3.3, we have

$$\sum_{l=1}^m c_l \left( \prod_{s=1}^{k_1} z_{n+1,i_{1,l},s} \prod_{t=1}^{k_2} z_{j_{2,l},t,0} \prod_{r=1}^{k_3} z_{j_{3,l},r,i_{3,l},r} \right) \in \mathcal{R}_3. \quad (4.71)$$

Let  $i_1, i_2, i_3 \in J'_1$  and  $j_1, j_2, j_3 \in J'_3$  (cf. (3.18)) such that  $i_1 < i_2 < i_3$  and  $j_1 < j_2 < j_3$ . Denote

$$\begin{aligned} \Lambda_{i_1, i_2, i_3}^{j_1, j_2, j_3} = & z_{j_1, i_1} z_{j_2, i_2} z_{j_3, i_3} + z_{j_1, i_2} z_{j_2, i_3} z_{j_3, i_1} + z_{j_1, i_3} z_{j_2, i_1} z_{j_3, i_2} \\ & - z_{j_1, i_1} z_{j_2, i_3} z_{j_3, i_2} - z_{j_1, i_2} z_{j_2, i_1} z_{j_3, i_3} - z_{j_1, i_3} z_{j_2, i_2} z_{j_3, i_1} \end{aligned} \quad (4.72)$$

Then we can write

$$h' = \phi \left( \sum_{s=1}^{m'} \Lambda_{i_1, s, i_2, s, i_3, s}^{j_1, s, j_2, s, j_3, s} g_s \right), \quad (4.73)$$

where  $g_s \in \mathcal{C}_{p-3}$ ,  $j_{1,s}, j_{2,s}, j_{3,s} \in J'_3$  and  $i_{1,s}, i_{2,s}, i_{3,s} \in J'_1$ .

Consider the correspondence between the terms in  $h'$  and the terms in  $\xi$ . When  $j_{1,s}, j_{2,s}, j_{3,s} \in J_3$  and  $i_{1,s}, i_{2,s}, i_{3,s} \in J_1$ ,  $\phi(\Lambda_{i_{1,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j_{3,s}})$  corresponds to  $\Delta_{i_{1,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j_{3,s}}$ . Note  $x_{j_{1,l,t}} x_{j_{1,l,t'}} = x_{j_{1,l,t'}} x_{j_{1,l,t}}$ , but  $E_{j_{1,l,t}, i_{1,l,t}} E_{j_{1,l,t'}, i_{1,l,t'}}$  may not be equal to  $E_{j_{1,l,t'}, i_{1,l,t}} E_{j_{1,l,t}, i_{1,l,t}}$ . So  $\prod_{t=1}^{k_{1,l}} z_{n+1, j_{1,l,t}}$  does not only correspond to  $\prod_{t=1}^{k_{1,l}} E_{j_{1,l,t}, i_{1,l,t}}$ , but also corresponds to  $\prod_{t=1}^{k_{1,l'}} E_{j_{1,l',t}, i_{1,l,t}}$  with

$$\{j_{1,l,t} \mid t \in \overline{1, k_1}\} = \{j_{1,l',t} \mid t \in \overline{1, k_1}\} = \mathcal{S}_1^x \left( \prod_{t=1}^{k_1} x_{j_{1,l,t}} \right). \quad (4.74)$$

(cf.(3.22)). According to Lemma 3.1,  $\prod_{t=1}^{k_1} z_{n+1, j_{1,l,t}}$  corresponds to

$$\prod_{t=1}^{k_1} E_{j_{1,l,t}, i_{1,l,t}} + \left\langle \Delta_{i_{1,l'}}^{j_{1,l'}} \mid (j_1, i_1), (j'_1, i'_1) \in L_1 \right\rangle \quad \text{in } S(\mathfrak{g}). \quad (4.75)$$

Similarly,  $\prod_{t=1}^{k_2} z_{j_{2,l,t}, 0}$  corresponds to

$$\prod_{t=1}^{k_2} E_{j_{2,l,t}, i_{2,l,t}} + \left\langle \Delta_{i_{2,l'}}^{j_{2,l'}} \mid (j_2, i_2), (j'_2, i'_2) \in L_2 \right\rangle \quad \text{in } S(\mathfrak{g}). \quad (4.76)$$

Fix  $y_{i'}$  in  $\prod_{t=1}^{k_2} y_{i_{2,l,t}}$  (cf. (4.59)). Then  $\Lambda_{0, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j_{3,s}} y_{i'}$  corresponds to  $\Delta_{i'_{2,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j_{3,s}}$  in the sense as (4.76) when  $i_{2,s}, i_{3,s} \in J_1$  and  $j_{1,s}, j_{2,s}, j_{3,s} \in J_3$ . Fix  $x_{j'}$  in  $\prod_{s=1}^{k_{1,l}} x_{j_{1,l,s}}$  (cf. (4.59)). Similarly,  $\Lambda_{i_{1,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, n+1} x_{j'}$  corresponds to  $\Delta_{i_{1,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j'}$  in the sense as (4.75) when  $i_{1,s}, i_{2,s}, i_{3,s} \in J_1$  and  $j_{1,s}, j_{2,s} \in J_3$ .

When  $j_1, i_3 \in J_2$ ,  $j_2, j_3 \in J_3$  and  $i_1, i_2 \in J_1$ , we recall (4.11) and have

$$\begin{aligned} \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} &\equiv E_{j_1, i_1} E_{j_2, i_2} E_{j_3, i_3} + E_{j_1, i_2} E_{j_2, i_3} E_{j_3, i_1} \\ &\quad - E_{j_1, i_2} E_{j_2, i_1} E_{j_3, i_3} - E_{j_1, i_1} E_{j_2, i_3} E_{j_3, i_2} \pmod{\langle I_{(1)} \rangle} \end{aligned} \quad (4.77)$$

as elements in  $S(\mathfrak{g})$ . Then  $\Lambda_{0, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, n+1} x_{i'} y_{j'}$  corresponds to  $\Delta_{i'_{2,s}, i_{2,s}, i_{3,s}}^{j_{1,s}, j_{2,s}, j'}$  in the sense as (4.77) when  $j_{2,s}, j_{3,s} \in J_3$  and  $i_{1,s}, i_{2,s} \in J_1$ . Hence

$$\xi \in \langle \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \mid j_1, j_2, j_3 \in J_2 \cup J_3, i_1, i_2, i_3 \in J_1 \cup J_2 \rangle + \langle I_{(1)}, \bar{I}_{(2)} \rangle \quad (4.78)$$

Next we want to verify that the above  $\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)}$  for  $j_1 < j_2 < j_3$  and  $i_1 < i_2 < i_3$ , which can always be obtained by re-indexing up to a sign. We divide it into six cases.

(1)  $j_1 \in J_2$  and  $i_3 \in J_2$  do not occur at the same time,  $j_1, j_2, j_3 \in J_2 \cup J_3$ ,  $i_1, i_2, i_3 \in J_1 \cup J_2$ .

In this case, we have

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} |_{\mathcal{A}} = 0. \quad (4.79)$$

For example, for  $i_1, i_2 \in J_1$ ,  $i_3 \in J_2$ ,  $j_1, j_2, j_3 \in J_3$ ,

$$E_{j_1, i_1} E_{j_2, i_2} E_{j_3, i_3} |_{\mathcal{A}} = (y_{j_1} y_{i_1} - x_{j_1} x_{i_1})(y_{j_2} y_{i_2} - x_{j_2} x_{i_2})(x_{j_3} \partial_{x_{i_3}} + y_{j_3} y_{i_3}) \quad (4.80)$$

Define an isomorphism  $\nu$  from  $\mathbb{F}[X_{J_1 \cup J_3}, Y_{\overline{1, n}}, \partial_{x_{n_1+1}}, \dots, \partial_{x_{n_2}}]$  to  $\mathcal{A}$  by

$$\nu(\partial_{x_{i_3}}) = -x_{i_3}, \quad \nu(x_{j'}) = x_{j'}, \quad \nu(y_j) = y_j \quad (4.81)$$

for  $i_3 \in J_2$ ,  $j \in \overline{1, n}$ ,  $j' \in J_1 \cup J_3$ . Then

$$\nu((y_{j_1}y_{i_1} - x_{j_1}x_{i_1})(y_{j_2}y_{i_2} - x_{j_2}x_{i_2})(x_{j_3}\partial_{x_{i_3}} + y_{j_3}y_{i_3})) \quad (4.82)$$

$$= (y_{j_1}y_{i_1} - x_{j_1}x_{i_1})(y_{j_2}y_{i_2} - x_{j_2}x_{i_2})(y_{j_3}y_{i_3} - x_{j_3}x_{i_3}) \quad (4.83)$$

We have already verified  $\nu(\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} |_{\mathcal{A}}) = 0$  in the proof of Lemma 3.3. So (4.79) holds.

(2)  $j_1, i_3 \in J_2$ ,  $j_2, j_3 \in J_3$  and  $i_1, i_2 \in J_1$ .

Let  $v_i = T(x^\alpha y^\beta) \in M_i$  with  $\mathfrak{d}(T(x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta) = i$ . Then

$$\begin{aligned} & \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(v_i) \\ & \in (E_{j_1, i_1} E_{j_2, i_2} E_{j_3, i_3} + E_{j_1, i_2} E_{j_2, i_3} E_{j_3, i_1} - E_{j_1, i_2} E_{j_2, i_1} E_{j_3, i_3} - E_{j_1, i_1} E_{j_2, i_3} E_{j_3, i_2})(v_i) + M_{i+2} \\ & \subset (x_{i_1}y_{i_2} - x_{i_2}y_{i_1})(x_{j_2}y_{j_3} - x_{j_3}y_{j_2})(x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i) + M_{i+2} \\ & \subset (E_{j_3, i_1} E_{j_2, i_2} - E_{j_3, i_2} E_{j_2, i_1})(x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i) + M_{i+2} \end{aligned} \quad (4.84)$$

According to (1.7), we have

$$[\tilde{\Delta}, x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}}] = 0 \quad (4.85)$$

So

$$\tilde{\Delta}((x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i)) = 0; \quad (4.86)$$

that is

$$((x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i)) \in M. \quad (4.87)$$

Furthermore,

$$\mathfrak{d}((x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i)) \leq i \quad (4.88)$$

Hence

$$(x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i) \in M_i. \quad (4.89)$$

Therefore,

$$\begin{aligned} & \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(v_i) \\ & \in (E_{j_3, i_1} E_{j_2, i_2} - E_{j_3, i_2} E_{j_2, i_1})((x_{j_1}\partial_{x_{i_3}} - y_{i_3}\partial_{y_{j_1}})(v_i)) + M_{i+2} \subset M_{i+2}. \end{aligned} \quad (4.90)$$

Moreover, we can prove as (4.49) that

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(M_i) \in M_{i+2} \quad (4.91)$$

for each  $i \in \mathbb{N}$ , thus

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)}. \quad (4.92)$$

(3)  $j_1, j_2, i_3 \in J_2$ ,  $j_3 \in J_3$  and  $i_1, i_2 \in J_1$ .

Recall (4.11) and  $v_i = T(x^\alpha y^\beta) \in M_i$  with  $\mathfrak{d}(T(x^\alpha y^\beta)) = \mathfrak{d}(x^\alpha y^\beta) = i$ . In this case,

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(v_i) \equiv \Delta_{i_1, i_2}^{j_1, j_2} E_{j_3, i_3}(v_i) \pmod{\langle I_{(1)} \rangle}. \quad (4.93)$$

Then

$$\begin{aligned}
& \Delta_{i_1, i_2}^{j_1, j_2} E_{j_3, i_3}(v_i) \\
&= (x_{i_2} y_{i_1} - x_{i_1} y_{i_2})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(x_{j_3} \partial_{x_{i_3}} + y_{j_3} y_{i_3})(v_i) \\
&= (x_{i_2} x_{j_3} y_{i_1} - x_{i_1} x_{j_3} y_{i_2})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}}) \partial_{x_{i_3}}(v_i) + \\
&\quad (x_{i_2} y_{i_1} y_{j_3} - x_{i_1} y_{i_2} y_{j_3})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}}) y_{i_3}(v_i) \\
&= (-E_{j_3, i_2} y_{i_1} + E_{j_3, i_1} y_{i_2})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}}) \partial_{x_{i_3}}(v_i) + \\
&\quad (E_{j_3, i_1} x_{i_2} - E_{j_3, i_2} x_{i_1})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}}) y_{i_3}(v_i) \\
&= E_{j_3, i_1}(y_{i_2} \partial_{x_{i_3}} + x_{i_2} \partial_{y_{i_3}})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(v_i) \\
&\quad - E_{j_3, i_2}(y_{i_1} \partial_{x_{i_3}} + x_{i_1} \partial_{y_{i_3}})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(v_i)
\end{aligned} \tag{4.94}$$

According to (1.7),

$$[\tilde{\Delta}, x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}}] = 0, \tag{4.95}$$

and

$$[\tilde{\Delta}, y_{i_1} \partial_{x_{i_3}} + x_{i_1} \partial_{y_{i_3}}] = 0, \tag{4.96}$$

Thus

$$\tilde{\Delta}((y_{i_2} \partial_{x_{i_3}} + x_{i_2} \partial_{y_{i_3}})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(v_i)) = 0 \tag{4.97}$$

Since

$$\mathfrak{d}((y_{i_2} \partial_{x_{i_3}} + x_{i_2} \partial_{y_{i_3}})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(v_i)) \leq i + 1, \tag{4.98}$$

we have

$$(y_{i_2} \partial_{x_{i_3}} + x_{i_2} \partial_{y_{i_3}})(x_{j_2} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j_2}})(v_i) \in M_{i+1}. \tag{4.99}$$

Therefore,

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(v_i) \in \Delta_{i_1, i_2}^{j_1, j_2} E_{j_3, i_3}(v_i) + M_{i+2} \subset M_{i+2}. \tag{4.100}$$

Furthermore, we can prove as (4.49) that

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}(M_i) \in M_{i+2} \tag{4.101}$$

for  $i \in \mathbb{N}$ . Then

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)}. \tag{4.102}$$

(4)  $j_1, i_2, i_3 \in J_2$ ,  $j_2, j_3 \in J_3$  and  $i_1 \in J_1$ .

It is symmetrical to the case (3), and so (4.102) still holds.

(5)  $j_1, j_2 \in J_2$ ,  $j_3 \in J_2 \cup J_3$  and  $i_1 \in J_1 \cup J_2$ ,  $i_2, i_3 \in J_2$ .

According to (4.11) and (4.16),

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in \langle I_{(1)} \rangle. \tag{4.103}$$

Naturally,

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)}. \tag{4.104}$$

(6)  $i_1, i_2, i_3 \in J_2$  and  $j_1, j_2, j_3 \in J_2 \cup J_3$ , or  $i_1, i_2, i_3 \in J_1 \cup J_2$  and  $j_1, j_2, j_3 \in J_2$ .

When  $i_1, i_2, i_3 \in J_2$ ,

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} = E_{j_3, i_1} \Delta_{i_2, i_3}^{j_1, j_2} - E_{j_3, i_2} \Delta_{i_1, i_3}^{j_1, j_2} + E_{j_3, i_3} \Delta_{i_1, i_2}^{j_1, j_2} \in \langle I_{(1)} \rangle. \quad (4.105)$$

Then

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)}. \quad (4.106)$$

The case of  $j_1, j_2, j_3 \in J_2$  is similar.

In summary,

$$\Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \in I_{(3)} \quad \text{for } i_1, i_2, i_3 \in J_1 \cup J_2, j_1, j_2, j_3 \in J_2 \cup J_3. \quad (4.107)$$

According to (4.78), we have proved that (4.55) holds. This completes the proof of the lemma.  $\square$

In order to calculate the associated variety of the irreducible module  $\mathcal{H}_{\langle \ell_1, \ell_2 \rangle}$  of all the cases in Proposition 1, we consider  $\ell_1, \ell_2 > 0$  and  $n_1 < n_2 = n$ . First we take

$$M_0 = \text{Span} \left\{ TN \begin{pmatrix} 0 & \ell_1 & 0 \\ k_{21} & k_{22} & 0 \end{pmatrix} \mid k_{21} + k_{22} = \ell_2 \right\}. \quad (4.108)$$

In particular, when  $n_1 + 1 = n_2 = n$ ,

$$M_0 = \text{Span} \left\{ T(x_n^{\ell_1} y^\beta) \mid x_n^{\ell_1} y^\beta \in \mathcal{A}, \sum_{s=1}^{n-1} \beta_s = \ell_2, \beta_n = 0 \right\}. \quad (4.109)$$

Set

$$\begin{aligned} V'_k &= \text{Span} \left\{ T(x^\alpha y^\beta) \mid x^\alpha y^\beta \in \mathcal{A}_{\langle \ell_1, \ell_2 \rangle}, \sum_{t=1}^{n_1} \alpha_t \leq k, \sum_{s=1}^n \beta_s = \ell_2, \alpha_{n_1+1} \beta_{n_1+1} = 0 \right\} \\ &= \text{Span} \left\{ TN \begin{pmatrix} t & \ell_1 + t & 0 \\ k_{21} & k_{22} & 0 \end{pmatrix} \mid 0 \leq t \leq k, k_{21} + k_{22} = \ell_2 \right\}. \end{aligned} \quad (4.110)$$

In particular, when  $n_1 + 1 = n_2 = n$ ,

$$V'_k = \text{Span} \left\{ T(x^\alpha y^\beta) \mid x^\alpha y^\beta \in \mathcal{A}_{\langle \ell_1, \ell_2 \rangle}, \sum_{t=1}^{n-1} \alpha_t \leq k, \sum_{s=1}^{n-1} \beta_s = \ell_2 \right\} \quad (4.111)$$

Define  $\mathfrak{d}' : \mathcal{A}_{\langle \ell_1, \ell_2 \rangle} \rightarrow \mathbb{N}$  as follows

$$\mathfrak{d}'(x^\alpha y^\beta) = \sum_{i \in J_1} \alpha_i, \quad (4.112)$$

$$\mathfrak{d}'(\lambda f) = \mathfrak{d}'(f), \quad \forall \lambda \in \mathbb{F} \setminus \{0\}, \quad (4.113)$$

$$\mathfrak{d}' \left( \sum_{i=1}^m f_i \right) = \max\{\mathfrak{d}'(f_1), \dots, \mathfrak{d}'(f_m)\}, \text{ where } f_i \text{ are monomials.} \quad (4.114)$$

Note that  $\mathfrak{d}'$  is different from  $\mathfrak{d}$  (cf. (2.46)), in order to ensure that  $\mathfrak{d}'(M_0) = \{0\}$ .

In this case,  $L = L_1 = J_2 \times J_1$  as (2.52). Similar to (2.48)-(2.53), the following properties of  $\mathfrak{d}'$  still hold:

$$\mathfrak{d}'(E_{i,j}(f)) \leq \begin{cases} \mathfrak{d}'(f) + 1 & \text{if } (i,j) \in L, \\ \mathfrak{d}'(f) & \text{if } (i,j) \notin L. \end{cases} \quad (4.115)$$

For each  $i \in \mathbb{N}$ ,

$$\mathfrak{d}'\left(\left(x_{n_1+1}y_{n_1+1}\right)^i \left(\tilde{\Delta} + \partial_{x_{n_1+1}}\partial_{y_{n_1+1}}\right)^i (x^\alpha y^\beta)\right) = \mathfrak{d}'(x^\alpha y^\beta) \text{ or } 0, \quad (4.116)$$

where  $\alpha_{n_1+1}\beta_{n_1+1} = 0$ . Then

$$\mathfrak{d}'(T(x^\alpha y^\beta)) = \mathfrak{d}'(x^\alpha y^\beta), \text{ where } \alpha_{n_1+1}\beta_{n_1+1} = 0. \quad (4.117)$$

**Lemma 4.4.** For  $k \in \mathbb{N}$ ,  $M_k = U_k(sl(n))(M_0) = V'_k$  holds.

*Proof.* We prove  $M_i = V'_i$  for any  $i \in \mathbb{N}$  by induction. According to (4.110),  $M_0 = V'_0$  holds. Suppose that  $M_i = V'_i$  with  $i \leq k-1$ . Consider  $i = k$ . Since (4.115) holds, we can prove that  $M_k \subset V'_k$  as (2.62)-(2.67) with  $V_k$  replaced by  $V'_k$ . Next we want to prove  $V'_k \subset M_k$ . Let  $f = T(x^\alpha y^\beta) \in V'_k$  with  $x^\alpha y^\beta \in \mathcal{A}_{(\ell_1, \ell_2)}$ ,  $\sum_{t=1}^{n_1} \alpha_t \leq k$ ,  $\sum_{s=1}^n \beta_s = \ell_2$ ,  $\alpha_{n_1+1}\beta_{n_1+1} = 0$ . When  $n_1 + 1 < n_2$ , we suppose  $\sum_{n_1+2 \leq i_2 \leq n} \alpha_{i_2} > 0$ . Then there exists  $i_1 \in J_1$  such that  $\alpha_{i_1} > 0$ . According to (2.7), we have

$$f = -E_{i_2, i_1}(T(x^{\alpha - \epsilon_{i_1} - \epsilon_{i_2}} y^\beta)) - T(\partial_{y_{i_2}} y^{\beta + \epsilon_{i_1}}) \in M_k. \quad (4.118)$$

On the other hand, when  $\sum_{n_1+2 \leq i_2 \leq n} \alpha_{i_2} = 0$ , we write

$$x^\alpha y^\beta = gh_1 h_2 x_{n_1+1}^{\ell_1+k} \quad (4.119)$$

with  $g \in X_{J_1}^k$ ,  $h_1 = \prod_{t=1}^m y_{s_t}^{\beta_{s_t}} \in Y_{J_1}^{k_{21}}$ ,  $h_2 \in Y_{J_2 \setminus \{n_1+1\}}^{k_{22}}$ ,  $\beta_{s_t} \neq 0$  for each  $t \in \overline{1, m}$ . In particular, when  $n_1 + 1 = n_2$ , we have  $h_2 = 1$  and  $k_{21} = \ell_2$  by (4.111). We write  $g = x_{i_0} g'$  for some  $i_0 \in J_1$ , and denote  $v'_0 = g' x_n^{\ell_1+k} h_1 h_2$ . Then  $f = T(x_{i_0} v'_0)$ . For  $t \in \overline{1, m}$ , we set

$$v_t = T\left(\frac{v'_0 x_{s_t} y_{i_0}}{y_{s_t}}\right) \in V'_k, \quad (4.120)$$

$$w_t = T\left(\frac{v'_0 y_{i_0}}{y_{s_t} x_{n_1+1}}\right) \in V'_{k-1} = M_{k-1}. \quad (4.121)$$

According to (4.111), we may assume that

$$-E_{n_1+1, s_t}(w_t) = \sum_{l=1}^{m'} T(\zeta_l), \quad (4.122)$$

where  $\zeta_l$  is a monomial in  $\mathcal{A}$  for  $j \in \overline{1, m'}$ ,  $T(\zeta_l) \in V'_k$  and  $\mathfrak{d}'(\zeta_l) \leq k$ . Since the monomials in  $T(\zeta_l)$  all have the common factor  $x_{n_1+1}y_{n_1+1}$  except for the first term  $\zeta_l$ . The monomials in  $\sum_{l=1}^{m'} T(\zeta_l)$  that do not have the common factor  $x_{n_1+1}y_{n_1+1}$  exactly are  $\zeta_1, \dots, \zeta_{m'}$ . By (4.121), we have

$$w_t \equiv \frac{v'_0 y_{i_0}}{y_{s_t} x_{n_1+1}} + \sum_{\substack{1 \leq j \leq m \\ j \neq t}} \frac{\beta_{s_j} v'_0 x_{s_j} y_{i_0} y_{n_1+1}}{(k + \ell_1) y_{s_j} y_{s_t}} + \frac{(\beta_{s_t} - 1) v'_0 x_{s_t} y_{i_0} y_{n_1+1}}{(k + \ell_1) y_{s_t}^2} \pmod{(x_{n_1+1} y_{n_1+1})^2}. \quad (4.123)$$

Thus

$$-E_{n_1+1,s_t}(w_t) \equiv \frac{v'_0 x_{s_t} y_{i_0}}{y_{s_t}} + \sum_{\substack{1 \leq j \leq m \\ j \neq t}} \frac{\beta_{s_j} v'_0 x_{s_j} y_{i_0}}{(k + \ell_1) y_{s_j}} + \frac{(\beta_{s_t} - 1) v'_0 x_{s_t} y_{i_0}}{(k + \ell_1) y_{s_t}} \pmod{x_{n_1+1} y_{n_1+1}}. \quad (4.124)$$

Therefore,

$$\begin{aligned} -E_{n_1+1,s_t}(w_t) &= T \left( \frac{v'_0 x_{s_t} y_{i_0}}{y_{s_t}} + \sum_{\substack{1 \leq j \leq m \\ j \neq t}} \frac{\beta_{s_j} v'_0 x_{s_j} y_{i_0}}{(k + \ell_1) y_{s_j}} + \frac{(\beta_{s_t} - 1) v'_0 x_{s_t} y_{i_0}}{(k + \ell_1) y_{s_t}} \right) \\ &= \frac{\beta_{s_t} + k + \ell_1 - 1}{k + \ell_1} v_t + \sum_{\substack{1 \leq j \leq m \\ j \neq t}} \frac{\beta_{s_j}}{k + \ell_1} v_j \end{aligned} \quad (4.125)$$

by (4.120).

We set the matrix

$$\mathbf{A} = \frac{1}{k + \ell_1} (\beta_{s_j} + \delta_{ij} \cdot (k + \ell_1 - 1))_{1 \leq i, j \leq m}. \quad (4.126)$$

Note that  $E_{n_1+1,s_t}(w_t) \in M_k$  for each  $t \in \overline{1, m}$ , and

$$- \begin{pmatrix} E_{n_1+1,s_1}(w_1) \\ E_{n_1+1,s_2}(w_2) \\ \vdots \\ E_{n_1+1,s_m}(w_m) \end{pmatrix} = \mathbf{A} \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix}. \quad (4.127)$$

Since

$$\det(\mathbf{A}) = \frac{(k + \ell_1 - 1)^{m-1}}{(k + \ell_1)^m} (k_{21} + \ell_1 + k - 1) \neq 0, \quad (4.128)$$

$\mathbf{A}$  is invertible. Thus  $v_t \in M_k$  for each  $t \in \overline{1, m}$ .

If  $i_0 \in \{s_1, \dots, s_m\}$ , there exist  $t_0 \in \overline{1, m}$  such that  $s_{t_0} = i_0$ . Then  $f = v_{t_0} \in M_k$ . Suppose that  $i_0 \notin \{s_1, \dots, s_m\}$ . Since  $T \left( \frac{v'_0}{x_{n_1+1}} \right) \in V'_{k-1} = M_{k-1}$ ,

$$E_{n_1+1,i_0} \left( T \left( \frac{v'_0}{x_{n_1+1}} \right) \right) \in M_k \subseteq V'_k. \quad (4.129)$$

Similarly, we have

$$T \left( \frac{v'_0}{x_{n_1+1}} \right) \equiv \frac{v'_0}{x_{n_1+1}} + \sum_{j=1}^m \frac{\beta_{s_j} v'_0 x_{s_j} y_{n_1+1}}{(k + \ell_1) y_{s_j}} \pmod{(x_{n_1+1} y_{n_1+1})^2}. \quad (4.130)$$

Then

$$-E_{n_1+1,i_0} T \left( \frac{v'_0}{x_{n_1+1}} \right) \equiv v'_0 x_{i_0} + \sum_{j=1}^m \frac{\beta_{s_j} v'_0 x_{s_j} y_{i_0}}{(k + \ell_1) y_{s_j}} \pmod{x_{n_1+1} y_{n_1+1}}. \quad (4.131)$$

Hence

$$-E_{n_1+1, i_0} T\left(\frac{v'_0}{x_{n_1+1}}\right) = T\left(v'_0 x_{i_0} + \sum_{j=1}^m \frac{\beta_{s_j} v'_0 x_{s_j} y_{i_0}}{(k + \ell_1) y_{s_j}}\right) = f + \sum_{j=1}^m \frac{\beta_{s_j}}{k + \ell_1} v_j. \quad (4.132)$$

Therefore,

$$f = -E_{n_1+1, i_0} T\left(\frac{v'_0}{x_{n_1+1}}\right) - \sum_{j=1}^m \frac{\beta_{s_j}}{k + \ell_1} v_j \in M_k. \quad (4.133)$$

In summary,  $f \in V'_k \Rightarrow f \in M_k$ . Recall we have already proved  $M_k \subset V'_k$ . Thus  $M_k = V'_k$ . We have proved the lemma by induction on  $k$ .  $\square$

Next we calculate the associated variety of  $M$  in this case.

**Lemma 4.5.** *If  $\ell_1, \ell_2 > 0$  and  $n_1 < n_2 = n$ ,*

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) = \mathcal{V}(\{E_{j,i}, \Delta_{i_1, i'_1}^{j_1, j'_1} \mid (j, i) \notin L; (j_1, i_1), (j'_1, i'_1) \in L\}). \quad (4.134)$$

*In particular, when  $\ell_1, \ell_2 > 0$  and  $n_1 + 1 = n_2 = n$ ,*

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) = \mathcal{V}(\{E_{j,i} \mid (j, i) \notin L\}). \quad (4.135)$$

*Proof.* Since

$$E_{j,i}(M_k) \subset M_k \quad \text{for } (j, i) \notin L, \quad (4.136)$$

we have  $\{E_{j,i} \mid (j, i) \notin L\} \subseteq I_{(1)}$ .  
For  $i \in \mathbb{1}, n_1$  and  $j \in n_1 + 2, n$ ,

$$E_{n_1+1, i}(T(x_{n_1+1}^{\ell_1} y_i^{\ell_2})) = -\frac{\ell_1 + \ell_2 + 1}{\ell_1 + 1} T(x_i x_{n_1+1}^{\ell_1+1} y_i^{\ell_2}) \notin M_0 \quad (4.137)$$

by (4.125) with  $s_t = i$ , and

$$E_{j,i}(T(x_{n_1+1}^{\ell_1} y_i^{\ell_2})) = -T(x_i x_j x_{n_1+1}^{\ell_1} y_i^{\ell_2}) \notin M_0 \quad (4.138)$$

by (2.7). Thus  $I_{(1)} = \{E_{j,i} \mid (j, i) \notin L\}$ .

For  $\xi \in \mathfrak{k}_{(p)}$  with  $p > 1$  (cf. (4.9)), we write

$$\xi = \sum_{s=1}^m a_s \prod_{t=1}^p E_{j_{s,t}, i_{s,t}} \quad \text{where } (j_{s,t}, i_{s,t}) \in L, a_s \in \mathbb{F}. \quad (4.139)$$

Fix  $v_0 \in M_0$ , we have

$$\begin{aligned} & \mathfrak{d}'\left(\left(\prod_{t=1}^p E_{j_{s,t}, i_{s,t}}\right)(v_0)\right) \\ &= \mathfrak{d}'\left(\left(\prod_{t=1}^p (-x_{i_{s,t}} x_{j_{s,t}} - y_{i_{s,t}} \partial_{y_{j_{s,t}}})\right)(v_0)\right) = p. \end{aligned} \quad (4.140)$$

Then  $\xi(v_0) = 0$  due to  $\xi(v_0) \in M_{p-1}$ .



Consider the partial sum of highest degree in  $\xi(v_0)$ :

$$\sum_{s=1}^m (-1)^p a_s \left( \prod_{t=1}^p (x_{i_s,t} x_{j_s,t}) \right) (v_0) = 0 \quad (4.141)$$

When  $n_1 + 1 = n_2$ , we have

$$\sum_{s=1}^m a_s \left( \prod_{t=1}^p x_{i_s,t} \right) = 0. \quad (4.142)$$

Hence  $\xi \in U_{p-1}(\mathfrak{g})$ . Therefore, for  $p > 1$ ,

$$I_{(p)} \equiv \{0\} \pmod{\langle I_{(1)} \rangle}. \quad (4.143)$$

Then

$$\text{Ann}_{S(\mathfrak{g})}(\overline{M}) = \langle E_{j,i} \mid (j,i) \notin L \rangle. \quad (4.144)$$

and (4.135) holds.

When  $n_1 + 1 < n_2$ , we have

$$\sum_{s=1}^m a_s \left( \prod_{t=1}^p x_{i_s,t} x_{j_s,t} \right) = 0. \quad (4.145)$$

According to the Lemma 3.1,

$$\xi \in \langle \Delta_{i_1, i'_1}^{j_1, j'_1} \mid (j_1, i_1), (j'_1, i'_1) \in L \rangle. \quad (4.146)$$

Recall (4.36), for  $v_i \in M_i$  and  $(j_1, i_1), (j'_1, i'_1) \in L$ ,

$$\left( \Delta_{i_1, i'_1}^{j_1, j'_1} \right)^{\ell_2+1} (v_i) = (x_{i'_1} y_{i_1} - x_{i_1} y_{i'_1})^{\ell_2+1} \left( x_{j'_1} \partial_{y_{j_1}} - x_{j_1} \partial_{y_{j'_1}} \right)^{\ell_2+1} (v_i) = 0, \quad (4.147)$$

since  $\sum_{j \in J_2} \deg_{y_j}(v_i) \leq \ell_2$ . Hence

$$\sqrt{I} = \langle E_{j,i}, \Delta_{i_1, i'_1}^{j_1, j'_1} \mid (j,i) \notin L; (j_1, i_1), (j'_1, i'_1) \in L \rangle \quad (4.148)$$

and (4.134) holds. □

Remind that we denoted the determinantal ideal  $I_t(S_1, S_2)$  to be the ideal generated by all the  $t$ -minors in the matrix  $(z_{i,j})_{i \in S_1, j \in S_2}$ , and  $\mathcal{V}_t(S_1, S_2) = \mathcal{V}(I_t)$  is called a *determinantal variety*, where  $S_1$  and  $S_2$  are subsets of  $\overline{1, n}$ . In particular,

$$I_2(S_1, S_2) = \langle \Delta_{i_1, i_2}^{j_1, j_2} \mid j_1, j_2 \in S_1; i_1, i_2 \in S_2 \rangle \quad (4.149)$$

and

$$I_3(S_1, S_2) = \langle \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \mid j_1, j_2, j_3 \in S_1; i_1, i_2, i_3 \in S_2 \rangle. \quad (4.150)$$

For convenience, we denote  $I_t(S_1, S_2) = 0$  and  $\mathcal{V}(I_t(S_1, S_2)) = \mathbb{A}^{|S_1|+|S_2|}$ , when  $t > |S_1|$  or  $t > |S_2|$ . Now we have our main theorem:

**Theorem 4.1.** *If  $n_1 < n_2$ ,  $\ell_1 \leq 0$  or  $\ell_2 \leq 0$ ,*

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}(I_3(J_2 \cup J_3, J_1 \cup J_2)) \cap \mathcal{V}(I_2(J_2, J_1)) \cap \mathcal{V}(I_2(J_3, J_2)) \cap \mathcal{V}(I_1(J_2, J_2)). \quad (4.151)$$

*If  $n_1 = n_2$ ,*

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}(I_3(J_3, J_1)). \quad (4.152)$$

*When  $n_1 < n_2 = n$  and  $\ell_1, \ell_2 > 0$ ,*

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}(I_2(J_2, J_1)). \quad (4.153)$$

*In particular, for the irreducible  $sl(n)$ -module  $\mathcal{H}_{(\ell_1, \ell_2)}$  in Proposition 1, the associated variety is covered by the above-mentioned cases.*

*Proof.* If  $\ell_1, \ell_2 > 0$  and  $n_1 < n_2 = n$ , then  $J_3 = \emptyset$ ,  $|J_1| = n_1$ ,  $|J_2| = n - n_1$  and

$$\begin{aligned} \mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) &= \mathcal{V}(\{E_{j,i}, \Delta_{i_1, i'_1}^{j_1, j'_1} \mid (j, i) \notin L; (j_1, i_1), (j'_1, i'_1) \in L\}) \\ &\cong \mathcal{V}(I_2(J_2, J_1)) \end{aligned} \quad (4.154)$$

as an affine variety by Lemma 4.5. Similarly, when  $\ell_1 \leq 0 \leq \ell_2$  and  $n_1 < n_2 = n$ , (4.154) still holds.

Suppose  $|J_1|, |J_2|, |J_3| \geq 2$ . According to Lemma 4.2 and Lemma 4.3, we have: if  $\ell_1 \leq 0$  or  $\ell_2 \leq 0$ ,

$$\begin{aligned} &\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \\ &= \mathcal{V} \left( \left\{ E_{i,j}, \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3}, \Delta_{k_1, k'_1}^{k_2, k'_2}, \Delta_{k_2, k'_2}^{k_3, k'_3} \mid \begin{array}{l} i_1, i_2, i_3 \in J_1 \cup J_2; j_1, j_2, j_3 \in J_2 \cup J_3; \\ (i, j) \notin L; k_s, k'_s \in J_s, s = 1, 2, 3 \end{array} \right\} \right) \\ &\cong \mathcal{V}(I_3(J_2 \cup J_3, J_1 \cup J_2) + I_2(J_2, J_1) + I_2(J_3, J_2) + I_1(J_2, J_2)); \end{aligned} \quad (4.155)$$

So (4.151) hold.

When  $n_1 < n_2$ ,  $|J_1| < 2$  or  $|J_2| = 1$  or  $|J_3| < 2$ , (4.32) and (4.78) still hold, but some terms in these two expressions are zero in this case. They correspond the cases:  $I_2(J_2, J_1) = 0$  if  $|J_2| \leq 1$  or  $|J_1| \leq 1$ ,  $I_2(J_3, J_2) = 0$  if  $|J_3| \leq 1$  or  $|J_2| \leq 1$ ,  $I_3(J_2 \cup J_3, J_1 \cup J_2) = 0$  if  $|J_2 \cup J_3| \leq 2$  or  $|J_1 \cup J_2| \leq 2$ . Hence (4.155) still hold.

Suppose  $n_1 = n_2$ . In this case,

$$\tilde{\Delta} = \sum_{i=1}^{n_1} x_i \partial_{y_i} + \sum_{s=n_2+1}^n y_s \partial_{x_s}, \quad (4.156)$$

and (2.8) no longer holds.

Suppose  $n_1 = n_2 < n$ . First we consider the case  $\ell_1 = -m_1, \ell_2 = -m_2 \leq 0$ . According to (6.6.52) in [38],

$$\begin{aligned} M &= \text{Span} \left\{ \left[ \prod_{r=1}^{n_1} x_r^{l_r} \right] \left[ \prod_{s=1}^{n-n_1} y_{n_1+s}^{k_s} \right] \left[ \prod_{r=1}^{n_1} \prod_{s=n_1+1}^n (x_r x_s - y_r y_s)^{l_{r,s}} \right] \right. \\ &\quad \left. \mid l_r, k_s, l_{r,s} \in \mathbb{N}; \sum_{r=1}^{n_1} l_r = m_1; \sum_{s=1}^{n-n_1} k_s = m_2 \right\}. \end{aligned} \quad (4.157)$$

We take

$$M_0 = \text{Span}\{X_{J_1}^{m_1} Y_{J_3}^{m_2}\} \quad (4.158)$$

(cf. (2.40) and (2.41)). Then by the arguments from (6.6.45) to (6.6.49) in [38], we can conclude that

$$M_k = \text{Span}\{X_{J_1}^{m_1} Y_{J_3}^{m_2} P^i \mid i \in \overline{0, k}\}. \quad (4.159)$$

In this case,  $L = L_3$  and we can calculate that

$$\begin{aligned} \mathcal{V}(\mathcal{H}_{(-m_1, -m_2)}) &= \mathcal{V}(\{E_{j,i}, \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \mid (j, i) \notin L; (j_s, i_s) \in L, s = 1, 2, 3\}) \\ &\cong \mathcal{V}(I_3(J_3, J_1)) \end{aligned} \quad (4.160)$$

by Lemma 3.3.

Next we consider the case  $1 < n_1 = n_2 < n$ ,  $\ell_1 \leq 0$  and  $\ell_2 \geq 0$ . We set  $\ell_1 = -m_1 - m_2$  and  $\ell_2 = m_2$  with  $m_1, m_2 \in \mathbb{N}$ . In this case,

$$\begin{aligned} M &= \mathcal{H}_{(-m_1 - m_2, m_2)} \\ &= \text{Span}\left\{ \left[ \prod_{r=1}^{n_1} x_r^{l_r} \right] \left[ \prod_{1 \leq p < q \leq n_1} (x_p y_q - x_q y_p)^{k_{p,q}} \right] \left[ \prod_{r=1}^{n_1} \prod_{s=n_1+1}^n (x_r x_s - y_r y_s)^{l_{r,s}} \right] \right. \\ &\quad \left. \mid l_r, k_{p,q}, l_{r,s} \in \mathbb{N}; \sum_{r=1}^{n_1} l_r = m_1; \sum_{1 \leq p < q \leq n_1} k_{p,q} = m_2 \right\} \end{aligned} \quad (4.161)$$

by (6.6.53) in [38]. Take

$$M_0 = \text{Span} \left\{ \left[ \prod_{1 \leq p < q \leq n_1} (x_p y_q - x_q y_p)^{k_{p,q}} \right] X_{J_1}^{m_1} \mid k_{p,q} \in \mathbb{N}, \sum_{1 \leq p < q \leq n_1} k_{p,q} = m_2 \right\}. \quad (4.162)$$

Then by (6.6.51) in [38], we can conclude that

$$M_k = \text{Span}\{M_0 P^i \mid i \leq k\}. \quad (4.163)$$

and

$$\begin{aligned} \mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) &= \mathcal{V}(\{E_{j,i}, \Delta_{i_1, i_2, i_3}^{j_1, j_2, j_3} \mid (j, i) \notin L; (j_s, i_s) \in L, s = 1, 2, 3\}) \\ &\cong \mathcal{V}(I_3(J_3, J_1)). \end{aligned} \quad (4.164)$$

holds. Symmetrically, when  $n_1 = n_2 < n$ ,  $\ell_1 \geq 0$  and  $\ell_2 \leq 0$ , the above equation still holds.

In summary, (4.152) holds if  $n_1 = n_2$ . This completes the proof of the theorem.  $\square$

In particular, when  $n_1 = n_2$  and  $|J_1|, |J_3| \geq 3$ ,

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_3(J_3, J_1) \quad (4.165)$$

is exactly a determinantal variety. When  $n_2 = n$ , the projective version of the associated variety

$$\mathcal{V}(\mathcal{H}_{(\ell_1, \ell_2)}) \cong \mathcal{V}_2(J_2, J_1) \quad (4.166)$$

is the well-known Serge variety in algebraic geometry.

The GK-dimension is equal to the dimension of the associated variety (as an affine variety). We can calculate that

$$\dim \mathcal{V}(M) = \begin{cases} 2n - 3, & \text{if } n_2 \neq n \text{ and } n_1 \neq n_2; \\ 2n - 4, & \text{if } 1 < n_1 = n_2 < n - 1; \\ n - 1, & \text{if } 1 = n_1 = n_2 \text{ or } n_1 = n_2 = n - 1 \text{ or } n_1 \neq n_2 = n, \end{cases} \quad (4.167)$$

which was obtained earlier by Bai in [2].

## References

- [1] J. Adams and D. Vogan, Associated varieties for real reductive groups, *Pure Appl. Math. Q.* **17** (2021), no. 4, 1191–1267.
- [2] Z. Bai, Gel'fand-Kirillov dimensions of the  $\mathbb{Z}^2$ -graded oscillator representations of  $sl(n)$ , *Acta Math. Sin. (Engl. Ser.)* **31** (2015), no. 6, 921–937.
- [3] Z. Bai, J. Ma and Y. Wang, On the annihilator variety of a highest weight module for classical Lie algebras, *arXiv:2304.03475v3 [math.RT]*, 7 Nov. 2023.
- [4] Z. Bai, M. Hunziker, X. Xie and R. Zierau, On the associated variety of a highest weight Harish-Chandra module, *arXiv:2402.08886v1 [math.RT]*, 14 Feb. 2024.
- [5] Z. Bai, W. Xiao and X. Xie, Gel'fand-Kirillov dimensions and associated varieties of highest weight modules, *Int. Math. Res. Not. IMRN* **2023**, no. 10, 8101–8142.
- [6] D. Barbasch and M. Bozicevic, The associated variety of an induced representation, *Proc. Amer. Math. Soc.* **127** (1999), no. 1, 279–288.
- [7] D. Barbasch and D. Vogan, The local structure of characters, *J. Funct. Anal.* **37** (1980), no. 1, 27–55.
- [8] D. Barbasch and D. Vogan, Primitive ideals and orbital integrals in complex classical groups, *Math. Ann.* **259**(1982), no. 2, 153–199.
- [9] J. Barshay, Determinantal varieties, monomial subgroups and algebras associated with ideals, *Proc. Amer. Math. Soc.* **40** (1973), 16–22.
- [10] M. Bernardara, M. Bolognesi and D. Faenzi, Homological projective duality for determinantal varieties, *Adv. Math.* **296** (2016), 181–209.
- [11] J. Bernstein, Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients. *Funkcional Anal. I Prilozen.* **5** (1971), no. 2, 1–16.
- [12] W. Borho and J-L Brylinski, Differential operators on homogeneous spaces. I. Irreducibility of the associated variety for annihilator of induced modules, *Invent. Math.* **69** (1982), no. 3, 437–476.
- [13] W. Borho and J. Jantzen, Über primitive Ideale in der Einhüllenden einer halbeinfachen Lie Algebra, *Invent. Math.* **39** (1977), no. 1, 1–53.

- [14] F. Catanese, Babbage’s conjecture, contact of surfaces, symmetric determinantal varieties, *Invent. Math.* **63** (1981), no. 3, 433–465.
- [15] C. de Concini, D. Eisenbud and C. Procesi, Young diagrams and determinantal varieties, *Invent. Math.* **56** (1980), no. 2, 129–165.
- [16] T. Enright and M. Hunziker, Resolutions and Hilbert series of determinantal varieties and unitary highest weight modules, *J. Algebra* **273** (2004), no. 2, 608–639.
- [17] T. Enright, M. Hunziker and M. Prueett, Diagrams of Hermitian type, highest weight modules, and syzygies of determinantal varieties, *Symmetry: representation theory and its applications*, 121–184, *Prog. Math.* **257**, Birkhäuser/Springer, New York, 2014.
- [18] P. Griffiths, Infinitesimal variations of Hodge structure. III. Determinantal varieties and the infinitesimal invariant of normal functions, *Compositio Math.* **50** (1983), no. 2-3, 267–364.
- [19] A. Gyoja and H. Yamashita Associated variety, Kostant-Sekiguchi correspondence, and locally free  $U(n)$ -action on Harish-Chandra modules, *J. Math. Soc. Japan* **51** (1999), no. 1, 129–149.
- [20] Y. Homma and M. Manabe, Determinantal Calabi-Yau varieties in Grassmannians and Givental I-functions, *J. High Energy Phys.* **2018**, no. 12, 046, *front matter+48pp*.
- [21] H. Jakobsen, An intrinsic classification of the unitarizable highest weight modules as well as their associated varieties, *Compositio Math.* **101** (1996), no. 3, 313–352.
- [22] A. Joseph, On classification of primitive ideals in the enveloping algebra of a semisimple Lie algebra, *Lie group representations, I (College Park, Md., 1982/1983)*, 30–76, *Lecture Notes in Math.* **1024**, Springer, Berlin, 1983.
- [23] A. Joseph, On the associated variety of a primitive ideal, *J. Algebra* **93** (1985), no. 2, 509–523.
- [24] A. Joseph, Annihilators and associated varieties of unitary highest weight modules, *Ann. Sci. École Norm. Sup. (4)* **25** (1992), no. 1, 1–45.
- [25] T. Kobayashi, Discrete decomposability of the restriction of  $A_q(\lambda)$  with respect to reductive subgroups III. Restriction of Harish-Chandra modules and associated varieties, *Invent. Math.* **131** (1998), no. 2, 229–256.
- [26] C. Luo and X. Xu,  $\mathbb{Z}^2$ -graded oscillator representations of  $sl(n)$ , *Comm. Algebra* **41** (2013), no. 8, 3147–3173.
- [27] H. Matumoto, Whittaker vectors and associated varieties, *Invent. Math.* **89**(1987), no. 1, 219–224.
- [28] W. McGovern, A triangular result for associated varieties of highest weight modules, *Comm. Algebra* **28** (2000), no. 4, 1835–1843.

- [29] A. Melnikov, Irreducibility of the associated varieties, *C. R. Acad. Sci. Paris Sér. I. Math.* **316** (1993), no. 1, 53–57.
- [30] I. Musson, Associated varieties for classical Lie superalgebras, *Hopf algebras and quantum groups (Brussels, 1998)*, 177-188, *Lecture Notes in Pure and Appl. Math.*, 209 Dekker, New York, 2000.
- [31] K. Nishiyama, P. Trapa and A. Wachi, Codimension one connectedness of the graph of associated varieties, *Tohoku Math. J. (2)* **68**(2010), no. 2, 199–239.
- [32] P. Trapa, Annihilators and associated varieties of  $A_q(\lambda)$  modules for  $U(p, q)$ , Singular Gelfand-Tsetlin modules of  $gl(n)$ , *Compositio Math.* **129** (2001), no. 1, 1–45.
- [33] D. Vogan, Gel’fand-Kirillov dimension for Harish-Chandra modules, *Invent. Math.* **48** (1978), no. 1, 75–98.
- [34] D. Vogan, Ordering of the primitive spectrum of a semisimple Lie algebra, *Math. Ann.* **248**(1980), no. 3, 195–203.
- [35] D. Vogan, Associated varieties and unipotent representations, *Harmonic analysis on reductive groups (Brunswick, ME. 1989)*, 315–388, *Prog. Math.* **101**, Birkhäuser Boston, MA, 1991.
- [36] G. Williamson, A reducible characteristic variety in type A, *Representations of Reductive Groups*, 517-532, *Progr. Math.* **312**, Birkhäuser/Springer, Cham, 2015.
- [37] K. Wong, Two-dimensional gauge dynamics and the topology of singular determinantal varieties, *J. High Energy Phys.* **2017**, no. 3, 132, *front matter+16pp*.
- [38] X. Xu, *Representations of Lie Algebras and Partial Differential Equations*, Springer, Singapore 2017, xxxvi+620pp.
- [39] H. Yamashita, Criteria of finiteness of restriction of  $U(\mathfrak{g})$ -modules to subalgebras and applications to Harish-Chandra modules. A study in relation to the associated varieties, *J. Funct. Anal.* **121** (1994), no. 2, 296-329.
- [40] H. Yamashita, Associated varieties and Gel’fand-Kirillov dimensions for the discrete series of a semisimple Lie group, *Proc. Japan Acad. Ser. A Math. Sci.* **70** (1994), no. 2, 50–55.
- [41] H. Yamashita, Description of the associated varieties for the discrete series representations of a semisimple Lie group—an elementary proof of differential operators of gradient type, *Comment. Math. Univ. St. Paul.* **47** (1998), no. 1, 35–52.