On the quantum numbers of the X(1880)

Qin-He Yang^{1,2,3}, Ling-Yun Dai^{1,4},^{*} and and Ulf-G. Meißner^{2,3,5†}

¹ School for Theoretical Physics, School of Physics and Electronics, Hunan University, Changsha 410082, China

² Helmholtz Institut für Strahlen- und Kernphysik and Bethe Center

for Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany

³Institute for Advanced Simulation (IAS-4), Forschungszentrum Jülich, D-52425 Jülich, Germany

⁴ Hunan Provincial Key Laboratory of High-Energy Scale Physics and Applications,

Hunan University, Changsha 410082, China and

⁵Peng Huanwu Collaborative Center for Research and Education, Beihang University, Beijing 100191, China

(Dated: December 11, 2024)

We study the properties of the X(1880), the structure around the $\bar{p}p$ threshold that appears in the

 $3(\pi^+\pi^-)$ invariant mass spectrum in the decay process of $J/\psi \to \gamma 3(\pi^+\pi^-)$. Nucleon-antinucleon rescattering is taken into account in our analysis, and the decay amplitude of $J/\psi \to \gamma 3(\pi^+\pi^-)$ can be obtained by the distorted wave Born approximation. With these amplitudes, we analyze the contributions to the X(1880) from different partial waves. Our analysis suggests that the X(1880) should be isoscalar 0^{-+} , and it is generated by the threshold behavior.

Introduction.- Over half a century, the quark model is believed to be the fundamental template for describing hadrons, which are composed of three quarks or a pair of quark-antiquark [1, 2]. With the advent of Quantum Chromodynamics, it became clear that other types of bound states should exist. In particular, the carriers of the strong force, the gluons, can form the hadrons in terms of hybrid and glueball, which is a kind of matter that can truly be called exotic. For recent studies addressing the role of gluons as constituents of hadrons, see, for example, Refs. [3–7]. Recently, a candidate for glueball was reported by the BESIII collaboration in Ref. [8], named X(1880). It is found in the $3(\pi^+\pi^-)$ invariant mass spectrum in the decay process of $J/\psi \to \gamma 3(\pi^+\pi^-)$, with much higher statistics than the earlier measurement [9]. Its mass and width are $M = 1882.1 \pm 1.7 \pm 0.7$ MeV and $\Gamma = 30.7 \pm 5.5 \pm 2.4$ MeV. The X(1880) attracts huge attention from the community, as it reveals a new structure of hdaron if the glueball nature is confirmed, see for example Refs. [10–16].

To study the properties of the resonance, one needs to know the quantum numbers of the X(1880) as a prerequisite. However, since there are many final states in the decay process of $J/\psi \to \gamma 3(\pi^+\pi^-)$, it is an arduous task for experimentalists performing a partial wave analysis to fix the quantum numbers of the X(1880). In this letter, we propose a simplified partial wave analysis to select the quantum numbers. Since the X(1880) is just around the $\bar{p}p$ threshold, a two-step process should work well for the interaction dynamics, namely $J/\psi \rightarrow$ $\gamma \bar{N}N \rightarrow \gamma 3(\pi^+\pi^-)$. The $\bar{N}N$ re-scattering plays an essential role in final-state interaction, and a series of comprehensive partial wave amplitudes of \overline{NN} scattering has already been obtained using chiral effective field theory (ChEFT) up to next-to-next-to-leading order $(N^{3}LO)$ [17]. With these amplitudes, one can construct the $J/\psi \to \gamma 3(\pi^+\pi^-)$ decay partial wave amplitudes through the distorted wave Born approximation

(DWBA) [18–20]. One by one, the partial wave amplitudes obtained can be applied to describe all the relevant experimental data sets, the invariant mass spectra of $J/\psi \rightarrow \gamma \bar{p}p$ and $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ and the crosssection of $p\bar{p} \rightarrow 3(\pi^+\pi^-)$. The quantum numbers of the X(1880) can be classified from the quality of fitting each partial wave amplitude to the data. Notice that the interference between different partial waves will not have such a distinct character as that of pure waves. The reason is that around the threshold, partial waves have a threshold factor, p^L . As an example, the square of the partial wave amplitudes will be either strongly suppressed or not for P- and S- waves, respectively.

Formalism.– As discussed above, the amplitude of the decay process $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ is obtained through $J/\psi \rightarrow \gamma N\bar{N} \rightarrow \gamma 3(\pi^+\pi^-)$ [21]. Thus, one needs three amplitudes, $J/\psi \rightarrow \gamma N\bar{N}$, $\bar{p}p \rightarrow 3(\pi^+\pi^-)$, and $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$. They are solved by a set of equations established via the DWBA,

$$F_{1}(Q) = A_{1}^{0}(p') + \int_{0}^{\infty} \frac{\mathrm{d}kk^{2}}{(2\pi)^{3}} A_{1}^{0}(k) \frac{1}{Q - 2E_{k} + i\epsilon} T(k, p'; E_{p'}),$$

$$F_{2}(Q) = A_{2}^{0}(p) + \int_{0}^{\infty} \frac{\mathrm{d}kk^{2}}{(2\pi)^{3}} T(p, k; E_{k}) \frac{1}{2E_{k} - Q + i\epsilon} A_{2}^{0}(k) .$$

$$F_{3}(Q) = A_{3}^{0}(Q) + \int_{0}^{\infty} \frac{\mathrm{d}kk^{2}}{(2\pi)^{3}} F_{1}(E_{k}) \frac{1}{Q - 2E_{k} + i\epsilon} A_{2}^{0}(k) ,$$

(1)

where the subscripts '1, 2, 3' represent the processes of $J/\psi \to \gamma N \bar{N}, N \bar{N} \to \gamma 3(\pi^+\pi^-)$, and $J/\psi \to \gamma 3(\pi^+\pi^-)$, respectively. T is the partial wave scattering amplitude of $N \bar{N} \to N \bar{N}$. Q is the $3(\pi^+\pi^-)$ or $N \bar{N}$ invariant mass for the different processes, p, p' are the center-of-mass momenta of initial and final $N \bar{N}$ system, and E is the energy of the nucleon, respectively. Following earlier works [17, 22], the transition amplitudes and annihilation po-

tential in Eq. (1) are parameterized as

$$\begin{aligned}
A_{1,2}^{0,S}(p) &= \tilde{C}_{1,2}^{S} + C_{1,2}^{S} p^{2} + D_{1,2}^{S} p^{4}, \\
A_{1,2}^{0,P}(p) &= C_{1,2}^{P} p + D_{1,2}^{P} p^{3}, \\
A_{1,2}^{0,D}(p) &= D_{1,2}^{D} p^{2}, \\
A_{3}^{0}(Q) &= \tilde{C}_{3} + C_{3} Q.
\end{aligned}$$
(2)

The superscripts S, P, D are short for different partial waves. The parameters in the annihilation potential are taken as real numbers to keep the same formalism as that of Ref. [17, 22]. For $A_3^0 (J/\psi \to \gamma 3(\pi^+\pi^-))$, it is written in a general formalism with lowest orders of momentum. The $\bar{N}N$ scattering amplitude T is solved by the Lippmann-Schiwinger equation (LSE) [17],

$$T_{N\bar{N}\to N\bar{N}}(p',p;E_p) = V_{N\bar{N}\to N\bar{N}}(p',p) + \int_0^\infty \frac{\mathrm{d}kk^2}{(2\pi)^3} \times V_{N\bar{N}\to N\bar{N}}(p',k) \frac{1}{2E_p - 2E_k + i\epsilon} T_{N\bar{N}\to N\bar{N}}(k,p;E_p) ,$$
(3)

with the potentials given by ChEFT up to N³LO [17]. Here, we will restrict our analysis to the region below $T_{\text{Lab}} \leq 120$ MeV, corresponding to $p_{\text{Lab}} \leq 490$ MeV. Notice that in this low-energy region, the $N\bar{N} \rightarrow N\bar{N}$ solution fits well the phase shifts of all the partial waves obtained from the PWA [23]. Especially, the lowest partial waves, ${}^{1}S_{0}$, ${}^{3}P_{0}$, ${}^{3}P_{1}$, ${}^{1}D_{2}$ and ${}^{3}P_{2}$, do not show a resonance-like behavior in the region of their most substantial variation. This makes all the LECs from ChEFT unchanged when the $T_{N\bar{N}\rightarrow N\bar{N}}$ is taken into Eq. (1) to produce the $J/\psi \rightarrow \gamma p\bar{p}$ and $N\bar{N} \rightarrow 3(\pi^{+}\pi^{-})$ amplitudes, leaving only a few parameters from the Born term to be fixed, i.e., \tilde{C}_{i} , C_{i} , and D_{i} , see Eq. (2).

With the amplitudes obtained in Eq. (1), one can analyze the $3(\pi^+\pi^-)$ and $p\bar{p}$ invariant mass spectra for the decay processes of $J/\psi \to \gamma 3(\pi^+\pi^-)$ and $J/\psi \to \gamma p\bar{p}$, and the cross section of the scattering process of $p\bar{p} \to 3(\pi^+\pi^-)$. According to parity and charge conjugation conservation, the permitted partial waves are 0^{-+} , 0^{++} , 1^{++} , 2^{-+} , and 2^{++} in the $3(\pi^+\pi^-)$ system for the J/ψ radiative decay, ignoring the higher partial waves with total angular momentum J > 2. Correspondingly, one needs to consider the 1S_0 , 3P_0 , 3P_1 , 1D_2 and 3P_2 waves in the $N\bar{N}$ system. One can take each partial wave individually to fit all the datasets of invariant mass spectra and cross sections. The partial wave that best describes all these data will be recognized as contributing to the structure around the $\bar{p}p$ threshold most.

On the experimental side, there are sufficient data points (74) in the energy region of $p_{\text{Lab}} \leq 490$, supporting the possibility of selecting the quantum numbers. There are 3 data points on the cross section of $p\bar{p} \rightarrow 3(\pi^+\pi^-)$ [24–26], 44 data points on the invariant mass spectra of $3(\pi^+\pi^-)$ in $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ (31 data points with high statistics) [8, 9], and 27 data points on the $\bar{p}p$ invariant mass spectra in $J/\psi \to \gamma p\bar{p}$ (12 data points with high statistics) [27, 28]. In contrast, we have 9, 7, and 5 parameters for the S-, P-, and D-wave cases. Also, there are five normalization factors for fitting the invariant mass spectra, in lack of the detection efficiency from the experimental measurements. Notice that the first parameter of the Born term in $J/\psi \to \gamma \bar{p}p$ has been fixed to be one, $\tilde{C}_1^P = C_1^P = D_1^D = 1$ as it is multiplied with the normalization factor for the detection efficiency.

Results and discussions. – As discussed above, we consider the lowest partial waves, ${}^{3}P_{0}(0^{++})$, ${}^{1}S_{0}(0^{-+})$, ${}^{3}P_{1}(1^{++})$, ${}^{1}D_{2}(2^{-+})$ and ${}^{3}P_{2}(2^{++})$, with both isoscalar and isovector channels. The fit quality for each partial wave is shown in Fig. 1. The results for isoscalar waves



FIG. 1. Comparison of N³LO fitting results for different partial waves. The fit results corresponding to the ${}^{1}S_{0}$, ${}^{3}P_{0}$, ${}^{3}P_{1}$, ${}^{1}D_{2}$ and ${}^{3}P_{2}$ partial waves are indicated by the purple dashed, black solid, green dot-dash, blue dot, and red dot-dash-dash lines, respectively. The data are taken from Refs. [8, 9, 24– 29]. The cut-off is chosen as R = 1.0 fm.

are in the left column, and the isovector ones are shown in the right column. The invariant mass spectra of the $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ and $J/\psi \rightarrow \gamma p\bar{p}$, and the cross section of $p\bar{p} \rightarrow 3(\pi^+\pi^-)$, are shown in the first, second, and third row, in order. To determine the quantum numbers of the structure around the $\bar{p}p$ threshold (called X(1880) for simplicity), we follow a simple strategy. That is, the partial wave with the correct quantum numbers should describe all the data well.

As can be found from the $3(\pi^+\pi^-)$ invariant mass spectra, the $I = 0^{3}P_{0}, {}^{3}P_{1}, {}^{1}D_{2}$ and the $I = 1 {}^{1}D_{2}$ waves can hardly describe the data, see the purple dashed, blue dotted, and green dot-dashed lines in the first row of Fig 1. One can exclude these quantum numbers. From the $p\bar{p}$ invariant mass spectra, one can find that all the waves, except for the I = 0 ${}^{1}S_{0}$, ${}^{3}P_{0}$ and I = 1 ${}^{1}S_{0}$ ones, can hardly describe the data. Among them, the I = 0 ${}^{3}P_{0}$ wave gives a much worse description of the data than the I = 0 ¹S₀-wave. From the $p\bar{p} \rightarrow 3(\pi^+\pi^-)$ cross section, one can find that only the I = 0 ${}^{1}S_{0}$ and I = 1 ${}^{1}S_{0}$ waves can describe the data well. This is unsurprising as the amplitudes of P- and D- waves are proportional to the momentum p^L , resulting in a vanishing cross section at the $\bar{p}p$ threshold. In short, only the I = 0 ¹S₀ wave survives by describing all the data well; the I = 1 $^{1}S_{0}$ wave is worse than the isoscalar one but much better than all other waves. Consequently, the X(1880) should be in the I = 0 ¹S₀ wave. The I = 1 ¹S₀ wave is the second choice but much worse than the former.

R (fm)	Isospin	${}^{1}S_{0}$	${}^{3}P_{0}$	${}^{3}P_{1}$	${}^{1}D_{2}$	${}^{3}P_{2}$
0.9	I = 0	1.96	8.94	41.92	67.39	21.25
	I = 1	4.26	35.10	30.76	69.05	29.98
1.0	I = 0	1.99	6.74	33.81	66.18	18.58
	I = 1	3.30	31.34	26.38	67.82	25.03
1.1	I = 0	2.04	6.34	25.70	64.78	16.40
	I = 1	3.32	28.00	21.20	66.45	20.85
1.2	I = 0	2.07	6.42	19.80	63.24	14.90
	I = 1	3.22	24.74	17.13	64.94	17.98

TABLE I. The χ^2 /d.o.f. for the differential partial waves, where d.o.f. represents the number of degrees of freedom.

To test the stability of our conclusion, we list all the χ^2 /d.o.f. for different cut-offs, R = 0.9, 1.0, 1.1, 1.2 fm in Table I. As can be found, the I = 0¹S₀ wave has the smallest χ^2 /d.o.f., around 2.0. The I = 1¹S₀ wave has the second smallest χ^2 /d.o.f., from 3.2 to 4.3. All other waves have much larger χ^2 /d.o.f.. This confirms that the X(1880) most likely has the quantum numbers $I J^{PC} = 0$ 0⁻⁺. Indeed, we have also tested the present analysis within $T_{\text{Lab}} \lesssim 100$ MeV, and the same conclusion is obtained.

After fixing the quantum numbers, one can analyze the contributions to the structure around the threshold from different parts as shown in Fig. 2. The Born term/background and loop contributions, corresponding to the first and second terms of Eq. (1), are shown as blue dashed and purple dash-dotted lines, respectively. As can be found, the structure in the $3(\pi^+\pi^-)$ variant mass



FIG. 2. Contributions from different parts of the $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ decay process. The solution is an isoscalar 1S_0 wave. The black solid, blue dashed, and purple dash-dotted lines indicate the contribution of the total, Born term (background), and the loop effect from $N\bar{N}$ rescattering, respectively.

spectrum is from the interference between the loop and background contributions. The background is smooth, and only the loop contribution, i.e., the $N\bar{N}$ rescattering effects, supplies a structure. It should be pointed out that from the $N\bar{N}$ scattering partial wave amplitudes, one can not find any poles around the threshold. This supports the statement that the X(1880) is generated by the threshold behavior.

Summary.– In this letter, we have presented a combined analysis on the decay process of the $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ and $J/\psi \rightarrow \gamma p\bar{p}$ reactions and the scattering process of $p\bar{p} \rightarrow 3(\pi^+\pi^-)$. The distorted wave Born approximation is applied to solve the amplitudes, with input of the $N\bar{N}$ scattering amplitudes given by chiral effective field theory up to N³LO. The isoscalar and isovector ${}^{1}S_{0}$, ${}^{3}P_{0}$, ${}^{3}P_{1}$, ${}^{1}D_{2}$ and ${}^{3}P_{2}$ partial waves are taken into the analysis one by one. Our result suggests that the quantum numbers of the structure around the $\bar{p}p$ threshold, the X(1880), discovered in the $3(\pi^+\pi^-)$ invariant mass spectrum of $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$ process, is most likely to be $I J^{PC} = 00^{-+}$. Further, the X(1880)is generated by the $\bar{N}N$ threshold effect. This can be checked by the future experiment.

Acknowledgements.– This work is supported by the National Natural Science Foundation of China (NSFC) with Grants No. 12322502, 12447186, 12335002, Joint Large Scale Scientific Facility Funds of the NSFC and Chinese Academy of Sciences (CAS) under Contract No. U1932110, Hunan Provincial Natural Science Foundation with Grant No. 2024JJ3004, Fundamental Research Funds for the central universities, and by the MKW NRW under the funding code NW21-024-A and by ERC EXOTIC (grant No. 101018170). The work of UGM was supported in part by the CAS President's International Fellowship Initiative (PIFI) (Grant No. 2025PD0022).

* dailingyun@hnu.edu.cn

- [†] meissner@hiskp.uni-bonn.de
- [1] M. Gell-Mann, Phys. Lett. 8, 214 (1964).
- [2] G. Zweig, (1964), 10.17181/CERN-TH-401.
- [3] S. D. Bass and P. Moskal, Rev. Mod. Phys. 91, 015003 (2019), arXiv:1810.12290 [hep-ph].
- [4] B. Ketzer, B. Grube, and D. Ryabchikov, Prog. Part. Nucl. Phys. **113**, 103755 (2020), arXiv:1909.06366 [hepex].
- [5] D. Vadacchino, in 39th International Symposium on Lattice Field Theory (2023) arXiv:2305.04869 [hep-lat].
- [6] M. R. Shepherd, J. J. Dudek, and R. E. Mitchell, Nature 534, 487 (2016), arXiv:1802.08131 [hep-ph].
- [7] A. Rodas *et al.* (JPAC), Phys. Rev. Lett. **122**, 042002 (2019), arXiv:1810.04171 [hep-ph].
- [8] M. Ablikim *et al.* (BESIII), Phys. Rev. Lett. **132**, 151901 (2024), arXiv:2310.17937 [hep-ex].
- [9] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. D 88, 091502 (2013).
- [10] S. G. Salnikov and A. I. Milstein, Nucl. Phys. B 1002, 116539 (2024), arXiv:2311.14309 [hep-ph].
- [11] M. Karliner and J. L. Rosner, (2024), arXiv:2406.05920 [hep-ph].
- [12] Y. Xiao, J.-X. Lu, and L.-S. Geng, (2024),

arXiv:2406.01292 [nucl-th].

- B.-Q. Ma, (2024), 10.1360/TB-2024-0578, arXiv:2406.19180 [hep-ph].
- [14] P.-Y. Niu, Z.-Y. Zhang, Y.-Y. Li, Q. Wang, and Q. Zhao, Phys. Rev. D **110**, 094020 (2024), arXiv:2408.14876 [hepph].
- [15] P. G. Ortega, D. R. Entem, F. Fernandez, and J. Segovia, (2024), arXiv:2410.12465 [hep-ph].
- [16] Z.-S. Jia, Z.-H. Zhang, F.-K. Guo, and G. Li, (2024), arXiv:2410.16873 [hep-ph].
- [17] L.-Y. Dai, J. Haidenbauer, and U.-G. Meißner, JHEP 07, 078 (2017).
- [18] X.-W. Kang, J. Haidenbauer, and U.-G. Meißner, Phys. Rev. D 91, 074003 (2015).
- [19] L.-Y. Dai, J. Haidenbauer, and U.-G. Meißner, Phys. Rev. D 98, 014005 (2018).
- [20] J. P. Dedonder, B. Loiseau, and S. Wycech, Phys. Rev. C 97, 065206 (2018).
- [21] Q.-H. Yang, D. Guo, and L.-Y. Dai, Phys. Rev. D 107, 034030 (2023), arXiv:2209.10101 [hep-ph].
- [22] X.-W. Kang, J. Haidenbauer, and U.-G. Meißner, JHEP 02, 113 (2014), arXiv:1311.1658 [hep-ph].
- [23] D. Zhou and R. G. E. Timmermans, Phys Rev C 86, 044003 (2012).
- [24] F. Sai, S. Sakamoto, and S. S. Yamamoto, Nucl. Phys. B 213, 371 (1983).
- [25] A. Bertin *et al.* (OBELIX Collaboration), Phys. Lett. B 369, 77 (1996).
- [26] E. Klempt, C. Batty, and J.-M. Richard, Phys. Rept. 413, 197 (2005).
- [27] J. Z. Bai *et al.* (BES Collaboration), Phys. Rev. Lett. **91**, 022001 (2003).
- [28] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **108**, 112003 (2012).
- [29] J. P. Alexander *et al.* (CLEO Collaboration), Phys. Rev. D 82, 092002 (2010).