

ARITHMETICITY AND GEOMETRICAL COMMENSURATORS

YANLONG HAO

ABSTRACT. This paper aims to characterize rank-one arithmetic and locally symmetric metrics in the coarsely geometric setting using coarse-geometric commensurators. We provide a positive answer in general under the Hilbert-Smith conjecture and unconditionally for finite volume negatively curved manifolds with finitely many cusps.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Arithmetic locally symmetric spaces are central topics in geometric group theory and are closely related to algebra, number theory, and combinatorics. Eberlein [7, 8] and Chen [5], characterized symmetric spaces H of non-compact type among all irreducible, simply-connected Riemannian manifold with non-positive sectional curvature by $\text{Isom}(H)$. Following Farb and Weinberger [9], Van Limbeek [28] extended this work to classify all closed manifolds (M, g) with non-discrete isometry group $\text{Isom}(\tilde{M}, g)$. Along the way, hidden symmetries [10] were used to detect all arithmetic closed locally symmetric manifolds.

On the other hand, there is a well-known result of Margulis says that a lattice Γ in a semisimple Lie group G is arithmetic if and only if its commensurate is dense in G .

In [12], the authors considered the coarsely geometric setting and, assuming the Hilbert-Smith conjecture, provided necessary and sufficient conditions for the arithmeticity of left-invariant hyperbolic metrics on hyperbolic groups, generalized the aforementioned result of Margulis. This paper analyzes the same question for general discrete subgroups of isometries of hyperbolic spaces.

A Γ action on a metric space (X, d) is called **metrically proper** if, for all $x \in X$ and a real number $R > 0$, the set $\{\gamma \in \Gamma / d(x, \gamma x) < R\}$ has only finitely many elements.

Let (X, d) be a good hyperbolic space, and let $\Gamma < \text{Isom}(X)$ be a discrete subgroup with $0 < \delta_\Gamma < \infty$ and of divergence type. As defined in [11], there exists a Bowen-Margulis-Sullivan type measure m_Γ^{BMS} on the unit tangent bundle $\partial^{(2)}X \times \mathbb{R}$. For more details, refer to section 2.D. For the property and definition of good hyperbolic places, see section 2.A and also [21]. Here, we call a subgroup of $\text{Isom}(X)$ discrete if it metrically properly acts on X .

It is important to note that there is no standard definition of volume in the coarsely geometric setting. Therefore, we adopt the following definition:

Date: December 11, 2024.

Definition 1.1. Γ is a $2\frac{1}{2}$ lattice if $0 < m_{\Gamma}^{\text{BMS}}(\partial^{(2)} X \times \mathbb{R}/\Gamma) < \infty$.

Consider an isometric Γ -action on X . Let p be a base point of X . Denote (Γ, d) as the metric space such that $d(\gamma_1, \gamma_2) = d(\gamma_1 p, \gamma_2 p)$ for all $\gamma_1, \gamma_2 \in \Gamma$. Therefore, the problem of classifying isometrical actions of Γ up to roughly isometry is equivalent to the problem of classifying left-invariant quasi-metric on Γ . In this paper, we only consider proper actions.

We call a left-invariant metric (Γ, d) **arithmetic** if there exists a symmetric space H of non-compact type and an embedding i of Γ into $\text{Isom}(\mathbf{H})$ as an arithmetic lattice, such that $(\Gamma, \lambda d)$ is roughly isometric to $(i(\Gamma)o, d_H)$ for some point $o \in \mathbf{H}$ for some $\lambda > 0$.

Now we turn to the second gradient of the paper. The purely group-theoretic concept related to the commensurator of a subgroup, known as the *abstract commensurator*

$$\text{AbCom}(\Gamma)$$

of a general countable group Γ , is defined as the set of equivalence classes $[\phi]$ of group isomorphisms $\phi : \Gamma' \rightarrow \Gamma''$ between finite index subgroups $\Gamma', \Gamma'' < \Gamma$. Two isomorphisms $\phi_i : \Gamma'_i \rightarrow \Gamma''_i$ ($i = 1, 2$), are considered equivalent $\phi_1 \sim \phi_2$ if there exist a finite index subgroup $\Gamma' < \Gamma'_1 \cap \Gamma'_2$ such that the restrictions of ϕ_1 and ϕ_2 to Γ' agree. The composition of (appropriate restrictions of) two such isomorphisms defines a group structure on $\text{AbCom}(\Gamma)$.

Definition 1.2. Let Γ be a countable group and d a metric on Γ as previously defined. The *coarsely-geometric commensurator group*, denote as $\text{GeCom}(\Gamma, d)$, is a subgroup of the abstract commensurator group $\text{AbCom}(\Gamma)$. It consists of classes $[\phi]$ of isomorphisms $\phi : \Gamma' \rightarrow \Gamma''$ that satisfy the condition:

$$\sup_{\gamma_1, \gamma_2 \in \Gamma'} |d(\phi(\gamma_1), \phi(\gamma_2)) - d(\gamma_1, \gamma_2)| < \infty,$$

for all $\gamma_1, \gamma_2 \in \Gamma'$. This group is independent of the choice of base point.

The main results presented in this paper are as follows.

Theorem A. *Let (X, d) be a good hyperbolic space. Suppose $\Gamma < \text{Isom}(X)$ is a non-elementary $2\frac{1}{2}$ lattice, and it acts metrically properly on X with compact limit set $\Lambda_X \Gamma$. Then there exists a finite normal subgroup N of Γ . If*

$$[\text{GeCom}(\Gamma, d) : \Gamma] = \infty,$$

Then there exists a finite normal subgroup N of Γ with quotient group $\bar{\Gamma} = \Gamma/N$ such that either

- (1) $\bar{\Gamma}$ is a lattice with a dense commensurator of a non-discrete, totally disconnected, locally compact group, or
- (2) $\bar{\Gamma}$ embeds as an arithmetic lattice in a rank-one simple real Lie group $G = \text{Isom}(\mathbf{H})$, and (Γ, d) is arithmetic.

Additionally, it is of interest to characterize locally symmetric metrics within the coarsely geometric framework.

Let $\text{rIsom}(X)$ denote the set of rough isometries on (X, d) , defined as:

$$\text{rIsom}(X) := \{f : X \rightarrow X \mid |f_*d - d| < \infty\}.$$

The group of roughly isometries is denoted as $\text{RIsom}(X) = \text{rIsom}(X) / \sim$, where $f \sim g$ if $|f - g| < \infty$.

An element $f \in \text{RIsom}(X)$ induces a homeomorphism $\partial f \in \text{Homeo}(\partial X)$. For every subgroup $\Gamma < \text{Isom}(X)$, we define

$$\begin{aligned} \text{H}_\Gamma(X) &= \{\partial f \mid f \in \text{RIsom}(X), \partial f(\Lambda\Gamma) = \Lambda\Gamma\}, \\ d\Gamma &= \{\partial\gamma \mid \gamma \in \Gamma\}. \end{aligned}$$

Theorem B. *Let (X, d) be a good hyperbolic space. Suppose that $\Gamma < \text{Isom}(X)$ is a $2\frac{1}{2}$ lattice and Γ acts metrically properly on X with compact limit set $\Lambda_X\Gamma$. Then there exists a finite normal subgroup N of Γ with quotient group $\bar{\Gamma} = \Gamma/N$. If the index $[\text{H}_\Gamma(X) : d\Gamma]$ is infinite. Then either*

- (1) $\bar{\Gamma}$ is a lattice of a non-discrete, totally disconnected, locally compact group,
or
- (2) $\bar{\Gamma}$ embeds as a lattice in a rank-one simple real Lie group $G = \text{Isom}(\mathbf{H})$,
and (Γ, d) is arithmetic.

1.A. **Hilbert-Smith conjecture.** While a positive solution to Hilbert's fifth problem has been found [29], the Hilbert-Smith conjecture remains elusive in most cases.

Conjecture 1.3 (Hilbert-Smith). *If a locally compact group acts continuously and faithfully on a connected manifold, it is a Lie group.*

Corollary 1.4. *Assume the Hilbert-Smith conjecture, under the condition of Theorem A and B, also assume that the limit set $\Lambda_X\Gamma$ is a manifold. Then case (1) in Theorems A and B cannot occur.*

The Hilbert-Smith conjecture has been established in a few instances. There is a concise proof for groups acting by Lipschitz maps on a Riemannian manifold [26]. John Pardon has solved the three-dimensional case [24].

Therefore, unconditionally, we obtain the following result.

Corollary 1.5. *In the statement of Theorem A and B, if Λ_X is a connected topological manifold of dimension no more than 3, then case (1) cannot occur.*

1.B. **Finite volume manifold.** Theorem A relies on Hilbert-Smith conjecture in general. However, under certain geometrical conditions, we may prove the result directly without using the Hilbert-Smith conjecture, leading to unconditional results.

Theorem C. *Let (M, g) be a finite-volume complete Riemannian manifold with cusps such that there exists $a, b > 0$ such that the sectional curvature of M satisfies $-a \leq \kappa \leq -b < 0$. Let $\Gamma = \pi_1(M)$. Assume that M has a finite Bowen-Margulis measure. Fix a point o in the Riemannian universal cover of M , \tilde{M} . Let d be the metric on Γ defined by the induced metric on the orbit Γo . If*

$$[\text{GeCom}(\Gamma, d) : \Gamma] = \infty,$$

(Γ, d) is rank-one arithmetic.

The paper is organized as follows: In Section 2, we recall some preliminaries. Section 3 shows that $2\frac{1}{2}$ -lattice is a lattice in a suitable locally compact group H . Then we study the structure of H in Section 4. Section 5 is the proof of Theorem A and B. Section 6 is devoted to the proof of Theorem C.

2. PRELIMINARIES

2.A. Good hyperbolic spaces. Gromov hyperbolic spaces play a fundamental role in modern mathematics. However, certain applications require stronger geometric properties. In [21], the concept of strong hyperbolic spaces is introduced. For our purposes, the following definition suffices.

Definition 2.1. A hyperbolic space X is said to be ϵ -good, where $\epsilon > 0$, if the following two properties hold for each basepoint $p \in X$:

- the Gromov product $(\cdot, \cdot)_p$ on X extends continuously to the bordification $X \cup \partial X$.
- The function $d_\epsilon := \exp(-\epsilon(\cdot, \cdot)_p)$ defines a metric on the boundary ∂X .

Example 2.2. If Γ is a hyperbolic group, then the Green metric defined by a symmetric and finitely supported random walk on Γ is both good hyperbolic and strong hyperbolic [21].

Another useful tool for the study of good hyperbolic spaces (X, d) is the Busemann function. Given $\xi \in \partial X$, the Busemann function $\beta_\xi : X \times X \rightarrow \mathbb{R}$ is defined as follows:

$$\beta_\xi(x, y) = \lim_{z \rightarrow \xi} d(x, z) - d(y, z).$$

The limit is independent of the choice of sequence z since X is a good hyperbolic space.

2.B. Cross-ratio. The classical cross-ratio on S^1 can be generalized to the boundary of good hyperbolic spaces [16, 22, 25]. Cross-ratios provide a unified interpretation of many geometric structures, making them a valuable tool for studying various spaces of representations.

Let Y be a space and $Y^{(4)}$ be the set of 4-tuples of pairwise distinct points in Y . A map $\mathbb{B} : Y^{(4)} \rightarrow \mathbb{R}$ is called a cross-ratio if it satisfies the following conditions:

$$\begin{aligned} \mathbb{B}(x, y, z, w) &= -\mathbb{B}(y, x, z, w); \\ \mathbb{B}(x, y, z, w) &= \mathbb{B}(z, w, x, y); \\ \mathbb{B}(x, y, z, w) &= \mathbb{B}(x, y, z, t) + \mathbb{B}(x, y, t, w); \\ \mathbb{B}(x, y, z, w) + \mathbb{B}(y, z, x, w) + \mathbb{B}(z, x, y, w) &= 0. \end{aligned}$$

For an ϵ -good hyperbolic space (X, d_X) , there exists a canonical cross-ratio on ∂X given by:

$$(2.1) \quad \mathbb{B}(\xi, \eta, \xi', \eta') = \ln \frac{d_\epsilon(\xi, \xi') d_\epsilon(\eta, \eta')}{d_\epsilon(\xi, \eta') d_\epsilon(\eta, \xi')}.$$

Otal observed that the definition in (2.1) is equivalent to

$$\mathbb{B}(\xi, \eta, \xi', \eta') = \lim_{(x, y, x', y') \rightarrow (\xi, \eta, \xi', \eta')} \frac{1}{2} [d_X(x, x') + d_X(y, y') - d_X(x, y') - d_X(y, x')].$$

Furthermore, the diagonal action of $\text{RIso}(X)$ on $\partial X^{(4)}$ preserves the cross-ratio according to the second definition.

2.C. Möbius homermorphisms. Let (K, d) be a compact metric space without isolated points, and consider Möbius self-homeomorphisms of K . Recall that the cross-ratio of a quadruple of distinct points in K is defined as:

$$\mathbb{B}(x, y, x', y') = \ln \frac{d(x, x')d(y, y')}{d(x, y')d(y, x')}.$$

A homeomorphism of K is called Möbius if it preserves the cross-ratio.

Following Sullivan [27], Möbius maps satisfy the *geometric mean-value property* [20]. i.e.,

A self homeomorphism g of K is Möbius if and only if there exists a positive continuous function on K , denoted g' , such that for all $x, y \in K$, we have:

$$g'(x)g'(y) = \frac{d(gx, gy)}{d(x, y)}.$$

Nica [20] has shown that the metric derivative g' is Lipschitz. Moreover, the group of Möbius maps is locally compact by [20, Proposition 5.6].

2.D. Bowen-Margulis-Sullivan measure. Discovered independently by Bowen and Margulis and reconstructed from the Patterson-Sullivan measure later, the Bowen-Margulis-Sullivan measure (BMS measure) of a negatively curved closed manifold contains critical geometric information. It serves as a tool to study the growth rate of primarily closed geodesics.

Furman [11] extended the BMS measure in a coarsely geometric setting. Here, we adopt a similar strategy. Let Γ be a group that acts metrically properly on a good hyperbolic space X , with limit set $\Lambda_X \Gamma$ and of divergence type. Assuming the critical exponent $0 < \delta < \infty$ and $\Lambda_X \Gamma$ is compact, we denote ν the Patterson-Sullivan measure on $\Lambda \Gamma$. For the construction of the Patterson-Sullivan measure, see [6, Theorem 15.4.9].

There is a $\Gamma \times \mathbb{R}$ action on $\partial^{(2)} \times \mathbb{R}$

$$(\gamma, t) \cdot (\xi, \eta, s) = (\gamma\xi, \gamma\eta, s + \frac{\beta_\xi(p, \gamma p) - \beta_\eta(p, \gamma p)}{2} + t),$$

where β is the Busemann function with base point $p \in X$. A straightforward computation shows that the measure $m_\Gamma^{\text{BMS}} = d_e^{-\frac{2\delta}{\epsilon}} \nu \times \nu \times dt$ is $\Gamma \times \mathbb{R}$ invariant. This is the measure used in Definition 1.1.

It is worth noting that it is likely the assumption of divergence type is not necessary. But we are not able to show this. Hence, all groups in this paper are of (generalized) divergence type.

3. $2\frac{1}{2}$ LATTICE IS A LATTICE

Consider a good hyperbolic space (X, d) and let $\Gamma < \text{Isom}(X)$ be a $2\frac{1}{2}$ lattice with a limit set $\Lambda\Gamma = S^{k-1}$. In this section, we establish that Γ is virtually a lattice of a locally compact group H . The properties of H are further analyzed in the next section.

3.A. Möbius homomorphism and H . The cross-ratio \mathbb{B} on ∂X , when restricted on the limit set $\Lambda_X\Gamma$, is Γ -invariant. Let

$$\text{Mob} := \{f \in \text{Homeo}(\Lambda_X\Gamma) \mid f^{(4)}\mathbb{B} = \mathbb{B}\}.$$

With respect to the metric d_ϵ , every $f \in \text{Mob}$ is an Möbius homeomorphism. It is important to note that when f is the boundary map of an isometry F , the metric derivative is related to the Busemann function as follows:

$$f'(\xi) = \exp(\epsilon\beta_\xi(p, F^{-1}p)).$$

Now we define the group H , which depends on m_Γ^{BMS} .

Let ν be the pattern-Sullivan measure on $\Lambda_X\Gamma$. We define H as:

$$H := \left\{ f \in \text{Homeo}(\Lambda_X\Gamma) \mid (f \times f)_*(d_\epsilon^{-\frac{2\delta}{\epsilon}} \nu \times \nu) = d_\epsilon^{-\frac{2\delta}{\epsilon}} \nu \times \nu \right\}.$$

By the definition of cross-ratio, all $f \in H$ preserve the cross-ratio almost everywhere with respect to the measure ν^4 . Since the cross-ratio is continuous, it follows that $f \in \text{Mob}$. Therefore, H is a closed subgroup of Mob and is therefore locally compact.

3.B. $d\Gamma$ is discrete. To prove that $d\Gamma$ is discrete in H , we combine the following three lemmas. Note that it is unclear that H is a subgroup of $\text{Homeo}(\partial X)$, therefore, the discreteness of Γ as a subgroup of $\text{Isom}(X)$ is not enough.

Lemma 3.1 and 3.2 are well-understood for proper spaces. Since we are not restricted to locally compact spaces, we give short proofs here.

Let $d : \text{Isom}(X, d) \rightarrow \text{Homeo}(\partial X)$ be the boundary map.

Lemma 3.1. *The kernel $F = \text{Ker}(d)$ has a bounded orbit.*

Proof. Take f, g and $h \in F$. Let $x = f(p)$, $y = g(p)$, $z = h(p)$. If $w_n \in X$ converge to $\xi \in \partial X$, then

$$2\lim_n (y, w_n)_x = d(x, y) + \beta_\xi(x, y) = d(x, y) + \frac{1}{\epsilon} \ln \left(\frac{(g^{-1})'(\xi)}{(f^{-1})'(\xi)} \right) = d(x, y).$$

Similarly, $2(z, \xi)_x = 2\lim_n (z, w_n)_x = d(x, z)$. By the hyperbolic inequality,

$$2(y, z)_x \geq \min\{d(x, y), d(x, z)\} - 2\delta.$$

In particular, taking $g = h = \text{Id}_X$, we have $d(f(p), p) \leq 2\delta$ for all $f \in F$. \square

Since Γ acts metrically properly on X , The kernel of the map $d : \Gamma \rightarrow d\Gamma$ is finite. The discreteness of $d\Gamma$ follows a similar strategy.

Lemma 3.2. *For any $K > 0$, the set $\Omega(K) := \{\gamma \in d\Gamma \mid \sup_{\xi \in \partial X} |\beta_\xi(\gamma p)| \leq K\}$ is finite. In particular, $d\Gamma$ is discrete in $\text{Homeo}(\partial X)$.*

Proof. Let $f \in \Omega(K)$. Then, for all $\xi \in \partial X$, we have $|\frac{1}{\epsilon} \ln(f^{-1})'(\xi)| \leq K$.

Fix a lift in Γ for f and denote it by f again. Let $y_0 = p$ $y_1 = f(p)$.

If b_n is a sequence in X converge to $\xi \in \partial X$. then, similar to before, for all $i = 1, 2,$

$$2 \lim_n (b_n, y_i)_p \geq d(y_i, p) - 2K.$$

Using the hyperbolic inequality, we can deduce that:

$$2(y_1, y_0)_p \geq \min\{d(p, y_0), d(p, y_1)\} - 2\delta - 2K.$$

This is equivalent to

$$d(y_1, y_0) - d(p, y_0) = d(y_1, p) \leq 2\delta + 2K$$

There orbit of p under $\Omega(K)$ is bounded. Therefore the set has only finitely many elements. \square

By restriction the action of $d\Gamma$ on ∂X to $\Lambda_X \Gamma$, we obtain a map $\psi : d\Gamma \rightarrow H$.

Lemma 3.3. *Ker(ψ) is finite and $\psi(d\Gamma)$ is discrete in H .*

Proof. Assume there is a sequence of elements $\{\gamma_n\}$ so that $\psi(\gamma_n)$ converges to the identity map. After taking a subsequence, we may assume that for all $\xi \in \Lambda_X \Gamma$, $\frac{1}{2} \leq (\gamma_n)'(\xi) \leq 2$.

Let $\eta \neq \zeta \in \Lambda_X \Gamma$. By the geometric mean-value property, we have:

$$\frac{1}{4} \leq \frac{d_\epsilon(\gamma_n \eta, \gamma_n \zeta)}{d_\epsilon(\eta, \zeta)} \leq 4.$$

Consider any point $\theta \in \partial X$. By the triangle inequality and the definition of the visual metric, we may assume (change η to ζ if necessary):

$$\frac{d_\epsilon(\eta, \zeta)}{8} \leq \frac{d_\epsilon(\gamma_n \eta, \gamma_n \zeta)}{2} \leq d_\epsilon(\gamma_n \theta, \gamma_n \eta) \leq 1.$$

Using the geometric mean-value property, we obtain the following:

$$\gamma_n'(\theta) = \frac{1}{\gamma_n'(\eta)} \frac{d_\epsilon^2(\gamma_n \theta, \gamma_n \eta)}{d_\epsilon^2(\theta, \eta)} \geq \frac{1}{2} \frac{d_\epsilon^2(\eta, \zeta)}{64}.$$

Similarly,

$$\gamma_n'(\theta) \leq \frac{2}{d_\epsilon^2(\eta, \zeta)}.$$

By Lemma 3.2, the sequence $\{\gamma_n\}$ contains only finitely many different elements and eventually constant. \square

The proof of Lemma 3.3 gives a useful criterion for precompactness.

Lemma 3.4. *Let A be a subset of H , if there exist two points $\xi \neq \eta \in \Lambda_X \Gamma$ such that*

- (1) *there exist $\kappa > 0$ such that $d(\gamma \xi, \gamma \eta) \geq \kappa$,*
- (2) *there exist $M > 0$ such that $\frac{1}{M} \leq \gamma'(\xi) \leq M$*

for all $\gamma \in A$. The A is precompact.

3.C. $d\Gamma$ is a lattice. We are now ready to prove the main theorem of this section.

Theorem 3.5. *The image $\psi(d\Gamma)$ is a lattice of H .*

Proof. In this proof, for simplicity, we assume $\Gamma = \psi(d\Gamma)$.

Similar as in [11], consider the abstract geodesic flow

$$H \rightarrow \text{Homeo}((\Lambda_X \Gamma)^{(2)} \times \mathbb{R}),$$

$$f(x, y, t) = (f(x), f(y), t + \frac{\log f'(x) - \log f'(y)}{2\delta_\Gamma}).$$

Then the H action is proper.

First, the quotient space $Y = (\Lambda_X \Gamma)^{(2)} \times \mathbb{R} / H$ is locally compact and second countable. The second countability is clear, and for local compactness, we can fix any point y in Y and a pre-image of y , say x . Taking a compact neighborhood C of x , Since the action is proper, we see that the projection map restricted to C is continuous, hence yielding a compact neighborhood of y .

By definition, there exists a probability measure m_Γ^{BMS} on $\bar{Y} = (\Lambda_X \Gamma)^{(2)} \times \mathbb{R} / \Gamma$. Let $p: \bar{Y} \rightarrow Y$ be the natural projection, and $\nu = p_*(m_\Gamma^{\text{BMS}})$. Since Y is a Randon space, there is disintegration $\{\mu_y / y \in Y\}$ on the fibre of p . The fiber of $y \in Y$ is $\Gamma \backslash H / K_y$, where K_y is the compact subgroup of H fixing y . For a general point y , μ_y induces a Γ -invariant measure ν_y on H / K_y , and provides a disintegration of $(\Lambda_X \Gamma)^{(2)} \rightarrow Y$.

$$\begin{array}{ccccc} H/K_y & \longrightarrow & (\Lambda_X \Gamma)^{(2)} \times \mathbb{R} & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow = \\ \Gamma \backslash H / K_y & \longrightarrow & \bar{Y} & \longrightarrow & Y \end{array}$$

Since m_Γ^{BMS} is Randon, ν_y is Randon. The fact that H preserve m_Γ^{BMS} ensures that ν_y is H -invariant ν -almost surely. Hence ν_y is Haar on H / K_y and Γ is a lattice in H . \square

4. STRUCTURE OF H

The structure of the group H can be determined by studying the property of the group Γ .

The amenable radical of a locally compact second countable group G , denoted as $\text{Rad}_{\text{am}}(G)$, is the largest closed normal amenable subgroup of G . The starting point for the structure of H is a consequence of Hilbert's fifth problem, which was first observed by Burger and Monod [4].

Theorem 4.1. [4, Theorem 3.3.3] *Every locally compact group G contains an open, normal, finite-index subgroup G_0 that contains $\text{Rad}_{\text{am}}(G)$. The quotient $G_0 / \text{Rad}_{\text{am}}(G)$ is topologically isomorphic to a direct product of a connected, center-free, semisimple, real Lie group without compact factors and a totally disconnected locally compact group with trivial amenable radical.*

To understand the structure of H , we focus on its amenable radical first, which is related to the (CAF) condition of Γ . The (CAF) condition states that every amenable commensurated subgroup of Γ is finite.

We utilize barycenter methods to show that Γ satisfies the (CAF) condition. Given a probability measure ν on $\Lambda_X\Gamma$, the L -barycenter $B_\nu(L)$ of ν is defined as the subset of X such that:

$$\left\{ x \in X \mid \int_{\xi \in \partial X} \beta_\xi(x, p) d\nu \leq \inf_{y \in X} \left\{ \int_{\xi \in \partial X} \beta_\xi(y, p) d\nu \right\} + L \right\}$$

Lemma 4.2. *If ν has no atoms, the set $B_\nu(L)$ is bounded for all $L > 0$.*

Proof. Let $F_\nu : X \rightarrow \mathbb{R}$ be the map defined as:

$$F_\nu(x) = 2 \int_{\xi \in \partial X} \beta_\xi(x, p) d\nu.$$

It suffices to show that $F_\nu(x) \rightarrow \infty$ as $d(x, p) \rightarrow \infty$.

Rewrite

$$F_\nu(x) = \int_{\xi \in \partial X} (d(x, p) - 2(\xi, x)_p) d\nu = d(x, p) - 2 \int_{\xi \in \partial X} (\xi, x)_p d\nu.$$

For each $x \in X$, let

$$E(x) := \left\{ \xi \in \Lambda_X\Gamma \mid (\xi, x)_p \geq \frac{d(x, p)}{3} \right\}.$$

Then

$$2 \int_{\xi \in \Lambda_X\Gamma} (\xi, x)_p d\nu \leq 2d(x, p)\nu(E(x)) + \frac{2d(x, p)}{3}(1 - \nu(E(x))).$$

Let $\xi \in E(x)$, then by hyperbolicity,

$$E(x) \subset \left\{ \zeta \in \Lambda_X\Gamma \mid (\zeta, \xi)_p \geq \frac{d(x, p)}{3} - \delta \right\}.$$

Since $\Lambda_X\Gamma$ is compact, and ν has no atoms, there exist a $K > 0$ so that $\nu(E(x)) \leq \frac{1}{8}$ when $d(x, p) \geq K$. It follows that for all x with $d(x, p) \geq K$, $F_\nu(x) \geq \frac{1}{6}d(x, p)$.

This completes the proof. \square

Lemma 4.3. *Γ satisfies the (CAF) condition.*

Proof. Let N be an amenable commensurated subgroup of Γ . Then there exists an N -invariant probability measure ν on the compact space $\Lambda_X\Gamma$.

Case 1: The support of the measure ν has points with a finite N -orbit. Note that the union of all finite N -orbit is Γ -invariant. Up to take a subgroup of finite index, N have at least 3 fixed points. From the geometric mean-value property and similar construction as in the proof of Lemma 3.2, the metric derivative of N is bounded from below and above. Hence N is finite by Lemma 3.2.

Case 2: The support of ν has no finite orbit point. Then ν has no atoms and by Lemma 4.2, $B_\nu(1)$ is bounded. Since N preserve the bounded subset $B_\nu(1)$, N must be finite.

This establishes the (CAF) condition of Γ . \square

In view of the following result, the amenable radical of H is compact.

Theorem 4.4 ([3], Theorem 5.2). *The amenable radical of a lattice envelope of a group with property (CAF) is compact.*

Remark 4.5. Even though the definition lattice envelope in [3] is assumed compact generated, the proof of Theorem 4.4 is independent of this condition.

Indeed, the amenable radical of H is trivial.

Lemma 4.6. *H has no compact radical.*

Proof. Assume K is a compact normal subgroup of H . Then there exists a constant κ such that for all $f \in K$ and $x \in \Lambda_X \Gamma$, $|\ln f'(x)| < \kappa$.

By considering a hyperbolic element $\gamma \in \Gamma$, we have that for all $f \in K$:

$$|\ln(\gamma^n f \gamma^{-n})'(\gamma^-)| < \kappa.$$

Since

$$\ln(\gamma^n f \gamma^{-n})'(\gamma^-) = \ln(\gamma^n)'(f(\gamma^-)) + \ln f'(\gamma^-) + \ln(\gamma^{-n})'(\gamma^-),$$

it indicates that $|\ln(\gamma^n)'(f(\gamma^-) - n\ell_\gamma)| < 2\kappa$ for all n . Taking a fixed $n > \frac{2\kappa}{\ell_\gamma}$, we have $d_\epsilon(\gamma^+, K\gamma^-) > 0$.

On the other hand, $K\gamma^- = \gamma K \gamma^{-1} \gamma^- = \gamma K \gamma^-$. Hence $K\gamma^- = \gamma^-$. Since this holds for all hyperbolic elements, $K = Id$. Therefore, H has no compact radical. \square

According to Theorem 4.1, H is virtually a product. Next Lemma states that one of the two factors is trivial.

Lemma 4.7. *H is not virtually a product of two nontrivial groups.*

Proof. Assume there exists a finite index subgroup H_0 of H such that $H_0 = H_1 \times H_2$. Let p_i be the projection $H \rightarrow H_i$, $i = 1, 2$. Replacing $\psi(d\Gamma)$ by $\psi(d\Gamma) \cap H_0$, the same argument as in Lemma 4.6 implies that the amenable radical of H_0 is trivial.

Let $\gamma \in \psi(d\Gamma) \cap H_0$ be a hyperbolic element. Since $p_i(\gamma)$ commute with γ , up to take square, $p_i(\gamma)$ fix γ^\pm pointwisely. Since for the metric derivative, we have $\gamma' = p_1(\gamma)' p_2(\gamma)'$. One of the two elements say, $p_1(\gamma)$ has a contraction fixed point at γ^+ . Then the fixed points of $p_1(\gamma)$ are exactly γ^\pm since it is γ -invariant. Now H_2 commute with $p_1(\gamma)$. Hence up to an index two subgroup, H_2 fixed γ^\pm . Consider the set $\text{Fix}(H_2)$. It is a H_0 -invariant closed subset of $\Lambda_X \Gamma$. Thus it is $\Lambda_X \Gamma$. In other words, H_2 is either trivial or $\mathbb{Z}/2\mathbb{Z}$.

If $H_2 = \mathbb{Z}/2\mathbb{Z}$, then H_2 is a compact radical of H . It is impossible. Therefore, H_2 is trivial. \square

Finally, we can classify the possible structures of H :

Theorem 4.8. *Let Γ act metrically properly on a good hyperbolic space (X, d) . Assuming Γ is a $2\frac{1}{2}$ lattice and its limit set $\Lambda\Gamma$ is compact, the image $\psi(d\Gamma)$ is a lattice of H as constructed in section 3. The group H is one of the following:*

- (1) *A discrete group.*
- (2) *A non-discrete, totally disconnected locally compact group with trivial amenable radical.*

(3) *A rank-one simple real Lie group $G = \text{Isom}(\mathbf{H})$.*

Proof. In view of Theorem 4.1. If H does not fall into cases (1) and (2), it is virtually a connected, center-free, semisimple real Lie group G without compact factors. Since $\text{Aut}(G) = G$, $H = G \times H'$ for some finite group H' . Since H' is in the amenable radical, it must be trivial, leading to $H = G$. And G is simple by Lemma 4.7.

Haettel's Theorem [15] states all isometric actions of lattices of higher rank Lie groups are elementary. This forces G to be of rank one. \square

5. PROOF OF THEOREM A AND B

The proofs of Theorem A and B rely on the following lemmas.

For any $h \in \text{GeCom}(\Gamma, d)$, it induces a roughly isometry on (Γ, d) , and thus a boundary map $\partial h \in \text{Homeo}(\Lambda_X \Gamma)$.

Lemma 5.1. *The kernel of the map $d : \text{GeCom}(\Gamma, d) \rightarrow \text{Homeo}(\Lambda_X \Gamma)$ is trivial, and the image is contained in H . Furthermore, we have:*

$$\begin{aligned} d(\text{GeCom}(\Gamma, d)) &\subset \text{Com}_H(d\Gamma), \\ [d(\text{GeCom}(\Gamma, d)) : d\Gamma] &= \infty. \end{aligned}$$

Proof. The $d(\text{GeCom}(\Gamma, d))$ preserves the Patterson-Sullivan measure class. It also preserves the cross-ratio. Therefore, $d(\text{GeCom}(\Gamma, d)) \subset H$.

Any $s \in \text{GeCom}(\Gamma, d)$ induces an element in $\text{AbCom}(d\Gamma)$, by the same proof as in [12, Proposition 2.3], which is compatible with the action on the boundary $\Lambda_X \Gamma$. Hence it is enough to show the kernel of d is finite.

Let $s \in \text{GeCom}(\Gamma, d)$ such that $d(s) = \text{Id}$. Consider a point-mass free measure ν on $\Lambda_X \Gamma$. It is clear that ν is invariant under s . Since s is a $(1, C)$ quasi-isometry of (Γ, d) , we have $s(B_\nu(1)) \subset B_\nu(1 + 2C)$. The same holds for $\gamma_*(\nu)$. Using the fact that $B_{\gamma_*(\nu)}(A) = \gamma B_\nu(A)$, we can find a constant M such that $d(s(\gamma), \gamma) < M$ for all $\gamma \in \Gamma$ by Lemma 4.2.

Let Γ_1 be the domain of s . Then, we have for all $\gamma_1 \in \Gamma$, $s(\gamma_1) = \gamma_1 \tau(\gamma_1)$ for some map τ with $d(\tau(\gamma_1), e) \leq M$. The equation $s(\gamma_1 \gamma_2) = s(\gamma_1) s(\gamma_2)$ can be rewritten as

$$\tau(\gamma_1 \gamma_2) = \gamma_2^{-1} \tau(\gamma_1) \gamma_2 \tau(\gamma_2).$$

Thus, we obtain $d(\tau(\gamma_1) \gamma_2, \gamma_2) = d(\gamma_2^{-1} \tau(\gamma_1) \gamma_2, e) \leq 2M$ for all $\gamma_1, \gamma_2 \in \Gamma_1$. Since Γ_1 is finite index subgroup, $\tau(\gamma_1) \in N = \text{Ker}(\Gamma \rightarrow d\Gamma)$.

Let $\Gamma_2 = \text{ker}(\Gamma \rightarrow \text{Aut}(N))$. Then τ (restricted to Γ_2) is a group homomorphism to the finite group N . Taking $\Gamma_3 = \text{ker } \tau$, we have $s = \text{Id}_{\Gamma_3} = \text{Id}_\Gamma$ as an equivalence class.

This completes the proof. \square

Lemma 5.1 rules out the case (1) in Theorem 4.8. And for Theorem B, case (1) in Theorem 4.8 is ruled out by conditions. Now in case (2), the Busemann cocycle of (Γ, d) extends to the rank-1 Lie group H . Rank-1 condition implies that the Busemann cocycle of H up to strict equivalence is 1 dimensional. For more details, see [18, Section 7]. Then Theorem A and B follow from the marked

length spectrum rigidity in coarse-geometrical setting, see [11] for cocompact case, and [17] for non-compact case.

6. FINITE VOLUME COMPLETE MANIFOLD

We will prove Theorem C in a stronger form, namely, Theorem 6.3. Before this, we will introduce some notation.

Let Γ be a group with bounded torsion, i.e. there exists $N > 0$, such that for all torsion elements in $\alpha\Gamma$, $\alpha^N = 1$. Assume Γ is hyperbolic relative to a finite collection of finitely generated infinite nilpotent groups $\{P_i\}_{i=1}^n$. In [14], Groves and Manning constructed a hyperbolic space with a proper Γ -action and $\{P_i\}_{i=1}^n$ are precisely the cusp groups. We call $(\Gamma, \{P_i\}_{i=1}^n)$ are *geometrically good* if Γ admit such a proper action on a proper good hyperbolic space (X, d) , Γ is a $2\frac{1}{2}$ -lattice for this action and P_i acts cocompactly on $\Lambda_X\Gamma \setminus \{\xi_i\}$ where ξ_i is the cusp point corresponds to P_i .

A typical example will be the fundamental group of the manifold in Theorem C.

Let $(\Gamma, \{P_i\}_{i=1}^n)$ be geometrically good and (X, d) be the correspondence Γ -space. By Theorem 4.8, Γ is a lattice of a locally compact group H . By the following lemma, if H is a totally disconnected group, Γ is a cocompact lattice of H .

Lemma 6.1 ([1], Corollary 4.11). *Let Γ be an unimodular, totally disconnected, locally compact group. Let $\Gamma < H$ be a closed subgroup of finite covolume. Let m_Γ denote the Haar measure of Γ . If there is an upper bound on the Haar measures of compact open subgroups in Γ , that is:*

$$\sup \{m_\Gamma(U) / U \text{ is a compact open subgroup in } H\} < \infty,$$

then H/Γ is compact.

Therefore, we have the following:

Lemma 6.2. *If there exists an element $h \in H$ such that there exists a cusp point $\xi \in \partial X$ with $h\xi = \xi$ and $h'(\xi) > 1$, then H is a Lie group.*

Proof. Assume H is not a lie group. Then by Theorem 4.8, either H is a finite extension of Γ , in which case, the h in the lemma does not exist, or H is a totally disconnected, non-discrete locally compact group.

Denote the fixed points of h on the boundary by ξ and η . Let

$$H_\xi^c = \{s \in H / s(\xi) = \xi, s'(\xi) = 1\}, \quad \Gamma_\xi^c = \Gamma \cap H_\xi^c.$$

Then since ξ is a cusp point, Γ_ξ^c is a discrete infinite virtually nilpotent subgroup of H_ξ^c . Notice that the orbit $H_\xi^c(\eta)$ is h -invariant. Now let Ξ be the orbit $H_\xi^c \cdot \eta$ with the Hamenstädt metric: [6, Proposition 3.6.19] $d'(\zeta, \zeta') = \frac{d_e(\zeta, \zeta')}{d_e(\zeta, \xi)d_e(\zeta', \xi)}$. Then H_ξ^c acts isometrically on Ξ . And h acts on Ξ via contraction with factor $\frac{1}{h'(\xi)^2}$.

Now the subspace $\Gamma_\xi^c(\eta)$ of Ξ is quasi-isometric to Γ_ξ^c by the construction, which is a nilpotent group. Since h is a contraction, $h^n(\Gamma_\xi^c(\eta))$ converge to the

asymptotic cone of Γ_ξ^c in Hasudorff-Gromov distance as pointed proper metric spaces. It follows that the asymptotic cone (see [13, 23]) of Γ_ξ^c embedding into Ξ via a Lipschiz map. Recall that this asymptotic cone is a manifold, hence $H_\xi^c(\eta)$ is not totally disconnected, A contradiction.

The results follow. \square

Theorem 6.3. *Let $(\Gamma, \{P_i\}_{i=1}^n)$ be geometrically good, and denote the orbit of Γ with induced metric by (Γ, d) . If*

$$[\text{GeCom}(\Gamma, d) : \Gamma] = \infty,$$

then (Γ, d) is rank-one arithmetic.

Proof. By Theorem 4.8, Γ is a lattice of a locally compact group H with three possibilities. By assumption, H is not a discrete group.

If H is a rank-one simple Lie group, we are done by the work of Margulis.

If H is a totally disconnected locally compact group. We will derive a contradiction. The construction is in two steps.

Step 1: Find a compact open subgroup fixed a cusp point.

For any cusp point ξ , denote

$$H_\xi = \{h \in H / h(\xi) = \xi\}.$$

By Lemma 6.2, $H_\xi = H_\xi^c$. Recall that Γ is a cocompact lattice of H . Let $K' < H$ be a compact open subgroup such that $K' \cap \Gamma = \{1\}$.

Let ξ be a cusp point of Γ and $\xi_i, 1 \leq i \leq m$ be a set of representatives of Γ nonequivalent cusps in the orbit $H\xi$, with $\xi_1 = \xi$. We also fix elements $h_i \in H$ so that $h_i\xi = \xi_i$.

By construction $d(\text{GeCom}(\Gamma, d))$ is dense in H . Let $k \in d(\text{GeCom}(\Gamma, d)) \cap K'$. Since $\text{GeCom}(\Gamma, d)$ map cusp point to cusp point. We have, there exist $1 \leq i \leq n$ and $\gamma \in \Gamma$, so that $\gamma k \xi_1 = \xi_i$. Therefore $\gamma k h_i^{-1} \xi_1 = \xi_i$. Now choose a compact fundamental domain Δ_i for Γ_{ξ_i} actions on $\partial X \setminus \{\xi_i\}$ and $\eta_i \in \Delta_i$. There exists $p_k \in \Gamma_{\xi_i}$ so that $p_k \gamma k h_i^{-1}(\eta_i) \in \Delta_i$. By the condition $(p_k \gamma k h_i^{-1})'(\xi_i) = 1$, and there are two almost fixed point, the set

$$A := \{p_k \gamma k h_i^{-1} / k \in d(\text{GeCom}(\Gamma, d)) \cap K'\}$$

is precompact by Lemma 3.4.

Thus the set $B = \{p_k \gamma / p_k \gamma k h_i^{-1} \in A\} \subset \cup_{i=1}^m A h_i K'$ is precompact. Note that Γ is a lattice in H , therefore, the set B is finite.

Now, $k h_i^{-1} \xi_1 = k \xi_1 = (p_k \gamma)^{-1} \xi_i$. Hence the orbit of ξ under $d(\text{GeCom}(\Gamma, d)) \cap K'$ is finite. Now the set is dense in K' , therefore, the orbit of ξ under K' is finite. Up to take a finite index subgroup, we have an open compact subgroup K of H which fixes the cusp point ξ .

Step 2: Asymptotic cone of nilpotent groups.

We consider the space Ξ with the Hamenstädt metric. Then K and Γ_ξ act isometrically on Ξ . By definition of geometrical goodness, Γ_ξ acts cocompactly on Ξ . Then Milnor-Švarc lemma implies that the orbit $\Gamma_\xi o$ with $o \in \Xi$ is quasi-isometric to a word metric.

Therefore, each $k \in K$ induced a quasi-isometry of a word metric on Γ_ξ . Notice that Γ_ξ is a nilpotent group. Recall the work of Gromov and Pansu on the asymptotic cone of nilpotent groups. We have that each k induced a group homeomorphism of the asymptotic cone of Γ_ξ , which is a Lie group N . Denote this map by

$$\chi : K \rightarrow \text{Aut}(N).$$

Note that $\chi(K)$ is a compact subgroup since all elements in K are isometry of Ξ .

On the other hand, N is the \mathbb{R} -Malcev completion of Γ_ξ [19]. Hence for all $k \in d(\text{GeCom}(\Gamma, d)) \cap K$, the induced map on finite index subgroups of Γ_ξ extends to N . Therefore,

$$\chi : d(\text{GeCom}(\Gamma, d)) \cap K \rightarrow \text{Aut}(N)$$

is injective. Therefore $\chi(K)$ is not discrete. A contradiction.

Hence, H is a Lie group. Following the same argument as in the proof of Theorem A, we have (Γ, d) is arithmetic.

This completes the proof. □

REFERENCES

- [1] U. Bader, P.-E. Caprace, T. Gelander, and S. Mozes, *Lattices in amenable groups*, *Fundamenta Mathematicae* **246** (2019), 217–255.
- [2] U. Bader and A. Furman, *Boundaries, rigidity of representations, and Lyapunov exponents*, arXiv preprint arXiv:1404.5107 (2014).
- [3] U. Bader, A. Furman, and R. Sauer, *Lattice envelopes*, *Duke Mathematical Journal* **169** (2020), no. 2, 213–278.
- [4] M. Burger and N. Monod, *Continuous bounded cohomology and applications to rigidity theory*, *Geometric And Functional Analysis* **12** (2002), no. 2, 219–280.
- [5] S.-S. Chen and P. Eberlein, *Isometry groups of simply connected manifolds of nonpositive curvature*, *Illinois Journal of Mathematics* **24** (1980), no. 1, 73–103.
- [6] T. Das, D. Simmons, and M. Urbański, *Geometry and dynamics in Gromov hyperbolic metric spaces*, Vol. 218, American Mathematical Soc., 2017.
- [7] P. Eberlein, *Lattices in spaces of nonpositive curvature*, *Annals of Mathematics* **111** (1980), no. 3, 435–476.
- [8] ———, *Isometry groups of simply connected manifolds of nonpositive curvature II*, *Acta Mathematica* **149** (1982), no. 1, 41–69.
- [9] B. Farb and S. Weinberger, *Hidden symmetries and arithmetic manifolds*, *Contemporary Mathematics* **387** (2005), 111.
- [10] ———, *Isometries, rigidity and universal covers*, *Annals of mathematics* **168** (2008), no. 3, 915–940.
- [11] A. Furman, *Coarse-geometric perspective on negatively curved manifolds and groups*, *Rigidity in dynamics and geometry*, 2002, pp. 149–166.
- [12] A. Furman and Y. Hao, *Marked length pattern and arithmeticity* (2023).
- [13] M. Gromov, *Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits)*, *Publications Mathématiques de l’IHÉS* **53** (1981), 53–78.
- [14] D. Groves and J. F. Manning, *Dehn filling in relatively hyperbolic groups*, *Israel Journal of Mathematics* **168** (2008), 317–429.
- [15] T. Haettel, V. Guirardel, and C. Horbez, *Hyperbolic rigidity of higher rank lattices*, *Annales Scientifiques de l’École Normale Supérieure*, 2020, pp. 439–468.
- [16] U. Hamenstädt, *Cocycles, Symplectic Structures and Intersection*, *Geometric and Functional Analysis* **1** (1999), no. 9, 90–140.

- [17] Y. Hao, *Marked length spectrum rigidity of group acts isometrically on a Gromov hyperbolic space*, arXiv preprint arXiv:2312.09019 (2023).
- [18] ———, *Marked length pattern rigidity for arithmetic manifolds*, Ergodic Theory and Dynamical Systems, posted on 2024, 1–25, DOI 10.1017/etds.2024.72.
- [19] A. I. Mal'cev, *On a class of homogeneous spaces*, Izv. Akad. Nauk SSSR Ser. Mat. **13** (1949), 9–32.
- [20] B. Nica, *Proper isometric actions of hyperbolic groups on L^p -spaces*, Compositio Mathematica **149** (2013), no. 5, 773–792.
- [21] B. Nica and J. Špakula, *Strong hyperbolicity*, Groups, Geometry, and Dynamics **10** (2016), no. 3, 951–964.
- [22] J.-P. Otal, *Sur la géométrie symplectique de l'espace des géodésiques d'une variété à courbure négative*, Revista matemática iberoamericana **8** (1992), no. 3, 441–456.
- [23] P. Pansu, *Croissance des boules et des géodésiques fermées dans les nilvariétés*, Ergodic Theory and Dynamical Systems **3** (1983), no. 3, 415–445.
- [24] J. Pardon, *The Hilbert–Smith conjecture for three-manifolds*, Journal of the American Mathematical Society **26** (2013), no. 3, 879–899.
- [25] F. Paulin, *Un groupe hyperbolique est déterminé par son bord*, Journal of the London Mathematical Society **54** (1996), no. 1, 50–74.
- [26] D. Repovš and E. V. Scepina, *A proof of the Hilbert–Smith conjecture for actions by Lipschitz maps*, Mathematische Annalen **308** (1997), 361–364.
- [27] D. Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, Publications Mathématiques de l'IHÉS **50** (1979), 171–202.
- [28] W. van Limbeek, *Riemannian manifolds with local symmetry*, Journal of Topology and Analysis **6** (2014), no. 02, 211–236.
- [29] L. Zippin and D. Montgomery, *Topological Transformation Groups.*, Interscience Publishers, 1955.

UNIVERSITY OF MICHIGAN

Email address: ylhao@umich.edu