

The spectral Einstein functional for the Dirac operator with torsion

Jin Hong^{a,b}, Yong Wang^{b,*}

^a*School of Mathematics and Statistics, Yili Normal University, Yining 835000, China*

^b*School of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, China*

Abstract

In this paper, we compute the spectral Einstein functional associated with the Dirac operator with torsion on even-dimensional spin manifolds without boundary.

Keywords: Dirac operator with torsion; spectral Einstein functional; noncommutative residue.

1. Introduction

Noncommutative residues are an important tool for studying noncommutative geometry. Therefore, much attention has been paid to noncommutative residues. In [2], Connors derived the analog of the conformal four-dimensional Polyakov action using noncommutative residues. Connors proved that the noncommutative residue on a compact manifold M coincides with the Dixmire trace on a pseudo-differential operator of order $\text{-dim} M$ [3]. Furthermore, Connors claimed that the noncommutative residue of the inverse square of the Dirac operator is equivalent to the Einstein-Hilbert action on [3, 4]. Kastler gave a direct proof of this theorem in [6], while Kalau and Walze obtained it simultaneously in [7] in the standard coordinate system. Based on the theory of noncommutative residues proposed by Wozki, Fedosov, etc, [8] the noncommutative residues of the classical elemental algebra of the Boutte de Montville calculus are constructed on compact manifolds of dimension $n > 2$. Using elliptic pseudo-differential operators and noncommutative residues is a natural way to study the spectral Einstein functional and the operator-theoretic interpretation of the gravitational action on bounded manifolds. Concerning the Dirac operator and the signature operator, Wang carried out the computation of noncommutative residues and succeeded in proving the Kastler-Kalau-Walze type theorem for manifolds with boundary [9–11].

Many noncommutative residue of the Dirac operator are studied [9–13]. In [15], Figueroa, et al. introduced a noncommutative integral based on the noncommutative residue [16]. In [17], Ackermann and Tolksdorf proved a generalized version of the well-known Lichnerowicz formula for the square of the most general Dirac operator with torsion D_T on an even-dimensional spin manifold associated with a metric connection with torsion. Wang, etc. [12] gave two kinds of operator-theoretic explanations of the gravitational action of Dirac operators with torsion in the case of 4-dimensional compact manifolds with flat boundary. In [13], Wang etc. gave some new spectral functionals which is the extension of spectral functionals to the noncommutative realm with torsion, and related them to the noncommutative residue for manifolds with boundary about Dirac operators with torsion. In [1], Dabrowski, et al. defined the spectral Einstein functional for a general spectral triple and for the noncommutative torus, they computed the spectral Einstein functional. Pfäffle and Stephan [18] considered orthogonal connections with arbitrary torsion on compact Riemannian manifolds and computed the spectral action. In [19], Pfäffle and Stephan considered compact Riemannian spin manifolds without boundary equipped with orthogonal connections, and investigated the induced Dirac operators. Iochum, et al. [20] derived a commutative spectral triple and studied the spectral action for a rather general geometric setting which includes the (skew-symmetric) torsion and the chiral bag

*Corresponding author.

Email addresses: jhong@nenu.edu.cn (Jin Hong), wangy581@nenu.edu.cn (Yong Wang)

conditions on the boundary. In [21], Pfäffle and Stephan gave a Lichnerowicz type formula for the Dirac operator with torsion. Hanisch, et al. [22] derived a formula for the gravitational part of the spectral action for Dirac operators on 4-dimensional manifolds with totally anti-symmetric torsion. In [23], Dabrowski, et al. examined the metric and Einstein bilinear functionals of differential forms for the Hodge–Dirac operator on an oriented, closed, even-dimensional Riemannian manifold. So it is natural to consider the spectral Einstein functional by the Dirac operator with torsion. The motivation of this paper is to compute the spectral Einstein functional associated with the Dirac operator with torsion on even-dimensional spin manifolds without boundary.

This paper is organized as follows. In Section 2, we give a brief exposition of the Dirac operator with torsion. Using the results in Sec.2, we obtain spectral Einstein functional associated with the Dirac operator with torsion on even-dimensional spin manifolds without boundary in the next section.

2. The Dirac operator with torsion

We give some definitions and basic notions which we will use in this paper.

Let M be a smooth compact oriented Riemannian n -dimensional manifolds without boundary and N be a vector bundle on M . We say that P is a differential operator of Laplace type, if it has locally the form

$$P = -(g^{ij}\partial_i\partial_j + A^i\partial_i + B), \quad (2.1)$$

where ∂_i is a natural local frame on TM , $(g^{ij})_{1 \leq i,j \leq n}$ is the inverse matrix associated to the metric matrix $(g_{ij})_{1 \leq i,j \leq n}$ on M , A^i and B are smooth sections of $\text{End}(N)$ on M (endomorphism). If P satisfies the form (2.1), then there is a unique connection ∇ on N and a unique endomorphism E such that

$$P = -[g^{ij}(\nabla_{\partial_i}\nabla_{\partial_j} - \nabla_{\nabla_{\partial_i}^L\partial_j}) + E],$$

where ∇^L is the Levi-Civita connection on M . Moreover (with local frames of T^*M and N), $\nabla_{\partial_i} = \partial_i + \omega_i$ and E are related to g^{ij} , A^i and B through

$$\begin{aligned} \omega_i &= \frac{1}{2}g_{ij}(A^j + g^{kl}\Gamma_{kl}^j \mathbf{id}), \\ E &= B - g^{ij}(\partial_i(\omega_j) + \omega_i\omega_j - \omega_k\Gamma_{ij}^k), \end{aligned}$$

where Γ_{kl}^j is the Christoffel coefficient of ∇^L .

Let ∇^T denote the metric connection

$$\langle \nabla_X^T Y, Z \rangle = \langle \nabla_X^L Y, Z \rangle + T(X, Y, Z),$$

where T is a there form.

Let M be an $n = 2m$ dimensional ($n \geq 3$) spin manifold, we can lift ∇^T to $\nabla^{S(TM),T}$ on $S(TM)$. The Dirac operator with torsion D_T is defined as:

$$\begin{aligned} D_T &= \sum_{j=1}^n c(e_j)\nabla_{e_j}^{S(TM),T} \\ &= \sum_{j=1}^n c(e_j) \left(e_j + \frac{1}{4} \sum_{l,t=1}^n \langle \nabla_{e_j}^T e_l, e_t \rangle c(e_l)c(e_t) \right) \\ &= D + \frac{1}{4} \sum_{j,l,t=1}^n T(e_j, e_l, e_t) c(e_j)c(e_l)c(e_t) \\ &= D + \frac{3}{2} \sum_{1 \leq j < l < t \leq n} T(e_j, e_l, e_t) c(e_j)c(e_l)c(e_t), \end{aligned}$$

where D is the Dirac operator induced by the Levi-Civita connection, and $c(e_j)$ be the Clifford action which satisfies the relation

$$c(e_i)c(e_j) + c(e_j)c(e_i) = -2g^M(e_i, e_j) = -2\delta_i^j.$$

By lemma 2.1 in [13] ,we get

$$D_T^2 = -g^{ab}(\nabla_{\partial_a}\nabla_{\partial_b} - \nabla_{\nabla_{\partial_a}^L\partial_b}) + E, \quad (2.2)$$

where

$$E = \frac{3}{2}dT + \frac{1}{4}s - \frac{3}{4}\|T\|^2, \quad (2.3)$$

and

$$\nabla_{\partial_a} = \partial_a - \tilde{T}_a = \partial_a + \frac{1}{4}\sum_{s,t=1}^n \langle \nabla_{\partial_a}^L e_s, e_t \rangle c(e_s)c(e_t) + \frac{3}{2}\sum_{1 \leq j < l \leq n} T(\partial_a, e_j, e_l)c(e_j)c(e_l). \quad (2.4)$$

So we have

$$\tilde{T}_a = -\frac{1}{4}\sum_{s,t=1}^n \langle \nabla_{\partial_a}^L e_s, e_t \rangle c(e_s)c(e_t) - \frac{3}{2}\sum_{1 \leq j < l \leq n} T(\partial_a, e_j, e_l)c(e_j)c(e_l). \quad (2.5)$$

In normal coordinates, \tilde{T}_a is expanded near $x = 0$ by Taylor expansion. By $\partial_l \langle \nabla_{\partial_a}^L e_s, e_t \rangle(x_0) = \frac{1}{2}R_{lats}(x_0)$ we get

$$\tilde{T}_a = T_a + T_{ab}x^b + O(x^2), \quad (2.6)$$

where

$$T_a = -\frac{3}{2}\sum_{1 \leq j < l \leq n} T(\partial_a, e_j, e_l)(x_0)c(e_j)c(e_l), \quad (2.7)$$

and

$$T_{ab} = -\frac{1}{8}\sum_{s,t=1}^n R_{bats}(x_0)c(e_s)c(e_t) - \frac{3}{2}\sum_{1 \leq j < l \leq n} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b}(x_0)c(e_j)c(e_l). \quad (2.8)$$

Lemma 2.1. [1] *The following identities hold:*

$$\sigma_{-2m}(\Delta_{T,E}^{-m}) = \|\xi\|^{-2m-2} \sum_{a,b,j,k=1}^{2m} \left(\delta_{ab} - \frac{m}{3}R_{ajbk}x^jx^k \right) \xi_a\xi_b + O(\mathbf{x}^3); \quad (2.9)$$

$$\sigma_{-2m-1}(\Delta_{T,E}^{-m}) = \frac{-2mi}{3}\|\xi\|^{-2m-2} \sum_{a,k=1}^{2m} Ric_{ak}x^k\xi_a - 2mi\|\xi\|^{-2m-2} \sum_{a,b=1}^{2m} (T_a\xi_a + T_{ab}x^b\xi_a) + O(\mathbf{x}^2); \quad (2.10)$$

$$\begin{aligned} \sigma_{-2m-2}(\Delta_{T,E}^{-m}) &= \frac{m(m+1)}{3}\|\xi\|^{-2m-4} \sum_{a,b=1}^{2m} Ric_{ab}\xi_a\xi_b \\ &\quad - 2m(m+1)\|\xi\|^{-2m-4} \sum_{a,b=1}^{2m} T_aT_b\xi_a\xi_b + m \sum_{a,b=1}^{2m} (T_aT_a - T_{aa})\|\xi\|^{-2m-2} \\ &\quad + 2m(m+1)\|\xi\|^{-2m-4} \sum_{a,b=1}^{2m} T_{ab}\xi_a\xi_b - mE\|\xi\|^{-2m-2} + O(\mathbf{x}), \end{aligned} \quad (2.11)$$

where R_{ajbk} and Ric_{ak} are the components of the Riemann and Ricci tensor.

By (2.3),(2.7),(2.8) and lemma 2.1, we get the following lemma.

Lemma 2.2. *General dimensional symbols about Dirac operator with torsion are given,*

$$\sigma_{-2m}(\Delta_{T,E}^{-m}) = \|\xi\|^{-2m-2} \sum_{a,b,j,k=1}^{2m} \left(\delta_{ab} - \frac{m}{3} R_{ajbk} x^j x^k \right) \xi_a \xi_b + O(\mathbf{x}^3); \quad (2.12)$$

$$\sigma_{-2m-1}(\Delta_{T,E}^{-m}) = \frac{-2m\sqrt{-1}}{3} \|\xi\|^{-2m-2} \sum_{a,b=1}^{2m} \text{Ric}_{ab} x^b \xi_a \quad (2.13)$$

$$\begin{aligned} & + 3m\sqrt{-1} \|\xi\|^{-2m-2} \sum_{a,j < l=1}^{2m} T(e_a, e_j, e_l)(x_0) c(e_j) c(e_l) \xi_a \\ & + \frac{m\sqrt{-1}}{4} \|\xi\|^{-2m-2} \sum_{a,b,t,s=1}^{2m} \text{R}_{bats}(x_0) c(e_s) c(e_t) x^b \xi_a \\ & + 3m\sqrt{-1} \|\xi\|^{-2m-2} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b}(x_0) c(e_j) c(e_l) x^b \xi_a + O(\mathbf{x}^2); \\ \sigma_{-2m-2}(\Delta_{T,E}^{-m}) = & \frac{m(m+1)}{3} \|\xi\|^{-2m-4} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b \quad (2.14) \\ & - \frac{9m(m+1)}{2} \|\xi\|^{-2m-4} \sum_{a,b,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)(x_0) T(e_b, e_{\hat{j}}, e_{\hat{l}})(x_0) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \xi_a \xi_b \\ & + \frac{9m}{4} \|\xi\|^{-2m-2} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)(x_0) T(e_a, e_{\hat{j}}, e_{\hat{l}})(x_0) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \\ & + \frac{3m}{2} \|\xi\|^{-2m-2} \sum_{a,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a}(x_0) c(e_j) c(e_l) \\ & - \frac{m(m+1)}{4} \|\xi\|^{-2m-4} \sum_{a,b,t,s=1}^{2m} \text{R}_{bats}(x_0) c(e_s) c(e_t) \xi_a \xi_b \\ & - 3m(m+1) \|\xi\|^{-2m-4} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b}(x_0) c(e_j) c(e_l) \xi_a \xi_b \\ & - m \|\xi\|^{-2m-2} \left(\frac{1}{4}s - \frac{3}{4}\|T\|^2 \right) \\ & - \frac{3m}{2} \|\xi\|^{-2m-2} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t)(x_0) c(e_i) c(e_j) c(e_k) c(e_t) + O(\mathbf{x}), \end{aligned}$$

where R_{ajbk} and Ric_{ak} are the components of the Riemann and Ricci tensor, s is the scalar curvature.

3. The spectral Einstein functional for the Dirac operator with torsion

For a pseudo-differential operator P , acting on sections of a spinor bundle over an even n -dimensional compact Riemannian spin manifold M , the analogue of the volume element in noncommutative geometry is the operator $D^{-n} =: ds^n$. And pertinent operators are realized as pseudodifferential operators on the spaces of sections. Extending previous definitions by Connes [14], a noncommutative integral was introduced in [15]

based on the noncommutative residue [16], combine (1.4) in [5] and [6], using the definition of the residue:

$$\int Pds^n := \text{Wres } PD^{-n} := \int_{S^*M} \text{tr} [\sigma_{-n}(PD^{-n})](x, \xi), \quad (3.1)$$

where $\sigma_{-n}(PD^{-n})$ denotes the $(-n)$ th order piece of the complete symbols of PD^{-n} , tr as shorthand of trace. This section is designed to get the metric functional and the spectral Einstein functional for the Dirac operator with torsion defined in [1].

Definition 3.1. [23] *If $(\mathcal{L}, D, \mathcal{H})$ is an n -summable spectral triple, let Ω_D^1 be the \mathcal{L} bimodule of one-forms generated by \mathcal{L} and $[D, \mathcal{L}]$. Moreover, assume that there exists a generalised algebra of pseudodifferential operators which contains \mathcal{L}, D , and $|D|^\ell$ for $\ell \in \mathbb{Z}$ with a tracial state Wres over this algebra (called a noncommutative residue), which identically vanishes on $T|D|^{-k}$ for any $k > n$ and a zero-order operator T (an operator in the algebra generated by \mathcal{L} and $\Omega^1(\mathcal{L})$). Then, for $u, w \in \Omega_D^1(\mathcal{L})$, we call*

$$\mathcal{A}(u, w) := \text{Wres}(uw|D|^{-n}), \quad (3.2)$$

the metric functional, and

$$\mathcal{B}(u, w) := \text{Wres}(u\{D, w\}D|D|^{-n}). \quad (3.3)$$

the Einstein functional.

Proposition 3.2. *Let M be an $n = 2m$ dimensional ($n \geq 3$) closed spin manifold and D_T a Dirac operator with torsion, then $(C^\infty(M), D_T, L^2(M, S(TM)))$ is an n -summable spectral triple.*

Theorem 3.3. *Let M be an $n = 2m$ dimensional ($n \geq 3$) spin manifold, the metric functional \mathcal{A}_{D_T} and the spectral Einstein functional \mathcal{B}_{D_T} are equal to*

$$\mathcal{A}_{D_T} = \text{Wres}\left(c(v)c(w)D_T^{-n}\right) = -2^m \frac{2\pi^m}{\Gamma(m)} \int_M g(v, w)d\text{Vol}_M; \quad (3.4)$$

$$\begin{aligned} \mathcal{B}_{D_T} &= \text{Wres}\left(c(v)(c(w)D_T + D_T c(w))D_T^{-n+1}\right) \\ &= 2^m \frac{2\pi^m}{\Gamma(m)} \int_M \left\{ -\frac{1}{6}\mathbb{G}(v, w) + \frac{73}{16}\|T\|^2 g(v, w) - \frac{25}{16} \sum_{j,l=1}^{2m} T(v, e_j, e_l)T(w, e_j, e_l) \right. \\ &\quad \left. + \frac{11}{4} \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) + \frac{17}{4} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \right\} d\text{Vol}_M, \end{aligned} \quad (3.5)$$

where $g(v, w) = \sum_{a=1}^n v_a w_a$ and $\mathbb{G}(v, w) = \text{Ric}(v, w) - \frac{1}{2}sg(v, w)$, $v = \sum_{a=1}^n v_a e_a$, $w = \sum_{b=1}^n w_b e_b$.

Proof. The proof of the \mathcal{A}_{D_T} formula for the measure function is obvious. Below we prove that the spectral Einstein function \mathcal{B}_{D_T} , splitting it into two parts as follows:

$$\mathcal{B}_1 = \text{Wres}\left(c(v)c(w)D_T^{-n+2}\right); \quad (3.6)$$

$$\mathcal{B}_2 = \text{Wres}\left(c(v)D_T c(w)D_T D_T^{-n}\right). \quad (3.7)$$

Part I) $\mathcal{B}_1 = \text{Wres}\left(c(v)c(w)D_T^{-n+2}\right)$, $\mathcal{P}_1 = c(v)c(w)$. Let $n = 2m$, by (3.1), we need to compute $\int_{S^*M} \text{tr} [\sigma_{-2m}(\mathcal{P}_1 D_T^{-2m+2})](x, \xi)$. Based on the algorithm yielding the principal symbol of a product of pseudo-differential operators in terms of the principal symbols of the factors, by (2.14) in lemma 2.2 we have

$$\begin{aligned}
\sigma_{-2m}(D_T^{-2m+2}) = & \frac{m(m-1)}{3} \|\xi\|^{-2m-2} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b \\
& - \frac{9m(m-1)}{2} \|\xi\|^{-2m-2} \sum_{a,b,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)(x_0) T(e_b, e_{\hat{j}}, e_{\hat{l}})(x_0) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \xi_a \xi_b \\
& + \frac{9(m-1)}{4} \|\xi\|^{-2m} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)(x_0) T(e_a, e_{\hat{j}}, e_{\hat{l}})(x_0) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \\
& + \frac{3(m-1)}{2} \|\xi\|^{-2m} \sum_{a,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a}(x_0) c(e_j) c(e_l) \\
& - \frac{m(m-1)}{4} \|\xi\|^{-2m-2} \sum_{a,b,t,s=1}^{2m} \text{R}_bats(x_0) c(e_s) c(e_t) \xi_a \xi_b \\
& - 3m(m-1) \|\xi\|^{-2m-2} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b}(x_0) c(e_j) c(e_l) \xi_a \xi_b \\
& - (m-1) \|\xi\|^{-2m} \left(\frac{1}{4} R^g - \frac{3}{4} \|T\|^2 \right) \\
& - \frac{3(m-1)}{2} \|\xi\|^{-2m} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t)(x_0) c(e_i) c(e_j) c(e_k) c(e_t) + O(\mathbf{x}).
\end{aligned} \tag{3.8}$$

By (3.8) and $\mathcal{P}_1 = c(v)c(w)$, we get

$$\begin{aligned}
\sigma_{-2m}(\mathcal{P}_1 D_T^{-2m+2}) = & \frac{m(m-1)}{3} \|\xi\|^{-2m-2} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b c(v) c(w) \\
& - \frac{9m(m-1)}{2} \|\xi\|^{-2m-2} \sum_{a,b,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_b, e_j, e_l)(x_0) T(e_a, e_{\hat{j}}, e_{\hat{l}})(x_0) c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \xi_a \xi_b \\
& + \frac{9(m-1)}{4} \|\xi\|^{-2m} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)(x_0) T(e_a, e_{\hat{j}}, e_{\hat{l}})(x_0) c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \\
& + \frac{3(m-1)}{2} \|\xi\|^{-2m} \sum_{a,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a}(x_0) c(v) c(w) c(e_j) c(e_l) \\
& - \frac{m(m-1)}{4} \|\xi\|^{-2m-2} \sum_{a,b,t,s=1}^{2m} \text{R}_bats(x_0) c(v) c(w) c(e_s) c(e_t) \xi_a \xi_b \\
& - 3m(m-1) \|\xi\|^{-2m-2} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b}(x_0) c(v) c(w) c(e_j) c(e_l) \xi_a \xi_b \\
& - (m-1) \|\xi\|^{-2m} \left(\frac{1}{4} R^g - \frac{3}{4} \|T\|^2 \right) c(v) c(w) \\
& - \frac{3(m-1)}{2} \|\xi\|^{-2m} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t)(x_0) c(v) c(w) c(e_i) c(e_j) c(e_k) c(e_t) + O(\mathbf{x}).
\end{aligned} \tag{3.9}$$

Below, we compute each term of $\int_{\|\xi\|=1} \text{tr}[\sigma_{-2m}(\mathcal{P}_1 D^{-2m+2})](x, \xi) \sigma(\xi)$ in turn. Based on the relation

of the Clifford action $\text{tr}(c(\mathcal{X})c(\mathcal{Y})) = -g(\mathcal{X}, \mathcal{Y})$ and $\int_{\|\xi\|=1} \xi_a \xi_b \sigma(\xi) = \frac{1}{n} \delta_a^b \text{Vol}(S^{n-1})$, we get the following equations.

(I-A)

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{m(m-1)}{3} \|\xi\|^{-2m-2} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b c(v) c(w) \right\} (x_0) \sigma(\xi) \\ &= -\frac{m-1}{6} s g(v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.10)$$

where s be the scalar curvature.

(I-B)

Let $v = \sum_{p=1}^{2m} v_p e_p$, $w = \sum_{q=1}^{2m} w_q e_q$, and based on the relation of the Clifford action and $\text{tr}\mathcal{X}\mathcal{Y} = \text{tr}\mathcal{Y}\mathcal{X}$, we can obtain the equality

$$\begin{aligned} & \text{tr} \left(\sum_{j \neq l, \hat{j} \neq \hat{l}=1}^{2m} c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \\ &= \sum_{j \neq l, \hat{j} \neq \hat{l}=1}^{2m} \left[v_l w_l \delta_j^{\hat{j}} - v_l w_j \delta_l^{\hat{j}} - v_{\hat{j}} w_l \delta_j^{\hat{l}} + v_{\hat{j}} w_j \delta_l^{\hat{l}} - v_l w_l \delta_j^{\hat{j}} + v_l w_{\hat{j}} \delta_j^{\hat{l}} + v_j w_{\hat{j}} \delta_l^{\hat{j}} - v_j w_j \delta_l^{\hat{l}} \right. \\ & \quad \left. - \delta_j^{\hat{l}} \delta_l^{\hat{j}} g(v, w) + \delta_j^{\hat{j}} \delta_l^{\hat{l}} g(v, w) \right] \text{tr}[id], \end{aligned} \quad (3.11)$$

where $v_l = g(v, e_l)$, $w_j = g(w, e_j)$, then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{9m(m-1)}{2} \|\xi\|^{-2m-2} \sum_{a,b,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_b, e_{\hat{j}}, e_{\hat{l}}) c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \xi_a \xi_b \right\} (x_0) \sigma(\xi) \\ &= -\frac{9(m-1)}{16} \sum_{a,b,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) \text{tr} \left(c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \text{Vol}(S^{n-1}) \\ &= -\frac{9(m-1)}{8} \sum_{a,j \neq l=1}^{2m} T^2(e_a, e_j, e_l) g(v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.12)$$

(I-C)

Similar to (I-B), one obtains

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{9(m-1)}{4} \|\xi\|^{-2m} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right\} (x_0) \sigma(\xi) \\ &= \frac{9(m-1)}{8} \sum_{a,j \neq l=1}^{2m} T^2(e_a, e_j, e_l) g(v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.13)$$

(I-D)

Since

$$\text{tr} \left(\sum_{j \neq l=1}^{2m} c(v) c(w) c(e_j) c(e_l) \right) = \sum_{j \neq l=1}^{2m} \left[v_j w_l + v_l w_j \right] \text{tr}[id], \quad (3.14)$$

and

$$\frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a}(x_0) = \nabla_{e_a}(T)(e_a, e_j, e_l)(x_0), \quad (3.15)$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{3(m-1)}{2} \|\xi\|^{-2m} \sum_{a,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a} c(v)c(w)c(e_j)c(e_l) \right\} (x_0) \sigma(\xi) \\ &= \frac{3(m-1)}{4} \sum_{a,j \neq l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a}(x_0) \text{tr} \left(c(v)c(w)c(e_j)c(e_l) \right) \text{Vol}(S^{n-1}) \\ &= \frac{3(m-1)}{4} \sum_{a,j \neq l=1}^{2m} \nabla_{e_a}(T)(e_a, e_j, e_l)(x_0) \left(v_j w_l + v_l w_j \right) \text{tr}[id] \text{Vol}(S^{n-1}) \\ &= 0. \end{aligned} \quad (3.16)$$

(I-E)
Since

$$\begin{aligned} & \int_{\|\xi\|=1} \left(\|\xi\|^{-2m-2} \sum_{a,b,t,s=1}^{2m} R_{bats} \xi_a \xi_b \right) (x_0) \sigma(\xi) \\ &= \frac{1}{2m} \sum_{a,t,s=1}^{2m} R_{aats} \text{Vol}(S^{n-1}) \\ &= 0, \end{aligned} \quad (3.17)$$

then

$$\int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m(m-1)}{4} \|\xi\|^{-2m-2} \sum_{a,b,t,s=1}^{2m} R_{bats} c(v)c(w)c(e_s)c(e_t) \xi_a \xi_b \right\} (x_0) \sigma(\xi) = 0. \quad (3.18)$$

(I-F)
Similar to (3.16)

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ -3m(m-1) \|\xi\|^{-2m-2} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b} c(v)c(w)c(e_j)c(e_l) \xi_a \xi_b \right\} (x_0) \sigma(\xi) \\ &= 0. \end{aligned} \quad (3.19)$$

(I-G)
Similarly to (I-A), we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ -(m-1) \|\xi\|^{-2m} \left(\frac{1}{4}s - \frac{3}{4}\|T\|^2 \right) c(v)c(w) \right\} (x_0) \sigma(\xi) \\ &= (m-1) \left(\frac{1}{4}s - \frac{3}{4}\|T\|^2 \right) g(v,w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.20)$$

Summing from (I-A) to (I-G) in turn, we get

$$\int_{\|\xi\|=1} \text{tr} \left\{ \sigma_{-2m} (c(v)c(w) D_T^{-2m+2}) \right\} (x_0) \sigma(\xi) \quad (3.21)$$

$$= \frac{m-1}{12} sg(v, w) \text{tr}[id] \text{Vol}(S^{n-1}) - \frac{3(m-1)}{4} \|T\|^2 g(v, w) \text{tr}[id] \text{Vol}(S^{n-1}).$$

Since $\text{tr}[id] = 2^m$ and $\text{Vol}(S^{n-1}) = \frac{2\pi^m}{\Gamma(m)}$, by (3.1) we obtain

$$\begin{aligned} \mathcal{B}_1 &= \text{Wres} \left(c(v)c(w)D_T^{-n+2} \right) \\ &= 2^m \frac{2\pi^m}{\Gamma(m)} \int_M \left\{ \frac{m-1}{12} sg(v, w) + \frac{73}{16} \|T\|^2 g(v, w) \right\} d\text{Vol}_M. \end{aligned} \quad (3.22)$$

Part II) $\mathcal{B}_2 = \text{Wres} \left(c(v)D_T c(w)D_T D_T^{-n} \right)$. $\mathcal{P}_2 = c(v)D_T c(w)D_T$, and $c(v)D_T := \mathcal{A}$, $c(w)D_T := \mathcal{B}$. Let $n = 2m$, by (3.1), we need to compute $\int_{S^* M} \text{tr} [\sigma_{-2m} (\mathcal{P}_1 D_T^{-2m})] (x, \xi)$. Based on the algorithm yielding the principal symbol of a product of pseudo-differential operators in terms of the principal symbols of the factors, we have

$$\begin{aligned} \sigma_{-2m} (\mathcal{A}\mathcal{B}D_T^{-2m}) &= \left\{ \sum_{|\alpha|=0}^{\infty} \frac{(-i)^{|\alpha|}}{\alpha!} \partial_\xi^\alpha [\sigma(\mathcal{A}\mathcal{B})] \partial_x^\alpha [\sigma(D_T^{-2m})] \right\}_{-2m} \\ &= \sigma_0(\mathcal{A}\mathcal{B})\sigma_{-2m}(D_T^{-2m}) + \sigma_1(\mathcal{A}\mathcal{B})\sigma_{-2m-1}(D_T^{-2m}) + \sigma_2(\mathcal{A}\mathcal{B})\sigma_{-2m-2}(D_T^{-2m}) \\ &\quad + (-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_2(\mathcal{A}\mathcal{B})] \partial_{x_j} [\sigma_{-2m-1}(D_T^{-2m})] + (-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_1(\mathcal{A}\mathcal{B})] \partial_{x_j} [\sigma_{-2m}(D_T^{-2m})] \\ &\quad - \frac{1}{2} \sum_{j,l=1}^{2m} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(\mathcal{A}\mathcal{B})] \partial_{x_j} \partial_{x_l} [\sigma_{-2m}(D_T^{-2m})]. \end{aligned} \quad (3.23)$$

Lemma 3.4. [13] The symbols of D_T are given

$$\begin{aligned} \sigma_0(D_T) &= -\frac{1}{4} \sum_{p,s,t=1}^{2m} w_{s,t}(e_p)c(e_p)c(e_s)c(e_t) + \frac{1}{4} \sum_{f<\alpha<\beta=1}^{2m} T(e_f, e_\alpha, e_\beta)c(e_f)c(e_\alpha)c(e_\beta); \\ \sigma_1(D_T) &= \sqrt{-1}c(\xi). \end{aligned}$$

By lemma 3.4 and $\mathcal{A} = c(v)D_T$, $\mathcal{B} = c(w)D_T$, we obtain the following lemma.

Lemma 3.5. The symbols of \mathcal{A} and \mathcal{B} are given

$$\begin{aligned} \sigma_0(\mathcal{A}) &= -\frac{1}{4} \sum_{p,s,t=1}^{2m} w_{s,t}(e_p)c(v)c(e_p)c(e_s)c(e_t) + \frac{1}{4} \sum_{f<\alpha<\beta=1}^{2m} T(e_f, e_\alpha, e_\beta)c(v)c(e_f)c(e_\alpha)c(e_\beta); \\ \sigma_1(\mathcal{A}) &= \sqrt{-1}c(v)c(\xi); \\ \sigma_0(\mathcal{B}) &= -\frac{1}{4} \sum_{p,s,t=1}^{2m} w_{s,t}(e_p)c(w)c(e_p)c(e_s)c(e_t) + \frac{1}{4} \sum_{f<\alpha<\beta=1}^{2m} T(e_f, e_\alpha, e_\beta)c(w)c(e_f)c(e_\alpha)c(e_\beta); \\ \sigma_1(\mathcal{B}) &= \sqrt{-1}c(w)c(\xi). \end{aligned}$$

Further, by the composition formula of pseudodifferential operators, we get the following lemma.

Lemma 3.6. The symbols of $\mathcal{A}\mathcal{B}$ are given

$$\sigma_0(\mathcal{A}\mathcal{B}) = \sigma_0(\mathcal{A})\sigma_0(\mathcal{B}) + (-i)\partial_{\xi_j} [\sigma_1(\mathcal{A})] \partial_{x_j} [\sigma_0(\mathcal{B})] + (-i)\partial_{\xi_j} [\sigma_0(\mathcal{A})] \partial_{x_j} [\sigma_1(\mathcal{B})] \quad (3.24)$$

$$\begin{aligned}
&= \frac{1}{16} \sum_{p,s,t,\hat{p},\hat{s},\hat{t}=1}^{2m} w_{s,t}(e_p) w_{\hat{s},\hat{t}}(e_{\hat{p}}) c(v) c(e_p) c(e_s) c(e_t) c(w) c(e_{\hat{p}}) c(e_{\hat{s}}) c(e_{\hat{t}}) \\
&\quad - \frac{1}{16} \sum_{f < \alpha < \beta, \hat{p}, \hat{s}, \hat{t}=1}^{2m} T(e_f, e_\alpha, e_\beta) w_{\hat{s},\hat{t}}(e_{\hat{p}}) c(v) c(e_f) c(e_\alpha) c(e_\beta) c(w) c(e_{\hat{p}}) c(e_{\hat{s}}) c(e_{\hat{t}}) \\
&\quad - \frac{1}{16} \sum_{p,s,t,\hat{f} < \hat{\alpha} < \hat{\beta}=1}^{2m} w_{s,t}(e_p) T(e_{\hat{f}}, e_{\hat{\alpha}}, e_{\hat{\beta}}) c(v) c(e_p) c(e_s) c(e_t) c(w) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}}) \\
&\quad + \frac{1}{16} \sum_{f < \alpha < \beta, \hat{f} < \hat{\alpha} < \hat{\beta}=1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_{\hat{f}}, e_{\hat{\alpha}}, e_{\hat{\beta}}) c(v) c(e_f) c(e_\alpha) c(e_\beta) c(w) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}}) \\
&\quad + \frac{1}{8} \sum_{j,p,s,t=1}^{2m} R_{jpst} c(v) c(dx_j) c(w) c(e_p) c(e_s) c(e_t) \\
&\quad - \frac{1}{4} \sum_{p,s,t,j,\gamma=1}^{2m} w_{s,t}(e_p) \partial x_j(w_\gamma) c(v) c(dx_j) c(e_\gamma) c(e_p) c(e_s) c(e_t) \\
&\quad + \frac{1}{4} \sum_{f < \alpha < \beta, j=1}^{2m} \frac{\partial T(e_f, e_\alpha, e_\beta)}{\partial x_j} c(v) c(dx_j) c(e_w) c(e_f) c(e_\alpha) c(e_\beta) \\
&\quad + \frac{1}{4} \sum_{f < \alpha < \beta, j, \gamma=1}^{2m} T(e_f, e_\alpha, e_\beta) \partial x_j(w_\gamma) c(v) c(dx_j) c(e_\gamma) c(e_f) c(e_\alpha) c(e_\beta);
\end{aligned}$$

$$\sigma_1(\mathcal{A}\mathcal{B}) = \sigma_1(\mathcal{A})\sigma_0(\mathcal{B}) + \sigma_0(\mathcal{A})\sigma_1(\mathcal{B}) + (-i)\partial_{\xi_j} [\sigma_1(\mathcal{A})] \partial_{x_j} [\sigma_1(\mathcal{B})] \quad (3.25)$$

$$\begin{aligned}
&= -\frac{\sqrt{-1}}{4} \sum_{p,s,t=1}^{2m} w_{s,t}(e_p) c(v) c(\xi) c(w) c(e_p) c(e_s) c(e_t) \\
&\quad + \frac{\sqrt{-1}}{4} \sum_{f < \alpha < \beta=1}^{2m} T(e_f, e_\alpha, e_\beta) c(v) c(\xi) c(w) c(e_f) c(e_\alpha) c(e_\beta) \\
&\quad - \frac{\sqrt{-1}}{4} \sum_{p,s,t=1}^{2m} w_{s,t}(e_p) c(w) c(\xi) c(v) c(e_p) c(e_s) c(e_t) \\
&\quad + \frac{\sqrt{-1}}{4} \sum_{f < \alpha < \beta=1}^{2m} T(e_f, e_\alpha, e_\beta) c(w) c(\xi) c(v) c(e_f) c(e_\alpha) c(e_\beta) \\
&\quad + \sqrt{-1} \sum_{j,\gamma=1}^{2m} \partial x_j(w_\gamma) c(v) c(dx_j) c(e_\gamma) c(\xi);
\end{aligned}$$

$$\sigma_2(\mathcal{A}\mathcal{B}) = \sigma_1(\mathcal{A})\sigma_1(\mathcal{B}) = -c(v)c(\xi)c(w)c(\xi). \quad (3.26)$$

Next, with (3.23), we compute each term of $\int_{\|\xi\|=1} \text{tr}[\sigma_{-2m}(\mathcal{AB}D_T^{-2m+2})](x, \xi)\sigma(\xi)$ in turn.

(II-1) For $\sigma_0(\mathcal{A}\mathcal{B})\sigma_{-2m}(D^{-2m})$:

According to (2.12) in lemma 2.2 and (3.24) in lemma 3.6, where $w_{s,t}(e_p) = 0$, we get

$$\begin{aligned}
&\sigma_0(\mathcal{A}\mathcal{B})\sigma_{-2m}(D^{-2m})(x_0) \\
&= \frac{\|\xi\|^2}{16} \sum_{f < \alpha < \beta, \hat{f} < \hat{\alpha} < \hat{\beta}=1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_{\hat{f}}, e_{\hat{\alpha}}, e_{\hat{\beta}}) c(v) c(e_f) c(e_\alpha) c(e_\beta) c(w) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}})
\end{aligned}$$

$$\begin{aligned}
& + \frac{\|\xi\|^2}{8} \sum_{j,p,s,t=1}^{2m} R_{jpst} c(v) c(dx_j) c(w) c(e_p) c(e_s) c(e_t) \\
& + \frac{\|\xi\|^2}{4} \sum_{f < \alpha < \beta, j=1}^{2m} \frac{\partial T(e_f, e_\alpha, e_\beta)}{\partial x_j} c(v) c(dx_j) c(e_w) c(e_f) c(e_\alpha) c(e_\beta) \\
& + \frac{\|\xi\|^2}{4} \sum_{f < \alpha < \beta, j, \gamma=1}^{2m} T(e_f, e_\alpha, e_\beta) \partial x_j (w_\gamma) c(v) c(dx_j) c(e_\gamma) c(e_f) c(e_\alpha) c(e_\beta).
\end{aligned}$$

(II-1-A)

Let $v = \sum_{p=1}^{2m} v_p e_p$, $w = \sum_{q=1}^{2m} w_q e_q$, and based on the relation of the Clifford action and $\text{tr} \mathcal{X} \mathcal{Y} = \text{tr} \mathcal{Y} \mathcal{X}$, we can obtain the equality

$$\begin{aligned}
& \text{tr} \left(\sum_{f \neq \alpha \neq \beta, \hat{f} \neq \hat{\alpha} \neq \hat{\beta} = 1}^{2m} c(v) c(e_f) c(e_\alpha) c(e_\beta) c(w) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}}) \right) \quad (3.28) \\
& = \sum_{f \neq \alpha \neq \beta, \hat{f} \neq \hat{\alpha} \neq \hat{\beta} = 1}^{2m} \left[v_f \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\beta w_{\hat{f}} - v_f \delta_f^{\hat{\beta}} \delta_{\hat{f}}^\beta w_{\hat{\alpha}} - v_f \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\beta w_{\hat{f}} + v_f \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\beta w_{\hat{\beta}} + v_f \delta_f^{\hat{f}} \delta_{\hat{\beta}}^\beta w_{\hat{\alpha}} - v_f \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\beta w_{\hat{\beta}} \right. \\
& \quad - v_\alpha \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\beta w_{\hat{f}} + v_\alpha \delta_f^{\hat{\beta}} \delta_{\hat{f}}^\beta w_{\hat{\alpha}} + v_\alpha \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\beta w_{\hat{f}} - v_\alpha \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\beta w_{\hat{\beta}} - v_\alpha \delta_f^{\hat{f}} \delta_{\hat{\beta}}^\beta w_{\hat{\alpha}} + v_\alpha \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\beta w_{\hat{\beta}} \\
& \quad + v_\beta \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha w_{\hat{f}} - v_\beta \delta_f^{\hat{\beta}} \delta_{\hat{f}}^\alpha w_{\hat{\alpha}} - v_\beta \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\alpha w_{\hat{f}} + v_\beta \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\alpha w_{\hat{\beta}} + v_\beta \delta_f^{\hat{f}} \delta_{\hat{\beta}}^\alpha w_{\hat{\alpha}} - v_\beta \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\alpha w_{\hat{\beta}} \\
& \quad + v_f \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha w_\beta - v_f \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\beta w_\alpha - v_f \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\alpha w_\beta + v_f \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\beta w_\alpha + v_f \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\alpha w_f - v_f \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha w_f \\
& \quad - v_\alpha \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha w_\beta + v_\alpha \delta_f^{\hat{\beta}} \delta_{\hat{f}}^\beta w_\alpha + v_\alpha \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\beta w_\beta - v_\alpha \delta_f^{\hat{f}} \delta_{\hat{\beta}}^\beta w_\alpha - v_\alpha \delta_f^{\hat{f}} \delta_{\hat{\beta}}^\alpha w_f + v_\alpha \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha w_f \\
& \quad + v_\beta \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\alpha w_\beta - v_\beta \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\beta w_\alpha - v_\beta \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\alpha w_\beta + v_\beta \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\beta w_\alpha + v_\beta \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\alpha w_f - v_\beta \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\alpha w_f \\
& \quad - \delta_f^{\hat{\beta}} \delta_{\hat{\alpha}}^\alpha \delta_{\hat{f}}^\beta g(v, w) + \delta_f^{\hat{\beta}} \delta_{\hat{f}}^\alpha \delta_{\hat{\alpha}}^\beta g(v, w) + \delta_f^{\hat{\alpha}} \delta_{\hat{\beta}}^\alpha \delta_{\hat{f}}^\beta g(v, w) - \delta_f^{\hat{\alpha}} \delta_{\hat{f}}^\alpha \delta_{\hat{\beta}}^\beta g(v, w) - \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\alpha \delta_{\hat{\beta}}^\beta g(v, w) + \delta_f^{\hat{f}} \delta_{\hat{\alpha}}^\alpha \delta_{\hat{\beta}}^\beta g(v, w) \Big] \text{tr}[id],
\end{aligned}$$

then

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ \frac{\|\xi\|^2}{16} \sum_{f < \alpha < \beta, \hat{f} < \hat{\alpha} < \hat{\beta} = 1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_{\hat{f}}, e_{\hat{\alpha}}, e_{\hat{\beta}}) c(v) c(e_f) c(e_\alpha) c(e_\beta) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}}) \right\} (x_0) \sigma(\xi) \quad (3.29) \\
& = \int_{\|\xi\|=1} \text{tr} \left\{ \frac{\|\xi\|^2}{16 \times 36} \sum_{f \neq \alpha \neq \beta, \hat{f} \neq \hat{\alpha} \neq \hat{\beta} = 1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_{\hat{f}}, e_{\hat{\alpha}}, e_{\hat{\beta}}) c(v) c(e_f) c(e_\alpha) c(e_\beta) c(e_{\hat{f}}) c(e_{\hat{\alpha}}) c(e_{\hat{\beta}}) \right\} (x_0) \sigma(\xi) \\
& = \sum_{f \neq \alpha \neq \beta = 1}^{2m} \left[\frac{1}{96} T^2(e_f, e_\alpha, e_\beta) g(v, w) - \frac{1}{16} T(v, e_\alpha, e_\beta) T(w, e_\alpha, e_\beta) \right] \text{tr}[id] \text{Vol}(S^{n-1}).
\end{aligned}$$

(II-1-B)

Similar to (I-B), it can be obtained that

$$\begin{aligned}
& \text{tr} \left(\sum_{j,p,t,s=1}^{2m} c(v) c(e_j) c(w) c(e_p) c(e_s) c(e_t) \right) \quad (3.30) \\
& = \sum_{j,p,t,s=1}^{2m} \left[-v_t w_p \delta_j^s - v_t w_j \delta_p^s + v_s w_p \delta_j^t + v_s w_j \delta_p^t - v_p w_s \delta_j^t + v_p w_t \delta_j^s \right]
\end{aligned}$$

$$- v_j w_t \delta_p^s + v_j w_s \delta_p^t + \delta_j^t \delta_p^s g(v, w) - \delta_j^s \delta_p^t g(v, w) \Big] \text{tr}[id],$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{\|\xi\|^2}{8} \sum_{j,p,s,t=1}^{2m} R_{jpsc} c(v) c(dx_j) c(w) c(e_p) c(e_s) c(e_t) \right\} (x_0) \sigma(\xi) \\ &= \left(\frac{1}{4} s g(v, w) - \frac{1}{2} \text{Ric}(v, w) \right) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.31)$$

(II-1-C)

Similarly, we have

$$\begin{aligned} & \text{tr} \left(\sum_{j,f \neq \alpha \neq \beta=1}^{2m} c(v) c(e_j) c(w) c(e_f) c(e_\alpha) c(e_\beta) \right) \\ &= \sum_{j,f \neq \alpha \neq \beta=1}^{2m} \left[-v_\beta w_f \delta_j^\alpha + v_\beta w_\alpha \delta_j^f + v_\alpha w_f \delta_j^\beta - v_\alpha w_\beta \delta_j^f - v_f w_\alpha \delta_j^\beta + v_f w_\beta \delta_j^\alpha \right] \text{tr}[id], \end{aligned} \quad (3.32)$$

and

$$\frac{\partial T(e_f, e_\alpha, e_\beta)}{\partial x_j}(x_0) = \nabla_{e_j}(T)(e_f, e_\alpha, e_\beta)(x_0), \quad (3.33)$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{\|\xi\|^2}{4} \sum_{f < \alpha < \beta, j=1}^{2m} \frac{\partial T(e_f, e_\alpha, e_\beta)}{\partial x_j} c(v) c(dx_j) c(e_w) c(e_f) c(e_\alpha) c(e_\beta) \right\} (x_0) \sigma(\xi) \\ &= \int_{\|\xi\|=1} \left\{ \frac{\|\xi\|^2}{24} \sum_{f \neq \alpha \neq \beta, j=1}^{2m} \frac{\partial T(e_f, e_\alpha, e_\beta)}{\partial x_j} \text{tr}[c(v) c(dx_j) c(e_w) c(e_f) c(e_\alpha) c(e_\beta)] \right\} (x_0) \sigma(\xi) \\ &= -\frac{1}{4} \sum_{j=1}^{2m} \nabla_{e_j}(T)(v, w, e_j) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.34)$$

(II-1-D)

Similarly, we have

$$\begin{aligned} & \text{tr} \left(\sum_{j,\gamma,f \neq \alpha \neq \beta=1}^{2m} c(v) c(e_j) c(e_\gamma) c(e_f) c(e_\alpha) c(e_\beta) \right) \\ &= \sum_{j,\gamma,f \neq \alpha \neq \beta=1}^{2m} \left[-v_\beta \delta_\gamma^f \delta_j^\alpha + v_\beta \delta_\gamma^\alpha \delta_j^f + v_\alpha \delta_\gamma^f \delta_j^\beta - v_\alpha \delta_\gamma^\beta \delta_j^f - v_f \delta_\gamma^\alpha \delta_j^\beta + v_f \delta_\gamma^\beta \delta_j^\alpha \right] \text{tr}[id], \end{aligned} \quad (3.35)$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \left\{ \frac{\|\xi\|^2}{4} \sum_{f < \alpha < \beta, j, \gamma=1}^{2m} T(e_f, e_\alpha, e_\beta) \partial x_j(w_\gamma) c(v) c(dx_j) c(e_\gamma) c(e_f) c(e_\alpha) c(e_\beta) \right\} (x_0) \sigma(\xi) \\ &= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{\|\xi\|^2}{24} \sum_{f \neq \alpha \neq \beta, j, \gamma=1}^{2m} T(e_f, e_\alpha, e_\beta) \partial x_j(w_\gamma) \text{tr}[c(v) c(dx_j) c(e_\gamma) c(e_f) c(e_\alpha) c(e_\beta)] \right\} (x_0) \sigma(\xi) \end{aligned} \quad (3.36)$$

$$= -\frac{1}{4} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \text{tr}[id] \text{Vol}(S^{n-1}).$$

Summing sequentially from **(II-1-A)** to **(II-1-D)** we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left[\sigma_0(\mathcal{AB}) \sigma_{-2m} (D^{-2m}) (x_0) \right] \sigma(\xi) \\ &= \left[\frac{1}{4} sg(v, w) - \frac{1}{2} \text{Ric}(v, w) \right] \text{tr}[id] \text{Vol}(S^{n-1}) \\ &+ \sum_{f \neq \alpha \neq \beta=1}^{2m} \left[\frac{1}{96} T^2(e_f, e_\alpha, e_\beta) g(v, w) - \frac{1}{16} T(v, e_\alpha, e_\beta) T(w, e_\alpha, e_\beta) \right] \text{tr}[id] \text{Vol}(S^{n-1}) \\ &- \frac{1}{4} \sum_{j=1}^{2m} \left[\nabla_{e_j}(T)(v, w, e_j) + T(v, \nabla_{e_j}^L w, e_j) \right] \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \quad (3.37)$$

(II-2) For $\sigma_1(\mathcal{AB}) \sigma_{-2m-1} (D^{-2m})$:

According to (2.13) in lemma 2.2 and (3.25) in lemma 3.6 , where $w_{s,t}(e_p)(x_0) = 0$, we get

$$\begin{aligned} & \sigma_1(\mathcal{AB}) \sigma_{-2m-1} (D^{-2m}) (x_0) \\ &= -\frac{3m}{4} \|\xi\|^{-2m-2} \sum_{f < \alpha < \beta, k < l=1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_a, e_k, e_l) \\ &\quad \times \left[c(v)c(\xi)c(w)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) + c(w)c(\xi)c(v)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) \right] \xi_a \\ &\quad - 3m \|\xi\|^{-2m-2} \sum_{j, \gamma, a, k < l=1}^{2m} \partial x_j(w_\gamma) T(e_a, e_k, e_l) c(v)c(dx_j)c(e_\gamma)c(\xi)c(e_k)c(e_l)\xi_a. \end{aligned} \quad (3.38)$$

(II-2-A)

Based on the relation of the Clifford action and $\text{tr} \mathcal{X} \mathcal{Y} = \text{tr} \mathcal{Y} \mathcal{X}$, we can obtain the equality

$$\begin{aligned} & \text{tr} \left(\sum_{f \neq \alpha \neq \beta, k \neq l, \eta=1}^{2m} c(v)c(e_\eta)c(w)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) + c(w)c(e_\eta)c(v)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) \right) \\ &= \sum_{f \neq \alpha \neq \beta, k \neq l, \eta=1}^{2m} \left[-2g(e_\eta, w) \text{tr}[c(v)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l)] \right. \\ &\quad \left. + 2g(v, w) \text{tr}[c(e_\eta)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l)] - 2g(v, e_\eta) \text{tr}[c(w)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l)] \right] \\ &= 2 \sum_{f \neq \alpha \neq \beta, k \neq l, \eta=1}^{2m} \left\{ w_\eta v_\beta \delta_f^l \delta_\alpha^k - w_\eta v_\beta \delta_f^k \delta_\alpha^l - w_\eta v_\alpha \delta_f^l \delta_\beta^k + w_\eta v_\alpha \delta_f^k \delta_\beta^l + w_\eta v_f \delta_\alpha^l \delta_\beta^k - w_\eta v_f \delta_\alpha^k \delta_\beta^l \right. \\ &\quad \left. + v_\eta w_\beta \delta_f^l \delta_\alpha^k - v_\eta w_\beta \delta_f^k \delta_\alpha^l - v_\eta w_\alpha \delta_f^l \delta_\beta^k + v_\eta w_\alpha \delta_f^k \delta_\beta^l + v_\eta w_f \delta_\alpha^l \delta_\beta^k - v_\eta w_f \delta_\alpha^k \delta_\beta^l \right. \\ &\quad \left. + \left[-\delta_\eta^\beta \delta_f^l \delta_\alpha^k + \delta_\eta^\beta \delta_f^k \delta_\alpha^l + \delta_\eta^\alpha \delta_f^l \delta_\beta^k - \delta_\eta^\alpha \delta_f^k \delta_\beta^l - \delta_\eta^f \delta_\alpha^l \delta_\beta^k + \delta_\eta^f \delta_\alpha^k \delta_\beta^l \right] g(v, w) \right\} \text{tr}[id], \end{aligned} \quad (3.39)$$

then

$$\int_{\|\xi\|=1} \text{tr} \left\{ -\frac{3m}{4} \|\xi\|^{-2m-2} \sum_{f < \alpha < \beta, k < l=1}^{2m} T(e_f, e_\alpha, e_\beta) T(e_a, e_k, e_l) \right\} \text{tr}[id] \text{Vol}(S^{n-1}) \quad (3.40)$$

$$\begin{aligned}
& \times c(v)c(\xi)c(w)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) + c(w)c(\xi)c(v)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l)\xi_a \Big\} (x_0)\sigma(\xi) \\
& = \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m}{16}\|\xi\|^{-2m-2} \sum_{f \neq \alpha \neq \beta, k \neq l, \eta=1}^{2m} T(e_f, e_\alpha, e_\beta)T(e_a, e_k, e_l) \right. \\
& \quad \left. \times c(v)c(e_\eta)c(w)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l) + c(w)c(e_\eta)c(v)c(e_f)c(e_\alpha)c(e_\beta)c(e_k)c(e_l)\xi_a\xi_\eta \right\} (x_0)\sigma(\xi) \\
& = \sum_{f \neq \alpha \neq \beta=1}^{2m} \left[-\frac{3}{8}T^2(e_f, e_\alpha, e_\beta)g(v, w) + \frac{3}{4}T(v, e_\alpha, e_\beta)T(w, e_\alpha, e_\beta) \right] \text{tr}[id]\text{Vol}(S^{n-1}).
\end{aligned}$$

(II-2-B)

Similarly to (I-B), it follows that

$$\begin{aligned}
& \text{tr} \left(\sum_{j, \gamma, k \neq l=1}^{2m} c(v)c(dx_j)c(e_\gamma)c(e_\eta)c(e_k)c(e_l) \right) \\
& = \sum_{j, \gamma, k \neq l=1}^{2m} \left[-v_l\delta_j^k\delta_\gamma^\eta + v_l\delta_j^\eta\delta_\gamma^k + v_k\delta_j^l\delta_\gamma^\eta - v_k\delta_j^\eta\delta_\gamma^l - v_\eta\delta_j^l\delta_\gamma^k + v_\eta\delta_j^k\delta_\gamma^l \right] \text{tr}[id],
\end{aligned} \tag{3.41}$$

then

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ -3m\|\xi\|^{-2m-2} \sum_{j, \gamma, a, k < l=1}^{2m} \partial x_j(w_\gamma)T(e_a, e_k, e_l)(v)c(dx_j)c(e_\gamma)c(e_\xi)c(e_k)c(e_l)\xi_a \right\} (x_0)\sigma(\xi) \\
& = \int_{\|\xi\|=1} \text{tr} \left\{ \frac{-3m}{2}\|\xi\|^{-2m-2} \sum_{\eta, j, \gamma, a, k \neq l=1}^{2m} \partial x_j(w_\gamma)T(e_a, e_k, e_l)c(v)c(dx_j)c(e_\gamma)c(e_\eta)c(e_k)c(e_l)\xi_a\xi_\eta \right\} (x_0)\sigma(\xi) \\
& = \frac{9}{2} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \text{tr}[id]\text{Vol}(S^{n-1}).
\end{aligned} \tag{3.42}$$

By summing (II-2-A) and (II-2-B), we get

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left[\sigma_1(\mathcal{A}\mathcal{B})\sigma_{-2m-1}(D^{-2m})(x_0) \right] \sigma(\xi) \\
& = \sum_{f \neq \alpha \neq \beta=1}^{2m} \left[-\frac{3}{8}T^2(e_f, e_\alpha, e_\beta)g(v, w) + \frac{3}{4}T(v, e_\alpha, e_\beta)T(w, e_\alpha, e_\beta) \right] \text{tr}[id]\text{Vol}(S^{n-1}) \\
& \quad + \frac{9}{2} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \text{tr}[id]\text{Vol}(S^{n-1}).
\end{aligned} \tag{3.43}$$

(II-3) For $\sigma_2(\mathcal{A}\mathcal{B})\sigma_{-2m-2}(D^{-2m})$:

According to (2.14) in lemma 2.2 and (3.26) in lemma 3.6, we get

$$\begin{aligned}
& \sigma_2(\mathcal{A}\mathcal{B})\sigma_{-2m-2}(D^{-2m})(x_0) \\
& = -\frac{m(m+1)}{3}\|\xi\|^{-2m-4} \sum_{a, b=1}^{2m} \text{Ric}_{ab}\xi_a\xi_b c(v)c(\xi)c(w)c(\xi) \\
& \quad + \frac{9m(m+1)}{2}\|\xi\|^{-2m-4} \sum_{a, b, j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l)T(e_b, e_{\hat{j}}, e_{\hat{l}})c(v)c(\xi)c(w)c(\xi)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})\xi_a\xi_b
\end{aligned} \tag{3.44}$$

$$\begin{aligned}
& -\frac{9m}{4}\|\xi\|^{-2m-2} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) c(v) c(\xi) c(w) c(\xi) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \\
& -\frac{3m}{2}\|\xi\|^{-2m-2} \sum_{j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a} c(v) c(\xi) c(w) c(\xi) c(e_j) c(e_l) \\
& + \frac{m(m+1)}{4}\|\xi\|^{-2m-4} \sum_{a,b,t,s=1}^{2m} R_{bats} c(v) c(\xi) c(w) c(\xi) c(e_s) c(e_t) \xi_a \xi_b \\
& + 3m(m+1)\|\xi\|^{-2m-4} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b} c(v) c(\xi) c(w) c(\xi) c(e_j) c(e_l) \xi_a \xi_b \\
& + m\|\xi\|^{-2m-2} (\frac{1}{4}R^g - \frac{3}{4}\|T\|^2) c(v) c(\xi) c(w) c(\xi) \\
& + \frac{3m}{2}\|\xi\|^{-2m-2} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t) c(v) c(\xi) c(w) c(\xi) c(e_i) c(e_j) c(e_k) c(e_t).
\end{aligned}$$

(II-3-A)

Based on the relation of the Clifford action and $\text{tr} \mathcal{X} \mathcal{Y} = \text{tr} \mathcal{Y} \mathcal{X}$, we get

$$\text{tr} \left(\sum_{f,g=1}^{2m} c(v) c(e_f) c(w) c(e_g) \right) = \sum_{f,g=1}^{2m} \left[v_g w_f - \delta_f^g g(v, w) + v_g w_f \right] \text{tr}[id], \quad (3.45)$$

then

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m(m+1)}{3} \|\xi\|^{-2m-4} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b c(v) c(\xi) c(w) c(\xi) \right\} (x_0) \sigma(\xi) \\
& = \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m(m+1)}{3} \|\xi\|^{-2m-4} \sum_{a,b,f,g=1}^{2m} \text{Ric}_{ab} \xi_a \xi_b \xi_f \xi_g c(v) c(e_f) c(w) c(e_g) \right\} (x_0) \sigma(\xi) \\
& = \left(\frac{m}{6} s g(v, w) - \frac{1}{3} \text{Ric}(v, w) \right) \text{tr}[id] \text{Vol}(S^{n-1}).
\end{aligned}$$

(II-3-B)

Based on the relation of the Clifford action and $\text{tr} \mathcal{X} \mathcal{Y} = \text{tr} \mathcal{Y} \mathcal{X}$, we get

$$\begin{aligned}
& \text{tr} \left(\sum_{f,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} c(v) c(e_f) c(w) c(e_f) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \quad (3.46) \\
& = \sum_{f,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} \left[2m \text{tr} \left(c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) - 2w_f \text{tr} \left(c(v) c(e_f) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \right],
\end{aligned}$$

and

$$\begin{aligned}
& \text{tr} \sum_{a,b,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} \left(c(v) c(e_a) c(w) c(e_b) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) + c(v) c(e_b) c(w) c(e_a) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \quad (3.47) \\
& = \sum_{a,b,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} \left[-2g(e_a, w) \text{tr} \left(c(v) c(e_b) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) + 2g(e_a, e_b) \text{tr} \left(c(v) c(w) c(e_j) c(e_l) c(e_{\hat{j}}) c(e_{\hat{l}}) \right) \right]
\end{aligned}$$

$$- 2g(w, e_b) \text{tr} \left(c(v)c(e_a)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}}) \right) \Big].$$

By (3.46), (3.47) and $\int_{\|\xi\|=1} \xi_a \xi_b \xi_f \xi_g \sigma(\xi) = \frac{1}{n(n+2)} (\delta_a^b \delta_f^g + \delta_a^f \delta_b^g + \delta_a^g \delta_b^f) \text{Vol}(S^{n-1})$, we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{9m(m+1)}{2} \|\xi\|^{-2m-4} \sum_{a,b,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_b, e_{\hat{j}}, e_{\hat{l}}) c(v)c(\xi)c(w)c(\xi)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})\xi_a \xi_b \right\} (x_0) \sigma(\xi) \\ &= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{9m(m+1)}{8} \sum_{a,b,f,g,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_b, e_{\hat{j}}, e_{\hat{l}}) c(v)c(e_f)c(w)c(e_g)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})\xi_a \xi_b \xi_f \xi_g \right\} (x_0) \sigma(\xi) \\ &= \frac{9}{32} \sum_{a,f,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) \text{tr}[c(v)c(e_f)c(w)c(e_g)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})] \text{Vol}(S^{n-1}) \\ &\quad + \frac{9}{32} \sum_{a,b,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_b, e_{\hat{j}}, e_{\hat{l}}) \text{tr}[c(v)c(e_a)c(w)c(e_b)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})] \text{Vol}(S^{n-1}) \\ &\quad + \frac{9}{32} \sum_{a,b,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_b, e_{\hat{j}}, e_{\hat{l}}) \text{tr}[c(v)c(e_b)c(w)c(e_a)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})] \text{Vol}(S^{n-1}) \\ &= \sum_{a,j \neq l=1}^{2m} \left[\frac{9}{8} (2-m) T^2(e_a, e_j, e_l) g(v, w) - \frac{9}{4} T(v, e_j, e_l) T(w, e_j, e_l) \right] \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \tag{3.48}$$

(II-3-C)

By (3.46) and $\int_{\|\xi\|=1} \xi_f \xi_g \sigma(\xi) = \frac{1}{n} \delta_f^g \text{Vol}(S^{n-1})$, we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{9m}{4} \|\xi\|^{-2m-2} \sum_{a,j < l, \hat{j} < \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) c(v)c(\xi)c(w)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}}) \right\} (x_0) \sigma(\xi) \\ &= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{9m}{8} \|\xi\|^{-2m-2} \sum_{a,f,g,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) c(v)c(e_f)c(w)c(e_g)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})\xi_f \xi_g \right\} (x_0) \sigma(\xi) \\ &= \frac{9}{16} \sum_{a,f,g,j \neq l, \hat{j} \neq \hat{l}=1}^{2m} T(e_a, e_j, e_l) T(e_a, e_{\hat{j}}, e_{\hat{l}}) \text{tr}[c(v)c(e_f)c(w)c(e_g)c(e_j)c(e_l)c(e_{\hat{j}})c(e_{\hat{l}})] \text{Vol}(S^{n-1}) \\ &= \sum_{a,j \neq l=1}^{2m} \frac{9}{8} (m-1) T^2(e_a, e_j, e_l) g(v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \tag{3.49}$$

(II-3-D)

Similarly to (I-B), we get

$$\begin{aligned} & \text{tr} \left(\sum_{f,j \neq l=1}^{2m} c(v)c(e_f)c(w)c(e_f)c(e_j)c(e_l) \right) \\ &= \sum_{f,j \neq l=1}^{2m} \left[2mv_l w_j - 2mv_j w_l - 2w_f v_l \delta_f^j + 2w_f v_j \delta_f^l \right] \text{tr}[id], \end{aligned} \tag{3.50}$$

then

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{3m}{2} \|\xi\|^{-2m-2} \sum_{j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a} c(v) c(\xi) c(w) c(\xi) c(e_j) c(e_l) \right\} (x_0) \sigma(\xi) \\
&= \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{3m}{4} \|\xi\|^{-2m-2} \sum_{j \neq l, f, g=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a} c(v) c(e_f) c(w) c(e_g) c(e_j) c(e_l) \xi_f \xi_g \right\} (x_0) \sigma(\xi) \\
&= -\frac{3}{8} \|\xi\|^{-2m-2} \sum_{j \neq l, f, g=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_a} \text{tr} \left(c(v) c(e_f) c(w) c(e_f) c(e_j) c(e_l) \right) \text{Vol}(S^{n-1}) \\
&= \frac{3}{2} (m-1) \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) \text{tr}[id] \text{Vol}(S^{n-1}).
\end{aligned} \tag{3.51}$$

(II-3-E)

Due to $\int_{\|\xi\|=1} \xi_a \xi_b \xi_f \xi_g \sigma(\xi) = \frac{1}{n(n+2)} (\delta_a^b \delta_f^g + \delta_a^f \delta_b^g + \delta_a^g \delta_b^f) \text{Vol}(S^{n-1})$ and $R_{aats} = 0$, we get

$$\begin{aligned}
& \int_{\|\xi\|=1} \left(\|\xi\|^{-2m-4} \sum_{a,b,f,g,t,s=1}^{2m} R_{bats} \xi_a \xi_b \xi_f \xi_g \right) \sigma(\xi) \\
&= \frac{1}{4m(m+1)} \sum_{a,b,f,g,t,s=1}^{2m} R_{bats} (\delta_a^f \delta_b^g + \delta_a^g \delta_b^f) \text{Vol}(S^{n-1}),
\end{aligned} \tag{3.52}$$

then

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ \frac{m(m+1)}{4} \|\xi\|^{-2m-4} \sum_{a,b,t,s=1}^{2m} R_{bats} c(v) c(\xi) c(w) c(\xi) c(e_s) c(e_t) \xi_a \xi_b \right\} (x_0) \sigma(\xi) \\
&= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{m(m+1)}{4} \|\xi\|^{-2m-4} \sum_{a,b,f,g,t,s=1}^{2m} R_{bats} c(v) c(e_f) c(w) c(e_g) c(e_s) c(e_t) \xi_a \xi_b \xi_f \xi_g \right\} (x_0) \sigma(\xi) \\
&= \frac{1}{16} \sum_{a,b,t,s=1}^{2m} R_{bats} \text{tr} \left(c(v) c(e_a) c(w) c(e_b) c(e_s) c(e_t) + c(v) c(e_a) c(w) c(e_b) c(e_s) c(e_t) \right) \text{Vol}(S^{n-1}) \\
&= \frac{1}{8} \sum_{a,b,t,s=1}^{2m} R_{bats} \left(-w_a v_t \delta_b^s + w_a v_s \delta_b^t + w_b v_t \delta_a^s - w_b v_s \delta_a^t \right) \text{Vol}(S^{n-1}) \\
&= 0.
\end{aligned} \tag{3.53}$$

(II-3-F)

By (3.30) and (3.33), we obtain

$$\begin{aligned}
& \int_{\|\xi\|=1} \text{tr} \left\{ 3m(m+1) \|\xi\|^{-2m-4} \sum_{a,b,j < l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b} c(v) c(\xi) c(w) c(\xi) c(e_j) c(e_l) \xi_a \xi_b \right\} (x_0) \sigma(\xi) \\
&= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{3m(m+1)}{2} \|\xi\|^{-2m-4} \sum_{a,b,f,g,j \neq l=1}^{2m} \frac{\partial T(\partial_a, e_j, e_l)}{\partial x_b} c(v) c(e_f) c(w) c(e_g) c(e_j) c(e_l) \xi_a \xi_b \xi_f \xi_g \right\} (x_0) \sigma(\xi) \\
&= \frac{3}{8} \sum_{a,f,j \neq l=1}^{2m} \nabla_{e_a}(T)(e_a, e_j, e_l) \text{tr} \left(c(v) c(e_f) c(w) c(e_f) c(e_j) c(e_l) \right) \text{Vol}(S^{n-1}) \\
&\quad + \frac{3}{8} \sum_{a,b,j \neq l=1}^{2m} \nabla_{e_b}(T)(e_a, e_j, e_l) \text{tr} \left(c(v) c(e_a) c(w) c(e_b) c(e_j) c(e_l) + c(v) c(e_b) c(w) c(e_a) c(e_j) c(e_l) \right) \text{Vol}(S^{n-1})
\end{aligned} \tag{3.54}$$

$$= -\frac{3}{2}(m+1)\nabla_{e_a}(T)(e_a, v, w)\text{tr}[id]\text{Vol}(S^{n-1}).$$

(II-3-G)

By (3.45), we obtain

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ m\|\xi\|^{-2m-2} \left(\frac{1}{4}\text{R}^g - \frac{3}{4}\|T\|^2 \right) c(v)c(\xi)c(w)c(\xi) \right\} (x_0)\sigma(\xi) \\ &= \int_{\|\xi\|=1} \text{tr} \left\{ m\|\xi\|^{-2m-2} \left(\frac{1}{4}\text{R}^g - \frac{3}{4}\|T\|^2 \right) c(v)c(e_f)c(w)c(e_g)\xi_f\xi_g \right\} (x_0)\sigma(\xi) \\ &= \frac{1}{2} \left(\frac{1}{4}\text{R}^g - \frac{3}{4}\|T\|^2 \right) [2g(v, w) - 2mg(v, w)]\text{tr}[id]\text{Vol}(S^{n-1}) \\ &= (1-m) \left(\frac{1}{4}s - \frac{3}{4}\|T\|^2 \right) g(v, w)\text{tr}[id]\text{Vol}(S^{n-1}). \end{aligned} \quad (3.55)$$

(II-3-H)

$$\sum_{i < j < k < t=1}^{2m} \text{tr} \left(c(v)c(e_f)c(w)c(e_f)c(e_i)c(e_j)c(e_k)c(e_t) \right) = 0, \quad (3.56)$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ \frac{3m}{2}\|\xi\|^{-2m-2} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t)c(v)c(\xi)c(w)c(\xi)c(e_i)c(e_j)c(e_k)c(e_t) \right\} (x_0)\sigma(\xi) \\ &= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{3m}{2}\|\xi\|^{-2m-2} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t)c(v)c(e_f)c(w)c(e_g)c(e_i)c(e_j)c(e_k)c(e_t)\xi_f\xi_g \right\} (x_0)\sigma(\xi) \\ &= \frac{3}{4} \sum_{i < j < k < t=1}^{2m} dT(e_i, e_j, e_k, e_t) \text{tr} \left(c(v)c(e_f)c(w)c(e_f)c(e_i)c(e_j)c(e_k)c(e_t) \right) \text{Vol}(S^{n-1}) \\ &= 0. \end{aligned} \quad (3.57)$$

Summing from (II-3-A) to (II-3-H), we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left[\sigma_2(\mathcal{AB})\sigma_{-2m-2}(D^{-2m})(x_0) \right] \sigma(\xi) \\ &= \left[\frac{m}{6}sg(v, w) - \frac{1}{3}\text{Ric}(v, w) \right] \text{tr}[id]\text{Vol}(S^{n-1}) \\ &+ \sum_{a \neq j \neq l=1}^{2m} \left[\frac{9}{8}T^2(e_a, e_j, e_l)g(v, w) - \frac{9}{4}T(v, e_j, e_l)T(w, e_j, e_l) \right] \text{tr}[id]\text{Vol}(S^{n-1}) \\ &- 3 \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w)\text{tr}[id]\text{Vol}(S^{n-1}) + (1-m) \left(\frac{1}{4}\text{R}^g - \frac{3}{4}\|T\|^2 \right) g(v, w)\text{tr}[id]\text{Vol}(S^{n-1}). \end{aligned} \quad (3.58)$$

(II-4) For $(-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_2(\mathcal{AB})] \partial_{x_j} [\sigma_{-2m-1}(D_T^{-2m})]$:

According to (2.13) in lemma 2.2 and (3.26) in lemma 3.6, we get

$$(-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_2(\mathcal{AB})] \partial_{x_j} [\sigma_{-2m-1}(D_T^{-2m})](x_0) \quad (3.59)$$

$$\begin{aligned}
&= \frac{2m}{3} \|\xi\|^{-2m-2} \sum_{a,b,j=1}^{2m} \text{Ric}_{ab} \xi_a \delta_j^b \left(c(v)c(dx_j) c(w)c(\xi) + c(v)c(\xi)c(w)c(dx_j) \right) \\
&\quad - \frac{m}{4} \|\xi\|^{-2m-2} \sum_{a,b,j,t,s=1}^{2m} \text{R}_{bats} \xi_a \delta_j^b \left(c(v)c(dx_j) c(w)c(\xi)c(e_s)c(e_t) + c(v)c(\xi)c(w)c(dx_j)c(e_s)c(e_t) \right) \\
&\quad - 3m \|\xi\|^{-2m-2} \sum_{a,b,j,\hat{j}<\hat{l}=1}^{2m} \frac{\partial T(\partial_a, e_{\hat{j}}, e_{\hat{l}})}{\partial x_b} \xi_a \delta_j^b \left(c(v)c(dx_j) c(w)c(\xi)c(e_{\hat{j}})c(e_{\hat{l}}) + c(v)c(\xi)c(w)c(dx_j)c(e_{\hat{j}})c(e_{\hat{l}}) \right).
\end{aligned}$$

(II-4- \mathbb{A})

Based on the relation of the Clifford action and $\text{tr} \mathcal{X} \mathcal{Y} = \text{tr} \mathcal{Y} \mathcal{X}$, we get

$$\sum_{a,b=1}^{2m} \text{tr} \left(c(v)c(e_b)c(w)c(e_a) + c(v)c(e_a)c(w)c(e_b) \right) = \sum_{a,b=1}^{2m} \left(2v_a w_b - 2\delta_a^b g(v,w) + 2v_b w_a \right) \text{tr}[id], \quad (3.60)$$

then

$$\begin{aligned}
&\int_{\|\xi\|=1} \text{tr} \left\{ \frac{2m}{3} \|\xi\|^{-2m-2} \sum_{a,b,j=1}^{2m} \text{Ric}_{ab} \xi_a \delta_j^b c(v)c(dx_j) c(w)c(\xi) + c(v)c(\xi)c(w)c(dx_j) \right\} (x_0) \sigma(\xi) \quad (3.61) \\
&= \int_{\|\xi\|=1} \text{tr} \left\{ \frac{2m}{3} \|\xi\|^{-2m-2} \sum_{a,b,f=1}^{2m} \text{Ric}_{ab} \xi_a \xi_f c(v)c(e_b)c(w)c(e_f) + c(v)c(e_f)c(w)c(e_b) \right\} (x_0) \sigma(\xi) \\
&= \frac{1}{3} \sum_{a,b=1}^{2m} \text{Ric}_{ab} \text{tr} \left(c(v)c(e_b)c(w)c(e_a) + c(v)c(e_a)c(w)c(e_b) \right) (x_0) \text{Vol}(S^{n-1}) \\
&= \frac{2}{3} \left(2 \text{Ric}(v,w) - sg(v,w) \right) \text{tr}[id] \text{Vol}(S^{n-1}).
\end{aligned}$$

(II-4- \mathbb{B})

For the same reason, we get

$$\sum_{a,b,s,t=1}^{2m} \text{R}_{bats} \text{tr} \left(c(v)c(e_b)c(w)c(e_a)c(e_s)c(e_t) + c(v)c(e_a)c(w)c(e_b)c(e_s)c(e_t) \right) = 0, \quad (3.62)$$

then

$$\begin{aligned}
&\int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m}{4} \|\xi\|^{-2m-2} \sum_{a,b,j,t,s=1}^{2m} \text{R}_{bats} \xi_a \delta_j^b \left(c(v)c(dx_j) c(w)c(\xi)c(e_s)c(e_t) + c(v)c(\xi)c(w)c(dx_j)c(e_s)c(e_t) \right) \right\} (x_0) \sigma(\xi) \quad (3.63) \\
&= \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{m}{4} \|\xi\|^{-2m-2} \sum_{a,b,t,s,f=1}^{2m} \text{R}_{bats} \xi_a \xi_f \left(c(v)c(e_b)c(w)c(e_f)c(e_s)c(e_t) + c(v)c(e_f)c(w)c(e_b)c(e_s)c(e_t) \right) \right\} (x_0) \sigma(\xi) \\
&= -\frac{1}{8} \sum_{a,b,t,s=1}^{2m} \text{R}_{bats} \text{tr} \left(c(v)c(e_b)c(w)c(e_a)c(e_s)c(e_t) + c(v)c(e_a)c(w)c(e_b)c(e_s)c(e_t) \right) (x_0) \text{Vol}(S^{n-1}) \\
&= 0.
\end{aligned}$$

(II-4- \mathbb{C})

By (3.15), we get

$$\sum_{a,b,\hat{j}\neq\hat{l}=1}^{2m} \nabla_{e_b}(T)(e_a, e_{\hat{j}}, e_{\hat{l}}) \text{tr} \left(c(v)c(e_b)c(w)c(e_a)c(e_{\hat{j}})c(e_{\hat{l}}) + c(v)c(e_a)c(w)c(e_b)c(e_{\hat{j}})c(e_{\hat{l}}) \right) \quad (3.64)$$

$$= -8 \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) \text{tr}[id],$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left\{ -3m \|\xi\|^{-2m-2} \sum_{a,b,j,\hat{j}<\hat{l}=1}^{2m} \frac{\partial T(\partial_a, e_{\hat{j}}, e_{\hat{l}})}{\partial x_b} \xi_a \delta_j^b \times \right. \\ & \quad \left. \times \left(c(v)c(dx_j)c(w)c(\xi)c(e_{\hat{j}})c(e_{\hat{l}}) + c(v)c(\xi)c(w)c(dx_j)c(e_{\hat{j}})c(e_{\hat{l}}) \right) \right\} (x_0) \sigma(\xi) \\ & = \int_{\|\xi\|=1} \text{tr} \left\{ -\frac{3m}{2} \sum_{a,b,f,\hat{j}\neq\hat{l}=1}^{2m} \frac{\partial T(\partial_a, e_{\hat{j}}, e_{\hat{l}})}{\partial x_b} \xi_a \xi_f \left(c(v)c(e_b)c(w)c(e_{\hat{j}})c(e_{\hat{l}}) + c(v)c(e_f)c(w)c(e_b)c(e_{\hat{j}})c(e_{\hat{l}}) \right) \right\} (x_0) \sigma(\xi) \\ & = -\frac{3}{4} \sum_{a,b,\hat{j}\neq\hat{l}=1}^{2m} \nabla_{e_b}(T)(e_a, e_{\hat{j}}, e_{\hat{l}}) \text{tr} \left(c(v)c(e_b)c(w)c(e_a)c(e_{\hat{j}})c(e_{\hat{l}}) + c(v)c(e_a)c(w)c(e_b)c(e_{\hat{j}})c(e_{\hat{l}}) \right) (x_0) \text{Vol}(S^{n-1}) \\ & = 6 \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \tag{3.65}$$

Summing from **(II-4-A)** to **(II-4-C)**, we get

$$\begin{aligned} & \int_{\|\xi\|=1} \text{tr} \left[(-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_2(\mathcal{AB})] \partial_{x_j} [\sigma_{-2m-1} (D_T^{-2m})] \right] (x_0) \sigma(\xi) \\ & = \frac{2}{3} \left(2 \text{Ric}(v, w) - sg(v, w) \right) \text{tr}[id] \text{Vol}(S^{n-1}) + 6 \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) \text{tr}[id] \text{Vol}(S^{n-1}). \end{aligned} \tag{3.66}$$

(II-5) For $(-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_1(\mathcal{AB})] \partial_{x_j} [\sigma_{-2m} (D_T^{-2m})]$:
According to (2.12) in lemma 2.2, we get

$$\begin{aligned} & \partial_{x_j} [\sigma_{-2m} (D_T^{-2m})] (x_0) \\ & = \partial_{x_j} \left[\|\xi\|^{-2m-2} \sum_{a,b,l,k=1}^{2m} \left(\delta_{ab} - \frac{m}{3} R_{albk} x^l x^k \right) \xi_a \xi_b \right] (x_0) \\ & = 0, \end{aligned} \tag{3.67}$$

then

$$\int_{\|\xi\|=1} \text{tr} \left[(-i) \sum_{j=1}^{2m} \partial_{\xi_j} [\sigma_1(\mathcal{AB})] \partial_{x_j} [\sigma_{-2m} (D_T^{-2m})] \right] (x_0) \sigma(\xi) = 0. \tag{3.68}$$

(II-6) For $-\frac{1}{2} \sum_{j,l=1}^{2m} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(\mathcal{AB})] \partial_{x_j} \partial_{x_l} [\sigma_{-2m} (D_T^{-2m})]$:
According to (2.12) in lemma 2.2 and (3.26) in lemma 3.6, we get

$$\begin{aligned} & -\frac{1}{2} \sum_{j,l=1}^{2m} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(\mathcal{AB})] \partial_{x_j} \partial_{x_l} [\sigma_{-2m} (D_T^{-2m})] (x_0) \\ & = -\frac{m}{6} \|\xi\|^{-2m-2} \sum_{a,b,j,l=1}^{2m} \left(R_{albj} + R_{ajbl} \right) \left[c(v)c(dx_l)c(w)c(dx_j) + c(v)c(dx_j)c(w)c(dx_l) \right] \xi_a \xi_b. \end{aligned} \tag{3.69}$$

Since

$$\begin{aligned} & \operatorname{tr} \sum_{a,j,l=1}^{2m} R_{alaj} \left[c(v)c(dx_l)c(w)c(dx_j) + c(v)c(dx_j)c(w)c(dx_l) \right] \\ &= \left(4\operatorname{Ric}(v,w) - 2sg(v,w) \right) \operatorname{tr}[id], \end{aligned}$$

then

$$\begin{aligned} & \int_{\|\xi\|=1} \operatorname{tr} \left\{ -\frac{1}{2} \sum_{j,l=1}^{2m} \partial_{\xi_j} \partial_{\xi_l} [\sigma_2(\mathcal{AB})] \partial_{x_j} \partial_{x_l} [\sigma_{-2m} (D_T^{-2m})] \right\} (x_0) \sigma(\xi) \\ &= \int_{\|\xi\|=1} \left\{ -\frac{m}{6} \|\xi\|^{-2m-2} \sum_{a,b,j,l=1}^{2m} \left(R_{albj} + R_{ajbl} \right) \operatorname{tr} \left[c(v)c(dx_l)c(w)c(dx_j) + c(v)c(dx_j)c(w)c(dx_l) \right] \xi_a \xi_b \right\} (x_0) \sigma(\xi) \\ &= -\frac{1}{6} \sum_{a,j,l=1}^{2m} R_{alaj} \operatorname{tr} \left[c(v)c(dx_l)c(w)c(dx_j) + c(v)c(dx_j)c(w)c(dx_l) \right] (x_0) \operatorname{Vol}(S^{n-1}) \\ &= -\frac{1}{3} \left(2\operatorname{Ric}(v,w) - sg(v,w) \right) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}). \end{aligned} \quad (3.70)$$

Summing from **(II-1)** to **(II-6)**, we get

$$\begin{aligned} & \int_{\|\xi\|=1} \operatorname{tr} \left\{ \sigma_{-2m} (\mathcal{AB} D_T^{-2m}) \right\} (x_0) \sigma(\xi) \\ &= -\frac{1}{6} \left(\operatorname{Ric}(v,w) - \frac{1}{2} sg(v,w) \right) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}) - \frac{m-1}{12} sg(v,w) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}) \\ &+ \frac{12m+61}{16} \|T\|^2 g(v,w) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}) - \frac{25}{16} \sum_{j,l=1}^{2m} T(v, e_j, e_l) T(w, e_j, e_l) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}) \\ &+ \frac{11}{4} \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}) + \frac{17}{4} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \operatorname{tr}[id] \operatorname{Vol}(S^{n-1}). \end{aligned} \quad (3.71)$$

Since $\operatorname{tr}[id] = 2^m$ and $\operatorname{Vol}(S^{n-1}) = \frac{2\pi^m}{\Gamma(m)}$, we obtain

$$\begin{aligned} \mathcal{B}_2 &= \operatorname{Wres} \left(c(v) D_T c(w) D_T D_T^{-n} \right) \\ &= 2^m \frac{2\pi^m}{\Gamma(m)} \int_M \left\{ -\frac{1}{6} \left(\operatorname{Ric}(v,w) - \frac{1}{2} sg(v,w) \right) - \frac{m-1}{12} sg(v,w) + \frac{12m+61}{16} \|T\|^2 g(v,w) \right. \\ &\quad \left. - \frac{25}{16} \sum_{j,l=1}^{2m} T(v, e_j, e_l) T(w, e_j, e_l) + \frac{11}{4} \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) + \frac{17}{4} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \right\} d\operatorname{Vol}_M. \end{aligned} \quad (3.72)$$

Hence Theorem 3.3 holds. \square

As an application of Theorem 3.3, we give the following definition.

Definition 3.7. For a closed oriented manifold M , the spectral Einstein functional with torsion on M is defined by

$$\mathcal{B}(v, w) := 2^m \frac{2\pi^m}{\Gamma(m)} \int_M \left\{ -\frac{1}{6} \mathbb{G}(v, w) + \frac{73}{16} \|T\|^2 g(v, w) - \frac{25}{16} \sum_{j,l=1}^{2m} T(v, e_j, e_l) T(w, e_j, e_l) \right\} \quad (3.73)$$

$$+ \frac{11}{4} \sum_{a=1}^{2m} \nabla_{e_a}(T)(e_a, v, w) + \frac{17}{4} \sum_{j=1}^{2m} T(v, \nabla_{e_j}^L w, e_j) \Big\} d\text{Vol}_M,$$

where $g(v, w) = \sum_{a=1}^n v_a w_a$ and $\mathbb{G}(v, w) = \text{Ric}(v, w) - \frac{1}{2}sg(v, w)$, $v = \sum_{a=1}^n v_a e_a$, $w = \sum_{b=1}^n w_b e_b$.

Acknowledgements

This work is sponsored by Natural Science Foundation of Xinjiang Uygur Autonomous Region 2024D01C341 and supported by the National Natural Science Foundation of China 11771070, 12061078. The authors thank the referee for his (or her) careful reading and helpful comments.

Declarations

- **Funding** The Natural Science Foundation of Xinjiang Uygur Autonomous Region 2024D01C341 and the National Natural Science Foundation of China 11771070, 12061078.
- **Conflict of interest** The authors declare no Conflict of interest.
- **Ethics approval and consent to participate** Ethics approval and consent to participate.
- **Consent for publication** All the authors agreed to publish this research.
- **Author contribution** All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by JH and YW. The first draft of the manuscript was written by JH and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

References

- [1] L. Dabrowski., A. Sitarz., P. Zalecki.: Spectral Metric and Einstein Functionals. *Advances in Mathematics*, 427, 109128, (2023).
- [2] A. Connes.: Quantized calculus and applications. XIth International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
- [3] A. Connes.: The action functional in Noncommutative geometry. *Comm. Math. Phys.* 117, 673-683, (1988).
- [4] A. Connes, J. Lott.: Particle models and Non-commutative geometry. *Nucl. Phys. B Proc.Supp* 18B, 29-47, (1990).
- [5] A. Connes., A H. Chamseddine.: Inner fluctuations of the spectral action. *J. Geom. Phys.* 2007, 57(1): 1-21.
- [6] D. Kastler.: The Dirac Operator and Gravitation. *Comm. Math. Phys.* 166, 633-643, (1995).
- [7] W. Kalau and M. Walze.: Gravity, Noncommutative geometry and the Wodzicki residue. *J. Geom. Physics*. 16, 327-344,(1995).
- [8] B. V. Fedosov, F. Gose, E. Leichtnam, E. Schrohe.: The noncommutative residue for manifolds with boundary. *J. Funct. Anal.* 142, 1-31, (1996).
- [9] Y. Wang.: Differential forms and the Wodzicki residue for Manifolds with Boundary. *J. Geom. Physics*. 56, 731-753, (2006).
- [10] Y. Wang.: Gravity and the Noncommutative Residue for Manifolds with Boundary. *Letters in Mathematical Physics*. 80, 37-56, (2007).
- [11] Y. Wang.: Lower-Dimensional Volumes and Kastler-kalau-Walze Type Theorem for Manifolds with Boundary . *Commun. Theor. Phys.* Vol 54, 38-42, (2010).
- [12] J. Wang., Y. Wang., C L. Yang., Dirac operators with torsion and the noncommutative residue for manifolds with boundary, *J. Geom. Phys.*, 2014, 81:92-111 .
- [13] J. Wang, Y. Wang, T. Wu, Dirac operators with torsion, spectral Einstein functionals and the noncommutative residue. *J. Math. Phys.* 64, 102505(2023).
- [14] A. Connes.: Noncommutative Geometry. Academic Press, San Diego. 1994.
- [15] H. Figueroa., J. M. Gracia-Bondía., F. Lizzi., J. C. Várilly.: A nonperturbative form of the spectral action principle in noncommutative geometry. *J. Geom. Phys.* 1998, 26(3-4): 329-339.
- [16] M. Wodzicki. Noncommutative residue I: Fundamentals, in K-theory, Arithmetic and Geometry, Yu. I. Manin, ed., Lecture Notes in Mathematics Vol. 1289 (Springer, Berlin, 1987).
- [17] T. Ackermann., J. Tolksdorf.: A generalized Lichnerowicz formula, the Wodzicki residue and gravity. *J. Geom. Physics.*,1996, 19:143-150.
- [18] F. Pfäffle., C. A. Stephan.: Chiral Asymmetry and the Spectral Action. *Commun. Math. Phys.* 321, 283-310 (2013).

- [19] F. Pfäffle., C. A. Stephan.: On gravity, torsion and the spectral action principle. *J. Funct. Anal.* 262, 1529–1565,(2012).
- [20] B. Iochum., C. Levy., D. Vassilevich.: Spectral Action for Torsion with and without Boundaries.*Commun. Math. Phys.* 310, 367–382,(2012).
- [21] F. Pfäffle., C. A. Stephan.: The Holst Action by the Spectral Action Principle. *Commun. Math. Phys.* 307, 261–273 (2011).
- [22] F. Hanisch., F. Pfäffle., C. A. Stephan.:The Spectral Action for Dirac Operators with Skew-Symmetric Torsion. *Commun. Math. Phys.* 300, 877–888 (2010).
- [23] L. Dabrowski., P. Zalecki., A. Sitarz.:Spectral Metric and Einstein Functionals for the Hodge-Dirac operator. *J. Noncommut. Geom.* (Online first) DOI 10.4171/JNCG/573(2024).