EQUIVALENCE CLASSES OF LOWER AND UPPER DESCENT WEAK BRUHAT INTERVALS

SEUNG-IL CHOI, SUN-YOUNG NAM, AND YOUNG-TAK OH

ABSTRACT. Let $\operatorname{Int}(n)$ denote the set of nonempty left weak Bruhat intervals in the symmetric group \mathfrak{S}_n . We investigate the equivalence relation $\stackrel{D}{\simeq}$ on $\operatorname{Int}(n)$, where $I \stackrel{D}{\cong} J$ if and only if there exists a descent-preserving poset isomorphism between I and J. For each equivalence class C of $(\operatorname{Int}(n), \stackrel{D}{\simeq})$, a partial order \preceq is defined by $[\sigma, \rho]_L \preceq [\sigma', \rho']_L$ if and only if $\sigma \preceq_R \sigma'$. Kim–Lee–Oh (2023) showed that the poset (C, \preceq) is isomorphic to a right weak Bruhat interval.

In this paper, we focus on lower and upper descent weak Bruhat intervals, specifically those of the form $[w_0(S), \sigma]_L$ or $[\sigma, w_1(S)]_L$, where $w_0(S)$ is the longest element in the parabolic subgroup \mathfrak{S}_S of \mathfrak{S}_n , generated by $\{s_i \mid i \in S\}$ for a subset $S \subseteq [n-1]$, and $w_1(S)$ is the longest element among the minimal-length representatives of left $\mathfrak{S}_{[n-1]\backslash S}$ -cosets in \mathfrak{S}_n . We begin by providing a poset-theoretic characterization of the equivalence relation \mathfrak{D}_{\sim} . Using this characterization, the minimal and maximal elements within an equivalence class C are identified when C is a lower or upper descent interval. Under an additional condition, a detailed description of the structure of (C, \preceq) is provided. Furthermore, for the equivalence class containing $[w_0(S), \sigma]_L$, an injective hull of $\mathsf{B}([w_0(S), \sigma]_L)$ is given, and for the equivalence class containing $[\sigma, w_1(S)]_L$, a projective cover of $\mathsf{B}([\sigma, w_1(S)]_L)$ is given. Here, $\mathsf{B}(I)$ denotes the weak Bruhat interval module of the 0-Hecke algebra associated with $I \in \operatorname{Int}(n)$. The results obtained are applied to investigate lower descent intervals arising from quotient modules and submodules of projective indecomposable modules of the 0-Hecke algebra.

CONTENTS

1.	Int	roduction	2
2.	Pre	eliminaries	L.
	2.1.	Compositions	Ц,
	2.2.	The weak Bruhat orders on the symmetric group	6
	2.3.	Modules of the 0-Hecke algebras from intervals and posets	7
3.	An	equivalence relation on $Int(n)$ and its poset-theoretic characterization	8
4.	Eq	uivalence classes of lower and upper descent weak Bruhat intervals	13
	4.1.	The canonical diagram posets	13

Date: December 30, 2024.

²⁰²⁰ Mathematics Subject Classification. 06A07, 20C08, 05E10, 05E05.

Key words and phrases. Weak Bruhat order, 0-Hecke algebra, labeled poset, Quasisymmetric function.

S.-I. CHOI, S.-Y. NAM, AND Y.-T. OH

	4.2.	The minimal and maximal elements of the equivalence class	18
5.	Lo	wer and upper descent intervals from quotient modules and submodules of	
	pro	bjective indecomposable $H_n(0)$ -modules	28
	5.1.	Lower descent intervals from quotient modules of projective indecomposable	
		$H_n(0)$ -modules	29
	5.2.	Upper descent intervals from submodules of projective indecomposable	
		$H_n(0)$ -modules I	41
	5.3.	Upper descent intervals from submodules of projective indecomposable	
		$H_n(0)$ -modules II	43
R	efere	nces	47

1. INTRODUCTION

Weak Bruhat intervals play a crucial role not only in the combinatorics of Coxeter groups but also in the representation theory of both generic and degenerate Hecke algebras associated with these groups.

For each positive integer n, let \mathfrak{S}_n denote the symmetric group on the set $[n] := \{1, 2, \ldots, n\}$. In this paper, we focus on weak Bruhat intervals in \mathfrak{S}_n . We define $\operatorname{Int}(n)$ to be the set of all nonempty weak Bruhat intervals in \mathfrak{S}_n . Unless explicitly stated otherwise, "weak Bruhat interval" will always refer to a left weak Bruhat interval throughout this paper. A central problem concerning $\operatorname{Int}(n)$ is the classification of its elements according to suitable equivalence relations.

We begin by introducing two significant and well-known equivalence relations. Let P_n denote the set of posets with ground set [n]. Each poset $P \in \mathsf{P}_n$ can be identified with the labeled poset (P, ω) with the labeling $\omega : P \to [n]$ given by $\omega(i) = i$. Consequently, to each poset $P \in \mathsf{P}_n$, one can associate the following generating function for its *P*-partitions:

$$K_P := \sum_{f:P\text{-partition}} x_1^{|f^{-1}(1)|} x_2^{|f^{-1}(2)|} \cdots$$

The first equivalence relation arises in the context of P-partition generating functions. For $P_1, P_2 \in \mathsf{P}_n$, define $P_1 \stackrel{K}{\simeq} P_2$ if $K_{P_1} = K_{P_2}$. The classification of posets in P_n with respect to this equivalence relation remains a long-standing open problem (for example, see [2, 23, 32, 28, 24, 20]). Additionally, Björner–Wachs demonstrated in [9, Theorem 6.8] that the intervals in $\operatorname{Int}(n)$ correspond precisely to regular posets in P_n under the mapping $P \mapsto$ $\Sigma_L(P)$, where $\Sigma_L(P)$ denotes the set $\{\sigma \in \mathfrak{S}_n \mid \sigma(i) \leq \sigma(j) \text{ for all } i, j \in [n] \text{ with } i \preceq_P j\}$. Through this correspondence, we can induce the equivalence relation \cong on $\operatorname{Int}(n)$, defined by $I \stackrel{K}{\cong} J$ if $P_I \stackrel{K}{\cong} P_J$, where P_I and P_J denote the unique regular posets in P_n such that $\Sigma_L(P_I) = I$ and $\Sigma_L(P_J) = J$, respectively. The classification of intervals in $\operatorname{Int}(n)$ under this equivalence relation also remains an open problem (see [17, Section 5]).

 $\mathbf{2}$

The second equivalence relation arises in the context of $H_n(0)$ -modules associated with the posets in P_n (for the definition of the 0-Hecke algebra $H_n(0)$, see Section 2.3). Let $P \in \mathsf{P}_n$. In [14, Definition 3.18], Duchamp-Hivert-Thibon defined a right $H_n(0)$ -module M_P associated with P. Building on this concept, Kim-Lee-Oh introduced a left $H_n(0)$ module \mathbf{M}_P associated with P in [18, Definition 2.8] through a slight modification of the original module. For $P_1, P_2 \in \mathsf{P}_n$, we define $P_1 \cong P_2$ if the modules \mathbf{M}_{P_1} and \mathbf{M}_{P_2} are isomorphic as left $H_n(0)$ -modules. The relation \cong is particularly important since it refines $\stackrel{K}{\cong}$. However, the classification of posets in P_n with respect to \cong remains an open problem. As noted earlier, according to [9, Theorem 6.8], we can define the equivalence relation \cong^M on $\operatorname{Int}(n)$ by stating that $I \cong J$ if $P_I \cong P_J$. On the other hand, Jung, Kim, Lee, and Oh [17] associated to each left weak Bruhat interval I an $H_n(0)$ -module $\mathsf{B}(I)$, referred to as the weak Bruhat interval module associated with I. By definition, $\mathsf{B}(I)$ is equal to M_{P_I} , and hence $I \cong^M J$ if and only if $\mathsf{B}(I) \cong \mathsf{B}(J)$ as $H_n(0)$ -modules (see Section 2.3). Although the set under consideration is restricted from P_n to $\operatorname{Int}(n)$, the classification of intervals in $\operatorname{Int}(n)$ under \cong remains an open problem as well (see [17, 18]).

We introduce the equivalence relation $\stackrel{D}{\simeq}$ on $\operatorname{Int}(n)$, which plays a central role in this study. This relation is defined as $I \stackrel{D}{\simeq} J$ if there exists a (left) descent-preserving poset isomorphism between I and J. Recently, Kim-Lee-Oh showed that $\stackrel{D}{\simeq}$ refines $\stackrel{M}{\simeq}$ and coincides with $\stackrel{M}{\simeq}$ on the subset of weak Bruhat intervals corresponding to regular Schurlabeled posets on [n]. Furthermore, they successfully classified weak Bruhat intervals within this subset with respect to $\stackrel{D}{\simeq}$. They also conjectured that $\stackrel{D}{\simeq}$ and $\stackrel{M}{\simeq}$ are, in fact, identical ([18, Theorem 5.5, Theorem 4.7, and Conjecture 7.2]). Recently, Yang-Yu [33] proved that this conjecture holds for all weak Bruhat interval modules in arbitrary finite Coxeter types beyond type A.

Given an equivalence class C of $(\operatorname{Int}(n), \stackrel{D}{\simeq})$, define a partial order \preceq on C by $[\sigma, \rho]_L \preceq [\sigma', \rho']_L$ if and only if $\sigma \preceq_R \sigma'$. It was shown in [18, Theorem 4.6] that (C, \preceq) forms an interval. Specifically, with min $C := [\sigma_0, \rho_0]_L$ and max $C := [\sigma_1, \rho_1]_L$, the poset (C, \preceq) is isomorphic to the right weak Bruhat interval $[\sigma_0, \sigma_1]_R$. The main purpose of the present paper is to investigate the equivalence classes of weak Bruhat intervals that are in distinguished form but not necessarily regular Schur-labeled. Specifically, these intervals take the form

$$[w_0(S), \rho]_L$$
 or $[\sigma, w_1(S)]_L$,

where $w_0(S)$ is the longest element in the parabolic subgroup \mathfrak{S}_S of \mathfrak{S}_n , generated by $\{s_i \mid i \in S\}$ for each subset S of [n-1], and $w_1(S)$ is the longest element in the set of minimal length representatives for left $\mathfrak{S}_{[n-1]\backslash S}$ -cosets. Björner–Wachs demonstrated in [8, Theorem 6.2] that for $S \subseteq S' \subseteq [n-1]$, the set $\{w \in \mathfrak{S}_n \mid S \subseteq \text{Des}_R(w) \subseteq S'\}$ is given by the weak Bruhat interval $[w_0(S), w_1(S')]_L$. Therefore, we refer to an interval of the form $[w_0(S), \rho]_L$ as a *lower descent interval* and an interval of the form $[\sigma, w_1(S)]_L$ as an *upper descent interval*. From the context of the representation theory of 0-Hecke algebras,

lower and upper descent intervals arise from suitable quotient modules and submodules from projective modules with basis $[w_0(S), w_1(S')]_L$ (see Section 5).

In Section 3, we provide a poset-theoretical characterization of the equivalence relation $\stackrel{D}{\simeq}$ on $\operatorname{Int}(n)$. More concretely, we show that for $I, J \in \operatorname{Int}(n), I \stackrel{D}{\simeq} J$ if and only if one of P_I and P_J can be obtained from the other by repeatedly applying label changes to pairs that are comparable but not in the covering relation (Theorem 3.8). This characterization facilitates a substantially enhanced understanding of the equivalence relation within a combinatorial framework.

In Section 4, we investigate the equivalence class C of a lower or upper descent interval.

In Section 4.1, given a diagram D with n boxes, we introduce two posets, $P_{F_D^{\downarrow}}$ and $P_{F_D^{\rightarrow}}$, referred to as the *canonical diagram posets* associated with D. In [12, Section 5.1], it is shown that every upper descent interval appears as $P_{F_D^{\rightarrow}}$ for some diagram D. Here, we show that every lower descent interval appears as $P_{F_D^{\downarrow}}$ for some diagram D (Proposition 4.9).

In Section 4.2, we characterize min C and max C by identifying posets $P, Q \in \mathsf{P}_n$ such that min $C = \Sigma_L(P)$ and max $C = \Sigma_L(Q)$. In particular, if C is the equivalence class of $[w_0(S), \rho]_L$ (respectively $[\sigma, w_1(S)]_L$), then min $C = [w_0(S), \rho]_L$ (respectively max $C = [\sigma, w_1(S)]_L$) (Theorem 4.12). For general lower or upper descent intervals, explicitly describing the poset structure of (C, \preceq) is challenging. However, under a suitable condition, we establish that $(C, \preceq) \cong (\mathsf{ST}(D), \leq)$ as posets, where D is a diagram related to the given interval and $(\mathsf{ST}(D), \leq)$ denotes the poset of standard tableaux on D. In this case, the equivalence class C of a lower descent interval has an upper descent interval as max C, and the equivalence class C of an upper descent interval has a lower descent interval as min C (Theorem 4.16). Using this property, we determine the injective hull of $\mathsf{B}([w_0(S), \rho]_L)$ and the projective cover of $\mathsf{B}([\sigma, w_1(S)]_L)$ (Corollary 4.19).

In Section 5, we examine lower and upper descent intervals arising from a projective indecomposable $H_n(0)$ -module. In the context of the representation theory of 0-Hecke algebras, lower and upper descent intervals arise from appropriately selected quotient modules and submodules of the projective modules with the basis $[w_0(S), w_1(S')]_R$, where $S \subseteq S' \subseteq [n-1]$.

In Section 5.1, we consider the $H_n(0)$ -modules arising from the sequence of surjective $H_n(0)$ -module homomorphisms

$$\mathbf{P}_{\alpha} \longrightarrow \mathcal{V}_{\alpha} \longrightarrow X_{\alpha} \longrightarrow \widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \longrightarrow \mathbf{F}_{\alpha}$$

given in [11, Corollary 4.6]. It was shown in [17] that all the modules in this series are, up to isomorphism, weak Bruhat interval modules and that the corresponding intervals are lower descent intervals. We show that all of them satisfy the condition in Theorem 4.16 and provide explicit descriptions of the equivalence classes C, along with the poset structures of (C, \preceq) (Proposition 5.4, Proposition 5.8 and Proposition 5.15). Combining these with

Corollary 4.19, we determine injective hulls of \mathcal{V}_{α} , X_{α} , and $\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}$ in a uniform manner (Corollary 5.6, Corollary 5.10, and Corollary 5.17). It should be noted that our injective hull of X_{α} is a projective indecomposable module, and that an injective hull of \mathcal{V}_{α} was already constructed in [10, Theorem 4.11] in a different manner.

In Section 5.2, by applying the anti-involution twist of $\theta \circ \chi$ to the above sequence, we obtain the sequence of injective $H_n(0)$ -module homomorphisms

$$\mathbf{F}_{\alpha^{c}} \longrightarrow \mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \longrightarrow \mathcal{R}X_{\alpha} \longrightarrow \mathcal{R}\mathcal{V}_{\alpha} \longrightarrow \mathbf{P}_{\alpha^{i}}$$

(see Lemma 5.18). We provide explicit descriptions of the equivalence classes, the poset structures, and the projective covers for these modules (Proposition 5.19 and Corollary 5.20).

In Section 5.3, we discuss the submodules \mathbf{Q}_{α}^{r} of projective $H_{n}(0)$ -modules that arise from the representation theory of 0-Hecke–Clifford algebras. Here, α ranges over the set of peak compositions of n. These modules are spanned by standard peak immaculate tableaux and are related to the quasisymmetric Schur Q-functions. Given a peak composition α of n, we provide an explicit description of the equivalence class, the poset structure, as well as a projective cover and injective hull for \mathbf{Q}_{α}^{r} (Theorem 5.22 and Theorem 5.24). As an importance consequence, we derive that \mathbf{Q}_{α}^{r} is indecomposable (Corollary 5.25).

2. Preliminaries

Throughout this paper, let n be a positive integer. Define [n] as $\{1, 2, ..., n\}$, and set $[0] := \emptyset$.

2.1. Compositions. A composition α of a nonnegative integer n, denoted by $\alpha \models n$, is a finite ordered list of positive integers $(\alpha_1, \alpha_2, \ldots, \alpha_l)$ satisfying $\sum_{i=1}^{l} \alpha_i = n$. We call lthe *length* of α and denote it by $\ell(\alpha)$. For convenience, we define the empty composition \emptyset to be the unique composition of size and length 0. If $\alpha_1 \ge \alpha_2 \ge \cdots \ge \alpha_l$, then we say that α is a *partition* of n.

Given $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_l) \models n$ and $I = \{i_1 < i_2 < \dots < i_p\} \subset [n-1]$, let


$$set(\alpha) := \{ \alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{l-1} \},\\comp(I) := (i_1, i_2 - i_1, \dots, n - i_p).$$

The set of compositions of n is in bijection with the set of subsets of [n-1] under the correspondence $\alpha \mapsto \operatorname{set}(\alpha)$ (or $I \mapsto \operatorname{comp}(I)$). Define

- α^{r} by the reverse composition $(\alpha_{l}, \ldots, \alpha_{1})$,
- α^{c} by the complement composition satisfying $set(\alpha^{c}) = [n-1] \setminus set(\alpha)$,
- α^{t} by the transpose composition $(\alpha^{r})^{c}$, and
- $\tilde{\alpha}$ by the partition obtained by sorting the parts of α in the weakly decreasing order.

And, for $I \subset [n-1]$, define $I^r := \operatorname{set}(\operatorname{comp}(I)^r)$ and $I^t := \operatorname{set}(\operatorname{comp}(I)^t)$.

Let $\alpha = (\alpha_1, \ldots, \alpha_l) \models n$. We define the *composition diagram* $\mathsf{cd}(\alpha)$ of α as a leftjustified array of n boxes where the *i*th row from the bottom has α_i boxes for $1 \le i \le l$. We also define the *ribbon diagram* $\mathsf{rd}(\alpha)$ of α as the connected skew diagram without 2×2 boxes, such that the *i*th column from the left has α_i boxes. For instance, if $\alpha = (1, 3, 2)$, then



2.2. The weak Bruhat orders on the symmetric group. The symmetric group \mathfrak{S}_n is generated by simple transpositions $s_i := (i, i + 1)$ with $1 \le i \le n - 1$. An expression of $\sigma \in \mathfrak{S}_n$ that uses the minimal number of simple transpositions is called a *reduced* expression for σ . This minimal number is denoted by $\ell(\sigma)$ and called the *length* of σ .

The *left descent set* and *right descent set* of a permutation σ are defined by

$$Des_{L}(\sigma) := \{i \in [n-1] \mid \ell(s_{i}\sigma) < \ell(\sigma)\} \text{ and} \\ Des_{R}(\sigma) := \{i \in [n-1] \mid \ell(\sigma s_{i}) < \ell(\sigma)\},$$

respectively. The left weak Bruhat order \leq_L and right weak Bruhat order \leq_R on \mathfrak{S}_n are defined to be the partial order on \mathfrak{S}_n whose covering relation \leq_L^c and \leq_R^c are given as follows:

$$\sigma \preceq_L^c s_i \sigma$$
 if and only if $i \notin \text{Des}_{\mathcal{L}}(\sigma)$ and
 $\sigma \preceq_R^c \sigma s_i$ if and only if $i \notin \text{Des}_{\mathcal{R}}(\sigma)$,

respectively. Given $\sigma, \rho \in \mathfrak{S}_n$, if $\sigma \preceq_L \rho$, then the *(left) weak Bruhat interval* from σ to ρ is defined by

$$[\sigma,\rho]_L := \{\gamma \in \mathfrak{S}_n \mid \sigma \preceq_L \gamma \preceq_L \rho\}$$

and if $\sigma \preceq_R \rho$, then the right weak Bruhat interval from σ to ρ is defined by

$$[\sigma,\rho]_R := \{\gamma \in \mathfrak{S}_n \mid \sigma \preceq_R \gamma \preceq_R \rho\}.$$

For $S \subseteq [n-1]$, let \mathfrak{S}_S be the parabolic subgroup of \mathfrak{S}_n generated by $\{s_i \mid i \in S\}$ and $w_0(S)$ the longest element in \mathfrak{S}_S . When S = [n-1], we simply write w_0 for $w_0(S)$. An element $w \in \mathfrak{S}_n$ can be written uniquely as w = zu, where $z \in \mathfrak{S}^S$ and \mathfrak{S}_S , with the property that $\ell(w) = \ell(z) + \ell(u)$. Here $\mathfrak{S}^S := \{z \in \mathfrak{S}_n \mid \text{Des}_R(z) \subseteq S^c\}$ is the set of minimal length representatives for left \mathfrak{S}_S -cosets, where $S^c = [n-1] \setminus S$.

Theorem 2.1. ([8, Theorem 6.2]) Given $S \subseteq S' \subseteq [n-1]$, the set $\{w \in \mathfrak{S}_n \mid S \subseteq Des_{\mathbb{R}}(w) \subseteq S'\}$ is exactly the weak Bruhat interval $[w_0(S), w_1(S')]_L$, where $w_0(S)$ is the longest element in \mathfrak{S}_S and $w_1(S')$ is the longest element in $\mathfrak{S}^{S'^c}$.

Since $w_1(S') = w_0 w_0(S'^c)$, Theorem 2.1 can be rewritten as

(2.1)
$$\{w \in \mathfrak{S}_n \mid S \subseteq \text{Des}_{\mathcal{L}}(w) \subseteq S'\} = [w_0(S), w_0(S'^c)w_0]_R.$$

2.3. Modules of the 0-Hecke algebras from intervals and posets. The 0-Hecke algebra $H_n(0)$ is the associative \mathbb{C} -algebra with 1 generated by the elements $\overline{\pi}_1, \overline{\pi}_2, \ldots, \overline{\pi}_{n-1}$ subject to the following relations:

(2.2)
$$\overline{\pi}_{i}^{2} = -\overline{\pi}_{i} \quad \text{for } 1 \leq i \leq n-1,$$
$$\overline{\pi}_{i}\overline{\pi}_{i+1}\overline{\pi}_{i} \equiv \overline{\pi}_{i+1}\overline{\pi}_{i}\overline{\pi}_{i+1} \quad \text{for } 1 \leq i \leq n-2,$$
$$\overline{\pi}_{i}\overline{\pi}_{j} \equiv \overline{\pi}_{j}\overline{\pi}_{i} \quad \text{if } |i-j| \geq 2.$$

Another set of generators consists of $\pi_i := \overline{\pi}_i + 1$ for i = 1, 2, ..., n - 1 with the same relations as above except that $\pi_i^2 = \pi_i$.

For any reduced expression $s_{i_1}s_{i_2}\cdots s_{i_p}$ for $\sigma \in \mathfrak{S}_n$, let $\overline{\pi}_{\sigma} := \overline{\pi}_{i_1}\overline{\pi}_{i_2}\cdots\overline{\pi}_{i_p}$ and $\pi_{\sigma} := \pi_{i_1}\pi_{i_2}\cdots\pi_{i_p}$. It is well known that these elements are independent of the choices of reduced expressions, and both $\{\overline{\pi}_{\sigma} \mid \sigma \in \mathfrak{S}_n\}$ and $\{\pi_{\sigma} \mid \sigma \in \mathfrak{S}_n\}$ are \mathbb{C} -bases for $H_n(0)$.

According to [27], there are 2^{n-1} pairwise inequivalent irreducible $H_n(0)$ -modules and 2^{n-1} pairwise inequivalent projective indecomposable $H_n(0)$ -modules, which are naturally indexed by compositions of n. For a composition α of n, let \mathbf{F}_{α} denote the 1-dimensional \mathbb{C} -vector space corresponding to the composition α of n, spanned by a vector v_{α} . For each $1 \leq i \leq n-1$, define an action of the generator π_i of $H_n(0)$ as follows:

$$\pi_i(v_\alpha) = \begin{cases} 0 & i \in \operatorname{set}(\alpha), \\ v_\alpha & i \notin \operatorname{set}(\alpha). \end{cases}$$

This module is the irreducible 1-dimensional $H_n(0)$ -module corresponding to α . And, the projective indecomposable $H_n(0)$ -module corresponding to α is given by the submodule $\mathbf{P}_{\alpha} = H_n(0)\pi_{w_0(\text{set}(\alpha)^c)}\overline{\pi}_{w_0(\text{set}(\alpha))}$ of the regular representation of $H_n(0)$.

One can construct modules of the 0-Hecke algebra using various combinatorial objects. This paper focuses on modules arising from weak Bruhat intervals and posets in P_n . First, we review the weak Bruhat interval $H_n(0)$ -modules introduced by Jung-Kim-Lee-Oh ([17]).

Definition 2.2. ([17, Definition 1]) Let $I \in Int(n)$.

(a) The weak Bruhat interval module associated with I, denoted by $\mathsf{B}(I)$, is the left $H_n(0)$ -module with $\mathbb{C}I$ as the underlying space and with the $H_n(0)$ -action defined by

$$\pi_i \cdot \gamma := \begin{cases} \gamma & \text{if } i \in \text{Des}_{\mathcal{L}}(\gamma), \\ 0 & \text{if } i \notin \text{Des}_{\mathcal{L}}(\gamma) \text{ and } s_i \gamma \notin I, \\ s_i \gamma & \text{if } i \notin \text{Des}_{\mathcal{L}}(\gamma) \text{ and } s_i \gamma \in I \end{cases}$$

for $1 \leq i \leq n-1$ and $\gamma \in I$.

(b) The negative weak Bruhat interval module associated with I, denote by $\overline{\mathsf{B}}(I)$, is the left $H_n(0)$ -module with $\mathbb{C}I$ as the underlying space and with the $H_n(0)$ -action defined by

$$\overline{\pi}_i \star \gamma := \begin{cases} -\gamma & \text{if } i \in \text{Des}_{\text{L}}(\gamma), \\ 0 & \text{if } i \notin \text{Des}_{\text{L}}(\gamma) \text{ and } s_i \gamma \notin I, \\ s_i \gamma & \text{if } i \notin \text{Des}_{\text{L}}(\gamma) \text{ and } s_i \gamma \in I \end{cases}$$

for $1 \leq i \leq n-1$ and $\gamma \in I$.

Indeed B(I) is the θ -twist of B(I). For the definition of the θ -twist, see Section 5.2.

Next, we review the left $H_n(0)$ -modules arising from posets. To begin with, for given $P \in \mathsf{P}_n$, we define

(2.3)
$$\Sigma_L(P) := \{ \sigma \in \mathfrak{S}_n \mid \sigma(i) \le \sigma(j) \text{ for all } i, j \in [n] \text{ with } i \preceq_P j \}.$$

Definition 2.3. ([18, Definition 2.8]) Let $P \in \mathsf{P}_n$. Define the $H_n(0)$ -module \mathbf{M}_P to be the underlying space $\mathbb{C}\Sigma_L(P)$ and with the $H_n(0)$ -action:

$$\pi_i \cdot \gamma := \begin{cases} \gamma & \text{if } i \in \text{Des}_{\mathcal{L}}(\gamma), \\ 0 & \text{if } i \notin \text{Des}_{\mathcal{L}}(\gamma) \text{ and } s_i \gamma \notin \Sigma_L(P), \\ \gamma s_i & \text{if } i \notin \text{Des}_{\mathcal{L}}(\gamma) \text{ and } s_i \gamma \in \Sigma_L(P) \end{cases}$$

for $i \in [n-1]$ and $\gamma \in \Sigma_L(P)$.

It should be noted that in the case where $\Sigma_L(P)$ is a left weak Bruhat interval, \mathbf{M}_P is identical to $\mathsf{B}(\Sigma_L(P))$.

3. An equivalence relation on Int(n) and its poset-theoretic Characterization

Let $\operatorname{Int}(n)$ denote the set of nonempty weak Bruhat intervals in \mathfrak{S}_n . This section provides a poset-theoretic characterization of the equivalence relation introduced in [18] on this set. We start by recalling the definition of this equivalence relation. For $I_1, I_2 \in \operatorname{Int}(n)$, a poset isomorphism $f: (I_1, \preceq_L) \to (I_2, \preceq_L)$ is called *descent-preserving* if

$$\operatorname{Des}_{\mathrm{L}}(\gamma) = \operatorname{Des}_{\mathrm{L}}(f(\gamma)) \text{ for all } \gamma \in I_1.$$

Define an equivalence relation $\stackrel{D}{\simeq}$ on $\operatorname{Int}(n)$ such that $I_1 \stackrel{D}{\simeq} I_2$ if there is a descent-preserving (poset) isomorphism between (I_1, \preceq_L) and (I_2, \preceq_L) . One of the key reasons for considering this equivalence relation is its role in refining the classification of the $H_n(0)$ -modules $\mathsf{B}(I)$ associated with $I \in \operatorname{Int}(n)$ up to isomorphism. Specifically, if $I_1 \stackrel{D}{\simeq} I_2$, then $\mathsf{B}(I_1) \cong \mathsf{B}(I_2)$ as $H_n(0)$ -modules. It was conjectured in [18, Conjecture 7.2] that the converse holds more generally in $\operatorname{Int}(n)$. It was proved in [18, Section 4 and Section 5] that the converse is true within the subset

 $\{\Sigma_L(P) \mid P \text{ is a regular Schur labeled poset on } [n]\} \subset \operatorname{Int}(n).$

Recently, Yang-Yu [33] proved that this conjecture holds for all weak Bruhat interval modules in arbitrary finite Coxeter types beyond type A. We begin with the following characterizations of $\stackrel{D}{\simeq}$.

Lemma 3.1. Let $I_1, I_2 \in Int(n)$. Then the following are equivalent.

- (a) $I_1 \stackrel{D}{\simeq} I_2$.
- (b) $B(I_1) \cong B(I_2)$ as $H_n(0)$ -modules.
- (c) $\overline{\mathsf{B}}(I_1) \cong \overline{\mathsf{B}}(I_2)$ as $H_n(0)$ -modules.

Proof. The equivalence (a) \Leftrightarrow (b) was established in [33, Theorem 4.11]. Furthermore, the equivalence (b) \Leftrightarrow (c) follows from [17, Theorem 4 (2)].

Every weak Bruhat interval corresponds to the set of linear extensions of a regular poset. We now recall the definition of regular posets.

Definition 3.2. ([9, p. 110]) A poset $P \in \mathsf{P}_n$ is said to be *regular* if the following holds: for all $x, y, z \in [n]$ with $x \leq_P z$, if x < y < z or z < y < x, then $x \leq_P y$ or $y \leq_P z$.

We denote by RP_n the set of all regular posets in P_n . The following theorem shows how regular posets can be characterized in terms of weak Bruhat intervals.

Theorem 3.3. ([9, Theorem 6.8]) Let U be a nonempty subset of \mathfrak{S}_n . Then, the following conditions are equivalent:

- (1) U is a weak Bruhat interval.
- (2) $U = \Sigma_L(P)$ for some $P \in \mathsf{RP}_n$.

Consider the map

$$\eta: \mathsf{P}_n \to \mathcal{P}(\mathfrak{S}_n), \quad P \mapsto \Sigma_L(P),$$

where $\mathcal{P}(\mathfrak{S}_n)$ is the power set of \mathfrak{S}_n . One can see that η is injective. Combining this with Theorem 3.3, we obtain a one-to-one correspondence

$$\eta|_{\mathsf{RP}_n} : \mathsf{RP}_n \to \operatorname{Int}(n), \quad P \mapsto \Sigma_L(P).$$

For $I \in \text{Int}(n)$, we denote by P_I the regular poset such that $\Sigma_L(P_I) = I$. Throughout this paper, we will identify any weak Bruhat interval $I \in \mathfrak{S}_n$ with the regular poset $P_I \in \mathsf{RP}_n$. Based on this identification, we provide a poset-theoretical characterization of the equivalence relation $\stackrel{D}{\simeq}$ on Int(n).

Definition 3.4. ([2, Definition 1.5]). An edge-decorated poset is a poset P such that each edge in its Hasse diagram is assigned to be either weak or strict.

Every poset $P \in \mathsf{P}_n$ can be equipped with the structure of an edge-decorated poset. This is achieved by viewing P as a labeled poset (P, ω) , where ω denotes the associated labeling. Following the convention for drawing Hasse diagrams of posets, we draw a bold edge (referred to as a *strict* edge) between x and y when $x \leq_P y$ and $\omega(x) > \omega(y)$, and a plain edge (referred to as a *weak* edge) when $x \leq_P y$ and $\omega(x) < \omega(y)$ in the Hasse diagram of P. Here \leq_P is used to denote the partial order of P and \leq the usual order on [n].

We say that two labeled posets (P, ω) and (Q, τ) are isomorphic, denoted by $(P, \omega) \stackrel{L}{\simeq} (Q, \tau)$, if there exists a poset isomorphism from P to Q that maps strict edges and weak edges in P to strict edges and weak edges in Q, respectively. Then it follows from the definition of (P, ω) -partition generating function that

$$(P,\omega) \stackrel{L}{\simeq} (Q,\tau) \Rightarrow (P,\omega) \stackrel{K}{\simeq} (Q,\tau)$$

(see [23, Lemma 3.6]).

The symmetric group \mathfrak{S}_n acts on P_n by composing the labeling with permutations. Specifically, for $\sigma \in \mathfrak{S}_n$ and $(P, \omega) \in \mathsf{P}_n$, the action is defined as

$$\sigma \cdot (P, \omega) = (P, \sigma \circ \omega).$$

Lemma 3.5. Let $P \in \mathsf{RP}_n$ and let $1 \le i \le n-1$.

- (a) If (i, i+1) is a comparable pair that is not in the covering relation in P, then $s_i \cdot P$ is equal to P as an edge-decorated poset and remains regular.
- (b) If (i, i + 1) is in the covering relation in P, then $s_i \cdot P$ is not equal to P as an edge-decorated poset, but it remains regular.
- (c) If (i, i+1) is an incomparable pair in P, then $s_i \cdot P$ equals P as an edge-decorated poset. Moreover, either $\Sigma_L(P) = \Sigma_L(s_i \cdot P)$ or $s_i \cdot P$ is not regular.

Proof. The assertions are straightforward when n = 1 or n = 2. Thus, we assume $n \ge 3$.

(a) Since i and i + 1 are comparable but not in the covering relation in P, it is evident that $s_i \cdot P$ is identical to P as edge-decorated posets.

To show that $s_i \cdot P$ remains regular, choose any triple (x, y, z) in $s_i \cdot P$ with $x \preceq_{s_i \cdot P} z$. If none of x, y, z are equal to i or i+1, the triple (x, y, z) remains unchanged between P and $s_i \cdot P$, so the regularity is preserved. If only one of x, y, z equals i or i+1, the regularity of P ensures that $s_i \cdot P$ remains regular as well. Otherwise, the following cases are possible:

- x = i < y = i + 1 < z,
- x < y = i < z = i + 1,
- z = i < y = i + 1 < x, or
- z < y = i < x = i + 1.

In all these cases, the assumption that (i, i+1) is a comparable pair in P guarantees that either $x \preceq_{s_i \cdot P} y$ or $y \preceq_{s_i \cdot P} z$. Therefore, $s_i \cdot P$ satisfies the regularity conditions.

(b) If $i \leq_P i + 1$ is in the covering relation in P, then $i + 1 \leq_{s_i \cdot P} i$ is in the covering relation in $s_i \cdot P$, making $s_i \cdot P$ distinct from P as an edge-decorated poset.

The regularity of $s_i \cdot P$ follows directly from the argument in (a), as the reversal of a covering relation does not disrupt regularity.

(c) Let (i, i + 1) be an incomparable pair in P. Since the relative orders in the Hasse diagrams of the connected components containing i and i + 1 in P remain unchanged, it follows that $s_i \cdot P$ is identical to P as an edge-decorated poset.

We now prove the second assertion. First, consider the case where every element $x \in P$ other than i, i + 1 is either incomparable to both i and i + 1, or comparable to both i and i + 1. In this case, the set of linear extensions remains unchanged, and thus $\Sigma_L(P) = \Sigma_L(s_i \cdot P)$. Next, suppose there exists an element $x \neq i, i + 1$ in P that is comparable to either i or i + 1, but not both. We focus on the case where $x \prec_P i$ or $i \prec_P x$, noting that the argument for the case where $x \prec_P i + 1$ or $i + 1 \prec_P x$ follows analogously. We have the following four subcases:

- $x \prec_P i$ and x < i: It holds that $x \prec_{s_i \cdot P} i + 1$ in $s_i \cdot P$ and x < i < i + 1. However, $x \not\preceq_{s_i \cdot P} i$ and $i \not\preceq_{s_i \cdot P} i + 1$ in $s_i \cdot P$. Thus $s_i \cdot P$ is not regular.
- $x \prec_P i$ and i + 1 < x: It holds that i < i + 1 < x. However, $x \not\preceq_P i + 1$ and $i + 1 \not\preceq_P i$ in $\cdot P$. This contradicts the assumption that P is regular.
- $i \prec_P x$ and x < i: It holds that $i + 1 \prec_{s_i \cdot P} x$ in $s_i \cdot P$ and x < i < i + 1. However, $i + 1 \not\preceq_{s_i \cdot P} i$ and $i \not\preceq_{s_i \cdot P} x$ in $s_i \cdot P$. Thus $s_i \cdot P$ is not regular.
- $i \prec_P x$ and i+1 < x: It holds that i < i+1 < x. However, $i \not\preceq_P i+1$ and $i+1 \not\preceq_P x$ in $\cdot P$. This contradicts the assumption that P is regular.

Remark 3.6. From the perspective of equivalence relations, we note that in Lemma 3.5 (a), $s_i \cdot P \stackrel{M}{\simeq} P$, and thus $s_i \cdot P \stackrel{K}{\simeq} P$; in (b), $s_i \cdot P \stackrel{K}{\simeq} P$, and thus $s_i \cdot P \stackrel{M}{\simeq} P$; and in (c), $s_i \cdot P \stackrel{K}{\simeq} P$.

Example 3.7. Consider the regular poset

$$P = 2 + 1 = 3 + 1 = 1$$

Note that (3, 4) is a comparable non-covering pair, (2, 3) is a covering pair, and (1, 2), (4, 5) are incomparable pairs in P. Now, consider the posets obtained by applying simple transpositions to P:

$$s_3 \cdot P = 5$$
 3 $s_2 \cdot P = 5$ 4 $s_1 \cdot P = 5$ 4 $s_1 \cdot P = 5$ 4 $s_2 \cdot P = 5$ 4 $s_1 \cdot P = 5$ 4 $s_2 \cdot P = 5$ 4 $s_3 \cdot P = 5$

We observe that the posets P, $s_3 \cdot P$, and $s_1 \cdot P$ are identical as edge-decorated posets. However, while $s_3 \cdot P$ is regular, $s_1 \cdot P$ is not. Additionally, $K_P \neq K_{s_2 \cdot P}$, since 12345 $\notin \Sigma_L(K_P)$.

We now present a poset-theoretic characterization of $\stackrel{D}{\simeq}$. For the proof, we introduce the notation

$$\mathcal{R}_{\rm st}(P) := \{ (x, y) \in [n]^2 \mid x \preceq_P y \text{ and } x \neq y \} \quad \text{for } P \in \mathsf{P}_n.$$

Theorem 3.8. Let $I, J \in \text{Int}(n)$. Then $I \stackrel{D}{\simeq} J$ if and only if $P_J = s_{i_r} \cdots s_{i_2} s_{i_1} \cdot P_I$ for some nonnegative integer r, where $(i_k, i_k + 1)$ is a comparable pair that is not in the covering relation in $s_{i_{k-1}} \cdots s_{i_2} s_{i_1} \cdot P_I$ for $1 \leq k \leq r$.

Proof. The assertion is straightforward when n = 1 or n = 2. Thus, we assume $n \ge 3$. In the trivial case where I = J, the assertion is clear, as for any weak Bruhat interval I in \mathfrak{S}_n , there exists a unique poset $P_I \in \mathsf{P}_n$ such that $\Sigma_L(P_I) = I$. Now, assume $I \stackrel{D}{\simeq} J$ with $I \ne J$. By [18, Theorem 4.6], for $1 \le i \le n-1$, the assertion is established by verifying the one-step equivalence: $I \stackrel{D}{\simeq} Is_i$ if and only if (i, i + 1) is a comparable pair that is not in the covering relation in P_I ; in this case $P_{Is_i} = s_i \cdot P_I$.

To begin with, we show that if (i, i+1) is a comparable pair in P_I that does not belong to the covering relation, then $P_{Is_i} = s_i \cdot P_I$ always holds. Suppose that $\Sigma_L(P_I) = [\sigma, \rho]_L$. Since P_I is regular and (i, i+1) is a comparable pair that is not in the covering relation in P_I , it follows from Lemma 3.5 that $s_i \cdot P_I$ is regular. So $\Sigma_L(s_i \cdot P_I)$ is a weak Bruhat interval, we put it by $[\sigma', \rho']_L$. We claim that $\sigma' = \sigma s_i$ and $\rho' = \rho s_i$.

For a given regular poset P, let $\Sigma_L(P) = [\delta, \eta]_L$. It is straightforward to verify that δ and η can be determined from P as follows: for $1 \leq k \leq n$,

(3.1)
$$\delta(k) = |\{x \mid x \leq_P k\} \cup \{x \mid x \text{ is incomparable to } k \text{ in } P \text{ and } x < k\}|,$$

$$\eta(k) = |\{x \mid x \preceq_P k\} \cup \{x \mid x \text{ is incomparable to } k \text{ in } P \text{ and } x > k\}|.$$

Applying this property to the posets P_I and $s_i \cdot P_I$, we observe that $\sigma'(i) = \sigma(i+1)$, $\rho'(i) = \rho(i+1)$, and all other entries remain unchanged. Therefore, we have

(3.2)
$$\Sigma_L(s_i \cdot P_I) = Is_i, \text{ equivalently } s_i \cdot P_I = P_{Is_i}.$$

Now, we prove the "if" direction. We have two cases.

• $i \prec_{P_I} i + 1$: From (2.3) it holds that $\gamma(i) < \gamma(i+1)$ for each $\gamma \in \Sigma_L(P)$. Since (i, i+1) is not in the covering relation in P_I , $\gamma(i) + 1 < \gamma(i+1)$. Then one can easily see that

$$\operatorname{Des}_{\mathrm{L}}(\gamma) = \operatorname{Des}_{\mathrm{L}}(\gamma s_i) \quad \text{for each } \gamma \in \Sigma_L(P_I).$$

Since $\Sigma_L(P_I) = I$ and $\Sigma_L(s_i \cdot P_I) = Is_i$, this implies a descent-preserving bijection $I \to Is_i, \ \gamma \mapsto \gamma s_i$. Hence, $I \simeq Is_i$.

• $i + 1 \prec_{P_I} i$: This case follows from a symmetric argument.

Next, we prove the "only if" direction. Suppose $I \cong^{D} Is_i$. Then there exists a descentpreserving bijection $f: I \to Is_i, \ \gamma \mapsto \gamma s_i$. Let $\Sigma_L(P_I) = [\sigma, \rho]_L$. We have two cases.

• $\rho(i) < \rho(i+1)$: Since $\sigma \preceq_L \rho$, it follows that $\sigma(i) < \sigma(i+1)$. By [9, Theorem 6.8], the strict relations in P_I are given by

$$\mathcal{R}_{\mathrm{st}}(P_I) = \{(x, y) \mid \sigma(x) < \sigma(y) \text{ and } \rho(x) < \rho(y)\}.$$

This implies $i \prec_{P_I} i + 1$. If i + 1 covers i, then it follows from (2.3) that $\sigma(i + 1) = \sigma(i) + 1$. It contradicts the descent-preserving map f. Hence, (i, i + 1) is a comparable pair but not in the covering relation in P_I .

• $\rho(i) > \rho(i+1)$: This case follows from a symmetric argument. This completes the proof.

4. Equivalence classes of lower and upper descent weak Bruhat intervals

In this section, we investigate the equivalence classes of lower and upper descent intervals, specifically weak Bruhat intervals of the forms

$$[w_0(S), \rho]_L$$
 or $[\sigma, w_1(S)]_L$, $(S \subseteq [n-1])$.

We first demonstrate that these intervals can be represented as $\Sigma_L(P)$ for some special posets $P \in \mathsf{P}_n$, which we refer to as *canonical diagram posets*. Next, given an equivalence class C, we provide the posets P and Q in P_n such that

min
$$C = \Sigma_L(P)$$
 and max $C = \Sigma_L(Q)$.

4.1. The canonical diagram posets. We recall the notion of canonical diagram posets as described in [12, Section 5.1]. Then, we examine their images under the typical involutions defined on P_n .

In this subsection, we consider an *n*-element subset of \mathbb{N}^2 , which we will treat as a diagram composed of *n* boxes located in the first quadrant. We identify each point (i, j) with the empty rectangle whose vertices are at (i - 1, j - 1), (i, j - 1), (i - 1, j), and (i, j) (see Example 4.8). Let \mathfrak{D}_n be the set of *n*-element subsets of \mathbb{N}^2 such that the corresponding diagram has no empty rows or columns within the smallest rectangle that can completely enclose the diagram. In particular, we consider the composition diagrams $\mathsf{cd}(\alpha)$ and the ribbon diagrams $\mathsf{rd}(\alpha)$ of size *n* as elements of \mathfrak{D}_n by positioning the lower leftmost box at (1, 1). Unless explicitly stated otherwise, we will assume throughout this section that $D \in \mathfrak{D}_n$.

Convention. In this paper, we regard a filling F of D with positive integers as a map

$$F: D \to \mathbb{Z}_{>0}, \quad (i,j) \mapsto F(i,j),$$

where F(i, j) denotes the entry in the box (i, j) of F. Given a filling F of D, we primarily utilize four reading words, denoted as $w_{TL}(F)$, $w_{LT}(F)$, $w_{LB}(F)$, and $w_{BL}(F)$, which are defined as follows:

- $w_{TL}(F)$ is the word obtained by reading the entries of F from top to bottom in each column, starting with the leftmost column.
- $w_{LT}(F)$ is the word obtained by reading the entries of F from left to right across each row, starting with the topmost row.
- $w_{LB}(F)$ is the word obtained by reading the entries of F from left to right across each row, starting with the bottommost row.
- $w_{BL}(F)$ is the word obtained by reading the entries of F from bottom to top in each column, starting with the leftmost column.

A filling F on D with positive integers is called *standard* if it contains the entries $1, 2, \ldots, n$, each appearing exactly once. Every word containing distinct entries from 1 to n is regarded as a permutation in \mathfrak{S}_n .

Definition 4.1. Let F be a standard filling of D.

(a) Let P_F denote the poset in P_n with partial order \preceq_{P_F} , defined by

 $i \leq_{P_F} j$ if and only if $x_i \leq x_j$ and $y_i \leq y_j$,

where (x_i, y_i) represents the position of *i* in the filling *F* for each $1 \le i \le n$.

(b) Define F_D^{\downarrow} (respectively F_D^{\rightarrow}) as the filling of D obtained by placing the integers $1, 2, \ldots, n$ sequentially, without repetition, down each column (respectively across each row), from top to bottom (respectively left to right), starting from the left-most column (respectively the uppermost row). We refer to $P_{F_D^{\downarrow}}$ and $P_{F_D^{\rightarrow}}$ as the canonical diagram posets associated with D.¹

Remark 4.2. In Definition 4.1(a), suppose P_F is regular. Let $\Sigma_L(P_F) = [\sigma, \rho]_L$. Using (3.1), one can obtain σ and ρ directly from F. To be precise, for $1 \le k \le n$,

 $\sigma(k) = |\{1 \le x \le n \mid x \text{ is lower-left of } k \text{ in } F\} \\ \cup \{1 \le x < k \mid x \text{ is strictly upper-left or strictly lower-right of } k \text{ in } F\}|, \text{ and } \rho(k) = |\{1 \le x \le n \mid x \text{ is lower-left of } k \text{ in } F\}$

 $\cup \{k < x \le n \mid x \text{ is strictly upper-left or strictly lower-right of } k \text{ in } F\}|.$

Since $P_{F_D^{\downarrow}}$ and $P_{F_D^{\rightarrow}}$ are regular posets, the sets $\Sigma_L(P_{F_D^{\downarrow}})$ and $\Sigma_L(P_{F_D^{\rightarrow}})$ form weak Bruhat intervals. Here, we investigate the properties of these intervals. To begin with, we introduce the necessary lemma and notations.

(i) Given a diagram $D \in \mathcal{D}_n$, let D^t be the diagram obtained by transposing the coordinates, that is, $D^t = \{(j, i) \mid (i, j) \in D\}$. Similarly, for a standard filling F on D, let F^t be the corresponding standard filling on D^t , defined by setting

$$F^{\mathrm{t}}(i,j) := F(j,i) \quad \text{for } (j,i) \in D.$$

Lemma 4.3. Let F be a standard filling on D. Then we have $P_F = P_{F^t}$.

Proof. To prove the assertion, we start by noting that $(F^t)^t = F$. Thus, it suffices to show that for all $1 \le i, j \le n$, we have that

$$i \preceq_{P_F} j \implies i \preceq_{P_{F^t}} j.$$

For $1 \leq i \leq n$, let (x_i, y_i) denote the position of i in the filling F. Suppose that $i \preceq_{P_F} j$. By the definition of P_F , this implies that

(4.1)
$$x_i \le x_j \quad \text{and} \quad y_i \le y_j.$$

In the filling F^{t} , the positions of i and j are transposed, with i now at (y_i, x_i) and j at (y_j, x_j) . The condition (4.1) can be rewritten as $y_i \leq y_j$ and $x_i \leq x_j$. Therefore, by the definition of the partial order for $P_{F^{t}}$, it follows that $i \leq_{P_{F^{t}}} j$.

¹Note that in [12], only the poset $P_{F_{D}}$ is referred to as the *canonical diagram poset associated with D*.

(ii) A filling T on D with entries 1, 2, ..., n is called a *standard tableau* on D if its entries are distinct and $T(i, j) \leq T(k, l)$ whenever $i \leq k$ and $j \leq l$. Let ST(D) denote the set of standard tableaux on D. Define

(4.2)
$$T'_D$$
 (respectively T_D)

to be the standard tableau on D obtained by sequentially placing the integers $1, 2, \ldots, n$ without repetition, along the columns (respectively rows), from bottom to top (respectively left to right), starting with the leftmost column (respectively the bottommost row).

(iii) For $P \in \mathsf{P}_n$, define \overline{P} as the poset in P_n where the order relation is given by

$$u \preceq_{\overline{P}} v \iff n+1-u \preceq_P n+1-v.$$

The map $\bar{}: \mathsf{P}_n \to \mathsf{P}_n, P \mapsto \overline{P}$, is clearly an involution. Furthermore, given a standard filling F on D, we denote by \overline{F} the standard filling obtained from F by replacing each entry i with n - i + 1 for all $1 \leq i \leq n$.

(iv) Let k and l denote the number of rows and columns of D, respectively. For $1 \le i \le k$ and $1 \le j \le l$, let r_i and c_j be the number of boxes in the *i*th row (from the top) and the *j*th column (from the left) of D, respectively. Then, define

$$r(D) := (r_1, r_2, \dots, r_k)$$
 and $c(D) := (c_1, c_2, \dots, c_l)$.

With the prerequisites (i)–(iv), we determine the minimal and maximal elements of the intervals $\Sigma_L(P_{F_D^{\downarrow}})$ and $\Sigma_L(P_{F_D^{\rightarrow}})$, which correspond to specific reading words derived from the standard tableaux.

Proposition 4.4. Let $D \in \mathfrak{D}_n$.

- (a) $\Sigma_L(P_{F_D^{\downarrow}}) = [\mathbf{w}_{\mathrm{LT}}(T'_D), \mathbf{w}_{\mathrm{LT}}(T_D)]_L$ and $\Sigma_L(P_{F_D^{\downarrow}}) = [\mathbf{w}_{\mathrm{TL}}(T'_D), \mathbf{w}_{\mathrm{TL}}(T_D)]_L$.
- (b) $w_{LT}(T_D) = w_1(\text{set}(\mathbf{r}(D)))$, thus $\Sigma_L(P_{F_D})$ is an upper descent interval.
- (c) $w_{\mathrm{TL}}(T'_D) = w_0(\operatorname{set}(\mathbf{c}(D))^{\mathrm{c}}), \text{ thus } \Sigma_L(P_{F_{\mathrm{c}}})$ is a lower descent interval.

Proof. (a) The first equality was established in [12, Theorem 5.3], so we will only prove the second. By Lemma 4.3, we have that $P_{F_D^{\downarrow}} = P_{(F_D^{\downarrow})^{t}}$. Applying $(F_D^{\downarrow})^{t} = (\overline{F_D^{\downarrow}})^{t}$ and $P_{\overline{F_D^{\downarrow}}} = \overline{P}_{F_D^{\downarrow}}$, to this equality yields that $P_{F_D^{\downarrow}} = \overline{P}_{F_D^{\downarrow}}$. On the other hand, from the first equality, it follows that

$$\Sigma_L(\overline{P}_{F_{D^{t}}}) = \Sigma_L(P_{F_{D^{t}}})w_0 = [\mathbf{w}_{\mathrm{LT}}(T_{D^{t}})w_0, \mathbf{w}_{\mathrm{LT}}(T'_{D^{t}})w_0]_L.$$

Observe that $w_{LT}(T_{D^t})w_0 = w_{TL}(T'_D)$ and $w_{LT}(T'_{D^t})w_0 = w_{TL}(T_D)$. Therefore,

$$\Sigma_L(P_{F_D^{\downarrow}}) = [\mathbf{w}_{\mathrm{TL}}(T'_D), \mathbf{w}_{\mathrm{TL}}(T_D)]_L.$$

(b) Suppose that k is the number of rows in D. For $1 \le i \le k-1$, let $\overline{r}_i := r_{i+1} + r_{i+2} + \cdots + r_k$. By the definitions of w_{LT} and T_D , we can express $w_{LT}(T_D)$ as follows:

$$w_{\rm LT}(T_D) = \underbrace{\overline{\mathbf{r}}_1 + 1 \ \overline{\mathbf{r}}_1 + 2 \ \cdots \ n}_{\text{1st row}} \underbrace{\overline{\mathbf{r}}_2 + 1 \ \overline{\mathbf{r}}_2 + 2 \ \cdots \ \overline{\mathbf{r}}_1}_{\text{2nd row}} \cdots \underbrace{1 \ 2 \ \cdots \ r_k}_{k \text{th row}}$$

Here, the notation "*i*th row" refers to the *i*th row of T_D when counted from the top. By definition, we have that $w_{LT}(T_D) = w_1(\text{set}(\mathsf{r}(D)))$. Thus, $\Sigma_L(P_{F_D})$ is an upper descent interval.

(c) This assertion can be proven in a manner similar to (b).

Next, we show that every lower descent interval is of the form $\Sigma_L(P_{F_D^{\downarrow}})$, and every upper descent interval is of the form $\Sigma_L(P_{F_D^{\rightarrow}})$. We begin by reviewing the result in [12, Section 5.1]. For $P \in \mathsf{P}_n$, let $\Sigma_R(P) := \{\sigma^{-1} \mid \sigma \in \Sigma_L(P)\}$.

Lemma 4.5. ([12, Theorem 5.6]) Given $\alpha \models n$ and $\rho \in \mathfrak{S}_n$ with $w_0(\operatorname{set}(\alpha)) \preceq_L \rho$, the diagram $E_{\alpha;\rho} \in \mathfrak{D}_n^{-2}$, constructed via [12, Algorithm 5.4], satisfies


$$\Sigma_R(P_{F_{E_{\alpha;\rho}}}) = \Xi([w_0(\alpha), \rho]_L),$$

where $\boldsymbol{\Xi}: \mathfrak{S}_n \to \mathfrak{S}_n$ is the bijection defined by $\boldsymbol{\Xi}(\gamma) = w_0 \gamma^{-1}$.

In Lemma 4.5, we let

$$(4.3) D_{\sigma;S} := E_{\operatorname{comp}(S)^{\mathsf{t}};\sigma w_0}.$$

Example 4.6. (cf. [12, Example 5.5]) Let $\sigma = 267935148$ and $S = \{4, 6\}$. Then comp $(S)^{t} = (1, 1, 2, 2, 1, 1, 1) \models 9$ and $\sigma w_{0} = 841539762$. For simplicity, let $\alpha := \text{comp}(S)^{t}$ and $\rho := \sigma w_{0}$. Then $D_{\sigma;S} = E_{\alpha;\rho}$. Since $w_{0}(\text{set}(\alpha)) = 321549876$ and $w_{0}(\alpha) \preceq_{L} \rho$, we can proceed with the construction of the diagram $E_{\alpha;\rho}$ using [12, Theorem 5.6]:



In what follows, we present an algorithm that, given $S \subseteq [n-1]$ and $\rho \in \mathfrak{S}_n$ satisfying $w_0(S) \preceq_L \rho$, produces a diagram $D \in \mathfrak{D}_n$. This diagram constitutes a slight modification of the algorithm described in [12, Algorithm 5.4].

Algorithm 4.7. Let $S \subseteq [n-1]$ and $w_0(S) \preceq_L \rho$. Step 1. Let $S^c = \{z_1, z_2, ..., z_p\}$, and $z_0 := 0$ and $z_{p+1} := n$. For $1 \le i \le p+1$, let

$$X_i(S;\rho) := \{\rho(r) \mid z_{i-1} + 1 \le r \le z_i\}$$

Step 2. Let $e := |\text{Des}_{L}(\rho)|$ and let $\text{Des}_{L}(\rho) = \{q_1 < q_2 < \cdots < q_e\}, q_0 := 0, \text{ and } q_{e+1} := n.$ Then, for $j = 1, 2, \dots, e+1$, set

$$Y_j(\rho) := \{ c \mid q_{j-1} + 1 \le c \le q_j \}$$

²It should be remarked that in [12, Theorem 5.6], the notation $D_{\alpha;\rho}$ is employed in place of $E_{\alpha;\rho}$.

Step 3. Let

$$D_{S;\rho} := \{(i,j) \mid X_i(S;\rho) \cap Y_j(\rho) \neq \emptyset\}.$$

Return $D_{S;\rho}$.

Example 4.8. Let $S = \{2, 5\} \subseteq 6$, and consider $\rho = 231564$. We apply Algorithm 4.7 to obtain the diagram $D_{S;\rho}$. The observation that $w_0(S) = 132465$ and $w_0(S) \preceq_L \rho$ gives us to proceed with the construction of $D_{S;\rho}$. Since $S^c = \{1, 3, 4\}$, Step 1 gives

$$X_1(S;\rho) = \{2\}, \quad X_2(S;\rho) = \{1,3\} \quad X_3(S;\rho) = \{5\} \text{ and } X_4(S;\rho) = \{4,6\}$$

With $\text{Des}_{L}(\rho) = \{1, 4\}, Step \ 2 \text{ gives}$

$$Y_1(\rho) = \{1\}, \quad Y_2(\rho) = \{2, 3, 4\}, \text{ and } Y_3(\rho) = \{5, 6\}.$$

Finally, Step 3 gives

$$D_{S;\rho} = \{(1,2), (2,1), (2,2), (3,3), (4,2), (4,3)\}.$$

The diagrams $D_{S;\rho}$ is represented graphically as follows:



Proposition 4.9. Let $S \subseteq [n-1]$.

(a) Let $\sigma \in \mathfrak{S}_n$ satisfy $w_1(S) \preceq_L \sigma$. The diagram $D_{\sigma;S}$, constructed using [12, Algorithm 5.4], satisfies

$$\Sigma_L(P_{F_{D_{\sigma,S}}}) = [\sigma, w_1(S)]_L.$$

(b) Let $\rho \in \mathfrak{S}_n$ satisfy $w_0(S) \preceq_L^{\circ,\circ} \rho$. The diagram $D_{S;\rho}$, constructed using Algorithm 4.7, satisfies

$$\Sigma_L(P_{F_{D_{S;\rho}}^{\downarrow}}) = [w_0(S), \rho]_L$$

Proof. (a) By Lemma 4.5, it follows that $\Sigma_L(P_{F_{E_{\alpha;\rho}}}) = [\rho w_0, w_0(\operatorname{set}(\alpha))w_0]_L$. Let $\rho w_0 = \sigma$ and $w_0(\operatorname{set}(\alpha))w_0 = w_1(S)$. Then, $\rho = \sigma w_0$ and

$$w_0(\operatorname{set}(\alpha)) = w_1(S)w_0 = w_0(S^{\operatorname{t}}).$$

Now, the assertion follows from the definition that $D_{\sigma;S} = E_{\text{comp}(S)^{t};\sigma w_{0}}$.

(b) The assertion can be proven using the same approach as in the proof of [12, Theorem 5.4], and thus we omit the proof. \Box

By combining Proposition 4.9 with Proposition 4.4(b) and (c), we obtain the equalities

$$\left\{ \Sigma_L(P_{F_D^{\perp}}) \mid D \in \mathfrak{D}_n \right\} = \left\{ [\sigma, w_1(S)]_L \mid S \subseteq [n-1], \sigma \preceq_L w_1(S) \right\},\\ \left\{ \Sigma_L(P_{F_D^{\perp}}) \mid D \in \mathfrak{D}_n \right\} = \left\{ [w_0(S), \rho]_L \mid S \subseteq [n-1], w_0(S) \preceq_L \rho \right\}.$$

4.2. The minimal and maximal elements of the equivalence class. We begin by recalling an essential structural theorem concerning an arbitrary equivalence class in $(\operatorname{Int}(n), \stackrel{D}{\simeq})$. Let *C* be an equivalence class in $(\operatorname{Int}(n), \stackrel{D}{\simeq})$. Define $\xi_C := \rho \sigma^{-1}$ for any $[\sigma, \rho]_L \in C$. By [18, Proposition 4.1], the permutation ξ_C is independent of the choice of representative $[\sigma, \rho]_L$ in *C*. Define a partial order \preceq on *C* by setting

$$[\sigma, \rho]_L \preceq [\sigma', \rho']_L$$
 if and only if $\sigma \preceq_R \sigma'$.

As shown in [18, Theorem 4.6], the set $\{\sigma \mid [\sigma, \rho]_L \in C\}$ forms a right weak Bruhat interval. Let σ_0 and σ_1 denote the minimal and maximal elements of this interval, respectively. By definition,

$$(C, \preceq) \cong [\sigma_0, \sigma_1]_R$$
 as posets.

The minimal and maximal elements of (C, \preceq) are denoted by min C and max C, respectively, so that min $C = [\sigma_0, \xi_C \sigma_0]_L$ and max $C = [\sigma_1, \xi_C \sigma_1]_L$.

In general, explicitly describing arbitrary equivalence classes remains an open problem. However, this issue was successfully addressed in [18] for the equivalence classes of $\Sigma_L(P)$, where P is a regular Schur-labeled poset on [n]. Here, when C is the equivalence class of a lower or upper descent interval, we provide the minimal and maximal elements of C.

We construct a filling F_D^{\nearrow} , derived from F_D^{\downarrow} . Consider the sequence of fillings

(4.4)
$$Z_0 = F_D^{\downarrow}, \ Z_1, \ Z_2, \dots, \ Z_n,$$

where each Z_i is obtained from Z_{i-1} through the following process.

Case 1: If there exists an entry x > i in Z_{i-1} satisfying the conditions:

- (i) x is positioned strictly above and to the right of i, and
- (ii) for each j = i, i + 1, ..., x 1, the entry j is positioned strictly below and weakly to the left of x, but not in the covering relation in $P_{Z_{i-1}}$,

then select the uppermost entry among such x's, denoted by **x**. Construct Z_i from Z_{i-1} by incrementing the entries $i, i + 1, ..., \mathbf{x} - 1$ by 1, and then swapping the original entry **x** with i.

Case 2: If no such x exists, set $Z_i := Z_{i-1}$.

It is clear that this process terminates in a finite number of steps. We define F_D^{\wedge} to be the final filling Z_n .

Example 4.10. Consider the case where

$$D = \square \qquad \text{and} \quad F_D^{\downarrow} = \square \boxed{\begin{array}{c}4 & 5\\1 & 2\\3\end{array}}$$

Then the sequence of fillings will be as follows:

$$Z_0 = Z_1 = Z_2 = F_D^{\downarrow}, \ Z_3 = \begin{array}{c} 3 & 5 \\ 1 & 2 \\ 4 \end{array}, \ Z_4 = \begin{array}{c} 3 & 4 \\ 1 & 2 \\ 5 \end{array}, \ Z_5 = \begin{array}{c} 3 & 4 \\ 1 & 2 \\ 5 \end{array}$$

As a step-by-step analysis, for i = 1 and i = 2, there is no x in Z_{i-1} satisfying conditions (i) and (ii), so $Z_1 = Z_2 = F_D^{\downarrow}$. For i = 3, the entry x = 4 satisfies the conditions, resulting in Z_3 . For i = 4, the entry x = 5 satisfies the conditions, producing Z_4 . For i = 5, the entry x = 6 satisfies the conditions, yielding Z_5 . Thus, the final filling is $F_D^{\nearrow} = Z_5$.

The filling F_D^{\nearrow} is evidently a standard filling on D. The following lemma pertains to the structure of F_D^{\nearrow} .

Lemma 4.11. Let $D \in \mathfrak{D}_n$.

- (a) The posets P_{FD} and P_{FD} are equal as edge-decorated posets.
 (b) If i ≤_{P_{FD}} i + 1, then (i, i + 1) is in the covering relation in P_{FD}.

Proof. (a) Let $\{Z_j\}_{0 \le j \le n}$ be the sequence generating F_D^{\nearrow} . We claim that for each $1 \le i \le n$, the posets $P_{Z_{i-1}}$ and P_{Z_i} maintain the same edge-decorated structure (for the definition of Z_i , see (4.4)). If **Case 1** is applied to Z_{i-1} , then the entries $i, i+1, \ldots, \mathbf{x}-1$ in Z_{i-1} are incremented by 1 based on conditions (i) and (ii). This increment does not alter their relative order. Additionally, the original entry \mathbf{x} is replaced by *i*. Since there is an entry greater than x positioned between x and x - 1 in $P_{Z_{i-1}}$, as ensured by conditions (i) and (ii), the relative order among these entries also remains unchanged. Thus, the claim holds in this case. If **Case 2** is applied, then $Z_i = Z_{i-1}$, so that the relative order and edge decorations clearly remain unchanged. Hence, the claim also holds.

For all $1 \leq i \leq n$, the posets $P_{Z_{i-1}}$ and P_{Z_i} retain the same edge-decorated structure. As a result, the final poset $P_{F_D^{\checkmark}}$ and the initial poset $P_{F_D^{\downarrow}}$ are identical as edge-decorated posets.

(b) Assume, for the sake of contradiction, that there exists a pair (i, i + 1) in $P_{F_D^{\nearrow}}$ such that $i \leq_{P_{F_D^{\nearrow}}} i + 1$, but (i, i + 1) is not in the covering relation in $P_{F_D^{\nearrow}}$. This implies that i+1 is strictly above and to the right of i in F_D^{\nearrow} , but there exists an entry $k \neq i, i+1$ within the smallest rectangle-shaped subdiagram containing the boxes for i and i + 1.

Consider the sequence $\{Z_j\}_{0 \le j \le n}$ generating F_D^{\nearrow} . By the construction of Z_j from Z_{j-1} , we know that the entries $1, 2, \ldots, i+1$ retain their position in $Z_{i+2}, Z_{i+3}, \ldots, Z_n$. Consequently, we have that

$$Z_i^{-1}(i) = (F_D^{\nearrow})^{-1}(i)$$
 and $Z_i^{-1}(i+1) = (F_D^{\nearrow})^{-1}(i+1).$

In this situation, the entry i + 1 satisfies conditions (i) and (ii). As a result, the entry i is placed in $(Z_i)^{-1}(i+1)$, equivalently i+1 is placed in $(Z_i)^{-1}(i)$. By the above argument, we have that

 $Z_i^{-1}(i) = (F_D^{\nearrow})^{-1}(i+1)$ and $Z_i^{-1}(i+1) = (F_D^{\nearrow})^{-1}(i)$,

that is, $i+1 \preceq_{P_{F_D^{\nearrow}}} i$. This contradicts the assumption, and therefore, no such pair (i, i+1)can exist in $P_{F_D^{\nearrow}}$ such that $i \preceq_{P_{F_D^{\nearrow}}} i+1$ without (i, i+1) being in the covering relation. \Box Given a diagram $D \in \mathfrak{D}_n$, let D^* be the diagram obtained by reflecting D across the line y = -x and shifting it appropriately to ensure it remains in \mathcal{D}_n . Explicitly, if D has r rows and c columns, then

$$D^* = \{ (r - j + 1, c - i + 1) \mid (i, j) \in D \}.$$

For a standard filling F on D, define F^* as the standard filling on D^* , given by

$$F^*(i,j) = F(r-j+1, c-i+1),$$

where r and c are the number of rows and columns of D, respectively. Using this notation, we define the filling $F_{D_{\sigma},s}^{\checkmark}$ as

(4.5)
$$F_{D_{\sigma;S}}^{\checkmark} := \left(F_{D_{S^{c};w_{0}\sigma}}^{\nearrow}\right)^{*}.$$

With these definitions in place, we are ready to state the main result of this subsection.

Theorem 4.12. Let S be a subset of [n-1]. Suppose that $\rho, \sigma \in \mathfrak{S}_n$ with $w_0(S) \preceq_L \rho$ and $\sigma \preceq_L w_1(S)$. Denote by $C_{S;\rho}$ and $C_{\sigma;S}$ the classes of the intervals $[w_0(S), \rho]_L$ and $[\sigma, w_1(S)]_L$, respectively. Then, the following results hold:

- (a) min $C_{S;\rho} = [w_0(S), \rho]_L$ and max $C_{S;\rho} = \Sigma_L(P_{F_{D_{S;\rho}}^{,\lambda}})$.
- (b) min $C_{\sigma;S} = \Sigma_L(P_{F_{D_{\sigma;S}}^{\swarrow}})$ and max $C_{\sigma;S} = [\sigma, w_1(S)]_L$.

Proof. (a) According to Proposition 4.9, $\Sigma_L(P_{F_{D_{S;\rho}}^{\downarrow}}) = [w_0(S), \rho]_L$. To establish the first equality, it suffices to show that min $C_{S;\rho} = \Sigma_L(P_{F_{D_{S;\rho}}^{\downarrow}})$. By Theorem 3.8 along with [18, Theorem 4.6], it is equivalent to verifying that $P_{F_{D_{S;\rho}}^{\downarrow}}$ contains no pair (i, i + 1) satisfying the following conditions:

- (i, i+1) is a comparable pair that is not in the covering relation, and
- i + 1 is less than i in the partial order of $P_{F_{D_{S;o}}^{\downarrow}}$.

This property follows directly from Definition 4.1.

Similarly, we can derive the second equality using Lemma 4.11.

(b) Since $w_0[\sigma, w_1(S)]_L = [w_0(S^c), w_0\sigma]_L$, it follows that

$$I \in C_{\sigma;S} \iff w_0 I \in C_{S^c;w_0\sigma}$$

and consequently,

 $I \stackrel{D}{\simeq} Is_i \text{ for } I \in C_{\sigma;S} \iff w_0 I \stackrel{D}{\simeq} w_0 Is_i \text{ for } w_0 I \in C_{S^c;w_0\sigma}.$

Here, we use the notation $\zeta U := \{\zeta \sigma \mid \sigma \in U\}$ for $U \subseteq \mathfrak{S}_n$ and $\zeta \in \mathfrak{S}_n$. This equivalence establishes that the map

$$C_{\sigma;S} \to C_{S^c;w_0\sigma}, \ I \mapsto w_0 I$$

is a bijection. Moreover, for $F \in \mathfrak{D}_n$, it holds that $w_0 \Sigma_L(P_F) = \Sigma_L(P_{F^*})$. By applying these two properties to $F_{D_{\sigma,S}}$, the desired result follows from (a) and (4.5).

Remark 4.13. For any $[\sigma, \rho]_L \in C_{S;\rho}$, we have $\text{Des}_L(\sigma) = \text{Des}_L(w_0(S)) = S$. This implies that σ cannot be of the form $w_0(J)$ for any $J \subset [n-1]$ other than S. Therefore, the class $C_{S;\rho}$ has a unique lower descent interval $[w_0(S), \rho]_L$. Similarly, for the same reason, $C_{\sigma;S}$ has a unique upper descent interval $[\sigma, w_1(S)]_L$.

Example 4.14. (a) Let n = 6, $S = \{2, 5\} \subseteq [5]$, and $\rho = 231564$. Then $w_0(S) \preceq_L \rho$. From Theorem 4.12(a) it follows that $\min C_{S;\rho} = [w_0(S), \rho]_L$ and $\max C_{S;\rho} = \Sigma_L(P_{F_{D_{S;\rho}}})$. It can be easily seen that


$$F_{D_{S;\rho}}^{\nearrow} = \begin{array}{c} 3 \\ 1 \\ 6 \end{array}$$

Now, by Remark 4.2, we obtain that $\min C_{S;\rho} = [132465, 231564]_L$ and $\max C_{S;\rho} = [134652, 235641]_L$.

Let us investigate $C_{S;\rho}$ in more detail. One can observe that P_{Z_j} $(1 \le j \le 4)$ are all the posets obtained from $F_{D_{S;\rho}}^{\downarrow}$ by applying label changes that satisfy the condition in Theorem 3.8, where


$$Z_1 := F_{D_{S;\rho}}^{\downarrow} = \begin{array}{c} 4 \\ 1 \\ 3 \\ 3 \end{array}, \quad Z_2 = \begin{array}{c} 3 \\ 1 \\ 4 \\ 4 \\ 4 \end{array}, \quad Z_3 = \begin{array}{c} 3 \\ 1 \\ 5 \\ 5 \\ 5 \\ 5 \\ 5 \end{array}, \text{ and } Z_4 := F_{D_{S;\rho}}^{\checkmark}.$$

Let $I_j := \Sigma_L(P_{Z_j})$ for $1 \le j \le 4$. By Theorem 3.8, $C_{S;\rho} = \{I_1, I_2, I_3, I_4\}$, and its poset structure is given as follows:



In this figure, the down arrow $\downarrow \cdot s_i$ on the left denotes the labeling change $i \leftrightarrow i+1$, and the down arrow $\downarrow \cdot s_i$ on the right denotes the right multiplication by s_i .

(b) Let n = 6, $S = \{1, 3, 4\} \subseteq [5]$, and $\sigma = 546213$. Since $\sigma \preceq_L w_1(S)$, it follows from [12, Algorithm 5.4] and (4.3) that we have



So, Theorem 4.12(b) yields that $\min C_{\sigma;S} = \Sigma_L(P_{F_{D_{\sigma;S}}^{\checkmark}})$ and $\max C_{\sigma;S} = [\sigma, w_1(S)]_L$. The fillings $F_{D_{\sigma;S}}^{\checkmark}$ and $F_{D_{\sigma;S}}^{\rightarrow}$ are given by

$$F_{D_{\sigma;S}}^{\checkmark} = \begin{array}{c} 1 \\ 2 & 6 \\ \hline 3 \\ 4 & 5 \end{array} \quad \text{and} \quad F_{D_{\sigma;S}}^{\rightarrow} = \begin{array}{c} 1 \\ 2 & 3 \\ \hline 4 \\ 5 & 6 \end{array}$$

From Remark 4.2, we compute

min $C_{\sigma;S} = [542136, 643125]_L$ and max $C_{\sigma;S} = [546213, 645312]_L$.

By comparing this result with (a), we observe the structure and relationships between the interval classes $C_{\sigma;S}$ and $C_{S^c;w_0\sigma}$ through their respective fillings and the poset representations of their elements.

For a complete understanding of the class C, it is essential to know not only min C and max C, but also the poset structure of (C, \preceq) . In the remainder of this subsection, we focus on lower and upper descent intervals that satisfy a specific condition. These intervals are particularly relevant to the 0-Hecke modules discussed in the next section.

A diagram $D \in \mathcal{D}_n$ contains a *strictly upper-right pair* if there are two boxes (x_1, y_1) and (x_2, y_2) in D satisfying:

(i) $x_1 < x_2$ and $y_1 < y_2$, and

(ii) no other boxes of D lie inside the smallest rectangle enclosing (x_1, y_1) and (x_2, y_2) . A diagram $D \in \mathcal{D}_n$ is free of a strictly upper-right configuration if it has no strictly upperright pairs. Pictorially, D is a diagram that does not contain a subdiagram of the following form (x_2, y_2)

(4.6)
$$(x_1, y_1)$$

For the sake of simplicity in notation, let D^x (or $(D)^x$) denote the diagram obtained by reflecting D across the x-axis and shifting it as needed, ensuring that it remains an element of \mathcal{D}_n . Similarly, for a filling F, let F^x be the filling obtained by reflecting F across the x-axis and shifting it upwards appropriately. To define a partial order on ST(D), we set $T \leq U$ if and only if $w_{TL}(T) \preceq_L w_{TL}(U)$. For $\zeta \in \mathfrak{S}_n$, let $\zeta \cdot T$ denote the tableau obtained from T by replacing each entry i with $\zeta(i)$ for $1 \leq i \leq n$. With these definitions, the following theorem can be stated.

Lemma 4.15. Let $D \in \mathfrak{D}_n$ and $T \in \mathsf{ST}(D^x)$.

(a) $T = s_{i_r} \cdots s_{i_2} s_{i_1} \cdot T'_{D^x}$ for some nonnegative integer r, where i_k is strictly upper-left of i + 1 in $s_{i_{k-1}} \cdots s_{i_2} s_{i_1} \cdot T'_{D_x}$ for $1 \le k \le r$. For the definition of T'_{D^x} , refer to (4.2).

(b) Suppose that D is free of a strictly upper-right configuration. Then, for $1 \le i \le n-1$, i is strictly upper-left of i+1 in T if and only if (i, i+1) is a comparable pair but not a covering relation in P_{T^x} .

Proof. (a) We aim to prove that $T = s_{i_r} \cdots s_{i_1} \cdot T'_{D^x}$, where each i_k satisfies the conditions stated in this lemma. To this end, we define an $H_n(0)$ -action on $\mathbb{C}\mathsf{ST}(D^x)$ as follows: For $1 \le i \le n-1$ and $T \in \mathsf{ST}(\mathcal{D}^x)$,

(4.7)
$$\pi_i \star T := \begin{cases} s_i \cdot T & \text{if } i \text{ is strictly upper-left of } i+1 \text{ in } T, \\ 0 & \text{if } i \text{ is lower-left of } i+1 \text{ in } T, \\ T & \text{otherwise.} \end{cases}$$

Note that $s_i \cdot T \in \mathsf{ST}(\mathcal{D}^x)$ whenever *i* is strictly upper-left of i + 1 in *T*. We need to check that the operators π_i satisfy the defining relations of the 0-Hecke algebra (see (4.7)).

We now verify that the operators π_i satisfy the defining relations of the 0-Hecke algebra:

- (i) $\pi_i^2 = \pi_i$: If *i* is strictly upper-left of i + 1 in *T*, then $\pi_i \star T = s_i \cdot T$. Since *i* is strictly lower-right of i + 1 in $s_i \cdot T$, it follows that $\pi_i \star (s_i \cdot T) = s_i \cdot T$. In all other cases, $\pi_i \star T = 0$ or $\pi_i \star T = T$, it is clear that $\pi_i^2 \star T = \pi_i \star T$.
- (ii) $\pi_i \pi_j = \pi_j \pi_i$ for |i j| > 1: If |i j| > 1, the sets $\{i, i + 1\}$ and $\{j, j + 1\}$ are disjoint, so the actions of π_i and π_j commute.
- (iii) $\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$: Consider the relative positions of i, i+1, i+2 in T:
 - i is strictly upper-left of i + 1.
 - If i + 2 is strictly lower-right of i + 1, then $\pi_i \pi_{i+1} \pi_i \star T = s_i s_{i+1} s_i \cdot T$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = s_{i+1} s_i s_{i+1} \cdot T$.
 - If i + 2 is strictly above and weakly right of i + 1, then $\pi_i \pi_{i+1} \pi_i \star T = \pi \pi_{i+1} \star (s_i \cdot T) = 0$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = 0$.
 - If i + 2 is strictly upper-left of i + 1 and strictly lower-right of i, then $\pi_i \pi_{i+1} \pi_i \star T = \pi \pi_{i+1} \star (s_i \cdot T) = \pi \star (s_{i+1} s_i \cdot T) = s_{i+1} s_i \cdot T$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star T = \pi_{i+1} \pi_i \star T = \pi_{i+1} \star (s_i \cdot T) = s_{i+1} s_i \cdot T$.
 - If i + 2 is upper-right of i, then $\pi_i \pi_{i+1} \pi_i \star T = \pi \pi_{i+1} \star (s_i \cdot T) = 0$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star T = \pi_{i+1} \star (s_i \cdot T) = 0.$
 - If i + 2 is strictly upper-left of i, then $\pi_i \pi_{i+1} \pi_i \star T = \pi \pi_{i+1} \star (s_i \cdot T) = \pi_i \star (s_i \cdot T) = s_i \cdot T$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star T = \pi_{i+1} \star (s_i \cdot T) = s_i \cdot T$.
 - *i* is lower-left of i + 1. In this case $\pi_i \pi_{i+1} \pi_i \star T = 0$.
 - If i + 2 is strictly lower-right of i, then i + 2 is strictly lower-right of i + 1. So $\pi_{i+1}\pi_i\pi_{i+1} \star T = \pi_{i+1}\pi_i \star (s_{i+1} \cdot T) = \pi_{i+1} \star (s_i s_{i+1} \cdot T) = 0$.
 - If i+2 is upper-right of i and strictly lower-right of i+1, then $\pi_{i+1}\pi_i\pi_{i+1}\star T = \pi_{i+1}\pi_i\star(s_{i+1}\cdot T) = 0$.
 - If i + 2 is upper-right of i + 1, then $\pi_{i+1}\pi_i\pi_{i+1} \star T = 0$.
 - If i + 2 is strictly upper-left of i + 1, then $\pi_{i+1}\pi_i\pi_{i+1} \star T = \pi_{i+1}\pi_i \star T = 0$.

- i is strictly lower-right of i + 1.
- If i+2 is strictly lower-right of i, then $\pi_i \pi_{i+1} \pi_i \star T = \pi_i \pi_{i+1} \star T = \pi_i \star (s_{i+1} \cdot T)$ $T) = s_i s_{i+1} \cdot T$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star (s_{i+1} \cdot T) = \pi_{i+1} \star (s_i s_{i+1} \cdot T) = s_i s_{i+1} \cdot T$.
- If i+2 is upper-right of i and strictly lower-right of i+1, then $\pi_i \pi_{i+1} \pi_i \star T = \pi_i \pi_{i+1} \star T = \pi_i \star (s_{i+1} \star T) = 0$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star (s_{i+1} \star T) = 0$.
- If i + 2 is strictly above and weakly left of i and strictly lower-right of i + 1, then $\pi_i \pi_{i+1} \pi_i \star T = \pi_i \pi_{i+1} \star T = \pi_i \star (s_{i+1} \star T) = s_{i+1} \cdot T$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = \pi_{i+1} \pi_i \star (s_{i+1} \cdot T) = \pi_{i+1} \star (s_{i+1} \cdot T) = s_{i+1} \cdot T$.
- If i + 2 is upper-right of i + 1, then $\pi_i \pi_{i+1} \pi_i \star T = \pi_i \pi_{i+1} \star T = 0$ and $\pi_{i+1} \pi_i \pi_{i+1} \star T = 0$.

- If i+2 is strictly upper-left of i+1, then $\pi_i \pi_{i+1} \pi_i \star T = T = \pi_{i+1} \pi_i \pi_{i+1} \star T$. In all configurations, we verify that $\pi_i \pi_{i+1} \pi_i \star T = \pi_{i+1} \pi_i \pi_{i+1} \star T$ holds.

These verifications confirm that the operators π_i satisfy the 0-Hecke algebra relations. Let M_{D^x} denote the resulting module. In $\mathsf{ST}(D^x)$, each tableau, except the tableau T'_{D^x} , contains a pair (i, i + 1) such that i is strictly lower-right of i + 1. This implies that M_D is generated by T'_D , which guarantees that every $T \in \mathsf{ST}(D^x)$ can be expressed as $T = s_{i_r} \cdots s_{i_2} s_{i_1} \cdot T'_{D^x}$, which each i_k satisfies the conditions stated in (a).

(b) We first address the "if" direction. By the given assumption, it follows from Definition 4.1 that i is positioned to the lower-left of i + 1 in T^x . Equivalently, i is positioned to the upper-left of i + 1 in T. Since T is a standard tableau, it is impossible for i and i + 1 to occupy the same column in T. Furthermore, as the shape of T^x is free of a strictly upper-right configuration, i and i + 1 cannot occupy the same row in T. Thus, i is strictly above and strictly to the left of i + 1 in T.

Next, we consider the "only if" direction. Under the assumption, i + 1 is strictly above and strictly to the right of i in T^x , implying that $i \leq_{T^x} i + 1$. Since the shape of T^x is free of a strictly upper-right configuration, there exists a box within the smallest rectangular subdiagram containing the boxes with i and i + 1. This ensures that (i, i + 1) forms a comparable pair that is not in the covering relation in P_{T^x} .

Theorem 4.16. Under the same hypothesis as in Theorem 4.12, we have the following.

(a) If $\mathcal{D} := D_{S;\rho}$ is free of a strictly upper-right configuration, then

$$\max C_{S;\rho} = \Sigma_L(P_{F_{\mathcal{D}}}) = [\mathbf{w}_{\mathrm{LT}}(T'_{\mathcal{D}}), w_1(\operatorname{set}(\mathbf{r}(\mathcal{D})))]_L.$$

Furthermore, $C_{S;\rho} = \{\Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{D}^x)\}, and (C_{S;\rho}, \preceq) \cong (\mathsf{ST}(\mathcal{D}^x), \leq) as$ posets.

(b) If $\mathcal{E} := D_{\sigma;S}$ is free of a strictly upper-right configuration, then

$$\min C_{\sigma;S} = \Sigma_L(P_{F_c^{\downarrow}}) = [w_0(\operatorname{set}(\mathsf{c}(D))^c), \operatorname{w}_{\operatorname{TL}}(T_{\mathcal{D}})]_L.$$

Furthermore, $C_{\sigma;S} = \{\Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{E}^x)\}$ and $(C_{\sigma;S}, \preceq) \cong (\mathsf{ST}(\mathcal{E}^x), \leq)$ as posets.

Proof. Since the proof for (b) is similar to that for (a), we provide the proof for (a) only.

Assume that the diagram \mathcal{D} is free of a strictly upper-right configuration. This assumption guarantees that during the construction of $F_{\mathcal{D}}^{\nearrow}$, the entry **x** chosen in **Case 1** is always the leftmost entry in the uppermost row among all entries greater than i - 1 in F_{i-1} . This ensures that $F_{\mathcal{D}}^{\nearrow} = F_{\mathcal{D}}^{\rightarrow}$. Based on Theorem 4.12(a), this leads to

$$\max C_{S;\rho} = \Sigma_L(P_{F_{\mathcal{D}}}).$$

Furthermore, by Proposition 4.4(a) and (b), we have

$$\Sigma_L(P_{F_{\mathcal{D}}}) = [\mathbf{w}_{\mathrm{LT}}(T'_{\mathcal{D}}), w_1(\operatorname{set}(\mathbf{r}(\mathcal{D})))]_L.$$

Combining these two equalities yields the first assertion.

Next, we prove the second assertion. Let us consider the map

$$\phi : \mathsf{ST}(\mathcal{D}^x) \to \operatorname{Int}(n), \quad T \mapsto \Sigma_L(P_{T^x}).$$

Claim 1. The diagram

$$\begin{array}{ccc} T & \stackrel{\phi}{\longrightarrow} & \Sigma_L(P_{T^x}) \\ s_i \cdot \downarrow & & \downarrow \cdot s_i \\ s_i \cdot T & \stackrel{\phi}{\longrightarrow} & \Sigma_L(P_{(s_i \cdot T)^x}) \end{array}$$

commutes. Here, *i* is strictly upper-left of i + 1 in *T*.

To verify this claim, consider the poset $P_{(s_i \cdot T)^x}$, which is obtained from P_{T^x} by swapping the labels *i* and *i* + 1. Since \mathcal{D} is free of a strictly upper-right configuration, we see that $P_{(s_i \cdot T)^x} \neq P_{T^x}$. By Lemma 3.5, if P_{T^x} is regular, then $P_{(s_i \cdot T)^x}$ is also regular. Since $P_{(s_i \cdot T)^x} = s_i \cdot P_{T^x}$, it follows from (3.2) that

$$\Sigma_L(P_{(s_i \cdot T)^x}) = \Sigma_L(s_i \cdot P_{T^x}) = \Sigma_L(P_{T^x})s_i.$$

Claim 2. The map ϕ is injective.

Assume that $\phi(T) = \phi(U)$ for $T, U \in \mathsf{ST}(\mathcal{D}^x)$. By Lemma 4.15(a), $T = \zeta \cdot T'_{\mathcal{D}^x}$ and $U = \zeta' \cdot T'_{\mathcal{D}^x}$ for some $\zeta, \zeta' \in \mathfrak{S}_n$. Hence,

$$\phi(T) = [w_0(S), \rho]_L \zeta^{-1} = [w_0(S), \rho]_L \zeta'^{-1} = \phi(U).$$

This implies that $\zeta = \zeta'$, and therefore T = U.

Claim 3. Im $(\phi) = C_{S;\rho}$, that is, $C_{S;\rho} = \{\Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{D}^x)\}.$

By Theorem 3.8, every interval in $C_{S;\rho}$ is of the form

$$[w_0(S),\rho]_L s_{i_1} s_{i_2} \cdots s_{i_r},$$

where $(i_k, i_k + 1)$ is a comparable pair that is not in the covering relation in $P_{[w_0(S),\rho]_L s_{i_1} s_{i_2} \cdots s_{i_{k-1}}}$ for $1 \leq k \leq r$. Observe that $\phi(T'_{\mathcal{D}^x}) = [w_0(S),\rho]_L \in C_{S;\rho}$. Let $T = s_{i_r} \cdots s_{i_2} s_{i_1} \cdot T'_{\mathcal{D}^x}$ for some nonnegative integer r, where i_k is strictly upper-left

of $i_k + 1$ in $s_{i_{k-1}} \cdots s_{i_2} s_{i_1} \cdot T'_{\mathcal{D}^x}$ for $1 \leq k \leq r$. Using *Claim 1* repeatedly, we derive that

$$\phi(T) = [w_0(S), \rho]_L s_{i_1} s_{i_2} \cdots s_{i_r}.$$

Now, the desired result follows from Lemma 4.15(b).

By Claim 2 and Claim 3, the map ϕ induces a bijection $\phi : \mathsf{ST}(\mathfrak{D}^x) \to C_{S;\rho}$. Let $T = s_{i_r} \cdots s_{i_2} s_{i_1} \cdot T'_{\mathfrak{D}^x}$ for some nonnegative integer r, where i_k is strictly upper-left of $i_k + 1$ in $s_{i_{k-1}} \cdots s_{i_2} s_{i_1} \cdot T'_{\mathfrak{D}^x}$ for $1 \leq k \leq r$. Similarly, let $U = s_{j_s} \cdots s_{j_2} s_{j_1} \cdot T'_{\mathfrak{D}^x}$ for some nonnegative integer s, where j_k is strictly upper-left of $j_k + 1$ in $s_{j_{k-1}} \cdots s_{j_2} s_{j_1} \cdot T'_{\mathfrak{D}^x}$ for $1 \leq k \leq s$. Suppose that $T \leq U$. Since $w_{\mathrm{TL}}(T'_{\mathfrak{D}^x}) = w_0(S)$, it follows from the definition of \leq that

$$s_{i_r}\cdots s_{i_2}s_{i_1}w_0(S) \preceq_L s_{j_s}\cdots s_{j_2}s_{j_1}w_0(S),$$

which implies that

(4.8)
$$w_0(S)s_{i_1}s_{i_2}\cdots s_{i_r} \preceq_R w_0(S)s_{j_1}s_{j_2}\cdots s_{j_s}$$

On the other hand, *Claim 1* says that

 $\phi(T) = [w_0(S), \rho]_L s_{i_1} s_{i_2} \cdots s_{i_r}$ and $\phi(U) = [w_0(S), \rho]_L s_{j_1} s_{j_2} \cdots s_{j_s}$.

Thus, (4.8) ensures that $\phi(T) \preceq_R \phi(U)$. Therefore, $\phi : (\mathsf{ST}(\mathcal{D}^x), \leq) \to (C_{S;\rho}, \preceq)$ is a poset isomorphism.

Example 4.17. Consider two subsets $S = \{2\}$ and $S' = \{3\}$ in [5], with permutations $\rho = 142563$ and $\sigma = 345126$ in \mathfrak{S}_6 . Observe that $w_0(S) = 132456 \leq_L \rho$ and $\sigma \leq_L w_1(S) = 456123$, indicating that both conditions hold. Following the construction outlined in Algorithm 4.7 and [12, Algorithm 5.4], we obtain the diagrams $D_{S;\rho}$ and $D_{\sigma;S}$ as the same diagram

$$\mathcal{D} := \{ (1,1), (2,1), (2,2), (3,2), (4,2), (5,1) \},$$

which can be visualized as \square . This implies that $\Sigma_L(P_{F_{\mathcal{D}}^{\downarrow}}) = [w_0(S), 142563]_L$ and $\Sigma_L(P_{F_{\mathcal{D}}^{\rightarrow}}) = [345126, w_1(S)]_L$, where the standard fillings $F_{\mathcal{D}}^{\downarrow}$ and $F_{\mathcal{D}}^{\rightarrow}$ are

$$F_{\mathcal{D}}^{\downarrow} = \begin{array}{c} 2 & 4 & 5 \\ \hline 1 & 3 & 6 \end{array} \quad \text{and} \quad F_{\mathcal{D}}^{\rightarrow} = \begin{array}{c} 1 & 2 & 3 \\ \hline 4 & 5 & 6 \end{array}$$


Note that \mathcal{D} is free of a strictly upper-right configuration. According to Theorem 4.16 we have that

$$\min C_{S;\rho} = \min C_{\sigma;S} = \Sigma_L(P_{F_{\mathcal{D}}^{\downarrow}}) \quad \text{and} \quad \max C_{S;\rho} = \max C_{\sigma;S} = \Sigma_L(P_{F_{\mathcal{D}}^{\rightarrow}}).$$

On the other hand, the elements of $\mathsf{ST}(\mathcal{D}^x)$ are

By comparing $C_{S;\rho}$ and $\mathsf{ST}(\mathfrak{D}^x)$ (see Figure 4.1), we conclude that

$$C_{S;\rho} = C_{\sigma;S'} = \{ \Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{D}^x) \}.$$



EQUIVALENCE CLASSES OF LOWER AND UPPER DESCENT WEAK BRUHAT INTERVALS 27

FIGURE 4.1. The elements of the class $C_{\{2\};142563}$ (or $C_{345126;\{3\}}$) and the set of standard fillings on \mathcal{D}

Remark 4.18. Suppose that $D_{S;\rho}$ is free of a strictly upper-right configuration. Then $C_{S;\rho}$ possesses a unique lower descent interval. This follows by applying Proposition 4.4 and Remark 4.13 to Theorem 4.16. Furthermore, the following identity can be derived:

$$(C_{S;\rho}, \preceq) \cong ([w_0(S), \mathbf{w}_{\mathrm{LT}}(T'_{D_{S'\circ}})]_R, \preceq_R)$$

by Theorem 4.12 and Theorem 4.16. However, the tableau descriptions in Theorem 4.16 provide additional combinatorial insight and are more convenient from this perspective. Similarly, if $D_{\sigma;S}$ is free of a strictly upper-right configuration, then $C_{\sigma;S}$ possesses a unique upper descent interval. Furthermore,

$$(C_{\sigma;S}, \preceq) \cong ([\mathbf{w}_{\mathrm{BL}}(T'_{D_{\sigma;S}}), w_1(S)]_R, \preceq_R).$$

It is quite interesting to observe that, in Theorem 4.16(a), $\max C_{S;\rho}$ is an upper descent interval, and in Theorem 4.16(b), $\min C_{\sigma;S}$ is a lower descent interval. In the following, we present a representation-theoretic interpretation of this observation. To do this, we introduce the notions of a projective cover and an injective hull of a finitely generated $H_n(0)$ -module.

Let A, B be finitely generated $H_n(0)$ -modules. A surjective $H_n(0)$ -module homomorphism $f: A \to B$ is called an *essential epimorphism* if an $H_n(0)$ -module homomorphism $g: X \to A$ is surjective whenever $f \circ g: X \to B$ is surjective. A *projective cover* of A is an essential epimorphism $f: P \to A$ with P projective, which always exists and is unique up

to isomorphism. Similarly, let M, N be finitely generated $H_n(0)$ -modules with $N \subsetneq M$. We say that M is an *essential extension* of N if $X \cap N \neq 0$ for all nonzero submodules X of M. An injective $H_n(0)$ -module homomorphism $\iota : M \to \mathbf{I}$ with \mathbf{I} injective is called an *injective hull* of M if \mathbf{I} is an essential extension of $\iota(M)$, which always exists and is unique up to isomorphism.

The 0-Hecke algebra is a Frobenius algebra, which implies that it is self-injective. Consequently, finitely generated projective and injective modules coincide (see [14, Proposition 4.1], [15, Proposition 4.1], and [5, Proposition 1.6.2]). Furthermore, as shown in [17, Section 3.2], each projective indecomposable module corresponds to a weak Bruhat interval, more precisely, for $\alpha \models n$,

$$\mathbf{P}_{\alpha} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^{\mathrm{c}}, w_1(\operatorname{set}(\alpha)^{\mathrm{c}}]_L) \text{ as } H_n(0)\text{-modules.}$$

With this preparation, we can derive the following result.

Corollary 4.19. Under the same hypothesis as in Theorem 4.12, we have the following. (a) If $\mathcal{D} := D_{S;o}$ is free of a strictly upper-right configuration, then

 $\mathsf{B}([w_0(\mathrm{Des}_{\mathrm{L}}(w_{\mathrm{BL}}(F_{\mathcal{D}}^{\rightarrow}))), w_1(\mathrm{set}(\mathsf{r}(\mathcal{D})))]_L)$

is an injective hull of $B([w_0(S), \rho]_L)$.

(b) If $\mathcal{E} := D_{\sigma;S}$ is free of a strictly upper-right configuration, then

 $\mathsf{B}([w_0(\operatorname{set}(\mathsf{c}(D))^{\mathrm{c}}), w_1(\operatorname{Des}_{\mathrm{L}}(\mathrm{w}_{\mathrm{LB}}(F_{\mathcal{E}}^{\downarrow})))]_L).$

is a projective cover of $\mathsf{B}([\sigma, w_1(S)]_L)$.

Proof. (a) Since \mathcal{D} is free of a strictly upper-right configuration, Theorem 4.16 implies that

 $\mathsf{B}([w_0(S),\rho]_L) \cong \mathsf{B}([w_{\mathrm{BL}}(F_{\mathcal{D}}^{\rightarrow})^{-1}, w_{\mathrm{LB}}(F_{\mathcal{D}}^{\rightarrow})^{-1}]_L) \quad \text{as } H_n(0)\text{-modules.}$

Define $I := \text{Des}_{L}(\mathbf{w}_{\text{BL}}(F_{\mathcal{D}}^{\rightarrow}))$ and $J := \text{Des}_{L}(\mathbf{w}_{\text{LB}}(F_{\mathcal{D}}^{\rightarrow}))$. By the definition of $F_{\mathcal{D}}^{\rightarrow}$, we observe that $I \subseteq J$ and $J = \text{set}(\mathbf{r}(\mathcal{D}))$. Since $\mathbf{w}_{\text{LB}}(F_{\mathcal{D}}^{\rightarrow})^{-1} = w_{1}(J)$, we have from [4, Theorem 4.6] that $\mathsf{B}([w_{0}(I), w_{1}(J)]_{L})$ is an injective hull of $\mathsf{B}([\mathbf{w}_{\text{BL}}(F_{\mathcal{D}}^{\rightarrow})^{-1}, \mathbf{w}_{\text{LB}}(F_{\mathcal{D}}^{\rightarrow})^{-1}]_{L})$. Finally, since the injective hull is preserved under $H_{n}(0)$ -module isomorphisms, the result follows.

(b) The proof for (b) follows in a similar manner to that of (a).

5. Lower and upper descent intervals from quotient modules and submodules of projective indecomposable $H_n(0)$ -modules

Let $S \subseteq S' \subseteq [n-1]$. In the context of the representation theory of 0-Hecke algebras, lower and upper descent intervals arise from appropriately selected quotient modules and submodules of the projective module with the basis $[w_0(S), w_1(S')]_L$:

$$\mathsf{B}([w_0(S), w_1(S')]_L) \cong \bigoplus_{\alpha} \mathbf{P}_{\alpha} \text{ as } H_n(0) \text{-modules},$$

where α ranges over compositions of n with $S \subseteq \operatorname{set}(\alpha)^c \subseteq S'$. Specifically, for each $\rho \in [w_0(S), w_1(S')]_L$, let M denote the submodule of $\mathsf{B}([w_0(S), w_1(S')]_L)$ generated by the set $[w_0(S), w_1(S')]_L \setminus [w_0(S), \rho]_L$, and let N represent the submodule generated by $[\rho, w_1(S')]_L$. Then, we obtain the following $H_n(0)$ -module isomorphisms:

$$\mathsf{B}([w_0(S), w_1(S')]_L)/M \cong \mathsf{B}([w_0(S), \rho]_L)$$

and

$$N \cong \mathsf{B}([\rho, w_1(S')]_L).$$

It should be noted that the latter modules were previously discussed in [14, Section 4.2].

In this section, the results presented in Section 4 are illustrated through an examination of lower descent intervals that arise from significant quotient modules and submodules of projective indecomposable $H_n(0)$ -modules.

5.1. Lower descent intervals from quotient modules of projective indecomposable $H_n(0)$ -modules. We begin by introducing the quotient modules under consideration. Let α be a composition of n.

- In [7], Berg-Bergeron-Saliola-Serrano-Zabrocki construct an indecomposable $H_n(0)$ module \mathcal{V}_{α} by defining a 0-Hecke action on the set $\mathsf{SIT}(\alpha)$ of standard immaculate tableaux of shape α . The image of this module under the quasisymmetric characteristic is the dual immaculate quasisymmetric function indexed by α .
- In [29], Searles constructs an indecomposable $H_n(0)$ -module X_{α} by defining a 0-Hecke action on the set $\mathsf{SET}(\alpha)$ of standard extended tableaux of shape α . The image of this module under the quasisymmetric characteristic is the extended Schur function indexed by α .
- In [11, Section 4.2], Choi–Kim–Nam–Oh construct an $H_n(0)$ -module $\widehat{\mathbf{S}}_{\alpha}$, which is generally not indecomposable, by defining a 0-Hecke action on the set $\mathsf{SYCT}(\alpha)$ of standard Young composition tableaux of shape α .³ The image of this module under the quasisymmetric characteristic is the Young quasisymmetric Schur function indexed by α . We here focus on the canonical submodule $\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}$ of $\widehat{\mathbf{S}}_{\alpha}$ defined in [11, page. 7767].

All definitions related to $SIT(\alpha)$, $SET(\alpha)$, $SYCT(\alpha)$, the fundamental quasisymmetric function F_{α} and the quasisymmetric characteristic ch can be found in [11, Section 2 and Section 4].

It was shown in [11, Corollary 4.6] that there exists a sequence of surjective $H_n(0)$ -homomorphisms among these modules, given by

(5.1)
$$\mathbf{P}_{\alpha} \longrightarrow \mathcal{V}_{\alpha} \longrightarrow X_{\alpha} \longrightarrow \widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \longrightarrow \mathbf{F}_{\alpha}.$$

³There are two remarks. First, a Young composition tableau of shape α is defined as a filling of $cd(\alpha)$. Second, although permuted Young composition tableaux are discussed in [11, Section 4.2], only the case $\sigma = id$ is considered in this work.

Jung-Kim-Lee-Oh [17] demonstrated that the intervals arising from these modules are lower descent intervals. In this paper, we show that these intervals satisfy the condition in Theorem 4.16, which allows us to provide an explicit description of (C, \preceq) .

Convention. For each \mathbf{Y}_{α} in Table 5.1 and Table 5.2, let $\mathcal{B}(\mathbf{Y}_{\alpha})$ denote the basis of tableaux associated with \mathbf{Y}_{α} . We assume the existence of a reading function

$$w: \mathcal{B}(\mathbf{Y}_{\alpha}) \to \mathfrak{S}_n, \quad T \mapsto w(T).$$

From this point onward, we will refer to the set $\{w(T) \mid T \in \mathcal{B}(\mathbf{Y}_{\alpha})\}\$ and its equivalence class as $\mathsf{RW}_w(\mathbf{Y}_{\alpha})$ and $C_{\mathbf{Y}_{\alpha}}$, respectively.

quasisymmetric functions $\{y_{\alpha} \mid \alpha \models n\}$	$H_n(0)$ -module \mathbf{Y}_{α}	$\operatorname{ch}(\mathbf{Y}_{lpha})$	tableau-basis $\mathcal{B}(\mathbf{Y}_{\alpha})$
ribbon Schur func. $\{s_{\alpha}\}$ ([31])	\mathbf{P}_{α} ([27])	$s_{lpha^{ m c}}$	standard ribbon tableaux of shape $rd(\alpha)$
dual immaculate func. $\{\mathfrak{S}^*_{\alpha}\}$ ([6])	\mathcal{V}_{lpha} ([7])	\mathfrak{S}^*_lpha	standard dual immaculate tableaux of shape $cd(\alpha)$
extended Schur func. $\{\mathcal{E}_{\alpha}\}$ ([1])	X_{α} ([29])	\mathcal{E}_{lpha}	standard extended tableaux of shape $cd(\alpha)$
Young quasisymm. Schur func. $\{\widehat{S}_{\alpha}\}$ ([21])	$\widehat{\mathbf{S}}_{lpha}$ ([11])	$\widehat{\mathcal{S}}_{lpha}$	standard Young composition tableaux of shape $cd(\alpha)$
fundamental quasisymm. func. $\{F_{\alpha}\}$ ([31])	\mathbf{F}_{α} ([27])	$F_{\alpha^{c}}$	The standard ribbon tableau T_{α}

TABLE 5.1. Quasisymmetric functions, associated $H_n(0)$ -modules, quasisymmetric characteristics, and tableau-bases.

5.1.1. \mathbf{P}_{α} . Recall that

$$\mathbf{P}_{\alpha} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), w_1(\operatorname{set}(\alpha)^{\mathrm{c}})]_L) \text{ as } H_n(0) \text{-modules.}$$

A standard ribbon tableau (SRT) of shape α is a filling T of the ribbon diagram $\mathsf{rd}(\alpha)$ with $\{1, 2, \ldots, n\}$ such that the entries are all distinct, the entries in each row increase from left to right, and the entries in each column increase from bottom to top. Let $\mathsf{SRT}(\alpha)$ be the set of SRTs of shape α . Let $T_{\alpha} \in \mathsf{SRT}(\alpha)$ be the standard ribbon tableau obtained by filling $\mathsf{rd}(\alpha)$ with the entries $1, 2, \ldots, n$ from bottom to top and from left to right. From [19, Lemma 5.2] (or [16, Section 3.2]) we have

(5.2)
$$\mathsf{RW}_{w_{\mathrm{TL}}}(\mathbf{P}_{\alpha}) = \{ w_{\mathrm{TL}}(T) \mid T \in \mathsf{SRT}(\alpha) \} = [w_0(\operatorname{set}(\alpha^{\mathrm{c}})), w_1(\operatorname{set}(\alpha^{\mathrm{c}}))]_L.$$


Proposition 5.1. Let α be a composition of n. Then

$$C_{\mathbf{P}_{\alpha}} = \{ [w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), w_1(\operatorname{set}(\alpha)^{\mathrm{c}})]_L \}.$$

Proof. Since $\mathsf{RW}_{w_{LT}}(\mathbf{P}_{\alpha}) = [w_0(\operatorname{set}(\alpha)^c), w_1(\operatorname{set}(\alpha)^c)]_L$, the diagram $D_{\operatorname{set}(\alpha)^c;w_1(\operatorname{set}(\alpha)^c)}$ constructed by Algorithm 4.7 is the ribbon diagram $\mathsf{rd}(\alpha)$. More precisely,

$$D_{\operatorname{set}(\alpha)^{c};w_{1}(\operatorname{set}(\alpha)^{c})} = \{(i, j+k_{i}) \mid 1 \le i \le \ell(\alpha), \ 1 \le j \le \alpha_{i}\},\$$

where $k_{\ell(\alpha)} := 0$ and $k_i := \sum_{i < r \le \ell(\alpha)} (\alpha_r - 1)$ for $1 \le i \le \ell(\alpha) - 1$. For example, if $\alpha = (1, 2, 1, 2, 1, 3)$, then



Now, the desired result follows from the property that

$$F_{D_{\operatorname{set}(\alpha)^{c}};w_{1}(\operatorname{set}(\alpha)^{c})}^{\downarrow} = F_{D_{\operatorname{set}(\alpha)^{c}};w_{1}(\operatorname{set}(\alpha)^{c})}^{\rightarrow}.$$

5.1.2. \mathbf{F}_{α} . It is well known that

$$\mathbf{F}_{\alpha} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^{c}), w_0(\operatorname{set}(\alpha)^{c})]_L) \text{ as } H_n(0) \text{-modules}$$

(for instance, see [17, Section 3.2]). By the definitions of \mathbf{F}_{α} and weak Bruhat interval modules in Section 2.3 and [17, Section 3.2], we see that $\mathbf{F}_{\alpha} \cong \mathsf{B}([\sigma,\sigma]_L)$ for all $\sigma \in \mathfrak{S}_n$ with $\operatorname{set}(\alpha)^c = \operatorname{Des}_L(\sigma)$.

Proposition 5.2. Let α be a composition of n. Then

$$C_{\mathbf{F}_{\alpha}} = \{ [\sigma, \sigma]_L \mid \sigma \in [w_0(\operatorname{set}(\alpha)^c), w_0(\operatorname{set}(\alpha))w_0]_R \}, \text{ and} \\ (C_{\mathbf{F}_{\alpha}}, \preceq) \cong (\mathsf{SRT}(\alpha), \leq) \quad (as \ posets).$$

Proof. The first assertions follow from (2.1). For the second assertion, consider the map

$$[w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), w_0(\operatorname{set}(\alpha))w_0]_R \to \mathsf{SRT}(\alpha), \ \sigma \mapsto T_{\sigma},$$

where T_{σ} denotes the SRT of shape α such that $w_{TL}(T_{\sigma}) = \sigma^{-1}$ (see [19, Lemma 5.2]). This map establishes an isomorphism between $(C_{\mathbf{F}_{\alpha}}, \preceq)$ and $(\mathsf{SRT}(\alpha), \leq)$, as required. \Box

5.1.3. \mathcal{V}_{α} . It was shown in [17, Section 3.2] that

$$\mathcal{V}_{\alpha} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^c), \operatorname{w}_{\operatorname{RB}}(\mathcal{T}'_{\alpha})]_L) \text{ as } H_n(0) \text{-modules.}$$


Here, \mathcal{T}'_{α} denotes the standard immaculate tableau of shape α , constructed by first filling the first column with entries $1, 2, \ldots, \ell(\alpha)$ from bottom to top. The remaining boxes are then filled with entries $\ell(\alpha) + 1, \ell(\alpha) + 2, \ldots, n$, moving left to right across each row, starting from the topmost row.

Let $D(\mathcal{V}_{\alpha}, \mathbf{w}_{RB})$ be the diagram consisting of

$$\{(i,1) \mid (1,i) \in \mathsf{cd}(\alpha)\} \cup \{(i,j+k_i) \mid (j,i) \in \mathsf{cd}(\alpha) \text{ and } j \ge 2\},\$$

where $k_{\ell(\alpha)} := 0$ and $k_i := \sum_{i < r \le \ell(\alpha)} (\alpha_r - 1)$ for $1 \le i < \ell(\alpha)$. For example, if $\alpha = (3, 2, 4)$, then





Lemma 5.3. (cf. [12, Theorem 5.6]) Let α be a composition. Then we have

$$\Sigma_L(P_{F_{D(\mathcal{V}_{\alpha}, \mathbf{w}_{\mathrm{RB}})}}) = [w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), \mathbf{w}_{\mathrm{RB}}(\mathcal{T}'_{\alpha})]_L.$$

Proof. By considering the definition of \mathcal{T}'_{α} , we observe that for $1 \leq i \leq \ell(\alpha)$,

$$(\mathcal{T}'_{\alpha})_{1,i} = i, \qquad (\mathcal{T}'_{\alpha})_{i,j} = j + k_i + (\ell(\alpha) - 1) \quad \text{for } 2 \le j \le \alpha_i, \quad \text{and} \\ \text{Des}_{\text{L}}(\mathbf{w}_{\text{RB}}(\mathcal{T}'_{\alpha})) = \{\ell(\alpha), \ell(\alpha) + 1, \dots, n-1\}.$$

Now, applying Algorithm 4.7 to $(set(\alpha)^c, w_{RB}(\mathcal{T}'_{\alpha}))$, we obtain the following sets:

$$X_{i}(\operatorname{set}(\alpha)^{c}; \operatorname{w}_{\operatorname{RB}}(\mathcal{T}_{\alpha}')) = \{i\} \cup \{j + k_{i} + (\ell(\alpha) - 1) \mid 2 \leq j \leq \alpha_{i}\} \text{ for } 1 \leq i \leq \ell(\alpha),$$

$$Y_{1}(\operatorname{w}_{\operatorname{RB}}(\mathcal{T}_{\alpha}')) = \{1, 2, \dots, \ell(\alpha)\}, \text{ and}$$

$$Y_{j}(\operatorname{w}_{\operatorname{RB}}(\mathcal{T}_{\alpha}')) = \{j + \ell(\alpha) - 1\} \text{ for } 2 \leq j \leq n - \ell(\alpha) + 1,$$

From this data, we can construct the desired diagram as required.

For an SIT \mathcal{T} , let $w_{RB1}(\mathcal{T})$ denote the reading word obtained from \mathcal{T} as follows:

(i) Begin by reading the entries in each row from right to left, starting with the bottommost row and moving upward, excluding the entries in the first column.

(ii) Then, read the entries in the first column from bottom to top.

Let \mathcal{T}_{α} be the SIT shape α obtained by filling $\mathsf{cd}(\alpha)$ with $1, 2, \ldots, n$ from left to right and from bottom to top. With these definitions, we have the following.

Proposition 5.4. Let α be a composition of n, let and $\mathcal{D} := D(\mathcal{V}_{\alpha}, \mathbf{w}_{RB})$.

- (a) $C_{\mathcal{V}_{\alpha}} = \{\Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{D}^x)\}.$ Furthermore, $\min C_{\mathcal{V}_{\alpha}} = [w_0(\operatorname{set}(\alpha)^c), \operatorname{w}_{\operatorname{RB}}(\mathcal{T}'_{\alpha})]_L \text{ and } \max C_{\mathcal{V}_{\alpha}} = [\operatorname{w}_{\operatorname{RB}1}(\mathcal{T}_{\alpha}), w_1([n-\ell(\alpha)])]_L.$
- (b) $(C_{\mathcal{V}_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{D}^x), \leq)$ as posets.

Proof. (a) By the construction of \mathcal{D} , it is evident that \mathcal{D} is free of a strictly upper-right configuration. Therefore, the first assertion follows directly from Theorem 4.16. For the second assertion, it follows from Lemma 5.3 that

$$\min C_{\mathcal{V}_{\alpha}} = [w_0(\operatorname{set}(\alpha)^{c}), \operatorname{w}_{\operatorname{RB}}(\mathcal{T}'_{\alpha})]_L.$$

To establish the final equality, consider

$$\max C_{\mathcal{V}_{\alpha}} = \Sigma_L(P_{F_{\mathcal{D}}}) = [\mathbf{w}_{\mathrm{LT}}(T_{\mathcal{D}}), \mathbf{w}_{\mathrm{LT}}(T_{\mathcal{D}})]_L.$$

Consider the tableau $T'_{\mathcal{D}}$. Its reading word, $w_{LT}(T'_{\mathcal{D}})$, can be expressed as

$$\underbrace{\alpha_1 \ \alpha_1 - 1 \ \cdots \ 2}_{\text{row } 1} \ \cdots \ \underbrace{n \ n - 1 \ \cdots \ n - \alpha_{\ell(\alpha)} + 2}_{\text{row } \ell(\alpha)} \ \underbrace{1 \ \alpha_1 + 1 \ \cdots \ \alpha_1 + \cdots + \alpha_{\ell(\alpha) - 1} + 1}_{\text{col. } 1}.$$

Here, 'row *i*' and 'col. 1' refer to the *i*th row and the first column in \mathcal{T}_{α} , respectively. Comparing this word with $w_{\text{RB1}}(\mathcal{T}_{\alpha})$, we have that $w_{\text{LT}}(\mathcal{T}_{\mathcal{D}}) = w_{\text{RB1}}(\mathcal{T}_{\alpha})$. On the other hand, we directly have that

$$w_{LT}(T_{\mathcal{D}}) = n \ n-1 \ \cdots \ \ell(\alpha) + 1 \ 1 \ 2 \ \cdots \ \ell(\alpha) = w_1([n-\ell(\alpha)]).$$

This completes the proof.

(b) Since \mathcal{D} is free of a strictly upper-right configuration, it follows from Theorem 4.12 and Theorem 4.16 that

$$(C_{\mathcal{V}_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{D}^x), \leq)$$
 as posets.

Example 5.5. We illustrate Proposition 5.4 using the diagram \mathcal{D} given in (5.4), where $\alpha = (3, 2, 4)$. Consider the tableaux

$$\mathcal{T}_{\alpha} = \begin{bmatrix} 6 & 7 & 8 & 9 \\ 4 & 5 & \\ 1 & 2 & 3 \end{bmatrix} \quad \text{and} \quad \mathcal{T}_{\alpha}' = \begin{bmatrix} 3 & 4 & 5 & 6 \\ 2 & 7 & \\ 1 & 8 & 9 \end{bmatrix},$$

then we have


$$w_{RB}(\mathcal{T}_{\alpha}) = 321549876, \quad w_{RB}(\mathcal{T}_{\alpha}) = 981726543 \text{ and}$$

 $w_{RB1}(\mathcal{T}'_{\alpha}) = 325987146, \quad w_{RB1}(\mathcal{T}'_{\alpha}) = 987654123.$

According to it follows from Proposition 5.4 that we compute

$$\min C_{\mathcal{V}_{\alpha}} = [w_0(\{1, 2, 4, 6, 7, 8\}), 981726543]_L \text{ and} \\ \max C_{\mathcal{V}_{\alpha}} = [325987146, w_1(\{1, 2, 3, 4, 5, 6\})]_L.$$


On the other hand, since \mathcal{D} is free of a strictly upper-right configuration, it follows from the standard fillings



that we have

min
$$C_{\mathcal{V}_{\alpha}} = \Sigma_L(P_{F_{\mathcal{D}}^{\downarrow}}) = [321549876, 981726543]_L$$
 and
max $C_{\mathcal{V}_{\alpha}} = \Sigma_L(P_{F_{\mathcal{D}}^{\rightarrow}}) = [325987146, 987654123]_L.$

Finally, the elements of $\mathsf{ST}(\mathcal{D}^x)$ are



By comparing $C_{\mathcal{V}_{\alpha}}$ and $\mathsf{ST}(\mathcal{D}^x)$, we observe that $(C_{\mathcal{V}_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{D}^x), \leq)$ as posets.

Combining Corollary 4.19 with Proposition 5.4 yields the following corollary.

Corollary 5.6. (cf. [10, Theorem 4.11]) Let α be a composition of n and $\mathcal{D} := D(\mathcal{V}_{\alpha}, w_{RB})$. Then

 $\mathsf{B}([w_0(I), w_1(\operatorname{set}(\mathsf{r}(\mathcal{D})))]_L)$

is an injective hull of \mathcal{V}_{α} , where $I = [n-1] \setminus \{\beta_1, \beta_2, \dots, \beta_{\ell(\alpha)-1}, n-\ell(\alpha), \dots, n-1\}$ and $\beta_i = \alpha_1 + \dots + \alpha_i - i$.

Proof. By Corollary 4.19 we know that $\mathsf{B}([w_0(\mathsf{Des}_{\mathsf{L}}(\mathsf{w}_{\mathsf{BL}}(F_{\mathcal{D}}^{\rightarrow}))), w_1(\mathsf{set}(\mathsf{r}(\mathcal{D})))]_L)$ is an injective hull of \mathcal{V}_{α} . By considering the diagram \mathcal{D} , it is straightforward to verify that $\mathsf{Des}_{\mathsf{L}}(\mathsf{w}_{\mathsf{BL}}(F_{\mathcal{D}}^{\rightarrow})) = [n-1] \setminus \{\beta_1, \ldots, \beta_{\ell(\alpha)-1}, n-\ell(\alpha), \ldots, n-1\}$, which completes the proof. \Box

It should be remarked that in [10, Theorem 4.11], an injective hull of \mathcal{V}_{α} was directly constructed via an injective map from standard immaculate tableaux to standard ribbon tableaux. Compared to this, the current method is more uniform.

5.1.4. X_{α} . It was shown in [17, Section 3.2] that

$$X_{\alpha} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^{c}), \operatorname{w}_{\operatorname{RB}}(\mathsf{T}'_{\alpha})]_L) \text{ as } H_n(0)\text{-modules.}$$

Here, T'_{α} denotes the standard extended tableau of shape α , constructed by filling with the entries $1, 2, \ldots, \ell(\alpha)$ in each column from bottom to top, starting from the leftmost column. Let $D(X_{\alpha}, w_{\mathrm{RB}})$ be the diagram $\mathsf{cd}(\alpha)^{\mathsf{t}}$, that is,

$$\{(j,i) \mid (i,j) \in \mathsf{cd}(\alpha)\}.$$

For example, if $\alpha = (3, 2, 4)$, then

Lemma 5.7. (cf. [12, Theorem 5.8]) Let α be a composition. Then we have

 $\stackrel{y}{\uparrow}$

$$\Sigma_L(P_{F_{D(X_{\alpha}, \mathbf{w}_{\mathrm{RB}})}^{\downarrow}}) = [w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), \mathbf{w}_{\mathrm{RB}}(\mathsf{T}'_{\alpha})]_L.$$

Proof. The assertion follows in a similar manner as in Lemma 5.3.

Let T_{α} be the standard extended tableau of shape α obtained by filling $\mathsf{cd}(\alpha)$ with $1, 2, \ldots, n$ in each row from left to right, starting from the bottommost row. Then we have the following.

Proposition 5.8. Let α be a composition of n, and let $\mathcal{E} := D(X_{\alpha}, w_{RB})$.

(a) $C_{X_{\alpha}} = \{\Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{E}^x)\}.$ Furthermore, $\min C_{X_{\alpha}} = [w_0(\operatorname{set}(\alpha)^c), \operatorname{w}_{\operatorname{RB}}(\mathsf{T}'_{\alpha})]_L \text{ and } \max C_{X_{\alpha}} = [\operatorname{w}_{\operatorname{BR}}(\mathsf{T}_{\alpha}), w_1(\operatorname{set}((\tilde{\alpha}')^r))]_L.$ (b) $(C_{X_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{E}^x), \leq) \text{ as posets.}$

Proof. (a) By the construction of \mathcal{E} , it is evident that the diagram \mathcal{E} is free of a strictly upper-right configuration. Therefore, by Theorem 4.12 and Theorem 4.16, the first assertion follows. The second assertion is a direct consequence of Lemma 5.7. To prove the final assertion, consider the equality

$$\max C_{X_{\alpha}} = \Sigma_L(P_{F_{\mathcal{E}}^{\rightarrow}}) = [\mathbf{w}_{\mathrm{LT}}(T_{\mathcal{E}}'), \mathbf{w}_{\mathrm{LT}}(T_{\mathcal{E}})]_L.$$

By definition, we observe that

$$\mathrm{w}_{\mathrm{LT}}(T'_{\mathcal{E}}) = \mathrm{w}_{\mathrm{BR}}((T'_{\mathcal{E}})^{\mathrm{t}}) \quad \mathrm{and} \quad \mathrm{w}_{\mathrm{LT}}(T_{\mathcal{E}}) = \mathrm{w}_{\mathrm{BR}}((T_{\mathcal{E}})^{\mathrm{t}}).$$

Since $(T'_{\mathcal{E}})^{t} = \mathsf{T}_{\alpha}$ and $(T_{\mathcal{E}})^{t} = \mathsf{T}'_{\alpha}$, we have

 $\max C_{X_{\alpha}} = [\mathbf{w}_{\mathrm{BR}}(\mathsf{T}_{\alpha}, \mathbf{w}_{\mathrm{BR}}(\mathsf{T}_{\alpha}')]_{L}.$

By the definitions of w_{LT} and $T_{\mathcal{E}}$, since

$$w_{LT}(T_{\mathcal{E}}) = \underbrace{\tilde{c}_{\tilde{\alpha}'_1} + 1 \cdots n}_{\text{row 1}} \underbrace{\tilde{c}_{\tilde{\alpha}'_2} + 1 \cdots \tilde{c}_{\tilde{\alpha}'_1}}_{\text{row 2}} \cdots \underbrace{1 \cdots c_1}_{\text{row } \tilde{\alpha}'_1},$$

where c_j is the number of boxes in the *j*th row of \mathcal{E} (counted from the top) and $\tilde{c}_j = \sum_{1 \le i \le j} c_i$, we conclude that $w_{\text{LT}}(T_{\mathcal{E}}) = w_1(\text{set}((\tilde{\alpha}')^r))$. This establishes the desired result.

(b) Since the diagram \mathcal{E} is free of a strictly upper-right configuration, it follows from Theorem 4.16(a) that

$$(C_{X_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{E}^x), \leq)$$
 as posets.

Example 5.9. Given $\alpha = (3, 2, 4)$, from the tableaux

$$\mathsf{T}_{\alpha} = \begin{array}{c|c} 6 & 7 & 8 & 9 \\ \hline 4 & 5 & \\ \hline 1 & 2 & 3 \end{array} \quad \text{and} \quad \mathsf{T}_{\alpha}' = \begin{array}{c|c} 3 & 6 & 8 & 9 \\ \hline 2 & 5 & \\ \hline 1 & 4 & 7 \end{array},$$

we have that

$$w_{RB}(T_{\alpha}) = 321549876, \quad w_{RB}(T'_{\alpha}) = 741529863 \text{ and} w_{BR}(T_{\alpha}) = 938257146, \quad w_{BR}(T'_{\alpha}) = 978456123.$$

According to Proposition 5.8, the minimal and maximal elements of $C_{X_{\alpha}}$ are

$$\min C_{X_{\alpha}} = [\mathbf{w}_{\mathrm{RB}}(\mathsf{T}_{\alpha}), \mathbf{w}_{\mathrm{RB}}(\mathsf{T}_{\alpha}')]_{L} \quad \text{and} \quad \max C_{X_{\alpha}} = [\mathbf{w}_{\mathrm{BR}}(\mathsf{T}_{\alpha}), \mathbf{w}_{\mathrm{BR}}(\mathsf{T}_{\alpha}')]_{L}.$$

On the other hand, we consider the right-hand diagram \mathcal{E} given in (5.3). The diagram \mathcal{E} is clearly free of a strictly upper-right configuration, so it follows from Theorem 4.12 that

$$F_{\mathcal{E}}^{\downarrow} = \begin{array}{|c|c|} \hline 6 \\ \hline 7 \\ \hline 2 & 4 & 8 \\ \hline 3 & 5 & 9 \end{array} \quad \text{and} \quad F_{\mathcal{E}}^{\nearrow} = F_{\mathcal{E}}^{\rightarrow} = \begin{array}{|c|} \hline 2 \\ \hline 4 & 5 & 6 \\ \hline 7 & 8 & 9 \end{array}$$

Then we compute

$$\min C_{X_{\alpha}} = \Sigma_L(P_{F_{\mathcal{E}}^{\downarrow}}) = [321549876, 74152983]_L = [w_0(\operatorname{set}(\alpha)^c), 74152983]_L \quad \text{and} \\ \max C_{X_{\alpha}} = \Sigma_L(P_{F_{\mathcal{E}}^{\rightarrow}}) = [938257146, 978456123]_L = [938257146, w_1(\operatorname{set}((\tilde{\alpha}')^r))]_L.$$

Corollary 5.10. Let α be a composition of n and $\mathcal{E} := D(X_{\alpha}, w_{RB})$. Then

 $\mathsf{B}([w_0(\operatorname{set}(\mathsf{r}(\mathcal{E}))), w_1(\operatorname{set}(\mathsf{r}(\mathcal{E})))]_L)$

is an injective hull of X_{α} . In particular, it is a projective indecomposable module.

36

Proof. By Corollary 4.19 we know that $\mathsf{B}([w_0(\mathrm{Des}_{\mathsf{L}}(\mathsf{w}_{\mathrm{BL}}(F_{\mathcal{E}}^{\rightarrow}))), w_1(\mathrm{set}(\mathsf{r}(\mathcal{E})))]_L)$ is an injective hull of X_{α} . By the definition of the diagram \mathcal{E} , we observe that \mathcal{E} contains no subdiagram of the form:



Using this observation and the definitions of $w_{\rm BL}$ and $F_{\mathcal{E}}^{\rightarrow}$, we see that the entry in the rightmost box of row r ($r \geq 2$) in $F_{\mathcal{E}}^{\rightarrow}$ is contained in $\mathrm{Des}_{\mathrm{L}}(w_{\mathrm{BL}}(F_{\mathcal{E}}^{\rightarrow}))$. Thus we have $\mathrm{Des}_{\mathrm{L}}(w_{\mathrm{BL}}(F_{\mathcal{E}}^{\rightarrow})) = \mathrm{set}(\tilde{\alpha}^{\mathrm{r}}) = \mathrm{set}(\mathsf{r}(\mathcal{E}))$, completing the proof.

5.1.5. $\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}$. The source tableau $\widehat{\tau}_{\alpha,\mathcal{C}}$ in the canonical class \mathcal{C} is constructed by filling $\mathsf{cd}(\alpha)$ with entries $1, 2, \ldots, n$ in each row from left to right, starting from the bottommost row. And, $\widehat{\tau}'_{\alpha,\mathcal{C}}$ denotes the sink tableau in \mathcal{C} , which can be obtained from the source tableau $\widehat{\tau}_{\alpha,\mathcal{C}}$ in C by the algorithm described in [3, Section 4.1] (or [12, Algorithm 5.13]). ⁴ It was shown in [17, Section 3.3] that

(5.5)
$$\widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \cong \mathsf{B}([w_0(\operatorname{set}(\alpha)^c), \mathsf{w}_{\operatorname{RB}}(\widehat{\tau}'_{\alpha,\mathcal{C}})]_L) \text{ as } H_n(0) \text{-modules.}$$

We here give an algorithm to the diagram from the pair $(w_0(\text{set}(\alpha)^c), w_{\text{RB}}(\hat{\tau}'_{\alpha,\mathcal{C}}))$.

Algorithm 5.11.

Step 1. For each $1 \leq i \leq \ell(\alpha)$, define

$$X_i := \{ (1, i), (2, i), \dots, (\alpha_i, i) \}.$$

Step 2. Define $\mathsf{BB}(\alpha)$ as the set of specific boxes in $\mathsf{cd}(\alpha)$ based on the following conditions:

- Include box (1, y) in column 1 if $1 \le y < \ell(\alpha)$ and $\alpha_y > 1$, or if $y = \ell(\alpha)$.
- Include box (x, y) for x > 1 if there is no box strictly above it in the same column or the column immediately to the left.

Define the order \leq on $\mathsf{BB}(\alpha)$ such that $(x, y) \leq (u, v)$ if either x = u = 1 and y < v, or if x < u.

Step 3. Arrange the elements of $BB(\alpha)$ in increasing order as

$$\mathsf{BB}(\alpha) = \{ (b_1, d_1) \le (b_2, d_2) \le \dots \le (b_l, d_l) \}.$$

Note that $b_1 = 1$. Define $Y_1 := \{(1, d_1), (1, d_1 - 1), \dots, (1, 1)\}$ and set $A_1 := \mathsf{cd}(\alpha) \setminus Y_1$.

Step 4. For each j = 2, 3, ..., l, construct the set Y_j based on the empty boxes in the diagram A_{j-1} as follows:

(i) If $b_{i+1} = 1$, then define

$$Y_{j+1} := \{ (1, d_{j+1}), (1, d_{j+1} - 1), \dots, (1, d_j + 1) \}.$$

Otherwise, set $Y_{j+1} := \{ (b_{j+1}, d_{j+1}) \}.$

⁴In [3, Section 4.1], the tableau $\hat{\tau}'_{\alpha,\mathcal{C}}$ is referred to as T_{sup} . Meanwhile, [12, Algorithm 5.13] is based on standard reverse composition tableaux (SRCTs). To adapt the algorithm for our case, it is necessary to apply the natural bijection between SRCTs and SYCTs

(ii) Let κ be the last box in Y_{j+1} , and let c be its column index. Check if there is an empty box in column c + 1 of A_j that is strictly below κ . If such a box exists, add the lowermost of these boxes to Y_{j+1} . Otherwise, stop the process for this Y_{j+1} .

Define $A_{j+1} := A \setminus Y_{j+1}$ using the updated Y_{j+1} .

Step 5. Using the sets X_i and Y_i , construct the final diagram $D_{\alpha,\mathcal{C}}$ by


$$(i, j) \in D_{\alpha, \mathcal{C}}$$
 if $X_i \cap Y_j \neq \emptyset$.

And, then return $D_{\alpha,\mathcal{C}}$.

Lemma 5.12. Given $\alpha \models n$, Algorithm 5.11 returns a diagram in \mathfrak{D}_n .

Proof. From Algorithm 5.11, every box in $cd(\alpha)$ is uniquely assigned to both a set X_i and a set Y_j . Since X_i forms the boxes in column i and Y_j forms the boxes in row j, it follows that $D_{\alpha,\mathcal{C}} \in \mathfrak{D}_n$.

Example 5.13. Applying Algorithm 5.11 to $\alpha = (2, 5, 1, 3, 3)$, one can see that



Lemma 5.14. Let α be a composition of n and let $\mathcal{C} := D(\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}, \mathbf{w}_{RB})$. Then we have

$$\Sigma_L(P_{F_{\mathcal{C}}^{\downarrow}}) = [w_0(\operatorname{set}(\alpha)^{\mathrm{c}}), \operatorname{w}_{\mathrm{RB}}(\hat{\tau}'_{\alpha,\mathcal{C}})]_L.$$

Proof. From Proposition 4.4(a) and (c), we have

$$\Sigma_L(P_{F_{\varphi}^{\downarrow}}) = [w_0(\operatorname{set}(\mathsf{c}(\mathfrak{C}))^{\mathrm{c}}, \operatorname{w}_{\operatorname{TL}}(T_{\mathfrak{C}})]_L.$$

According to Algorithm 5.11, the boxes in X_i $(1 \le i \le \ell(\alpha))$ are arranged in the *i*th column of \mathcal{C} , which implies that $c(\mathcal{C}) = \alpha$. Thus, it follows that $w_0(\operatorname{set}(c(\mathcal{C}))^c) = w_0(\operatorname{set}(\alpha)^c)$.

Next, we show that $w_{TL}(T_c) = w_{RB}(\hat{\tau}'_{\alpha})$. Recall the construction of the tableau $\hat{\tau}'_{\alpha}$ as described in [3, Section 4.1], as well as the definition of T_c . Comparing this method with Algorithm 5.11, one can easily observe that the entries in the *i*th row of $\hat{\tau}'_{\alpha}$ from right to left correspond precisely to those in the *i*th column of T_c from top to bottom (refer to Example 5.13). This implies that $w_{TL}(T_c) = w_{RB}(\hat{\tau}'_{\alpha})$.

Given a standard filling T with 1, 2, ..., n, let \overline{T} be the filling obtained from T by allocating $\overline{T}_{i,j} = n + 1 - T_{i,j}$. For an SYCT $\tau \in C$ and a standard filling R of the same shape, let $w_R(\tau)$ denote the word obtained by reading the entries of τ according to the order specified by R. Here, the order of R refers to the sequence dictated by the boxes labeled $1, 2, \ldots, n$ in R.

Proposition 5.15. Let α be a composition of n. Let $\mathcal{C} := D_{\alpha,\mathcal{C}}$.

(a) C_{ŝ_{α,C}} = {Σ_L(P_{T^x}) | T ∈ ST(C^x)}. Furthermore,
min C_{ŝ_{α,C}} = [w₀(set(α)^c), w_{RB}(τ'_{α,C})]_L and max C_{ŝ_{α,C}} = [w_{τ'_{α,C}}(τ̂_{α,C}), w₁(set(r(C)))]_L.
(b) (C_{ŝ_{α,C}}, ≤) ≅ (ST(C^x), ≤) as posets.

Proof. (a) First, we prove that \mathcal{C} is free of a strictly upper-right configuration. To do this, we recall the set $\mathsf{BB}(\alpha)$ in Step 3 of Algorithm 5.11. Letting $l_{\alpha} = |\mathsf{BB}(\alpha)|$, we have

$$\mathsf{BB}(\alpha) = \{ (b_1, d_1) \le (b_2, d_2) \le \dots \le (b_{l_\alpha}, d_{l_\alpha}) \}.$$

For $1 \leq j \leq l_{\alpha}$, define the subdiagram $\mathcal{C}|_{[j]}$ of \mathcal{C} by $(u, v) \in \mathcal{C}|_{[j]}$ if $X_u \cap Y_v \neq \emptyset$ for $1 \leq u \leq \ell(\alpha)$ and $1 \leq v \leq j$. For the definition of X_i and Y_k , see Algorithm 5.11. By definition, it follows that $\mathcal{C} = \mathcal{C}|_{[l_{\alpha}]}$. The proof proceeds by induction on j. If j = 1, then $\mathcal{C}|_{[1]} = \{(1,1),(1,2),\ldots,(1,b_1)\}$, which is evidently free of a strictly upper-right configuration. Now, assume that for all $1 \leq j < l_{\alpha}$, $\mathcal{C}|_{[j]}$ is free of a strictly upper-right configuration. Suppose, for contradiction, that $\mathcal{C}|_{[j+1]}$ has a strictly upper-right pair. Then there are two pairs $(x_1, j') \in \mathcal{C}|_{[j]}$ and $(x_2, j+1) \in \mathcal{C}|_{[j+1]}$ such that

- (i) $x_1 < x_2, j' \le j < j+1$, and
- (ii) no box in $\mathcal{C}|_{[j+1]}$ lies within the smallest rectangular subdiagram containing (x_1, j') and $(x_2, j+1)$.

Note that the boxes (x_1, j') and $(x_2, j+1)$ are constructed from $(X_{x_1}, Y_{j'})$ and (X_{x_2}, Y_{j+1}) , respectively. Let c_1 and c_2 be the column indices of boxes in $cd(\alpha)$ corresponding to $(X_{x_1}, Y_{j'})$ and (X_{x_2}, Y_{j+1}) , respectively. Since $x_1 < x_2$, we have two cases:

- $c_1 < c_2$: Here, $(c_2 1, x_2)$ lies in $cd(\alpha)$ and belongs to Y_q for some $j' \le q < j + 1$. This implies that (x_2, q) is a box of the rectangular diagram determined by (x_1, j') and $(x_2, j + 1)$, contradicting the assumption.
- $c_1 > c_2$: Let x'_2 be the largest row index such that $(c_2, x'_2) \in \mathsf{cd}(\alpha)$ and $x'_2 < x_2$. Here, (c_2, x'_2) belongs to Y_q for some $j' \leq q < j + 1$. This again implies that (x_2, q) is a box of the rectangular diagram determined by (x_1, j') and $(x_2, j + 1)$, leading to a contradiction.

Thus, $\mathcal{C}|_{[j+1]}$ is free of a strictly upper-right configuration. Thus, by induction, it follows that $\mathcal{C}|_l = \mathcal{C}$ is free of a strictly upper-right configuration. Consequently, we derive the first equality

$$C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \{ \Sigma_L(P_{T^x}) \mid T \in \mathsf{ST}(\mathcal{C}^x) \}.$$

The second equality follows directly from Theorem 4.16(b) and Lemma 5.14. Finally, recall that

$$\max C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \Sigma_L(P_{F_{\mathcal{C}}^{\rightarrow}}) = [\mathsf{w}_{\mathrm{LT}}(T_{\mathcal{C}}'), w_1(\operatorname{set}(\mathsf{r}(\mathcal{C})))]_L$$

as shown in Theorem 4.16(a). Therefore, the third equality follows from the equality

$$\mathbf{w}_{\mathrm{LT}}(T'_{\mathcal{C}}) = \mathbf{w}_{\overline{\hat{\tau}'_{\alpha,\mathcal{C}}}}(\hat{\tau}_{\alpha,\mathcal{C}}).$$

(b) This assertion directly follows from Theorem 4.16.

Example 5.16. Let $\alpha = (2, 5, 1, 3, 3) \models 14$. We validate Proposition 5.15(a) using the tableaux $\hat{\tau}_{\alpha,\mathcal{C}}, \hat{\tau}'_{\alpha,\mathcal{C}}$, and $\overline{\hat{\tau}'_{\alpha,\mathcal{C}}}$:

$$\hat{\tau}_{\alpha,\mathcal{C}} = \begin{array}{c} 121314\\ 91011\\ 8\\ \hline 34567\\ 12 \end{array} \qquad \hat{\tau}'_{\alpha,\mathcal{C}} = \begin{array}{c} 91213\\ 6811\\ 5\\ \hline 5\\ 3471014\\ 12 \end{array} \qquad \bar{\tau}'_{\alpha,\mathcal{C}} = \begin{array}{c} 632\\ 974\\ 10\\ \hline 1211851\\ 1413 \end{array}$$

From the definitions of w_{RB} and $w_{\overline{\hat{\tau}'_{\alpha,c}}}$, we compute that

$$\begin{split} [\mathbf{w}_{\mathrm{RB}}(\hat{\tau}_{\alpha,\mathcal{C}}), \mathbf{w}_{\mathrm{RB}}(\hat{\tau}'_{\alpha,\mathcal{C}})]_{L} &= [w_{0}(\{1,3,4,5,6,9,10,12,13\}), 2\,1\,14\,10\,7\,4\,3\,5\,11\,8\,6\,13\,12\,9]_{L}, \\ [\mathbf{w}_{\overline{\hat{\tau}'_{\alpha,\mathcal{C}}}}(\hat{\tau}_{\alpha,\mathcal{C}}), \mathbf{w}_{\overline{\hat{\tau}'_{\alpha,\mathcal{C}}}}(\hat{\tau}'_{\alpha,\mathcal{C}})]_{L} &= [7\,14\,13\,11\,6\,12\,10\,5\,9\,8\,4\,3\,2\,1, w_{1}(\{1,2,5,8,11,13\})]_{L}. \end{split}$$

On the other hand, since the diagram \mathcal{C} associated with the composition α and \mathcal{C} , as described in Example 5.13 is free of a strictly upper-right configuration, it follows from Theorem 4.16 that the minimal and maximal elements of $C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}}$ are

 $\min C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \Sigma_L(P_{F_{\mathcal{C}}^{\downarrow}}) = [w_0(\{1,3,4,5,6,9,10,12,13\}), 2\,1\,14\,10\,7\,4\,3\,5\,11\,8\,6\,13\,12\,9]_L,$ $\max C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \Sigma_L(P_{F_{\mathcal{C}}^{\rightarrow}}) = [7\,14\,13\,11\,6\,12\,10\,5\,9\,8\,4\,3\,2\,1, w_1(\{1,2,5,8,11,13\})]_L.$

Here,

$$F_{c}^{\downarrow} = \begin{array}{c} 3 \\ 12 \\ 12 \\ 5 \\ 1014 \\ 6 \\ 8 \\ 11 \\ 1 \\ 2 \end{array} \quad \text{and} \quad F_{c}^{\checkmark} = F_{c}^{\rightarrow} = \begin{array}{c} 1 \\ 2 \\ 3 \\ 6 \\ 7 \\ 8 \\ 9 \\ 1011 \\ 12 \\ 14 \end{array}$$

Thus, by comparing these results, we confirm that

 $\min C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = [\mathbf{w}_{\mathrm{RB}}(\hat{\tau}_{\alpha,\mathcal{C}}), \mathbf{w}_{\mathrm{RB}}(\hat{\tau}_{\alpha,\mathcal{C}}')]_{L} \quad \text{and} \quad \max C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = [\mathbf{w}_{\overline{\tau}_{\alpha,\mathcal{C}}'}(\hat{\tau}_{\alpha,\mathcal{C}}), \mathbf{w}_{\overline{\tau}_{\alpha,\mathcal{C}}'}(\hat{\tau}_{\alpha,\mathcal{C}}')]_{L}.$ We have the following from Corollary 4.19.

Corollary 5.17. Let α be a composition of n and $\mathcal{C} := D(\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}, \mathbf{w}_{RB})$. Then $\mathbf{P}([\mathbf{w}, (\mathbf{D}_{\mathsf{S}_{\alpha}}, (\mathbf{w}_{\mathsf{R}_{\alpha}}, (\mathbf{F}_{\mathsf{S}_{\alpha}}^{\rightarrow}))) = \mathbf{w}(\mathsf{sot}(\mathbf{r}(\mathcal{C})))]$

$$\mathsf{B}([w_0(\mathrm{Des}_{\mathrm{L}}(\mathrm{w}_{\mathrm{BL}}(F_{\mathfrak{C}}^{\rightarrow}))), w_1(\mathrm{set}(\mathsf{r}(\mathfrak{C})))]_L)$$

is an injective hull of $\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}$.

5.2. Upper descent intervals from submodules of projective indecomposable $H_n(0)$ -modules I. We begin by introducing the submodules under consideration. Let α be a composition of n.

- In [26, Section 6], Niese–Sundaram–van Willigenburg–Vega–Wang constructs an indecomposable $H_n(0)$ -module \mathcal{RV}_{α}^5 by defining a 0-Hecke action on the set of standard dual immaculate tableaux of shape α . The image of this module under the quasisymmetric characteristic is the row-strict dual immaculate quasisymmetric function indexed by α .
- In [26, Section 7], Niese–Sundaram–van Willigenburg–Vega–Wang construct an indecomposable $H_n(0)$ -module $\mathcal{R}X_{\alpha}$ ⁵ by defining a 0-Hecke action on the set of *standard extended tableaux of shape* α . The image of this module under the quasisymmetric characteristic is the *row-strict extended Schur function* indexed by α .
- In [3, Section 3], Bardwell–Searles construct an $H_n(0)$ -module $\mathcal{R}\widehat{\mathbf{S}}_{\alpha}{}^6$ by defining a 0-Hecke action on the set of *standard Young row-strict tableaux of shape* α . The image of this module under the quasisymmetric characteristic is the *Young row-strict quasisymmetric Schur function* indexed by α .

quasisymmetric functions $\{y_{\alpha} \mid \alpha \models n\}$	$H_n(0)$ -module \mathbf{Y}_{α}	$\operatorname{ch}(\mathbf{Y}_{lpha})$	$\textbf{tableau-basis} \ \mathcal{B}(\mathbf{Y}_{\alpha})$
row-strict dual immaculate func. $\{\mathcal{RG}^*_{\alpha}\}$ ([25])	\mathcal{RV}_{lpha} ([26])	$\mathcal{R}\mathfrak{S}^*_lpha$	standard dual immaculate tableaux of shape $cd(\alpha)$
row-strict extended Schur func. $\{\mathcal{RE}_{\alpha}\}$ ([25])	$\mathcal{R}X_{\alpha}$ ([26])	\mathcal{RE}_lpha	standard extended tableaux of shape $cd(\alpha)$
row-strict Young quasisymm. Schur func. $\{\mathcal{R}_{\alpha}\}$ ([22])	$\mathcal{R}\widehat{\mathbf{S}}_{lpha}$ ([3])	\mathcal{R}_{lpha}	standard Young composition tableaux of shape $cd(\alpha)$

TABLE 5.2. Quasisymmetric functions, associated $H_n(0)$ -modules, quasisymmetric characteristics, and tableau-bases

To begin with, we observe that these modules can be obtained from the quotient modules in Section 5.1 by applying an anti-involution. Let us review the definitions of (anti-) involution twists. Given an automorphism μ of $H_n(0)$ and a left $H_n(0)$ -module M, we define $\mu[M]$ by the left $H_n(0)$ -module with the same underlying space as M and with the action \cdot_{μ} defined by

$$h \cdot_{\mu} v := \mu(h) \cdot v$$
 for $h \in H_n(0)$ and $v \in M$.

We define $\mathbf{T}^+_{\mu} : H_n(0)$ -mod $\to H_n(0)$ -mod to be the covariant functor, called the μ -twist, sending a left $H_n(0)$ -module M to $\mu[M]$ and an $H_n(0)$ -module homomorphism $f : M \to N$ to $\mathbf{T}^+_{\mu}(f) : \mu[M] \to \mu[N]$ defined by $\mathbf{T}^+_{\mu}(f)(v) = f(v)$ for $v \in M$.

⁵In [26], the modules \mathcal{V}_{α} , $\mathcal{R}\mathcal{V}_{\alpha}$ and $\mathcal{R}X_{\alpha}$ are denoted by \mathcal{W}_{α} , \mathcal{V}_{α} and \mathcal{Z}_{α} , respectively.

⁶In [3], the module $\mathcal{R}\widehat{\mathbf{S}}_{\alpha}$ is denoted by \mathbf{R}_{α} .

Similarly, given an anti-automorphism ν of $H_n(0)$ and a left $H_n(0)$ -module M, we define $\nu[M]$ by the left $H_n(0)$ -module with M^* , the dual space of M, as the underlying space and with the action \cdot^{ν} defined by

(5.6)
$$(h \cdot^{\nu} \delta)(v) := \delta(\nu(h) \cdot v) \text{ for } h \in H_n(0), \, \delta \in M^*, \text{ and } v \in M.$$

We define \mathbf{T}_{ν}^{-} : $H_n(0)$ -mod $\rightarrow H_n(0)$ -mod to be the contravariant functor, called the ν -twist, sending an $H_n(0)$ -module M to $\nu[M]$ and an $H_n(0)$ -module homomorphism f: $M \rightarrow N$ to $\mathbf{T}_{\nu}^{-}(f) : \nu[N] \rightarrow \nu[M]$ defined by $\mathbf{T}_{\nu}^{-}(f)(\delta) = \delta \circ f$.

In this subsection, we consider two involutions ϕ , θ and an anti-involution χ on $H_n(0)$ defined by

$$\phi(\overline{\pi}_i) = \overline{\pi}_{n-i}, \quad \theta(\overline{\pi}_i) = -\pi_i, \text{ and } \chi(\overline{\pi}_i) = \overline{\pi}_i \quad (1 \le i \le n-1).$$

These (anti-)involutions were introduced by Fayers [15, Proposition 3.2] and commute with each other.

Lemma 5.18. (cf. [17, Table 2 and Section 4]) Let α be a composition of n. Then we have the following $H_n(0)$ -module isomorphisms:

(5.7)
$$\theta \circ \chi[\mathbf{P}_{\alpha}] \cong \mathbf{P}_{\alpha^{t}}, \quad \theta \circ \chi[\mathbf{F}_{\alpha}] \cong \mathbf{F}_{\alpha^{c}}, and$$

(5.8)
$$\mathcal{RV}_{\alpha} \cong \boldsymbol{\theta} \circ \boldsymbol{\chi}[\mathcal{V}_{\alpha}], \quad \mathcal{RX}_{\alpha} \cong \boldsymbol{\theta} \circ \boldsymbol{\chi}[X_{\alpha}], \quad \mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \cong \boldsymbol{\theta} \circ \boldsymbol{\chi}[\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}].$$

Furthermore, we have a sequence of injective $H_n(0)$ -module homomorphisms

(5.9)
$$\mathbf{F}_{\alpha^{c}} \longrightarrow \mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}} \longrightarrow \mathcal{R}X_{\alpha} \longrightarrow \mathcal{R}\mathcal{V}_{\alpha} \longrightarrow \mathbf{P}_{\alpha^{t}}$$

Proof. The isomorphisms in (5.7) and (5.8) can be found in [17, Table 2] and [17, Section 4], respectively. And, the series in (5.9) is obtained from (5.1) by applying the duality functor $\mathbf{T}^+_{\theta} \circ \mathbf{T}^-_{\chi}$.

It was shown in [17, Table 1] that $\theta \circ \chi[\mathsf{B}([\sigma, \rho]_L)] \cong \mathsf{B}([\rho w_0, \sigma w_0]_L)$ as $H_n(0)$ -modules. Since the modules in (5.1) are, up to isomorphism, of the form $\mathsf{B}([w_0(\operatorname{set}(\alpha)^c), -]_L))$, this result implies that the modules in (5.9) are, up to isomorphism, of the form $\mathsf{B}([-, w_1(\operatorname{set}(\alpha^r))]_L))$. We are now ready to state the main result of this subsection.

Proposition 5.19. Let α be a composition of n. Let $\mathcal{D} := D(\mathcal{V}_{\alpha}, \mathbf{w}_{RB}), \mathcal{E} := D(X_{\alpha}, \mathbf{w}_{RB}),$ and $\mathcal{C} := D(\widehat{\mathbf{S}}_{\alpha, \mathcal{C}}, \mathbf{w}_{RB})$. Then we have the following.

- (a) $C_{\mathcal{R}\mathcal{V}_{\alpha}} = \{Iw_0 \mid I \in C_{\mathcal{V}_{\alpha}}\}, and (C_{\mathcal{R}\mathcal{V}_{\alpha}}, \preceq) \cong (\mathsf{ST}((\mathcal{D}^{\mathsf{t}})^x), \leq) as posets.$
- (b) $C_{\mathcal{R}X_{\alpha}} = \{ Iw_0 \mid I \in C_{X_{\alpha}} \}, and (C_{\mathcal{R}X_{\alpha}}, \preceq) \cong (\mathsf{ST}((\mathcal{E}^t)^x), \leq) as posets.$
- (c) $C_{\mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \{ Iw_0 \mid I \in C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} \}, and (C_{\mathcal{R}\widehat{\mathbf{S}}_{\alpha}}, \preceq) \cong (\mathsf{ST}((\mathcal{C}^t)^x), \leq) as posets.$

Here, Iw_0 denotes the set $\{\zeta w_0 \mid \zeta \in I\}$.

Proof. By combining the isomorphism $\theta \circ \chi[\mathsf{B}([\sigma, \rho]_L)] \cong \mathsf{B}([\rho w_0, \sigma w_0]_L)$ with (5.8), we deduce the following:

$$C_{\mathcal{R}\mathcal{V}_{\alpha}} = \{ Iw_0 \mid I \in C_{\mathcal{V}_{\alpha}} \} \quad C_{\mathcal{R}X_{\alpha}} = \{ Iw_0 \mid I \in C_{X_{\alpha}} \} \quad C_{\mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} = \{ Iw_0 \mid I \in C_{\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}} \}$$

Additionally, from [12, Theorem 3.6] it follows that $\theta \circ \chi(\mathbf{M}_P) \cong \mathbf{M}_{\overline{P}}$ for $P \in \mathsf{P}_n$. Moreover, the proof of Proposition 4.4(a) establishes that $\overline{P}_{F_D^{\downarrow}} = P_{F_D^{\downarrow}}$ for all $D \in \mathfrak{D}_n$. These results, combined with Proposition 5.4, Proposition 5.8, and Proposition 5.15, confirm the desired poset isomorphisms.

The following corollary follows from Corollary 5.6, Corollary 5.10 and Corollary 5.17. Corollary 5.20. Let α be a composition of n. Let $\mathcal{D} := D(\mathcal{V}_{\alpha}, \mathbf{w}_{\text{RB}}), \mathcal{E} := D(X_{\alpha}, \mathbf{w}_{\text{RB}}),$ and $\mathcal{C} := D(\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}, \mathbf{w}_{\text{RB}})$. Then we have the following.

- (a) $\mathsf{B}([w_1(\operatorname{set}(\mathsf{r}(\mathcal{D}))^{\mathrm{t}}), w_1(I^{\mathrm{t}})]_L)$ is a projective cover of \mathcal{RV}_{α} . Here, I denotes the set in Corollary 5.6(a).
- (b) $\mathsf{B}([w_0(\operatorname{set}(\mathsf{r}(\mathcal{E}))^{\mathrm{t}}), w_1(\operatorname{set}(\mathsf{r}(\mathcal{E}))^{\mathrm{t}})]_L)$ is a projective cover of $\mathcal{R}X_{\alpha}$.
- (c) $\mathsf{B}([w_0(\operatorname{set}(\mathsf{r}(\mathcal{C}))^{\mathrm{t}}), w_1(\operatorname{Des}_{\mathrm{L}}(w_{\mathrm{BL}}(F_{\mathcal{C}}^{\rightarrow}))^{\mathrm{t}})]_L)$ is a projective cover of $\mathcal{R}\widehat{\mathbf{S}}_{\alpha,\mathcal{C}}$.

5.3. Upper descent intervals from submodules of projective indecomposable $H_n(0)$ -modules II. In this subsection, we discuss the submodules of a projective indecomposable $H_n(0)$ -module that are related to the representation theory of 0-Hecke–Clifford algebras.

Assume that α is a peak composition of n, meaning that $\alpha_i \neq 1$ for $1 \leq i < \ell(\alpha)$. A standard peak immaculate tableau (SPIT) of shape α is a standard immaculate tableau (SIT) \mathcal{T} of shape α such that, for each $1 \leq k \leq n$, the subdiagram of $\mathsf{cd}(\alpha)$ consisting of boxes filled with entries $\leq k$ forms the diagram of a peak composition. Let $\mathsf{SPIT}(\alpha)$ denote the set of SPITs of shape α . In [30], Searles introduced a 0-Hecke–Clifford module structure on $\mathsf{SPIT}(\alpha)$. Here, however, we consider only the 0-Hecke module structure on $\mathsf{SPIT}(\alpha)$.

Recall the module \mathbf{Q}_{α}^{r} , where the $H_{n}(0)$ -action on the \mathbb{C} -span of $\mathsf{SPIT}(\alpha)$ is defined as follows: for each $i = 1, 2, \ldots, n-1$ and $\mathcal{T} \in \mathsf{SPIT}(\alpha)$,

(5.10)
$$\overline{\pi}_i \cdot \mathcal{T} = \begin{cases} -\mathcal{T} & \text{if } i \text{ is weakly above } i+1 \text{ in } \mathcal{T}, \\ 0 & \text{if } i \text{ and } i+1 \text{ are in the first column of } \mathcal{T}, \\ s_i \cdot \mathcal{T} & \text{otherwise.} \end{cases}$$

We refer to [13, Section 3.2] for the details.⁷

⁷It was shown in [30] that the image of the 0-Hecke–Clifford module $\mathbf{Q}_{\alpha}^{r} \uparrow_{H_{n}(0)}^{HCl_{n}(0)}$ under the peak quasisymmetric characteristic is given by $\widetilde{Q}_{\alpha} = \sum_{\mathcal{T} \in \mathsf{SPIT}(\alpha)} K_{\operatorname{Peak}(\operatorname{Des}_{L}(\mathsf{w}_{\mathrm{LT}}(\mathcal{T})))}$, where $K_{\operatorname{Peak}(\operatorname{Des}_{L}(\mathsf{w}_{\mathrm{LT}}(\mathcal{T})))}$ is the peak quasisymmetric function associated to $\operatorname{Peak}(\operatorname{Des}_{L}(\mathsf{w}_{\mathrm{LT}}(\mathcal{T})))$. This function is referred to as the quasisymmetric Schur Q-function indexed by α . We refer to [13, Section 2] for the undefined notations.

Let \mathcal{T}_{α}'' be the SPIT of shape α whose entries in column 1 are the first $\ell(\alpha)$ odd numbers, whose entries in column 2 are the first $\ell(\alpha)-1$ or $\ell(\alpha)$ even numbers (depending on whether or not the last part of α is equal to 1), and whose entries in subsequent rows from top to bottom are the remaining numbers, increasing consecutively up each row. Then the following lemma shows that $\mathsf{RW}_{w_{\text{LT}}}(\mathbf{Q}_{\alpha}^{r})$ forms an upper descent interval.

Lemma 5.21. ([13, Lemma 3.18]) Let α be a peak composition of n. Then we have

$$\mathbf{Q}_{\alpha}^{r} \cong \overline{\mathsf{B}}([\mathsf{w}_{\mathrm{LT}}(\mathcal{T}_{\alpha}''), w_{1}(\mathrm{set}(\alpha)^{\mathrm{r}})]_{L}) \quad as \ H_{n}(0) \text{-modules}.$$


Let $D(\mathbf{Q}_{\alpha}^{r}, \mathbf{w}_{\mathrm{LT}})$ be the diagram whose *j*th row is

$$\begin{cases} \{(j+x-1,j) \mid 1 \le x \le \alpha_{\ell(\alpha)}\} & \text{if } j = \ell(\alpha), \\ \{(j,j), (j+1,j)\} \cup \{(k_j+x,j) \mid 3 \le x \le \alpha_j\} & \text{if } j < \ell(\alpha). \end{cases}$$

Here, $k_j := (\ell(\alpha)-1) + \sum_{j < r \le \ell(\alpha)} (\alpha_r - 2)$ for $1 \le j \le \ell(\alpha) - 2$. For example, if $\alpha = (3, 2, 4, 2)$ and $\beta = (3, 2, 3, 1)$, then the corresponding diagrams $D(\mathbf{Q}^r_{\alpha}, \mathbf{w}_{\mathrm{LT}})$ and $D(\mathbf{Q}^r_{\beta}, \mathbf{w}_{\mathrm{LT}})$ are

- $\{(1,1), (2,1), (2,2), (3,2), (3,3), (4,3), (4,4), (5,4), (6,3), (7,3), (8,1)\}$ and
- $\{(1,1), (2,1), (2,2), (3,2), (3,3), (4,3), (4,4), (5,3), (6,1)\},\$

respectively. Graphically, these diagrams are represented as



From Proposition 4.4 and Lemma 5.21 it follows that

(5.12)
$$\mathsf{RW}_{\mathsf{w}_{\mathrm{LT}}}(\mathbf{Q}_{\alpha}^{r}) = \Sigma_{L}(P_{F_{D}^{\rightarrow}(\mathbf{Q}_{\alpha}^{r},\mathsf{w}_{\mathrm{TT}})})$$

For an SPIT \mathcal{T} , we let $w_{dr}(\mathcal{T})$ the reading word obtained from \mathcal{T} by reading the entries at (1, j), (2, j - 1) for $1 \leq j \leq \ell(\alpha)$ in the first two columns, and then continuing to read the entries top to bottom in each subsequent row, with the rows read in order from left to right. For a positive integer k, let $2[k] := \{2, 4, \ldots, 2k\}$.

Theorem 5.22. Let α be a peak composition of n. Let $C_{\mathbf{Q}_{\alpha}^{r}}$ be the equivalence class of $[w_{\mathrm{LT}}(\mathcal{T}_{\alpha}''), w_{1}(\operatorname{set}(\alpha^{\mathrm{r}}))]_{L}$ and $\mathfrak{G} := D(\mathbf{Q}_{\alpha}^{r}, w_{\mathrm{LT}}).$

(a)
$$C_{\mathbf{Q}_{\alpha}^{r}} = \{\Sigma_{L}(P_{T^{x}}) \mid T \in \mathsf{ST}(\mathfrak{G}^{x})\}.$$
 Furthermore,
 $\min C_{\mathbf{Q}_{\alpha}^{r}} = [w_{0}(2[\ell(\alpha) - 1]), w_{\mathrm{dr}}(\mathcal{T}_{\alpha})]_{L} \quad and \quad \max C_{\mathbf{Q}_{\alpha}^{r}} = [w_{\mathrm{LT}}(\mathcal{T}_{\alpha}''), w_{1}(\operatorname{set}(\alpha^{\mathrm{r}}))]_{L}$
(b) $C_{\mathbf{Q}_{\alpha}^{r}} \cong (\mathsf{ST}(\mathfrak{G}^{x}), \leq) \text{ as posets.}$

Proof. (a) By the construction of the diagram \mathcal{G} , it is clear that \mathcal{G} is free of a strictly upper-right configuration. Therefore, by Theorem 4.16, we have the first assertion that

$$C_{\mathbf{Q}_{\alpha}^{r}} = \{ \Sigma_{L}(P_{T^{x}}) \mid T \in \mathsf{ST}(\mathcal{G}^{x}) \}$$

For the second assertion, let $\min C_{\mathbf{Q}_{\alpha}^{r}} = [\sigma_{0}, \rho_{0}]_{L}$. Since $\operatorname{set}(\mathsf{c}(\mathcal{G})) = \{1, 3, \ldots, 2\ell(\alpha) - 1, 2\ell(\alpha), \ldots, n-1\}$, it follows from Theorem 4.16 that

$$\sigma_0 = w_0(\operatorname{set}(\mathsf{c}(\mathcal{G}))^{\mathrm{c}}) = w_0(2[\ell(\alpha) - 1]).$$

For ρ_0 , Theorem 4.12 and Proposition 4.4 imply that $\rho_0 = w_{TL}(T_g)$, where T_g is the standard tableau on \mathcal{G} (see (4.2)). Writing $w_{TL}(T_g)$ explicitly, we have

$$w_{\mathrm{TL}}(T_{\mathfrak{G}}) = 1 \underbrace{\alpha_{1} + 1 \ 2}_{\mathrm{col.} \ 2} \cdots \underbrace{\mathbf{s}_{\ell(\alpha)-1}(\alpha) + 1 \ \mathbf{s}_{\ell(\alpha)-2}(\alpha) + 2}_{\mathrm{col.} \ \ell(\alpha)} \underbrace{\mathbf{s}_{\ell(\alpha)-1}(\alpha) + 2 \ \mathbf{s}_{\ell(\alpha)-1}(\alpha)}_{\mathrm{row} \ \ell(\alpha)} \underbrace{\mathbf{s}_{\ell(\alpha)-2}(\alpha) + 2 \ \cdots \ \mathbf{s}_{\ell(\alpha)-1}(\alpha)}_{\mathrm{row} \ \ell(\alpha)} \cdots \underbrace{3 \ 4 \ \cdots \ \alpha_{1}}_{\mathrm{row} \ 1}.$$

Here, $\mathbf{s}_j(\alpha) := \sum_{1 \le r \le j} \alpha_r$ for $1 \le j < \ell(\alpha)$. From the definitions of w_{dr} and \mathcal{T}_{α} , we observe that the entries in column i ($2 \le i \le \ell(\alpha)$) correspond to $\mathcal{T}_{\alpha}(1,i)$ and $\mathcal{T}_{\alpha}(2,i-1)$, while the entries in row i ($1 \le i \le \ell(\alpha)$) correspond to the entries in row i of \mathcal{T}_{α} . This gives that $w_{\mathrm{TL}}(T_{\mathfrak{g}}) = w_{\mathrm{dr}}(\mathcal{T}_{\alpha})$. Thus we have

$$\min C_{\mathbf{Q}_{\alpha}^{r}} = [w_0(2[\ell(\alpha) - 1]), \mathbf{w}_{\mathrm{dr}}(\mathcal{T}_{\alpha})]_L$$

For the third assertion, let $\max C_{\mathbf{Q}_{\alpha}^{r}} = [\sigma_{1}, \rho_{1}]_{L}$. By Theorem 4.12 and Proposition 4.4, we have $\sigma_{1} = w_{\mathrm{BL}}(T'_{\mathrm{g}})$, where T'_{g} is the standard tableau on \mathcal{G} . Explicitly, $w_{\mathrm{BL}}(T'_{\mathrm{g}})$ is written as

$$\underbrace{k \ k+1 \cdots k+\alpha_{\ell(\alpha)}-1}_{\operatorname{row} \ell(\alpha)} \underbrace{k-2 \ k-1 \ k+\alpha_{\ell(\alpha)} \ k+\alpha_{\ell(\alpha)}+1 \ \cdots}_{\operatorname{row} 1} \cdots \underbrace{1 \ 2 \ \cdots n}_{\operatorname{row} 1},$$

where $k := 2\ell(\alpha) - 1$. Observing the correspondence with \mathcal{T}''_{α} , we have that $w_{\mathrm{BL}}(T'_{\mathfrak{G}}) = w_{\mathrm{LT}}(\mathcal{T}''_{\alpha})$. For ρ_1 , since $\operatorname{set}(\mathsf{r}(\mathfrak{G})) = \{\alpha_{\ell(\alpha)}, \alpha_{\ell(\alpha)} + \alpha_{\ell(\alpha)-1}, \ldots, \alpha_{\ell(\alpha)} + \cdots + \alpha_1\}$, it follows from Theorem 4.16 that $\rho_1 = w_1(\operatorname{set}(\mathsf{r}(\mathfrak{G}))) = w_1(\operatorname{set}(\alpha^r))$. Thus, we have

$$\max C_{\mathbf{Q}_{\alpha}^{r}} = [\mathbf{w}_{\mathrm{LT}}(\mathcal{T}_{\alpha}''), w_{1}(\mathrm{set}(\alpha^{\mathrm{r}}))]_{L}$$

(b) By the construction of \mathcal{G} , it is clear that \mathcal{G} is free of a strictly upper-right configuration. Therefore, by Theorem 4.12 and Theorem 4.16, we have

$$(C_{\mathbf{Q}^r_{\alpha}}, \preceq) \cong (\mathsf{ST}(\mathcal{G}^x), \leq)$$
 as posets.

Example 5.23. Let $\alpha = (3, 2, 3, 1) \models 9$. Consider the tableaux

$$\mathcal{T}_{\alpha}^{"} = \begin{array}{|c|c|c|} \hline 7 \\ \hline 5 & 6 & 8 \\ \hline 3 & 4 \\ \hline 1 & 2 & 9 \end{array} \quad \text{and} \quad \mathcal{T}_{\alpha} = \begin{array}{|c|} \hline 9 \\ \hline 6 & 7 & 8 \\ \hline 4 & 5 \\ \hline 1 & 2 & 3 \end{array}$$

Using Theorem 5.22, we compute the following:

$$\min C_{\mathbf{Q}_{\alpha}^{r}} = [w_{0}(\{2, 4, 6\}), 142659783]_{L} \text{ and} \\ \max C_{\mathbf{Q}_{\alpha}^{r}} = [756834129, w_{1}(\{1, 4, 6\})]_{L}.$$

Thus, $C_{\mathbf{Q}_{\alpha}^{r}}$ is the equivalence class of $[756834129, w_{1}(\{1, 4, 6\})]_{L}$.

On the other hand, the diagram $D_{756834129;\{1,4,6\}}$ corresponds to the diagram \mathcal{G} shown in (5.11). Notably, \mathcal{G} is free of a strictly upper-right configuration. Given the standard fillings

$$F_{\mathcal{G}}^{\downarrow} = \begin{array}{c} 6 \\ 4 \\ 7 \\ 8 \\ 1 \\ 3 \\ 9 \end{array} \quad \text{and} \quad F_{\mathcal{G}}^{\rightarrow} = \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 7 \\ 8 \\ 9 \end{array}$$

it follows from Theorem 4.16 and Remark 4.2 that we confirm

min $C_{\mathbf{Q}_{\alpha}^{r}} = [132547689, 142659783]_{L}$ and max $C_{\mathbf{Q}_{\alpha}^{r}} = [756834129, 967845123]_{L}$.

To further explore $C_{\mathbf{Q}_{\alpha}^{r}}$, we note that $P_{F^{x}}$ corresponds to all the posets obtained from $F \in \mathsf{ST}(\mathfrak{G}^{x})$ by Theorem 5.22(b). By computational enumeration, it is found that there are precisely 594 elements in $\mathsf{ST}(\mathfrak{G}^{x})$.

We close this section by providing the projective cover and the injective hull of \mathbf{Q}_{α}^{r} .

Theorem 5.24. Let α be a peak composition of n and let $\mathcal{G} := D(\mathbf{Q}_{\alpha}^{r}, \mathbf{w}_{\mathrm{LT}})$.

- (a) $\overline{\mathsf{B}}([w_0(2[\ell(\alpha)-1]), w_1(\mathrm{Des}_{\mathrm{L}}(\mathsf{w}_{\mathrm{LB}}(F_{\mathrm{g}}^{\downarrow})))]_L)$ is a projective cover of \mathbf{Q}_{α}^r .
- (b) $\overline{\mathsf{B}}([w_0(\operatorname{set}(\alpha^{\mathrm{r}})), w_1(\operatorname{set}(\alpha^{\mathrm{r}}))]_L)$ is an injective hull of \mathbf{Q}_{α}^r . In particular, it is a projective indecomposable module.

Proof. Since \mathcal{G} is free of a strictly upper-right configuration, it follows from Corollary 4.19 that

$$\overline{\mathsf{B}}([w_0(\operatorname{set}(\mathsf{c}(\mathfrak{G}))^{\mathrm{c}}), w_1(\operatorname{Des}_{\mathrm{L}}(\mathrm{w}_{\mathrm{LB}}(F_{\mathfrak{G}}^{\downarrow})))]_L)$$

is a projective cover of \mathbf{Q}_{α}^{r} . Observing that

 $\operatorname{set}(\mathsf{c}(\mathfrak{G})) = \{1, 3, \dots, 2\ell(\alpha) - 1, 2\ell(\alpha), \dots, n-1\},\$

we deduce the assertion (a).

Similarly, using Corollary 4.19, we also find that

 $\overline{\mathsf{B}}([w_0(\mathrm{Des}_{\mathrm{L}}(\mathrm{w}_{\mathrm{BL}}(F_{\mathfrak{G}}^{\rightarrow}))), w_1(\mathrm{set}(\mathsf{r}(\mathfrak{G})))]_L)$

is an injective hull of \mathbf{Q}_{α}^{r} . Since

 $\mathrm{Des}_{\mathrm{L}}(\mathrm{w}_{\mathrm{BL}}(F_{\mathfrak{G}}^{\rightarrow})) = \{\alpha_{\ell(\alpha)}, \alpha_{\ell(\alpha)} + \alpha_{\ell(\alpha)-1}, \dots, \alpha_{\ell(\alpha)} + \dots + \alpha_{2}\} = \mathrm{set}(\mathsf{r}(\mathfrak{G})),$

the assertion (b) follows. Moreover, this injective hull coincides with a projective indecomposable module. $\hfill \Box$

As an important consequence of Theorem 5.24(b), we derive the following corollary.

Corollary 5.25. Let α be a peak composition of n. Then the $H_n(0)$ -module \mathbf{Q}_{α}^r is indecomposable.

References

- [1] S. Assaf and D. Searles. Kohnert polynomials. *Experiment. Math.*, pages 1–27, 2019.
- [2] J.-C. Aval, K. Djenabou, and P. R. W. McNamara. Quasisymmetric functions distinguishing trees. Algebraic Combinatorics, 6(3):595–614, 2023.
- [3] J. Bardwell and D. Searles. 0-Hecke modules for Young row-strict quasisymmetric Schur functions. European Journal of Combinatorics, 102:103494, 2022.
- [4] J. Bardwell and D. Searles. Weak Bruhat interval modules of finite-type 0-Hecke algebras and projective covers. arXiv preprint arXiv:2311.10068, 2023.
- [5] D. J. Benson. Representations and cohomology. I, volume 30 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1991. Basic representation theory of finite groups and associative algebras.
- [6] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki. A lift of the Schur and Hall–Littlewood bases to non-commutative symmetric functions. *Canad. J. Math.*, 66(3):525–565, 2014.
- [7] C. Berg, N. Bergeron, F. Saliola, L. Serrano, and M. Zabrocki. Indecomposable modules for the dual immaculate basis of quasisymmetric functions. *Proc. Amer. Math. Soc.*, 143(3):991–1000, 2015.
- [8] A. Björner and M. L. Wachs. Generalized quotients in Coxeter groups. Trans. Amer. Math. Soc., 308(1):1–37, 1988.
- [9] A. Björner and M. L. Wachs. Permutation statistics and linear extensions of posets. J. Combin. Theory Ser. A, 58(1):85–114, 1991.
- [10] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. Homological properties of 0-Hecke modules for dual immaculate quasisymmetric functions. *Forum Math. Sigma*, 10:e91, 2022.
- [11] S.-I. Choi, Y.-H. Kim, S.-Y. Nam, and Y.-T. Oh. The projective cover of tableau-cyclic indecomposable H_n(0)-modules. Trans. Amer. Math. Soc., 375(11):7747–7782, 2022.
- [12] S.-I. Choi, Y.-H. Kim, and Y.-T. Oh. Poset modules of the 0-Hecke algebras and related quasisymmetric power sum expansions. *European Journal of Combinatorics*, 120:103965, 2024.
- [13] S.-I. Choi, S.-Y. Nam, and Y.-T. Oh. Quasisymmetric Schur Q-functions and peak Young quasisymmetric Schur functions. arXiv preprint, arXiv:2405.05867v1 [math.CO], 2024.
- [14] G. Duchamp, F. Hivert, and J.-Y. Thibon. Noncommutative symmetric functions. VI. Free quasisymmetric functions and related algebras. *Internat. J. Algebra Comput.*, 12(5):671–717, 2002.
- [15] M. Fayers. 0-Hecke algebras of finite Coxeter groups. J. Pure Appl. Algebra, 199(1-3):27-41, 2005.
- [16] J. Huang. A tableau approach to the representation theory of 0-Hecke algebras. Ann. Comb., 20(4):831–868, 2016.
- [17] W.-S. Jung, Y.-H. Kim, S.-Y. Lee, and Y.-T. Oh. Weak Bruhat interval modules of the 0-Hecke algebra. Math. Z., 301(4):3755–3786, 2022.

- [18] Y.-H. Kim, S.-Y. Lee, and Y.-T. Oh. Regular Schur labeled skew shape posets and their 0-Hecke modules. *Forum Math. Sigma*, 12:e1110, 2024.
- [19] D. Krob and J.-Y. Thibon. Noncommutative symmetric functions. IV. Quantum linear groups and Hecke algebras at q = 0. J. Algebraic Combin., 6(4):339–376, 1997.
- [20] R. I. Liu and M. Weselcouch. P-partition generating function equivalence of naturally labeled posets. J. Combin. Theory Ser. A, 170:105136, 31, 2020.
- [21] K. Luoto, S. Mykytiuk, and S. van Willigenburg. An introduction to quasisymmetric Schur functions. SpringerBriefs in Mathematics. Springer, New York, 2013.
- [22] S. K. Mason and E. Niese. Skew row-strict quasisymmetric Schur functions. J. Algebraic Combin., 42(3):763–791, 2015.
- [23] P. R. McNamara and R. E. Ward. Equality of P-partition generating functions. Ann. Comb., 18(3):489–514, 2014.
- [24] P. R. W. McNamara and S. van Willigenburg. Towards a combinatorial classification of skew Schur functions. Trans. Amer. Math. Soc., 361(8):4437–4470, 2009.
- [25] E. Niese, S. Sundaram, S. van Willigenburg, J. Vega, and S. Wang. Row-strict dual immaculate functions. Adv. Appl. Math., 149:102540, 2023.
- [26] E. Niese, S. Sundaram, S. van Willigenburg, J. Vega, and S. Wang. 0-Hecke modules for row-strict dual immaculate functions. *Trans. Amer. Math. Soc.*, 377(4):2525–2582, 2024.
- [27] P. N. Norton. 0-Hecke algebras. J. Austral. Math. Soc. Ser. A, 27(3):337–357, 1979.
- [28] V. Reiner, K. M. Shaw, and S. van Willigenburg. Coincidences among skew Schur functions. Adv. Math., 216(1):118–152, 2007.
- [29] D. Searles. Indecomposable 0-Hecke modules for extended Schur functions. Proc. Amer. Math. Soc., 148(5):1933–1943, 2020.
- [30] D. Searles. Diagram supermodules for 0-Hecke–Clifford algebras. arXiv preprint, arXiv:2202.12022 [math.RT], 2022.
- [31] R. Stanley. Enumerative combinatorics. Vol. 2, volume 62 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999.
- [32] R. P. Stanley. Ordered structures and partitions, volume 119. American Mathematical Soc., 1972.
- [33] H. Yang and H. Yu. Classification of weak Bruhat interval modules of 0-Hecke algebras. arXiv preprint, arXiv:2410.07990v1 [math.RT], 2024.

CENTER FOR QUANTUM STRUCTURES IN MODULES AND SPACES, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, REPUBLIC OF KOREA

Email address: ignatioschoi@snu.ac.kr

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, REPUBLIC OF KOREA *Email address*: synam.math@gmail.com

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, REPUBLIC OF KOREA *Email address*: ytoh@sogang.ac.kr