Non-topological Solitons of Dark Matter and Gravitational Wave Signals

Alexander Libanov^{1, *}

¹Institute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary Prospect 7a, Moscow 117312, Russia

(Dated: December 13, 2024)

Within the framework of General Relativity, it can be shown that gravitational waves are radiated with the merger of massive compact objects. Such gravitational wave signals are observed on Earth on various detectors, in particular, on Laser Interferometer Gravitational Wave Observatory (LIGO) and Virgo. During the operation of these detectors, many events have been detected. Those events are associated with the merger of massive compact objects, however, the nature of some merging objects has not yet been reliably established. This work considers non-topological solitons of dark matter – Q-balls, as candidates for the role of massive compact objects. In this work one of the simplest models of Q-balls, the mechanism of their birth during a phase transition in the early Universe and the mechanism of their mass gaining during the evolution of the Universe, which is based on their mutual merger are considered. As a result, it is analyzed whether Q-balls of dark matter can be candidates for the role of massive compact objects.

I. INTRODUCTION

It follows from the General Theory of Relativity that gravitational waves can be radiated during the merger of massive compact objects [1-6]. Such objects are usually claimed by neutron stars or black holes [7]. Nevertheless, with the more and more new signals, unusual data began to arise [2-4, 8-10].

On August 14th, 2019, at 21:10:39 UTC, GW190814 was detected using LIGO and Virgo. GW190814 is a gravitational wave event received from the merger of two massive compact objects with masses of $22.3 - 24.3 M_{\odot}$ and $2.50 - 2.67 \,\mathrm{M_{\odot}}$ respectively [4]. This event indicates that the smaller object is located at the mass boundary between a light black hole and a massive neutron star [11]. It is noteworthy that no electromagnetic signal was received from this event [12]. There are still discussions about the nature of the second object. One possible scenario is a merger of a black hole and a neutron star. In this case strong restrictions are imposed on the radius of the neutron star. The other possible scenario is a merger of two black holes [9], [13]. However, there are models that suggest that the smaller object is exotic: a gravastar, a quark star, or a boson star [14-17].

On January 5, 2020, at 16:24:26 UTC and on January 15, 2020 at 4:23:10 UTC, signals GW200105 and GW200115 were received using LIGO and Virgo, respectively. GW200105 received as a result of rotation and merger of a binary system, the masses of objects in which are $8.9^{+1.2}_{-1.5}$ M_{\odot} and $1.9^{+0.3}_{-0.2}$ M_{\odot}, and GW200115 is a result of rotation and merger of a binary system, the masses of objects in which are 5.7^{+1.8}_{-2.1} M_{\odot} and $1.5^{+0.7}_{-0.3}$ M_{\odot}. The masses of the smaller objects are very different from the mass of the second component of the system, from which the GW190814 signal was received. Moreover, these events were again not accompanied by

an electromagnetic signal. Hypothetically, these objects could be dark matter stars [11], [18].

The astrophysical understanding of stellar evolution suggests that stellar-mass black holes may not have masses less than $\sim 5 \ M_{\odot}$ and neutron stars are expected to have a maximum mass of $\sim 3 M_{\odot}$ [19–24]. It is also worth noting that LIGO and Virgo distinguish objects primarily by their masses that is why the nature of merging objects is difficult to determine only by the signature of gravitational waves. These lead to the fact that there is a mass gap between heavy neutron stars and light black holes, equal to $\sim 3-5 M_{\odot}$ [8], [10]. Thus, the problem arises of explaining some events like GW190814, GW200105 and GW200115, since some objects in such systems that generate gravitational wave signals lie in the mass gap for black holes or violate the equation of state of neutron stars. Moreover, there is still a problem of dark matter in cosmology, since there are a sufficient number of candidates that have not yet been experimentally confirmed [25-27].

Some works suggest that dark matter stars may be components of binary systems of compact objects [28-30]. According to this possible solution to the problems of unusual gravitational wave signals, mass-gap and cold dark matter is considered in this work. The model of non-topological solitons – Q-balls of dark matter – may be a candidate for the components of such binary systems. Firstly, using this model, one can try to explain the unusual gravitational wave signals received by LIGO and Virgo, without modifying the equation of state of neutron stars. Secondly, Q-balls may be candidates for the role of cold dark matter [31]. Thus, the work considers the parameters of Q-balls, their mass and radius, the mechanism of formation of cosmological Q-balls, simple mechanisms for increasing their mass due to merger with other cosmological Q-balls is proposed.

^{*} e-mail: libanov.am18@physics.msu.ru

II. PHYSICS OF Q-BALLS

A. Q-ball parameters

The Q-ball is a non-topological soliton. There are many different models that allow the existence of Q-balls [32], [33]. This work considers one of the simplest models of Q-balls – the Friedberg-Lee-Sirlin theory. This theory consists of two scalar fields, one of which is real and the other is complex, and the reason for the existence and stability of the Q-ball is the presence of a charge in the Q-ball. In the Friedberg-Lee-Sirlin theory, it is assumed that particles of the χ field acquire mass during interaction with some additional scalar field φ [34–36]. The Lagrangian of Friedberg-Lee-Sirlin theory is

$$\mathcal{L} = \frac{1}{2} (\partial_{\mu} \varphi)^2 - U(\varphi) + (\partial_{\mu} \chi)^* \partial_{\mu} \chi - k^2 \varphi^2 \chi^* \chi, \quad (1)$$
$$U(\varphi) = (\varphi^2 - v^2)^2,$$

where χ is a complex scalar field, φ is real scalar field, $k = h/\lambda^{1/4}$ in notations of Ref. [36] and v are some constants [34]. The mass of the field χ in a vacuum is

$$m_{\chi} = kv, \tag{2}$$

This theory has U(1)-symmetry:

$$\chi \to e^{i\alpha}\chi$$

and, therefore, there is a conserved charge Q. The state with the lowest energy with a sufficiently large Q is a spherical Q-ball with conditions:

$$\begin{cases} \varphi = 0, \ x < R, \\ \varphi = v, \ x > R, \end{cases}$$

where R is the radius of the Q-ball.

In the approximation of a sufficiently large Q, the radius R and the energy E of the Q-ball are determined by the balance of the energy of the massless field χ located in the potential well and the potential energy of the massive field φ inside. Then the energy is

$$E(R) = \frac{\pi Q}{R} + \frac{4\pi}{3}R^{3}U_{0},$$
(3)

where

$$U_0 = U(0) - U(v) = v^4.$$

By differentiating (3) by R and equating the derivative to zero, which is the condition for the minimum energy, an expression for the radius of the Q-ball is obtained

$$R_Q = \left(\frac{Q}{4}\right)^{1/4} \frac{1}{v}.\tag{4}$$

Now, if we substitute (4) into (3), we can obtain an expression for the mass of the Q-ball

$$m_Q = \frac{4\sqrt{2}\pi}{3} v Q^{3/4},\tag{5}$$

assuming that the mass of the Q-ball corresponds to the minimum energy of the Q-ball [34], [36].

It is worth noting that there is a stability condition for the Q-ball, which gives a limit on the minimum charge:

$$Q_{min} = \frac{m_Q}{m_\chi}.$$
 (6)

B. First order phase transition

It is assumed that the early expanding flat Universe is uniformly filled with particles of the fields φ and χ , and cosmological Q-balls are born as a result of the Ist order phase transition, which occurs due to the cooling of the Universe to some critical temperature T_c , and $T_c \sim v$. During the phase transition, bubbles of a new phase are formed ($\varphi = \varphi_c$), which, in turn, merge. Thus, in some volume there remains one bubble of the old phase ($\varphi = 0$), which collects particles of the χ field under the assumption of sufficient massiveness of the particles of the χ field, and which is the future cosmological Qball (Fig. 1). This phase transition model does not im-



FIG. 1. A schematic representation of the I-st order phase transition in the early Universe. The white area is the area of the old phase ($\varphi = 0$), the blue area is the area of the new phase ($\varphi = \varphi_c$) [36]. As can be seen from the figure, at some point in time, one area of the old phase remains in the allocated volume, which, for simplicity, is considered spherical within the framework of this work.

pose special restrictions on the area of appearance of new phase bubbles during the phase transition, so, theoretically, new phase bubbles can appear in the remaining bubbles of the old phase, therefore, it is assumed that a cosmological Q-ball is formed if and only if a new phase bubble does not form in the remaining bubble of the old phase [37], [38].

The physics of the phase transition imposes a restriction on the charge of the cosmological Q-ball from above. In the framework of the work, dark matter is represented by particles of the χ field, and there is an asymmetry [36]

$$\frac{n_{\chi} - n_{\bar{\chi}}}{s} = \frac{n_Q Q}{s} = \eta_{\chi},\tag{7}$$

where n_{χ} is the concentration of particles of the field χ , $n_{\bar{\chi}}$ is the concentration of antiparticles of the field χ , n_Q is the concentration of Q-balls in the Universe, s is the density of entropy, and $\eta_{\chi} \leq 1$ is the parameter responsible for the asymmetry of the particles of the field χ . This parameter is determined by the dynamics at earlier stages in interactions with other fields, but in the realities of this work it will be considered free [39], [40].

It is possible to estimate the volume of the remaining bubble of old phase, and hence estimate the volume in which the χ field particles are collected, using the birth rate of new phase bubbles and taking into account the probability of the birth of a new phase bubble in a remaining bubble of the old phase during a phase transition (i.e. estimate volume from which the cosmological Q-ball is born):

$$V_{\star} = \xi \left(\frac{uA^{1/2}M_{pl}^{*}}{T_{c}^{2}L^{3/2}}\right)^{3},$$
(8)

It is worth noting that the charge of the cosmological Qball is determined by the number of charged particles of the χ field located in volume V_{\star} , therefore, it is possible to estimate the charge of the cosmological Q-ball:

$$Q_{\star} = \eta_{\chi} \xi \frac{2\pi^2 g_{\star}}{45} \left(\frac{u A^{1/2} M_{pl}^*}{L^{3/2} T_c} \right)^3, \qquad (9)$$

where $T_c \sim v$, $M_{Pl}^* = M_{Pl}/(1.66\sqrt{g_*})$, $g_* \sim 100$ is the effective number of degrees of freedom at the phase transition temperature, u – the velocity of the bubble walls of the new phase during the phase transition; $\xi \sim 1$, $L \sim 100$, $A \sim 1$ – constants related to the calculation method (for more detailed calculations and physics see Ref. [36]). Thus, the charge of the cosmological Q-ball lies within

$$Q_{min} < Q < Q_{\star},\tag{10}$$

where Q_{min} is defined by (6) and Q_{\star} is defined by (9).

C. Distribution of Q-balls according to their charges

In the previous section, it was assumed that all cosmological Q-balls are born with a charge of Q_{\star} . Now it is proposed to find the distribution of such Q-balls by their charges, taking into account (10). To do this, we need to find the probability of the birth of a cosmological Qball with a charge greater than some \bar{Q} . Obviously, the probability of a new phase bubble appearing in a bubble of the old phase of volume V is proportional to the volume of the considered area and the characteristic time of collapse of the bubble, since the longer the time and volume, the greater the chance of formation of a new phase bubble in the area [41].

$$F_b = V\Gamma \frac{R}{u}.$$
 (11)

Since it is assumed that a cosmological Q-ball is born from a bubble of the old phase before a bubble of the new one appears in it, the following estimation can be given by

$$V_{\star}\Gamma\frac{R_{\star}}{u} \sim 1. \tag{12}$$

This estimation takes place, because when the maximum volume of the considered area of the old phase is reached, the probability of a bubble of a new phase appearing in it tends to unity.

Next, we need to find the probability of the birth of a Q-ball with a charge greater than \bar{Q} . From the definition of probability

$$F = 1 - F_b. \tag{13}$$

This expression can be divided by (12):

$$F = 1 - \frac{V\Gamma R/u}{V_{\star}\Gamma R_{\star}/u}.$$
(14)

Let the particles of the field χ be evenly distributed throughout the early Universe, then $Q \sim V$, and therefore (14) takes the form

$$F = 1 - \left(\frac{Q}{Q_{\star}}\right)^{4/3} = \int \frac{dP}{dQ} dQ, \qquad (15)$$

where dP/dQ is the probability of the birth of a Q-ball in the range from Q to Q + dQ. To find this probability, it is necessary to consider the probability of the birth of a Q-ball with a charge greater than \bar{Q} and with a charge of $\bar{Q} + dQ$ and find their difference

$$F(\bar{Q} + dQ) - F(\bar{Q}) = \int_{\bar{Q} + dQ}^{Q_{\star}} \frac{dP}{dQ} dQ - \int_{\bar{Q}}^{Q_{\star}} \frac{dP}{dQ} dQ.$$

Hence, the distribution of cosmological Q-balls by their charges is

$$\frac{dP}{dQ} = -\frac{F(\bar{Q} + dQ) - F\bar{Q}}{dQ} = -\frac{dF}{dQ}.$$
 (16)

Thus,

$$n(Q) \sim \alpha \int_{Q_{min}}^{Q} \frac{dP}{dQ} dQ, \qquad (17)$$

where α is determined from the normalization condition

$$\alpha \int_{Q_{min}}^{Q_{\star}} Q \frac{dP}{dQ} dQ = Q_{\star}$$

and is equal to 7/4. Here and further, it is assumed that $Q_{min} \ll Q_{\star}$. Now we can get the final expression for the distribution (17)

$$n(Q) \sim \frac{7}{4} \left(\frac{Q}{Q_{\star}}\right)^{4/3}.$$
(18)

III. ESTIMATION OF THE POTENTIAL PARAMETER

In this section, the Lagrangian (1) parameter v will be estimated. As discussed above, Q-balls are dark matter, and therefore it is necessary to satisfy certain conditions.

Firstly, the mean cross-section of the interaction of bulk dark matter should not exceed $1 \text{ cm}^2/\text{g}$ [41–45].

$$\langle \bar{\sigma} \rangle_b = \bar{\sigma}_\star \int_0^1 \frac{x^{-1/4} x^{3/4} (1-x)}{x^{3/4} (1-x)} dx \approx 1.3 \bar{\sigma}_\star \lesssim 1 \text{ cm}^2/\text{g},$$
(10)

where $x = Q/Q_{\star}, \, \bar{\sigma}_{\star} = \bar{\sigma}(Q_{\star}).$

The mean cross-section is

$$\bar{\sigma}(Q) = \frac{\pi R_Q^2}{m_Q} = \frac{3}{8\sqrt{2}} v^{-3} Q^{-1/4}, \qquad (20)$$

Then from (19), taking into account (9) and (20), we can get the lower limit for v

$$v_{min} \gtrsim \frac{1.07 \cdot 10^{-7} u^{2/3}}{\eta_{\chi}^{1/9}} \text{ GeV.}$$
 (21)

Secondly, the present energy density of Q-balls should not exceed the present energy density of dark matter. The energy density of Q-balls obtained using (5) and (18)

$$\rho = \int_0^{Q_\star} m_Q dn(Q) \sim Q^{25/12}.$$
 (22)

This function has a maximum at the upper limit of integration Q_{\star} , therefore, for a rough estimate, it is assumed that cosmological Q-balls with Q_{\star} charge dominate in dark matter.

There is a connection between the energy density of Q-balls and the cross-section, given in [41]. Using this ratio, we can find the upper limit on v

$$\rho_{DM} \gtrsim \frac{4\sqrt{2}\pi}{3} v \cdot Q_{\star}^{-1/4} \eta_{\chi} s_0, \qquad (23)$$

where $\rho_{DM} = 10^{-6} \text{ GeV/cm}^3$ is the modern density of dark matter, $s_0 = 3 \cdot 10^3 \text{ cm}^{-3}$ is the modern density of entropy. Thus, the upper limit on v is

$$v_{max} \lesssim \frac{5.6 \cdot u^{3/7}}{\eta_{\chi}^{3/7}} \text{ GeV.}$$
 (24)

Dependence of the maximum and minimum values of von the asymmetry is shown in the Fig. 2. In some works it is proposed to estimate the asymmetry of the particles of the field χ as a quark asymmetry $3\Delta_B \approx 2.7 \cdot 10^{-10}$, and the velocity of cosmological Q-balls as $u \sim 1$. In the case of such estimations, v is [36]

$$1.2 \text{ keV} \lesssim v \lesssim 70.4 \text{ TeV}.$$
 (25)

Now, accordingly, within the framework of this estimation, it is possible to find restrictions on the mass of cosmological Q-balls

$$3 \cdot 10^{-5} \text{ kg } \lesssim m_{\star} \lesssim 8 \cdot 10^{8} \text{ kg.}$$
 (26)



FIG. 2. Dependence of cosmological constraints on the parameter v of the Lagrangian (1) on the magnitude of the asymmetry of the particles of the field $\chi \eta_{\chi}$ in the case of the velocity of cosmological Q-balls u = 1 and $T_c = v$. v_{min} is the lower limit and is represented in keV (blue dashed line), v_{max} is the upper limit and is represented in GeV (red line).

IV. Q-BALLS MERGING

It follows from (26) that cosmological Q-balls cannot claim to be exotic objects of stellar mass, therefore it is necessary to consider their inelastic interaction, which will occur with a probability of ~ 50% [46], [47].

A. Flat expanding Universe

In this model, it is assumed that after the phase transition shown in Fig. 1, the early flat expanding Universe is uniformly filled with cosmological Q-balls with the same charges (9) and the same velocities u. It is proposed to select one cosmological Q-ball, which will absorb other cosmological Q-balls during the evolution of the Universe, and based on this simple mechanism to obtain an estimation of the mass of the present Q-ball.

At the initial stage, the velocity of the cosmological Qball (it is also the velocity of the wall of the region from which the cosmological Q-ball is born during the phase transition) u, the asymmetry η_{χ} of the particles of the field χ , the Lagrangian parameter v and the phase transition temperature $T_c \sim v$ are considered free parameters limited only by simple physical considerations. Thus, the velocity of the cosmological Q-ball u should not exceed the speed of light

the asymmetry of the χ field is limited as follows from the definition of asymmetry (7)

$$0 \lesssim \eta_{\chi} \lesssim 1;$$

The parameter v, in addition to the restrictions shown in Fig. 2, is limited from above by the temperature of the Planck epoch, and from below by the temperature of the present Universe. This is also true for the phase transition temperature T_c [48].

$$10^{18 \div 19} \text{ GeV} \gtrsim v \sim T_c \gtrsim 2.3 \times 10^{-13} \text{ GeV} \approx 2.7 \text{ K}.$$

Firstly, the model assumes that the early flat expanding Universe is uniformly filled with cosmological Q-balls, which are born in bubbles of a new phase of the volume V_{\star} , which, in turn, is determined by the expression (8). Due to this fact at the initial stage there is on average one cosmological Q-ball in this volume. Secondly, due to the evolution of the Universe, the concentration of Q-balls is not a constant value (here and further, the concentration change from the interaction of the selected Q-ball with cosmological Q-balls is considered insignificant). To account for changes in concentration due to the expansion of the Universe, it is necessary to introduce a scale factor, which is determined from the Friedman equation and is equal to

$$a = a_0 \left(\frac{\Omega_M}{\Omega_\Lambda}\right)^{1/3} (\sinh[\frac{3}{2}\sqrt{\Omega_\Lambda}H_0t])^{2/3}, \qquad (27)$$

where $\Omega_M = 0.24$, $\Omega_{\Lambda} = 0.76$, $H_0 = 2.2 \times 10^{-18} \text{ s}^{-1}$ [38], [49]. Normalization of the scale factor a_0 is determined from the condition

$$a(t_0) = 1$$

where $t_0 = 14$ Gyr is the present epoch of the Universe. Then, we can find the concentration of cosmological Qballs in the universe:

$$n_{\star} = \frac{1}{V_{\star}a^3(t)}.$$
 (28)

It is assumed that cosmological Q-balls with velocities of the order of u merge with a selected Q-ball, therefore their interaction is determined by the geometric crosssection of the selected Q-ball, which is defined as follows

$$\sigma = \frac{\pi R_Q^2(t)}{2},\tag{29}$$

where $R_Q(t)$ is determined from the equation (4) and obviously grows over time due to the absorption of the selected Q-ball of cosmological Q-balls. The probability of an inelastic interaction of bulk dark matter is taken into account here, equal to ~ 50% [46], [47].

Now we can make an equation describing change of the charge (and, accordingly, mass) during the interaction of the selected Q-ball with cosmological Q-balls. Let the selected cosmological Q-ball absorb charge ΔQ in time Δt . Obviously, the greater the concentration n of cosmological Q-balls in a given region and the larger the geometric cross-section of the interaction σ , the selected Q-ball absorbs the greater charge

$$\Delta Q = \sigma n u \Delta t, \tag{30}$$

where u is the velocity of the cosmological Q-ball, equal to the velocity of the walls during collapse. Now we need to combine the equations (28) and (29) and make the limit transition

$$\Delta Q \to 0, \ \Delta t \to 0.$$

Thus, the law of charge changing of the selected Q-ball is obtained during the absorption of cosmological Q-balls by it in time

$$\dot{Q} = Q_{\star}\sigma(Q)u\frac{1}{V_{\star}a^3(t)},\tag{31}$$

where $\sigma(Q)$ is the geometric cross-section of the interaction of Q-balls, determined by the equation (29), a(t)is the scale factor determined by the equation (27). For the correct formulation of the mathematical problem, it is necessary to add one boundary condition, which is determined from a simple physical consideration: at the time when the phase transition occurred $t_c \sim 0$ (hereafter, for simplicity, it is assumed that the phase transition occurs in the early Universe), the charge of the selected Q-ball

$$Q(t_c) = Q_\star. \tag{32}$$

Combining (31) and (32), we can get a correct mathematical problem

$$\begin{cases} \dot{Q} = uQ_{\star}\sigma(Q)n_{\star}, \\ Q(t_c) = Q_{\star}. \end{cases}$$
(33)

The solution, taking into account the boundary condition, is

$$Q(t) = \left(-\frac{\pi Q_{\star} u \sqrt{\Omega_{\Lambda}}}{12 v^2 V_{\star} a_0^3 \Omega_M H_0} \coth\left[\frac{3}{2}\sqrt{\Omega_{\Lambda}}H_0 t\right] + C\right)^2,$$
(34)

where C is the integration constant equal to

$$C = \frac{1}{2} \left(2\sqrt{Q_{\star}} + \frac{\pi\sqrt{\Omega_{\Lambda}}Q_{\star}u \coth\left[\frac{3}{2}\sqrt{\Omega_{\Lambda}}H_{0}t_{c}\right]}{6v^{2}V_{\star}a_{0}^{3}\Omega_{M}H_{0}} \right), \quad (35)$$

where t_c is the time of the phase transition, and Q_{\star} is defined by the expression (9), m_{\star} is defined by the expression (5) with substitution into it (9) and V_{\star} is defined by the expression (8).

In any physically meaningful configuration of the free parameters v, u, T_c and η_{χ} the solution (34) is almost time-independent, and therefore, taking into account (5), it is impossible to obtain the selected Q-ball of stellar mass, and free cosmological Q-balls in the case of a flat expanding Universe almost do not interact.

B. Galaxies

As it was shown in the previous section, free cosmological Q-balls almost do not interact with the selected Q-ball, and, as a result, the selected Q-ball almost does not change its mass during the evolution of the Universe. In fact, this is primarily due to the presence of a scale factor (28), which strongly suppresses the interaction of cosmological Q-balls with a selected Q-ball.

It is known that the expansion of the Universe does not affect gravitationally bound structures in the Universe, for example, galaxies. However, it is known that galaxies are formed primarily from dark matter [50]. Let all dark matter be represented by cosmological Q-balls. In this case, it can be assumed that after the phase transition, cosmological Q-balls participate in the formation of galaxies and, therefore, are in the gravitational potential of galaxies (the mechanisms of galaxy formation and cosmological Q-balls entering the gravitational potential of galaxies are not discussed in detail in this work) [51]. This model assumption allows us to get rid of the scale factor (28) and, as a result, locally get rid of the suppression of the interaction of cosmological Q-balls with the selected Q-ball. Further, being in the gravitational potential of the galaxy, cosmological Q-balls begin to be absorbed by the selected Q-ball, which, in turn, is expected to acquire a mass of the order of the mass of the Sun, so that such Q-balls can fill the mass gap between light black holes and massive neutron stars.

Now it is necessary to discuss in detail the formulation of the law of interaction of cosmological Q-balls with a selected Q-ball within the framework of a model of their interaction in the gravitational potential of galaxies, which will be constructed by analogy with the law (33). Firstly, we need to change the concentration (28). In addition to the absence of a scale factor, the volume in which one cosmological Q-ball is located also changes. This is due to the fact that dark matter in galaxies is unevenly distributed. In this work, it is assumed that cosmological Q-balls of dark matter obey the Navarro–Frank-White profile. This profile is

$$\rho(r) = \frac{\rho_0}{\frac{r}{R_s}(1 + \frac{r}{R_s})^2},$$
(36)

where $\rho_0 = 2 \text{ GeV/cm}^3$, $R_s = 8.5 \text{ kpc}$, r is the distance from the center of the galaxy (hereafter the characteristic parameters for the Milky Way are used, since this is the most studied galaxy) [52]. Then the concentration is

$$n(r) = \frac{\rho(r)}{m_{\star}}.$$
(37)

Secondly, Q-balls are located in the gravitational field of the galaxy and represent cold dark matter. This imposes some restrictions on the velocity of the Q-balls (but not on the velocity of the walls during the phase transition!). The characteristic orbital velocity of stars in the Milky Way $u_{\star} \approx 220$ km/s. A natural assumption is made here that Q-balls in the gravitational potential of the galaxy move at velocities characteristic of stars in this potential.

Thirdly, due to the fact that the phase transition occurs due to the cooling of the Universe, additional restrictions appear on the Lagrangian parameter v and the temperature of the phase transition T_c , due to the fact that the phase transition must occur before the formation of the first galaxies. Since the first galaxies appear at time $t \sim 100$ Myr $\div 1$ Gyr, the following restriction can be set

$$10^{18 \div 19} \text{ GeV} \gtrsim v \sim T_c \gtrsim 10^{-12} \text{ GeV},$$
 (38)

where the lower limit corresponds to the temperature of the Universe at time $t \sim 1$ Gyr [38].

Now we can start setting a mathematical problem in the same way as in (33). The cross-section does not depend on model assumptions, therefore it has the form (29). The characteristic velocity of Q-balls in a galaxy is equal, as discussed above, to the characteristic orbital velocity of stars in the gravitational potential of galaxies $u_{\star} \approx 220$ km/s. The concentration is given by the expression (37), taking into account the density distribution of dark matter in the galaxy (36). Due to the fact that cosmological Q-balls are not absorbed by the selected Q-ball before they enter the gravitational potential of the galaxy, it is convenient to solve the equation within $t \in [0; 13]$ Gyr, where it is taken into account that the first galaxies appear at time 1 Gyr. Thus, a differential equation similar to (33) is

$$\begin{cases} \dot{Q} = Q_{\star}\sigma(Q)u_{\star}n, \ t \in [0;13] \text{ Gyr}, \\ Q(0) = Q_{\star}. \end{cases}$$
(39)

This equation is easily solved by separating the variables, and the solution takes the form

$$Q(t,r) = \left(\frac{Q_{\star}\pi u_{\star}\rho(r)}{8v^2m_{\star}}(t+\sqrt{Q_{\star}})\right)^2,$$
 (40)

where Q_{\star} is defined by the expression (9), m_{\star} is defined by the expression (5) with substitution (9) into it.

V. SOLUTION ANALYSIS

In this section, we analyze the solution (40) obtained within the framework of the model of merging cosmological Q-balls with a selected Q-ball in the gravitational potential of the galaxy.

To obtain the total mass of the selected Q-ball, the solution (40) must be substituted into the expression for the mass (5). Thus, the final expression for the mass is

$$m_Q(v, u, \eta_{\chi}, u_{\star}, T_c, r) =$$

$$= \frac{\sqrt{3}\pi^{5/2}}{32 \cdot 2^{3/4}} v^{-7/2} Q_{\star}^{3/4} u_{\star}^{3/2} \rho^{3/2} (t + \sqrt{Q_{\star}})^{3/2}, \qquad (41)$$

where is $Q_{\star} = Q_{\star}(v, u, \eta_{\chi}, T_c)$ is defined by the expression (9), and $\rho = \rho(r)$ from the expression (36). The main motivation of this work is to try to explain gravitational wave signals, therefore it is necessary to satisfy the following limitation

$$m_Q(v, u, \eta_\chi, u_\star, T_c, r) \gtrsim 1 M_\odot. \tag{42}$$

As a first approximation, there are many sets of free parameters that satisfy this restriction, however, the work presents several restrictions on free parameters, in particular (19), (23) and (38), which must be satisfied. In this case, the set of free parameters is somewhat narrowed, but, nevertheless, suitable solutions still exist. The following values can be called the most successful set of parameters

$$\begin{cases} v \approx 7.5 \times 10^{-9} \text{ GeV}, \\ u = 1, \\ \eta_{\chi} = 1, \\ u_{\star} = 0.0007, \\ T_c \approx 1.5 \times 10^{-11} \text{ GeV}. \end{cases}$$
(43)

It can be seen that although there is some difference between v and T_c , which, nevertheless, can be explained by inaccuracies of the model, the restriction (38) is satisfied, and forging of Q-balls occurs at the time $t_c = 10 \text{ s} \div 3 \times 10^5$ years since the Big Bang (this corresponds to the photon epoch) [38].

The mass limit (42) is satisfied. From the form of (41) taking into account (36) and (43), it can be understood that the further away the selected Q-ball is from the center of the galaxy, the smaller its mass. For example, in the case of r = 0.05 kpc, the mass is

$$m_Q \Big|_{r=0.05 \text{ kpc}} \approx 1 \text{ M}_{\odot}.$$
 (44)

The Fig. 3 demonstrates the evolution of mass as a function of time t and distance from the center of the galaxy r. The Fig. 4 demonstrates the mass profile of the selected Q-ball as a function of the distance from the center of the galaxy r at the time t = 13 Gyr. It can be seen that the main interaction of the selected Q-ball with cosmological Q-balls occurs in the central part of the galaxy.

In this work, it is assumed to explain the presence of unusual gravitational wave signals using Q-balls of stellar mass, therefore, in the first approximation, it is necessary that the total number of Q-balls of stellar mass does not exceed the limit on the number of neutron stars in the galaxy, which is estimated as 10^8 [53]. Otherwise, LIGO/Virgo will observe an excessive number of gravitational wave events. It can be seen from Fig. 3 and Fig. 4 that the population of Q-balls of stellar mass is located at a distance of $r = 0 \div 0.05$ kpc. Then the number of Q-balls of stellar mass is

$$N_Q^{stellar} = \int_0^{0.05 \text{ kpc}} \frac{4\pi r^2 \rho(r)}{m_Q(r)} dr \approx 4 \times 10^6, \qquad (45)$$

which is several orders of magnitude less than the limit on the number of neutron stars in the galaxy, and therefore this limit is satisfied. It is worth noting that the mass of the selected Q-ball in the galaxy cannot be less than m_{\star} , therefore the total number of Q-balls of all masses in the



FIG. 3. The evolution of mass (41) depending on the distance from the center of the galaxy r and time t. The red curve corresponds to the situation when the selected Q-ball is located at a distance of 0.02 kpc. The blue curve corresponds to the situation when the selected Q-ball is located at a distance of 0.05 kpc. The green curve corresponds to the situation when the selected Q-ball is located at a distance of 0.10 kpc. The purple curve corresponds to the situation when the selected Q-ball is located at a distance of 0.25 kpc. The pink curve corresponds to the situation when the selected Q-ball is located at a distance of 0.50 kpc.



FIG. 4. The mass profile of the selected Q-ball (41) depending on the distance from the center of the galaxy r at time t = 13 kpc (red curve).

galaxy is

$$N_Q^{total} = \int_0^{19 \text{ kpc}} \frac{4\pi r^2 \rho(r)}{m_Q(r)} dr + \int_{19 \text{ kpc}}^{200 \text{ kpc}} \frac{4\pi r^2 \rho(r)}{m_\star} dr \sim 10^{17},$$
(46)

where the upper limit of integration in the second term is determined from the radius of the dark matter halo, and the upper limit of integration in the first term and the lower in the second is determined from the equation

$$m_Q(r_{max}) = m_\star \Rightarrow r_{max} \approx 19 \text{ kpc.}$$
 (47)

Thus, the masses of all Q-balls lie within

$$m_{\star} \approx 4 \times 10^{-6} \,\mathrm{M}_{\odot} \le m_Q(r) \lesssim 4 \,\mathrm{M}_{\odot}.$$
 (48)

Checking the limitation of the average cross-section of the interaction of bulk dark matter, taking into account (19), gives the following result

$$\langle \bar{\sigma} \rangle_b \approx 3 \text{ cm}^2/\text{g} \sim 1 \text{ cm}^2/\text{g}.$$
 (49)

This values have an equal order, which is acceptable within the framework of the model due to the presence of a large number of inaccuracies. Moreover, since this restriction, in its essence, is a restriction from below on the parameter v of the Lagrangian (1), the restriction (23) is automatically satisfied.

The last parameter that we have to check is the radius of the Q-ball, which is set by the expression (4). So, the radius of the selected Q-ball with a mass of 4 M_{\odot} , taking into account (40) is

$$R_Q = \sqrt{2}vQ^{1/4} \Big|_{r=0.02 \text{ kpc}} \sim 10^{10} \text{ km},$$
 (50)

which corresponds to the radius of the Solar system. The radius of a cosmological Q-ball with parameters (43) and accounting (9) is

$$R_Q = \sqrt{2}v Q_\star^{1/4} \approx 8 \times 10^8 \text{ km.}$$
 (51)

The radius of the selected Q-ball R_Q as a function of the distance from the center of the galaxy r is shown in the Fig. 5. The main parameters of the selected Q-ball in the present epoch are presented in the Table I.



FIG. 5. The radius of the selected Q-ball R_Q as a function of the distance from the center of the galaxy r (red curve).

VI. SUMMARY AND DISCUSSIONS

It is worth noting that it could be assumed that cosmological Q-balls are born during a phase transition immediately with masses of the order of one mass of the Sun. In this case, the need to compile the law of interaction of cosmological Q-balls with a selected Q-ball (33) has disappeared, but any physically meaningful configuration of the free parameters v, u, T_c and η_{χ} prohibits this scenario. According to (26), cosmological Q-balls cannot claim to be compact objects of stellar mass, therefore it was necessary to consider their interaction.

The simplest case assumes that some selected cosmological Q-ball absorbs other cosmological Q-balls in the flat expanding Universe. The trivial equation of such interaction (31) shows that for any physically meaningful set of free parameters that include the Lagrangian (1) parameter v, the asymmetry η_{χ} of the particles of the field χ , the velocity of the walls of the bubble of the new phase u (and, accordingly, the velocity of free cosmological Qballs), the temperature of the phase transition T_c , the selected cosmological Q-ball is unable to gain any significant mass. First of all, this is due to the expansion of the Universe, due to which the "gas" of cosmological Q-balls becomes extremely sparse. Thus, within the framework of such a trivial approximation, it is impossible to explain the unusual gravitational wave events received by LIGO and Virgo.

Due to the fact that the expansion of the Universe suppresses the interaction of cosmological Q-balls, it was proposed to consider the merger of cosmological Q-balls in gravitationally bound objects – galaxies. This makes it possible to neglect the expansion of the Universe, but adds new restrictions, for example, on the epoch of the phase transition, the concentration of cosmological Qballs and their velocity. A simple equation (39) was compiled, the solution (40) of which leads to the following results: selected cosmological Q-ball is able to gain the necessary mass of the order of the mass of the Sun, however, with any physically meaningful set of free parameters, such a Q-ball has a radius of the order of the radius of the Solar System. Such a value of the present radius of the selected Q-ball is significantly larger than the Schwarzschild radius for one mass of the Sun, which leads to the fact that the object turns out to be too "loose", which contradicts the requirement of compactness.

It is also worth discussing some inaccuracies of optimal free parameters. Firstly, there is a difference between the parameter v of the Lagrangian (1) and the phase transition temperature T_c . This difference in the framework of this work is considered insignificant, since the estimate of $v \sim T_c$ is quite rough and for clarification it requires analyzing the physics of the phase transition in more detail, which was not the purpose of the work. Secondly, it is easy to notice that the optimal parameter v does not correspond well to the constraint shown in the Fig. 2. It is worth noting that this restriction is derived from restrictions on the cross-section of the interaction of the

TABLE I. The main parameters of a selected Q-ball in the modern era.

The optimal parameters:	$v \sim 10^{-8} { m GeV}$	$T_c \sim 10^{-11} { m GeV}$	$\eta_{\chi} = 1$	u = 1	$u_{\star} = 0.0007$	$m_{\star} \sim 10^{-6} \ \mathrm{M}_{\odot}$
$\overline{r(\mathrm{kpc})}$	0.02	0.05	1	10	19	200
$M_Q({ m M}_{\odot})$	4	1	0.008	10^{-5}	10^{-6}	10^{-6}
$R_Q(m km)$	10^{11}	10^{10}	10^{10}	10^{9}	10^{9}	10^{9}

bulk dark matter and the energy density of dark matter in the present Universe, therefore, first of all, it is worth relying on these known quantities. Indeed, despite the fact that the restriction shown in the Fig. 2 is performed poorly, the restriction on the cross-section of the bulk dark matter (49) is performed well enough, which allows using such an optimal parameter with good accuracy. Thirdly, estimates of the volume from which the cosmological Q-ball (8) and its charge (9) are also inaccurate. In addition to the inaccuracies associated with calculations, which are described in detail in Ref. [36], certain inaccuracies are carried by the parameter g_* , which is responsible for the effective number of degrees of freedom and which depends on temperature. Clarifying this parameter is a separate work, therefore, it was decided to use the value proposed in [36] for estimations. Fourth, the Milky Way was chosen as a typical galaxy, as it is the most studied. In fact, there is a vast population of different types of galaxies that may differ significantly from ours in their configuration. This leads to the fact that the concentration of cosmological Q-balls in galaxies (37) can differ significantly depending on the galaxy, which may affect the final result.

In addition to the analysis of the considered models, other important results were also obtained in this work. The distribution of cosmological Q-balls by their charges (18), presented in [41], was clarified. Also, as already mentioned, a restriction was obtained on the parameter v of the Lagrangian (1), which is shown in the Fig. 2. Estimates of the populations of such Q-balls in the dark matter halo (45) and (46) were also obtained. Given that the mass of the selected cosmological Q-ball decreases rapidly with distance to the center of the galaxy, the population can be roughly divided into two groups: Qballs of stellar mass, which are located at a distance of up to 0.5 kpc from the center of the galaxy and the number of which is estimated as (45) and Q-balls of small masses (of the order of the mass of the cosmological Q-ball m_{\star}), which are located at distances greater than 0.5 kpc. It is worth noting that these populations, taking into account (48), are consistent with the results of observing microlensing events [54], as a result of which they may be of potential interest in other sections of astrophysics and cosmology.

Thus, despite the fact that the presented simple models do not allow Q-balls of dark matter to be candidates for the role of compact massive objects, this does not mean that Q-balls of dark matter cannot explain the unusual gravitational wave signals received by LIGO and Virgo. Only one simple model of Friedberg-Lee-Searlen Q-balls was considered in this work, however, there are other models that admit the existence of non-topological solitons, which can give a different result even within the framework of the proposed merger models [32], [33]. However, this study may help to narrow the scope of the search for solutions to the problem of unusual gravitational wave signals and add new limitations for future research.

ACKNOWLEDGMENTS

The Author is grateful to Sergey Troitsky for the original idea and for fruitful discussions on the manuscript. This work was supported by the Russian Science Foundation (RSF) grant 22-12-00215.

- A. Le Tiec and J. Novak, *Theory of gravitational waves*, in *An Overview of Gravitational Waves*, p. 1–41, World Scientific, (Feb., 2017), DOI.
- [2] LIGO SCIENTIFIC, VIRGO collaboration, GW190521: A Binary Black Hole Merger with a Total Mass of 150M_☉, Phys. Rev. Lett. **125** (2020) 101102 [2009.01075].
- [3] LIGO SCIENTIFIC, VIRGO collaboration, Properties and Astrophysical Implications of the 150 M_☉ Binary Black Hole Merger GW190521, Astrophys. J. Lett. **900** (2020) L13 [2009.01190].
- [4] LIGO SCIENTIFIC, VIRGO collaboration, GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object, Astrophys. J. Lett. 896 (2020) L44 [2006.12611].
- [5] LIGO SCIENTIFIC, VIRGO collaboration, Population Properties of Compact Objects from the Second LIGO-Virgo Gravitational-Wave Transient Catalog, Astrophys. J. Lett. **913** (2021) L7 [2010.14533].
- [6] E. J. Baxter, D. Croon, S. D. McDermott and J. Sakstein, Find the Gap: Black Hole Population Analysis with an Astrophysically Motivated Mass Function, Astrophys. J. Lett. **916** (2021) L16 [2104.02685].
- [7] A. Buonanno and B. S. Sathyaprakash, Sources of gravitational waves: Theory and observations, 2015.
- [8] A. Królak and P. Verma, Recent Observations of Gravitational Waves by LIGO and Virgo Detectors, Universe 7 (2021) 137.

- [9] R. Beradze and M. Gogberashvili, Unexpected ligo events and the mirror world, Monthly Notices of the Royal Astronomical Society 503 (2021) 2882–2886.
- [10] S. E. Woosley and A. Heger, The Pair-Instability Mass Gap for Black Holes, Astrophys. J. Lett. 912 (2021) L31 [2103.07933].
- [11] S. Wystub, Y. Dengler, J.-E. Christian and J. Schaffner-Bielich, Constraining exotic compact stars composed of bosonic and fermionic dark matter with gravitational wave events, 2110.12972.
- [12] R. Abbott et al, GW190814: Gravitational waves from the coalescence of a 23 solar mass black hole with a 2.6 solar mass compact object, The Astrophysical Journal Letters 896 (2020) L44.
- [13] H. Müller and B. D. Serot, Relativistic mean-field theory and the high-density nuclear equation of state, Nuclear Physics A 606 (1996) 508–537.
- [14] P. O. Mazur and E. Mottola, Gravitational vacuum condensate stars, Proceedings of the National Academy of Sciences 101 (2004) 9545–9550.
- [15] I. Bombaci, A. Drago, D. Logoteta, G. Pagliara and I. Vidaña, Was gw190814 a black hole-strange quark star system?, Physical Review Letters 126 (2021).
- [16] Z. Cao, L.-W. Chen, P.-C. Chu and Y. Zhou, Gw190814: Circumstantial evidence for up-down quark star, Physical Review D 106 (2022).
- [17] D. J. Kaup, Klein-gordon geon, Phys. Rev. 172 (1968) 1331.
- [18] R. Abbott et al, Observation of gravitational waves from two neutron star-black hole coalescences, The Astrophysical Journal Letters 915 (2021) L5.
- [19] C. D. Bailyn, R. K. Jain, P. Coppi and J. A. Orosz, The mass distribution of stellar black holes, The Astrophysical Journal 499 (1998) 367–374.
- [20] F. Özel, D. Psaltis, R. Narayan and J. E. McClintock, The black hole mass distribution in the galaxy, The Astrophysical Journal 725 (2010) 1918–1927.
- [21] W. M. Farr, N. Sravan, A. Cantrell, L. Kreidberg, C. D. Bailyn, I. Mandel et al., *The mass distribution of stellar-mass black holes*, *The Astrophysical Journal* **741** (2011) 103.
- [22] F. Özel, D. Psaltis, R. Narayan and A. Santos Villarreal, On the mass distribution and birth masses of neutron stars, The Astrophysical Journal 757 (2012) 55.
- [23] B. Kiziltan, A. Kottas, M. De Yoreo and S. E. Thorsett, The neutron star mass distribution, The Astrophysical Journal 778 (2013) 66.
- [24] J. Antoniadis, T. M. Tauris, F. Ozel, E. Barr, D. J. Champion and P. C. C. Freire, *The millisecond pulsar* mass distribution: Evidence for bimodality and constraints on the maximum neutron star mass, 2016.
- [25] V. C. Rubin and W. K. Ford, Jr., Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions, Astrophys. J. 159 (1970) 379.
- [26] G. Bertone and T. Tait, M. P., A new era in the search for dark matter, Nature 562 (2018) 51 [1810.01668].
- [27] K. Freese, Review of observational evidence for dark matter in the universe and in upcoming searches for dark stars, EAS Publications Series 36 (2009) 113–126.
- [28] B. K. K. Lee, M.-c. Chu and L.-M. Lin, Could the gw190814 secondary component be a bosonic dark matter admixed compact star?, The Astrophysical Journal 922 (2021) 242.

- [29] F. Di Giovanni, N. Sanchis-Gual, P. Cerdá-Durán and J. A. Font, Can fermion-boson stars reconcile multimessenger observations of compact stars?, Physical Review D 105 (2022).
- [30] C. Pacilio, M. Vaglio, A. Maselli and P. Pani, Gravitational-wave detectors as particle-physics laboratories: Constraining scalar interactions with a coherent inspiral model of boson-star binaries, Physical Review D 102 (2020).
- [31] A. Kusenko and P. J. Steinhardt, Q ball candidates for selfinteracting dark matter, Phys. Rev. Lett. 87 (2001) 141301 [astro-ph/0106008].
- [32] M. I. Tsumagari, The physics of q-balls, 2009. 10.48550/ARXIV.0910.3845.
- [33] S. R. Coleman, *Q-balls*, *Nucl. Phys. B* **262** (1985) 263.
- [34] R. Friedberg, T. D. Lee and A. Sirlin, Class of scalar-field soliton solutions in three space dimensions, *Phys. Rev. D* 13 (1976) 2739.
- [35] R. Friedberg, T. Lee and A. Sirlin, Gauge-field non-topological solitons in three space-dimensions (ii), Nuclear Physics B 115 (1976) 32.
- [36] E. Krylov, A. Levin and V. Rubakov, Cosmological phase transition, baryon asymmetry and dark matter Q-balls, Phys. Rev. D 87 (2013) 083528 [1301.0354].
- [37] A. H. Guth and E. J. Weinberg, Cosmological consequences of a first-order phase transition in the su₅ grand unified model, Phys. Rev. D 23 (1981) 876.
- [38] V. Rubakov and D. Gorbunov, Introduction To The Theory Of The Early Universe: Hot Big Bang Theory. World Scientific Publishing Company, 2011.
- [39] M. Stuke, Particle asymmetries in the early universe, Progress in Particle and Nuclear Physics 66 (2011) 220–225.
- [40] K. Zarembo, Lepton asymmetry of the universe and charged quark-gluon plasma, Physics Letters B 493 (2000) 375–379.
- [41] S. Troitsky, Supermassive dark-matter Q-balls in galactic centers?, JCAP 11 (2016) 027 [1510.07132].
- [42] M. Vogelsberger, J. Zavala and A. Loeb, Subhaloes in self-interacting galactic dark matter haloes: Self-interacting galactic dark matter haloes, Monthly Notices of the Royal Astronomical Society 423 (2012) 3740–3752.
- [43] M. Rocha, A. H. G. Peter, J. S. Bullock, M. Kaplinghat, S. Garrison-Kimmel, J. Oñorbe et al., Cosmological simulations with self-interacting dark matter – i. constant-density cores and substructure, Monthly Notices of the Royal Astronomical Society 430 (2013) 81–104.
- [44] J. Zavala, M. Vogelsberger and M. G. Walker, Constraining self-interacting dark matter with the milky way's dwarf spheroidals, Monthly Notices of the Royal Astronomical Society: Letters 431 (2013) L20–L24.
- [45] O. D. Elbert, J. S. Bullock, S. Garrison-Kimmel, M. Rocha, J. Oñorbe and A. H. G. Peter, Core formation in dwarf haloes with self-interacting dark matter: no fine-tuning necessary, Monthly Notices of the Royal Astronomical Society 453 (2015) 29–37.
- [46] T. Multamaki and I. Vilja, Q ball collisions in the MSSM: Gauge mediated supersymmetry breaking, Phys. Lett. B 484 (2000) 283 [hep-ph/0005162].
- [47] T. Multamaki and I. Vilja, Q Ball collisions in the MSSM: Gravity mediated supersymmetry breaking, Phys. Lett. B 482 (2000) 161 [hep-ph/0003270].

- [49] N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont,
 C. Baccigalupi, M. Ballardini et al., *Planck2018 results:* Vi. cosmological parameters, Astronomy & Astrophysics 641 (2020) A6.
- [50] S. J., D. C. A. and D. I., Galaxy formation, Proceedings of the International School of Physics "Enrico Fermi" 186 (2014) 137–187.
- [51] C. J. Conselice, The evolution of galaxy structure over

cosmic time, Annual Review of Astronomy and Astrophysics **52** (2014) 291–337.

- [52] G. Bertone, Particle Dark Matter: Observations, Models and Searches. Cambridge University Press, 2010.
- [53] N. Sartore, E. Ripamonti, A. Treves and R. Turolla, Galactic neutron stars: I. space and velocity distributions in the disk and in the halo, Astronomy and Astrophysics 510 (2010) A23.
- [54] P. Mróz et al., No massive black holes in the Milky Way halo, Nature 632 (2024) 749 [2403.02386].