

# Differential uniformity and costacyclic code from some power mapping

Yuehui Cui, Jinquan Luo

School of Mathematics and Statistics, Central China Normal University, Wuhan 430079, China

## Abstract

In this paper, we study the differential properties of  $x^d$  over  $\mathbb{F}_{p^n}$  with  $d = p^{2l} - p^l + 1$ . By studying the differential equation of  $x^d$  and the number of rational points on some curves over finite fields, we completely determine differential spectrum of  $x^d$ . Then we investigate the  $c$ -differential uniformity of  $x^d$ . We also calculate the value distribution of a class of exponential sum related to  $x^d$ . In addition, we obtain a class of six-weight consta-cyclic codes, whose weight distribution is explicitly determined. Part of our results is a complement of the works shown in [3, 4] which mainly focus on cross correlations.

**Keywords:** Differential spectrum,  $c$ -differential uniformity, Constacyclic code, Exponential sum, Weight distribution.

**2010 Mathematics Subject Classification:** 94A60, 11T06, 11T23.

## 1 Introduction

Let  $\mathbb{F}_{p^n}$  be the finite field with  $p^n$  elements, where  $p$  is a prime and  $n$  is a positive integer. Let  $\mathbb{F}_{p^n}^*$  denote the multiplicative group of the finite field  $\mathbb{F}_{p^n}$ . For any function  $f : \mathbb{F}_{p^n} \rightarrow \mathbb{F}_{p^n}$ , the derivative function of  $f$  in respect to any element  $a \in \mathbb{F}_{p^n}$  is a function from  $\mathbb{F}_{p^n}$  to  $\mathbb{F}_{p^n}$  defined by

$$D_a f(x) = f(x + a) - f(x), \forall x \in \mathbb{F}_{p^n}.$$

For any  $a, b \in \mathbb{F}_{p^n}$ , let

$$\delta(a, b) = \#\{x \in \mathbb{F}_{p^n} \mid D_a f(x) = b\}.$$

The differential uniformity of  $f$  is defined as

$$\delta = \max \{\delta(a, b) \mid a \in \mathbb{F}_{p^n}^*, b \in \mathbb{F}_{p^n}\}.$$

If  $\delta = 1$ , then  $f$  is called a perfect nonlinear (PN) function. If  $\delta = 2$ , then  $f$  is called an almost perfect nonlinear (APN) function. The readers can refer to [8, 9, 11, 14] for more information on PN and APN

---

*Email addresses* :luojinquan@mail.ccn.u.edu.cn (Jinquan Luo); hfcyh1@163.com (Yuehui Cui).

functions. In recent years, the design of Substitution box (S-box) that can effectively resist differential attacks has become a hot topic in cryptographic research. Studies have shown that the power functions with low differential uniformity offer significant advantages in S-box design. Therefore, researching the power functions with low differential uniformity and exploring their application in S-box design has become an important research direction in the field of cryptography. For related details, we refer to [5, 6]. When  $f(x) = x^d$  is a power mapping,

$$(x+a)^d - x^d = b \Leftrightarrow a^d \left( \left( \frac{x}{a} + 1 \right)^d - \left( \frac{x}{a} \right)^d \right) = b$$

implying that  $\delta(a, b) = \delta(1, \frac{b}{a^d})$  for all  $a \in \mathbb{F}_{p^n}^*$  and  $b \in \mathbb{F}_{p^n}$ . Thus, the differential properties of  $f(x) = x^d$  are completely determined by the values of  $\delta(1, b)$  when  $b$  runs through  $\mathbb{F}_{p^n}$ .

**Definition 1.** Let  $f(x) = x^d$  be a power function from  $\mathbb{F}_{p^n}$  to  $\mathbb{F}_{p^n}$  with differential uniformity  $\delta$ . Denote

$$\omega_i = \# \{b \in \mathbb{F}_{p^n} \mid \delta(1, b) = i\},$$

where  $0 \leq i \leq \delta$ . The differential spectrum of  $f(x) = x^d$  is defined as

$$\mathbb{DS} = \{\omega_i \mid 0 \leq i \leq \delta\}.$$

Sometimes we ignore the zeros in  $\mathbb{DS}$ . We also have two basic identities [2].

$$\sum_{i=0}^{\delta} \omega_i = p^n \text{ and } \sum_{i=0}^{\delta} i\omega_i = p^n. \quad (1)$$

These two identities are useful in calculating the differential spectrum of  $f$ . However, determining the differential spectrum of a power mapping remains theoretically challenging. We list some results in table 1. Differential spectrum of a power mapping has many applications, such as cross-correlation distribution, Walsh transformation,  $t$ -designs and coding theory. Readers can refer to [7, 10-13, 15] for more information.

To define the capacity of a particular function to withstand multiplicative differential attack [16], Ellingsen et al. [17] proposed the concept of  $c$ -differential uniformity, which is described as follows.

**Definition 2.** Let  $a, c \in \mathbb{F}_{p^n}$ . For any function  $f : \mathbb{F}_{p^n} \rightarrow \mathbb{F}_{p^n}$ , the (multiplicative)  $c$ -derivative of  $f$  in respect to any element  $a$  is defined as

$${}_c D_a f(x) = f(x+a) - cf(x).$$

Let

$${}_c \Delta_f(a, b) = \# \{x \in \mathbb{F}_{p^n} \mid {}_c D_a f(x) = b\}.$$

We call

$${}_c \Delta_f = \max \{ {}_c \Delta_f(a, b) \mid a, b \in \mathbb{F}_{p^n} \text{ and } (a, c) \neq (0, 1) \},$$

the  $c$ -differential uniformity of  $f$ . If  ${}_c \Delta_f = \delta$ , then we say  $f$  is differentially  $(c, \delta)$ -uniform.

If  ${}_c \Delta_f = 1$ , then  $f$  is called a perfect  $c$ -nonlinear (PcN) function. If  ${}_c \Delta_f = 2$ , then  $f$  is called an almost perfect  $c$ -nonlinear (APcN) function. Functions exhibiting low  $c$ -differential uniformity demonstrate strong resistance to certain forms of differential cryptanalysis. In order to calculate the  $c$ -differential uniformity of a power function  $f(x) = x^d$  over  $\mathbb{F}_{p^n}$  with  $c \neq 1$ , the following result was introduced.

$d$	Conditions	$\delta$	References
$2^t + 1$	$\gcd(t, n) = s$	$2^s$	[2,21]
$2^n - 2$	$n \geq 2$	2 or 4	[2,21]
$2^{2t} - 2^t + 1$	$\gcd(t, n) = s, \frac{n}{s}$ odd	$2^s$	[2]
$2^{2k} + 2^k + 1$	$n = 4k$	4	[2,22]
$2^t - 1$	$t = 3, n - 2$	6 or 8	[23]
$2^t - 1$	$t = n/2, n/2 + 1, n$ even	$2^{\frac{n}{2}} - 2$ or $2^{\frac{n}{2}}$	[2]
$2^t - 1$	$t = (n - 1)/2, (n + 3)/2, n$ odd	6 or 8	[24]
$2^{3l} + 2^{2l} + 2^l - 1$	$n = 4l$	$2^{2l}$	[25]
$2^m + 2^{(m+1)/2} + 1$	$n = 2m, m \geq 5$ odd	8	[26]
$2^{m+1} + 3$	$n = 2m, m \geq 5$ odd	8	[26]
$\frac{2^m - 1}{2^k + 1} + 1$	$n = 2m, \gcd(k, m) = 1$	$2^m$	[27]
$2 \cdot 3^{\frac{n-1}{2}} + 1$	$n$ odd	4	[7]
$\frac{p^k + 1}{2}$	$\gcd(k, n) = e$	$\frac{p^e - 1}{2}$ or $p^e + 1$	[28]
$\frac{p^n + 1}{p^m + 1} + \frac{p^n - 1}{2}$	$p \equiv 3 \pmod{4}, m \mid n, n$ odd	$\frac{p^m + 1}{2}$	[28]
$p^{2k} - p^k + 1$	$p$ odd, $\gcd(k, n) = e, \frac{n}{e}$ odd	$p^e + 1$	[29,30]
$p^n - 3$	$p$ odd, any $n$	$\leq 5$	[31,32]
$p^m + 2$	$n = 2m, p \geq 3$	4	[33]
$\frac{p^n + 3}{2}$	$p \geq 5, p^n \equiv 1 \pmod{4}$	3	[34]
$\frac{5^n - 3}{2}$	$n \geq 2$	4 or 5	[35]
$\frac{p^n - 3}{2}$	$p^n \geq 7, p^n \equiv 3 \pmod{4}, p^n \neq 27$	2 or 3	[36]
$k(p^m - 1)$	odd $p, n = 2m, \gcd(k, p^m + 1) = 1$	$p^m - 2$	[37]
$\frac{p^n + 3}{2}$	$p^n \equiv 3 \pmod{4}, p \neq 3$	2 or 4	[38]
$2p^m - 1$	$n = 2m, p$ odd	$p^m$	[39]
$\frac{p^n + 1}{4} + \frac{p^n - 1}{2}$	$p^n \equiv 3 \pmod{8}$ or $p^n \equiv 7 \pmod{8}$	$\leq 4$	[40,41]
$\frac{p^n + 1}{4}$	$p^n \equiv 3 \pmod{8}$ or $p^n \equiv 7 \pmod{8}$	$\leq 4$	[40,41]
$d(p^k + 1) \equiv 2 \pmod{p^n - 1}$	$\gcd(n, k) = e, \frac{n}{e}$ odd	$\frac{p^e + 1}{2}, \frac{p^e - 1}{2}$ or $p^e + 1$	[42]
$p^{2l} - p^l + 1$	$n = 4l, \text{ any prime } p$	$p^{2l} - p^l$	This paper

Table 1: Some power functions  $f(x) = x^d$  over  $\mathbb{F}_{p^n}$  with known differential spectrum.

**Lemma 1.** ([18]) *Let  $f(x) = x^d$  be a power function over  $\mathbb{F}_{p^n}$ . Then*

$${}_c\Delta_f = \max \{ \{ {}_c\Delta_f(1, b) : b \in \mathbb{F}_{p^n} \} \cup \{ \gcd(d, p^n - 1) \} \}.$$

The rest of this paper is organized as follows. In section 2, we introduce some notations and auxiliary tools. In section 3, we determine the differential spectrum of  $x^{p^{2l}-p^l+1}$ . We also investigate the  $c$ -differential uniformity of the power mapping  $x^{p^{2l}-p^l+1}$ . In section 4, we calculate the value distribution of a class of exponential sum. As an application, a class of six-weight cyclic codes were presented. The conclusive remarks are given in section 5.

For convenience, we fix the following notation in this paper:

$\mathbb{F}_{p^n}$	finite field of $p^n$ elements
$\mathbb{F}_{p^n}^*$	multiplicative group of $\mathbb{F}_{p^n}$
$\psi$	a primitive element of $\mathbb{F}_{p^n}$
$\text{Tr}_1^n$	trace function from $\mathbb{F}_{p^n}$ to $\mathbb{F}_p$
$\zeta_p$	primitive complex $p$ -th root of unity

## 2 Preliminaries

Before stating the main results, we introduce some notations. Let  $e$  be a positive integer and define

$$\mu_e = \{x \in \mathbb{F}_{p^n} \mid x^e = 1\}.$$

If  $x \in \mathbb{F}_{p^n}^*$  and  $x = \psi^t$  with  $0 \leq t < p^n - 1$ , we denote  $\text{ind}_\psi(x) = t$ . Let  $\mathbb{F}_{p^n}^\sharp = \mathbb{F}_{p^n} \setminus \{0, -1\}$ . We denote

$$S_{r,s} = \{x \in \mathbb{F}_{p^n}^\sharp : \text{ind}_\psi(x+1) \equiv r \pmod{p^l+1}, \text{ind}_\psi(x) \equiv s \pmod{p^l+1}\},$$

where  $0 \leq r, s \leq p^l$ ,  $n = 4l$ . Obviously,

$$\bigsqcup_{0 \leq i, j \leq p^l} S_{i,j} = \mathbb{F}_{p^n}^\sharp$$

(here  $\bigsqcup$  indicates disjoint union).

The following result plays a crucial role in this paper.

**Lemma 2.** ([1, Theorem 1]) *Let  $p$  be a prime number,  $n = 2ks$  with  $k$  and  $s$  positive integers. Let  $n_1$  and  $n_2$  be two positive integers,  $t = \gcd(n_1, n_2)$ ,  $\text{lcm}(n_1, n_2) \mid p^s + 1$ . Let  $0 \leq r_1 \leq n_1 - 1$  and  $0 \leq r_2 \leq n_2 - 1$ . Let  $N_{p^n}(\chi)$  be the number of  $\mathbb{F}_{p^n}$ -rational points on the affine curve*

$$\chi : \alpha x^{n_1} + \beta y^{n_2} + 1 = 0.$$

For  $\alpha \in C_{1,r_1} = \left\{ \psi^{r_1+n_1 w} \mid 0 \leq w \leq \frac{p^{n_1}-1}{n_1} - 1 \right\}$ ,  $\beta \in C_{2,r_2} = \left\{ \psi^{r_2+n_2 w} \mid 0 \leq w \leq \frac{p^{n_2}-1}{n_2} - 1 \right\}$ . Then we have

- (i) if  $r_1 = r_2 = 0$ , then  $N_{p^n}(\chi) = p^n + (-1)^{k-1}((n_1 - 1)(n_2 - 1) + 1 - t)p^{\frac{n}{2}} - t + 1$ .
- (ii) if  $r_1 = 0$ ,  $r_2 \neq 0$ ,  $t \nmid r_2$ , then  $N_{p^n}(\chi) = p^n + (-1)^k(n_1 - 2)p^{\frac{n}{2}} + 1$ .
- (iii) if  $r_1 \neq 0$ ,  $r_2 = 0$ ,  $t \nmid r_1$ , then  $N_{p^n}(\chi) = p^n + (-1)^k(n_2 - 2)p^{\frac{n}{2}} + 1$ .
- (iv) if  $r_1 \neq 0$ ,  $r_2 \neq 0$  and  $t \nmid r_1 - r_2$ , then  $N_{p^n}(\chi) = p^n + (-1)^{k-1}2p^{\frac{n}{2}} + 1$ .

(v) if  $r_1 \neq 0$ ,  $r_2 \neq 0$  and  $t \mid r_1 - r_2$ , then  $N_{p^n}(\chi) = p^n + (-1)^k(t-2)p^{\frac{n}{2}} - t + 1$ .

**Lemma 3.** ([43]). Let  $n = 2m$  be an even positive integer and  $a, b \in \mathbb{F}_{2^n}^*$ . Then the quadratic equation  $x^2 + ax + b = 0$  has both solutions in  $\mu_{2^{m+1}}$  if and only if

$$b = a^{1-2^m} \text{ and } \text{Tr}_1^m(a^{-2^m-1}) = 1.$$

### 3 Differential properties of $f(x) = x^{p^{2l}-p^l+1}$ over $\mathbb{F}_{p^n}$

#### 3.1 Differential spectrum

Let  $d = p^{2l} - p^l + 1$ , where  $n = 4l$ ,  $l$  is a positive integer and  $p$  is a prime number. In this subsection, we determine the differential spectrum of  $f(x) = x^d$  over  $\mathbb{F}_{p^n}$ , since the differential spectrum of  $x^{p^{3l}-p^{2l}+p^l}$  and  $x^{p^{2l}-p^l+1}$  are identical. To determine the differential spectrum of  $f$ , we mainly study the number of solutions in  $\mathbb{F}_{p^n}$  of

$$(x+1)^d - x^d = b \tag{2}$$

for any  $b \in \mathbb{F}_{p^n}$ . Denote by  $\delta(b)$  the number of solutions of (2) in  $\mathbb{F}_{p^n}$ . We have the following result.

**Lemma 4.** With the notations above, we have

(i)  $\delta(1) = p^l$ .

(ii) if  $b^{p^l+1} = 1$  and  $b \neq 1$ , then  $\delta(b) = p^{2l} - p^l$ .

(iii) if  $b^{p^l+1} \neq 1$ , then  $\delta(b) = 0$  or  $2$ .

*Proof.* Let  $\Delta(x) = (x+1)^d - x^d$  and  $\alpha = \psi^{\frac{p^n-1}{p^l+1}}$ . If  $x \in \mathbb{F}_{p^n}^\#$ , we can distinguish  $(p^l+1)^2$  disjoint cases:

	Set	Equation (2)	$x + 1$	$x$
Case 1	$S_{0,0}$	$1 = b$	To Be Determined	To Be Determined
Case 2	$S_{0,1}$	$x + 1 - \alpha x = b$	$\frac{b-\alpha}{1-\alpha}$	$\frac{b-1}{1-\alpha}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $p^l + 1$	$S_{0,p^l}$	$x + 1 - \alpha^{p^l} x = b$	$\frac{b-\alpha^{p^l}}{1-\alpha^{p^l}}$	$\frac{b-1}{1-\alpha^{p^l}}$
Case $p^l + 2$	$S_{1,0}$	$\alpha(x + 1) - x = b$	$\frac{b-1}{\alpha-1}$	$\frac{b-\alpha}{\alpha-1}$
Case $p^l + 3$	$S_{1,1}$	$\alpha = b$	To Be Determined	To Be Determined
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $2(p^l + 1)$	$S_{1,p^l}$	$\alpha(x + 1) - \alpha^{p^l} x = b$	$\frac{b-\alpha^{p^l}}{\alpha-\alpha^{p^l}}$	$\frac{b-\alpha}{\alpha-\alpha^{p^l}}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $p^l(p^l + 1) + 1$	$S_{p^l,0}$	$\alpha^{p^l}(x + 1) - x = b$	$\frac{b-1}{\alpha^{p^l}-1}$	$\frac{b-\alpha^{p^l}}{\alpha^{p^l}-1}$
Case $p^l(p^l + 1) + 2$	$S_{p^l,1}$	$\alpha^{p^l}(x + 1) - \alpha x = b$	$\frac{b-\alpha}{\alpha^{p^l}-\alpha}$	$\frac{b-\alpha^{p^l}}{\alpha^{p^l}-\alpha}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $(p^l + 1)^2$	$S_{p^l,p^l}$	$\alpha^{p^l} = b$	To Be Determined	To Be Determined

Table 2: Equation (2) with  $x \in \mathbb{F}_{p^n}^\#$

(i) Note that  $\Delta(x) = 1$  has no solution in  $\bigsqcup_{1 \leq i \leq p^l} S_{i,i}$ . If  $\Delta(x) = 1$  has one solution  $x_1 \in S_{j_1, j_2}$ , where  $j_1 \neq j_2$  and  $j_1, j_2 \neq 0$ , the

$$x_1 = \frac{1 - \alpha^{j_1}}{\alpha^{j_1} - \alpha^{j_2}} \text{ and } x_1 + 1 = \frac{1 - \alpha^{j_2}}{\alpha^{j_1} - \alpha^{j_2}}.$$

Notice that  $\alpha^i \in \mu_{p^l+1}$  for  $i = 1, 2, \dots, p^l$  which implies  $\alpha^{ip^l} = \alpha^{ip^{3l}} = \alpha^{-i}$  and  $\alpha^{ip^{2l}} = \alpha^i$ . Then we have

$$x_1^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{1 - \alpha^{j_1}}{\alpha^{j_1} - \alpha^{j_2}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \alpha^{2j_2} = \alpha^{j_2},$$

$$(x_1 + 1)^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{1 - \alpha^{j_2}}{\alpha^{j_1} - \alpha^{j_2}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \alpha^{2j_1} = \alpha^{j_1}.$$

But this contradicts  $j_1 \neq j_2$ . So  $\Delta(x) = 1$  has no solution in  $\mathbb{F}_{p^n}^\# \setminus S_{0,0}$ . If  $x \in S_{0,0}$ , then  $\Delta(x) = 1$  is equivalent to the system of equations

$$\begin{cases} y - x = 1, \\ \text{ind}_\psi(y) = \text{ind}_\psi(x) \equiv 0 \pmod{p^l + 1}. \end{cases} \quad (3)$$

By Lemma 2, we have  $p^l - 2$  solutions to (3). We also get  $\Delta(0) = 1$  and  $\Delta(-1) = 1$ . In total,  $\delta(1) = p^l$ .

(ii) We only show the case  $b = \alpha$ . The other cases for  $b = \alpha^k$ ,  $2 \leq k \leq p^l$  are similar and we omit them. Note that  $\Delta(x) = \alpha$  has no solution in  $\bigsqcup_{\substack{0 \leq i \leq p^l \\ i \neq 1}} S_{i,i}$ . If  $\Delta(x) = \alpha$  has one solution  $y_1 \in S_{k_1, k_2}$  with

$k_1 \neq k_2$  and  $k_1, k_2 \neq 1$ , then

$$y_1 = \frac{\alpha - \alpha^{k_1}}{\alpha^{k_1} - \alpha^{k_2}} \text{ and } y_1 + 1 = \frac{\alpha - \alpha^{k_2}}{\alpha^{k_1} - \alpha^{k_2}}.$$

Then we obtain

$$\begin{aligned} y_1^{p^{3l} - p^{2l} + p^l - 1} &= \left( \frac{\alpha - \alpha^{k_1}}{\alpha^{k_1} - \alpha^{k_2}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{\alpha^{p^l} \alpha^{k_1} - 1}{\alpha - \alpha^{k_1}} \right)^2 \alpha^{2k_2} = \alpha^{k_2}, \\ (y_1 + 1)^{p^{3l} - p^{2l} + p^l - 1} &= \left( \frac{\alpha - \alpha^{k_2}}{\alpha^{k_1} - \alpha^{k_2}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{\alpha^{p^l} \alpha^{k_2} - 1}{\alpha - \alpha^{k_2}} \right)^2 \alpha^{2k_1} = \alpha^{k_1}, \end{aligned}$$

which yields

$$\alpha^{k_1} + \alpha^{k_2} = \alpha^{-2} \alpha^{k_1} \alpha^{k_2} + 2\alpha - 1. \quad (4)$$

Taking  $p^{l+1}$ -th power on both sides of (4), it implies

$$\alpha^{k_1} \alpha^{k_2} = \alpha^2, \quad (5)$$

which yields

$$\begin{cases} \alpha^{k_1} \alpha^{k_2} = \alpha^2, \\ \alpha^{k_1} + \alpha^{k_2} = 2\alpha. \end{cases}$$

Hence,  $\alpha^{k_1} = \alpha^{k_2} = \alpha$ , which contradicts  $k_1 \neq k_2$ . So  $\Delta(x) = \alpha$  has no solution in  $\mathbb{F}_{p^n}^\# \setminus S_{1,1}$ . If  $x \in S_{1,1}$ , then  $\Delta(x) = \alpha$  is equivalent to the system of equations

$$\begin{cases} \psi y - \psi x = 1, \\ \text{ind}_\psi(y) = \text{ind}_\psi(x) \equiv 0 \pmod{p^l + 1}. \end{cases} \quad (6)$$

By Lemma 2, we have  $p^{2l} - p^l$  solutions to (6). We also get  $\Delta(0) \neq \alpha$  and  $\Delta(-1) \neq \alpha$ . In total,  $\delta(\alpha) = p^{2l} - p^l$ .

(iii) Note that

$$d(p^l + 1) \equiv p^l + 1 \pmod{p^{4l} - 1}. \quad (7)$$

Using (2) and (7), one obtains

$$(x + 1)^{p^l + 1} = (x^d + b)^{p^l + 1}. \quad (8)$$

According to (8), we have

$$x^d + b = \gamma_1(x + 1), \quad (9)$$

for some  $\gamma_1 \in \mu_{p^l + 1}$ . Taking  $(p^l + 1)$ -th power on both sides of (9), we have

$$x^{p^l + 1} = (\gamma_1(x + 1) - b)^{p^l + 1}.$$

Then we obtain

$$\gamma_2 x = \gamma_1(x+1) - b, \quad (10)$$

for some  $\gamma_2 \in \mu_{p^l+1}$ . When  $\gamma_1 = \gamma_2$ , we obtain  $b = \gamma_1 \in \mu_{p^l+1}$ . This situation has been discussed in (i). When  $\gamma_1 \neq \gamma_2$ , we have  $x = \frac{b-\gamma_1}{\gamma_1-\gamma_2}$  which is a solution of (2) if and only if

$$\left(\frac{b-\gamma_2}{\gamma_1-\gamma_2}\right)^d - \left(\frac{b-\gamma_1}{\gamma_1-\gamma_2}\right)^d = b. \quad (11)$$

Notice that  $\gamma_i \in \mu_{p^l+1}$  for  $i = 1, 2$  which implies  $\gamma_i^{p^l} = \gamma_i^{p^{3l}} = \gamma_i^{-1}$  and  $\gamma_i^{p^{2l}} = \gamma_i$ . Then by a detailed calculation one can obtain that

$$(b-\gamma_i)^d = \frac{b^{p^{3l}+p^l} - (b^{p^{3l}} + b^{p^l})\gamma_i^{p^l} + \gamma_i^{2p^l}}{b^{p^{2l}} - \gamma_i}$$

for any  $b^{p^{2l}} \neq \gamma_i$  and from which one can also have

$$(\gamma_1 - \gamma_2)^d = \frac{\gamma_1 - \gamma_2}{\gamma_1^2 \gamma_2^2}.$$

Then (11) can be rewritten as

$$\begin{aligned} & (\gamma_2 - \gamma_1)b^{p^{3l}+p^l} + (b^{p^{3l}+p^{2l}} + b^{p^{2l}+p^l} + b) \left(\frac{1}{\gamma_1} - \frac{1}{\gamma_2}\right) + (b^{p^{3l}} + b^{p^l}) \left(\frac{\gamma_1}{\gamma_2} - \frac{\gamma_2}{\gamma_1}\right) \\ & + (b^{p^{2l}} + b^{p^{2l}+1}) \left(\frac{1}{\gamma_2^2} - \frac{1}{\gamma_1^2}\right) + \frac{\gamma_2}{\gamma_1^2} - \frac{\gamma_1}{\gamma_2^2} - b^{2p^{2l}+1} \left(\frac{1}{\gamma_1\gamma_2^2} - \frac{1}{\gamma_2\gamma_1^2}\right) = 0. \end{aligned} \quad (12)$$

Taking  $p^{2l}$ -th power on both sides of (12) and subtracting these two equations lead to

$$(b^{p^{2l}} - b) \left[ (b^{p^{3l}} + b^{p^l} - 1) \left(\frac{1}{\gamma_1} - \frac{1}{\gamma_2}\right) + \left(\frac{1}{\gamma_2^2} - \frac{1}{\gamma_1^2}\right) + b^{p^{2l}+1} \left(\frac{1}{\gamma_1\gamma_2^2} - \frac{1}{\gamma_2\gamma_1^2}\right) \right] = 0. \quad (13)$$

*Case 1:*  $b^{p^{2l}} \neq b$ . Then by (13) one gets

$$\gamma_1 + \gamma_2 = (b^{p^{3l}} + b^{p^l} - 1)(\gamma_1\gamma_2) - b^{p^{2l}+1}. \quad (14)$$

Taking  $p^{l+1}$ -th power on both sides of (14), it implies

$$(b^{p^{3l}} - 1)(1 - b^{p^l})(\gamma_1\gamma_2)^2 + (b^{p^{2l}} - 1)(b - 1)\gamma_1\gamma_2 = 0. \quad (15)$$

Then we have

$$\begin{cases} \gamma_1\gamma_2 = (b^{p^{3l}} - 1)(1 - b^{p^l})^{p^l-1}, \\ \gamma_1 + \gamma_2 = (b^{p^{3l}} + b^{p^l} - 1)(b^{p^{3l}} - 1)(1 - b^{p^l})^{p^l-1} - b^{p^{2l}+1}, \end{cases}$$

i.e.,  $\gamma_1, \gamma_2$  are two solutions of

$$x^2 - ((b^{p^{3l}} + b^{p^l} - 1)(b^{p^{3l}} - 1)(1 - b^{p^l})^{p^l-1} - b^{p^{2l}+1})x + (b^{p^{3l}} - 1)(1 - b^{p^l})^{p^l-1} = 0.$$



Hence, when  $\gamma_1 \neq \gamma_2$ ,  $b \notin \mu_{p^l+1}$  and  $b^{p^{2l}} \neq b$ , the equation (2) has either no solution or two solutions.

*Case 2:*  $b^{p^{2l}} = b$ . For a fixed  $b$ , if  $\Delta(x_2) = b$ , where

$$x_2 = \frac{b - \alpha^{j_3}}{\alpha^{j_3} - \alpha^{j_4}} \in S_{j_3, j_4},$$

then we obtain

$$\begin{aligned} x_2^{p^{3l} - p^{2l} + p^l - 1} &= \left( \frac{b - \alpha^{j_3}}{\alpha^{j_3} - \alpha^{j_4}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{b^{p^l} \alpha^{j_3} - 1}{b - \alpha^{j_3}} \right)^2 \alpha^{2j_4} = \alpha^{j_4}, \\ (x_2 + 1)^{p^{3l} - p^{2l} + p^l - 1} &= \left( \frac{b - \alpha^{j_4}}{\alpha^{j_3} - \alpha^{j_4}} \right)^{p^{3l} - p^{2l} + p^l - 1} = \left( \frac{b^{p^l} \alpha^{j_4} - 1}{b - \alpha^{j_4}} \right)^2 \alpha^{2j_3} = \alpha^{j_3}, \end{aligned}$$

which yields

$$\alpha^{j_3} + \alpha^{j_4} = b^{2p^l} \alpha^{j_3} \alpha^{j_4} + 2b - 1. \quad (16)$$

Taking  $p^{l+1}$ -th power on both sides of (16), it implies

$$\alpha^{j_3} \alpha^{j_4} = (b^{p^l} - 1)^{2(p^l - 1)}. \quad (17)$$

Then by (16) and (17) one can obtain

$$\begin{cases} \alpha^{j_3} \alpha^{j_4} = (b^{p^l} - 1)^{2(p^l - 1)}, \\ \alpha^{j_3} + \alpha^{j_4} = b^{2p^l} (b^{p^l} - 1)^{2(p^l - 1)} + 2b - 1, \end{cases}$$

i.e.,  $\alpha^{j_3}, \alpha^{j_4}$  are two solutions of

$$x^2 - (b^{2p^l} (b^{p^l} - 1)^{2(p^l - 1)} + 2b - 1)x + (b^{p^l} - 1)^{2(p^l - 1)} = 0.$$

Hence, when  $\gamma_1 \neq \gamma_2$ ,  $b \notin \mu_{p^l+1}$  and  $b^{p^{2l}} = b$ , the equation (2) has either no solution or two solutions. This completes the proof.  $\square$

We determine the differential spectrum of  $f$  in the following result.

**Theorem 1.** *Let  $f(x) = x^d$  be a power mapping defined over  $\mathbb{F}_{p^n}$ , where  $d = p^{3l} - p^{2l} + p^l$ ,  $n = 4l$ . Then the differential spectrum of  $f$  is given by*

$$\mathbb{DS} = \left\{ \omega_0 = \frac{p^n + p^{3l} - p^{2l} - p^l - 2}{2}, \omega_2 = \frac{p^n - p^{3l} + p^{2l} - p^l}{2}, \omega_{p^l} = 1, \omega_{p^{2l} - p^l} = p^l \right\}.$$

*Proof.* From (1) and Lemma 4, we obtain

$$\omega_0 + \omega_2 + \omega_{e_1} + \omega_{e_2} = p^n \quad (18)$$

$$2 \cdot \omega_2 + p^l \cdot \omega_{p^l} + (p^{2l} - p^l) \cdot \omega_{p^{2l} - p^l} = p^n \quad (19)$$

$$\omega_{p^l} = 1 \quad (20)$$

$$\omega_{p^{2l} - p^l} = p^l. \quad (21)$$

Solving the system equations consisting of (18)-(21) we obtain the result.  $\square$

### 3.2 $c$ -differential uniformity of $x^{p^{2l}-p^l+1}$ over $\mathbb{F}_{p^n}$

We give the following result on the  $c$ -differential uniformity of  $f(x) = x^{p^{3l}-p^{2l}+p^l}$  for  $c \in \mathbb{F}_{p^n} \setminus \mu_{p^l+1}$ , where  $n = 4l$ . For  $c \in \mathbb{F}_{p^n} \setminus \mu_{p^l+1}$ , the  $c$ -differential uniformity of  $x^{p^{2l}-p^l+1}$  and  $x^{p^{3l}-p^{2l}+p^l}$  are the same.

**Theorem 2.** *Let  $d = p^{3l} - p^{2l} + p^l$ . For  $c \in \mathbb{F}_{p^n} \setminus \mu_{p^l+1}$ , we have  ${}_c\Delta_f \leq (p^l + 1)^2$ .*

*Proof.* For a fixed  $c \in \mathbb{F}_{p^n} \setminus \mu_{p^l+1}$ , we mainly study the number of solutions in  $\mathbb{F}_{p^n}$  of

$$\Delta(x) = (x+1)^d - cx^d = b \quad (22)$$

for any  $b \in \mathbb{F}_{p^n}$ . Denote by  $\delta_c(b)$  the number of solutions of (22) in  $\mathbb{F}_{p^n}$ . Let  $\alpha = \psi^{(p^n-1)/(p^l+1)}$ . If  $x \in \mathbb{F}_{p^n}^\#$ , we can distinguish  $(p^l + 1)^2$  disjoint cases:

	Set	Equation (22)	$x+1$	$x$
Case 1	$S_{0,0}$	$x+1 - cx = b$	$\frac{b-c}{1-c}$	$\frac{b-1}{1-c}$
Case 2	$S_{0,1}$	$x+1 - \alpha cx = b$	$\frac{b-\alpha c}{1-\alpha c}$	$\frac{b-1}{1-\alpha c}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $p^l + 1$	$S_{0,p^l}$	$x+1 - \alpha^{p^l} cx = b$	$\frac{b-\alpha^{p^l} c}{1-\alpha^{p^l} c}$	$\frac{b-1}{1-\alpha^{p^l} c}$
Case $p^l + 2$	$S_{1,0}$	$\alpha(x+1) - cx = b$	$\frac{b-c}{\alpha-c}$	$\frac{b-\alpha}{\alpha-c}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $2p^l + 2$	$S_{1,p^l}$	$\alpha(x+1) - \alpha^{p^l} cx = b$	$\frac{b-\alpha^{p^l} c}{\alpha-\alpha^{p^l} c}$	$\frac{b-\alpha}{\alpha-\alpha^{p^l} c}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $p^{2l} + p^l + 1$	$S_{p^l,0}$	$\alpha^{p^l}(x+1) - cx = b$	$\frac{b-c}{\alpha^{p^l}-c}$	$\frac{b-\alpha^{p^l}}{\alpha^{p^l}-c}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
Case $(p^l + 1)^2$	$S_{p^l,p^l}$	$\alpha^{p^l}(x+1) - \alpha^{p^l} cx = b$	$\frac{b-\alpha^{p^l} c}{\alpha^{p^l}-\alpha^{p^l} c}$	$\frac{b-\alpha^{p^l}}{\alpha^{p^l}-\alpha^{p^l} c}$

Table 3: Equation (22) with  $x \in \mathbb{F}_{p^n}^\#$

From Table 3, we have  $\delta_c(b) \leq (p^l + 1)^2$  for  $b \neq 1, c$  since  $\Delta(0) = 1$  and  $\Delta(-1) = c$ . Note that  $\Delta(x) = 1$  has no solution in  $\bigsqcup_{0 \leq i \leq p^l} S_{0,i}$ , and  $\Delta(x) = c$  has no solution in  $\bigsqcup_{0 \leq i \leq p^l} S_{i,0}$  and so,  $\delta_c(1), \delta_c(c) \leq (p^l + 1)^2$ . Then, according to  $\gcd(d, p^n - 1) \leq 3$  and lemma 1, we get  ${}_c\Delta_f \leq (p^l + 1)^2$ .  $\square$

## 4 On Exponential sum

In [3, 4], Hellesteth determined the value distribution of the cross correlation function for  $d = p^{3l} - p^{2l} + p^l$  with  $n = 4l$ ,  $p^l \not\equiv 2 \pmod{3}$ . So what happens if  $p^l \equiv 2 \pmod{3}$ ? In this section, we give some relevant

conclusions. We fix  $n = 4l$ , where  $l$  is a positive integer. Let  $d_1 = p^{3l} - p^{2l} + p^l$ , where  $p^l \equiv 2 \pmod{3}$ . Define

$$S_{d_1}(u, v) = \sum_{x \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(ux^{d_1} - vx)}, \quad (23)$$

where  $(u, v) \in \mathbb{F}_{p^n} \times \mathbb{F}_{p^n}$ ,  $\zeta_p = e^{\frac{2\pi\sqrt{-1}}{p}}$  is a primitive complex  $p$ -th root of unity, and  $\text{Tr}_1^n(x) = \sum_{i=0}^{n-1} x^{p^i}$  denotes the absolute trace mapping from  $\mathbb{F}_{p^n}$  to  $\mathbb{F}_p$ . Define

$$\begin{aligned} C_0 &= \left\{ x \in \mathbb{F}_{p^n} \mid x = y^{p^l+1} \text{ for some } y \in \mathbb{F}_{p^n} \right\}, \\ C_\infty &= \{0\}, \\ C_1 &= \mathbb{F}_{p^n} \setminus (C_0 \cup C_\infty). \end{aligned}$$

In order to determine the values of  $S_{d_1}(u, v)$ . We need one result without proof for which the readers can refer to [19].

**Lemma 5.** *Suppose  $n = 4l$  and  $\zeta_p$  be a complex  $p$ -th root of unity. Then*

$$\sum_{y \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(ay^{p^l+1})} = \begin{cases} p^n & \text{if } a \in C_\infty, \\ -p^{3l} & \text{if } a \in C_0, \\ p^{2l} & \text{if } a \in C_1. \end{cases}$$

The following result is essential to determine the value of  $S_{d_1}(u, v)$ .

**Lemma 6.** *Let  $d_1 = p^{3l} - p^{2l} + p^l$ , where  $p^l \equiv 2 \pmod{3}$ . For each fixed  $u \in \mathbb{F}_{p^n}^*$ , we have*

$$\{S_{d_1}(u, v) : v \in \mathbb{F}_{p^n}^*\} = \begin{cases} \{S_{d_1}(1, v) : v \in \mathbb{F}_{p^n}^*\}, & \text{if } u \text{ is a cube in } \mathbb{F}_{p^n}^*, \\ \{S_{d_1}(\psi, v) : v \in \mathbb{F}_{p^n}^*\}, & \text{if } u \text{ is a non-cube in } \mathbb{F}_{p^n}^*. \end{cases}$$

*Proof.* Since  $\gcd(d_1, p^n - 1) = 3$ , there exist two integers  $u_1$  and  $v_1$  such that  $u_1 d_1 + v_1(p^n - 1) = 3$ . Then for any  $x \in \mathbb{F}_{p^n}^*$ , we have  $x^3 = x^{u_1 d_1}$ . Moreover, note that every  $u \in \mathbb{F}_{p^n}^*$  can be written uniquely as  $u = \psi^{3k_1+j}$  for some  $0 \leq k_1 < (p^n - 1)/3$  and  $0 \leq j \leq 2$ . Replacing  $x$  by  $\psi^{-k_1} u_1 x$  in  $S_{d_1}(u, v)$  gives

$$S_{d_1}(u, v) = \begin{cases} S_{d_1}(1, v\psi^{-k_1} u_1), & \text{if } j = 0, \\ S_{d_1}(\psi, v\psi^{-k_1} u_1), & \text{if } j = 1. \end{cases}$$

If  $j = 2$ , denote  $p = 3k_2 + 2$ ,  $y = \psi^{(k_1 - k_2)u_1} x$  and  $z = y^{p^{n-1}}$ , then

$$\begin{aligned} S_{d_1}(u, v) &= \sum_{x \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(ux^{d_1} - vx)} \\ &= \sum_{y \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(\psi^p y^{d_1} - v\psi^{(k_2 - k_1)u_1} y)} \\ &= \sum_{z \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(\psi z^{d_1} - v p^{n-1} \psi^{p^{n-1}(k_2 - k_1)u_1} z)} \\ &= S_{d_1}(\psi, v p^{n-1} \psi^{p^{n-1}(k_2 - k_1)u_1}). \end{aligned}$$

Because  $v\psi^{-ku_1}$  and  $v^{p^{n-1}}\psi^{p^{n-1}(k_2-k_1)u_1}$  all run through  $\mathbb{F}_{p^n}^*$  as  $v$  runs through  $\mathbb{F}_{p^n}^*$ , we can obtain the result.  $\square$

We define

$$n_k(u, v) = \#\{j \mid u\psi^{d_1 j} - v\psi^j \in C_k, 0 \leq j \leq p^l\}$$

for  $k = 0, 1, \infty$  and  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ .

**Lemma 7.** *Let  $n = 4l$ , where  $l$  is a positive integer. Let  $d_1 = p^{3l} - p^{2l} + p^l$ , where  $p^l \equiv 2 \pmod{3}$ . Let  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ , then the following holds*

(i)  $n_\infty(u, v) = 1$  if  $(\frac{v}{u})^{p^l+1} = 1$  and  $n_\infty(u, v) = 0$  otherwise.

(ii)  $n_0(u, v) + n_\infty(u, v) \leq 3$ .

(iii) If  $u = 1$  and  $n_\infty(u, v) = 1$ , then  $n_0(u, v) = 0$  for  $v = 1, \psi^{(d_1-1)\frac{p^l+1}{3}}, \psi^{2(d_1-1)\frac{p^l+1}{3}}$  and  $n_0(u, v) = 1$  otherwise.

(iv) If  $u = \psi$  and  $n_\infty(u, v) = 1$ , then  $n_0(u, v) = 1$ .

*Proof.* (i) If  $u\psi^{d_1 j} - v\psi^j = 0$ , then  $\psi^{(d_1-1)j} = \frac{v}{u}$ . Note that

$$d_1 - 1 = p^{3l} - p^{2l} + p^l - 1 = \frac{p^n - 1}{p^l + 1}.$$

It follows that  $n_\infty(u, v) = 1$  if  $(\frac{v}{u})^{p^l+1} = 1$  and  $n_\infty(u, v) = 0$  otherwise.

(ii) Let  $T(u, v) = n_0(u, v) + n_\infty(u, v)$ . Since there are  $n_\infty(u, v)$  solutions corresponding to  $ux^{d_1} - vx = 0$  and  $n_0(u, v)$  solutions corresponding to  $ux - vx^{d_1} \in C_0$ . Then  $T(u, v)$  is the number of solutions of

$$(ux^{d_1} - vx)^{p^{3l}+p^l} - (ux^{d_1} - vx)^{p^{2l}+1} = 0$$

where  $x = \psi^j$ ,  $0 \leq j \leq p^l$ . Straightforward calculation implies that  $T(u, v)$  is the number of solutions of

$$u^{p^{2l}+1}x^{3(d_1-1)} - (v^{p^{3l}+p^l} + v^{p^{2l}}u + vu^{p^{2l}})x^{2(d_1-1)} + (v^{p^{2l}+1} + v^{p^{3l}}u^{p^l} + v^{p^l}u^{p^{3l}})x^{d_1-1} - u^{p^{3l}+p^l} = 0,$$

where  $x = \psi^j$ ,  $0 \leq j \leq p^l$ . Therefore, by letting  $y = x^{d_1-1}$ , we obtain that  $T(u, v)$  is the number of solutions to

$$\begin{cases} u^{p^{2l}+1}y^3 - (v^{p^{3l}+p^l} + v^{p^{2l}}u + vu^{p^{2l}})y^2 + (v^{p^{2l}+1} + v^{p^{3l}}u^{p^l} + v^{p^l}u^{p^{3l}})y - u^{p^{3l}+p^l} = 0. \\ y^{p^l+1} = 1. \end{cases}$$

It follows that  $n_0(u, v) + n_\infty(u, v) \leq 3$ .

(iii) In the case  $n_\infty(u, v) = 1$ , it follows from (1) that  $(\frac{v}{u})^{p^l+1} = 1$ . This implies that  $u^{p^l+1} = v^{p^l+1}$ .

*Case 1:*  $u = v = 1$ . Then  $T(u, v)$  is the number of common solutions of  $y^{p^l+1} = 1$  and

$$y^3 - 3y^2 + 3y - 1 = (y - 1)^3 = 0.$$

So, we obtain  $T(u, v) = n_0(u, v) + n_\infty(u, v) = 1$ .

*Case 2:*  $v = \psi^{(d_1-1)j}$ ,  $u = \psi^{(d_1-1)j}$ , where  $d_1 - 1 = \frac{p^l-1}{p^l+1}$  for  $1 \leq j \leq p^l$ . Then  $T(u, v)$  is the number of common solutions of  $y^{p^l+1} = 1$  and

$$\begin{aligned} & y^3 - (\psi^{-2j(d_1-1)} + 2\psi^{j(d_1-1)})y^2 + (\psi^{2j(d_1-1)} + 2\psi^{-j(d_1-1)})y - 1 \\ &= (y - \psi^{-2j(d_1-1)})(y - \psi^{j(d_1-1)})^2 \\ &= 0. \end{aligned}$$

Then  $n_0(u, v) = 0$  for  $j = \frac{p^l+1}{3}, \frac{2(p^l+1)}{3}$  and  $n_0(u, v) = 1$  otherwise.

(iv) In the case  $n_\infty(u, v) = 1$ , it follows from (1) that  $(\frac{v}{u})^{p^l+1} = 1$ . This implies that  $u^{p^l+1} = v^{p^l+1}$ .

*Case 1:*  $v = u = \psi$ . Then  $T(a, b)$  is the number of common solutions of  $y^{p^l+1} = 1$  and

$$\begin{aligned} & \psi^{p^{2l}+1}y^3 - (\psi^{p^{3l}+p^l} + 2\psi^{p^{2l}+1})y^2 + (\psi^{p^{2l}+1} + 2\psi^{p^{3l}+p^l})y - \psi^{p^{3l}+p^l} \\ &= (\psi^{p^{2l}+1}y - \psi^{p^{3l}+p^l})(y - 1)^2 \\ &= 0. \end{aligned}$$

So, we have  $T(u, v) = n_0(u, v) + n_\infty(u, v) = 2$ .

*Case 2:*  $v = \psi^{(d_1-1)j}u = \psi^{(d_1-1)j}\psi$ , where  $d_1 - 1 = \frac{p^{4l}-1}{p^l+1}$ ,  $j = 1, 2, \dots, p^l$ . Then  $T(u, v)$  is the number of common solutions of  $y^{p^l+1} = 1$  and

$$\begin{aligned} & \psi^{p^{2l}+1}y^3 - (\psi^{-2j(d_1-1)}\psi^{p^{3l}+p^l} + 2\psi^{j(d_1-1)}\psi^{p^{2l}+1})y^2 \\ &+ (\psi^{2j(d_1-1)}\psi^{p^{2l}+1} + 2\psi^{-j(d_1-1)}\psi^{p^{3l}+p^l})y - \psi^{p^{3l}+p^l} \\ &= (\psi^{p^{2l}+1}y - \psi^{-2j(d_1-1)}\psi^{p^{3l}+p^l})(y - \psi^{j(d_1-1)})^2 \\ &= 0. \end{aligned}$$

So, we get  $T(u, v) = n_0(u, v) + n_\infty(u, v) = 2$ . □

Sum it up,  $S_{d_1}(u, v)$  can be explicitly determined.

**Theorem 3.** *Let  $p$  be a prime number and  $p^l \equiv 2 \pmod{3}$ . Let  $d_1 = p^{3l} - p^{2l} + p^l$ . When  $(u, v)$  runs through  $\mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ ,  $S_{d_1}(u, v)$  takes values in*

$$\{-2p^{2l}, -p^{2l}, 0, p^{2l}, p^{3l} - p^{2l}, p^{3l}\}.$$

*Proof.* Note that every  $x \in \mathbb{F}_{p^n}^*$  can be written as  $x = \psi^{(p^l+1)k+j}$ , where  $0 \leq k < \frac{p^n-1}{p^l+1} = K$  and

$0 \leq j \leq p^l$ . From  $\psi^{(p^l+1)d} = \psi^{p^l+1}$  and Lemma 5 we have

$$\begin{aligned}
S_{d_1}(u, v) &= \sum_{x \in \mathbb{F}_{p^n}} \zeta_p^{\text{Tr}_1^n(ux^{d_1} - vx)} \\
&= \sum_{j=0}^{p^l} \sum_{k=0}^K \zeta_p^{\text{Tr}_1^n(u\psi^{(p^l+1)d_1 k + dj} - v\psi^{(p^l+1)k+j})} + 1 \\
&= \sum_{j=0}^{p^l} \sum_{k=0}^K \zeta_p^{\text{Tr}_1^n(\psi^{(p^l+1)k}(u\psi^{d_1 j} - v\psi^j))} + 1 \\
&= \frac{1}{p^l + 1} (p^{4l} n_\infty(u, v) - p^{3l} n_0(u, v) + p^{2l} n_1(u, v)).
\end{aligned}$$

It is therefore sufficient to find the distribution of  $(n_\infty(u, v), n_0(u, v), n_1(u, v))$  when  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ . According to Lemma 7, when  $u = 1$  or  $\psi$ , the triplet  $(n_\infty(u, v), n_0(u, v), n_1(u, v))$  and the value of  $S_{d_1}(u, v)$  have the following possibilities:

$n_\infty(u, v)$	$n_0(u, v)$	$n_1(u, v)$	$S_{d_1}(u, v)$
0	0	$p^l + 1$	$-1 + p^{2l}$
0	1	$p^l$	$-1$
0	2	$p^l - 1$	$-1 - p^{2l}$
0	3	$p^l - 2$	$-1 - 2p^{2l}$
1	0	$p^l$	$-1 + p^{3l}$
1	1	$p^l - 1$	$-1 + p^{3l} - p^{2l}$

By Lemma 6, the desired result follows. □

To determine the multiplicity on each value of  $S_{d_1}(u, v)$  for  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ . We also need several moment identities.

**Lemma 8.** *For the exponential sum  $S_{d_1}(u, v)$ , the following identities hold:*

$$(i) \quad \sum_{u, v \in \mathbb{F}_{p^n}} S_{d_1}(u, v) = p^{2n}.$$

$$(ii) \quad \sum_{u, v \in \mathbb{F}_{p^n}} S_{d_1}(u, v)^2 = p^{3n}.$$

$$(iii) \quad \sum_{u, v \in \mathbb{F}_{p^n}} S_{d_1}(u, v)^3 = p^{13l} + p^{3n} - p^{9l}.$$

*Proof.* The proof of (i) and (ii) is obvious and we omit it here. For (iii), we can calculate

$$\sum_{u, v \in \mathbb{F}_{p^n}} S_{d_1}(u, v)^3 = p^{2n} \cdot (3p^n - 2 + (p^n - 1) \cdot M_1)$$

where  $M_1$  is the number of solutions to

$$\begin{cases} y_1 + y_2 + 1 = 0, \\ y_1^{d_1} + y_2^{d_1} + 1 = 0, \end{cases}$$

satisfying  $y_1 y_2 \neq 0$ . Since  $d_1 = p^{3l} - p^{2l} + p^l$ , this equals the number of solutions of

$$y^{p^{3l}+p^l}(-y-1)^{p^{2l}} + (y+1)^{p^{3l}+p^l}y^{p^{2l}} - y^{p^{2l}}(y+1)^{p^{2l}} = 0$$

in  $\mathbb{F}_{p^n} \setminus \{0, -1\}$ , and therefore

$$(y^{p^l} - y)^{p^{2l}+p^l} = 0$$

which gives  $M_1 = p^l - 2$ . Then

$$\sum_{u, v \in \mathbb{F}_{p^n}} S_{d_1}(u, v)^3 = p^{2n}(3p^n - 2 + (p^n - 1)(p^l - 2)) = p^{13l} + p^{3n} - p^{9l}.$$

□

Based on Theorem 3, Lemmas 7 and 8, the value distribution of  $S_{d_1}(u, v)$  is given as follows.

**Theorem 4.** *Let  $p$  is a prime number and  $p^l \equiv 2 \pmod{3}$ . Let  $d_1 = p^{3l} - p^{2l} + p^l$ . Then the value distribution of the exponential sum  $S_{d_1}(u, v)$  with  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$  can be described as follows.*

$$S_{d_1}(u, v) = \begin{cases} -2p^{2l}, & \text{occurs } (p^{8l} - 3p^{7l} + 3p^{6l} - p^{5l} - p^{4l} + 3p^{3l} - 3p^{2l} + p^l)/6 \text{ times,} \\ -p^{2l}, & \text{occurs } p^{7l} - p^{6l} - p^{3l} + p^{2l} \text{ times,} \\ 0, & \text{occurs } (p^{8l} - p^{7l} + p^{6l} - p^{5l} - 3p^{4l} + p^{3l} - p^{2l} + p^l + 2)/2 \text{ times,} \\ p^{2l}, & \text{occurs } (p^{8l} - p^{5l} - p^{4l} + p^l)/3 \text{ times,} \\ p^{3l} - p^{2l}, & \text{occurs } p^l(p^{4l} - 1) \text{ times,} \\ p^{3l}, & \text{occurs } p^{4l} - 1 \text{ times.} \end{cases}$$

*Proof.* When  $(u, v)$  runs through  $\mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ , the possible values of  $S_d(u, v)$  are confirmed in Theorem 3. Suppose  $-2p^{2l}$  occurs  $E_0$  times,  $-p^{2l}$  occurs  $E_1$  times,  $0$  occurs  $E_2$  times,  $p^{2l}$  occurs  $E_3$  times,  $p^{3l} - p^{2l}$  occurs  $E_4$  times,  $p^{3l}$  occurs  $E_5$  times. Then firstly we have

$$E_0 + E_1 + E_2 + E_3 + E_4 + E_5 = p^n(p^n - 1). \quad (24)$$

Secondly, by Lemma 8 we can derive

$$-2E_0 - E_1 + E_3 + (p^l - 1) \cdot E_4 + p^l \cdot E_5 = p^{6l} - p^{2l} \quad (25)$$

$$4E_0 + E_1 + E_3 + (p^l - 1)^2 \cdot E_4 + p^{2l} \cdot E_5 = p^{2n} - p^n \quad (26)$$

$$-8E_0 - E_1 + E_3 + (p^l - 1)^3 \cdot E_4 + p^{3l} \cdot E_5 = p^{7l} - p^{3l}. \quad (27)$$

From Lemmas 6, 7 and Theorem 3, we can derive

$$E_4 = p^l(p^{4l} - 1) \quad (28)$$

$$E_5 = (p^{4l} - 1). \quad (29)$$

Solving the system equations consisting of (24)-(29) we obtain the value distribution of  $S_{d_1}(u, v)$  for  $(u, v) \in \mathbb{F}_{p^n}^* \times \mathbb{F}_{p^n}$ . □

Let  $n = 4l$  and  $p^l \equiv 2 \pmod{3}$ . Let  $h_1(x)$  and  $h_2(x)$  be the minimal polynomials of  $\psi^{-1}$  and  $\psi^{-(p^{3l}-p^{2l}+p^l)}$  over  $\mathbb{F}_p$ , respectively. Let  $\mathcal{C}$  be the cyclic code with parity-check polynomial  $h_1(x)h_2(x)$ . By Delsarte's Theorem [20], the cyclic code  $\mathcal{C}$  can be expressed as

$$\mathcal{C} = \left\{ c_{u,v} = \left( \text{Tr}_1^n (u\psi^{i(p^{3l}-p^{2l}+p^l)} + v\psi^i) \right)_{i=0}^{p^n-2} \mid u, v \in \mathbb{F}_{p^n} \right\}.$$

Denote code  $\mathcal{C}'$  by

$$\mathcal{C}' = \left\{ c'_{u,v} = \left( \text{Tr}_1^n (u\psi^{j(p^{3l}-p^{2l}+p^l)} + v\psi^j) \right)_{j=0}^{\frac{p^n-p}{p-1}} \mid u, v \in \mathbb{F}_{p^n} \right\}.$$

Based on the result in Theorem 4, the weight distribution of  $\mathcal{C}'$  is determined as follows.

**Theorem 5.** *The  $\mathcal{C}'$  is a  $\left[ \frac{p^n-1}{p-1}, 2n, p^{n-1} - p^{3l-1} \right]$  cyclic code. Furthermore, the weight distribution of  $\mathcal{C}'$  is given in Table 4.*

Weight	Number of codewords
0	1
$p^{n-1} - p^{3l-1}$	$p^n - 1$
$p^{n-1} - p^{3l-1} + p^{2l-1}$	$p^l(p^n - 1)$
$p^{n-1} - p^{2l-1}$	$(p^{2n} - p^{5l} - p^n + p^l)/3$
$p^{n-1}$	$(p^{2n} - p^{7l} + p^{6l} - p^{5l} - p^n + p^{3l} - p^{2l} + p^l)/2$
$p^{n-1} + p^{2l-1}$	$p^{7l} - p^{6l} - p^{3l} + p^{2l}$
$p^{n-1} + 2p^{2l-1}$	$(p^{2n} - 3p^{7l} + 3p^{6l} - p^{5l} - p^n + 3p^{3l} - 3p^{2l} + p^l)/6$

Table 4: The weight distribution of  $\mathcal{C}'$

*Proof.* It is obvious that dimension of  $\mathcal{C}$  is  $2n$ . For the codeword  $c_{u,v}$  of  $\mathcal{C}$ , we also calculate its Hamming weight by using exponential sum:

$$\begin{aligned} \omega_H(c_{u,v}) &= p^n - 1 - \# \left\{ x \in \mathbb{F}_{p^n}^* \mid \text{Tr}_1^n (ux^{p^{3l}-p^{2l}+p^l} + vx) = 0 \right\} \\ &= p^n - \frac{1}{p} \sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_{p^n}} \omega_p^{y \text{Tr}_1^n (ux^{p^{3l}-p^{2l}+p^l} + vx)} \\ &= p^{n-1}(p-1) - \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_{p^n}} \omega_p^{\text{Tr}_1^n (u(xy)^{p^{3l}-p^{2l}+p^l} + vxy)} \\ &= p^{n-1}(p-1) - \frac{p-1}{p} \sum_{x \in \mathbb{F}_{p^n}} \omega_p^{\text{Tr}_1^n (ux^{p^{3l}-p^{2l}+p^l} + vx)}. \end{aligned}$$



Therefore, the weight distribution of cyclic code  $\mathcal{C}$  is directly determined by the value distribution of the exponential sum  $\sum_{x \in \mathbb{F}_{p^n}} \omega_p^{\text{Tr}_1^n(ux^{p^{3l}-p^{2l}+p^l}+vx)}$ . If  $u = 0$ , we have

$$\omega_H(c_{0,v}) = \begin{cases} 0 & \text{if } v = 0, \\ p^{n-1}(p-1) & \text{if } v \neq 0. \end{cases}$$

If  $u \neq 0$ , the value distribution of  $\omega_H(c_{u,v})$  follows from Theorem 4. Let  $\beta = \psi^{\frac{p^n-1}{p-1}}$  be a primitive element in  $\mathbb{F}_p$ . Note that

$$\text{Tr}_1^n \left( u\psi^{(j+k\frac{p^n-1}{p-1})(p^{3l}-p^{2l}+p^l)} + v\psi^{j+k\frac{p^n-1}{p-1}} \right) = \beta^k \cdot \text{Tr}_1^n \left( u\psi^{j(p^{3l}-p^{2l}+p^l)} + v\psi^j \right), \quad 0 \leq k \leq p-2$$

implies

$$c_{u,v} = (c'_{u,v}, \beta c'_{u,v}, \dots, \beta^{p-2} c'_{u,v}),$$

which yields the weight distribution of  $\mathcal{C}'$ . □

**Remark 1.**  $\mathcal{C}'$  is  $\beta^{-1}$ -constacyclic code.

In what follows, we provide some examples to verify our results.

**Example 1.** Let  $p = 5$ ,  $n = 4$ . Then  $\mathcal{C}'$  is a 5-ary cyclic code with parameters  $[156, 8, 100]$ . The weight enumerator of  $\mathcal{C}'$  is

$$1 + 624z^{100} + 3120z^{105} + 128960z^{120} + 162240z^{125} + 62400z^{130} + 33280z^{135}.$$

**Example 2.** Let  $p = 2$ ,  $n = 12$ . Then  $\mathcal{C}'$  is a binary cyclic code with parameters  $[4095, 24, 1792]$ . The weight enumerator of  $\mathcal{C}'$  is

$$1 + 4095z^{1792} + 32760z^{1824} + 5580120z^{2016} + 7452900z^{2048} + 1834560z^{2080} + 1872780z^{2112}.$$

**Example 3.** Let  $p = 11$ ,  $n = 4$ . Then  $\mathcal{C}'$  is a 11-ary cyclic code with parameters  $[1464, 8, 1210]$ . The weight enumerator of  $\mathcal{C}'$  is

$$1 + 14640z^{1210} + 161040z^{1221} + 71394400z^{1320} + 98234400z^{1331} + 17714400z^{1342} + 26840000z^{1353}.$$

## 5 Conclusion

In this paper, we completely calculate the differential spectrum of  $x^{p^{2l}-p^l+1}$ . We investigate the  $c$ -differential uniformity of the power mapping  $x^{p^{2l}-p^l+1}$ . We also calculate the value distribution of a class of exponential sum. In addition, we obtain a class of six-weight  $p$ -ary  $\beta^{-1}$ -constacyclic cyclic codes whose weight distribution can be explicitly determined.

## References

- [1] Wang, L., Luo, J. Rational points and zeta functions of some curves over finite fields. *Science China Mathematics* **53**, 2855-2863 (2010)
- [2] Blondeau, C., Canteaut, A., Charpin, P.: Differential properties of power functions. *Int. J. Inf. Coding Theory* **1**(2), 149-170 (2010)
- [3] Helleseht, T.: A note on the cross-correlation function between two binary maximal length linear sequence. *Discrete Math.* **23**(2) 301-307 (1978)
- [4] Helleseht, T.: Pairs of m-sequences with a six-valued cross-correlation. In: No, J.S., Song, H.Y., Helleseht, T., Kumar, P.V. (eds.) *Mathematical Properties of Sequences and Other Combinatorial Structures*, 1-6. Boston (2003)
- [5] Boura, C., Canteaut, A., Jean, J., Suder, V.: Two notions of differential equivalence on Sboxes. *Des. Codes Cryptogr* **87**(2), 185-202 (2019)
- [6] Nyberg, K.: Differentially uniform mappings for cryptography. In *Workshop on the Theory and Application of Cryptographic Techniques*. pp. 55-64. Berlin, Heidelberg: Springer Berlin Heidelberg. (1993)
- [7] Dobbertin, H., Helleseht, T., Kumar, P. V., Martinsen, H.: Ternary m-sequences with three-valued cross-correlation function: new decimations of Welch and Niho type. *IEEE Trans. Inf. Theory* **47**(4), 1473-1481 (2001)
- [8] Bracken C., Byrne E., Markin N., McGuire G.: New families of quadratic almost perfect nonlinear trinomials and multinomials. *Finite Fields and Appl* **14**(3), 703-714 (2008)
- [9] Budaghyan L., Carlet C., Helleseht T., Li N., Sun B.: On upper bounds for algebraic degrees of APN functions. *IEEE Trans. Inf. Theory* **64**(6), 4399-4411 (2017)
- [10] Tang, C., Ding, C., Xiong, M.: Codes, Differentially  $\delta$ -Uniform Functions, and  $t$ -Designs. *IEEE Trans. Inf. Theory* **66**(6), 3691-3703 (2019)
- [11] Carlet, C.: *Boolean functions for cryptography and coding theory*. Cambridge University Press, Cambridge (2021)
- [12] Carlet, C., Charpin, P., Zinoviev, V.: Codes, bent functions and permutations suitable for DES-like cryptosystems. *Des. Codes Cryptogr* **15**, 125-156 (1998)
- [13] Charpin, P., Peng, J.: Differential uniformity and the associated codes of cryptographic functions. *Adv. Math. Commun* **13**(4), 579-600 (2019)
- [14] Taniguchi, H.: On some quadratic APN functions. *Des. Codes Cryptogr* **87**(9), 1973-1983 (2019)
- [15] Carlet, C.: Characterizations of the differential uniformity of vectorial functions by the Walsh transform. *IEEE Trans. Inf. Theory* **64**(9), 6443-6453 (2017)
- [16] Borisov, N., Chew, M., Johnson, R., Wagner, D.: Multiplicative differentials. In *Fast Software Encryption: 9th International Workshop, FSE 2002 Leuven, Belgium, February 4-6, 2002 Revised Papers* 9. pp. 17-33. Springer Berlin Heidelberg. (2002)

- [17] Ellingsen, P., Felke, P., Riera, C., St?nic?, P., Tkachenko, A.: C-differentials, multiplicative uniformity, and (almost) perfect c-nonlinearity. *IEEE Trans. Inf. Theory* **66**(9), 5781-5789 (2020)
- [18] Mesnager, S., Riera, C., St?nic?, P., Yan, H., Zhou, Z.: Investigations on c-(almost) perfect nonlinear functions. *IEEE Trans. Inf. Theory* **67**(10), 6916-6925 (2021)
- [19] Baumert, L. D., McEliece, R. J.: Weights of irreducible cyclic codes. *Information and control*. **20**(2) 158-175 (1972)
- [20] Delsarte P.: On subfield subcodes of modified Reed-Solomon codes. *IEEE Trans. Inf. Theory* **21**(5), 575-576 (1975)
- [21] Eddahmani, S., Mesnager, S.: Explicit values of the DDT, the BCT, the FBCT, and the FBDT of the inverse, the gold, and the Bracken-Leander S-boxes. *Cryptogr. Commun* **14**(6), 1301-1344 (2022)
- [22] Xiong, M., Yan, H.: A note on the differential spectrum of a differentially 4-uniform power function. *Finite Fields and Their Applications*, 48, 117-125 (2017)
- [23] Blondeau, C., Canteaut, A., Charpin, P.: Differential properties of  $x \mapsto x^{2^t-1}$ . *IEEE Trans. Inf. Theory* **57**(12), 8127-8137 (2011)
- [24] Blondeau, C., Perrin, L.: More differentially 6-uniform power functions. *Des. Codes Cryptogr* **73**, 487-505 (2014)
- [25] Li, N., Wu, Y., Zeng, X., Tang, X.: A note on the differential spectrum of a class of power functions. In *2022 10th International Workshop on Signal Design and Its Applications in Communications (IWSDA)* (pp. 1-5). IEEE. (2022)
- [26] Xiong, M., Yan, H., Yuan, P.: On a conjecture of differentially 8-uniform power functions. *Des. Codes Cryptogr*, 86, 1601-1621 (2018)
- [27] Xie, X., Mesnager, S., Li, N., He, D., Zeng, X.: On the Niho type locally-APN power functions and their boomerang spectrum. *IEEE Trans. Inf. Theory* **69**(6), 4056-4064 (2022)
- [28] Choi, S. T., Hong, S., No, J. S., Chung, H.: Differential spectrum of some power functions in odd prime characteristic. *Finite Fields and Their Applications*, 21, 11-29 (2013)
- [29] Lei, L., Ren, W., Fan, C.: The differential spectrum of a class of power functions over finite fields. *Advances in Mathematics of Communications*. **15**(3), 525-537 (2021)
- [30] Yan, H., Zhou, Z., Weng, J., Wen, J., Helleseht, T., Wang, Q.: Differential spectrum of Kasami power permutations over odd characteristic finite fields. *IEEE Trans. Inf. Theory* **65**(10), 6819-6826 (2019)
- [31] Xia, Y., Zhang, X., Li, C., Helleseht, T.: The differential spectrum of a ternary power mapping. *Finite Fields Appl.* 64 1-16 (2020)
- [32] Yan, H., Xia, Y., Li, C., Helleseht, T., Xiong, M., Luo, J.: The Differential Spectrum of the Power Mapping  $x^{p^n-3}$ . *IEEE Trans. Inf. Theory* **68**(8) (2022)
- [33] Man, Y., Xia, Y., Li, C., Helleseht, T.: On the differential properties of the power mapping  $x^{p^m+2}$ . *Finite Fields Appl.* 84, 102100 (2022)

- [34] Jiang, S., Li, K., Li, Y., Qu, L.: (2022). Differential spectrum of a class of power functions. *Journal of Cryptologic Research.* **9**(3), 484-495 (2022)
- [35] Yan, H., Li, C.: Differential spectra of a class of power permutations with characteristic 5. *Des. Codes Cryptogr.* **89**, 1181-1191 (2021)
- [36] Yan, H., Mesnager, S., Tan, X.: On a class of APN power functions over odd characteristic finite fields: Their differential spectrum and c-differential properties. *Discrete Mathematics.* **347**(4), 113881 (2024)
- [37] Hu, Z., Li, N., Xu, L., Zeng, X., Tang, X.: The differential spectrum and boomerang spectrum of a class of locally-APN functions. *Des. Codes Cryptogra* **91**(5), 1695-1711 (2023)
- [38] Yan, H., Mesnager, S., Tan, X.: The complete differential spectrum of a class of power permutations over odd characteristic finite fields. *IEEE Trans. Inf. Theory* **69**(11), 7426-7438 (2023)
- [39] Yan, H., Li, Z.: A note on the differential spectrum of a class of power mappings with Niho exponent. *Cryptography and Communications* **14**(5), 1081-1089 (2022)
- [40] Tan, X., Yan, H. Differential spectrum of a class of APN power functions. *Des. Codes Cryptogr* **91**(8), 2755-2768 (2023)
- [41] Bao, F., Xia, Y., Chen, S., Li, C., Hellesteth, T.: The differential spectrum of a power permutation. *Advances in Mathematics of Communications.* (2024)
- [42] Tian, S., Cheng, Y.: Differential spectra for a class of power functions over  $\mathbb{F}_{p^n}$ . *J. Syst. Sci. Math. Sci.* **37**(5), 1351-1367 (2017)
- [43] Tu, Z., Zeng, X., Li, C., Hellesteth, T.: A class of new permutation trinomials. *Finite Fields and Their Appl.* **50** 178-195 (2018)