

A KASTLER-KALAU-WALZE TYPE THEOREM FOR THE J -TWIST OF THE DIRAC OPERATOR WITH TORSION

JIN HONG SIYAO LIU YONG WANG*

ABSTRACT. In this paper, we give a Lichnerowicz type formula for the J -twist of the Dirac operator with torsion. And we prove a Kastler-Kalau-Walze type theorem for the J -twist of the Dirac operator with torsion on 4-dimensional and 6-dimensional almost product Riemannian spin manifold with boundary.

1. INTRODUCTION

Noncommutative residues were first proposed in [1, 2]. Noncommutative residues are an important tool for studying noncommutative geometry. Therefore, much attention has been paid to noncommutative residues. Connes put forward that the noncommutative residue of the square of the inverse of the Dirac operator was proportioned to the Einstein-Hilbert action, which is called the Kastler-Kalau-Walze theorem now[3, 4]. Kastler gave a brute-force proof of this theorem[5]. In the same time, Kalau-Walze and Ackermann proved this theorem by using the normal coordinates system and the heat kernel expansion, respectively[6, 7]. The result of Connes was extended to the higher dimensional case [8]. Wang generalized the Connes' results to the case of manifolds with boundary in [9, 10], and proved the Kastler-Kalau-Walze type theorem for the Dirac operator and the signature operator on lower-dimensional manifolds with boundary. In [11], for the Dirac operator, Wang computed $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{-1}]$, in these cases the boundary term vanished. Most of the operators studied in the literature have the leading symbol $\sqrt{-1}c(\xi)$ [12, 13, 14, 15, 16].

Earlier some preliminaries and lemmas about the Dirac operator D and the J -twist are given in [17]. In [18, 19], the author got estimates on the higher eigenvalues of the Dirac operator on locally reducible Riemannian manifolds by the J -twist of the Dirac operator. It can be obtained by simple calculations that the leading symbol of the J -twist D_J of the Dirac operator is not $\sqrt{-1}c(\xi)$. Liu and Wang proved a Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator on 3-dimensional, 4-dimensional and 6-dimensional almost product Riemannian spin manifold with boundary in [22, 23]. On

2010 Mathematics Subject Classification: 53C40; 53C42.

Key words and phrases: Dirac operator; the J -twist of the Dirac operator with torsion; Kastler-Kalau-Walze type theorems.

* *Corresponding author.*

the other hand, Jean-Michel Bismut [24] proved a local index theorem for Dirac operators on a Riemannian manifold M associated with connections on TM which have non zero torsion. In [25], Ackermann and Tolksdorf proved a generalized version of the well-known Lichnerowicz formula for the square of the most general Dirac operator with torsion D_T on an even-dimensional spin manifold associated to a metric connection with torsion. Wang etc.[26] computed the lower dimensional volume $\widetilde{\text{Wres}}[\pi^+(D_T^*)^{-p_1} \circ \pi^+(D_T)^{-p_2}]$ and got a Kastler-Kalau-Walze type theorems associated with Dirac operators with torsion on compact manifolds with boundary. So it is natural to consider the J -twist of the Dirac operator with torsion which is the common generalization of J-twisted Dirac operator and the Dirac operator with torsion. And it is natural to give the Kastler-Kalau-Walze type theorem for the J-twisted Dirac operator with torsion. The motivation of this paper is to give the Kastler-Kalau-Walze type theorem for the J-twisted Dirac operator with torsion.

The paper is structured as follows. In Section 2, this paper will firstly introduce the basic notions of the almost product Riemannian manifold and the J -twist of the Dirac operator with torsion. We also give a Lichnerowicz type formula for the J -twist of the Dirac operator with torsion and a Kastler-Kalau-Walze type theorem for the J -twist of the Dirac operator with torsion on n -dimensional almost product Riemannian spin manifold without boundary. Next, we prove that the Kastler-Kalau-Walze type theorem on 4-dimensional and 6-dimensional almost product Riemannian spin manifold with boundary for the J -twist of the Dirac operator with torsion.

2. THE J -TWIST OF THE DIRAC OPERATOR WITH TORSION AND ITS LICHNEROWICZ FORMULA

We give some definitions and basic notions which we will use in this paper.

Let M be a smooth compact oriented Riemannian n -dimensional manifolds without boundary and N be a vector bundle on M . We say that P is a differential operator of Laplace type, if it has locally the form

$$(2.1) \quad P = -(g^{ij}\partial_i\partial_j + A^i\partial_i + B),$$

where ∂_i is a natural local frame on TM , $(g^{ij})_{1 \leq i, j \leq n}$ is the inverse matrix associated to the metric matrix $(g_{ij})_{1 \leq i, j \leq n}$ on M , A^i and B are smooth sections of $\text{End}(N)$ on M (endomorphism). If P satisfies the form (2.1), then there is a unique connection ∇ on N and a unique endomorphism E such that

$$P = -[g^{ij}(\nabla_{\partial_i}\nabla_{\partial_j} - \nabla_{\nabla_{\partial_i}^L\partial_j}) + E],$$

where ∇^L is the Levi-Civita connection on M . Moreover (with local frames of T^*M and N), $\nabla_{\partial_i} = \partial_i + \omega_i$ and E are related to g^{ij} , A^i and B through

$$\begin{aligned} \omega_i &= \frac{1}{2}g_{ij}(A^i + g^{kl}\Gamma_{kl}^j \text{id}), \\ E &= B - g^{ij}(\partial_i(\omega_j) + \omega_i\omega_j - \omega_k\Gamma_{ij}^k), \end{aligned}$$

where Γ_{kl}^j is the Christoffel coefficient of ∇^L .

Let M be a n -dimensional ($n \geq 3$) oriented compact Riemannian manifold with a Riemannian metric g^M . We recall that the Dirac operator with torsion D_T is locally given as following:

$$D_T = \sum_{i,j=1}^n g^{ij} c(\partial_i) \nabla_{\partial_j}^{S(TM),T} = \sum_{i=1}^n c(e_i) \nabla_{e_i}^{S(TM),T},$$

where $c(e_i)$ be the Clifford action which satisfies the relation

$$c(e_i)c(e_j) + c(e_j)c(e_i) = -2g^M(e_i, e_j) = -2\delta_i^j,$$

$$\nabla_{e_j}^{S(TM),T} = e_j + \frac{1}{4} \sum_{l,t=1}^n \langle \nabla_{e_j}^T e_l, e_t \rangle c(e_l)c(e_t),$$

$$\langle \nabla_{e_j}^T e_l, e_t \rangle = \langle \nabla_{e_j}^L e_l, e_t \rangle + T(e_j, e_l, e_t)$$

and

$$\sigma^j = \frac{1}{4} \sum_{l,t=1}^n \langle \nabla_{e_j}^L e_l, e_t \rangle c(e_l)c(e_t),$$

where T is a there form.

Let J be a $(1, 1)$ -tensor field on (M, g^M) such that $J^2 = \text{id}$,

$$g^M(J(X), J(Y)) = g^M(X, Y),$$

for all vector fields $X, Y \in \Gamma(TM)$. Here id stands for the identity map. (M, g^M, J) is an almost product Riemannian manifold. We can also define on almost product Riemannian spin manifold the following J -twist of the Dirac operator with torsion $D_{J,T}$ by

$$(2.2) \quad D_{J,T} := \sum_{j=1}^n c[e_j] \nabla_{J(e_j)}^{S(TM),T} = \sum_{j=1}^n c[J(e_j)] \nabla_{e_j}^{S(TM),T}.$$

$$(2.3) \quad T_J := \frac{1}{4} \sum_{j,l,t=1}^n T(e_j, e_l, e_t) c[J(e_j)] c(e_l)c(e_t).$$

Let $\xi = \sum_j \xi_j dx_j$, $\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k$, we denote that

$$\Gamma^k = \sum_{i,j} g^{ij} \Gamma_{ij}^k; \quad \sigma^j = \sum_i g^{ij} \sigma_i; \quad \partial^j = \sum_i g^{ij} \partial_i.$$

By (2.4), (2.5) in [17]

$$(2.4) \quad D_{J,T}^2 = -\frac{1}{8} \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l) c(e_i) c(e_j) c(e_k) c(e_l) - g^{ij} \partial_i \partial_j - 2\sigma^j \partial_j$$

$$\begin{aligned}
& + \Gamma^k \partial_k - g^{ij} [\partial_i (\sigma_j) + \sigma_i \sigma_j - \Gamma_{ij}^k \sigma_k] + \frac{1}{4} s + \sum_{\alpha, \beta=1}^n c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J) e_\beta] \nabla_{e_\beta}^S \\
& + \sum_{i,j} \{c[J(e_i)] \langle e_i, dx_l \rangle T_J \partial_l + T_J c[J(e_i)] \langle e_i, dx_l \rangle \partial_l\} \\
& + \sum_{i,j} \{c[J(e_i)] \langle e_i, dx_l \rangle \partial_l (T_J) + c[J(e_i)] \sigma_i T_J + T_J c[J(e_i)] \sigma_i\} + T_J^2.
\end{aligned}$$

where s is the scalar curvature, $e_i^* = g^M(e_i, \cdot)$ and $\langle X, dx_k \rangle = g^M(X, (dx_k)^*)$, for a vector field X . Then we have

$$\begin{aligned}
(2.5) \quad (\omega_i)_{D_{J,T}^2} &= \sigma_i - \frac{1}{2} \sum_{\alpha, p=1}^n g_{ip} c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J)(dx_p)^*] \\
&\quad - \frac{1}{2} \sum_{\alpha, p} g_{ip} c[J(e_\alpha)] \langle e_\alpha, dx_p \rangle T_J - \frac{1}{2} \sum_{\alpha, p} g_{ip} T_J c[J(e_\alpha)] \langle e_\alpha, dx_p \rangle,
\end{aligned}$$

(2.6)

$$\begin{aligned}
E_{D_{J,T}^2} &= \frac{1}{8} \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l) c(e_i) c(e_j) c(e_k) c(e_l) + \sum_{i,j=1}^n g^{ij} [\partial_i (\sigma_j) + \sigma_i \sigma_j - \Gamma_{ij}^k \sigma_k] \\
&\quad - \sum_{\alpha, j=1}^n c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J)(dx_j)^*] \sigma_j - \sum_{i,j=1}^n g^{ij} \left\{ \partial_i \left(\sigma_j - \frac{1}{2} \sum_{\nu, l=1}^n g_{jl} c[J(e_\nu)] c[(\nabla_{e_\nu}^L J)(dx_l)^*] \right. \right. \\
&\quad \left. \left. - \frac{1}{2} \sum_{\nu, l} g_{il} c[J(e_\nu)] \langle e_\nu, dx_l \rangle T_J - \frac{1}{2} \sum_{\nu, l} g_{il} T_J c[J(e_\nu)] \langle e_\nu, dx_l \rangle \right) \right. \\
&\quad \left. + \left(\sigma_i - \frac{1}{2} \sum_{\alpha, p=1}^n g_{ip} c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J)(dx_p)^*] - \frac{1}{2} \sum_{\alpha, p=1}^n g_{ip} c[J(e_\alpha)] \langle e_\alpha, dx_p \rangle T_J \right. \right. \\
&\quad \left. \left. - \frac{1}{2} \sum_{\alpha, p=1}^n g_{ip} T_J c[J(e_\alpha)] \langle e_\alpha, dx_p \rangle \right) \times \left(\sigma_j - \frac{1}{2} \sum_{\nu, l=1}^n g_{jl} c[J(e_\nu)] c[(\nabla_{e_\nu}^L J)(dx_l)^*] \right. \right. \\
&\quad \left. \left. - \frac{1}{2} \sum_{\nu, l=1}^n g_{jl} c[J(e_\nu)] \langle e_\nu, dx_l \rangle T_J - \frac{1}{2} \sum_{\nu, l} g_{jl} T_J c[J(e_\nu)] \langle e_\nu, dx_l \rangle \right) \right. \\
&\quad \left. - \sum_{k=1}^n \left(\sigma_k - \frac{1}{2} \sum_{\mu, h=1}^n g_{kh} c[J(e_\mu)] c[(\nabla_{e_\mu}^L J)(dx_h)^*] - \frac{1}{2} \sum_{\mu, h=1}^n g_{kh} c[J(e_\mu)] \langle e_\mu, dx_h \rangle T_J \right. \right. \\
&\quad \left. \left. - \frac{1}{2} \sum_{\mu, h=1}^n g_{kh} T_J c[J(e_\mu)] \langle e_\mu, dx_h \rangle \right) \Gamma_{ij}^k \right\} - \frac{1}{4} s - T_J^2 - \sum_{i=1}^n c[J(e_i)] \sigma_i T_J - \sum_{i=1}^n T_J c[J(e_i)] \sigma_i
\end{aligned}$$

$$- \sum_{\alpha,j}^n c[J(e_\alpha)] \langle e_\alpha, dx_j \rangle \partial_j(T_J).$$

Since E is globally defined on M , taking normal coordinates at x_0 , we have $\sigma^i(x_0) = 0$, $\Gamma^k(x_0) = 0$, $g^{ij}(x_0) = \delta_i^j$, $\partial^j(x_0) = e_j$, $\partial^j[c(\partial_j)](x_0) = 0$, $\nabla_{e_j}^L e_k(x_0) = 0$ and $\nabla_Y^S(c(X)) = c(\nabla_Y^L X)$, a simple calculation shows that

$$\begin{aligned} & \frac{1}{2} \sum_{\nu,j=1}^n \partial^j(c[J(e_\nu)]c[(\nabla_{e_\nu}^L J)(dx_j)^*])(x_0) \\ &= \frac{1}{2} \sum_{\nu,j=1}^n c[\nabla_{e_j}^L(J)e_\nu]c[(\nabla_{e_\nu}^L J)e_j](x_0) + \frac{1}{2} \sum_{\nu,j=1}^n c[J(e_\nu)]c[(\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j](x_0), \end{aligned}$$

so that

$$\begin{aligned} E_{D_{J,T}^2}(x_0) &= \frac{1}{8} \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l)c(e_i)c(e_j)c(e_k)c(e_l) \\ &+ \frac{1}{2} \sum_{\nu,j=1}^n c[\nabla_{e_j}^L(J)e_\nu]c[(\nabla_{e_\nu}^L J)e_j] + \frac{1}{2} \sum_{\nu,j=1}^n c[J(e_\nu)]c[(\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j] \\ &- \frac{1}{4} \sum_{\alpha,\nu,j=1}^n c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_j]c[J(e_\nu)]c[(\nabla_{e_\nu}^L J)e_j] - \frac{1}{4}s - T_J^2 \\ &- \frac{1}{4} \sum_{\hat{j},\hat{l},\hat{\alpha},j=1}^n [J(e_j)]T(e_{\hat{j}}, e_{\hat{l}}, e_{\hat{\alpha}})c[\nabla_{\partial_j}^L(J)e_{\hat{j}}]c(e_{\hat{l}})c(e_{\hat{\alpha}}) \\ &+ \frac{1}{4} \sum_{\hat{j},\hat{l},\hat{\alpha},j=1}^n T(e_{\hat{j}}, e_{\hat{l}}, e_{\hat{\alpha}}) \left\{ c[\nabla_{e_j}^L(J)e_j]c[J(e_{\hat{j}})]c(e_{\hat{l}})c(e_{\hat{\alpha}}) + c[J(e_j)]c[\nabla_{e_j}^L(J)e_{\hat{j}}]c(e_{\hat{l}})c(e_{\hat{\alpha}}) \right\} \\ &- \frac{1}{4} \sum_{\alpha,j=1}^n c[J(e_\alpha)]c[\nabla_{e_\alpha}^L(J)e_j] \left\{ c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\} \\ &- \frac{1}{4} \sum_{v,\alpha,j=1}^n c[J(dx_j)]T_J \left\{ c[J(e_v)]c[\nabla_{e_v}^L(J)e_j] + c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\} \\ &- \frac{1}{4} \sum_{v,\alpha,j=1}^n T_Jc[J(dx_j)] \left\{ c[J(e_v)]c[\nabla_{e_v}^L(J)e_j] + c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\}. \end{aligned}$$

We get the following Lichnerowicz formulas:

(2.7)

$$D_{J,T}^2 = -g^{ij}(\nabla_{\partial_i}\nabla_{\partial_j} - \nabla_{\nabla_{\partial_i}^L\partial_j}) - \frac{1}{8} \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l)c(e_i)c(e_j)c(e_k)c(e_l) + \frac{1}{4}s + T_J^2$$

$$\begin{aligned}
& -\frac{1}{2} \sum_{\nu,j=1}^n c[\nabla_{e_j}^L(J)e_\nu]c[(\nabla_{e_\nu}^L J)e_j] - \frac{1}{2} \sum_{\nu,j=1}^n c[J(e_\nu)]c[(\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j] \\
& + \frac{1}{4} \sum_{\alpha,\nu,j=1}^n c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_j]c[J(e_\nu)]c[(\nabla_{e_\nu}^L J)e_j] \\
& - \frac{1}{4} \sum_{\hat{j},\hat{l},\hat{\alpha},j=1}^n [J(e_j)]T(J(e_{\hat{j}}), e_{\hat{l}}, e_{\hat{\alpha}})c[\nabla_{e_{\hat{j}}}^L(J)e_{\hat{j}}]c(e_{\hat{l}})c(e_{\hat{\alpha}}) \\
& + \frac{1}{4} \sum_{\hat{j},\hat{l},\hat{\alpha},j=1}^n T(J(e_{\hat{j}}), e_{\hat{l}}, e_{\hat{\alpha}}) \left\{ c[\nabla_{e_{\hat{j}}}^L(J)e_{\hat{j}}]c[J(e_{\hat{j}})]c(e_{\hat{l}})c(e_{\hat{\alpha}}) + c[J(e_j)]c[\nabla_{e_j}^L(J)e_j]c(e_{\hat{l}})c(e_{\hat{\alpha}}) \right\} \\
& - \frac{1}{4} \sum_{\alpha,j=1}^n c[J(e_\alpha)]c[\nabla_{e_\alpha}^L(J)e_j] \left\{ c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\} \\
& - \frac{1}{4} \sum_{v,\alpha,j=1}^n c[J(dx_j)]T_J \left\{ c[J(e_v)]c[\nabla_{e_v}^L(J)e_j] + c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\} \\
& - \frac{1}{4} \sum_{v,\alpha,j=1}^n T_Jc[J(dx_j)] \left\{ c[J(e_v)]c[\nabla_{e_v}^L(J)e_j] + c[J(dx_j)]T_J + T_Jc[J(dx_j)] \right\}.
\end{aligned}$$

Now, let's calculate the specification of $D_{J,T}^3$.

(2.8)

$$\begin{aligned}
D_{J,T}^3 &= \sum_{i=1}^n c[J(e_i)] \langle e_i, dx_l \rangle (-g^{ij} \partial_l \partial_i \partial_j) + \sum_{i=1}^n c[J(e_i)] \langle e_i, dx_l \rangle \left[-(\partial_l g^{ij}) \partial_i \partial_j - g^{ij} (2\sigma_i \partial_j \right. \\
& - \Gamma_{ij}^k \partial_k) \partial_l + \sum_{\alpha,\beta,k=1}^n c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_\beta] \langle e_\beta, dx_k \rangle \partial_l \partial_k \left. \right] + \sum_{i=1}^n c[J(e_i)] \sigma_i (-g^{ij} \partial_i \partial_j) \\
& - T_J g^{ij} \partial_i \partial_j + \sum_{i=1}^n c[J(e_i)] \langle e_i, dx_l \rangle (c[J(e_i)] T_J + T_J c[J(e_i)]) \partial_i \partial_j \\
& + \sum_{i=1}^n c[J(e_i)] \langle e_i, dx_l \rangle \left[-2(\partial_l g^{ij}) \sigma_i \partial_j - 2g^{ij} (\partial_l \sigma_i) \partial_j + (\partial_l g^{ij}) \Gamma_{ij}^k \partial_k + g^{ij} (\partial_l \Gamma_{ij}^k) \partial_k \right. \\
& + \sum_{\alpha,\beta,k=1}^n \partial_l (c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_\beta]) \langle e_\beta, dx_k \rangle \partial_k + \sum_{\alpha,\beta,k=1}^n c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_\beta] (\partial_l \langle e_\beta, dx_k \rangle) \partial_k \\
& \left. + (\partial_l c[J(e_i)]) T_J + c[J(e_i)] (\partial_l T_J) + (\partial_l T_J) c[J(e_i)] + T_J (\partial_l c[J(e_i)]) \right] \\
& + \sum_{i=1}^n c[J(e_i)] \langle e_i, dx_l \rangle \partial_l \left[-\frac{1}{8} \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l) c(e_i) c(e_j) c(e_k) c(e_l) - g^{ij} ((\partial_i \sigma_j) \right.
\end{aligned}$$

$$\begin{aligned}
& + \sigma_i \sigma_j - \Gamma_{ij}^k \sigma_k) + \frac{1}{4} s + \sum_{\alpha, \beta, k=1}^n c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J) e_\beta] \langle e_\beta, dx_k \rangle \sigma_k + c[J(e_j)] (\partial_j T_J) \\
& + c[J(e_\alpha)] \sigma_\alpha T_J + T_J c[J(e_\alpha)] \sigma_\alpha + T_J^2 \left. + \left[\sum_{i=1}^n c[J(e_i)] \sigma_i + T_J \right] \right. \\
& \times \left[\frac{1}{4} s + (c[J(e_j)] T_J + T_J c[J(e_j)]) \partial_j - \frac{1}{8} \sum_{i, j, k, l=1}^n R(J(e_i), J(e_j), e_k, e_l) c(e_i) c(e_j) c(e_k) c(e_l) \right. \\
& - 2\sigma^j \partial_j + \Gamma^k \partial_k - g^{ij} ((\partial_i \sigma_j) + \sigma_i \sigma_j - \Gamma_{ij}^k \sigma_k) + \sum_{\alpha, \beta, k=1}^n c[J(e_\alpha)] c[(\nabla_{e_\alpha}^L J) e_\beta] \langle e_\beta, dx_k \rangle (\partial_k + \sigma_k) \\
& \left. \left. + c[J(e_j)] (\partial_j T_J) + c[J(e_\alpha)] \sigma_\alpha T_J + T_J c[J(e_\alpha)] \sigma_\alpha + T_J^2 \right] \right].
\end{aligned}$$

According to the detailed descriptions in [7], we know that the noncommutative residue of a generalized laplacian $\tilde{\Delta}$ is expressed as

$$(n-2)\Phi_2(\tilde{\Delta}) = \pi^{-\frac{n}{2}} \Gamma\left(\frac{n}{2}\right) \widetilde{res}(\tilde{\Delta}^{-\frac{n}{2}+1}),$$

where $\Phi_2(\tilde{\Delta})$ denotes the integral over the diagonal part of the second coefficient of the heat kernel expansion of $\tilde{\Delta}$. Now let $\tilde{\Delta} = D_{J,T}^2$. Since $D_{J,T}^2$ is a generalized laplacian, we can suppose $D_{J,T}^2 = \Delta - E$, then, we have

$$(2.9) \quad \text{Wres}(D_{J,T}^2)^{-\frac{n-2}{2}} = \frac{(n-2)\pi^{\frac{n}{2}}}{(\frac{n}{2}-1)!} \int_M \text{tr}(-\frac{1}{6}s + E_{D_{J,T}^2}) d\text{Vol}_M,$$

where Wres denote the noncommutative residue, tr denote tr. By

$$(2.10) \quad \text{tr}[c(X)c(Y)] = -g^M(X, Y)\text{tr}[\text{id}];$$

$$(2.11) \quad \text{tr}[c(X)c(Y)c(Z)c(W)] = g^M(X, W)g^M(Y, Z)\text{tr}[\text{id}] - g^M(X, Z)g^M(Y, W)\text{tr}[\text{id}] \\ + g^M(X, Y)g^M(Z, W)\text{tr}[\text{id}];$$

$$(2.12) \quad \text{tr}[c(X)c(Y)c(Z)c(W)c(V)c(U)] \\ = -g^M(X, U)g^M(Y, V)g^M(Z, W)\text{tr}[\text{id}] + g^M(X, U)g^M(Y, W)g^M(Z, V)\text{tr}[\text{id}] \\ - g^M(X, U)g^M(Y, Z)g^M(W, V)\text{tr}[\text{id}] + g^M(X, V)g^M(Y, U)g^M(Z, W)\text{tr}[\text{id}] \\ - g^M(X, V)g^M(Y, W)g^M(Z, U)\text{tr}[\text{id}] + g^M(X, V)g^M(Y, Z)g^M(W, U)\text{tr}[\text{id}] \\ - g^M(X, W)g^M(Y, U)g^M(Z, V)\text{tr}[\text{id}] + g^M(X, W)g^M(Y, V)g^M(Z, U)\text{tr}[\text{id}] \\ - g^M(X, W)g^M(Y, Z)g^M(V, U)\text{tr}[\text{id}] + g^M(X, Z)g^M(Y, U)g^M(W, V)\text{tr}[\text{id}] \\ - g^M(X, Z)g^M(Y, V)g^M(W, U)\text{tr}[\text{id}] + g^M(X, Z)g^M(Y, W)g^M(V, U)\text{tr}[\text{id}] \\ - g^M(X, Y)g^M(Z, U)g^M(W, V)\text{tr}[\text{id}] + g^M(X, Y)g^M(Z, V)g^M(W, U)\text{tr}[\text{id}]$$

$$-g^M(X, Y)g^M(Z, W)g^M(V, U)\text{tr}[\text{id}]$$

where $X, Y, Z, W, V, U \in \Gamma(TM)$. We calculate that

$$\begin{aligned}
(2.13) \quad & \sum_{i,j,k,l=1}^n \text{tr}[R(J(e_i), J(e_j), e_k, e_l)c(e_i)c(e_j)c(e_k)c(e_l))] \\
&= \sum_{i,j,k,l=1}^n R(J(e_i), J(e_j), e_k, e_l)\text{tr}[c(e_i)c(e_j)c(e_k)c(e_l)] \\
&= \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i)\text{tr}[\text{id}] - \sum_{i,j=1}^n R(J(e_i), J(e_j), e_i, e_j)\text{tr}[\text{id}] \\
&= 2 \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i)\text{tr}[\text{id}],
\end{aligned}$$

$$(2.14) \quad \sum_{\nu,j=1}^n \text{tr}[c[\nabla_{e_j}^L(J)e_\nu]c[(\nabla_{e_\nu}^L J)e_j]] = - \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j)\text{tr}[\text{id}],$$

$$\begin{aligned}
(2.15) \quad & \sum_{\nu,j=1}^n \text{tr}[c[J(e_\nu)]c[(\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j]] \\
&= - \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j)\text{tr}[\text{id}],
\end{aligned}$$

$$\begin{aligned}
(2.16) \quad & \sum_{\alpha,\nu,j=1}^n \text{tr}[c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)e_j]c[J(e_\nu)]c[(\nabla_{e_\nu}^L J)e_j]] \\
&= \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j)g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu))\text{tr}[\text{id}] \\
&\quad - \sum_{\nu,j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j)\text{tr}[\text{id}] \\
&\quad + \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j)g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j)\text{tr}[\text{id}].
\end{aligned}$$

By applying the formulae shown in (2.9)-(2.16), we get:

Theorem 2.1. *If M is a n -dimensional almost product Riemannian spin manifold without boundary, we have the following:*

$$\text{Wres}(D_{J,T}^{-n+2}) =$$

$$\begin{aligned}
& \frac{(n-2)\pi^{\frac{n}{2}}}{(\frac{n}{2}-1)!} \int_M 2^{\frac{n}{2}} \left\{ \left(\frac{1}{4} \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) - \frac{1}{2} \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right. \right. \\
& - \frac{1}{2} \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \\
& - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \\
& - \frac{1}{4} \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) + \frac{1}{4} \sum_{\nu,j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{12}s \\
& - \frac{1}{2} \sum_{j,m,p} T(e_m, e_p, J(e_m)) g^M((\nabla_{e_j}^L J)e_j, e_p) - \frac{1}{2} \sum_{j,m,\alpha} T(e_m, J(e_j), J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_m)) \\
& - \frac{1}{2} \sum_{j,m,\alpha} T(e_m, J(e_j), J(e_m)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\alpha)) + \frac{1}{2} \sum_{m,p,\alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_m, e_p) \\
& - \frac{1}{2} \sum_{m,p,\alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_m}^L J)e_\alpha, e_p) - \frac{1}{4} \sum_{j,p,m} T(e_j, e_p, J(e_m)) T(e_m, e_p, J(e_j)) \\
& \left. - \frac{1}{2} \sum_{j,l,\alpha} T^2(J(e_j), e_l, e_\alpha) \right\} d\text{Vol}_M,
\end{aligned}$$

where s is the scalar curvature.

3. A KASTLER-KALAU-WALZE TYPE THEOREM FOR 4-DIMENSIONAL MANIFOLDS WITH BOUNDARY

Firstly, we need the basic notions of Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary that will be used throughout the paper. For the details, see Ref.[11]. And we use the same definitions and symbols as in [11].

Let $U \subset M$ be a collar neighborhood of ∂M which is diffeomorphic with $\partial M \times [0, 1)$. By the definition of $h(x_n) \in C^\infty([0, 1))$ and $h(x_n) > 0$, there exists $\widehat{h} \in C^\infty((-\varepsilon, 1))$ such that $\widehat{h}|_{[0,1)} = h$ and $\widehat{h} > 0$ for some sufficiently small $\varepsilon > 0$. Then there exists a metric g' on $\widetilde{M} = M \cup_{\partial M} \partial M \times (-\varepsilon, 0]$ which has the form on $U \cup_{\partial M} \partial M \times (-\varepsilon, 0]$

$$g' = \frac{1}{\widehat{h}(x_n)} g^{\partial M} + dx_n^2,$$

such that $g'|_M = g$. We fix a metric g' on the \widetilde{M} such that $g'|_M = g$.

Definition 3.1. [11] *Lower dimensional volumes of almost product spin manifolds with boundary are defined by*

$$\text{Vol}_n^{(p_1, p_2)} M := \widetilde{\text{Wres}}[\pi^+ D_{J,T}^{-p_1} \circ \pi^+ D_{J,T}^{-p_2}],$$

where $\widetilde{\text{Wres}}$ denote the noncommutative residue for manifolds with boundary as in [20].

We can get the spin structure on \widetilde{M} by extending the spin structure on M . Let D be the Dirac operator associated to g' on the spinors bundle $S(T\widetilde{M})$.

By [11], we get

$$(3.1) \quad \widetilde{\text{Wres}}[\pi^+ D_{J,T}^{-p_1} \circ \pi^+ D_{J,T}^{-p_2}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-n}(D_{J,T}^{-p_1-p_2})]\sigma(\xi)dx + \int_{\partial M} \Phi$$

and

$$(3.2) \quad \Phi = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{tr}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D_{J,T}^{-p_1})(x', 0, \xi', \xi_n)] \\ \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D_{J,T}^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx',$$

where the sum is taken over $r+l-k-|\alpha|-j-1=-n$, $r \leq -p_1$, $l \leq -p_2$.

Since $[\sigma_{-n}(D_{J,T}^{-p_1-p_2})]|_M$ has the same expression as $\sigma_{-n}(D_{J,T}^{-p_1-p_2})$ in the case of manifolds without boundary, so locally we can compute the first term by [5], [6], [11], [21].

We firstly compute

$$(3.3) \quad \widetilde{\text{Wres}}[\pi^+ D_{J,T}^{-1} \circ \pi^+ D_{J,T}^{-1}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)}[\sigma_{-4}(D_{J,T}^{-2})]\sigma(\xi)dx + \int_{\partial M} \Psi,$$

where

$$(3.4) \quad \Psi = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \text{tr}_{S(TM)}[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D_{J,T}^{-1}) \\ (x', 0, \xi', \xi_n) \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D_{J,T}^{-1})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx',$$

the sum is taken over $r+l-k-j-|\alpha|-1=-4$, $r \leq -1$, $l \leq -1$.

Then we give the interior of $\widetilde{\text{Wres}}[\pi^+ D_{J,T}^{-1} \circ \pi^+ D_{J,T}^{-1}]$,

$$(3.5) \quad \int_M \int_{|\xi|=1} \text{tr}_{S(TM)}[\sigma_{-4}(D_{J,T}^{-2})]\sigma(\xi)dx = 2\pi^2 \\ \int_M \left\{ \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) - 2 \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right. \\ \left. - 2 \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \right. \\ \left. - \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \right\}$$

$$\begin{aligned}
& - \sum_{\alpha, \nu, j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) + \sum_{\nu, j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{3}s \\
& - 2 \sum_{j, m, p} T(e_m, e_p, J(e_m)) g^M((\nabla_{e_j}^L J)e_j, e_p) - 2 \sum_{j, m, \alpha} T(e_m, J(e_j), J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_m)) \\
& - 2 \sum_{j, m, \alpha} T(e_m, J(e_j), J(e_m)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\alpha)) + 2 \sum_{m, p, \alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_m, e_p) \\
& - 2 \sum_{m, p, \alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_m}^L J)e_\alpha, e_p) - \sum_{j, p, m} T(e_j, e_p, J(e_m)) T(e_m, e_p, J(e_j)) \\
& - 2 \sum_{j, l, \alpha} T^2(J(e_j), e_l, e_\alpha) \Big\} d\text{Vol}_M
\end{aligned}$$

where $\Omega_n = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$.

We begin by computing $\int_{\partial M} \Psi$. Since, some operators have the following symbols.

Lemma 3.2. *The following identities hold:*

$$\begin{aligned}
\sigma_1(D_{J,T}) &= \sqrt{-1}c[J(\xi)]; \\
\sigma_0(D_{J,T}) &= -\frac{1}{4} \sum_{i, j, k=1}^n \omega_{j,k}(e_i) c[J(e_i)] c(e_j) c(e_k) + \frac{1}{4} \sum_{u \neq s \neq t=1}^n T_{vst} a_u^v c(e_u) c(e_s) c(e_t) \\
&\quad - \frac{1}{2} \sum_{v, u, t=1}^n T_{vut} a_u^v c(e_t),
\end{aligned}$$

where $T_{vst} := T(e_v, e_s, e_t)$.

Write

$$D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha; \quad \sigma(D_{J,T}) = p_1 + p_0; \quad \sigma(D_{J,T}^{-1}) = \sum_{j=1}^\infty q_{-j}.$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned}
1 &= \sigma(D_{J,T} \circ D_{J,T}^{-1}) = \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha [\sigma(D_{J,T})] D_x^\alpha [\sigma(D_{J,T}^{-1})] \\
&= (p_1 + p_0)(q_{-1} + q_{-2} + q_{-3} + \cdots) \\
&\quad + \sum_j (\partial_{\xi_j} p_1 + \partial_{\xi_j} p_0) \left[(D_{J,T})_{x_j} q_{-1} + (D_{J,T})_{x_j} q_{-2} + (D_{J,T})_{x_j} q_{-3} + \cdots \right] \\
&= p_1 q_{-1} + \left[p_1 q_{-2} + p_0 q_{-1} + \sum_j \partial_{\xi_j} p_1 (D_{J,T})_{x_j} q_{-1} \right] + \cdots,
\end{aligned}$$

so

$$q_{-1} = p_1^{-1}; \quad q_{-2} = -p_1^{-1} [p_0 p_1^{-1} + \sum_j \partial_{\xi_j} p_1 (D_{J,T})_{x_j} (p_1^{-1})].$$

Lemma 3.3. *The following identities hold:*

$$\begin{aligned}\sigma_{-1}(D_{J,T}^{-1}) &= \frac{ic[J(\xi)]}{|\xi|^2}; \\ \sigma_{-2}(D_{J,T}^{-1}) &= \frac{c[J(\xi)]\sigma_0(D_{J,T})c[J(\xi)]}{|\xi|^4} + \frac{c[J(\xi)]}{|\xi|^6} \sum_{j=1}^n c[J(dx_j)] \left[\partial_{x_j}(c[J(\xi)])|\xi|^2 - c[J(\xi)]\partial_{x_j}(|\xi|^2) \right].\end{aligned}$$

When $n = 4$, then $\text{tr}[\text{id}] = 4$, since the sum is taken over $r + l - k - j - |\alpha| - 1 = -4$, $r \leq -1, l \leq -1$, then we have the following five cases:

case a) I) $r = -1, l = -1, k = j = 0, |\alpha| = 1$

By applying the formula shown in (3.4), we can calculate

$$\Psi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_x^{\alpha} \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'.$$

By Lemma 3.3, for $i < n$, $J(dx_p) = \sum_{h=1}^n a_h^p dx_h$ then

An easy computation shows that

$$\begin{aligned}\partial_{\xi_n} \partial_{x_i} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} &= - \frac{2i\xi_n}{(1 + \xi_n^2)^2} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_i}(a_h^p) c(dx_h) \\ &\quad + \frac{i(1 - \xi_n^2)}{(1 + \xi_n^2)^2} \sum_{h=1}^n \partial_{x_i}(a_h^n) c(dx_h).\end{aligned}$$

Likewise,

$$\begin{aligned}\pi_{\xi_n}^+ \partial_{\xi_i} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} &= \frac{1}{2(\xi_n - i)} c[J(dx_i)] - \frac{1}{2(\xi_n - i)^2} \xi_i c[J(dx_n)] \\ &\quad + \frac{2i - \xi_n}{2(\xi_n - i)^2} \sum_{q=1}^{n-1} \xi_i \xi_q c[J(dx_q)].\end{aligned}$$

On account of the above result,

We note that $i < n$, $\int_{|\xi'|=1} \{\xi_{i_1} \cdots \xi_{i_{2d+1}}\} \sigma(\xi') = 0$, then we have

$$\begin{aligned}(3.6) \quad \Psi_1 &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_x^{\alpha} \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx' \\ &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(1 - \xi_n^2)}{2(\xi_n - i)^3 (\xi_n + i)^2} \sum_{\beta=1}^n \sum_{i=1}^{n-1} a_{\beta}^i \partial_{x_i}(a_{\beta}^n) \text{tr}[\text{id}] d\xi_n \sigma(\xi') dx' \\ &\quad + \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i(2i - \xi_n)(1 - \xi_n^2)}{2(\xi_n - i)^4 (\xi_n + i)^2} \sum_{\beta=1}^n \sum_{i,q=1}^{n-1} \xi_i \xi_q a_{\beta}^q \partial_{x_i}(a_{\beta}^n) \text{tr}[\text{id}] d\xi_n \sigma(\xi') dx'\end{aligned}$$

$$\begin{aligned}
& + \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{i\xi_n}{(\xi_n - i)^4(\xi_n + i)^2} \sum_{\beta=1}^n \sum_{i,p=1}^{n-1} \xi_i \xi_p a_\beta^n \partial_{x_i}(a_\beta^p) \text{tr}[\text{id}] d\xi_n \sigma(\xi') dx' \\
& = -\frac{\pi}{6} \sum_{\beta=1}^n \sum_{i=1}^{n-1} a_\beta^i \partial_{x_i}(a_\beta^n) \text{tr}[\text{id}] \Omega_3 dx' - \frac{\pi}{6} \sum_{\beta=1}^n \sum_{i=1}^{n-1} a_\beta^n \partial_{x_i}(a_\beta^i) \text{tr}[\text{id}] dx'.
\end{aligned}$$

From [5, 22], we have $\int_{|\xi'|=1} \xi_i \xi_j \sigma(\xi') = \frac{\Omega_3}{3} \delta_i^j$, and $\sum_{\beta=1}^n \sum_{i=1}^{n-1} a_\beta^i \partial_{x_i}(a_\beta^n) = \sum_{\beta,i=1}^n a_\beta^i \partial_{x_i}(a_\beta^n)$, then

$$\Psi_1 = 0.$$

case a) II) $r = -1$, $l = -1$, $k = |\alpha| = 0$, $j = 1$

Using (3.4), we get

$$\Psi_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n}^2 \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'.$$

It is easy to check that

$$\begin{aligned}
\pi_{\xi_n}^+ \partial_{x_n} \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} & = \frac{1}{2(\xi_n - i)} \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p \partial_{x_n}(a_h^p) c(dx_h) + \frac{i}{2(\xi_n - i)} \sum_{h=1}^n \partial_{x_n}(a_h^n) c(dx_h) \\
& + \frac{1}{2(\xi_n - i)} \sum_{p,h=1}^{n-1} \xi_p a_h^p \partial_{x_n}(c(dx_h)) + \frac{i}{2(\xi_n - i)} \sum_{h=1}^{n-1} a_h^n \partial_{x_n}(c(dx_h)) \\
& + \frac{2i - \xi_n}{4(\xi_n - i)^2} h'(0) \sum_{h=1}^n \sum_{p=1}^{n-1} \xi_p a_h^p c(dx_h) - \frac{1}{4(\xi_n - i)^2} h'(0) \sum_{h=1}^n a_h^n c(dx_h),
\end{aligned}$$

where $\sum_{h=1}^{n-1} \partial_{x_n}(c(dx_h)) = \sum_{h=1}^{n-1} \frac{1}{2} h'(0) c(dx_h)$.

By calculation, we have

$$\partial_{\xi_n}^2 \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} = \frac{-2i + 6i\xi_n^2}{(\xi_n - i)^3(\xi_n + i)^3} \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_\beta^i c(dx_\beta) + \frac{-6i\xi_n + 2i\xi_n^3}{(\xi_n - i)^3(\xi_n + i)^3} \sum_{\beta=1}^n a_\beta^n c(dx_\beta).$$

Consequently,

$$\begin{aligned}
(3.7) \quad \Psi_2 & = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n}^2 \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx' \\
& = \frac{\pi}{48} \sum_{\beta=1}^n \sum_{i=1}^{n-1} a_\beta^i \partial_{x_n}(a_\beta^i) \text{tr}[\text{id}] \Omega_3 dx' + \frac{\pi}{16} \sum_{\beta=1}^n a_\beta^n \partial_{x_n}(a_\beta^n) \text{tr}[\text{id}] \Omega_3 dx' \\
& + \frac{\pi}{96} \sum_{i,\beta=1}^{n-1} (a_\beta^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{\pi}{32} \sum_{\beta=1}^{n-1} (a_\beta^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx'
\end{aligned}$$

$$-\frac{5\pi}{192} \sum_{\beta=1}^n \sum_{i=1}^{n-1} (a_{\beta}^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{3\pi}{64} \sum_{\beta=1}^n (a_{\beta}^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx'.$$

case a) III) $r = -1$, $l = -1$, $j = |\alpha| = 0$, $k = 1$

By (3.4), we calculate that

$$\Psi_3 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'.$$

Hence, we have

$$(3.8) \quad \begin{aligned} \Psi_3 = & -\frac{\pi}{48} \sum_{\beta=1}^n \sum_{i=1}^{n-1} a_{\beta}^i \partial_{x_n} (a_{\beta}^i) \text{tr}[\text{id}] \Omega_3 dx' - \frac{\pi}{96} \sum_{i,\beta=1}^{n-1} (a_{\beta}^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ & + \frac{5\pi}{192} \sum_{\beta=1}^n \sum_{i=1}^{n-1} (a_{\beta}^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{\pi}{16} \sum_{\beta=1}^n a_{\beta}^n \partial_{x_n} (a_{\beta}^n) \text{tr}[\text{id}] \Omega_3 dx' \\ & - \frac{\pi}{32} \sum_{\beta=1}^{n-1} (a_{\beta}^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{3\pi}{64} \sum_{\beta=1}^n (a_{\beta}^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx'. \end{aligned}$$

In combination with the calculation,

$$\Psi_1 + \Psi_2 + \Psi_3 = 0.$$

case b) $r = -2$, $l = -1$, $k = j = |\alpha| = 0$

Similarly, we get

$$\Psi_4 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-2}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'.$$

We first compute

$$\begin{aligned} \sigma_{-2}(D_{J,T}^{-1})(x_0) = & \frac{c[J(\xi)] \sigma_0(D_{J,T})(x_0) c[J(\xi)]}{|\xi|^4} - \frac{c[J(\xi)]}{|\xi|^6} h'(0) |\xi'|^2 c[J(dx_n)] c[J(\xi)] \\ & + \frac{c[J(\xi)]}{|\xi|^4} \left[\sum_{j,p,h=1}^n \xi_p \partial_{x_j} (a_h^p) c[J(dx_j)] c(dx_h) + \sum_{p=1}^n \sum_{h=1}^{n-1} \xi_p a_h^p c[J(dx_n)] \partial_{x_n} (c(dx_h)) \right], \end{aligned}$$

where

$$\begin{aligned} \sigma_0(D_{J,T})(x_0) = & -\frac{1}{4} h'(0) \sum_{\mu=1}^n \sum_{\nu=1}^{n-1} a_{\mu}^{\nu} c(dx_{\mu}) c(dx_n) c(dx_{\nu}) + \frac{1}{4} \sum_{u \neq s \neq t, v=1}^n T_{vst} a_u^v c(e_u) c(e_s) c(e_t) \\ & - \frac{1}{2} \sum_{v,u,t=1}^n T_{vut} a_u^v c(e_t). \end{aligned}$$

For the sake of convenience in writing, we denote

$$(3.9) \quad \sigma_{-2}(D_{J,T}^{-1})(x_0)|_{|\xi'|=1} := B_0(x_0) + B_1(x_0) + B_2(x_0),$$

$$\begin{aligned} B_0(x_0) &= -\frac{1}{4}h'(0) \sum_{\mu=1}^n \sum_{\nu=1}^{n-1} \frac{c[J(\xi)]a_\nu^\mu c(dx_\mu)c(dx_n)c(dx_\nu)c[J(\xi)]}{(1+\xi_n^2)^2} - \frac{c[J(\xi)]}{(1+\xi_n^2)^3}h'(0)c[J(dx_n)]c[J(\xi)] \\ &\quad + \frac{c[J(\xi)]}{(1+\xi_n^2)^2} \left[\sum_{j,p,h=1}^n \xi_p \partial_{x_j}(a_h^p)c[J(dx_j)]c(dx_h) + \sum_{p=1}^n \sum_{h=1}^{n-1} \xi_p a_h^p c[J(dx_n)]\partial_{x_n}(c(dx_h)) \right]; \\ B_1(x_0) &= \frac{1}{4} \sum_{u \neq s \neq t, v=1}^n \frac{c[J(\xi)]T_{vst}a_u^v c(e_u)c(e_s)c(e_t)c[J(\xi)]}{(1+\xi_n^2)^2}; \\ B_2(x_0) &= -\frac{1}{2} \sum_{v,u,t=1}^n \frac{c[J(\xi)]T_{vut}a_u^v c(e_t)c[J(\xi)]}{(1+\xi_n^2)^2}, \end{aligned}$$

means that

$$\pi_{\xi_n}^+ \sigma_{-2}(D_{J,T}^{-1})(x_0)|_{|\xi'|=1} = \pi_{\xi_n}^+(B_0(x_0)) + \pi_{\xi_n}^+(B_1(x_0)) + \pi_{\xi_n}^+(B_2(x_0)).$$

By computation, we have

$$\begin{aligned} (3.10) \quad & -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ B_0(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})(x_0)] d\xi_n \sigma(\xi') dx' \\ &= -\frac{\pi}{48} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} ((a_\nu^n)^2 (a_l^i)^2 - (a_l^i)^2 a_\nu^n a_n^n) \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ &\quad - \frac{\pi}{48} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} ((a_\beta^i)^2 a_l^j \partial_{x_j}(a_l^i) - a_l^i a_\beta^j a_\beta^i \partial_{x_j}(a_l^i) + a_l^i a_l^j a_\beta^i \partial_{x_j}(a_\beta^n)) \text{tr}[\text{id}] \Omega_3 dx' \\ &\quad - \frac{\pi}{48} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} (a_\beta^n a_l^j a_\beta^i \partial_{x_j}(a_l^i) - a_l^i a_\beta^j a_\beta^i \partial_{x_j}(a_l^i) + a_l^i a_l^j a_\beta^i \partial_{x_j}(a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\ &\quad + \frac{\pi}{24} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} (a_\beta^i a_l^j a_\beta^n \partial_{x_j}(a_l^i) - a_l^i a_\beta^j a_\beta^n \partial_{x_j}(a_l^i) + a_l^i a_l^j a_\beta^n \partial_{x_j}(a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\ &\quad - \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ &\quad - \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{5\pi}{192} \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 (a_\beta^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \end{aligned}$$

$$-\frac{\pi}{128} \sum_{\beta,l=1}^n (a_\beta^n)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{5\pi}{128} \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx'.$$

By computation, we have

(3.11)

$$\begin{aligned} \pi_{\xi_n}^+(B_1(x_0)) &= \pi_{\xi_n}^+ \sum_{u \neq s \neq t, v=1}^n \frac{c[J(\xi)] T_{vst} a_u^v c(e_u) c(e_s) c(e_t) c[J(\xi)]}{4(1 + \xi_n^2)^2} \\ &= \pi_{\xi_n}^+ \left(\frac{\xi_n^2}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\ &\quad + \pi_{\xi_n}^+ \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} a_l^n a_r^\alpha a_u^v \xi_\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\ &\quad + \pi_{\xi_n}^+ \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} a_l^q a_r^n a_u^v \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\ &\quad + \pi_{\xi_n}^+ \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} T_{vst} a_l^q a_r^\alpha a_u^v \xi_\alpha \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\ &= -\frac{i\xi_n}{16(\xi_n - i)^2} \sum_{l,r,v,u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\ &\quad - \frac{i}{16(\xi_n - i)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} a_l^n a_r^\alpha a_u^v \xi_\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\ &\quad - \frac{i}{16(\xi_n - i)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} a_l^q a_r^n a_u^v \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\ &\quad - \frac{i\xi_n + 2}{16(\xi_n - i)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} T_{vst} a_l^q a_r^\alpha a_u^v \xi_\alpha \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r). \end{aligned}$$

Thus, we have

(3.12)

$$\begin{aligned} &\text{tr}[\pi_{\xi_n}^+ B_1(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) |_{|\xi'|=1} \\ &= \frac{-i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i=1}^{n-1} T_{vst} \xi_i a_l^n a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\ &\quad + \frac{-\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,\alpha=1}^{n-1} T_{vst} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \end{aligned}$$

$$\begin{aligned}
& + \frac{-\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{i, q=1}^{n-1} T_{vst} \xi_i \xi_q a_l^q a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{i\xi_n(i\xi_n + 2)}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{i, q, \alpha=1}^{n-1} T_{vst} \xi_i \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{\xi_n - \xi_n^3}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{1 - \xi_n^2}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{1 - \xi_n^2}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} \xi_q a_l^q a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i(1 - \xi_n^2)(i\xi_n + 2)}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{q, \alpha=1}^{n-1} T_{vst} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)].
\end{aligned}$$

By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, we have the equality:

$$\begin{aligned}
(3.13) \quad & \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& = \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \left[-\delta_u^l \delta_s^\beta \delta_t^r + \delta_u^l \delta_s^r \delta_t^\beta + \delta_u^\beta \delta_s^l \delta_t^r - \delta_u^\beta \delta_s^r \delta_t^l - \delta_u^r \delta_s^l \delta_t^\beta + \delta_u^r \delta_s^\beta \delta_t^l \right] \text{tr}[id].
\end{aligned}$$

By $i < n$, $\int_{|\xi'|=1} \{\xi_{i_1} \cdots \xi_{i_{2d+1}}\} \sigma(\xi') = 0$, $\int_{|\xi'|=1} \xi_i \xi_j \sigma(\xi') = \frac{4\pi}{3} \delta_i^j$, then we have

$$(3.14) \quad -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ B_1(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx' = 0.$$

By means of similar calculations, we have

$$\begin{aligned}
(3.15) \quad & \text{tr}[\pi_{\xi_n}^+ B_2(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0)|_{|\xi'|=1} \\
& = \frac{i\xi_n^2}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i=1}^{n-1} T_{vut} \xi_i a_l^n a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{\xi_n}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)]
\end{aligned}$$

$$\begin{aligned}
& + \frac{\xi_n}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i\xi_n(i\xi_n + 2)}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, q, \alpha=1}^{n-1} T_{vut} \xi_i \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{\xi_n - \xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n T_{vut} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{1 - \xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{\alpha=1}^{n-1} T_{vut} \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{1 - \xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{q=1}^{n-1} T_{vut} \xi_q a_l^q a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{i(1 - \xi_n^2)(i\xi_n + 2)}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)].
\end{aligned}$$

By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, we have the equality:

$$(3.16) \quad \sum_{l, t, r, \beta=1}^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] = \sum_{l, t, r, \beta=1}^n [\delta_l^\beta \delta_t^r - \delta_l^r \delta_t^\beta + \delta_l^t \delta_r^\beta] \text{tr}[id].$$

By $i < n$, $\int_{|\xi'|=1} \{\xi_{i_1} \cdots \xi_{i_{2d+1}}\} \sigma(\xi') = 0$, $\int_{|\xi'|=1} \xi_i \xi_j \sigma(\xi') = \frac{\Omega_3}{3} \delta_i^j$, then we have

$$\begin{aligned}
(3.17) \quad & - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ B_2(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx' \\
& = - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \frac{\xi_n}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{u, v, t, r=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_t^n a_r^\alpha a_u^v a_\beta^i \right. \\
& + \frac{\xi_n}{16(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{u, v, t, l=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_t^n a_u^v a_\beta^i - \frac{\xi_n - \xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{u, v, t, r=1}^n T_{vut} a_t^n a_r^n a_u^v a_\beta^n \\
& \left. - \frac{i(1 - \xi_n^2)(i\xi_n + 2)}{8(\xi_n - i)^2(1 + \xi_n^2)^2} \sum_{u, v, t, l=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_t^\alpha a_u^v a_\beta^n \right\} \text{tr}[id] d\xi_n \sigma(\xi') dx' \\
& = \frac{\pi}{96} \sum_{u, v, t, r=1}^n \sum_{i=1}^{n-1} T_{vut} a_t^n (a_r^i)^2 a_u^v \Omega_3 \text{tr}[id] dx' + \frac{\pi}{96} \sum_{u, v, t, l=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_t^n a_u^v \Omega_3 \text{tr}[id] dx'
\end{aligned}$$

$$+ \frac{\pi}{48} \sum_{u,v,t,l=1}^n \sum_{q=1}^{n-1} T_{vut}(a_l^q)^2 a_u^v a_t^n \Omega_3 \text{tr}[id] dx'.$$

Hence, we have

$$\begin{aligned} \Psi_4 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-2}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{\pi}{48} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} ((a_\nu^n)^2 (a_l^i)^2 - (a_l^i)^2 a_\nu^n) \text{tr}[id] \Omega_3 h'(0) dx' \\ &\quad - \frac{\pi}{48} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} ((a_\beta^i)^2 a_l^j \partial_{x_j}(a_l^n) - a_l^i a_\beta^j a_\beta^i \partial_{x_j}(a_l^n) + a_l^i a_\beta^j a_\beta^i \partial_{x_j}(a_\beta^n)) \text{tr}[id] \Omega_3 dx' \\ &\quad - \frac{\pi}{48} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} (a_\beta^n a_l^j a_\beta^i \partial_{x_j}(a_l^i) - a_l^n a_\beta^j a_\beta^i \partial_{x_j}(a_l^i) + a_l^n a_\beta^j a_\beta^i \partial_{x_j}(a_\beta^i)) \text{tr}[id] \Omega_3 dx' \\ &\quad + \frac{\pi}{24} \sum_{l,j,\beta=1}^n \sum_{i=1}^{n-1} (a_\beta^i a_l^j a_\beta^n \partial_{x_j}(a_l^i) - a_l^i a_\beta^j a_\beta^n \partial_{x_j}(a_l^i) + a_l^i a_\beta^j a_\beta^n \partial_{x_j}(a_\beta^i)) \text{tr}[id] \Omega_3 dx' \\ &\quad - \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \text{tr}[id] \Omega_3 h'(0) dx' - \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[id] \Omega_3 h'(0) dx' \\ &\quad - \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[id] \Omega_3 h'(0) dx' + \frac{5\pi}{192} \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 (a_\beta^n)^2 \text{tr}[id] \Omega_3 h'(0) dx' \\ &\quad - \frac{\pi}{128} \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_\beta^n)^2 (a_l^n)^2 \text{tr}[id] \Omega_3 h'(0) dx' + \frac{5\pi}{128} \sum_{\beta,l=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 (a_l^n)^2 \text{tr}[id] \Omega_3 h'(0) dx' \\ &\quad + \frac{\pi}{96} \sum_{u,v,t,r=1}^n \sum_{i=1}^{n-1} T_{vut} a_t^n (a_r^i)^2 a_u^v \Omega_3 \text{tr}[id] dx' + \frac{\pi}{96} \sum_{u,v,t,l=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_t^n a_u^v \Omega_3 \text{tr}[id] dx' \\ &\quad + \frac{\pi}{48} \sum_{u,v,t,l=1}^n \sum_{q=1}^{n-1} T_{vut}(a_l^q)^2 a_u^v a_t^n \Omega_3 \text{tr}[id] dx'. \end{aligned}$$

case c) $r = -1$, $l = -2$, $k = j = |\alpha| = 0$

We calculate

$$\Psi_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-2}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'.$$

It is evident that

$$(3.18) \quad \pi_{\xi_n}^+ \left(\frac{ic[J(\xi)]}{|\xi|^2} \right) (x_0)|_{|\xi'|=1} = \frac{1}{2(\xi_n - i)} \sum_{\beta=1}^n \sum_{i=1}^{n-1} \xi_i a_\beta^i c(dx_\beta) + \frac{i}{2(\xi_n - i)} \sum_{\beta=1}^n a_\beta^n c(dx_\beta).$$

Same classification as in case b (3.9), we have

$$(3.19) \quad \begin{aligned} & -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} B_0](x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} ((a_\nu^n)^2 (a_l^i)^2 - (a_l^i)^2 a_\nu^n a_n^n) \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ &+ \frac{\pi}{48} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} ((a_\beta^i)^2 a_l^j \partial_{x_j} (a_l^n) - a_l^i a_\beta^j a_\beta^i \partial_{x_j} (a_l^n) + a_l^i a_l^j a_\beta^i \partial_{x_j} (a_\beta^n)) \text{tr}[\text{id}] \Omega_3 dx' \\ &+ \frac{\pi}{48} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} (a_\beta^n a_l^j a_\beta^i \partial_{x_j} (a_l^i) - a_l^n a_\beta^j a_\beta^i \partial_{x_j} (a_l^i) + a_l^n a_l^j a_\beta^i \partial_{x_j} (a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\ &- \frac{\pi}{24} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} (a_\beta^i a_l^j a_\beta^n \partial_{x_j} (a_l^i) - a_l^i a_\beta^j a_\beta^n \partial_{x_j} (a_l^i) + a_l^i a_l^j a_\beta^n \partial_{x_j} (a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\ &+ \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_l^i)^2 (a_\nu^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ &+ \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{5\pi}{192} \sum_{\beta, l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 (a_\beta^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\ &+ \frac{\pi}{128} \sum_{\beta, l=1}^n (a_\beta^n)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{5\pi}{128} \sum_{\beta, l=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx'. \end{aligned}$$

By computation, we have

$$(3.20) \quad \begin{aligned} \partial_{\xi_n} (B_1(x_0)) &= \partial_{\xi_n} \sum_{u \neq s \neq t, v=1}^n \frac{c[J(\xi)] T_{vst} a_u^v c(e_u) c(e_s) c(e_t) c[J(\xi)]}{4(1 + \xi_n^2)^2} \\ &= \partial_{\xi_n} \left(\frac{\xi_n^2}{4(1 + \xi_n^2)^2} \sum_{l, r, v, u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\ &+ \partial_{\xi_n} \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l, r, v, u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} a_l^n a_r^\alpha a_u^v \xi_\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \end{aligned}$$

$$\begin{aligned}
& + \partial_{\xi_n} \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} a_l^q a_r^n a_u^v \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\
& + \partial_{\xi_n} \left(\frac{\xi_n}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} T_{vst} a_l^q a_r^\alpha a_u^v \xi_\alpha \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right) \\
& = \frac{\xi_n}{2(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\
& + \frac{1 - \xi_n^2}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} a_l^n a_r^\alpha a_u^v \xi_\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\
& + \frac{1 - \xi_n^2}{4(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} a_l^q a_r^n a_u^v \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \\
& - \frac{\xi_n}{2(1 + \xi_n^2)^2} \sum_{l,r,v,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} T_{vst} a_l^q a_r^\alpha a_u^v \xi_\alpha \xi_q c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r).
\end{aligned}$$

Thus, we have

(3.21)

$$\begin{aligned}
& \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} B_1](x_0)|_{|\xi|^'=1} \\
& = \frac{\xi_n}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i=1}^{n-1} T_{vst} \xi_i a_l^n a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\
& + \frac{1 - \xi_n^2}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,\alpha=1}^{n-1} T_{vst} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\
& + \frac{1 - \xi_n^2}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,q=1}^{n-1} T_{vst} \xi_i \xi_q a_l^q a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\
& - \frac{\xi_n}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,q,\alpha=1}^{n-1} T_{vst} \xi_i \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\
& + \frac{i\xi_n}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)] \\
& + \frac{i - i\xi_n^2}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} T_{vst} \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) c(dx_\beta)]
\end{aligned}$$

$$\begin{aligned}
& + \frac{i - i\xi_n^2}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{q=1}^{n-1} T_{vst} \xi_q a_l^q a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i\xi_n}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{q, \alpha=1}^{n-1} T_{vst} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)].
\end{aligned}$$

According to (4.3), we have

$$(3.22) \quad -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} B_1](x_0) d\xi_n \sigma(\xi') dx' = 0$$

By means of similar calculations, we have

$$\begin{aligned}
(3.23) \quad \partial_{\xi_n}(B_2(x_0)) & = \partial_{\xi_n} \sum_{u=s, v, t=1}^n \frac{c[J(\xi)] T_{vut} a_u^v c(e_t) c[J(\xi)]}{2(1 + \xi_n^2)^2} \\
& = -\frac{\xi_n}{(1 + \xi_n^2)^2} \sum_{l, r, v, u, t=1}^n T_{vut} a_l^n a_r^n a_u^v c(dx_l) c(dx_t) c(dx_r) \\
& \quad - \frac{1 - \xi_n^2}{2(1 + \xi_n^2)^2} \sum_{l, r, v, u, t=1}^n \sum_{\alpha=1}^{n-1} T_{vut} a_l^n a_r^\alpha a_u^v \xi_\alpha c(dx_l) c(dx_t) c(dx_r) \\
& \quad - \frac{1 - \xi_n^2}{2(1 + \xi_n^2)^2} \sum_{l, r, v, u, t=1}^n \sum_{q=1}^{n-1} T_{vut} a_l^q a_r^n a_u^v \xi_q c(dx_l) c(dx_t) c(dx_r) \\
& \quad + \frac{\xi_n}{(1 + \xi_n^2)^2} \sum_{l, r, v, u, t=1}^n \sum_{\alpha, q=1}^{n-1} T_{vut} a_l^q a_r^\alpha a_u^v \xi_\alpha \xi_q c(dx_l) c(dx_t) c(dx_r).
\end{aligned}$$

Then

(3.24)

$$\begin{aligned}
& \text{tr}[\pi_{\xi_n}^+ B_2(x_0) \times \partial_{\xi_n} \sigma_{-1}(D_{J,T}^{-1})](x_0)|_{|\xi'|=1} \\
& = -\frac{\xi_n}{2(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i=1}^{n-1} T_{vut} \xi_i a_l^n a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& \quad - \frac{1 - \xi_n^2}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& \quad - \frac{1 - \xi_n^2}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& \quad + \frac{\xi_n}{2(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{i, q, \alpha=1}^{n-1} T_{vut} \xi_i \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)]
\end{aligned}$$

$$\begin{aligned}
& - \frac{i\xi_n}{2(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n T_{vut} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i - i\xi_n}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{\alpha=1}^{n-1} T_{vut} \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i - i\xi_n}{4(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{q=1}^{n-1} T_{vut} \xi_q a_l^q a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{\xi_n}{2(\xi_n - i)(1 + \xi_n^2)^2} \sum_{\beta, u, v, t, l, r=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)].
\end{aligned}$$

According to (3.16), we have

(3.25)

$$\begin{aligned}
& - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} B_2](x_0) d\xi_n \sigma(\xi') dx' \\
& = - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ - \frac{1 - \xi_n^2}{4(\xi_n - i)^3 (\xi_n + i)^2} \sum_{u, v, t, r=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_t^n a_r^\alpha a_u^v a_i^i \right. \\
& - \frac{1 - \xi_n^2}{4(\xi_n - i)^3 (\xi_n + i)^2} \sum_{u, v, l, r=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_t^n a_u^v a_i^i \\
& - \frac{i\xi_n}{2(\xi_n - i)^3 (\xi_n + i)^2} \sum_{u, v, t, r=1}^n T_{vut} a_t^n (a_r^n)^2 a_u^v \\
& \left. - \frac{i\xi_n}{2(\xi_n - i)^3 (\xi_n + i)^2} \sum_{u, v, t, l=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_l^\alpha a_u^v a_t^n \right\} \text{tr}[id] d\xi_n \sigma(\xi') dx' \\
& = - \frac{\pi}{48} \sum_{u, v, t, r=1}^n \sum_{i=1}^{n-1} T_{vut} a_t^n a_u^v (a_r^i)^2 \text{tr}[id] \Omega_3 dx' - \frac{\pi}{48} \sum_{u, v, l, r=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_t^n a_u^v \text{tr}[id] \Omega_3 dx' \\
& - \frac{\pi}{16} \sum_{u, v, t, r=1}^n T_{vut} a_t^n (a_r^n)^2 a_u^v \text{tr}[id] \Omega_3 dx' - \frac{\pi}{48} \sum_{u, v, t, l=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_u^v a_t^n \text{tr}[id] \Omega_3 dx'.
\end{aligned}$$

Consequently

(3.26)

$$\Psi_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-2}(D_{J,T}^{-1})](x_0) d\xi_n \sigma(\xi') dx'$$

$$\begin{aligned}
&= \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} ((a_\nu^n)^2 (a_l^i)^2 - (a_l^i)^2 a_\nu^n) \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\
&+ \frac{\pi}{48} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} ((a_\beta^i)^2 a_l^j \partial_{x_j} (a_l^n) - a_l^i a_\beta^j a_\beta^i \partial_{x_j} (a_l^n) + a_l^i a_l^j a_\beta^i \partial_{x_j} (a_\beta^n)) \text{tr}[\text{id}] \Omega_3 dx' \\
&+ \frac{\pi}{48} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} (a_\beta^n a_l^j a_\beta^i \partial_{x_j} (a_l^i) - a_l^n a_\beta^j a_\beta^i \partial_{x_j} (a_l^i) + a_l^n a_l^j a_\beta^i \partial_{x_j} (a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\
&- \frac{\pi}{24} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} (a_\beta^i a_l^j a_\beta^n \partial_{x_j} (a_l^i) - a_l^i a_\beta^j a_\beta^n \partial_{x_j} (a_l^i) + a_l^i a_l^j a_\beta^n \partial_{x_j} (a_\beta^i)) \text{tr}[\text{id}] \Omega_3 dx' \\
&+ \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' + \frac{\pi}{96} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\
&+ \frac{\pi}{48} \sum_{l=1}^n \sum_{\nu, i=1}^{n-1} (a_\nu^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{5\pi}{192} \sum_{\beta, l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 (a_\beta^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\
&+ \frac{\pi}{128} \sum_{\beta, l=1}^n (a_\beta^n)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' - \frac{5\pi}{128} \sum_{\beta, l=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 (a_l^n)^2 \text{tr}[\text{id}] \Omega_3 h'(0) dx' \\
&- \frac{\pi}{48} \sum_{u, v, t, r=1}^n \sum_{i=1}^{n-1} T_{vut} a_t^n a_u^v (a_r^i)^2 \text{tr}[\text{id}] \Omega_3 dx' - \frac{\pi}{48} \sum_{u, v, l, r=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_t^n a_u^v \text{tr}[\text{id}] \Omega_3 dx' \\
&- \frac{\pi}{16} \sum_{u, v, t, r=1}^n T_{vut} a_t^n (a_r^n)^2 a_u^v \text{tr}[\text{id}] \Omega_3 dx' - \frac{\pi}{48} \sum_{u, v, t, l=1}^n \sum_{i=1}^{n-1} T_{vut} (a_l^i)^2 a_u^v a_t^n \text{tr}[\text{id}] \Omega_3 dx'.
\end{aligned}$$

In summary,

$$\Psi = \Psi_1 + \Psi_2 + \Psi_3 + \Psi_4 + \Psi_5 = 0.$$

Thus, the following theorem is obtained.

Theorem 3.4. *Let M be a 4-dimensional almost product Riemannian spin manifold with the boundary ∂M and the metric g^M as above, $D_{J,T}$ be the J -twist of the Dirac operator on \widetilde{M} , then*

(3.27)

$$\begin{aligned}
&\widetilde{\text{Wres}}[\pi^+ D_{J,T}^{-1} \circ \pi^+ D_{J,T}^{-1}] \\
&= \int_M 2\pi^2 \left\{ \sum_{i, j=1}^n R(J(e_i), J(e_j), e_j, e_i) - 2 \sum_{\nu, j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right\}
\end{aligned}$$

$$\begin{aligned}
& - 2 \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \\
& - \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j)g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \\
& - \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j)g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) + \sum_{\nu,j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{3}s \\
& - 2 \sum_{j,m,p} T(e_m, e_p, J(e_m))g^M((\nabla_{e_j}^L J)e_j, e_p) - 2 \sum_{j,m,\alpha} T(e_m, J(e_j), J(e_\alpha))g^M((\nabla_{e_\alpha}^L J)e_j, J(e_m)) \\
& - 2 \sum_{j,m,\alpha} T(e_m, J(e_j), J(e_m))g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\alpha)) + 2 \sum_{m,p,\alpha} T(e_m, e_p, J(e_\alpha))g^M((\nabla_{e_\alpha}^L J)e_m, e_p) \\
& - 2 \sum_{m,p,\alpha} T(e_m, e_p, J(e_\alpha))g^M((\nabla_{e_m}^L J)e_\alpha, e_p) - \sum_{j,p,m} T(e_j, e_p, J(e_m))T(e_m, e_p, J(e_j)) \\
& - 2 \sum_{j,l,\alpha} T^2(J(e_j), e_l, e_\alpha) \Big\} d\text{Vol}_M
\end{aligned}$$

where s is the scalar curvature.

4. A KASTLER-KALAU-WALZE TYPE THEOREM FOR 6-DIMENSIONAL MANIFOLDS WITH BOUNDARY

Lemma 4.1. *The following identities hold:*

$$\begin{aligned}
\sigma_3(D_{J,T}^3) &= \sqrt{-1}c[J(\xi)]|\xi|^2; \\
\sigma_2(D_{J,T}^3) &= \sum_{i,j,l=1}^n c[J(dx_l)]\partial_l(g^{ij})\xi_i\xi_j + c[J(\xi)](2\sigma^k - \Gamma^k)\xi_k - \sum_{\alpha=1}^n c[J(\xi)]c[J(e_\alpha)]c[(\nabla_{e_\alpha}^L J)(\xi^*)] \\
& - \frac{1}{4}|\xi|^2 \sum_{s,t,l=1}^n \omega_{s,t}(e_l)c[J(e_l)]c(e_s)c(e_t) - \sum_{j,l=1}^n c[J(e_l)]c[J(e_j)]T_J\xi_l\xi_j \\
& - \sum_{j,l=1}^n c[J(e_l)]T_Jc[J(e_j)]\xi_l\xi_j + T_J|\xi|^2,
\end{aligned}$$

where $\xi^* = \sum_{\beta=1}^n \langle e_\beta, \xi \rangle e_\beta$.

Write

$$D_x^\alpha = (-i)^{|\alpha|}\partial_x^\alpha; \quad \sigma(D_{J,T}^3) = p_3 + p_2 + p_1 + p_0; \quad \sigma(D_{J,T}^{-3}) = \sum_{j=3}^\infty q_{-j}.$$

By the composition formula of pseudodifferential operators, we have

$$1 = \sigma(D_{J,T} \circ D_{J,T}^{-1}) = \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha [\sigma(D_{J,T})] D_x^\alpha [\sigma(D_{J,T}^{-1})]$$

$$\begin{aligned}
&= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \cdots) \\
&\quad + \sum_j (\partial_{\xi_j} p_3 + \partial_{\xi_j} p_2 + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0)(D_{x_j} q_{-3} + D_{x_j} q_{-4} + D_{x_j} q_{-5} + \cdots) \\
&= p_3 q_{-3} + (p_3 q_{-4} + p_2 q_{-3} + \sum_j \partial_{\xi_j} p_3 D_{x_j} q_{-3}) + \cdots,
\end{aligned}$$

so

$$q_{-3} = p_3^{-1}; \quad q_{-4} = -p_3^{-1}[p_2 p_3^{-1} + \sum_j \partial_{\xi_j} p_3 D_{x_j} (p_3^{-1})].$$

Lemma 4.2. *The following identities hold:*

$$\begin{aligned}
\sigma_{-3}(D_{J,T}^{-3}) &= \frac{ic[J(\xi)]}{|\xi|^4}; \\
\sigma_{-4}(D_{J,T}^{-3}) &= \frac{c[J(\xi)]\sigma_2(D_{J,T}^3)c[J(\xi)]}{|\xi|^8} \\
&\quad + \frac{c[J(\xi)]}{|\xi|^{10}} \sum_{j=1}^n \left[c[J(dx_j)]|\xi|^2 + 2\xi_j c[J(\xi)] \right] \left[\partial_{x_j}(c[J(\xi)])|\xi|^2 - c[J(\xi)]\partial_{x_j}(|\xi|^2) \right].
\end{aligned}$$

When $n = 6$, then $\text{tr}[\text{id}] = 8$, since the sum is taken over $r + l - k - j - |\alpha| - 1 = -6$, $r \leq -1, l \leq -3$, then we have the following five cases:

case $\hat{\mathbf{a}}$ I) $r = -1, l = -3, k = j = 0, |\alpha| = 1$

By applying the formula shown in (3.2), we can calculate

$$\hat{\Psi}_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx'.$$

By Lemma 3.3 and Lemma 4.2, for $i < n$, $J(dx_p) = \sum_{h=1}^n a_h^p dx_h$. Then by a simple calculation we get

$$\hat{\Psi}_1 = 0.$$

case $\hat{\mathbf{a}}$ II) $r = -1, l = -3, k = |\alpha| = 0, j = 1$

Using (3.2), we get

$$\hat{\Psi}_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx'.$$

A simple calculation yields

$$\hat{\Psi}_2 = \frac{\pi}{64} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_n} (a_h^i) \text{tr}[\text{id}] \Omega_4 dx' + \frac{3\pi}{64} \sum_{h=1}^n a_h^n \partial_{x_n} (a_h^n) \text{tr}[\text{id}] \Omega_4 dx'$$

$$\begin{aligned}
& + \frac{\pi}{128} \sum_{h,i=1}^{n-1} (a_h^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' + \frac{3\pi}{128} \sum_{h=1}^{n-1} (a_h^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& - \frac{7\pi}{320} \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{3\pi}{64} \sum_{h=1}^n (a_h^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.
\end{aligned}$$

case $\hat{\mathbf{a}}$ III) $r = -1$, $l = -3$, $j = |\alpha| = 0$, $k = 1$

By (3.2), we calculate that

$$\hat{\Psi}_3 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx'.$$

Hence, we have

$$\begin{aligned}
\hat{\Psi}_3 & = -\frac{\pi}{64} \sum_{h=1}^n \sum_{i=1}^{n-1} a_h^i \partial_{x_n} (a_h^i) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' - \frac{\pi}{128} \sum_{i,h=1}^{n-1} (a_h^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& + \frac{21\pi}{640} \sum_{h=1}^n \sum_{i=1}^{n-1} (a_h^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{3\pi}{64} \sum_{h=1}^n a_h^n \partial_{x_n} (a_h^n) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' \\
& - \frac{3\pi}{128} \sum_{h=1}^{n-1} (a_h^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' + \frac{9\pi}{128} \sum_{h=1}^n (a_h^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.
\end{aligned}$$

In combination with the calculation,

$$\hat{\Psi}_1 + \hat{\Psi}_2 + \hat{\Psi}_3 = \frac{7\pi}{640} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' + \frac{3\pi}{128} \sum_{l=1}^n (a_l^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.$$

case $\hat{\mathbf{b}}$ $r = -1$, $l = -4$, $k = j = |\alpha| = 0$

By (3.2), we calculate that

$$\begin{aligned}
\hat{\Psi}_4 & = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-4})](x_0) d\xi_n \sigma(\xi') dx' \\
& = i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times \sigma_{-3}(D_{J,T}^{-4})](x_0) d\xi_n \sigma(\xi') dx'.
\end{aligned}$$

From Lemma 4.1 and Lemma 4.2 we get

$$\begin{aligned}
\sigma_{-4}(D_{J,T}^{-3}) & = \frac{c[J(\xi)] \sigma_2(D_{J,T}^3) c[J(\xi)]}{|\xi|^8} \\
& + \frac{c[J(\xi)]}{|\xi|^{10}} \sum_{j=1}^n \left[c[J(dx_j)] |\xi|^2 + 2\xi_j c[J(\xi)] \right] \left[\partial_{x_j} (c[J(\xi)]) |\xi|^2 - c[J(\xi)] \partial_{x_j} (|\xi|^2) \right]
\end{aligned}$$

$$\begin{aligned}
&= \frac{c[J(\xi)]\sigma_2(D_J^3)c[J(\xi)]}{|\xi|^8} \\
&\quad + \frac{c[J(\xi)]}{|\xi|^{10}} \sum_{j=1}^n \left[c[J(dx_j)]|\xi|^2 + 2\xi_j c[J(\xi)] \right] \left[\partial_{x_j}(c[J(\xi)])|\xi|^2 - c[J(\xi)]\partial_{x_j}(|\xi|^2) \right] \\
&\quad + \frac{c[J(\xi)]T_J c[J(\xi)]}{|\xi|^6} - \frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)]c[J(e_l)]c[J(e_j)]T_J c[J(\xi)]\xi_l \xi_j \\
&\quad - \frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)]c[J(e_l)]T_J c[J(e_j)]c[J(\xi)]\xi_l \xi_j \\
&:= Q_0 + Q_1 + Q_2
\end{aligned}$$

$$\begin{aligned}
Q_0 &:= \frac{c[J(\xi)]\sigma_2(D_J^3)c[J(\xi)]}{|\xi|^8} \\
&\quad + \frac{c[J(\xi)]}{|\xi|^{10}} \sum_{j=1}^n \left[c[J(dx_j)]|\xi|^2 + 2\xi_j c[J(\xi)] \right] \left[\partial_{x_j}(c[J(\xi)])|\xi|^2 - c[J(\xi)]\partial_{x_j}(|\xi|^2) \right];
\end{aligned}$$

$$Q_1 := \frac{c[J(\xi)]T_J c[J(\xi)]}{|\xi|^6};$$

$$Q_2 := -\frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)]c[J(e_l)]c[J(e_j)]T_J c[J(\xi)]\xi_l \xi_j - \frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)]c[J(e_l)]T_J c[J(e_j)]c[J(\xi)]\xi_l \xi_j.$$

A simple calculation gives

$$\begin{aligned}
(4.1) \quad & i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times Q_0](x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{\pi}{16} \sum_{i=1}^{n-1} (a_n^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' - \frac{\pi}{16} \sum_{i=1}^{n-1} a_i^i a_n^n \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
&\quad + \frac{11\pi}{128} \sum_{l=1}^n (a_n^l)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' - \frac{\pi}{64} \sum_{i=1}^{n-1} (a_n^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
&\quad - \frac{23\pi}{320} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' + \frac{\pi}{64} \sum_{i,j=1}^{n-1} (a_j^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
&\quad + \frac{\pi}{64} \sum_{l,\alpha=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 g^M(J(dx_\alpha), (\nabla_{e_\alpha}^L J) e_n) \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
&\quad - \frac{\pi}{16} \sum_{i=1}^{n-1} g^M(J(dx_i), (\nabla_{e_n}^L J) e_i) \text{tr}[\text{id}] \Omega_4 h'(0) dx'
\end{aligned}$$

$$\begin{aligned}
& + \frac{\pi}{16} \sum_{i=1}^{n-1} g^M (J(dx_n), (\nabla_{e_i}^L J) e_i) \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& - \frac{\pi}{64} \sum_{l,\alpha=1}^n (a_l^n)^2 g^M (J(dx_\alpha), (\nabla_{e_\alpha}^L J) e_n) \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& + \frac{\pi}{8} \sum_{l=1}^n \sum_{i=1}^{n-1} a_l^i \partial_{x_i} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.
\end{aligned}$$

(4.2)

$$\begin{aligned}
Q_1(x_0)|_{|\xi'|=1} &= \frac{c[J(\xi)] T_J c[J(\xi)]}{|\xi|^6} (x_0)|_{|\xi'|=1} \\
&= \frac{1}{4|\xi|^6} \left[\sum_{\alpha,q,v,l,r,u \neq s \neq t=1}^n \xi_q \xi_\alpha T_{vst} a_u^v a_l^q a_r^\alpha c(dx_l) c(e_u) c(e_s) c(e_t) c(dx_r) \right. \\
&\quad \left. - 2 \sum_{\alpha,q,l,r,u,t,v=1}^n \xi_q \xi_\alpha T_{vut} a_u^v a_l^q a_r^\alpha c(dx_l) c(e_t) c(dx_r) \right] (x_0)|_{|\xi'|=1} \\
&= \frac{\xi_n^2}{4(1+\xi_n^2)^3} \sum_{v,l,r,u \neq s \neq t=1}^n T_{vst} a_u^v a_l^n a_r^n c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) (x_0) \\
&\quad + \frac{\xi_n}{4(1+\xi_n^2)^3} \sum_{v,l,r,u \neq s \neq t=1}^n \sum_{\alpha=1}^{n-1} \xi_\alpha T_{vst} a_u^v a_l^n a_r^\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) (x_0) \\
&\quad + \frac{\xi_n}{4(1+\xi_n^2)^3} \sum_{v,l,r,u \neq s \neq t=1}^n \sum_{q=1}^{n-1} \xi_q T_{vst} a_u^v a_l^q a_r^n c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) (x_0) \\
&\quad + \frac{1}{4(1+\xi_n^2)^3} \sum_{v,l,r,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} \xi_q \xi_\alpha T_{vst} a_u^v a_l^q a_r^\alpha c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) (x_0),
\end{aligned}$$

and

$$\begin{aligned}
(4.3) \quad & \sum_{\beta,l,r,u \neq s \neq t=1}^n \operatorname{tr} \left[c(dx_\beta) c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right] \\
&= \sum_{\beta,l,r,u \neq s \neq t=1}^n \left[-\delta_\beta^t \delta_l^s \delta_u^r + \delta_\beta^t \delta_l^u \delta_s^r + \delta_\beta^s \delta_l^t \delta_u^r - \delta_\beta^s \delta_l^u \delta_t^r - \delta_\beta^u \delta_l^t \delta_s^r + \delta_\beta^u \delta_l^s \delta_t^r \right].
\end{aligned}$$

Obtained from (4.2) and (4.3)

$$(4.4) \quad i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times Q_1](x_0) d\xi_n \sigma(\xi') dx'$$

$$\begin{aligned}
&= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} -\frac{\xi_n}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{v,\beta,l,r,u \neq s \neq t=1}^n \sum_{i,\alpha=1}^{n-1} T_{vst} \xi_i \xi_\alpha a_\beta^i a_l^n a_r^\alpha a_u^v \\
&\quad \times \text{tr} \left[c(dx_\beta) c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right] \\
&\quad - \frac{\xi_n}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{v,\beta,l,r,u \neq s \neq t=1}^n \sum_{i,q=1}^{n-1} T_{vst} \xi_i \xi_q a_\beta^i a_l^q a_r^n a_u^v \\
&\quad \times \text{tr} \left[c(dx_\beta) c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right] \\
&\quad - \frac{i \xi_n^2}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{v,\beta,l,r,u \neq s \neq t=1}^n T_{vst} a_\beta^n a_l^n a_r^n a_u^v \\
&\quad \times \text{tr} \left[c(dx_\beta) c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right] \\
&\quad - \frac{i}{8(\xi_n - i)(1 + \xi_n^2)^2} \sum_{v,\beta,l,r,u \neq s \neq t=1}^n \sum_{\alpha,q=1}^{n-1} T_{vst} \xi_\alpha \xi_q a_\beta^n a_l^q a_r^\alpha a_u^v \\
&\quad \times \text{tr} \left[c(dx_\beta) c(dx_l) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right] (x_0) d\xi_n \sigma(\xi') dx' \\
&= 0.
\end{aligned}$$

$$\begin{aligned}
(4.5) \quad Q_2(x_0)|_{|\xi'|=1} &= \left[-\frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)] c[J(e_l)] c[J(e_j)] T_J c[J(\xi)] \xi_l \xi_j \right. \\
&\quad \left. - \frac{1}{|\xi|^8} \sum_{j,l=1}^n c[J(\xi)] c[J(e_l)] T_J c[J(e_j)] c[J(\xi)] \xi_l \xi_j \right] (x_0)|_{|\xi'|} \\
&= -\frac{1}{(1 + \xi_n^2)^4} \sum_{q,\alpha,l,j,v,r,u \neq s \neq t=1}^n \xi_q \xi_\alpha \xi_l \xi_j T_{vst} a_p^q a_r^\alpha a_k^l a_h^j a_u^v \times \\
&\quad \left[c(dx_p) c(dx_k) c(dx_h) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right. \\
&\quad \left. + c(dx_p) c(dx_k) c(dx_u) c(dx_s) c(dx_t) c(dx_h) c(dx_r) \right] (x_0),
\end{aligned}$$

and

$$(4.6) \quad \text{Tr} \mathfrak{N} := \sum_{\beta,p,k,h,r,u \neq s \neq t=1}^n \text{tr} \left[c(dx_\beta) c(dx_p) c(dx_k) c(dx_h) c(dx_u) c(dx_s) c(dx_t) c(dx_r) \right]$$

$$\begin{aligned}
& + c(dx_\beta)c(dx_p)c(dx_k)c(dx_u)c(dx_s)c(dx_t)c(dx_h)c(dx_r) \Big] \\
& = \sum_{\beta,p,k,h,r,u \neq s \neq t=1}^n \left\{ -2g(dx_h \cdot dx_t) \text{tr} \left[c(dx_\beta)c(dx_p)c(dx_k)c(dx_u)c(dx_s)c(dx_r) \right] \right. \\
& \quad + 2g(dx_h \cdot dx_s) \text{tr} \left[c(dx_\beta)c(dx_p)c(dx_k)c(dx_u)c(dx_t)c(dx_r) \right] \\
& \quad \left. - 2g(dx_h \cdot dx_u) \text{tr} \left[c(dx_\beta)c(dx_p)c(dx_k)c(dx_s)c(dx_t)c(dx_r) \right] \right\}.
\end{aligned}$$

Obtained from (4.5) and (4.6)

$$\begin{aligned}
(4.7) \quad & i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D_{J,T}^{-1}) \times Q_2](x_0) d\xi_n \sigma(\xi') dx' \\
& = i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \frac{\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,q=1}^{n-1} T_{vst} \xi_i \xi_q a_\beta^i a_p^q a_r^n a_k^n a_h^n a_u^v \text{Tr} \mathfrak{N} \right. \\
& \quad + \frac{\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,\alpha=1}^{n-1} T_{vst} \xi_i \xi_\alpha a_\beta^i a_p^\alpha a_r^n a_k^n a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,l=1}^{n-1} T_{vst} \xi_i \xi_l a_\beta^i a_p^n a_r^n a_k^l a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,j=1}^{n-1} T_{vst} \xi_i \xi_j a_\beta^i a_p^n a_r^n a_k^n a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,q,\alpha,l=1}^{n-1} T_{vst} \xi_i \xi_q \xi_\alpha \xi_l a_\beta^i a_p^q a_r^\alpha a_k^l a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,q,\alpha,j=1}^{n-1} T_{vst} \xi_i \xi_q \xi_\alpha \xi_j a_\beta^i a_p^q a_r^\alpha a_k^n a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,q,l,j=1}^{n-1} T_{vst} \xi_i \xi_q \xi_l \xi_j a_\beta^i a_p^q a_r^n a_k^l a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& \quad + \frac{\xi_n}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,\alpha,l,j=1}^{n-1} T_{vst} \xi_i \xi_\alpha \xi_l \xi_j a_\beta^i a_p^n a_r^\alpha a_k^l a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& \quad \left. + \frac{i \xi_n^4}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{i,\alpha,l,j=1}^{n-1} T_{vst} a_\beta^n a_p^n a_r^n a_k^n a_h^n a_u^v \text{Tr} \mathfrak{N} \right\}
\end{aligned}$$

$$\begin{aligned}
& + \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{q,\alpha=1}^{n-1} T_{vst} \xi_q \xi_\alpha a_\beta^n a_p^q a_r^\alpha a_k^n a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& + \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{q,l=1}^{n-1} T_{vst} \xi_q \xi_l a_\beta^n a_p^q a_r^n a_k^l a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& + \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{q,j=1}^{n-1} T_{vst} \xi_q \xi_j a_\beta^n a_p^q a_r^n a_k^n a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& + \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{\alpha,l=1}^{n-1} T_{vst} \xi_\alpha \xi_l a_\beta^n a_p^n a_r^\alpha a_k^l a_h^n a_u^v \text{Tr} \mathfrak{N} \\
& + \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{\alpha,j=1}^{n-1} T_{vst} \xi_\alpha \xi_j a_\beta^n a_p^n a_r^\alpha a_k^n a_h^j a_u^v \text{Tr} \mathfrak{N} \\
& + \left. \frac{i\xi_n^2}{8(\xi_n - i)^2(1 + \xi_n^2)^4} \sum_{l,j=1}^{n-1} T_{vst} \xi_l \xi_j a_\beta^n a_p^n a_r^l a_k^j a_h^n a_u^v \text{Tr} \mathfrak{N} \right\} (x_0) d\xi_n \sigma(\xi') dx' \\
& = 0.
\end{aligned}$$

Combining (4.1) and (4.4) and (4.7) yields

$$\begin{aligned}
\hat{\Psi}_4 & = \frac{\pi}{16} \sum_{i=1}^{n-1} (a_n^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' - \frac{\pi}{16} \sum_{i=1}^{n-1} a_i^i a_n^n \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& + \frac{11\pi}{128} \sum_{l=1}^n (a_n^l)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' - \frac{\pi}{64} \sum_{i=1}^{n-1} (a_n^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& - \frac{23\pi}{320} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_i^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' + \frac{\pi}{64} \sum_{i,j=1}^{n-1} (a_j^i)^2 \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& + \frac{\pi}{64} \sum_{l,\alpha=1}^n \sum_{i=1}^{n-1} (a_l^i)^2 g^M (J(dx_\alpha), (\nabla_{e_\alpha}^L J) e_n) \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& - \frac{\pi}{16} \sum_{i=1}^{n-1} g^M (J(dx_i), (\nabla_{e_n}^L J) e_i) \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& + \frac{\pi}{16} \sum_{i=1}^{n-1} g^M (J(dx_n), (\nabla_{e_i}^L J) e_i) \text{tr}[\text{id}] \Omega_4 h'(0) dx' \\
& - \frac{\pi}{64} \sum_{l,\alpha=1}^n (a_l^n)^2 g^M (J(dx_\alpha), (\nabla_{e_\alpha}^L J) e_n) \text{tr}[\text{id}] \Omega_4 h'(0) dx'
\end{aligned}$$

$$+ \frac{\pi}{8} \sum_{l=1}^n \sum_{i=1}^{n-1} a_l^i \partial_{x_i} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.$$

case $\hat{\mathbf{c}}$ $r = -2$, $l = -3$, $k = j = |\alpha| = 0$
By (3.18), we calculate that

$$\hat{\Psi}_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\pi_{\xi_n}^+ \sigma_{-2}(D_{J,T}^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx'$$

Same classification as in case b (3.9) By computation, we have

$$(4.8) \quad \begin{aligned} & -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\pi_{\xi_n}^+ B_0(x_0) \times \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -\frac{\pi}{64} \sum_{l,j,\beta=1}^n (a_\beta^n)^2 a_l^j \partial_{x_j} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' - \frac{\pi}{64} \sum_{l,j,\beta=1}^n (a_\beta^i)^2 a_l^j \partial_{x_j} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' \\ &+ \frac{\pi}{256} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_l^n)^2 a_i^j a_n^n \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' + \frac{3\pi}{256} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_l^i)^2 a_\nu^\nu a_n^n \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\ &- \frac{\pi}{32} \sum_{l,j=1}^n a_l^j \partial_{x_j} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' + \frac{\pi}{256} \sum_{l,\beta=1}^n (a_\beta^n)^2 (a_l^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\ &+ \frac{49\pi}{1280} \sum_{l,\beta=1}^n \sum_{i=1}^{n-1} (a_l^n)^2 (a_\beta^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{3\pi}{256} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_l^n)^2 (a_i^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\ &- \frac{5\pi}{256} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{\pi}{64} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_l^n)^2 (a_\nu^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \end{aligned}$$

According to (4.3) and (3.9), the following equation holds

$$(4.9) \quad \begin{aligned} & -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \operatorname{tr}[\pi_{\xi_n}^+ B_1(x_0) \times \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \\ & \left\{ -\frac{\xi_n}{4(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,\alpha=1}^{n-1} T_{vst} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^\nu a_\beta^i \operatorname{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \right\} \\ & - \frac{\xi_n}{4(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta,l,r,v,u \neq s \neq t=1}^n \sum_{i,q=1}^{n-1} T_{vst} \xi_i \xi_q a_l^n a_r^q a_u^\nu a_\beta^i \operatorname{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \end{aligned}$$

$$\begin{aligned}
& + \frac{\xi_n - 3\xi_n^3}{16(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n T_{vst} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \\
& - \frac{i(1 - 3\xi_n^2)(i\xi_n + 2)}{16(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u \neq s \neq t=1}^n \sum_{q, \alpha=1}^{n-1} T_{vst} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_u)c(dx_s)c(dx_t)c(dx_r)c(dx_\beta)] \Big\} \\
& (x_0) d\xi_n \sigma(\xi') dx' \\
& = 0.
\end{aligned}$$

(4.10)

$$\begin{aligned}
& - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{tr}[\pi_{\xi_n}^+ B_2(x_0) \times \partial_{\xi_n} \sigma_{-3}(D_{J,T}^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
& = - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \frac{\xi_n}{2(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_l^n a_r^\alpha a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \right. \\
& + \frac{\xi_n}{2(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_r^n a_u^v a_\beta^i \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& + \frac{i(1 - 3\xi_n^2)(i\xi_n + 2)}{8(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_r^\alpha a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \\
& \left. - \frac{\xi_n - 3\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n T_{vut} a_l^n a_r^n a_u^v a_\beta^n \text{tr}[c(dx_l)c(dx_t)c(dx_r)c(dx_\beta)] \right\} (x_0) d\xi_n \sigma(\xi') dx' \\
& = - i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \frac{\xi_n}{2(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{i, \alpha=1}^{n-1} T_{vut} \xi_i \xi_\alpha a_t^n a_r^\alpha a_u^v a_l^i \right. \\
& + \frac{\xi_n}{2(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{i, q=1}^{n-1} T_{vut} \xi_i \xi_q a_l^q a_t^n a_u^v a_l^i \\
& - \frac{i(1 - 3\xi_n^2)(i\xi_n + 2)}{8(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n \sum_{q, \alpha=1}^{n-1} T_{vut} \xi_q \xi_\alpha a_l^q a_l^\alpha a_u^v a_t^n \\
& \left. - \frac{\xi_n - 3\xi_n^3}{8(\xi_n - i)^2(1 + \xi_n^2)^3} \sum_{\beta, l, r, v, u, t=1}^n T_{vut} a_t^n a_r^n a_u^v a_r^n \right\} (x_0) d\xi_n \sigma(\xi') dx' \\
& = 0.
\end{aligned}$$

(4.8)-(4.10)

$$\hat{\Psi}_5 = -\frac{\pi}{64} \sum_{l, j, \beta=1}^n (a_\beta^n)^2 a_l^j \partial_{x_j} (a_l^n) \text{tr}[\text{id}] \Omega_4 dx' - \frac{\pi}{64} \sum_{l, j, \beta=1}^n \sum_{i=1}^{n-1} (a_\beta^i)^2 a_l^j \partial_{x_j} (a_l^n) \text{tr}[\text{id}] \Omega_4 dx'$$

$$\begin{aligned}
& + \frac{\pi}{256} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_l^n)^2 a_i^i a_n^n \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' + \frac{3\pi}{256} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_l^i)^2 a_\nu^\nu a_n^n \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& - \frac{\pi}{32} \sum_{l,j=1}^n a_l^j \partial_{x_j} (a_l^n) \operatorname{tr}[\operatorname{id}] \Omega_4 dx' + \frac{\pi}{256} \sum_{l,\beta=1}^n (a_\beta^n)^2 (a_l^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& + \frac{49\pi}{1280} \sum_{l,\beta=1}^n \sum_{i=1}^{n-1} (a_l^n)^2 (a_\beta^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{3\pi}{256} \sum_{l=1}^n \sum_{i=1}^{n-1} (a_i^n)^2 (a_l^n)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' \\
& - \frac{5\pi}{256} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_\nu^n)^2 (a_l^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx' - \frac{\pi}{64} \sum_{l=1}^n \sum_{\nu,i=1}^{n-1} (a_l^n)^2 (a_\nu^i)^2 \operatorname{tr}[\operatorname{id}] \Omega_4 h'(0) dx'.
\end{aligned}$$

By simple calculation

$$\sum_{i=1}^{n-1} g^M (J(dx_i), (\nabla_{e_n}^L J) e_i) (x_0) = \sum_{i=1}^{n-1} g^M (J(dx_i), \nabla_{e_n}^L J(e_i) - J(\nabla_{e_n}^L e_i)) (x_0) = 0,$$

by similar calculations, we can get

$$\begin{aligned}
& \sum_{i=1}^{n-1} g^M (J(dx_n), (\nabla_{e_i}^L J) e_i) (x_0) = 0; \\
& \sum_{\alpha=1}^n g^M (J(dx_\alpha), (\nabla_{e_\alpha}^L J) e_n) (x_0) = 0.
\end{aligned}$$

In summary,

$$\Psi = \Psi_1 + \Psi_2 + \Psi_3 + \Psi_4 + \Psi_5 = -\frac{\pi}{2} (1 - \langle J(e_6), e_6 \rangle^2) h'(0) \Omega_4 dx'.$$

Theorem 4.3. *Let M be a 6-dimensional almost product Riemannian spin manifold with the boundary ∂M and the metric g^M as above, $D_{J,T}$ be the J -twist of the Dirac operator with torsion on \widetilde{M} , then*

(4.11)

$$\begin{aligned}
& \widetilde{\operatorname{Wres}}[\pi^+ D_{J,T}^{-1} \circ \pi^+ D_{J,T}^{-1}] \\
& = \int_M 4\pi^2 \left\{ \sum_{i,j=1}^n R(J(e_i), J(e_j), e_j, e_i) - 2 \sum_{\nu,j=1}^n g^M(\nabla_{e_j}^L(J)e_\nu, (\nabla_{e_\nu}^L J)e_j) \right. \\
& \quad - 2 \sum_{\nu,j=1}^n g^M(J(e_\nu), (\nabla_{e_j}^L(\nabla_{e_\nu}^L(J)))e_j - (\nabla_{\nabla_{e_j}^L e_\nu}^L(J))e_j) \\
& \quad \left. - \sum_{\alpha,\nu,j=1}^n g^M(J(e_\alpha), (\nabla_{e_\nu}^L J)e_j) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\nu)) \right\}
\end{aligned}$$

$$\begin{aligned}
& - \sum_{\alpha, \nu, j=1}^n g^M(J(e_\alpha), (\nabla_{e_\alpha}^L J)e_j) g^M(J(e_\nu), (\nabla_{e_\nu}^L J)e_j) + \sum_{\nu, j=1}^n g^M((\nabla_{e_\nu}^L J)e_j, (\nabla_{e_\nu}^L J)e_j) - \frac{5}{3}s \\
& - 2 \sum_{j, m, p} T(e_m, e_p, J(e_m)) g^M((\nabla_{e_j}^L J)e_j, e_p) - 2 \sum_{j, m, \alpha} T(e_m, J(e_j), J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_m)) \\
& - 2 \sum_{j, m, \alpha} T(e_m, J(e_j), J(e_m)) g^M((\nabla_{e_\alpha}^L J)e_j, J(e_\alpha)) + 2 \sum_{m, p, \alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_\alpha}^L J)e_m, e_p) \\
& - 2 \sum_{m, p, \alpha} T(e_m, e_p, J(e_\alpha)) g^M((\nabla_{e_m}^L J)e_\alpha, e_p) - \sum_{j, p, m} T(e_j, e_p, J(e_m)) T(e_m, e_p, J(e_j)) \\
& - 2 \sum_{j, l, \alpha} T^2(J(e_j), e_l, e_\alpha) \Big\} d\text{Vol}_M - \frac{\pi}{2} \int_{\partial M} (1 - \langle J(e_6), e_6 \rangle^2) h'(0) \Omega_4 d\text{Vol}_{\partial M}
\end{aligned}$$

where s is the scalar curvature.

5. DECLARATIONS

Ethics approval and consent to participate No applicable.

Consent for publication No applicable.

Availability of data and material The authors confirm that the data supporting the findings of this study are available within the article.

Competing interests The authors declare no conflict of interest.

Funding Sponsored by Natural Science Foundation of Xinjiang Uygur Autonomous-Region 2024D01C341 and the National Natural Science Foundation of China 11771070, 12061078.

Authors' contributions All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Jin Hong, Siyao Liu and Yong Wang. The first draft of the manuscript was written by Jin Hong and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

REFERENCES

- [1] Guillemin V W., A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. Adv. Math., 1985, 55(2): 131-160.
- [2] Wodzicki M., Local invariants of spectral asymmetry. Invent. Math., 1995, 75(1): 143-178.

- [3] Connes A., Quantized calculus and applications. 11th International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, 1995.
- [4] Connes A., The action functional in noncommutative geometry. *Comm. Math. Phys.*,1998, 117: 673-683.
- [5] Kastler D., The Dirac operator and gravitation. *Comm. Math. Phys.*,1995, 166: 633-643.
- [6] Kalau W., Walze M., Gravity, noncommutative geometry and the Wodzicki residue. *J. Geom. Phys.*,1995, 16:327-344.
- [7] Ackermann T.,A note on the Wodzicki residue. *J. Geom. Phys.*,1996, 20:404-406.
- [8] Ugalde W J., Differential forms and the Wodzicki residue. *J. Geom. Phys.*,2008, 58: 1739-1751.
- [9] Wang Y., Differential forms and the Wodzicki residue for manifolds with boundary. *J. Geom. Phys.*,2006, 56: 731-753.
- [10] Wang Y., Differential forms the noncommutative residue for manifolds with boundary in the non-product case. *Lett. Math. Phys.*,2006, 77: 41-51.
- [11] Wang Y., Gravity and the noncommutative residue for manifolds with boundary. *Lett. Math. Phys.*,2007, 80:37-56.
- [12] Wang Y., Lower-dimensional volumes and Kastler-Kalau-Walze type theorem for manifolds with boundary. *Commun. Theor. Phys.*, 54: 38-42.
- [13] Wang J., Wang Y., The Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. *J. Math. Phys.*,2015, 56: 052501.
- [14] Wei S., Wang Y., Modified Novikov operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. *Adv. Math. Phys.*,2020: 1-28.
- [15] Wu T., Wang J., Wang Y., Dirac-Witten operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. *J. Nonlinear Math. Phys.*,2022, 29: 1-40.
- [16] Wang Y., A Kastler-Kalau-Walze type theorem and the spectral action for perturbations of Dirac operators on manifolds with boundary. *Abstr. Appl. Anal.*,2014, 17: 286-299.
- [17] Kim E C., Lower bounds of the Dirac eigenvalues on compact Riemannian spin manifolds with locally product structure. *arXiv:math/0402427*.
- [18] Chen Y., The Dirac operator on locally reducible Riemannian manifolds. *J. Geom. Phys.*,2019, 139: 17-24.
- [19] Chen Y., The Dirac operator on manifold admitting parallel one-form. *J. Geom. Phys.*,2017, 117: 214-221.
- [20] Fedosov B V., Golse F., Leichtnam E., Schrohe E., The noncommutative residue for manifolds with boundary. *J. Funct. Anal.*,1996, 142: 1-31.
- [21] Ponge R., Noncommutative geometry and lower dimensional volumes in Riemannian geometry. *Lett. Math. Phys.*,2008, 83(1):19-32.
- [22] Liu S., Wang Y., A Kastler-Kalau-Walze type theorem for the J -twist D_J of the Dirac operator. *J. Nonlinear Math. Phys.*,2023, 30(2): 743-749.
- [23] Liu S., Wang Y., The J -twist D_J of the Dirac operator and the Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. *Int. J. Geom. Methods Mod. Phys.*,2023, 20(11):2350188.
- [24] Bismut J M., A local index theorem for non Kähler manifolds. *Math. Ann.*,1989, 284: 681-699.
- [25] Ackermann T., Tolksdorf J.: A generalized Lichnerowicz formula, the Wodzicki residue and gravity. *J. Geom. Physics.*,1996, 19:143-150.
- [26] Wang J., Wang Y., Yang C L., Dirac operators with torsion and the noncommutative residue for manifolds with boundary, *J. Geom. Phys.*,2014, 81:92-111 .

J. Hong

School of Mathematics and Statistics, Yili Normal University, Yining 835000, China

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China

Email: jhong@nenu.edu.cn

S. Liu

School of Mathematics and Statistics, Changchun University of Science and Technology, Changchun 130022, China

Email: liusy719@nenu.edu.cn

Y. Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China

Email: wangy581@nenu.edu.cn