

Categorification of modules and construction of schemes

Abhishek Banerjee ^{*} Subhajit Das [†] Surjeet Kour [‡]

Abstract

We use categorification of module structures to study algebraic geometry over symmetric monoidal categories. This brings together the relative algebraic geometry over symmetric monoidal categories developed by Toën and Vaquié, along with the theory of module categories over monoidal categories. We obtain schemes over a datum $(\mathcal{C}, \mathcal{M})$, where $(\mathcal{C}, \otimes, 1)$ is a symmetric monoidal category and \mathcal{M} is a module category over \mathcal{C} . One of our main tools is using the datum $(\mathcal{C}, \mathcal{M})$ to give a Grothendieck topology on the category of affine schemes over $(\mathcal{C}, \otimes, 1)$ that we call the “spectral \mathcal{M} -topology.” This consists of “fpqc \mathcal{M} -coverings” with certain special properties. We also give a counterpart for a construction of Connes and Consani by presenting a notion of scheme over a composite datum consisting of a \mathcal{C} -module category \mathcal{M} and the category of commutative monoids with an absorbing element.

MSC(2020) Subject Classification: 14A30, 18M05

Keywords : symmetric monoidal categories, module categories, fpqc coverings

1 Introduction

In this paper, we use categorification of module structures to study algebraic geometry over symmetric monoidal categories. The relative algebraic geometry over a symmetric monoidal category $(\mathcal{C}, \otimes, 1)$ was developed by Toën and Vaquié [32]. Module categories over tensor categories have been studied in various contexts by a number of authors (see, for instance, Bénabou [4], Janelidze and Kelly [19], Ostrik [26], Street [28]). We develop a notion of scheme over a datum $(\mathcal{C}, \mathcal{M})$, where \mathcal{C} is a symmetric monoidal category and \mathcal{M} is a module category over \mathcal{C} .

The idea of doing algebraic geometry over symmetric monoidal categories has roots in the work of Deligne [13], as also in that of Hakim [18]. In [32], Toën and Vaquié gave a notion of scheme over a symmetric monoidal category $(\mathcal{C}, \otimes, 1)$ that is purely categorical, along with notions of Zariski immersions, Zariski coverings and fpqc coverings. By choosing the category $(\mathcal{C}, \otimes, 1)$ appropriately, one opens up a number of new geometries by taking \mathcal{C} for instance to be the category of sets, or commutative monoids, or the category of symmetric spectra. When \mathcal{C} is taken to be the category of \mathbb{Z} -modules, one recovers the usual notion of a scheme over $\text{Spec}(\mathbb{Z})$. There is also a homotopy version of the theory in [32], obtained by taking \mathcal{C} to be a symmetric monoidal model category. The latter has been developed into a theory of homotopical algebraic geometry using higher categorical structures by Toën and Vezzosi in [30], [31]. The ideas of [32] have also been explored for instance in [1], [2], [3], [23], [24] (see also related work by Connes and Consani in [9], [10], [11], [12]).

We know that a symmetric monoidal category $(\mathcal{C}, \otimes, 1)$ may be seen as the categorification of a commutative ring. In the framework of Toën and Vaquié [32], the category of affine schemes is taken to be the opposite of that of commutative monoid objects in $(\mathcal{C}, \otimes, 1)$. Thereafter, the theory in [32] is developed by using objects in \mathcal{C} equipped with module actions of commutative monoid objects in $(\mathcal{C}, \otimes, 1)$. This suggests that we should be able to study a

^{*}Department of Mathematics, Indian Institute of Science, Bangalore. Email: abhishekbannerjee1313@gmail.com.

[†]Department of Mathematics, Indian Institute of Science, Bangalore. Email: subhajitdas@iisc.ac.in.

[‡]Department of Mathematics, Indian Institute of Technology, Delhi. Email: koursurjeet@gmail.com.

similar algebraic geometry by categorifying not just the notion of a commutative monoid, but also that of modules over it. Therefore, the starting point of this paper is to consider a datum $(\mathcal{C}, \mathcal{M})$, where \mathcal{C} is a symmetric monoidal category and \mathcal{M} is a category equipped with a functor

$$\boxtimes : \mathcal{C} \times \mathcal{M} \longrightarrow \mathcal{M} \quad (1.1)$$

satisfying certain conditions similar to that of a module over a ring (see Section 2 for details). We mention here that the framework of module categories over monoidal categories has proved to be extremely versatile, with applications in various fields such as the theory of subfactors (see [5], [25]), weak Hopf algebras (see [26], [29]) and in conformal field theory (see [17]). This theory becomes particularly rich when the monoidal category has certain additional structures, such as rigidity or (multi-)fusion structures (see for instance, [14], [15]). As such, module categories over monoidal categories have increasingly become a topic of study in their own right. For more on this subject, we refer the reader for instance to [16, § 5].

We start with a datum $(\mathcal{C}, \mathcal{M})$ as in (1.1). Given a commutative monoid object a in $(\mathcal{C}, \otimes, 1)$, we consider the category $\mathbf{Mod}_{\mathcal{M}}(a)$ of a -modules in \mathcal{M} . The objects of $\mathbf{Mod}_{\mathcal{M}}(a)$ are pairs (m, ρ) where $m \in \mathcal{M}$ and $\rho : a \boxtimes m \longrightarrow m$ is a morphism satisfying conditions similar to a module action (see Definition 2.1). We then build up the key notions of coverings by using properties of the adjoint functors of “restriction of scalars” and “extension of scalars” between these module categories.

One of the main tools in our paper is the use of the datum $(\mathcal{C}, \mathcal{M})$ to define a Grothendieck topology on the category $Aff_{\mathcal{C}}$ of affine schemes over $(\mathcal{C}, \otimes, 1)$, which we refer to as the “spectral \mathcal{M} -topology.” This consists of “fpqc \mathcal{M} -coverings” with certain special properties that we explain in Section 3. We obtain schemes over $(\mathcal{C}, \mathcal{M})$ using coverings by means of affine schemes in the category of sheaves for the spectral \mathcal{M} -topology on $Aff_{\mathcal{C}}$. We will typically refer to these as \mathcal{M} -schemes. The category of \mathcal{M} -schemes appears to have good properties in that it is closed under pullbacks, disjoint unions and any scheme may be recovered as the quotient of an equivalence relation on the disjoint union of affine schemes. By considering simplicial \mathcal{M} -schemes, we are also able to obtain a good theory of descent data corresponding to a morphism of \mathcal{M} -schemes. We also incorporate change of base functors between categories of schemes over respective data $(\mathcal{C}, \mathcal{M})$ and $(\mathcal{C}', \mathcal{M}')$ as in (1.1) connected by a lax functor and an adjunction between the symmetric monoidal categories \mathcal{C} and \mathcal{C}' .

In the final part of this paper, we suggest a notion of scheme over a composite datum consisting of a \mathcal{C} -module category \mathcal{M} and the category $CMon_0$ of commutative monoids with an absorbing element. This is inspired by the work of Connes and Consani [9], who presented a notion of scheme over a gluing of the category of commutative monoids with that of commutative rings, by means of adjoint functors. We mention here that in future, we hope to extend our methods to the homotopical context, where \mathcal{C} and \mathcal{M} additionally carry model structures.

We now describe the paper in more detail. We begin with a module category \mathcal{M} over \mathcal{C} as in (1.1). Let $Comm(\mathcal{C})$ denote the category of commutative monoid objects in \mathcal{C} . For $a \in Comm(\mathcal{C})$, the category $\mathbf{Mod}_{\mathcal{M}}(a)$ of a -modules with values in \mathcal{M} is described in Section 2. For a morphism $\alpha : a \longrightarrow b$ in $Comm(\mathcal{C})$, we construct an adjoint pair (α^*, α_*) of extension and restriction of scalars between the categories $\mathbf{Mod}_{\mathcal{M}}(a)$ and $\mathbf{Mod}_{\mathcal{M}}(b)$. Some of the material in Section 2 is possibly well known, but we record it here in order to fix notation. Further, we show that the assignment $a \mapsto \mathbf{Mod}_{\mathcal{M}}(a)$ determines a pseudo-functor from $Comm(\mathcal{C})$ to the category of categories.

As in [32], we take the category $Aff_{\mathcal{C}}$ of affine schemes to be the opposite of the category $Comm(\mathcal{C})$ of commutative monoid objects in \mathcal{C} . For any $a \in Comm(\mathcal{C})$, we let $Spec(a) \in Aff_{\mathcal{C}} = Comm(\mathcal{C})^{op}$ denote the corresponding affine scheme. Similarly, for a morphism $\alpha : a \longrightarrow b$ in $Comm(\mathcal{C})$, we denote by $\alpha^{op} : Spec(b) \longrightarrow Spec(a)$ the corresponding morphism of affine schemes in $Aff_{\mathcal{C}}$.

In Section 3, we define the topology on $Aff_{\mathcal{C}}$. We say that a morphism $\alpha : a \longrightarrow b$ in $Comm(\mathcal{C})$ is \mathcal{M} -flat if $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ is exact. We will say that a collection $\{\alpha_i^{op} : Spec(a_i) \longrightarrow Spec(a)\}_{i \in I}$ of \mathcal{M} -flat morphisms is an fpqc \mathcal{M} -cover for $Spec(a)$ if there exists a finite subset $J \subseteq I$ such that the collection of functors $\{\alpha_j^* : \mathbf{Mod}_{\mathcal{M}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(a_j)\}_{j \in J}$ is conservative (see Definition 3.3). If $\alpha : a \longrightarrow b$ is \mathcal{M} -flat and an epimorphism of finite type in $Comm(\mathcal{C})$, we will say that α^{op} is a spectral immersion relative to \mathcal{M} . A spectral

\mathcal{M} -cover $\{\alpha_i^{op} : Spec(a_i) \rightarrow Spec(a)\}_{i \in I}$ is an fpqc \mathcal{M} -cover such that each α_i^{op} is a spectral immersion relative to \mathcal{M} . The main result of Section 3 is that both the fpqc \mathcal{M} -coverings as well as the spectral \mathcal{M} -coverings define Grothendieck topologies on $Aff_{\mathcal{C}}$ (see Proposition 3.7). From then onwards, we set $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ be the category of sheaves on $Aff_{\mathcal{C}}$ with respect to the spectral \mathcal{M} -topology.

We say that the \mathcal{C} -module category (\mathcal{M}, \boxtimes) is subcanonical if the fpqc \mathcal{M} -topology (and hence, the spectral \mathcal{M} -topology) on $Aff_{\mathcal{C}}$ is subcanonical, i.e., for each $Spec(a) \in Aff_{\mathcal{C}}$, the representable presheaf $Aff_{\mathcal{C}}(_, Spec(a))$ on $Aff_{\mathcal{C}}$ is a sheaf for the fpqc \mathcal{M} -topology on $Aff_{\mathcal{C}}$. In this situation, an affine scheme $Spec(a) \in Aff_{\mathcal{C}}$ may be treated as an object of $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. For most of this paper, we will assume that (\mathcal{M}, \boxtimes) is subcanonical. We mention that later on in Section 6, we have collected a number of examples of subcanonical module categories arising in various natural ways such as from diagonal actions, functor categories, representation categories of monoids and from directed graphs.

In Section 4, we give the definition of Zariski open immersion relative to \mathcal{M} for a morphism $f : F \rightarrow G$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ (see Definition 4.2). We show that this notion is stable under base change and closed under composition in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. By an \mathcal{M} -scheme (see Definition 4.8), we mean a sheaf $F \in Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ for the spectral \mathcal{M} -topology on $Aff_{\mathcal{C}}$ such that there is an epimorphism

$$\coprod_{i \in I} X_i \rightarrow F \quad (1.2)$$

in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ where each X_i is an affine scheme and each $X_i \rightarrow F$ is a Zariski open immersion relative to \mathcal{M} . We show that the full subcategory $Sch(\mathcal{C})_{\mathcal{M}}$ of $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ consisting of \mathcal{M} -schemes is closed under coproducts and pullbacks. We also provide a description (see Theorem 4.14) of \mathcal{M} -schemes in terms of quotients of disjoint unions of affine schemes over an equivalence relation satisfying certain properties.

In Section 5, we consider symmetric monoidal categories $(\mathcal{C}, \otimes, 1)$ and $(\mathcal{D}, \otimes, 1)$ which are connected by an adjunction $(\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}, \mathbf{A} : \mathcal{D} \rightarrow \mathcal{C})$ that satisfies certain monoidal properties. We also consider a \mathcal{C} -module category (\mathcal{M}, \boxtimes) , a \mathcal{D} -module category (\mathcal{N}, \boxplus) and a lax \mathcal{C} -linear functor

$$(\mathbb{L}, \Gamma) : \mathbf{B}_*(\mathcal{N}) = (\mathcal{N}, \boxplus^{\mathbf{B}}) \rightarrow (\mathcal{M}, \boxtimes) \quad (1.3)$$

where $\boxplus^{\mathbf{B}}$ denotes the restriction of the \mathcal{D} -module action on \mathcal{N} to a \mathcal{C} -module action by means of the functor $\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}$. In that case, we have an adjoint pair (see Theorem 5.3)

$$(\widetilde{\mathbf{A}}_! : Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \rightarrow Sh(Aff_{\mathcal{D}})_{\mathcal{N}}, \quad \widetilde{\mathbf{B}}_! : Sh(Aff_{\mathcal{D}})_{\mathcal{N}} \rightarrow Sh(Aff_{\mathcal{C}})_{\mathcal{M}}) \quad (1.4)$$

Additionally, if \mathcal{M} and \mathcal{N} are subcanonical, then we obtain a change of base functor $\widetilde{\mathbf{A}}_! : Sch(\mathcal{C})_{\mathcal{M}} \rightarrow Sch(\mathcal{D})_{\mathcal{N}}$ at the level of schemes. In Section 7, we study a category $Desc(F \xrightarrow{f} S)$ of descent data for a morphism $f : F \rightarrow S$ of \mathcal{M} -schemes. More precisely, we associate a simplicial \mathcal{M} -scheme $\mathbb{X}(f)$ to the morphism $f : F \rightarrow S$ and show (see Proposition 7.6) that the category of descent data is equivalent to that of simplicial \mathcal{M} -schemes that are cartesian over $\mathbb{X}(f)$.

In [9], Connes and Consani present a notion of a scheme over the field \mathbb{F}_1 with one element. For this, they begin by gluing together the category of commutative rings with that of commutative monoids by means of an adjunction. Then, a scheme over \mathbb{F}_1 (see [9, Definition 4.7]) consists of a \mathbb{Z} -scheme, a scheme over the category of commutative monoids and a natural transformation which gives a bijection when applied to a field. Our objective in Section 8 is to present a counterpart of this construction. For this, we glue together the category $Comm(\mathcal{C})$ with the category $CMon_0$ of commutative monoids with an absorbing element. By considering units in the monoid $\mathcal{C}(1, c)$ for $c \in Comm(\mathcal{C})$, we obtain a collection $fld(\mathcal{C})$ of objects that approximate the idea of a “field” in \mathcal{C} . We conclude by defining the notion of $(\mathcal{M}, CMon_0)$ -scheme, which consists of an \mathcal{M} -scheme, a scheme over $CMon_0$ in the sense of [9] and a natural transformation that gives a bijection when applied to any $c \in fld(\mathcal{C})$ (see Definition 8.3).

Acknowledgements: Authors A.B. and S.K. are grateful to the Research in Pairs program at the CIRM in Marseille, where part of this paper was written. Author S.D. was supported by Prime Minister’s Research Fellowship PMRF-21-4890.03.

2 Categorical actions and the pseudo-functor $\mathbf{Mod}_{\mathcal{M}}$

Throughout, we let $(\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete. We recall (see [8], [19]) that a left \mathcal{C} -module category $(\mathcal{M}, \boxtimes, \lambda, l)$ consists of a bifunctor $\boxtimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ and a collection of isomorphisms

$$\begin{aligned}\lambda &= \left(\lambda_{a,b,m} : a \boxtimes (b \boxtimes m) \xrightarrow{\sim} (a \otimes b) \boxtimes m \mid a, b \in \mathcal{C}, m \in \mathcal{M} \right) \\ l &= \left(l_m : 1 \boxtimes m \xrightarrow{\sim} m \mid m \in \mathcal{M} \right)\end{aligned}\tag{2.1}$$

which are natural and subject to certain coherence axioms. Right \mathcal{C} -module categories are defined similarly. Let $\mathcal{M} = (\mathcal{M}, \boxtimes)$ and $\mathcal{M}' = (\mathcal{M}', \boxtimes')$ be left \mathcal{C} -module categories. A lax \mathcal{C} -linear functor $(L, J) : \mathcal{M} \rightarrow \mathcal{M}'$ consists of a functor $L : \mathcal{M} \rightarrow \mathcal{M}'$ together with a natural transformation $J = \{J_{c,m} : c \boxtimes' L(m) \rightarrow L(c \boxtimes m) : (c, m) \in \mathcal{C} \times \mathcal{M}\}$ satisfying certain coherence axioms (see [8], [19]). Additionally, if J is a natural isomorphism, then (L, J) is called a strong \mathcal{C} -linear functor. We denote the category of left \mathcal{C} -module categories and lax \mathcal{C} -linear functors by $\mathcal{C} - \mathbf{Mod}^{lx}$.

From now onwards, we fix a left \mathcal{C} -module category $(\mathcal{M}, \boxtimes, \lambda, l)$ such that the underlying category \mathcal{M} is both complete and cocomplete. Additionally, for every $a \in \mathcal{C}, m \in \mathcal{M}$, we assume that the functors $a \boxtimes _ : \mathcal{M} \rightarrow \mathcal{M}$ and $_ \boxtimes m : \mathcal{C} \rightarrow \mathcal{M}$ preserve colimits. We let $\mathbf{Comm}(\mathcal{C})$ denote the category of commutative monoid objects in \mathcal{C} .

Definition 2.1. Let $a = (a, \mu : a \otimes a \rightarrow a, \iota : 1 \rightarrow a) \in \mathbf{Comm}(\mathcal{C})$. A left a -module in \mathcal{M} is a pair (m, ρ) consisting of an object $m \in \mathcal{M}$ and a morphism $\rho : a \boxtimes m \rightarrow m$ such that

$$\rho \circ (\mu \boxtimes 1_m) \circ \lambda_{a,a,m} = \rho \circ (1_a \boxtimes \rho) : a \boxtimes (a \boxtimes m) \rightarrow m \quad \rho \circ (\iota \boxtimes 1_m) = l_m : 1 \boxtimes m \rightarrow m \tag{2.2}$$

A morphism $\psi : (m, \rho) \rightarrow (m', \rho')$ of left a -modules is a morphism $\psi : m \rightarrow m'$ in \mathcal{M} such that $\rho' \circ (1_a \boxtimes \psi) = \psi \circ \rho$. For $a \in \mathbf{Comm}(\mathcal{C})$, we let $\mathbf{Mod}_{\mathcal{M}}(a)$ denote the category of left a -modules.

Remark 2.2. For $a \in \mathbf{Comm}(\mathcal{C})$, we may consider the associated monad $\mathbf{T} = a \boxtimes _$ on \mathcal{M} . Then, the Eilenberg-Moore category of \mathbf{T} -algebras is the category $\mathbf{Mod}_{\mathcal{M}}(a)$. Hence, the forgetful functor $U_a : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathcal{M}$ reflects isomorphisms and creates limits. Further, since the endofunctor underlying \mathbf{T} preserves colimits, U_a also creates colimits. In particular, since \mathcal{M} is complete and cocomplete, it follows that $\mathbf{Mod}_{\mathcal{M}}(a)$ is complete and cocomplete and U_a preserves both limits and colimits.

Let $\alpha : (a, \mu_a, \iota_a) \rightarrow (b, \mu_b, \iota_b)$ be a morphism in $\mathbf{Comm}(\mathcal{C})$. We consider the restriction of scalars $\alpha_* : \mathbf{Mod}_{\mathcal{M}}(b) \rightarrow \mathbf{Mod}_{\mathcal{M}}(a)$, given by $(m, \rho) \mapsto (m, \rho \circ (\alpha \boxtimes 1_m))$. The extension of scalars $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ may now be defined as follows: for $(m, \rho) \in \mathbf{Mod}_{\mathcal{M}}(a)$, we consider the object

$$b \boxtimes_a m := \mathit{Coeq} \left(b \boxtimes (a \boxtimes m) \begin{array}{c} \xrightarrow{1_b \boxtimes \rho} \\ \xrightarrow{\lambda_{b,a,m}} \end{array} (b \otimes a) \boxtimes m \xrightarrow{(\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_m} b \boxtimes m \right) \tag{2.3}$$

in \mathcal{M} . We denote the canonical epimorphism $b \boxtimes m \rightarrow b \boxtimes_a m$ by $\mathit{coeq}_{\alpha, (m, \rho)}$. Using Remark 2.2, it follows that (2.3) is also a coequalizer in $\mathbf{Mod}_{\mathcal{M}}(b)$.

Lemma 2.3. There is a canonical morphism $\tilde{\rho} : b \boxtimes (b \boxtimes_a m) \rightarrow b \boxtimes_a m$ making $b \boxtimes_a m$ into a left b -module.

Proof. We consider the following diagram :

$$\begin{array}{ccccccc}
b \boxtimes (b \boxtimes (a \boxtimes m)) & \xrightarrow{1_b \boxtimes (1_b \boxtimes \rho)} & b \boxtimes ((b \otimes a) \boxtimes m) & \xrightarrow{1_b \boxtimes ((\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_m)} & b \boxtimes (b \boxtimes m) & \xrightarrow{1_b \boxtimes coeq} & b \boxtimes (b \boxtimes_a m) \\
\downarrow \lambda_{b,b,a \boxtimes m} & & & & \downarrow \lambda_{b,b,m} & & \downarrow 1_b \boxtimes (b \boxtimes_a m) \\
(b \otimes b) \boxtimes (a \boxtimes m) & \xrightarrow{\lambda_{b \otimes b, a, m}} & ((b \otimes b) \otimes a) \boxtimes m & \xrightarrow{\sim} & (b \otimes (b \otimes a)) \boxtimes m & \xrightarrow{(1_b \otimes (\mu_b \circ (1_b \otimes \alpha))) \boxtimes 1_m} & (b \otimes b) \boxtimes m \longrightarrow b \boxtimes (b \boxtimes_a m) \\
\downarrow \mu_b \boxtimes 1_{a \boxtimes m} & & & & \downarrow \mu_b \boxtimes 1_m & & \downarrow \text{dashed} \\
b \boxtimes (a \boxtimes m) & \xrightarrow{\lambda_{b,a,m}} & (b \otimes a) \boxtimes m & \xrightarrow{1_b \boxtimes \rho} & b \boxtimes m & \xrightarrow{coeq} & b \boxtimes_a m \\
& & & & & & \downarrow \text{dashed}
\end{array} \quad (2.4)$$

Since $b \boxtimes _$ preserves colimits, it is evident that the top row of (2.4) is a coequalizer.

(1) Because of the naturality of the isomorphisms in λ , the associativity constraints $\lambda_{b,b,a \boxtimes m}$ and $\lambda_{b,b,m}$ satisfy $\lambda_{b,b,m} \circ (1_b \boxtimes (1_b \boxtimes \rho)) = (1_{b \otimes b} \boxtimes \rho) \circ \lambda_{b,b,a \boxtimes m}$.

(2) Using the pentagon axiom in \mathcal{M} and the naturality of λ , we see that the $\lambda_{b,b,a \boxtimes m}$ and $\lambda_{b,b,m}$ fit into a commutative square

$$\begin{array}{ccccc}
b \boxtimes (b \boxtimes (a \boxtimes m)) & \xrightarrow{1_b \boxtimes \lambda_{b,a,m}} & b \boxtimes ((b \otimes a) \boxtimes m) & \xrightarrow{1_b \boxtimes ((\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_m)} & b \boxtimes (b \boxtimes m) \\
\downarrow \lambda_{b,b,a \boxtimes m} & & & & \downarrow \lambda_{b,b,m} \\
(b \otimes b) \boxtimes (a \boxtimes m) & \xrightarrow{\lambda_{b \otimes b, a, m}} & ((b \otimes b) \otimes a) \boxtimes m & \xrightarrow{\sim} & (b \otimes (b \otimes a)) \boxtimes m \xrightarrow{(1_b \otimes (\mu_b \circ (1_b \otimes \alpha))) \boxtimes 1_m} (b \otimes b) \boxtimes m
\end{array}$$

(3) Using the functoriality of \boxtimes , we see that $(1_b \boxtimes \rho) \circ (\mu_b \boxtimes 1_{a \boxtimes m}) = (\mu_b \boxtimes 1_m) \circ (1_{b \otimes b} \boxtimes \rho)$.

(4) Using the naturality of λ , the pentagon axiom in \mathcal{C} , the associativity of μ_b and the functoriality of \boxtimes , we have the commutative diagram

$$\begin{array}{ccccc}
(b \otimes b) \boxtimes (a \boxtimes m) & \xrightarrow{\lambda_{b \otimes b, a, m}} & ((b \otimes b) \otimes a) \boxtimes m & \xrightarrow{\sim} & (b \otimes (b \otimes a)) \boxtimes m \xrightarrow{(1_b \otimes (\mu_b \circ (1_b \otimes \alpha))) \boxtimes 1_m} (b \otimes b) \boxtimes m \\
\downarrow \mu_b \boxtimes 1_{a \boxtimes m} & & & & \downarrow \mu_b \boxtimes 1_m \\
b \boxtimes (a \boxtimes m) & \xrightarrow{\lambda_{b,a,m}} & (b \otimes a) \boxtimes m & \xrightarrow{(\mu_b \circ (1_b \boxtimes \alpha)) \boxtimes 1_m} & b \boxtimes m
\end{array}$$

Using (1) and (2), it follows that the diagrams in the top and middle rows of 2.4 are naturally isomorphic. Hence, the coequalizer of the middle row is $(b \otimes b) \boxtimes m \xrightarrow{(1_b \boxtimes coeq) \circ \lambda_{b,b,n}^{-1}} b \boxtimes (b \boxtimes_a m)$.

Using (3) and (4), there is an induced morphism $\tilde{\rho} : b \boxtimes (b \boxtimes_a m) \longrightarrow b \boxtimes_a m$. It may be verified that $\tilde{\rho}$ makes $b \boxtimes_a m$ into a left b -module. \square

Theorem 2.4. *Let $\alpha : (a, \mu_a, \iota_a) \longrightarrow (b, \mu_b, \iota_b)$ be a morphism in $\text{Comm}(\mathcal{C})$. Then, the assignment $(m, \rho) \mapsto (b \boxtimes_a m, \tilde{\rho})$ extends to a functor $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ which is left adjoint to the restriction of scalars $\alpha_* : \mathbf{Mod}_{\mathcal{M}}(b) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(a)$.*

Proof. Let $(m, \rho) \in \mathbf{Mod}_{\mathcal{M}}(a)$, $(n, \tau) \in \mathbf{Mod}_{\mathcal{M}}(b)$. We need to show that there is a natural isomorphism

$$\mathbf{Mod}_{\mathcal{M}}(b)((b \boxtimes_a m, \tilde{\rho}), (n, \tau)) \cong \mathbf{Mod}_{\mathcal{M}}(a)((m, \rho), \alpha_*((n, \tau))) \quad (2.5)$$

Let $\phi : (b \boxtimes_a m, \tilde{\rho}) \rightarrow (n, \tau)$ be a morphism in $\mathbf{Mod}_{\mathcal{M}}(b)$. We consider $\bar{\phi} : m \rightarrow n$ given by the composite

$$m \cong 1 \boxtimes m \xrightarrow{\iota_b \boxtimes 1_m} b \boxtimes m \xrightarrow{coeq_{\alpha, (m, \rho)}} b \boxtimes_a m \xrightarrow{\phi} n \quad (2.6)$$

It is easy to see that $\bar{\phi} \in \mathbf{Mod}_{\mathcal{M}}(a)((m, \rho), \alpha_*((n, \tau)))$. We check the naturality of the collection of maps

$$\{\mathbf{Mod}_{\mathcal{M}}(b)((b \boxtimes_a m, \tilde{\rho}), (n, \tau)) \rightarrow \mathbf{Mod}_{\mathcal{M}}(a)((m, \rho), \alpha_*((n, \tau))), \quad \phi \mapsto \bar{\phi} \mid (n, \tau) \in \mathbf{Mod}_{\mathcal{M}}(b)\} \quad (2.7)$$

Let $\xi : (n, \tau) \rightarrow (n', \tau')$ be any morphism in $\mathbf{Mod}_{\mathcal{M}}(b)$. For any $\phi \in \mathbf{Mod}_{\mathcal{M}}(b)((b \boxtimes_a m, \tilde{\rho}), (n, \tau))$, we have

$$\overline{\xi \circ \phi} = \left(m \xrightarrow{\sim} 1 \boxtimes m \xrightarrow{\iota_b \boxtimes 1_m} b \boxtimes m \xrightarrow{coeq} b \boxtimes_a m \xrightarrow{\phi} n \xrightarrow{\xi} n' \right) = \xi \circ \bar{\phi}. \quad (2.8)$$

This proves the naturality of the collection (2.7).

Now let $\psi : (m, \rho) \rightarrow \alpha_*((n, \tau))$ be a morphism in $\mathbf{Mod}_{\mathcal{M}}(a)$. We note that $\tau \circ (\alpha \boxtimes \psi) = \psi \circ \rho$. Hence,

$$\begin{aligned} \tau \circ (1_b \boxtimes \psi) \circ (1_b \boxtimes \rho) &= \tau \circ (1_b \boxtimes (\psi \circ \rho)) \\ &= \tau \circ (1_b \boxtimes (\tau \circ (\alpha \boxtimes \psi))) \\ &= \tau \circ (1_b \boxtimes \tau) \circ (1_b \boxtimes (\alpha \boxtimes \psi)) \\ &= \tau \circ (\mu_b \boxtimes 1_n) \circ \lambda_{b, b, n} \circ (1_b \boxtimes (\alpha \boxtimes \psi)) \quad [\text{the associativity of } \tau] \\ &= \tau \circ (\mu_b \boxtimes 1_n) \circ ((1_b \otimes \alpha) \boxtimes \psi) \circ \lambda_{b, a, m} \quad [\text{the naturality of } \lambda] \\ &= \tau \circ ((\mu_b \circ (1_b \otimes \alpha)) \boxtimes \psi) \circ \lambda_{b, a, m} \\ &= \tau \circ (1_b \boxtimes \psi) \circ ((\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_m) \circ \lambda_{b, a, m} \end{aligned} \quad (2.9)$$

By the universal property of the coequalizer (2.3), there exists a unique morphism $\underline{\psi} : b \boxtimes_a m \rightarrow n$ such that the triangle

$$\begin{array}{ccc} b \boxtimes m & \xrightarrow{coeq_{\alpha, (m, \rho)}} & b \boxtimes_a m \\ & \searrow \tau \circ (1_b \boxtimes \psi) & \swarrow \underline{\psi} \\ & n & \end{array} \quad (2.10)$$

commutes. To show that $\underline{\psi}$ is a morphism in $\mathbf{Mod}_{\mathcal{M}}(b)$, we consider the diagram

$$\begin{array}{ccccc} (b \otimes b) \boxtimes m & \xrightarrow{(1_b \boxtimes coeq_{\alpha, (m, \rho)}) \circ \lambda_{b, b, m}^{-1}} & b \boxtimes (b \boxtimes_a m) & \xrightarrow{1_b \boxtimes \underline{\psi}} & b \boxtimes n \\ \mu_b \boxtimes 1_m \downarrow & & \tilde{\rho} \downarrow & & \downarrow \tau \\ b \boxtimes m & \xrightarrow{coeq_{\alpha, (m, \rho)}} & b \boxtimes_a m & \xrightarrow{\underline{\psi}} & n \end{array} \quad (2.11)$$

It follows from the diagram (2.4) that the left hand square in (2.11) commutes. Using (2.10) and the naturality of λ , it may be verified that $\tau \circ (1_b \boxtimes \underline{\psi}) \circ (1_b \boxtimes coeq_{\alpha, (m, \rho)}) \circ \lambda_{b, b, m}^{-1} = \underline{\psi} \circ coeq_{\alpha, (m, \rho)} \circ (\mu_b \boxtimes 1_m)$, which shows that the outer square in (2.11) commutes. We now note that the morphism

$$(1_b \boxtimes coeq_{\alpha, (m, \rho)}) \circ \lambda_{b, b, m}^{-1} : (b \otimes b) \boxtimes m \rightarrow b \boxtimes (b \boxtimes_a m) \quad (2.12)$$

is a coequalizer and hence an epimorphism. It now follows from the diagram (2.11) that $\tau \circ (1_b \boxtimes \underline{\psi}) = \underline{\psi} \circ \tilde{\rho}$. This shows that $\underline{\psi} : (b \boxtimes_a m, \tilde{\rho}) \rightarrow (n, \tau)$ is a morphism in $\mathbf{Mod}_{\mathcal{M}}(b)$.

It is straightforward to check that the two assignments $\phi \mapsto \bar{\phi}$ and $\psi \mapsto \underline{\psi}$ are mutual inverses. It follows by [7, Proposition 6.7.2] that there is a unique functor $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ which is left adjoint to α_* and such that $\alpha^*((m, \rho)) = (b \boxtimes_a m, \tilde{\rho})$ for every $(m, \rho) \in \mathbf{Mod}_{\mathcal{M}}(a)$. This completes the proof. \square

It may be verified that for a morphism $\phi : (m, \rho) \longrightarrow (m', \rho')$ in $\mathbf{Mod}_{\mathcal{M}}(a)$, the induced morphism $1_b \boxtimes_a \phi := \alpha^*(\phi)$ satisfies $\text{coeq}_{\alpha, (m', \rho')} \circ (1_b \boxtimes_a \phi) = (1_b \boxtimes_a \phi) \circ \text{coeq}_{\alpha, (m, \rho)}$. We shall denote the unit and counit of the adjunction (α^*, α_*) by η_α and ε_α respectively.

Let the following be a pushout square in $\text{Comm}(\mathcal{C})$.

$$\begin{array}{ccc} b' & \xleftarrow{\alpha'} & b \\ \beta' \uparrow & & \uparrow \beta \\ a' & \xleftarrow{\alpha} & a \end{array} \quad (2.13)$$

We recall (see for instance, [32, § 2]) that the pushout b' is given by $b \otimes_a a'$.

Lemma 2.5. *Let $\alpha : a \longrightarrow a', \beta : a \longrightarrow b$ and $\gamma : c \longrightarrow b$ be morphisms in $\text{Comm}(\mathcal{C})$. Then,*

- (1) $(a' \otimes_a b) \boxtimes m \cong a' \boxtimes_a (b \boxtimes m)$ as left a' -modules, and this is natural in $m \in \mathcal{M}$.
- (2) $(a' \otimes_a b) \boxtimes_c m \cong a' \boxtimes_a (b \boxtimes_c m)$ as left a' -modules, and this is natural in $(m, \rho) \in \mathbf{Mod}_{\mathcal{M}}(c)$.

Proof. (1) Since $-- \boxtimes m : \mathcal{C} \longrightarrow \mathcal{M}$ preserves colimits, we have

$$(a' \otimes_a b) \boxtimes m \cong \text{Coeq} \left(a' \boxtimes (a \boxtimes (b \boxtimes m)) \cong (a' \otimes_a a \otimes b) \boxtimes m \xrightarrow[\begin{smallmatrix} ((\mu_{a'} \circ (1_{a'} \otimes \alpha)) \otimes 1_b) \boxtimes 1_m \end{smallmatrix}]{\begin{smallmatrix} (1_{a'} \otimes (\mu_b \circ (\beta \otimes 1_b))) \boxtimes 1_m \end{smallmatrix}} (a' \otimes b) \boxtimes m \cong a' \boxtimes (b \boxtimes m) \right) \cong a' \boxtimes_a (b \boxtimes m) \quad (2.14)$$

in \mathcal{M} . Using Remark 2.2, it may be verified that (2.14) holds in $\mathbf{Mod}_{\mathcal{M}}(a')$.

(2) We consider the diagram

$$\begin{array}{ccc} b \boxtimes (c \boxtimes m) & \xrightarrow[((\mu_b \circ (1_b \otimes \gamma)) \boxtimes 1_m) \circ \lambda_{b, c, m}]{1_b \boxtimes \rho} & b \boxtimes m \end{array} \quad (2.15)$$

in $\mathbf{Mod}_{\mathcal{M}}(b)$. Applying β_* to (2.15), we obtain a diagram in $\mathbf{Mod}_{\mathcal{M}}(a)$. Using Remark 2.2, it follows that the coequalizer of this diagram in $\mathbf{Mod}_{\mathcal{M}}(a)$ is $b \boxtimes m \xrightarrow{\text{coeq}_{\gamma, (m, \rho)}} b \boxtimes_c m$. Since $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(a')$ is a left adjoint, it preserves coequalizers. Hence,

$$\begin{array}{ccc} a' \boxtimes_a (b \boxtimes (c \boxtimes m)) & \xrightarrow[1_{a'} \boxtimes (((\mu_b \circ (1_b \otimes \gamma)) \boxtimes 1_m) \circ \lambda_{b, c, m})]{1_{a'} \boxtimes_a (1_b \boxtimes \rho)} & a' \boxtimes_a (b \boxtimes m) \xrightarrow{1_{a'} \boxtimes \text{coeq}_{\gamma, (m, \rho)}} a' \boxtimes_a (b \boxtimes_c m) \end{array} \quad (2.16)$$

is a coequalizer in $\mathbf{Mod}_{\mathcal{M}}(a')$. Using part (1), it follows that $(a' \otimes_a b) \boxtimes_c m \cong a' \boxtimes_a (b \boxtimes_c m)$ as left a' -modules. \square

We denote by Cat the 2-category of categories (see, for instance, [6, § 7]). The following result will be used in later sections.

Proposition 2.6. (1) *The assignment*

$$\begin{array}{ccc} \mathbf{Mod}_{\mathcal{M}} : \text{Comm}(\mathcal{C}) & \longrightarrow & \text{Cat} \\ (\text{Comm}(\mathcal{C}) \ni a \mapsto \mathbf{Mod}_{\mathcal{M}}(a)) & & ((\alpha : a \longrightarrow b) \mapsto (\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(b))) \end{array} \quad (2.17)$$

is a pseudo-functor.

(2) *For every $\alpha : a \longrightarrow b$ in $\text{Comm}(\mathcal{C})$, the restriction of scalars $\alpha_* : \mathbf{Mod}_{\mathcal{M}}(b) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(a)$ is conservative, i.e., a morphism ϕ in $\mathbf{Mod}_{\mathcal{M}}(b)$ is an isomorphism if and only if $\alpha_*(\phi)$ is an isomorphism in $\mathbf{Mod}_{\mathcal{M}}(a)$.*

(3) *For any pushout square (2.13) in $\text{Comm}(\mathcal{C})$, there is a natural isomorphism $\alpha^* \circ \beta_* \xrightarrow{\sim} \beta'_* \circ \alpha'^*$.*

Proof. (1) It is easy to see that for any $a \in \text{Comm}(\mathcal{C})$, $(1_a)^* \cong 1_{\mathbf{Mod}_{\mathcal{M}}(a)}$. Now, let $a \xrightarrow{\alpha} b \xrightarrow{\beta} c$ be morphisms in $\text{Comm}(\mathcal{C})$. We consider $\beta : b \rightarrow c$, $1_b : b \rightarrow b$ and $\alpha : a \rightarrow b$. Using Lemma 2.5, it follows that

$$(\beta \circ \alpha)^* \cong c \boxtimes_a - \cong (c \otimes_b b) \boxtimes_a - \cong c \boxtimes_b (b \boxtimes_a -) \cong \beta^* \circ \alpha^* \quad (2.18)$$

It can be checked that $\mathbf{Mod}_{\mathcal{M}}$ satisfies the coherence axioms for a pseudo-functor.

(2) This follows from the fact that for every $c \in \mathcal{C}$, the forgetful functor $\mathbf{Mod}_{\mathcal{M}}(c) \rightarrow \mathcal{M}$ reflects isomorphisms.

(3) Applying Lemma 2.5 to the morphisms $\alpha : a \rightarrow a'$, $\beta : a \rightarrow b$, $1_b : b \rightarrow b$, it follows that

$$\beta'_* \circ \alpha'^* = b' \boxtimes_b - = (a' \otimes_a b) \boxtimes_b - \cong a' \boxtimes_a (b \boxtimes_b -) \cong a' \boxtimes_a - = \alpha^* \circ \beta_* \quad (2.19)$$

□

3 The topology on $\text{Aff}_{\mathcal{C}}$

We set $\text{Aff}_{\mathcal{C}} := \text{Comm}(\mathcal{C})^{op}$. For any object $a \in \text{Comm}(\mathcal{C})$, the corresponding object in $\text{Aff}_{\mathcal{C}}$ will be denoted by $\text{Spec}(a)$. For any morphism $\alpha : a \rightarrow b$ in $\text{Comm}(\mathcal{C})$, the corresponding morphism in $\text{Aff}_{\mathcal{C}}$ will be denoted by $\alpha^{op} : \text{Spec}(b) \rightarrow \text{Spec}(a)$. In this section, we will introduce the topologies on $\text{Aff}_{\mathcal{C}}$ relative to \mathcal{M} .

Definition 3.1. A morphism $\alpha : a \rightarrow b$ in $\text{Comm}(\mathcal{C})$ is \mathcal{M} -flat if the extension of scalars $\alpha^* : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ preserves finite limits.

Lemma 3.2. \mathcal{M} -flatness is stable under pushouts and closed under compositions in $\text{Comm}(\mathcal{C})$.

Proof. Using Proposition 2.6, it is straightforward to check that the composition of \mathcal{M} -flat morphisms is \mathcal{M} -flat. We now consider the following pushout square in $\text{Comm}(\mathcal{C})$ such that α is \mathcal{M} -flat.

$$\begin{array}{ccc} b' = b \otimes_a a' & \xleftarrow{\alpha'} & b \\ \beta'_* \uparrow & & \uparrow \beta \\ a' & \xleftarrow{\alpha} & a \end{array} \quad (3.1)$$

We need to show that α' is \mathcal{M} -flat, i.e., $\alpha'^* : \mathbf{Mod}_{\mathcal{M}}(b) \rightarrow \mathbf{Mod}_{\mathcal{M}}(b \otimes_a a')$ preserves finite limits. Let $\mathbb{D} : \mathcal{I} \rightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ be a diagram, where \mathcal{I} is a finite category. There is a canonical morphism

$$\alpha'^* \left(\lim_{\mathcal{I}} \mathbb{D} \right) \rightarrow \lim_{\mathcal{I}} (\alpha'^* \circ \mathbb{D}) \quad (3.2)$$

in $\mathbf{Mod}_{\mathcal{M}}(b \otimes_a a')$. Applying β'_* to (3.2), using Proposition 2.6 and the fact that β_* and β'_* preserve limits, we have a morphism

$$\alpha^* \left(\lim_{\mathcal{I}} (\beta_* \circ \mathbb{D}) \right) \cong \alpha^* \left(\beta_* \left(\lim_{\mathcal{I}} \mathbb{D} \right) \right) \cong \beta'_* \left(\alpha'^* \left(\lim_{\mathcal{I}} \mathbb{D} \right) \right) \rightarrow \beta'_* \left(\lim_{\mathcal{I}} (\alpha'^* \circ \mathbb{D}) \right) \cong \lim_{\mathcal{I}} (\beta'_* \circ \alpha'^* \circ \mathbb{D}) \cong \lim_{\mathcal{I}} (\alpha^* \circ (\beta_* \circ \mathbb{D})) \quad (3.3)$$

Since α is \mathcal{M} -flat, the morphism in (3.3) must be an isomorphism. Further since β'_* is conservative, it follows that the morphism in (3.2) is an isomorphism. Hence, α' is \mathcal{M} -flat. □

For an object $a \in \text{Comm}(\mathcal{C})$, we consider the coslice category $a/\text{Comm}(\mathcal{C})$, or the category of “ a -algebras.” We recall that objects of $a/\text{Comm}(\mathcal{C})$ are the morphisms in $\text{Comm}(\mathcal{C})$ with domain a . Let $\alpha : a \rightarrow b$ be a morphism in $\text{Comm}(\mathcal{C})$. We also recall (see [32, § 2.3]) that α is of finite type if for any filtered diagram $\mathbb{D} : \mathcal{I} \rightarrow a/\text{Comm}(\mathcal{C})$, the canonical map

$$\text{colim}_{i \in \mathcal{I}} a/\text{Comm}(\mathcal{C})(b, \mathbb{D}(i)) \rightarrow a/\text{Comm}(\mathcal{C})(b, \text{colim}_{i \in \mathcal{I}} \mathbb{D}(i)) \quad (3.4)$$

is an isomorphism.

Definition 3.3. A family of morphisms $\{\alpha_i^{op} : \text{Spec}(a_i) \rightarrow \text{Spec}(a)\}_{i \in I}$ in $\text{Aff}_{\mathcal{C}}$ is an fpqc \mathcal{M} -cover for $\text{Spec}(a)$ if

- (1) For each $i \in I$, the morphism $\alpha_i : a \rightarrow a_i$ is \mathcal{M} -flat.
- (2) There exists a finite subset $J \subseteq I$ such that the collection of functors

$$\{\alpha_j^* = (a_j \boxtimes_a -) : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(a_j)\}_{j \in J} \quad (3.5)$$

is conservative, i.e., a morphism ϕ in $\mathbf{Mod}_{\mathcal{M}}(a)$ is an isomorphism if and only if $\alpha_j^*(\phi)$ is an isomorphism in $\mathbf{Mod}_{\mathcal{M}}(a_j)$ for each $j \in J$.

Definition 3.4. A morphism $\alpha^{op} : \text{Spec}(b) \rightarrow \text{Spec}(a)$ in $\text{Aff}_{\mathcal{C}}$ is a spectral immersion relative to \mathcal{M} if α is \mathcal{M} -flat and an epimorphism of finite type in $\text{Comm}(\mathcal{C})$.

Lemma 3.5. Spectral immersions relative to \mathcal{M} are stable under pullbacks and closed under compositions in $\text{Aff}_{\mathcal{C}}$.

Proof. We need to show that \mathcal{M} -flat epimorphisms of finite type are stable under pushouts and compositions in $\text{Comm}(\mathcal{C})$. We consider a pushout square in $\text{Comm}(\mathcal{C})$ such that $\beta : a \rightarrow b$ is of finite type.

$$\begin{array}{ccc} b' = b \otimes_a a' & \xleftarrow{\alpha'} & b \\ \beta' \uparrow & & \uparrow \beta \\ a' & \xleftarrow{\alpha} & a \end{array} \quad (3.6)$$

We claim that $\beta' : a' \rightarrow b' = b \otimes_a c$ is of finite type. Let $\mathbb{D} : \mathcal{I} \rightarrow a'/\text{Comm}(\mathcal{C})$ be a filtered diagram. We consider the functor $r_\alpha : a'/\text{Comm}(\mathcal{C}) \rightarrow a/\text{Comm}(\mathcal{C})$ that takes $(a' \rightarrow c) \mapsto (a \xrightarrow{\alpha} a' \rightarrow c)$. We note that the forgetful functors $\pi_a : a/\text{Comm}(\mathcal{C}) \rightarrow \text{Comm}(\mathcal{C})$ and $\pi_{a'} : a'/\text{Comm}(\mathcal{C}) \rightarrow \text{Comm}(\mathcal{C})$ create filtered colimits and are conservative. Since $\pi_a \circ r_\alpha = \pi_{a'}$, it follows that r_α preserves filtered colimits. Further, r_α has a left adjoint

$$\begin{aligned} p_\alpha : a/\text{Comm}(\mathcal{C}) &\rightarrow a'/\text{Comm}(\mathcal{C}) \\ (a \rightarrow c) &\mapsto (a' \rightarrow a' \otimes_a c) \end{aligned} \quad (3.7)$$

Since β is of finite type and r_α preserves filtered colimits, we have

$$\begin{aligned} \text{colim}_{i \in \mathcal{I}} a'/\text{Comm}(\mathcal{C})(a' \otimes_a b, \mathbb{D}(i)) &\cong \text{colim}_{i \in \mathcal{I}} a/\text{Comm}(\mathcal{C})(b, r_\alpha(\mathbb{D}(i))) \\ &\cong a/\text{Comm}(\mathcal{C})(b, \text{colim}_{i \in \mathcal{I}} r_\alpha(\mathbb{D}(i))) \\ &\cong a/\text{Comm}(\mathcal{C})(b, r_\alpha(\text{colim}_{i \in \mathcal{I}} \mathbb{D}(i))) \cong a'/\text{Comm}(\mathcal{C})(a' \otimes_a b, \text{colim}_{i \in \mathcal{I}} \mathbb{D}(i)) \end{aligned} \quad (3.8)$$

This shows that $\beta' : a' \rightarrow a' \otimes_a b$ is of finite type. It may be verified that a composition of morphisms of finite type is a morphism of finite type. Using Lemma 3.2, it follows that \mathcal{M} -flat epimorphisms of finite type are stable under pushouts and closed under compositions in $\text{Comm}(\mathcal{C})$. \square

Definition 3.6. A family of morphisms $\{\alpha_i^{op} : \text{Spec}(a_i) \rightarrow \text{Spec}(a)\}_{i \in I}$ in $\text{Aff}_{\mathcal{C}}$ is a spectral \mathcal{M} -cover for $\text{Spec}(a)$ if it is an fpqc \mathcal{M} -cover and for each $i \in I$, α_i^{op} is a spectral immersion relative to \mathcal{M} .

Proposition 3.7. (1) The assignment $\text{Aff}_{\mathcal{C}} \ni \text{Spec}(a) \mapsto \{\text{fpqc } \mathcal{M}\text{-covers for } \text{Spec}(a)\}$ is a basis for a Grothendieck topology on $\text{Aff}_{\mathcal{C}}$.

(2) The assignment $\text{Aff}_{\mathcal{C}} \ni \text{Spec}(a) \mapsto \{\text{spectral } \mathcal{M}\text{-covers for } \text{Spec}(a)\}$ is a basis for a Grothendieck topology on $\text{Aff}_{\mathcal{C}}$.

Proof. (1) If $\alpha^{op} : Spec(b) \rightarrow Spec(a)$ is an isomorphism, it is clear that the singleton $\{\alpha^{op}\}$ is an fpqc \mathcal{M} -cover. Let $\{\alpha_i^{op} : Spec(a_i) \rightarrow Spec(a)\}_{i \in I}$ be an fpqc \mathcal{M} -cover and $\beta^{op} : Spec(b) \rightarrow Spec(a)$ be a morphism in $Aff_{\mathcal{C}}$. We consider the pushout square

$$\begin{array}{ccc} a_i \otimes_a b & \xleftarrow{\alpha'_i} & b \\ \beta'_i \uparrow & & \uparrow \beta \\ a_i & \xleftarrow{\alpha_i} & a \end{array} \quad (3.9)$$

in $Comm(\mathcal{C})$ for each $i \in I$. We claim that the family of pullbacks $\{(\alpha'_i)^{op} : Spec(a_i) \times_{Spec(a)} Spec(b) \cong Spec(a_i \otimes_a b) \rightarrow Spec(b)\}_{i \in I}$ is an fpqc \mathcal{M} -cover for $Spec(b)$. Using Lemma 3.2, it follows that each $\alpha'_i : b \rightarrow a_i \otimes_a b$ is \mathcal{M} -flat. Further, there exists a finite subset $J \subseteq I$ such that the following collection of functors is conservative.

$$\{\alpha_j^* = (a_j \boxtimes_a -) : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(a_j)\}_{j \in J} \quad (3.10)$$

We claim that the functor $(\alpha_j^*)_{j \in J} = ((a_j \otimes_a b) \boxtimes_b -)_{j \in J} : \mathbf{Mod}_{\mathcal{M}}(b) \rightarrow \prod_{j \in J} \mathbf{Mod}_{\mathcal{M}}(a_j \otimes_a b)$ is conservative. It suffices to show that the composite functor $(\beta_{j*}' \circ \alpha_j'^*)_{j \in J}$

$$\mathbf{Mod}_{\mathcal{M}}(b) \xrightarrow{(\alpha_j'^*)_{j \in J}} \prod_{j \in J} \mathbf{Mod}_{\mathcal{M}}(a_j \otimes_a b) \xrightarrow{\prod_{j \in J} \beta_{j*}'} \prod_{j \in J} \mathbf{Mod}_{\mathcal{M}}(a_j) \quad (3.11)$$

is conservative. Using Proposition 2.6, we have $(\beta_{j*}' \circ \alpha_j'^*)_{j \in J} \cong (\alpha_j^* \circ \beta_*)_{j \in J} = (\alpha_j^*)_{j \in J} \circ \beta_*$. Since β_* is conservative, it now follows that the functor in (3.11) is conservative.

Using Proposition 2.6 and Lemma 3.2, it may be verified that composition of fpqc \mathcal{M} -covers must be an fpqc \mathcal{M} -cover. This completes the proof.

(2) The result follows from part (1) and Lemma 3.5. \square

We will refer to the topology on $Aff_{\mathcal{C}}$ coming from part (1) of Propostion 3.7 as the fpqc \mathcal{M} -topology and denote it by $fpqc_{\mathcal{M}}$. The associated topology on $Aff_{\mathcal{C}}$ in part (2) of Propostion 3.7 will be called the spectral \mathcal{M} -topology and denoted by $spc_{\mathcal{M}}$.

We denote the category of presheaves of sets on $Aff_{\mathcal{C}}$ by $PSh(Aff_{\mathcal{C}})$ and let $Sh(Aff_{\mathcal{C}}, fpqc_{\mathcal{M}})$ be the full subcategory of $PSh(Aff_{\mathcal{C}})$ whose objects are the sheaves with respect to the fpqc \mathcal{M} -topology. The category of sheaves on $Aff_{\mathcal{C}}$ with respect to the spectral \mathcal{M} -topology will be denoted by $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. We note that we have full subcategories

$$Sh(Aff_{\mathcal{C}}, fpqc_{\mathcal{M}}) \subseteq Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \subseteq PSh(Aff_{\mathcal{C}}) \quad (3.12)$$

We will often use the term “affine scheme” to mean either an object $Spec(a) \in Aff_{\mathcal{C}}$ or the corresponding presheaf on $Aff_{\mathcal{C}}$. Similarly, we will use $\alpha^{op} : Spec(a) \rightarrow Spec(a')$ to denote both a morphism in $Aff_{\mathcal{C}}$ and the corresponding morphism in $PSh(Aff_{\mathcal{C}})$.

4 Schemes relative to \mathcal{M}

From now onwards, we will say that the left \mathcal{C} -module category (\mathcal{M}, \boxtimes) is subcanonical if the fpqc \mathcal{M} -topology (and hence, the spectral \mathcal{M} -topology) on $Aff_{\mathcal{C}}$ is subcanonical, i.e., for each $Spec(a) \in Aff_{\mathcal{C}}$, the representable presheaf $Aff_{\mathcal{C}}(_, Spec(a))$ on $Aff_{\mathcal{C}}$ is a sheaf for the fpqc \mathcal{M} -topology on $Aff_{\mathcal{C}}$. In this section, we present the main definition of this paper, that of a scheme relative to \mathcal{M} . We assume throughout this section that the left \mathcal{C} -module category \mathcal{M} is subcanonical. We begin with the following lemma.

Lemma 4.1. *For every spectral \mathcal{M} -cover $\{\alpha_i^{op} : Spec(a_i) \rightarrow Spec(a)\}_{i \in I}$, the canonical morphism*

$$\prod_{i \in I} Aff_{\mathcal{C}}(_, Spec(a_i)) \xrightarrow{\prod_{i \in I} Aff_{\mathcal{C}}(_, \alpha_i^{op}) = \prod_{i \in I} \alpha_i^{op}} Aff_{\mathcal{C}}(_, Spec(a)) \quad (4.1)$$

is an epimorphism in $Sh(Aff_C)_\mathcal{M}$.

Proof. Let p, q be morphisms in $Sh(Aff_C)_\mathcal{M}$ such that the compositions

$$\begin{aligned} \coprod_{i \in I} Aff_C(_, Spec(a_i)) &\xrightarrow{\coprod_{i \in I} \alpha_i^{op}} Aff_C(_, Spec(a)) \xrightarrow{p} F \\ \coprod_{i \in I} Aff_C(_, Spec(a_i)) &\xrightarrow{\coprod_{i \in I} \alpha_i^{op}} Aff_C(_, Spec(a)) \xrightarrow{q} F \end{aligned} \quad (4.2)$$

are equal. Then, (4.2) implies that for each $i \in I$, we have $p \circ \alpha_i^{op} = q \circ \alpha_i^{op}$ in $Sh(Aff_C)_\mathcal{M}$. Let p' (resp. q') be the element of $F(Spec(a))$ corresponding under Yoneda lemma to p (resp. q). It follows that for each $i \in I$, $p \circ \alpha_i^{op} = q \circ \alpha_i^{op}$ corresponds under Yoneda Lemma to the element $F(\alpha_i^{op})(p') = F(\alpha_i^{op})(q') \in F(Spec(a_i))$. Since $F \in Sh(Aff_C)_\mathcal{M}$ and $\{\alpha_i^{op} : Spec(a_i) \rightarrow Spec(a)\}_{i \in I}$ is a spectral \mathcal{M} -cover, we see that the map

$$F(Spec(a)) \rightarrow \prod_{i \in I} F(Spec(a_i)) \quad x \mapsto (F(\alpha_i^{op})(x))_{i \in I} \quad (4.3)$$

must be injective. It follows that $p' = q'$ and hence $p = q$. \square

Definition 4.2. (1) Let $X := Aff_C(_, Spec(a))$ be an affine scheme and $F \subseteq X$ a subsheaf in $Sh(Aff_C)_\mathcal{M}$. We say that $F \subseteq X$ is a Zariski open relative to \mathcal{M} if there is a family $\{Spec(a_i) \rightarrow Spec(a)\}_{i \in I}$ of spectral immersions relative to \mathcal{M} such that the induced morphism $\coprod_{i \in I} Aff_C(_, Spec(a_i)) \rightarrow Aff_C(_, Spec(a)) = X$ has image F , i.e., we have

$$\coprod_{i \in I} Aff_C(_, Spec(a_i)) \xrightarrow{epi} F \hookrightarrow Aff_C(_, Spec(a)) = X \quad (4.4)$$

in $Sh(Aff_C)_\mathcal{M}$.

(2) A morphism $F \rightarrow G$ in $Sh(Aff_C)_\mathcal{M}$ is a Zariski open immersion relative to \mathcal{M} if for each $X \in Aff_C$ and each morphism $X \rightarrow G$, the induced morphism $F \times_G X \rightarrow X$ in $Sh(Aff_C)_\mathcal{M}$ is a monomorphism whose image is Zariski open relative to \mathcal{M} in X .

Lemma 4.3. (1) Let $G \rightarrow H$ be a morphism in $Sh(Aff_C)_\mathcal{M}$. Then, the functor $_ \times_H G : Sh(Aff_C)_\mathcal{M}/H \rightarrow Sh(Aff_C)_\mathcal{M}/G$ sending $(F \rightarrow H) \mapsto (F \times_H G \rightarrow G)$ preserves colimits.

(2) Let $\coprod_{i \in I} F_i \rightarrow H$ be an epimorphism in $Sh(Aff_C)_\mathcal{M}$. If $G \rightarrow H$ is a morphism in $Sh(Aff_C)_\mathcal{M}$, then the induced morphism $\coprod_{i \in I} F_i \times_H G \rightarrow G$ is an epimorphism in $Sh(Aff_C)_\mathcal{M}$.

Proof. (1) It is easy to verify that the analogous result holds in the category of sets. Since colimits in $PSh(Aff_C)$ are computed objectwise, we see that the functor $_ \times_H G : PSh(Aff_C)/H \rightarrow PSh(Aff_C)/G$ sending $(F \rightarrow H) \mapsto (F \times_H G \rightarrow G)$ preserves colimits. Let $(\cdot)_\mathcal{M}^{++} : PSh(Aff_C) \rightarrow Sh(Aff_C)_\mathcal{M}$ be the sheafification functor which is left adjoint to the inclusion $Sh(Aff_C)_\mathcal{M} \hookrightarrow PSh(Aff_C)$. In particular, the functor $(\cdot)_\mathcal{M}^{++}$ preserves colimits. Since sheafification is obtained from filtered colimits (see for instance, [22, § 3.5]), we know that it also preserves finite limits. The result is now clear.

(2) It follows from part (1) that $\coprod_i F_i \times_H G \cong \coprod_i (F_i \times_H G)$ in $Sh(Aff_C)_\mathcal{M}/G$. Since $\coprod_i F_i \rightarrow H$ is an epimorphism in $Sh(Aff_C)_\mathcal{M}$, it induces an epimorphism $(\coprod_i F_i \rightarrow H) \rightarrow (H \xrightarrow{id} H)$ in $Sh(Aff_C)_\mathcal{M}/H$. Using part (1), it follows that the morphism $(\coprod_i F_i \times_H G \cong \coprod_i (F_i \times_H G) \rightarrow G) \rightarrow (G \xrightarrow{id} G)$ is an epimorphism in $Sh(Aff_C)_\mathcal{M}/G$ and hence $\coprod_{i \in I} F_i \times_H G \rightarrow G$ is an epimorphism in $Sh(Aff_C)_\mathcal{M}$. This completes the proof. \square

Lemma 4.4. A Zariski open immersion $f : F \rightarrow G$ relative to \mathcal{M} is a monomorphism in $Sh(Aff_C)_\mathcal{M}$.

Proof. Let $p : H \rightarrow F, q : H \rightarrow F$ be morphisms in $Sh(Aff_C)_\mathcal{M}$ such that $f \circ p = f \circ q$. Suppose that $p \neq q$. Then, there exists $Spec(a) \in Aff_C$ with $p_{Spec(a)} \neq q_{Spec(a)}$. Hence, there must be an element $x \in H(Spec(a))$ with $p_{Spec(a)}(x) \neq q_{Spec(a)}(x)$. We set $Y = Aff_C(_, Spec(a))$. Let $h : Y \rightarrow H$ (resp. $g : Y \rightarrow G$) be the morphism

corresponding under Yoneda lemma to the element $x \in H(\text{Spec}(a))$ (resp. $f_{\text{Spec}(a)}(p_{\text{Spec}(a)}(x)) \in G(\text{Spec}(a))$). It follows that

$$f \circ p \circ h = g \quad (4.5)$$

Further, since $p_{\text{Spec}(a)}(x) \neq q_{\text{Spec}(a)}(x)$, we have

$$p \circ h \neq q \circ h \quad (4.6)$$

We consider the pullback squares in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccc} F \times_G Y & \xrightarrow{f'} & Y \\ \downarrow g' & & \downarrow g \\ F & \xrightarrow{f} & G \end{array} \quad \begin{array}{ccc} H \times_G Y & \xrightarrow{t} & Y \\ \downarrow g'' & & \downarrow g \\ H & \xrightarrow{f \circ p = f \circ q} & G \end{array} \quad (4.7)$$

Using the universal property of $F \times_G Y$, the pair $(p \circ g'' : H \times_G Y \rightarrow F, t : H \times_G Y \rightarrow Y)$ (resp. $(q \circ g'' : H \times_G Y \rightarrow F, t : H \times_G Y \rightarrow Y)$) induces a unique morphism

$$p' : H \times_G Y \rightarrow F \times_G Y, \quad (\text{resp. } q' : H \times_G Y \rightarrow F \times_G Y) \quad (4.8)$$

such that $p \circ g'' = g' \circ p'$ and $f' \circ p' = t$ (resp. $q \circ g'' = g' \circ q'$ and $f' \circ q' = t$).

By Definition 4.2(2), since f is a Zariski open immersion relative to \mathcal{M} , we see that f' is a monomorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Hence, the equality $f' \circ p' = t = f' \circ q'$ implies that $p' = q'$. It follows that

$$p \circ g'' = g' \circ p' = g' \circ q' = q \circ g'' \quad (4.9)$$

Using (4.5) and the universal property of $H \times_G Y$, the pair $(h : Y \rightarrow H, 1_Y : Y \rightarrow Y)$ induces a unique morphism $u : Y \rightarrow H \times_G Y$ such that $t \circ u = 1_Y$ and $g'' \circ u = h$. It follows from (4.9) that

$$p \circ h = p \circ g'' \circ u = q \circ g'' \circ u = q \circ h \quad (4.10)$$

which contradicts (4.6). Hence $p = q$. This proves the result. \square

Lemma 4.5. *In $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$, Zariski open immersions relative to \mathcal{M} are stable under base change and closed under composition.*

Proof. It is easy to see that Zariski open immersions relative to \mathcal{M} are stable under base change.

Suppose that $f : F \rightarrow G$ and $g : G \rightarrow H$ are Zariski open immersions relative to \mathcal{M} . We need to show that $g \circ f$ is a Zariski open immersion relative to \mathcal{M} . Let $h : X \rightarrow H$ be a morphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ where $X \in Aff_{\mathcal{C}}$. We consider the following pullback diagram

$$\begin{array}{ccccc} F \times_H X \cong F \times_G (G \times_H X) & \xrightarrow{f'} & G \times_H X & \xrightarrow{g'} & X \\ \downarrow h'' & & \downarrow h' & & \downarrow h \\ F & \xrightarrow{f} & G & \xrightarrow{g} & H \end{array} \quad (4.11)$$

It follows from Lemma 4.4 that both f and g are monomorphisms. Hence, $g' \circ f' : F \times_H X \cong F \times_G (G \times_H X) \rightarrow G \times_H X \rightarrow X$ is a monomorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. It remains to show that $F \times_H X \subseteq X$ is Zariski open relative to \mathcal{M} .

Since $g : G \rightarrow H$ is a Zariski open immersion relative to \mathcal{M} , we see that $g' : G \times_H X \rightarrow X$ is a monomorphism whose image is Zariski open relative to \mathcal{M} . Hence, there exists a family $\{\alpha_i : X_i \rightarrow X\}_{i \in I}$ of spectral immersions

relative to \mathcal{M} such that $\coprod_i \alpha_i : \coprod_i X_i \rightarrow X$ has image $G \times_H X$ in $Sh(Aff_C)_{\mathcal{M}}$. For each $i' \in I$, we consider the composition $X_{i'} \rightarrow \coprod_{i \in I} X_i \rightarrow G \times_H X$ and form the following pullback diagram in $Sh(Aff_C)_{\mathcal{M}}$

$$\begin{array}{ccccc} F \times_G X_{i'} \cong (F \times_H X) \times_{G \times_H X} X_{i'} & \xrightarrow{\quad} & F \times_H X & \xrightarrow{h''} & F \\ \downarrow & & \downarrow f' & & \downarrow f \\ X_{i'} & \xrightarrow{\quad} & G \times_H X & \xrightarrow{h'} & G \end{array} \quad (4.12)$$

Since Zariski open immersions relative to \mathcal{M} are stable under pullbacks, the morphism $f' : F \times_H X \rightarrow G \times_H X$ is a Zariski open immersion relative to \mathcal{M} . It follows that the morphism

$$F \times_G X_{i'} \cong (F \times_G (G \times_H X)) \times_{G \times_H X} X_{i'} \rightarrow X_{i'} \quad (4.13)$$

in (4.12) is a monomorphism whose image is Zariski open relative to \mathcal{M} . Hence, there is a family $\{Y_{i'j} \rightarrow X_{i'}\}_{j \in J_{i'}}$ of spectral immersions relative to \mathcal{M} such that the image of the canonical morphism $\coprod_{j \in J_{i'}} Y_{i'j} \rightarrow X_{i'}$ in $Sh(Aff_C)_{\mathcal{M}}$ is $F \times_G X_{i'}$. Using Lemma 3.5 and Lemma 4.3, it may be verified that

$$\{Y_{i'j} \rightarrow X_{i'} \rightarrow X\}_{j \in J_{i'}, i' \in I} \quad (4.14)$$

is a family of spectral immersions relative to \mathcal{M} such that the image of the canonical morphism $\coprod_{j \in J_{i'}, i' \in I} Y_{i'j} \rightarrow \coprod_{i' \in I} X_{i'} \rightarrow X$ is $F \times_H X$. This shows that $F \times_H X \subseteq X$ is Zariski open relative to \mathcal{M} . This completes the proof. \square

Proposition 4.6. *Let $X \in Aff_C$ be an affine scheme.*

- (1) *Let $f : F \rightarrow X$ be a Zariski open immersion relative to \mathcal{M} . Then, the image of f is Zariski open relative to \mathcal{M} .*
- (2) *Let $F \in Sh(Aff_C)_{\mathcal{M}}$ be a subsheaf of X that is Zariski open relative to \mathcal{M} . Then, the inclusion $F \hookrightarrow X$ is a Zariski open immersion relative to \mathcal{M} .*

Proof. (1) The proof is straightforward.

(2) Since $F \subseteq X$ is Zariski open relative to \mathcal{M} , there exists a family $\{X_i \rightarrow X\}_{i \in I}$ of spectral immersions relative to \mathcal{M} such that the image of the canonical morphism $\coprod_i X_i \rightarrow X$ in $Sh(Aff_C)_{\mathcal{M}}$ is F . Let $Z \in Aff_C$ and $Z \rightarrow X$ be a morphism in $Sh(Aff_C)_{\mathcal{M}}$. For each $i' \in I$, we have a pullback diagram in $Sh(Aff_C)_{\mathcal{M}}$

$$\begin{array}{ccccc} X_{i'} \times_X Z & \longrightarrow & F \times_X Z & \hookrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ X_{i'} & \longrightarrow & F & \hookrightarrow & X \end{array} \quad (4.15)$$

where the morphism $X_{i'} \rightarrow F$ is the composition $X_{i'} \rightarrow \coprod_{i \in I} X_i \rightarrow F$. We note that each $X_{i'} \times_X Z$ is affine. Using Lemma 3.5, it follows that each composition

$$X_{i'} \times_X Z \rightarrow F \times_X Z \rightarrow Z \quad (4.16)$$

in (4.15) is a spectral immersion relative to \mathcal{M} . Further, by Lemma 4.3, the induced morphism $\coprod_{i' \in I} X_{i'} \times_X Z \rightarrow Z$ has image $F \times_X Z$. Hence, $F \times_X Z \subseteq Z$ is Zariski open relative to \mathcal{M} . This shows that the inclusion $F \hookrightarrow X$ is a Zariski open immersion relative to \mathcal{M} . \square

Our next result shows that spectral immersions relative to \mathcal{M} are also Zariski open immersions relative to \mathcal{M} in $Sh(Aff_C)_{\mathcal{M}}$.

Proposition 4.7. *Let $\alpha^{op} : \text{Spec}(a) \rightarrow \text{Spec}(b)$ be a spectral immersion relative to \mathcal{M} . Then α^{op} considered as a morphism $Z = \text{Aff}_C(_, \text{Spec}(a)) \rightarrow \text{Aff}_C(_, \text{Spec}(b)) = Y$ is a Zariski open immersion relative to \mathcal{M} in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$.*

Proof. Let $X = \text{Spec}(c)$ be an affine scheme and $g : X \rightarrow Y$ be a morphism in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$. Using Lemma 3.5, the morphism

$$Z \times_Y X = \text{Spec}(a \otimes_b c) \cong \text{Spec}(a) \times_{\text{Spec}(b)} \text{Spec}(c) \rightarrow \text{Spec}(c) = X \quad (4.17)$$

is a spectral immersion relative to \mathcal{M} in Aff_C , and in particular a monomorphism in Aff_C . Since the Yoneda embedding $\text{Aff}_C \hookrightarrow \text{PSh}(\text{Aff}_C)$ preserves limits, the morphism

$$Z \times_Y X \cong \text{Aff}_C(_, \text{Spec}(a) \times_{\text{Spec}(b)} \text{Spec}(c)) \cong \text{Aff}_C(_, \text{Spec}(a \otimes_b c)) \rightarrow \text{Aff}_C(_, \text{Spec}(c)) \cong X \quad (4.18)$$

is a monomorphism in $\text{PSh}(\text{Aff}_C)$, and hence in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$. Further, since the morphism in (4.17) is a spectral immersion relative to \mathcal{M} , the subsheaf $Z \times_Y X \subseteq X$ is Zariski open relative to \mathcal{M} by Definition 4.2(1). Hence, α^{op} is a Zariski open immersion relative to \mathcal{M} . \square

We are ready to define the notion of a scheme relative to \mathcal{M} .

Definition 4.8. (1) *Let $F \in \text{Sh}(\text{Aff}_C)_{\mathcal{M}}$. An affine Zariski \mathcal{M} -covering of F is a family $\{X_i \rightarrow F\}_{i \in I}$ of morphisms in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$ where*

- (i) *For each $i \in I$, $X_i \in \text{Aff}_C$ is an affine scheme and $X_i \rightarrow F$ is a Zariski open immersion relative to \mathcal{M} .*
- (ii) *The induced morphism $\coprod_{i \in I} X_i \rightarrow F$ is an epimorphism in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$.*

(2) *An object $F \in \text{Sh}(\text{Aff}_C)_{\mathcal{M}}$ is an \mathcal{M} -scheme if it admits an affine Zariski \mathcal{M} -covering.*

It is clear that every affine $X \in \text{Aff}_C$ is an \mathcal{M} -scheme. We let $\text{Sch}(\mathcal{C})_{\mathcal{M}}$ be the full subcategory of $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$ whose objects are \mathcal{M} -schemes. We now prove some properties of $\text{Sch}(\mathcal{C})_{\mathcal{M}}$.

Lemma 4.9. (1) *Let $X, Y \in \text{Aff}_C$ be affine schemes and $Y \rightarrow X \leftarrow G$ be morphisms in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$. If G is an \mathcal{M} -scheme, then $F := Y \times_X G$ in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$ is an \mathcal{M} -scheme.*

(2) *Suppose that $F \rightarrow F_0$ is a morphism in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$ where $F_0 \in \text{Sch}(\mathcal{C})_{\mathcal{M}}$. If $\{g_i : X_i \rightarrow F_0\}_{i \in I}$ is an affine Zariski \mathcal{M} -covering of F_0 such that each pullback $X_i \times_{F_0} F$ is an \mathcal{M} -scheme, then F is an \mathcal{M} -scheme.*

Proof. (1) Since G is an \mathcal{M} -scheme, it has an affine Zariski \mathcal{M} -covering $\{h_i : X_i \rightarrow G\}_{i \in I}$. For each $i \in I$, we have a pullback diagram in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$

$$\begin{array}{ccccc} X_i \times_X Y \cong X_i \times_G F & \longrightarrow & F & \longrightarrow & Y \\ \downarrow & & \downarrow & & \downarrow \\ X_i & \xrightarrow{h_i} & G & \longrightarrow & X \end{array} \quad (4.19)$$

It is clear that $X_i \times_G F \cong X_i \times_X Y$ is affine. Since $h_i : X_i \rightarrow G$ is a Zariski open immersion relative to \mathcal{M} , it follows from Lemma 4.5 that the morphism $X_i \times_G F \rightarrow F$ in (4.19) is a Zariski open immersion relative to \mathcal{M} . Further, by Lemma 4.3, the induced morphism

$$\coprod_{i \in I} X_i \times_G F \rightarrow F \quad (4.20)$$

is an epimorphism in $\text{Sh}(\text{Aff}_C)_{\mathcal{M}}$. It follows that $\{X_i \times_X Y \cong X_i \times_G F \rightarrow F\}_{i \in I}$ is an affine Zariski \mathcal{M} -covering of F . This shows that F is an \mathcal{M} -scheme.

(2) For each $i \in I$, we have a pullback square in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccc} X_i \times_{F_0} F & \xrightarrow{g'_i} & F \\ \downarrow & & \downarrow \\ X_i & \xrightarrow{g_i} & F_0 \end{array} \quad (4.21)$$

By assumption, there exists an affine Zariski \mathcal{M} -covering $\{U_{ij} \rightarrow X_i \times_{F_0} F\}_{j \in J_i}$ of $X_i \times_{F_0} F$. Since the morphism $g_i : X_i \rightarrow F_0$ is a Zariski open immersion relative to \mathcal{M} , it follows from Lemma 4.5 that for each $j \in J_i$, the composition

$$U_{ij} \rightarrow X_i \times_{F_0} F \xrightarrow{g'_i} F \quad (4.22)$$

is a Zariski open immersion relative to \mathcal{M} .

For each $i \in I$, since the morphism $\coprod_{j \in J_i} U_{ij} \rightarrow X_i \times_{F_0} F$ induced by the affine Zariski \mathcal{M} -covering $\{U_{ij} \rightarrow X_i \times_{F_0} F\}_{j \in J_i}$ is an epimorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$, the induced morphism

$$\coprod_{i \in I, j \in J_i} U_{ij} \rightarrow \coprod_{i \in I} X_i \times_{F_0} F \quad (4.23)$$

is an epimorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Further, by Lemma 4.3, the induced morphism $\coprod_{i \in I} X_i \times_{F_0} F \xrightarrow{\coprod_{i \in I} g'_i} F$ is an epimorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. It follows that $\{U_{ij} \rightarrow X_i \times_{F_0} F \xrightarrow{g'_i} F\}_{i \in I, j \in J_i}$ is an affine Zariski \mathcal{M} -covering of F . \square

Lemma 4.10. *Let F be an \mathcal{M} -scheme and let $F_0 \hookrightarrow F$ be a Zariski open immersion relative to \mathcal{M} in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Then, F_0 is an \mathcal{M} -scheme.*

Proof. Since F is an \mathcal{M} -scheme, there exists an affine Zariski \mathcal{M} -covering $\{g_i : X_i \rightarrow F\}_{i \in I}$. For each $i \in I$, we have a pullback square in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccc} F_0 \times_F X_i & \xrightarrow{p_i} & X_i \\ \downarrow g'_i & & \downarrow g_i \\ F_0 & \hookrightarrow & F \end{array} \quad (4.24)$$

Since $F_0 \hookrightarrow F$ is a Zariski open immersion relative to \mathcal{M} , the morphism $p_i : F_0 \times_F X_i \rightarrow X_i$ is a monomorphism whose image is Zariski open relative to \mathcal{M} in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. By Definition 4.2(1), there is a family

$$\{h_{ij} : U_{ij} \rightarrow X_i\}_{j \in J_i} \quad (4.25)$$

of spectral immersions relative to \mathcal{M} where each $U_{ij} \in Aff_{\mathcal{C}}$ is affine and such that $\coprod_{j \in J_i} U_{ij} \xrightarrow{\coprod_{j \in J_i} h_{ij}} X_i$ has image $F_0 \times_F X_i$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. It follows that the family $\{U_{ij} \rightarrow F_0 \times_F X_i \rightarrow F_0\}_{i \in I, j \in J_i}$ is an affine Zariski \mathcal{M} -covering of F_0 . Hence, F_0 is an \mathcal{M} -scheme. \square

Lemma 4.11. *Let $\{F_i : i \in I\}$ be a family of objects in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ and let $F := \coprod_{i \in I} F_i$. Then, for each $i' \in I$, the canonical morphism $F_{i'} \rightarrow F$ is a Zariski open immersion relative to \mathcal{M} .*

Proof. Let $X = Spec(a) \in Aff_{\mathcal{C}}$ be an affine and

$$f : X \rightarrow F = \coprod_{i \in I} F_i \quad (4.26)$$

be a morphism in $Sh(Aff_C)_M$. We have a pullback diagram in $Sh(Aff_C)_M$

$$\begin{array}{ccc} F_{i'} \times_F X & \longrightarrow & X \\ \downarrow & & \downarrow f \\ F_{i'} & \longrightarrow & F \end{array} \quad (4.27)$$

It is clear that $F_{i'} \rightarrow F$ is a monomorphism in $Sh(Aff_C)_M$. Hence, $F_{i'} \times_F X \rightarrow X$ is a monomorphism in $Sh(Aff_C)_M$. We claim that its image is Zariski open relative to M .

Under Yoneda Lemma, the morphism f in (4.26) corresponds to an element of $F(Spec(a))$. We note that the coproduct $F = \coprod_{i \in I} F_i$ in $Sh(Aff_C)_M$ is the sheafification of the coproduct of the family $\{F_i\}_{i \in I}$ in $PSH(Aff_C)$. Using the description of sheafification as a filtered colimit (see for instance, [22, § 3.5]) there exists a spectral M -cover $\{\alpha_k^{op} : X_k \rightarrow X\}_{k \in K}$ such that for each $k \in K$, there is an element $u(k) \in I$ along with a factorization

$$\begin{array}{ccc} X_k & \xrightarrow{\alpha_k^{op}} & X \\ f_k \downarrow & & \downarrow f \\ F_{u(k)} & \longrightarrow & F \end{array} \quad (4.28)$$

It follows that there exists $g_k : X_k \rightarrow F_{u(k)} \times_F X$ such that the following diagram commutes for each $k \in K$

$$\begin{array}{ccccc} & & X_k & & \\ & \swarrow f_k & \downarrow g_k & \searrow \alpha_k^{op} & \\ F_{u(k)} & \longleftarrow & F_{u(k)} \times_F X & \longrightarrow & X \end{array} \quad (4.29)$$

Using Lemma 4.3, we have $X \cong F \times_F X \cong \coprod_{i \in I} (F_i \times_F X)$. Now since $\{\alpha_k^{op} : X_k \rightarrow X\}_{k \in K}$ is a spectral M -cover, it follows from Lemma 4.1 that the induced morphism

$$\coprod_{i \in I} \coprod_{u(k)=i} X_k \cong \coprod_{k \in K} X_k \xrightarrow{\coprod_{k \in K} \alpha_k^{op} = \coprod_{i \in I} \coprod_{u(k)=i} g_k} X \cong \coprod_{i \in I} (F_i \times_F X) \quad (4.30)$$

is an epimorphism in $Sh(Aff_C)_M$. We note that the following is a pullback square in $Sh(Aff_C)_M$

$$\begin{array}{ccc} \coprod_{u(k)=i'} X_k & \xrightarrow{\coprod_{u(k)=i'} g_k} & F_{i'} \times_F X \\ \downarrow & & \downarrow \\ \coprod_{i \in I} \coprod_{u(k)=i} X_k & \xrightarrow{\coprod_{i \in I} \coprod_{u(k)=i} g_k} & \coprod_{i \in I} (F_i \times_F X) \cong X \end{array} \quad (4.31)$$

Using Lemma 4.3, it follows that the image of $\coprod_{u(k)=i'} \alpha_k^{op} : \coprod_{u(k)=i'} X_k \rightarrow X$ is $F_{i'} \times_F X$. This completes the proof. \square

Proposition 4.12. *The subcategory $Sch(C)_M$ of $Sh(Aff_C)_M$ is closed under*

- (1) *coproducts.*
- (2) *pullbacks.*

Proof. (1) Let $\{F_i\}_{i \in I}$ be a family of \mathcal{M} -schemes. We need to show that the coproduct $F := \coprod_{i \in I} F_i$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ is an \mathcal{M} -scheme. For each $i \in I$, there exists an affine Zariski \mathcal{M} -covering $\{U_{ij} \rightarrow F_i\}_{j \in J_i}$. Since the induced morphism $\coprod_{j \in J_i} U_{ij} \rightarrow F_i$ is an epimorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$, the induced morphism

$$\coprod_{i \in I} \coprod_{j \in J_i} U_{ij} \rightarrow F = \coprod_{i \in I} F_i \quad (4.32)$$

is an epimorphism in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Using Lemma 4.11 and Lemma 4.5, it follows that $\{U_{ij} \rightarrow F_i \rightarrow F\}_{i \in I, j \in J_i}$ is an affine Zariski \mathcal{M} -covering for F . Hence, F is an \mathcal{M} -scheme.

(2) Let $F \xrightarrow{f} H \xleftarrow{g} G$ be morphisms in $Sch(\mathcal{C})_{\mathcal{M}}$. We claim that $F \times_H G \in Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ is an \mathcal{M} -scheme. Since F and G are \mathcal{M} -schemes, there exist affine Zariski \mathcal{M} -coverings $\{X_i \rightarrow F\}_{i \in I}$ and $\{Y_j \rightarrow G\}_{j \in J}$. For each $i \in I$ and $j \in J$, we consider the following commutative diagram in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccccc} X_i \times_H Y_j \cong (X_i \times_H G) \times_G Y_j & \longrightarrow & Y_j & & \\ \downarrow & & \downarrow & & \\ X_i \times_H G \cong X_i \times_F (F \times_H G) & \longrightarrow & F \times_H G & \xrightarrow{f'} & G \\ \downarrow & & \downarrow g' & & \downarrow g \\ X_i & \longrightarrow & F & \xrightarrow{f} & H \end{array} \quad (4.33)$$

Using Lemma 4.9(2), to show that $F \times_H G$ is an \mathcal{M} -scheme, it suffices to show that each $X_i \times_H G$ is an \mathcal{M} -scheme. By a further application of Lemma 4.9(2) to the composition

$$X_i \times_H G \rightarrow F \times_H G \rightarrow G \quad (4.34)$$

in 4.33, we see that it suffices to show that each $(X_i \times_H G) \times_G Y_j \cong X_i \times_H Y_j$ is an \mathcal{M} -scheme. Hence, we may assume that $F = X$ and $G = Y$ are affine. We will show that the pullback in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ of the diagram $X \xrightarrow{f} H \xleftarrow{g} Y$ is an \mathcal{M} -scheme.

Let $\{h_k : Z_k \rightarrow H\}_{k \in K}$ be an affine Zariski \mathcal{M} -covering of the \mathcal{M} -scheme H . For each $k \in K$, we consider the following pullback diagram in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccccc} Z_k \times_H X & \xrightarrow{f_k} & Z_k & \xleftarrow{g_k} & Z_k \times_H Y \\ h_k^X \downarrow & & \downarrow h_k & & \downarrow h_k^Y \\ X & \xrightarrow{f} & H & \xleftarrow{g} & Y \end{array} \quad (4.35)$$

Since $h_k : Z_k \rightarrow H$ is a Zariski open immersion relative to \mathcal{M} , we see that $h_k^X : Z_k \times_H X \rightarrow X$ and $h_k^Y : Z_k \times_H Y \rightarrow Y$ are monomorphisms whose images are Zariski open relative to \mathcal{M} . By definition, there exists a family

$$\{U_{ki} \rightarrow X\}_{i \in I_k} \quad (\text{resp. } \{V_{kj} \rightarrow Y\}_{j \in J_k}) \quad (4.36)$$

of spectral immersions relative to \mathcal{M} such that the image of the canonical morphism $\coprod_{i \in I_k} U_{ki} \rightarrow X$ (resp. $\coprod_{j \in J_k} V_{kj} \rightarrow Y$) is $Z_k \times_H X$ (resp. $Z_k \times_H Y$). Using Lemma 4.3, it follows that the induced morphisms

$$\coprod_{k \in K, i \in I_k} U_{ki} \rightarrow \coprod_{k \in K} Z_k \times_H X \xrightarrow{\coprod_{k \in K} h_k^X} X \quad \text{and} \quad \coprod_{k \in K, j \in J_k} V_{kj} \rightarrow \coprod_{k \in K} Z_k \times_H Y \xrightarrow{\coprod_{k \in K} h_k^Y} Y \quad (4.37)$$

are epimorphisms in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Hence using Proposition 4.7, $\{U_{ki} \rightarrow X\}_{k \in K, i \in I_k}$ (resp. $\{V_{kj} \rightarrow Y\}_{k \in K, j \in J_k}$) is an affine Zariski \mathcal{M} -covering of X (resp. Y). Further, for each $k \in K, i' \in I_k$ and $j' \in J_k$, we have a commutative

diagram

$$\begin{array}{ccccccc}
U_{ki'} & \longrightarrow & \coprod_{i \in I_k} U_{ki} & \twoheadrightarrow & Z_k \times_H X & \xrightarrow{f_k} & Z_k \xleftarrow{g_k} Z_k \times_H Y \ll \coprod_{j \in J_k} V_{kj} \longleftarrow V_{kj'} \\
\downarrow & & & & \downarrow h_k & & \downarrow \\
X & \xrightarrow{\quad f \quad} & & & H & \xleftarrow{\quad g \quad} & Y
\end{array} \tag{4.38}$$

We now consider the following commutative diagram in $Sh(Aff_C)_M$

$$\begin{array}{ccccc}
U_{ki'} \times_H V_{kj'} \cong (U_{ki'} \times_H Y) \times_Y V_{kj'} & \longrightarrow & V_{kj'} & & \\
\downarrow & & \downarrow & \searrow & \\
U_{ki'} \times_H Y \cong U_{ki'} \times_X (X \times_H Y) & \longrightarrow & X \times_H Y & \xrightarrow{f'} & Y \\
\downarrow & & \downarrow g' & & \downarrow g \\
U_{ki'} & \longrightarrow & X & \xrightarrow{f} & H \\
& & & & \nwarrow h_k \\
& & & & Z_k
\end{array} \tag{4.39}$$

where the morphism $U_{ki'} \rightarrow Z_k$ (resp. $V_{kj'} \rightarrow Z_k$) is the composition

$$U_{ki'} \longrightarrow \coprod_{i \in I_k} U_{ki} \twoheadrightarrow Z_k \times_H X \xrightarrow{f_k} Z_k \quad (\text{resp. } V_{kj'} \longrightarrow \coprod_{j \in J_k} V_{kj} \twoheadrightarrow Z_k \times_H Y \xrightarrow{g_k} Z_k) \tag{4.40}$$

from (4.38). Using Lemma 4.9(2), to show that $X \times_H Y$ is an \mathcal{M} -scheme, it suffices to show that each $(U_{ki'} \times_H Y) \times_Y V_{kj'} \cong U_{ki'} \times_H V_{kj'}$ is an \mathcal{M} -scheme. Using Lemma (4.4), we note that $h_k : Z_k \rightarrow H$ is a monomorphism in $Sh(Aff_C)_M$. Hence, $U_{ki'} \times_H V_{kj'} \cong U_{ki'} \times_{Z_k} V_{kj'}$, which is an affine scheme. This completes the proof. \square

Lemma 4.13. *Let $Y = Spec(a) \in Aff_C$ be an affine, $F \in Sh(Aff_C)_M$ and $\{H_i\}_{i \in I}$ be a family of objects in $Sh(Aff_C)_M$ with $H := \coprod_{i \in I} H_i \in Sh(Aff_C)_M$. Let $f : H \rightarrow F$ be an epimorphism and $g : Y \rightarrow F$ be a morphism in $Sh(Aff_C)_M$. Then, there exists a family $\{\alpha_k^{op} : Y_k \rightarrow Y\}_{k \in K}$ of spectral immersions relative to \mathcal{M} such that*

(i) *the morphism $\coprod_{k \in K} \alpha_k^{op} : \coprod_{k \in K} Y_k \rightarrow Y$ is an epimorphism in $Sh(Aff_C)_M$.*

(ii) *for each $k \in K$, there is an element $w(k) \in I$ along with a factorization*

$$\begin{array}{ccc}
Y_k & \xrightarrow{\alpha_k^{op}} & Y \\
\downarrow & & \downarrow g \\
H_{w(k)} & \longrightarrow & H = \coprod_{i \in I} H_i \xrightarrow{f} F
\end{array} \tag{4.41}$$

Proof. Applying Yoneda Lemma, the morphism g corresponds to an element of $F(Spec(a))$. Since $f : H \rightarrow F$ is an epimorphism, it follows from [22, § 3.7, Corollary 5] that there is a spectral \mathcal{M} -cover $\{\beta_j^{op} : V_j \rightarrow Y\}_{j \in J}$ such that for each $j \in J$, there is a factorization

$$\begin{array}{ccc}
V_j & \xrightarrow{\beta_j^{op}} & Y \\
g_j \downarrow & & \downarrow g \\
\coprod_{i \in I} H_i = H & \xrightarrow{f} & F
\end{array} \tag{4.42}$$

For each $i \in I$ and $j \in J$, we consider the following pullback square in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccc} H_i \times_H V_j & \longrightarrow & V_j \\ \downarrow & & \downarrow g_j \\ H_i & \longrightarrow & H \end{array} \quad (4.43)$$

Using Lemma 4.11, we note that the morphism $H_i \times_H V_j \longrightarrow V_j$ in (4.43) is a monomorphism whose image is Zariski open relative to \mathcal{M} in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. For $i \in I$, $j \in J$, by choosing spectral immersions $\{Y_k^{ij} \longrightarrow V_j\}_{k \in K_{ij}}$ whose image is the same as that of $H_i \times_H V_j \longrightarrow V_j$, the result now follows from Lemma 3.5, Lemma 4.1 and Lemma 4.3. \square

The next proposition gives a characterization of the subcategory $Sch(\mathcal{C})_{\mathcal{M}}$ of $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ in terms of quotients.

Theorem 4.14. *An object $F \in Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ is an \mathcal{M} -scheme if and only if there exists a family $\{U_i\}_{i \in I}$ of affines and an equivalence relation $R \hookrightarrow H \times H$ on $H := \coprod_{i \in I} U_i$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ such that*

(i) *For any $i, j \in I$, the composition $R_{i,j} := R \times_{H \times H} (U_i \times U_j) \xrightarrow{r_{i,j}} U_i \times U_j \longrightarrow U_i$ is a Zariski open immersion relative to \mathcal{M} .*

(ii) *For each $i \in I$, the subobject $R_{i,i} \hookrightarrow U_i \times U_i$ is the image of the diagonal morphism $U_i \xrightarrow{(1_{U_i}, 1_{U_i})} U_i \times U_i$.*

(iii) *$F \cong H/R$, i.e*

$$F = \text{Coeq} \left(R \hookrightarrow H \times H \begin{array}{c} \xrightarrow{\pi_1} \\ \xrightarrow{\pi_2} \end{array} H \right) \quad (4.44)$$

in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$, where $\pi_1, \pi_2 : H \times H \longrightarrow H$ are the two projections.

Proof. Suppose that F is an \mathcal{M} -scheme. Then, F has an affine Zariski \mathcal{M} -covering $\{f_i : U_i \longrightarrow F\}_{i \in I}$. We set $H := \coprod_{i \in I} U_i$. Let R be the following pullback in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$

$$\begin{array}{ccc} R := H \times_F H & \xrightarrow{p_1} & H = \coprod_{i \in I} U_i \\ p_2 \downarrow & & \downarrow \coprod_{i \in I} f_i \\ H = \coprod_{i \in I} U_i & \xrightarrow{\coprod_{i \in I} f_i} & F \end{array} \quad (4.45)$$

It may be verified that $R \xrightarrow{(p_1, p_2)} H \times H$ is an equivalence relation on H . For each $i, j \in I$, let $R_{i,j} := R \times_{H \times H} (U_i \times U_j) \xrightarrow{r_{i,j}} U_i \times U_j$ denote the canonical morphism to $U_i \times U_j$. It may be verified that $R_{i,j}$ may also be written as the pullback

$$\begin{array}{ccccc} U_i \times_F U_j \cong R_{i,j} & \xrightarrow{r_{i,j}} & U_i \times U_j & \longrightarrow & U_j \\ r_{i,j} \downarrow & & & & \downarrow f_j \\ U_i \times U_j & & & & \\ \downarrow & & & & \downarrow \\ U_i & \xrightarrow{f_i} & & & F \end{array} \quad (4.46)$$

Since $f_j : U_j \longrightarrow F$ is a Zariski open immersion relative to \mathcal{M} , it follows from Lemma 4.5 that the composition $R_{i,j} \xrightarrow{r_{i,j}} U_i \times U_j \longrightarrow U_i$ in (4.46) is a Zariski open immersion relative to \mathcal{M} . Hence, R satisfies condition (i). Taking $i = j$ in 4.46, we see that there is a morphism $s_i : U_i \longrightarrow R_{i,i}$ such that

$$\left(U_i \xrightarrow{(1_{U_i}, 1_{U_i})} U_i \times U_i \right) = \left(U_i \xrightarrow{s_i} R_{i,i} \xrightarrow{r_{i,i}} U_i \times U_i \right) \quad (4.47)$$

Using Lemma 4.4, it may be verified that s_i is an isomorphism. Hence, R satisfies condition (ii). Since $H \rightarrow F$ is an epimorphism, condition (iii) follows from [22, § 4.7, Theorem 8].

Conversely, let $\{U_i\}_{i \in I}$ be a family of affines and $R \hookrightarrow H \times H$ be an equivalence relation on $H = \coprod_{i \in I} U_i$ satisfying conditions (i), (ii) and (iii). Let $f : H \rightarrow F$ be the canonical epimorphism associated to the coequalizer in (4.44).

For each $i \in I$, let $f_i : U_i \rightarrow F$ be the composition $U_i \rightarrow H \xrightarrow{f} F$. Using condition (iii), it may be verified that for each $i, j \in I$, we have a pullback square as in (4.46) in $Sh(Aff_C)_{\mathcal{M}}$. We claim that

$$\{f_i : U_i \rightarrow F\}_{i \in I} \quad (4.48)$$

is an affine Zariski \mathcal{M} -covering. Since $f : H = \coprod_{i \in I} U_i \rightarrow F$ is an epimorphism, it suffices to show that $f_i : U_i \rightarrow F$ is a Zariski open immersion relative to \mathcal{M} for each $i \in I$. Since the monomorphism $U_i \xrightarrow{(1_{U_i}, 1_{U_i})} U_i \times U_i$ has a retraction, it follows from condition (ii) that

$$U_i \cong R_{i,i} \cong U_i \times_F U_i \quad (4.49)$$

Hence, $f_i : U_i \rightarrow F$ is a monomorphism.

Let $Y = Spec(a) \in Aff_C$ be affine and $g : Y \rightarrow F$ be a morphism in $Sh(Aff_C)_{\mathcal{M}}$. We consider the following pullback square in $Sh(Aff_C)_{\mathcal{M}}$

$$\begin{array}{ccc} U_i \times_F Y & \longrightarrow & Y \\ \downarrow & & \downarrow g \\ U_i & \xrightarrow{f_i} & F \end{array} \quad (4.50)$$

It is clear that the morphism $U_i \times_F Y \rightarrow Y$ in (4.50) is a monomorphism. We claim that its image is Zariski open relative to \mathcal{M} . Using Lemma 4.13, it follows that there is a family $\{\alpha_k^{op} : Y_k \rightarrow Y\}_{k \in K}$ of spectral immersions relative to \mathcal{M} such that the morphism $\coprod_{k \in K} \alpha_k^{op} : \coprod_{k \in K} Y_k \rightarrow Y$ is an epimorphism in $Sh(Aff_C)_{\mathcal{M}}$ and such that for each $k \in K$, there is an element $w(k) \in I$ along with a factorization

$$\begin{array}{ccc} Y_k & \xrightarrow{\alpha_k^{op}} & Y \\ h_k \downarrow & & \downarrow g \\ U_{w(k)} & \xrightarrow{f_{w(k)}} & F \end{array} \quad (4.51)$$

Since the square

$$\begin{array}{ccccc} R_{i,w(k)} & \xrightarrow{r_{i,w(k)}} & U_i \times U_{w(k)} & \longrightarrow & U_{w(k)} \\ r_{i,w(k)} \downarrow & & & & \downarrow f_{w(k)} \\ U_i \times U_{w(k)} & & & & \\ \downarrow & & & & \downarrow \\ U_i & \xrightarrow{f_i} & & & F \end{array} \quad (4.52)$$

is a pullback in $Sh(Aff_C)_{\mathcal{M}}$, it follows from (4.51) that

$$(U_i \times_F Y) \times_Y Y_k \cong (U_i \times_F U_{w(k)}) \times_{U_{w(k)}} Y_k \cong R_{i,w(k)} \times_{U_{w(k)}} Y_k \quad (4.53)$$

We now have the following pullback square in $Sh(Aff_C)_{\mathcal{M}}$

$$\begin{array}{ccccc} (U_i \times_F Y) \times_Y Y_k \cong R_{i,w(k)} \times_{U_{w(k)}} Y_k & \longrightarrow & Y_k & & \\ \downarrow & & \downarrow h_k & & \\ R_{i,w(k)} & \xrightarrow{r_{i,w(k)}} & U_i \times U_{w(k)} & \longrightarrow & U_{w(k)} \end{array} \quad (4.54)$$

Since by condition (i), the morphism $R_{i,w(k)} \xrightarrow{r_{i,w(k)}} U_i \times U_{w(k)} \rightarrow U_{w(k)}$ is a Zariski open immersion relative to \mathcal{M} , it follows that the morphism $(U_i \times_F Y) \times_Y Y_k \cong R_{i,w(k)} \times_{U_{w(k)}} Y_k \rightarrow Y_k$ in (4.54) is a monomorphism whose image is Zariski open relative to \mathcal{M} . Hence, there exists a family $\{Z_{ikj} \rightarrow Y_k\}_{j \in J_{ik}}$ of spectral immersions relative to \mathcal{M} such that the image of the induced morphism $\coprod_{j \in J_{ik}} Z_{ikj} \rightarrow Y_k$ is $(U_i \times_F Y) \times_Y Y_k$. It follows from Lemma 3.5 and Lemma 4.3 that $\{Z_{ikj} \rightarrow Y_k \xrightarrow{\alpha_k^{op}} Y\}_{k \in K, j \in J_{ik}}$ is a family of spectral immersions relative to \mathcal{M} such that the image of the induced morphism $\coprod_{k \in K, j \in J_{ik}} Z_{ikj} \rightarrow Y$ is $U_i \times_F Y$. Hence, $U_i \times_F Y$ is Zariski open relative to \mathcal{M} . This shows that $f_i : U_i \rightarrow F$ is a Zariski open immersion relative to \mathcal{M} . The proof is now complete. \square

5 Change of Base

Throughout this section, we assume that $(\mathcal{C}, \otimes, 1)$ and $(\mathcal{D}, \otimes, 1)$ are closed symmetric monoidal categories which are both complete and cocomplete. We fix an adjunction

$$(\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}, \mathbf{A} : \mathcal{D} \rightarrow \mathcal{C}) \quad (5.1)$$

with unit $\eta : id_{\mathcal{C}} \rightarrow \mathbf{A}\mathbf{B}$ and counit $\varepsilon : \mathbf{B}\mathbf{A} \rightarrow id_{\mathcal{D}}$. We further assume that the left adjoint \mathbf{B} has a strong symmetric monoidal structure i.e. there is a natural isomorphism

$$(\mathbf{B}(a) \otimes \mathbf{B}(b) \xrightarrow{\sim} \mathbf{B}(a \otimes b) : a, b \in \mathcal{C}) \quad (5.2)$$

and an isomorphism $1 \xrightarrow{\sim} \mathbf{B}(1)$ subject to certain coherence axioms (see for instance, [21, § 11.2]). In that case, it may be verified that the right adjoint \mathbf{A} has a lax symmetric monoidal structure given by maps (see for instance, [21, § 11.2])

$$(\mathbf{A}(a) \otimes \mathbf{A}(b) \rightarrow \mathbf{A}(a \otimes b) : a, b \in \mathcal{D}) \quad (5.3)$$

It then follows that the adjunction in (5.1) induces an adjunction

$$(\mathbf{B} : \text{Comm}(\mathcal{C}) \rightarrow \text{Comm}(\mathcal{D}), \mathbf{A} : \text{Comm}(\mathcal{D}) \rightarrow \text{Comm}(\mathcal{C})) \quad (5.4)$$

or equivalently, an adjunction

$$(\mathbf{A}^{op} : \text{Aff}_{\mathcal{D}} \rightarrow \text{Aff}_{\mathcal{C}}, \mathbf{B}^{op} : \text{Aff}_{\mathcal{C}} \rightarrow \text{Aff}_{\mathcal{D}}) \quad (5.5)$$

Hence, there is an adjunction $(\mathbf{A}_!, \mathbf{B}_!)$

$$\mathbf{A}_! = - \circ \mathbf{A} : \text{PSh}(\text{Aff}_{\mathcal{C}}) \rightarrow \text{PSh}(\text{Aff}_{\mathcal{D}}), \mathbf{B}_! = - \circ \mathbf{B} : \text{PSh}(\text{Aff}_{\mathcal{D}}) \rightarrow \text{PSh}(\text{Aff}_{\mathcal{C}}) \quad (5.6)$$

We also fix a left \mathcal{C} -module category (\mathcal{M}, \boxtimes) (resp. a left \mathcal{D} -module category (\mathcal{N}, \boxplus)) such that \mathcal{M} (resp. \mathcal{N}) is both complete and cocomplete and the bifunctor $\boxtimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ (resp. $\boxplus : \mathcal{D} \times \mathcal{N} \rightarrow \mathcal{N}$) preserves colimits in both variables.

Lemma 5.1. (1) The functor $\mathbf{B}^{op} : \text{Aff}_{\mathcal{C}} \rightarrow \text{Aff}_{\mathcal{D}}$ preserves limits.

(2) Suppose that $\mathbf{B}^{op} : \text{Aff}_{\mathcal{C}} \rightarrow \text{Aff}_{\mathcal{D}}$ takes fpqc \mathcal{M} -covers to fpqc \mathcal{N} -covers and $\mathbf{A} : \mathcal{D} \rightarrow \mathcal{C}$ preserves filtered colimits. Then $\mathbf{B}^{op} : \text{Aff}_{\mathcal{C}} \rightarrow \text{Aff}_{\mathcal{D}}$ takes spectral immersions relative to \mathcal{M} to spectral immersions relative to \mathcal{N} and takes spectral \mathcal{M} -covers to spectral \mathcal{N} -covers.

Proof. (1) The result is clear from the adjunction in (5.5).

(2) It follows from (5.4) that $\mathbf{B} : \text{Comm}(\mathcal{C}) \rightarrow \text{Comm}(\mathcal{D})$ preserves colimits and hence epimorphisms. Let $\alpha : a \rightarrow b$ be an \mathcal{M} -flat morphism in $\text{Comm}(\mathcal{C})$. Then $\{\alpha^{op} : \text{Spec}(b) \rightarrow \text{Spec}(a), 1_{\text{Spec}(a)} : \text{Spec}(a) \rightarrow \text{Spec}(a)\}$ is an fpqc \mathcal{M} -cover in $\text{Aff}_{\mathcal{C}}$. Since \mathbf{B}^{op} takes fpqc \mathcal{M} -covers to fpqc \mathcal{N} -covers, we see that $\{\mathbf{B}^{op}(\alpha^{op}) = \mathbf{B}(\alpha)^{op}, 1_{\text{Spec}(\mathbf{B}(a))}\}$ is

an fpqc \mathcal{N} -cover in $Aff_{\mathcal{D}}$. In particular, $\mathbf{B}(\alpha)$ is \mathcal{N} -flat, i.e., $\mathbf{B} : Comm(\mathcal{C}) \rightarrow Comm(\mathcal{D})$ takes \mathcal{M} -flat morphisms to \mathcal{N} -flat morphisms.

We now show that $\mathbf{B} : Comm(\mathcal{C}) \rightarrow Comm(\mathcal{D})$ preserves morphisms of finite type. We note that since $\mathbf{A} : \mathcal{D} \rightarrow \mathcal{C}$ preserves filtered colimits, so does the induced functor $\mathbf{A} : Comm(\mathcal{D}) \rightarrow Comm(\mathcal{C})$. Let $\alpha : a \rightarrow b$ be a morphism of finite type in $Comm(\mathcal{C})$. We need to show that $\mathbf{B}(\alpha) : \mathbf{B}(a) \rightarrow \mathbf{B}(b)$ is of finite type. Let $\mathbb{D} : \mathcal{I} \rightarrow \mathbf{B}(a)/Comm(\mathcal{D})$ be a filtered diagram. It may be verified that (5.4) induces an adjunction $(\widehat{\mathbf{B}}, \widehat{\mathbf{A}})$

$$\begin{aligned} \widehat{\mathbf{B}} : a/Comm(\mathcal{C}) &\rightarrow \mathbf{B}(a)/Comm(\mathcal{D}) & (a \xrightarrow{\beta} c) &\mapsto (\mathbf{B}(a) \xrightarrow{\mathbf{B}(\beta)} \mathbf{B}(c)) \\ \widehat{\mathbf{A}} : \mathbf{B}(a)/Comm(\mathcal{D}) &\rightarrow a/Comm(\mathcal{C}) & (\mathbf{B}(a) \xrightarrow{\gamma} d) &\mapsto (a \xrightarrow{\eta_a} \mathbf{A}\mathbf{B}(a) \xrightarrow{\mathbf{A}(\gamma)} \mathbf{A}(d)) \end{aligned} \quad (5.7)$$

We note that we have a commutative square

$$\begin{array}{ccc} \mathbf{B}(a)/Comm(\mathcal{D}) & \xrightarrow{\widehat{\mathbf{A}}} & a/Comm(\mathcal{C}) \\ \downarrow & & \downarrow \\ Comm(\mathcal{D}) & \xrightarrow{\mathbf{A}} & Comm(\mathcal{C}) \end{array} \quad (5.8)$$

where the vertical arrows in (5.8) are the forgetful functors. Since $\mathbf{A} : Comm(\mathcal{D}) \rightarrow Comm(\mathcal{C})$ and the forgetful functors in (5.8) preserve filtered colimits and the functor $a/Comm(\mathcal{C}) \rightarrow \mathcal{C}$ is conservative, it follows from (5.8) that $\widehat{\mathbf{A}} : \mathbf{B}(a)/Comm(\mathcal{D}) \rightarrow a/Comm(\mathcal{C})$ preserves filtered colimits. Since $\alpha : a \rightarrow b$ is a morphism of finite type in $Comm(\mathcal{C})$, we have

$$\begin{aligned} \mathbf{B}(a)/Comm(\mathcal{D})(\mathbf{B}(\alpha), \operatorname{colim}_{i \in \mathcal{I}} \mathbb{D}(i)) &\cong \mathbf{B}(a)/Comm(\mathcal{D})(\widehat{\mathbf{B}}(\alpha), \operatorname{colim}_{i \in \mathcal{I}} \mathbb{D}(i)) \\ &\cong a/Comm(\mathcal{C})(\alpha, \widehat{\mathbf{A}}(\operatorname{colim}_{i \in \mathcal{I}} \mathbb{D}(i))) \\ &\cong a/Comm(\mathcal{C})(\alpha, \operatorname{colim}_{i \in \mathcal{I}} \widehat{\mathbf{A}}(\mathbb{D}(i))) \\ &\cong \operatorname{colim}_{i \in \mathcal{I}} a/Comm(\mathcal{C})(\alpha, \widehat{\mathbf{A}}(\mathbb{D}(i))) \\ &\cong \operatorname{colim}_{i \in \mathcal{I}} \mathbf{B}(a)/Comm(\mathcal{D})(\widehat{\mathbf{B}}(\alpha), \mathbb{D}(i)) \\ &\cong \operatorname{colim}_{i \in \mathcal{I}} \mathbf{B}(a)/Comm(\mathcal{D})(\mathbf{B}(\alpha), \mathbb{D}(i)) \end{aligned} \quad (5.9)$$

Hence, $\mathbf{B} : Comm(\mathcal{C}) \rightarrow Comm(\mathcal{D})$ preserves morphisms of finite type. It follows that $\mathbf{B}^{op} : Aff_{\mathcal{C}} \rightarrow Aff_{\mathcal{D}}$ takes spectral immersions relative to \mathcal{M} to spectral immersions relative to \mathcal{N} . Finally, since $\mathbf{B}^{op} : Aff_{\mathcal{C}} \rightarrow Aff_{\mathcal{D}}$ takes fpqc \mathcal{M} -covers to fpqc \mathcal{N} -covers, it follows that \mathbf{B}^{op} also takes spectral \mathcal{M} -covers to spectral \mathcal{N} -covers. This concludes the proof. \square

Since the functor $\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}$ has a strong symmetric monoidal structure, we now recall (see for instance, [8, Proposition 3.6.1]) that the left \mathcal{D} -action $\boxplus : \mathcal{D} \times \mathcal{N} \rightarrow \mathcal{N}$ restricts to the left \mathcal{C} -action $\boxplus^{\mathbf{B}} : \mathcal{C} \times \mathcal{N} \rightarrow \mathcal{N}$ given by

$$\mathcal{C} \times \mathcal{N} \xrightarrow{\mathbf{B} \times 1_{\mathcal{N}}} \mathcal{D} \times \mathcal{N} \xrightarrow{\boxplus} \mathcal{N} \quad (5.10)$$

We will denote the left \mathcal{C} -module category $(\mathcal{N}, \boxplus^{\mathbf{B}})$ by $\mathbf{B}_*(\mathcal{N})$. Since $\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}$ preserves colimits and $\boxplus : \mathcal{D} \times \mathcal{N} \rightarrow \mathcal{N}$ preserves colimits in both variables, it follows that $\boxplus^{\mathbf{B}} : \mathcal{C} \times \mathcal{N} \rightarrow \mathcal{N}$ preserves colimits in both variables. Further, for every $a \in Comm(\mathcal{C})$, we have $\mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(a) = \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a))$. We now fix a lax \mathcal{C} -linear functor

$$(\mathbb{L}, \Gamma) : \mathbf{B}_*(\mathcal{N}) = (\mathcal{N}, \boxplus^{\mathbf{B}}) \rightarrow (\mathcal{M}, \boxtimes) \quad (5.11)$$

We recall (see for instance, [8, Lemma 3.3.7]) that for each $a \in \text{Comm}(\mathcal{C})$, there is an induced functor

$$\begin{aligned} \mathbb{L}^a : \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a)) &= \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(a) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(a) \\ (n, a \boxplus^{\mathbf{B}} n \xrightarrow{\rho} n) &\mapsto \left(\mathbb{L}(n), a \boxtimes \mathbb{L}(n) \xrightarrow{\Gamma_{a,n}} \mathbb{L}(a \boxplus^{\mathbf{B}} n) \xrightarrow{\mathbb{L}(\rho)} \mathbb{L}(n) \right) \end{aligned} \quad (5.12)$$

By abuse of notation, we will typically denote \mathbb{L}^a simply by \mathbb{L} .

Lemma 5.2. *Let $\alpha : a = (a, \mu_a, \iota_a) \longrightarrow (b, \mu_b, \iota_b) = b$ be a morphism in $\text{Comm}(\mathcal{C})$. Then,*

(1) *For each $n = (n, \rho) \in \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a))$, there is a canonical morphism*

$$\theta_{\alpha, \mathbf{B}, \mathbb{L}, n} : b \boxtimes_a \mathbb{L}(n) \longrightarrow \mathbb{L}(\mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n) \quad (5.13)$$

in $\mathbf{Mod}_{\mathcal{M}}(b)$.

(2) *If \mathbb{L} preserves coequalizers and $(\mathbb{L}, \Gamma) : \mathbf{B}_*(\mathcal{N}) \longrightarrow (\mathcal{M}, \boxtimes)$ is strong \mathcal{C} -linear, i.e., Γ is a natural isomorphism, then for each $(n, \rho) \in \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a))$, the morphism in (5.13) is an isomorphism in $\mathbf{Mod}_{\mathcal{M}}(b)$.*

Proof. (1) We note that as in (2.3), $\mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n$ is given by the following coequalizer in $\mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(b)) = \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(b)$

$$\mathbf{B}(b) \boxplus (\mathbf{B}(a) \boxplus n) \xrightarrow[\sim]{1_{\mathbf{B}(b)} \boxplus \rho} (\mathbf{B}(b) \otimes \mathbf{B}(a)) \boxplus n \xrightarrow[\psi]{} \mathbf{B}(b) \boxplus n \xrightarrow{\text{coeq}_{\mathbf{B}(\alpha), (n, \rho)}} \mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n \quad (5.14)$$

where ψ is the composition

$$(\mathbf{B}(b) \otimes \mathbf{B}(a)) \boxplus n \cong \mathbf{B}(b \otimes a) \boxplus n \xrightarrow{\mathbf{B}(\mu_b \circ (1_b \otimes \alpha)) \boxplus 1_n} \mathbf{B}(b) \boxplus n \quad (5.15)$$

Applying $\mathbb{L} : \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(b)) = \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(b) \longrightarrow \mathbf{Mod}_{\mathcal{M}}(b)$ to (5.14) gives the following commutative diagram

$$\begin{array}{ccc} \mathbb{L}((\mathbf{B}(b) \otimes \mathbf{B}(a)) \boxplus n) & \xleftarrow{\sim} & \mathbb{L}(\mathbf{B}(b) \boxplus (\mathbf{B}(a) \boxplus n)) \xrightarrow{\mathbb{L}(1_{\mathbf{B}(b)} \boxplus \rho)} \mathbb{L}(\mathbf{B}(b) \boxplus n) \\ \mathbb{L}(\psi) \downarrow & & \downarrow \mathbb{L}(\text{coeq}_{\mathbf{B}(\alpha), (n, \rho)}) \\ \mathbb{L}(\mathbf{B}(b) \boxplus n) & \xrightarrow{\mathbb{L}(\text{coeq}_{\mathbf{B}(\alpha), (n, \rho)})} & \mathbb{L}(\mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n) \end{array} \quad (5.16)$$

in $\mathbf{Mod}_{\mathcal{M}}(b)$. It may be verified that

$$\Gamma_{b, \mathbf{B}(a) \boxplus n} \circ (1_b \boxtimes \Gamma_{a, n}) : b \boxtimes (a \boxtimes \mathbb{L}(n)) \longrightarrow \mathbb{L}(\mathbf{B}(b) \boxplus (\mathbf{B}(a) \boxplus n)), \quad \Gamma_{b, n} : b \boxtimes \mathbb{L}(n) \longrightarrow \mathbb{L}(\mathbf{B}(b) \boxplus n) \quad (5.17)$$

are morphisms in $\mathbf{Mod}_{\mathcal{M}}(b)$. We now have the following commutative diagrams in $\mathbf{Mod}_{\mathcal{M}}(b)$

$$\begin{array}{ccc} b \boxtimes (a \boxtimes \mathbb{L}(n)) & \xrightarrow{1_b \boxtimes (\mathbb{L}(\rho) \circ \Gamma_{a, n})} & b \boxtimes \mathbb{L}(n) \\ \Gamma_{b, \mathbf{B}(a) \boxplus n} \circ (1_b \boxtimes \Gamma_{a, n}) \downarrow & & \downarrow \Gamma_{b, n} \\ \mathbb{L}(\mathbf{B}(b) \boxplus (\mathbf{B}(a) \boxplus n)) & \xrightarrow{\mathbb{L}(1_{\mathbf{B}(b)} \boxplus \rho)} & \mathbb{L}(\mathbf{B}(b) \boxplus n) \end{array} \quad (5.18)$$

$$\begin{array}{ccc} b \boxtimes (a \boxtimes \mathbb{L}(n)) & \xrightarrow{\sim} & (b \otimes a) \boxtimes \mathbb{L}(n) \xrightarrow{(\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_{\mathbb{L}(n)}} b \boxtimes \mathbb{L}(n) \\ \Gamma_{b, \mathbf{B}(a) \boxplus n} \circ (1_b \boxtimes \Gamma_{a, n}) \downarrow & & \downarrow \Gamma_{b, n} \\ \mathbb{L}(\mathbf{B}(b) \boxplus (\mathbf{B}(a) \boxplus n)) & \xrightarrow{\sim} & \mathbb{L}((\mathbf{B}(b) \otimes \mathbf{B}(a)) \boxplus n) \xrightarrow{\mathbb{L}(\psi)} \mathbb{L}(\mathbf{B}(b) \boxplus n) \end{array} \quad (5.19)$$

It follows from (5.16) and the universal property of the coequalizer

$$b \boxtimes_a \mathbb{L}(n) = \text{Coeq} \left(b \boxtimes (a \boxtimes \mathbb{L}(n)) \xrightarrow[\sim]{} (b \otimes a) \boxtimes \mathbb{L}(n) \xrightarrow[\substack{1_b \boxtimes (\mathbb{L}(\rho) \circ \Gamma_{a,n}) \\ (\mu_b \circ (1_b \otimes \alpha)) \boxtimes 1_{\mathbb{L}(n)}}]{} b \boxtimes \mathbb{L}(n) \right) \quad (5.20)$$

that there is a morphism $\theta_{\alpha, \mathbf{B}, \mathbb{L}, n} : b \boxtimes_a \mathbb{L}(n) \rightarrow \mathbb{L}(\mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n)$ in $\mathbf{Mod}_{\mathcal{M}}(b)$.

(2) The result is clear from the proof of (1). \square

Theorem 5.3. *Suppose that we are given the following data:*

(a) *An adjunction $(\mathbf{B} : \mathcal{C} \rightarrow \mathcal{D}, \mathbf{A} : \mathcal{D} \rightarrow \mathcal{C})$ between closed symmetric monoidal categories \mathcal{C} and \mathcal{D} such that the left adjoint \mathbf{B} has a strong symmetric monoidal structure.*

(b) *A left \mathcal{C} -module category (\mathcal{M}, \boxtimes) , a left \mathcal{D} -module category (\mathcal{N}, \boxplus) and a lax \mathcal{C} -linear functor*

$$(\mathbb{L}, \Gamma) : \mathbf{B}_*(\mathcal{N}) = (\mathcal{N}, \boxplus^{\mathbf{B}}) \rightarrow (\mathcal{M}, \boxtimes) \quad (5.21)$$

satisfying conditions described above.

Suppose also that $\mathbf{A} : \mathcal{D} \rightarrow \mathcal{C}$ preserves filtered colimits and the functor $\mathbb{L} : \mathbf{B}_(\mathcal{N}) \rightarrow \mathcal{M}$ is conservative and preserves finite limits. Further, suppose that for any \mathcal{M} -flat morphism $a \rightarrow b$ in $\text{Comm}(\mathcal{C})$ and any $n \in \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a))$, the canonical morphism in (5.13)*

$$b \boxtimes_a \mathbb{L}(n) \rightarrow \mathbb{L}(\mathbf{B}(b) \boxplus_{\mathbf{B}(a)} n) \quad (5.22)$$

is an isomorphism in $\mathbf{Mod}_{\mathcal{M}}(b)$. Then,

(1) *The functor $\mathbf{B}^{op} : \text{Aff}_{\mathcal{C}} \rightarrow \text{Aff}_{\mathcal{D}}$ takes fpqc \mathcal{M} -covers to fpqc \mathcal{N} -covers.*

(2) *The functor $\mathbf{B}_! = _ \circ \mathbf{B} : PSh(\text{Aff}_{\mathcal{D}}) \rightarrow PSh(\text{Aff}_{\mathcal{C}})$ in (5.6) restricts to a functor $\widetilde{\mathbf{B}}_! : Sh(\text{Aff}_{\mathcal{D}})_{\mathcal{N}} \rightarrow Sh(\text{Aff}_{\mathcal{C}})_{\mathcal{M}}$ which has a left adjoint $\widetilde{\mathbf{A}}_! : Sh(\text{Aff}_{\mathcal{C}})_{\mathcal{M}} \rightarrow Sh(\text{Aff}_{\mathcal{D}})_{\mathcal{N}}$.*

Additionally, if \mathcal{M} and \mathcal{N} are subcanonical, then

(3) *The functor $\widetilde{\mathbf{A}}_!$ restricts to a functor $Sch(\mathcal{C})_{\mathcal{M}} \rightarrow Sch(\mathcal{D})_{\mathcal{N}}$ sending $F \mapsto \widetilde{\mathbf{A}}_!(F)$.*

For any affine $X = \text{Spec}(a) \in \text{Aff}_{\mathcal{C}}$, $\widetilde{\mathbf{A}}_!(X)$ is canonically isomorphic to the affine scheme $\mathbf{B}^{op}(\text{Spec}(a)) \in \text{Aff}_{\mathcal{D}}$.

Proof. (1) Since $\mathbb{L} : \mathbf{B}_*(\mathcal{N}) \rightarrow \mathcal{M}$ is conservative and preserves finite limits, it follows from Remark 2.2 that for each $a \in \text{Comm}(\mathcal{C})$, the functor $\mathbb{L} = \mathbb{L}^a : \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(a) \rightarrow \mathbf{Mod}_{\mathcal{M}}(a)$ in (5.12) is also conservative and preserves finite limits. Let $\{\alpha_i^{op} : \text{Spec}(a_i) \rightarrow \text{Spec}(a)\}_{i \in I}$ be an fpqc \mathcal{M} -cover in $\text{Aff}_{\mathcal{C}}$. We need to show that

$$\{\mathbf{B}^{op}(\alpha_i^{op}) : \mathbf{B}^{op}(\text{Spec}(a_i)) \rightarrow \mathbf{B}^{op}(\text{Spec}(a))\}_{i \in I} \quad (5.23)$$

is an fpqc \mathcal{N} -cover in $\text{Aff}_{\mathcal{D}}$. By definition, each $\alpha_i : a \rightarrow a_i$ is \mathcal{M} -flat and there exists a finite subset $J \subseteq I$ such that the functor $((\alpha_j)^*)_{j \in J} = (a_j \boxtimes_a _)_{j \in J} : \mathbf{Mod}_{\mathcal{M}}(a) \rightarrow \prod_{j \in J} \mathbf{Mod}_{\mathcal{M}}(a_j)$ is conservative. For each $i \in I$, we know that $\alpha_i : a \rightarrow a_i$ is \mathcal{M} -flat and hence the following square

$$\begin{array}{ccc} \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(a) = \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a)) & \xrightarrow{\mathbb{L}^a} & \mathbf{Mod}_{\mathcal{M}}(a) \\ \mathbf{B}(\alpha_i)^* \downarrow & & \downarrow \alpha_i^* \\ \mathbf{Mod}_{\mathbf{B}_*(\mathcal{N})}(a_i) = \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a_i)) & \xrightarrow{\mathbb{L}^{a_i}} & \mathbf{Mod}_{\mathcal{M}}(a_i) \end{array} \quad (5.24)$$

commutes up to a natural isomorphism. Since $\mathbb{L}^a, \mathbb{L}^{a_i}$ preserve finite limits, α_i is \mathcal{M} -flat (i.e., α_i^* preserves finite limits) and \mathbb{L}^{a_i} is conservative, it follows from (5.24) that $\mathbf{B}(\alpha_i)^*$ preserves finite limits, i.e., $\mathbf{B}(\alpha_i)$ is \mathcal{N} -flat.

Further, since the square (5.24) commutes up to a natural isomorphism for each $i \in J \subseteq I$, the following diagram

$$\begin{array}{ccc} \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a)) & \xrightarrow{\mathbb{L}^a} & \mathbf{Mod}_{\mathcal{M}}(a) \\ (\mathbf{B}(\alpha_j)^*)_{j \in J} \downarrow & & \downarrow (\alpha_j^*)_{j \in J} \\ \prod_{j \in J} \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a_j)) & \xrightarrow{\prod_{j \in J} \mathbb{L}^{a_j}} & \prod_{j \in J} \mathbf{Mod}_{\mathcal{M}}(a_j) \end{array} \quad (5.25)$$

also commutes up to a natural isomorphism. Since \mathbb{L}^a , $(\alpha_j^*)_{j \in J}$ and $\prod_{j \in J} \mathbb{L}^{a_j}$ are conservative, it follows from (5.25) that

$$(\mathbf{B}(\alpha_j)^*)_{j \in J} : \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a)) \longrightarrow \prod_{j \in J} \mathbf{Mod}_{\mathcal{N}}(\mathbf{B}(a_j)) \quad (5.26)$$

is conservative. Hence, $\{\mathbf{B}^{op}(\alpha_i^{op}) : \mathbf{B}^{op}(\text{Spec}(a_i)) \longrightarrow \mathbf{B}^{op}(\text{Spec}(a))\}_{i \in I}$ is an fpqc \mathcal{N} -cover in $Aff_{\mathcal{D}}$.

(2) Using part (1) and Lemma 5.1, it follows that $\mathbf{B}^{op} : Aff_{\mathcal{C}} \longrightarrow Aff_{\mathcal{D}}$ takes spectral \mathcal{M} -covers to spectral \mathcal{N} -covers. Since \mathbf{B}^{op} preserves pullbacks, it follows that the functor $\mathbf{B}_! = _ \circ \mathbf{B} : PSh(Aff_{\mathcal{D}}) \longrightarrow PSh(Aff_{\mathcal{C}})$ in (5.6) restricts to a functor

$$\widetilde{\mathbf{B}}_! : Sh(Aff_{\mathcal{D}})_{\mathcal{N}} \longrightarrow Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \quad (5.27)$$

We define the functor $\widetilde{\mathbf{A}}_! : Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \longrightarrow Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$ as the composition

$$Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \hookrightarrow PSh(Aff_{\mathcal{C}}) \xrightarrow{\mathbf{A}_! = _ \circ \mathbf{A}} PSh(Aff_{\mathcal{D}}) \xrightarrow{(-)_{\mathcal{N}}^{++}} Sh(Aff_{\mathcal{D}})_{\mathcal{N}} \quad (5.28)$$

where $(-)_{\mathcal{N}}^{++}$ is the sheafification functor associated to the spectral \mathcal{N} -topology on $Aff_{\mathcal{D}}$. Since $(-)_{\mathcal{N}}^{++}$ has a right adjoint and $(\mathbf{A}_!, \mathbf{B}_!)$ is an adjoint pair as in (5.6), we have

$$\begin{aligned} Sh(Aff_{\mathcal{D}})_{\mathcal{N}}(\widetilde{\mathbf{A}}_!(F), G) &= Sh(Aff_{\mathcal{D}})_{\mathcal{N}}(\mathbf{A}_!(F)_{\mathcal{N}}^{++}, G) \\ &\cong PSh(Aff_{\mathcal{D}})(\mathbf{A}_!(F), G) \\ &\cong PSh(Aff_{\mathcal{C}})(F, \mathbf{B}_!(G)) = Sh(Aff_{\mathcal{C}})_{\mathcal{M}}(F, \widetilde{\mathbf{B}}_!(G)) \end{aligned} \quad (5.29)$$

for $F \in Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$ and $G \in Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$.

(3) Since \mathcal{M} is subcanonical, $Aff_{\mathcal{C}}$ is a full subcategory of $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. For $\text{Spec}(a) \in Aff_{\mathcal{C}}$, we have natural isomorphisms

$$\begin{aligned} \widetilde{\mathbf{A}}_!(Aff_{\mathcal{C}}(_, \text{Spec}(a))) &= (Aff_{\mathcal{C}}(\mathbf{A}^{op}(_), \text{Spec}(a)))_{\mathcal{N}}^{++} \\ &\cong (Aff_{\mathcal{D}}(_, \mathbf{B}^{op}(\text{Spec}(a))))_{\mathcal{N}}^{++} \quad [\text{using the adjunction in (5.5)}] \\ &\cong Aff_{\mathcal{D}}(_, \mathbf{B}^{op}(\text{Spec}(a))) \quad [\text{since } \mathcal{N} \text{ is subcanonical}] \end{aligned} \quad (5.30)$$

Hence, $\widetilde{\mathbf{A}}_! : Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \longrightarrow Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$ restricts to $\mathbf{B}^{op} : Aff_{\mathcal{C}} \longrightarrow Aff_{\mathcal{D}}$ on $Aff_{\mathcal{C}} \subset Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$.

Now, let $F \in Sch(\mathcal{C})_{\mathcal{M}} \subseteq Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. We need to show that $\widetilde{\mathbf{A}}_!(F) \in Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$ is an \mathcal{N} -scheme. It follows from Proposition 4.14 that there exists a family $\{U_i\}_{i \in I}$ of affines in $Aff_{\mathcal{C}}$ and an equivalence relation $R \subseteq H \times H$ on $H := \coprod_{i \in I} U_i$ such that $F \cong H/R$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$. Further, for each $i, j \in I$, the composition $R_{i,j} := R \times_{H \times H} (U_i \times U_j) \xrightarrow{r_{i,j}} U_i \times U_j \longrightarrow U_i$ is a Zariski open immersion relative to \mathcal{M} and the subobject $R_{i,i} \subseteq U_i \times U_i$ is the image of $U_i \xrightarrow{(1_{U_i}, 1_{U_i})} U_i \times U_i$ in $Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$.

Since limits of presheaves are computed objectwise, we see that $\mathbf{A}_! = _ \circ \mathbf{A} : PSh(Aff_{\mathcal{C}}) \longrightarrow PSh(Aff_{\mathcal{D}})$ preserves limits. Further, since the sheafification functor $(-)_{\mathcal{N}}^{++}$ preserves finite limits, it follows from (5.28) that $\widetilde{\mathbf{A}}_! : Sh(Aff_{\mathcal{C}})_{\mathcal{M}} \longrightarrow Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$ preserves finite limits and in particular, monomorphisms. Also, since $\widetilde{\mathbf{A}}_!$ is a

left adjoint, it preserves colimits and in particular, epimorphisms. Hence, $\widetilde{\mathbf{A}}_!(R) \subseteq \widetilde{\mathbf{A}}_!(H) \times \widetilde{\mathbf{A}}_!(H)$ is an equivalence relation on $\widetilde{\mathbf{A}}_!(H)$ and

$$\widetilde{\mathbf{A}}_!(F) \cong \widetilde{\mathbf{A}}_!(H) / \widetilde{\mathbf{A}}_!(R) \cong \left(\prod_{i \in I} \widetilde{\mathbf{A}}_!(U_i) \right) / \widetilde{\mathbf{A}}_!(R) \quad (5.31)$$

in $Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$. We will verify that the equivalence relation $\widetilde{\mathbf{A}}_!(R)$ satisfies the conditions of Proposition 4.14. Using (5.30), it follows that for each $i \in I$, $\widetilde{\mathbf{A}}_!(U_i) = \mathbf{B}^{op}(U_i) \in Aff_{\mathcal{D}}$ is an affine. Further, $\widetilde{\mathbf{A}}_!(R_{i,i}) \subseteq \widetilde{\mathbf{A}}_!(U_i) \times \widetilde{\mathbf{A}}_!(U_i)$ is the image of $(1_{\widetilde{\mathbf{A}}_!(U_i)}, 1_{\widetilde{\mathbf{A}}_!(U_i)}) : \widetilde{\mathbf{A}}_!(U_i) \rightarrow \widetilde{\mathbf{A}}_!(U_i) \times \widetilde{\mathbf{A}}_!(U_i)$ in $Sh(Aff_{\mathcal{D}})_{\mathcal{N}}$.

Now, for each $i, j \in I$, since $R_{i,j} \xrightarrow{r_{i,j}} U_i \times U_j \rightarrow U_i$ is a Zariski open immersion relative to \mathcal{M} , it follows from Proposition 4.6 that its image is Zariski open relative to \mathcal{M} . Hence, there exists a family $\{\beta_{ijk}^{op} : Y_{ijk} \rightarrow U_i\}_{k \in K_{ij}}$ of spectral immersions relative to \mathcal{M} such that the image of the induced morphism $\prod_{k \in K_{ij}} \beta_{ijk}^{op} : \prod_{k \in K_{ij}} Y_{ijk} \rightarrow U_i$ is $R_{i,j}$. Since $\widetilde{\mathbf{A}}_!$ preserves colimits and finite limits and restricts to the functor $\mathbf{B}^{op} : Aff_{\mathcal{C}} \rightarrow Aff_{\mathcal{D}}$ on the subcategory $Aff_{\mathcal{C}} \subset Sh(Aff_{\mathcal{C}})_{\mathcal{M}}$, it follows from part (1) and Lemma 5.1 that

$$\left\{ \widetilde{\mathbf{A}}_!(Y_{ijk}) \cong \mathbf{B}^{op}(Y_{ijk}) \xrightarrow{\widetilde{\mathbf{A}}_!(\beta_{ijk}^{op}) = \mathbf{B}^{op}(\beta_{ijk}^{op})} \mathbf{B}^{op}(U_i) = \widetilde{\mathbf{A}}_!(U_i) \right\}_{k \in K_{ij}} \quad (5.32)$$

is a family of spectral immersions relative to \mathcal{N} such that the image of the induced morphism $\prod_{k \in K_{ij}} \widetilde{\mathbf{A}}_!(Y_{ijk}) \rightarrow \widetilde{\mathbf{A}}_!(U_i)$ is $\widetilde{\mathbf{A}}_!(R_{i,j})$. Hence, the image of the monomorphism

$$\widetilde{\mathbf{A}}_! \left(R_{i,j} \xrightarrow{r_{i,j}} U_i \times U_j \rightarrow U_i \right) = \left(\widetilde{\mathbf{A}}_!(R_{i,j}) \xrightarrow{\widetilde{\mathbf{A}}_!(r_{i,j})} \widetilde{\mathbf{A}}_!(U_i \times U_j) \cong \widetilde{\mathbf{A}}_!(U_i) \times \widetilde{\mathbf{A}}_!(U_j) \rightarrow \widetilde{\mathbf{A}}_!(U_i) \right) \quad (5.33)$$

is Zariski open relative to \mathcal{N} . Since \mathcal{N} is subcanonical, it follows from Proposition 4.6 that the monomorphism in (5.33) is a Zariski open immersion relative to \mathcal{N} . Using Proposition 4.14, it follows that $\widetilde{\mathbf{A}}_!(F)$ is an \mathcal{N} -scheme. This concludes the proof. \square

Corollary 5.4. *Let \mathcal{E} be a closed symmetric monoidal category which is both complete and cocomplete. Let $(\mathcal{P}, \boxtimes^{\mathcal{P}})$ (resp. $(\mathcal{Q}, \boxtimes^{\mathcal{Q}})$) be a left \mathcal{E} -module category which is both complete and cocomplete and such that $\boxtimes^{\mathcal{P}} : \mathcal{E} \times \mathcal{P} \rightarrow \mathcal{P}$ (resp. $\boxtimes^{\mathcal{Q}} : \mathcal{E} \times \mathcal{Q} \rightarrow \mathcal{Q}$) preserves colimits in both variables. Let $\mathbb{S} : \mathcal{Q} \rightarrow \mathcal{P}$ be a lax \mathcal{E} -linear functor which is conservative and preserves finite limits. Further, suppose that for every \mathcal{P} -flat morphism $\alpha : a \rightarrow b$ in $Comm(\mathcal{E})$ and $n \in \mathbf{Mod}_{\mathcal{Q}}(a)$, the canonical morphism $\theta_{\alpha, id_{\mathcal{E}}, \mathbb{S}, n} : b \boxtimes_a^{\mathcal{P}} \mathbb{S}(n) \rightarrow \mathbb{S}(b \boxtimes_a^{\mathcal{Q}} n)$ is an isomorphism in $\mathbf{Mod}_{\mathcal{P}}(b)$.*

If \mathcal{Q} is subcanonical, then so is \mathcal{P} and for each $F \in Sch(\mathcal{E})_{\mathcal{P}} \subset PSh(Aff_{\mathcal{E}})$, the sheafification $(F)_{\mathcal{Q}}^{++}$ of F with respect to the spectral \mathcal{Q} -topology is a \mathcal{Q} -scheme.

Proof. The result follows from an application of Theorem 5.3 to the adjoint pair $(id_{\mathcal{E}}, id_{\mathcal{E}})$. \square

6 Examples

In this section, we give a number of examples of our theory.

Example 6.1. Let $(\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete. Then \mathcal{C} may be treated as a left \mathcal{C} -module category. It follows from [32, Corollary 2.11] that representable presheaves on $Aff_{\mathcal{C}}$ are also fpqc sheaves, which means that \mathcal{C} is subcanonical as a \mathcal{C} -module category in the sense of Section 4. It is also clear that schemes over \mathcal{C} treated as a left \mathcal{C} -module category are precisely the schemes relative to $(\mathcal{C}, \otimes, 1)$ in the sense of Toën and Vaquié [32].

Example 6.2. Let $(\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete. Let $n \in \mathbb{N}$. We note that the product category $\mathcal{C}^n = \underbrace{\mathcal{C} \times \dots \times \mathcal{C}}_{n \text{ -times}}$ is both complete and cocomplete and has a left \mathcal{C} -module

category structure $\overset{n}{\otimes} : \mathcal{C} \times \mathcal{C}^n \longrightarrow \mathcal{C}^n$ given by

$$c \overset{n}{\otimes} (a_1, \dots, a_n) := (c \otimes a_1, \dots, c \otimes a_n), \quad c \in \mathcal{C}, (a_1, \dots, a_n) \in \mathcal{C}^n \quad (6.1)$$

Since $(\mathcal{C}, \otimes, 1)$ is closed, it is easy to see that $\overset{n}{\otimes} : \mathcal{C} \times \mathcal{C}^n \longrightarrow \mathcal{C}^n$ preserves colimits in both variables. We will denote the left $(\mathcal{C}, \otimes, 1)$ -module category $(\mathcal{C}^n, \overset{n}{\otimes})$ simply by \mathcal{C}^n . For each $a \in \text{Comm}(\mathcal{C})$, the category $\mathbf{Mod}_{\mathcal{C}^n}(a) \cong (\mathbf{Mod}_{\mathcal{C}}(a))^n$. Further, for each morphism $\alpha : a \longrightarrow b$ in $\text{Comm}(\mathcal{C})$, it may be verified that the extension of scalars $b \overset{n}{\otimes}_a _ : \mathbf{Mod}_{\mathcal{C}^n}(a) \longrightarrow \mathbf{Mod}_{\mathcal{C}^n}(b)$ associated to \mathcal{C}^n is given by,

$$\begin{aligned} b \overset{n}{\otimes}_a _ : \mathbf{Mod}_{\mathcal{C}^n}(a) &\cong (\mathbf{Mod}_{\mathcal{C}}(a))^n \longrightarrow (\mathbf{Mod}_{\mathcal{C}}(b))^n \cong \mathbf{Mod}_{\mathcal{C}^n}(b) \\ (m_1, m_2, \dots, m_n) &\mapsto (b \otimes_a m_1, b \otimes_a m_2, \dots, b \otimes_a m_n) \end{aligned} \quad (6.2)$$

It follows that $\alpha : a \longrightarrow b$ is \mathcal{C}^n -flat if and only if it is \mathcal{C} -flat. Using (6.2), it may be verified that fpqc (resp. spectral) \mathcal{C}^n -covers are the same as fpqc (resp. spectral) \mathcal{C} -covers. Hence, it follows from Example 6.1 that \mathcal{C}^n is subcanonical. Further, \mathcal{C}^n -schemes are the same as \mathcal{C} -schemes.

Example 6.3. More generally, let $(\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete. We fix a set J , considered as a discrete category. It is clear that the functor category $[J, \mathcal{C}] = \prod_J \mathcal{C}$ is both complete and cocomplete. We note that $[J, \mathcal{C}]$ has a canonical left $(\mathcal{C}, \otimes, 1)$ -module category structure $\overset{J}{\otimes} : \mathcal{C} \times [J, \mathcal{C}] \longrightarrow [J, \mathcal{C}]$ defined as follows : for each $c \in \mathcal{C}$ and $M \in [J, \mathcal{C}]$

$$c \overset{J}{\otimes} M : J \longrightarrow \mathcal{C}, \quad (J \ni j \mapsto c \otimes M(j) \in \mathcal{C}) \quad (6.3)$$

Since colimits in functor categories are computed objectwise and $\otimes : \mathcal{C} \times \mathcal{C} \longrightarrow \mathcal{C}$ preserves colimits in both variables, hence so does the bifunctor $\overset{J}{\otimes} : \mathcal{C} \times [J, \mathcal{C}] \longrightarrow [J, \mathcal{C}]$ given by (6.3). We will denote the left $(\mathcal{C}, \otimes, 1)$ -module category $([J, \mathcal{C}], \overset{J}{\otimes})$ simply by $[J, \mathcal{C}]$.

It is clear that for each $a \in \text{Comm}(\mathcal{C})$, $\mathbf{Mod}_{[J, \mathcal{C}]}(a) \cong [J, \mathbf{Mod}_{\mathcal{C}}(a)]$. Further, for each morphism $\alpha : a \longrightarrow b$ in $\text{Comm}(\mathcal{C})$, it may be verified that the extension of scalars $b \overset{J}{\otimes}_a _ : \mathbf{Mod}_{[J, \mathcal{C}]}(a) \longrightarrow \mathbf{Mod}_{[J, \mathcal{C}]}(b)$ associated to $[J, \mathcal{C}]$ is the composition

$$\mathbf{Mod}_{[J, \mathcal{C}]}(a) \cong [J, \mathbf{Mod}_{\mathcal{C}}(a)] \xrightarrow{\alpha^* \circ _} [J, \mathbf{Mod}_{\mathcal{C}}(b)] \cong \mathbf{Mod}_{[J, \mathcal{C}]}(b) \quad (6.4)$$

where $\alpha^* = b \otimes_a _ : \mathbf{Mod}_{\mathcal{C}}(a) \longrightarrow \mathbf{Mod}_{\mathcal{C}}(b)$ is the extension of scalars associated to \mathcal{C} . Since limits in functor categories are computed objectwise, it follows that $\alpha : a \longrightarrow b$ in $\text{Comm}(\mathcal{C})$ is $[J, \mathcal{C}]$ -flat if and only if it is \mathcal{C} -flat. Using (6.4), it may also be verified that fpqc (resp. spectral) $[J, \mathcal{C}]$ -covers are the same as fpqc (resp. spectral) \mathcal{C} -covers in $\text{Aff}_{\mathcal{C}}$. It again follows from Example 6.1 that $[J, \mathcal{C}]$ is subcanonical. Further, $[J, \mathcal{C}]$ -schemes are the same as \mathcal{C} -schemes.

Remark 6.4. Let (\mathcal{M}, \boxtimes) be a subcanonical left $(\mathcal{C}, \otimes, 1)$ -module category. Using the same arguments as that of Example 6.3, it may be verified that the bifunctor $\overset{J}{\boxtimes} : \mathcal{C} \times [J, \mathcal{M}] \longrightarrow [J, \mathcal{M}]$ defined by

$$c \overset{J}{\boxtimes} M : J \longrightarrow \mathcal{M}, \quad (J \ni j \mapsto c \boxtimes M(j) \in \mathcal{M}) \quad (c \in \mathcal{C} \text{ and } M \in [J, \mathcal{M}]) \quad (6.5)$$

makes $[J, \mathcal{M}]$ into a subcanonical $(\mathcal{C}, \otimes, 1)$ -module category. Further, $[J, \mathcal{M}]$ -schemes are the same as \mathcal{M} -schemes.

In what follows, we will denote the category of presheaves of sets on an arbitrary small category \mathcal{Z} by $\widehat{\mathcal{Z}} := PSh(\mathcal{Z})$. We note that the category $\widehat{\mathcal{Z}}$ is both complete and cocomplete. For $F, G \in \widehat{\mathcal{Z}}$, let $F \bullet G \in \widehat{\mathcal{Z}}$ denote their product in $PSh(\mathcal{Z})$. We recall (see for instance, [22, § 1.6]) that $(\widehat{\mathcal{Z}}, \bullet, 1)$ is a cartesian closed monoidal category where $1 \in \widehat{\mathcal{Z}}$ denotes the terminal object. It is clear from Example 6.1 that the left $(\widehat{\mathcal{Z}}, \bullet, 1)$ -module category $(\widehat{\mathcal{Z}}, \bullet)$ is subcanonical.

Proposition 6.5. *Let $\Phi : \mathcal{X} \rightarrow \mathcal{Y}$ be an essentially surjective functor between small categories. Then, the bifunctor*

$$\boxtimes : \widehat{\mathcal{Y}} \times \widehat{\mathcal{X}} \rightarrow \widehat{\mathcal{X}}, \quad (F, H) \mapsto ((F \circ \Phi^{op}) \bullet H) \in \widehat{\mathcal{X}} \quad (6.6)$$

preserves colimits in both variables and makes $\widehat{\mathcal{X}}$ into a left $(\widehat{\mathcal{Y}}, \bullet, 1)$ -module category. Further, $(\widehat{\mathcal{X}}, \boxtimes)$ is subcanonical and for any $(\widehat{\mathcal{X}}, \boxtimes)$ -scheme F on $Aff_{(\widehat{\mathcal{Y}}, \bullet, 1)}$, the sheafification $(F)_{(\widehat{\mathcal{Y}}, \bullet)}^{++}$ of F with respect to the spectral $(\widehat{\mathcal{Y}}, \bullet)$ -topology on $Aff_{(\widehat{\mathcal{Y}}, \bullet, 1)}$ is a $(\widehat{\mathcal{Y}}, \bullet)$ -scheme.

Proof. We consider the functor

$$\widehat{\Phi} : \widehat{\mathcal{Y}} = PSh(\mathcal{Y}) \rightarrow PSh(\mathcal{X}) = \widehat{\mathcal{X}}, \quad \widehat{\mathcal{Y}} \ni F \mapsto F \circ \Phi^{op} \in \widehat{\mathcal{X}} \quad (6.7)$$

Since limits and colimits in presheaf categories are computed objectwise, we note that $\widehat{\Phi} = _ \circ \Phi^{op}$ preserves limits and colimits. In particular, $\widehat{\Phi}$ preserves finite products so that there is a natural isomorphism

$$\delta = \left(\delta_{F,G} : \widehat{\Phi}(F) \bullet \widehat{\Phi}(G) \xrightarrow{\sim} \widehat{\Phi}(F \bullet G) \mid F, G \in \widehat{\mathcal{Y}} \right) \quad (6.8)$$

It may now be verified that the bifunctor

$$\boxtimes : \widehat{\mathcal{Y}} \times \widehat{\mathcal{X}} \rightarrow \widehat{\mathcal{X}}, \quad (F, H) \mapsto \left(\widehat{\Phi}(F) \bullet H \right) = ((F \circ \Phi^{op}) \bullet H) \in \widehat{\mathcal{X}} \quad (6.9)$$

preserves colimits in both variables and makes $\widehat{\mathcal{X}}$ into a left $(\widehat{\mathcal{Y}}, \bullet, 1)$ -module category. We need to show that $(\widehat{\mathcal{X}}, \boxtimes)$ is subcanonical. We note that the natural isomorphism in (6.8)

$$\left(F \boxtimes \widehat{\Phi}(G) = \widehat{\Phi}(F) \bullet \widehat{\Phi}(G) \xrightarrow{\delta_{F,G}} \widehat{\Phi}(F \bullet G) \mid F, G \in \widehat{\mathcal{Y}} \right) \quad (6.10)$$

makes $\widehat{\Phi} : (\widehat{\mathcal{Y}}, \bullet) \rightarrow (\widehat{\mathcal{X}}, \boxtimes)$ into a strong $(\widehat{\mathcal{Y}}, \bullet, 1)$ -linear functor. Since $\widehat{\Phi}$ preserves coequalizers, it follows from Lemma 5.2(2) (with $\mathcal{C} = \mathcal{D} = (\widehat{\mathcal{Y}}, \bullet, 1)$, $\mathbf{B} = \mathbf{A} = id_{\widehat{\mathcal{Y}}}$, $(\mathcal{M}, \boxtimes) = (\widehat{\mathcal{X}}, \boxtimes)$, $(\mathcal{N}, \boxplus) = (\widehat{\mathcal{Y}}, \bullet)$ and $(\mathbb{L}, \Gamma) = (\widehat{\Phi}, \delta)$) that for any morphism $\alpha : F \rightarrow G$ in $Comm(\widehat{\mathcal{Y}})$ and any $H \in \mathbf{Mod}_{\widehat{\mathcal{Y}}}(F)$, the canonical morphism

$$\theta_{\alpha, id_{\widehat{\mathcal{Y}}}, \widehat{\Phi}, H} : G \boxtimes_F \widehat{\Phi}(H) \rightarrow \widehat{\Phi}(G \bullet_F H) \quad (6.11)$$

is an isomorphism in $\mathbf{Mod}_{\widehat{\mathcal{X}}}(G)$. Since $\Phi : \mathcal{X} \rightarrow \mathcal{Y}$ is essentially surjective, it follows that $\widehat{\Phi}$ is conservative. Using Corollary 5.4 (with $\mathcal{E} = \widehat{\mathcal{Y}}$, $(\mathcal{P}, \boxtimes^{\mathcal{P}}) = (\widehat{\mathcal{X}}, \boxtimes)$, $(\mathcal{Q}, \boxtimes^{\mathcal{Q}}) = (\widehat{\mathcal{Y}}, \bullet)$ and $\mathbb{S} = (\widehat{\Phi}, \delta)$) and the fact that $(\widehat{\mathcal{Y}}, \bullet)$ is subcanonical as a $\widehat{\mathcal{Y}}$ -module category, the result follows. \square

Let A be an ordinary monoid with identity element e , considered as a one-object category. We recall that the category $PSh(A)$ is the category $\mathbf{Rep}(A)$ of representations (right actions) of A on sets. An object of $\mathbf{Rep}(A)$ is a pair (U, ρ) , where U is a set and $\rho : U \times A \rightarrow U$ is the action $u.a := \rho(u, a)$.

Example 6.6. Let $\sigma : M \rightarrow N$ be a morphism of monoids. Then, σ is an essentially surjective functor between the one-object categories associated to M and N . Using Proposition 6.5, it follows that the bifunctor $\boxtimes : \mathbf{Rep}(N) \times \mathbf{Rep}(M) \rightarrow \mathbf{Rep}(M)$ given by

$$(S, \rho_S) \boxtimes (T, \rho_T) = (S \times T, \rho) \in \mathbf{Rep}(M), \quad (S, \rho_S) \in \mathbf{Rep}(N), (T, \rho_T) \in \mathbf{Rep}(M) \quad (6.12)$$

where

$$\rho : (S \times T) \times M \longrightarrow (S \times T), \quad ((s, t), m) \mapsto (\rho_S(s, \sigma(m)), \rho_T(t, m)) = (s \cdot \sigma(m), t \cdot m) \in S \times T \quad (6.13)$$

preserves colimits in both variables and makes $\mathbf{Rep}(M)$ into a subcanonical left $(\mathbf{Rep}(N), \bullet, 1)$ -module category. Further, for any $F \in Sch(\mathbf{Rep}(N))_{\mathbf{Rep}(M)}$, the sheafification $(F)_{\mathbf{Rep}(N)}^{++}$ of F with respect to the spectral $(\mathbf{Rep}(N), \bullet)$ -topology on $Aff_{\mathbf{Rep}(N)}$ is a $\mathbf{Rep}(N)$ -scheme.

Example 6.7. We recall (see for instance, [21, § 2.7]) that a directed graph G is a tuple (V, E, s, t) where V and E are sets and $s : E \longrightarrow V$ (resp. $t : E \longrightarrow V$) is a function, called source (resp. target). A morphism $G = (V, E, s, t) \longrightarrow G' = (V', E', s', t')$ of directed graphs is a pair

$$(f : V \longrightarrow V', g : E \longrightarrow E') \quad (6.14)$$

of functions such that $s' \circ g = f \circ s$ and $t' \circ g = f \circ t$. The category of directed graphs and graph morphisms is denoted by $Digrph$.

Let M be a monoid and $m, n \in M$. We consider the bifunctor

$$\overset{(m,n)}{\boxtimes} : \mathbf{Rep}(M) \times Digrph \longrightarrow Digrph, \quad ((U, \rho), (V, E, s, t)) \mapsto (U \times V, U \times E, s', t') \quad (6.15)$$

where

$$\begin{aligned} s' : U \times E &\longrightarrow U \times V, & (u, a) &\mapsto (\rho(u, m), s(a)) = (u \cdot m, s(a)) \\ t' : U \times E &\longrightarrow U \times V, & (u, a) &\mapsto (\rho(u, n), t(a)) = (u \cdot n, t(a)) \end{aligned} \quad (6.16)$$

We claim that $(Digrph, \overset{(m,n)}{\boxtimes})$ is a subcanonical left $\mathbf{Rep}(M)$ -module category. To see this, we consider the category

$$\mathcal{X} = \left(\begin{array}{ccc} & d_0 & \\ 0 & \xrightarrow{\quad} & 1 \\ & d_1 & \end{array} \right) \quad (6.17)$$

with exactly two objects $0, 1$ and two non-identity arrows d_0 and d_1 . It may be verified that the functor

$$PSh(\mathcal{X}) = \widehat{\mathcal{X}} \longrightarrow Digrph, \quad \widehat{\mathcal{X}} \ni G \mapsto (G(0), G(1), G(d_0^{op}), G(d_1^{op})) \in Digrph \quad (6.18)$$

is an isomorphism of categories. Let \mathcal{Y} be the one-object category associated to the monoid M . Let $\Phi_{m,n} : \mathcal{X} \longrightarrow \mathcal{Y}$ be the functor which sends 0 and 1 to the unique object of \mathcal{Y} and the arrow d_0 (resp. d_1) to m (resp. n). It may be verified that the functor $\widehat{\Phi_{m,n}} = _ \circ \Phi_{m,n}^{op} : PSh(\mathcal{Y}) \longrightarrow PSh(\mathcal{X})$ is given by

$$_ \circ \Phi_{m,n}^{op} : PSh(\mathcal{Y}) = \mathbf{Rep}(M) \longrightarrow Digrph \cong PSh(\mathcal{X}), \quad (U, \rho) \mapsto (U, U, s_U, t_U) \quad (6.19)$$

where

$$\begin{aligned} s_U : U &\longrightarrow U, & u &\mapsto \rho(u, m) = u \cdot m \\ t_U : U &\longrightarrow U, & u &\mapsto \rho(u, n) = u \cdot n \end{aligned} \quad (6.20)$$

Using (6.15), it follows that for $(U, \rho) \in \mathbf{Rep}(M)$ and $G \in Digrph$

$$\widehat{\Phi_{m,n}}(U, \rho) \bullet G = (U, \rho) \overset{(m,n)}{\boxtimes} G \quad (6.21)$$

where $\bullet : Digrph \times Digrph \longrightarrow Digrph$ is the cartesian monoidal structure on $Digrph \cong PSh(\mathcal{X})$. Since $\Phi_{m,n} : \mathcal{X} \longrightarrow \mathcal{Y}$ is essentially surjective, it follows from Proposition 6.5 that $(Digrph, \overset{(m,n)}{\boxtimes})$ is a subcanonical left $\mathbf{Rep}(M)$ -module category.

Example 6.8. In particular, let M be the monoid with exactly one element e . It is clear that the category $\mathbf{Rep}(M) = PSh(M)$ is the cartesian monoidal category Set . It follows from Example 6.7 that the bifunctor

$$\overset{(e,e)}{\boxtimes} : Set \times Digrph \longrightarrow Digrph, \quad (S, G) \mapsto (S, S, 1_S, 1_S) \bullet G \in Digrph \quad (6.22)$$

makes $Digrph$ into a subcanonical left Set -module category.

7 Descent in terms of Simplicial \mathcal{M} -Schemes

Throughout this section, let $\mathcal{C} = (\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete. We let $\mathcal{M} = (\mathcal{M}, \boxtimes)$ be a subcanonical left \mathcal{C} -module category which is both complete and cocomplete and such that $\boxtimes : \mathcal{C} \times \mathcal{M} \longrightarrow \mathcal{M}$ preserves colimits in both variables.

We introduce some notation. Let $f : F \longrightarrow S$ be a morphism between \mathcal{M} -schemes. Using Proposition 4.12, we note that $f : F \longrightarrow S$ induces a base change functor $\widehat{f} : Sch(\mathcal{C})_{\mathcal{M}}/S \longrightarrow Sch(\mathcal{C})_{\mathcal{M}}/F$, $(H \longrightarrow S) \mapsto (H \times_S F \longrightarrow F)$. Let $n \in \mathbb{N}$. Using Proposition 4.12, we can consider the limit $\underbrace{F \times_S \dots \times_S F}_{n\text{-times}}$ in $Sch(\mathcal{C})_{\mathcal{M}}$ and we denote the projection to the i -th component by $\pi_i^n : \underbrace{F \times_S \dots \times_S F}_{n\text{-times}} \longrightarrow F$ for $0 \leq i \leq n-1$.

It follows from the universal property of $F \times_S F$ that for each $0 \leq i < j \leq 2$, there is a morphism $p_{ij} : F \times_S F \times_S F \longrightarrow F \times_S F$ induced by the morphisms $\pi_i^3, \pi_j^3 : F \times_S F \times_S F \longrightarrow F$. It may be verified that $\pi_1^2 \circ p_{02} = \pi_1^2 \circ p_{12}$, $\pi_0^2 \circ p_{02} = \pi_0^2 \circ p_{01}$ and $\pi_1^2 \circ p_{01} = \pi_0^2 \circ p_{12}$. In a manner similar to [27, Tag023U], we consider the following definition.

Definition 7.1. A descent datum relative to a morphism $f : F \longrightarrow S$ of \mathcal{M} -schemes is a pair (V, ϕ) where $V \in Sch(\mathcal{C})_{\mathcal{M}}/F$ and

$$\phi : \widehat{\pi_0^2}(V) = V \times_F (F \times_S F) \cong V \times_S F \xrightarrow{\sim} F \times_S V \cong (F \times_S F) \times_F V = \widehat{\pi_1^2}(V) \quad (7.1)$$

is an isomorphism in $Sch(\mathcal{C})_{\mathcal{M}}/(F \times_S F)$ such that the following diagram in $Sch(\mathcal{C})_{\mathcal{M}}/(F \times_S F \times_S F)$ is commutative

$$\begin{array}{ccc} \widehat{p_{01}}(\widehat{\pi_0^2}(V)) \cong \widehat{p_{02}}(\widehat{\pi_0^2}(V)) & \xrightarrow{\widehat{p_{02}}(\phi)} & \widehat{p_{02}}(\widehat{\pi_1^2}(V)) \cong \widehat{p_{12}}(\widehat{\pi_1^2}(V)) \\ & \searrow \widehat{p_{01}}(\phi) \quad \nearrow \widehat{p_{12}}(\phi) & \\ & \widehat{p_{01}}(\widehat{\pi_1^2}(V)) \cong \widehat{p_{12}}(\widehat{\pi_0^2}(V)) & \end{array} \quad (7.2)$$

A morphism $g : (V, \phi) \longrightarrow (V', \phi')$ of descent data relative to $f : F \longrightarrow S$ is a morphism $g : V \longrightarrow V'$ in $Sch(\mathcal{C})_{\mathcal{M}}/F$ such that the following square in $Sch(\mathcal{C})_{\mathcal{M}}/(F \times_S F)$ is commutative :

$$\begin{array}{ccc} \widehat{\pi_0^2}(V) \cong V \times_S F & \xrightarrow{\phi} & F \times_S V \cong \widehat{\pi_1^2}(V) \\ \widehat{\pi_0^2}(g) \downarrow & & \downarrow \widehat{\pi_1^2}(g) \\ \widehat{\pi_0^2}(V') \cong V' \times_S F & \xrightarrow{\phi'} & F \times_S V' \cong \widehat{\pi_1^2}(V') \end{array} \quad (7.3)$$

The category of descent data relative to $f : F \longrightarrow S$ will be denoted by $Desc(F \xrightarrow{f} S)$.

Remark 7.2. It may be verified that if $diag : F \longrightarrow F \times_S F$ is the diagonal morphism, then for $(V, \phi) \in Desc(F \longrightarrow S)$, $\widehat{diag}(\phi) : \widehat{diag}(\pi_0^2(V)) \cong V \longrightarrow \widehat{diag}(\pi_1^2(V)) \cong V$ is the identity morphism 1_V .

We let Δ be the standard simplex category (see for instance [20, § 1]) with objects $[n] := \{0, 1, \dots, n\}$, $n \geq 0$. A simplicial \mathcal{M} -scheme is a functor $\mathbb{F} : \Delta^{op} \rightarrow \text{Sch}(\mathcal{C})_{\mathcal{M}}$. This consists of a sequence of \mathcal{M} -schemes $\mathbb{F}_0, \mathbb{F}_1, \mathbb{F}_2, \dots$ along with face morphisms $d_j^n = d_j^n(\mathbb{F}) : \mathbb{F}_n \rightarrow \mathbb{F}_{n-1}$ (resp. degeneracy morphisms $s_j^n = s_j^n(\mathbb{F}) : \mathbb{F}_n \rightarrow \mathbb{F}_{n+1}$) of \mathcal{M} -schemes for each $0 \leq j \leq n$, satisfying the standard simplicial identities. We denote by $SSch(\mathcal{C})_{\mathcal{M}}$ the category of simplicial \mathcal{M} -schemes.

For a simplicial \mathcal{M} -scheme \mathbb{F} , let $\mathcal{D}(\mathbb{F})$ denote the category of pairs (V, ϕ) where $V \in \text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_0$ and

$$\phi : \widehat{d}_1^1(V) = V \times_{\mathbb{F}_0, d_1^1} \mathbb{F}_1 \xrightarrow{\sim} \mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} V = \widehat{d}_0^1(V) \quad (7.4)$$

is an isomorphism in $\text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_1$ such that $\widehat{s}_0^0(\phi) = 1_V$ and such that $\widehat{d}_1^2(\phi) = \widehat{d}_0^2(\phi) \circ \widehat{d}_2^2(\phi)$ in $\text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_2$. A morphism $g : (V, \phi) \rightarrow (V', \phi')$ in $\mathcal{D}(\mathbb{F})$ is a morphism $g : V \rightarrow V'$ in $\text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_0$ such that the following square in $\text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_1$ is commutative

$$\begin{array}{ccc} \widehat{d}_1^1(V) = V \times_{\mathbb{F}_0, d_1^1} \mathbb{F}_1 & \xrightarrow{\phi} & \mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} V = \widehat{d}_0^1(V) \\ \widehat{d}_1^1(g) \downarrow & & \downarrow \widehat{d}_0^1(g) \\ \widehat{d}_1^1(V') = V' \times_{\mathbb{F}_0, d_1^1} \mathbb{F}_1 & \xrightarrow{\phi'} & \mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} V' = \widehat{d}_0^1(V') \end{array} \quad (7.5)$$

Let $\alpha : \mathbb{G} \rightarrow \mathbb{F}$ be a morphism in $SSch(\mathcal{C})_{\mathcal{M}}$. Similar to [27, Tag 0249], we say that α is cartesian if for every morphism $\tau : [n] \rightarrow [m]$ in Δ , the following commutative diagram is a pullback square in $\text{Sch}(\mathcal{C})_{\mathcal{M}}$

$$\begin{array}{ccc} \mathbb{G}_m & \xrightarrow{\alpha_m} & \mathbb{F}_m \\ \mathbb{G}(\tau) \downarrow & & \downarrow \mathbb{F}(\tau) \\ \mathbb{G}_n & \xrightarrow{\alpha_n} & \mathbb{F}_n \end{array} \quad (7.6)$$

We say that \mathbb{G} is cartesian over \mathbb{F} if there is a cartesian morphism $\mathbb{G} \rightarrow \mathbb{F}$ of simplicial \mathcal{M} -schemes. For a simplicial \mathcal{M} -scheme \mathbb{F} , we denote by $\text{Cart}(\mathbb{F})$, the subcategory of $SSch(\mathcal{C})_{\mathcal{M}}/\mathbb{F}$ whose objects are cartesian morphisms $\mathbb{G} \rightarrow \mathbb{F}$ and whose morphisms are cartesian morphisms $\mathbb{G} \rightarrow \mathbb{H}$ in $SSch(\mathcal{C})_{\mathcal{M}}/\mathbb{F}$.

Lemma 7.3. *Let $\mathbb{F} \rightarrow \mathbb{G}$ and $\mathbb{G} \rightarrow \mathbb{H}$ be morphisms of simplicial \mathcal{M} -schemes. If $\mathbb{G} \rightarrow \mathbb{H}$ is cartesian, then the composition $\mathbb{F} \rightarrow \mathbb{G} \rightarrow \mathbb{H}$ is cartesian if and only if $\mathbb{F} \rightarrow \mathbb{G}$ is cartesian.*

Proof. The result is clear from general properties of pullback squares. \square

It follows from Lemma 7.3 that if \mathbb{F} is a simplicial \mathcal{M} -scheme, then $\text{Cart}(\mathbb{F})$ is a full subcategory of $SSch(\mathcal{C})_{\mathcal{M}}/\mathbb{F}$.

Lemma 7.4. *Let \mathbb{F} be a simplicial \mathcal{M} -scheme. Then, the category $\text{Cart}(\mathbb{F})$ is equivalent to the category $\mathcal{D}(\mathbb{F})$.*

Proof. Let $f : \mathbb{G} \rightarrow \mathbb{F}$ be an object in $\text{Cart}(\mathbb{F})$. Since f is cartesian, the following are pullback squares in $\text{Sch}(\mathcal{C})_{\mathcal{M}}$.

$$\begin{array}{ccc} \mathbb{G}_1 & \xrightarrow{f_1} & \mathbb{F}_1 \\ d_1^1(\mathbb{G}) \downarrow & & \downarrow d_1^1(\mathbb{F}) \\ \mathbb{G}_0 & \xrightarrow{f_0} & \mathbb{F}_0 \end{array} \quad \begin{array}{ccc} \mathbb{G}_1 & \xrightarrow{f_1} & \mathbb{F}_1 \\ d_0^1(\mathbb{G}) \downarrow & & \downarrow d_0^1(\mathbb{F}) \\ \mathbb{G}_0 & \xrightarrow{f_0} & \mathbb{F}_0 \end{array} \quad (7.7)$$

We define $\phi : G_0 \times_{\mathbb{F}_0, d_1^1} \mathbb{F}_1 = \mathbb{G}_1 \rightarrow \mathbb{G}_1 = \mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} \mathbb{G}_0$ to be the identity morphism $1_{\mathbb{G}_1}$. It follows from general properties of pullback squares that we have a functor $\Theta : \text{Cart}(\mathbb{F}) \rightarrow \mathcal{D}(\mathbb{F})$, $(\mathbb{G} \xrightarrow{f} \mathbb{F}) \mapsto (\mathbb{G}_0 \xrightarrow{f_0} \mathbb{F}_0, \mathbb{G}_1 \xrightarrow{1_{\mathbb{G}_1}} \mathbb{G}_1)$. We claim that Θ is an equivalence of categories.

For any $0 \leq i \leq j \leq k$, let $\tau_i^k : [0] \rightarrow [k]$ be given by $0 \mapsto i$ and $\tau_{ij}^k : [1] \rightarrow [k]$ be given by $0 \mapsto i, 1 \mapsto j$. Suppose that $(V, \phi) \in \mathcal{D}(\mathbb{F})$. We define $\mathbb{G}^{(V, \phi)} : \Delta^{op} \rightarrow \text{Sch}(\mathcal{C})_{\mathcal{M}}$ by $\mathbb{G}_n^{(V, \phi)} := \mathbb{F}_n \times_{\mathbb{F}(\tau_n^n), \mathbb{F}_0} V$. Let $\lambda : [n] \rightarrow [m]$ be a morphism in Δ . We note that

$$d_0^1 \circ \mathbb{F}(\tau_{\lambda(n)m}^m) = \mathbb{F}(\tau_m^m) \quad d_1^1 \circ \mathbb{F}(\tau_{\lambda(n)m}^m) = \mathbb{F}(\tau_{\lambda(n)}^m) \quad \lambda \circ \tau_n^n = \tau_{\lambda(n)}^m \quad (7.8)$$

Using (7.8), we have

$$\begin{aligned} \mathbb{F}_m \times_{\mathbb{F}(\tau_{\lambda(n)m}^m), \mathbb{F}_1} (\mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} V) &\cong \mathbb{F}_m \times_{\mathbb{F}(\tau_m^m), \mathbb{F}_0} V = \mathbb{G}_m^{(V, \phi)} \\ \mathbb{F}_m \times_{\mathbb{F}(\tau_{\lambda(n)m}^m), \mathbb{F}_1} (\mathbb{F}_1 \times_{d_1^1, \mathbb{F}_0} V) &\cong \mathbb{F}_m \times_{\mathbb{F}(\tau_{\lambda(n)}^m), \mathbb{F}_0} V = \mathbb{F}_m \times_{\mathbb{F}(\lambda), \mathbb{F}_n} \mathbb{G}_n^{(V, \phi)} \end{aligned} \quad (7.9)$$

Since $\phi : \widehat{d}_1^1(V) \xrightarrow{\sim} \widehat{d}_0^1(V)$ is an isomorphism in $\text{Sch}(\mathcal{C})_{\mathcal{M}}/\mathbb{F}_1$, we may define $\mathbb{G}(\lambda) : \mathbb{G}_m^{(V, \phi)} \rightarrow \mathbb{G}_n^{(V, \phi)}$ as the composition

$$\mathbb{G}_m^{(V, \phi)} \cong \mathbb{F}_m \times_{\mathbb{F}(\tau_{\lambda(n)m}^m), \mathbb{F}_1} (\mathbb{F}_1 \times_{d_0^1, \mathbb{F}_0} V) \xrightarrow{\widehat{\mathbb{F}(\tau_{\lambda(n)m}^m)(\phi)}^{-1}} \mathbb{F}_m \times_{\mathbb{F}(\tau_{\lambda(n)m}^m), \mathbb{F}_1} (\mathbb{F}_1 \times_{d_1^1, \mathbb{F}_0} V) \cong \mathbb{F}_m \times_{\mathbb{F}(\lambda), \mathbb{F}_n} \mathbb{G}_n^{(V, \phi)} \rightarrow \mathbb{G}_n^{(V, \phi)} \quad (7.10)$$

where $\mathbb{F}_m \times_{\mathbb{F}(\lambda), \mathbb{F}_n} \mathbb{G}_n^{(V, \phi)} \rightarrow \mathbb{G}_m^{(V, \phi)}$ is the canonical morphism associated to $\mathbb{F}_m \times_{\mathbb{F}(\lambda), \mathbb{F}_n} \mathbb{G}_n^{(V, \phi)}$. Using properties of ϕ , it may be verified that $\mathbb{G}^{(V, \phi)}$ is a simplicial \mathcal{M} -scheme and that the family $f^{(V, \phi)} = \left(\mathbb{G}_n^{(V, \phi)} = \mathbb{F}_n \times_{\mathbb{F}(\tau_n^n), \mathbb{F}_0} V \rightarrow \mathbb{F}_n \right)_{[n] \in \Delta^{op}}$ of canonical morphisms is a cartesian morphism from $\mathbb{G}^{(V, \phi)}$ to \mathbb{F} . We now have a functor

$$\Omega : \mathcal{D}(\mathbb{F}) \rightarrow \text{Cart}(\mathbb{F}), \quad (V, \phi) \mapsto (\mathbb{G}^{(V, \phi)} \xrightarrow{f^{(V, \phi)}} \mathbb{F}) \quad (7.11)$$

It may be verified that Ω is a quasi-inverse for $\Theta : \text{Cart}(\mathbb{F}) \rightarrow \mathcal{D}(\mathbb{F})$. \square

Definition 7.5. Let $f : F \rightarrow S$ be a morphism of \mathcal{M} -schemes. The simplicial \mathcal{M} -scheme associated to f is the functor

$$\mathbb{X}(f) : \Delta^{op} \rightarrow \text{Sch}(\mathcal{C})_{\mathcal{M}}, \quad [n] \mapsto \underbrace{F \times_S \dots \times_S F}_{(n+1) \text{ times}} \quad \Delta([n], [m]) \ni \lambda \mapsto (\mathbb{X}(f))(\lambda) \quad (7.12)$$

where $(\mathbb{X}(f))(\lambda) : \underbrace{F \times_S \dots \times_S F}_{(m+1) \text{ times}} \rightarrow \underbrace{F \times_S \dots \times_S F}_{(n+1) \text{ times}}$ is induced by the family $\left\{ \underbrace{F \times_S \dots \times_S F}_{(m+1) \text{ times}} \xrightarrow{\pi_{\lambda(i)}^{m+1}} F : i \in [n] \right\}$.

Proposition 7.6. For a morphism $f : F \rightarrow S$ of \mathcal{M} -schemes, the categories $\text{Cart}(\mathbb{X}(f))$ and $\text{Desc}(F \xrightarrow{f} S)$ are equivalent.

Proof. We note that for the simplicial \mathcal{M} -scheme $\mathbb{X}(f)$, $d_1^1 = \pi_0^2$ and $d_0^1 = \pi_1^2$ as morphisms $\mathbb{X}(f)_1 = F \times_S F \rightarrow F = \mathbb{X}(f)_0$. Similarly, $d_1^2 = p_{02}, d_0^2 = p_{12}$ and $d_2^2 = p_{01}$ as morphisms $\mathbb{X}(f)_2 = F \times_S F \times_S F \rightarrow F \times_S F = \mathbb{X}(f)_1$. Further, $s_0^0 : \mathbb{X}(f)_0 = F \rightarrow \mathbb{X}(f)_1 = F \times_S F$ is the diagonal morphism *diag*. It follows from Definition 7.1 and Remark 7.2 that $\mathcal{D}(\mathbb{X}(f)) = \text{Desc}(F \xrightarrow{f} S)$ as categories. The result is now clear from Lemma 7.4. \square

8 Gluing Schemes over an adjunction

In [9], Connes and Consani present a notion of scheme over a gluing together of the category of commutative monoids with that of commutative rings, using the usual adjunction. Accordingly, a scheme over a “field with one element” in the sense of [9, Definition 4.7] consists of a usual scheme over $\text{Spec}(\mathbb{Z})$, a scheme over the category of commutative monoids, and a compatibility condition. In this final section, we present a counterpart of their construction.

Throughout this section, we let $(\mathcal{C}, \otimes, 1)$ be a closed symmetric monoidal category which is both complete and cocomplete and has a zero object $\mathbf{0}$. Since \mathcal{C} is closed, for each $c \in \mathcal{C}$, we have $\mathbf{0} \otimes c \cong \mathbf{0} \cong c \otimes \mathbf{0}$. For each $c, c' \in \mathcal{C}$, we denote by $0_{c,c'}$, the zero morphism $c \rightarrow \mathbf{0} \rightarrow c'$. We note that composing and tensoring with zero morphisms gives zero morphisms. Because \mathcal{C} has a zero object, it is clear that kernels and cokernels exist in \mathcal{C} .

Let $CMon_0 \subset Comm(Set)$ denote the category whose objects are commutative monoids $(M, \bullet, e, 0)$ with identity element e and an absorbing element 0 and whose morphisms are monoid homomorphisms that preserve the absorbing element. Let $*$ denote the singleton set. We recall (see for instance, [7, § 6.4]) that the natural transformation

$$\xi = (\xi_{a,b} : \mathcal{C}(1, a) \times \mathcal{C}(1, b) \rightarrow \mathcal{C}(1, a \otimes b), (f, g) \mapsto f \otimes g)_{(a,b) \in \mathcal{C} \times \mathcal{C}} \quad (8.1)$$

and the map $*$ $\rightarrow \mathcal{C}(1, 1)$ choosing the identity morphism id_1 form a lax symmetric monoidal structure on the functor $\mathcal{C}(1, _) : \mathcal{C} \rightarrow Set$. Hence, there is an induced functor

$$\mathcal{C}(1, _) : Comm(\mathcal{C}) \rightarrow CMon_0 \subset Comm(Set), \quad a \mapsto \mathcal{C}(1, a) \quad (8.2)$$

In detail, for each $(a, \mu_a : a \otimes a \rightarrow a, \iota_a : 1 \rightarrow a) \in Comm(\mathcal{C})$, the hom-set $\mathcal{C}(1, a)$ has the following structure

$$\iota_a \in \mathcal{C}(1, a), \quad 0_{1,a} \in \mathcal{C}(1, a), \quad \mathcal{C}(1, a) \times \mathcal{C}(1, a) \xrightarrow{\xi_{a,a}} \mathcal{C}(1, a \otimes a) \xrightarrow{\mu_a \circ -} \mathcal{C}(1, a) \quad (8.3)$$

which makes it into a commutative monoid with absorbing element $0_{1,a}$.

Lemma 8.1. *Let $(M, \bullet, e, 0) \in CMon_0$. For each $m \in M$, let $i_m^M : 1 \rightarrow \coprod_M 1$ be the m^{th} -coprojection in \mathcal{C} . Then, the following object in \mathcal{C}*

$$\mathcal{C}[M] := Coker(i_0) = Coeq \left(1 \begin{array}{c} \xrightarrow{i_0^M} \\ \xrightarrow{0_{1, \coprod_M 1}} \end{array} \coprod_M 1 \right) \quad (8.4)$$

has the structure of a commutative monoid object in \mathcal{C} .

Proof. Since $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ preserves colimits in both variables, hence there is a natural isomorphism

$$\left(\left(\coprod_A 1 \right) \otimes \left(\coprod_B 1 \right) \right) \xrightarrow{\sim} \coprod_{A \times B} 1 \mid A, B \in Set \quad (8.5)$$

We recall (see for instance, [7, § 6.4]) that along with the isomorphism $1 \xrightarrow{id_1} 1 \cong \coprod_* 1$, the isomorphisms in (8.5) give a strong symmetric monoidal structure on the functor $\coprod_{(_)} 1 : Set \rightarrow \mathcal{C}$, $A \mapsto \coprod_A 1$. Since $(M, \bullet, e, 0) \in CMon_0$, $\coprod_M 1$ has the structure of a commutative monoid object in \mathcal{C} , given by

$$\begin{aligned} \mu^M : \left(\coprod_M 1 \right) \otimes \left(\coprod_M 1 \right) &\cong \coprod_{M \times M} 1 \xrightarrow{\coprod \bullet} \coprod_M 1 \\ i_e^M : 1 &\rightarrow \coprod_M 1 \end{aligned} \quad (8.6)$$

Since $0 \in M$ is an absorbing element, there is a commutative diagram in Set

$$\begin{array}{ccc} M \cong M \times * & \xrightarrow{id_M \times 0} & M \times M \\ \downarrow & & \bullet \downarrow \\ * & \xrightarrow{0} & M \end{array} \quad (8.7)$$

By applying $\coprod_{(_) } 1 : \text{Set} \rightarrow \mathcal{C}$ on (8.7) and using the naturality of (8.5), we get the following commutative diagram in \mathcal{C}

$$\begin{array}{ccc} \coprod_{M \times *} 1 \cong (\coprod_M 1) \otimes 1 & \xrightarrow{id \otimes i_0^M} & (\coprod_M 1) \otimes (\coprod_M 1) \cong \coprod_{M \times M} 1 \\ \downarrow & & \downarrow \mu^M \\ 1 & \xrightarrow{i_0^M} & \coprod_M 1 \end{array} \quad (8.8)$$

It is clear that the morphism $(\coprod_M 1) \otimes 1 \rightarrow 1$ in (8.8) makes the unit object 1 into a left $\coprod_M 1$ -module such that $i_0^M : 1 \rightarrow \coprod_M 1$ is a morphism of left $\coprod_M 1$ -modules. Using Remark 2.2, it follows that $\mathcal{C}[M]$ in (8.4) is the cokernel of $i_0^M : 1 \rightarrow \coprod_M 1$ in $\mathbf{Mod}_{\mathcal{C}}(\coprod_M 1)$. Hence, there is a morphism $\rho^M : (\coprod_M 1) \otimes \mathcal{C}[M] \rightarrow \mathcal{C}[M]$ in \mathcal{C} making $\mathcal{C}[M]$ into a left $\coprod_M 1$ -module. Moreover, we have

$$\rho^M \circ (id \otimes coker(i_0^M)) = coker(i_0^M) \circ \mu^M \quad (8.9)$$

where $coker(i_0^M) : \coprod_M 1 \rightarrow \mathcal{C}[M]$ is the canonical morphism in \mathcal{C} (and hence in $\mathbf{Mod}_{\mathcal{C}}(\coprod_M 1)$) associated to the cokernel in (8.4). Since $\coprod_{\mathcal{C}} \mathcal{C}[M] : \mathcal{C} \rightarrow \mathcal{C}$ preserves colimits, $\mathcal{C}[M] \otimes \mathcal{C}[M] \cong Coker(i_0^M \otimes id_{\mathcal{C}[M]})$ in \mathcal{C} . Using (8.9), it may be verified that the composition

$$1 \otimes \mathcal{C}[M] \xrightarrow{i_0^M \otimes id_{\mathcal{C}[M]}} \left(\coprod_M 1 \right) \otimes \mathcal{C}[M] \xrightarrow{\rho^M} \mathcal{C}[M] \quad (8.10)$$

in \mathcal{C} is the zero morphism. Hence, it follows from the universal property of the cokernel $\mathcal{C}[M] \otimes \mathcal{C}[M] \cong Coker(i_0^M \otimes id_{\mathcal{C}[M]})$ that there is a unique morphism $\theta^M : \mathcal{C}[M] \otimes \mathcal{C}[M] \rightarrow \mathcal{C}[M]$ such that the following triangle in \mathcal{C}

$$\begin{array}{ccc} (\coprod_M 1) \otimes \mathcal{C}[M] & \xrightarrow{coker(i_0^M) \otimes id_{\mathcal{C}[M]}} & \mathcal{C}[M] \otimes \mathcal{C}[M] \\ \rho^M \downarrow & \swarrow \theta^M & \\ \mathcal{C}[M] & & \end{array} \quad (8.11)$$

is commutative. Using (8.6), (8.9) and (8.11), it may be verified that the morphisms $\theta^M : \mathcal{C}[M] \otimes \mathcal{C}[M] \rightarrow \mathcal{C}[M]$ and $1 \xrightarrow{i_e} \coprod_M 1 \xrightarrow{coker(i_0^M)} \mathcal{C}[M]$ make $\mathcal{C}[M]$ into a commutative monoid in \mathcal{C} such that $coker(i_0^M) : \coprod_M 1 \rightarrow \mathcal{C}[M]$ is a morphism in $Comm(\mathcal{C})$. \square

Proposition 8.2. *The functor $\mathcal{C}(1, _) : Comm(\mathcal{C}) \rightarrow CMon_0$ has a left adjoint $\mathcal{C}[_] : CMon_0 \rightarrow Comm(\mathcal{C})$, given on objects by $(M, \bullet, e, 0) \mapsto \left(\mathcal{C}[M], \theta^M, 1 \xrightarrow{i_e^M} \coprod_M 1 \xrightarrow{coker(i_0^M)} \mathcal{C}[M] \right)$.*

Proof. Using [7, Proposition 6.7.2], it suffices to check that for every $M = (M, \bullet, e, 0) \in CMon_0$, there is a bijection

$$Comm(\mathcal{C})(\mathcal{C}[M], a) \xrightarrow{\sim} CMon_0(M, \mathcal{C}(1, a)) \quad (8.12)$$

which is natural in $a = (a, \mu_a, \iota_a) \in Comm(\mathcal{C})$. Let $a \in Comm(\mathcal{C})$ and $\alpha : \mathcal{C}[M] \rightarrow a$ be a morphism in $Comm(\mathcal{C})$. Since $coker(i_0^M) : \coprod_M 1 \rightarrow \mathcal{C}[M]$ is a morphism in $Comm(\mathcal{C})$, $\alpha \circ coker(i_0^M)$ is a morphism in $Comm(\mathcal{C})$. Hence, using (8.3) and (8.6), it follows that the map

$$\hat{\alpha} : M \rightarrow \mathcal{C}(1, a), \quad m \mapsto \alpha \circ coker(i_0^M) \circ i_m^M \quad (8.13)$$

is a monoid homomorphism. We note that $\hat{\alpha}(0) = \alpha \circ coker(i_0^M) \circ i_0^M = 0_{1,a}$ and hence, $\hat{\alpha}$ is a morphism in $CMon_0$.

Conversely, given a morphism $\beta : M \rightarrow \mathcal{C}(1, a)$ in $CMon_0$, it may be verified that the induced morphism $\coprod_{m \in M} \beta(m) : \coprod_M 1 \rightarrow a$ is in $Comm(\mathcal{C})$. We note that $(\coprod_{m \in M} \beta(m)) \circ i_0^M = \beta(0) = 0_{1,a}$ in \mathcal{C} . It follows from (8.4) that there is a unique morphism

$$\tilde{\beta} : \mathcal{C}[M] \rightarrow a, \quad \text{such that} \quad \coprod_{m \in M} \beta(m) = \tilde{\beta} \circ coker(i_0^M) \quad (8.14)$$

in \mathcal{C} . Hence, $\tilde{\beta} \circ coker(i_0^M) \circ i_e^M = \coprod_{m \in M} \beta(m) \circ i_e^M = \beta(e) = \iota_a$. Using the fact that $\coprod_{m \in M} \beta(m)$ is a morphism in $Comm(\mathcal{C})$, it may be verified that $\mu_a \circ (\beta \otimes \tilde{\beta}) = \tilde{\beta} \circ \theta^M$. Hence $\tilde{\beta} : \mathcal{C}[M] \rightarrow a$ is a morphism in $Comm(\mathcal{C})$.

Using (8.13) and (8.14), it is clear that the map $Comm(\mathcal{C})(\mathcal{C}[M], a) \rightarrow CMon_0(M, \mathcal{C}(1, a))$, $\alpha \mapsto \hat{\alpha}$ is a bijection with inverse map sending $\beta \mapsto \tilde{\beta}$. This completes the proof. \square

We now recall (see [9, § 4.1]) how to glue together categories using an adjunction. Let $(L : \mathcal{A} \rightarrow \mathcal{B}, R : \mathcal{B} \rightarrow \mathcal{A})$ be an adjunction between categories \mathcal{A} and \mathcal{B} . By definition, there is a natural isomorphism

$$\Phi = \left\{ \Phi_{a,b} : Hom_{\mathcal{B}}(La, b) \xrightarrow{\sim} Hom_{\mathcal{A}}(a, Rb) \mid a \in \mathcal{A}, b \in \mathcal{B} \right\} \quad (8.15)$$

One can form (see [9, § 4.1]) the category $\mathcal{A} \cup_{L,R} \mathcal{B}$ which has the following data :

- (1) $Ob(\mathcal{A} \cup_{L,R} \mathcal{B}) = Ob(\mathcal{A}) \amalg Ob(\mathcal{B})$.
- (2) For any $a \in \mathcal{A}, b \in \mathcal{B}$, $Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(b, a) := \emptyset$ and $Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(a, b) := Hom_{\mathcal{B}}(La, b) \cong Hom_{\mathcal{A}}(a, Rb)$. For $a, a' \in \mathcal{A}$ and $b, b' \in \mathcal{B}$, $Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(a', a) = \mathcal{A}(a', a)$ and $Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(b', b) = \mathcal{B}(b', b)$.
- Let $\circ_{\mathcal{A}}$ and $\circ_{\mathcal{B}}$ denote the compositions of \mathcal{A} and \mathcal{B} , respectively.
- (3) Let $a, a' \in \mathcal{A}, b, b' \in \mathcal{B}$ and $f \in Hom_{\mathcal{A}}(a', a)$, $g \in Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(a, b)$ and $h \in Hom_{\mathcal{B}}(b, b')$. One defines

$$\begin{aligned} g \circ f &:= g \circ_{\mathcal{B}} L(f) \in Hom_{\mathcal{B}}(La', b) = Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(a', b) \\ h \circ g &:= h \circ_{\mathcal{B}} g \in Hom_{\mathcal{B}}(La, b') = Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(a, b') \end{aligned} \quad (8.16)$$

We recall (see [9, Proposition 4.1]) that $j_{\mathcal{A}} : \mathcal{A} \hookrightarrow \mathcal{A} \cup_{L,R} \mathcal{B}$ and $j_{\mathcal{B}} : \mathcal{B} \hookrightarrow \mathcal{A} \cup_{L,R} \mathcal{B}$ are full subcategories of the category $\mathcal{A} \cup_{L,R} \mathcal{B}$. For every $b \in \mathcal{B}$, there is a canonical morphism $\delta_b \in Hom_{\mathcal{A} \cup_{L,R} \mathcal{B}}(Rb, b) = Hom_{\mathcal{B}}(LRb, b)$ given by,

$$\delta_b := \Phi_{Rb,b}^{-1}(1_{Rb}) \quad (8.17)$$

It is clear that $\delta = (\delta_b)_{b \in \mathcal{B}}$ is a natural transformation $j_{\mathcal{A}} \circ R \rightarrow j_{\mathcal{B}}$.

Using Proposition 8.2 and the above construction, we obtain the category $CMon_0 \cup_{\mathcal{C}[_], \mathcal{C}(1, _)} Comm(\mathcal{C})$ along with a natural transformation $\delta : j_{CMon_0} \circ \mathcal{C}(1, _) \rightarrow j_{Comm(\mathcal{C})}$ as in (8.17). We consider the following class

$$fld(\mathcal{C}) = \{c \in Comm(\mathcal{C}) : \mathcal{C}(1, c) \setminus \{0_{1,c}\} \text{ is the group of units of the monoid } \mathcal{C}(1, c) \text{ in (8.3)}\} \quad (8.18)$$

which may be seen as the collection of fields in \mathcal{C} .

Let (\mathcal{M}, \boxtimes) be a subcanonical left \mathcal{C} -module category where \mathcal{M} is both complete and cocomplete and $\boxtimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ preserves colimits in both variables. Motivated by the ideas in [9, § 4.3], we make the following definition.

Definition 8.3. An $(\mathcal{M}, CMon_0)$ -scheme is a functor $F : CMon_0 \cup_{\mathcal{C}[_], \mathcal{C}(1, _)} Comm(\mathcal{C}) \rightarrow Set$ such that

- (1) The restriction $F_{\mathcal{C}}$ of F to $Comm(\mathcal{C})$ is an \mathcal{M} -scheme.
- (2) The restriction \underline{F} of F to $CMon_0$ is a $CMon_0$ -scheme, in the sense of [9].
- (3) For each object $c \in fld(\mathcal{C})$, the map $F(\delta_c) : \underline{F}(\mathcal{C}(1, c)) \rightarrow F_{\mathcal{C}}(c)$ is a bijection of sets.

References

- [1] A. Banerjee, *Derived schemes and the field with one element*, J. Math. Pures Appl. (9) **97** (2012), no. 3, 189–203.
- [2] ———, *Schemes over symmetric monoidal categories and torsion theories*, J. Pure Appl. Algebra **220** (2016), no. 9, 3017–3047.
- [3] ———, *On Noetherian schemes over $(C, \otimes, 1)$ and the category of quasi-coherent sheaves*, Indiana Univ. Math. J. **70** (2021), no. 1, 81–119.
- [4] J. Bénabou, *Introduction to bicategories*, Reports of the Midwest Category Seminar, Lecture Notes in Math., No. 47, Springer, Berlin-New York, 1967, pp. 1–77.
- [5] J. Böckenhauer, D. E. Evans, and Y. Kawahigashi, *Chiral structure of modular invariants for subfactors*, Comm. Math. Phys. **210** (2000), no. 3, 733–784.
- [6] F. Borceux, *Handbook of categorical algebra. 1*, Encyclopedia of Mathematics and its Applications, vol. 50, Cambridge University Press, Cambridge, 1994. Basic category theory.
- [7] ———, *Handbook of categorical algebra. 2*, Encyclopedia of Mathematics and its Applications, vol. 51, Cambridge University Press, Cambridge, 1994. Categories and structures.
- [8] M. Capucci and B. Gavranović, *Actegories for the Working Amthematician*, arXiv:2203.16351 (2022).
- [9] A. Connes and C. Consani, *Schemes over \mathbb{F}_1 and zeta functions*, Compos. Math. **146** (2010), no. 6, 1383–1415.
- [10] ———, *On the notion of geometry over \mathbb{F}_1* , J. Algebraic Geom. **20** (2011), no. 3, 525–557.
- [11] ———, *Absolute algebra and Segal’s Γ -rings: au dessous de $\overline{\text{Spec}(\mathbb{Z})}$* , J. Number Theory **162** (2016), 518–551.
- [12] ———, *On absolute algebraic geometry the affine case*, Adv. Math. **390** (2021), Paper No. 107909, 44.
- [13] P. Deligne, *Catégories tannakiennes*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 111–195.
- [14] P. Etingof, D. Nikshych, and V. Ostrik, *On fusion categories*, Ann. of Math. (2) **162** (2005), no. 2, 581–642.
- [15] ———, *Fusion categories and homotopy theory*, Quantum Topol. **1** (2010), no. 3, 209–273. With an appendix by Ehud Meir.
- [16] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Mathematical Surveys and Monographs, vol. 205, American Mathematical Society, Providence, RI, 2015.
- [17] J. Fuchs and C. Schweigert, *Category theory for conformal boundary conditions*, Vertex operator algebras in mathematics and physics (Toronto, ON, 2000), Fields Inst. Commun., vol. 39, Amer. Math. Soc., Providence, RI, 2003, pp. 25–70.
- [18] M. Hakim, *Topos annelés et schémas relatifs*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. Band 64, Springer-Verlag, Berlin-New York, 1972.
- [19] G. Janelidze and G. M. Kelly, *A note on actions of a monoidal category*, Theory Appl. Categ. **9** (2001/02), 61–91. CT2000 Conference (Como).
- [20] M. Land, *Introduction to infinity-categories*, Compact Textbooks in Mathematics, Birkhäuser/Springer, Cham, 2021.
- [21] S. Mac Lane, *Categories for the working mathematician*, 2nd ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.
- [22] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic*, Universitext, Springer-Verlag, New York, 1994.
- [23] F. Marty, *Relative Zariski open objects*, J. K-Theory **10** (2012), no. 1, 9–39.
- [24] ———, *Smoothness in relative geometry*, J. K-Theory **12** (2013), no. 3, 461–491.
- [25] A. Ocneanu, *Paths on Coxeter diagrams: From platonic solids and singularities to minimal models and subfactors*, Lectures on Operator Theory, Fields Inst. Monogr., Amer. Math. Soc., 2000, pp. 243–323.
- [26] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups **8** (2003), no. 2, 177–206.
- [27] *The Stacks project*, stacks.math.columbia.edu.
- [28] R. Street, *Polynomials as spans*, Cah. Topol. Géom. Différ. Catég. **61** (2020), no. 2, 113–153.
- [29] K. Szlachányi, *Finite quantum groupoids and inclusions of finite type*, Mathematical physics in mathematics and physics (Siena, 2000), Fields Inst. Commun., vol. 30, Amer. Math. Soc., Providence, RI, 2001, pp. 393–407.
- [30] B. Toën and G. Vezzosi, *Homotopical algebraic geometry. I. Topos theory*, Adv. Math. **193** (2005), no. 2, 257–372.
- [31] ———, *Homotopical algebraic geometry. II. Geometric stacks and applications*, Mem. Amer. Math. Soc. **193** (2008), no. 902, x+224.
- [32] B. Toën and M. Vaquié, *Au-dessous de $\text{Spec } \mathbb{Z}$* , J. K-Theory **3** (2009), no. 3, 437–500.