

STABILITY AND SINGULAR SET OF THE TWO-CONVEX LEVEL SET FLOW

SIAO-HAO GUO

ABSTRACT. The level set flow of a mean-convex closed hypersurface is stable off singularities, in the sense that the level set flow of the perturbed hypersurface would be close in the smooth topology to the original flow wherever the latter is regular. To study the behavior near singularities, we further assume that the initial hypersurface is two-convex and that the flow has finitely many singular times. In this case, the singular set of the flow would have finitely many connected components, each of which is either a point or a compact C^1 embedded curve. Then under additional conditions, we show that near each connected component of the singular set of the original flow, the perturbed flow would have “the same type” of singular set as that of the singular component.

CONTENTS

1. Introduction	1
2. <u>Stability of mean-convex LSF off singularities</u>	5
2.1. Estimates for initial hypersurface	5
2.2. Classical theory of MCF	8
2.3. Interior estimates for mean-convex LSF	11
2.4. Smooth convergence off singularities	20
3. <u>Singular set of two-convex LSF</u>	25
3.1. Round points	25
3.2. Cylindrical points	26
3.3. Singular set at the first singular time	35
3.4. Singular set at subsequent singular times	36
4. <u>Types of singular components</u>	41
4.1. Vanishing type	43
4.2. Splitting type	44
4.3. Bumpy type	54
4.4. Appendix: Two-convexity	59
References	60

1. INTRODUCTION

Let Σ_0 be a closed connected (smoothly embedded) hypersurface in \mathbb{R}^n with $n \geq 3$. There exists a unique number $T_1 \in (0, \infty)$, called the first singular time, so that the mean curvature flow (MCF) $\{\Sigma_t\}_{t \in [0, T_1]}$ starting from Σ_0 at time 0

The research was partially supported by the grant 112-2115-M-002-016-MY3 of the National Science and Technology Council of Taiwan.

cannot be smoothly extended any further; in fact, the second fundamental form of Σ_t blows up as $t \nearrow T_1$ (see [H1]). The **level set flow** (LSF) $\{\Sigma_t\}_{t \in [0, \infty)}$ starting from Σ_0 at time 0 is a continuation of the MCF past the first singular time via the “level set method” (cf. [OS], [CGG], [ES]). Such a flow is uniquely determined by Σ_0 and agrees with the MCF for $t \in [0, T_1)$. The flow would vanish after a finite time; the unique number $T_{ext} \in (0, \infty)$ is called the “time of extinction” provided the flow does not vanish until after time T_{ext} .

In this paper we assume that Σ_0 is **mean-convex**, namely, its mean curvature is positive with respect to the inward unit normal vector. In this case, the LSF $\{\Sigma_t\}$ would stay in Ω_0 for $t > 0$, where Ω_0 is the open connected set bounded by Σ_0 (see the Jordan-Brouwer separation theorem in [GP]), and it would sweep out Ω_0 monotonically. Specifically, for each point $p \in \Omega_0$, the flow would pass through p at some moment and then it would never return. Let us denote by $u(p)$ the unique time when the flow $\{\Sigma_t\}$ passes through p .¹ As such, the LSF can be described as the level sets of the function u as follows:

$$(1.1) \quad \Sigma_t = \{x \in \bar{\Omega}_0 : u(x) = t\}, \quad t \geq 0.$$

Such a function u , called the **arrival time function** of the flow, is actually Lipschitz continuous on $\bar{\Omega}_0$; moreover, it is the unique viscosity solution² to the following Dirichlet problem (cf. [ES]):

$$(1.2) \quad \begin{aligned} - \left(\mathbf{I} - \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) \cdot \nabla^2 u &= 1 \quad \text{in } \Omega_0, \\ u &= 0 \quad \text{on } \partial\Omega_0 = \Sigma_0. \end{aligned}$$

In this case, the flow is called a **mean-convex LSF**. Note that the superlevel set

$$(1.3) \quad \Omega_t := \{x \in \Omega_0 : u(x) > t\}, \quad t > 0$$

is contracting in the sense that

$$(1.4) \quad t_1 < t_2 \Rightarrow \Omega_{t_1} \supset \bar{\Omega}_{t_2}.$$

Note also that

$$T_{ext} = \max_{\Omega_0} u.$$

A point $p \in \Omega_0$ is called a **regular point** of $\{\Sigma_t\}$ provided the flow is locally smooth and monotonic³ near the point p for t close to $u(p)$; by (1.1) and the implicit function theorem, this is equivalent to saying that the arrival time function u is locally smooth near p with $\nabla u(p) \neq 0$. In that case, u would satisfy (1.2) in the classical sense near p and $\{\Sigma_t\}$ would be a MCF near p for t close to $u(p)$, whose mean curvature with respect to the unit normal $\frac{\nabla u}{|\nabla u|}$ would be

$$(1.5) \quad H = -\nabla \cdot \frac{\nabla u}{|\nabla u|} = \frac{-1}{|\nabla u|} \left(\mathbf{I} - \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right) \cdot \nabla^2 u = \frac{1}{|\nabla u|};$$

in particular, $\{\Sigma_t\}$ is mean-convex with respect to $N = \frac{\nabla u}{|\nabla u|}$.

On the other hand, if a point $p \in \Omega_0$ is not a regular point of $\{\Sigma_t\}$, it would be called a **singular point** of the flow. The set of all singular points of $\{\Sigma_t\}$, denoted by \mathcal{S} , is of Hausdorff dimension at most $n - 2$ (cf. [W1]). As $\{\mathcal{H}^{n-1} \lfloor \Sigma_t\}$

¹When $p \in \partial\Omega_0 = \Sigma_0$, set $u(p) = 0$.

²See [ES] for the precise definition of viscosity solutions to equation (1.2).

³That is, $\partial_t x \cdot N > 0$ for some choice of locally smooth unit normal vector field N of $\{\Sigma_t\}$.

is a Brakke flow (cf. [I1], [W1]), we may consider the tangent flows at the singular points (cf. [I2]). It turns out that the tangent flows at each singular point is unique; it is either a shrinking sphere or a shrinking k -cylinder⁴ (cf. [W2], [W4], [CM2]). Following [G2], singular points are called **round points** and **k -cylindrical points** if the tangent flows are shrinking spheres and shrinking k -cylinders, respectively.

By [CM4], at every singular point of $\{\Sigma_t\}$ the arrival time function u is twice differentiable and ∇u vanishes. As a corollary, singular points of $\{\Sigma_t\}$ are precisely the critical points of u , namely,

$$(1.6) \quad \mathcal{S} = \{x \in \Omega_0 : \nabla u(x) = 0\}.$$

By Lemma 2.10, the singular set \mathcal{S} is a compact set.

In this paper we would like to investigate the stability of mean-convex LSF. Based on the uniqueness theorem of viscosity solutions to the Dirichlet problem (1.2) in [ES] and also the smooth estimates for a mean-convex LSF in [HK], we have the following result:

Theorem 1.1. *Given a sequence of closed connected hypersurfaces $\{\Sigma_0^k\}_{k \in \mathbb{N}}$ tending in the C^4 topology to a mean-convex closed connected hypersurface Σ_0 , i.e.,*

$$\Sigma_0^k \xrightarrow{C^4} \Sigma_0 \quad \text{as } k \rightarrow \infty,$$

their respective LSFs $\{\Sigma_t^k\}$, $k \in \mathbb{N}$, would converge locally smoothly to the mean-convex LSF $\{\Sigma_t\}$ in $(\Omega_0 \setminus \mathcal{S}) \times (0, \infty)$ as $k \rightarrow \infty$.

Note that when k is large, Σ_0^k would also be mean-convex and so $\{\Sigma_t^k\}$ is a mean-convex LSF that will stay in Ω_0^k , where Ω_0^k is the region bounded by Σ_0^k . Proof of Theorem 1.1 will be given in Section 2.4.

As a consequence of Theorem 1.1, suppose that $\{\Sigma_t\}$ is regular for $t \in [a, b]$, where $0 < a < b < T_{ext}$; in other words, $\{\Sigma_t\}_{t \in [a, b]}$ is a mean-convex MCF. Then $\{\Sigma_t^k\}_{t \in [a, b]}$ would also be regular when k is large and converge smoothly to $\{\Sigma_t\}_{t \in [a, b]}$ as $k \rightarrow \infty$ (see Corollary 2.22). Notice that in the case where $b < T_1$, this is the well-known stability theorem for MCF. So what's new is when $a > T_1$.

What's more, when k is sufficiently large, Theorem 1.1 implies that the singular set \mathcal{S}^k of the mean-convex LSF $\{\Sigma_t^k\}$ would be contained in an arbitrarily small neighborhood of \mathcal{S} (see Corollary 2.19).

In order to compare the flows near \mathcal{S} , we need to work out the structure of the singular set first. To this end, in what follows we shall focus on the case where Σ_0 is **two-convex**,⁵ which is a condition stronger than the mean-convexity⁶ but weaker than the convexity. The motivation is that by [CHN] (see also [HK]), the flow $\{\Sigma_t\}$ would be uniformly two-convex at every regular point with respect to the unit normal $\frac{\nabla u}{|\nabla u|}$ (see (4.28)); consequently, every singular point of $\{\Sigma_t\}$ must be either a round point or a 1-cylindrical point, the latter of which will henceforth be abbreviated to a **cylindrical point**. Such a flow will be referred to as a **two-convex LSF**.

⁴That is, up to a rotation, the self-shrinking of $S^{\frac{n-k-1}{\sqrt{2(n-k-1)}}} \times \mathbb{R}^k$, where $k \in \{1, \dots, n-2\}$.

⁵ $\kappa_1 + \kappa_2 > 0$, where $\kappa_1 \leq \kappa_2 \leq \dots \leq \kappa_{n-1}$ denote the principal curvatures of a closed hypersurface with respect to the inward unit normal vector field.

⁶When $n = 3$, two-convex is just mean-convex.

Note that $t \in (0, T_{ext}]$ is called a singular time of the flow $\{\Sigma_t\}$ provided

$$\mathcal{S} \cap \Sigma_t \neq \emptyset;$$

by (1.6), a singular time of $\{\Sigma_t\}$ amounts to a critical value of u . Otherwise, t would be called a regular time of the flow, i.e., a regular value of u . It is conjectured in [CM4] that a mean-convex LSF has finitely many singular times (see [AAG] for the rotationally symmetric examples). Under the hypothesis of finitely many singular times, we prove the following theorem on the basis of [CM2] and [CM3]:

Theorem 1.2. *If the LSF $\{\Sigma_t\}$ starting from a two-convex closed connected hypersurface Σ_0 has finitely many singular times*

$$T_1 \leq \dots \leq T_{ext},$$

then its singular set \mathcal{S} is a finite disjoint union of points and/or compact C^1 embedded curves.

Note that by (1.6) and the mean value theorem, u is constant on each connected component of \mathcal{S} , meaning that singularities on each component of \mathcal{S} occur at the same time. Theorem 1.2 is related to a speculation in [CM3], which conjectures that the spacetime singular set has only finitely many connected components. The proof of Theorem 1.2 will be given in Section 3.4.

Recall that singular points of $\{\Sigma_t\}$ are critical points of u . In general, critical points of a function can be classified into local maximum points, local minimum points, and saddle points. In the case of the arrival time function u , there are no local minimum points (see (1.2)). Moreover, by Corollary 3.1 and Corollary 3.2, a point $p \in \Omega_0$ is a saddle point of u if and only if p is not only a cylindrical point of $\{\Sigma_t\}$ but also a boundary point of the superlevel set $\Omega_{u(p)}$ (see (1.3)). In that case, owing to the asymptotically cylindrical behavior of $\{\Sigma_t\}$ near p (see Section 3.2), for any small $\phi > 0$ there exists $r = r(\phi) > 0$ so that

$$(1.7) \quad \Omega_{u(p)} \cap B_r(p) \subset \mathcal{C}_\phi(p),$$

where $\mathcal{C}_\phi(p)$ is a double cone with vertex p , axis parallel to the tangent cylinder of $\{\Sigma_t\}$ at p , and angle ϕ (see (3.3)); hence, $\Omega_{u(p)}$ has a cusp singularity at p .

By contrast, if a point $p \in \Omega_0$ is a local maximum point of u , then $\Omega_{u(p)}$ vanishes near p . According to Remark 3.5, when a connected component of the singular set \mathcal{S} (see Theorem 1.2) is a curve, all its interior points must be local maximum points of u . Thus, whether the flow has the same kind of behavior near these curves depends crucially on their endpoints. We then classify the curves in Theorem 1.2 into the following three types:

- (1) **Vanishing type:** if either the curve has no endpoints (i.e., the curve is closed) or both of its endpoints are local maximum points of u ;
- (2) **Splitting type:** if both endpoints of the curve are saddle points of u ;
- (3) **Bumpy type:** if one endpoint of the curve is a local maximum point of u and the other is a saddle point of u .

When a connected component of \mathcal{S} is a point, it can be regarded as a “degenerate” curve and so it can also be classified into the above three categories (see the comment following Definition 4.1).

Now let us write

$$(1.8) \quad \mathcal{S} = \bigsqcup_j \mathcal{S}_j,$$

where each \mathcal{S}_j is a connected component of \mathcal{S} (called a **singular component** for short). Let $\hat{\delta} > 0$ be a sufficiently small constant such that (4.1), (4.2), and (4.3) hold. As mentioned earlier, when k is large, the singular set \mathcal{S}^k of the flow $\{\Sigma_t^k\}$ in Theorem 1.1 would satisfy

$$(1.9) \quad \mathcal{S}^k \subset \bigsqcup_j \mathcal{S}_j^{\hat{\delta}},$$

where $\mathcal{S}_j^{\hat{\delta}}$ is the $\hat{\delta}$ -neighborhood of \mathcal{S}_j . In addition, the singular times of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ would approximate to $u(\mathcal{S}_j)$ (see Proposition 2.16). In the following theorem we show that under certain conditions, the flow $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ would have the same type of singular set as \mathcal{S}_j .

Theorem 1.3. *Let $\{\Sigma_t\}$ be the two-convex LSF in Theorem 1.2 with singular set (1.8), and let $\{\Sigma_t^k\}$, $k \in \mathbb{N}$, be the LSFs in Theorem 1.1. Given a small constant $\hat{\delta} > 0$ fulfilling (4.1), (4.2), and (4.3), suppose that for every sufficiently large k the following hold:*

- (1) *For each j , there is at most one singular time of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ (see Assumption 4.3);*
- (2) *When the singular component \mathcal{S}_j is of the splitting type, the flow $\{\Sigma_t\}$ “does not get into the singular set near the endpoints” (see Assumption 4.12);*
- (3) *When the singular component \mathcal{S}_j is of the bumpy type, there are singularities of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ (see Assumption 4.16).*

Then there exists $k_{\hat{\delta}} \in \mathbb{N}$ so that for every $k \geq k_{\hat{\delta}}$, $\{\Sigma_t^k\}$ is a two-convex LSF with singular set (1.9); moreover, in each $\mathcal{S}_j^{\hat{\delta}}$, $\{\Sigma_t^k\}$ has precisely one singular component, which is of the same type as \mathcal{S}_j .

The proof of Theorem 1.3 will be divided into three parts according to the type of \mathcal{S}_j : Proposition 4.6 (in Section 4.1) is when \mathcal{S}_j belongs to the vanishing type, Proposition 4.14 (in Section 4.2) is when \mathcal{S}_j belongs to the splitting type, and Proposition 4.19 (in Section 4.3) is when \mathcal{S}_j belongs to the bumpy type.

2. STABILITY OF MEAN-CONVEX LSF OFF SINGULARITIES

The objective of this section is to establish Theorem 1.1. To accomplish that, we manage to obtain uniform estimates for the mean-convex LSFs away from singularities. This can be divided into two parts: the estimates near the boundary (i.e., near the initial time 0) in Section 2.2 using the classical theory of MCF, and the interior estimates at the regular points in Section 2.3 based on the noncollapsing property of mean-convex LSF (cf. [HK]). What we do in Section 2.1 is to give estimates of the noncollapsing constant and the entropy bound, which are essential in Section 2.3 and also Section 3.2 (for the cylindrical scales). Lastly, in Section 2.4 we employ these uniform estimates combined with the uniqueness theorem of viscosity solutions to the Dirichlet problem (1.2) to prove Theorem 1.1.

2.1. Estimates for initial hypersurface. We shall estimate the noncollapsing constant α in Section 2.1.1 and the entropy bound λ in Section 2.1.2 of the initial hypersurface Σ_0 . Note that the constants α and λ will be preserved by the mean-convex LSF due to the maximum principle and Huisken’s monotonicity formula, respectively (see the comments following Proposition 2.3 and Proposition 2.5).

2.1.1. *Noncollapsing.* Recall that Σ_0 is called α -**noncollapsing** for some $\alpha > 0$ provided for every $p \in \Sigma_0$, there exist a pair of open balls of radius $\alpha/H(p)$ on each side⁷ of Σ_0 that kiss at p (cf. [An], [ALM], [HK]). Such a constant α always exists and can be estimated as follows.

Lemma 2.1. *There exists a constant $\vartheta(n) \in (0, \frac{1}{2}]$ with the following property: If Σ is a hypersurface in \mathbb{R}^n such that $0 \in \Sigma$, $T_0\Sigma = \mathbb{R}^{n-1} \times \{0\}$ and that for some $r > 0$, $\Sigma \cap B_r$ is a graph*

$$x^n = f(x^1, \dots, x^{n-1})$$

over (some part of) $T_0\Sigma$ with

$$\max_{\Sigma \cap B_r} r |A| \leq 1,$$

where A is the second fundamental form of Σ . Then

$$B_+ := B_{\vartheta(n)r}(0, \vartheta(n)r), \quad B_- := B_{\vartheta(n)r}(0, -\vartheta(n)r)$$

are contained in B_r and located on the upside and downside of $\Sigma \cap B_r$, respectively.

Proof. Note that $f(0) = 0$ and $\nabla f(0) = 0$. Let $\rho \in (0, \frac{r}{2}]$ be a radius so that the local graph function f is defined on B_ρ^{n-1} with $|\nabla f| \leq \varepsilon$, where $\varepsilon \in (0, 1)$ is a small constant to be determined. With the local graph parametrization, the metric and the second fundamental form of Σ can be written as

$$g_{ij} = \delta_{ij} + \partial_i f \partial_j f, \quad A_{ij} = \frac{\partial_{ij} f}{\sqrt{1 + |\nabla f|^2}}.$$

Let

$$K = \max_{\Sigma \cap B_r} |A|.$$

Then it follows from

$$|A_{ij} v^i v^j| \leq K g_{ij} v^i v^j \quad \forall v \in \mathbb{R}^{n-1}$$

that on B_ρ^{n-1} we have for every $v \in \mathbb{R}^{n-1}$,

$$(2.1) \quad |\partial_{ij} f v^i v^j| \leq \sqrt{1 + |\nabla f|^2} K (\delta_{ij} + \partial_i f \partial_j f) v^i v^j \leq 2K |v|^2$$

provided $\varepsilon = \varepsilon(n) \in (0, 1)$ is sufficiently small. In particular, we obtain

$$|\nabla f| \leq \hat{C}(n) K \rho \quad \text{on } B_\rho^{n-1},$$

where $\hat{C}(n) \geq 1$ is a constant. As a result, we deduce that the ‘‘optimal’’ radius ρ in the above must be bounded below by

$$(2.2) \quad \hat{\rho} := \frac{\varepsilon(n)}{2\hat{C}(n)K} \geq \frac{\varepsilon(n)}{2\hat{C}(n)} r.$$

In addition, by Taylor’s theorem we have

$$f(x') = \int_0^1 \partial_{ij} f(sx') x^i x^j (1-s) ds$$

⁷One of the balls is on the inside, i.e., Ω_0 , and the other is on the outside, i.e., $\mathbb{R}^n \setminus \bar{\Omega}_0$, where Ω_0 is the region bounded by Σ_0 .

for $x' = (x^1, \dots, x^{n-1}) \in B_{\hat{\rho}}^{n-1}$. It follows from (2.1) that

$$(2.3) \quad |f(x')| \leq K |x'|^2 \leq \frac{1}{r} |x'|^2 \leq \frac{\hat{C}(n)}{\varepsilon(n)r} |x'|^2 \quad \forall x' \in B_{\hat{\rho}}^{n-1}.$$

Set

$$\vartheta(n) = \frac{\varepsilon(n)}{2\hat{C}(n)}.$$

By (2.2) and (2.3), the two open balls B_+ and B_- are contained in $B_r \cap \{|x'| < \vartheta(n)r\}$ and located on each side of the graph $x^n = f(x')$. \square

In order to apply Lemma 2.1 to every point of Σ_0 , we introduce the following definition:

Definition 2.2. For a closed hypersurface Σ , let \mathfrak{R} be the set of every $r \in (0, K^{-1}]$, where

$$K = \max_{\Sigma} |A|,$$

such that that for every $p \in \Sigma$, $\Sigma \cap B_r(p)$ is a graph over (some part of) $T_p \Sigma$. Note that $\mathfrak{R} \neq \emptyset$ due to the embeddedness and compactness of Σ . Then define

$$\text{rad } \Sigma = \sup \mathfrak{R}.$$

The following proposition follows from applying Lemma 2.1 to every point of Σ_0 .

Proposition 2.3. *Let ι, κ be positive constants such that*

$$\text{rad } \Sigma_0 \geq \iota, \quad \min_{\Sigma_0} H \geq \kappa.$$

Then there is a constant $\alpha = \alpha(n, \iota, \kappa) > 0$ such that Σ_0 is α -noncollapsing.

According to [HK], the α -noncollapsing property of Σ_0 can be preserved by the mean-convex LSF in the sense that near every regular point $p \in \Omega_0$, there exist two open balls of radius

$$\frac{\alpha}{H(p)} = \alpha |\nabla u(p)|$$

kissing at p , one of which is inside $\Omega_{u(p)}$ and the other is outside $\bar{\Omega}_{u(p)}$.

2.1.2. *Entropy.* The area ratios of Σ_0 can be estimated as follows.

Lemma 2.4. *Let $K, \mathfrak{A}, \mathfrak{D}$ be positive constants such that*

$$\max_{\Sigma_0} H \leq K, \quad \mathcal{H}^{n-1}(\Sigma_0) \leq \mathfrak{A}, \quad \text{diam } \Sigma_0 \leq \mathfrak{D}.$$

Then for every $p \in \mathbb{R}^n$ and $r > 0$ we have

$$\frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq 2^{n-1} e^{K(\mathfrak{D}+1)} \mathfrak{A}.$$

Proof. Set $R = \mathfrak{D} + 1$.

Case 1: For $p \in \Sigma_0$ and $r \in (0, R]$, by the monotonicity formula (cf. [Al]) we have

$$(2.4) \quad \frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq e^{K(R-r)} \frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_R(p))}{R^{n-1}} \leq e^{KR} \frac{\mathcal{H}^{n-1}(\Sigma_0)}{R^{n-1}}.$$

Case 2: For $p \in \Sigma_0$ and $r > R$, we have

$$\frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq \frac{\mathcal{H}^{n-1}(\Sigma_0)}{R^{n-1}}.$$

⁸The constant α can be chosen to be any number in $(0, \vartheta(n)\iota\kappa)$.

Case 3: For $p \notin \Sigma_0$ and $r \in (0, \frac{1}{2}R]$, either $\Sigma_0 \cap B_r(p) = \emptyset$, in which case we have $\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p)) = 0$, or we can find $q \in \Sigma_0 \cap B_r(p)$ so that (2.4) yields

$$\frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq \frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_{2r}(q))}{r^{n-1}} \leq 2^{n-1} e^{KR} \frac{\mathcal{H}^{n-1}(\Sigma_0)}{R^{n-1}}.$$

Case 4: For $p \notin \Sigma_0$ and $r > \frac{1}{2}R$, we have

$$\frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq \frac{\mathcal{H}^{n-1}(\Sigma_0)}{(\frac{1}{2}R)^{n-1}} = 2^{n-1} \frac{\mathcal{H}^{n-1}(\Sigma_0)}{R^{n-1}}.$$

Therefore, for every $p \in \mathbb{R}^n$ and $r > 0$ we have

$$\frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}} \leq 2^{n-1} e^{KR} \frac{\mathcal{H}^{n-1}(\Sigma_0)}{R^{n-1}} \leq 2^{n-1} e^{K(\mathfrak{D}+1)} \mathfrak{A}.$$

□

Recall that the entropy of Σ_0 is defined as (cf. [CM1])

$$E[\Sigma_0] = \sup_{p \in \mathbb{R}^n, r > 0} F\left(\frac{1}{r}(\Sigma_0 - p)\right),$$

where F is the Gaussian area, namely,

$$F(\Sigma) := \int_{\Sigma} \frac{1}{(4\pi)^{\frac{n-1}{2}}} e^{-\frac{|x|^2}{4}} d\mathcal{H}^{n-1}(x).$$

As the area ratios of Σ_0 are uniformly bounded, Σ_0 has finite entropy (cf. [E]); moreover, by a simple calculation we have

$$(2.5) \quad E[\Sigma_0] \leq C(n) \sup_{p \in \mathbb{R}^n, r > 0} \frac{\mathcal{H}^{n-1}(\Sigma_0 \cap B_r(p))}{r^{n-1}}.$$

Combining (2.5) with Lemma 2.4 give the following proposition.

Proposition 2.5. *Let $K, \mathfrak{A}, \mathfrak{D}$ be positive constants such that*

$$\sup_{\Sigma_0} H \leq K, \quad \mathcal{H}^{n-1}(\Sigma_0) \leq \mathfrak{A}, \quad \text{diam } \Sigma_0 \leq \mathfrak{D}.$$

Then there exists a constant $\lambda = \lambda(n, K, \mathfrak{A}, \mathfrak{D}) > 0$ so that $E[\Sigma_0] \leq \lambda$.

Since $\{\mathcal{H}^{n-1}[\Sigma_t]\}_{t \geq 0}$ is a Brakke flow (cf. [I1], [W1]), it follows from Huisken's monotonicity formula (cf. [H2], [I2]) that the entropy is nonincreasing along $\{\Sigma_t\}$; in particular,

$$E[\Sigma_t] \leq \lambda \quad \forall t \geq 0.$$

2.2. Classical theory of MCF. In this subsection we shall get smooth estimates for $\{\Sigma_t\}$ when t is close to 0, which corresponds to the estimates at points near $\partial\Omega_0 = \Sigma_0$. As the LSF $\{\Sigma_t\}$ is a MCF for $t \in [0, T_1)$, where T_1 is the first singular time, such estimates are primarily based on the classical theory of MCF. The main results are Proposition 2.6, Proposition 2.8, (2.15), and (2.16).

Recall that one of the ways to estimate the MCF near the initial time is to describe the flow as a evolving graph over the initial hypersurface and then study the graph function via the PDE it satisfies. Specifically, when t is sufficiently close to 0, Σ_t can be parametrized as a normal graph of $v_t = v(\cdot, t)$ over Σ_0 , namely,

$$(2.6) \quad x_t = x(\cdot, t) = x_0 + v_t N_0,$$

where x_0 and N_0 are the position vector and the inward unit normal vector of Σ_0 , respectively. Thus, $\{\Sigma_t\}$ being a MCF starting from Σ_0 at time 0 corresponds to that the function v satisfying a certain quasilinear parabolic PDE with $v(\cdot, 0) = 0$. More precisely, under the condition that

$$(2.7) \quad |\nabla v| + \frac{|v|}{\text{rad } \Sigma_0} \leq \varepsilon(n),$$

where ∇ is the Levi-Civita connection on Σ_0 , the norm $|\cdot|$ is defined by the induced metric g_0 on Σ_0 , and $\text{rad } \Sigma_0$ is defined in Definition 2.2, the PDE is given by

$$(2.8) \quad \partial_t v = H_0 + a^{ij} \nabla_{ij} v + |A_0|^2 v + b,$$

with

$$\begin{aligned} a^{ij} &= g_0^{ij} + 2A_0^{ij} v + Q^{ij}(\nabla v, A_0 v), \\ b &= v \nabla A_0 * \nabla v + A * Q(\nabla v, A_0 v) + v \nabla A_0 * Q(\nabla v, A_0 v) \end{aligned}$$

(see Lemma 3.14 in [G1]), where

- (1) H_0 is the mean curvature of Σ_0 ;
- (2) $\nabla_{ij} v$ is the Hessian of v on Σ_0 ;
- (3) the notation Q means some kind of analytic function/tensor that is at least “quadratic” (in the form of contraction via the metric g_0) in its arguments;
- (4) the notation $*$ means some kind of contraction of tensors.

The short time existence of MCF, the uniqueness of MCF, the stability of MCF, etc., all follow from studying (2.8) using the classical theory of parabolic PDE (cf. [HP]). The following proposition is one of the direct results.

Proposition 2.6. *Let n, ι, K, K', K'' be positive constants such that*

$$(2.9) \quad \text{rad } \Sigma_0 \geq \iota, \quad \max_{\Sigma_0} |A| \leq K, \quad \max_{\Sigma_0} |\nabla A| \leq K', \quad \max_{\Sigma_0} |\nabla^2 A| \leq K''.$$

Then there exists $\dot{T} = \dot{T}(n, \iota, K, K', K'') > 0$ so that $\{\Sigma_t\}_{0 \leq t \leq 2\dot{T}}$ is a MCF;⁹ moreover, for every $t \in [0, 2\dot{T}]$, the hypersurface Σ_t stays in a tubular neighborhood of Σ_0 and is a normal graph of $v(\cdot, t)$ over Σ_0 with $v \in C^\infty(\Sigma_0 \times [0, 2\dot{T}])$ satisfying (2.7), (2.8), and

$$(2.10) \quad |\nabla^2 v| \leq C(n, \iota, K, K', K'').$$

Furthermore, if $\{\Sigma_0^k\}_{k \in \mathbb{N}}$ is a sequence of closed connected hypersurfaces tending in the C^4 topology to Σ_0 as $k \rightarrow \infty$, then for every sufficiently large k the following hold:

- (1) *The LSF $\{\Sigma_t^k\}$ starting from Σ_0^k at time 0 is a MCF during $t \in [0, 2\dot{T}]$; in addition, the flow $\{\Sigma_t^k\}_{0 \leq t \leq 2\dot{T}}$ converges in the C^2 topology to $\{\Sigma_t\}_{0 \leq t \leq 2\dot{T}}$ as $k \rightarrow \infty$.*
- (2) *The initial hypersurface Σ_0^k is mean-convex;¹⁰ hence the associated arrival time function $u^k : \Omega_0^k \rightarrow [0, \infty)$ of $\{\Sigma_t^k\}$ is defined, where Ω_0^k is the open connected region bounded by Σ_0^k .*

By virtue of the smooth compactness of MCF, the sense of convergence in Proposition 2.6 can be improved away from the initial time as follows.

⁹In particular, $T_1 > 2\dot{T}$.

¹⁰When Σ_0 is two-convex, Σ_0^k would be two-convex when k is large (see Proposition 4.22).

Corollary 2.7. *In Proposition 2.6, for any given $\tau \in (0, \dot{T})$, the MCF $\{\Sigma_t^k\}_{t \in [\tau, 2\dot{T}]}$ would converge smoothly to $\{\Sigma_t\}_{t \in [\tau, 2\dot{T}]}$ as $k \rightarrow \infty$.*

Proof. Choose a large open ball \mathcal{B} in \mathbb{R}^n so that

$$\Sigma_t \subset \mathcal{B} \quad \forall t \in [0, 2\dot{T}].$$

By Proposition 2.6, there exists $k_0 \in \mathbb{N}$ so that for every $k \geq k_0$, the MCF $\{\Sigma_t^k\}_{t \in [0, 2\dot{T}]}$ satisfies

$$\begin{aligned} \Sigma_t^k &\subset \mathcal{B} \quad \forall 0 \leq t \leq 2\dot{T}, \\ \max_{0 \leq t \leq 2\dot{T}} \max_{\Sigma_t^k} |A| &\leq 2 \max_{0 \leq t \leq 2\dot{T}} \max_{\Sigma_t} |A|, \\ \mathcal{H}^n(\Sigma_0^k) &\leq 2 \mathcal{H}^n(\Sigma_0). \end{aligned}$$

Fix $\tau \in (0, \dot{T})$. It follows from the compactness theorem for MCF (cf. [H2]) that any subsequence of

$$\left\{ \left\{ \Sigma_t^k \right\}_{t \in [\tau, 2\dot{T}]} \right\}_{k \geq k_0}$$

has a smoothly convergent subsequence, whose limit by Proposition 2.6 is indeed $\{\Sigma_t\}_{t \in [\tau, 2\dot{T}]}$. Thus, the whole sequence $\left\{ \left\{ \Sigma_t^k \right\}_{t \in [\tau, 2\dot{T}]} \right\}_{k \geq k_0}$ must converge smoothly to $\{\Sigma_t\}_{t \in [\tau, 2\dot{T}]}$ as $k \rightarrow \infty$, completing the proof. \square

Due to the maximum principle, the mean-convexity is preserved by the MCF (cf. [M]) and hence $\{\Sigma_t\}_{0 \leq t \leq 2\dot{T}}$ moves monotonically toward the inside. Below we estimate how far the flow can be away from $\partial\Omega_0 = \Sigma_0$.

Proposition 2.8. *Let κ be a positive constant such that*

$$(2.11) \quad \min_{\Sigma_0} H \geq \kappa.$$

Then the normal graph of $v_t = v(\cdot, t)$ in Proposition 2.6 moves monotonically toward the inside with

$$\partial_t v \geq \kappa$$

on $\Sigma_0 \times [0, 2\dot{T}]$. In particular,

$$\text{dist}(\Omega_{\dot{T}}, \Sigma_0) \geq \kappa \dot{T}.$$

Proof. Along a MCF we have $\partial_t x \cdot N = H$ (cf. [M]). In the normal graph parametrization (2.6), this means that

$$\partial_t v N_0 \cdot N = H.$$

Then it follows from the maximum principle for the mean curvature along MCF that

$$(2.12) \quad |\partial_t v| \geq |\partial_t v N_0 \cdot N| = H \geq \kappa$$

on $\Sigma_0 \times [0, 2\dot{T}]$ (cf. [M]). In particular, $\partial_t v \neq 0$ everywhere on $\Sigma_0 \times [0, 2\dot{T}]$. On the other hand, since

$$\partial_t v(\cdot, 0) = H_0 \geq \kappa,$$

by the intermediate value theorem we deduce that $\partial_t v(\cdot, t) > 0$ for every $t \in [0, 2\dot{T}]$; hence by (2.12) we obtain $\partial_t v \geq \kappa$ on $\Sigma_0 \times [0, 2\dot{T}]$. \square

Note that Proposition 2.6 yields the estimates of the second fundamental form of $\{\Sigma_t\}_{0 \leq t \leq 2\dot{T}}$ and its covariant derivatives. More specifically, by Lemma 3.14 in [G1], under the condition (2.7), for every $t \in [0, 2\dot{T}]$ we have

$$(2.13) \quad \max_{\Sigma_t} |A| \leq C(n) \left(\max_{\Sigma_0} |A| + \max_{\Sigma_0} |\nabla^2 v(\cdot, t)| + \max_{\Sigma_0} |v(\cdot, t)| \max_{\Sigma_0} |\nabla A| \right)$$

Let κ be the constant in (2.11). Note that (2.7) implies

$$(2.14) \quad \frac{\kappa}{\sqrt{n-1}} |v| \leq \frac{1}{\sqrt{n-1}} H_0 |v| \leq |A_0 v| \leq \frac{|v|}{\text{rad } \Sigma_0} \leq \varepsilon(n).$$

Thus, (2.10), (2.13), and (2.14) yield

$$(2.15) \quad \max_{0 \leq t \leq 2\dot{T}} \max_{\Sigma_t} |A| \leq C(n, \iota, \kappa, K, K', K''),$$

where the notations are as defined in Proposition 2.6 and Proposition 2.8. Then it follows the maximum principle (for instance, see Proposition 6.2 in [G1]) that

$$\max_{\Sigma_t} |\nabla A| + \max_{\Sigma_t} |\nabla^2 A| \leq C \left(n, \max_{0 \leq \tau \leq 2\dot{T}} \max_{\Sigma_\tau} |A|, \max_{\Sigma_0} |\nabla A|, \max_{\Sigma_0} |\nabla^2 A| \right)$$

for $0 \leq t \leq 2\dot{T}$. So by (2.15) we obtain

$$(2.16) \quad \max_{0 \leq t \leq 2\dot{T}} \max_{\Sigma_t} |\nabla A| + \max_{0 \leq t \leq 2\dot{T}} \max_{\Sigma_t} |\nabla^2 A| \leq C(n, \iota, \kappa, K, K', K'').$$

2.3. Interior estimates for mean-convex LSF. In this subsection we shall obtain smooth estimates for $\{\Sigma_t\}$ at regular points that are away from $\partial\Omega_0 = \Sigma_0$. Such estimates are based on the noncollapsing property of mean-convex LSF (cf. [HK]), saying that the smooth scale (see Definition 2.11) at a regular point p is bounded below by, up to a multiplicative constant,

$$H(p)^{-1} = |\nabla u(p)|.$$

(see Proposition 2.12, Proposition 2.13, and Corollary 2.15). In the end of the subsection there is an estimate of the Hessian $\nabla^2 u$ (see Proposition 2.14), which is pivotal in getting the convergence of $\nabla u^{k's}$ ¹¹ in Section 2.4 (see Proposition 2.18).

To begin with, we give an estimate of $|\nabla u|$ from above in Proposition 2.9, which will be used in Section 2.4 to prove the convergence of u^k 's (see Proposition 2.16). In [ES] there is already a gradient estimate for u , which depends implicitly on Σ_0 . Here we improve the estimate such that it depends more explicitly on Σ_0 . Proposition 2.9 can also be regarded as a generalization of the maximum principle for the mean curvature along a mean-convex LSF.

Proposition 2.9. *Let κ be a positive constant such that*

$$\min_{\Sigma_0} H \geq \kappa.$$

Then the gradient of the arrival time function u satisfies

$$\sup_{\Omega_0} |\nabla u| \leq \kappa^{-1}.$$

As a consequence, at every regular point $p \in \Omega_0$ we have

$$H(p) = |\nabla u(p)|^{-1} \geq \kappa.$$

¹¹ u^k is the arrival time function of $\{\Sigma_t^k\}$, see Proposition 2.6.

Proof. By [ES], for every $\epsilon \in (0, 1]$ there is $u^\epsilon \in C^\infty(\bar{\Omega}_0)$ satisfying the ϵ -elliptic regularization of (1.2), that is,

$$(2.17) \quad - \left(\mathbf{I} - \frac{\nabla u^\epsilon \otimes \nabla u^\epsilon}{|\nabla u^\epsilon|^2 + \epsilon^2} \right) \cdot \nabla^2 u^\epsilon = 1 \quad \text{in } \Omega_0,$$

$$u^\epsilon = 0 \quad \text{on } \Sigma_0;$$

and as $\epsilon \searrow 0$

$$(2.18) \quad u^\epsilon(x) \xrightarrow{C^0} u(x) \quad \text{on } \bar{\Omega}_0.$$

Moreover, there exists a constant $C \geq 1$ such that for every $x \in \bar{\Omega}_0$,

$$(2.19) \quad \frac{1}{C} \text{dist}(x, \Sigma_0) \leq u^\epsilon(x) \leq C \text{dist}(x, \Sigma_0),$$

$$|\nabla u^\epsilon(x)| \leq C.$$

The geometric interpretation of the ϵ -elliptic regularization is that

$$(2.20) \quad \Gamma_t^\epsilon := \{(x, x^{n+1}) \in \bar{\Omega}_0 \times \mathbb{R} : x^{n+1} = \epsilon^{-1}u^\epsilon(x) - \epsilon^{-1}t\}, \quad t \geq 0$$

is a translating MCF in higher dimensional space, which moves downward (along the negative x^{n+1} -direction) with speed ϵ^{-1} ; in other words, $\Gamma_0^\epsilon = \{x^{n+1} = \epsilon^{-1}u^\epsilon(x)\}$ is a translating soliton. Note that its mean curvature with respect to the downward unit normal

$$(2.21) \quad \frac{(\nabla \epsilon^{-1}u^\epsilon, -1)}{\sqrt{|\nabla \epsilon^{-1}u^\epsilon|^2 + 1}} = \frac{(\nabla u^\epsilon, -\epsilon)}{\sqrt{|\nabla u^\epsilon|^2 + \epsilon^2}}$$

is

$$(2.22) \quad -\nabla \cdot \frac{\nabla \epsilon^{-1}u^\epsilon}{\sqrt{|\nabla \epsilon^{-1}u^\epsilon|^2 + 1}} = \frac{-1}{\sqrt{|\nabla \epsilon^{-1}u^\epsilon|^2 + 1}} \left(\mathbf{I} - \frac{\nabla \epsilon^{-1}u^\epsilon \otimes \nabla \epsilon^{-1}u^\epsilon}{|\nabla \epsilon^{-1}u^\epsilon|^2 + 1} \right) \cdot \nabla^2 \epsilon^{-1}u^\epsilon$$

$$= \frac{-1}{\sqrt{|\nabla u^\epsilon|^2 + \epsilon^2}} \left(\mathbf{I} - \frac{\nabla u^\epsilon \otimes \nabla u^\epsilon}{|\nabla u^\epsilon|^2 + \epsilon^2} \right) \cdot \nabla^2 u^\epsilon = \frac{1}{\sqrt{|\nabla u^\epsilon|^2 + \epsilon^2}}.$$

By (2.18), as $\epsilon \searrow 0$

$$(2.23) \quad u^\epsilon(x) - \epsilon x^{n+1} \xrightarrow{C_{loc}^0} u(x) \quad \text{on } \bar{\Omega}_0 \times \mathbb{R}_+,$$

meaning that the arrival time function of $\{\Gamma_t^\epsilon\}$ (which can be obtained by solving the equation in (2.20) for t) converges locally uniformly to the arrival time function of the ‘‘cylindrical’’ mean-convex LSF

$$(2.24) \quad \Gamma_t := \Sigma_t \times \mathbb{R}_+ = \{(x, x^{n+1}) \in \bar{\Omega}_0 \times \mathbb{R}_+ : u(x) = t\}, \quad t \geq 0.$$

In fact, the convergence is locally smooth for small $t > 0$ and away from $x^{n+1} = 0$ (as was mentioned in section 4.1 in [HK]). To see that, firstly let us get a uniform bound for the local area as follows. By (2.19), Proposition 2.8, coarea formula,

and the fact that the total area is non-increasing along MCF $\{\Sigma_t\}_{t \in [0, \dot{T}]}$,¹² for each $0 < \epsilon \leq \frac{\kappa}{2C} \dot{T}$ we have

$$\begin{aligned}
 & \mathcal{H}^n(\Gamma_0^\epsilon \cap \{0 < x^{n+1} < 2\}) \\
 &= \int_{0 < u^\epsilon < 2\epsilon} \sqrt{|\nabla \epsilon^{-1} u^\epsilon|^2 + 1} dx = \frac{1}{\epsilon} \int_{0 < u^\epsilon < 2\epsilon} \sqrt{|\nabla u^\epsilon|^2 + \epsilon^2} dx \\
 &\leq \frac{1}{\epsilon} \int_{0 < u^\epsilon < 2\epsilon} \sqrt{C^2 + \epsilon^2} dx \leq \frac{1}{\epsilon} \sqrt{C^2 + \epsilon^2} \mathcal{H}^n \{x \in \Omega_0 : 0 < \text{dist}(x, \Sigma_0) < 2C\epsilon\} \\
 &\leq \frac{1}{\epsilon} \sqrt{C^2 + \epsilon^2} \mathcal{H}^n \left\{ x \in \Omega_0 : 0 < u(x) < \frac{2C}{\kappa} \epsilon \right\} \\
 &= \frac{1}{\epsilon} \sqrt{C^2 + \epsilon^2} \int_0^{\frac{2C}{\kappa} \epsilon} \int_{\Sigma_t} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1} dt \\
 &\leq \frac{2C}{\kappa} \sqrt{C^2 + 1} K \mathcal{H}^{n-1}(\Sigma_0),
 \end{aligned}$$

where

$$K = \sup_{0 < u < \dot{T}} \frac{1}{|\nabla u|} = \max_{0 \leq u \leq \dot{T}} \max_{\Sigma_t} H.$$

Let $\delta \in (0, \frac{1}{2} \dot{T})$ be a sufficiently small constant (depending on n, \dot{T}). It follows from the local area bound (see Proposition 4.9 in [E]) and Brakke's compactness theorem (cf. [I1]) that for some sequence $\epsilon_i \searrow 0$, the MCF $\{\Gamma_t^{\epsilon_i}\}_{t \in (\frac{1}{16}\delta, \frac{31}{16}\delta)}$ converge in the weak topology to an integral Brakke flow $\{\mu_t\}_{t \in (\frac{1}{16}\delta, \frac{31}{16}\delta)}$ in $\Omega_0 \times (\frac{1}{16}, \frac{31}{16})$. In view of (2.23) we have $\text{spt } \mu_t \subset \Gamma_t$; it then follows from the one-sided minimization property of mean-convex MCF (see 3.9 in [W1] and Remark 2.5 in [HK]) that μ_t is of unit density. Thus, we have $\mu_t \leq \mathcal{H}^n \llcorner \Gamma_t$. Basing on the curvature estimate for MCF (see Theorem 2.1 in [G1] for the modification of [W3] and Theorem 5.6 in [E]), we would get local smooth estimates for $\{\Gamma_t^{\epsilon_i}\}_{t \in (\frac{1}{4}\delta, \frac{7}{4}\delta)}$ in $\Omega_0 \times (\frac{1}{4}, \frac{7}{4})$ (see Proposition 3.22 in [E]) for i sufficiently large. Upon passing to a subsequence, the smooth compactness theorem gives that the MCF $\{\Gamma_t^{\epsilon_i}\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ converge locally smoothly to a MCF $\{\tilde{\Gamma}_t\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ in $\Omega_0 \times (\frac{1}{2}, \frac{3}{2})$. Since for each $t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)$, $\tilde{\Gamma}_t$ is an embedded hypersurface contained in the embedded hypersurface Γ_t , we deduce that $\tilde{\Gamma}_t = \Gamma_t$ in $\Omega_0 \times (\frac{1}{2}, \frac{3}{2})$. Therefore, $\{\Gamma_t^{\epsilon_i}\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ converge locally smoothly to $\{\Gamma_t\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ in $\Omega_0 \times (\frac{1}{2}, \frac{3}{2})$.

By (2.24), the infimum mean curvature of $\{\Gamma_t\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ in $\Omega_0 \times (\frac{1}{2}, \frac{3}{2})$ is

$$\inf_{\frac{1}{2}\delta < t < \frac{3}{2}\delta} \min_{\Sigma_t} H \geq \kappa$$

by the maximum principle for mean curvature (cf. [M]). By (2.20) and (2.22), the infimum mean curvature of $\{\Gamma_t^{\epsilon_i}\}_{t \in (\frac{1}{2}\delta, \frac{3}{2}\delta)}$ in $\Omega_0 \times (\frac{1}{2}, \frac{3}{2})$ is

$$\inf_{\frac{1}{2}\delta + \frac{1}{2}\epsilon_i < u^{\epsilon_i} < \frac{3}{2}\delta + \frac{3}{2}\epsilon_i} \frac{1}{\sqrt{|\nabla u^{\epsilon_i}|^2 + \epsilon_i^2}}.$$

¹²See Proposition 2.6 for the definition of \dot{T} .

By the smooth convergence in the last paragraph, given $\sigma > 0$ there exists $i_\sigma \in \mathbb{N}$ so that

$$\inf_{\frac{1}{2}\delta + \frac{1}{2}\epsilon_i < u^{\epsilon_i} < \frac{3}{2}\delta + \frac{3}{2}\epsilon_i} \frac{1}{\sqrt{|\nabla u^{\epsilon_i}|^2 + \epsilon^2}} \geq \frac{\kappa}{1 + \sigma} \quad \forall i \geq i_\sigma$$

In addition, due to (2.18) we may assume that

$$\left\{ \frac{1}{2}\delta + \frac{1}{2}\epsilon_i < u^{\epsilon_i} < \frac{3}{2}\delta + \frac{3}{2}\epsilon_i \right\} \supset \{u = \delta\} = \Sigma_\delta \quad \forall i \geq i_\sigma$$

and hence

$$\min_{\Sigma_\delta} \frac{1}{\sqrt{|\nabla u^{\epsilon_i}|^2 + \epsilon^2}} \geq \frac{\kappa}{1 + \sigma} \quad \forall i \geq i_\sigma,$$

which gives that

$$(2.25) \quad \max_{\Sigma_\delta} |\nabla u^{\epsilon_i}| \leq \max_{\Sigma_\delta} \sqrt{|\nabla u^{\epsilon_i}|^2 + \epsilon^2} \leq \frac{1 + \sigma}{\kappa} \quad \forall i \geq i_\sigma.$$

On the other hand, differentiating (2.17) with respect to x^k following by multiplying $\partial_k u^\epsilon$ yields

$$-a_\epsilon^{ij}(\nabla u^\epsilon) \partial_{ij} (|\nabla u^\epsilon|^2) - \frac{\partial a_\epsilon^{ij}}{\partial \xi^l}(\nabla u^\epsilon) \partial_{ij} u^\epsilon \partial_l (|\nabla u^\epsilon|^2) = -2a_\epsilon^{ij}(\nabla u^\epsilon) \partial_{ik} u^\epsilon \partial_{jk} u^\epsilon \leq 0,$$

where $a_\epsilon^{ij}(\xi) = \delta^{ij} - \frac{\xi^i \otimes \xi^j}{|\xi|^2 + \epsilon^2}$. It follows from the maximum principle that

$$(2.26) \quad \max_{\Omega_\delta} |\nabla u^\epsilon| = \max_{\partial\Omega_\delta = \Sigma_\delta} |\nabla u^\epsilon|$$

Thus, combining (2.25) with (2.26) gives

$$(2.27) \quad \max_{\Omega_\delta} |\nabla u^{\epsilon_i}| \leq \frac{1 + \sigma}{\kappa} \quad \forall i \geq i_\sigma.$$

Finally, given a point $p \in \Omega_0$, we have the following three cases to consider:

Case 1: $p \in \Omega_0 \setminus \Omega_\delta$. Then p must be a regular point of the flow since $u(p) \in (0, T_1)$ is a regular time. Moreover, since $\{\Sigma_t\}_{t \in [0, T_1]}$ is a MCF, the maximum principle for the mean curvature gives

$$H(p) = \frac{1}{|\nabla u(p)|} \geq \kappa$$

(cf. [M]). So $|\nabla u(p)| \leq \kappa^{-1}$.

Case 2: $p \in \Omega_\delta$ is a singular point of the flow. In this case we have $\nabla u(p) = 0$.

Case 3: $p \in \Omega_\delta$ is a regular point of the flow. Let $v = \frac{\nabla u(p)}{|\nabla u(p)|}$. Then by the mean value theorem, (2.18), and (2.27) we have

$$\frac{1}{s} |u(p + sv) - u(p)| = \lim_{i \rightarrow \infty} \frac{1}{s} |u^{\epsilon_i}(p + sv) - u^{\epsilon_i}(p)| \leq \frac{1 + \sigma}{\kappa}$$

for every $0 < s < \text{dist}(p, \Sigma_\delta)$. Letting $s \searrow 0$ gives

$$|\nabla u(p)| = |\nabla u(p) \cdot v| \leq \frac{1 + \sigma}{\kappa}$$

Since $\sigma > 0$ is arbitrary, we have $|\nabla u(p)| \leq \kappa^{-1}$. \square

Next, we will be devoted to the smooth estimates for $\{\Sigma_t\}_{t > \hat{T}}$ on its regular part. Before that, we have the following simple observation.

Lemma 2.10. *The singular set \mathcal{S} of the flow $\{\Sigma_t\}$ is a closed set (and hence is compact).*

Proof. Given a sequence $\{p_i\}_{i \in \mathbb{N}} \subset \mathcal{S}$ tending to a point $p \in \mathbb{R}^n$, note that by Proposition 2.6, we have $p_i \in \Omega_{\dot{T}}$ (i.e., $u(p_i) > \dot{T}$) for every $i \in \mathbb{N}$. Thus, $p \in \bar{\Omega}_{\dot{T}}$.

On the other hand, recall that \mathcal{S} is the set of critical points of u . So $\nabla u(p_i) = 0$ for every i . By the continuity of ∇u on $\bar{\Omega}_{\dot{T}}$ (cf. [CM4]), we obtain

$$\nabla u(p) = \lim_{i \rightarrow \infty} \nabla u(p_i) = 0.$$

Therefore $p \in \mathcal{S}$. □

The set of all regular points in Ω_0 is obviously an open set. In fact, let $p \in \Omega_0$ be a regular point of $\{\Sigma_t\}$, since u is smooth near p with $\nabla u(p) \neq 0$, the implicit function theorem yields that there is a neighborhood of p where the level sets of u are all regular and they jointly constitute a MCF by (1.2). This motivates the following definition.

Definition 2.11. Let $p \in \Omega_0$ be a regular point of $\{\Sigma_t\}$. When a radius $r > 0$ is sufficiently small, we have $B_r(p) \subset \Omega_0 \setminus \mathcal{S}$ ¹³ and that every level set $\Sigma_t = \{u = t\}$ in $B_r(p)$ is a $\varepsilon(n)$ -Lipschitz graph over $T_p \Sigma_{u(p)}$, where $\varepsilon(n) \in (0, 1)$ is a small constant as required by Lemma 3.10 in [G1] and Lemma 2.1 (see (2.1)). The supremum of all such radii is called the **smooth scale** of $\{\Sigma_t\}$ at p .

In the following proposition we estimates the smooth scales from below.

Proposition 2.12. *Let $\iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}$ be positive constants such that*

$$\begin{aligned} \text{rad } \Sigma_0 &\geq \iota, \quad \min_{\Sigma_0} H \geq \kappa, \\ \max_{\Sigma_0} |A| &\leq K, \quad \max_{\Sigma_0} |\nabla A| \leq K', \quad \max_{\Sigma_0} |\nabla^2 A| \leq K'', \\ \mathcal{H}^{n-1}(\Sigma_0) &\leq \mathfrak{A}, \quad \text{diam } \Sigma_0 \leq \mathfrak{D}. \end{aligned}$$

Then there exists $\gamma = \gamma(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}) > 0$ so that the smooth scale of $\{\Sigma_t\}$ at a regular point $p \in \Omega_{\dot{T}}$ is greater than $\gamma |\nabla u(p)|$, where $\dot{T} > 0$ is the constant in Proposition 2.6.

Proof. By Proposition 2.3 and [HK], $\{\Sigma_t\}$ is α -noncollapsing, where $\alpha = \alpha(n, \iota, \kappa) > 0$; by Proposition 2.5 and [H2], the entropy of $\{\Sigma_t\}$ is bounded above by $\lambda = \lambda(n, K, \mathfrak{A}, \mathfrak{D}) > 0$.

Let $p \in \Omega_{\dot{T}}$ be a regular point of the flow. By Proposition 2.8 we have

$$B_{\kappa \dot{T}}(p) \subset \Omega_0;$$

additionally, Proposition 2.9 implies

$$\begin{aligned} \kappa \dot{T} &= \kappa \dot{T} H(p) |\nabla u(p)| \geq \kappa^2 \dot{T} |\nabla u(p)|, \\ \sqrt{\dot{T}} &= \sqrt{\dot{T}} H(p) |\nabla u(p)| \geq \kappa \sqrt{\dot{T}} |\nabla u(p)|, \end{aligned}$$

which gives that

$$\min \left\{ |\nabla u(p)|, \kappa \dot{T}, \sqrt{\dot{T}} \right\} \geq \min \left\{ 1, \kappa^2 \dot{T}, \kappa \sqrt{\dot{T}} \right\} |\nabla u(p)|.$$

¹³Note that u is smooth in $B_r(p)$ with $\nabla u \neq 0$; it follows that (1.2) would be satisfied in the classical sense on $B_r(p)$ and hence the level sets of u in $B_r(p)$ form a MCF.

It follows from Haslhofer-Kleiner regularity theorem (see Theorem 1.8, Remark 2.7, and Corollary 3.3 in [HK]) that $\{\Sigma_t\}_{t \in (u(p) - \rho^2, u(p) + \rho^2)}$ is regular in $B_\rho(p)$, where

$$\rho = \eta^2 |\nabla u(p)|$$

with

$$\eta = \frac{\min \left\{ 1, \kappa^2 \dot{T}, \kappa \sqrt{\bar{T}} \right\}}{C(n, \alpha, \lambda)} \in (0, 1),$$

and that

$$(2.28) \quad \sup_{t \in (u(p) - \rho^2, u(p) + \rho^2)} \sup_{\Sigma_t \cap B_\rho(p)} \rho |A| \leq 1,$$

$$(2.29) \quad \sup_{t \in (u(p) - \rho^2, u(p) + \rho^2)} \sup_{\Sigma_t \cap B_\rho(p)} \rho^2 |\nabla A| \leq 1,$$

$$(2.30) \quad \inf_{t \in (u(p) - \rho^2, u(p) + \rho^2)} \inf_{\Sigma_t \cap B_\rho(p)} H |\nabla u(p)| \geq \frac{1}{C(n)}.$$

Let

$$(2.31) \quad x(t) : \left(u(p) - \frac{\rho^2}{2\sqrt{n-1}}, u(p) + \frac{\rho^2}{2\sqrt{n-1}} \right) \rightarrow B_{\frac{\rho}{2}}(p)$$

be the ‘‘trajectory’’ of p along the MCF $\{\Sigma_t\}$, namely,

$$x(u(p)) = p, \quad x(t) \in \Sigma_t, \quad \frac{d}{dt}x(t) = \vec{H}(x(t)).$$

Notice that in (2.31) we have used the fact that

$$(2.32) \quad \left| \frac{d}{dt}x(t) \right| = H(x(t)) \leq \sqrt{n-1} |A(x(t))| \leq \frac{\sqrt{n-1}}{\rho}.$$

Let $\varepsilon = \varepsilon(n) > 0$ be the small constant as specified in Definition 2.11. By (2.28), and the α -noncollapsing property of the flow, there exists $\sigma = \sigma(n, \alpha, \varepsilon) \in (0, \frac{1}{2})$ so that $\Sigma_t \cap B_{\sigma\rho}(x(t))$ is a $\frac{\varepsilon}{3}$ -Lipschitz graph over $T_{x(t)}\Sigma_t$. Using the evolution of the unit normal vector (cf. [M]) and (2.29) we get

$$(2.33) \quad \left| \frac{d}{dt}N(x(t)) \right| = |-\nabla H(x(t))| \leq C(n) |\nabla A(x(t))| \leq \frac{C(n)}{\rho^2}.$$

By (2.32) and (2.33), there is $\theta = \theta(n, \sigma, \varepsilon) \in (0, 1)$ so that

$$\Sigma_t \cap B_{\frac{1}{2}\sigma\rho}(p) \subset \Sigma_t \cap B_{\sigma\rho}(x(t))$$

is a $\frac{\varepsilon}{2}$ -Lipschitz graph over $T_p\Sigma_{u(p)}$ for every $t \in (u(p) - \theta\rho^2, u(p) + \theta\rho^2)$ and that

$$(2.34) \quad \sqrt{n-1}\theta \leq \frac{\varepsilon\sigma}{4\hat{C}},$$

where $\hat{C} = \hat{C}(n)$ is the constant in the proof of Lemma 2.1.

Moreover, by (2.28), (2.34), and the argument in the proof of Lemma 2.1 (with slight modifications), we infer that the domains of the aforementioned local graphs over $T_p\Sigma_{u(p)}$ all contain a $(n-1)$ -dimensional ball of radius $\frac{\varepsilon\sigma}{4\hat{C}}\rho$ centered at p .

For ease of notations, let us assume that $p = 0$, $T_p \Sigma_{u(p)} = \mathbb{R}^{n-1} \times \{0\}$, and $\frac{\nabla u(p)}{|\nabla u(p)|} = (0, 1)$. Then

$$\mathcal{D} := \bigsqcup_{t \in (u(p) - \theta\rho^2, u(p) + \theta\rho^2)} \Sigma_t \cap B_{\frac{1}{2}\sigma\rho}(p)$$

contains the graph $x^{n+1} = f(x', t)$ satisfying

$$(2.35) \quad f(0, u(p)) = 0, \quad \nabla f(0, u(p)) = 0, \quad |\nabla f(x', t)| \leq \frac{\varepsilon}{2},$$

where

$$(2.36) \quad x' = (x^1, \dots, x^{n-1}) \in B_{\frac{\varepsilon\sigma}{4C}\rho}^{n-1}, \quad t \in (u(p) - \theta\rho^2, u(p) + \theta\rho^2).$$

It follows from the regularity of the flow and the equation of MCF (cf. [M]) that

$$(2.37) \quad \frac{\partial_t f}{\sqrt{1 + |\nabla f|^2}} = \nabla \cdot \frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} = H.$$

Note by (2.30) we have $H \geq \frac{\eta^2}{C(n)}\rho^{-1}$, which gives that

$$(2.38) \quad \partial_t f = \sqrt{1 + |\nabla f|^2} H \geq H \geq \frac{\eta^2}{C(n)}\rho^{-1}.$$

By (2.35), (2.36), and (2.38), we conclude that

$$\mathcal{D} \supset B_{\min\{\frac{\theta\eta^2}{2\varepsilon C(n)}, \frac{\varepsilon\sigma}{4C}\}\rho}^{n-1} \times B_{\frac{\theta\eta^2}{2C(n)}\rho}^1 \supset B_{\gamma|\nabla u(p)|}^n,$$

where $\gamma = \gamma(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D})$. \square

All the higher order covariant derivatives of the second fundamental form of Σ_t follow immediately by [EH].

Proposition 2.13. *In Proposition 2.12, by taking a smaller constant if necessary, we may assume that*

$$(2.39) \quad \gamma = \gamma(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}) \leq \frac{1}{2} \left(\sqrt{1 + 4\kappa^2 \dot{T}} - 1 \right),$$

where $\dot{T} = \dot{T}(n, \iota, K, K', K'') > 0$ is the constant in Proposition 2.6. Then for every regular point $p \in \Omega_{\dot{T}}$, the flow $\{\Sigma_t\}$ satisfies

$$\sup_{\Sigma_t \cap B_r(p)} r^{m+1} |\nabla^m A| \leq C(n, m) \quad \forall m \geq 0,$$

where $r = \frac{1}{2}\gamma|\nabla u(p)|$.

Proof. Fix a regular point $p \in \Omega_{\dot{T}}$ and let $r = \frac{1}{2}\gamma|\nabla u(p)|$. By Proposition 2.9, for every $x \in B_{2r}(p) \subset \Omega_0$ (see Proposition 2.12) we have

$$|u(x) - u(p)| \leq \kappa^{-1}|x - p| < 2\kappa^{-1}r = \kappa^{-1}\gamma|\nabla u(p)|,$$

which yields that

$$(2.40) \quad \begin{aligned} u(x) &> u(p) - \kappa^{-1}\gamma|\nabla u(p)| > \dot{T} - \kappa^{-1}\gamma|\nabla u(p)| \\ &= \left(\dot{T}H(p) - \kappa^{-1}\gamma \right) |\nabla u(p)| \geq \left(\kappa\dot{T} - \kappa^{-1}\gamma \right) |\nabla u(p)| \geq 4r^2. \end{aligned}$$

Note that the last inequality comes from Proposition 2.9 and (2.39).

Since $\{\Sigma_t\}_{t \geq 0}$ is a MCF in $B_{2r}(p)$ with

$$0 < (N \cdot N(p))^{-1} \leq \sqrt{1 + \varepsilon^2(n)}$$

(see Definition 2.11 and Proposition 2.12), the Ecker-Huisken smooth estimates for MCF (cf. [EH]) gives that

$$\sup_{\Sigma_t \cap B_r(p)} r^{m+1} |\nabla^m A| \leq C(n, m) \quad \forall m \in \mathbb{N} \cup \{0\}$$

for every $t > 4r^2$. The conclusion follows immediately by noting that (2.40) gives that $\Sigma_t \cap B_{2r}(p) = \emptyset$ for $t \in [0, 4r^2]$. \square

We are in a position to give an estimate of the Hessian $\nabla^2 u$, which is essential to Proposition 2.18 in Section 2.4.

Proposition 2.14. *Let $\iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}$ be positive constants such that*

$$\begin{aligned} \text{rad } \Sigma_0 &\geq \iota, \quad \min_{\Sigma_0} H \geq \kappa, \\ \max_{\Sigma_0} |A| &\leq K, \quad \max_{\Sigma_0} |\nabla A| \leq K', \quad \max_{\Sigma_0} |\nabla^2 A| \leq K'', \\ \mathcal{H}^{n-1}(\Sigma_0) &\leq \mathfrak{A}, \quad \text{diam } \Sigma_0 \leq \mathfrak{D}. \end{aligned}$$

Then the Hessian of the arrival time function u satisfies

$$\sup_{\Omega_0} |\nabla^2 u| \leq C(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}).$$

Proof. If $x \in \Omega_0$ is regular point of the flow, say $x \in \Sigma_t$ with $t = u(x)$. Then let $N(x) = \frac{\nabla u(x)}{|\nabla u(x)|}$ be the unit normal vector and $\{e_1, \dots, e_{n-1}\}$ be an orthonormal basis of $T_x \Sigma_t$, by the calculations in [CM4] we have

$$(2.41) \quad \begin{aligned} -\nabla^2 u \cdot (e_i \otimes e_j) &= \frac{A(e_i, e_j)}{H}, \\ -\nabla^2 u \cdot (N \otimes e_i) &= \frac{\nabla H \cdot e_i}{H^2}, \\ -\nabla^2 u \cdot (N \otimes N) &= \frac{\Delta H}{H^3} + \frac{|A|^2}{H^2}. \end{aligned}$$

where A denotes the second fundamental form of Σ_t and Δ is the Laplace-Beltrami operator on Σ_t . Let \dot{T} be as given in Proposition 2.6. Given a point $p \in \Omega_0$, below we divide into three cases to estimate $|\nabla^2 u(p)|$.

Case 1: $p \in \Omega_0 \setminus \Omega_{\dot{T}}$, i.e., $u(p) \in (0, \dot{T}]$. By (2.15), (2.16), and Proposition 2.9, we have

$$\begin{aligned} \frac{|A(p)|}{H(p)} &\leq \frac{C(n, \iota, \kappa, K, K', K'')}{\kappa}, \\ \frac{|\nabla H(p)|}{H^2(p)} &\leq C(n) \frac{|\nabla A(p)|}{H^2(p)} \leq \frac{C(n, \iota, \kappa, K, K', K'')}{\kappa^2}, \\ \frac{|\Delta H(p)|}{H^3(p)} &\leq C(n) \frac{|\nabla^2 A(p)|}{H^3(p)} \leq \frac{C(n, \iota, \kappa, K, K', K'')}{\kappa^3}. \end{aligned}$$

It follows by (2.41) that $|\nabla^2 u(p)| \leq C(n, \iota, \kappa, K, K', K'')$.

Case 2: $p \in \Omega_{\bar{t}}$ is a regular point. By Definition 2.11, Proposition 2.12, and Proposition 2.13, we have

$$\begin{aligned} \frac{|A(p)|}{H(p)} &= |A(p)| |\nabla u(p)| \leq \frac{C(n)}{\gamma(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D})}, \\ \frac{|\nabla H(p)|}{H^2(p)} &\leq C(n) |\nabla A(p)| |\nabla u(p)|^2 \leq \frac{C(n)}{\gamma^2(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D})}, \\ \frac{|\Delta H(p)|}{H^3(p)} &\leq C(n) |\nabla^2 A(p)| |\nabla u(p)|^3 \leq \frac{C(n)}{\gamma^3(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D})}. \end{aligned}$$

It follows by (2.41) that $|\nabla^2 u(p)| \leq C(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D})$.

Case 3: $p \in \Omega_{\bar{t}}$ is a singular point. By [CM4],

$$-\nabla^2 u(p) = \frac{1}{n-1} I_n$$

if p is a round point and

$$-\nabla^2 u(p) = \frac{1}{n-2} \mathcal{O} \begin{pmatrix} I_{n-1} & 0 \\ 0 & 0 \end{pmatrix} \mathcal{O}^{-1},$$

if p is a cylindrical point, where

$$\mathcal{O} = [N(p), e_1, \dots, e_{n-1}]$$

is the orthogonal matrix such that e_n is an unit vector along the axial direction of the tangent cylinder at p and $\{e_1, \dots, e_{n-1}\}$ is an orthonormal basis of the orthogonal complement of $\text{span}\{N(p), e_{n-1}\}$. In either case, $|\nabla^2 u(p)| \leq C(n)$. \square

The following corollary is based on Lemma 3.10 in [G1], Proposition 2.13, and Proposition 2.14. It will be used in Lemma 2.21 in Section 2.4.

Corollary 2.15. *Let $p \in \Omega_{\bar{t}}$ be a regular point of the flow $\{\Sigma_t\}$ and choose*

$$(2.42) \quad r \in \left[\frac{1}{4} \gamma |\nabla u(p)|, \frac{1}{2} \gamma |\nabla u(p)| \right],$$

where $\gamma > 0$ is the constant in Proposition 2.12 and Proposition 2.13.

If $\Sigma_t \cap B_{2r}(p)$ is a $2\varepsilon(n)$ -Lipschitz graph of $f(\cdot, t)$ over some hyperplane Π (which can be chosen as, but not restricted to, $T_p \Sigma_u(p)$ by Proposition 2.12 and (2.42)). Then the graph of $f(\cdot, t)$ restricted in $B_r(p)$ would satisfy

$$(2.43) \quad r^{m+2l-1} |\partial_t^l \nabla^m f| \leq C(n, m, l) \quad \text{whenever } m + 2l \geq 2$$

and

$$r \partial_t f \geq \frac{\gamma}{4\Lambda},$$

where $\Lambda = \Lambda(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}) > 0$ is a constant.

Proof. By Lemma 3.10 in [G1] and Proposition 2.13, we have

$$(2.44) \quad r^{m-1} |\nabla^m f| \leq C(n, m) \quad \forall m \geq 2.$$

Since $\{\Sigma_t\}$ is regular and graphical in $B_{2r}(p)$ by Definition 2.11, Proposition 2.12, and (2.42), the function f would satisfy

$$(2.45) \quad \partial_t f = \sqrt{1 + |\nabla f|^2} \nabla \cdot \frac{\nabla f}{\sqrt{1 + |\nabla f|^2}} = \left(I - \frac{\nabla f \otimes \nabla f}{1 + |\nabla f|^2} \right) \cdot \nabla^2 f$$

(cf. [M]). Using (2.44) and keeping on differentiating (2.45) with respect to t , we then obtain (2.43).

Additionally, by Definition 2.11 and Proposition 2.12, we have $B_{2r}(p) \subset \Omega_0 \setminus \mathcal{S}$. It follows from Proposition 2.14, and (2.42) that

$$|\nabla u(x)| \leq |\nabla u(p)| + C(n) \|\nabla^2 u\|_{C(\bar{B}_r(p))} r \leq \Lambda |\nabla u(p)| \quad \forall x \in B_r(p),$$

where $\Lambda = \Lambda(n, \iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}) > 0$ is a constant. Thus, by (2.42) and (2.45) we obtain

$$r \partial_t f = r \sqrt{1 + |\nabla f|^2} H \geq rH \geq \frac{\gamma}{4} \frac{|\nabla u(p)|}{|\nabla u|} \geq \frac{\gamma}{4\Lambda}.$$

□

2.4. Smooth convergence off singularities. Let $\iota, \kappa, K, K', K'', \mathfrak{A}, \mathfrak{D}$ be positive constants such that

$$(2.46) \quad \begin{aligned} \text{rad } \Sigma_0 &> \iota, \quad \min_{\Sigma_0} H > \kappa, \\ \max_{\Sigma_0} |A| &< K, \quad \max_{\Sigma_0} |\nabla A| < K', \quad \max_{\Sigma_0} |\nabla^2 A| < K'', \\ \mathcal{H}^{n-1}(\Sigma_0) &< \mathfrak{A}, \quad \text{diam } \Sigma_0 < \mathfrak{D}. \end{aligned}$$

Let $\{\Sigma_0^k\}_{k \in \mathbb{N}}$ be a sequence of closed connected hypersurfaces tending to Σ_0 in the C^4 topology (see Proposition 2.6). Due to the convergence, we may assume for simplicity that (2.46) hold with Σ_0^k in place of Σ_0 for every $k \in \mathbb{N}$ and that Proposition 2.6 holds for every $k \in \mathbb{N}$.

With the uniform estimates from Section 2.2 and Section 2.3, in this subsection we shall be equipped to prove Theorem 1.1. The critical part is to show that u^k (i.e., the arrival time function of $\{\Sigma_t^k\}$, see Proposition 2.6) converges locally in the C^1 topology to u (see Proposition 2.16 and Proposition 2.18). The convergence signifies that if p is a regular point of $\{\Sigma_t\}$, because of

$$\lim_{k \rightarrow \infty} |\nabla u^k(p)| = |\nabla u(p)|$$

and that the smooth scale of $\{\Sigma_t^k\}$ at p depends on $|\nabla u^k(p)|$ (see Proposition 2.12), it is not surprising that $\{\Sigma_t^k\}$ would converge locally smoothly to $\{\Sigma_t\}$ near p as $k \rightarrow \infty$. This is the core of Corollary 2.21, from which Theorem 1.1 follows.

Let us start with the uniform convergence of the arrival time functions. Note that $u : \bar{\Omega}_0 \rightarrow [0, \infty)$ can be extended continuously to \mathbb{R}^n simply by interpreting it to be zero on $\mathbb{R}^n \setminus \bar{\Omega}_0$. The same is true for every $u^k : \bar{\Omega}_0^k \rightarrow [0, \infty)$. The following proposition is based on Proposition 2.9 and the uniqueness of viscosity solutions to the Dirichlet problem (1.2) (cf. [ES]).

Proposition 2.16. *Upon identifying the arrival time function u^k with its natural continuous extension (by interpreting it to be zero outside $\bar{\Omega}_0^k$), u^k converges uniformly to u in \mathbb{R}^n as $k \rightarrow \infty$.*

Proof. Choose $R > 0$ large so that $\bar{\Omega}_0$ is contained in the open ball B_R . For convenience, let us assume that for every $k \in \mathbb{N}$, Σ_0^k is also contained in B_R . Note that by [ES] we have

$$(2.47) \quad \Sigma_t^k \subset B_{\sqrt{R^2 - 2(n-1)t}} \quad \forall t \geq 0,$$

which yields that

$$(2.48) \quad \max_{\mathbb{R}^n} u^k = \max_{\bar{\Omega}_0^k} u^k \leq \frac{R^2}{2(n-1)}.$$

Given $\epsilon \in (0, \dot{T})$, where $\dot{T} > 0$ is as given in Proposition 2.6, choose $\delta > 0$ so small that $\Omega_{\epsilon/2} \subset \Omega_0$ is outside the 4δ -tubular-neighborhood of Σ_0 . By Proposition 2.6, there exists $k_\epsilon \in \mathbb{N}$ so that for every $k \geq k_\epsilon$, Σ_0^k is contained in the δ -tubular-neighborhood of Σ_0 and that $\Omega_{\epsilon/2}^k \subset \Omega_0$ ¹⁴ is outside the 3δ -neighborhood of Σ_0 . Then for each $x_0 \in \mathbb{R}^n$, we claim that

$$(2.49) \quad |u^k(x) - u^k(x_0)| \leq \epsilon \quad \forall x \in B_{\min\{\delta, \kappa\epsilon\}}(x_0), \quad k \geq k_\epsilon.$$

Let $k \geq k_\epsilon$. Below we divide into three cases to verify:

Case 1: x_0 belongs to the 2δ -tubular-neighborhood of Σ_0 (so $x_0 \notin \Omega_{\epsilon/2}^k$). Then for every $x \in B_\delta(x_0)$, it is within the 3δ -tubular-neighborhood of Σ_0 and hence falls outside $\Omega_{\epsilon/2}^k$; thus,

$$|u^k(x) - u^k(x_0)| \leq u^k(x) + u^k(x_0) \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Case 2: $x_0 \in \Omega_0$ is outside the 2δ -tubular-neighborhood of Σ_0 (so $x_0 \in \Omega_0^k$). Then for every $x \in B_{\min\{\delta, \kappa\epsilon\}}(x_0)$, it is in Ω_0 and outside the δ -tubular-neighborhood of Σ_0 and so falls within Ω_0^k . Thus, $B_{\min\{\delta, \kappa\epsilon\}}(x_0) \subset \Omega_0^k$ and for every $x \in B_{\min\{\delta, \kappa\epsilon\}}(x_0)$, by (2.46) and Proposition 2.9 we have

$$|u^k(x) - u^k(x_0)| \leq \kappa^{-1}|x - x_0| \leq \epsilon.$$

Case 3: $x_0 \notin \Omega_0$ is outside the 2δ -tubular-neighborhood of Σ_0 (so $x_0 \notin \Omega_0^k$). For every $x \in B_\delta(x_0)$, it is outside Ω_0 and the δ -tubular-neighborhood of Σ_0 and hence falls outside Ω_0^k ; thus,

$$|u^k(x) - u^k(x_0)| = 0.$$

By (2.47), (2.48), and (2.49), Arzelà-Ascoli compactness theorem implies that for any subsequence of $\{u^k\}_{k \in \mathbb{N}}$ there would exist a further subsequence that converges uniformly to some continuous function $\hat{u} : \mathbb{R}^n \rightarrow [0, \infty)$. Notice that \hat{u} vanishes on $\mathbb{R}^n \setminus \bar{\Omega}_0$, which (by the continuity of \hat{u}) implies that $\hat{u}|_{\Sigma_0} = 0$. Moreover, by the argument in the proof of Theorem 2.7 in [ES], \hat{u} would satisfy the equation (1.2) in the viscosity sense on Ω_0 . Thus, by the uniqueness of viscosity solutions to the Dirichlet problem of (1.2) (see Theorem 7.5 in [ES]), we deduce that $\hat{u} = u$.

By the last paragraph, any subsequence of $\{u^k\}_{k \in \mathbb{N}}$ has a further subsequence converging uniformly to u . Therefore, the whole sequence $\{u^k\}_{k \in \mathbb{N}}$ must converge uniformly to u . \square

We immediately have the following corollary.

Corollary 2.17. *Let T_{ext}^k be the time of extinction of $\{\Sigma_t^k\}$. Then we have*

$$\lim_{k \rightarrow \infty} T_{ext}^k = \lim_{k \rightarrow \infty} \max_{\Omega_0^k} u^k = \max_{\Omega_0} u = T_{ext}.$$

Given the convergence of the arrival time functions in Proposition 2.16, we can proceed to prove the convergence of gradients with the help of Proposition 2.9 and Proposition 2.14.

¹⁴ $\Omega_{\epsilon/2}^k = \{u^k > \frac{\epsilon}{2}\}$ is the region bounded by $\Sigma_{\epsilon/2}^k$.

Proposition 2.18. ∇u^k converges locally uniformly to ∇u in Ω_0 as $k \rightarrow \infty$.

Proof. Let \mathcal{B} be any open ball in \mathbb{R}^n such that $\bar{\mathcal{B}} \subset \Omega_0$. Choose $k_{\mathcal{B}} \in \mathbb{N}$ sufficiently large so that $\bar{\mathcal{B}} \subset \Omega_0^k$ for every $k \geq k_{\mathcal{B}}$. By Proposition 2.9 and Proposition 2.14, any subsequence of $\{\nabla u^k|_{\bar{\mathcal{B}}}\}_{k \geq k_{\mathcal{B}}}$ has a further subsequence, say $\{\nabla u^{k_j}|_{\bar{\mathcal{B}}}\}_{j \in \mathbb{N}}$, that converges uniformly to some continuous vector-valued function $F = (F^1, \dots, F^n)$ on $\bar{\mathcal{B}}$ by Arzelà-Ascoli compactness theorem. Note that for every function $\zeta \in C_c^1(\mathcal{B})$, by Proposition 2.16 there holds

$$\begin{aligned} \int_{\mathcal{B}} F^i \zeta \, dx &= \lim_{j \rightarrow \infty} \int_{\mathcal{B}} \partial_i u^{k_j} \zeta \, dx = - \lim_{j \rightarrow \infty} \int_{\mathcal{B}} u^{k_j} \partial_i \zeta \, dx \\ &= - \int_{\mathcal{B}} u \partial_i \zeta \, dx = \int_{\mathcal{B}} \partial_i u \zeta \, dx \quad \forall i \in \{1, \dots, n\}, \end{aligned}$$

we deduce that $F = \nabla u|_{\bar{\mathcal{B}}}$. Therefore, the whole sequence $\{\nabla u^k|_{\bar{\mathcal{B}}}\}_{k \geq k_{\mathcal{B}}}$ must converges uniformly to $\nabla u|_{\bar{\mathcal{B}}}$. \square

What follow are two corollaries concerning the singular sets and singular times of $\{\Sigma_t^k\}$, respectively. Note that by Proposition 2.6 and Proposition 2.8, we may assume that the singular set of the mean-convex LSF $\{\Sigma_t^k\}$ is contained in $\Omega_{\dot{T}}$ when k is large.

Corollary 2.19. *Given an open set U strictly contained in $\Omega_0 \setminus \mathcal{S}$, there is $k_U \in \mathbb{N}$ so that for every $k \geq k_U$, the flow $\{\Sigma_t^k\}$ is regular in U .*

As an illustration, if $U = \Omega_{\dot{T}} \setminus \bar{\mathcal{S}}^\delta$ with $\delta > 0$, where

$$\mathcal{S}^\delta := \{x : \text{dist}(x, \mathcal{S}) < \delta\},$$

then for every $k \geq k_U$, the singular set of $\{\Sigma_t^k\}$ would be contained in $\bar{\mathcal{S}}^\delta$.

Proof. Let U be an open set such that $\bar{U} \subset \Omega_0 \setminus \mathcal{S}$. Firstly, choose $k_0 \in \mathbb{N}$ sufficiently large so that $\bar{U} \subset \Omega_0^k$ for $k \geq k_0$. It follows from Proposition 2.18 that there is $k_U \geq k_0$ so that

$$\min_{\bar{U}} |\nabla u^k| \geq \frac{1}{2} \min_{\bar{U}} |\nabla u| > 0,$$

which yields that the flow $\{\Sigma_t^k\}$ is regular in U . \square

Corollary 2.20. *Suppose that the flow $\{\Sigma_t\}$ is regular during $t \in [a, b]$, where $0 < a < b < T_{ext}$, then there exists $k_{[a,b]} \in \mathbb{N}$ so that for every $k \geq k_{[a,b]}$, the flow $\{\Sigma_t^k\}$ is regular during $t \in [a, b]$.*

Proof. By Lemma 2.10 and the continuity of u , the set of singular times is compact (and hence closed). So there exists

$$0 < a' < a < b < b' < T_{ext}$$

such that

$$\{a' \leq u \leq b'\} \subset \Omega_0 \setminus \mathcal{S}.$$

Then $U := \{a' < u < b'\}$ is an open set strictly contained in $\Omega_0 \setminus \mathcal{S}$. By Proposition 2.16, there is $k_0 \in \mathbb{N}$ so that for $k \geq k_0$,

$$\{a \leq u^k \leq b\} \subset \{a' < u < b'\} = U.$$

It follows from Corollary 2.19 that there exists $k_U \geq k_0$ so that for every $k \geq k_U$, the flow $\{\Sigma_t^k\}$ is regular in U and so $\{\Sigma_t^k\}$ is regular during $t \in [a, b]$. \square

On account of the uniform estimates for the flows (see Proposition 2.12, Proposition 2.13, and Corollary 2.15) and the local C^1 convergence of the arrival time functions (see Proposition 2.16 and Proposition 2.18), we are now able to prove the following lemma, which is the very essence of Theorem 1.1.

Lemma 2.21. *The flow $\{\Sigma_t^k\}$ converges locally smoothly to $\{\Sigma_t\}$ in $(\Omega_{\dot{T}} \setminus \mathcal{S}) \times (0, \infty)$ as $k \rightarrow \infty$.*

Proof. Fix $p \in \Omega_{\dot{T}} \setminus \mathcal{S}$. By Proposition 2.16 and Proposition 2.18 we have

$$(2.50) \quad \lim_{k \rightarrow \infty} u^k(p) = u(p) > \dot{T}$$

$$(2.51) \quad \lim_{k \rightarrow \infty} \nabla u^k(p) = \nabla u(p) \neq 0.$$

So when k is large, $p \in \Omega_{\dot{T}}^k$ is a regular point of $\{\Sigma_t^k\}$. Let $r = \frac{1}{3}\gamma|\nabla u(p)|$. By Definition 2.11, Proposition 2.12, and (2.51), there exists $k_0 \in \mathbb{N}$ so that when $k \geq k_0$,

$$\frac{2}{3}|\nabla u(p)| \leq |\nabla u^k(p)| \leq \frac{4}{3}|\nabla u(p)|$$

and every level hypersurface $\Sigma_t^k = \{u^k = t\}$ in $B_{2r}(p)$ is a $2\varepsilon(n)$ -Lipschitz graph of $f^k(\cdot, t)$ over $T_p\Sigma_{u(p)}$. Also, $\Sigma_t = \{u = t\}$ in $B_{2r}(p)$ is certainly a $\varepsilon(n)$ -Lipschitz graph of $f(\cdot, t)$ over $T_p\Sigma_{u(p)}$. Then by Corollary 2.15 the graph of $f^k(\cdot, t)$ restricted in $B_r(p)$ would satisfy

$$(2.52) \quad r^{m+2l-1} |\partial_t^l \nabla^m f^k| \leq C(n, m, l) \quad \text{whenever } m + 2l \geq 2,$$

$$(2.53) \quad |\nabla f^k| \leq 2\varepsilon(n) \leq 2,$$

$$(2.54) \quad a \leq r \partial_t f^k \leq b,$$

where $a = \frac{\gamma}{4\Lambda}$ and $b = b(n)$ are positive constants.

For ease of notations, let us assume that $p = 0$, $T_p\Sigma_{u(p)} = \mathbb{R}^{n-1} \times \{0\}$, and $\frac{\nabla u(p)}{|\nabla u(p)|} = (0, 1)$. So for $k \geq k_0$, Σ_t^k in $B_{2r}(p)$ can be described as $x^n = f^k(x', t)$, where $x' = (x^1, \dots, x^{n-1})$. For instance, since $0 \in \Sigma_{u^k(0)}^k$, we have

$$(2.55) \quad f^k(0, u^k(0)) = 0.$$

From (2.53), (2.54), and (2.55), we deduce that the domain of $f^k(\cdot, t)$ contains $\bar{B}_{\frac{a}{16b}r}^{n-1}$ for every $t \in [u^k(0) - \frac{r^2}{4b}, u^k(0) + \frac{r^2}{4b}]$; moreover,

$$(2.56) \quad \begin{aligned} \frac{a}{8b}r &\leq f^k\left(x', u^k(0) + \frac{r^2}{4b}\right) \leq \frac{r}{2}, \\ -\frac{r}{2} &\leq f^k\left(x', u^k(0) - \frac{r^2}{4b}\right) \leq -\frac{a}{8b}r, \end{aligned}$$

for every $x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}$. Notice that by (2.54) we have

$$(2.57) \quad f^k\left(x', u^k(0) - \frac{r^2}{4b}\right) \leq f^k(x', t) \leq f^k\left(x', u^k(0) + \frac{r^2}{4b}\right).$$

for every $x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}$ and $t \in [u^k(0) - \frac{r^2}{4b}, u^k(0) + \frac{r^2}{4b}]$.

For each $\delta \in (0, \frac{1}{4b})$, let

$$\mathcal{D}_\delta = \bar{B}_{\frac{a}{16b}r}^{n-1} \times \left[u(0) + \delta r^2 - \frac{r^2}{4b}, u(0) - \delta r^2 + \frac{r^2}{4b} \right].$$

It follows from (2.50), (2.52), (2.53), (2.56), (2.57) that any subsequence of $\{f^k|_{\mathcal{D}_\delta}\}$ has a further subsequence that converges smoothly to some function \tilde{f} on \mathcal{D}_δ . We claim that $\tilde{f} = f|_{\mathcal{D}_\delta}$. To prove that, fix

$$t_0 \in \left[u(0) + \delta r^2 - \frac{r^2}{4b}, u(0) - \delta r^2 + \frac{r^2}{4b} \right]$$

and let $\epsilon > 0$ be any sufficiently small number. Note that (2.54) holds with f in place of f^k , so we have

$$\begin{aligned} \{x^n > f(x', t_0 + \epsilon) : x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}\} \cap B_r &\subset \{u > t_0 + \epsilon\}, \\ \{x^n < f(x', t_0 - \epsilon) : x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}\} \cap B_r &\subset \{u < t_0 - \epsilon\}. \end{aligned}$$

By Proposition 2.16, there is $k_\epsilon \in \mathbb{N}$ so that for $k \geq k_\epsilon$ we have

$$\{u > t_0 + \epsilon\} \subset \{u^k > t_0\}, \quad \{u < t_0 - \epsilon\} \subset \{u^k < t_0\},$$

which yields that

$$f(x', t_0 - \epsilon) \leq f^k(x', t_0) \leq f(x', t_0 + \epsilon) \quad \forall x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}, k \geq k_\epsilon.$$

By the squeeze theorem, $f^k(x', t_0) \rightarrow f(x', t_0)$ for every $x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}$, proving the claim.

Therefore, we obtain

$$f^k \xrightarrow{C^\infty} f \quad \text{on } \mathcal{D}_\delta.$$

Note that (2.52), (2.53), (2.54), (2.55), (2.56), and (2.57) hold with f and u in place of f^k and u^k , respectively. Now choose $\delta > 0$ sufficiently small so that

$$\begin{aligned} \frac{a}{12b}r &\leq f\left(x', u(0) - \delta r^2 + \frac{r^2}{4b}\right) \leq \frac{r}{2}, \\ -\frac{r}{2} &\leq f\left(x', u(0) + \delta r^2 - \frac{r^2}{4b}\right) \leq -\frac{a}{12b}r \end{aligned}$$

for every $x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}$. Then when k is large we have

$$\begin{aligned} \frac{a}{16b}r &\leq f^k\left(x', u(0) - \delta r^2 + \frac{r^2}{4b}\right) \leq \frac{r}{2}, \\ -\frac{r}{2} &\leq f^k\left(x', u(0) + \delta r^2 - \frac{r^2}{4b}\right) \leq -\frac{a}{16b}r \end{aligned}$$

for every $x' \in \bar{B}_{\frac{a}{16b}r}^{n-1}$.

Finally, we conclude that the flow $\{\Sigma_t^k\}$ converges smoothly to the flow $\{\Sigma_t\}$ in $\bar{B}_{\frac{a}{16b}r}^{n-1} \times [-\frac{r}{2}, \frac{r}{2}]$ for $t \in [u(0) + \delta r^2 - \frac{r^2}{4b}, u(0) - \delta r^2 + \frac{r^2}{4b}]$ as $k \rightarrow \infty$; when k is large, the flows $\{\Sigma_t^k\}$ and $\{\Sigma_t\}$ are empty in $\bar{B}_{\frac{a}{16b}r}^{n-1} \times [-\frac{a}{16b}r, \frac{a}{16b}r]$ for $t \notin [u(0) + \delta r^2 - \frac{r^2}{4b}, u(0) - \delta r^2 + \frac{r^2}{4b}]$. \square

Now we are in a position to prove Theorem 1.1.

Proof. (of Theorem 1.1) Note that $\mathcal{S} \subset \Omega_{\dot{T}}$ by Proposition 2.6. In view of Lemma 2.21, it suffices to show that $\{\Sigma_t^k\}$ converge locally smoothly to $\{\Sigma_t\}$ in $\Omega_0 \setminus \Omega_{\dot{T}}$ as $k \rightarrow \infty$.

Fix $p \in \Omega_0 \setminus \Omega_{\dot{T}}$, let

$$\tau = \frac{1}{2}u(p) \in (0, \dot{T})$$

and choose $r > 0$ so that

$$B_{2r}(p) \subset \left\{ \frac{3}{2}\tau < u < \frac{3}{2}\dot{T} \right\}.$$

By Proposition 2.16, when k is large we have

$$\left\{ \frac{3}{2}\tau < u < \frac{3}{2}\dot{T} \right\} \subset \left\{ \tau < u^k < 2\dot{T} \right\};$$

thus, by Corollary 2.7 the flow $\{\Sigma_t^k\}$ converges locally smoothly to $\{\Sigma_t\}$ in $B_r(p)$ as $k \rightarrow \infty$. \square

As a consequence, Corollary 2.20 can be improved as follows.

Corollary 2.22. *If the flow $\{\Sigma_t\}$ is regular during $t \in [a, b]$, where $0 < a < b < T_{ext}$, then $\{\Sigma_t^k\}_{a \leq t \leq b}$ is a MCF when k is large and converges smoothly to $\{\Sigma_t\}_{a \leq t \leq b}$ as $k \rightarrow \infty$.*

Proof. As in the proof of Corollary 2.20, choose

$$0 < a' < a < b < b' < T_{ext}$$

and $k_0 \in \mathbb{N}$ such that

$$\{a \leq u^k \leq b\} \subset \{a' < u < b'\} \subset \Omega_0 \setminus \mathcal{S}$$

for every $k \geq k_0$. Then the conclusion follows from Theorem 1.1 that the flow $\{\Sigma_t^k\}$ converges locally smoothly to $\{\Sigma_t^k\}$ in $\{a' \leq u \leq b'\}$. \square

3. SINGULAR SET OF TWO-CONVEX LSF

From now on, the initial hypersurface Σ_0 is assumed to be two-convex. It follows from [CHN] that the singular set \mathcal{S} of $\{\Sigma_t\}$ would consist of round points and/or cylindrical points (i.e., 1-cylindrical points, see [G2] for the definitions).

In this section we aim to establish Theorem 1.2. Firstly, in Section 3.1 and Section 3.2 we apply the theories in [CM2] and [CM3] to investigate the asymptotic behavior of the flow $\{\Sigma_t\}$ and the singular set \mathcal{S} near a round point and a cylindrical point, respectively. Then on the basis of these findings, in Section 3.3 we study the structure of the singular set at the first singular time T_1 under the assumption that that T_1 is an isolated singular time. Lastly, in Section 3.4 we give Proposition 3.13, by which we can apply the result in Section 3.3 recursively to prove Theorem 1.2.

3.1. Round points. In this subsection we will briefly review the behavior of mean-convex LSF near a round point and then conclude with Corollary 3.1.

Now let p be a round point of $\{\Sigma_t\}$. By Brakke's regularity theorem (cf. [B], [I1]), in a small neighborhood $U \subset \Omega_0$ of p , Σ_t would be asymptotically spherical around p for every $t < u(p)$. Choose t_0 less than and sufficiently close to $u(p)$ such that

$$\hat{\Sigma}_{t_0} := \Sigma_{t_0} \cap U$$

is a convex closed connected hypersurface. Let $\hat{\Omega}_{t_0}$ be the region bounded by $\hat{\Sigma}_{t_0}$.¹⁵ As u satisfy (1.2) in the viscosity sense on $\hat{\Omega}_{t_0}$ with the boundary condition $u = t_0$ on $\partial\hat{\Omega}_{t_0} = \hat{\Sigma}_{t_0}$, it follows from the existence and uniqueness theorem in [ES] for solutions to the Dirichlet problem (1.2) that

$$\left\{ \Sigma_t \cap \hat{\Omega}_{t_0} \right\}_{t > t_0}$$

is indeed the mean-convex LSF starting from $\hat{\Sigma}_{t_0}$ at time t_0 .

On the other hand, by [H1] the MCF $\left\{ \tilde{\Sigma}_t \right\}_{t \geq t_0}$ starting from $\hat{\Sigma}_{t_0}$ at time t_0 would shrink monotonically into a point as $t \nearrow \tilde{T}$, where \tilde{T} is the first singular time of $\left\{ \tilde{\Sigma}_t \right\}$. By [ES],

$$\Sigma_t \cap \hat{\Omega}_{t_0} = \tilde{\Sigma}_t$$

for every $t \in [t_0, \tilde{T})$. Thus, we infer that $\Sigma_t \cap \hat{\Omega}_{t_0}$ shrinks to the round point p as $t \nearrow u(p) = \tilde{T}$.

Furthermore, when $u(p) = T_1$ (i.e., the first singular time of the flow), then by the connectedness of Σ_t for $t \in [0, T_1)$, $\Sigma_t \cap \hat{\Omega}_{t_0}$ must be the only component of Σ_t for every t less than and sufficiently close to T_1 . In other words, the whole Σ_t would be asymptotically spherical around p and shrinks to p as $t \nearrow T_1$.

Corollary 3.1. *The round point p of $\{\Sigma_t\}$ is an isolated singular point of the flow and a local maximum point of u . Moreover, if $u(p)$ is the first singular time of the flow, then p would be the unique singular point of $\{\Sigma_t\}$ and a global maximum point of u ; that is to say, $\{\Sigma_t\}$ would shrink to the point p at time $u(p)$ and then vanish.*

3.2. Cylindrical points. We shall begin this subsection with extracting some crucial facts concerning cylindrical points from [CM2] and [CM3] (see also Section 2 in [G2]), and then proceed to prove the main results of this subsection, Proposition 3.9 and Corollary 3.10, which give the local structure of the singular set at the first singular time near a cylindrical point under certain assumptions. The key to establish Proposition 3.9 is to prove two critical lemmas: Lemma 3.6 and Lemma 3.7, which are based upon the properties of cylindrical points referred at the beginning of the subsection.

Now let p be a cylindrical point. For ease of notations (which is valid only for this subsection), let us perform a rigid motion to the flow in space and do a translation in time so as to assume that p is the origin, the singular time $u(p)$ is 0, and that the tangent flow at 0 is $\{\sqrt{-t}\mathcal{C}\}_{t < 0}$, where

$$\mathcal{C} := \frac{S^{n-2}}{\sqrt{2(n-2)}} \times \mathbb{R}.$$

In this case, it is convenient to adopt the following coordinates:

$$x = (y, z) \in \mathbb{R}^{n-1} \times \mathbb{R}.$$

By [CM2] (see also [CM3]), given small positive constants ϕ and ϵ , there are $\delta > 0$ and $L \geq 1$ (depending on $n, \lambda, \phi, \epsilon$) so that if for some $t_0 < 0$ the rescaled level

¹⁵Note that $\hat{\Omega}_{t_0}$ is an open neighborhood of p in Ω_0 .

¹⁶See Proposition 2.5.

set

$$(3.1) \quad \left(\frac{1}{\sqrt{-2t_0}} \Sigma_{2t_0} \right) \cap B_L$$

is δ -close in the C^m topology to some cylinder congruent to \mathcal{C} in B_L , where $m \geq 2$ is an absolute constant (depending on n, λ), then for every $t \in [t_0, 0)$, the rescaled level set

$$\left(\frac{1}{\sqrt{-t}} \Sigma_t \right) \cap B_M$$

would be ϵ -close in the C^m topology¹⁷ to \mathcal{C} in B_M , where

$$M = \frac{\sqrt{2(n-2)}}{\sin \phi};$$

namely, $\frac{1}{\sqrt{-t}} \Sigma_t$ in B_M can be parametrized as a normal graph of a function $f_t(\omega, z)$ over \mathcal{C} whose C^m norm is less than ϵ , where $(\omega, z) \in S^{n-2} \times \mathbb{R}$ are the cylindrical coordinates of \mathcal{C} .¹⁸ After undoing the rescaling, Σ_t in $B_{M\sqrt{-t}}$ can be parametrized as

$$(3.2) \quad x_t(\omega, z) = \left(\left(\sqrt{2(n-2)} + f_t(\omega, z) \right) \sqrt{-t} \omega, z \right), \quad \omega \in S^{n-2}, |z| \lesssim M\sqrt{-t}.$$

Note that such $t_0 < 0$ exists due to Brakke's regularity theorem (cf. [B], [I1]).

Let

$$r = \sqrt{2(n-2)(-t_0)}.$$

It follows that in $B_r \setminus \mathcal{E}_\phi$, where¹⁹

$$(3.3) \quad \mathcal{E}_\phi = \{ |y| \leq |z| \tan \phi \},$$

$\Sigma_t = \{u = t\}$ would be ϵ -close "relative to the scale $\sqrt{-t}$ "²⁰ in the C^m topology to

$$\sqrt{-t} \mathcal{C} = \left\{ |y| = \sqrt{2(n-2)(-t)} \right\}$$

for every $t < 0$. Such a number r is called a (ϕ, ϵ) -**cylindrical scale** of the flow $\{\Sigma_t\}$ at the cylindrical point 0.

Before getting into more details about the cylindrical points from [CM2] and [CM3], let us digress for a moment to discuss the cylindrical points as critical points of the arrival time function u . This is indispensable for Section 4.

Corollary 3.2. *The cylindrical point 0 would never be a local minimum point of u in view of*

$$(3.4) \quad \sup_{B_r \setminus \mathcal{E}_\phi} u \leq u(0) = 0;$$

thus, it is either a local maximum point or a saddle point of u .

As a consequence, the cylindrical point 0 is a saddle point of u if and only if it is not a local maximum point of u , that is, for every $\varepsilon > 0$ there is $x \in B_\varepsilon$ such

¹⁷The closeness in the higher order topology would follow from [EH] and interpolation.

¹⁸That is, \mathcal{C} can be typically parametrized as

$$x(\omega, z) = \left(\sqrt{2(n-2)} \omega, z \right), \quad \omega \in S^{n-2}, z \in \mathbb{R}.$$

¹⁹ \mathcal{E}_ϕ is a solid double cone with apex 0, axis $\{0\} \times \mathbb{R}$ (i.e., the z -axis), and angle ϕ .

²⁰In the sense that after rescaling by the factor $\frac{1}{\sqrt{-t}}$, the two "normalized" hypersurfaces would be ϵ -close in the C^m topology.

that $u(x) > u(0) = 0$ (in other words, the cylindrical point 0 can be approached by points from the superlevel set $\{u > u(0) = 0\}$).

Recall that from the asymptotically cylindrical behavior of the flow $\{\Sigma_t\}$ in $B_r \setminus \mathcal{C}_\phi$, we have

$$\{u > u(0) = 0\} \cap B_r \subset \mathcal{C}_\phi \setminus \{0\}.$$

Note that $\mathcal{C}_\phi \setminus \{0\}$ has two components: the upside and the downside. When the cylindrical points 0 is a saddle point, depending on whether it can be approached by points in $\{u > u(0) = 0\}$ from only one side or both sides, we have the following definition:

Definition 3.3. When the cylindrical point 0 is a saddle point of u , we call it a **one-sided saddle point** of u provided it can be approached by points in $\{u > u(0) = 0\}$ from only one side; in other words, it must be a local maximum point on either the upside or the downside, say the upside, in the sense that there exists $\varepsilon_0 > 0$ so that $u(0) = 0$ is the maximum value of u on $\bar{B}_{\varepsilon_0}^{n-1} \times [0, \varepsilon_0]$.

When the saddle point 0 can be approached by points in $\{u > u(0) = 0\}$ from both sides, it would be called a **two-sided saddle point** of u .

Below we have a remark, which will be used in the discussion of singular components of the bumpy type in Section 4.

Remark 3.4. Suppose that the cylindrical point 0 is a local maximum point of u on the upside, namely, there exists $\varepsilon_0 \in (0, r)$ so that $u(0) = 0$ is the maximum value of u on $\bar{B}_{\varepsilon_0}^{n-1} \times [0, \varepsilon_0]$. By (3.4) we actually have

$$u \leq u(0) = 0 \quad \text{on } B_r \cap \{0 \leq z \leq \varepsilon_0\}.$$

Assume further that there is no other singular points of $\{\Sigma_t\}$ in $B_r \cap \{0 \leq z \leq \varepsilon_0\}$ except the cylindrical point 0. Then we have

$$u < u(0) = 0 \quad \text{on } B_r \cap \{0 \leq z < \varepsilon_0\}.$$

Because otherwise there would exist $(\tilde{y}, \tilde{z}) \in (B_r \setminus \{0\}) \cap \{0 \leq z < \varepsilon_0\}$ such that $u(\tilde{y}, \tilde{z}) = 0$. In view of the asymptotically cylindrical behavior of $\{\Sigma_t\}$, we infer that

$$(\tilde{y}, \tilde{z}) \in (B_r \setminus \{0\}) \cap \{0 \leq z < \varepsilon_0\} \cap \mathcal{C}_\phi \subset B_r \cap \{0 < z < \varepsilon_0\},$$

which yields that (\tilde{y}, \tilde{z}) is an interior local maximum point of u (and hence a critical point of u). This is in contradiction to the assumption that the cylindrical point 0 is the only singular point of $\{\Sigma_t\}$ in $B_r \cap \{0 \leq z \leq \varepsilon_0\}$. Therefore, (in view of the continuity of u at the cylindrical point 0) we have

$$(3.5) \quad \max_{B_r \cap \{z=z_0\}} u < u(0) \quad \forall z_0 \in (0, \varepsilon_0); \quad \max_{B_r \cap \{z=z_0\}} u \rightarrow u(0) \quad \text{as } z_0 \searrow 0.$$

Now let us proceed to the further details regarding the cylindrical points from [CM2] and [CM3]. Firstly, owing to Brakke's regularity theorem, we can rechoose t_0 in (3.1) even closer to 0 so that

$$\left(\frac{1}{\sqrt{-2t_0}} \{u = 2t_0\} \right) \cap B_{2L}$$

is $\frac{\delta}{2}$ -close in the C^m topology to \mathcal{C} in B_{2L} . Applying the theorems in [CM2] to every cylindrical point near 0 (if any) yields the following: there exists $\rho > 0$ (depending

on $n, \lambda, \phi, \epsilon, |t_0|$, and the Lipschitz constant of u)²¹ so that every cylindrical point in $\bar{B}_\rho^{n-1} \times [-\rho, \rho]$ has a uniform (ϕ, ϵ) -cylindrical scale $r = \frac{1}{2}\sqrt{2(n-2)(-t_0)}$.²²

By [CM3] (see also [CM5]), let $\theta = \theta(n) \in (0, 1)$ be an appropriately small constant, there would be small constants $\phi = \phi(n)$ and $\epsilon = \epsilon(n)$ so that every cylindrical point (y_0, z_0) in $\bar{B}_\rho^{n-1} \times [-\rho, \rho]$ has a uniform $(\phi(n), \epsilon(n))$ -cylindrical scale²³ and is contained in a graph $y = \psi(z)$, where

$$\psi : [-\rho, \rho] \rightarrow \bar{B}_\rho^{n-1}$$

is a θ -Lipschitz function with $\psi(0) = 0$,²⁴ and the direction of the 1-dimensional axis of the tangent cylinder at (y_0, z_0) would be θ -close to the direction of the z -axis.

Moreover, in case there exist $-\rho \leq a < b \leq \rho$ so that every point on the curve segment

$$(3.6) \quad \{(y, z) : y = \psi(z), z \in [a, b]\}$$

is a cylindrical point,²⁵ then $\psi \in C^1[a, b]$ and that there are no other singular points in $\bar{B}_\rho^{n-1} \times [a, b]$. Furthermore, in this case the tangent line to the C^1 curve segment at each point would be the 1-dimensional axis of the tangent cylinder at the corresponding point. Notice that

$$u(\psi(z), z) = u(0) \quad \forall z \in [a, b]$$

since ∇u vanishes along the curve (3.6). Consequently, $u(0) = 0$ is the unique singular time of the flow in $\bar{B}_\rho^{n-1} \times [a, b]$.

Remark 3.5. In the above scenario, for each $z_0 \in [a, b]$, the flow is asymptotically cylindrical in $B_r(\psi(z_0), z_0) \setminus \mathcal{C}_\phi^{z_0}$, where $\mathcal{C}_\phi^{z_0}$ is the solid cone with apex $(\psi(z_0), z_0)$, axis aligned with the axis of the tangent cylinder at $(\psi(z_0), z_0)$ (whose direction is θ -close to the direction of the z -axis), and angle ϕ . Particularly, we deduce that

$$\max_{\bar{B}_\rho^{n-1}} u(\cdot, z_0) = u(0) = 0 \quad \forall z_0 \in [a, b].$$

Therefore, $u(0) = 0$ is the maximum value of u on $\bar{B}_\rho^{n-1} \times [a, b]$. In the case where $a = 0$ and $b \in (0, \rho]$, the cylindrical point 0 would be either a local maximum point or a one-sided saddle point of u such that it is a local maximum point on the upside (see Definition 3.3).

Next, we are going to prove two critical lemmas, Lemma 3.6 and Lemma 3.7, from which the main results of this subsection (Proposition 3.9 and Corollary 3.10) follow. The intuition behind Lemma 3.6 is that the ‘‘neck’’ between two cylindrical points should not break before the first singular time.

²¹The intention is to ensure that for any cylindrical point $q \in \bar{B}_\rho^{n-1} \times [-\rho, \rho]$ there holds

$$\frac{|q| + \sqrt{|u(q)|}}{\sqrt{-t_0}}$$

is sufficiently small (depending on n, m, δ).

²²By Footnote 21, r is much larger than ρ , so we may assume that $\bar{B}_{2\rho}^{n-1} \times [-2\rho, 2\rho] \subset B_r$.

²³In the rest of the paper, this would be simply referred to as a cylindrical scale.

²⁴With the θ -Lipschitz condition of ψ and $\psi(0) = 0$, we actually have that

$$\psi(z) \in \bar{B}_{\theta\rho}^{n-1} \subset B_\rho^{n-1}$$

for every $z \in [-\rho, \rho]$.

²⁵By the last paragraph, it is equivalent to say that for every $z \in [a, b]$, there exists a cylindrical point in $B_\rho^{n-1} \times \{z\}$.

Lemma 3.6. *Let the notations be as stated in the above. Note that the parameters ϕ, ϵ, θ are small and that the scale $r \sim \sqrt{-t_0} \gg \rho$.*

Suppose that $u(0) = 0$ is the first singular time of the flow $\{\Sigma_t\}$ and that the cylindrical point 0 is a maximum point of u in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ for some $\varrho \in (0, \rho]$. If there exists another cylindrical point $(y_, z_*) \in \bar{B}_\varrho^{n-1} \times (0, \varrho]$, then the singular set of $\{\Sigma_t\}$ in $\bar{B}_\varrho^{n-1} \times [0, z_*]$ would be a C^1 embedded curve that is composed of cylindrical points and is given by*

$$\{(y, z) : y = \psi(z), z \in [0, z_*]\} \subset \{u = 0\}.$$

Proof. First of all, note that the hypotheses implies that $t = 0$ would be the unique singular time of the flow in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$; particularly, $u(y_*, z_*) = 0$. In addition, in view of the paragraphs between Remark 3.4 and Remark 3.5, to prove the lemma, it suffices to show that for every $z \in (0, z_*)$ there exists a cylindrical point in $B_\varrho^{n-1} \times \{z\}$.

Observe that if $u(y_0, z_0) = 0$ for some $(y_0, z_0) \in B_\varrho^{n-1} \times (0, z_*)$, then it must be a local maximum point of u since $u(0) = 0$ is the maximum value of u on $\bar{B}_\varrho^{n-1} \times [0, \varrho]$; hence (y_0, z_0) would be a singular point of the flow, which by Corollary 3.1 is indeed a cylindrical point. Thus, our goal is to show that for every $z \in (0, z_*)$ there exists a point in $B_\varrho^{n-1} \times \{z\}$ at which u vanishes.

Next, since Σ_t approximates to the cylinder $\sqrt{-t}\mathcal{C}$ in $B_r \setminus \mathcal{C}_\phi$ for $t < 0$, we can choose $t_1 \in (t_0, 0)$ so that

$$(3.7) \quad \Sigma_{t_1} \cap (\bar{B}_\varrho^{n-1} \times \{z\}) \subset \left(B_\varrho^{n-1} \setminus \bar{B}_{z \tan \phi}^{n-1} \right) \times \{z\} \quad \forall z \in [-2\varrho, 2\varrho].$$

Let Θ be the ‘‘tubular’’ closed region in $B_\varrho^{n-1} \times [0, z_*]$ that is bounded from below, above, and lateral by respectively $z = 0$, $z = z_*$, and $\Sigma_{t_1} \approx \sqrt{-t_1}\mathcal{C}$. Consider the set \mathfrak{T} consisting of $t \in [t_1, 0)$ for which there exists a continuous curve on $\Sigma_t \cap \Theta$ that goes from the bottom $z = 0$ to the top $z = z_*$. It is clear that $t_1 \in \mathfrak{T}$ since $\Sigma_{t_1} \approx \sqrt{-t_1}\mathcal{C}$ in Θ . More generally, notice that we can actually find some $t_2 \in (t_1, 0)$ so that for every $t \in [t_1, t_2]$, the condition (3.7) holds with t in place of t_1 (based on the asymptotically cylindrical behavior of the flow in $B_r \setminus \mathcal{C}_\phi$), which yields that $[t_1, t_2] \subset \mathfrak{T}$.

Let

$$(3.8) \quad \hat{T} = \sup \{t \in [t_1, 0) : [t_1, t] \subset \mathfrak{T}\}.$$

We claim that $\hat{T} = 0$. If so, fix $z_0 \in (0, z_*)$, then for every $t \in [t_1, 0) \subset \mathfrak{T}$, as there exists a continuous path on $\Sigma_t \cap \Theta$ going from the bottom to the top and hence must passing through

$$(3.9) \quad \Theta_{z_0} := \Theta \cap \{z = z_0\},$$

we can find a point $(y_t, z_0) \in \Sigma_t \cap \Theta_{z_0}$, which gives that $u(y_t, z_0) = t$. By the compactness of Θ_{z_0} and the continuity of u , we then obtain a point

$$(y_0, z_0) \in \Theta_{z_0} \subset B_\varrho^{n-1} \times \{z_0\}$$

satisfying $u(y_0, z_0) = 0$, proving the lemma. In the rest of the proof, we shall verify the claim.

Suppose the contrary that $\hat{T} < 0$. Let $\hat{\Theta}$ be the extended region of Θ by adding two conical layers Δ_0 and Δ_* in the bottom and top, respectively. The two layers are defined as follows: let $\varphi \in (0, \phi]$ be a small constant (to be determined); let Δ_0 be the closed region that is bounded from below, above, and lateral

by respectively $\{z = -|y| \tan \varphi\}$, $\{z = |y| \tan \varphi\}$, and $\Sigma_{t_1} \approx \sqrt{-t_1} \mathcal{C}$;²⁶ let Δ_* be the closed region that is bounded from below, above, and lateral by respectively $\{z - z_* = -|y - y_*| \tan \varphi\}$, $\{z - z_* = |y - y_*| \tan \varphi\}$, and $\Sigma_{t_1} \approx \sqrt{-t_1} \mathcal{C}$.²⁷ Now let us choose $\varphi \in (0, \phi]$ sufficiently small (depending also on z_*) so that Δ_0 and Δ_* are disjoint.

Recall that both of the cylindrical points 0 and (y_*, z_*) have cylindrical scale r , so the flow is asymptotically cylindrical in both $B_r \setminus \mathcal{C}_\phi$ and $B_r(y_*, z_*) \setminus \mathcal{C}_\phi^*$, where \mathcal{C}_ϕ^* is the counterpart of \mathcal{C}_ϕ for the cylindrical point (y_*, z_*) , namely, \mathcal{C}_ϕ^* is the solid cone with apex (y_*, z_*) , axis aligned with the axis of the tangent cylinder at (y_*, z_*) (whose direction is θ -close to the direction of the z -axis), and angle ϕ . Particularly, the flow is asymptotically cylindrical in both Δ_0 and Δ_* .

In view of $\Sigma_t \cap \Theta_0 \approx \sqrt{-t} \mathcal{C} \cap \Theta_0$ for every $t \in [t_1, \hat{T}/2]$, and that $\partial\Delta_0 \setminus \Sigma_{t_1}$ ²⁸ is contained in $z = \pm |y| \tan \varphi$, there is a uniform distance between $\Sigma_t \cap \Theta_0$ and $\partial\Delta_0 \setminus \Sigma_{t_1}$ for every $t \in [t_1, \hat{T}/2]$. The same is true if we replace Θ_0 and Δ_0 by respectively Θ_{z_*} and Δ_* . Specifically, there is $\sigma > 0$ so that

$$(3.10) \quad \text{dist}(\Sigma_t \cap \Theta_0, \partial\Delta_0 \setminus \Sigma_{t_1}) \geq \sigma, \quad \text{dist}(\Sigma_t \cap \Theta_{z_*}, \partial\Delta_* \setminus \Sigma_{t_1}) \geq \sigma$$

for every $t \in [t_1, \hat{T}/2]$.

In addition, the asymptotically cylindrical behavior of the flow in Δ_0 and Δ_* implies that there is not any singular points in

$$(\Delta_0 \cup \Delta_*) \cap \left\{ t_1 \leq u \leq \hat{T}/2 \right\}.$$

Also, recall that the only singular time of the flow in

$$\Theta \subset \bar{B}_\varrho^{n-1} \times [0, \varrho]$$

is 0. Thus, we infer that $\hat{\Theta} \cap \left\{ t_1 \leq u \leq \hat{T}/2 \right\}$ is a compact set consisting of regular points (so $\nabla u \neq 0$), which yields that

$$(3.11) \quad K := \max_{\hat{\Theta} \cap \{t_1 \leq u \leq \hat{T}/2\}} |\nabla u|^{-1}$$

is a finite positive number.

Now choose $t_3 \in [t_2, \hat{T})$ and $t_4 \in (\hat{T}, \hat{T}/2)$ so that

$$(3.12) \quad t_4 - t_3 < \frac{\sigma}{K}.$$

Since $t_3 \in \mathfrak{T}$, there exists a continuous curve

$$\gamma_{t_3} : [0, 1] \rightarrow \Sigma_{t_3} \cap \Theta$$

such that

$$(3.13) \quad \gamma_{t_3}(0) \in \Sigma_{t_3} \cap \Theta_0, \quad \gamma_{t_3}(1) \in \Sigma_{t_3} \cap \Theta_{z_*}.$$

Let Φ_τ be the (local) flow generated by the vector field $\frac{\nabla u}{|\nabla u|^2}$.²⁹ Because $\gamma_{t_3}[0, 1] \subset \Sigma_{t_3} \cap \Theta$ is a compact set consisting of regular points (see the last paragraph), Φ_τ

²⁶Because $\Sigma_{t_1} \cap \Delta_0 \subset \Sigma_{t_1} \cap (B_r \setminus \mathcal{C}_\phi)$.

²⁷Because $\Sigma_{t_1} \cap \Delta_* \subset \Sigma_{t_1} \cap (B_r \setminus \mathcal{C}_\phi)$ by (3.7).

²⁸It is the union of the top and bottom boundaries of Δ_0 .

²⁹For a regular point x , $\tau \mapsto \Phi_\tau(x)$ is the unique integral curve of the vector field $\frac{\nabla u}{|\nabla u|^2}$ such that $\Phi_0(x) = x$, which has short time existence. Note that if $u(x) = t$, then $u(\Phi_\tau(x)) = t + \tau$.

acts on $\gamma_{t_3} [0, 1]$ for $\tau \geq 0$ sufficiently small with $\Phi_\tau \circ \gamma_{t_3}$ being a continuous curve lying on $\Sigma_{t_3+\tau} \cap \text{int } \hat{\Theta}$. In fact, by noticing that the curve $\Phi_\tau \circ \gamma_{t_3}(s)$ will never hit the lateral boundary of $\hat{\Theta}$ (which is contained in Σ_{t_1}) and taking (3.10), (3.11), and (3.12) into account, we infer that so long as $\tau \in [0, t_4 - t_3]$, Φ_τ can act on $\gamma_{t_3} [0, 1]$ with

$$\gamma_{t_3+\tau}^{(0)} := \Phi_\tau \circ \gamma_{t_3} : [0, 1] \rightarrow \Sigma_{t_3+\tau} \cap \text{int } \hat{\Theta}$$

being a continuous curve that, in view of (3.10), (3.11), (3.12), and (3.13), satisfies

$$(3.14) \quad \gamma_{t_3+\tau}^{(0)}(0) \in \Delta_0, \quad \gamma_{t_3+\tau}^{(0)}(1) \in \Delta_*.$$

To finish the proof, we need to modify $\gamma_t^{(0)}$ for each $t \in (t_3, t_4]$ so as to make it stay on $\Sigma_t \cap \Theta$ (instead of $\Sigma_t \cap \hat{\Theta}$), start from somewhere on $\Sigma_t \cap \Theta_0$, and end up on $\Sigma_t \cap \Theta_{z^*}$. If so, we would have $[t_1, t_4] \subset \mathfrak{I}$, which gives the desired contradiction that $t_4 > \hat{T}$ (see (3.8)).

The aforementioned modification of $\gamma_t^{(0)}$ for each $t \in (t_3, t_4]$ will be done in two steps. In the first step we would deal with the “bottom” part. Specifically, we shall find a continuous map $\mathfrak{J}_t : \Sigma_t \cap \hat{\Theta} \rightarrow \Sigma_t \cap (\Theta \cup \Delta_*)$ (as will be seen in the next paragraph) that pushes the portions of $\gamma_t^{(0)}$ lying on $\Sigma_t \cap \Delta_0 \setminus \Theta$ continuously to $\Sigma_t \cap \Delta_0 \cap \Theta$ and keeps the rest invariant, so we would obtain a new continuous curve:

$$\mathfrak{J}_t \circ \gamma_t^{(0)} : [0, 1] \rightarrow \Sigma_t \cap (\Theta \cup \Delta_*).$$

Notice that $\mathfrak{J}_t \circ \gamma_t^{(0)}(0)$ might not be on $\Sigma_t \cap \Theta_0$; instead, by (3.14) we have

$$\mathfrak{J}_t \circ \gamma_t^{(0)}(0) \in \Sigma_t \cap \Theta \cap \Delta_0.$$

Since $\Sigma_t \cap \Theta \cap \Delta_0 \approx \sqrt{-t} \mathcal{C} \cap \Theta \cap \Delta_0$, we can find a curve $\varsigma_t : [0, 1] \rightarrow \Sigma_t \cap \Theta \cap \Delta_0$ so that

$$\varsigma_t(0) \in \Sigma_t \cap \Theta_0, \quad \varsigma_t(1) = \mathfrak{J}_t \circ \gamma_t^{(0)}(0).$$

Then jointing this two paths together gives a new curve

$$\gamma_t^{(1)}(s) = \begin{cases} \varsigma_t(2s), & s \in [0, \frac{1}{2}] \\ \mathfrak{J}_t \circ \gamma_t^{(0)}(2s-1), & s \in [\frac{1}{2}, 1] \end{cases}$$

satisfying

$$\begin{aligned} \gamma_t^{(1)} &: [0, 1] \rightarrow \Sigma_t \cap (\Theta \cup \Delta_*) \\ \gamma_t^{(1)}(0) &\in \Sigma_t \cap \Theta_0, \quad \gamma_t^{(1)}(1) \in \Sigma_t \cap \Delta_*, \end{aligned}$$

completing the first step. In the second step, we deal with the “top” part using essentially the same trick as in the first step, so we are able to transform $\gamma_t^{(1)}$ into a new curve $\gamma_t^{(2)}$ satisfying

$$\begin{aligned} \gamma_t^{(2)} &: [0, 1] \rightarrow \Sigma_t \cap \Theta \\ \gamma_t^{(2)}(0) &\in \Sigma_t \cap \Theta_0, \quad \gamma_t^{(2)}(1) \in \Sigma_t \cap \Theta_{z^*}. \end{aligned}$$

Finally, $\gamma_t = \gamma_t^{(2)}$ is the desired continuous curve.

The last piece to complete the entire proof is to construct the map

$$\mathfrak{J}_t : \Sigma_t \cap \hat{\Theta} \rightarrow \Sigma_t \cap (\Theta \cup \Delta_*)$$

for each $t \in (t_3, t_4]$ as mentioned in the last paragraph. As $\Sigma_t \cap \Delta_0 \approx \sqrt{-t} \mathcal{C} \cap \Delta_0$, we can parametrize $\Sigma_t \cap \Delta_0$ using the cylindrical coordinates (ω, z) defined in (3.2). Then we define a map (between submanifolds)

$$\mathfrak{J}_t : \Sigma_t \cap \Delta_0 \rightarrow \Sigma_t \cap \Delta_0 \cap \Theta$$

in terms of local coordinates as

$$\mathfrak{J}_t(\omega, z) = (\omega, z_+),$$

where $z_+ = \max\{z, 0\}$. It is not hard to see that the map \mathfrak{J}_t can be extended on $\Sigma_t \cap \hat{\Theta}$ in a continuous manner such that $\mathfrak{J}_t = \text{id}$ on $\Sigma_t \cap \hat{\Theta} \setminus \Delta_0$. \square

In contrast to Lemma 3.6, in which the cylindrical point 0 is assumed to be a local maximum point of u on the upside, in Lemma 3.7 we consider the case where the cylindrical point 0 is not a local maximum point on the upside. The additional assumption that $u(0) = 0$ is locally the unique singular time on the upside is of vital importance for the lemma.

Lemma 3.7. *Let notations be as defined in Lemma 3.6. Suppose that 0 is the first singular time of the flow $\{\Sigma_t\}$ and that the cylindrical point 0 is not a local maximum point of u in $\bar{B}_\rho^{n-1} \times [0, \rho]$ in the sense that for every $\varepsilon > 0$, there is $x \in B_\varepsilon \cap (\bar{B}_\rho^{n-1} \times [0, \rho])$ such that $u(x) > u(0) = 0$.*

If there exists $\varrho \in (0, \rho]$ so that $u(0) = 0$ is the unique singular time of the flow in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ for some $\varrho \in (0, \rho]$, then the cylindrical point 0 would be the unique singular point of the flow in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$.

Proof. Suppose the contrary that there is another singular point (y_*, z_*) in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$. Note that $u(y_*, z_*) = 0$ by hypothesis and that (y_*, z_*) must be a cylindrical point by Corollary 3.1. Moreover, in view of the asymptotically cylindrical behavior of the flow in $B_r \setminus \mathcal{C}_\phi$, we infer that $(y_*, z_*) \in \mathcal{C}_\phi$ with $z_* > 0$.

Recall that every cylindrical point in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ is located on a θ -Lipschitz graph $y = \psi(z)$ and has a cylindrical scale r . It follows that the flow is asymptotically cylindrical in $B_r(y_*, z_*) \setminus \mathcal{C}_\phi^*$, where \mathcal{C}_ϕ^* is the solid cone with apex (y_*, z_*) , axis aligned with the axis of the tangent cylinder at (y_*, z_*) (whose direction is θ -close to the direction of the z -axis), and angle ϕ . Thus, we obtain

$$(3.15) \quad \bar{B}_\varrho^{n-1} \times \{0\} \subset (B_r \setminus \mathcal{C}_\phi) \cup \{0\} \subset \{u \leq u(0) = 0\},$$

$$(3.16) \quad \bar{B}_\varrho^{n-1} \times \{z_*\} \subset (B_r(y_*, z_*) \setminus \mathcal{C}_\phi^*) \cup \{(y_*, z_*)\} \subset \{u \leq u(y_*, z_*) = 0\}.$$

Now (using the asymptotically cylindrical behavior of the flow in $B_r \setminus \mathcal{C}_\phi$) choose $t_1 \in (t_0, 0)$ such that

$$\Sigma_{t_1} \cap (\bar{B}_\varrho^{n-1} \times \{z\}) \subset \left(B_\varrho^{n-1} \setminus \bar{B}_{z \tan \phi}^{n-1} \right) \times \{z\} \quad \forall z \in [0, \varrho].$$

Let Θ be the closed region in $B_\varrho^{n-1} \times [0, z_*]$ that is bounded from below, above, and lateral by respectively $z = 0$, $z = z_*$, and $\Sigma_{t_1} \approx \sqrt{-t_1} \mathcal{C}$. Since $u(0) = 0$ is not a local maximum value of u on $\bar{B}_\varrho^{n-1} \times [0, \varrho]$, we have

$$\hat{T} = \max_{\Theta} u > 0.$$

Choose $p \in \Theta$ so that $u(p) = \hat{T}$. In view of (3.15), (3.16), and that the lateral boundary of Θ is contained in Σ_{t_1} , p is not on $\partial\Theta$ and therefore is an interior

maximum point (and hence a critical point) of u . Thus, \hat{T} is another singular time of the flow, contradicting the hypothesis. \square

Before moving on to the primary conclusions of this subsection, let us make the following remark, which will be used in the discussion of singular components of the splitting/bumpy type in Section 4.

Remark 3.8. In Lemma 3.7, note that the cylindrical point 0 must be either a two-sided saddle point of u or a one-sided saddle point of u such that it is a local maximum point on the downside (see Definition 3.3). Because of the asymptotically cylindrical behavior of $\{\Sigma_t\}$ in $B_r \setminus \mathcal{C}_\phi$, we have $\Sigma_\tau \approx \sqrt{-\tau}\mathcal{C}$ in B_r , where

$$\tau = u(0) - \frac{r^2}{4(n-2)} = -\frac{r^2}{4(n-2)}.$$

By the continuity of u at the cylindrical point 0, we can find $\hat{r} \in (0, r)$ so that

$$(3.17) \quad \min_{\mathcal{C}_\phi \cap \{0 \leq z \leq \hat{r}\}} u > \tau.$$

Moreover, since the cylindrical point 0 is not a local maximum point of u on the top, we can choose $\hat{\varepsilon} \in (0, \hat{r})$ such that

$$\hat{t} := \max_{\hat{\mathcal{T}} \cap \{z = \hat{\varepsilon}\}} u > u(0) = 0,$$

where

$$(3.18) \quad \hat{\mathcal{T}} := \bar{\Omega}_\tau \cap B_r \cap \{0 \leq z \leq \hat{\varepsilon}\}$$

is a ‘‘tubular’’ closed region bounded by Σ_τ , $\{z = 0\}$, and $\{z = \hat{\varepsilon}\}$ from the lateral, bottom, and top, respectively. By the asymptotically cylindrical behavior of $\{\Sigma_t\}$ in $B_r \setminus \mathcal{C}_\phi$ and the intermediate value theorem, we then deduce that

$$(3.19) \quad \Sigma_t \cap \{z = \hat{\varepsilon}\} \cap \hat{\mathcal{T}} \neq \emptyset \quad \forall t \in [\tau, \hat{t}],$$

$$(3.20) \quad \Sigma_t \cap \hat{\mathcal{T}} \subset \text{int } \mathcal{C}_\phi \quad \forall t \in (0, \hat{t}].$$

In case where $u(0) = 0$ is the unique singular time of $\{\Sigma_t\}$ in \bar{B}_r^+ , $\Sigma_t \cap \hat{\mathcal{T}}$ would be a hypersurface contained in $\text{int } \mathcal{C}_\phi$ for every $t \in (0, \hat{t}]$.

Now we are all set to prove the main result of this subsection.

Proposition 3.9. *With the same notations as in Lemma 3.6 and Lemma 3.7. Assume that $u(0) = 0$ is the first singular time of the flow $\{\Sigma_t\}$ and also the unique singular time of the flow in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ for some $\varrho \in (0, \rho]$.*

Then either the cylindrical point 0 is an isolated³⁰ singular point in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$, or there exists $z_ \in (0, \varrho]$ so that the singular set of the flow in $\bar{B}_\varrho^{n-1} \times [0, z_*]$ is a C^1 embedded curve that is comprised of cylindrical points and is defined by*

$$\{(y, z) : y = \psi(z), z \in [0, z_*]\}.$$

Proof. If the cylindrical point 0 is an isolated singular point in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$, then we are done. So let us assume that 0 is not an isolated singular point in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$; particularly, 0 is not the only singular points in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$. So by Lemma 3.7 we

³⁰It means that 0 is the unique singular point in $B_\varepsilon \cap (\bar{B}_\varrho^{n-1} \times [0, \varrho])$ for some $\varepsilon > 0$.

infer that the cylindrical point 0 must be a local maximum point of u in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ in the sense that for some $\tilde{\varrho} \in (0, \varrho]$ we have

$$u(x) \leq u(0) = 0 \quad \forall x \in \bar{B}_{\tilde{\varrho}}^{n-1} \times [0, \tilde{\varrho}].$$

Namely, the cylindrical point 0 is a maximum point of u on $\bar{B}_{\tilde{\varrho}}^{n-1} \times [0, \tilde{\varrho}]$.

Since 0 is not an isolated singular point in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$, we can find a singular point $(y_*, z_*) \neq 0$ in $\bar{B}_{\tilde{\varrho}}^{n-1} \times [0, \tilde{\varrho}]$, which has to be a cylindrical point by Corollary 3.1. Recalling that all cylindrical points in $\bar{B}_\varrho^{n-1} \times [0, \varrho]$ must lie on the graph $y = \psi(z)$, we deduce that $z_* > 0$ (because $\psi(0) = 0$). Then it follows from Lemma 3.6 that the singular set of the flow in $\bar{B}_\varrho^{n-1} \times [0, z_*]$ is a C^1 embedded curve consisting of purely cylindrical points that is given by

$$\{(y, z) : y = \psi(z), z \in [0, z_*]\} \subset \mathcal{S} \cap \Sigma_0.$$

□

Since the analogous result of Proposition 3.9 holds for the flow on the downside $\bar{B}_\varrho^{n-1} \times [-\varrho, 0]$ as well. Combining them together yields the following corollary.

Corollary 3.10. *With the same notations as in Proposition 3.9. If $u(0) = 0$ is the first singular time of the flow $\{\Sigma_t\}$ and also the unique singular time of the flow in $\bar{B}_\varrho^{n-1} \times [-\varrho, \varrho]$ for some $\varrho > 0$.*

Then either the cylindrical point 0 is an isolated singular point in $\bar{B}_\varrho^{n-1} \times [-\varrho, \varrho]$, or there exist $z_^+ \in [0, \varrho]$ and $z_*^- \in [-\varrho, 0]$, at least one of which is nonzero, so that the singular set of the flow in $\bar{B}_\varrho^{n-1} \times [z_*^-, z_*^+]$ is a C^1 embedded curve consisting of purely cylindrical points given by*

$$\{(y, z) : y = \psi(z), z \in [z_*^-, z_*^+]\};$$

moreover, in case that one of $\{z_^-, z_*^+\}$ is 0, say $z_*^- = 0$, then there exists $\varepsilon > 0$ so that the cylindrical point 0 is the unique singular point in $B_\varepsilon \cap (\bar{B}_\varrho^{n-1} \times [-\varrho, 0])$.*

3.3. Singular set at the first singular time. In this subsection we shall consider the structure of the singular set at the first singular time T_1 under the assumption that T_1 is an isolated singular time of the flow. The main result (see the following proposition) is based on Corollary 3.1 (in Section 3.1) and Corollary 3.10 (in Section 3.2). This will be utilized in Section 3.4 to prove Theorem 1.2.

Proposition 3.11. *Suppose either $T_1 = T_{ext}$, or that $T_1 < T_{ext}$ is an isolated singular time of the flow $\{\Sigma_t\}$, that is, there exists a “second” singular time $T_2 \in (T_1, T_{ext}]$ so that $\{\Sigma_t\}$ is regular during $t \in (T_1, T_2)$. Then the singular set of $\{\Sigma_t\}$ at the first singular time, i.e.,*

$$\mathcal{S} \cap \Sigma_{T_1},$$

has finitely many connected components, each of which is either a single (round or cylindrical) point, or a compact C^1 embedded curve (with or without boundary) consisting of cylindrical points.

Proof. Firstly, recall that \mathcal{S} is a closed subset of Ω_0 (see Lemma 2.10) and note that $\Sigma_{T_1} = \{u = T_1\}$ is a closed subset of Ω_0 owing to the continuity of u . Consequently, $\mathcal{S} \cap \Sigma_{T_1}$ is a closed subset of Ω_0 and hence a compact set.

Secondly, if $\mathcal{S} \cap \Sigma_{T_1}$ has a round point, then by Corollary 3.1, this round point would be the only element of $\mathcal{S} \cap \Sigma_{T_1}$ and hence the proposition would be proved.

So for the rest of the proof let us assume that $\mathcal{S} \cap \Sigma_{T_1}$ comprises purely cylindrical points.

Fix $p \in \mathcal{S} \cap \Sigma_{T_1}$. Note that we can find $r > 0$ so that T_1 is the only singular time of the flow in $B_r(p)$. This is obviously true when $T_1 = T_{ext}$; in the case where $T_1 < T_{ext}$, since $\{u \geq T_2\}$ is a compact set in Ω_0 and $p \notin \{u \geq T_2\}$, there exists $r > 0$ such that $B_r(p) \cap \{u \geq T_2\} = \emptyset$, which means that

$$(3.21) \quad B_r(p) \subset \{u < T_2\},$$

and so there are no other singular times than T_1 in $B_r(p)$. Thus, it follows from Corollary 3.10 that there is $\varepsilon \in (0, r]$ so that one of the following holds:

- (1) $\mathcal{S} \cap B_\varepsilon(p) = \{p\}$;
- (2) $\mathcal{S} \cap B_\varepsilon(p)$ is a C^1 embedded curve with p being an interior point;
- (3) $\mathcal{S} \cap B_\varepsilon(p)$ is a C^1 embedded curve with p being a boundary point.

Notice that $\mathcal{S} \cap B_\varepsilon(p) = \mathcal{S} \cap \Sigma_{T_1} \cap B_\varepsilon(p)$ by (3.21).

Therefore, by the compactness of $\mathcal{S} \cap \Sigma_{T_1}$, we infer that $\mathcal{S} \cap \Sigma_{T_1}$ is a finite disjoint union of points and/or compact C^1 embedded curves (with or without boundary). \square

3.4. Singular set at subsequent singular times. We shall prove Theorem 1.2 in the end of this subsection. The idea of the proof is as follows. By Proposition 3.13 (which will be proved in this subsection), the superlevel set of the arrival time function u at the first singular time T_1 is a finite disjoint union of open connected sets, namely,

$$\Omega_{T_1} = \{u > T_1\} = \bigsqcup_{i=1}^{m_1} \Omega_{T_1}^{(i)},$$

and so

$$\Sigma_t = \bigsqcup_{i=1}^{m_1} \left(\Sigma_t \cap \Omega_{T_1}^{(i)} \right), \quad t > T_1;$$

moreover, each $\Sigma_t \cap \Omega_{T_1}^{(i)}$ is itself a two-convex LSF for $t \in (T_1, \infty)$ and a MCF of closed connected hypersurfaces for $t \in (T_1, T_2)$, where T_2 is the second singular time of $\{\Sigma_t\}$. It follows that the singular set of $\{\Sigma_t\}$ at time T_2 is indeed the union

of the singular sets of $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t > T_1}$'s at their "first" singular time, where \hat{i} 's are

those indices such that $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t > T_1}$ does become singular at $t = T_2$. In this

way, Proposition 3.11 from Section 3.3 can be applied to each $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t > T_1}$ to study the singular set of $\{\Sigma_t\}$ at time T_2 . Such process can be repeated to study the singular set of $\{\Sigma_t\}$ at all subsequent singular times $T_1 < T_2 < \dots < T_{ext}$.

To prove Proposition 3.13, we need the following lemma.

Lemma 3.12. *For each regular time $t \in (0, T_{ext})$, $\Sigma_t = \{x \in \Omega_0 : u(x) = t\}$ is a finite disjoint union of two-convex³¹ closed connected hypersurfaces*

$$\Sigma_t^{(1)}, \dots, \Sigma_t^{(m)};$$

³¹With respect to the inward unit normal.

moreover, $\Omega_t = \{x \in \Omega_0 : u(x) > t\}$ is the disjoint union of

$$\Omega_t^{(1)}, \dots, \Omega_t^{(m)},$$

where $\Omega_t^{(i)}$ is the open connected set bounded by $\Sigma_t^{(i)}$ for $i \in \{1, \dots, m\}$.

Proof. Let $t \in (0, T_{ext})$ be a regular value of u . By the implicit function theorem, Σ_t is a closed hypersurface in Ω_0 . As Σ_t is compact, it has finitely many connected components, say $\Sigma_t^{(1)}, \dots, \Sigma_t^{(m)}$, each of which is a closed connected hypersurface in Ω_0 . By the Jordan-Brouwer separation theorem (see [GP]), each $\Sigma_t^{(i)}$ would bound a region $\Omega_t^{(i)}$ contained³² in Ω_0 . Because Σ_t is two-convex with respect to $\frac{\nabla u}{|\nabla u|}$ (cf. [CHN]), so is $\Sigma_t^{(i)}$. Note that the unit normal $\frac{\nabla u}{|\nabla u|}$ is either entirely inward or entirely outward for each $\Sigma_t^{(i)}$. In view of the fact that every closed hypersurface has a point where it is convex with respect to the inward unit normal vector, we conclude that $\frac{\nabla u}{|\nabla u|}$ is entirely inward for each $\Sigma_t^{(i)}$.

To see that $\Omega_t^{(i)} \subset \Omega_t$, firstly note that because the vector field $\frac{\nabla u}{|\nabla u|}$ on $\Sigma_t^{(i)}$ points toward $\Omega_t^{(i)}$, we are able to find a small tubular open neighborhood U of $\Sigma_t^{(i)}$ so that $\Omega_t^{(i)} \cap U \subset \Omega_t$. Were $\Omega_t^{(i)} \setminus \Omega_t \neq \emptyset$, we would have

$$p \in \Omega_t^{(i)} \setminus \Omega_t \subset \bar{\Omega}_t^{(i)} \setminus U$$

so that

$$u(p) = \min_{\bar{\Omega}_t^{(i)} \setminus U} u \leq t.$$

It would follow that $p \in \Omega_t^{(i)}$ with

$$u(p) = \min_{\bar{\Omega}_t^{(i)}} u;$$

particularly, p is a critical point of u . However, u has no local minimum points (see Corollary 3.1 and Corollary 3.2). Thus we obtain $\Omega_t^{(i)} \subset \Omega_t$.

Recall that both $\Omega_t^{(i)}$ and $\mathbb{R}^n \setminus \bar{\Omega}_t^{(i)}$ are open connected (and hence path-connected) by the Jordan-Brouwer separation theorem. When $m > 1$, for every $j \neq i$,

$$\begin{aligned} \Sigma_t^{(i)} \cap \Sigma_t^{(j)} &= \emptyset, \\ \Sigma_t^{(i)} \cap \Omega_t^{(j)} &\subset \Sigma_t \cap \Omega_t = \emptyset, \end{aligned}$$

giving that

$$\Sigma_t^{(i)} \subset \mathbb{R}^n \setminus \bar{\Omega}_t^{(j)}.$$

Notice that $\mathbb{R}^n \setminus \bar{\Omega}_t^{(j)}$ is open connected. Then it follows from the path-connectedness of $\Omega_t^{(i)}$ that

$$\bar{\Omega}_t^{(i)} \subset \mathbb{R}^n \setminus \bar{\Omega}_t^{(j)},$$

that is, $\bar{\Omega}_t^{(i)} \cap \bar{\Omega}_t^{(j)} = \emptyset$.

To finish the proof, we have to show that

$$\Omega_t \subset \Omega_t^{(1)} \cup \dots \cup \Omega_t^{(m)}.$$

³²The mod 2 winding number of $\Sigma_t^{(i)}$ around every point in $\mathbb{R}^n \setminus \bar{\Omega}_0$ is 0 (which can be seen by taking a point far away from $\Sigma_t^{(i)}$ and noting that $\mathbb{R}^n \setminus \bar{\Omega}_0$ is path-connected with $(\mathbb{R}^n \setminus \bar{\Omega}_0) \cap \Sigma_t^{(i)} = \emptyset$), so $\Omega_t^{(i)}$ is contained in Ω_0 .

Suppose the contrary that there is $p \in \Omega_t$ such that $p \notin \Omega_t^{(i)}$ for every i . Let

$$R = \sup \{r > 0 : B_r(p) \subset \Omega_t\}.$$

Then $B_R(p) \cap \bar{\Omega}_t^{(i)} = \emptyset$ for every i and $\partial B_R(p) \cap \Sigma_t \neq \emptyset$. Choose $q \in \partial B_R(p) \cap \Sigma_t$. Then $q \in \partial B_R(p) \cap \Sigma_t^{(i_0)}$ for some i_0 . Since $\nabla u(q) \neq 0$ points toward $\Omega_t^{(i_0)}$, there exists $\delta > 0$ so that

$$B_\delta(q) \setminus \bar{\Omega}_t^{(i_0)} \subset \{u < t\},$$

contradicting that $\emptyset \neq B_R(p) \cap B_\delta(q) \subset \Omega_t$. \square

Proposition 3.13. *Suppose that $T_1 < T_{ext}$ is an isolated singular time of $\{\Sigma_t\}$, that is, there exists a singular time $T_2 \in (T_1, T_{ext}]$ so that the flow is regular during the time period (T_1, T_2) . Then Ω_{T_1} is a finite disjoint union of open connected sets*

$$\Omega_{T_1}^{(1)}, \dots, \Omega_{T_1}^{(m_1)}.$$

Moreover, for each $i \in \{1, \dots, m_1\}$,

$$\Sigma_t^{(i)} := \Sigma_t \cap \Omega_{T_1}^{(i)}, \quad t > T_1$$

is a two-convex MCF of closed connected hypersurfaces for $t \in (T_1, T_2)$ and a two-convex LSF for $t \in (T_1, \infty)$.³³ Notice that as $\Sigma_t \subset \Omega_{T_1}$ for $t > T_1$, we have

$$\Sigma_t = \bigsqcup_{i=1}^{m_1} \Sigma_t^{(i)} \quad \forall t > T_1.$$

Proof. Fix $t_1 \in (T_1, T_2)$. By Lemma 3.12, Σ_{t_1} is a finite disjoint union of closed connected hypersurfaces $\Sigma_{t_1}^{(1)}, \dots, \Sigma_{t_1}^{(m_1)}$ that are two-convex with respect to the inward normal; moreover, Ω_{t_1} is the disjoint union of $\Omega_{t_1}^{(1)}, \dots, \Omega_{t_1}^{(m_1)}$, where $\Omega_{t_1}^{(i)}$ is the open connected set bounded by $\Sigma_{t_1}^{(i)}$.

Since u satisfies (1.2) in the viscosity sense on $\Omega_{t_1}^{(i)}$ with the boundary condition $u = t_1$ on $\partial\Omega_{t_1}^{(i)} = \Sigma_{t_1}^{(i)}$, the existence and uniqueness theorem in [ES] yields that

$$\Sigma_t^{(i)} := \Sigma_t \cap \Omega_{t_1}^{(i)} \quad t > t_1$$

is indeed the mean-convex LSF starting from $\Sigma_{t_1}^{(i)}$ at time t_1 . In addition, since the flow $\{\Sigma_t\}$ is a two-convex MCF for $t \in (T_1, T_2)$, the flow $\{\Sigma_t^{(i)}\}$ is also a two-convex MCF for $t \in [t_1, T_2)$.

It follows that for every $t \in [t_1, T_2)$, each $\Sigma_t^{(i)}$ is a closed connected hypersurface and hence is exactly a connected component of Σ_t . Let $\Omega_t^{(i)}$ be the open connected set bounded by $\Sigma_t^{(i)}$ for $t \in (t_1, T_2)$. Since $\{\Sigma_t^{(i)}\}_{t \in [t_1, T_2)}$ is a mean-convex MCF, $\{\Omega_t^{(i)}\}_{t \in [t_1, T_2)}$ is contracting. By Lemma 3.12, Ω_t is the disjoint union of $\Omega_t^{(1)}, \dots, \Omega_t^{(m_1)}$ for $t \in [t_1, T_2)$.

Replacing $t_1 \in (T_1, T_2)$ in the above by a decreasing sequence tending to T_1 , we can extend³⁴ the flow $\{\Sigma_t^{(i)}\}$ backward in time so as to obtain that $\{\Sigma_t^{(i)}\}_{t \in (T_1, T_2)}$

³³In the sense that for every $t_1 \in (T_1, T_2)$, $\{\Sigma_t^{(i)}\}_{t \geq t_1}$ is the LSF starting from $\Sigma_{t_1}^{(i)}$.

³⁴On the basis of the uniqueness theorem for MCF and LSF, and also the semigroup property of LSF operator (cf. [ES]).

is a two-convex MCF of closed connected hypersurfaces, that $\{\Sigma_t^{(i)}\}_{t>T_1}$ is a two-convex LSF, and that Σ_t is the disjoint union of $\Sigma_t^{(1)}, \dots, \Sigma_t^{(m_1)}$ for every $t \in (T_1, T_2)$. Also, $\Omega_t^{(i)}$ is defined for every $t \in (T_1, T_2)$ as the open connected set bounded by $\Sigma_t^{(i)}$. We then have that $\{\Omega_t^{(i)}\}_{t \in (T_1, T_2)}$ is contracting and that Ω_t is the disjoint union of $\Omega_t^{(1)}, \dots, \Omega_t^{(m_1)}$ for $t \in (T_1, T_2)$.

For each $i \in \{1, \dots, m_1\}$ let

$$\Omega_{T_1}^{(i)} = \bigcup_{t \in (T_1, T_2)} \Omega_t^{(i)}.$$

Since $\Omega_t^{(i)}$ is open connected for every $t \in (T_1, T_2)$, so is $\Omega_{T_1}^{(i)}$. When $m_1 > 1$, for every $j \neq i$ we would have $\Omega_{T_1}^{(i)} \cap \Omega_{T_1}^{(j)} = \emptyset$; otherwise, choose $p \in \Omega_{T_1}^{(i)} \cap \Omega_{T_1}^{(j)}$, then by the contracting property of $\Omega_t^{(i)}$ and $\Omega_t^{(j)}$, there would exist $t_1 \in (T_1, T_2)$ so that $p \in \Omega_{t_1}^{(i)} \cap \Omega_{t_1}^{(j)}$, contradicting that $\Omega_{t_1}^{(i)} \cap \Omega_{t_1}^{(j)} = \emptyset$. In addition, note that

$$\begin{aligned} \Omega_{T_1} &= \{u > T_1\} = \bigcup_{t \in (T_1, T_2)} \{u > t\} = \bigcup_{t \in (T_1, T_2)} \Omega_t \\ &= \bigcup_{t \in (T_1, T_2)} \bigcup_{i=1}^{m_1} \Omega_t^{(i)} = \bigcup_{i=1}^{m_1} \bigcup_{t \in (T_1, T_2)} \Omega_t^{(i)} = \bigcup_{i=1}^{m_1} \Omega_{T_1}^{(i)}. \end{aligned}$$

Lastly, given a sequence $\{t_k\}_{k \in \mathbb{N}} \subset (T_1, T_2)$ such that $t_k \searrow T_1$, by construction $\{\Sigma_t^{(i)}\}_{t \geq t_k}$ is a mean-convex LSF starting from $\Sigma_{t_k}^{(i)}$ at time t_k , so we have

$$\Sigma_t^{(i)} \subset \Omega_{t_k}^{(i)} \quad \forall t > t_k$$

for each $i \in \{1, \dots, m_1\}$. Since $\Omega_{t_k}^{(i)} \cap \Omega_{t_k}^{(j)} = \emptyset$ whenever $i \neq j$ and that Σ_t is the disjoint union of $\Sigma_t^{(1)}, \dots, \Sigma_t^{(m_1)}$, we infer that

$$\Sigma_t^{(i)} = \Sigma_t \cap \Omega_{t_k}^{(i)}$$

so long as $t > t_k$. Now given $t > T_1$, we can choose $k_0 \in \mathbb{N}$ such that $t > t_{k_0}$, then we have

$$\Sigma_t^{(i)} = \bigcup_{k=k_0}^{\infty} \Sigma_t \cap \Omega_{t_k}^{(i)} = \Sigma_t \cap \Omega_{T_1}^{(i)}$$

for every $i \in \{1, \dots, m_1\}$. □

With Proposition 3.11 and Proposition 3.13, we are able to prove Theorem 1.2.

Proof. (of Theorem 1.2) If $T_1 < T_{ext}$, choose $t_1 \in (T_1, T_2)$. Note that

$$\Sigma_t = \{u = t\} \subset \{u > T_1\} = \Omega_{T_1} \quad \forall t \geq t_1.$$

By Proposition 3.13, Ω_{T_1} is a finite disjoint union of open connected sets

$$\Omega_{T_1}^{(1)}, \dots, \Omega_{T_1}^{(m_1)};$$

moreover, for each $i \in \{1, \dots, m_1\}$,

$$\Sigma_t \cap \Omega_{T_1}^{(i)}, \quad t \geq t_1$$

is the (two-convex) LSF³⁵ starting at time t_1 from $\Sigma_{t_1} \cap \Omega_{T_1}^{(i)}$, which is a two-convex closed connected hypersurface. Note that for each $i \in \{1, \dots, m_1\}$, all the singular times of the flow $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t \geq t_1}$ must be contained in $\{T_2 \leq \dots \leq T_{ext}\}$, and that at time T_2 , there must be some $i \in \{1, \dots, m_1\}$ so that $\Sigma_{T_2} \cap \Omega_{T_1}^{(i)}$ has singular points.³⁶ Then it follows from Proposition 3.11 that for each $i \in \{1, \dots, m_1\}$, the singular set of the LSF $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t \geq t_1}$ at the time T_2 , which is the set of singular points in $\Sigma_{T_2} \cap \Omega_{T_1}^{(i)}$, is either empty or a finite disjoint union of points and/or compact C^1 embedded curves (with or without boundary). Because the singular points of Σ_{T_2} is indeed the union of singular points of $\Sigma_{T_2} \cap \Omega_{T_1}^{(i)}$ for $i \in \{1, \dots, m_1\}$, we infer that the singular set of the flow $\{\Sigma_t\}$ at the second singular time T_2 is a finite disjoint union of points and/or compact C^1 embedded curves.

If $T_2 < T_{ext}$, choose $t_2 \in (T_2, T_3)$. Note that $\Sigma_t \subset \Omega_{T_2} \subset \Omega_{T_1}$ for $t \geq t_2$. For each $i \in \{1, \dots, m_1\}$, by Proposition 3.13, $\Omega_{T_2} \cap \Omega_{T_1}^{(i)}$ would be a finite disjoint union of open connected sets.³⁷ Since Ω_{T_1} is the disjoint union of $\Omega_{T_1}^{(1)}, \dots, \Omega_{T_1}^{(m_1)}$, the set Ω_{T_2} is a finite disjoint union of open connected sets

$$\Omega_{T_2}^{(1)}, \dots, \Omega_{T_2}^{(m_2)}.$$

Note that for each $j \in \{1, \dots, m_2\}$, there exists $i \in \{1, \dots, m_1\}$ such that $\Omega_{T_2}^{(j)}$ is a connected component of $\Omega_{T_2} \cap \Omega_{T_1}^{(i)}$; it follows from Proposition 3.13 that

$$\left(\Sigma_t \cap \Omega_{T_1}^{(i)} \right) \cap \Omega_{T_2}^{(j)} = \Sigma_t \cap \Omega_{T_2}^{(j)}, \quad t \geq t_2$$

is the (two-convex) LSF starting at time t_2 from $\Sigma_{t_2} \cap \Omega_{T_2}^{(j)}$, which is a two-convex closed connected hypersurface. Since

$$\Sigma_t = \bigsqcup_{j=1}^{m_2} \Sigma_t \cap \Omega_{T_2}^{(j)} \quad \forall t \geq t_2,$$

the singular times of the flow $\left\{ \Sigma_t \cap \Omega_{T_2}^{(j)} \right\}_{t \geq t_2}$ must be contained in $\{T_3 \leq \dots \leq T_{ext}\}$

for every j ; at time T_3 , there must be some $j \in \{1, \dots, m_2\}$ so that $\Sigma_{T_3} \cap \Omega_{T_2}^{(j)}$ has singular points. So it follows from Proposition 3.11 that for each $j \in \{1, \dots, m_2\}$, the singular set of the LSF $\left\{ \Sigma_t \cap \Omega_{T_2}^{(j)} \right\}_{t \geq t_2}$ at the time T_3 , which is the set of singular points in $\Sigma_{T_3} \cap \Omega_{T_2}^{(j)}$, is either empty or a finite disjoint union of points and/or compact C^1 embedded curves. Thus, the singular set of the flow $\{\Sigma_t\}$ at the third singular time T_3 is a finite disjoint union of points and/or compact C^1 embedded curves.

The conclusion would follow after repeating this process for finitely many times. \square

³⁵The restriction of u on $\bar{\Omega}_{t_1}^{(i)}$ serves as the arrival time function of the two-convex LSF.

³⁶For such i 's, T_2 is actually the first singular time of the flow $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t \geq t_1}$.

³⁷If the LSF $\left\{ \Sigma_t \cap \Omega_{T_1}^{(i)} \right\}_{t \geq t_1}$ does not become singular at time T_2 , then $\Omega_{T_2} \cap \Omega_{T_1}^{(i)}$ is itself an open connected set, which is bounded by the closed connected hypersurface $\Sigma_{T_2} \cap \Omega_{T_1}^{(i)}$.

4. TYPES OF SINGULAR COMPONENTS

In this section the two-convex LSF $\{\Sigma_t\}$ is assumed to have finitely many singular times in order that Theorem 1.2 holds. Let $\{\Sigma_t^k\}$, $k \in \mathbb{N}$, be the LSFs in Theorem 1.1. In view of Proposition 4.22 (in Section 4.4), we may assume for simplicity that $\{\Sigma_t^k\}$ is a two-convex LSF for every $k \in \mathbb{N}$.

Recall that when k is large, by Theorem 1.1 (see also Corollary 2.19) $\{\Sigma_t^k\}$ is regular and close in the smooth topology to $\{\Sigma_t\}$ away from \mathcal{S} . The target of this section is to analyze the singular set of $\{\Sigma_t^k\}$ near \mathcal{S} with results in Theorem 1.3. To achieve that, we will firstly classify the singular components (i.e., connected components of the singular set) in Definition 4.1. Then we will make adequate assumptions, including Assumption 4.3, Assumption 4.12 (in Section 4.2), and Assumption 4.16 (in Section 4.3), so as to ensure that near each singular component of $\{\Sigma_t\}$, the flow $\{\Sigma_t^k\}$ would have exactly the same type of singular set as that singular component. The proof of Theorem 1.3 is composed of three parts: Proposition 4.6 in Section 4.1, Proposition 4.14 in Section 4.2, and Proposition 4.19 in Section 4.3.

To start with, let us write the singular set \mathcal{S} of $\{\Sigma_t\}$ as a finite disjoint union of singular components, namely,

$$\mathcal{S} = \bigsqcup_j \mathcal{S}_j,$$

where each singular component \mathcal{S}_j is either a point or a compact C^1 embedded curve (with or without boundary). Note that u is constant on each \mathcal{S}_j ,³⁸ that is to say, every singularity of \mathcal{S}_j occurs at the same time. In addition, when \mathcal{S}_j is a curve, by Remark 3.5 every interior point of \mathcal{S}_j is a local maximum point of u , and every boundary point (i.e., endpoint) of \mathcal{S}_j is either a local maximum point or a one-sided saddle point of u . We then classify singular components, according to their endpoints, into the following three types:

Definition 4.1. A singular component of $\{\Sigma_t\}$ belongs to

- (1) the **vanishing type** - if it is a single local maximum point of u , or a compact C^1 embedded curve whose endpoints are both local maximum points of u , or a closed C^1 embedded curve (i.e., no endpoints);
- (2) the **splitting type** - if it is either a single two-sided saddle point of u , or a compact C^1 embedded curve whose endpoints are both one-sided saddle points of u ;
- (3) the **bumpy type** - if it is either a single one-sided saddle point of u , or a compact C^1 embedded curve with one endpoint being a one-sided saddle point while the other being a local maximum point of u .

The reason why the two-sided saddle points and the one-sided saddle points are classified as the splitting type and the bumpy type, respectively, is as follows. Recall that by Definition 3.3, a saddle point p is two-sided or one-sided depends on how many “sides” the point can be approached by the superlevel set $\Omega_{u(p)} = \{u > u(p)\}$. From this perspective, and considering the fact that a splitting curve can be (and can only be) approached by the superlevel set near the two “ends,” a two-sided saddle point can be regarded as a “degenerate” curve of the splitting type. Likewise,

³⁸When $\mathcal{S}_j = \{x(t) : t \in [a, b]\}$ is a curve, we have $\frac{d}{dt} u(x(t)) = \nabla u(x(t)) \cdot x'(t) = 0$.

a one-sided saddle point can be viewed as a degenerate curve of the bumpy type because the superlevel set can be approached from only one side/end.

Now let $\hat{\delta} > 0$ be a sufficiently small constant such that the following hold:

- For every $j \neq j'$,

$$(4.1) \quad \mathcal{S}_j^{\hat{\delta}} \cap \mathcal{S}_{j'}^{\hat{\delta}} = \emptyset,$$

where

$$\mathcal{S}_j^{\hat{\delta}} = \left\{ x \in \mathbb{R}^n : \text{dist}(x, \mathcal{S}_j) < \hat{\delta} \right\}$$

is the $\hat{\delta}$ -neighborhood of \mathcal{S}_j .

- For every $l \geq 1$,

$$(4.2) \quad T_{l-1} + \hat{\delta} < T_l - \hat{\delta},$$

where $0 = T_0 < T_1 \leq \dots \leq T_m = T_{ext}$ are the set of singular times of $\{\Sigma_t\}$.

- If $u(\mathcal{S}_j) = T_l$ (i.e., \mathcal{S}_j occurs at time T_l), then

$$(4.3) \quad \overline{\mathcal{S}_j^{\hat{\delta}}} \subset \Omega_{T_{l-1}}^{(i)},$$

where $\Omega_{T_{l-1}}^{(i)}$ is some connected component of $\Omega_{T_{l-1}}$ (see Proposition 3.13).

The following proposition is a collection of Corollary 2.17, Corollary 2.19 (see also (4.1)), and Corollary 2.22 (see also Proposition 3.13 and (4.2)).

Proposition 4.2. *Given $\epsilon > 0$ and $0 < \delta < \hat{\delta}$, there exists $k_{\epsilon, \delta} \in \mathbb{N}$ so that for every $k \geq k_{\epsilon, \delta}$ the following hold:*

- (1) *The extinction time T_{ext}^k of $\{\Sigma_t^k\}$ satisfies*

$$T_{ext}^k \in (T_{ext} - \delta, T_{ext} + \delta).$$

- (2) *The singular set \mathcal{S}^k of $\{\Sigma_t^k\}$ satisfies*

$$\mathcal{S}^k \subset \bigsqcup_j \mathcal{S}_j^{\hat{\delta}}.$$

- (3) *For every $l \geq 1$, $\{\Sigma_t^k\}_{T_{l-1} + \delta \leq t \leq T_l - \delta}$ is a two-convex MCF that is ϵ -close in the C^m topology to $\{\Sigma_t\}_{T_{l-1} + \delta \leq t \leq T_l - \delta}$, where $m = m(n, \lambda)$ is the constant in Section 3.2.*

We will make good use of the above proposition in Section 4.1, Section 4.2, and Section 4.3 with ϵ and δ chosen sufficiently small subject to the asymptotic behavior of the flow $\{\Sigma_t\}$ near each of its singular component. Throughout this section we assume that

Assumption 4.3. *When k is sufficiently large, the flow $\{\Sigma_t^k\}$ has at most one singular time in $\mathcal{S}_j^{\hat{\delta}}$ for every j ; in particular, Theorem 1.2 and Definition 4.1 are applicable to $\{\Sigma_t^k\}$ as well.*

Two supplementary assumptions will be made for the splitting case (see Assumption 4.12) in Section 4.2 and the bumpy case (see Assumption 4.16) in Section 4.3.

Remark 4.4. On account of (4.3), each singular component \mathcal{S}_j of $\{\Sigma_t\}$ is indeed the singular set of the flow

$$(4.4) \quad \left\{ \Sigma_t \cap \Omega_{T_{l-1}}^{(i)} \right\}_{t > T_{l-1}}$$

at its “first” singular time (see also the exposition at the beginning of Section 3.4). Thus, throughout Section 4, upon replacing the flow by some of its connected component (e.g., (4.4)) if necessary, we may assume without loss of generality that every singular component under discussion occurs at the first singular time.

4.1. Vanishing type. In this subsection we shall give a criteria (see Proposition 4.5) to distinguish between the vanishing type and the splitting/bumpy type. Then we will prove the “stability” of singular components of the vanishing type in Proposition 4.6.

Let us begin by recalling that a singular component \mathcal{S}_j of $\{\Sigma_t\}$ is of the vanishing type if and only if it comprises local maximum points of u (see Definition 4.1). In the following proposition, we show that the flow $\{\Sigma_t\}$ would shrink to \mathcal{S}_j at time $T_1 = u(\mathcal{S}_j)$ (see Remark 4.4) and then vanish completely. A typical example is when \mathcal{S}_j is a single round point (see Corollary 3.1).

Proposition 4.5. *A singular component \mathcal{S}_j of $\{\Sigma_t\}$ is of the vanishing type if and only if $T_1 = T_{ext}$ (in the setting of Remark 4.4); in that case, $\mathcal{S}_j = \mathcal{S}$ (that is, \mathcal{S} has only one component and hence is connected).*

In other words, a singular component \mathcal{S}_j of $\{\Sigma_t\}$ is of the splitting/bumpy type if and only if $T_1 < T_{ext}$.

Proof. Let \mathcal{S}_j be a singular component of the vanishing type. By Definition 4.1, \mathcal{S}_j consists of local maximum points of u and is compact, so there exists $\delta > 0$ sufficiently small such that

- (1) the δ -neighborhood \mathcal{S}_j^δ of \mathcal{S}_j is strictly contained in Ω_0 ;
- (2) in \mathcal{S}_j^δ there are no other singular points of the flow $\{\Sigma_t\}$ than \mathcal{S}_j ;
- (3) $u \leq T_1$ in \mathcal{S}_j^δ .

Then the value of u at every point in $\bar{\mathcal{S}}_j^{\delta/2} \setminus \mathcal{S}_j$ is strictly less than T_1 ; otherwise, that point would be a local maximum point of u and hence a singular point of $\{\Sigma_t\}$, contradicting the second condition in the above. In particular, we have

$$\tau := \max_{\partial \mathcal{S}_j^{\delta/2}} u < T_1.$$

Note that for every $t \in (\tau, T_1)$, $\Sigma_t \cap \mathcal{S}_j^{\delta/2}$ is a nonempty (by the intermediate value theorem) closed hypersurface since $t \in (0, T_1)$ is a regular value of u and

$$\Sigma_t \cap \partial \mathcal{S}_j^{\delta/2} = \emptyset.$$

As Σ_t is connected, we infer that $\Sigma_t \subset \mathcal{S}_j^{\delta/2}$. It follows that $\Omega_t \subset \mathcal{S}_j^{\delta/2}$ and hence

$$T_{ext} = \max_{\Omega_t} u \leq \max_{\mathcal{S}_j^{\delta/2}} u = T_1.$$

Thus, $T_1 = T_{ext}$. Additionally, since

$$\mathcal{S} = \mathcal{S} \cap \Omega_t \subset \mathcal{S} \cap \mathcal{S}_j^{\delta/2} = \mathcal{S}_j,$$

we conclude that $\mathcal{S}_j = \mathcal{S}$.

Conversely, if $T_1 = T_{ext}$, then every component of \mathcal{S} is obviously of the vanishing type since \mathcal{S} consists of global maximum points of u . Let \mathcal{S}_j be one of the components of \mathcal{S} . Then by the above argument we obtain $\mathcal{S}_j = \mathcal{S}$. \square

The following result is based on Proposition 4.2 and Proposition 4.5.

Proposition 4.6. *If \mathcal{S}_j is a singular component of $\{\Sigma_t\}$ that belongs to the vanishing type, then for every sufficiently large k , in $\mathcal{S}_j^{\hat{\delta}}$ there is precisely one singular component of $\{\Sigma_t^k\}$, which is of the vanishing type.*

Proof. By the argument in the proof of Proposition 4.5, there is $\delta > 0$ sufficiently small that

$$\bar{\Omega}_{T_1-\delta} \subset \mathcal{S}_j^{\hat{\delta}} \subset \overline{\mathcal{S}_j^{\hat{\delta}}} \subset \Omega_{\delta}.$$

By Proposition 4.2, when k is large, $\{\Sigma_t^k\}_{\delta \leq t \leq T_1-\delta}$ is a two-convex MCF of closed connected hypersurfaces so that

$$\bar{\Omega}_{T_1-\delta}^k \subset \mathcal{S}_j^{\hat{\delta}} \subset \overline{\mathcal{S}_j^{\hat{\delta}}} \subset \Omega_{\delta}^k.$$

In view of Assumption 4.3,

$$T_{ext}^k = \max_{\Omega_{T_1-\delta}} u^k$$

is the unique singular time of the flow $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$. Then applying Proposition 4.5 to the LSF $\{\Sigma_t^k\}_{t \geq \delta}$ gives that the singular set of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ is connected and of the vanishing type. \square

4.2. Splitting type. In this subsection we will be devoted to show the “stability” of singular components of the splitting type (see Proposition 4.13 and Proposition 4.14) under Assumption 4.12. In view of Proposition 4.5, it can be assumed throughout this (and also the next) subsection that $T_1 < T_{ext}$. Additionally, note that a singular component of the splitting/bumpy type consists of cylindrical points.

Prior to beginning the subsection, let us give a brief overview as follows. Let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ belonging to the splitting type. When k is large, we firstly prove in Proposition 4.7 that every cylindrical point of $\{\Sigma_t^k\}$ near \mathcal{S}_j has a uniform (i.e., independent of position and k) cylindrical scale and then in Remark 4.8 and Remark 4.9 show that $\Sigma_{\tau_j}^k$ locally looks like a “tube” around \mathcal{S}_j for some $\tau_j < T_1 = u(\mathcal{S}_j)$. In Proposition 4.10 we demonstrate that Ω_t “splits” near \mathcal{S}_j for $t > T_1$, which, under Assumption 4.12, yields that $\{\Sigma_t^k\}$ must have singularities near \mathcal{S}_j (see Lemma 4.11). Lastly, in Proposition 4.13 and Proposition 4.14, we conclude that $\{\Sigma_t^k\}$ would have precisely one singular component near \mathcal{S}_j , which is of the splitting type.

Now let us start with the following proposition. Note that the “cylindrical scale” therein refers to the $(\phi(n), \epsilon(n))$ -cylindrical scale in Section 3.2 (see the exposition between Remark 3.4 and Remark 3.5).

Proposition 4.7. *Let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ consisting of cylindrical points. Then there exists $0 < r_j < \hat{\delta}$ with the following properties:*

- *Every cylindrical point of $\{\Sigma_t\}$ on \mathcal{S}_j has a uniform cylindrical scale r_j .*
- *When k is large, every cylindrical point q of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$, if any, has a cylindrical scale r_j ; moreover, every other cylindrical points of $\{\Sigma_t^k\}$ in*

$B_{r_j}(q)$ would be located in a small Lipschitz graph over the axis of the tangent cylinder of $\{\Sigma_t^k\}$ at q .

- In the case where \mathcal{S}_j is a curve with endpoints, we have

$$(4.5) \quad \bar{B}_{3r_j}^+(p) \cap \mathcal{S}_j = \{p\}$$

for each endpoint p of \mathcal{S}_j , where $B_{3r_j}^+(p)$ is one of the half balls cut from $B_{3r_j}(p)$ by the hyperplane \mathcal{P} passing through p and orthogonal to the tangent cylinder of $\{\Sigma_t\}$ at p , and $\bar{B}_{3r_j}^+(p)$ is the closure of $B_{3r_j}^+(p)$.

Proof. Firstly, let $\psi, \delta > 0$ be small constants (to be determined), we claim that there is $0 < r_j < \delta$ so that every cylindrical point of $\{\Sigma_t\}$ on \mathcal{S}_j has a uniform (ψ, δ) -cylindrical scale $3r_j$. This is obviously true if \mathcal{S}_j is just a single point. When \mathcal{S}_j is a curve, say

$$\mathcal{S}_j = \{\Gamma(s) : s \in [0, 1]\},$$

then by Section 3.2, for every $s \in [0, 1]$ there is an open neighborhood \mathcal{O}_s of $\Gamma(s)$ so that every cylindrical point of $\{\Sigma_t\}$ in \mathcal{O}_s has a uniform (ψ, δ) -cylindrical scale \mathcal{R}_s . By the compactness of \mathcal{S}_j , there exist

$$0 \leq s_1 < \cdots < s_m \leq 1$$

so that

$$\mathcal{S}_j \subset \mathcal{O}_{s_1} \cup \cdots \cup \mathcal{O}_{s_m}.$$

Then every cylindrical point of $\{\Sigma_t\}$ on \mathcal{S}_j has a uniform (ψ, δ) -cylindrical scale $3r_j$, where

$$r_j := \frac{1}{3} \min \left\{ \mathcal{R}_{s_1}, \dots, \mathcal{R}_{s_m}, \delta \right\},$$

proving the claim. Note also that in the case where \mathcal{S}_j is a curve with endpoints, (4.5) holds by choosing r_j even smaller.

By Proposition 2.16 and Proposition 4.2, given a small constant $\sigma \in (0, 1)$ (to be determined), there exists $k_0 \in \mathbb{N}$ so that for every $k \geq k_0$ the following hold:

- the singular set of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ is contained in $\mathcal{S}_j^{\sigma r_j}$;
- $|T_{1,j}^k - T_1|^{\frac{1}{2}} \leq \sigma r_j$, where $T_{1,j}^k$ is the unique singular time of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$;
- $\frac{1}{\sqrt{T_1-t}}(\Sigma_t^k - p)$ is σ -close in C^m -topology to $\frac{1}{\sqrt{T_1-t}}(\Sigma_t - p)$ in $B_{1/\sigma}$ for every $p \in \mathcal{S}_j$ and $t \in \left[T_1 - \frac{r_j^2}{n-2}, T_1 - \frac{r_j^2}{8(n-2)} \right]$.

Then by Section 3.2 there are small positive constants ψ, δ, σ (depending on n and λ)³⁹ so that

- every cylindrical point of $\{\Sigma_t\}$ on \mathcal{S}_j has a uniform $(\phi(n), \epsilon(n))$ -cylindrical scale r_j ;
- when $k \geq k_0$, every cylindrical point of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$, if any, has a uniform $(\phi(n), \epsilon(n))$ -cylindrical scale r_j ; moreover, for each cylindrical point q of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$, every other cylindrical point of $\{\Sigma_t^k\}$ in $B_{r_j}(q)$ would be located in a small Lipschitz graph over the axis of the tangent cylinder of $\{\Sigma_t^k\}$ at q .

□

³⁹See Proposition 2.5 for the uniform estimate of the entropy of $\{\Sigma_t^k\}$ when k is large.

Based on the asymptotically cylindrical behavior of the flow within the cylindrical scale (see Section 3.2), we then have the following remarks.

Remark 4.8. Let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ consisting of cylindrical points. By Proposition 4.7, every cylindrical point of $\{\Sigma_t\}$ on \mathcal{S}_j has a cylindrical r_j . It follows that for every $p \in \mathcal{S}_j$, Σ_{τ_j} is asymptotically cylindrical in $B_{r_j}(p)$, where

$$\tau_j := T_1 - \frac{r_j^2}{4(n-2)};$$

consequently, $\Sigma_{\tau_j} \cap \mathcal{S}_j^{r_j}$ is a **tube-like hypersurface** around \mathcal{S}_j . By Proposition 4.2, when k is large, $\Sigma_{\tau_j}^k$ would be close to Σ_{τ_j} so that in $\mathcal{S}_j^{r_j}$ it is also a tube-like hypersurface around \mathcal{S}_j (and also every singular component of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{r_j}$, if any).

Remark 4.9. Suppose in Remark 4.8 we assume that

$$\mathcal{S}_j = \{\Gamma(s) : s \in [-1, 1]\}$$

is a curve of the splitting type, namely, its endpoints are one-sided saddle points. Since $\mathcal{S}_j^{\hat{\delta}} \cap \mathcal{S}_{j'}^{\hat{\delta}} = \emptyset$ for $j' \neq j$ and that $r_j < \hat{\delta}$ (see Proposition 4.7), the first singular time T_1 is indeed the unique singular time of $\{\Sigma_t\}$ in $\mathcal{S}_j^{r_j}$. Following Remark 3.8, near the endpoints⁴⁰ we can choose two hyperplanes \mathcal{P}_+ and \mathcal{P}_- ⁴¹ that are orthogonal to the tangent cylinders of $\{\Sigma_t\}$ at $\Gamma(1)$ and $\Gamma(-1)$, respectively, such that

$$\mathcal{P}_+ \cap \mathcal{S}_j^{r_j} \cap \mathcal{S}_j = \emptyset, \quad \mathcal{P}_- \cap \mathcal{S}_j^{r_j} \cap \mathcal{S}_j = \emptyset$$

and that the tube-like connected hypersurface Σ_{τ_j} together with \mathcal{P}_+ and \mathcal{P}_- bound a closed **tubular region** \mathcal{T}_j in $\mathcal{S}_j^{r_j}$ with the following properties:

- (1) $\mathcal{S}_j \subset \text{int } \mathcal{T}_j$.
- (2) Let $\tilde{\mathcal{P}}_+$ and $\tilde{\mathcal{P}}_-$ ⁴² parallel to \mathcal{P}_+ and \mathcal{P}_- , respectively, so that

$$\tilde{\mathcal{P}}_+ \cap \mathcal{T}_j \cap \mathcal{S}_j = \{\Gamma(1)\}, \quad \tilde{\mathcal{P}}_- \cap \mathcal{T}_j \cap \mathcal{S}_j = \{\Gamma(-1)\}.$$

Note that the tube \mathcal{T}_j is cut by $\tilde{\mathcal{P}}_+$ and $\tilde{\mathcal{P}}_-$ into three closed pieces,⁴³

$$\mathcal{T}_j = \hat{\mathcal{T}}_j^+ \cup \mathcal{T}_j^* \cup \hat{\mathcal{T}}_j^-,$$

where \mathcal{T}_j^* is the truncated tubular closed region bounded by Σ_{τ_j} (from the lateral) and the two caps $\tilde{\mathcal{P}}_+ \cap \mathcal{T}_j$ and $\tilde{\mathcal{P}}_- \cap \mathcal{T}_j$ (from the two ends); $\hat{\mathcal{T}}_j^+$ is the tubular region bounded by Σ_{τ_j} and the caps $\mathcal{P}_+ \cap \mathcal{T}_j$ and $\tilde{\mathcal{P}}_+ \cap \mathcal{T}_j$; and $\hat{\mathcal{T}}_j^-$ is the tubular region bounded by Σ_{τ_j} and the caps $\mathcal{P}_- \cap \mathcal{T}_j$ and $\tilde{\mathcal{P}}_- \cap \mathcal{T}_j$. Note also that $\mathcal{S}_j \subset \mathcal{T}_j^*$ and that

$$\Gamma(1) \in \hat{\mathcal{T}}_j^+ \subset \bar{B}_{r_j}^+(\Gamma(1)),$$

$$\Gamma(-1) \in \hat{\mathcal{T}}_j^- \subset \bar{B}_{r_j}^-(\Gamma(-1)),$$

⁴⁰Here $\Gamma(1)$ and $\Gamma(-1)$ correspond to the origin in Remark 3.8. Note that the orientation is chosen in such a way that the curve is on the downside $z \leq 0$.

⁴¹Here \mathcal{P}_+ and \mathcal{P}_- correspond to $z = \varepsilon$ in (3.18).

⁴²Here $\tilde{\mathcal{P}}_+$ and $\tilde{\mathcal{P}}_-$ correspond to $z = 0$ in (3.18).

⁴³Here $\hat{\mathcal{T}}_j^+$ and $\hat{\mathcal{T}}_j^-$ correspond to $\hat{\mathcal{T}}$ in (3.18).

where $\bar{B}_{r_j}^+(\Gamma(1))$ and $\bar{B}_{r_j}^-(\Gamma(-1))$ are the closures of the open half-balls⁴⁴ $B_{r_j}^+(\Gamma(1))$ and $B_{r_j}^-(\Gamma(-1))$, respectively, and are defined analogously as in (4.5).

(3) There exists a time $\mathring{t}_j > T_1$ so that for every $t \in [\tau_j, \mathring{t}_j]$,⁴⁵

$$\Sigma_t \cap \mathcal{P}_+ \cap \mathcal{T}_j \neq \emptyset, \quad \Sigma_t \cap \mathcal{P}_- \cap \mathcal{T}_j \neq \emptyset.$$

(4) For every $t \in (T_1, \mathring{t}_j]$,

$$\Sigma_t \cap \hat{\mathcal{T}}_j^+ \subset \text{int } \mathcal{C}_\phi^+, \quad \Sigma_t \cap \hat{\mathcal{T}}_j^- \subset \text{int } \mathcal{C}_\phi^-,$$

where \mathcal{C}_ϕ^+ and \mathcal{C}_ϕ^- are the cones⁴⁶ defined analogously as in (3.3) for the endpoints.

(5) There hold⁴⁷

$$(4.6) \quad \min_{\mathcal{C}_\phi^+ \cap \hat{\mathcal{T}}_j^+} u > \tau_j, \quad \min_{\mathcal{C}_\phi^- \cap \hat{\mathcal{T}}_j^-} u > \tau_j.$$

By the asymptotically cylindrical behavior of each point on \mathcal{S}_j within the cylindrical scale r_j and (4.6), there exists $\tau'_j > \tau_j$ so that for $t \in [\tau_j, \tau'_j]$, $\bar{\Omega}_t \cap \mathcal{T}_j$ is also a tubular closed connected region bounded by the tube-like hypersurface $\Sigma_t \cap \mathcal{T}_j$ and the two caps $\mathcal{P}_+ \cap \mathcal{T}_j, \mathcal{P}_- \cap \mathcal{T}_j$.

Note that even in the case where \mathcal{S}_j is a single two-sided saddle points (which can be regarded as a degenerate curve with one-sided endpoints, see Definition 4.1), the aforementioned tubular region can be defined analogously.⁴⁸

Furthermore, by Proposition 2.16, Proposition 4.2, and Remark 4.8, when k is large, $\Sigma_{\tau_j}^k$ and the two hyperplanes \mathcal{P}_+ and \mathcal{P}_- would also bound a closed tubular region \mathcal{T}_j^k in \mathcal{S}_j^k ; in addition, for $t \in [\tau_j, \frac{1}{2}(\tau_j + \tau'_j)]$, $\bar{\Omega}_t^k \cap \mathcal{T}_j^k$ is a tubular closed connected region bounded by $\Sigma_t^k \cap \mathcal{T}_j^k, \mathcal{P}_+ \cap \mathcal{T}_j^k$, and $\mathcal{P}_- \cap \mathcal{T}_j^k$.

The following proposition indicates that when \mathcal{S}_j is a singular component of the splitting type, the superlevel set $\Omega_t = \{u > t\}$ becomes separated in a tubular neighborhood of \mathcal{S}_j for $t > T_1 = u(\mathcal{S}_j)$.

Proposition 4.10. *Let the notations be as defined in Remark 4.9. If*

$$\mathcal{S}_j = \{\Gamma(s) : s \in [-1, 1]\}$$

is a singular component of $\{\Sigma_t\}$ that belongs to the splitting type, then there exists $\mathring{t}_j \in (T_1, T_2)$ so that for every $t \in (T_1, \mathring{t}_j]$,

$$\begin{aligned} \bar{\Omega}_t \cap \mathcal{T}_j &\subset B_{\frac{3}{2}r_j}^+(\Gamma(1)) \cup B_{\frac{3}{2}r_j}^-(\Gamma(-1)), \\ \Sigma_t \cap \mathcal{P}_+ \cap \mathcal{T}_j &\neq \emptyset, \quad \Sigma_t \cap \mathcal{P}_- \cap \mathcal{T}_j \neq \emptyset. \end{aligned}$$

Note that

$$\begin{aligned} B_{\frac{3}{2}r_j}^+(\Gamma(1)) \cap B_{\frac{3}{2}r_j}^-(\Gamma(-1)) &= \emptyset, \\ \mathcal{P}_+ \cap \mathcal{T}_j &\subset B_{\frac{3}{2}r_j}^+(\Gamma(1)), \quad \mathcal{P}_- \cap \mathcal{T}_j \subset B_{\frac{3}{2}r_j}^-(\Gamma(-1)). \end{aligned}$$

⁴⁴Here $B_{r_j}^+(\Gamma(1))$ and $B_{r_j}^-(\Gamma(-1))$ correspond to $B_r \cap \{z > 0\}$ in Remark 3.8.

⁴⁵See (3.19).

⁴⁶Here \mathcal{C}_ϕ^+ and \mathcal{C}_ϕ^- correspond to \mathcal{C}_ϕ in (3.20).

⁴⁷See (3.17).

⁴⁸In that case, $\tilde{\mathcal{P}}_+ = \tilde{\mathcal{P}}_-$ and $\mathcal{T}_j^* = \hat{\mathcal{T}}_j^+ \cap \tilde{\mathcal{P}}_+$.

Moreover, the same results hold when $\mathcal{S}_j = \{p\}$ is a single two-sided saddle point. In that case, set

$$\Gamma(1) = \Gamma(-1) = p$$

and let $B_{\frac{3}{2}r_j}^+(\Gamma(1))$ and $B_{\frac{3}{2}r_j}^-(\Gamma(-1))$ be the upper and lower open half-balls of $B_{\frac{3}{2}r_j}(p)$, respectively (see Remark 3.8 and Remark 4.9).

Proof. The proofs for the case where \mathcal{S}_j is a single two-sided saddle point and the case where \mathcal{S}_j is a curve with one-sided saddle endpoints are in the same spirit, so without loss of generality let us assume that

$$\mathcal{S}_j = \{\Gamma(s) : s \in [-1, 1]\}$$

is a curve with one-sided saddle endpoints .

By Proposition 4.7 we have

$$(4.7) \quad \bar{B}_{\frac{3}{2}r_j}^+(\Gamma(1)) \cap \mathcal{S}_j = \{\Gamma(1)\}, \quad \bar{B}_{\frac{3}{2}r_j}^-(\Gamma(-1)) \cap \mathcal{S}_j = \{\Gamma(-1)\},$$

where the half balls are defined as in Remark 4.9. Now let us apply the results in Remark 4.9. There exists a time $\hat{t}_j \in (T_1, T_2)$ so that for every $t \in (\tau_j, \hat{t}_j]$,

$$\Sigma_t \cap \mathcal{P}_+ \cap \mathcal{T}_j \neq \emptyset, \quad \Sigma_t \cap \mathcal{P}_- \cap \mathcal{T}_j \neq \emptyset.$$

Note that the caps satisfy

$$\mathcal{P}_+ \cap \mathcal{T}_j = \mathcal{P}_+ \cap \hat{\mathcal{T}}_j^+ \subset \mathcal{P}_+ \cap \bar{B}_{r_j}^+(\Gamma(1)) \subset B_{\frac{3}{2}r_j}^+(\Gamma(1)),$$

$$\mathcal{P}_- \cap \mathcal{T}_j = \mathcal{P}_- \cap \hat{\mathcal{T}}_j^- \subset \mathcal{P}_- \cap \bar{B}_{r_j}^-(\Gamma(-1)) \subset B_{\frac{3}{2}r_j}^-(\Gamma(-1)).$$

Moreover, for every $t \in (T_1, \hat{t}_j]$,

$$(4.8) \quad \bar{\Omega}_t \cap \hat{\mathcal{T}}_j^+ \subset \text{int } \mathcal{C}_\phi^+ \cap \bar{B}_{r_j}^+(\Gamma(1)) \subset \text{int } \mathcal{C}_\phi^+ \cap B_{\frac{3}{2}r_j}^+(\Gamma(1)),$$

$$\bar{\Omega}_t \cap \hat{\mathcal{T}}_j^- \subset \text{int } \mathcal{C}_\phi^- \cap \bar{B}_{r_j}^-(\Gamma(-1)) \subset \text{int } \mathcal{C}_\phi^- \cap B_{\frac{3}{2}r_j}^-(\Gamma(-1)).$$

In view of the asymptotically cylindrical behavior of $\{\Sigma_t\}$ about each point on \mathcal{S}_j within the cylindrical scale r_j , we deduce that for every $t \in (T_1, \hat{t}_j]$,

$$\bar{\Omega}_t \cap \mathcal{T}_j^* = \emptyset$$

(see Remark 3.5) and so

$$\bar{\Omega}_t \cap \mathcal{T}_j = \left(\bar{\Omega}_t \cap \hat{\mathcal{T}}_j^+ \right) \cup \left(\bar{\Omega}_t \cap \hat{\mathcal{T}}_j^- \right);$$

it follows from (4.8) that

$$\bar{\Omega}_t \cap \mathcal{T}_j = \left(\bar{\Omega}_t \cap \hat{\mathcal{T}}_j^+ \right) \cup \left(\bar{\Omega}_t \cap \hat{\mathcal{T}}_j^- \right) \subset B_{\frac{3}{2}r_j}^+(\Gamma(1)) \cup B_{\frac{3}{2}r_j}^-(\Gamma(-1)).$$

Note that by (4.7) we have

$$B_{\frac{3}{2}r_j}^+(\Gamma(1)) \cap B_{\frac{3}{2}r_j}^-(\Gamma(-1)) = \emptyset.$$

□

Recall that in the end of Remark 4.9, it is mentioned that when k is large, $\bar{\Omega}_{\tau_j}^k \cap \mathcal{T}_j^k$ is a connected solid tube. The following lemma says that if $\bar{\Omega}_{t_*}^k \cap \mathcal{T}_j^k$ becomes disconnected for some $t_* > \tau_j$, then under some conditions such as (4.9), the flow $\{\Sigma_t^k\}$ would become singular in \mathcal{T}_j^k at some time $\hat{t} \in (\tau_j, t_*)$.

Lemma 4.11. *Let k be a large integer so that for t close to τ_j , $\Sigma_t^k \cap \mathcal{S}_j^{r_j}$ is a tube-like hypersurface around \mathcal{S}_j and $\bar{\Omega}_t^k \cap \mathcal{T}_j^k$ is a tubular closed connected region (see Remark 4.9). If there exists $t_* > \tau_j$ so that the following hold:*

- (1) $\Sigma_{t_*}^k \cap \mathcal{T}_j^k$ has no singular points of the flow $\{\Sigma_t^k\}$.
- (2) $\bar{\Omega}_{t_*}^k \cap \mathcal{T}_j^k$ is disconnected in such a way that every connected component of $\bar{\Omega}_{t_*}^k \cap \mathcal{T}_j^k$ intersects at most one of the caps (i.e., $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\mathcal{P}_- \cap \mathcal{T}_j^k$); specifically, there exists one component that intersects $\mathcal{P}_+ \cap \mathcal{T}_j^k$, and there exists another component that does not intersect $\mathcal{P}_+ \cap \mathcal{T}_j^k$.
- (3) The unit normal vector $\frac{\nabla u^k}{|\nabla u^k|}$ satisfies

$$(4.9) \quad \begin{aligned} \frac{\nabla u^k}{|\nabla u^k|} \cdot \nu_+ &\neq -1 \text{ on } \mathcal{P}_+ \cap \mathcal{T}_j^k \cap \{\tau_j < u^k < t_*\}, \\ \frac{\nabla u^k}{|\nabla u^k|} \cdot \nu_- &\neq -1 \text{ on } \mathcal{P}_- \cap \mathcal{T}_j^k \cap \{\tau_j < u^k < t_*\}, \end{aligned}$$

where ν_+ and ν_- are the outward (i.e., pointing toward the outside of \mathcal{T}_j^k) unit normal vector of the caps $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\mathcal{P}_- \cap \mathcal{T}_j^k$, respectively.

Then there is $\hat{t} \in (\tau_j, t_*)$ such that $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$ has singular points of $\{\Sigma_t^k\}$.

Proof. First of all, note that the set of regular values of $u^k|_{\mathcal{T}_j^k}$, namely,

$$(4.10) \quad \{t : \nabla u^k(x) \neq 0 \text{ whenever } x \in \mathcal{T}_j^k \text{ with } u(x) = t\}$$

is open because the set of critical values of $u^k|_{\mathcal{T}_j^k}$ is compact (see Lemma 2.10 or Assumption 4.3). Note also that the minimum of u^k on any compact set can only be attained on the boundary since u^k has no interior local minimum points (see Corollary 3.1 and Corollary 3.2).

Let $\bar{\Omega}_t^{k,+}$ be the union connected components of $\bar{\Omega}_t^k \cap \mathcal{T}_j^k$ that intersect $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and let $\bar{\Omega}_t^{k,-}$ be the union the remaining connected components of $\bar{\Omega}_t^k \cap \mathcal{T}_j^k$. Define \mathfrak{T} as the set of time $t \in [\tau_j, t_*)$ such that

- $\Sigma_t^k \cap \mathcal{T}_j^k$ has no singular points of $\{\Sigma_t^k\}$;
- $\bar{\Omega}_t^{k,+} \cap \mathcal{P}_- \cap \mathcal{T}_j^k = \emptyset$ and $\bar{\Omega}_t^{k,-} \cap \mathcal{P}_+ \cap \mathcal{T}_j^k = \emptyset$.

Let

$$(4.11) \quad \hat{t} = \inf \{\tilde{t} \in (\tau_j, t_*) : [\tilde{t}, t_*] \subset \mathfrak{T}\}.$$

By the assumptions and (4.10), it is not hard to see that $\hat{t} \in (\tau_j, t_*)$. We claim that $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$ must have singular points of $\{\Sigma_t^k\}$.

Suppose the contrary that $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$ has no singular points of $\{\Sigma_t^k\}$. Then every time $t \in [\hat{t}, t_*)$ would be a regular value of $u^k|_{\mathcal{T}_j^k}$; in fact, by (4.10) we can even find $\hat{t}' < \hat{t}$ so that

$$(4.12) \quad r := \min_{\{\hat{t}' \leq u^k \leq t_*\} \cap \mathcal{T}_j^k} |\nabla u^k| > 0.$$

Note that $\bar{\Omega}_{\hat{t}}^k \cap \text{int } \mathcal{T}_j^k$ is contained in $\bar{\Omega}^+ \cup \bar{\Omega}^-$, where

$$\bar{\Omega}^+ := \bigcup_{t \in (\hat{t}, t_*)} \bar{\Omega}_t^{k,+}, \quad \bar{\Omega}^- := \bigcup_{t \in (\hat{t}, t_*)} \bar{\Omega}_t^{k,-}.$$

Moreover, it follows from (4.12), Definition 2.11, Proposition 2.12, Proposition 2.13, and Corollary 2.15 that there exists a sequence $t_i \searrow \hat{t}$ such that

$$(4.13) \quad \overline{\partial\bar{\Omega}_{t_i}^{k,+} \cap \text{int } \mathcal{T}_j^k} \xrightarrow{C^\infty} \Sigma^+, \quad \overline{\partial\bar{\Omega}_{t_i}^{k,-} \cap \text{int } \mathcal{T}_j^k} \xrightarrow{C^\infty} \Sigma^- \quad \text{as } i \rightarrow \infty,$$

where Σ^+ and Σ^- are hypersurfaces contained in $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$ with boundary on $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\mathcal{P}_- \cap \mathcal{T}_j^k$. We have

$$\overline{\partial\Omega^+ \cap \text{int } \mathcal{T}_j^k} \subset \Sigma^+, \quad \overline{\partial\Omega^- \cap \text{int } \mathcal{T}_j^k} \subset \Sigma^-.$$

Furthermore, it is true that

$$(4.14) \quad \bar{\Omega}^+ \cap \mathcal{P}_- \cap \mathcal{T}_j^k = \emptyset, \quad \bar{\Omega}^- \cap \mathcal{P}_+ \cap \mathcal{T}_j^k = \emptyset.$$

Because if one of them were false, say $\bar{\Omega}^+ \cap \mathcal{P}_- \cap \mathcal{T}_j^k \neq \emptyset$, then we would have $q \in \bar{\Omega}^+ \cap \mathcal{P}_- \cap \mathcal{T}_j^k$. By (4.13) and the condition that

$$\bar{\Omega}_{t_i}^{k,+} \cap \mathcal{P}_- \cap \mathcal{T}_j^k = \emptyset \quad \forall i,$$

$\bar{\Omega}^+$ would “touch” $\mathcal{P}_- \cap \mathcal{T}_j^k$ at q since Σ^+ is on one side⁴⁹ of the cap $\mathcal{P}_- \cap \mathcal{T}_j^k$ with $q \in \Sigma^+ \cap \mathcal{P}_- \cap \mathcal{T}_j^k$, which yields that

$$\frac{\nabla u^k(q)}{|\nabla u^k(q)|} \cdot \nu_- = -1,$$

which is in contradiction with (4.9).

Most importantly, we should have

$$\liminf_{i \rightarrow \infty} \text{dist} \left(\bar{\Omega}_{t_i}^{k,+}, \bar{\Omega}_{t_i}^{k,-} \right) = 0;$$

otherwise, upon passing to a subsequence, there would exist $\varrho > 0$ such that

$$\text{dist} \left(\bar{\Omega}_{t_i}^{k,+}, \bar{\Omega}_{t_i}^{k,-} \right) \geq \varrho \quad \forall i,$$

which yields that $\bar{\Omega}^+$ and $\bar{\Omega}^-$ would be disjoint, which would lead to a contradiction with the choice of \hat{t} in view of (4.12). Thus, for each i we can choose $p_i^+ \in \partial\bar{\Omega}_{t_i}^{k,+} \cap \text{int } \mathcal{T}_j^k$ and $p_i^- \in \partial\bar{\Omega}_{t_i}^{k,-} \cap \text{int } \mathcal{T}_j^k$ so that

$$\liminf_{i \rightarrow \infty} |p_i^+ - p_i^-| = 0.$$

By passing to a subsequence, we may assume that

$$(4.15) \quad \lim_{i \rightarrow \infty} p_i^+ = \lim_{i \rightarrow \infty} p_i^- = p.$$

By hypothesis p is a regular point of $\{\Sigma_{\hat{t}}^k\}$ (since $u^k(p) = \hat{t}$ is a regular value of $u^k|_{\mathcal{T}_j^k}$); additionally, p is neither on the lateral boundary of \mathcal{T}_j^k (because $u^k(p) = \hat{t} > \tau_j$) nor on the caps (in view of (4.14)). Thus, we can choose $r > 0$ sufficiently small so that $B_r(p) \subset \text{int } \mathcal{T}_j^k$ and that every level set of u^k in $B_r(p)$ is a small Lipschitz graph over $T_p \Sigma_{\hat{t}}^k$ (see Definition 2.11).

By (4.15), when i is large the points p_i^+ and p_i^- would be in $B_r(p)$. Since $B_r(p) \cap \{u^k \geq t_i\}$ is path-connected, there exists a path in $B_r(p) \cap \{u^k \geq t_i\}$ that joining p_i^+ and p_i^- together. This is in contradiction to the definitions of $\bar{\Omega}_{t_i}^{k,+}$ and $\bar{\Omega}_{t_i}^{k,-}$. Therefore, we conclude that $\{\Sigma_{\hat{t}}^k\}$ must have singular points on $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$. \square

⁴⁹The inside of the tube \mathcal{T}_j^k .

Lemma 4.11 will be used in the proofs of Proposition 4.13 and Proposition 4.14. In order that the condition (4.9) holds so as to apply Lemma 4.11, we make the following assumption for every splitting component of $\{\Sigma_t\}$.

Assumption 4.12. *Whenever \mathcal{S}_j is a singular component of $\{\Sigma_t\}$ that belongs to the splitting type, we assume that the flow $\{\Sigma_t\}$ “does not get into \mathcal{S}_j near the endpoints” in the sense that*

$$\min_{\mathcal{P}_+ \cap \mathcal{T}_j} \frac{\nabla u}{|\nabla u|} \cdot \nu_+ > -1, \quad \min_{\mathcal{P}_- \cap \mathcal{T}_j} \frac{\nabla u}{|\nabla u|} \cdot \nu_- > -1,$$

where ν_+ and ν_- are the outward (i.e., pointing toward the outside of \mathcal{T}_j) unit normal vector of the caps $\mathcal{P}_+ \cap \mathcal{T}_j$ and $\mathcal{P}_- \cap \mathcal{T}_j$, respectively. Note that ∇u vanishes at no points on the caps $\mathcal{P}_+ \cap \mathcal{T}_j$ and $\mathcal{P}_- \cap \mathcal{T}_j$ (see Remark 4.9) and that

$$\frac{\nabla u}{|\nabla u|} \cdot \nu_{\pm} \approx 0$$

on the part of $\mathcal{P}_{\pm} \cap \mathcal{T}_j$ that is close to the lateral boundary $\Sigma_{\tau_j} \cap \mathcal{T}_j$ due to the asymptotically cylindrical behavior of the flow $\{\Sigma_t\}$.

It follows from Proposition 2.18 that when k is large,

$$\min_{\mathcal{P}_+ \cap \mathcal{T}_j^k} \frac{\nabla u^k}{|\nabla u^k|} \cdot \nu_+ > -1, \quad \min_{\mathcal{P}_- \cap \mathcal{T}_j^k} \frac{\nabla u^k}{|\nabla u^k|} \cdot \nu_- > -1.$$

We are now in a position to prove the main results of this subsection.

Proposition 4.13. *Let \mathcal{S}_j be a singular component of the flow $\{\Sigma_t\}$ that belongs to the splitting type. Then for every sufficiently large k , there exist singular components of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ and they all belong to the splitting/bumpy type; moreover, if one of the singular components of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ is of the splitting type, then it would be the unique singular component of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$.*

Proof. Let k be a sufficiently large integer. By Proposition 2.16, Proposition 4.2, Proposition 4.10, Assumption 4.12, and Lemma 4.11, there exists

$$\hat{t} \in \left(\tau_j, \frac{1}{2} (T_1 + \hat{t}_j) \right)$$

so that $\Sigma_{\hat{t}}^k \cap \mathcal{T}_j^k$ has singular points of $\{\Sigma_t^k\}$, where $\hat{t}_j \in (T_1, T_2)$ is the time given in Proposition 4.10. It follows from Assumption 4.3 that

$$\hat{t} = T_{1,j}^k$$

is the unique singular time of $\{\Sigma_t^k\}$ in $\mathcal{T}_j^k \subset \mathcal{S}_j^{\hat{\delta}}$. Note that by Corollary 2.19 and Remark 4.9, all the singular points of $\{\Sigma_t^k\}$ in $\mathcal{S}_j^{\hat{\delta}}$ would be contained in $\text{int } \mathcal{T}_j^k$.

By Assumption 4.3 and Proposition 4.5 (see also Proposition 4.2), there are finitely many connected components of the singular set of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k , each of which belongs to the splitting or bumpy types (in particular, there are no round points of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k).

Now assume that there is a singular component $\hat{\mathcal{S}}_j^k$ of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k which belongs to the splitting type. By Proposition 4.7, any point $q \in \hat{\mathcal{S}}_j^k$ has a cylindrical scale r_j and every other cylindrical point of $\{\Sigma_t^k\}$ in $B_{r_j}(q)$ is located in a small Lipschitz graph over the axis of the tangent cylinder of $\{\Sigma_t^k\}$ at q . Applying Remark 4.8 and

Remark 4.9 (see also Assumption 4.3) to the singular component \dot{S}_j^k of $\{\Sigma_t^k\}$, we can find two hyperplanes $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ near the endpoints⁵⁰ of \dot{S}_j^k that are orthogonal to the tangent cylinders of $\{\Sigma_t^k\}$ at the endpoints, respectively, such that

- there are no singular points of $\{\Sigma_t^k\}$ on $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ (see Proposition 4.7);
- $\Sigma_{\tau_j}^k$ is a tube-like hypersurface around \dot{S}_j^k and it together with the two caps, $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$, bound a “truncated” tubular closed region $\dot{\mathcal{T}}_j^k \subset \mathcal{T}_j^k$ satisfying $\dot{S}_j^k \subset \text{int } \dot{\mathcal{T}}_j^k$;
- \dot{S}_j^k is the unique singular component of $\{\Sigma_t^k\}$ in $\dot{\mathcal{T}}_j^k$ (see Proposition 4.7).

Next, we would like to prove (by contradiction) that \dot{S}_j^k is indeed the only singular component of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k (instead of only in $\dot{\mathcal{T}}_j^k$). For convenience, let us assume that the “orientation” is chosen in such a way that $\dot{\mathcal{P}}_+^k$ is adjacent to \mathcal{P}_+ and $\dot{\mathcal{P}}_-^k$ is adjacent to \mathcal{P}_- . Thus, the tube \mathcal{T}_j^k would be cut by $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ into three closed pieces:

$$\mathcal{T}_j^k = \hat{\mathcal{T}}_j^{k+} \cup \dot{\mathcal{T}}_j^k \cup \hat{\mathcal{T}}_j^{k-},$$

where $\hat{\mathcal{T}}_j^{k+}$ is the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ (from the two ends); likewise, $\hat{\mathcal{T}}_j^{k-}$ is the closed region bounded by $\Sigma_{\tau_j}^k$, $\mathcal{P}_- \cap \mathcal{T}_j^k$, and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$.

Suppose the contrary that there exists another singular component \ddot{S}_j^k in \mathcal{T}_j^k . Since $\ddot{S}_j^k \cap \dot{\mathcal{T}}_j^k = \emptyset$, we may assume without loss of generality that $\ddot{S}_j^k \subset \hat{\mathcal{T}}_j^{k+}$. Following Remark 4.9, let $\tilde{\mathcal{P}}_+^k$ be the hyperplane that passes through the endpoint of \ddot{S}_j^k (the one that is near the side of $\dot{\mathcal{P}}_+^k$) and is parallel to $\dot{\mathcal{P}}_+^k$ (i.e., orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at the endpoint). Fix a point $p \in \ddot{S}_j^k$ and let $\ddot{\mathcal{P}}$ be the hyperplane passing through p and orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at p . Define $\ddot{\mathcal{T}}_j^k$ as the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\ddot{\mathcal{P}} \cap \mathcal{T}_j^k$ (from the two ends). Notice that $u^k = \tau_j < T_{1,j}^k$ on the lateral boundary of the tube $\ddot{\mathcal{T}}_j^k$ and that

$$\max_{\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k} u^k = \max_{\ddot{\mathcal{P}} \cap \mathcal{T}_j^k} u^k = T_{1,j}^k,$$

in light of the asymptotically cylindrical behavior of $\{\Sigma_t^k\}$ (see (3.4)).

On the other hand, applying Remark 3.8 to the saddle endpoint of \dot{S}_j^k on $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$, we would find $\hat{t}_j^k > T_{1,j}^k$ so that

$$\Sigma_t^k \cap \ddot{\mathcal{T}}_j^k \neq \emptyset \quad \forall t \in (T_{1,j}^k, \hat{t}_j^k].$$

Let $\hat{\Sigma}_{\hat{t}_j^k}^k$ be any connected component of $\Sigma_{\hat{t}_j^k}^k \cap \ddot{\mathcal{T}}_j^k$. Note that $\hat{\Sigma}_{\hat{t}_j^k}^k$ is away from the boundary $\partial \ddot{\mathcal{T}}_j^k$ since

$$\max_{\partial \ddot{\mathcal{T}}_j^k} u^k = T_{1,j}^k.$$

⁵⁰In the case where \dot{S}_j^k is a single point, $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ would two parallel hyperplanes on each “side” of the point, see Remark 3.8 and Remark 4.9.

Thus, $\hat{\Sigma}_{\dot{t}_j^k}^k$ would be a closed hypersurface in $\ddot{\mathcal{T}}_j^k$ and hence enclose an open region $\hat{\Omega}_{\dot{t}_j^k}^k$ in $\ddot{\mathcal{T}}_j^k$. As $u^k = \dot{t}_j^k$ on $\partial\hat{\Omega}_{\dot{t}_j^k}^k = \hat{\Sigma}_{\dot{t}_j^k}^k$, the maximum value

$$\tilde{T}_{1,j}^k := \max_{\hat{\Omega}_{\dot{t}_j^k}^k} u^k \geq \dot{t}_j^k > T_{1,j}^k$$

would be attained at some (interior) point in $\hat{\Omega}_{\dot{t}_j^k}^k$, which would be a critical point of u^k . It follows that $\tilde{T}_{1,j}^k$ is another singular time of $\{\Sigma_t^k\}$ in $\mathcal{T}_j^k \subset \mathcal{S}_j^\delta$, contradicting Assumption 4.3. Therefore, there are no other singular components in \mathcal{T}_j^k . \square

One last step to establish the ‘‘stability’’ of singular components of the splitting tube is to rule out the existence of the bumpy components in Proposition 4.13.

Proposition 4.14. *In fact, in Proposition 4.13 there are no singular components of $\{\Sigma_t^k\}$ that belong to the bumpy type. Therefore, $\{\Sigma_t^k\}$ has precisely one singular component in \mathcal{S}_j^δ , which is of the splitting type.*

Proof. Suppose that there is a singular component $\mathring{\mathcal{S}}_j^k$ of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k belonging to the bumpy type. When $\mathring{\mathcal{S}}_j^k$ is a curve, let p be the local maximum endpoint of $\mathring{\mathcal{S}}_j^k$ (see Definition 4.1) and let us assume (without loss of generality) that p is ‘‘adjacent’’ to \mathcal{P}_+ (so the other saddle endpoint is adjacent to \mathcal{P}_-). When $\mathring{\mathcal{S}}_j^k$ is a single one-sided saddle point, let p be $\mathring{\mathcal{S}}_j^k$ itself and let us assume that u^k is a local maximum point ‘‘on the side toward \mathcal{P}_+ ’’ (see Definition 3.3).

Recall that \mathcal{S}_j is a singular component of $\{\Sigma_t\}$ that belongs to the splitting type. By Proposition 4.10 there exists $\dot{t}_j \in (T_1, T_2)$ so that

$$\bar{\Omega}_t \cap \mathcal{P}_+ \cap \mathcal{T}_j \supset \Sigma_t \cap \mathcal{P}_+ \cap \mathcal{T}_j \neq \emptyset \quad \forall t \in (T_1, \dot{t}_j].$$

By Proposition 2.16 and Proposition 4.2, if k is sufficiently, we may assume that $\dot{t}_j > T_{1,j}^k$ and that

$$\bar{\Omega}_{\frac{1}{2}(T_{1,j}^k + \dot{t}_j)} \cap \mathcal{P}_+ \cap \mathcal{T}_j^k \neq \emptyset,$$

where $T_{1,j}^k \approx T_1$ is the unique singular time of $\{\Sigma_t^k\}$ in $\mathcal{T}_j^k \subset \mathcal{S}_j^\delta$ (see Proposition 4.13). Then applying the intermediate value theorem to $u^k|_{\mathcal{P}_+ \cap \mathcal{T}_j^k}$ implies that

$$(4.16) \quad \Sigma_t^k \cap \mathcal{P}_+ \cap \mathcal{T}_j^k \neq \emptyset \quad \forall t \in \left[\tau_j, \frac{1}{2}(T_{1,j}^k + \dot{t}_j) \right].$$

In addition, by Assumption 4.12 there holds

$$(4.17) \quad \min_{\mathcal{P}_+ \cap \mathcal{T}_j^k} \frac{\nabla u^k}{|\nabla u^k|} \cdot \nu_+ > -1$$

where ν_+ is the unit normal vector of $\mathcal{P}_+ \cap \mathcal{T}_j^k$ that points toward the outside of the tube \mathcal{T}_j^k .

On the other hand, using Remark 3.4 and Proposition 4.7, we could find two parallel hyperplanes, $\tilde{\mathcal{P}}_+^k$ and $\mathring{\mathcal{P}}_+^k$, with the following properties:

- $\tilde{\mathcal{P}}_+^k$ passes through p and is orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at p ; $\mathring{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ is close to $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and is ‘‘between’’ $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\mathcal{P}_+ \cap \mathcal{T}_j^k$.

- There are no singular points of $\{\Sigma_t^k\}$ on $\mathring{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and

$$\bar{t} := \max_{\mathring{\mathcal{P}}_+^k \cap \mathcal{T}_j^k} u^k \in (\tau_j, u^k(p))$$

(see (3.5)). Note that $u^k(p) = T_{1,j}^k$.

In view of the asymptotically cylindrical behavior of $\{\Sigma_t^k\}$ on $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \setminus \{p\}$, we infer that

$$(4.18) \quad \Sigma_t^k \cap \tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \neq \emptyset \quad \forall t \in [\tau_j, T_{1,j}^k)$$

and that

$$(4.19) \quad \frac{\nabla u^k}{|\nabla u^k|} \cdot \tilde{\nu} \approx 0 \quad \text{on } \tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \cap \{\tau_j \leq u^k < T_{1,j}^k\},$$

where $\tilde{\nu}$ is a unit normal vector of $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$.

Now define $\mathring{\mathcal{T}}_j^k$ as the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ (from the two ends). Note that $\mathring{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ would cut $\mathring{\mathcal{T}}_j^k$ into two pieces. Let

$$t_* = \frac{1}{2}(\bar{t} + T_{1,j}^k) < T_{1,j}^k,$$

which is a regular value of $u^k|_{\mathring{\mathcal{T}}_j^k}$ by Assumption 4.3. Note that by (4.16) and (4.18) we have

$$\bar{\Omega}_{t_*}^k \cap \mathcal{P}_+ \cap \mathcal{T}_j^k \neq \emptyset, \quad \bar{\Omega}_{t_*}^k \cap \tilde{\mathcal{P}}_+^k \cap \mathring{\mathcal{T}}_j^k \neq \emptyset.$$

Moreover, since $u^k \leq \bar{t}$ on $\mathring{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$, we have

$$\bar{\Omega}_{t_*}^k \cap \mathring{\mathcal{P}}_+^k \cap \mathring{\mathcal{T}}_j^k = \emptyset.$$

Thus, every connected component of $\bar{\Omega}_{t_*}^k \cap \mathring{\mathcal{T}}_j^k$ intersects at most one of the caps, i.e., $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$. Applying Lemma 4.11 to the flow $\{\Sigma_t^k\}$ on the tube $\mathring{\mathcal{T}}_j^k$ (noting that we have (4.17) and (4.19)), we deduce that $\Sigma_{\hat{t}}^k \cap \mathring{\mathcal{T}}_j^k$ contains a singular point of $\{\Sigma_t^k\}$ for some $\hat{t} \in (\tau_j, t_*)$. In particular, $\hat{t} < T_{1,j}^k$ is another singular time of $\{\Sigma_t^k\}$ in $\mathring{\mathcal{T}}_j^k \subset \mathcal{S}_j^{\hat{t}}$. This is in contradiction with Assumption 4.3. Therefore, there are no singular components of $\{\Sigma_t^k\}$ that belong to the bumpy type. \square

4.3. Bumpy type. In this subsection we shall prove the ‘‘stability’’ of singular components of the bumpy type under Assumption 4.16. Specifically, let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ that belongs to the bumpy type, when k is large we assume that $\{\Sigma_t^k\}$ has singular components near \mathcal{S}_j (see Assumption 4.16) and show that all these singular components must be of the bumpy type (see Proposition 4.18) and that there is indeed only one singular component of $\{\Sigma_t^k\}$ near \mathcal{S}_j (see Proposition 4.19). Note that $u(\mathcal{S}_j) = T_1 < T_{ext}$ by Proposition 4.5.

Let us begin with the following remark, which is the counterpart of Remark 4.9 in Section 4.2 for the bumpy case.

Remark 4.15. When \mathcal{S}_j is a curve of the bumpy type, say

$$\mathcal{S}_j = \{\Gamma(s) : s \in [-1, 1]\}.$$

Let us assume without loss of generality that $\Gamma(1)$ is the local maximum point of u and $\Gamma(-1)$ is the one-sided saddle point of u . By Remark 4.8, $\Sigma_{\tau_j} \cap \mathcal{S}_j^{\tau_j}$ is a

tube-like hypersurface around \mathcal{S}_j . Near the endpoints $\Gamma(1)$ and $\Gamma(-1)$, by Remark 3.4 and Remark 3.8, we can find two hyperplanes, \mathcal{P}_+ and \mathcal{P}_- , that are orthogonal to the tangent cylinders of $\{\Sigma_t\}$ at $\Gamma(1)$ and $\Gamma(-1)$, respectively, such that⁵¹

$$\mathcal{P}_+ \cap \mathcal{S}_j^{r_j} \cap \mathcal{S}_j = \emptyset, \quad \mathcal{P}_- \cap \mathcal{S}_j^{r_j} \cap \mathcal{S}_j = \emptyset$$

and that the tube-like connected hypersurface Σ_{τ_j} together with \mathcal{P}_+ and \mathcal{P}_- bound a tubular closed region \mathcal{T}_j in $\mathcal{S}_j^{r_j}$ with the following properties:

- (1) $\mathcal{S}_j \subset \text{int } \mathcal{T}_j$.
- (2) Let $\tilde{\mathcal{P}}_+$ and $\tilde{\mathcal{P}}_-$ be two hypersurfaces parallel to \mathcal{P}_+ and \mathcal{P}_- ,⁵² respectively, such that

$$\tilde{\mathcal{P}}_+ \cap \mathcal{T}_j \cap \mathcal{S}_j = \{\Gamma(1)\}, \quad \tilde{\mathcal{P}}_- \cap \mathcal{T}_j \cap \mathcal{S}_j = \{\Gamma(-1)\}.$$

- (3) On the cap $\mathcal{P}_+ \cap \mathcal{T}_j$ we have⁵³

$$\bar{t}_j := \max_{\mathcal{P}_+ \cap \mathcal{T}_j} u \in (\tau_j, T_1).$$

- (4) There exists a time $\hat{t}_j > T_1$ so that⁵⁴

$$\Sigma_t \cap \mathcal{P}_- \cap \mathcal{T}_j \neq \emptyset \quad \forall t \in [\tau_j, \hat{t}_j].$$

In the case where \mathcal{S}_j is a single one-sided saddle point (which can be regarded as a “degenerate” curve with a local maximum endpoint and a one-sided saddle endpoint, see the comment following Definition 4.1), the aforementioned results still hold. In this case, the two hyperplanes, \mathcal{P}_+ and \mathcal{P}_- , are parallel (since both are orthogonal to the tangent cylinder of $\{\Sigma_t\}$ at \mathcal{S}_j) and close to each other; \mathcal{P}_+ is assumed to be on the side in which u is a local maximum point (see Definition 3.3). Note that $\tilde{\mathcal{P}}_+ = \tilde{\mathcal{P}}_-$ is the hyperplane that passes through \mathcal{S}_j and orthogonal to the tangent cylinder of $\{\Sigma_t\}$ at \mathcal{S}_j .

Furthermore, by Proposition 2.16, Proposition 4.2, and Remark 4.8, when k is large, $\Sigma_{\tau_j}^k$ and the two hyperplanes \mathcal{P}_+ and \mathcal{P}_- would also bound a closed tubular region \mathcal{T}_j^k in $\mathcal{S}_j^{r_j}$; in addition, let

$$\bar{t}_j^k = \max_{\mathcal{P}_+ \cap \mathcal{T}_j^k} u^k$$

and $T_{1,j}^k$ be the unique singular time of $\{\Sigma_t^k\}$ in \mathcal{S}_j^δ , then we have

$$|\bar{t}_j^k - \bar{t}_j| + |T_{1,j}^k - T_1| \leq \frac{1}{3} \min \{T_1 - \bar{t}_j, \bar{t}_j - \tau_j\}.$$

It follows that

$$(4.20) \quad \bar{t}_j^k \in (\tau_j, T_{1,j}^k).$$

Unlike the vanishing and splitting cases, we assume (instead of proving) the existence of singularities of $\{\Sigma_t^k\}$ near a bumpy component of $\{\Sigma_t\}$ as follows.

Assumption 4.16. *In case \mathcal{S}_j is a singular component of $\{\Sigma_t\}$ that belongs to the bumpy type, we assume that for every sufficiently large k , the flow $\{\Sigma_t^k\}$ has singularities in \mathcal{S}_j^δ .*

⁵¹ \mathcal{P}_+ corresponds to $\{z = z_0\}$ in Remark 3.4 with $z_0 > 0$ sufficiently close to 0; \mathcal{P}_- corresponds to $\{z = \varepsilon\}$ in Remark 3.8.

⁵² $\tilde{\mathcal{P}}_+$ corresponds to $\{z = 0\}$ in Remark 3.4; $\tilde{\mathcal{P}}_-$ corresponds to $\{z = 0\}$ in Remark 3.8.

⁵³See (3.5).

⁵⁴See (3.19).

Next, for the singular components of $\{\Sigma_t^k\}$ in \mathcal{S}_j^δ , we are going to characterize their types in Proposition 4.18 and prove the uniqueness of the singular components in Proposition 4.19. To streamline the proofs, in the following remark we set up the requisite notations and provide a preliminary description of the singular components of $\{\Sigma_t^k\}$ in \mathcal{S}_j^δ .

Remark 4.17. Let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ that belongs to the bumpy type. When k is large, by Corollary 2.19 and Remark 4.15, all the singular points of $\{\Sigma_t^k\}$ in \mathcal{S}_j^δ would be contained in $\text{int } \mathcal{T}_j^k$. Moreover, by Assumption 4.3 and Proposition 4.5 (see also Proposition 4.2), there are finitely many connected components of the singular set of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k , each of which belongs to the splitting/bumpy type (in particular, there are no round points of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k).

Let $\dot{\mathcal{S}}_j^k$ be a singular component of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k . By Proposition 4.7, any point $q \in \dot{\mathcal{S}}_j^k$ has a cylindrical scale r_j and every other cylindrical point of $\{\Sigma_t^k\}$ in $B_{r_j}(q)$ is located in a small Lipschitz graph over the axis of the tangent cylinder of $\{\Sigma_t^k\}$ at q . Applying Remark 4.9 (if $\dot{\mathcal{S}}_j^k$ is of the splitting type) / Remark 4.15 (if $\dot{\mathcal{S}}_j^k$ is of the bumpy type) to the singular component $\dot{\mathcal{S}}_j^k$ of $\{\Sigma_t^k\}$, we can find two hyperplanes $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ near the endpoints of $\dot{\mathcal{S}}_j^k$ that are orthogonal to the tangent cylinders of $\{\Sigma_t^k\}$ at the endpoints, respectively, such that

- (1) there are no singular points of $\{\Sigma_t^k\}$ on $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ (see Proposition 4.7);
- (2) $\Sigma_{\tau_j}^k$ is a tube-like hypersurface around $\dot{\mathcal{S}}_j^k$ and it together with the two caps, $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$, bound a “truncated” tubular closed region $\hat{\mathcal{T}}_j^k \subset \mathcal{T}_j^k$ satisfying $\dot{\mathcal{S}}_j^k \subset \text{int } \hat{\mathcal{T}}_j^k$;
- (3) $\dot{\mathcal{S}}_j^k$ is the only singular component of $\{\Sigma_t^k\}$ in $\hat{\mathcal{T}}_j^k$ (see Proposition 4.7).

For convenience, let us assume that the “orientation” is chosen in such a way that $\dot{\mathcal{P}}_+^k$ is adjacent to \mathcal{P}_+ and $\dot{\mathcal{P}}_-^k$ is adjacent to \mathcal{P}_- . Thus, the tube \mathcal{T}_j^k would be cut by $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ into three closed pieces:

$$\mathcal{T}_j^k = \hat{\mathcal{T}}_j^{k+} \cup \hat{\mathcal{T}}_j^k \cup \hat{\mathcal{T}}_j^{k-},$$

where $\hat{\mathcal{T}}_j^{k+}$ is the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ (from the two ends); likewise, $\hat{\mathcal{T}}_j^{k-}$ is the closed region bounded by $\Sigma_{\tau_j}^k$, $\mathcal{P}_- \cap \mathcal{T}_j^k$, and $\dot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$.

Proposition 4.18. *Let \mathcal{S}_j be a singular component of $\{\Sigma_t\}$ that belongs to the bumpy type and let $\dot{\mathcal{S}}_j^k$ be any singular component of $\{\Sigma_t^k\}$ in \mathcal{S}_j^δ as stated in Remark 4.17. Then*

- if the singular component $\dot{\mathcal{S}}_j^k$ is a curve, the endpoint that is adjacent to \mathcal{P}_+ must be a local maximum point of u^k ;
- if $\dot{\mathcal{S}}_j^k$ is a single point, $\dot{\mathcal{S}}_j^k$ must be a local maximum point of u^k “on the side toward \mathcal{P}_+ ” (see Definition 3.3).

In either case, $\dot{\mathcal{S}}_j^k$ is of the bumpy type.

Proof. Suppose the contrary that

- when \mathcal{S}_j is a curve: the endpoint p that is adjacent to \mathcal{P}_+ is a one-sided saddle point of u^k ;

- when $\dot{\mathcal{S}}_j^k$ is a single point: $p = \dot{\mathcal{S}}_j^k$ is not a local maximum point of u^k on the side toward \mathcal{P}_+ .

Recall that the choice of the two hyperplanes $\dot{\mathcal{P}}_+^k$ and $\dot{\mathcal{P}}_-^k$ in Remark 4.17 is based on applying Remark 4.9/Remark 4.15 to the singular component $\dot{\mathcal{S}}_j^k$ of $\{\Sigma_t^k\}$. By either of the remarks, there exists a time $\dot{t}_j^k > T_{1,j}^k$ so that

$$(4.21) \quad \Sigma_t^k \cap \dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \neq \emptyset \quad \forall t \in [\tau_j, \dot{t}_j^k].$$

On the other hand, recall that by Remark 4.15 we have

$$(4.22) \quad \bar{t}_j^k = \max_{\mathcal{P}_+ \cap \mathcal{T}_j^k} u^k \in (\tau_j, T_{1,j}^k).$$

Let $\tilde{\mathcal{P}}_+^k$ be the hyperplane passing through p and orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at p . Note that $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ is parallel (and close) to $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and is “between” $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$. In view of the asymptotically cylindrical behavior of $\{\Sigma_t^k\}$ on $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \setminus \{p\}$, we have (see (3.4))

$$(4.23) \quad \max_{\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k} u^k = T_{1,j}^k.$$

Now define $\check{\mathcal{T}}_j^k$ as the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ (from the two ends). Notice that by (4.21) we have

$$\Sigma_{\dot{t}_j^k}^k \cap \check{\mathcal{T}}_j^k \neq \emptyset$$

and that by (4.22), (4.23), and that $u^k = \tau_j$ on the lateral, we have

$$\Sigma_{\bar{t}_j^k}^k \cap \partial\check{\mathcal{T}}_j^k = \emptyset.$$

As \dot{t}_j^k is a regular value⁵⁵ of $u^k|_{\check{\mathcal{T}}_j^k}$, $\Sigma_{\dot{t}_j^k}^k \cap \check{\mathcal{T}}_j^k$ is a closed hypersurface, which encloses an open region $\check{\Omega}_{\dot{t}_j^k}^k \subset \check{\mathcal{T}}_j^k$. It follows that

$$\tilde{T}_{1,j}^k := \max_{\check{\Omega}_{\dot{t}_j^k}^k} u^k \geq \dot{t}_j^k > T_{1,j}^k$$

would be attained at some interior point of $\check{\mathcal{T}}_j^k$, which would be a critical point of u^k . Consequently, $\tilde{T}_{1,j}^k$ is another singular time of $\{\Sigma_t^k\}$ in $\check{\mathcal{T}}_j^k \subset \mathcal{S}_j^{\delta}$, contradicting Assumption 4.3. Therefore, when $\dot{\mathcal{S}}_j^k$ is a curve, p must be a local maximum point of u^k ; when $\dot{\mathcal{S}}_j^k$ is a single point, p is a local maximum point of u^k on the side toward \mathcal{P}_+ . \square

Proposition 4.19. *In Proposition 4.18, there is actually only one singular component of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k . Therefore, $\{\Sigma_t^k\}$ has precisely one singular component in \mathcal{S}_j^{δ} , which is of the bumpy type.*

Proof. Let $\dot{\mathcal{S}}_j^k$ be a singular component of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k as stated in Proposition 4.18. Suppose the contrary that there exists another singular component $\check{\mathcal{S}}_j^k$ of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k . Since $\check{\mathcal{S}}_j^k \cap \dot{\mathcal{T}}_j^k = \emptyset$, we may assume without loss of generality that $\check{\mathcal{S}}_j^k \subset \hat{\mathcal{T}}_j^{k+}$ (see Remark 4.17).

⁵⁵ $T_{1,j}^k$ is the only critical value of $u^k|_{\check{\mathcal{T}}_j^k}$ by Assumption 4.3.

Let q be the endpoint of $\dot{\mathcal{S}}_j^k$ that is adjacent to $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$; in case $\dot{\mathcal{S}}_j^k$ is a single one-sided saddle point, then q is $\dot{\mathcal{S}}_j^k$ itself. By Proposition 4.18, q is a one-sided saddle point of u^k such that it is not a local maximum point of u^k on the side toward $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$.

By Remark 4.15 and Remark 4.17, we can find two hyperplanes $\ddot{\mathcal{P}}_+^k$ and $\ddot{\mathcal{P}}_-^k$ near the endpoints of $\dot{\mathcal{S}}_j^k$ that are orthogonal to the tangent cylinders of $\{\Sigma_t^k\}$ at the endpoints, respectively, such that the following hold:

- (1) $\ddot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ are between $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$; the orientation is chosen in such a way that $\ddot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ is adjacent to $\mathcal{P}_+ \cap \mathcal{T}_j^k$ and $\ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ is adjacent to $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$.
- (2) There are no singular points of $\{\Sigma_t^k\}$ on $\ddot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$.
- (3) There exists a time $\ddot{t}_j^k > T_{1,j}^k$ so that

$$(4.24) \quad \Sigma_t^k \cap \ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k \neq \emptyset \quad \forall t \in [\tau_j, \ddot{t}_j^k].$$

Let $\tilde{\mathcal{P}}_-^k$ be the hyperplane passing through q and orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at q . Note that $\tilde{\mathcal{P}}_-^k$ is parallel and close to $\ddot{\mathcal{P}}_-^k$ and that $\tilde{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ is between $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$. In view of the asymptotically cylindrical behavior of $\{\Sigma_t^k\}$ on $\tilde{\mathcal{P}}_-^k \cap \mathcal{T}_j^k \setminus \{q\}$, we have (see (3.4))

$$(4.25) \quad \max_{\tilde{\mathcal{P}}_-^k \cap \mathcal{T}_j^k} u^k = T_{1,j}^k.$$

Let p be the endpoint $\dot{\mathcal{S}}_j^k$ that is adjacent to $\dot{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$. Let $\tilde{\mathcal{P}}_+^k$ be the hyperplane passing through p and orthogonal to the tangent cylinder of $\{\Sigma_t^k\}$ at p . Note that $\tilde{\mathcal{P}}_+^k$ is parallel and close to $\dot{\mathcal{P}}_+^k$. In view of the asymptotically cylindrical behavior of $\{\Sigma_t^k\}$ on $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k \setminus \{p\}$, we have (see (3.4))

$$(4.26) \quad \max_{\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k} u^k = T_{1,j}^k.$$

Now define $\widehat{\mathcal{T}}_j^k$ as the tubular closed region bounded by $\Sigma_{\tau_j}^k$ (from the lateral) and the two caps $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ (from the two ends). Note that $\ddot{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$ is between $\tilde{\mathcal{P}}_+^k \cap \mathcal{T}_j^k$ and $\tilde{\mathcal{P}}_-^k \cap \mathcal{T}_j^k$. It follows from (4.24) that

$$\Sigma_{\ddot{t}_j^k}^k \cap \widehat{\mathcal{T}}_j^k \neq \emptyset.$$

Note that

$$\Sigma_{\ddot{t}_j^k}^k \cap \partial \widehat{\mathcal{T}}_j^k = \emptyset$$

by (4.25), (4.26), and that $u^k = \tau_j$ on the lateral. Since \ddot{t}_j^k is a regular value⁵⁶ of $u^k|_{\widehat{\mathcal{T}}_j^k}$, $\Sigma_{\ddot{t}_j^k}^k \cap \widehat{\mathcal{T}}_j^k$ is a closed hypersurface, which encloses an open region $\widehat{\Omega}_{\ddot{t}_j^k}^k$ in $\widehat{\mathcal{T}}_j^k$. Thus,

$$\widehat{T}_{1,j}^k := \max_{\widehat{\Omega}_{\ddot{t}_j^k}^k} u^k \geq \ddot{t}_j^k > T_{1,j}^k$$

⁵⁶ $T_{1,j}^k$ is the only critical value of $u^k|_{\widehat{\mathcal{T}}_j^k}$.

would be attained at some interior point of $\widehat{\mathcal{T}}_j^k$, which would be a critical point of u^k . Consequently, $\widetilde{T}_{1,j}^k$ is another singular time of $\{\Sigma_t^k\}$ in $\widehat{\mathcal{T}}_j^k \subset \mathcal{S}_j^{\delta}$, contradicting Assumption 4.3. Thus, \mathcal{S}_j^k is the only one singular component of $\{\Sigma_t^k\}$ in \mathcal{T}_j^k . \square

4.4. Appendix: Two-convexity. The purpose of this appendix is to show that in Theorem 1.3, as Σ_0 is two-convex, Σ_0^k would also be two-convex when k is large (see Proposition 4.22). To this end, let us begin with the following lemma, which is a direct result of the min-max theorem in linear algebra.

Lemma 4.20. *Let V be a finite dimensional vector space with an inner product g . Let S be a self-adjoint operator on the inner product space (V, g) with eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots$. Let A be the associated symmetric bilinear form of S , namely,*

$$A(v, w) = g(Sv, w) \quad \forall v, w \in V.$$

Given a pair of linearly independent vectors v_1 and v_2 in V , let $\mathbf{G}(v_1, v_2)$ and $\mathbf{A}(v_1, v_2)$ be the 2×2 matrices whose (k, l) -components are given by respectively

$$\mathbf{G}_{kl}(v_1, v_2) = g(v_k, v_l), \quad \mathbf{A}_{kl}(v_1, v_2) = A(v_k, v_l), \quad \text{for } k, l \in \{1, 2\}.$$

Then

- (1) *the symmetric matrix $\mathbf{G}(v_1, v_2)$ is positive definite;*
- (2) *whenever $\text{span}\{\tilde{v}_1, \tilde{v}_2\} = \text{span}\{v_1, v_2\}$, we have*

$$\text{trace}(\mathbf{G}^{-1}\mathbf{A})(\tilde{v}_1, \tilde{v}_2) = \text{trace}(\mathbf{G}^{-1}\mathbf{A})(v_1, v_2).$$

Thus, $\mathbf{G}^{-1}\mathbf{A}$ can be regarded as a function defined on the Grassmannian $Gr_2(V)$, i.e., the set of all 2-dimensional vector subspaces of V . Most importantly,

$$\lambda_1 + \lambda_2 = \min_{Gr_2(V)} \text{trace}(\mathbf{G}^{-1}\mathbf{A}).$$

The following corollary follows from applying Lemma 4.20 to the Weingarten map of a hypersurface Σ on the tangent space at each point so as to obtain an expression of $\kappa_1 + \kappa_2$, which will be used in Proposition 4.22. Note that as the function $\mathbf{G}^{-1}\mathbf{A}(v_1, v_2)$ in Lemma 4.20 is defined on the Grassmannian, we may assume that the two linearly independent tangent vectors,

$$\begin{aligned} v_1 &= v_1^1 \partial_1 + \dots + v_1^{n-1} \partial_{n-1} \simeq (v_1^1, \dots, v_1^{n-1}), \\ v_2 &= v_2^1 \partial_1 + \dots + v_2^{n-1} \partial_{n-1} \simeq (v_2^1, \dots, v_2^{n-1}), \end{aligned}$$

are orthonormal with respect to the dot product in order to have the compactness.

Corollary 4.21. *Let Σ be a hypersurface and let $x = x(\xi)$ be a local parametrization of Σ . Let $g_{ij}(\xi)$ and $A_{ij}(\xi)$ be the components of the metric and the second fundamental form of Σ with respect to the local parametrization $x(\xi)$, respectively.*

Let $\mathbb{V} \subset \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ be the set of all pairs of orthonormal (with respect to the dot product) vectors in \mathbb{R}^{n-1} . For each ξ , define two matrix-valued functions $\mathbf{G}(\xi; \cdot)$ and $\mathbf{A}(\xi; \cdot)$ on \mathbb{V} as follows: Given

$$\{v_1 = (v_1^1, \dots, v_1^{n-1}), \quad v_2 = (v_2^1, \dots, v_2^{n-1})\} \in \mathbb{V},$$

let $\mathbf{G}(\xi; v_1, v_2)$ and $\mathbf{A}(\xi; v_1, v_2)$ be the 2×2 matrices whose (k, l) -components are given by respectively

$$\mathbf{G}_{kl}(\xi; v_1, v_2) = g_{ij}(\xi) v_k^i v_l^j, \quad \mathbf{A}_{kl}(\xi; v_1, v_2) = A_{ij}(\xi) v_k^i v_l^j, \quad \text{for } k, l \in \{1, 2\}.$$

Then we have

$$(4.27) \quad \kappa_1(\xi) + \kappa_2(\xi) = \min_{\mathbb{V}} \text{trace}(\mathbf{G}^{-1}\mathbf{A})(\xi; \cdot),$$

where $\kappa_1(\xi)$ and $\kappa_2(\xi)$ are the smallest two principal curvatures of Σ at $x(\xi)$. Also, note that by the compactness of \mathbb{V} and the continuity of $\mathbf{G}^{-1}\mathbf{A}$, $\kappa_1 + \kappa_2$ is a continuous function.

Since Σ_0 is a two-convex closed hypersurface, the function $\kappa_1 + \kappa_2$ is bounded below by a positive constant on Σ_0 . By virtue of (4.27), we get the following result.

Proposition 4.22. *Let β be a positive constant such that*

$$\min_{\Sigma_0} (\kappa_1 + \kappa_2) > \beta \max_{\Sigma_0} H.$$

Since

$$\Sigma_0^k \xrightarrow{C^4} \Sigma_0 \quad \text{as } k \rightarrow \infty,$$

when k is large, Σ_0^k would be β -uniformly convex in the sense that

$$(4.28) \quad \kappa_1 + \kappa_2 \geq \beta H > 0.$$

It then follows from [CHN] that the LSF $\{\Sigma_t^k\}$ is β -uniformly two-convex in the sense that (4.28) holds at every regular point (with respect to the unit normal vector field $\frac{\nabla u^k}{|\nabla u^k|}$).

REFERENCES

- [Al] W.K. Allard, On the first variation of a varifold, *Ann. of Math. (2)* 95 (1972), 417–491.
- [An] B. Andrews, Noncollapsing in mean-convex mean curvature flow, *Geom. Topol.* 16 (2012), no.3, 1413–1418.
- [AAG] S. Altschuler, S. Angenent, and Y. Giga, Mean curvature flow through singularities for surfaces of rotation, *J. Geom. Anal.* 5 (1995), no.3, 293–358.
- [ALM] B. Andrews, M. Langford, and J. McCoy, Non-collapsing in fully non-linear curvature flows, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 30 (2013), no.1, 23–32.
- [B] K. Brakke, *The Motion of a surface by its mean curvature*, Mathematical Notes, vol. 20, Princeton University Press, Princeton, 1978.
- [CGG] Y.G. Chen, Y. Giga, and S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Differential Geom.* 33 (1991), no. 3, 749–786.
- [CHN] J. Cheeger, R. Haslhofer, and A. Naber, Quantitative stratification and the regularity of mean curvature flow, *Geom. Funct. Anal.* 23 (2013), no. 3, 828–847.
- [CM1] T.H. Colding and W.P. Minicozzi II, Generic mean curvature flow I: generic singularities, *Ann. of Math. (2)* 175 (2012), no. 2, 755–833.
- [CM2] T.H. Colding and W.P. Minicozzi II, Uniqueness of blowups and Łojasiewicz inequalities, *Ann. of Math. (2)* 182 (2015), no. 1, 221–285.
- [CM3] T.H. Colding and W.P. Minicozzi II, The singular set of mean curvature flow with generic singularities, *Invent. Math.* 204 (2016), no. 2, 443–471.
- [CM4] T.H. Colding and W.P. Minicozzi II, Differentiability of the arrival time, *Comm. Pure Appl. Math.* 69 (2016), no. 12, 2349–2363.
- [CM5] T.H. Colding and W.P. Minicozzi II, Regularity of the level set flow, *Comm. Pure Appl. Math.* 71 (2018), no. 4, 814–824.
- [E] K. Ecker, *Regularity theory for mean curvature flow*, Progress in Nonlinear Differential Equations and their Applications, 57. Birkhäuser Boston, Inc., Boston, MA, 2004. xiv+165 pp. ISBN: 0-8176-3243-3.
- [EH] K. Ecker and G. Huisken, Interior estimates for hypersurfaces moving by mean curvature, *Invent. Math.* 105 (1991), no.3, 547–569.
- [ES] L.C. Evans and J. Spruck, Motion of level sets by mean curvature I, *J. Differential Geom.* 33 (1991), no. 3, 635–681.

- [G1] S.-H. Guo, Asymptotic behavior and stability of mean curvature flow with a conical end, *Adv. Math.* 375 (2020), 107408, 63 pp.
- [G2] S.-H. Guo, Singular set and curvature blow-up rate of the level set flow, *J. Reine Angew. Math.* 807 (2024), 1–29.
- [GP] V. Guillemin and A. Pollack, *Differential topology*, Reprint of the 1974 original. AMS Chelsea Publishing, Providence, RI, 2010. xviii+224 pp. ISBN: 978-0-8218-5193-7 58-01 (57-01).
- [H1] G. Huisken, Flow by mean curvature of convex surfaces into spheres, *J. Differential Geom.* 20 (1984), no. 1, 237–266.
- [H2] G. Huisken, Asymptotic behavior for singularities of the mean curvature flow, *J. Differential Geom.* 31 (1990), no. 1, 285–299.
- [HK] R. Haslhofer and B. Kleiner, Mean curvature flow of mean convex hypersurfaces, *Comm. Pure Appl. Math.* 70 (2017), no. 3, 511–546.
- [HP] G. Huisken and A. Polden, *Geometric evolution equations for hypersurfaces*, Calculus of variations and geometric evolution problems (Cetraro, 1996), Springer-Verlag, Berlin, 1999, pp. 45–84.
- [I1] T. Ilmanen, Elliptic regularization and partial regularity for motion by mean curvature. *Mem. Amer. Math. Soc.* 108 (1994), no. 520, x+90 pp.
- [I2] T. Ilmanen, *Singularities of mean curvature flow of surfaces*, 1995, preprint.
- [M] C. Mantegazza, *Lecture notes on mean curvature flow*, *Progr. Math.*, 290 Birkhäuser/Springer Basel AG, Basel, 2011. xii+166 pp.
- [OS] S. Osher and J. Sethian, Fronts propagating with curvature-dependent speed: Algorithms based on Hamilton-Jacobi formulations, *Journal of Computational Physics*, Volume 79, Issue 1, 1988, Pages 12-49.
- [W1] B. White, The size of the singular set in mean curvature flow of mean-convex sets, *J. Amer. Math. Soc.* 13 (2000), no. 3, 665–695.
- [W2] B. White, The nature of singularities in mean curvature flow of mean-convex sets, *J. Amer. Math. Soc.* 16 (2003), no. 1, 123–138.
- [W3] B. White, A local regularity theorem for mean curvature flow, *Ann. of Math. (2)* 161 (2005), no. 3, 1487–1519.
- [W4] B. White, Subsequent singularities in mean-convex mean curvature flow, *Calc. Var. Partial Differential Equations* 54 (2015), no. 2, 1457–1468.

Department of Mathematics, National Taiwan University, Taipei 106, Taiwan.

E-mail: shguo@ntu.edu.tw