

A unital in the generalized hexagon of order two, and an exceptional isomorphism between finite groups of Lie type.

Markus J. Stroppel

December 13, 2024

Abstract

We construct a model of the Hermitian unital of order 3 (obtained from the non-degenerate hermitian form in three variables over the field of order 9) inside the octonion algebra over the field of order 2. This construction is invariant under the automorphism group of that algebra, and explains the known isomorphism from the finite group of exceptional Lie type onto the group of semi-similitudes of the hermitian form.

Mathematics Subject Classification (MSC 2000): 20D06, 17A75, 17A36, 17A45, 51E12,

Keywords: classical group, exceptional isomorphism, group of Lie type, unital, generalized hexagon, generalized polygon, octonion, composition algebra, subalgebra, automorphism

Introduction

We consider the (necessarily split) octonion algebra \mathbb{O} over the finite field \mathbb{F}_q with q elements. The automorphism group $\text{Aut}(\mathbb{O})$ is a finite group of Lie type, namely $G_2(q)$. For $q > 2$, this is a simple group ([6], [7], see [27, Ch. 4]). For the smallest case $q = 2$, it is known (e.g., see [27, 4.4.4]) that this group is not simple, in fact its commutator group has index 2, and is a simple group isomorphic to $\text{PSU}(3, \mathbb{F}_9 | \mathbb{F}_3)$. That group is a subgroup of index 2 in the full group of automorphisms of the Hermitian unital of order 3 (see Section 5).

We are going to construct an incidence structure isomorphic to that Hermitian unital. Our construction will be invariant under $\text{Aut}(\mathbb{O})$, and we obtain an isomorphism from that group onto the automorphism group $\text{PTU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ of the Hermitian unital. See the discussion in Section 2 below for more details on the group $\text{PTU}(3, \mathbb{F}_9 | \mathbb{F}_3)$.

Aiming at an elementary treatment, we will only assume basic information about composition algebras. In particular, we include a determination of the order of $G_2(q)$, see 3.5 below.

1 The octonion algebra, and its automorphisms

Let \mathbb{F} be any field. The split octonion algebra over \mathbb{F} is obtained by the following doubling process (known as the Cayley-Dickson process, see [24, Sect. 1.5]). We start with the split quaternion algebra $\mathbb{H} = \mathbb{F}^{2 \times 2}$ over \mathbb{F} , cp. [24, p. 19]. The standard involution on \mathbb{H} maps $a = \begin{pmatrix} a_{00} & a_{01} \\ a_{10} & a_{11} \end{pmatrix}$ to $\bar{a} = \begin{pmatrix} a_{11} & -a_{01} \\ a_{01} & a_{00} \end{pmatrix}$. The determinant is a multiplicative quadratic form N on \mathbb{H} , and $\det(a)\mathbf{1} = a\bar{a} = \bar{a}a$, where $\mathbf{1}$ is the identity matrix in \mathbb{H} . The trace of a is obtained as $T(a)\mathbf{1} = a + \bar{a}$, the polar form of N is $(a|b) := N(a+b) - N(a) - N(b) = \bar{a}b + \bar{b}a = T(\mathbf{1}, \bar{a}b)\mathbf{1}$.

We form the direct sum of \mathbb{H} with a copy $\mathbb{H}w$. For $a, b, x, y \in \mathbb{H}$, the product is given by $(a+xw)(b+yw) := ab + \bar{y}x + (ya + x\bar{b})w$, the norm is extended by $N(a+xw) := N(a) - N(x)$, and $\overline{a+xw} := \bar{a} - xw$ extends the standard involution.

We identify \mathbb{F} with $\mathbb{F}\mathbf{1}$ from now on. Then $N(a+xw) = \overline{(a+xw)}(a+xw)$, and $w^2 = 1$. Note that $\mathbb{H}w = \mathbb{H}^\perp$, and that the standard involution is an anti-automorphism: we have $\overline{uv} = \bar{v}\bar{u}$ for all $u, v \in \mathbb{O}$. As in any unary composition algebra, we have $(su|v) = (u|\bar{s}v)$ and $(us|v) = (u|v\bar{s})$, for all $s, u, v \in \mathbb{O}$ (see [24, 1.3.2]).

We write $\text{Aut}(\mathbb{O})$ for the group of all algebra automorphisms of \mathbb{O} . The norm form N is the unique multiplicative quadratic form on \mathbb{O} , so every element of $\text{Aut}(\mathbb{O})$ is an isometry of N .

For any subset $M \subseteq \mathbb{O}$, let $\langle M \rangle$ denote the (not necessarily unary) subalgebra generated by M . We suppress curly braces, writing $\langle x \rangle := \langle \{x\} \rangle$ and $\langle x, y \rangle := \langle \{x, y\} \rangle$ etc..

1.1 Proposition ([18, 4.4]). *If $a, b \in \mathbb{O} \setminus \mathbb{F}$ have the same norm and trace then there is $\alpha \in \text{Aut}(\mathbb{O})$ with $a^\alpha = b$. \square*

1.2 Definition. Let \mathcal{L} denote the set of all 2-dimensional subalgebras of \mathbb{O} with trivial multiplication. In particular, let $Q_0 := \langle p_0w, n_0w \rangle = \mathbb{F}p_0w + \mathbb{F}n_0w$, then $Q_0 \in \mathcal{L}$, and \mathcal{L} is the $\text{Aut}(\mathbb{O})$ -orbit of Q_0 (see [18, 5.9]). The elements of \mathcal{L} serve as the lines in Schellekens' description [23, p. 207] for the split Cayley hexagon; the points are the elements of $\mathcal{P} := \{\mathbb{F}n \mid n \in \mathbb{O}, n^2 = 0 \neq n\}$. A different description of that generalized hexagon is given in [17, Sect. 4]: points and lines are the subalgebras of dimension 5 and 6, respectively, in \mathbb{O} .

The following results may be found in [24, Section 2.1, 1.7.3].

1.3 Lemma. *Let H, H' be subalgebras of \mathbb{O} , and assume $H \cong \mathbb{H} \cong H'$.*

- (a) *For any $v \in H^\perp$ with $N(v) \neq 0$, we have $H^\perp = Hv = vH$, and $(a+xw)(b+yw) = ab - N(v)\bar{y}x + (ya + x\bar{b})v$ holds for all $a, b, x, y \in H$.*

(b) *There is an automorphism of \mathbb{O} that maps H onto H' .*

(c) *The stabilizer of H in $\text{Aut}(\mathbb{O})$ consists of all maps $\alpha_{s,t}^{H,v} : a + xv \mapsto s^{-1}as + (ts^{-1}xs)v$ with $s \in H^\times := \{z \in H \mid N(z) \neq 0\} = \text{GL}(2, \mathbb{F})$ and $t \in \{z \in H \mid N(z) = 1\} = \text{SL}(2, \mathbb{F})$; here $a, x \in H$. \square*

For each $x \in \mathbb{O}$, we have $x^2 - T(x)x + N(x) = 0$. So $\langle x \rangle$ is a subalgebra of dimension at most 2; we have $\langle x \rangle = Fx$ if $x \in F$ or if $N(x) = 0$, and $\langle x \rangle = F + Fx$ if $N(x) \neq 0$. From $\bar{x} \in F + Fx$ and $yx = \overline{\bar{x}y}$ we infer that $\langle 1, x, y \rangle = F + Fx + Fy + Fxy$.

1.4 Definitions. Let \mathcal{D} denote the set of all 2-dimensional subalgebras of \mathbb{O} such that the restriction $N|_D$ is anisotropic, and has non-trivial polarization. Thus the elements of \mathcal{D} are separable quadratic extension fields of \mathbb{F} . If \mathbb{F} is a field of finite order q then each element of \mathcal{D} is isomorphic to the field \mathbb{F}_{q^2} of order q^2 . Choose $\delta \in \mathbb{F}_q$ such that the polynomial $m_\delta(x) := x^2 - x - \delta$ is irreducible over \mathbb{F}_q ; then each $D \in \mathcal{D}$ contains two roots u_D, \bar{u}_D of m_δ . Note that $N(u_D) = -\delta$ and $T(u_D) = 1$, so $\bar{u}_D = 1 - u_D$ and $u_D^2 = u + \delta$. Explicitly, $u := \begin{pmatrix} 0 & 1 \\ \delta & 1 \end{pmatrix} \in \mathbb{H}$ is a root of m_δ , and $\langle u \rangle \in \mathcal{D}$.

By Artin's Theorem (see [24, Prop. 1.5.2] or [22, Thm. 3.1, p. 29]), the unary subalgebra generated by two elements is associative. For $c \in \mathbb{O}$, this implies that the restriction of the multiplication turns \mathbb{O} into a left module over $\langle c \rangle$.

1.5 Notation. For $D \in \mathcal{D}$ we obtain that \mathbb{O} is a vector space of dimension 4 over D . On that vector space, the map $g : \mathbb{O} \times \mathbb{O} \rightarrow D : (x, y) \mapsto (u - \bar{u})^{-1}((x|uy) - \bar{u}(x|y))$ is a non-degenerate hermitian form, cp. [3, 2.9] or [14]. Note also that $g(x, x) = N(x)$.

The restriction \hat{g} of g to D^\perp is non-degenerate, as well.

2 The Hermitian unital of order three, and its automorphisms

In the projective plane $\text{PG}(2, \mathbb{F}_9)$, we have the Hermitian unital: using homogeneous coordinates from \mathbb{F}_9^3 and the (essentially unique) non-degenerate hermitian form h on \mathbb{F}_9^3 , the points are those one-dimensional subspaces of \mathbb{F}_9^3 such that the restriction of the form is zero; the blocks of the unital are the secants—induced by subspaces of dimension 2 such that the restriction of the form is not degenerate (but isotropic, as \mathbb{F}_9 is finite).

One knows ([5, p. 104], [1, 2.1, 2.2, see also p. 29]) that this Hermitian unital has $3^3 + 1 = 28$ points, each block contains $3 + 1 = 4$ of these points, and there are $3^2 = 9$ blocks through each one of these points. In other words, that Hermitian unital is a unital of order 3.

There are many isomorphism classes of unitals of order 3 (see [2]). Among these, the class of the Hermitian unital is characterized by the fact that the automorphism group is very large. In

particular, among all finite unitals, the Hermitian unitals and the Ree unitals are the only ones that admit a group of automorphisms that is doubly transitive on the set of points, see [16]. (The proof of that result uses the classification of finite simple groups.) Among the unitals of order 3, an exhaustive computer search [19] has shown that the Hermitian and the Ree unital, respectively, of order 3 are the only ones that admit a point-transitive group of automorphisms.

In our proof of 6.4 below, we identify the Hermitian unital by the properties that it admits all translations and does not contain any O’Nan configuration, see [25].

Specializing a general result [21] about Hermitian unitals, we obtain that the automorphism group of the Hermitian unital of order 3 is the group $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ induced by the group $\text{GU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ of all semi-similitudes of the Hermitian form. As $\text{Aut}(\mathbb{F}_9)$ has order 2, the group $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ induced by the group $\text{GU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ of all \mathbb{F}_9 -linear similitudes has index 2 in $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$.

We note that every multiplier of a similitude in $\text{GU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ is a norm of the quadratic extension, and obtained as the multiplier of a scalar multiple of the identity. So $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3) = \text{PU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ is induced by the group $\text{U}(3, \mathbb{F}_9 | \mathbb{F}_3)$ of all isometries of the Hermitian form. The multiplicative group of \mathbb{F}_9 is cyclic of order 8, and contains no elements of order 3. Therefore, the homomorphism from $\text{U}(3, \mathbb{F}_9 | \mathbb{F}_3)$ onto the group $\text{PU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ induced on the projective plane $\text{PG}(2, \mathbb{F}_9)$ has a kernel of order 8, but its restriction to the special unitary group $\text{SU}(3, \mathbb{F}_9 | \mathbb{F}_3) := \text{U}(3, \mathbb{F}_9 | \mathbb{F}_3) \cap \text{SL}(3, \mathbb{F}_9)$ is injective, and $\text{SU}(3, \mathbb{F}_9 | \mathbb{F}_3) \cong \text{PSU}(3, \mathbb{F}_9 | \mathbb{F}_3) \cong \text{PU}(3, \mathbb{F}_9 | \mathbb{F}_3)$.

The order of $\text{PU}(3, \mathbb{F}_{q^2} | \mathbb{F}_q)$ is $(q^3 + 1)q^3(q^2 - 1)$, see [13, Corollary to Theorem 2.50, p. 63]. We may also specialize the general formula $|\text{SU}(n, \mathbb{F}_{q^2} | \mathbb{F}_q)| = q^{(n^2-n)/2} \prod_{k=2}^n (q^k - (-1)^k)$; see [26, p. 118] or [8, 11.29]. For $q = 3$, this gives $|\text{SU}(3, \mathbb{F}_9 | \mathbb{F}_3)| = 3^3(3^2 - 1)(3^3 + 1) = 3^3 \cdot 8 \cdot 28 = 2^5 \cdot 3^3 \cdot 7$. So $|\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)| = 2^6 \cdot 3^3 \cdot 7 = 12096$.

3 A linear space

In this section, we assume that $\mathbb{F} = \mathbb{F}_q$ is the finite field of order q , for arbitrary q . Then \mathcal{D} consists of all subalgebras of the octonion algebra \mathbb{O} over \mathbb{F}_q that are isomorphic to \mathbb{F}_{q^2} . For any two elements $D, E \in \mathcal{D}$, the subalgebra $\langle D \cup E \rangle$ is associative, and has dimension at most 4. Moreover, it is a vector space over D ; with $\dim_D \langle D \cup E \rangle > 1$. So the dimensions are $\dim_D \langle D \cup E \rangle = 2$ and $\dim \langle D \cup E \rangle = 4$.

There are only two possibilities for the isomorphism type of $\langle D \cup E \rangle$.

3.1 Lemma. *Let D, E be two elements in \mathcal{D} . Then either $\langle D \cup E \rangle$ is a quaternion algebra, or there exists an element $z \in D^\perp$ with $z^2 = 0 \neq z$ and $\langle D \cup E \rangle = D + Dz$.*

Proof. We already know that $X := \langle D \cup E \rangle$ has dimension 4. The quadratic extension $D | \mathbb{F}_q$ is separable, so the polarization of the restriction $N|_D$ of the norm is not degenerate. Consider

$Q := X^\perp \cap X$. For $x, y \in X$ and $z \in Q$ we have $(x|yz) = (\bar{y}x|z) = 0$, so $XQ \subseteq Q$. In particular, we have $DQ \subseteq Q$, and Q is a vector space over D . Note also that $Q \subseteq D^\perp \subseteq 1^\perp$ implies $\bar{z} = -z$ for each $z \in Q$.

If $Q = \{0\}$ then the polarization of the restriction $N|_X$ is not degenerate. So X is a composition algebra, and then a quaternion algebra.

If $Q \neq \{0\}$ then $\dim_D Q = 1$. For $z \in Q$ we find $N(z) = -z^2 \in Q \cap \mathbb{F}_q$, and $D^\perp \cap \mathbb{F}_q = \{0\}$ yields $z^2 = 0$. For $a \in D$, we then also find $(az)z = az^2 = 0$, and use $0 = z\bar{a} + a\bar{z} = z\bar{a} - az$ to obtain $z(az) = z^2\bar{a} = 0$. So $QQ = \{0\}$, and $X = D + Dz$. \square

3.2 Remarks. As we assume a finite ground field in the present section, the quaternion algebras in 3.1 are split ones. The arguments in the proof of 3.1 are valid over infinite ground fields, as well. However, the elements of \mathcal{D} may then belong to different isomorphism types, and there may be non-split quaternion algebras.

If $\langle D \cup E \rangle$ is not a quaternion algebra then Dz is a 2-dimensional subalgebra with trivial multiplication (so $Dz \in \mathcal{L}$, see 1.2), and contained in D^\perp .

3.3 Definitions. Let \mathcal{H} be the set consisting of all quaternion subalgebras in \mathbb{O} , and \mathcal{X} the set of all subalgebras of the form $D + Dz$, where $D \in \mathcal{D}$ and $z \in D^\perp$ with $z^2 = 0 \neq z$. For $D \in \mathcal{D}$ and $Z \in \mathcal{Y} \in \{\mathcal{X}, \mathcal{H}\}$, we put $\mathcal{D}_Z := \{D \in \mathcal{D} \mid D < Z\}$ and $\mathcal{Y}_D := \{Y \in \mathcal{Y} \mid D < Y\}$. We form the incidence structures $\mathcal{U} := (\mathcal{D}, \mathcal{X}, <)$ and $\mathcal{V} := (\mathcal{D}, \mathcal{X} \cup \mathcal{H}, <)$.

3.4 Proposition. *The group $\text{Aut}(\mathbb{O})$ acts by automorphisms both on \mathcal{U} and on \mathcal{V} .*

(a) *Each one of the sets \mathcal{D} , \mathcal{X} , and \mathcal{H} is an orbit under $\text{Aut}(\mathbb{O})$, of length*

$$\begin{aligned} |\mathcal{D}| &= \frac{1}{2} q^3 (q^2 + q + 1)(q - 1), \\ |\mathcal{X}| &= \frac{1}{2} q (q^2 + q + 1)(q - 1)(q + 1)(q^2 - q + 1), \\ |\mathcal{H}| &= q^4 (q^2 + q + 1)(q^2 - q + 1), \text{ respectively.} \end{aligned}$$

(b) *For $D \in \mathcal{D}$, $X \in \mathcal{X}$, and $H \in \mathcal{H}$, we have*

$$|\mathcal{X}_D| = q^3 + 1, \quad |\mathcal{H}_D| = q^2(q^2 - q + 1), \quad |\mathcal{D}_X| = q^2, \quad |\mathcal{D}_H| = \frac{1}{2}q(q - 1).$$

Proof. Clearly, the group $\text{Aut}(\mathbb{O})$ acts on each one of the sets \mathcal{D} , \mathcal{X} , \mathcal{H} , and preserves inclusion. It is known (see [18]) that these sets are orbits under $\text{Aut}(\mathbb{O})$.

In order to determine $|\mathcal{D}|$ we fix $\delta \in \mathbb{F}$ such that m_δ is irreducible (see 1.4), and note that each $D \in \mathcal{D}$ contains exactly two elements with norm $-\delta$ and trace 1. If $a + xw$ is such an element then $T(a) = 1$ and $N(x) = N(a) + \delta$. If $N(a) = -\delta$ then a is a conjugate of $u = \begin{pmatrix} 0 & 1 \\ \delta & 1 \end{pmatrix}$ in \mathbb{H} , and $N(x) = 0$. We have $q^2 - q$ possible choices for a because the centralizer of u in \mathbb{H} is $C = \langle u \rangle$, and there are $q^3 + q^2 - q$ many possibilities for x in this case. If $N(a) \neq -\delta$ then $N(x) \neq 0$. There remain $q^3 - (q^2 - q)$ possible choices for a , and then $(q^2 - 1)q$ many choices for x with

$N(x) = N(a) + \delta$. In total, this gives $(q^2 - q)(q^3 + q^2 - q) + (q^3 - q^2 + q)(q^3 - q) = q^6 - q^3$, and $|\mathcal{D}| = \frac{1}{2}q^3(q^3 - 1)$, as claimed.

We determine $|\mathcal{D}_H|$ and $|\mathcal{H}_D|$ next. Without loss, we may assume $H = \mathbb{H}$ and $D = C$. The elements of $\mathcal{D}_{\mathbb{H}}$ form a single conjugacy class under $\mathbb{H}^\times = \mathrm{GL}(2, \mathbb{F}_q)$. The normalizer of C in $\mathrm{GL}(2, \mathbb{F}_q)$ induces $\mathrm{Gal}(C|F)$ on C ; the kernel of the action is C^\times . So the orbit $\mathcal{D}_{\mathbb{H}}$ has length $\frac{1}{2}(q-1)q$.

In order to find $|\mathcal{H}_C|$, we note first that $\mathbb{H} = C + Cj$, where $j := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in C^\perp \cap \mathbb{H}$ has norm -1 . So $C^\perp = Cj + \mathbb{H}w$. We determine the set A of all elements of norm -1 in C^\perp . Any such element is of the form $aj + xw$ with $a \in C$, $x \in \mathbb{H}$, and such that $-1 = N(aj + xw) = -N(a) - N(x)$. If $N(a) = 0$ then $a = 0$, and $N(x) = 1$ yields $x \in \mathrm{SL}(2, \mathbb{F}_q)$; there are $(q+1)q(q-1)$ many possible choices. If $N(a) = 1$ then $a \in C \cap \mathrm{SL}(2, \mathbb{F}_q) \cong C_{q+1}$, and $N(x) = 0$. This gives $(q+1)(q^3 + q^2 - q)$ many choices. Finally, if $N(a) \notin \{0, 1\}$ then $N(x) = 1 - N(a) \notin \{1, 0\}$. There are $(q+1)(q-2)$ choices for a , and $(q^2 - 1)q$ many choices for x . Summing up, we obtain that A has $(q+1)q^2(q^2 - q + 1)$ elements in total. For a given $i \in A$, each element of $Ci \cap A$ gives the same element $C + Ci \in \mathcal{H}_C$. So $|\mathcal{H}_C| = q^2(q^2 - q + 1)$, as claimed. The value for $|\mathcal{H}|$ now follows from the equation $|\mathcal{H}| \cdot |\mathcal{D}_H| = |\mathcal{D}| \cdot |\mathcal{H}_D|$ which is obtained by counting the flags in $(\mathcal{D}, \mathcal{H}, <)$ in two ways.

In order to find $|\mathcal{D}_X|$, we may assume $X = C + Cz$, with $C = \langle u \rangle$ as in 1.4, and $z \in C^\perp$ with $z^2 = 0 \neq z$. We search for elements $a + bz$ with $a, b \in C$ such that $1 = T(u) = T(a + bz) = T(a)$ and $-\delta = N(u) = N(a + bz) = N(a)$. Then $a \in \{u, \bar{u}\}$, while $b \in C$ is arbitrary. This gives $2q^2$ generators for elements of \mathcal{D}_X . Each element of \mathcal{D}_X contains exactly two of those generators, and $|\mathcal{D}_X| = q^2$ is proved.

The number $|\mathcal{X}_C|$ is obtained by counting the elements of $\{z \in C^\perp \mid z^2 = 0 \neq z\}$. If $aj + xw$ is such an element (with $a \in C$ and $x \in \mathbb{H}$) then $0 = N(aj + xw) = -N(a) - N(x)$ yields $N(x) = -N(a)$. For $a = 0$ we obtain $N(x) = 0 \neq x$; there are $q^3 + q^2 - q - 1 = (q^2 - 1)(q + 1)$ possible choices. For $a \neq 0$ we have $N(a) \neq 0$, and $|\mathrm{SL}(2, \mathbb{F}_q)| = (q^2 - 1)q$ possible choices for x . This gives a total of $(q^2 - 1)(q + 1 + (q^2 - 1)q) = (q^2 - 1)(q^3 + 1)$ non-trivial nilpotent elements in C^\perp . In each $Y \in \mathcal{X}_C$, there are $|C^\times| = q^2 - 1$ such elements. So $|\mathcal{X}_C| = q^3 + 1$, as claimed. Finally, we obtain the number $|\mathcal{X}|$ from $|\mathcal{X}| \cdot |\mathcal{D}_X| = |\mathcal{D}| \cdot |\mathcal{X}_D|$. \square

3.5 Corollary. *The order of $\mathrm{Aut}(\mathbb{O})$ is $|\mathcal{H}| \cdot |\mathrm{Aut}(\mathbb{O})_{\mathbb{H}}| = q^6(q^6 - 1)(q^2 - 1)$.* \square

3.6 Remark. At least if \mathbb{F} is finite of odd order, we interpret \mathcal{V} in terms of the split Cayley hexagon and its embedding (see [23]) in the projective space \mathbb{P} consisting of all vector subspaces of 1^\perp , as follows. The point set of the hexagon is the quadric $\mathcal{P} := \{\mathbb{F}z \mid z^2 = 0 \neq z \in \mathbb{O}\}$. The set \mathcal{L} of lines consists of those lines in \mathbb{P} that are subalgebras with trivial multiplication.

Let $\mathbb{F}^\square := \{s^2 \mid s \in \mathbb{F}\}$. For $K \in \mathcal{D} \cup \mathcal{X} \cup \mathcal{H}$ let \check{K} denote the intersection of K with 1^\perp . The $\frac{1}{2}q^3(q^3 - 1)$ points \check{D} with $D \in \mathcal{D}$ are those points of \mathbb{P} with $N(\check{D}) \cap (-\mathbb{F}^\square) = \{0\}$;

outside \mathcal{P} , we also have $\frac{1}{2}q^3(q^3 + 1)$ points S with $N(S) = -\mathbb{F}^\square$. Each $X \in \mathcal{X}$ is of the form $X = D + L$ with $D \in \mathcal{D}$ and $L = X^\perp \cap X \in \mathcal{L}$. Conversely, for each $L \in \mathcal{L}$ and $D \in \mathcal{D}$ with $L \leq D^\perp$ we have $D + L \in \mathcal{X}$. So $\check{X} = \check{D} + L$ is the span of a line L of the hexagon with a point outside \mathcal{P} orthogonal to L . Finally, each quaternion subalgebra $H \in \mathcal{H}$ corresponds to a subspace \check{H} that intersects \mathcal{P} in a point-regulus of the hexagon (cp. [17, proof of 5.6]).

4 The stabilizer of a subfield

For $D \in \mathcal{D}$, we use the hermitian form \hat{g} on D^\perp , as introduced in 1.5.

4.1 Definitions. Consider a separable quadratic field extension $\mathbb{E}|\mathbb{F}$, and a non-degenerate hermitian form h on \mathbb{E}^3 , with respect to the generator of $\text{Gal}(\mathbb{E}|\mathbb{F})$ which we denote by $s \mapsto \bar{s}$. We define the graph Γ with vertex set $S := \{\mathbb{E}v \mid v \in \mathbb{E}^3, h(v, v) \neq 0\}$; two vertices $\mathbb{E}x$ and $\mathbb{E}y$ are adjacent if the restriction of h to $\mathbb{E}x + \mathbb{E}y$ is not degenerate.

4.2 Lemma. *The graph Γ is connected if \mathbb{F} has more than two elements.*

Proof. Consider vertices $\mathbb{E}x, \mathbb{E}y \in S$ that are not adjacent. We will show that there is a path of length 3 from $\mathbb{E}x$ to $\mathbb{E}y$, unless \mathbb{F} has only two elements.

Without loss of generality, we replace h by a suitable scalar multiple and obtain $h(x, x) = 1$. There exists $n \in \mathbb{E}x + \mathbb{E}y$ with $h(x, n) = 0 = h(y, n)$ and $n \neq 0$. So $\mathbb{E}x + \mathbb{E}y = \mathbb{E}x + \mathbb{E}n$, there are $\alpha, \beta \in \mathbb{E}$ such that $y = \alpha x + \beta n$, and $\beta \neq 0$. Replacing n by a suitable multiple we may assume $\beta = 1$. As h is not degenerate, there exists $z' \in x^\perp$ with $h(z', n) = 1$. Since the trace of $\mathbb{E}|\mathbb{F}$ is surjective, there exists $s \in \mathbb{E}$ with $s + \bar{s} = 1 - h(z', z')$. For $z := z' + sn$, we obtain $h(z, z) = h(z', z') + sh(n, z') + h(z', n)\bar{s} + h(sn, sn) = h(z', z') + s + \bar{s} = 1$ while still $h(x, z) = 0$ and $h(n, z) = 1$. With respect to the basis x, n, z , the form h is thus described by the Gram matrix $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$.

We obtain $h(y, y) = \alpha\bar{\alpha}$; recall that this is not zero because $\mathbb{E}y \in S$. For an arbitrary element $v' \in y^\perp \setminus \mathbb{E}n$, there exists $\gamma \in \mathbb{E}$ such that $\mathbb{E}v' = \mathbb{E}v$, with $v = (1, \gamma, -\bar{\alpha})$. We compute $h(v, v) = 1 - \bar{\alpha}\bar{\gamma} - \gamma\alpha + \bar{\alpha}\alpha$ and $h(v, z) = \gamma - \bar{\alpha}$. The restriction of h to $\mathbb{E}v + \mathbb{E}z$ is described by the Gram matrix $\begin{pmatrix} h(v, v) & \gamma - \bar{\alpha} \\ \bar{\gamma} - \alpha & 1 \end{pmatrix}$; it is degenerate only if the determinant

$$\Delta := h(v, v) - (\gamma - \bar{\alpha})(\bar{\gamma} - \alpha) = 1 - \bar{\alpha}\bar{\gamma} - \gamma\alpha + \bar{\alpha}\alpha - (\gamma - \bar{\alpha})(\bar{\gamma} - \alpha) = 1 - \gamma\bar{\gamma}$$

is zero. We want to choose γ in such a way that both Δ and $h(v, v)$ are non-zero: then $\mathbb{E}v$ is a vertex adjacent to $\mathbb{E}z$, and $(\mathbb{E}x, \mathbb{E}z, \mathbb{E}v, \mathbb{E}y)$ is a path of length 3 in the graph Γ , as required.

If $\bar{\alpha}\alpha \neq -1$ we just take $\gamma = 0$. If $\bar{\alpha}\alpha = -1$ then $h(v, v) = -(\bar{\alpha}\bar{\gamma} + \alpha\gamma)$. As multiplication by α is a bijection of \mathbb{E} and the trace is an \mathbb{F} -linear surjection from \mathbb{E} onto \mathbb{F} , the set of $\gamma \in \mathbb{E}$

with $\overline{\alpha\gamma} + \alpha\gamma \neq 0$ is the complement of an \mathbb{F} -affine line in \mathbb{E} , and not empty. If \mathbb{F} has more than two elements then there is more than one square in \mathbb{F}^\times , and we can adjust γ so that $\bar{\gamma}\gamma \neq 1$; then $\Delta \neq 0$. If $|\mathbb{F}| = 3$ then $\gamma := \alpha^{-1}$ yields $h(v, v) = -(\overline{\alpha\gamma} + \alpha\gamma) = -2 = 1 \neq 0$ and $\Delta = 1 - \bar{\gamma}\gamma = 2 \neq 0$. \square

4.3 Remarks. If \mathbb{F} is a finite field of order q then Γ is the complement of the confluence graph of the Hermitian unital of order q . The case $q = 2$ is a true exception: here the unital (of order 2) is isomorphic to the affine plane of order 3, the adjacency relation is parallelity of lines, and there are 4 connected components (viz., parallel classes).

4.4 Lemma. Consider $D \in \mathcal{D}$ and u_D as in 1.4. If $\beta \in \text{Aut}(\mathbb{O})_{u_D}$ leaves Di invariant for some $i \in D^\perp$ with $N(i) \neq 0$ then $\det_D(\beta) = 1$. In particular, if $\alpha \in \text{Aut}(\mathbb{O})_{u_D}$ leaves invariant a quaternion subalgebra containing u_D then $\det_D(\alpha) = 1$.

Proof. The set $H := D + Di$ forms a quaternion subalgebra (see 1.3). Choose $v \in H^\perp$ with norm $N(v) \neq 0$.

By our hypothesis, there exists $s \in D$ such that β maps i to si . We have $N(s) = 1$ because β preserves the norm. Hilbert's Theorem 90 (see [15, 4.31]) asserts that there exists $c \in D$ with $c^{-1}\bar{c} = s^{-1}$. From 1.3(c) we know that there is an automorphism $\alpha := \alpha_{c,s}^{H,v}$ mapping $a+bi+xv$ (with $a, b \in D$ and $x \in H$) to $c^{-1}(a+bi)c + (sc^{-1}xc)v = a + c^{-1}\bar{c}bi + (\bar{c}^{-1}xc)v = a + s^{-1}bi + (\bar{c}^{-1}xc)v$. In particular, the image si of i under β is mapped to i , and the product $\gamma := \beta\alpha$ fixes i . Moreover, we have $v^\alpha = sv$, and $(iv)^\alpha = iv$. With respect to the D -basis $1, i, v, iv$, the map α is thus described by a diagonal matrix, with determinant $\det_C(\alpha) = s^{-1}s = 1$.

Now γ fixes both 1 and i . Thus H^\perp is invariant under γ ; and there are elements $a, e \in D$ such that $v^\gamma = av + e(iv) = av + (ie)v$. Then $(iv)^\gamma = iv^\gamma = i(av) + i((ie)v) = (ai)v + (iei)v = (i\bar{a})v + (\bar{e}i^2)v = -\bar{e}N(i)v + \bar{a}(iv)$. As $\gamma \in \text{Aut}(\mathbb{O})$ preserves the norm, we have $N(v) = N(v^\gamma) = N(av + (ie)v) = (N(a) + N(ie))N(v)$, and $1 = N(a) + N(ie) = a\bar{a} + e\bar{e}N(i) = \det_D(\gamma)$ follows. This implies $\det_D(\beta) = 1$.

Finally, assume that $\alpha \in \text{Aut}(\mathbb{O})_u$ leaves invariant a quaternion algebra H containing D . Then $D^\perp \cap H$ is invariant, and spanned by an element of non-trivial norm, so the previous result applies. \square

4.5 Lemma. Consider $D \in \mathcal{D}$, and a root u_D of m_δ in D .

- (a) The stabilizer $\text{Aut}(\mathbb{O})_{u_D}$ of u_D is a subgroup of $\text{SU}(C^\perp, \hat{g})$.
- (b) If \mathbb{F} is finite of order q then $\text{Aut}(\mathbb{O})_{u_D} = \text{SU}(D^\perp, \hat{g}) \cong \text{SU}(3, \mathbb{F}_{q^2} | \mathbb{F}_q)$. In particular, $\text{Aut}(\mathbb{O})_{u_D}$ acts doubly transitively on the set $\mathcal{X}_D = \{Dx \mid x \in D^\perp \setminus \{0\}, N(x) = 0\}$.

Proof. The stabilizer $\text{Aut}(\mathbb{O})_D$ of the subalgebra D acts on D^\perp by semi-similitudes of \hat{g} , and induces a subgroup of $\Gamma\text{U}(C^\perp, \hat{g})$. That action is faithful because D^\perp generates the algebra \mathbb{O} . The stabilizer $\text{Aut}(\mathbb{O})_u$ induces similitudes of \tilde{g} . From $g(x, x) = \text{N}(x)$ we then obtain that the multipliers are trivial, and $\text{Aut}(\mathbb{O})_u$ induces a subgroup of the group $\text{U}(C^\perp, \hat{g})$ of isometries. We are going to show that $\text{Aut}(\mathbb{O})_u$ is contained in $\Sigma := \text{Aut}(\mathbb{O})_u \cap \text{SU}(D^\perp, \hat{g})$.

We consider first the case where $q = 2$. Without loss of generality, we may assume $D = C$ and $u_D = u$, see 1.3(b). We use $j = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in C^\perp \cap \mathbb{H}$ and $n = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \in \mathbb{H}$. For any other nilpotent element $aj + xw \in C^\perp = Cj + \mathbb{H}w$, we have $0 = \text{N}(aj + xw) = \text{N}(a) + \text{N}(x)$.

Consider $c \in C^\times$ and $t \in \mathbb{H}$ with $\text{N}(t) = 1$, i.e., $t \in \text{SL}(2, \mathbb{F}_2)$. Then $\alpha_{c,1}^{\mathbb{H},w}$ leaves \mathbb{H} invariant, and belongs to Σ by 4.4. The set $\Xi := \{\alpha_{c,t}^{\mathbb{H},w} \mid c \in C^\times, t \in \text{SL}(2, \mathbb{F}_2)\}$ forms a subgroup of order 18 in Σ . The stabilizer Ξ_{nw} is generated by $\alpha_{1,u_j}^{\mathbb{H},w}$, and has order 2. The orbit of nw under Ξ thus has length 9; it contains all ¹nilpotent elements of the form xw with $\text{N}(x) = 0 \neq x$. There remain the nilpotent elements of the form $aj + xw$ with $a \in C^\times$ and $x \in \text{SL}(2, \mathbb{F}_2)$. The stabilizer Ξ_{j+w} is trivial, so the orbit of $j + w$ under Ξ contains all the remaining non-trivial nilpotent elements of C^\perp .

Now consider the quaternion algebra $H := C + Cw$, we have $j \in H^\perp$ and $\mathbb{O} = H + Hj$. The automorphism $\alpha_{1,w}^{H,j} \in \Sigma$ maps $w + j$ from the Ξ -orbit of length 18 to $w + wj = (1 + j)w$ in the Ξ -orbit of length 9. So the orbits are fused into a single Σ -orbit, and it remains to study the stabilizer of nw in $\text{Aut}(\mathbb{O})_u$. If $\gamma \in \text{Aut}(\mathbb{O})_u$ fixes nw then it leaves $(C + Cnw)^\perp$ invariant. Pick $x \in (C + C(nw))^\perp \setminus C(nw)$. Then $\text{N}(x) \neq 0$, and the restriction of γ to $C(nw) + Cx$ is described by a triangular matrix with diagonal entries 1, s , for some $s \in C^\times$. If $s \neq 1$ then x may be chosen as an eigenvector for s , and 4.4 yields $\det_C(\gamma) = 1$. If $s = 1$ we pick $y \in C^\perp \setminus (nw)^\perp$, then $1, nw, x, y$ is a C -basis for \mathbb{O} . We write the image of y as $y^\gamma = ey + z$ with $e \in C$ and $z \in (nw)^\perp$. Now $0 \neq g(nw, y) = g((nw)^\gamma, y^\gamma) = g(nw, ey + z) = g(nw, y)e$. This yields $e = 1$ and then $\det_C(\gamma) = 1$.

For the case of a general finite field \mathbb{F} of order $q > 2$, we use the graph Γ introduced in 4.1. For vertices Dx and Dy , we may assume $\text{N}(x) = \text{N}(y)$ because the extension $D|\mathbb{F}$ has surjective norm form. If Dx and Dy are adjacent, we use a non-isotropic $z \in (D + Dx + Dy)^\perp$ to obtain a quaternion algebra $H := D + Dz$. Then $x, y \in H^\perp$, so $H^\perp = Hx$ and there exists $t \in H$ with $y = tx$. We have $\text{N}(x) = \text{N}(y) = \text{N}(tx)$, so $\text{N}(t) = 1$ and $\alpha_{1,t}^{H,x} \in \Sigma$ maps Dx to Dy . By 4.2, the graph Γ is connected, and we find that Σ is transitive on the set of vertices

¹ In fact, it contains xw for

$$x \in \left\{ \begin{array}{l} n = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} n = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} n = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \\ nu = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} nu = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} nu = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \\ n\bar{u} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} n\bar{u} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} n\bar{u} = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \end{array} \right\}.$$

of Γ . So $\text{Aut}(\mathbb{O})_{u_D}$ is the product of Σ and the stabilizer of Dx in $\text{Aut}(\mathbb{O})_{u_D}$. That stabilizer is contained in Σ by 4.4, and we have proved that $\text{Aut}(\mathbb{O})_{u_D}$ coincides with Σ .

Specializing the general formula $|\text{SU}(n, \mathbb{F}_{q^2} | \mathbb{F}_q)| = q^{(n^2-n)/2} \prod_{k=2}^n (q^k - (-1)^k)$ from [26, p. 118] or [8, 11.29], we obtain $|\text{SU}(C^\perp, \hat{g})| = |\text{SU}(3, \mathbb{F}_{q^2} | \mathbb{F}_q)| = q^3(q^3 + 1)(q^2 - 1)$.

The order of $\text{Aut}(\mathbb{O}) = \text{G}_2(q)$ is $|\text{G}_2(q)| = q^6(q^6 - 1)(q^2 - 1)$, see 3.5. From 3.4 we know $|\mathcal{D}| = \frac{1}{2}q^3(q^3 - 1)$. Therefore, we obtain $|\text{Aut}(\mathbb{O})_D| = 2q^3(q^3 + 1)(q^2 - 1)$, and $|\text{Aut}(\mathbb{O})_{u_D}| = q^3(q^3 + 1)(q^2 - 1) = |\text{SU}(3, \mathbb{F}_{q^2} | \mathbb{F}_q)| = |\text{SU}(C^\perp, \hat{g})|$, and $\text{Aut}(\mathbb{O})_{u_D} = \text{SU}(C^\perp, \hat{g})$, as claimed.

The action of $\text{Aut}(\mathbb{O})_{u_D}$ on $\{Dx \mid x \in D^\perp \setminus \{0\}, N(x) = 0\}$ is equivalent to the action of $\text{SU}(3, \mathbb{F}_{q^2} | \mathbb{F}_q)$ on the set of points of the Hermitian unital of order q . That action is doubly transitive ([21, 1.4], see [13, Thm. 2.50]). \square

5 A unital of unitals

In the present section, we concentrate on the case where $q := |\mathbb{F}| = 2$. Thus \mathcal{D} is the set of all subalgebras of \mathbb{O} that are isomorphic to the field \mathbb{F}_4 with 4 elements. As in 1.4, we consider $u = \begin{pmatrix} 0 & 1 \\ -\delta & 1 \end{pmatrix}$, necessarily with $\delta = 1$.

From 3.4 we know that $\mathcal{U} = (\mathcal{D}, \mathcal{X}, <)$ is a linear space with parameters $|\mathcal{D}| = 28$ and $|\mathcal{D}_X| = 4$ for each $X \in \mathcal{X}$. So \mathcal{U} is a unital of order 3. (The elements of \mathcal{H} are omitted because $|\mathcal{D}_H| = 1$ if $|\mathbb{F}| = 2$.)

We will show that \mathcal{U} admits all translations, and that \mathcal{U} contains no O’Nan configurations. Then [25] yields that \mathcal{U} is in fact isomorphic to the Hermitian unital of order 3.

5.1 Notation. In $\mathbb{H} = \mathbb{F}_2^{2 \times 2}$, we have $p_0 := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $n_0 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $m_0 := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Note that $\overline{p_0} = 1 - p_0$, while $n_0 = \overline{n_0}$ and $m_0 = \overline{m_0}$ belong to $1^\perp \cap \mathbb{H} = \{a \in \mathbb{H} \mid \text{T}(a) = 0\}$.

Clearly, the group $\text{Aut}(\mathbb{O})$ acts by automorphisms of the incidence structure \mathcal{U} .

5.2 Lemma. Consider $D \in \mathcal{D}$.

- (a) For $c, d \in D$ and $x \in D^\perp$, define $\tau_c: D + D^\perp = \mathbb{O} \rightarrow \mathbb{O}$ by $\tau_c(d + x) = d + cx$. Then $T_{[D]} := \{\tau_c \mid c \in D \setminus \{0\}\}$ is a subgroup of $\text{Aut}(\mathbb{O})_D$, and acts trivially on the set of all D -subspaces in D^\perp .
- (b) For each $X \in \mathcal{X}_D$, the stabilizer $\text{Aut}(\mathbb{O})_X$ is doubly transitive on \mathcal{D}_X .
- (c) The stabilizer $\text{Aut}(\mathbb{O})_D$ acts doubly transitively on \mathcal{X}_D .
- (d) The group $\text{Aut}(\mathbb{O})$ is doubly transitive on \mathcal{D} .

Proof. We note that $D \cong \mathbb{F}_4$ (by our standing assumption), so $\bar{c} = c^2$ holds for each $c \in D$. Since the orthogonal space D^\perp is not totally singular, we find $j \in D^\perp$ with $N(j) \neq 0$ (so actually $N(j) = 1$). Then $H := D + Dj$ is a quaternion algebra, and we have $jc = \bar{c}j$ for each $c \in D$. Pick $v \in H^\perp$ with $N(v) = 1$, so $\mathbb{O} = H + Hv$.

Consider $c \in D \setminus \{0\}$; then $N(c) = 1$, and $\alpha_{c,c}^{H,v}$ is an automorphism of \mathbb{O} (see 1.3(c)). For $d \in D$, that automorphism maps d to $c^{-1}dc = d$, and maps dj to $c^{-1}dj c = c^{-1}d\bar{c}j = c^{-1}c^2dj = c(dj)$. For $y \in H$, it maps $z = yv$ to $(cc^{-1}yc)v = c(yv) = cz$, and maps $uz = u(yv) = (yu)v$ to $(cc^{-1}yuc)v = (yuc)v = c((yu)z) = c(uz)$. So $\alpha_{c,c}^{H,v} = \tau_c$, and assertion (a) follows. As any two points in \mathcal{D} are joined by an element of \mathcal{X} , we also obtain that $\text{Aut}(\mathbb{O})$ is transitive on \mathcal{D} .

For $X \in \mathcal{X}_D$, the group $T_{[D]}$ is transitive on $\mathcal{D}_X \setminus \{D\}$. For E in that set, the group $T_{[E]}$ is transitive on $\mathcal{D}_X \setminus \{E\}$, and assertion (b) follows.

Assertion (c) has been noted in 4.5. Together with assertion (b), it yields that the stabilizer $\text{Aut}(\mathbb{O})_D$ is transitive on $\mathcal{D} \setminus \{D\}$, and assertion (d) follows from transitivity of $\text{Aut}(\mathbb{O})$ on \mathcal{D} (see 3.4). \square

5.3 Theorem. *The action of $\text{Aut}(\mathbb{O})$ on \mathcal{U} is faithful.*

Proof. It is known (e.g., see [18, 5.9]) that the set \mathcal{L} forms a single orbit under $\text{Aut}(\mathbb{O})$. We have $X = C + C(n_0w) \in \mathcal{X}$, and $X^\perp \cap X = C(n_0w) \in \mathcal{L}$. Also, we have $L := \langle n_0w, n_0 \rangle \in \mathcal{L}$, so there exists $Y \in \mathcal{X}$ with $L = Y^\perp \cap Y$.

If $\alpha \in \text{Aut}(\mathbb{O})$ acts trivially on \mathcal{U} then α fixes X, Y , and $(X^\perp \cap X) \cap (Y^\perp \cap Y) = Fn_0w$. The latter space consists of 0 and n_0w , so α fixes n_0w . As $\text{Aut}(\mathbb{O})$ is transitive on the set of non-central elements of given norm and trace in \mathbb{O} (see 1.1), that argument shows that every nilpotent element is fixed by α . Every element of norm 0 and trace 1 is in the orbit of $p_0 = n_0m_0$, every element of norm 1 and trace 1 is in the orbit of $u = n_0 + m_0 + m_0n_0$, and every non-central element of norm 1 and trace 0 is in the orbit of $n_0 + m_0$. So \mathbb{O} is generated by its nilpotent elements, and α is trivial on \mathbb{O} . This shows that the action of $\text{Aut}(\mathbb{O})$ on \mathcal{U} is faithful. \square

5.4 Remark. For each $D \in \mathcal{D}$, the set \mathcal{X}_D consists of the 9 points of the Hermitian unital $(U_C, \mathcal{S}_C, \epsilon)$ of order 2, see [17, 5.12], cp. also 7.4 below. This justifies the title of the present section: the blocks of the incidence structure $\mathcal{U} = (\mathcal{D}, \mathcal{X}, <)$ are unitals of order 2 in the split Cayley hexagon (and $\mathcal{U} = (\mathcal{D}, \mathcal{X}, <)$ is itself a unital, of order 3).

6 Identification of the unital

6.1 Lemma. *Let D, E be two elements of \mathcal{D} , and pick elements $d = u_D \in D, e = u_E \in E$ of order 3, respectively. Then the set $D^\times E^\times = \{1, d, d^2, e, e^2, de, d^2e, de^2, d^2e^2\}$ contains two involutions, and every other element of $(D^\times E^\times) \setminus \{1\}$ has order 3. In particular, we have either*

$$\mathcal{D}_{\langle D \cup E \rangle} = \{\langle d \rangle, \langle e \rangle, \langle de \rangle, \langle d^2e^2 \rangle\}, \quad \text{or} \quad \mathcal{D}_{\langle D \cup E \rangle} = \{\langle d \rangle, \langle e \rangle, \langle d^2e \rangle, \langle de^2 \rangle\}.$$

Proof. By 5.2(d), we may assume $D = C = \langle u \rangle$ and $E = \langle u + z \rangle$, with some nilpotent element $z \in C^\perp$. Note that $z \in C^\perp$ implies $zu = \bar{u}z = u^2z$. We choose $d := u \in C$ and $e := u + z \in E$ (the other possible choices u^2 or $(u + z)^2 = u^2 + z$ amount to permutations of $D^\times E^\times$ that do not affect the result).

Now $de = u^2 + uz$, $d^2e = 1 + u^2z$, $de^2 = 1 + uz$, $d^2e^2 = u + u^2z$. Elements of $1 + C^\times z$ have trace 0 and norm 1, and thus are involutions. The elements of $\{u, u^2\} + Cz$ have trace 1 and norm 1. So they have order 3. The points in the joining block are $C = \langle d \rangle = \langle u \rangle$, $\langle e \rangle = \langle u + z \rangle$, $\langle de \rangle = \langle u^2 + uz \rangle$, and $\langle d^2e^2 \rangle = \langle u + u^2z \rangle$, cp. 5.2(a). \square

6.2 Example. We use $p_0 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $n_0 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $m_0 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ as introduced in 5.1 but abbreviate $x := x_0$ throughout. We have $C = \langle u \rangle \in \mathcal{D}$. For $\alpha, \beta \in \{0, 1, 2\}$, the subalgebras $D_\alpha := \langle u + (nu^\alpha)w \rangle$ and $E_\beta := \langle u + (mu^\beta)w \rangle$ also belong to \mathcal{D} . The subalgebras $X := C + C((nu)w) = \langle C \cup D_0 \rangle$ and $Y := C + C((mu)w) = \langle C \cup E_0 \rangle$ belong to \mathcal{X} ; we have $\mathcal{D}_X = \{C, D_0, D_1, D_2\}$ and $\mathcal{D}_Y = \{C, E_0, E_1, E_2\}$ by 5.2. The sets $\{D_0, D_1, D_2\}$ and $\{E_0, E_1, E_2\}$, respectively, are orbits under the group $T_{[C]}$ introduced in 5.2(a).

We compute the square of $(u + (nu^\alpha)w)$ as $u^2 + u((nu^\alpha)w) + ((nu^\alpha)w)u + ((nu^\alpha)w)^2 = u^2 + (nu^\alpha(u + \bar{u})w) = u^2 + (nu^\alpha)w$.

Let $Z_{\alpha,\beta} := \langle D_\alpha \cup E_\beta \rangle$ be the block joining D_α and E_β . In order to find the two points in $Z_{\alpha,\beta} \setminus \{D_\alpha, E_\beta\}$, we follow 6.1, and form the products

$$\begin{aligned} z_{\alpha,\beta}^{\varepsilon,\varphi} &:= (u + (nu^\alpha)w)^\varepsilon (u + (mu^\beta)w)^\varphi &= u^\varepsilon u^\varphi + \overline{mu^\beta} nu^\alpha + (mu^\beta u^\varepsilon + nu^\alpha \overline{u^\varphi})w \\ & &= u^{\varepsilon+\varphi} + u^{2\beta} m n u^\alpha + (mu^{\beta+\varepsilon} + nu^{\alpha+2\varphi})w \end{aligned}$$

for $\varepsilon, \varphi \in \{1, 2\}$ and $\alpha, \beta \in \{0, 1, 2\}$. For given (α, β) , the two points in question are generated by elements $z_{\alpha,\beta}^{\varepsilon,\varphi}$ of trace 1. The trace of $u^{2\beta} m n u^\alpha = u^{2\beta} \bar{p} u^\alpha$ equals 0 if $(\alpha, \beta) \in \{(0, 2), (1, 0), (2, 1)\}$. In all other cases, that trace equals 1.

If $u^{2\beta} \bar{p} u^\alpha$ has trace 0 we choose $\varepsilon = \varphi \in \{1, 2\}$; then $\mathrm{T}(z_{\alpha,\beta}^{\varepsilon,\varphi}) = \mathrm{T}(u^{2\varepsilon} + u^{2\beta} \bar{p} u^\alpha) = 1$; the two points are $\langle z_{\alpha,\beta}^{1,1} \rangle$ and $\langle z_{\alpha,\beta}^{2,2} \rangle$.

If $u^{2\beta} \bar{p} u^\alpha$ has trace 1 we choose $\varepsilon, \varphi \in \{1, 2\}$ such that $\varepsilon + \varphi = 3$, then $\overline{u^\varphi} = \bar{u}^{2\varepsilon} = u^\varepsilon$, and $\mathrm{T}(z_{\alpha,\beta}^{\varepsilon,\varphi}) = 1$, as well. In these cases, the two points are $\langle z_{\alpha,\beta}^{1,2} \rangle$ and $\langle z_{\alpha,\beta}^{2,1} \rangle$.

6.3 Proposition. *There are no O’Nan configurations in \mathcal{U} .*

Proof. Aiming at a contradiction, we assume that there exists an O’Nan configuration in \mathcal{U} . We use the elements introduced in 5.1 but abbreviate $g := g_0$ throughout, for $g \in \{m, n, p\}$.

By 5.2(c), we may assume that $C = \langle u \rangle$ as in 1.4 is a point of the configuration, and that the blocks $X := \langle u, u + nw \rangle$ and $Y := \langle u, u + mw \rangle$ belong to the configuration. Then we are in the situation studied in 6.2.




Figure 1: An O'Nan configuration, hypothesized.

The configuration has further points D_α, D_γ in X , and E_β, E_δ in Y such that the sixth point is $J := \langle D_\alpha \cup E_\beta \rangle \cap \langle D_\gamma \cup E_\delta \rangle$. (See Figure 1.)

Applying an element of $T_{[C]}$ (see 5.2(a)), we may assume $\gamma = 0$. From 6.2 we then know $J = \langle z_{\alpha,\beta}^{\varepsilon,\varphi} \rangle = \langle z_{0,\delta}^{\varepsilon',\varphi'} \rangle$. So we are searching for $\alpha \in \{1, 2\}$ and $\beta \in \{0, 1, 2\} \setminus \{\delta\}$ such that $z_{\alpha,\beta}^{\varepsilon,\varphi}$ equals either $z_{0,\delta}^{\varepsilon',\varphi'}$ or its square $1 + z_{0,\delta}^{\varepsilon',\varphi'}$, with suitable $\varepsilon, \varphi, \varepsilon', \varphi'$, see 6.2.

Using the direct sum decomposition $\mathbb{O} = \mathbb{H} + \mathbb{H}w$, we write $z_{0,\delta}^{\varepsilon',\varphi'} = a + xw$ and $z_{\alpha,\beta}^{\varepsilon,\varphi} = b + yw$ with $a = u^{\varepsilon'+\varphi'} + u^{2\delta}\bar{p}$, $x = mu^{\delta+\varepsilon'} + nu^{2\varphi'}$, $b = u^{\varepsilon+\varphi} + u^{2\beta}\bar{p}u^\alpha$, $y = mu^{\beta+\varepsilon} + nu^{\alpha+2\varphi}$ in \mathbb{H} . We compute a , and find suitable (α, β) with $b \in \{a, a^2\}$; recall that $a^2 = \bar{a} = a + 1$. Then we check that $x \neq y$ holds for all such suitable (α, β) .

We find a, x as in the following table:

$(0, \delta)$	$(0, 0)$		$(0, 1)$		$(0, 2)$	
(ε', φ')	$(1, 2)$	$(2, 1)$	$(1, 2)$	$(2, 1)$	$(1, 1)$	$(2, 2)$
a	p		$p + n$		$m + \bar{p}$	$p + m$
x	$1 + m$	$1 + n$	1	u	u	$1 + m$

For α, β with $\alpha \neq 0$ we determine $b = u^{\varepsilon+\varphi} + u^{2\beta}\bar{p}u^\alpha$ and $y = mu^{\beta+\varepsilon} + nu^{\alpha+2\varphi}$:

(α, β)	$(1, 0)$		$(1, 1)$		$(1, 2)$		$(2, 0)$		$(2, 1)$		$(2, 2)$	
(ε, φ)	$(1, 1)$	$(2, 2)$	$(1, 2)$	$(2, 1)$	$(1, 2)$	$(2, 1)$	$(1, 2)$	$(2, 1)$	$(1, 1)$	$(2, 2)$	$(1, 2)$	$(2, 1)$
b	$n + \bar{p}$	$p + n$	$m + \bar{p}$		\bar{p}		$p + m$		p	\bar{p}	$n + \bar{p}$	
y	\bar{u}	$1 + n$	$1 + n$	$n + m$	u	\bar{u}	\bar{u}	1	1	$n + m$	$n + m$	$1 + m$

Thus we obtain that no point J exists as required, and there are no O'Nan configurations in \mathcal{U} , as claimed. \square

6.4 Theorem. *The unital $\mathcal{U} = (\mathcal{D}, \mathcal{X}, <)$ is isomorphic to the Hermitian unital of order 3.*

Proof. For $D \in \mathcal{D}$, we know from 5.2(a) that there exists a subgroup $T_{[D]} \leq \text{Aut}(\mathbb{O})$ fixing D and each D -subspace of D^\perp , and acting transitively on $\mathcal{D}_X \setminus \{D\}$, for each $X \in \mathcal{X}_D$. So $T_{[D]}$ induces a group of translations with center D on the unital \mathcal{U} (in the sense of [10], [11], [12]), and \mathcal{U} is a unital admitting all translations. In 6.3 we have shown that \mathcal{U} does not contain any O’Nan configuration. Now [25] says that \mathcal{U} is a Hermitian unital. \square

7 An isomorphism of groups

As a corollary to 6.4 and 5.3, we obtain a known result from group theory ([7], cp. [27, 4.4.4]):

7.1 Theorem. *The automorphism group $G_2(2) = \text{Aut}(\mathbb{O})$ of the octonion algebra \mathbb{O} over \mathbb{F}_2 is isomorphic to the group $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$. In fact, the action of $\text{Aut}(\mathbb{O})$ on the set \mathcal{D} of subalgebras isomorphic to \mathbb{F}_4 is equivalent to the doubly transitive action of $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ on the set of points of the hermitian unital of order 3.*

Proof. From 5.3 we know that $\text{Aut}(\mathbb{O})$ acts faithfully on \mathcal{U} ; this gives an injective homomorphism from $\text{Aut}(\mathbb{O})$ to $\text{Aut}(\mathcal{U}) \cong \text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$. It remains to show that the orders of the two groups are the same. For any prime power q , one has $|G_2(q)| = q^6(q^6 - 1)(q^2 - 1)$, see 3.5. So $|G_2(2)| = 2^6 \cdot 63 \cdot 3 = 2^6 \cdot 3^3 \cdot 7 = 12\,096$. This coincides with the order of $|\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)|$, see Section 2. \square

Note that the group $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ is not simple; the subgroup $\text{PU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ has index 2, and is normal. Thus our explicit isomorphism from $\text{Aut}(\mathbb{O})$ onto $\text{PGU}(3, \mathbb{F}_9 | \mathbb{F}_3)$ corroborates the known fact that $\text{Aut}(\mathbb{O}) \cong G_2(2)$ is not simple in the case $q = 2$.

7.2 Remarks. Our arguments in 6.3 could be replaced by an application of computer-based results. From 5.3 we know that $\text{Aut}(\mathbb{O})$ acts faithfully on \mathcal{U} ; this action is doubly transitive on the set \mathcal{D} of points (see 5.2).

An exhaustive computer search [19] for unitals of order 3 with non-trivial automorphism groups has produced (among 4466 isomorphism types in total) only two unitals with point-transitive groups, namely, the Hermitian unital and the Ree unital (see [19, p. 266]). The full automorphism group of the Ree unital of order 3 is the group of twisted Lie type ${}^2G_2(3)$, and has order $2^3 \cdot 3^3 \cdot 7 = 1\,512$. (See [20], [9] for the automorphism group of a Ree unital of arbitrary finite order.) Thus we obtain that \mathcal{U} is isomorphic to the Hermitian unital.

Much earlier than [19], a computer search by Brouwer (see [4, Proposition, p. 1]) has also characterized the Hermitian unital of order 3 by the fact that it does not contain O’Nan configurations. Brouwer [4] reports on the results of a computer search for unitals of order 3 that

admit an automorphism of order 7; he found 8 isomorphism types. According to [19], the number of isomorphism types actually is 11.

We also remark that there are deep (and much more general) results that allow the recognition of the unital. From [16] we infer that the Hermitian unitals and the Ree unitals are the only unitals with doubly transitive automorphism group; as above, the Ree unital is excluded by the size of its full automorphism group. The more recent result in [10] characterizes Hermitian unitals by the fact that they allow all translations. Both results use the classification of finite simple groups.

Together with our observation 4.5, the isomorphism 7.1 immediately yields the following.

7.3 Corollary. *The stabilizer $\text{Aut}(\mathbb{O})_C$ of the subalgebra $C = \mathbb{F}_2 + \mathbb{F}_2u \cong \mathbb{F}_4$ is isomorphic to the stabilizer of an isotropic point in $\text{PGU}(3, \mathbb{F}_9|\mathbb{F}_3)$, and that stabilizer is isomorphic to $\Gamma\text{U}(3, \mathbb{F}_4|\mathbb{F}_2) \cong \text{AGL}(2, \mathbb{F}_3)$. \square*

The (unique) unital of order 2 (which is isomorphic to the affine plane of order 3) can be constructed in the Hermitian unital \mathcal{U} of order 3, as follows:

7.4 Proposition. *Pick any point $D \in \mathcal{D}$, and let \mathcal{B} consist of all those 3-element subsets of \mathcal{X}_D such that no element of $\mathcal{X} \setminus \mathcal{X}_D$ meets each member of that subset. Then $(\mathcal{X}_D, \mathcal{B}, \in)$ is isomorphic to the affine plane of order 3.*

Proof. We follow O’Nan [21, Section 3] on his way to construct the circle geometry (i.e., the inversive plane) of order 3: The point set of that inversive plane is obtained by adding a new element ∞ to \mathcal{X}_C . We describe the set \mathcal{C} of circles. The set $\mathcal{C} \setminus \mathcal{C}_\infty$ of circles not through ∞ consists of those sets of 4 blocks in \mathcal{X}_D that meet a given block not through C .

The remaining circles are obtained by adding ∞ to each 3-element subset of \mathcal{X}_D that is not contained in any element of $\mathcal{C} \setminus \mathcal{C}_\infty$ (as constructed before). So $\mathcal{C}_\infty = \mathcal{B}$; those 3-element subsets consist of three blocks through C such that no block in $\mathcal{X} \setminus \mathcal{X}_D$ meets all three of them.

The discussion in [21, Sections 3,4] shows that $(\mathcal{X}_D \cup \{\infty\}, \mathcal{C})$ is isomorphic to the inversive plane of order 3. Then $\mathcal{P} = (\mathcal{X}_D, \mathcal{C}_\infty)$ is the affine plane of order 3. \square

For O’Nan, it was a major task to show that the set \mathcal{C}_∞ is invariant under all automorphisms of the Hermitian unital. In the small case at hand, this is obvious because we can describe the elements of \mathcal{C}_∞ quite easily.

For general q , the Hermitian unital of order q can be found in terms of lines in the split Cayley hexagon of order q , see [17].

References

- [1] S. G. Barwick and G. Ebert, *Unitals in projective planes*, Springer Monographs in Mathematics, Springer, New York, 2008, doi:10.1007/978-0-387-76366-8. MR 2440325. Zbl 1156.51006.
- [2] A. Betten, D. Betten, and V. D. Tonchev, *Unitals and codes*, Discrete Math. **267** (2003), no. 1-3, 23–33, doi:10.1016/S0012-365X(02)00600-3. MR 1991559. Zbl 71024.05011.
- [3] A. Blunck, N. Knarr, B. Stroppel, and M. J. Stroppel, *Transitive groups of similitudes generated by octonions*, J. Group Theory **21** (2018), no. 6, 1001–1050, doi:10.1515/jgth-2018-0018. MR 3871471. Zbl 1439.20059.
- [4] A. E. Brouwer, *Some unitals on 28 points and their embeddings in projective planes of order 9*, in: *Geometries and groups (Berlin, 1981)*, Lecture Notes in Math. 893, pp. 183–188, Springer, Berlin, 1981. MR 655065. Zbl 0557.51002.
- [5] P. Dembowski, *Finite geometries*, Ergebnisse der Mathematik und ihrer Grenzgebiete 44, Springer-Verlag, Berlin, 1968, doi:10.1007/978-3-642-62012-6. MR 0233275. Zbl 0865.51004.
- [6] L. E. Dickson, *Theory of linear groups in an arbitrary field*, Trans. Amer. Math. Soc. **2** (1901), no. 4, 363–394, doi:10.2307/1986251. MR 1500573. JfM 32.0131.03.
- [7] L. E. Dickson, *A new system of simple groups*, Math. Ann. **60** (1905), no. 1, 137–150, doi:10.1007/BF01447497. MR 1511290. JfM 36.0206.01.
- [8] L. C. Grove, *Classical groups and geometric algebra*, Graduate Studies in Mathematics 39, American Mathematical Society, Providence, RI, 2002, doi:10.1090/gsm/039. MR 1859189. Zbl 0990.20001.
- [9] T. Grundhöfer, B. Krinn, and M. J. Stroppel, *Non-existence of isomorphisms between certain unitals*, Des. Codes Cryptogr. **60** (2011), no. 2, 197–201, doi:10.1007/s10623-010-9428-2. MR 2795696. Zbl 05909195.
- [10] T. Grundhöfer, M. J. Stroppel, and H. Van Maldeghem, *Unitals admitting all translations*, J. Combin. Des. **21** (2013), no. 10, 419–431, doi:10.1002/jcd.21329. MR 3090721. Zbl 1276.05021.
- [11] T. Grundhöfer, M. J. Stroppel, and H. Van Maldeghem, *A non-classical unital of order four with many translations*, Discrete Math. **339** (2016), no. 12, 2987–2993, doi:10.1016/j.disc.2016.06.008. MR 3533345. Zbl 1357.51001.

-
- [12] T. Grundhöfer, M. J. Stroppel, and H. Van Maldeghem, *Moufang sets generated by translations in unitals*, J. Combin. Des. **30** (2022), no. 2, 91–104, doi:10.1002/jcd.21813, arxiv:2008.11445. MR 4373602. Zbl 7799526.
- [13] D. R. Hughes and F. C. Piper, *Projective planes*, Graduate Texts in Mathematics 6, Springer-Verlag, New York, 1973. MR 0333959. Zbl 0484.51011.
- [14] N. Jacobson, *A note on hermitian forms*, Bull. Amer. Math. Soc. **46** (1940), 264–268, doi:10.1090/S0002-9904-1940-07187-3. MR 0001957. Zbl 0024.24503. JfM 66.0048.03.
- [15] N. Jacobson, *Basic algebra. I*, W. H. Freeman and Company, New York, 2nd ed., 1985. MR 780184. Zbl 0557.16001.
- [16] W. M. Kantor, *Homogeneous designs and geometric lattices*, J. Combin. Theory Ser. A **38** (1985), no. 1, 66–74, doi:10.1016/0097-3165(85)90022-6. MR 773556. Zbl 0559.05015.
- [17] N. Knarr and M. J. Stroppel, *Split cayley hexagons via subalgebras of octonion algebras*, Manuscript (2024), submitted.
- [18] N. Knarr and M. J. Stroppel, *Subalgebras of octonion algebras*, J. Algebra **664** (2025), 42–74, doi:10.1016/j.jalgebra.2024.10.004. MR 4812310.
- [19] V. Krčadinac, *Steiner 2-designs $S(2, 4, 28)$ with nontrivial automorphisms*, Glas. Mat. Ser. III **37(57)** (2002), no. 2, 259–268. MR 1951530. Zbl 1023.05021.
- [20] H. Lüneburg, *Some remarks concerning the Ree groups of type (G_2)* , J. Algebra **3** (1966), 256–259, doi:10.1016/0021-8693(66)90014-7. MR 0193136. Zbl 0135.39401.
- [21] M. E. O’Nan, *Automorphisms of unitary block designs*, J. Algebra **20** (1972), 495–511, doi:10.1016/0021-8693(72)90070-1. MR 0295934. Zbl 0241.05013.
- [22] R. D. Schafer, *An introduction to nonassociative algebras*, Pure and Applied Mathematics 22, Academic Press, New York, 1966. MR 0210757. Zbl 0145.25601.
- [23] G. J. Schellekens, *On a hexagonal structure. I*, Nederl. Akad. Wetensch. Proc. Ser. A 65 = Indag. Math. **24** (1962), 201–217, doi:10.1016/S1385-7258(62)50019-X. MR 0143075. Zbl 0105.13001.
- [24] T. A. Springer and F. D. Veldkamp, *Octonions, Jordan algebras and exceptional groups*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2000, doi:10.1007/978-3-662-12622-6. MR 1763974. Zbl 1087.17001.

- [25] M. J. Stroppel, *Unitals without O’Nan configurations are classical if they admit all translations*, Note Mat. (2024), to appear.
- [26] D. E. Taylor, *The geometry of the classical groups*, Sigma Series in Pure Mathematics 9, Heldermann Verlag, Berlin, 1992. MR 1189139. Zbl 0767.20001.
- [27] R. A. Wilson, *The finite simple groups*, Graduate Texts in Mathematics 251, Springer-Verlag London Ltd., London, 2009, doi : 10.1007/978-1-84800-988-2. MR 2562037. Zbl 1203.20012.

Markus J. Stroppel

LExMath
Fakultät 8
Universität Stuttgart
70550 Stuttgart
stroppel@mathematik.uni-stuttgart.de