

SINGULAR METRICS WITH NONNEGATIVE SCALAR CURVATURE AND RCD

XIANZHE DAI, CHANGLIANG WANG, LIHE WANG, AND GUOFANG WEI

ABSTRACT. We show that a uniformly Euclidean metric with isolated singularity on $M^n = T^n \# M_0$, where $4 \leq n \leq 7$ or $n \geq 4$, M_0 spin, and nonnegative scalar curvature on the smooth part is Ricci flat and extends smoothly over the singularity. This confirms Schoen's Conjecture in these cases. The key to the proof is to show that the space has nonnegative synthetic Ricci curvature, i.e., an $RCD(0, n)$ space. Our result also holds when the singular set consists of a finite union of submanifolds (of possibly different dimensions) intersecting transversally under additional assumption on the co-dimension and the location of the singular set.

CONTENTS

1. Introduction	1
2. Removable singularity via RCD	6
2.1. RCD spaces and uniformly Euclidean metric	6
2.2. Gradient estimate of eigenfunctions	9
2.3. Proof of Theorems 1.2, 1.5 and 1.6	12
Appendix A. Proof of Theorem 2.6	15
References	18

1. INTRODUCTION

The existence problem of Riemannian metrics with positive scalar curvature is a fundamental topic in Riemannian geometry. In the smooth setting, it has been well studied and many important results have been established. Kazdan-Warner [23] and Schoen [32] proved that on a closed manifold, there exists a smooth metric with positive scalar curvature iff its Yamabe constant (aka σ -constant or Schoen constant) is positive. Moreover, their rigidity result says that on a closed manifold with a nonpositive Yamabe constant,

XD was partially supported by Simons Foundation.

CW was partially supported by the Fundamental Research Funds for the Central Universities and Shanghai Pilot Program for Basic Research.

GW was partially supported by NSF DMS grant 2403557.

any metric with nonnegative scalar curvature must be Ricci flat. Recall that, for a closed manifold M , its Yamabe constant $\sigma(M)$ is defined as

$$\sigma(M) := \sup\{Y(M, [g_0]) \mid [g_0] \text{ is a conformal class of smooth metrics on } M\},$$

where

$$Y(M, [g_0]) := \inf \left\{ \int_M \text{Sc}_g d\text{vol}_g \mid g \in [g_0], \text{Vol}_g(M) = 1 \right\}$$

and Sc_g denotes the scalar curvature of a smooth metric g . The Yamabe constant $\sigma(M)$ is a diffeomorphism invariant of M . For example, for a torus T^n , $\sigma(T^n) = 0$ as a consequence of the Geroch conjecture established by Schoen-Yau [30] and Gromov-Lawson [16].

For simply connected manifolds of dimension ≥ 5 , the problem of what closed manifolds should admit metrics of positive scalar curvature is fully understood in terms of the second Stiefel-Whitney class (which determines whether the manifold is spin or not) and α -invariant [15, 36].

Motivated by general relativity, the Geroch conjecture says that there is no smooth metric of positive scalar curvature on T^n . This was first proved by Schoen-Yau [30] in dimensions $n \leq 7$ using the minimal surface method, and later by Gromov-Lawson [16, 17] in arbitrary dimensions using the Dirac operator method. More generally, Gromov-Lawson [17] proved that a Λ^2 -enlargeable manifold does not admit any complete metric with positive scalar curvature. Moreover, one has the rigidity result which says that the metric with nonnegative scalar curvature on such manifolds must be flat. In particular, for any closed spin manifold M^n , $T^n \# M$ is Λ^2 -enlargeable, and as a result, $\sigma(T^n \# M) \leq 0$. Recently, these results have been generalized to allow M to be non-compact by Chodosh-Li [8] for dimensions ≤ 7 , Wang-Zhang [39] for spin manifolds of arbitrary dimensions, and also Lesourd-Unger-Yau [24] in dimension three,

Inspired by the investigation of weak notions of nonnegative scalar curvature, e.g., Gromov's polyhedral comparison theory [18], and the positive mass theorem for singular metrics, it is natural to study the forementioned Kazdan-Warner result and Geroch conjecture for singular metrics. A natural class of such singular metrics is that of uniformly Euclidean metrics. Following Li-Mantoulidis [25], we say a measurable symmetric 2-tensor g on a closed manifold M to be a *uniformly Euclidean metric* (also called L^∞ metric), if

$$(1.1) \quad \Lambda^{-1}\bar{g} \leq g \leq \Lambda\bar{g}$$

holds a.e. on M for some smooth metric \bar{g} on M and constant $\Lambda > 0$. A point is a smooth point for g if there is an open neighborhood in which g is smooth. Otherwise, we will call it a singular point. The singular set is the collection of singular point; it is closed by definition as its complement, called the smooth part of g , is an open set. We say g has positive (nonnegative resp.) scalar curvature if its scalar curvature on the smooth part is positive (nonnegative resp.).

There are two basic problems in this direction. First of all, just like the smooth case, one has the existence/non-existence problem of singular uniformly Euclidean metrics with positive scalar curvature on a closed manifold with nonpositive Yamabe constant. Secondly, related to the rigidity part, a new challenge here is the smooth extension/removable singularity problem of uniformly Euclidean metrics with nonnegative scalar curvature. Note that once the metric can be smoothly extended over the singular set, the rigidity result directly follows from the Kazdan-Warner result. For uniformly Euclidean metrics with conical singularities along a submanifold of codimension 2, Li-Mantoulidis [25] obtained an affirmative answer to both problems, provided the angle of the cone $\leq 2\pi$, while they also provided counterexample when the angle of the cone $> 2\pi$. In the case of singular set having codimension ≥ 3 , Schoen has the following conjecture.

Conjecture 1.1 (Conjecture 1.5 in [25]). Suppose M is a closed smooth manifold whose Yamabe constant is nonpositive, and g is a uniformly Euclidean metric on M that is smooth away from a closed embedded submanifold $S \subset M$ with codimension ≥ 3 . Then g extends smoothly to M , and hence is Ricci flat. In other words,

$$\begin{aligned} & \text{Sc}_g \geq 0 \text{ on } M \setminus S, \quad \sigma(M) \leq 0, \text{ and } \text{codim}(S) \geq 3 \\ \implies & g \text{ extends smoothly to } M \text{ and } \text{Ric}_g \equiv 0. \end{aligned}$$

Li-Mantoulidis [25] proved this conjecture for 3-dimensional M . They used a removal singularity result of Smith-Yang [35] for 3-dimensional Einstein metrics. On the other hand, very recently, Cecchini-Frenck-Zeidler [3] constructed counterexamples to Conjecture 1.1 on simply connected closed manifold of dimension ≥ 8 . In contrast, the conjecture could still be true on manifolds with certain topology. Dai-Sun-Wang [10] proved Conjecture 1.1 for metrics with isolated conical singularities on $T^n \# M_0$, including a smooth extension and rigidity result, provided that $3 \leq n \leq 7$ or M_0 is spin. On $T^n \# M_0$, the non-existence problem of general uniformly Euclidean metrics with the singular set having co-dimension $\geq \frac{n}{2} + 1$ can be answered affirmatively in a similar way. Kazaras [21] proved that, on a four-dimensional enlargeable closed smooth manifold, a uniformly Euclidean metric g that is smooth away from a smoothly embedded finite 1-complex and has nonnegative scalar curvature on the smooth part must be Ricci flat on the smooth part. Cecchini-Frenck-Zeidler [3] proved that a uniformly Euclidean metric on a closed spin manifold with the fundamental group satisfying some condition on Rosenberg index must be Ricci flat on the smooth part, provided that the singular set consisting of finitely many points and the scalar curvature is nonnegative on the smooth part. Wang-Xie [38] also proved a similar result on manifolds admitting a Lipschitz map to T^n with nonzero degree and allowing the singular set having codim $\geq \frac{n}{2} + 1$. Shi-Wang-Wu-Zhu [34] defined open Schoen-Yau-Schick manifolds and proved that there is no complete metric with positive scalar curvature on such manifolds and consequently proved that a uniformly Euclidean metric with nonnegative scalar curvature and a singular set of codimension $\geq \frac{n}{2} + 1$ on a closed Schoen-Yau-Schick manifold must be Ricci flat on the smooth part. Kazaras

[21], Cecchini-Frenck-Zeidler [3], Wang-Xie [38] and Shi-Wang-Wu-Zhu [33] obtained the Ricci flatness of the metrics on the smooth part, but did not establish a smooth extension result.

The main result of this paper is a partial affirmative solution of Conjecture 1.1 for manifolds which are connected sums with the torus.

Theorem 1.2. *Let $M^n = T^n \# M_0$, $n \geq 4$, be a connected closed manifold. Assume either $4 \leq n \leq 7$ or M_0 is spin. Suppose*

- 1). *g is a uniformly Euclidean metric on M that is smooth away from a closed, embedded submanifold S with codimension $\geq \max(4, \frac{n}{2} + 1)$.*
- 2). *There is a $(n - 1)$ -sphere dividing $M = T^n \# M_0$ into the T^n -part and M_0 -part, such that each component of S lies either in the T^n -part or M_0 -part. Further, if a component of S lies in the torus part, we assume that it is contained in a topological ball.*
- 3). *The scalar curvature $Sc_g \geq 0$ on $M \setminus S$.*

Then g extends smoothly to M (possibly with a change of differential structure) and (M^n, g) is isometric to a flat torus.

In particular, this confirms Schoen's Conjecture for isolated singularity when $M^n = T^n \# M_0$, $4 \leq n \leq 7$ or $n \geq 4$ and M_0 is spin.

Remark 1.3. As pointed out in [3], one may need to change differential structure when extending the metric. This is also the case for other well-known removable singularity results. Our result also holds for singular set which consists of a finite union of submanifolds (of possibly different dimensions) intersecting transversally.

Remark 1.4. One may think of this result as a removable singularity theorem. Unlike the well-known removable singularity results for Einstein metrics or Yang-Mills instantons, our result is not local but rely on the global topological structure.

A novel and crucial ingredient in the proof of Theorem 1.2 is the following result, which is of independent interest. It says that a uniformly Euclidean metric on a closed manifold smooth away from a closed singular set of codimension 4 or higher and with Ricci curvature bounded below by K on the smooth part is $\text{RCD}(K, n)$, i.e., its Ricci curvature is bounded below by K in the weak sense.

Theorem 1.5. *Let M^n be a connected closed manifold of dimension $n \geq 4$, and g a uniformly Euclidean metric on M that is smooth away from a closed set S with codimension ≥ 4 . If, for some constant K , the Ricci curvature $\text{Ric}_g \geq K$ on the smooth part, then (M^n, g) is a $\text{RCD}(K, n)$ space.*

As an interesting application, we obtain the following removable singularity result. In contrast to the local nature of the previous results of this type, our result is global, depending crucially on the global topological assumption.

Theorem 1.6. *Let M^n be a connected closed manifold of dimension $n \geq 4$ with infinite fundamental group. If g is a uniformly Euclidean metric on M with isolated singularity and nonnegative Ricci curvature on the smooth part, then (M^n, g) extends smoothly to M and thus $\text{Ric}_g \geq 0$. Moreover, a finite cover \hat{M} of M is diffeomorphic to $T^k \times X$, for some $1 \leq k \leq n$ and X a closed simply connected manifold. Furthermore, there is a locally isometrically trivial fibration $\hat{M} \rightarrow T^k$ with fiber X .*

The main technical component of the proof of Theorem 1.5 is a gradient estimate for eigenfunctions. It consists of two steps. First, using De Giorgi-Nash-Moser estimate and Cheng-Yau gradient estimate, we obtain an initial gradient estimate for eigenfunctions, which blows up at the singular part of g . Nevertheless, this enables us to establish an L^2 bound on the Hessian of the eigenfunctions, using the uniform Euclidean and codimension assumptions. Then, by Bochner formula and Kato's inequality, a differential inequality for the norm of the gradient of an eigenfunction is shown to hold over the whole manifold in the weak sense, thanks to the L^2 Hessian bound. The final gradient estimate then follows by the Moser iteration. This is similar in spirit to the strategy used in [28, 2] for stratified spaces.

This paper is organized as follows. In Section 2, we prove Theorem 1.5 and then apply it to prove Theorems 1.2 and 1.6. More precisely, in Subsection 2.1, we recall a definition of RCD space and describe a metric measure structure induced by a uniformly Euclidean metric as in Conjecture 1.1. We also prove that such metric measure spaces satisfy the infinitesimal Hilbertian condition and Sobolev-to-Lipschitz property. In Subsection 2.2, we derive a gradient estimate for the eigenfunctions of the Laplace operator, which plays an important role in the proof of the Bakry-Émery (BE, for short) condition for uniformly Euclidean metrics. Based on these, in Subsection 2.3, we prove Theorem 1.5, which follows from [19, Theorem 3.7]; see also Bertrand-Ketterer-Mondello-Richard's work on stratified spaces [2], as well as related work of Jiang-Sheng-Zhang [20] and Szekelyhidi [37]. By applying Theorem 1.5, we then prove Theorems 1.2 and 1.6, and give several interesting examples as their applications. These examples illustrate how naturally our approach of using metric measure spaces deals with the possible change of differential structures, as well as situations outside Theorem 1.5 our results still apply. Finally, in Appendix A, we provide a detailed proof of Theorem 2.6.

Acknowledgements: This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930, while XD and GW were in residence at the Simons Laufer Mathematical Sciences Institute (formerly MSRI) in Berkeley, California, during the Fall semester of 2024. XD and GW thank SLMATH for providing a stimulating environment and acknowledge useful discussions with Rick Schoen, Vincent Guedj, Xinyu Zhu, Conghan Dong. XD and CW would like to thank Jian Wang and Simone Cecchini for valuable conversations. Finally we would like to thank Gabor Szekelyhidi for very fruitful communication.

2. REMOVABLE SINGULARITY VIA RCD

2.1. RCD spaces and uniformly Euclidean metric. We recall a definition of the RCD(K, N) condition for metric measure spaces, see [1, 11, 13].

Definition 2.1. *A metric measure space (X, d, m) is said to be an RCD(K, N) space for some $K \in \mathbb{R}$ and $N \in [1, \infty]$, if*

- (1) *there exists $x_0 \in X$ and a constant $C > 0$ such that $\int_X e^{-Cd(x, x_0)} dm < +\infty$,*
- (2) *(Infinitesimally Hilbertian condition) the Sobolev space $W^{1,2}(X, d, m)$ is a Hilbert space,*
- (3) *(Sobolev-to-Lipschitz property) for any $f \in W^{1,2}(X, d, m)$ satisfying $|\nabla f| \in L^\infty(M)$, it has a Lipschitz representative \tilde{f} with $\text{Lip}(\tilde{f}) \leq \|\nabla f\|_{L^\infty(X)}$,*
- (4) *(BE(K, N) condition) for any $f \in D(\Delta)$, and any test function $\varphi \in L^\infty(X) \cap D(\Delta)$ with $\varphi \geq 0$ and $\Delta\varphi \in L^\infty(X)$, the weak Bochner inequality*

$$(2.1) \quad \frac{1}{2} \int_X |\nabla f|^2 \Delta\varphi dm \geq \int_X \varphi \left(\frac{1}{N} (\Delta f)^2 + \langle \nabla f, \nabla \Delta f \rangle + K |\nabla f|^2 \right) dm$$

holds.

In this paper, we focus on a closed connected smooth manifold M^n with a uniformly Euclidean metric g as in Conjecture 1.1, that is, g is smooth away from a closed singular set S with codimension ≥ 3 . We will denote by $\Omega := M \setminus S$ the smooth part of the metric and by $k := \dim(S)$.

We describe a metric measure structure on such (M^n, g) as follows. We say that a piecewise C^1 curve $\gamma : [a, b] \rightarrow M$ is *admissible* if its image $\gamma([a, b])$ intersects with S at only finitely many points. For such a curve, its length is defined as $L_g(\gamma) = \int_a^b |\gamma'(t)|_g dt$. Then, the distance between any $x, y \in M$ is defined as

$$d_g(x, y) := \inf \{ L_g(\gamma) \mid \gamma : [a, b] \rightarrow M \text{ is admissible curve with } \gamma(a) = x, \gamma(b) = y \}.$$

The volume measure $d\mu_g$ is defined in the usual way. A direct consequence of the uniformly Euclidean condition is

$$(2.2) \quad \Lambda^{-\frac{n}{2}} d\mu_{\bar{g}} \leq d\mu_g \leq \Lambda^{\frac{n}{2}} d\mu_{\bar{g}},$$

where $d\mu_{\bar{g}}$ is the volume measure of the smooth metric \bar{g} on M . As a result, on a sufficiently small tubular neighborhood $T_\epsilon(S)$ of the singular set S , we have that locally,

$$(2.3) \quad d\mu_g \sim r^{n-k-1} dr d\theta^{n-k-1} d\mu_{\bar{g}_S},$$

where $r = r(x) := d(x, S)$, $d\theta^{n-k-1}$ is the volume measure on \mathbb{S}^{n-k-1} , and $d\mu_{\bar{g}_S}$ is the volume measure on S with the restricted metric \bar{g}_S . Here \sim means bounded above and below by the right hand side (not precise asymptotic). By applying this volume measure estimate, we prove the following zero capacity property of S .

Lemma 2.2 (Zero capacity property). *Let M^n , $n \geq 3$, be a closed manifold, and g a uniformly Euclidean metric on M with singular set a closed embedded submanifold of codimension ≥ 3 . There exists a family of functions $\psi_\delta \in C_0^\infty(\Omega)$, $\delta \in (0, \delta_0)$ for some $\delta_0 > 0$, such that the following two properties hold:*

- (1) *for any compact subset $A \subset \Omega := M \setminus S$, $\psi_\delta|_A \equiv 1$ hold for all sufficiently small δ ,*
- (2) *$0 \leq \psi_\delta \leq 1$ for all δ and $|\nabla \psi_\delta| \leq \frac{C}{\delta}$. Moreover,*

$$(2.4) \quad \int_{\Omega} |\nabla \psi_\delta|^2 d\mu_g \rightarrow 0, \quad \text{as } \delta \rightarrow 0.$$

Proof. Take a smooth cut-off function $\eta : [0, \infty) \rightarrow [0, 1]$ such that

$$(2.5) \quad \eta = \begin{cases} 0, & \text{in } [0, \frac{1}{2}], \\ 1, & \text{in } [1, \infty), \end{cases}$$

and $|\eta'| \leq C$ on $[0, \infty)$ for some constant C . Let $r(x) := d(x, S)$, and $\psi_\delta(x) := \eta(\frac{r}{\delta})$. Clearly, $|\nabla \psi_\delta| \leq \frac{C}{\delta}$ and $\text{supp}(\nabla \psi_\delta) \subset A_{\frac{\delta}{2}, \delta}(p) := \{x \in M \mid \frac{\delta}{2} \leq d(x, S) \leq \delta\}$. By combining with the estimate of the volume measure in (2.3), this implies

$$(2.6) \quad \int_{\Omega} |\nabla \psi_\delta|^2 d\mu_g \leq \frac{C^2}{\delta^2} \int_{A_{\frac{\delta}{2}, \delta}(p)} d\mu_g \leq \frac{C'}{\delta^2} \int_{\frac{\delta}{2}}^{\delta} r^{n-k-1} dr \leq C'' \delta^{n-k-2} \rightarrow 0,$$

as $\delta \rightarrow 0$, since $n - k \geq 3$. This completes the proof. \square

By using Lemma 2.2, one can show that the Sobolev space $H^{1,2}(M)$ defined as the $H^{1,2}$ -completion of $C_0^\infty(M \setminus S)$ coincides with the Sobolev space $W^{1,2}(M, d_g, d\mu_g)$ on the metric measure space $(M, d_g, d\mu_g)$, see Remark 3.2 and Proposition 3.3 in [19], for details. In particular, $(M, d_g, d\mu_g)$ satisfies the infinitesimally Hilbertian condition in Definition 2.1.

For $f \in W^{1,2}(M)$, if there exists a function $u \in W^{1,2}(M)$ such that

$$\int_M u v d\mu_g = - \int_M \langle df, dv \rangle_g d\mu_g$$

holds for any test function $v \in C_0^\infty(M \setminus S)$, then $\Delta f := u$ is said to be the Laplace of f , and the domain of the Laplace operator is

$$D(\Delta) := \{f \in W^{1,2}(M) \mid \Delta f \in W^{1,2}(M)\}.$$

For proving the Sobolev-to-Lipschitz property for uniformly Euclidean metrics, we need the following result about the length of admissible curves.

Lemma 2.3. *Let M^n , $n \geq 3$, be a closed connected manifold, and g a uniformly Euclidean metric on M as in Conjecture 1.1. Let $\gamma : [a, b] \rightarrow M$ be an admissible curve on M . For any $\epsilon > 0$, there exists an admissible curve γ_ϵ with the same endpoints as γ such that the image of γ_ϵ is contained in the regular part $M \setminus S$ probably except the endpoints, and $L_g(\gamma_\epsilon) \leq L_g(\gamma) + \epsilon$.*

Proof. Because γ intersects with S at only finitely many points and the length is additive, without loss of generality, we assume that γ intersects with S only at an interior point p . Then we only need to modify the curve γ near p . For that, we take a small ball $B_\delta(p)$ centered at p with radius δ with respect to the smooth metric \bar{g} . Let $\gamma(t_1)$ be the first point on γ that intersects $\partial B_\delta(p)$, and $\gamma(t_2)$ be the last such point, where $[t_1, t_2] \subset (a, b)$. For sufficiently small $\delta > 0$, we can take a smooth curve c in $B_\delta(p)$ that connects $\gamma(t_1)$ and $\gamma(t_2)$, has length $< \frac{\epsilon}{\sqrt{\Lambda}}$ with respect to \bar{g} , and does not intersect with S . Now, define γ_ϵ to be the concatenation of $\gamma|_{[a, t_0]}$, c and $\gamma|_{[t_1, b]}$. Then the image of γ_ϵ intersects with S at most at the endpoints. Moreover, $L_g(\gamma_\epsilon) \leq L_g(\gamma) + L_g(c) \leq L_g(\gamma) + \epsilon$. \square

Lemma 2.4. *Let M^n , $n \geq 3$, be a closed connected manifold, and g a uniformly Euclidean metric on M as in Conjecture 1.1. Any $f \in W^{1,2}(M)$ with gradient $\nabla f \in L^2(M) \cap L^\infty(M)$ has a Lipschitz representative \tilde{f} whose Lipschitz constant $\leq \|\nabla f\|_{L^\infty(M)}$.*

Proof. First note that the smooth part $\Omega := M \setminus S$ is path connected, since the singular set S has codimension ≥ 3 . Take any two points $x, y \in \Omega$. By Lemma 2.3, for any $\epsilon > 0$, there exists a piecewise smooth curve $\gamma_\epsilon \subset \Omega$ from x to y such that the length $L_g(\gamma_\epsilon)$ of γ with respect to g

$$(2.7) \quad L_g(\gamma) \leq (1 + \epsilon)d_g(x, y).$$

Then by locally Lipschitz property of f in Ω with Lipschitz constant $\leq \|\nabla f\|_{L^\infty(M)}$, and applying the fundamental theorem of calculus to $u \circ \gamma$, we have

$$|f(x) - f(y)| \leq (1 + \epsilon)\|\nabla f\|_{L^\infty(M)}d_g(x, y).$$

Thus, by letting $\epsilon \rightarrow 0$, we have that u is uniformly Lipschitz on Ω with Lipschitz constant $\leq \|\nabla f\|_{L^\infty(M)}$. Finally, because Ω is dense in M , u has a unique Lipschitz continuous extension \tilde{f} on M , which equals f almost everywhere on M . \square

The uniformly Euclidean property directly implies the following L^2 -strong compactness property.

Lemma 2.5 (L^2 -strong compactness). *Let M^n , $n \geq 3$, be a closed manifold, and g a uniformly Euclidean metric on M . The inclusion map $i : W^{1,2}(M, g) \hookrightarrow L^2(M, g)$ is compact.*

Proof. By the definition, the Sobolev space $W^{1,2}(M, g)$ and $L^2(M, g)$ depend only on the quasi-isometry class of g and hence is equivalent to $W^{1,2}(M, \bar{g})$ for a smooth metric \bar{g} . Clearly $W^{1,2}(M, \bar{g})$ embeds compactly in $L^2(M, \bar{g})$ on a compact manifold. \square

Using the L^2 -strong compactness in Lemma 2.5 and the standard argument in functional analysis, one obtains that the spectrum of the Laplace operator Δ_g consists of discrete eigenvalues with finite multiplicities, and normalized eigenfunctions form an orthonormal basis of $L^2(M, g)$.

2.2. Gradient estimate of eigenfunctions. For proving the $BE(K, n)$ condition for uniformly Euclidean metrics, as in Theorem 3.7 in [19], we expand a Sobolev function in terms of eigenfunctions of Δ_g , and a gradient estimate of eigenfunctions plays a crucial role in the proof. In this subsection, we derive some estimates for gradient and Hessian of the eigenfunctions.

Let M^n be a closed manifold, and g a uniformly Euclidean metric on M that is smooth away from a closed embedded submanifold S . As M is a closed manifold and g is uniformly Euclidean, for any $q \in \Omega$, $0 < 2r < d(q, S)$, the gradient estimate of Cheng-Yau [7] still applies on $B_r(q)$, provided one has a Ricci curvature lower bound on $B_{2r}(q)$. The following combines the versions in [4] and [26].

Theorem 2.6. *Let (M^n, g) be as above and $\Omega := M \setminus S$ the regular part of g . If u is a positive function defined on the closed ball $B(q, 2r) \subset \Omega$ that satisfies $\Delta u = K(u)$ and moreover $\text{Ric}_g \geq -(n-1)H^2$ ($H \geq 0$) on $B(q, 2r)$. Then on $B(q, r)$,*

$$\frac{|\nabla u(x)|}{u} \leq \max \left\{ \sup_{B(q, 2r)} \left(\frac{|K(u)|}{u} \right)^{\frac{1}{2}}, \right. \\ \left. (n-1)H + C_1(n)r^{-1} + C_2(n, H)r^{-\frac{1}{2}} + \sup_{B(q, 2r)} \left[(n-1) \left| K' - \frac{n+1}{n-1} \frac{K}{u} \right| \right]^{\frac{1}{2}} \right\}.$$

Since this version is not in the literature, for completeness we provide a proof in the appendix.

Here, as an application, we deduce the following crucial estimate.

Lemma 2.7. *Let (M^n, g) be as above.*

(1) *There exists $0 < \alpha = \alpha(n, \Lambda) < 1$, such that the $W^{1,2}$ -eigenfunctions u of the Laplace operator Δ_g satisfy $u \in C^\alpha(M)$. Moreover, if λ is the eigenvalue,*

$$(2.8) \quad [u]_{C^\alpha(M)} \leq C(n, \Lambda, \lambda, \bar{g}) \|u\|_{L^\infty(M)} \leq C \|u\|_{L^2(M)}.$$

(2) *Assume that $\text{Ric}_g \geq K$ on Ω for some constant K . For all $x \in \Omega$ with $r(x) = d(x, S) \leq 1$, we have the gradient estimate*

$$(2.9) \quad |\nabla u|(x) \leq \frac{C(n, \Lambda, \lambda, \bar{g}, K)}{r(x)^{1-\alpha}} \|u\|_{L^2(M)},$$

for the same α in (2.8).

Proof. Since M is uniformly Euclidean the first statement (1) is the De Giorgi-Nash-Moser estimate, see Chapter 8 in [14], [27] for example.

For (2), in order to apply Theorem 2.6 we do a shift to get a positive solution. Namely fix any $y \in \Omega$ and let $r(y) = 2s > 0$. Then $v = u - m + s \geq s$ with $m = \inf\{u : B_{2s}(y)\}$ is a positive solution of

$$(2.10) \quad -\Delta v = \lambda v + \lambda(m - s)$$

in the ball $B_{2s}(y) \subset \Omega$. Since $\text{Ric} \geq K$ on Ω , by Theorem 2.6, we have on $B_s(y)$

$$\begin{aligned} |\nabla u| &= |\nabla v| \leq v \left((n-1)K + C(n)s^{-1} + C(n, K)s^{-\frac{1}{2}} + \left[(n-1)(2\lambda + \lambda \frac{|m|+s}{s}) \right]^{1/2} \right) \\ &= v s^{-1} \left((n-1)Ks + C(n) + C(n, K)\sqrt{s} + \sqrt{(n-1)s\lambda(3s+|m|)^{1/2}} \right). \end{aligned}$$

By definition, there is a $q \in \overline{B_{2s}(y)}$ such that

$$|v(y)| \leq |u(y) - u(q)| + s.$$

Now by (2.8),

$$|u(y) - u(q)| \leq Cs^\alpha \|u\|_{L^2(M)}.$$

Hence, for $s \leq 1$,

$$|\nabla u(y)| \leq C(n, \Lambda, \lambda, \bar{g}, K)s^\alpha s^{-1} \|u\|_{L^2(M)}.$$

Our estimate follows since $y \in \Omega$ is arbitrary. \square

By applying the gradient estimate in Lemma 2.7, we prove the L^2 -integrability of the Hessian of eigenfunctions as in the following lemma.

Lemma 2.8. *Let M^n be a closed manifold, and g a uniformly Euclidean metric on M that is smooth away from a closed embedded submanifold S with codimension $n - k \geq 4$, and $\text{Ric}_g \geq K$ on Ω . For any $W^{1,2}$ -eigenfunction u of Δ_g , $|\text{Hess } u| \in L^2(M)$.*

Proof. Let u be a $W^{1,2}$ -eigenfunction of Δ with eigenvalue λ , i.e. $-\Delta u = \lambda u$ in the weak sense. By standard elliptic regularity theory, $u \in C^\infty(\Omega)$. Bochner formula and $\text{Ric}_g \geq K$ on Ω imply

$$(2.11) \quad \frac{1}{2}\Delta|\nabla u|^2 \geq |\text{Hess } u|^2 + (K - \lambda)|\nabla u|^2 \quad \text{on } \Omega.$$

Multiplying the inequality (2.11) by ψ_δ^2 in Lemma 2.2, and then integrating it over M gives

$$(2.12) \quad \frac{1}{2} \int_M |\nabla u|^2 \Delta \psi_\delta^2 d\mu_g \geq \int_M \psi_\delta^2 (|\text{Hess } u|^2 + (K - \lambda)|\nabla u|^2) d\mu_g.$$

For the left hand side,

$$\begin{aligned} \int_M |\nabla u|^2 \Delta \psi_\delta^2 d\mu_g &= -4 \int_M |\nabla u| \psi_\delta \langle \nabla |\nabla u|, \nabla \psi_\delta \rangle d\mu_g \\ &\leq \int_M \psi_\delta^2 |\text{Hess } u|^2 d\mu_g + 4 \int_M |\nabla u|^2 |\nabla \psi_\delta|^2 d\mu_g. \end{aligned}$$

Plugging this into (2.12) and rearranging the equation give

$$\frac{1}{2} \int_M \psi_\delta^2 |\text{Hess } u|^2 d\mu_g \leq \int_M 2|\nabla u|^2 |\nabla \psi_\delta|^2 d\mu_g - (K - \lambda) \int_M \psi_\delta^2 |\nabla u|^2 d\mu_g.$$

Then by applying estimates in Lemmas 2.2 and 2.7 and equation (2.3) for the integrand in the first term on the right hand side, we obtain

$$\begin{aligned} \frac{1}{2} \int_M \psi_\delta^2 |\text{Hess } u|^2 d\mu_g &\leq C \int_{\frac{\delta}{2}}^\delta r^{2\alpha+n-k-5} dr - (K - \lambda) \int_M \psi_\delta^2 |\nabla u|^2 d\mu_g \\ &\leq \tilde{C} \delta^{2\alpha-4+n-k} - (K - \lambda) \int_M \psi_\delta^2 |\nabla u|^2 d\mu_g. \end{aligned}$$

Now because $n - k \geq 4$, by letting $\delta \rightarrow 0$, we obtain

$$(2.13) \quad \int_M |\text{Hess } u|^2 d\mu_g \leq 2(\lambda - K) \int_M |\nabla u|^2 d\mu_g < +\infty.$$

□

By using Lemma 2.8, similar as in [28] on page 56 for eigenfunctions on stratified spaces, we prove

Lemma 2.9. *Let (M^n, g) as in Lemma 2.8. For any $W^{1,2}$ -eigenfunction u of Δ_g , $|\nabla u| \in L^\infty(M)$.*

Proof. Let u be a $W^{1,2}$ -eigenfunction of Δ with eigenvalue λ . Then $u \in C^\infty(\Omega)$. For simplicity of notation, we denote $v := |\nabla u|$. By (2.11), Kato's inequality $|\nabla|\nabla u|| \leq |\text{Hess } u|$, one can show that there exists a positive constant c such that

$$(2.14) \quad -\Delta v \leq cv \text{ in the barrier sense on } \Omega.$$

Indeed, let $v_\epsilon = \sqrt{|\nabla u|^2 + \epsilon^2} - \epsilon$. By the Kato's inequality, one has

$$(2.15) \quad |\nabla v_\epsilon|^2 = \frac{|\nabla u|^2 \cdot |\nabla|\nabla u||^2}{|\nabla u|^2 + \epsilon^2} \leq \frac{|\nabla u|^2 \cdot |\text{Hess } u|^2}{|\nabla u|^2 + \epsilon^2} \leq |\text{Hess } u|^2.$$

Moreover, putting $c := \max\{\lambda - K, 1\} > 0$, and using (2.11) and (2.15), one has

$$\begin{aligned} \frac{1}{2} \Delta(v_\epsilon + \epsilon)^2 &= \frac{1}{2} \Delta|\nabla u|^2 \geq |\text{Hess } u|^2 - (\lambda - K)|\nabla u|^2 \\ &\geq |\text{Hess } u|^2 - c((v_\epsilon + \epsilon)^2 - \epsilon^2) \\ &\geq |\nabla v_\epsilon|^2 - c(v_\epsilon + \epsilon)^2 \text{ on } \Omega. \end{aligned}$$

Thus,

$$(v_\epsilon + \epsilon) \Delta(v_\epsilon + \epsilon) + |\nabla v_\epsilon|^2 \geq |\nabla v_\epsilon|^2 - c(v_\epsilon + \epsilon)^2.$$

Consequently, $\Delta v_\epsilon \geq -c v_\epsilon - c\epsilon$ for any $\epsilon > 0$.

If $x \in \Omega$ and $v(x) \neq 0$, then v is smooth at x . The calculation above with $\epsilon = 0$ shows that $\Delta v \geq -cv$.

If $v(x_0) = 0$, then $v_\epsilon(x_0) = 0 = v(x_0)$ and $v_\epsilon \leq v$. Hence, $\Delta v \geq -cv$ in the barrier sense at x_0 .

We now claim that the inequality in (2.14) holds in the weak sense on the whole manifold M . Then, with the help of the Sobolev inequality in Lemma 2.5, a Moser iteration argument implies $v = |\nabla u| \in L^\infty(M)$.

In the rest of the proof, we show that the inequality in (2.14) holds in the weak sense on M , that is, for any $\varphi \in W^{1,2}(M)$, $\varphi \geq 0$,

$$(2.16) \quad \int_M \langle \nabla v, \nabla \varphi \rangle d\mu_g \leq c \int_M v \varphi d\mu_g.$$

Because $C^\infty(M)$ is dense in $W^{1,2}(M)$, we only need to establish the inequality (2.16) for $\varphi \in C^\infty(M)$, $\varphi \geq 0$. For such φ and ψ_δ in Lemma 2.2, we multiply the inequality in (2.14) by $\psi_\delta \varphi$, and integrate over M . Then by integrating by parts and rearranging the inequality, we obtain

$$(2.17) \quad \int_M \psi_\delta \langle \nabla v, \nabla \varphi \rangle d\mu_g \leq c \int_M \psi_\delta \varphi v d\mu_g - \int_M \varphi \langle \nabla v, \nabla \psi_\delta \rangle d\mu_g.$$

For the second term on the right hand side, because the smooth function φ on the compact manifold M is bounded, we have

$$\left| \int_M \varphi \langle \nabla v, \nabla \psi_\delta \rangle d\mu_g \right| \leq C \left| \int_M \langle \nabla v, \nabla \psi_\delta \rangle d\mu_g \right|.$$

Then by applying Cauchy-Schwarz inequality and the Kato's inequality, we obtain

$$\left| \int_M \varphi \langle \nabla v, \nabla \psi_\delta \rangle d\mu_g \right| \leq C \|\nabla v\|_{L^2(M)} \|\nabla \psi_\delta\|_{L^2(M)} \leq C \|\text{Hess } u\|_{L^2(M)} \|\nabla \psi_\delta\|_{L^2(M)} \rightarrow 0,$$

as $\delta \rightarrow 0$, since $\|\text{Hess } u\|_{L^2(M)}$ is finite by Lemma 2.8 and $\|\psi_\delta\|_{L^2(M)} \rightarrow 0$ by Lemma 2.2. Thus, by letting $\delta \rightarrow 0$ in (2.17), we prove the inequality (2.16) for nonnegative smooth function φ , so for nonnegative test functions $\varphi \in W^{1,2}(M)$. \square

2.3. Proof of Theorems 1.2, 1.5 and 1.6. With our work from the previous subsection, Theorem 1.5 follows from [19, Theorem 3.7], see also [2, 20]. For completeness, we give a detailed proof here.

Proof of Theorem 1.5. It suffices to check four conditions in Definition 2.1. Condition (1) follows from the compactness of M and the uniform Euclidean property of g . Condition (2) is also clear. Condition (3) follows from Lemma 2.4.

It remains to check the BE(K, n) condition in (4). Let $\Omega := M \setminus S$, $\{u_i\}_{i=1}^{+\infty}$ be the orthonormal basis of $L^2(M)$ consisting of eigenfunctions of Δ . For any $f \in W^{1,2}(M)$ and $N \in \mathbb{N}$, let $f_N := \sum_{i=1}^N a_i u_i$, where $a_i := \int_M f u_i d\mu_g$. As before, $k = \dim S$, and by assumption $n - k \geq 4$.

Bochner formula and $\text{Ric}_g \geq K$ on Ω imply

$$(2.18) \quad \frac{1}{2} \Delta |\nabla f_N|^2 \geq |\text{Hess } f_N|^2 + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2 \quad \text{on } \Omega.$$

By Lemma 2.8 and Lemma 2.9, for any eigenfunction u_i , we have

$$\int_M |\nabla |\nabla u_i|^2|^2 d\mu_g = 4 \int_M |\nabla u_i|^2 |\nabla |\nabla u_i|^2|^2 d\mu_g \leq 4 \int_M |\nabla u_i|^2 |\text{Hess } u_i|^2 d\mu_g < +\infty.$$

Thus, $|\nabla u_i|^2 \in W^{1,2}(M)$ for all $i \in \mathbb{N}$, so $|\nabla f_N|^2 \in W^{1,2}(M)$. Also $|\text{Hess } f_N| \in L^2(M)$ from Lemma 2.8.

For any test function $\varphi \in L^\infty(M) \cap D(\Delta)$ with $\varphi \geq 0$ and $\Delta\varphi \in L^\infty(M)$, by multiplying the inequality (2.18) by $\varphi\psi_\delta$, and integration by parts, we have

$$(2.19) \quad \begin{aligned} & -\frac{1}{2} \int_M \langle \nabla(\varphi\psi_\delta), \nabla |\nabla f_N|^2 \rangle d\mu_g \\ & \geq \int_M \varphi\psi_\delta (|\text{Hess } f_N|^2 + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2) d\mu_g. \end{aligned}$$

By Lemma 2.2, $\psi_\delta \rightarrow 1$ in $L^2(M)$ and $\nabla\psi_\delta \rightarrow 0$ in $L^2(M)$ as $\delta \rightarrow 0$, and since $|\nabla f_N|^2 \in W^{1,2}(M)$, we have

$$-\frac{1}{2} \int_M \langle \nabla(\varphi\psi_\delta), \nabla |\nabla f_N|^2 \rangle \rightarrow -\frac{1}{2} \int_M \langle \nabla\varphi, \nabla |\nabla f_N|^2 \rangle d\mu_g = \frac{1}{2} \int_M |\nabla f_N|^2 \Delta\varphi d\mu_g,$$

as $\delta \rightarrow 0$. Moreover, by the dominated convergence theorem, we have

$$\begin{aligned} & \int_M \varphi\psi_\delta (|\text{Hess } f_N|^2 + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2) d\mu_g \\ & \rightarrow \int_M \varphi (|\text{Hess } f_N|^2 + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2) d\mu_g. \end{aligned}$$

Thus, by letting $\delta \rightarrow 0$ in (2.19), we have

$$\begin{aligned} \frac{1}{2} \int_M |\nabla f_N|^2 \Delta\varphi d\mu_g & \geq \int_M \varphi (|\text{Hess } f_N|^2 + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2) d\mu_g \\ & \geq \int_M \varphi \left(\frac{(\Delta f_N)^2}{n} + \langle \nabla \Delta f_N, \nabla f_N \rangle + K |\nabla f_N|^2 \right) d\mu_g. \end{aligned}$$

Finally, letting $N \rightarrow \infty$, we obtain that

$$\frac{1}{2} \int_M |\nabla f|^2 \Delta\varphi d\mu_g \geq \int_M \varphi \left(\frac{(\Delta f)^2}{n} + \langle \nabla \Delta f, \nabla f \rangle + K |\nabla f|^2 \right) d\mu_g$$

for any test function $\varphi \in L^\infty(M) \cap D(\Delta)$ with $\varphi \geq 0$ and $\Delta\varphi \in L^\infty(M)$. This completes the proof. \square

Theorem 1.5 provides the crucial ingredient to the proof of Theorem 1.2.

Proof of Theorem 1.2. First of all, we note that, under the assumptions for M_0 in Theorem 1.2, there is no uniformly Euclidean metric g on $M := T^n \# M_0$ smooth away from a closed embedded submanifold S with co-dimension $\geq \frac{n}{2} + 1$, which has nonnegative

scalar curvature on the smooth part and positive scalar curvature at a point. This non-existence result can be established by using the conformal blow up technique dating back to Schoen's final resolution of Yamabe problem [31], which reduces the problem to the non-existence results of Chodosh-Li [8] and Wang-Zhang [39] for complete metrics. This was recently used in studies of uniformly Euclidean metric with positive scalar curvature by Li-Mantoulidis [25], Kazaras [21], Wang-Xie [38] and others, and also in studies of positive mass theorem and positive scalar curvature with conical singularity by Dai-Sun-Wang [9, 10]. Since the method carries over in a similar way, we refer to these references for details. Roughly speaking, one uses the Green's function of the conformal Laplacian but with the scalar curvature suitably truncated to conformally blow up the singular metric. This produces a complete metric with positive scalar curvature on $(T^n \# M_0) \setminus S$, leading to a contradiction with non-existence results of Chodosh-Li [8] and Wang-Zhang [39]. We only emphasize a few things here. Firstly, for uniformly Euclidean metrics, the Green's function can be solved by Littman-Stampacchia-Weinberger [27] and Schoen-Yau [30]. Also, the condition that the co-dimension of $S \geq \frac{n}{2} + 1$ guarantees the completeness of the blow up metric, see Proposition 2.4 in Wang-Xie [38]. And finally, the assumption on the location and size of the singular set, Condition 2), guarantees that $M \setminus S$ can again be written as $T^n \# M_1$ for suitable M_1 , and hence the results of Chodosh-Li [8] and Wang-Zhang [39] can be applied.

Then by Theorem B in Kazdan [22], the metric g must be Ricci flat. Therefore, Theorem 1.5 implies that $(T^n \# M_0, d_g, d\mu_g)$ is a RCD(0, n) space. Consequently, we conclude that $(T^n \# M_0, d_g)$ is isometric to a flat torus T^n by Corollary 1.4 in Mondino-Wei [29], and in particular, g extends smoothly to $T^n \# M_0$ (possibly with a change of differential structure). This completes the proof. \square

Proof of Theorem 1.6. By Theorem 1.5, (M, g) is RCD(0, n) and the splitting theorem of Gigli [12] applies. Since $\pi_1(M)$ is infinite, the universal cover \tilde{M} splits as $\mathbb{R}^k \times X$ for some $1 \leq k \leq n$ and X compact simply connected metric measure space. Now if \tilde{X} has singularity, it will give rise to non-isolated singularity for M . Since (M, g) has only isolated singularity, X must be smooth and our result follows, see [6, Theorem 9.2] and [5] for more details. \square

We will end this section with several examples.

Example 1: This example is motivated by Cecchini-Frenck-Zeidler [3]; see [3] for related discussion. Let $M = T^n \# \Sigma$, where Σ is an exotic sphere of dimension n . Consider g a uniformly Euclidean metric on M with an isolated singularity at a point in Σ and nonnegative scalar curvature on the smooth part. Then the metric measure space (M, g) induced by g does not remember the original smooth structure, and hence it is isometric to the flat torus T^n , which can be thought as the connected sum of the torus with the

standard sphere.

Example 2: Let $M = T^n$ and $T^k \subset M$ a linear sub-torus where $n - k \geq 4$. If g is a uniformly Euclidean metric on M with singular set $S = T^k$ and nonnegative scalar curvature on the smooth part. Then by Gromov-Lawson [15], $M \setminus S$ is Λ^2 -enlargeable, and hence cannot carry complete metrics of positive scalar metric. It follows by the conformal blowup method as in the proof of Theorem 1.2, g must be Ricci flat on the smooth part. Then by Theorem 1.6, g extends smoothly over S . This example illustrates the situation where our result still applies even if the singular set is not contained in a topological ball. See [15, Example 6.9] for more examples.

Example 3: Consider a closed Schoen-Yau-Schick (SYS) manifold M of dimension $4 \leq n \leq 7$ (here we use the definition given by Shi-Wang-Wu-Zhu [33]). By the very recent work of Shi-Wang-Wu-Zhu [33, Corollary 1.9], a uniformly Euclidean metric with isolated singularity on M and nonnegative scalar curvature on the smooth part must be Ricci flat on the smooth part. On the other hand, a SYS manifold has infinite fundamental group. By Theorem 1.6, the metric extends smoothly, in other words, isolated singularity is removable. Similarly, by Cecchini-Frenck-Zeidler [3, Theorem C and Remark 1.9], a uniformly Euclidean metric with isolated singularity on a closed enlargeable manifold must be Ricci flat on the smooth part. Then because a closed enlargeable manifold has infinite fundamental group, again the singularity is removable.

APPENDIX A. PROOF OF THEOREM 2.6

The main ingredients are Bochner's formula, the maximal principle, cut-off functions and the Laplacian comparison.

Proof of Theorem 2.6. : Let $h = \log u$. Then $\nabla h = \frac{\nabla u}{u}$, $\Delta h = \frac{\Delta u}{u} - \left| \frac{\nabla u}{u} \right|^2 = \frac{K(u)}{u} - |\nabla h|^2$. By the Bochner formula we have

$$\begin{aligned} \frac{1}{2} \Delta |\nabla h|^2 &= |\text{Hess } h|^2 + \langle \nabla h, \nabla(\Delta h) \rangle + \text{Ric}(\nabla h, \nabla h) \\ \text{(A.1)} \quad &\geq |\text{Hess } h|^2 + \langle \nabla h, \nabla \frac{K(u)}{u} \rangle - \langle \nabla h, \nabla(|\nabla h|^2) \rangle - (n-1)H^2 |\nabla h|^2. \end{aligned}$$

For the Hessian term one could use the Schwarz inequality

$$\text{(A.2)} \quad |\text{Hess } h|^2 \geq \frac{(\Delta h)^2}{n}.$$

Indeed, when $H = 0$, this estimate is enough. For $H > 0$, using this estimate one would get $(n(n-1))^{1/2}$ instead of $n-1$ for the coefficient of H in (2.8). To get the best constant for $H > 0$, note that (A.2) is only optimal when Hessian at all directions are same. Harmonic functions in the model spaces are radial functions, so their Hessian along the radial direction would be different from the spherical directions. Therefore one computes

the norm by separating the radial direction and uses Schwarz inequality in the spherical directions. Let $\{e_i\}$ be an orthonormal basis with $e_1 = \frac{\nabla h}{|\nabla h|}$ (we only need to prove the case $|\nabla h| \neq 0$), the potential radial direction, denote $h_{ij} = \text{Hess } h(e_i, e_j)$. We compute,

$$\begin{aligned}
|\text{Hess } h|^2 &= h_{11}^2 + 2 \sum_{j=2}^n h_{1j}^2 + \sum_{i,j \geq 2} h_{ij}^2 \\
&\geq h_{11}^2 + 2 \sum_{j=2}^n h_{1j}^2 + \frac{(\Delta h - h_{11})^2}{n-1} \\
&= h_{11}^2 + 2 \sum_{j=2}^n h_{1j}^2 + \frac{(|\nabla h|^2 + h_{11} - \frac{K(u)}{u})^2}{n-1} \\
&\geq \frac{n}{n-1} \sum_{j=1}^n h_{1j}^2 + \frac{1}{n-1} |\nabla h|^4 + \frac{2}{n-1} h_{11} |\nabla h|^2 - \frac{2}{n-1} \frac{K(u)}{u} (h_{11} + |\nabla h|^2).
\end{aligned}$$

Now $h_{1j} = \frac{1}{|\nabla h|} \langle \nabla_{e_j} \nabla h, \nabla h \rangle = \frac{1}{2|\nabla h|} e_j(|\nabla h|^2)$ and $h_{11} = \frac{1}{2|\nabla h|^2} \langle \nabla |\nabla h|^2, \nabla h \rangle$. Therefore (A.3)

$$|\text{Hess } h|^2 \geq \frac{n}{4(n-1)} \frac{|\nabla |\nabla h|^2|^2}{|\nabla h|^2} + \frac{|\nabla h|^4 + \langle \nabla |\nabla h|^2, \nabla h \rangle}{n-1} - \frac{2}{n-1} \frac{K(u)}{u} (h_{11} + |\nabla h|^2).$$

Note that when $K(u) = 0$, i.e. u is harmonic, then equality holds if and only if $\text{Hess } h$ are same on the level set of h , i.e.

$$\text{Hess } h = -\frac{|\nabla h|^2}{n-1} \left(g - \frac{1}{|\nabla h|^2} dh \otimes dh \right).$$

By plugging (A.3) into (A.1) we get

$$\begin{aligned}
\frac{1}{2} \Delta |\nabla h|^2 &\geq \frac{n}{4(n-1)} \frac{|\nabla |\nabla h|^2|^2}{|\nabla h|^2} + \frac{|\nabla h|^4}{n-1} - \frac{n-2 + \frac{K(u)}{u|\nabla h|^2}}{n-1} \langle \nabla |\nabla h|^2, \nabla h \rangle \\
\text{(A.4)} \quad &+ (K' - \frac{n+1}{n-1} \frac{K}{u}) |\nabla h|^2 - (n-1) H^2 |\nabla h|^2.
\end{aligned}$$

If $|\nabla h|^2$ achieves a local maximum inside $B(q, 2R)$ then we are done. Assume $|\nabla h|(q_0)$ is the maximum for some $q_0 \in B(q, 2R)$, then $\nabla |\nabla h|^2(q_0) = 0$, $\Delta |\nabla h|^2(q_0) \leq 0$. Plug these into (A.4) gives

$$0 \geq \frac{|\nabla h|^4}{n-1} + (K' - \frac{n+1}{n-1} \frac{K}{u}) |\nabla h|^2 - (n-1) H^2 |\nabla h|^2.$$

Hence $|\nabla h|^2 \leq (n-1)^2 H^2 + (n-1) \sup_{B(q, 2R)} |K' - \frac{n+1}{n-1} \frac{K}{u}|$.

In general the maximum could occur at the boundary and one has to use a cut-off function to force the maximum is achieved in the interior. Let $f : [0, 2R] \rightarrow [0, 1]$ be a

smooth function with

$$(A.5) \quad f|_{[0,R]} \equiv 1, \quad \text{supp } f \subset [0, 2R),$$

$$(A.6) \quad -cR^{-1}f^{1/2} \leq f' \leq 0,$$

$$(A.7) \quad |f''| \leq cR^{-2},$$

where $c > 0$ is a universal constant. Let $\phi : \overline{B(q, 2R)} \rightarrow [0, 1]$ with $\phi(x) = f(r(x))$, where $r(x) = d(x, q)$ is the distance function. Set $G = \phi|\nabla h|^2$. Then G is nonnegative on M and has compact support in $B(q, 2R)$. Therefore it achieves its maximum at some point $q_0 \in B(q, 2R)$. We can assume $r(q_0) \in [R, 2R)$ since if $q_0 \in B(q, R)$, then $|\nabla h|^2$ achieves maximal at q_0 on $B(q, R)$ and previous argument applies.

If q_0 is not a cut point of q then ϕ is smooth at q_0 and we have

$$(A.8) \quad \Delta G(q_0) \leq 0, \quad \nabla G(q_0) = 0.$$

At the smooth point of r , $\nabla G = \nabla\phi|\nabla h|^2 + \phi\nabla|\nabla h|^2$,

$$\Delta G = \Delta\phi|\nabla h|^2 + 2\langle\nabla\phi, \nabla|\nabla h|^2\rangle + \phi\Delta|\nabla h|^2.$$

Using (A.4), (A.8) and express $|\nabla h|^2, \nabla|\nabla h|^2$ in terms of $G, \nabla G$ we get, At the maximal point q_0 of G ,

$$(A.9) \quad \begin{aligned} 0 \geq \Delta G(q_0) &\geq \frac{\Delta\phi}{\phi}G - 2\frac{|\nabla\phi|^2}{\phi^2}G + \frac{n}{2(n-1)}\frac{|\nabla\phi|^2}{\phi^2}G + \frac{2}{n-1}\frac{G^2}{\phi} \\ &+ \frac{2(n-2 + \frac{K(u)}{u|\nabla h|^2})}{n-1}\langle\nabla h, \nabla\phi\rangle\frac{G}{\phi} + 2(K' - \frac{n+1}{n-1}\frac{K}{u})G - 2(n-1)H^2G. \end{aligned}$$

Since $\phi(x) = f(r(x))$ is a radial function, $|\nabla\phi| = |f'|$, by (A.6),

$$\langle\nabla h, \nabla\phi\rangle \geq -|\nabla h||\nabla\phi| = -G^{\frac{1}{2}}\frac{|\nabla\phi|}{\phi^{1/2}} \geq -G^{\frac{1}{2}}\frac{c}{R}.$$

Also $\Delta\phi = f'\Delta r + f''$. Since $f' \leq 0$, by the Laplacian comparison theorem and (A.7)

$$\Delta\phi \geq f'\Delta_H r - cR^{-2},$$

where $\Delta_H r = (n-1)H \coth(Hr)$, which is $\leq (n-1)H \coth(HR)$ on $[R, 2R]$. Hence

$$\begin{aligned} \Delta\phi &\geq -cR^{-1}((n-1)H \coth(HR) + R^{-1}) \\ &\geq -cR^{-1}[(n-1)(2R^{-1} + 4H) + R^{-1}] \end{aligned}$$

Multiply (A.9) by $\frac{(n-1)\phi}{G}$ and plug these in, we get

$$\begin{aligned} 0 &\geq 2G - 2(n-2 + |\frac{K(u)}{u|\nabla h|^2}|)\frac{c}{R}G^{\frac{1}{2}} \\ &\quad - \frac{c}{R}\left(\frac{(n-1)(2n-1)}{R} + 4(n-1)^2H + \left(\frac{3n}{2} - 2\right)\frac{c}{R}\right) \end{aligned}$$

$$-2(n-1) \left| K' - \frac{n+1}{n-1} \frac{K}{u} \right| - 2(n-1)^2 H^2.$$

Now if $|\nabla h|^2 \leq \frac{|K(u)|}{u}$, we are done. So we can assume $-\frac{|K(u)|}{u|\nabla h|^2} \geq -1$. In this case we have

$$\begin{aligned} 0 \geq & 2G - 2(n-1) \frac{c}{R} G^{\frac{1}{2}} - \frac{c}{R} \left(\frac{(n-1)(2n-1)}{R} + 4(n-1)^2 H + \left(\frac{3n}{2} - 2 \right) \frac{c}{R} \right) \\ & - 2(n-1) \left| K' - \frac{n+1}{n-1} \frac{K}{u} \right| - 2(n-1)^2 H^2. \end{aligned}$$

Solving this quadratic inequality gives

$$(G(q_0))^{\frac{1}{2}} \leq (n-1)H + C_1(n)R^{-1} + C_2(n, H)R^{-1/2} + \left((n-1) \left| K' - \frac{n+1}{n-1} \frac{K}{u} \right| \right)^{1/2}.$$

Therefore

$$\begin{aligned} \sup_{B(q,R)} |\nabla h| &= \sup_{B(q,R)} G^{1/2} \leq \sup_{B(q,2R)} G^{1/2} \\ &\leq (n-1)H + C_1(n)R^{-1} + C_2(n, H)R^{-1/2} + \left[(n-1) \left| K' - \frac{n+1}{n-1} \frac{K}{u} \right| \right]^{1/2}. \end{aligned}$$

If q_0 is in the cut locus of q , use the upper barrier $r_{q_0, \epsilon}(x)$ for $r(x)$ and let $\epsilon \rightarrow 0$ gives the same estimate. \square

REFERENCES

- [1] Luigi Ambrosio, Andrea Mondino, and Giuseppe Savaré. Nonlinear diffusion equations and curvature conditions in metric measure spaces. *Mem. Amer. Math. Soc.*, 262(1270):v+121, 2019.
- [2] Jérôme Bertrand, Christian Ketterer, Ilaria Mondello, and Thomas Richard. Stratified spaces and synthetic Ricci curvature bounds. *Ann. Inst. Fourier (Grenoble)*, 71(1):123–173, 2021.
- [3] Simone Cecchini, Georg Frencck, and Rudolf Zeidler. Positive scalar curvature with point singularities. *arXiv e-prints*, page arXiv:2407.20163, July 2024.
- [4] Jeff Cheeger. *Degeneration of Riemannian metrics under Ricci curvature bounds*. Lezioni Fermiane. [Fermi Lectures]. Scuola Normale Superiore, Pisa, 2001.
- [5] Jeff Cheeger and Detlef Gromoll. The splitting theorem for manifolds of nonnegative Ricci curvature. *J. Diff. Geom.*, 6:119–128, 1971.
- [6] Jeff Cheeger and Detlef Gromoll. On the structure of complete manifolds of nonnegative curvature. *Ann. Math.*, 96(3):413–443, 1972.
- [7] S. Y. Cheng and S. T. Yau. Differential equations on Riemannian manifolds and their geometric applications. *Comm. Pure Appl. Math.*, 28(3):333–354, 1975.
- [8] Otis Chodosh and Chao Li. Generalized soap bubbles and the topology of manifolds with positive scalar curvature. *Ann. of Math. (2)*, 199(2):707–740, 2024.
- [9] Xianzhe Dai, Yukai Sun, and Changliang Wang. Positive mass theorem for asymptotically flat manifolds with isolated conical singularities. *Sci. China Math.*, 10.1007/s11425-024-2325-6:1–16, 2024.
- [10] Xianzhe Dai, Yukai Sun, and Changliang Wang. Positive scalar curvature and isolated conical singularity. *arXiv:2412.02941*, 2024.
- [11] Matthias Erbar, Kazumasa Kuwada, and Karl-Theodor Sturm. On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces. *Invent. Math.*, 201(3):993–1071, 2015.
- [12] Nicola Gigli. An overview of the proof of the splitting theorem in spaces with non-negative Ricci curvature. *Anal. Geom. Metr. Spaces*, 2(1):169–213, 2014.

- [13] Nicola Gigli. On the differential structure of metric measure spaces and applications. *Mem. Amer. Math. Soc.*, 236(1113):vi+91, 2015.
- [14] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [15] Mikhael Gromov and H. Blaine Lawson, Jr. The classification of simply connected manifolds of positive scalar curvature. *Ann. of Math. (2)*, 111(3):423–434, 1980.
- [16] Mikhael Gromov and H. Blaine Lawson, Jr. Spin and scalar curvature in the presence of a fundamental group. I. *Ann. of Math. (2)*, 111(2):209–230, 1980.
- [17] Mikhael Gromov and H. Blaine Lawson, Jr. Positive scalar curvature and the Dirac operator on complete Riemannian manifolds. *Inst. Hautes Études Sci. Publ. Math.*, (58):83–196, 1983.
- [18] Misha Gromov. Dirac and Plateau billiards in domains with corners. *Cent. Eur. J. Math.*, 12:1109–1156, 2014.
- [19] Shouhei Honda. Bakry-émery conditions on almost smooth metric measure spaces. *Anal. Geom. Metr. Spaces*, 6(1):129–145, 2018.
- [20] Wenshuai Jiang, Weimin Sheng, and Huaiyu Zhang. Removable singularity of positive mass theorem with continuous metrics. *Math. Z.*, 302(2):839–874, 2022.
- [21] Demetre Kazaras. Desingularizing positive scalar curvature 4-manifolds. *Math. Ann.* <https://doi.org/10.1007/s00208-024-02829-5>, 2024.
- [22] Jerry L. Kazdan. Deformation to positive scalar curvature on complete manifolds. *Math. Ann.*, 261(2):227–234, 1982.
- [23] Jerry L. Kazdan and F. W. Warner. Scalar curvature and conformal deformation of Riemannian structure. *J. Differential Geometry*, 10:113–134, 1975.
- [24] Martin Lesourd, Ryan Unger, and Shing-Tung Yau. The positive mass theorem with arbitrary ends. *J. Differential Geom.*, 128(1):257–293, 2024.
- [25] Chao Li and Christos Mantoulidis. Positive scalar curvature with skeleton singularities. *Math. Ann.*, 374:99–131, 2019.
- [26] Peter Li and Jiaping Wang. Complete manifolds with positive spectrum. II. *J. Differential Geom.*, 62(1):143–162, 2002.
- [27] W. Littman, G. Stampacchia, and H. F. Weinberger. Regular points for elliptic equations with discontinuous coefficients. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)*, 17:43–77, 1963.
- [28] Ilaria Mondello. The Yamabe problem on stratified spaces. *HAL Id : tel-01204671*, 2015.
- [29] Andrea Mondino and Guofang Wei. On the universal cover and the fundamental group of an RCD $^*(K, N)$ -space. *J. Reine Angew. Math.*, 753:211–237, 2019.
- [30] R. Schoen and S. T. Yau. On the structure of manifolds with positive scalar curvature. *Manuscripta Math.*, 28(1-3):159–183, 1979.
- [31] Richard Schoen. Conformal deformation of a Riemannian metric to constant scalar curvature. *J. Differential Geom.*, 20(2):479–495, 1984.
- [32] Richard M. Schoen. Variational theory for the total scalar curvature functional for Riemannian metrics and related topics. In *Topics in calculus of variations (Montecatini Terme, 1987)*, volume 1365 of *Lecture Notes in Math.*, pages 120–154. Springer, Berlin, 1989.
- [33] Yuguang Shi, Jian Wang, Runzhang Wu, and Jintian Zhu. On open manifolds admitting no complete metric with positive scalar curvature. *arXiv:2404.01660v2*, to appear at *Annales de L’Institut Fourier*, 2024.
- [34] Yuguang Shi, Wenlong Wang, Guodong Wei, and Jintian Zhu. On the fill-in of nonnegative scalar curvature metrics. *Math. Ann.*, 379(1-2):235–270, 2021.
- [35] P. D. Smith and Deane Yang. Removing point singularities of Riemannian manifolds. *Trans. Amer. Math. Soc.*, 333(1):203–219, 1992.

- [36] Stephan Stolz. Simply connected manifolds of positive scalar curvature. *Ann. of Math. (2)*, 136(3):511–540, 1992.
- [37] Gábor Székelyhidi. Singular Kähler-Einstein metrics and RCD spaces. *arXiv:2408.10747*, 2024.
- [38] Jinmin Wang and Zhizhang Xie. Scalar curvature rigidity of spheres with subsets removed and L^∞ metrics. *arXiv e-prints*, page arXiv:2407.21312, July 2024.
- [39] Xiangsheng Wang and Weiping Zhang. On the generalized Geroch conjecture for complete spin manifolds. *Chin. Ann. Math. Ser. B*, 43(6):1143–1146, 2022.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA CA93106, USA
Email address: dai@math.ucsb.edu

SCHOOL OF MATHEMATICAL SCIENCES AND INSTITUTE FOR ADVANCED STUDY, KEY LABORATORY OF INTELLIGENT COMPUTING AND APPLICATIONS(MINISTRY OF EDUCATION), TONGJI UNIVERSITY, SHANGHAI 200092, CHINA

Email address: wangchl@tongji.edu.cn

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IA 52242-1419 USA
Email address: lihe-wang@uiowa.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA CA93106, USA
Email address: wei@math.ucsb.edu