

Eikonal 2.0: General Solutions for Modified Eikonal Equations

Iryna YEHORCHENKO

Institute of Mathematics of NAS Ukraine, 3 Tereshchenkivs'ka Str., Kyiv-4, Ukraine

E-mail: iyegorch@imath.kiev.ua

Institute of Mathematics, PAN, Poland

iyehorchenko@impan.pl

Abstract

We find implicit general solutions for modified eikonal equations $u_a u_a = F(u_t)$, where lower indices at dependent variables designate derivatives, $a = 1, 2, \dots, n$, and summation is implied over the repeated indices.

We will consider general solutions and symmetries of these equations, and relations of these equations to reduction procedures of higher-order PDE.

1 Introduction

This paper is a prolongation of the paper [1], one in the series intended as comprehensive study of the eikonal equations and systems that include them.

We will consider here a class of nonlinear first-order partial differential equations (PDE) in the following form:

$$u_a u_a = F(u_t), \quad (1)$$

where in general we have one real-valued dependent variable $u = u(x_0, x_1, \dots, x_n)$, a time variable x_0 and n space variables x_1, \dots, x_n . Thus class includes both the standard eikonal equation and the Hamilton-Jacobi equation. Here we do not consider cases when $F(u_t) = \text{const}$.

I would like to point out that u and other dependent variable take values in a real space, as the complex-valued dependent variables require different treatment and a special study.

The intended audience of this paper is wider than "the symmetry analysis people", so many detailed arrangements and calculations that are obvious for the above group are written explicitly. The results in this paper I believe at the moment of writing to be new will be adduced at the end. I mentioned only the literature what was immediately relevant for the paper. Further we will mostly adhere to notations introduced in [1].

Group classification of the class

$$u_a u_a = F(t, u, u_t), \quad (2)$$

the class (1) we consider being its subclass, was performed in [2]. We adduce here this classification.

	F	Basis Symmetry Operators
0	$F(t, u, u_t)$	∂_a, J_{ab}
1	$e^{\delta t} f(u, u_t), \delta \in \{0; 1\}$	$\partial_a, J_{ab}, 2\partial_t - \delta x_a \partial_a$
2	$e^u h(u_t)$	$\partial_a, J_{ab}, \partial_t, 2\partial_u - x_a \partial_a$
3	$ u ^{2-\delta} h(u_t), \delta \neq 2$	$\partial_a, J_{ab}, \partial_t, 2t\partial_t + 2u\partial_u + \delta x_a \partial_a$
4	$h(u_t)$	$\partial_a, J_{ab}, \partial_t, \partial_u, D$
5	e^{u_t}	$\partial_a, J_{ab}, \partial_t, \partial_u, D, x_a \partial_a - 2t\partial_u$
6	$ u_t ^\beta, \beta \neq 0, 1, 2$	$\partial_a, J_{ab}, \partial_t, \partial_u, D, (\beta - 2)x_a \partial_a - 2u\partial_u$
7	u_t^2	$2g_{\mu\nu}c^\mu(u)x_\nu x_\varkappa \partial_\varkappa - c^\varkappa(u)g_{\mu\nu}x_\mu x_\nu \partial_\varkappa + g_{\mu\nu}b^{\mu\varkappa}(u)x_\nu \partial_\varkappa +$ $+d(u)x_\varkappa \partial_\varkappa + a^\varkappa(u)\partial_\varkappa + \eta(u)\partial_u$
8	$\varepsilon_2 u_t^2 + \varepsilon_1$	$\partial_a, J_{ab}, \partial_t, \partial_u, D, J_{ua} = u\partial_a + \varepsilon_1 x_a \partial_u, J_{ta} = t\partial_a + \varepsilon_2 x_a \partial_t,$ $J_{ut} = u\partial_t - \varepsilon_1 \varepsilon_2 t \partial_u, K_a = 2x_a D - s^2 \partial_a, K_u = 2uD +$ $\varepsilon_1 s^2 \partial_u, K_t = 2tD + \varepsilon_2 s^2 \partial_t, \text{ where } s^2 = x_a x_a - \varepsilon_1 u^2 - \varepsilon_2 t^2$
9	$\varepsilon_2 e^u u_t^2 + \varepsilon_1$	$\partial_a, J_{ab}, \partial_t, t\partial_t + 2\partial_u, (t^2 - 4\varepsilon_1 \varepsilon_2 e^u)\partial_t + 4t\partial_u$
10	$\cos^{-2} u u_t^2 + 1$	$\partial_a, J_{ab}, \partial_t, \cos t \tan u \partial_t - \sin t \partial_u, \sin t \tan u \partial_t + \cos t \partial_u$
11	$\pm(\cos^{-2} u u_t^2 - 1)$	$\partial_a, J_{ab}, \partial_t, \cosh t \tan u \partial_t + \sinh t \partial_u, \sinh t \tan u \partial_t +$ $\cosh t \partial_u$
12	$\cosh^{-2} u u_t^2 + 1$	$\partial_a, J_{ab}, \partial_t, \cosh t \tanh u \partial_t - \sinh t \partial_u, \sinh t \tanh u \partial_t -$ $\cosh t \partial_u$

$F = F(t, u, u_t)$, $f = f(u, u_t)$, $h = h(u_t)$ are arbitrary smooth functions of their arguments, δ is constant, $\varepsilon_1, \varepsilon_2 = \pm 1$, $(\varepsilon_1, \varepsilon_2) \neq (-1, -1)$, $J_{ab} = x_a \partial_b - x_b \partial_a$, $D = t\partial_t + u\partial_u + x_a \partial_a$. In the case 7, $g_{\mu\nu}$ is metric tensor of Minkowsky space $\mathbb{R}^{1,n}$, i.e. $g_{00} = -g_{11} = -g_{22} = -g_{33} = 1$, $g_{\mu\nu} = 0$, $\mu \neq \nu$; $c^\mu, b^{\mu\varkappa}, d, a^\varkappa, \eta$ are arbitrary smooth functions of the variable u (for operators of such form to form an algebra, it is necessary to require that these functions were infinitely differentiable or real analytical); the indices μ, ν and \varkappa take values from 0 to 3.

Let us note that the known Hamilton-Jacobi and eikonal equations, and the relativistic Hamilton equation are selected as representatives of the classes of equivalent equations having the widest symmetry.

2 General Solutions

In consideration of the general solutions for the eikonal equations, we will follow the idea, given in [3], where also general solutions for the standard eikonal equations $u_\mu u_\mu = 1$ and $u_\mu u_\mu = 0$ for $n = 3$ are adduced.

Here we generalize the relevant formulae the modified eikonal equations and for the arbitrary number of space variables.

We used ideas for finding general solutions outlined e.g. in [3]. The main tool of this approach is to use contact transformations that are nonlocal transformations involving dependent and independent variables, and also first derivatives of dependent variables.

We will apply the following format of such contact transformations, the indices at independent variables x'_μ , x_μ are their indices indicating their respective numbers, and the indices at dependent variables u'_μ , u_μ designate their derivatives with respect to the relevant independent variables x'_μ or x_μ .

$$\begin{aligned}x'_\mu &= \varphi_\mu(x_\mu, u, u_\mu), \\u' &= \Phi(x_\mu, u, u_\mu), \\u'_\mu &= \Phi_\mu(x_\mu, u, u_\mu).\end{aligned}\tag{3}$$

Here x'_μ , u' are new transformed variables, u'_μ are transformed first derivatives, φ_μ , Φ and Φ_μ are sufficiently smooth functions on their variables. Note that indices at Φ_μ are just numerating indices, not derivatives.

We can apply the partial derivative operators to (1) and conclude linear dependence of the resulting equations considered as linear equations for the first derivatives of u , and that the rank of the matrix of the second derivatives of the function u may take values from 0 to n , where n is the number of space variables.

The equation $u_a u_a = F(u_t)$ has real solutions for which the rank of the matrix of the second derivatives is zero, all first derivatives u_μ are constants, and u is linearly dependent on x_μ :

$$u = c_\mu x_\mu + c,$$

where c_μ and c are arbitrary constants such that $c_a c_a = F(c_0)$, when $F(c_0)$ is non-negative.

In the case when the rank of U is equal to n , we can apply the contact transformations

$$\begin{aligned}H &= x_a u_{x_a} - u, & y_a &= u_{x_a}, & t &= y_0; \\H_{y_a} &= x_a, & H_{y_0} &= -u_t;\end{aligned}$$

and obtain the following equation for the new function H :

$$y_a y_a = F(H_{y_0}).$$

We can linearize it with respect to H_{y_0} : $H_{y_0} = \Phi(y_b y_b)$, where Φ is a inverse function to F . Its solution is

$$H = y_0 \Phi(y_a y_a) - \Psi(y_d),$$

where $\Psi(y_d)$ is a sufficiently smooth arbitrary function. Further we use parameters τ_d instead of y_d . An n -parameter general solution ($\text{rank}(U) = n$):

$$\begin{aligned}u &= -x_a \tau_a + t \Phi(\tau_b \tau_b) + \Psi(\tau_d), \\x_a - t \Phi_{\tau_a}(\tau_b \tau_b) - \Psi_{\tau_a}(\tau_d) &= 0\end{aligned}\tag{4}$$

where $\Psi(\tau_d)$ is a sufficiently smooth arbitrary function on n parameters τ_d .

The formula (4) includes arbitrary functions, so all solutions (4) cannot be equivalent to symmetry solutions.

In general (not always), formulas for the general solutions give wider classes of solutions than the Lie symmetry method. As to the zero-rank general solution, all such solutions are symmetry solutions.

General solutions depending on k parameters may be taken as follows:

$$\begin{aligned} u &= -x_b\tau_b + t\Phi(\tau_d\tau_d + w_m w_m) + w_m x_{k+m} + \Psi(\tau_d), \\ x_b - t\Phi_{\tau_b}(\tau_d\tau_d + w_m w_m) - w_m\tau_b x_{k+m} - \Psi_{\tau_b}(\tau_d) &= 0, \end{aligned}$$

where $\Psi(\tau_d)$ is a sufficiently smooth arbitrary function on n parameters τ_d .

3 Examples for Specific Equations

We will write an example using one of the equations from the group classification table:

$$u_a u_a = \exp(u_t),$$

or

$$u_t = \ln(u_a u_a). \tag{5}$$

Equation (5) has real solutions for which the rank of the matrix of the second derivatives is zero, all first derivatives u_a and u_t are constants, and u is linearly dependent on x_μ :

$$u = c_\mu x_\mu + c,$$

where c_μ and c are arbitrary constants such that $c_a c_a = \exp(c_0)$.

An n -parameter general solution ($\text{rank}(U) = n$):

$$\begin{aligned} u &= -x_a\tau_a + t \ln(\tau_b\tau_b) + \Psi(\tau_d), \\ x_a - t \frac{2\tau_a}{\tau_b\tau_b} - \Psi_{\tau_a}(\tau_d) &= 0, \end{aligned}$$

where $\Psi(\tau_d)$ is a sufficiently smooth arbitrary function on n parameters τ_d .

General solutions depending on k parameters may be taken as follows:

$$\begin{aligned} u &= -x_b\tau_b + t \ln(\tau_d\tau_d + w_m w_m) + w_m x_{k+m} + \Psi(\tau_d), \\ x_b - t \frac{2\tau_b}{\tau_c\tau_c} - w_m\tau_b x_{k+m} - \Psi_{\tau_b}(\tau_d) &= 0, \end{aligned}$$

where $\Psi(\tau_d)$ is a sufficiently smooth arbitrary function on n parameters τ_d .

4 New Results, Conclusions and Further Work

The new results of this paper are general solutions of modified eikonal equations for arbitrary number of space variables.

There are multiple papers searching for numerical solutions of the boundary problems for eikonal and similar first-order partial differential equations. On my opinion, it would be interesting to look for these using the general solutions.

Further work will include study of the coupled modified eikonal equations in higher dimensions, relevant conditional and hidden symmetries, and looking for higher-order equations with the same or similar symmetry properties as modified eikonal equations.

5 Acknowledgements

The first and foremost my acknowledgements go to the Armed Forces of Ukraine and to the Territorial Defence Forces of Ukrainian Regions due to whom I am alive and is able to work.

Please remember that Russia is an aggressor country and still plans to kill all Ukrainians not going to be their subordinates, and Russian scientists contribute to killings despite words about peace from a tiny portion of them; if even by trying to persuade naïve people that Russia is a civilised country.

This research was mostly completed during my work at the Institute of Mathematics of the Polish Academy of Sciences with the grant of the Narodowe Centrum Nauki (Poland), Grant No.2017/26/A/ST1/00189. I would like to thank the Institute of Mathematics of the Polish Academy of Sciences for their hospitality and grant support, to the National Academy of Sciences of the USA and the National Centre of Science of Poland for their grant support.

Further work was supported by a grant from the Simons Foundation (1290607, I.A.Y).

References

- [1] Yehorchenko I.A., Eikonal 1.0, arXiv:2212.09914
- [2] Popovych R.O., Yehorchenko I.A., Group Classification of Generalized Eikonal Equations. Ukrainian Mathematical Journal 53, 1841—1850 (2001).
- [3] Fushchych W.I., Zhdanov R.Z., Revenko I.V., Compatibility and solution of non-linear d'Alembert-Hamilton equations, Preprint 90.39, Institute of Mathematics of Acad.Sci. Ukr. SSR, Kyiv, 1990, 65 p.