FRACTIONAL SOBOLEV-CHOQUARD CRITICAL SYSTEMS WITH HARDY TERM AND WEIGHTED SINGULARITIES

RONALDO B. ASSUNÇÃO, OLÍMPIO H. MIYAGAKI, AND RAFAELLA F. S. SIQUEIRA

ABSTRACT. In this paper, we consider a fractional *p*-Laplacian system of equations in the entire space \mathbb{R}^N with doubly critical singular nonlinearities involving a local critical Sobolev term together with a nonlocal Choquard critical term; the problem also includes a homogeneous singular Hardy term; moreover, all the nonlinearities involve singular critical weights. To prove the main result we use a refinement of Sobolev inequality that is related to Morrey space because our problem involves doubly critical exponentss and a version of the Caffarelli-Kohn-Nirenberg inequality. With the help of these results, we provide sufficient conditions under which a weak nontrivial solution to the problem exists via variational methods.

CONTENTS

1.	Introduction and main results	1
2.	Historical background	5
3.	Existence of solutions for auxiliary minimization problems	7
4.	Existence of Palais-Smale sequence	14
5.	Proof of Theorem 1	18
References		22

1. INTRODUCTION AND MAIN RESULTS

In the present work, we consider the following fractional *p*-Laplacian system of equations in the entire space \mathbb{R}^N featuring doubly critical nonlinearities, involving a local critical Sobolev term together with a nonlocal Choquard critical term; the problem also includes a homogeneous Hardy term; additionally, all terms have critical singular weights. More precisely, we deal with the problem

(1)

$$\begin{cases} (-\Delta)_{p,\theta}^{s}u - \gamma_{1}\frac{|u|^{p-2}u}{|x|^{sp+\theta}} = \left[I_{\mu} * F(\cdot, u)\right](x)f(x, u) + \frac{|u|^{p_{s}^{*}(\beta,\theta)-2}u}{|x|^{\beta}} + \frac{\eta a}{a+b}\frac{|u|^{a-2}u|v|^{b}}{|x|^{\beta}} \\ (-\Delta)_{p,\theta}^{s}v - \gamma_{2}\frac{|v|^{p-2}v}{|x|^{sp+\theta}} = \left[I_{\mu} * F(\cdot, v)\right](x)f(x, v) + \frac{|v|^{p_{s}^{*}(\beta,\theta)-2}v}{|x|^{\beta}} + \frac{\eta b}{a+b}\frac{|u|^{a}|v|^{b-2}v}{|x|^{\beta}} \end{cases}$$

Date: Belo Horizonte, December 16, 2024.

Key words and phrases. Systems of fractional p-Laplacian operator, doubly critical singular problem, variational methods, weighted Sobolev and Morrey spaces, Caffarelli-Kohn-Nirenberg inequality.

Rafaella F. S. Siqueira was partially supported by CNPq/Brazil.

²⁰¹⁰ Mathematics Subject Classification. Primary: 35B33; 35J92; 35R11. Secondary: 35A23, 35B38, 35J20.

Olímpio H. Miyagaki was supported by Grant 2022/16407-1—São Paulo Research Foundation (FAPESP) and Grant 303256/2022-2—CNPq/Brazil.

where 0 < s < 1; $0 < \alpha$, $\beta < sp + \theta < N$; $0 < \mu < N$; $2\delta + \mu < N$; $\eta \in \mathbb{R}^+$; $\gamma_1, \gamma_2 < \gamma_H$ with the best fractional Hardy constant γ_H to be defined below (without lost of generality, to simplify the notation we can consider the only parameter $\gamma = \gamma_1 = \gamma_2$); the Hardy-Sobolev and Stein-Weiss upper critical fractional exponents (this latter also called Hardy-Littlewood-Sobolev upper critical exponent) are respectively defined by

$$p_s^*(\beta, \theta) = \frac{p(N-\beta)}{N-sp-\theta}$$
 and $p_s^{\sharp}(\delta, \theta, \mu) = \frac{p(N-\delta-\mu/2)}{N-sp-\theta};$

to simplify the notation, we write $p_s^*(\beta, \theta) = p_s^*$ and $p_s^{\sharp}(\delta, \theta, \mu) = p_s^{\sharp}$, additionally $a + b = p_s^*(\beta, \theta)$. Moreover, $I_{\mu}(x) = |x|^{-\mu}$ is the Riesz potential of order μ ; the functions $f, F \colon \mathbb{R}^N \times \mathbb{R} \to \mathbb{R}$ are respectively defined by

(2)
$$f(x,t) = f_{\delta,\theta,\mu}(x,t) = \frac{|t|^{p_s^{\sharp}-2}t}{|x|^{\delta}}$$
 and $F(x,t) = F_{\delta,\theta,\mu}(x,t) = \frac{|t|^{p_s^{\sharp}}}{|x|^{\delta}},$

that is, $F_{\delta,\theta,\mu}(x,t) = p_s^{\sharp} \int_0^{|t|} f_{\delta,\theta,\mu}(x,\tau) \,\mathrm{d}\tau$; and the term with convolution integral,

$$[I_{\mu} * F_{\delta,\theta,\mu}(\cdot, u)](x) \coloneqq \int_{\mathbb{R}^N} \frac{|u(y)|^{p_s^{\sharp}}}{|x-y|^{\mu}|y|^{\delta}} \,\mathrm{d}y\,,$$

is known as Choquard type nonlinearity.

Let us now introduce the spaces of functions that are meaningful to our considerations. Throughout this work, we denote the norm of the weighted Lebesgue space $L^p(\mathbb{R}^N, |x|^{-\lambda})$ by

$$\|u\|_{L^p(\mathbb{R}^N;|x|^{-\lambda})} \coloneqq \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{\lambda}} \,\mathrm{d}x\right)^{\frac{1}{p}}$$

for any $0 \leq \lambda < N$ and $1 \leq p < +\infty$.

We say that a Lebesgue measurable function $u \colon \mathbb{R}^N \to \mathbb{R}$ belongs to the weighted Morrey space $L_M^{p,\gamma+\lambda}(\mathbb{R}^N,|x|^{-\lambda})$ if

$$\|u\|_{L^{p,\gamma+\lambda}_{M}(\mathbb{R}^{n},|x|^{-\lambda})} \coloneqq \sup_{x \in \mathbb{R}^{N}, R \in \mathbb{R}_{+}} \left\{ \left(R^{\gamma+\lambda-N} \int_{B_{R}(x)} \frac{|u|^{p}}{|x|^{\lambda}} \mathrm{d}x \right)^{\frac{1}{p}} \right\} < +\infty,$$

where $1 \leq p < +\infty; \gamma, \lambda \in \mathbb{R}_+$, and $0 < \gamma + \lambda < N$.

Our concerns involve the homogeneous fractional Sobolev-Slobodeckij space $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ defined as the completion of the space $C_0^{\infty}(\mathbb{R}^N)$ with respect to the Gagliardo seminorm given by

$$u \mapsto [u]_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)} \coloneqq \left(\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^p}{|x|^{\theta_1} |x - y|^{N + sp} |y|^{\theta_2}} \mathrm{d}x \mathrm{d}y \right)^{\frac{1}{p}},$$

i.e., $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N) = \overline{C^{\infty}_0(\mathbb{R}^N)}^{[\cdot]}$. We can equip the homogeneous fractional Sobolev space $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ with the norm

$$\begin{aligned} \|u\|_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^{N})} &\coloneqq \left(\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x\mathrm{d}y - \gamma \int_{\mathbb{R}^{N}} \frac{|u|^{p}}{|x|^{sp+\theta}} \mathrm{d}x \right)^{\frac{1}{p}} \\ &\coloneqq \left([u]^{p}_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^{n})} - \gamma \|u\|^{p}_{L^{p}(\mathbb{R}^{N};|x|^{-sp-\theta})} \right)^{\frac{1}{p}}. \end{aligned}$$

Here, we assume that $\gamma < \gamma_H$, where the best fractional Hardy constant is defined by

$$\gamma_H \coloneqq \inf_{\substack{u \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^n) \\ u \neq 0}} \frac{[u]^p_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^n)}}{\|u\|^p_{L^p(\mathbb{R}^n; |x|^{-sp-\theta})}}.$$

This turns the space $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ into a Banach space; moreover, this space is uniformly convex; in particular, it is reflexive and separable.

For simplicity, hereafter we denote the Cartesian product space of two Banach spaces $W = \dot{W}^{s,p}_{\theta}(\mathbb{R}^N) \times \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$, endowed with the norm

$$||(u,v)||_W \coloneqq \left(||u||_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)}^p + ||v||_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)}^p \right)^{1/p}.$$

Intuitively, problem (1) is understood as showing the existence of a pair $(u, v) \in W$ such that

$$\begin{cases} (-\Delta)_{p,\theta}^{s}u - \gamma \frac{|u|^{p-2}u}{|x|^{sp+\theta}} = \left(\int_{\mathbb{R}^{N}} \frac{|u|^{p_{s}^{\sharp}}}{|x-y|^{\mu}|y|^{\delta}} \mathrm{d}y\right) \frac{|u|^{p_{s}^{\sharp}-2}u}{|x|^{\delta}} + \frac{|u|^{p_{s}^{\ast}-2}u}{|x|^{\beta}} + \frac{\eta a}{a+b} \frac{|u|^{a-2}u|v|^{b}}{|x|^{\beta}} + \frac{\eta a}{(a+b)^{2}} \frac{|u|^{a}}{|x|^{\beta}} + \frac{\eta a}{(a+b)^{2}} \frac{|u|^$$

where the fractional *p*-Laplacian operator is defined for $\theta = \theta_1 + \theta_2$, $x \in \mathbb{R}^N$, and any function $u \in C_0^{\infty}(\mathbb{R}^N)$, as

(3)
$$(-\Delta)_{p,\theta}^{s} u(x) \coloneqq \text{p.v.} \int_{\mathbb{R}^{N}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x|^{\theta_{1}} |x - y|^{N+sp} |y|^{\theta_{2}}} \,\mathrm{d}y \,,$$

and p.v. is the Cauchy's principal value. This operator is the prototype of nonlinear nonlocal elliptic operator and can also be defined on smooth functions by

$$(-\Delta)_{p,\theta}^s u(x) \coloneqq 2\lim_{\varepsilon \to 0} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))}{|x|^{\theta_1} |x - y|^{N+sp} |y|^{\theta_2}} \,\mathrm{d}y$$

This definition is consistent, up to a normalization constant $C = C(N, s, \theta)$, with the usual definition of the linear fractional Laplacian operator $(-\Delta)^s$ for p = 2 and $\theta = 0$.

Our main goal in this work is to show that problem (1) admits at least a weak solution, by which term we mean a function $(u, v) \in W$ such that

$$\begin{split} &\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(\phi_{1}(x) - \phi_{1}(y))}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x\mathrm{d}y \\ &+ \iint_{\mathbb{R}^{2N}} \frac{|v(x) - v(y)|^{p-2}(v(x) - v(y))(\phi_{2}(x) - \phi_{2}(y))}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x\mathrm{d}y \\ &- \gamma_{1} \int_{\mathbb{R}^{N}} \frac{|u|^{p-2}u\phi_{1}}{|x|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \frac{|v|^{p-2}v\phi_{2}}{|x|^{sp+\theta}} \mathrm{d}x \\ &= \iint_{\mathbb{R}^{2N}} \frac{|u(x)|^{p_{s}^{\sharp}-2}|u(y)|^{p_{s}^{\sharp}}u(x)\phi_{1}(x)}{|x|^{\delta}|x - y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y \\ &+ \iint_{\mathbb{R}^{2N}} \frac{|v(x)|^{p_{s}^{\sharp}-2}|v(y)|^{p_{s}^{\sharp}}v(x)\phi_{2}(x)}{|x|^{\delta}} \mathrm{d}x\mathrm{d}y \\ &+ \int_{\mathbb{R}^{N}} \frac{|u|^{p_{s}^{*}(\beta,\theta)-2}u\phi_{1}(x)}{|x|^{\beta}} \mathrm{d}x + \int_{\mathbb{R}^{N}} \frac{|v|^{p_{s}^{*}(\beta,\theta)-2}v\phi_{2}(x)}{|x|^{\beta}} \mathrm{d}x \end{split}$$

$$+\int_{\mathbb{R}^N}\frac{\eta a|u|^{a-2}u\phi_1|v|^b}{|x|^{\beta}}\mathrm{d}x+\int_{\mathbb{R}^N}\frac{\eta b|u|^a|v|^{b-2}v\phi_2}{|x|^{\beta}}\mathrm{d}x$$

for any pair of test functions $(\phi_1, \phi_2) \in W$. Now we define the energy functional $I: W \to \mathbb{R}$ by

$$I(u,v) = \frac{1}{p} \Big[\iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{|v(x) - v(y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \Big] - \frac{\gamma_{1}}{p} \int_{\mathbb{R}^{N}} \frac{|u|^{p}}{|x|^{sp+\theta}} \mathrm{d}x - \frac{\gamma_{2}}{p} \int_{\mathbb{R}^{N}} \frac{|v|^{p}}{|x|^{sp+\theta}} \mathrm{d}x - \frac{1}{2p_{s}^{\sharp}(\delta, \theta, \mu)} \Big[\iint_{\mathbb{R}^{2N}} \frac{|u(x)|^{p_{s}^{\sharp}}|u(y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x - y|^{\mu}|y|^{\delta}} \mathrm{d}x \mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{|v(x)|^{p_{s}^{\sharp}}|v(y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x - y|^{\mu}|y|^{\delta}} \mathrm{d}x \mathrm{d}y \Big] (4) \qquad - \frac{1}{p_{s}^{*}} \Big[\int_{\mathbb{R}^{N}} \frac{|u|^{p_{s}^{*}}}{|x|^{\beta}} \mathrm{d}x + \int_{\mathbb{R}^{N}} \frac{|v|^{p_{s}^{*}}}{|x|^{\beta}} \mathrm{d}x \Big] - \frac{1}{p_{s}^{*}} \int_{\mathbb{R}^{N}} \frac{\eta |u|^{a}|v|^{b}}{|x|^{\beta}} \mathrm{d}x.$$

For the parameters in the previously specified intervals, the energy functional I is well defined and is continuously differentiable, i.e., $I \in C^1(\dot{W}^{s,p}_{\theta}(\mathbb{R}^N);\mathbb{R})$; moreover, a nontrivial critical point of the energy functional I is a nontrivial weak solution to problem (1).

Theorem 1. Problem (1) has at least a nontrivial weak solution provided that 0 < s < 1; $0 < \alpha, \beta < sp + \theta < N$; $0 < \mu < N$; $a + b = p_s^*(\beta, \theta)$; $\eta \in \mathbb{R}^+$ and $\gamma_1, \gamma_2 < \gamma_H$.

In this work we also consider the following variants of problem (1), namely one problem with a Hardy potential and double Sobolev type nonlinearities,

(5)
$$\begin{cases} (-\Delta)_{p,\theta}^{s} u - \gamma_1 \frac{|u|^{p-2} u}{|x|^{sp+\theta}} = \sum_{k=1}^{k=2} \frac{|u|^{p_s^*-2} u}{|x|^{\beta}} + \frac{\eta a}{a+b} \frac{|u|^{a-2} u|v|^{b}}{|x|^{\beta_k}} \\ (-\Delta)_{p,\theta}^{s} v - \gamma_2 \frac{|v|^{p-2} v}{|x|^{sp+\theta}} = \sum_{k=1}^{k=2} \frac{|v|^{p_s^*-2} v}{|x|^{\beta_k}} + \frac{\eta b}{a+b} \frac{|u|^a |v|^{b-2} v}{|x|^{\beta_k}} \end{cases}$$

and another one with a Hardy potential and double Choquard type nonlinearities, (6)

$$\begin{cases} (-\Delta)_{p,\theta}^{s} u - \gamma_{1} \frac{|u|^{p-2} u}{|x|^{sp+\theta}} = \sum_{k=1}^{k=2} \left[I_{\mu_{k}} * F_{\delta,\theta,\mu_{k}}(\cdot,u) \right](x) f_{\delta,\theta,\mu_{k}}(x,u) + \frac{\eta a}{a+b} \frac{|u|^{a-2} u|v|^{b}}{|x|^{\beta_{k}}} \\ (-\Delta)_{p,\theta}^{s} v - \gamma_{2} \frac{|v|^{p-2} v}{|x|^{sp+\theta}} = \sum_{k=1}^{k=2} \left[I_{\mu_{k}} * F_{\delta,\theta,\mu_{k}}(\cdot,v) \right](x) f_{\delta,\theta,\mu_{k}}(x,v) + \frac{\eta b}{a+b} \frac{|u|^{a} |v|^{b-2} v}{|x|^{\beta_{k}}} \end{cases}$$

The notion of weak solution to problems (5) and (6) can be defined in the same way as that for problem (1), i.e., we multiply the differential equations by a pair of test functions and use a kind of integration by parts. Then we recognize these expressions as the derivatives of an energy functional which, under the appropriate hypotheses on the parameters, is continuously differentiable. This means that weak solutions to these problems are critical points of the appropriate energy functional. By adapting the method used in the proof of Theorem 1 we deduce the following result.

Theorem 2. Problems (5) and (6) have at least a nontrivial weak solution under similar assumptions as in Theorem 1, i.e., 0 < s < 1; $0 < \alpha_k$, $\beta_k < sp + \theta < N$; $0 < \mu_k < N$; $a + b = p_s^*(\beta_k, \theta)$; $\eta \in \mathbb{R}^+$ and $\gamma_1, \gamma_2 < \gamma_H$ for $k \in \{1, 2\}$.

Notation. For $\rho \in \mathbb{R}_+$, we define $B_{\rho}(x) \coloneqq \{y \in \mathbb{R}^N : |x-y| < \rho\}$, the open ball centered at x with radius ρ . The constant ω_N denotes the volume of the unit ball in \mathbb{R}^N . The arrows \rightarrow and \rightarrow denote the strong convergence and the weak convergence, respectively. Given the functions $f, g \colon \mathbb{R}^N \to \mathbb{R}$, we recall that f = O(g) if there is a constant $C \in \mathbb{R}_+$ such that $|f(x)| \leq C|g(x)|$ for all $x \in \mathbb{R}^N$; and f = o(g) as $x \to x_0$ if $\lim_{x \to x_0} |f(x)|/|g(x)| = 0$. The pair r and r' denote Hölder conjugate exponents, i.e., 1/r + 1/r' = 1 or r + r' = rr'. The positive and negative parts of a function ϕ are denoted by $\phi_{\pm} \coloneqq \max\{\pm \phi, 0\}$. Throughout this paper, we will use the following notations: $tz \coloneqq t(u, v) = (tu, tv)$ for all $(u, v) \in W$ and $t \in \mathbb{R}$; (u, v) is said to be nonnegative in \mathbb{R}^N if $u \ge 0$ and $v \ge 0$ in \mathbb{R}^N ; (u, v) is said to be positive in \mathbb{R}^N if u > 0 and v > 0 in \mathbb{R}^N . Finally, $C \in \mathbb{R}_+$ denotes a universal constant that may change from line to line; when it is relevant, we will add subscripts to specify the dependence of certain parameters.

2. HISTORICAL BACKGROUND

The fractional Laplacian. There are many equivalent definitions of the fractional Laplacian. In our case, on the Euclidean space \mathbb{R}^N of dimension $N \ge 1$, for $\theta = \theta_1 + \theta_2$ and the above specified intervals for the parameters, we define the nonlocal elliptic *p*-Laplacian operator with the help of the Cauchy's principal value integral as in (3).

For problems with two nonlinearities involving the Laplacian operator, see Filipucci, Pucci & Robert [13]. For similar problems involving the fractional Laplacian, see Servadei & Valdinoci [30], Ghoussoub & Shakerian [16], Chen & Squassina [11] Chen [8, 9], Assunção, Silva & Miyagaki [3]. For a survey paper on the subject of fractional Sobolev spaces, see Di Nezza, Palatucci & Valdinoci [12]; see also Molica Bisci, Rădulescu & Servadei [23].

The Choquard equation. On the Euclidean space \mathbb{R}^N of dimension $N \ge 1$ and for $x \in \mathbb{R}^N$, the equation $-\Delta u + V(x)u = (I_{\mu} * |u|^q)|u|^{q-2}u$ was introduced by Choquard in the case N = 3 and q = 2 to model a certain approximation to Hartree-Fock theory of onecomponent plasma and to describe a electron trapped in its own hole. Also in this situation it finds physical significance in the work by Frölich and Pekar on the description of the quantum mechanics of a polaron at rest. When $V(x) \equiv 1$, the groundstate solutions exist if $2^{\flat} := 2(N - \mu/2))/N < q < 2(N - \mu/2)/(N - 2s) := 2^{\sharp}$ due to the mountain pass lemma or the method of the Nehari manifold, while there are no nontrivial solution if $q = 2^{\flat}$ or if $q = 2^{\sharp}$ as a consequence of the Pohozaev identity.

In general, the associated Schrödinger-type evolution equation $i\partial_t \psi = \Delta \psi + (I_\mu * |\psi|^2)\psi$ is a model for large systems of atoms with an attractive interaction that is weaker and has a longer range than that of the nonlinear Schrödinger equation. Standing wave solutions of this equation are solutions to the Choquard equation. For more information on the various results related to the non-fractional Choquard-type equations and their variants see Moroz & Van Schaftingen [24] and Mukherjee & Sreenadh [26].

The Morrey spaces. The study of Morrey spaces is motivated by many reasons. Initially, these spaces were introduced by Morrey in order to understand the regularity of solutions to elliptic partial differential equations. Regularity theorems, which allow one to conclude higher regularity of a function that is a solution of a differential equation together with a lower regularity of that function, play a central role in the theory of partial differential equations. One example of this kind of regularity theorem is a version of the Sobolev embedding theorem which states that $W^{j+m,p}(\Omega) \subset C^{j,\lambda}(\overline{\Omega})$ for $0 < \lambda \leq m - N/p$, where $j \in \mathbb{N}$ and $\Omega \subset \mathbb{R}^N$ is a Lipschitz domain.

Morrey spaces can complement the boundedness properties of operators that Lebesgue spaces can not handle. In line with this, many authors study the boundedness of various integral operators on Morrey spaces. The theory of Morrey spaces may come in useful when the Sobolev embedding theorem is not readily available. For more information on Morrey spaces, see Gantumur [15] and Sawano [29].

Systems of fractional elliptic equations. The subject of two or more fractional elliptic equations have been widely studied in recent years. We devote this section on briefly glimpsing the results that have already been proved in the context of existence, non-existence, uniqueness and multiplicity of solutions to systems of fractional elliptic equations.

Liu & Wang [21] gave a sufficient condition on large coupling coefficients for the existence of a nontrivial ground state solution in a system of nonlinear Schrödinger equation; they also considered bound state solutions. Chen & Deng [10] investigated the existence of two nontrivial solutions to the fractional p-Laplacian system involving concave-convex nonlinearities via the Nehari method. Chen [7] obtained the existence of infinitely many nonnegative solutions for a class of the quasilinear Schrödinger system in \mathbb{R}^N in the Laplacian setting and investigate the multiplicity of solutions for a p-Kirchhoff system driven by a nonlocal integro-differential operator with zero Dirichlet boundary data. Xiang, Zhang & Rădulescu [32] studied the multiplicity of solutions for a p-Kirchhoff system driven by a nonlocal integro-differential operator with zero Dirichlet boundary data. Chen & Squassina [11] used Nehari manifold techniques to obtain the existence of multiple solutions to a fractional *p*-Laplacian system involving critical concave-convex nonlinearities. Fiscella, Pucci & Saldi [14], using several variational methods, dealt with the existence of nontrivial nonnegative solutions of Schrödinger–Hardy systems driven by two possibly different fractional *p*-Laplacian operators. The main features of this paper is the presence of the Hardy term and the fact that the nonlinearities do not necessarily satisfy the Ambrosetti-Rabinowitz condition. Wang, Zhang & Zhang [31] are interested in a fractional Laplacian system in the whole space \mathbb{R}^N , which involves critical Sobolev-type nonlinearities and critical Hardy-Sobolev-type nonlinearities. Yang [33] considered the existence of nontrivial weak solutions to a doubly critical system involving fractional Laplacian in \mathbb{R}^N with subcritical weight. More recently, Lu & Shen [22] studied a critical fractional p-Laplacian system with homogeneous nonlinearity; they used a concentration compactness principle associated with fractional p-Laplacian system for the fractional order Sobolev spaces in bounded domains, which is significantly more difficult to prove than in the case of a single fractional *p*-Laplacian equation and is of independent interest.

Most of the existing results have been developed for systems with two equations. For a general system, Lin & Wei [19, 20] studied a system with several coupled nonlinear Schrödinger equations in the whole space up to three dimensions which has some applications in nonlinear optics. The existence of ground state solutions may depend on the coupling constants that model the interaction between the components of the system. If a constant is positive, the interaction is attractive; otherwise, it is repulsive. When all the constants are positive and some associated matrix is positively definite, they proved the existence of a radially symmetric ground state solution; however, if all the constants are negative, our if one of them is negative and the matrix is positively definite, there is no ground state solution. They also obtained the existence of a bound state solution which is non-radially symmetric in the three dimensional case.

Our contribution to the problem and some of its difficulties. The present work is motivated by Assunção, Miyagaki & Siqueira [4], Wang, Zhang & Zhang [31] and Yang [33]. Our existence result can be regarded as an extension and improvement of the corresponding existence results in these works. More precisely, we will extend the result in [4] to a system of coupled equations in \mathbb{R}^N with the general fractional *p*-Laplacian with p > 1and $\theta = \theta_1 + \theta_2$ not necessarily zero. Moreover, we use a refinement of Sobolev inequality that is related to Morrey space because our problem involves doubly critical exponents. As one can expect, the nonlocality of the fractional *p*-Laplacian makes it more difficult to study. In our case, one of the main difficulties when dealing with this problem is the lack of compactness of Sobolev embedding theorem for the critical exponent. Therefore, we have to develop a precise analysis of the level of the Palais-Smale sequences obtained with the application of the mountain pass theorem and study their behavior concerning strong convergence of one of its scaled subsequences.

3. EXISTENCE OF SOLUTIONS FOR AUXILIARY MINIMIZATION PROBLEMS

We begin this section by introducing two important and sharp Rayleigh-Ritz constants. The first one is related to the Gagliardo seminorm, the Hardy term, and a convolution involving the upper Stein-Weiss exponent,

(7)
$$S^{\sharp} \coloneqq \inf_{(u,v)\in W\setminus\{0\}} \frac{\|(u,v)\|_W^p}{(Q^{\sharp}(u,v))^{\frac{p}{2p_s^{\sharp}(\delta,\theta,\mu)}}}$$

where the quadratic form $Q^{\sharp} \colon W \to \mathbb{R}$ is given by

(8)
$$Q^{\sharp}(u,v) \coloneqq \iint_{\mathbb{R}^{2N}} \frac{|u(x)|^{p_{s}^{\sharp}(\delta,\theta,\mu)}|u(y)|^{p_{s}^{\sharp}(\delta,\theta,\mu)} + |v(x)|^{p_{s}^{\sharp}(\delta,\theta,\mu)}|v(y)|^{p_{s}^{\sharp}(\delta,\theta,\mu)}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \,\mathrm{d}x \,\mathrm{d}y.$$

The second one is related to the Gagliardo seminorm, ther Hardy term, and the norm in the critical weighted Lebesgue space, that is, the Sobolev constant,

(9)
$$S^* \coloneqq \inf_{(u,v) \in W \setminus \{0\}} \frac{\|(u,v)\|_W^p}{(Q^*(u,v))^{\frac{p}{p_s^*(\beta,\theta)}}}$$

where the quadratic form $Q^* \colon W \to \mathbb{R}$ is given by

(10)
$$Q^*(u,v) \coloneqq \int_{\mathbb{R}^N} \frac{|u(x)|^{p_s^*(\beta,\theta)} + |v(x)|^{p_s^*(\beta,\theta)} + \eta |u(x)|^a |v(x)|^b}{|x|^\beta} \,\mathrm{d}x$$

For general $p \neq 2$, the explicit formula for the extremal functions for the *p*-fractional Sobolev inequality is not known yet, though it is conjectured that it is of the form

$$U(x) = \frac{C}{(1 + |x|^{\frac{p}{p-1}})^{\frac{N-sp}{p}}}$$

up to translation and scaling. However, there is a result about the asymptotic behavior of U, as seen in Brasco, Mosconi & Squassina [5] and Mosconi, Perera, Squassina & Yang [25].

One of the first major difficulties that we encounter is the lack of an explicit formula for a minimizer of the quantity S^* . There is a conjecture about the minimizers which states that they have the form

$$U(x) = \frac{c}{[1 + (|x - x_0|/\varepsilon))^{\frac{p}{p-1}}]^{\frac{N-sp}{p}}}$$

where $C \neq \mathbb{R}^N$, $x_0 \in \mathbb{R}^N$, and $\varepsilon \in \mathbb{R}_+$. This conjecture was proved by Lieb [18] in the case p = 2; however, for $p \neq 2$ it is not even known if these functions are minimizers. To overcome this difficulty, we will work with some asymptotic estimates for minimizers

recently obtained by Brasco, Mosconi & Squassina [6]; see also Mosconi, Perera, Squassina & Yang [25].

In the next lemma we present some embeddings of Morrey spaces into weighted Lebesgue spaces; the proof is straightforward.

Lemma 3. The following fundamental properties are true. (1) $L^{p\rho}(\mathbb{R}^N, |y|^{-\rho\lambda}) \hookrightarrow L^{p,\gamma+\lambda}(\mathbb{R}^N, |y|^{-\lambda})$ for $\rho = \frac{N}{\gamma+\lambda} > 1$. (2) For any $p \in (1, +\infty)$, we have $L^{p,\gamma+\lambda}(\mathbb{R}^N, |y|^{-\lambda}) \hookrightarrow L^{1,\frac{\gamma}{p}+\frac{\lambda}{p}}(\mathbb{R}^N, |y|^{-\frac{\lambda}{p}})$. (3) For $1 \leq p < +\infty$ and $\gamma + \lambda = N$, we have $L^{p,N}_M(\mathbb{R}^N, |y|^{-\lambda}) = L^p(\mathbb{R}^N, |y|^{-\lambda})$, *i.e.*,

- (3) For $1 \leq p < +\infty$ and $\gamma + \lambda = N$, we have $L_M^{p,N}(\mathbb{R}^N, |y|^{-\lambda}) = L^p(\mathbb{R}^N, |y|^{-\lambda})$, i.e., $L_M^{p,N}(\mathbb{R}^N, |y|^{-\lambda})$ and $L^p(\mathbb{R}^N, |y|^{-\lambda})$ are continuously embedded in each other.
- (4) If if we assume that $s \in (0,1)$ and $0 < \alpha < sp + \theta < N$, for $1 \leq q < p_s^*(\alpha,\theta)$ and $r = \frac{\alpha}{p_s^*(\alpha,\theta)}$, it holds

(11)
$$\dot{W}^{s,p}_{\theta}(\mathbb{R}^{N}) \hookrightarrow L^{p^{*}_{s}(\alpha,\theta)}(\mathbb{R}^{N},|y|^{-\alpha}) \hookrightarrow L^{q,\frac{(N-sp-\theta)q}{p}+qr}(\mathbb{R}^{N},|y|^{-pr})$$

and the norms in these spaces share the same dilation invariance. (5) For any $q \in [1, p_s^*(0, \theta)), \dot{W}_{\theta}^{s, p}(\mathbb{R}^N) \hookrightarrow L^{p_s^*(0, \theta)}(\mathbb{R}^N) \hookrightarrow L^{q, \frac{(N-sp-\theta)q}{p}}(\mathbb{R}^N).$

For more properties of Lebesgue spaces, integral inequalities and boundedness properties of the operators in generalized Morrey spaces, see Sawano [29].

Next, we use the following versions of the fractional Hardy-Sobolev and Caffarelli-Kohn-Nirenberg inequalities; see Nguyen & Squassina [27, Theorem 1.1]; see also Abdellaoui & Bentifour [1].

Lemma 4. Let $N \ge 1, p \in (1, +\infty), s \in (0, 1), 0 \le \alpha \le sp + \theta < N, \theta, \theta_1, \theta_2, \beta \in \mathbb{R}$ be such that $\theta = \theta_1 + \theta_2$. If $1/p_s^*(\alpha, \theta) - \alpha/Np_s^*(\alpha, \theta) > 0$, then there exists a positive constant $C(N, \alpha, \theta)$ such that

(12)
$$\left(\int_{\mathbb{R}^N} \frac{|u|^{p_s^*(\alpha,\theta)}}{|x|^{\alpha}} \mathrm{d}x\right)^{\frac{p}{p_s^*(\alpha,\theta)}} \leqslant C(N,\alpha,\theta) \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^p}{|x|^{\theta_1}|x - y|^{N+sp}|y|^{\theta_2}} \mathrm{d}x\mathrm{d}y$$

for all $u \in W^{s,p}_{\theta}(\mathbb{R}^N)$.

We note that the norm $\|\cdot\|$ is comparable with the Gagliardo seminorm $[\cdot]_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)}$ as stated in the next result.

Corollary 5. Under the hypotheses of Lemma 4, if $\gamma < \gamma_H$ then

(13)
$$\begin{pmatrix} 1 - \frac{\gamma_+}{\gamma_H} \end{pmatrix} \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^p}{|x|^{\theta_1} |x - y|^{N+sp} |y|^{\theta_2}} \mathrm{d}x \mathrm{d}y \\ \leqslant ||u||^p \leqslant \left(1 + \frac{\gamma_-}{\gamma_H}\right) \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^p}{|x|^{\theta_1} |x - y|^{N+sp} |y|^{\theta_2}} \mathrm{d}x \mathrm{d}y$$

where $\gamma_{\pm} = \max\{\pm\gamma, 0\}.$

We can deduce another useful inequality.

Corollary 6. Under the hypotheses of Lemma 4 we have

(14)
$$\iint_{\mathbb{R}^{2N}} \frac{|u(x)|^{p_s^{\sharp}(\delta,\theta,\mu)}|u(y)|^{p_s^{\sharp}(\delta,\theta,\mu)}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y \leqslant C(N,\delta,\mu) \|u\|^{2p_s^{\sharp}(\delta,\theta,\mu)}_{\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)}$$

Based on the embeddings (11) we establish an improved weighted fractional Caffarelli-Kohn-Nirenberg inequality whose proof can be found in [4]; see also [28].

Lemma 7. (Fractional Caffarelli-Kohn-Nirenberg inequality) Let $s \in (0,1)$ and $0 < \beta < sp + \theta < N$. Then there exists $C = C(N, s, \beta) > 0$ such that for any $\zeta \in (\overline{\zeta}, 1)$ and for any $q \in [1, p_s^*)$, for all $u \in \dot{W}_{\theta}^{s,p}(\mathbb{R}^N)$ it holds

(15)
$$\left(\int_{\mathbb{R}^{N}} \frac{|u(y)|^{p_{s}^{*}}}{|y|^{\beta}} \mathrm{d}y\right)^{\frac{1}{p_{s}^{*}}} \leq \|u\|_{\dot{W}^{s,p}_{\theta}}^{\zeta} \|u\|_{L^{q,\frac{N-sp-\theta}{p}q+qr}(\mathbb{R}^{N},|y|^{-qr})}^{1-\zeta}$$

where $\bar{\zeta} = \max\{p/p_s^*, (p_s^*(0,\theta) - 1)/p_s^*\} > 0$ and $r = \beta/p_s^*$.

Now we state a result about local convergence; see [17, Lemma 2.3].

Lemma 8. Let $s \in (0,1)$ and $0 < r < s + \frac{\theta}{p} < \frac{N}{p}$. If $\{u_k\}$ is a bounded sequence in $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ and $u_k \rightharpoonup u$ in $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$, then as $k \rightarrow +\infty$,

$$\frac{u_k}{|x|^{r+\frac{\theta r}{sp}}} \to \frac{u}{|x|^{r+\frac{\theta r}{sp}}} \text{ in } L^p_{\text{loc}}(\mathbb{R}^N).$$

Proposition 9. For $s \in (0, 1)$ the best constants S^{\sharp} and S^{*} verify the following items.

- (1) If $0 < \alpha < sp + \theta < N$, $\mu \in (0, N)$ and $\gamma < \gamma_H$, then S^{\sharp} is attained in W;
- (2) If $0 < \beta < sp + \theta < N$ and $\gamma < \gamma_H$, then S^* is attained in W.

Proof. (1) If $0 < \alpha < sp + \theta < N$ and $\gamma < \gamma_H$, let $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$ be a minimizing sequence of S^{\sharp} such that

(16)
$$Q^{\sharp}(u_k, v_k) = 1, \qquad ||(u_k, v_k)||^p \to S^{\sharp}$$

as $k \to +\infty$. Recall that $r = \frac{\alpha}{p_s^*(\alpha,\theta)}$.

Claim 1. We have
$$C_1 \leq ||u_k||_{L_M^{q,\frac{N-sp-\theta}{p}q+qr}(\mathbb{R}^N,|y|^{-pr})} \leq C_2$$
.

Proof. We just use the embeddings (11) and the Caffarelli-Kohn-Nirenberg's inequality (15). For more details, see [4, Eq. (4.2), p. 23] \Box

For any $k \in \mathbb{N}$ large enough, we may find $\lambda_k > 0$ and $x_k \in \mathbb{R}^N$ such that

(17)
$$\lambda_{k}^{-sp-\theta+pr} \int_{B_{\lambda_{k}(x_{k})}} \frac{|u_{k}(y)|^{p}}{|y|^{pr}} \mathrm{d}y > ||u_{k}||_{L_{M}^{p,N-sp-\theta+pr}(\mathbb{R}^{N},|y|^{-pr})}^{p} - \frac{C}{2k} \ge C > 0$$

for constants $C \in \mathbb{R}_+$.

Our goal is to pass to the limit as $k \to +\infty$ in the minimizing sequence. To do this, we adapt the Levy's concentration principle; more precisely, we create another sequence $\{(\tilde{u}_k, \tilde{v}_k)\}_{k \in \mathbb{N}} \subset W$ that will help us to control the radius and the centers of these balls. Let

$$\tilde{u}_k(x) = \lambda_k^{\frac{N-sp-\theta}{p}} u_k(\lambda_k x) \text{ and } \tilde{v}_k(x) = \lambda_k^{\frac{N-sp-\theta}{p}} v_k(\lambda_k x)$$

be the appropriate scaling for the class of problems that we consider and define $\tilde{x}_k := x/\lambda_k$ and $\tilde{x}_k := x/\lambda_k$. Then, using the change of variables $y = \lambda_k x$ with $dy = \lambda_k^N dx$ we have,

(18)
$$\int_{B_1\left(\frac{x_k}{\lambda_k}\right)} \frac{|\tilde{u}_k(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C > 0 \quad \text{and} \quad \int_{B_1\left(\frac{x_k}{\lambda_k}\right)} \frac{|\tilde{v}_k(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C > 0.$$

Now we claim that S^{\sharp} is invariant under the previously defined dilation.

In fact, $Q^{\sharp}(\tilde{u}_k, \tilde{v}_k) = 1$. To show this property, we use the change of variables $\bar{x} = \lambda_k x$ and $\bar{y} = \lambda_k y$, we have

$$\begin{split} Q^{\sharp}(\tilde{u}_{k},\tilde{v}_{k}) &= \iint_{\mathbb{R}^{2N}} \frac{|u_{k}(x)|^{p_{s}^{\sharp}}|u_{k}(y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{|v_{k}(y)|^{p_{s}^{\sharp}}|v_{k}(y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y \\ &= \iint_{\mathbb{R}^{2N}} \lambda_{ks}^{\frac{N-sp-\theta}{p}2p_{s}^{\sharp}} \frac{|u_{k}(\lambda_{k}x)|^{p_{s}^{\sharp}}|u_{k}(\lambda_{k,1}y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y \\ &+ \iint_{\mathbb{R}^{2N}} \lambda_{k}^{\frac{N-sp-\theta}{p}2p_{s}^{\sharp}} \frac{|v_{k}(\lambda_{k}x)|^{p_{s}^{\sharp}}|v_{k}(\lambda_{k}y)|^{p_{s}^{\sharp}}}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x\mathrm{d}y \\ &= \iint_{\mathbb{R}^{2N}} \frac{|u_{k}(\bar{x})|^{p_{s}^{\sharp}}|u_{k}(\bar{y})|^{p_{s}^{\sharp}}}{|\bar{x}|^{\delta}|\bar{x}-\bar{y}|^{\mu}|\bar{y}|^{\delta}} \mathrm{d}\bar{x}\mathrm{d}\bar{y} + \iint_{\mathbb{R}^{2N}} \frac{|v_{k}(\bar{x})|^{p_{s}^{\sharp}}|v_{k}(\bar{y})|^{p_{s}^{\sharp}}}{|\bar{x}|^{\delta}|\bar{x}-\bar{y}|^{\mu}|\bar{y}|^{\delta}} \mathrm{d}\bar{x}\mathrm{d}\bar{y} \end{split}$$

Furthermore, $\|(\tilde{u}_k, \tilde{v}_k)\|^p \to S^{\sharp}$. In fact, we know that $\{(\tilde{u}_k, \tilde{v}_k)\}_{k \in \mathbb{N}}$ is a minimizing sequence for S^{\sharp} . Using the same change of variables $\bar{x} = \lambda_k x$ and $\bar{y} = \lambda_k y$, we obtain

$$\begin{split} \|(\tilde{u}_{k},\tilde{v}_{k})\|^{p} &= \iint_{\mathbb{R}^{2N}} \frac{|u_{k}(x) - u_{k}(y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{|v_{k}(x) - v_{k}(y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \\ &- \gamma_{1} \int_{\mathbb{R}^{N}} \frac{|u_{k}|^{p}}{|\bar{x}|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \frac{|v_{k}|^{p}}{|\bar{x}|^{sp+\theta}} \mathrm{d}\bar{x} \\ &= \iint_{\mathbb{R}^{2N}} \lambda_{k,1}^{N-sp-\theta} \frac{|u_{k}(\lambda_{k}x) - u_{k}(\lambda_{k}y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \\ &+ \iint_{\mathbb{R}^{2N}} \lambda_{k}^{N-sp-\theta} \frac{|v_{k}(\lambda_{k}x) - v_{k}(\lambda_{k}y)|^{p}}{|x|^{\theta_{1}}|x - y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \\ &- \gamma_{1} \int_{\mathbb{R}^{N}} \lambda_{k}^{N-sp-\theta} \frac{|u_{k}|^{p}}{|x|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \lambda_{k,2}^{N-sp-\theta} \frac{|v_{k}|^{p}}{|x|^{sp+\theta}} \mathrm{d}x \\ &= \iint_{\mathbb{R}^{2N}} \frac{|u_{k}(\bar{x}) - u_{k}(\bar{y})|^{p}}{|\bar{x}|^{\theta_{1}}|\bar{x} - \bar{y}|^{N+sp}|\bar{y}|^{\theta_{2}}} \mathrm{d}\bar{x} \mathrm{d}\bar{y} + \iint_{\mathbb{R}^{2N}} \frac{|v_{k}(\bar{x}) - v_{k}(\bar{y})|^{p}}{|\bar{x}|^{\theta_{1}+\theta}} \mathrm{d}x \\ &= \iint_{\mathbb{R}^{2N}} \frac{|u_{k}(\bar{x}) - u_{k}(\bar{y})|^{p}}{|\bar{x}|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \frac{|v_{k}|^{p}}{|\bar{x}|^{\theta_{1}}|\bar{x} - \bar{y}|^{N+sp}|\bar{y}|^{\theta_{2}}} \mathrm{d}\bar{x} \mathrm{d}\bar{y} \\ &- \gamma_{1} \int_{\mathbb{R}^{N}} \frac{|u_{k}|^{p}}{|\bar{x}|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \frac{|v_{k}|^{p}}{|\bar{x}|^{sp+\theta}} \mathrm{d}\bar{x} \\ &= \|(u_{k}, v_{k})\|^{p}. \end{split}$$

And since $||(u_k, v_k)||^p \to S^{\sharp}$ as $k \to +\infty$, we deduce that $||(\tilde{u}_k, \tilde{v}_k)||^p \to S^{\sharp}$ as $k \to +\infty$. In this way, the sequence $\{(\tilde{u}_k, \tilde{v}_k)\}_{k \in \mathbb{N}} \subset W$ is also a minimizing sequence for S^{\sharp} such that we have

(19)
$$Q^{\sharp}(\tilde{u}_k, \tilde{v}_k) = 1, \qquad \|(\tilde{u}_k, \tilde{v}_k)\|^p \to S^{\sharp}.$$

Consider $\{\tilde{x}_k\} = \{\frac{x_k}{\lambda_k}\}$, from inequality (18) together with Hölder's inequality

$$0 < C \leqslant \int_{B_1(\tilde{x}_k)} \frac{|\tilde{u}_k(x)|^p}{|x|^{pr}} \mathrm{d}x$$
$$\leqslant \left(\int_{B_1(\tilde{x}_k)} 1\mathrm{d}x\right)^{1-\frac{p}{p_s^*(\alpha,\theta)}} \left(\int_{B_1(\tilde{x}_k)} \left(\frac{|\tilde{u}_k(x)|^p}{|x|^{pr}}\right)^{\frac{p_s^*(\alpha,\theta)}{p}} \mathrm{d}x\right)^{\frac{p}{p_s^*(\alpha,\theta)}}$$

$$\leqslant C \left(\int_{B_1(\tilde{x}_k)} \frac{|\tilde{u}_k(x)|^{p_s^*(\alpha,\theta)}}{|x|^{\alpha}} \mathrm{d}x \right)^{\frac{p}{p_s^*(\alpha,\theta)}}.$$

Therefore,

(20)
$$\left(\int_{B_1(\tilde{x}_k)} \frac{|\tilde{u}_k(x)|^{p^*_s(\alpha,\theta)}}{|x|^{\alpha}} \mathrm{d}x\right)^{\frac{p}{p^*_s(\alpha,\theta)}} \ge C > 0$$

We claim that the sequence $\{\tilde{x}_k\} \subset \mathbb{R}^N$ of the centers of the balls is bounded. We argue by contradiction and suppose that $|\tilde{x}_k| \to +\infty$ as $k \to +\infty$; then for any $x \in B_1(\tilde{x}_k)$, we have $|x| \ge |\tilde{x}_k| - 1$ for $k \in \mathbb{N}$ large enough. By Hölder's inequality, we obtain

$$\begin{split} \int_{B_1(\tilde{x}_k)} \frac{|\tilde{u}_k(x)|^{p_s^*(\alpha,\theta)}}{|x|^{\alpha}} \mathrm{d}x &\leq \frac{1}{(|\tilde{x}_k|-1)^{\alpha}} \int_{B_1(\tilde{x}_k)} |\tilde{u}_k(x)|^{p_s^*(\alpha,\theta)} \mathrm{d}x \\ &\leq \frac{C}{(|\tilde{x}_k|-1)^{\alpha}} \left(\int_{B_1(\tilde{x}_k)} |\tilde{u}_k(x)|^{p_s^*(0,\theta)} \mathrm{d}x \right)^{\frac{p_s^*(\alpha,\theta)}{p_s^*(0,\theta)}} \\ &\leq \frac{C}{(|\tilde{x}_k|-1)^{\alpha}} \|\tilde{u}_k(x)\|_{L^{p_s^*(\alpha,\theta)}}^{p_s^*(\alpha,\theta)} \\ &\leq \frac{C}{(|\tilde{x}_k|-1)^{\alpha}} \|u_k(x)\|_{W_{\theta}^{s,p}(\mathbb{R}^N)}^{p_s^*(\alpha,\theta)} \\ &\leq \frac{C}{(|\tilde{x}_k|-1)^{\alpha}} \to 0 \qquad (k \to +\infty) \end{split}$$

where we used the boundedness of the minimizing sequence $\{\tilde{u}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$. This is a contradiction with inequality (20) and this implies that the sequence $\{\tilde{x}_k\}\subset\mathbb{R}^N$ is bounded.

From inequality (18) and the boundedness of the sequence $\{\tilde{x}_k\} \subset \mathbb{R}^N$ of the centers of the balls, we may find R > 0 such that $B_R(0)$ contains all balls of center \tilde{x}_k and radius 1; moreover, with

(21)
$$\int_{B_R(0)} \frac{|\tilde{u}_k(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C_1 > 0.$$

Similar computations can also be done with respect to the sequence of functions $\{\tilde{v}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$. Therefore, we obtain $\|(\tilde{u}_k, \tilde{v}_k)\| = \|(u_k, v_k)\| \leq C$ for $k \in \mathbb{N}$ large enough and there exists a function pair $(\tilde{u}, \tilde{v}) \in W$ such that

(22)
$$(\tilde{u}_k, \tilde{v}_k) \rightharpoonup (\tilde{u}, \tilde{v})$$
 weakly in $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N) \times \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$

(23)
$$(\tilde{u}_k, \tilde{v}_k) \to (\tilde{u}, \tilde{v})$$
 a.e. on $\mathbb{R}^N \times \mathbb{R}^N$,

as $k \to +\infty$, up to subsequences. According to Lemma 8, we have

$$\left(\frac{\tilde{u}_k}{|x|^r}, \frac{\tilde{v}_k}{|x|^r}\right) \to \left(\frac{\tilde{u}}{|x|^r}, \frac{\tilde{v}}{|x|^r}\right) \text{ in } L^p_{\text{loc}}(\mathbb{R}^N) \times L^p_{\text{loc}}(\mathbb{R}^N);$$

hence,

$$\int_{B_R(0)} \frac{|\tilde{u}(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C_1 > 0,$$

and we deduce that $\tilde{u} \neq 0$. Similarly, we can get $\tilde{v} \neq 0$.

We can verify in the same way as we did in [4, Lemma 2.6, p. 16] that

(24)
$$1 = Q^{\sharp}(\tilde{u}_k, \tilde{v}_k) = Q^{\sharp}(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v}) + Q^{\sharp}(\tilde{u}, \tilde{v}) + o(1).$$

By a Brézis-Lieb's lemma, we obtain

(25)
$$\int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{u}_{k} - \tilde{u} \right|^{p_{s}^{\sharp}}}{\left| y \right|^{\delta}} \right) \frac{\left| \tilde{u}_{k} - \tilde{u} \right|^{p_{s}^{\sharp}}}{\left| x \right|^{\delta}} \mathrm{d}x + \int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{u} \right|^{p_{s}^{\sharp}}}{\left| y \right|^{\delta}} \right) \frac{\left| \tilde{u} \right|^{p_{s}^{\sharp}}}{\left| x \right|^{\delta}} \mathrm{d}x \\ = \int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{u}_{k} \right|^{p_{s}^{\sharp}}}{\left| y \right|^{\delta}} \right) \frac{\left| \tilde{u}_{k} \right|^{p_{s}^{\sharp}}}{\left| x \right|^{\delta}} \mathrm{d}x + o(1),$$

and

(26)
$$\int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{v}_{k} - \tilde{v} \right|^{p_{s}^{\sharp}}}{|y|^{\delta}} \right) \frac{\left| \tilde{v}_{k} - \tilde{v} \right|^{p_{s}^{\sharp}}}{|x|^{\delta}} \mathrm{d}x + \int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{v} \right|}{|y|^{\delta}} \right) \frac{\left| \tilde{v} \right|}{|x|^{\delta}} \mathrm{d}x \\ = \int_{\mathbb{R}^{N}} \left(I_{\mu} * \frac{\left| \tilde{v}_{k} \right|^{p_{s}^{\sharp}}}{|y|^{\delta}} \right) \frac{\left| \tilde{v}_{k} \right|^{p_{s}^{\sharp}}}{|x|^{\delta}} \mathrm{d}x + o(1),$$

Therefore, by definition (7), by weak convergence $(\tilde{u}_k, \tilde{v}_k) \rightharpoonup (\tilde{u}, \tilde{v})$ in W together with the Brézis–Lieb lemma and by the estimate (24), we have

$$S^{\sharp} = \lim_{k \to \infty} \|(\tilde{u}_{k}, \tilde{v}_{k})\|^{p}$$

= $\|(\tilde{u}, \tilde{v})\|^{p} + \lim_{k \to \infty} \|(\tilde{u}_{k} - \tilde{u}, \tilde{v}_{k} - \tilde{v})\|^{p}$
$$\geq S^{\sharp} (Q^{\sharp}(\tilde{u}, \tilde{v}))^{\frac{p}{p_{s}^{\sharp}}} + S^{\sharp} \left(\lim_{k \to \infty} Q^{\sharp} (\tilde{u}_{k} - \tilde{u}, \tilde{v}_{k} - \tilde{v})\right)^{\frac{p}{p_{s}^{\sharp}}}$$

$$\geq S^{\sharp} (Q^{\sharp}(\tilde{u}, \tilde{v}) + \lim_{k \to \infty} Q^{\sharp} (\tilde{u}_{k} - \tilde{u}, \tilde{v}_{k} - \tilde{v}))^{\frac{p}{p^{\sharp}}}$$

= $S^{\sharp},$

where in the last but one passage above we used the inequality

 $(27) (a+b)^q \leqslant a^q + b^q,$

valid for all $a, b \in \mathbb{R}^*_+$ and 0 < q < 1. So we have equality in all passages, that is,

(28)
$$Q^{\sharp}(\tilde{u},\tilde{v}) = 1, \qquad \lim_{k \to \infty} Q^{\sharp}(\tilde{u}_k - \tilde{u},\tilde{v}_k - \tilde{v}) = 0,$$

since $\tilde{u}, \tilde{v} \neq 0$. It turns out that, since

$$S^{\sharp} = \|(\tilde{u}, \tilde{v})\|^p + \lim_{k \to \infty} \|(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\|^p,$$

then

$$S^{\sharp} = \|(\tilde{u}, \tilde{v})\|^p$$
 and $\lim_{k \to \infty} \|(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\|^p = 0.$

Finally, by inequality

$$\begin{split} \iint_{\mathbb{R}^{2N}} \frac{||\tilde{u}(x)| - |\tilde{u}(y)||^p}{|x|^{\theta_1}|x - y|^{N+sp}|y|^{\theta_2}} \mathrm{d}x \mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{||\tilde{v}(x)| - |\tilde{v}(y)||^p}{|x|^{\theta_1}|x - y|^{N+sp}|y|^{\theta_2}} \mathrm{d}x \mathrm{d}y \\ &\leqslant \iint_{\mathbb{R}^{2N}} \frac{|\tilde{u}(x) - \tilde{u}(y)|^p}{|x|^{\theta_1}|x - y|^{N+sp}|y|^{\theta_2}} \mathrm{d}x \mathrm{d}y + \iint_{\mathbb{R}^{2N}} \frac{|\tilde{v}(x) - \tilde{v}(y)|^p}{|x|^{\theta_1}|x - y|^{N+sp}|y|^{\theta_2}} \mathrm{d}x \mathrm{d}y. \end{split}$$

we deduce that $(|\tilde{u}|, |\tilde{v}|) \in W$ is also a minimizer for S^{\sharp} ; so we can assume that $\tilde{u} \ge 0, \tilde{v} \ge 0$. Thus, S^{\sharp} is achieved by a non-negative function in the case $0 < \alpha < sp + \theta$ and $\gamma < \gamma_H$.

(2) For $0 < \beta < sp + \theta < N$ and $\gamma < \gamma_H$, let $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$ be a minimizing sequence for S^* such that

(29)
$$Q^*(u_k, v_k) = 1, \qquad ||(u_k, v_k)||^p \to S^*$$

as $k \to +\infty$.

Now we claim that S^* is invariant under the previously defined dilation. Let

$$\tilde{u}_k(x) = \lambda_{k,1}^{\frac{N-sp-\theta}{p}} u_k(\lambda_{k,1}x), \qquad \qquad \tilde{v}_k(x) = \lambda_{k,2}^{\frac{N-sp-\theta}{p}} v_k(\lambda_{k,2}x)$$

and $\tilde{x}_k = \frac{x_k}{\lambda_{k,i}}$ for $i \in \{1, 2\}$ as in the previous case. In this way, the sequence $\{(\tilde{u}_k, \tilde{v}_k)\}_{k \in \mathbb{N}} \subset W$ is also a minimizing sequence for S^* such that we have

$$Q^*(\tilde{u}_k, \tilde{v}_k) = 1, \qquad \|(\tilde{u}_k, \tilde{v}_k)\|^p \to S^*$$

We have already shown that $\|(\tilde{u}_k, \tilde{v}_k)\| = \|(u_k, v_k)\|$ for every $k \in \mathbb{N}$. Hence, $\|(\tilde{u}_k, \tilde{v}_k)\|^p \to S^*$.

We claim that the sequence $\{\tilde{x}_k\} \subset \mathbb{R}^N$ is bounded and the proof follows the same steps already presented. From this boundedness and inequality (18), we may find R > 0 such that $B_R(0)$ contains all the unitary balls $B_1(\tilde{x}_k)$ centered in \tilde{x}_k and

(30)
$$\int_{B_R(0)} \frac{|v_k(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C_1 > 0.$$

Since $||v_k|| = ||u_k|| \leq C$, there exists a $v \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ such that

(31)
$$v_k \rightharpoonup v \quad \text{in } \dot{W}^{s,p}_{\theta}(\mathbb{R}^N), \qquad v_k \rightarrow v \text{ a.e. } \text{ on } \mathbb{R}^N,$$

as $k \to +\infty$, up to subsequences. According to Lemma 8, we have

$$\frac{v_k}{|x|^r} \to \frac{v}{|x|^r} \quad \text{in } L^p_{\text{loc}}(\mathbb{R}^N),$$

as $k \to +\infty$, where $r = \frac{\beta}{p_s^*}$. Therefore,

$$\int_{B_R(0)} \frac{|v(x)|^p}{|x|^{pr}} \mathrm{d}x \ge C_1 > 0,$$

and we deduce that $v \neq 0$.

We may verify by a variant of Brézis–Lieb Lemma [4, Lemma 2.2, p. 13] that, if $q = p_s^*(\beta, \theta)$ and $\delta = \beta$, then

$$1 = \int_{\mathbb{R}^N} \frac{|v_k|^{p_s^*}}{|x|^{\beta}} \mathrm{d}x = \int_{\mathbb{R}^N} \frac{|v_k - v|^{p_s^*}}{|x|^{\beta}} \mathrm{d}x + \int_{\mathbb{R}^N} \frac{|v|^{p_s^*}}{|x|^{\beta}} \mathrm{d}x + o(1).$$

By definition (9) and by weak convergence $(\tilde{u}_k, \tilde{v}_k) \rightharpoonup (\tilde{u}, \tilde{v})$ in W, we deduce that

$$S^* = \lim_{k \to \infty} \|(\tilde{u}_k, \tilde{v}_k)\|^p$$

= $\|(\tilde{u}, \tilde{v})\|^p + \lim_{k \to \infty} \|(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\|^p$
$$\geq S^* (Q^*(\tilde{u}, \tilde{v}))^{\frac{p}{p_s^*}} + S^* \left(\lim_{k \to \infty} Q^*(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\right)^{\frac{p}{p_s^*}}$$

$$\geq S^* (Q^*(\tilde{u}, \tilde{v}) + \lim_{k \to \infty} Q^*(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v}))^{\frac{p}{p_s^*}}$$

= S^*

where we used the inequality (27). So we have equality in all passages, that is,

(32)
$$Q^*(\tilde{u}, \tilde{v}) = 1, \qquad \lim_{k \to \infty} Q^*(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v}) = 0,$$

since $\tilde{u}, \tilde{v} \neq 0$. It turns out that, since $S^* = \|(\tilde{u}, \tilde{v})\|^p + \lim_{k \to \infty} \|(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\|^p$, then $S^* = \|(\tilde{u}, \tilde{v})\|^p$ and $\lim_{k \to \infty} \|(\tilde{u}_k - \tilde{u}, \tilde{v}_k - \tilde{v})\|^p = 0$.

As in the previous case, we deduce that $|(\tilde{u}, \tilde{v})| \in W$ is also a minimizer for S^* is achieved by a non-negative function in the case $0 < \beta < sp + \theta$ and $\gamma < \gamma_H$.

4. EXISTENCE OF PALAIS-SMALE SEQUENCE

We shall now use the minimizers of S^{\sharp} and S^{*} obtained in Proposition 9 to prove the existence of a nontrivial weak solution for equation (1). Recall the definition (4) of the energy functional associated to roblem (1). The fractional Sobolev and fractional Hardy-Sobolev inequalities imply that $I \in C^{1}(W, \mathbb{R})$ and that

$$\begin{split} \langle I'(u,v), (\phi_{1},\phi_{2}) \rangle \\ &= \iint_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^{p-2}(u(x)-u(y))(\phi_{1}(x)-\phi_{1}(y))}{|x|^{\theta_{1}}|x-y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \\ &+ \iint_{\mathbb{R}^{2N}} \frac{|v(x)-v(y)|^{p-2}(v(x)-v(y))(\phi_{2}(x)-\phi_{2}(y))}{|x|^{\theta_{1}}|x-y|^{N+sp}|y|^{\theta_{2}}} \mathrm{d}x \mathrm{d}y \\ &- \gamma_{1} \int_{\mathbb{R}^{N}} \frac{|u|^{p-2}u\phi_{1}}{|x|^{sp+\theta}} \mathrm{d}x - \gamma_{2} \int_{\mathbb{R}^{N}} \frac{|v|^{p-2}v\phi_{2}}{|x|^{sp+\theta}} \mathrm{d}x \\ &- \iint_{\mathbb{R}^{2N}} \frac{|u(x)|^{p_{s}^{\sharp}-2}|u(y)|^{p_{s}^{\sharp}}u(x)\phi_{1}(x)}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x \mathrm{d}y - \iint_{\mathbb{R}^{2N}} \frac{|v(x)|^{p_{s}^{\sharp}-2}|v(y)|^{p_{s}^{\sharp}}v(x)\phi_{2}(x)}{|x|^{\delta}|x-y|^{\mu}|y|^{\delta}} \mathrm{d}x \mathrm{d}y \\ &- \int_{\mathbb{R}^{N}} \frac{|u|^{p_{s}^{\ast}-2}u(x)\phi_{1}}{|x|^{\beta}} \mathrm{d}x - \int_{\mathbb{R}^{N}} \frac{|v|^{p_{s}^{\ast}-2}v(x)\phi_{2}}{|x|^{\beta}} \mathrm{d}x \\ &+ \int_{\mathbb{R}^{N}} \frac{\eta a|u|^{a-2}u\phi_{1}|v|^{b}}{|x|^{\beta}} \mathrm{d}x + \int_{\mathbb{R}^{N}} \frac{\eta b|u|^{a}|v|^{b-2}v\phi_{2}}{|x|^{\beta}} \mathrm{d}x. \end{split}$$

Note that a nontrivial critical point of I is a nontrivial weak solution to equation (1).

Recall that a Palais-Smale sequence for the energy functional I at the level $c \in \mathbb{R}$ is a sequence $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$ such that

(33)
$$\lim_{k \to +\infty} I(u_k, v_k) = c \quad \text{and} \quad \lim_{k \to +\infty} I'(u_k, v_k) = 0 \quad \text{strongly in } W'.$$

This sequence is referred to as a $(PS)_c$ sequence.

Now we state a result that ensures the existence of a Palais-Smale sequence for the energy functional.

Proposition 10. Let $s \in (0,1), 0 < \alpha, \beta < sp + \theta < N, \mu \in (0,N)$ and $\gamma < \gamma_H$. Consider the functional $I: W \to \mathbb{R}$ defined in (4) on the Banach space W. Then there exists a $(PS)_c$ sequence $\{(u_k, v_k)\} \subset W$ for I at some level $c \in (0, c^*)$, where

(34)
$$c^* \coloneqq \min\left\{ \left(\frac{1}{p} - \frac{1}{2p_s^{\sharp}(\delta, \theta, \mu)}\right) S^{\sharp \frac{2p_s^{\sharp}}{2p_s^{\sharp}(\delta, \theta, \mu) - p}}, \left(\frac{1}{p} - \frac{1}{p_s^{*}(\beta, \theta)}\right) S^{* \frac{p_s^{*}(\beta, \theta)}{p_s^{*}(\beta, \theta) - p}} \right\}.$$

To prove Proposition 10 we need the following version of the mountain pass theorem by Ambrosetti & Rabinowitz [2]. **Lemma 11.** (Montain Pass Lemma) Let $(W, \|\cdot\|)$ be a Banach space and let $I \in C^1(W, \mathbb{R})$ a functional such that the following conditions are satisfied:

(1) I(0,0) = 0;

(2) There exist $\rho > 0$ and r > 0 such that $I(u, v) \ge \rho$ for all $u, v \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ with ||(u, v)|| = r;

(3) There exist $e \in W$ with ||e|| > r such that I(e) < 0; define

$$c \coloneqq \inf_{g \in \Gamma} \sup_{t \in [0,1]} I(g(t)),$$

where

$$\Gamma := \left\{ g \in C^0([0,1], \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)) \colon g(0) = 0, g(e) < 0 \right\}.$$

Then $c \ge \rho > 0$, and there exists a $(PS)_c$ sequence $\{(u_k, v_k)\} \subset W$ for I at level c, i.e.,

$$\lim_{k \to +\infty} I(u_k, v_k) = c \quad \text{and} \quad \lim_{k \to +\infty} I'(u_k, v_k) = 0 \quad \text{strongly in } W'$$

The proof of Proposition 10 follows from the next two lemmas.

Lemma 12. The functional I verifies the assumptions of Lemma 11.

Proof. Clearly, we have I(0,0) = 0. We now verify the second assumption of Lemma 11. Recalling the definition (10) of the quadratic form Q^* and using inequality (14), for any $(u, v) \in W$ we obtain

$$\begin{split} I(u,v) &\ge \frac{1}{p} \| (u,v) \|_{W}^{p} - \frac{C}{2p_{s}^{\sharp}} \Big[\| u \|_{\dot{W}_{\theta}^{s,p}(\mathbb{R}^{N})}^{2p_{s}^{\sharp}} + \| v \|_{\dot{W}_{\theta}^{s,p}(\mathbb{R}^{N})}^{2p_{s}^{\sharp}} \Big] - \frac{1}{p_{s}^{*}} Q^{*}(u,v) \\ &\ge \frac{1}{p} \| (u,v) \|_{W}^{p} - \frac{C}{2p_{s}^{\sharp}} \Big[\| u \|_{\dot{W}_{\theta}^{s,p}(\mathbb{R}^{N})}^{p} + \| v \|_{\dot{W}_{\theta}^{s,p}(\mathbb{R}^{N})}^{p} \Big] \frac{2p_{s}^{\sharp}}{p} - \frac{1}{p_{s}^{*}} Q^{*}(u,v) \\ &\ge \frac{1}{p} \| (u,v) \|_{W}^{p} - C_{1} \| (u,v) \|^{2p_{s}^{\sharp}} - C_{2} \| (u,v) \|^{p_{s}^{*}}. \end{split}$$

Since $s \in (0,1), 0 < \alpha, \beta < sp + \theta < N$ and $\mu \in (0,N)$, we have that $p_s^*(\beta,\theta) > p$ and $2p_s^{\sharp} > p_s^*(\alpha,\theta) > p$. Therefore, there exists r > 0 small enough such that

$$\inf_{\|(u,v)\|=r} I(u,v) > \rho,$$

so item (2) of Lemma 11 are satisfied.

For $(u, v) \in W$ and $t \in \mathbb{R}_+$, we have

$$I(tu, tv) = \frac{t^p}{p} ||(u, v)||^p - \frac{t^{2p_s^{\sharp}}}{2p_s^{\sharp}} Q^{\sharp}(u, v) - \frac{t^{p_s^{\star}}}{p_s^{\star}} Q^{\star}(u, v);$$

since $2p_s^{\sharp} > p_s^*(\alpha, \theta) > p$, we deduce that

$$\lim_{t \to +\infty} I(tu, tv) = -\infty \quad \text{for any } (u, v) \in W.$$

Consequently, for any fixed $e \in W$, there exists $t_e > 0$ such that $||t_e e|| > r$ and $I(t_e e) < 0$. Thus, item (3) of Lemma 11 is satisfied.

From Lemma 12 above, we guarante by Lemma 11 the existence of a Palais-Smale sequence $\{(u_k, v_k)\} \subset W$ such that

$$\lim_{k \to +\infty} I(u_k, v_k) = c \quad \text{and} \quad \lim_{k \to +\infty} I'(u_k, v_k) = 0 \quad \text{strongly in } W'.$$

Moreover, by the definition of c we deduce that $c \ge \rho > 0$. Therefore c > 0 for all function $(u, v) \in W \setminus \{(0, 0)\}.$

Lemma 13. Suppose that $\mu \in (0, N)$ and that $0 < \alpha < sp + \theta$. Then there exists $(u, v) \in W \setminus \{(0, 0)\}$ such that $c \in (0, c^*)$, where c^* is defined in (34).

Proof. Using Proposition 9, we obtain the minimizers $(u_1, v_1) \in W$ for S^{\sharp} and $(u_2, v_2) \in W$ for S^* , respectively. Thus, there exist a function $(e_1, e_2) \in W$ defined by

$$(e_1, e_2) = \begin{cases} (u_1, v_1), & \text{if } \frac{2p_s^{\sharp} - p}{2pp_s^{\sharp}} S_{\mu}(N, s, \gamma, \alpha)^{\frac{2p_{\mu}^{\sharp}(\alpha, \theta)}{2p_{\mu}^{\sharp}(\alpha, \theta) - p}} \leqslant \frac{sp - \beta}{p(N - \beta)} \Lambda(N, s, \gamma, \beta)^{\frac{N - \beta}{sp - \beta}} \\ (u_2, v_2), & \text{if } \frac{2p_s^{\sharp}(\delta, \theta, \mu) - p}{2pp_s^{\sharp}} S_{\mu}(N, s, \gamma, \alpha)^{\frac{2p_{\mu}^{\sharp}(\alpha, \theta)}{2p_{\mu}^{\sharp}(\alpha, \theta) - p}} > \frac{sp - \beta}{p(N - \beta)} \Lambda(N, s, \gamma, \beta)^{\frac{N - \beta}{sp - \beta}} \end{cases}$$

such that $||(e_1, e_2)|| > r$ and $I(e_1, e_2) < 0$. We can define

$$c \coloneqq \inf_{g \in \Gamma} \sup_{t \in [0,1]} I(g(t)),$$

where

$$\Gamma \coloneqq \left\{ g \in C^0([0,1], \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)) \colon g(0) = 0, g(e_1, e_2) < 0 \right\}$$

Clearly, we have that c > 0. For the case where $e = (u_1, v_1)$, we can deduce that

$$0 < c < \frac{2p_s^{\sharp} - p}{2pp_s^{\sharp}} S_{\mu}(N, s, \gamma, \alpha)^{\frac{p_s^{\sharp}}{p_{\mu}^{\star}(\alpha, \theta) - 1}}.$$

In fact, for all $t \ge 0$, by the definition of the functional I, we have that

$$I(tu_1, tv_1) \leqslant \frac{t^p}{p} \| (u_1, v_1) \|^p - \frac{t^{2p_s^{\sharp}}}{2p_s^{\sharp}} Q^{\sharp}(u_1, v_1) \eqqcolon f_1(t).$$

It is easy to see that $f'_1(t) = t^{p-1}[||(u_1, v_1)||^p - t^{2p_s^{\sharp}-p}Q^{\sharp}(u_1, v_1)]$. So, $f'_1(\tilde{t}) = 0$ for

(35)
$$\tilde{t} = \left(\frac{\|(u_1, v_1)\|^p}{Q^{\sharp}(u_1, v_1)}\right)^{\frac{1}{2p_s^{\sharp} - p}}$$

and this is a point of maximum for f_1 . Besides of that, this maximum value is

$$f_1(\tilde{t}) = \left[\frac{1}{p} - \frac{1}{2p_s^{\sharp}}\right] \frac{\|(u_1, v_1)\|^{\frac{2pp_s^{\sharp}}{2p_s^{\sharp} - p}}}{Q^{\sharp}(u_1, v_1)^{\frac{p}{2p_s^{\sharp} - p}}} = \left[\frac{2p_s^{\sharp} - p}{2pp_s^{\sharp}}\right] S^{\sharp \frac{2p_s^{\sharp}}{2p_s^{\sharp} - p}}.$$

Therefore,

(36)
$$\sup_{t \ge 0} I(tu_1, tv_1) \leqslant \sup_{t \ge 0} f_1(t) = \frac{2p_s^{\sharp} - p}{2pp_s^{\sharp}(\delta, \theta, \mu)} S_{\sharp}^{\frac{2p_s^{\sharp}}{2p_s^{\sharp} - \mu}}$$

The equality does not hold in (36); otherwise, we would have that $\sup_{t\geq 0} I(tu_1, tv_1) = \sup_{t\geq 0} f_1(t)$. Let $t_1 > 0$ be the point where $\sup_{t\geq 0} I(tu_1, tv_1)$ is attained. We have

$$f_1(t_1) - \frac{t_1^{p_s^*(\beta,\theta)}}{p_s^*(\beta,\theta)} Q^*(u_1,v_1) = f_1(\tilde{t})$$

which means that $f_1(t_1) > f_1(\tilde{t})$, since $t_1 > 0$. This contradicts the fact that \tilde{t} is the unique maximum point for f_1 . Thus, we have strict inequality in (36), that is,

(37)
$$\sup_{t \ge 0} I(tu_1, tv_1) < \sup_{t \ge 0} f_1(t) = \frac{2p_{\mu}^{\sharp}(\alpha, \theta) - p}{2pp_{\mu}^{\sharp}(\alpha, \theta)} S^{\sharp \frac{2p_{\mu}^{\sharp}(\alpha, \theta)}{2p_{\mu}^{\sharp}(\alpha, \theta) - p}}.$$

Therefore, $0 < c < \frac{2p_{\mu}^{\sharp}(\alpha,\theta) - p}{2pp_{\mu}^{\sharp}(\alpha,\theta)} S^{\sharp \frac{2p_{\mu}^{\sharp}(\alpha,\theta)}{2p_{\mu}^{\sharp}(\alpha,\theta) - p}}$. Similarly, for the case of $e = (u_2, v_2)$, we can verify that

(38)
$$\sup_{t \ge 0} I(tu_2, tv_2) < \frac{sp - \beta}{p(N - \beta)} \Lambda(N, s, \gamma, \beta)^{\frac{N - \beta}{sp - \beta}}.$$

In fact, for all $t \ge 0$, by functional I definition we have that

$$I(tu_2, tv_2) \leqslant \frac{t^p}{p} \| (u_2, v_2) \|^p - \frac{t^{p_s^*}}{p_s^*} Q^*(u, v) \coloneqq g_1(t).$$

It is easy to see that $g'_1(t) = t^{p-1} \left[\|(u_2, v_2)\|^p - t^{p_s^* - p} Q^*(u, v) \right]$. So, $g_1(\tilde{t}) = 0$ for

$$\tilde{t} = \left(\frac{\|(u_2, v_2)\|^p}{Q^*(u, v)}\right)^{\frac{1}{p_s^* - p}}$$

and this is a point of maximum for g_1 . Besides of that, this maximum value is

$$g_1(\tilde{t}) = \left[\frac{1}{p} - \frac{1}{p_s^*}\right] \left(\frac{\|(u_2, v_2)\|^p}{Q^{*\frac{p}{p_s^*}}}\right)^{\frac{p_s}{p_s^* - p}} = \frac{sp + \theta - \beta}{p(N - \beta)} S^{*\frac{N - \beta}{sp + \theta - \beta}}$$

Therefore,

(39)
$$\sup_{t \ge 0} I(tu_2, tv_2) \le \sup_{t \ge 0} g_1(t) = \frac{sp + \theta - \beta}{p(N - \beta)} S^* \frac{N - \beta}{sp + \theta - \beta}.$$

The equality does not hold in (39), otherwise, we would have that $\sup I(tu_2, tv_2) =$ $\sup_{t\geq 0} g_1(t)$. Let $t_1 > 0$, where $\sup_{t\geq 0} I(tu_2, tv_2)$ is attained. We have

$$g_1(t_1) - \frac{t_1^{2p_s^\sharp}}{2p_s^\sharp} Q^\sharp(u_2, v_2) = g_1(\tilde{t})$$

which means that $g_1(t_1) > g_1(\tilde{t})$, since $t_1 > 0$. This contradicts the fact that \tilde{t} is the unique maximum point for $g_1(t)$. Thus

(40)
$$\sup_{t \ge 0} I(tu_2, tv_2) < \sup_{t \ge 0} g_1(t) = \frac{sp + \theta - \beta}{p(N - \beta)} S^* \frac{N - \beta}{sp + \theta - \beta}$$

Therefore, $0 < c < \frac{sp+\theta-\beta}{p(N-\beta)}S^{*\frac{N-\beta}{sp+\theta-\beta}}$.

From the definition (34) of c^* and from inequalities (37) and (40), we have

$$0 < c < c^* \coloneqq \min\left\{ \left(\frac{1}{p} - \frac{1}{2p_s^{\sharp}}\right) S^{\sharp \frac{2p_s^{\sharp}}{2p_s^{\sharp} - p}}, \left(\frac{1}{p} - \frac{1}{p_s^{*}}\right) S^{* \frac{p_s^{*}}{p_s^{*} - p}} \right\}.$$

The lemma is proved.

Proof of Proposition 10. Follows immediately from Lemmas 12 and 13.

5. Proof of Theorem 1

The existence of a solution will follow from the proof of the Theorem 1.

Proof of Theorem 1. Suppose that $s \in (0, 1), 0 < \alpha, \beta < sp + \theta, \mu \in (0, N)$ and $\gamma < \gamma_H$. Let $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$ be a Palais-Smale sequence $(PS)_c$ as in Proposition 10, i.e.,

$$I(u_k, v_k) \to c, \ I'(u_k, v_k) \to 0$$
 strongly in W' as $k \to +\infty$

Then

(41)
$$I(u_k, v_k) = \frac{1}{p} ||(u_k, v_k)||^p - \frac{1}{2p_s^{\sharp}} Q^{\sharp}(u_k, u_k) - \frac{1}{p_s^{\ast}} Q^{\ast}(u_k, v_k) = c + o(1)$$

and

(42)
$$\langle I'(u_k, v_k), (u_k, v_k) \rangle = \|(u_k, v_k)\|^p - Q^{\sharp}(u_k, v_k) - Q^*(u_k, v_k) = o(1).$$

From (41) and (42), if $2p_s^{\sharp} \ge p_s^* > p$, we have

$$\begin{aligned} c + o(1) \|(u_k, v_k)\| &= I(u_k, v_k) - \frac{1}{p_s^*} \langle I'(u_k, v_k), (u_k, v_k) \rangle \\ &= \frac{p_s^* - p}{p \cdot p_s^*} \|(u_k, v_k)\|^p + \left(\frac{1}{p_s^*} - \frac{1}{2p_s^\sharp}\right) Q^\sharp(u_k, v_k) \\ &\geqslant \frac{p_s^* - p}{p \cdot p_s^*} \|(u_k, v_k)\|^p. \end{aligned}$$

Again from (41) and (42), if $p_s^* > 2p_s^{\sharp} > p$, we have

$$\begin{aligned} c + o(1) \| (u_k, v_k) \| &= I(u_k, v_k) - \frac{1}{2p_s^{\sharp}} \langle I'(u_k, v_k), (u_k, v_k) \rangle \\ &= \frac{2p_s^{\sharp} - p}{p \cdot 2p_s^{\sharp}} \| (u_k, v_k) \|^p + \left(\frac{1}{2p_s^{\sharp}} - \frac{1}{p_s^{*}}\right) Q^*(u_k, v_k) \\ &\geqslant \frac{2p_s^{\sharp} - p}{p \cdot 2p_s^{\sharp}} \| (u_k, v_k) \|^p. \end{aligned}$$

Thus, $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$ is a bounded sequence; so from the estimate (42) there exists a subsequence, still denoted by $\{(u_k, v_k)\}_{k \in \mathbb{N}} \subset W$, such that $||(u_k, v_k)||^p \to b, Q^*(u_k, v_k) \to d_1, Q^{\sharp}(u_k, u_k) \to d_2$, as $k \to +\infty$; additionally, $b = d_1 + d_2$. By the definitions of S^{\sharp} and S^* , we get

$$d_1^{\frac{p}{p_s^*}}S^* \leqslant b = d_1 + d_2, \qquad d_2^{\frac{1}{p_s^*}}S^{\sharp} \leqslant b = d_1 + d_2.$$

From the first inequality we have $d_1^{\frac{p}{p_s^*}}S^* - d_1 \leq d_2$, that is

(43)
$$d_1^{\frac{p}{p_s^*}} \left(S^* - d_1^{\frac{p^* - p}{p_s^*}} \right) \leqslant d_2.$$

And from the second inequality we have $d_2^{\frac{1}{p_s}}S^{\sharp} - d_2 \leq d_1$, that is,

(44)
$$d_{2}^{\frac{1}{p_{s}^{\sharp}}} \left(S^{\sharp} - d_{2}^{\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}}} \right) \leqslant d_{1}.$$

Claim 2. We have

$$S^* - d_1^{\frac{p_s^* - p}{p_s^*}} > 0, \qquad S^{\sharp} - d_2^{\frac{p_s^{\sharp} - 1}{p_s^{\sharp}}} > 0.$$

Proof. In fact, since $c + o(1) ||(u_k, v_k)|| = I(u_k, v_k) - \frac{1}{p} \langle I'(u_k, v_k), (u_k, v_k) \rangle$, we have

$$I(u_k, v_k) - \frac{1}{p} \langle I'(u_k, v_k), (u_k, v_k) \rangle = \left(\frac{1}{p} - \frac{1}{2p_{\mu}^{\sharp}(\alpha, \theta)}\right) Q^{\sharp}(u_k, u_k) + \left(\frac{1}{p} - \frac{1}{p_s^*}\right) Q^*(u_k, v_k)$$
$$= c + o(1) ||(u_k, v_k)||.$$

Passing to the limit as $k \to +\infty$, we get

(45)
$$\left(\frac{1}{p} - \frac{1}{p_s^*}\right) d_1 + \left(\frac{1}{p} - \frac{1}{2p_\mu^\sharp(\alpha, \theta)}\right) d_2 = c;$$

so,

$$d_1 \leqslant \left(\frac{1}{p} - \frac{1}{p_s^*}\right)^{-1} c = \frac{p(N - \beta)}{sp + \theta - \beta} c, \qquad d_2 \leqslant \left(\frac{1}{p} - \frac{1}{2p_{\mu}^{\sharp}(\alpha, \theta)}\right)^{-1} c = \frac{2pp_s^{\sharp}}{2p_s^{\sharp} - p} c.$$

Using these upper bounds for d_1, d_2 and the fact $0 < c < c^*$, we have

$$S^* - d_1^{\frac{p_s^* - p}{p_s^*}} \ge S^* - \left[\frac{p(N - \beta)}{sp + \theta - \beta} c\right]^{\frac{p_s^* - p}{p_s^*}} > S^* - \left[\frac{p(N - \beta)}{sp + \theta - \beta} c^*\right]^{\frac{p_s^* - p}{p_s^*}}$$
$$\ge S^* - \left[\frac{p(N - \beta)}{sp + \theta - \beta} \cdot \frac{(sp + \theta - \beta)}{p(N - \beta)} S^* \frac{N - \beta}{(sp + \theta - \beta)}\right]^{\frac{p_s^* - p}{p_s^*}} = S^* - S^* = 0.$$

Similarly,

$$\begin{split} S^{\sharp} - d_{2}^{\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}}} \geqslant S^{\sharp} - \left[\frac{2pp_{s}^{\sharp}}{2p_{s}^{\sharp}-p}c\right]^{\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}}} > S^{\sharp} - \left[\frac{2pp_{s}^{\sharp}}{2p_{s}^{\sharp}-p}c^{*}\right]^{\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}}} \\ \geqslant S^{\sharp} \left[\frac{2pp_{s}^{\sharp}}{(2p_{s}^{\sharp}-p)} \cdot \frac{(2p_{s}^{\sharp}-p)}{2pp_{s}^{\sharp}}S^{\sharp\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}-1}}\right]^{\frac{p_{s}^{\sharp}-1}{p_{s}^{\sharp}}} = S^{\sharp} - S^{\sharp} = 0. \end{split}$$

This concludes the proof of the claim.

Following up, inequalities (43) and (44) imply, respectively, that

$$\left[S^* - \left(\frac{p(N-\beta)}{sp+\theta-\beta}c\right)^{\frac{p_s^*-p}{p_s^*}}\right] d_1^{\frac{p}{p_s^*}} \leqslant \left[S^* - d_1^{\frac{p_s^*-p}{p_s^*}}\right] d_1^{\frac{p}{p_s^*}} \leqslant d_2$$

and

$$\left[S^{\sharp} - \left(\frac{2pp_{s}^{\sharp}}{2p_{s}^{\sharp} - p}c\right)^{\frac{p_{s}^{\sharp} - 1}{p_{s}^{\sharp}}}\right] d_{2}^{\frac{1}{p_{s}^{\sharp}}} \leqslant \left[S^{\sharp} - d_{2}^{\frac{p_{s}^{\sharp} - 1}{p_{s}^{\sharp}}}\right] d_{2}^{\frac{1}{p_{s}^{\sharp}}} \leqslant d_{1}.$$

If $d_1 = 0$ and $d_2 = 0$, then (45) implies that c = 0, which is in contradiction with c > 0. Therefore, $d_1 > 0$ and $d_2 > 0$ and we can choose $\epsilon_0 > 0$ such that $d_1 \ge \epsilon_0 > 0$ and $d_2 \ge \epsilon_0 > 0$; moreover, there exists a $K \in \mathbb{N}$ such that

$$Q^*(u_k, v_k) > \frac{\epsilon_0}{2}, \qquad Q^{\sharp}(u_k, u_k) > \frac{\epsilon_0}{2}$$

for every k > K. The embeddings (11), and the improved Sobolev inequality (15) imply that there exist $C_1, C_2 > 0$ such that

$$0 < C_2 \leqslant \|u_k\|_{L^{p,N-sp-\theta+pr}_M(\mathbb{R}^N,|y|^{-pr})} \leqslant C_1,$$

where $r = \frac{\alpha}{p_*^*(\alpha,\theta)}$. For any k > K, we may find $\lambda_k > 0$ and $x_k \in \mathbb{R}^N$ such that

$$\lambda_k^{(N-sp-\theta+pr)-N} \int_{B_{\lambda_k}(x_k)} \frac{|u_k(y)|^p}{|y|^{pr}} \mathrm{d}y > \|u_k\|_{L^{p,N-sp-\theta+pr}_M(\mathbb{R}^N,|y|^{-pr})} - \frac{C}{2k} \ge \tilde{C} > 0.$$

Now we define the sequence $\{\tilde{u}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ by $\tilde{u}_k(x) = \lambda_k^{\frac{N-sp-\theta}{p}}u_k(\lambda_k x)$ and the sequence $\{\tilde{v}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ by $\tilde{v}_k(x) = \lambda_k^{\frac{N-sp-\theta}{p}}v_k(\lambda_k x)$. As we have already shown, $\|\tilde{u}_k\| = \|u_k\| \leq C$ and $\|\tilde{v}_k\| = \|v_k\| \leq C$ for every $k \in \mathbb{N}$; so, there exist $u \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ and $v \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ such that, after passage to subsequence, still denoted in the same way,

$$\tilde{u}_k \rightharpoonup u \text{ in } \dot{W}^{s,p}_{\theta}(\mathbb{R}^N) \qquad \text{and} \quad \tilde{v}_k \rightharpoonup v \text{ in } \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$$

as $k \to +\infty$. In a fashion similar to the proof of [3, Proposition 2-1], we can prove that $u \neq 0$ and $v \neq 0$.

In addition, the boundedness of the sequences $\{\tilde{u}_k\}_{k\in\mathbb{N}} \subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ and $\{\tilde{v}_k\}_{k\in\mathbb{N}} \subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ implies that the sequences $\{|\tilde{u}_k|^{p_s^*-2}\tilde{u}_k\}_{k\in\mathbb{N}} \subset L^{\frac{p_s^*}{p_s^*-1}}(\mathbb{R}^N, |x|^{-\beta})$ and $\{|\tilde{v}_k|^{p_s^*-2}\tilde{v}_k\}_{k\in\mathbb{N}} \subset L^{\frac{p_s^*}{p_s^*-1}}(\mathbb{R}^N, |x|^{-\beta})$ are bounded too. In fact, by embeddings (11), we obtain

$$\int_{\mathbb{R}^N} \frac{\left| |\tilde{u}_k|^{p_s^* - 2} \cdot \tilde{u}_k \right|^{\frac{p_s}{p_s^* - 1}}}{|x|^\beta} \mathrm{d}x = \int_{\mathbb{R}^N} \frac{|\tilde{u}_k|^{p_s^*}}{|x|^\beta} \mathrm{d}x < C.$$

and

$$\int_{\mathbb{R}^N} \frac{\left| |\tilde{v}_k|^{p_s^* - 2} \cdot \tilde{v}_k \right|^{\frac{p_s}{p_s^* - 1}}}{|x|^\beta} \mathrm{d}x = \int_{\mathbb{R}^N} \frac{|\tilde{v}_k|^{p_s^*}}{|x|^\beta} \mathrm{d}x < C.$$

Then, after passage to subsequence, still denoted in the same way, we deduce that

(46)
$$|\tilde{u}_k|^{p_s^*-2}\tilde{u}_k \rightarrow |u|^{p_s^*-2}u \quad \text{in } L^{\frac{p_s^*}{p_s^*-1}}(\mathbb{R}^N, |x|^{-\beta})$$

and

(47)
$$|\tilde{v}_k|^{p_s^*-2}\tilde{v}_k \rightharpoonup |v|^{p_s^*-2}v \quad \text{in } L^{\frac{p_s^*}{p_s^*-1}}(\mathbb{R}^N, |x|^{-\beta})$$

as $k \to +\infty$.

For any $\phi_1, \phi_2 \in L^{p_s^*(\alpha,\theta)}(\mathbb{R}^N, |x|^{-\alpha})$, Lemma 2.5 in [4] implies that

(48)
$$\lim_{k \to \infty} \int_{\mathbb{R}^N} \left[I_{\mu} * F_{\alpha}(\cdot, \tilde{u}_k) \right](x) f_{\alpha}(x, \tilde{u}_k) \phi_1(x) dx$$
$$= \int_{\mathbb{R}^N} \left[I_{\mu} * F_{\alpha}(\cdot, u) \right](x) f_{\alpha}(x, u) \phi_1(x) dx$$

and

(49)
$$\lim_{k \to \infty} \int_{\mathbb{R}^N} \left[I_{\mu} * F_{\alpha}(\cdot, \tilde{v}_k) \right](x) f_{\alpha}(x, \tilde{v}_k) \phi_2(x) \mathrm{d}x$$
$$= \int_{\mathbb{R}^N} \left[I_{\mu} * F_{\alpha}(\cdot, v) \right](x) f_{\alpha}(x, v) \phi_2(x) \mathrm{d}x.$$

Since $\dot{W}^{s,p}_{\theta}(\mathbb{R}^N) \hookrightarrow L^{p^*_s(\alpha,\theta)}(\mathbb{R}^N, |x|^{-\alpha})$, (48) and (49) hold for any $\phi_1, \phi_2 \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$.

Finally, we need to check that $\{\tilde{u}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ and $\{\tilde{v}_k\}_{k\in\mathbb{N}}\subset \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ are also a $(PS)_c$ sequence for the functional I at energy level c. Do to this, we note that the norms

in $L^{p_s^*(\alpha,\theta)}(\mathbb{R}^N, |x|^{-\alpha})$ are invariant under the special dilatation $\tilde{u}_k = \lambda_k^{\frac{N-sp-\theta}{p}} u_k(\lambda_k x)$ and $\tilde{v}_k = \lambda_k^{\frac{N-sp-\theta}{p}} v_k(\lambda_k x)$. In fact

$$\|\tilde{u}_k\|_{L^{p_s^*(\alpha,\theta)}}^{p_s^*(\alpha,\theta)} = \int_{\mathbb{R}^N} \frac{\lambda_k^{\frac{N-sp-\theta}{p}} p_s^*(\alpha,\theta)}{|x|^{\alpha}} |u_k(\lambda_k x)|^{p_s^*(\alpha,\theta)}}{|x|^{\alpha}} \mathrm{d}x = \int_{\mathbb{R}^N} \frac{|u_k(\bar{x})|^{p_s^*(\alpha,\theta)}}{|\bar{x}|^{\alpha}} \mathrm{d}\bar{x} = \|u_k\|_{L^{p_s^*(\alpha,\theta)}}^{p_s^*(\alpha,\theta)}$$

and

$$\|\tilde{v}_{k}\|_{L^{p_{s}^{*}(\beta,\theta)}}^{p_{s}^{*}(\beta,\theta)} = \int_{\mathbb{R}^{N}} \frac{\lambda_{k}^{\frac{N-sp-\theta}{p}} p_{s}^{*}(\beta,\theta)}{|x|^{\beta}} |v_{k}(\lambda_{k}x)|^{p_{s}^{*}(\beta,\theta)}}{dx} = \int_{\mathbb{R}^{N}} \frac{|v_{k}(\bar{x})|^{p_{s}^{*}(\beta,\theta)}}{|\bar{x}|^{\beta}} d\bar{x} = \|v_{k}\|_{L^{p_{s}^{*}(\beta,\theta)}}^{p_{s}^{*}(\beta,\theta)}.$$

Additionally, we have

$$\int_{\mathbb{R}^N} \frac{\eta |\tilde{u}_k(x)|^a |\tilde{v}_k(x)|^b}{|x|^\beta} \mathrm{d}x = \int_{\mathbb{R}^N} \frac{\eta \lambda_k^{\frac{N-sp-\theta}{p}(a+b)} |u_k(\lambda_k x)|^a |v_k(\lambda_k x)|^b}{|x|^\beta} \mathrm{d}x$$
$$= \int_{\mathbb{R}^N} \frac{\eta |u_k(\bar{x})|^a |v_k(\bar{x})|^b}{|\bar{x}|^\beta} \mathrm{d}\bar{x}.$$

Thus, we have

$$\lim_{k \to +\infty} I(\tilde{u}_k, \tilde{v}_k) = c$$

Moreover, for all $\phi_1, \phi_2 \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$, we have $\phi_{1,k}(x) = \lambda_k^{\frac{N-sp-\theta}{p}} \phi_1(x/\lambda_k) \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$ and $\phi_{2,k}(x) = \lambda_k^{\frac{N-sp-\theta}{p}} \phi_2(x/\lambda_k) \in \dot{W}^{s,p}_{\theta}(\mathbb{R}^N)$. From $I'(u_k, v_k) \to 0$ in W' as $k \to +\infty$, we can deduce that

$$\lim_{k \to +\infty} \langle I'(\tilde{u}_k, \tilde{v}_k), (\phi_1, \phi_2) \rangle = \lim_{k \to +\infty} \langle I'(u_k, v_k), (\phi_1, \phi_2) \rangle = 0.$$

Thus (46), (47), (48) and (49) lead to

$$\langle I'(u,v), (\phi_1, \phi_2) \rangle = \lim_{k \to +\infty} \langle I'(\tilde{u}_k, \tilde{v}_k), (\phi_1, \phi_2) \rangle = 0.$$

Hence (u, v) is a nontrivial weak solution to problem 1.

Proof of Theorem 2. The proof follows the same steps of the proof of Theorem 1. Here we only remark that for problem (5) with a Hardy potential and double Sobolev type nonlinearities we have to define the value below which we can recover the compactness of the Palais-Smale sequences by

$$c^* := \min_{k \in \{1,2\}} \left\{ \left(\frac{1}{p} - \frac{1}{p_s^*} \right) S^*(N, s, \gamma, \beta_k)^{\frac{p_s^*(\beta_k, \theta)}{p_s^*(\beta_k, \theta) - p}} \right\}.$$

Similarly, for problem (6) with a Hardy potential and double Choquard type nonlinearities we have to define the corresponding number by

$$c^* \coloneqq \min_{k \in \{1,2\}} \left\{ \left(\frac{1}{p} - \frac{1}{2p_s^{\sharp}(\delta,\theta,\mu_k)}\right) S^{\sharp}(N,s,\gamma,\beta_k)^{\frac{2p_s^{\sharp}(\delta,\theta,\mu_k)}{2p_s^{\sharp}(\delta,\theta,\mu_k) - p}} \right\}.$$

The details are omitted.

References

- B. Abdellaoui and R. Bentifour, "Caffarelli-Kohn-Nirenberg type inequalities of fractional order with applications," J. Funct. Anal., vol. 272, no. 10, pp. 3998–4029, 2017. [Online]. Available: https://doi.org/10.1016/j.jfa.2017.02.007 8
- [2] A. Ambrosetti and P. H. Rabinowitz, "Dual variational methods in critical point theory and applications," *Journal of functional Analysis*, vol. 14, no. 4, pp. 349–381, 1973. 14
- [3] R. B. Assunção, J. C. Silva, and O. H. Miyagaki, "A fractional *p*-Laplacian problem with multiple critical Hardy–Sobolev nonlinearities," *Milan Journal of Mathematics*, vol. 88, no. 1, pp. 65–97, 2020. 5, 20
- [4] R. B. Assunção, O. H. Miyagaki, and R. F. S. Siqueira, "Fractional Sobolev-Chocard critical equation with Hardy term and weighted singularities," 2023, https://arxiv.org/abs/2311.00852. 6, 7, 8, 9, 12, 13, 20
- [5] L. Brasco, S. Mosconi, and M. Squassina, "Optimal decay of extremals for the fractional Sobolev inequality," *Calculus of Variations and Partial Differential Equations*, vol. 55, pp. 1–32, 2016. 7
- [6] —, "Optimal decay of extremals for the fractional Sobolev inequality," Calc. Var. Partial Differential Equations, vol. 55, no. 2, pp. Art. 23, 32, 2016. [Online]. Available: https://doi.org/10.1007/s00526-016-0958-y 8
- [7] C. Chen, "Infinitely many solutions to a class of quasilinear Schrödinger system in rⁿ," Applied Mathematics Letters, vol. 52, pp. 176–182, 2016.
- [8] W. Chen, "Fractional elliptic problems with two critical Sobolev-Hardy exponents," *Electron. J. Differential Equations*, pp. Paper No. 22, 12, 2018.
- [9] ——, "Existence of solutions for fractional p-Kirchhoff type equations with a generalized Choquard nonlinearity," *Topol. Methods Nonlinear Anal.*, vol. 54, no. 2, pp. 773–791, 2019. [Online]. Available: https://doi.org/10.12775/tmna.2019.069 5
- [10] W. Chen and S. Deng, "The nehari manifold for a fractional p-laplacian system involving concaveconvex nonlinearities," Nonlinear Analysis: Real World Applications, vol. 27, pp. 80–92, 2016.
- [11] W. Chen and M. Squassina, "Critical nonlocal systems with concave-convex powers," Advanced Nonlinear Studies, vol. 16, no. 4, pp. 821–842, 2016. 5, 6
- [12] E. Di Nezza, G. Palatucci, and E. Valdinoci, "Hitchhiker's guide to the fractional Sobolev spaces," Bull. Sci. Math., vol. 136, no. 5, pp. 521–573, 2012. [Online]. Available: https://doi.org/10.1016/j.bulsci.2011.12.004 5
- [13] R. Filippucci, P. Pucci, and F. Robert, "On a p-Laplace equation with multiple critical nonlinearities," J. Math. Pures Appl. (9), vol. 91, no. 2, pp. 156–177, 2009. [Online]. Available: https://doi.org/10.1016/j.matpur.2008.09.008 5
- [14] A. Fiscella, P. Pucci, and S. Saldi, "Existence of entire solutions for Schrödinger-hardy systems involving two fractional operators," *Nonlinear Analysis*, vol. 158, pp. 109–131, 2017.
- [15] T. Gantumur, "An introduction to the Morrey and Campanato spaces," https://www.math.mcgill.ca/gantumur/math580f12/MorreyCampanato.pdf, January 2012. 6
- [16] N. Ghoussoub and S. Shakerian, "Borderline variational problems involving fractional Laplacians and critical singularities," Adv. Nonlinear Stud., vol. 15, no. 3, pp. 527–555, 2015. [Online]. Available: https://doi.org/10.1515/ans-2015-0302 5
- [17] G. Li and T. Yang, "The existence of a nontrivial weak solution to a double critical problem involving a fractional Laplacian in ℝⁿ with a Hardy term," Acta Mathematica Scientia, vol. 40, no. 6, pp. 1808–1830, oct 2020. [Online]. Available: https://doi.org/10.1007 9
- [18] E. H. Lieb, "Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities," Ann. of Math. (2), vol. 118, no. 2, pp. 349–374, 1983. [Online]. Available: https://doi.org/10.2307/2007032
- [19] T.-C. Lin and J. Wei, "Ground state of n coupled nonlinear Schrödinger equations in \mathbb{R}^n , $n \leq 3$," Communications in mathematical physics, vol. 255, pp. 629–653, 2005. 6
- [20] —, "Erratum: "Ground state of N coupled nonlinear Schrödinger equations in \mathbb{R}^n , $n \leq 3$ " [Comm. Math. Phys. **255** (2005), no. 3, 629–653; mr2135447]," Comm. Math. Phys., vol. 277, no. 2, pp. 573–576, 2008. [Online]. Available: https://doi.org/10.1007/s00220-007-0365-5_6
- [21] Z. Liu and Z.-Q. Wang, "Ground states and bound states of a nonlinear Schrödinger system," Advanced Nonlinear Studies, vol. 10, no. 1, pp. 175–193, 2010. 6

- [22] G. Lu and Y. Shen, "Existence of solutions to fractional *p*-Laplacian systems with homogeneous nonlinearities of critical Sobolev growth," Adv. Nonlinear Stud., vol. 20, no. 3, pp. 579–597, 2020.
 [Online]. Available: https://doi.org/10.1515/ans-2020-2098 6
- [23] G. Molica Bisci, V. Rădulescu, and R. Servadei, Variational methods for nonlocal fractional problems, ser. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 2016, vol. 162, with a foreword by Jean Mawhin. [Online]. Available: https://doi.org/10.1017/CBO9781316282397 5
- [24] V. Moroz and J. Van Schaftingen, "A guide to the Choquard equation," J. Fixed Point Theory Appl., vol. 19, no. 1, pp. 773–813, 2017. [Online]. Available: https://doi.org/10.1007/s11784-016-0373-15
- [25] S. Mosconi, K. Perera, M. Squassina, and Y. Yang, "The Brezis-Nirenberg problem for the fractional *p*-Laplacian," *Calc. Var. Partial Differential Equations*, vol. 55, no. 4, pp. Art. 105, 25, 2016. [Online]. Available: https://doi.org/10.1007/s00526-016-1035-27, 8
- [26] T. Mukherjee and K. Sreenadh, "Fractional Choquard equation with critical nonlinearities," NoDEA Nonlinear Differential Equations Appl., vol. 24, no. 6, pp. Paper No. 63, 34, 2017. [Online]. Available: https://doi.org/10.1007/s00030-017-0487-15
- [27] H.-M. Nguyen and M. Squassina, "Fractional Caffarelli-Kohn-Nirenberg inequalities," J. Funct. Anal., vol. 274, no. 9, pp. 2661–2672, 2018. [Online]. Available: https://doi.org/10.1016/j.jfa.2017.07.007 8
- [28] G. Palatucci and A. Pisante, "Improved Sobolev embeddings, profile decomposition, and concentration-compactness for fractional Sobolev spaces," *Calc. Var. Partial Differential Equations*, vol. 50, no. 3-4, pp. 799–829, 2014. [Online]. Available: https://doi.org/10.1007/s00526-013-0656-y 8
- [29] Y. Sawano, "A thought on generalized Morrey spaces," J. Indones. Math. Soc., vol. 25, no. 3, pp. 210–281, 2019. 6, 8
- [30] R. Servadei and E. Valdinoci, "Mountain pass solutions for non-local elliptic operators," Journal of Mathematical Analysis and Applications, vol. 389, no. 2, pp. 887–898, 2012. 5
- [31] L. Wang, B. Zhang, and H. Zhang, "Fractional Laplacian system involving doubly critical nonlinearities in rⁿ," *Electronic Journal of Qualitative Theory of Differential Equations*, vol. 2017, no. 57, pp. 1–17, 2017. 6
- [32] M. Xiang, B. Zhang, and V. D. Rădulescu, "Multiplicity of solutions for a class of quasilinear Kirchhoff system involving the fractional p-laplacian," *Nonlinearity*, vol. 29, no. 10, p. 3186, 2016.
- [33] T. Yang, "On doubly critical coupled systems involving fractional Laplacian with partial singular weight," *Mathematical Methods in the Applied Sciences*, vol. 44, no. 17, pp. 13448–13467, 2021.

Ronaldo B. Assunção

DEPARTAMENTO DE MATEMÁTICA — UNIVERSIDADE FEDERAL DE MINAS GERAIS, UFMG Email address: ronaldo@mat.ufmg.br

Olímpio H. Miyagaki

DEPARTAMENTO DE MATEMÁTICA — UNIVERSIDADE FEDERAL DE SÃO CARLOS, UFSCAR Email address: ohmiyagaki@gmail.com

RAFAELLA F. S. SIQUEIRA

DEPARTAMENTO DE MATEMÁTICA — CENTRO FEDERAL DE EDUCAÇÃO TECNOLÓGICA, CEFET *Email address:* rafaella.siqueira@cefetmg.br