

Kardar-Parisi-Zhang scaling in time-crystalline matter

Romain Daviet,^{*} Carl Philipp Zelle,^{*} Armin Asadollahi, and Sebastian Diehl
Institut für Theoretische Physik, Universität zu Köln, 50937 Cologne, Germany

We discuss the universal behavior linked to the Goldstone mode associated with the spontaneous breaking of time-translation symmetry in many-body systems, in which the order parameter traces out a limit cycle. We show that this universal behavior is closely tied to Kardar-Parisi-Zhang physics, which can strongly affect the scaling properties in all dimensions. Our work predicts the relevance of KPZ in numerous systems such as nonreciprocal phases in active matter, active magnets, driven-dissipative quantum systems, and synchronization of oscillators.

Introduction – Time-translation symmetry is a fundamental symmetry of matter in and out of equilibrium. Breaking it spontaneously is impossible in thermodynamic equilibrium governed by a bounded Hamiltonian [1, 2], but it can occur far from equilibrium. Since time translations form a continuous symmetry, breaking it spontaneously gives rise to a gapless mode as a consequence of the Goldstone theorem [3–6]. Because time-translation invariance is generically an exact symmetry, this Goldstone mode must be a robust origin of universal scaling throughout extended regions of parameter space in a plethora of nonequilibrium systems.

Breaking of time-translation symmetry can occur both in quantum and classical nonequilibrium systems. A particular incarnation are dynamical limit cycles – or continuous time crystals – where the macroscopic order parameter retains a periodic motion. This phenomenon has recently gained much attention in engineered quantum systems, ranging from ultracold Bose condensates immersed into optical cavities [7–11] over dissipative Bose condensates such as exciton-polariton systems [12–14], magnon condensates [15, 16] and continuous dissipative time crystals [11, 17–20]. But limit cycles in many-body systems are also relevant in classical systems [21–26], in particular through synchronization of oscillators relevant in many different contexts, such as biology or chemistry [27, 28]. Recently, realizations of such phases have gained a lot of interest in active matter set ups, in particular in systems with nonreciprocal interactions [5, 29–37], and also through generalization to ‘active’ magnetic systems [38–41].

In this work, we distill the universal behavior connected to spontaneous time-translation symmetry breaking in driven open systems, with an emphasis on limit-cycle phases. Two key factors determine their universal physics. First, the limit cycle periodicity itself dictates that the Goldstone mode of time translations [4, 5, 42] is an angular variable. Second, the necessarily nonequilibrium character of the underlying dynamics, together with the continuous growth of the limit-cycle phase, implies that this Goldstone mode is subject to the nonlinearity of the paradigmatic Kardar-Parisi-Zhang (KPZ) equation [43–45]. The Goldstone mode dynamics is thus governed by a compact KPZ equation [46–48]. The uni-

versal scaling behavior is then expected to show both a regime governed by the KPZ universality class, but also regimes dominated by the presence of vortex defects allowed by the compactness of the limit-cycle variable [49–52]. These theoretical findings are complemented by numerical simulations of extended van der Pol systems in one and two spatial dimensions.

The compact KPZ equation has been discussed previously in a realization via the phase variable of a broken quantum mechanical phase rotation symmetry. Here, the origin is generally different and does not even require a continuous internal symmetry on the microscopic level. In fact, a number of previous instances of the emergence of KPZ in driven open (quantum) systems can be rationalized based on the time-translation symmetry breaking mechanism, as we argue below.

Time-translation symmetry breaking – A system is said to be time-translation invariant if its generator of dynamics does not depend explicitly on time. Time-translation symmetry is then broken spontaneously, if in the stable state there is an observable collective degree of freedom, that has a time-dependent expectation value $\langle \phi(t, \mathbf{x}) \rangle = \varphi(t)$, or equivalently

$$\langle \phi(t, \mathbf{x}) \phi(0, 0) \rangle \rightarrow \varphi(t) \varphi(0) \text{ for } |\mathbf{x}|, |t| \rightarrow \infty. \quad (1)$$

A limit cycle thus necessitates both, a macroscopic condensation phenomenon as a prerequisite as in usual symmetry breaking scenarios, and then a synchronization at large distances and times in the presence of fluctuations. Such a condensation is only possible in a thermodynamically large system, and typically occurs in dimensions larger or equal to three. However, in low dimensions, symmetry breaking and Goldstone modes are well known to be, in and out of equilibrium, fruitful starting points around which to expand in order to understand the long-wavelength physics [53–55].

Our approach encompasses both quantum and classical systems driven out of equilibrium. In the more general quantum case, the system is microscopically described by a Lindbladian, or equivalently by a Schwinger-Keldysh generating functional [56, 57], which takes the form (see supplemental material (SM) for details [58])

$$Z[\tilde{\mathbf{J}}, \mathbf{J}] = \int \mathcal{D}\phi \mathcal{D}\tilde{\phi} \exp(iS[\phi, \tilde{\phi}] + i\tilde{\mathbf{J}}\phi + i\mathbf{J}\tilde{\phi}), \quad (2)$$

where $\mathbf{J}, \tilde{\mathbf{J}}$ are source terms, which are needed to extract response and correlations functions, and $\phi, \tilde{\phi}$ are multi-component fields, which can be chosen real without loss of generality [59]. ϕ is the order parameter field, while $\tilde{\phi}$ is its corresponding response field. For the description of collective effects in generic driven open quantum systems, a semiclassical limit can be taken [57, 60], in which the Schwinger-Keldysh action reduces to the Martin-Siggia-Rose-Janssen-De Dominicis action (MSRJD) [61–63]. While still able to capture effects of quantum mechanical origin like phase coherence, this description is formally equivalent to classical stochastic Langevin dynamics.

In this formulation, a system is time-translation symmetric, if the microscopic action $S[\phi, \tilde{\phi}]$ does not depend on time explicitly but only implicitly through $\phi(t), \tilde{\phi}(t)$. The system spontaneously breaks (weak) time-translation symmetry, if the noise-averaged expectation value of the classical field is explicitly time dependent, $\partial_t \langle \phi(t) \rangle = \partial_t \varphi(t) \neq 0$ [64]. In the case of a time crystal, $\langle \phi(t) \rangle$ traces out a closed orbit with period T , which can be parametrized by an angular variable $\theta_0 \in [0, 2\pi)$, $\varphi(t + \theta_0 \bar{T})$ ($\bar{T} = \frac{T}{2\pi}$), see also Fig. 1. Discrete time translations by T remain unbroken. This is analogous to an ordinary crystal breaking continuous spatial translations down to discrete ones, but the time crystal is only possible out of equilibrium [1, 2].

In particular, if $\varphi(t)$ is a valid stable state — a solution of the equation of motion — so is $\varphi(t + \theta_0 \bar{T})$. Formally, we thus have a continuous manifold of equivalent stable states, with translations between the different states mediated by θ_0 . These transitions are gapless, since the Keldysh action for different θ_0 is degenerate, constituting the soft Goldstone mode (see SM for a more formal derivation). Again, the argument is analogous to the phonon in an ordinary crystal. We will refer to the Goldstone mode of broken time translations as the “chronon”.

To construct the chronon excitations, according to the usual procedure for Goldstone modes, we now promote the global symmetry transformation to a local, slowly varying one: The constant time shift becomes $\theta_0 \rightarrow \theta(t, \mathbf{x})$, which varies on time scales much larger than the limit cycle period T . In total we can parametrize spatial fluctuations around the limit cycle within the path integral as

$$\phi(t, \mathbf{x}) = \varphi(t + \bar{T}\theta(t, \mathbf{x}), \mathbf{x}) + \mathbf{N}(t, \mathbf{x}), \quad (3)$$

where $\theta(t, \mathbf{x})$ corresponds to local, gapless fluctuations along the limit cycle and $\mathbf{N}(t, \mathbf{x})$ collects the gapped excitations perpendicular to it. Clearly, the chronon is represented by a compact $SO(2)$ Goldstone mode due to the limit cycle periodicity, see again Fig. 1 for a visualization.

Effective field theory for the Goldstone mode – To capture the effective dynamics of the chronon, we need to integrate out the gapped fluctuations $\mathbf{N}(t, \mathbf{x})$ and their respective noises. Before doing this explicitly, we




FIG. 1. Visualization of a generic limit cycle $\varphi_0(t)$. The Goldstone mode $\theta(t, \mathbf{x})$ corresponds to fluctuations along the limit cycle in the direction $\hat{e}_\theta = \dot{\varphi}_0(t)/\|\dot{\varphi}_0(t)\|$, while $\hat{e}_N(t, \mathbf{x})$ collects all remaining, directions perpendicular to the limit cycle, fluctuations along which are gapped. Note that in higher dimensional phase spaces, the limit cycle is still a one-dimensional closed curve. One can think of the chronon θ as the angle-like variable parametrising this curve. This does not exclude oscillations of single real degrees of freedom, as their phase space is constituted of $\phi = (\phi, \partial_t \phi)^T$.

construct the remaining action $S[\tilde{\theta}(t, \mathbf{x}), \varphi(t + \theta(t, \mathbf{x})\bar{T})]$ based on symmetry. Since we are expanding around a time-dependent stable state, this will lead to an explicitly time-dependent action. Because $\theta(t, \mathbf{x})$ is fluctuating on timescales much larger than the period of the limit cycle $\tau \gg T$, we can eliminate these time-dependencies by averaging over a period T . Afterwards, time-translation invariance of the original model implies invariance under a constant shift $\theta(t, \mathbf{x}) + c$ and the effective field theory for the chronon is gapless, see SM for details [58]. To leading order, it has to take the form

$$S = \int_{t, \mathbf{x}} \tilde{\theta} \left(\partial_t \theta - Z \nabla^2 \theta + \frac{g}{2} (\nabla \theta)^2 \right) - D \tilde{\theta}^2, \quad (4)$$

which is equivalent to the KPZ equation, as seen upon passing from the MSRJD/Keldysh path integral to the stochastic Langevin formulation [57, 60, 65].

The quadratic sector here describes a noisy, diffusive Goldstone mode. The only symmetry allowed interaction is the KPZ coupling $\sim g\theta(\nabla\theta)^2$, which can be relevant and can therefore impact the long time and distance scaling behavior. In addition, there is a notion of directionality – the phase θ grows continuously in the direction the limit cycle is tracing out, see Fig. 1. This excludes a reflection symmetry $\theta \rightarrow -\theta$, and thus the KPZ coupling is symmetry allowed and will be generated under coarse graining.

Scaling predictions – This framework allows us to predict the scaling behavior in time crystals. In low dimensions $d = 1, 2$, the KPZ coupling is relevant, and we expect strong fluctuations linked to the non-diffusive behavior of the chronon, measured by the phase correlator $C_\theta(t, \mathbf{x}) = \langle (\theta(t, \mathbf{x}) - \theta(0, 0))^2 \rangle$. Let us discuss how this impacts generic correlation functions. Since

the limit cycle is periodic with period $T = 2\pi/\Omega$, we can Fourier expand the order parameter field as $\varphi(t) = \sum_n \mathbf{a}_n \cos(n\Omega t + \delta_n)$, and fluctuations $\theta(t, \mathbf{x})$ along the limit cycle thus lead to $\phi(t, \mathbf{x}) = \sum_n \mathbf{a}_n \cos((n\Omega t + \theta(t, \mathbf{x}) + \delta_n))$. By means of a cumulant expansion, the correlation function of these fluctuations thus reads

$$\begin{aligned} C(t, \mathbf{x}) &= \langle \phi(t, \mathbf{x}) \cdot \phi(0, 0) \rangle \\ &= \sum_n \mathbf{a}_n^2 \cos(n\Omega t + \delta_n) e^{-\frac{n^2}{2} \langle (\theta(t, \mathbf{x}) - \theta(0, 0))^2 \rangle_c} \\ &\sim e^{-\frac{1}{2} \langle (\theta(t, \mathbf{x}) - \theta(0, 0))^2 \rangle_c}, \end{aligned} \quad (5)$$

where in the last line, we specify the leading scaling of the envelope of the correlation function. If the phase fluctuations are within the KPZ universality class, this means that

$$\ln C(t, \mathbf{x}) = -A|\mathbf{x}|^{2\chi} \mathcal{C}(t/\mathbf{x}^z) \Rightarrow \begin{cases} C(t, 0) \sim -At^{2\beta} \\ C(0, \mathbf{x}) \sim -B|\mathbf{x}|^{2\chi} \end{cases}, \quad (6)$$

where χ is the roughness exponent, $\beta = z\chi$ with z the dynamical critical exponent, and \mathcal{C} a universal scaling function [52, 66–69]. The exponents are known exactly in one dimension $d = 1$: $\beta = 1/3$, $\chi = 1/2$, and approximately in higher dimensions. Specifically, $\beta \approx 0.24$ and $\chi \approx 0.39$ [70] in $d = 2$.

Adding to these smooth long-wavelength KPZ fluctuations, there is a second nonlinear effect: The phase compactness allows for the existence of topological defects, space-time vortices and solitons in $d = 1$ [51, 71], and (spatial or space-time) vortices in $d = 2$ [49, 50, 52]. Once present, these defects will cause exponential decay of the correlation functions beyond a scale L_v set by their mean separation. In turn, $L_v \sim e^{\Delta/\sigma}$, where Δ is the activation action of a defect, and σ the noise level, for not too large nonlinearities. Out of equilibrium, these defects are always activated in $d = 1, 2$ [49, 50, 71], i.e. Δ does not scale with system size. However, at low noise level and on intermediate scales $|\mathbf{x}|, t^{1/z} \ll L_v$, one can observe KPZ scaling.

Chronons in the many-body van der Pol system – As a concrete and paradigmatic example, we consider an array of classical or quantum van der Pol oscillators in d spatial dimensions [73–79]. This model has no continuous internal symmetry which could be broken spontaneously, and naively, one then might not expect any Goldstone mode to occur. It can be considered as an effective model for generic time-translation symmetry breaking phases arising from different contexts such as nonreciprocal field theories as well as coherently driven and dissipative Bose condensates. In the continuum limit, it is given by

$$(\partial_t^2 + (2\gamma + u\phi^2 - Z_1\nabla^2)\partial_t + \omega_0^2 - Z_2\nabla^2)\phi + \xi = 0, \quad (7)$$




FIG. 2. Autocorrelation function $C(x = 0, t)$ obtained by numerical simulations of Eq. (7) discretized for a van der Pol chain of length 1024, averaged over 1000 noise realizations. The parameters are $r = -\gamma = Z_2 = u = 1$, $Z_1 = 3$, and $D = 1$. The autocorrelation function displays rapid oscillations (see inset), but its envelope (red line) shows the expected KPZ scaling at sufficiently large scales as shown through the fit for $t \in [10^2, 10^3]$ (black dotted line), which yields $\beta = 0.31$, which is in very good agreement with the theoretical value. The right bottom inset shows a space-time vortex of $(\phi, \partial_t \phi)$ at higher noise, $D = 2$.

where $\phi(t, \mathbf{x})$ is a real scalar field, and ξ a Gaussian white noise, $\langle \xi(t, \mathbf{x}) \xi(t', \mathbf{x}') \rangle = D\delta(t - t')\delta(\mathbf{x} - \mathbf{x}')$. It features limit cycle oscillations in the regime where the damping is replaced by an antidamping $\gamma < 0$. Performing the adiabatic elimination of the transverse fluctuations and temporal averaging as anticipated above indeed yields a KPZ equation for the phase field, see SM for details [58]

$$\partial_t \theta - Z\nabla^2 \theta + \frac{g}{2}(\nabla \theta)^2 + \xi = 0, \quad (8)$$

where all parameters can be expressed in terms of the limit cycle solutions of the van der Pol oscillator.

This explicitly bridges the microscopic physics of the spatially extended van der Pol system to the macroscopic physics of the chronon, and thus universal scaling behavior. To confirm the predicted KPZ scaling in this system, we perform numerical simulations of the Langevin dynamics (7) in $d = 1$ and $d = 2$. We solve the stochastic differential equation discretized in real space using the Euler–Maruyama method, with a time step equal to $10^{-2}\bar{T}$. The correlation function displays the predicted KPZ scaling in time in $d = 1$, Fig. 2, which has been also observed independently in [80], as well as topological defects. In two dimensions, at low noise level we observe KPZ scaling to excellent accuracy, see Fig. 3. At higher noise level, the topological defects proliferate, and decays are exponential.

The chronon in various platforms – We finally point out several instances of systems where the chronon is ex-




FIG. 3. Simulations of the van der Pol equation in $d = 2$ for a square lattice of size 256×256 , and $r = -\gamma = Z_2 = u = 1$, $Z_1 = 1.25$, averaged over 1000 realizations. (a) The decay of the equal-time correlation displays KPZ scaling with exponent $2\chi \approx 0.76$ for $D = 2$, as shown by the fit (black dashed line). The scaling at large $|\mathbf{x}|$ is cut off by the finite system size. When $D = 2.7$, the scaling is exponential, and is associated with spatial vortices, see inset for a snapshot from the corresponding simulation. (b) The autocorrelation, for $D = 2$, also shows KPZ scaling and the fit (dashed line) yields $\beta = 0.24$. The inset shows the correlation function for different \mathbf{x} plotted in a scaling form, using the fitted exponents. The scaling is consistent with Eq. (6), and other KPZ simulations [52, 70, 72].

pected to emerge, or has already been observed. Driven-dissipative condensates are typically described by a complex, rotating order parameter $\psi_0 = \sqrt{\rho_0} e^{-i\Omega t}$, breaking an internal $U(1)$ symmetry next to time translations [12, 57, 81]. However, both symmetries act identically on the order parameter, namely as a phase shift. The transformation generated by one of the symmetries can be compensated by a properly chosen transformation of the other. Consequently, there is only one independent generator broken, and only one Goldstone mode emerging. This mode has been theoretically demonstrated and experimentally observed to display the behavior of a compact KPZ mode [46, 52, 82]. But, time-translation symmetry breaking also explains and predicts the emergence of KPZ scaling in systems where such an internal continuous symmetry is broken down explicitly either fully, or to discrete ones, such as in coherently driven condensates [83] ($U(1) \searrow \mathbb{Z}_2$) or non-reciprocal magnetic models [30, 84] ($SO(2) \searrow \mathbb{Z}_4$).

Similarly, a stable traveling wave state [21] $\langle \phi(t, \mathbf{x}) \rangle = \phi_0 \cos(\Omega t - \mathbf{q}_0 \cdot \mathbf{x})$ is also predicted to display KPZ-scaling in its fluctuations, since time- and space translations again act identically on the order parameter. Instances here are encountered in magnon condensates [16, 85], or in nonreciprocal active matter with conserved order parameter [31–34, 86]. The latter were indeed predicted to display KPZ physics [87].

Conclusion and outlook – The breaking of the external continuous symmetry of time translations gives rise to a soft mode, the chronon, in turn unleashing universal

scaling behavior in extended parameter regimes. Since time-translation symmetry is not affected by interactions or microscopic anisotropies potentially spoiling internal symmetries, the laid out mechanism is very robust. We have exemplified this general mechanism for the paradigmatic example of extended systems of van der Pol oscillators both analytically and numerically. This dramatically enlarges the range of platforms offering prospects for an experimental realization of the KPZ phase including its roughening transition in three dimensions, in any noisy system that displays synchronization.

We have focused on time-crystalline phases where time-translation symmetry is broken down to a periodic motion, leading to an $SO(2)$ Goldstone mode. The original KPZ equation describing a growing noisy interface with collective variable $\langle \phi(t, \mathbf{x}) \rangle = vt$, arising for instance in unbounded potentials, can likewise be embedded in the time-translation symmetry breaking scenario. Formally, the only difference lies in a linear or a periodic growth of a collective variable, taking values in \mathbb{R} or in $\mathbb{R}/\mathbb{Z} \simeq SO(2)$. This results in fluctuations described by a compact or non-compact Goldstone mode, respectively.

In systems realizing time-crystalline order, as in ‘active’ [38, 39, 41] and non-reciprocal spin systems [29, 40] the respective phases may in addition break internal spin rotation symmetries [5, 6], or space translation symmetry [21, 88]. In general, this leads to additional Goldstone modes that couple to the chronon. Further soft modes can also emerge from conserved quantities giving rise to out-of-equilibrium hydrodynamics. Our theory immedi-

ately predicts that the Goldstone mode of time translations will still be subject to KPZ-like nonlinearities. The couplings to the additional soft modes as well as potentially modified noise structures may lead to new universal behavior establishing novel nonthermal phases of matter with distinct scaling laws. The systematic field theoretic study within nonequilibrium nonlinear σ -models represents an intriguing avenue for future research.

Acknowledgments— We thank Y. Avni, L. Canet, L. Delacretaz, M. Fruchart, R. Hanai, J. Lang, A. Rosch, V. Vitelli for useful discussions. We acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) CRC 1238 project C04 number 277146847.

-
- * These authors contributed equally to this work.
Emails: zelle@thp.uni-koeln.de daviet@thp.uni-koeln.de
- [1] P. Bruno, Impossibility of spontaneously rotating time crystals: A no-go theorem, *Physical Review Letters* **111**, 070402 (2013).
 - [2] H. Watanabe and M. Oshikawa, Absence of quantum time crystals, *Physical Review Letters* **114**, 251603 (2015).
 - [3] T. Hayata and Y. Hidaka, Diffusive nambu-goldstone modes in quantum time-crystals, [arXiv:1808.07636 \[hep-th\]](https://arxiv.org/abs/1808.07636) (2018).
 - [4] M. Hongo, S. Kim, T. Noumi, and A. Ota, Effective field theory of time-translational symmetry breaking in nonequilibrium open system, *Journal of High Energy Physics* **2019**, 1 (2019).
 - [5] C. P. Zelle, R. Daviet, A. Rosch, and S. Diehl, Universal phenomenology at critical exceptional points of nonequilibrium $O(N)$ models, *Physical Review X* **14**, 021052 (2024).
 - [6] R. Daviet, C. P. Zelle, A. Rosch, and S. Diehl, Nonequilibrium criticality at the onset of time-crystalline order, *Physical Review Letters* **132**, 167102 (2024).
 - [7] F. Piazza and H. Ritsch, Self-ordered limit cycles, chaos, and phase slippage with a superfluid inside an optical resonator, *Physical Review Letters* **115**, 163601 (2015).
 - [8] E. T. Owen, J. Jin, D. Rossini, R. Fazio, and M. J. Hartmann, Quantum correlations and limit cycles in the driven-dissipative heisenberg lattice, *New Journal of Physics* **20**, 045004 (2018).
 - [9] N. Dogra, M. Landini, K. Kroeger, L. Hruby, T. Donner, and T. Esslinger, Dissipation-induced structural instability and chiral dynamics in a quantum gas, *Science* **366**, 1496 (2019).
 - [10] B. Buča, C. Booker, and D. Jaksch, Algebraic theory of quantum synchronization and limit cycles under dissipation, *SciPost Physics* **12**, [10.21468/scipostphys.12.3.097](https://arxiv.org/abs/10.21468/scipostphys.12.3.097) (2022).
 - [11] B. Buča and D. Jaksch, Dissipation induced nonstationarity in a quantum gas, *Physical Review Letters* **123**, 260401 (2019).
 - [12] I. Carusotto and C. Ciuti, Quantum fluids of light, *Reviews of Modern Physics* **85**, 299 (2013).
 - [13] L. M. Sieberer, S. D. Huber, E. Altman, and S. Diehl, Dynamical critical phenomena in driven-dissipative systems, *Physical Review Letters* **110**, 195301 (2013).
 - [14] R. Hanai, A. Edelman, Y. Ohashi, and P. B. Littlewood, Non-hermitian phase transition from a polariton bose-einstein condensate to a photon laser, *Physical Review Letters* **122**, 185301 (2019).
 - [15] S. O. Demokritov, V. E. Demidov, O. Dzyapko, G. A. Melkov, A. A. Serga, B. Hillebrands, and A. N. Slavin, Bose-einstein condensation of quasi-equilibrium magnons at room temperature under pumping, *Nature* **443**, 430 (2006).
 - [16] P. Nowik-Boltyk, O. Dzyapko, V. E. Demidov, N. G. Berloff, and S. O. Demokritov, Spatially non-uniform ground state and quantized vortices in a two-component bose-einstein condensate of magnons, *Scientific Reports* **2**, [10.1038/srep00482](https://arxiv.org/abs/10.1038/srep00482) (2012).
 - [17] F. Iemini, A. Russomanno, J. Keeling, M. Schirò, M. Dalmonte, and R. Fazio, Boundary time crystals, *Physical Review Letters* **121**, 035301 (2018).
 - [18] O. Scarlatella, R. Fazio, and M. Schirò, Emergent finite frequency criticality of driven-dissipative correlated lattice bosons, *Physical Review B* **99**, 064511 (2019).
 - [19] H. Keßler, J. G. Cosme, M. Hemmerling, L. Mathey, and A. Hemmerich, Emergent limit cycles and time crystal dynamics in an atom-cavity system, *Physical Review A* **99**, 053605 (2019).
 - [20] P. Kongkhambut, J. Skulte, L. Mathey, J. G. Cosme, A. Hemmerich, and H. Keßler, Observation of a continuous time crystal, *Science* **377**, 670 (2022).
 - [21] M. C. Cross and P. C. Hohenberg, Pattern formation outside of equilibrium, *Reviews of Modern Physics* **65**, 851 (1993).
 - [22] T. Risler, J. Prost, and F. Jülicher, Universal critical behavior of noisy coupled oscillators, *Physical Review Letters* **93**, 175702 (2004).
 - [23] T. Risler, J. Prost, and F. Jülicher, Universal critical behavior of noisy coupled oscillators: A renormalization group study, *Physical Review E* **72**, 016130 (2005).
 - [24] L. Guislain and E. Bertin, Nonequilibrium phase transition to temporal oscillations in mean-field spin models, *Physical Review Letters* **130**, 207102 (2023).
 - [25] L. Guislain and E. Bertin, Discontinuous phase transition from ferromagnetic to oscillating states in a nonequilibrium mean-field spin model, *Physical Review E* **109**, 034131 (2024).
 - [26] J. Meibohm and M. Esposito, Small-amplitude synchronization in driven potts models, *Physical Review E* **110**, 044114 (2024).
 - [27] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence* (Springer Berlin Heidelberg, 1984).
 - [28] A. Pikovsky, M. Rosenblum, and J. Kurths, *Synchronization: A Universal Concept in Nonlinear Sciences* (Cambridge University Press, 2001).
 - [29] M. Fruchart, R. Hanai, P. B. Littlewood, and V. Vitelli, Non-reciprocal phase transitions, *Nature* **592**, 363 (2021).
 - [30] Y. Avni, M. Fruchart, D. Martin, D. Seara, and V. Vitelli, The non-reciprocal ising model, [arXiv:2311.05471 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2311.05471) (2023).
 - [31] S. Saha, J. Agudo-Canalejo, and R. Golestanian, Scalar active mixtures: The nonreciprocal cahn-hilliard model, *Physical Review X* **10**, 041009 (2020).
 - [32] Z. You, A. Baskaran, and M. C. Marchetti, Nonreciprocity as a generic route to traveling states, *Proceedings of*

- the National Academy of Sciences **117**, 19767 (2020).
- [33] T. Frohoff-Hülsmann, J. Wrembel, and U. Thiele, Suppression of coarsening and emergence of oscillatory behavior in a cahn-hilliard model with nonvariational coupling, *Physical Review E* **103**, 042602 (2021).
- [34] F. Brauns and M. C. Marchetti, Nonreciprocal pattern formation of conserved fields, *Physical Review X* **14**, 021014 (2024).
- [35] T. Suchanek, K. Kroy, and S. A. Loos, Irreversible mesoscale fluctuations herald the emergence of dynamical phases, *Physical Review Letters* **131**, 258302 (2023).
- [36] R. Hanai, Nonreciprocal frustration: Time crystalline order-by-disorder phenomenon and a spin-glass-like state, *Physical Review X* **14**, 011029 (2024).
- [37] L. Guislain and E. Bertin, Collective oscillations in a three-dimensional spin model with non-reciprocal interactions, [arXiv:2405.13925 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2405.13925) (2024).
- [38] N. del Ser, L. Heinen, and A. Rosch, Archimedean screw in driven chiral magnets, *SciPost Phys.* **11**, 009 (2021).
- [39] N. del Ser and V. Lohani, Skyrmion jellyfish in driven chiral magnets, *SciPost Phys.* **15**, 065 (2023).
- [40] R. Hanai, D. Ootsuki, and R. Tazai, Photoinduced non-reciprocal magnetism, [arXiv preprint arXiv:2406.05957](https://arxiv.org/abs/2406.05957) (2024).
- [41] D. Hardt, R. Doostani, S. Diehl, N. del Ser, and A. Rosch, Active magnetic matter: Propelling ferrimagnetic domain walls by dynamical frustration, [arXiv:2405.14320](https://arxiv.org/abs/2405.14320) (2024).
- [42] C.-K. Chan, T. E. Lee, and S. Gopalakrishnan, Limit-cycle phase in driven-dissipative spin systems, *Physical Review A* **91**, 051601 (2015).
- [43] M. Kardar, G. Parisi, and Y.-C. Zhang, Dynamic scaling of growing interfaces, *Physical Review Letters* **56**, 889 (1986).
- [44] J. Krug, Origins of scale invariance in growth processes, *Advances in Physics* **46**, 139 (1997).
- [45] K. A. Takeuchi, An appetizer to modern developments on the kardar-parisi-zhang universality class, *Physica A: Statistical Mechanics and its Applications* **504**, 77 (2018).
- [46] E. Altman, L. M. Sieberer, L. Chen, S. Diehl, and J. Toner, Two-Dimensional Superfluidity of Exciton Polaritons Requires Strong Anisotropy, *Phys. Rev. X* **5**, 011017 (2015).
- [47] L. He, L. M. Sieberer, E. Altman, and S. Diehl, Scaling properties of one-dimensional driven-dissipative condensates, *Physical Review B* **92**, 155307 (2015).
- [48] R. Lauter, A. Mitra, and F. Marquardt, From kardar-parisi-zhang scaling to explosive desynchronization in arrays of limit-cycle oscillators, *Physical Review E* **96**, 012220 (2017).
- [49] L. M. Sieberer, G. Wachtel, E. Altman, and S. Diehl, Lattice duality for the compact kardar-parisi-zhang equation, *Physical Review B* **94**, 104521 (2016).
- [50] G. Wachtel, L. M. Sieberer, S. Diehl, and E. Altman, Electrodynamical duality and vortex unbinding in driven-dissipative condensates, *Physical Review B* **94**, 104520 (2016).
- [51] F. Vercesi, Q. Fontaine, S. Ravets, J. Bloch, M. Richard, L. Canet, and A. Minguzzi, Phase diagram of one-dimensional driven-dissipative exciton-polariton condensates, *Physical Review Research* **5**, 043062 (2023).
- [52] K. Deligiannis, Q. Fontaine, D. Squizzato, M. Richard, S. Ravets, J. Bloch, A. Minguzzi, and L. Canet, Kardar-parisi-zhang universality in discrete two-dimensional driven-dissipative exciton polariton condensates, *Physical Review Research* **4**, 043207 (2022).
- [53] A. Altland and B. Simons, *Condensed Matter Field Theory* (Cambridge University Press, 2023).
- [54] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena (4th edition)*, International Series of Monographs on Physics, Vol. 113 (Clarendon Press, Oxford, 2002).
- [55] A. Beekman, L. Rademaker, and J. van Wezel, An introduction to spontaneous symmetry breaking, *SciPost Physics Lecture Notes* **10.21468/scipostphyslectnotes.11** (2019).
- [56] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, 2011).
- [57] L. M. Sieberer, M. Buchhold, and S. Diehl, Keldysh field theory for driven open quantum systems, *Reports on Progress in Physics* **79**, 096001 (2016).
- [58] See Supplemental Material at [URL], which includes Refs. [89–93], for additional details on the mapping to the KPZ equation.
- [59] In particular, for a complex order parameter, ϕ collects its real and imaginary parts.
- [60] L. M. Sieberer, M. Buchhold, J. Marino, and S. Diehl, Universality in driven open quantum matter, [arXiv:2312.03073 \[cond-mat.stat-mech\]](https://arxiv.org/abs/2312.03073) (2023).
- [61] P. C. Martin, E. D. Siggia, and H. A. Rose, Statistical dynamics of classical systems, *Physical Review A* **8**, 423 (1973).
- [62] H.-K. Janssen, On a lagrangean for classical field dynamics and renormalization group calculations of dynamical critical properties, *Zeitschrift für Physik B Condensed Matter and Quanta* **23**, 377 (1976).
- [63] C. D. Dominicis, Techniques de rénormalisation de la théorie des champs et dynamique des phénomènes critiques, *Le Journal de Physique Colloques* **37**, C1 (1976).
- [64] In the Keldysh framework, symmetries come with a fine structure often referred to as weak and strong or classical and quantum symmetries [57, 60]. In an open system, the strong time-translation symmetry is always broken (energy is not conserved), and the symmetry of interest is the weak version.
- [65] A. Kamenev, *Field theory of non-equilibrium systems* (Cambridge University Press, 2023).
- [66] U. C. Täuber, *Critical Dynamics* (Cambridge University Press, 2014).
- [67] M. Prähofer and H. Spohn, Universal distributions for growth processes in 1 + 1 dimensions and random matrices, *Physical Review Letters* **84**, 4882 (2000).
- [68] M. Prähofer and H. Spohn, Exact scaling functions for one-dimensional stationary KPZ growth, *Journal of Statistical Physics* **115**, 255 (2004).
- [69] T. Kloss, L. Canet, and N. Wschebor, Nonperturbative renormalization group for the stationary kardar-parisi-zhang equation: Scaling functions and amplitude ratios in 1+1, 2+1, and 3+1 dimensions, *Physical Review E* **86**, 051124 (2012).
- [70] A. Pagnani and G. Parisi, Numerical estimate of the kardar-parisi-zhang universality class in (2+1) dimensions, *Physical Review E* **92**, 010101 (2015).
- [71] L. He, L. M. Sieberer, and S. Diehl, Space-time vortex driven crossover and vortex turbulence phase transition in one-dimensional driven open condensates, *Physical Review Letters* **118**, 085301 (2017).
- [72] F. Helluin, D. Pinto-dias, Q. Fontaine, S. Ravets,

- J. Bloch, A. Minguzzi, and L. Canet, Phase diagram and universal scaling regimes of two-dimensional exciton-polariton bose-einstein condensates, [arXiv:2411.04311 \[cond-mat.quant-gas\]](#) (2024).
- [73] B. van der Pol, A theory of the amplitude of free and forced triode vibrations, *Radio Review* **1**, 701 (1920).
- [74] S. Walter, A. Nunnenkamp, and C. Bruder, Quantum synchronization of a driven self-sustained oscillator, *Physical Review Letters* **112**, 094102 (2014).
- [75] S. Walter, A. Nunnenkamp, and C. Bruder, Quantum synchronization of two van der pol oscillators, *Annalen der Physik* **527**, 131 (2014).
- [76] T. E. Lee and H. R. Sadeghpour, Quantum synchronization of quantum van der pol oscillators with trapped ions, *Physical Review Letters* **111**, 234101 (2013).
- [77] S. Dutta and N. R. Cooper, Critical response of a quantum van der pol oscillator, *Physical Review Letters* **123**, 250401 (2019).
- [78] L. B. Arosh, M. C. Cross, and R. Lifshitz, Quantum limit cycles and the rayleigh and van der pol oscillators, *Physical Review Research* **3**, 013130 (2021).
- [79] A. Cabot, G. L. Giorgi, and R. Zambrini, Nonequilibrium transition between dissipative time crystals, *PRX Quantum* **5**, 030325 (2024).
- [80] R. Gutiérrez and R. Cuerno, Kardar-Parisi-Zhang fluctuations in the synchronization dynamics of limit-cycle oscillators, *Phys. Rev. Res.* **6**, 033324 (2024).
- [81] I. S. Aranson and L. Kramer, The world of the complex ginzburg-landau equation, *Reviews of Modern Physics* **74**, 99 (2002).
- [82] Q. Fontaine, D. Squizzato, F. Baboux, I. Amelio, A. Lemaître, M. Morassi, I. Sagnes, L. Le Gratiet, A. Harouri, M. Wouters, I. Carusotto, A. Amo, M. Richard, A. Minguzzi, L. Canet, S. Ravets, and J. Bloch, Kardar-parisi-zhang universality in a one-dimensional polariton condensate, *Nature* **608**, 687 (2022).
- [83] O. K. Diessel, S. Diehl, and A. Chiocchetta, Emergent kardar-parisi-zhang phase in quadratically driven condensates, *Physical Review Letters* **128**, 070401 (2022).
- [84] K. Blom, U. Thiele, and A. Godec, *Interplay between local and global order controls bifurcations in nonreciprocal systems without a conservation law* (2024), [arXiv:2407.20312 \[cond-mat.stat-mech\]](#).
- [85] S. M. Rezende, Theory of coherence in bose-einstein condensation phenomena in a microwave-driven interacting magnon gas, *Physical Review B* **79**, 174411 (2009).
- [86] T. Suchanek, K. Kroy, and S. A. M. Loos, Entropy production in the nonreciprocal cahn-hilliard model, *Physical Review E* **108**, 064610 (2023).
- [87] G. Pisegna, S. Saha, and R. Golestanian, Emergent polar order in non-polar mixtures with non-reciprocal interactions, [arXiv:2404.05396 \[cond-mat.soft\]](#) (2024).
- [88] D. Nigro, D. Trypogeorgos, A. Gianfrate, D. Sanvitto, I. Carusotto, and D. Gerace, Supersolidity of polariton condensates in photonic crystal waveguides, [arXiv:2407.06671 \[cond-mat.quant-gas\]](#) (2024).
- [89] M. D. Schwartz, *Quantum Field Theory and the Standard Model* (Cambridge University Press, 2014).
- [90] N. Dupuis, *Field Theory of Condensed Matter and Ultracold Gases* (WORLD SCIENTIFIC (EUROPE), 2022).
- [91] F. Verhulst, *Nonlinear Differential Equations and Dynamical Systems* (Springer Berlin Heidelberg, 1996).
- [92] T. Sun, H. Guo, and M. Grant, Dynamics of driven interfaces with a conservation law, *Physical Review A* **40**, 6763 (1989).
- [93] D. E. Wolf and J. Villain, Growth with surface diffusion, *Europhysics Letters (EPL)* **13**, 389 (1990).

Supplemental Material for ‘Kardar-Parisi-Zhang scaling in time-crystalline matter’

BRIEF OVERVIEW: LINDBLAD-KELDYSH THEORY VS. CLASSICAL STOCHASTIC SYSTEMS

Here we give a brief contextualization of Eq. (2) in the main text, starting from driven open many-body quantum systems and then connecting to classical stochastic systems [S60]. The former are often well described by a Lindbladian,

$$\partial_t \hat{\rho} = \int_{\mathbf{x}} \left(-i[\hat{h}, \hat{\rho}] + \sum_{\alpha} (\hat{l}_{\alpha} \hat{\rho} \hat{l}_{\alpha}^{\dagger} - \frac{1}{2} \{ \hat{l}_{\alpha}^{\dagger} \hat{l}_{\alpha}, \hat{\rho} \}) \right), \quad (\text{S1})$$

with Hamiltonian density $\hat{h} = \hat{h}[\hat{\psi}^{\dagger}(\mathbf{x}), \hat{\psi}(\mathbf{x})]$ and Lindblad operators $\hat{l}_{\alpha} = \hat{l}_{\alpha}[\hat{\psi}^{\dagger}(\mathbf{x}), \hat{\psi}(\mathbf{x})]$ depending on the canonical field operators $\hat{\psi}^{\dagger}, \hat{\psi}$. For example, \hat{h} can describe kinetic energy and local interactions of bosons, and $\hat{l}_1 = \sqrt{\gamma_l} \hat{\psi}(\mathbf{x}), \hat{l}_2 = \sqrt{\gamma_p} \hat{\psi}^{\dagger}(\mathbf{x}), \hat{l}_3 = \sqrt{\kappa} \hat{\psi}^2(\mathbf{x})$ describe single-body loss, single-particle incoherent pump, and two-body loss, respectively, with the associated rates absorbed into the operators. Having order parameter theories in mind, we restrict to bosons here.

By the usual Trotterization procedure, this theory can be reformulated as an equivalent Lindblad-Keldysh functional integral,

$$\begin{aligned} Z &= \int \mathcal{D}(\psi_{\pm}^*, \psi_{\pm}) \exp(iS[\psi_{\pm}^*, \psi_{\pm}]), \\ S &= \int_{t, \mathbf{x}} \left(\sum_{\sigma} \sigma (\psi_{\sigma}^* i \partial_t \psi_{\sigma} - h_{\sigma}) + \sum_{\alpha} (l_{+, \alpha} l_{-, \alpha}^* - \frac{1}{2} (l_{+, \alpha}^* l_{+, \alpha} + l_{-, \alpha}^* l_{-, \alpha})) \right), \end{aligned} \quad (\text{S2})$$

where $h_{\pm} = h_{\pm}[\psi_{\pm}^*, \psi_{\pm}]$ etc., with the \pm index tracking the left (+) and right (-) action of operators in the Lindblad equation.

A change of basis $\psi = (\psi_+ + \psi_-)/\sqrt{2}, \tilde{\psi} = (\psi_+ - \psi_-)/\sqrt{2}$ introduces order parameter (or ‘classical’) and response (or ‘quantum’) fields. This transformation is particularly useful in generic driven open problems with (i) a finite Markovian noise level for all degrees of freedom, and (ii) gapless modes, as in the case of the Goldstone modes discussed in the main text: In this case, in the low frequency regime well below the noise level, a semiclassical limit can be taken, which amounts to a truncation up to second order in $\tilde{\psi}$, while the full dependence of ψ is kept. In this limit, the Lindblad-Keldysh theory reduces to the Martin-Siggia-Rose-Janssen-De Dominicis (MSRJD) theory, which in turn is equivalent to stochastic Langevin equations.

The above action is expressed in terms of complex field variables. While this is natural in the context of quantum systems, clearly the action can be re-expressed in real fields according to $\psi = (q + ip)\sqrt{2}, \tilde{\psi} = (\tilde{q} + i\tilde{p})\sqrt{2}$. Classical stochastic systems are often described in real coordinates from the start; for a unified approach, we work with general real vector fields in the main text, for which the Schwinger-Keldysh or MSRJD functional takes the form

$$Z[\tilde{\mathbf{J}}, \mathbf{J}] = \int \mathcal{D}\phi \mathcal{D}\tilde{\phi} \exp(iS[\phi, \tilde{\phi}] + i\tilde{\mathbf{J}}\phi + i\mathbf{J}\tilde{\phi}), \quad (\text{S3})$$

in presence of external sources.

NAMBU-GOLDSTONE MODE OF TIME TRANSLATIONS

Because of time-translation symmetry, the functional integral is invariant under infinitesimal time translations $t \rightarrow t + \epsilon, \phi' \rightarrow \phi + \epsilon \partial_t \phi$ whose only generator is ∂_t . From the Goldstone theorem [S89, S90], we therefore get a Goldstone mode from its breaking whenever the order parameter is time-dependent [S5, S6]

$$\int_{t'} (G^R)_{i,j}^{-1}(t, t', \mathbf{q} = 0) \partial_{t'} \varphi_j(t') = 0, \quad (\text{S4})$$

where G^R is the response to the external source J . As long as $\partial_t \varphi \neq 0$, this implies that the response function possesses a pole at vanishing momentum. If we can go into a frame that is comoving with the limit cycle where $\varphi(t) = v \hat{e}_{\parallel}$, we have

$$(G^R)_{\parallel}^{-1}(\omega = 0, \mathbf{q} = 0) = 0. \quad (\text{S5})$$

the Goldstone mode corresponds to a pole in the response function at vanishing frequency and momenta. At the linear level, the existence of such a mode is ensured by the Floquet theorem.

EFFECTIVE GOLDSTONE DYNAMICS FOR LIMIT CYCLES

Symmetry based construction

In the symmetry broken phase, time-translation symmetry is still realized, albeit in a nonlinear way. It acts in the following way on the fields:

$$\theta(t, \mathbf{x}) \rightarrow \theta(t + t_0, \mathbf{x}) + \frac{t_0}{\bar{T}}, \quad \tilde{\theta}(t, \mathbf{x}) \rightarrow \tilde{\theta}(t + t_0, \mathbf{x}). \quad (\text{S6})$$

Since a constant θ corresponds to a symmetry operation, the associated variation of the action has to vanish, $S[\tilde{\theta}(t + \theta, \mathbf{x}), \varphi(t + \bar{T}\theta)] - S[\tilde{\theta}(t, \mathbf{x}), \varphi(t)] = 0$. Therefore, all terms that appear in the effective action S_θ that are linear in $\tilde{\theta}(t, \mathbf{x})$ have to involve at least one derivative acting on $\theta(t, \mathbf{x})$. This is analogous to the usual shift symmetry of an internal $U(1)$ Goldstone mode. However, due to the external nature of time-translation symmetry, there is additional θ dependencies. These arise from acting with differential operators on the fluctuation around the limit-cycle stable state $\varphi(t + \theta(t, \mathbf{x}))\bar{T}$ and therefore always appear as $f(t + \theta\bar{T})$ for some function f . Clearly, this also leads to explicit time dependencies. From a symmetry perspective, this is a consequence of the fact that the combination $t + \theta(t, \mathbf{x})\bar{T}$ is also invariant under a time translation [S4]. Altogether, the action takes the form

$$S_\theta[\tilde{\theta}, \theta] = \int_{t, \mathbf{x}} \tilde{\theta} (\partial_t - Z[t + \theta\bar{T}]\nabla^2) + g[t + \theta\bar{T}]\tilde{\theta}(\nabla\theta)^2 - 2D[t + \theta\bar{T}]\tilde{\theta}^2, \quad (\text{S7})$$

But, the functions Z, D and g that carry these explicit time dependencies are periodic functions with the limit-cycle period \bar{T} . Considering timescales much larger $\tau \gg \bar{T}$, one can safely eliminate these time dependencies by averaging over one period \bar{T} . This makes the action only implicitly time-dependent, and the KPZ action of the main text is recovered.

Some constraints apply on the response field part of the action. First, the action must vanish for $\tilde{\theta} = 0$ to respect conservation of probability [S56]. In addition, the action (S7) describes the dynamics if θ is not a conserved field. Indeed, around a limit cycle, we do not expect the Goldstone fluctuations to be conserved in general, in particular in driven open systems. However, there may be special cases, where θ is subject to a conserved noise, which could lead to different variant of the KPZ universality class [S66, S92, S93]. Finally, we note that all the other operators that are allowed by symmetry are irrelevant in the low-energy regime near the Gaussian theory, $g = 0$, and they will not change the universal physics.

Explicit construction for the van der Pol oscillator

We now explicitly construct the effective dynamics in a limit-cycle phase for a two-dimensional order parameter $\phi = (\phi_1, \phi_2)$. The construction can equivalently be done at the level of the action, or on the equivalent stochastic equation of motion. For concreteness, we will use the van der Pol oscillator as an explicit example

$$(\partial_t^2 + (2\gamma + u\phi^2 - Z_1\nabla^2)\partial_t + \omega_0^2 + \lambda\phi^2 - Z_2\nabla^2)\phi + \xi = 0, \quad (\text{S8})$$

where $(\phi_1, \phi_2) = (\phi, \pi = \partial_t\phi)$, but the following formulas are easily leveraged to an arbitrary case. For negative damping $\gamma < 0$, its mean-field solution is a limit cycle, which we denote $f(t)$, and time-translation symmetry is spontaneously broken.

As shown in Eq. (3) of the main text, the idea is to work in the moving frame that locally has one of its basis vectors along the limit-cycle solution, the so-called Frenet frame. For the van der Pol case, the tangent vector $\hat{\mathbf{e}}_\theta(t)$ and normal vector $\hat{\mathbf{e}}_N(t)$ to the limit-cycle mean-field solution $\varphi(t) = (f(t), \dot{f}(t))$ are given by

$$\hat{\mathbf{e}}_\theta(t) = \frac{\dot{\varphi}(t)}{\|\dot{\varphi}(t)\|} = \frac{(\dot{f}(t), \ddot{f}(t))^T}{f(t)^2 + \dot{f}(t)^2}, \quad \hat{\mathbf{e}}_N(t) = \frac{(-\ddot{f}(t), \dot{f}(t))^T}{f(t)^2 + \dot{f}(t)^2}. \quad (\text{S9})$$

The fluctuations around the limit cycle can then be parameterized as

$$\begin{pmatrix} \phi(t, \mathbf{x}) \\ \pi(t, \mathbf{x}) \end{pmatrix} = \begin{pmatrix} f(t + \theta(t, \mathbf{x})\bar{T}) \\ \dot{f}(t + \theta(t, \mathbf{x})\bar{T}) \end{pmatrix} + N(t, \mathbf{x})\hat{\mathbf{e}}_N(t + \theta(t, \mathbf{x})\bar{T}). \quad (\text{S10})$$

This ensures the gaplessness of $\theta(t, \mathbf{x})$, because a constant θ simply transports along the limit cycle (along $\hat{\mathbf{e}}_\theta$ for an infinitesimal θ). For an order parameter that has more than two components, the construction can be generalized by identifying the $M - 1$ dimensional hyperspace, which is orthogonal to the limit cycle at every time t .

This parametrization can then be inserted into the action or Langevin dynamics. Once projected onto the two vectors $\hat{\mathbf{e}}_\theta$ and $\hat{\mathbf{e}}_N$, the latter takes the form

$$\partial_t N - Z_N(t + \theta\bar{T})\nabla^2 N + m_N(t + \theta\bar{T})N + Y_N(t + \theta\bar{T})\nabla^2 \theta + g_N(t + \theta\bar{T})(\nabla\theta)^2 + D_N(t + \theta\bar{T})\xi_N = 0, \quad (\text{S11})$$

$$\partial_t \theta - Z_\theta(t + \theta\bar{T})\nabla^2 \theta + Y_\theta(t + \theta\bar{T})\nabla^2 N + m_\theta(t + \theta\bar{T})N + g_\theta(t + \theta\bar{T})(\nabla\theta)^2 + D_\theta(t + \theta\bar{T})\xi_\theta = 0, \quad (\text{S12})$$

where we neglected all nonlinear terms in N , noticing that its fluctuations are gapped, while the gapless Goldstone nature of θ is evident. Again, all the explicit t dependencies are functions of $f(t)$ and its derivatives. In particular, we have

$$Z_\theta = \bar{T}\ddot{f}(t)\frac{(Z_2\dot{f}(t) + Z_1\ddot{f}(t))}{\dot{f}(t)^2 + \ddot{f}(t)^2}, \quad \text{and } g_\theta = -\bar{T}^2\frac{\ddot{f}(t)(Z_2\dot{f}(t) + Z_1\ddot{f}(t))}{\dot{f}(t)^2 + \ddot{f}(t)^2}. \quad (\text{S13})$$

Now, the Goldstone mode fluctuates on all timescales and we are in particular interested in long times $t \gg \bar{T}$, where $f(t)$ oscillates very quickly. It is therefore justified to average the right hand side over time following the average method [S91]. Such a procedure is described in the noiseless case in [S21], and leads to a so-called phase equation. Neglecting the gapped fluctuations of the phase, this procedure yields the KPZ equation (8) of the main text with coefficients $g = \bar{T}^{-1} \int_0^T dt g_\theta(t)$ and $Z = \bar{T}^{-1} \int_0^T dt Z_\theta(t)$. To compute the corrections coming from the gapped degree of freedom, we can use the first equation to adiabatically eliminate N , i.e., solving the first equation for $N(\theta)$. This is equivalent to performing the Gaussian integral over N and its response field in the path-integral formalism.

This derivation shows how KPZ dynamics for a gapless chronon emerges in a limit-cycle phase in any dimensions without broken internal continuous symmetries: it solely depends on the breaking of time-translation symmetry.