ON A BOUNDARY PAIR OF A DISSIPATIVE OPERATOR

RYTIS JURŠĖNAS

ABSTRACT. The aim of this brief note is to demonstrate that the boundary pair of a dissipative operator is determined by the unitary boundary pair of its symmetric part.

1. INTRODUCTION

Let T be a maximal dissipative operator in a Hilbert space $(\mathcal{H},\langle\,\cdot\,,\,\cdot\,\rangle)$. According to [\[8,](#page-7-0) Lemma 3.1] there is a Hilbert space $(\mathscr{E},\langle \cdot, \cdot \rangle_{\mathscr{E}})$ and an operator $\Gamma_{01} : \mathscr{H} \to \mathscr{E}, \mathscr{D}_{\Gamma_{01}} = \mathscr{D}_T$, which is bounded in graph norm of T , has dense range in \mathscr{E} , and such that

(1)
$$
\langle x, Ty \rangle - \langle Tx, y \rangle = i \langle \Gamma_{01} x, \Gamma_{01} y \rangle_{g} \quad (x, y \in \mathcal{D}_T).
$$

And similarly for the maximal dissipative operator $(-T^*)$.

Following [\[1,](#page-7-1) [2\]](#page-7-2) the pair $(\mathscr{E}, \Gamma_{01})$ is a boundary pair of T.

Here we show that:

- a) The maximality condition and the density of the domain of T are not needed for (1) to hold.
- b) $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ can be replaced by the Krein space $(\mathcal{H}, [\cdot, \cdot])$.
- c) Up to an isometric isomorphism $\mathscr E$ is determined by the closure of the sesquilinear form γ_N .

d) Given \mathcal{E} , the operator Γ_{01} is defined uniquely by the boundary value space of the symmetric part of T. With T in (1) one associates the symmetric positive and closable form

$$
\gamma_T[x, y] = -i(\langle x, Ty \rangle - \langle Tx, y \rangle)
$$
 on \mathcal{D}_T .

The quadratic form $\gamma_T [x] = \gamma_T [x, x]$. Its kernel, $\{x \in \mathcal{D}_T | \operatorname{Im}(x, Tx) = 0\}$, defines the symmetric part S of T, whose orthogonal complement in T is the dissipative restriction $N = T | (T + iI)^{-1}(\mathcal{N}_i); \mathcal{N}_i = \text{Ker}_i S^*$ is the deficiency subspace. Since the positive closable form $\gamma_N = \gamma_T$ on \mathcal{D}_N has trivial kernel, $(\mathcal{D}_N, (\cdot, \cdot)_N)$ with scalar product $(\cdot, \cdot)_N = \gamma_N[\cdot, \cdot]$ becomes an inner product space; its completion $(\mathcal{D}_{\bar{\gamma}_N}, (\cdot, \cdot)_{N}^{\times})$, where $\bar{\gamma}_N$ is the closure of γ_N , is isometrically isomorphic to \mathscr{E} , the latter being an intrinsic completion of the positive G-space $\Gamma_s(\mathcal{D}_T)$ in the Krein $\begin{pmatrix} 0 & -iI' \\ iI' & 0 \end{pmatrix}$ $\iota_{i,j}^{(0)}$ $-iI'$)-space $\mathcal{H}' \oplus \mathcal{H}'$, where Γ_s is the operator part of a boundary relation Γ in a unitary boundary pair (\mathcal{H}', Γ) for S^* , with the closed domain $\mathcal{D}_{\Gamma} = S^*$, and which in operator case is identified with \mathcal{D}_{S^*} ; such a boundary relation exists by [\[10,](#page-7-3) Proposition 3.7].

Given $\mathcal E$ in [\(1\)](#page-0-0) as described above, the operator $\Gamma_{01} = V\Gamma_s \mathcal{Q}_T$, where the operator V is closed continuous and semi-unitary from the pre-Hilbert space $\Gamma_s(\mathcal{D}_T)$ to \mathcal{E} , and which maps $\Gamma_s(\mathcal{D}_T)$ bijectively onto a dense lineal in \mathscr{E} ; the existence of V is due to the boundedness of the form γ_T in graph norm. Letting in particular $\mathcal{E} = \mathcal{D}_{\bar{\gamma}_N}$ gives

$$
\Gamma_{01} = U(T + iI)^{-1} P_i(T + iI)
$$

for some semi-unitary $U: \mathcal{D}_{\gamma_N} \to \mathcal{D}_{\bar{\gamma}_N}$, where P_i is an orthogonal projection in \mathcal{H} onto \mathcal{N}_i . (In this case the operator $\sqrt{F_b} = U(T + iI)^{-1}P_i$ in the proof of [\[8,](#page-7-0) Lemma 3.1].)

Formalized statements a)–d) are now given in the next section.

2. Main theorem

Let $(\mathcal{H}, [\cdot, \cdot])$ be a Krein space; an inner product is conjugate-linear in its first factor. Let the space of graphs $\mathcal{H}_{\Gamma} = \mathcal{H} \oplus \mathcal{H}$ be the Krein space $(\mathcal{H}_{\Gamma}, [\cdot, \cdot]_{\Gamma})$ with an indefinite metric (*cf.* [\[3,](#page-7-4) Section 2.1])

$$
[(x_1,y_1), (x_2,y_2)]_{\Gamma} = -i([x_1, y_2] - [y_1, x_2]).
$$

Date: December 16, 2024.

²⁰¹⁰ Mathematics Subject Classification. 47B50, 47B44, 46C20.

Key words and phrases. Dissipative operator, boundary pair, unitary boundary pair, Krein space.

Given a canonical symmetry J in $\mathcal{H}, (\mathcal{H}_{\Gamma}, [\cdot, \cdot]_{\Gamma})$ is a J_{Γ} -space with the canonical symmetry $J_{\Gamma}(x, y)$ = $(-iJy, iJx).$

Given a lineal (i.e. a linear subset) T in \mathcal{H}_{Γ} (i.e. a relation in \mathcal{H}), its J_Γ-orthogonal complement (i.e. the J-adjoint in H) is denoted by T^c ; if H is a Hilbert space, $T^c = T^*$.

Although here we mainly work with operators, it is convenient in many cases to identify an operator T with its graph, a relation with trivial indefiniteness (multivalued part) $\text{Ind } T$.

The domain, kernel, range of a relation (operator) T are denoted by \mathcal{D}_T , Ker T (also Ker_{λ} T = Ker(T − λI), $\lambda \in \mathbb{C}$, \mathcal{R}_T respectively. If necessary, the notions extend naturally to relations (operators) acting from one space to another space.

With an operator T in $\mathcal H$ one associates the symmetric sesquilinear form

$$
\gamma_T[x, y] = -i([x, Ty] - [Tx, y]) \text{ on } \mathcal{D}_{\gamma_T} = \mathcal{D}_T
$$

and the corresponding quadratic form $\gamma_T[x] = \gamma_T[x, x]$.

An operator T in $(\mathcal{H}, [\cdot, \cdot])$ is dissipative (symmetric) if Im[x, Tx] ≥ 0 (Im[x, Tx] = 0) for all $x \in \mathcal{D}_T$. An operator T is dissipative (symmetric) iff a lineal T in \mathcal{H}_{Γ} , *i.e.* the graph of T, is non-negative (neutral).

Any dissipative operator can be extended into a maximal one. A maximal dissipative operator in $\mathcal H$ (*i.e.* a maximal non-negative subspace in \mathcal{H}_{Γ}) is necessarily densely defined. A closed dissipative operator T is maximal iff such is the operator $(-T^c)$.

Let $(\mathcal{H}', \langle \cdot, \cdot \rangle')$ be a Hilbert space and let the space of graphs $\mathcal{H}'_{\Gamma} = \mathcal{H}' \oplus \mathcal{H}'$ be the Krein space $(\mathcal{H}_{\Gamma}^{\prime}, [\cdot, \cdot]_{\Gamma}^{\prime})$ with an indefinite metric

$$
[(u_1, v_1), (u_2, v_2)]'_{\Gamma} = -i(\langle u_1, v_2 \rangle' - \langle v_1, u_2 \rangle')
$$

and canonical symmetry $J'_{\Gamma}(u, v) = (-iv, iu)$.

The main result is stated as follows.

Theorem 1. A closed dissipative operator in a Krein space $(\mathcal{H}, [\cdot, \cdot])$ with canonical symmetry J has a boundary pair $(\mathscr{E}, \Gamma_{01})$; that is, there is a Hilbert space $(\mathscr{E}, \langle \cdot, \cdot \rangle_{\mathscr{E}})$ and an operator $\Gamma_{01} : \mathscr{H} \to \mathscr{E}, \mathscr{D}_{\Gamma_{01}} = \mathscr{D}_T$, which is bounded in graph norm of T , has dense range in \mathcal{E} , and such that

(2)
$$
[x, Ty] - [Tx, y] = i \langle \Gamma_{01} x, \Gamma_{01} y \rangle_{g} \quad (x, y \in \mathcal{D}_T).
$$

Specifically:

a) Γ_{01} is of the form

$$
\Gamma_{01} = V\Gamma_s E
$$
, $Ex = \begin{pmatrix} x \\ Tx \end{pmatrix}$, $\mathcal{D}_E = \mathcal{D}_T$

where Γ_s is the operator part of a boundary relation Γ in a ubp (\mathcal{H}',Γ) for $S^c = \mathfrak{D}_{\Gamma}$, the closed symmetric operator S in H is defined by

$$
S = T | \operatorname{Ker} \gamma_T \quad (\operatorname{Ker} \gamma_T = \{ x \in \mathcal{D}_T | \gamma_T [x] = 0 \})
$$

and V is a closed semi-unitary injective operator from the pre-Hilbert space $(\Gamma_s(T), [\cdot, \cdot]_{\Gamma})$ to $\&$.

b) Up to an isometric isomorphism ε is the completion $(\mathfrak{D}_{\bar{\gamma}_N}, (\cdot, \cdot)_{\bar{N}})$ of the pre-Hilbert space $(\mathcal{D}_N, (\cdot, \cdot)_N)$, where

$$
N = T | (JT + iI)^{-1} (\mathcal{N}_i \cap \mathcal{R}_{JT + iI}), \quad \mathcal{N}_i = \text{Ker}_i JS^c,
$$

 $(x, y)_N = \gamma_N [x, y]$ $(x, y \in \mathcal{D}_N)$; $\gamma_N = \gamma_T$ on \mathcal{D}_N

and $\bar{\gamma}_N$ is the closure of the positive sesquilinear form γ_N .

c) Particularly therefore one can always choose the boundary pair with $\mathcal{E} = \mathcal{D}_{\bar{\gamma}_N}$ and

$$
\Gamma_{01} = U(JT + iI)^{-1} P_{i}(JT + iI)
$$

where P_i is an orthogonal projection in $(\mathcal{H}, [\cdot, J \cdot])$ onto $\mathcal{N}_i \cap \mathcal{R}_{JT+iI}$, and U is a closed semi-unitary operator $(\mathcal{D}_N, (\cdot, \cdot)_N) \to (\mathcal{D}_{\bar{\gamma}_N}, (\cdot, \cdot)_{\bar{N}})$, which maps \mathcal{D}_N bijectively onto a dense lineal in $\mathcal{D}_{\bar{\gamma}_N}$.

Any closed symmetric operator S in $\mathcal H$ has a unitary boundary pair (ubp) $(\mathcal H', \Gamma)$, where Γ is a unitary relation from a J_{Γ} -space to a J'_{Γ} -space, such that $\bar{\mathcal{D}}_{\Gamma} = S^c$; an overbar refers to the closure. Without loss of generality one chooses a ubp with the closed domain, $\mathcal{D}_{\Gamma} = S^c$.

If S has equal defect numbers dim $\mathcal{N}_i = \dim \mathcal{N}_{-i}$ ($\mathcal{N}_{-i} = \text{Ker}_{-i} JS^c$), a ubp (\mathcal{H}', Γ) can be chosen as an ordinary boundary triple (obt), *i.e.* a ubp with Γ surjective.

The above classic facts on boundary value spaces of symmetric relations, and much more, can be found in [\[5,](#page-7-5) [6,](#page-7-6) [10,](#page-7-3) [11,](#page-7-7) [12,](#page-7-8) [13\]](#page-7-9), and we skip the details.

For the theory of operators in indefinite inner product spaces we refer to monographs [\[3,](#page-7-4) [7\]](#page-7-10). We only mention basic notions used in the theorem. An operator U from a W_1 -space $(\mathcal{H}_1, [\cdot, \cdot]_1)$ to a W_2 -space $(\mathcal{H}_2, [\cdot, \cdot]_2)$ is isometric if $[Ux_1, Ux_1]_2 = [x_1, x_1]_1$ $(x_1 \in \mathcal{D}_U); U$ is semi-unitary if it is isometric and $\mathcal{D}_U = \mathcal{H}_1$; U is unitary if it is semi-unitary and $\mathcal{R}_U = \mathcal{H}_2$. The spaces \mathcal{H}_1 and \mathcal{H}_2 are isometrically isomorphic if there is a unitary operator U which is injective (and hence bijective).

A W-space becomes a G-space if its Gram operator has trivial kernel. The Gram operator of the G-space $(\mathscr{L}_T, [\cdot, \cdot]_{\Gamma}')$ in a) reads

$$
G = P_{\mathcal{L}_T} J'_{\Gamma} | \mathcal{L}_T
$$

where $P_{\mathcal{L}_T}$ is an orthogonal projection in $[\cdot, J'_\Gamma \cdot]'_\Gamma$ onto \mathcal{L}_T .

The normed spaces $\mathcal{H}_1, \mathcal{H}_2$ are isomorphic if there is a closed continuous and continuously invertible bijection, *i.e.* a homeomorphism, $U: \mathcal{H}_1 \to \mathcal{H}_2$. Particularly isometrically isomorphic normed spaces are isomorphic. Throughout all normed spaces are (pre-)Hilbert spaces via polarization.

3. Proofs

Lemma 2. Given a subspace T in $(\mathcal{H}_{\Gamma}, [\cdot, \cdot]_{\Gamma})$, there is a Hilbert space $(\mathcal{H}', \langle \cdot, \cdot \rangle')$ and a bounded operator $\begin{pmatrix} \Gamma_0 \\ \Gamma_1 \end{pmatrix} \in \mathcal{B}(T, \mathcal{H}_{\Gamma}')$ such that

(3)
$$
[x, y'] - [x', y] = \langle \Gamma_0 \hat{x}, \Gamma_1 \hat{y} \rangle' - \langle \Gamma_1 \hat{x}, \Gamma_0 \hat{y} \rangle'
$$

for all $\hat{x} = \begin{pmatrix} x \\ x' \end{pmatrix} \in T$, $\hat{y} = \begin{pmatrix} y \\ y' \end{pmatrix} \in T$.

Specifically, if Γ_s is the operator part of a boundary relation Γ in a ubp (\mathcal{H}', Γ) for $S^c = \mathcal{D}_{\Gamma}$, where $S = T \cap T^c$ is the isotropic part of T, then $\binom{\Gamma_0}{\Gamma_1} = \Gamma_s | T$ and is a closed continuous and unitary operator from the indefinite inner product space $(T, [\cdot, \cdot]_{\Gamma})$ to the G-space $(\mathscr{L}_T, [\cdot, \cdot]_{\Gamma}')$ with

$$
\mathcal{L}_T = \Gamma_s(T) = \mathcal{R}_{\left(\begin{smallmatrix} \Gamma_0 \\ \Gamma_1 \end{smallmatrix}\right)}.
$$

A subspace is a closed lineal. Observe that in the lemma T is an arbitrary closed relation in H . For the column notation of operators we refer to [\[14\]](#page-7-11).

Proof. Let $S = T \cap T^c$, the isotropic part of T in \mathcal{H}_{Γ} ; hence there is a ubp (\mathcal{H}', Γ) for $S^c = \mathcal{D}_{\Gamma}(\supseteq T)$. That Γ is isometric explicitly reads

$$
[\hat{x}, \hat{y}]_{\Gamma} = [\hat{u}, \hat{v}]_{\Gamma}'
$$

for all $(\hat{x}, \hat{u}) \in \Gamma$, $(\hat{y}, \hat{v}) \in \Gamma$.

That Γ is unitary implies particularly it is a closed relation. Let Γ_s be the operator part of Γ; then Γ_s | T is the operator part of a closed relation $\Gamma | T$, and is therefore a bounded on $\mathcal{D}_{\Gamma | T} = T$ isometric operator $\mathscr{H}_{\Gamma} \to \mathscr{H}_{\Gamma}'$.

The range $\Gamma_s(T)$ is a subspace in \mathcal{H}'_{Γ} , since such is $\Gamma(T)$.

Since Ind Γ is an isotropic subspace in \mathcal{R}_{Γ} , the Green's identity implies

$$
[\hat{x}, \hat{y}]_{\Gamma} = [(\Gamma_s | T)\hat{x}, (\Gamma_s | T)\hat{y}]_{\Gamma}'
$$

for all $\hat{x} \in T$, $\hat{y} \in T$.

To accomplish the proof it remains to observe, first, that the operator Γ_s | T can be given the (graph) form

$$
\Gamma_s \,|\, T = \begin{pmatrix} \Gamma_0 \\ \Gamma_1 \end{pmatrix}
$$

where operators $\Gamma_0 = \pi_0 \Gamma_s | T$ and $\Gamma_1 = \pi_1 \Gamma_s | T$ and

$$
\pi_0: \mathcal{H}'_{\Gamma} \to \mathcal{H}', \quad \begin{pmatrix} u \\ u' \end{pmatrix} \mapsto u \, ;
$$

$$
\pi_1: \mathcal{H}'_{\Gamma} \to \mathcal{H}', \quad \begin{pmatrix} u \\ u' \end{pmatrix} \mapsto u'
$$

and, second, that $(\mathscr{L}_T, [\cdot, \cdot]'_T)$ is a non-degenerate subspace, because its isotropic part is $\Gamma_s(S)$, while $S = \text{Ker } \Gamma_s$.

Let T be a closed operator in a J-space \mathcal{H} , and consider the Hilbert space

$$
\mathcal{H}_T = (\mathcal{D}_T, \langle \cdot, \cdot \rangle_T)
$$

with scalar product

$$
\langle x, y \rangle_T = \langle x, y \rangle + \langle Tx, Ty \rangle, \quad \langle x, y \rangle = [x, Jy]
$$

and graph norm

$$
||x||_T = (||x||^2 + ||Tx||^2)^{1/2}, \quad ||x|| = \langle x, x \rangle^{1/2}.
$$

The form γ_T is bounded on \mathcal{H}_T , $|\gamma_T[x]| \leq 2||x||_T^2$, by Cauchy–Schwarz, so Reisz's representation theorem tells us that

$$
\gamma_T[x,y] = \langle x \, , \, Fy \rangle_T
$$

for some self-adjoint operator F in \mathcal{H}_T , which is bounded, $F \in \mathcal{B}(\mathcal{H}_T)$. We state this as a separate lemma.

Lemma 3. Let T be a closed operator in H . Then there is a self-adjoint operator $F \in \mathcal{B}(\mathcal{H}_T)$ such that $[x, Ty] - [Tx, y] = i\langle x, Fy\rangle_T \quad (x, y \in \mathcal{D}_T).$

Let T be a closed operator in $\mathcal{H}, S(= T | \text{Ker } \gamma_T)$ be its symmetric part, and define the closed operator restriction

$$
N=T\,|\,\mathcal{H}_T\ominus\mathcal{D}_S\,.
$$

Note: $T = S \oplus N$ (orthogonal componentwise sum); $\mathcal{D}_S \cap \mathcal{D}_N = \{0\}.$

Let Γ_s and $\mathcal{L}_T = \Gamma_s(T) = \Gamma_s(N)$ be as in Lemma [2.](#page-2-0) Then

$$
\varphi = E^{-1}(\Gamma_s \,|\, N)^{-1}
$$

is a bijection $\mathcal{L}_T \to \mathcal{D}_N$.

If T in Lemma [3](#page-3-0) is dissipative, then $F \geq 0$ and

$$
\gamma_T[x] = \|\sqrt{F}\,x\|_T^2 \quad (x \in \mathcal{D}_T)
$$

and a self-adjoint operator $\sqrt{F} \in \mathcal{B}(\mathcal{H}_T)$ is parametrized by

$$
\sqrt{F} = V_F \Gamma_s E \,, \quad V_F = \sqrt{F} \varphi \,.
$$

Because

$$
||V_F\hat{u}||_T^2 = [\hat{u}, \hat{u}]_{\Gamma}' \quad (\hat{u} \in \mathcal{L}_T)
$$

the operator V_F with

$$
\mathcal{D}_{V_F}=\mathcal{L}_T\,,\quad \mathcal{R}_{V_F}=\mathcal{R}_{\sqrt{F}}
$$

is closed continuous, injective, and semi-unitary from the positive G-space $(\mathscr{L}_T, [\cdot, \cdot]'_\Gamma)$ to the Hilbert subspace $(\bar{\mathcal{R}}_{\sqrt{F}}, \langle \cdot, \cdot \rangle_T)$ of \mathcal{H}_T ; \mathcal{L}_T is non-negative and then positive in \mathcal{H}'_Γ since T is nonnegative in \mathcal{H}_Γ .

With T as previously, let $(\mathscr{E}, \langle \cdot, \cdot \rangle_{\mathscr{E}})$ be a Hilbert space isometrically isomorphic to the Hilbert space $(\bar{\mathcal{R}}_{\sqrt{F}}, \langle \cdot, \cdot \rangle_T)$, and let $U_F : \bar{\mathcal{R}}_{\sqrt{F}} \to \mathscr{E}$ be the (standard) unitary operator. Put

$$
\Gamma_{01} = U_F \sqrt{F} = V \Gamma_s E \,, \quad V = U_F V_F
$$

and note that the lineal $\mathcal{R}_{\Gamma_{01}} = \mathcal{R}_V$ is dense in \mathcal{E} .

The latter combined with Lemmas [2,](#page-2-0) [3](#page-3-0) proves, first, the existence of a boundary pair $(\mathscr{E}, \Gamma_{01})$ in Theorem [1](#page-1-0) and, second, the form of Γ_{01} in a).

Remarks 4. a) As a simple corollary of [\(2\)](#page-1-1):

Real eigenvalues of a closed dissipative operator, T , are precisely those of its symmetric part, S .

To see this use $2(\text{Im }\lambda)[x, x] = ||\Gamma_{01}x||_{\mathcal{E}}^2$ $(x \in \text{Ker}_{\lambda} T, \lambda \in \sigma_p(T))$ and $\text{Ker } \Gamma_{01} = \mathcal{D}_S$.

b) As it appears from [\(2\)](#page-1-1), in applications it is enough to have established the domain restriction of Γ_s to T , which usually is done via (3) .

c) Consider a pre-Hilbert space $(\mathcal{R}_F, (\cdot, \cdot)_F)$ with scalar product

 $(u_x, u_y)_F = \langle x, F y \rangle_T$; $u_x = Fx$, $u_y = Fy$; $x, y \in \mathcal{H}_T$.

Its completion is by [\[9,](#page-7-12) Lemma 2.4] a Hilbert space $(\mathcal{R}_{\sqrt{F}}, (\cdot, \cdot)_{F}^{\sim})$ with scalar product

$$
(\sqrt{F}x \, , \, \sqrt{F}y)^\sim_F = \langle x \, , \, P_Fy\rangle_T \quad \, (x,y \in \mathcal{H}_T)
$$

where P_F is an orthogonal projection in \mathcal{H}_T onto $\bar{\mathcal{R}}_{\sqrt{F}}$, which is continuously embedded in the Hilbert space \mathcal{H}_T . Let ι be the corresponding embedding and let $\iota^* \colon \mathcal{H}_T \to \mathcal{R}_{\sqrt{F}}$ be its adjoint, *i.e.* $(u, \iota^*x)_{F}^{\sim} =$ $\langle \iota u, x \rangle_T (x \in \mathcal{H}_T, u \in \mathcal{R}_{\sqrt{F}}).$ Then by [\[9,](#page-7-12) Theorem 2.7] $F = u^*$.

Next we characterize $\mathcal E$ in Theorem [1](#page-1-0) b).

An intrinsic completion of the positive G-space \mathcal{L}_T in \mathcal{H}'_{Γ} is the completion with respect to the intrinsic norm

$$
|\hat{u}|_{\mathcal{L}_T} = ([\hat{u}, \hat{u}]_{\Gamma}')^{1/2} \quad (\hat{u} \in \mathcal{L}_T).
$$

Such a completion of \mathcal{L}_T is unique in the sense that any other intrinsic completion is isometrically isomorphic to the given one, with an isomorphism which acts as the identity on \mathcal{L}_T ; see [\[9\]](#page-7-12) for details.

Lemma 5. $(\mathscr{E}, \langle \cdot, \cdot \rangle_{\mathscr{E}})$ is an intrinsic completion of $(\mathscr{L}_T, [\cdot, \cdot]_{\Gamma}^{\prime}).$

Proof. Let $(\mathscr{E}, \langle \cdot, \cdot \rangle_{\mathscr{E}})$ be an intrinsic completion of $(\mathscr{L}_T, [\cdot, \cdot]_{\Gamma})$. Then by [\[7,](#page-7-10) Theorem V.2.1] $(\mathscr{E},\langle \cdot, \cdot \rangle_{\mathscr{E}})$ is a Hilbert space, and there is a (necessarily closed) semi-unitary operator from the G-space \mathcal{L}_T to the Hilbert space \mathcal{E} , which maps \mathcal{L}_T bijectively onto a dense lineal in \mathcal{E} ; V in Theorem [1](#page-1-0) a) has the required properties.

Any other operator Γ'_{01} : $\mathcal{H} \to \mathcal{E}'$, $\mathcal{D}_{\Gamma'_{01}} = \mathcal{D}_T$, which is bounded in graph norm of T, has dense range in a Hilbert space $(\mathscr{E}', \langle \cdot, \cdot \rangle_{\mathscr{E}'})$, and such that

$$
[x, Ty] - [Tx, y] = \mathbf{i} \langle \Gamma'_{01} x, \Gamma'_{01} y \rangle_{g}, \quad (x, y \in \mathcal{D}_T)
$$

is of the form $\Gamma'_{01} = V'\Gamma_{01}$ for some unitary operator $V': \mathscr{E} \to \mathscr{E}'$; the proof is analogous to [\[8,](#page-7-0) Lemma 3.3]. Therefore $\mathscr{E}' = V'(\mathscr{E})$ is just another intrinsic completion of the G-space, isometrically isomorphic to \mathscr{E} . \Box

An equivalent statement in Lemma [5](#page-4-0) is that a Banach space $(\mathscr{E}, \|\cdot\|_{\mathscr{E}})$ with norm $\|\cdot\|_{\mathscr{E}}$ induced by the scalar product $\langle \cdot, \cdot \rangle_{\mathcal{E}}$ is a completion of the normed space $(\mathcal{L}_T, |\cdot|_{\mathcal{L}_T})$.

The norm

$$
|\cdot|_{\mathcal{L}_T} = |\varphi \cdot|_N \quad \text{on} \quad \mathcal{L}_T
$$

or equivalently

$$
|\cdot|_N = |\varphi^{-1} \cdot |_{\mathcal{L}_T} \quad \text{on} \quad \mathcal{D}_N
$$

where the norm

$$
|\cdot|_N = \sqrt{\gamma_N[\cdot]}
$$
 (with $\gamma_N = \gamma_T$ on $\mathcal{D}_{\gamma_N} = \mathcal{D}_N$).

That is, φ is an isometric isomorphism from the normed space $(\mathscr{L}_T, |\cdot|_{\mathscr{L}_T})$ to the normed space $(\mathscr{D}_N, |\cdot|_N)$, which therefore extends to an isometric isomorphism from a completion $(\mathscr{E}, \|\cdot\|_{\mathscr{E}})$ of $(\mathscr{L}_T, |\cdot|_{\mathscr{L}_T})$ to a completion of $(\mathcal{D}_N, |\cdot|_N)$.

The next lemma accomplishes the proof of Theorem [1](#page-1-0) b).

Lemma 6. $(\mathscr{E}, \|\cdot\|_{\mathscr{E}})$ is isometrically isomorphic to a completion of the normed space $(\mathscr{D}_N, |\cdot|_N)$. Moreover, the space $(\mathfrak{D}_{\bar{\gamma}_N}, |\cdot|_{\widetilde{N}})$ with norm

$$
|x|_N^{\sim} = \sqrt{\bar{\gamma}_N[x]} \quad (x \in \mathcal{D}_{\bar{\gamma}_N})
$$

is the completion of $(\mathfrak{D}_N, |\cdot|_N)$.

Proof. The form γ_N is closable, since $\gamma_N[x] \leq 2||x|| ||Nx||$ $(x \in \mathcal{D}_{\gamma_N})$.

Since $(\mathcal{D}_{\bar{\gamma}_N}, \|\cdot\|_{\bar{\gamma}_N})$ is the Banach space with norm $||x||_{\bar{\gamma}_N} = (|x|_N^{\sim 2} + ||x||^2)^{1/2}$ $(x \in \mathcal{D}_{\bar{\gamma}_N})$, each sequence (x_n) in $\mathcal{D}_{\bar{\gamma}_N}$ converges to $x \in \mathcal{D}_{\bar{\gamma}_N}$ in norm $\|\cdot\|_{\bar{\gamma}_N}$, and then also in norm $|\cdot|_{N}^{\sim}$. In particular therefore a Cauchy sequence (x_n) in $(\mathcal{D}_{\bar{\gamma}_N}, |\cdot|_N^{\sim})$ is convergent.

That $(\mathcal{D}_{\bar{\gamma}_N}, |\cdot|_{\widetilde{N}})$ is the completion of $(\mathcal{D}_{\gamma_N}, |\cdot|_N)$ is seen from the definition: the lineal $\mathcal{D}_{\bar{\gamma}_N}$ is the set of limits in $(\mathcal{H}, \|\cdot\|)$ of Cauchy sequences (x_n) in $(\mathcal{D}_{\gamma_N}, |\cdot|_N)$; *i.e.* $(\mathcal{D}_{\gamma_N}, |\cdot|_N^{\sim}) \subseteq (\mathcal{D}_{\tilde{\gamma}_N}, |\cdot|_N^{\sim})$ densely. \Box

Remarks 7. As a simple corollary:

a) dim $\mathscr{E} = \dim \mathscr{D}_{\bar{\gamma}_N}$; hence $(\mathscr{L}_T, |\cdot|_{\mathscr{L}_T})$ is a Banach space iff $\gamma_N = \bar{\gamma}_N$.

b) The normed spaces $(\mathcal{R}_{\sqrt{F}}, \|\cdot\|_T)$ and $(\mathcal{D}_N, |\cdot|_N)$ and $(\mathcal{L}_T, |\cdot|_{\mathcal{L}_T})$ are pairwise isometrically isomor-

phic.
(Since Ker $\sqrt{F} = \mathcal{D}_S$ and since $\mathcal{D}_S \cap \mathcal{D}_N$ is trivial, $(\sqrt{F} | \mathcal{D}_N)^{-1}$ is such an isometric isomorphism $\mathcal{R}_{\sqrt{F}} \to \mathcal{D}_N$ by $\gamma_N = \gamma_T$ on \mathcal{D}_N .)

The above Banach spaces are Hilbert spaces via $\gamma_N[\cdot,\cdot]$. Particularly $(\mathcal{D}_{\bar{\gamma}_N},(\cdot,\cdot)_{\bar{N}})$ with scalar product $(\cdot, \cdot)_{N}^{\sim} = \bar{\gamma}_{N}[\cdot, \cdot]$ is the Hilbert space.

Lastly, in order to see Theorem [1](#page-1-0) c) use that $\varphi \Gamma_s E$ is 0 on \mathcal{D}_S and I on \mathcal{D}_N , and then that $\mathcal{D}_N =$ $(JT + iI)^{-1}(\mathcal{N}_i \cap \mathcal{R}_{JT+iI}).$

Remark 8. With the boundary pair in Theorem [1](#page-1-0) c) U extends by continuity to a unitary operator in $(\mathscr{D}_{\bar{\gamma}_N}, (\cdot, \cdot)_{\widetilde{N}})$, which can be set to *I*.

4. Criterion for completeness

Let T be a closed dissipative operator in a Krein space \mathcal{H} . When is the form γ_N closed? Some of the equivalent conditions are: $\mathcal{R}_G = \mathcal{L}_T$; \mathcal{L}_T is uniformly positive; $\mathcal{L}_T[\hat{+}]\mathcal{L}_T^{[\perp]} = \mathcal{H}'_T$. If at least one holds, then V in Theorem [1](#page-1-0) a) is a unitary operator from the Hilbert space $(\mathscr{L}_T, [\cdot, \cdot]'_\Gamma)$ to the Hilbert space $(\mathscr{E}, \langle\;\cdot\;,\;\cdot\;\rangle_{\mathscr{E}}).$

For example

$$
\mathcal{R}_{\Gamma_{01}} = \mathcal{R}_V = \mathcal{E} \quad \text{if} \quad \dim \mathcal{H}' < \infty
$$

since a finite-dimensional non-degenerate subspace is a Krein space (cf. [\[4,](#page-7-13) Lemma 2.4]); in our case \mathcal{L}_T is positive, hence a Hilbert space.

The Hilbert space \mathcal{H}' is finite-dimensional if e.g. S has equal finite defect numbers.

In practice one vaguely determines Γ_0 and Γ_1 , and then rigorously verifies [\(3\)](#page-2-1). Therefore suppose Γ_0 , $Γ_1$ are already known: We give a necessary and sufficient condition for $γ_N$ to be closed in terms of these operators.

Define operator restrictions S_1 , N_1 of T, which satisfy $T = S_1 \widehat{\oplus} N_1$, by

$$
S_1 = \text{Ker} \Gamma_1, \quad N_1 = T \, | \, \mathcal{H}_T \ominus \mathcal{D}_{S_1} \, .
$$

Define also the operator Θ_{01} in \mathcal{H}' by

$$
\Theta_{01} = (\Gamma_0 \,|\, N_1)(\Gamma_1 \,|\, N_1)^{-1} \quad (\mathfrak{D}_{\Theta_{01}} = \mathfrak{R}_{\Gamma_1}).
$$

It is a closed operator, because the inverse relation $\Theta_{01}^{-1} = \Gamma_s(N_1)$ in \mathcal{H}' is a subspace in \mathcal{H}'_{Γ} ; Θ_{01}^{-1} is an operator iff $S_0 \cap N_1 = \{0\}$, where $S_0 = \text{Ker } \Gamma_0$.

The adjoint Θ_{01}^* in \mathcal{H}' of Θ_{01} is a closed relation in \mathcal{H}' (Ind $\Theta_{01}^* = \mathcal{R}_{\Gamma_1}^{\perp}$) with dense domain.

Proposition 9. The following are equivalent:

a) $(\mathcal{R}_{\binom{\Gamma_0}{\Gamma_1}},[\cdot,\cdot]_{\Gamma})$ is complete, i.e. the form γ_N is closed;

- b) $\left\| \left[(\Gamma_1 i\Gamma_0) \right] N \right\| (\Gamma_1 + i\Gamma_0) \| N \|^{-1} \| < 1;$
- c) $\mathcal{R}_{\Gamma_1} + (\Gamma_0(S_1))^{\perp} = \mathcal{H}' = (\Theta_{01}^* \Theta_{01})((\Gamma_0(S_1))^{\perp}) + \Gamma_0(S_1).$

In b) $\|\cdot\|$ is the operator sup-norm, and in c) the action of the relation $\Theta_{01}^* - \Theta_{01}$ on the orthogonal complement $(\Gamma_0(S_1))^{\perp}$ means the action on the lineal $\mathfrak{D}_{\Theta_{01}} \cap \mathfrak{D}_{\Theta_{01}^*} \cap (\Gamma_0(S_1))^{\perp}$.

Proof. a) \Leftrightarrow b) This is the statement that a positive subspace \mathscr{L}_T in \mathscr{H}'_{Γ} with the angular operator K is uniformly positive iff $||K|| < 1$. It remains to remark, first, that K in \mathcal{H}_{Γ} with domain $iI | \mathcal{R}_{\Gamma_1+i\Gamma_0}$ maps $\begin{pmatrix} u \\ iu \end{pmatrix}$ to $\begin{pmatrix} 2iv-u \\ iu+2v \end{pmatrix}$, where $u = (\Gamma_1 + i\Gamma_0)\hat{x}$, $v = \Gamma_0\hat{x}$, $\hat{x} \in T$, and, second, that $\text{Ker}(\Gamma_1 + i\Gamma_0) = S$.

a) \Leftrightarrow c) Viewing \mathscr{L}_T as a relation in \mathscr{H}' , the equality in $\mathscr{L}_T[\hat{+}]\mathscr{L}_T^{[\hat{\perp}]} \subseteq \mathscr{H}'_{\Gamma}$ (where $\mathscr{L}_T^{[\hat{\perp}]}$ is the J'_{Γ} orthogonal complement of \mathcal{L}_T) holds iff

$$
\mathcal{R}_{\mathcal{L}_T} + \mathcal{R}_{\mathcal{L}_T^{[\perp]}} = \mathcal{H}' = \text{Ker}(\mathcal{L}_T[\hat{+}] \mathcal{L}_T^{[\perp]}).
$$

Because $\text{Ker}(\mathcal{L}_T[\hat{+}]\mathcal{L}_T^{[\perp]}) = \mathcal{R}_{(\mathcal{L}_T^{[\perp]})^{-1}\mathcal{L}_{T}-I}$, so it suffices to show that

$$
\mathcal{L}_T^{[\perp]} = [\Theta_{01}^* \, | \, \mathcal{D}_{\Theta_{01}^*} \cap (\Gamma_0(S_1))^{\perp}]^{-1}
$$

for the rest is a routine computation.

But the above $\mathscr{L}_T^{[\perp]}$ is easily seen from $(u, v) \in \mathscr{L}_T \Leftrightarrow u = \Theta_{01}v + \xi$, some $v \in \mathscr{R}_{\Gamma_1}$ and $\xi \in \Gamma_0(S_1)$. \square

Remark 10. If e.g. (\mathcal{H}', Γ) is an obt for S^c , this does not necessarily imply $\mathcal{R}_{\Gamma_1} = \mathcal{H}'$, since by definition Γ_1 is $\pi_1 \Gamma$ restricted to $T \subseteq S^c$.

5. Example of application

Let Ω be a measurable subset of $[0, \infty)$, which is not a null set, and let ι be the continuous inclusion of $L^2(\Omega)$ in $L^2(0,\infty)$: $\tilde{y} = iy = y$ a.e. on Ω and $\tilde{y} = 0$ a.e. on $\Omega^c = [0,\infty) \setminus \Omega$. Define the lineal $\mathscr{L}_{\Omega} =$ ${y \in L^2(\Omega) | \tilde{y} \in H^2(0, \infty)$; $\tilde{y}'(0) = h\tilde{y}(0)$, where h is a real number. Let q be a measurable bounded complex-valued function on Ω , with Im $q(t) > 0$ for a.e. $t \in \Omega$.

Proposition 11. Let $L = \left(-\frac{d^2}{dt^2} + q(t)\right) | \mathcal{L}_{\Omega}$, a closed densely defined dissipative operator in $L^2(\Omega)$. Then the operator norm of its Cayley transform $||(L - iI)(L + iI)^{-1}|| = 1$.

For a dissipative L, clearly $(L - iI)(L + iI)^{-1}$ is contractive, so the point here is that the sup-norm is precisely 1.

Proof. Denote by the same q a trivial extension of q to $[0, \infty)$; one can equally assume that q on $[0, \infty)$ is as in [\[8,](#page-7-0) Examples 3.4, 4.14]. In the Hilbert space $\mathcal{H} = L^2(0,\infty)$ the maximal dissipative operator $T = \tau = -\frac{d^2}{dt^2} + q(t)$ on $x \in H^2(0, \infty)$, $x'(0) = hx(0)$. The quadratic form $\gamma_T[x] = 2\|\sqrt{\text{Im }q} x\|_{L^2(\Omega)}^2$ $(x \in \mathcal{D}_T)$, $\text{so } \mathfrak{D}_S = \text{Ker} \gamma_T \text{ reads } \{x \in \mathfrak{D}_T | x = 0 \text{ a.e. on } \Omega\}, \text{ i.e. } \mathfrak{D}_S = \iota_c(\mathcal{L}'_{\Omega^c}), \mathcal{L}'_{\Omega^c} = \{y \in L^2(\Omega^c) | \iota_c y \in \mathfrak{D}_T\},$ where ι_c is the continuous inclusion of $L^2(\Omega^c)$ in $L^2(0,\infty)$: $\iota_c y = 0$ a.e. on Ω and $\iota_c y = y$ a.e. on Ω^c . Then $\mathcal{D}_N = \iota(\mathcal{L}_\Omega)$. In order to see the latter, let $\Delta_N = \{x \in \mathcal{D}_T | x = 0$ a.e. on $\Omega^c\}$. On the one hand $\mathcal{D}_S \cap \Delta_N = \{0\}, \mathcal{D}_S \dotplus \Delta_N = \mathcal{D}_T$; on the other hand $\mathcal{D}_S \dotplus \mathcal{D}_N = \mathcal{D}_T$, $\Delta_N \subseteq \mathcal{D}_N = \mathcal{H}_T \ominus \mathcal{D}_S$, so $\Delta_N = \mathcal{D}_N$ with

$$
(y, \, ty_1)_N = \gamma_L[y, y_1] \quad (y, y_1 \in \mathcal{L}_{\Omega}).
$$

Since $\mathcal{D}_T \subseteq \mathcal{H}$ densely, $\mathcal{L}_{\Omega} \subseteq L^2(\Omega)$ densely; hence $\mathcal{D}_{\bar{\gamma}_N} = \iota(L^2(\Omega))$ with

 $(uy, \iota y_1)_{N}^{\sim} = \langle y, y_1 \rangle_{L^2(\Omega, 2 \operatorname{Im} q(t)dt)} \quad (y, y_1 \in L^2(\Omega)).$

(Because the norm $\|\cdot\|_{L^2(\Omega,2 \text{Im } q(t)dt)} = \|\sqrt{2 \text{Im } q} \cdot \|_{L^2(\Omega)}$ is equivalent to $\|\cdot\|_{L^2(\Omega)}$, the Hilbert space $(\mathcal{D}_{\bar{\gamma}_N}, (\cdot, \cdot)_{\widetilde{N}})$ is isomorphic to $L^2(\Omega) = L^2(\Omega, dt)$.)

From the above, since $(\iota y, \iota y_1)_N = (\iota y, \iota y_1)_N^{\infty}$ for $y, y_1 \in \mathcal{L}_{\Omega}$, one puts in [\(3\)](#page-2-1) $\Gamma_0 \hat{x} = y$, $\Gamma_1 \hat{x} = Ly$, where $\hat{x} = Ex, x = x_S + iy, x_S \in \mathcal{D}_S, y \in \mathcal{L}_{\Omega}$, and then applies Proposition [9](#page-5-0) a), b), which accomplishes the proof. \Box

Remark 12. An alternative proof is given by computing a ubp (\mathcal{H}', Γ) for S^* , which we sketch here for illustrative purposes.

The operator $H: L^2(\Omega) \oplus L^2(\Omega^c) \to \mathcal{H}, (\frac{y}{y_c}) \mapsto \iota y + \iota_c y_c$ defines an isometric isomorphism, and $H^{-1}TH = (\frac{L}{\rho} \rho)$ in $L^2(\Omega) \oplus L^2(\Omega^c)$, where $L_c = \tau | \mathcal{L}'_{\infty}$ in $L^2(\Omega^c)$ is closed densely defined symmetri L_0^L $\stackrel{0}{\downarrow}$ in $L^2(\Omega) \oplus L^2(\Omega^c)$, where $L_c = \tau | \mathscr{L}_{\Omega^c}$ in $L^2(\Omega^c)$ is closed densely defined symmetric. Let L_c^* denote the adjoint operator and let (\mathcal{H}'', Γ') be a ubp for $L_c^* = \mathfrak{D}_{\Gamma'}$. Then the adjoint relation S^* in \mathcal{H} is given by

$$
H^{-1}S^*H = \begin{pmatrix} L^2(\Omega) \oplus L^2(\Omega) & 0 \\ 0 & L_c^* \end{pmatrix}
$$

and a ubp (\mathcal{H}', Γ) for $S^* = \mathcal{D}_{\Gamma}$ can be given the form: $\mathcal{H}' = \mathcal{H}'' \oplus L^2(\Omega)$ with

$$
\Gamma = \left\{ \left(\left(H \begin{pmatrix} y \\ y_c \end{pmatrix}, H \begin{pmatrix} y_1 \\ L_c^* y_c \end{pmatrix} \right), \left(\begin{pmatrix} u \\ y \end{pmatrix}, \begin{pmatrix} v \\ y_1 \end{pmatrix} \right) \right) \middle| y, y_1 \in L^2(\Omega) ;
$$

$$
\left(\begin{pmatrix} y_c \\ L_c^* y_c \end{pmatrix}, \begin{pmatrix} u \\ v \end{pmatrix} \right) \in \Gamma' \right\}.
$$

By replacing Γ' by its operator part one constructs the operator part Γ_s of Γ ; hence the operator $\Gamma_s | T = \binom{\Gamma_0}{\Gamma_1}$ maps $(H\left(\begin{smallmatrix} y\\ y_c \end{smallmatrix}\right), H\left(\begin{smallmatrix} Ly\\ L_c y_c \end{smallmatrix}\right)) \in T$ to $\left(\left(\begin{smallmatrix} 0\\ y \end{smallmatrix}\right), \left(\begin{smallmatrix} 0\\ Ly \end{smallmatrix}\right)\right) \in \mathscr{L}_T$.

Similar analysis can be done for complex h, Im $h > 0$, by replacing $L^2(\Omega, 2 \text{Im } q(t) dt)$ by $L^2(\Omega \cup$ ${0}$, $2 \operatorname{Im} h \, \delta + 2 \operatorname{Im} q(t) dt$, where δ is the Dirac measure supported at 0.

REFERENCES

- [1] Yu. Arlinskii, Families of symmetric operators with trivial domains of their squares, Compl. Anal. Oper. Theory 17 (2023), no. 120.
- [2] , Squares of symmetric operators, Compl. Anal. Oper. Theory 18 (2024), no. 161.
- [3] T. Azizov and I. Iokhvidov, Linear Operators in Spaces with an Indefinite Metric, John Wiley & Sons. Inc., 1989.
- [4] D. Baidiuk, V. Derkach, and S. Hassi, Unitary boundary pairs for isometric operators in Pontryagin spaces and generalized coresolvents, Compl. Anal. Oper. Theory 15 (2021), no. 32.
- [5] J. Behrndt, V. A. Derkach, S. Hassi, and H. de Snoo, A realization theorem for generalized Nevanlinna families, Operators and Matrices 5 (2011), no. 4, 679–706.
- [6] J. Behrndt, S. Hassi, and H. de Snoo, Boundary Value Problems, Weyl Functions, and Differential Operators, Monographs in Mathematics, Volume 108, Birkhauser, Switzerland AG, 2020.
- [7] J. Bognár, *Indefinite inner product spaces*, Springer-Verlag Berlin Heidelberg New York, 1974.
- [8] M. Brown, M. Marletta, S. Naboko, and I. Wood, The functional model for maximal dissipative operators (translation form): An approach in the spirit of operator knots, Trans. Amer. Math. Soc. 373 (2020), no. 6, 4145–4187.
- [9] B. Curgus and H. Langer, *Continuous embeddings, completions and complementation in Krein spaces*, Radovi Matematički 12 (2003), 37–79.
- [10] V. Derkach, S. Hassi, M. Malamud, and H. de Snoo, *Boundary relations and their Weyl families*, Trans. Amer. Math. Soc. 358 (2006), no. 12, 5351–5400.
- [11] V. A. Derkach and M. M. Malamud, Extension theory of symmetric operators and boundary value problems, Vol. 104, Institute of Mathematics of NAS of Ukraine, Kiev, 2017 (in Russian).
- [12] V. Derkach, S. Hassi, M. Malamud, and H. de Snoo, Boundary triplets and Weyl functions. Recent developments, Operator Methods for Boundary Value Problems, London Math. Soc. Lecture Note Series, 2012, pp. 161–220.
- [13] V. Derkach, S. Hassi, and M. M. Malamud, Generalized boundary triples, I. Some classes of isometric and unitary boundary pairs and realization problems for subclasses of Nevanlinna functions, Math. Nachr. 293 (2020), no. 7, 1278–1327.
- [14] S. Hassi, J. Labrousse, and H. de Snoo, Operational calculus for rows, columns, and blocks of linear relations, Advances in Operator Theory 5 (2020), no. 3, 1193–1228.

Vilnius University, Institute of Theoretical Physics and Astronomy, Sauletekio ave. 3, 10257 Vilnius, Lithuania ˙ Email address: rytis.jursenas@tfai.vu.lt