On signs of eigenvalues of modular forms satisfying Ramanujan conjecture

Nagarjuna Chary Addanki

Introduction

Siegel modular forms of genus n and weight k of level N are holomorphic functions on the Siegel upper half space \mathbb{H}_n that satisfy the modularity condition with respect to congruence subgroups of $\operatorname{Sp}_{2n}(\mathbb{Q})$. We denote a congruence subgroup of genus n and level N by $\Gamma^{(n)}(N)$. Let $M_k(\Gamma^{(n)}(N))$ denote the space of Siegel modular forms of weight k, genus n over $\Gamma^{(n)}(N)$ and $\operatorname{S}_k(\Gamma^{(n)}(N))$ denote the subspace of cuspidal forms. The space of cusp forms has a special basis called Hecke eigenforms. They arise as eigenvectors with respect to operators called the Hecke operators. For each positive integer m there is a Hecke operator associated to it, denoted by T(m). For a Hecke eigenform F, let $\lambda_F(m)$ denote the eigenvalue of T(m). For a normalised eigenform these eigenvalues are real. Hence the behavior of signs of the eigenvalues can be studied.

[8, Theorem 5] proved that for two normalized Hecke eigenforms $F \in S_{k_1}(\Gamma^{(1)}(N_1))$ and $G \in S_{k_2}(\Gamma^{(1)}(N_2))$, if $\operatorname{sign}(\lambda_F(p^r)) = \operatorname{sign}(\lambda_G(p^r))$ for almost all p and r then F = G. Thus two genus 1 modular forms can be compared by studying the signs of the eigenvalues. In case of genus 2, the space $S_k(\Gamma^{(2)}(1))$ decomposes into two subspaces, mutually orthogonal to each other. The first subspace is known as the Maass subspace and it is generated by Saito-Kurokawa lifts. Saito-Kurokawa lifts are modular forms of genus 2 constructed using a form of genus 1 as explained in [9]. Breulmann, in [5], showed that $F \in S_k(\Gamma^{(2)}(1))$ is a Saito-Kurokawa lift if and only if $\lambda_F(m) > 0$ for all $m \ge 1$. Kohnen, in [7], showed that a Hecke eigenform $F \in S_k(\Gamma^{(2)}(1))$ is in the orthogonal complement of the Maass space if and only if there are infinitely many sign changes in the sequence $\{\lambda_F(m)\}_{m\ge 1}$. These results underscore the significance of analyzing the signs of Hecke eigenvalues. In this article, we focus on the eigenvalues of the modular forms of genus 2 with level. Ikeda lifts, which are generalizations of the Saito-Kurokawa lifts to a higher genus, show a similar property. In [1] we proved that for a genus 4 Ikeda lift F, for a fixed r $\lambda_F(p^r) \ge 0$ for all sufficiently large p.

Pitale and Schmidt in [10] proved that, for a $F \in S_k(\Gamma_0^{(2)}(N))$ and in the orthogonal compliment of the Maass subspace, there are infinitely many prime numbers p such that the sequence of Hecke eigenvalues $\{\lambda_F(p^r)\}_{r\geq 1}$ has infinitely many sign changes. Theorem 4 of [6] proves that, under a specific condition, if $F \in S_{k_1}(\Gamma^{(2)}(1))$ and $G \in S_{k_2}(\Gamma^{(2)}(1))$ are in orthogonal complement of their respective Maass subspaces then for a set of primes of positive density, $\lambda_F(p)\lambda_G(p) < 0$. In this article, we use the techniques used in [6] to prove a similar result for Siegel modular forms with level that satisfy the Ramanujan conjecture. The main result is

Theorem 0.1. Let $F \in S_{k_1}(\Gamma^{(2)}(N_1))$ and $G \in S_{k_2}(\Gamma^{(2)}(N_2))$ be two Hecke eigenforms that satisfy the Ramanujan conjecture. Let π_F and π_G be cuspidal automorphic representations of $GSp_4(\mathbb{A}_Q)$ associated with F and G respectively. Assume that if

$$L(s, \pi_F, \text{spin}) = L(s, \pi_1)L(s, \pi_2) \text{ and } L(s, \pi_G, \text{spin}) = L(s, \tau_1)L(s, \tau_2)$$

for some cuspidal automorphic representations π_1, π_2, τ_1 and τ_2 over $\operatorname{GL}_2(\mathbb{A}_{\mathbb{Q}})$ then all representations are pairwise non isomorphic. Also, assume that for some $c \in (0, 4)$ and $\alpha > 15/16$,

$$\#\{p \le x : |\lambda_G(p)| > c\} \ge \alpha \frac{x}{\log x}$$

for sufficiently large x. Then the set of primes $\{p : \lambda_F(p)\lambda_G(p) < 0\}$ has a positive density.

The main result is on the signs of $\lambda_F(p)$. When the local factor of the spin L-function of a Hecke eigenform F is written as a Dirichlet series, the coefficient of p^{-s} is the eigenvalue $\lambda_F(p)$. Hence, to study the properties of $\lambda_F(p)$ it is sufficient to study the coefficient p^{-s} of the L-function. We extensively use the prime number theorem stated as Theorem 3 of [14] for asymptotic behavior of the coefficients.

Outline of the paper: In the first section of the article, we talk about basics of Siegel modular forms, automorphic representation associated with a modular form, and give a brief description of representations associated to the modular forms satisfying the Ramanujan conjecture. This section gives a description of the different types of L-functions to be expected for a eigenform satisfying the Ramanujan conjecture. In Section 2, we show the relation between the eigenvalue $\lambda_F(p)$ and the coefficient of p^{-s} of the L-function. Using Theorem 3 of [14], we prove few technical results that would be used for the main result. In the final section, we prove the main result and explain the assumptions made in the theorem.

1 Automorphic representations

For any ring R, let

$$\operatorname{GSp}_{2n}(R) = \left\{ g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \operatorname{GL}_{2n}(R) : {}^{t}gJg = \mu(g)J, \ J = \begin{pmatrix} 0 & 1_{n} \\ -1_{n} & 0 \end{pmatrix} \right\}$$

where μ is the similated homomorphism, 1_n is identity matrix of size n and $A, B, C, D \in M_n(R)$.

$$\operatorname{Sp}_{2n}(R) \coloneqq \{g \in \operatorname{GSp}_{2n}(R) : \mu(g) = 1\}.$$

Let N be a positive integer. Principal congruence subgroup of level N and genus n is defined to be the subgroup

$$\{g \in \operatorname{Sp}_{2n}(\mathbb{Z}) : g \equiv 1_{2n} (\text{mod } N)\}.$$

Congruence subgroup of level N and genus n is a finite indexed subgroup of $\text{Sp}_{2n}(\mathbb{Z})$ containing the principal congruence subgroup.

Let $\Gamma^{(n)}(N)$ denote a congruence subgroup of level N and genus n. A Siegel modular form F, of genus n, weight k with respect to $\Gamma^{(n)}(N)$, is a holomorphic function on the Siegel upper half space

$$\mathbb{H}_n \coloneqq \{ Z : Z \in M_n(\mathbb{C}), \ {}^t Z = Z \text{ and } \operatorname{Im}(Z) > 0 \}$$

satisfying the following two conditions.

1. Modularity condition

$$F((AZ+B)(CZ+D)^{-1}) = \det(CZ+D)^k F(Z) \quad \forall \ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma^{(n)}(N) \text{ and } Z \in \mathbb{H}_n$$

2. For n = 1, F(Z) is bounded on $\{Z = X + iY : Y \ge Y_0\} \forall Y_0 > 0$.

Holomorphy and modularity imply that a Siegel modular form has a Fourier expansion of the form

$$F(Z) = \sum_{\substack{T=T^t, \ T \ge 0\\T \text{ half integral}}} A(T) e^{2\pi i tr(TZ)}.$$

Siegel modular forms over $\Gamma^{(n)}(N)$ are generally called Siegel modular forms with level. Let $M_k(\Gamma^{(n)}(N))$ denote the space of Siegel modular forms of genus n and weight k over $\Gamma^{(n)}(N)$. F is called cuspidal if A(T) = 0 unless T > 0 and let $S_k(\Gamma^{(n)}(N))$ denote the subspace of cusp forms. This article focuses on the cusp forms of the genus 2 with level.

In case of genus 2 there are 4 congruence subgroups. They are

1. Borel congruence subgroup

$$B(N) = \operatorname{Sp}_4(\mathbb{Z}) \cap \begin{bmatrix} \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} \end{bmatrix}$$

2. Siegel congruence subgroups

$$\Gamma_0^2(N) = \operatorname{Sp}_4(\mathbb{Z}) \cap \begin{bmatrix} \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \end{bmatrix}$$

3. Klingen congruence subgroup

$$Q(N) = \operatorname{Sp}_4(\mathbb{Z}) \cap \begin{bmatrix} \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} \end{bmatrix}$$

4. Paramodular congruence subgroup

$$K(N) = \operatorname{Sp}_4(\mathbb{Q}) \cap \begin{bmatrix} \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & N^{-1}\mathbb{Z} \\ \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} \end{bmatrix}$$

Let $\Gamma^{(2)}(N)$ represent one of the four congruence subgroups above. For each $\Gamma^{(2)}(N)$ we can find an open compact subgroup $K_{\mathfrak{f}}$ of $\operatorname{GSp}_4(\mathbb{A}_{\mathbb{Q}})$ such that $\Gamma^{(2)}(N) = \operatorname{GSp}_4(\mathbb{Q}) \cap \operatorname{GSp}_4(\mathbb{R})^+ K_{\mathfrak{f}}$. Here $\operatorname{GSp}_4(\mathbb{R})^+$ is a subgroup of $\operatorname{GSp}_4(\mathbb{R})$ consisting of matrices with positive similitude. In the case of the congruence subgroups of genus 2, we describe the construction of $K_{\mathfrak{f}}$ below.

For a fixed N, let r_p denote a positive integer such that $p^{r_p}|N$ and $p^{r_p+1} \nmid N$.

1. If
$$\Gamma^{(2)}(N) = B(N)$$
 then $K_{\mathfrak{f}} = \prod_{p \mid N} B_{\mathfrak{p}}(p^{r_p}) \prod_{p \nmid N} \mathrm{GSp}_4(\mathbb{Z}_p)$ where

$$B_{\mathfrak{p}}(p^{r_p}) = \operatorname{Sp}_4(\mathbb{Z}_p) \cap \begin{bmatrix} \mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^{r_p}\mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^{r_p}\mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix}$$

2. If $\Gamma^{(2)}(N) = \Gamma^2_0(N)$ then $K_{\mathfrak{f}} = \prod_{p \mid N} \Gamma^2_{0,\mathfrak{p}}(p^{r_p}) \prod_{p \nmid N} \mathrm{GSp}_4(\mathbb{Z}_p)$ where

$$\Gamma_{0,\mathfrak{p}}^{2}(p^{r_{p}}) = \operatorname{Sp}_{4}(\mathbb{Z}_{p}) \cap \begin{bmatrix} \mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} \\ \mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} \\ p^{r_{p}}\mathbb{Z}_{p} & p^{r_{p}}\mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} \\ p^{r_{p}}\mathbb{Z}_{p} & p^{r_{p}}\mathbb{Z}_{p} & \mathbb{Z}_{p} & \mathbb{Z}_{p} \end{bmatrix}$$

3. If $\Gamma^{(2)}(N) = Q(N)$ then $K_{\mathfrak{f}} = \prod_{p \mid N} Q_{\mathfrak{p}}(p^{r_p}) \prod_{p \nmid N} \mathrm{GSp}_4(\mathbb{Z}_p)$ where

$$Q_{\mathfrak{p}}(p^{r_p}) = \operatorname{Sp}_4(\mathbb{Z}_p) \cap \begin{bmatrix} \mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ \mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^{r_p}\mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & p^{r_p}\mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix}$$

4. If $\Gamma^{(2)}(N) = K(N)$ then $K_{\mathfrak{f}} = \prod_{p \mid N} K_{\mathfrak{p}}(p^{r_p}) \prod_{p \nmid N} \mathrm{GSp}_4(\mathbb{Z}_p)$ where

$$K_{\mathfrak{p}}(p^{r_p}) = \operatorname{Sp}_4(\mathbb{Q}) \cap \begin{bmatrix} \mathbb{Z}_p & p^{r_p} \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p & p^{-r} \mathbb{Z}_p \\ \mathbb{Z}_p & p^{r_p} \mathbb{Z}_p & \mathbb{Z}_p & \mathbb{Z}_p \\ p^{r_p} \mathbb{Z}_p & p^{r_p} \mathbb{Z}_p & p^{r_p} \mathbb{Z}_p & \mathbb{Z}_p \end{bmatrix}$$

For each cusp form, there is an associated automorphic representation over $\operatorname{GSp}_4(\mathbb{A}_{\mathbb{Q}})$. For a fixed positive integer N, let $F \in S_k(\Gamma^{(2)}(N))$ be a cusp form, $\Gamma^{(2)}(N)$ be any one of the four congruence subgroups defined above and K_f be the open compact group such that $\Gamma^{(2)}(N) = \operatorname{GSp}_4(\mathbb{Q}) \cap \operatorname{GSp}_4(\mathbb{R})^+ K_f$. The Strong Approximation Theorem for $\operatorname{GSp}(\mathbb{A}_{\mathbb{Q}})$ states that

$$\mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}}) \cong \mathrm{GSp}_4(\mathbb{Q})(\mathrm{GSp}_4(\mathbb{R})^+ K_{\mathfrak{f}}).$$

It implies that, given $g \in \mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$ there exists $g_q \in \mathrm{GSp}_4(\mathbb{Q})$, $g_\infty \in \mathrm{GSp}_4(\mathbb{R})^+$, $k \in K_{\mathfrak{f}}$ such that $g = g_q(g_\infty k)$. An automorphic form associated with F is a function on $\mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$ denoted by ϕ_F and defined as follows: For $g \in \mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$,

$$\phi_F(g) \coloneqq \mu(g_\infty)^k \det(C_\infty I_2 + D_\infty)^{-k} F\left((A_\infty I_2 + B_\infty)(C_\infty I_2 + D_\infty)^{-1}\right)$$

where

$$I_2 = \begin{bmatrix} i & 0\\ 0 & i \end{bmatrix} \quad \text{and} \quad g_{\infty} = \begin{bmatrix} A_{\infty} & B_{\infty}\\ C_{\infty} & D_{\infty} \end{bmatrix}$$

From the fact that $F \in S_k(\Gamma^{(2)}(N))$ and the strong approximation theorem, it follows that ϕ_F is well defined. It can also be shown that $\phi_F \in L^2(Z(\mathbb{A}_{\mathbb{Q}})\mathrm{GSp}_4(\mathbb{Q})\backslash\mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}}))$. Given $h \in \mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$, we define right translation of ϕ_F by

$$h.\phi_F(g) \coloneqq \phi_F(gh).$$

Let V_F denote the subspace of $L^2(Z(\mathbb{A}_{\mathbb{Q}})\mathrm{GSp}_4(\mathbb{Q}))$ generated by $h.\phi_F$ for $h \in \mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$. The group $\mathrm{GSp}_4(\mathbb{A}_{\mathbb{Q}})$ acts on V_F by right translation. This action is defined as the representation associated with F and is denoted by π_F . More details on the construction of π_F can be found in Section 4 of [4] and Section 3.2 of [11].

Since the representation is trivial on the center of $\text{GSp}_4(\mathbb{A}_{\mathbb{Q}})$, it can be seen as a representation of $\text{PGSp}_4(\mathbb{A}_{\mathbb{Q}})$. Using the exceptional isomorphism, $\text{PGSp}_4(\mathbb{A}_{\mathbb{Q}}) \cong \text{SO}_5(\mathbb{A}_{\mathbb{Q}})$, π_F can be extended to a representation of $\text{SO}_5(\mathbb{A}_{\mathbb{Q}})$. Hence, given F, we can attach a representation of $\text{SO}_5(\mathbb{A}_{\mathbb{Q}})$. Theorem 1.3.2 of [3] gives a classification of all such representations. In Section 2.2 of [12], Schmidt explains the classification specific to the case of modular forms of the genus 2. In this case, there are 6 distinct classes. In this article, we focus on modular forms that satisfy the generalized Ramanujan conjecture.

Generalized Ramanujan Conjecture: Let $F \in S_k(\Gamma^{(n)}(N))$ be a Hecke eigenform with Satake-*p*-parameters $\alpha_{0,p}^{(F)}, \alpha_{1,p}^{(F)}, ..., \alpha_{n,p}^{(F)}$. A prime *p* is called unramified if $p \nmid N$. GRC states that for all the unramified primes *p*, the Satake-*p*-parameters satisfy

$$|\alpha_{i,p}| = 1$$
 for $i = 1, 2, ..., n$.

[12, Prop 2.1] proves that G and Y are the only classes that satisfy the Ramanujan conjecture.

• General type, $(\mathbf{G}) : F \in S_k(\Gamma^{(2)}(N))$ is said to be of type \mathbf{G} , if there exists a cuspidal automorphic representation π of $\operatorname{GL}_4(\mathbb{A}_{\mathbb{Q}})$ such that

$$L(s, \pi_F, spin) = L(s, \pi)$$

• Yoshida type, $(\mathbf{Y}) : F \in S_k(\Gamma^{(2)}(N))$ is said to be of type \mathbf{Y} , if there exists two cuspidal automorphic representations π_1 , π_2 of $\operatorname{GL}_2(\mathbb{A}_{\mathbb{Q}})$ such that

$$L(s, \pi_F, spin) = L(s, \pi_1)L(s, \pi_2).$$

An example of such modular forms are the Yoshida lifts.

Definition 1 (Yoshida lifts). Let $f \in S_{k_1}(\Gamma_0(N_1))$ and $g \in S_{k_2}(\Gamma_0(N_2))$ be two Hecke eigen newforms where

$$\Gamma_0(N) = \{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z}) : \ c \equiv 0 (\operatorname{mod} N) \}.$$

 $F \in S_k(\Gamma^{(2)}(N))$ is said to be a Yoshida lift of f and g, if π_F is irreducible and

$$L(s, \pi_F, \operatorname{spin}) = L(s, \pi_f)L(s, \pi_g)$$

2 First eigenvalue

For each $g \in \mathrm{GSp}_{2n}(\mathbb{Q})^+ \cap M_{2n}(\mathbb{Z})$ such that $\mathrm{gcd}(\mu(g), N) = 1$ we can associate a Hecke operator T(g) on $M_k(\Gamma^{(n)}(N))$. Let $\Gamma = \mathrm{Sp}_{2n}(\mathbb{Z})$, for $F \in M_k(\Gamma^{(n)}(N))$,

$$T(g)F := \sum_{i} F|_{k}g_{i} \quad \text{where } \Gamma^{(n)}(N)g\Gamma^{(n)}(N) = \sqcup_{i}\Gamma^{(n)}(N)g_{i}, \quad g_{i} = \begin{pmatrix} A_{i} & B_{i} \\ C_{i} & D_{i} \end{pmatrix}$$

and $F|_{k}g_{i}(Z) = \mu(g)^{nk - \frac{n(n+1)}{2}} \det(C_{i}Z + D_{i})^{-k}F((A_{i}Z + B_{i})(C_{i}Z + D_{i})^{-1}).$

For a positive integer m such that gcd(m, N) = 1,

$$T(m) \coloneqq \sum_{g: \mu(g) = m} T(g).$$

In Theorem 4.7 of [2], it is proved that there exists a basis for $M_k(\Gamma^{(n)}(N))$ which are eigenforms with respect to all Hecke operators T(p) such that $p \nmid N$. For a Hecke eigenform $F \in S_k(\Gamma^{(n)}(N))$, denote $\mu_F(g)$ as the eigenvalue of the operator T(g). Classically $\mu_F(g)$ can be expressed in terms of Satake p – parameters. For any g with $\mu(g) = p^r$, depending on F there are n + 1 complex numbers $(a_{0,p}^{(F)}, a_{1,p}^{(F)}, \ldots, a_{n,p}^{(F)})$ satisfying

$$\mu_F(g) = \left(p^{nk - \frac{n(n+1)}{4}} a_{0,p}^{(F)}\right)^r \sum_i \prod_{j=1}^n (a_{j,p}^{(F)} p^{-j})^{d_{ij}} \text{ where } \Gamma^{(n)}(N) g \Gamma^{(n)}(N) = \bigsqcup_i \Gamma^{(n)}(N) g_i, \quad (1)$$
$$g_i = \begin{pmatrix} A_i & B_i \\ 0 & D_i \end{pmatrix} \text{ and } D_i = \begin{pmatrix} p^{d_{i1}} & * \\ & \ddots & \vdots \\ 0 & \dots & p^{d_{in}} \end{pmatrix}.$$

The complex numbers $a_{j,p}^{(F)}$ for $0 \le j \le n$ are called the Satake p parameters of F.

Lemma 2.1. If $F \in S_k(\Gamma^{(2)}(N))$ is a Hecke eigenform and $a_{0,p}^{(F)}$, $a_{1,p}^{(F)}$, $a_{2,p}^{(F)}$ are the Satake-pparameters then

$$\mu_F(p) = p^{2k - \frac{3}{2}} (a_{0,p}^{(F)} + a_{0,p}^{(F)} a_{1,p}^{(F)} + a_{0,p}^{(F)} a_{2,p}^{(F)} + a_{0,p}^{(F)} a_{1,p}^{(F)} a_{2,p}^{(F)}).$$

Г₁

Proof. For the Hecke operator T(p) we have the following decomposition

$$T(p) = \Gamma_2 \begin{bmatrix} 1_2 & \\ & p1_2 \end{bmatrix} \Gamma_2 = \Gamma_2 \begin{bmatrix} p1_2 & \\ & 1_2 \end{bmatrix} \sqcup \bigsqcup_{a \in \mathbb{Z}/p\mathbb{Z}} \Gamma_2 \begin{bmatrix} 1 & a & \\ & p & \\ & & 1 \end{bmatrix} \sqcup \bigsqcup_{\alpha, d \in \mathbb{Z}/p\mathbb{Z}} \Gamma_2 \begin{bmatrix} 1 & a & b \\ & 1 & b & d \\ & & p \end{bmatrix} \cdot \begin{bmatrix} p & & \\ & 1 & a & \\ & & p & \\ & & p & \end{bmatrix} \sqcup \bigsqcup_{a, b, d \in \mathbb{Z}/p\mathbb{Z}} \Gamma_2 \begin{bmatrix} 1 & a & b & \\ & 1 & b & d & \\ & & p & \\ & & p & \end{bmatrix} \cdot$$

There are four kinds of right cosets in the above decomposition,

$$g_{1} = \begin{bmatrix} p_{1_{2}} \\ & 1_{2} \end{bmatrix}, \ g_{2} = \begin{bmatrix} 1 & * & * \\ & p & \\ & & & 1 \end{bmatrix}, \ g_{3} = \begin{bmatrix} p & & & \\ * & 1 & * \\ & & & p \end{bmatrix}, \ g_{4} = \begin{bmatrix} 1 & * & * \\ & 1 & * & * \\ & & p & \\ & & & p \end{bmatrix}.$$

With D'_i s,

$$D_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, D_2 = \begin{bmatrix} p & 0 \\ 0 & 1 \end{bmatrix}, D_3 = \begin{bmatrix} 1 & * \\ 0 & p \end{bmatrix}, D_4 = \begin{bmatrix} p & 0 \\ 0 & p \end{bmatrix}.$$

 $\mu_F(p)$ can be calculated using the formula in (1) and evaluating the contribution of each D_i . For D_1 , $d_{1,1} = d_{1,2} = 0$. Hence the contribution to $\mu_F(p)$ is $p^{2k-\frac{3}{2}}a_{0,p}^{(F)}$. For D_2 , $d_{2,1} = 1$ and $d_{2,2} = 0$, and there are p number of cosets of these kind. Adding each cosets contribution to the eigenvalue we get

$$p^{2k-\frac{3}{2}}a_{0,p}^{(F)}pa_{1,p}^{(F)}p^{-1} = p^{2k-\frac{3}{2}}a_{0,p}^{(F)}a_{1,p}^{(F)}.$$

Similarly the contribution of D'_{3} s and D'_{4} s comes out to be $p^{2k-\frac{3}{2}}a^{(F)}_{0,p}a^{(F)}_{2,p}$ and $p^{2k-\frac{3}{2}}a^{(F)}_{0,p}a^{(F)}_{2,p}a^{(F)}_{2,p}a^{(F)}_{2,p}$ respectively. Adding everything, it follows that

$$\mu_F(p) = p^{2k - \frac{3}{2}} (a_{0,p}^{(F)} + a_{0,p}^{(F)} a_{1,p}^{(F)} + a_{0,p}^{(F)} a_{2,p}^{(F)} + a_{0,p}^{(F)} a_{1,p}^{(F)} a_{2,p}^{(F)}).$$

Let π_F be the automorphic representation associated with F and $L(s, \pi_F)$ be the corresponding spin L-function. $\pi_F = \otimes_p' \pi_p$ where π_p is an unramified representation of $\mathrm{GSp}_4(\mathbb{Q}_p)$ for all primes $p \nmid N$. And $L(s, \pi_F) = \prod L_p(s, \pi_{F,p})$, where $L_p(s, \pi_{F,p})$ are called the local L-factors. There exists complex numbers $a_{1,p}^p$, $a_{2,p}$, $a_{3,p}$ and $a_{4,p}$ such that

$$L_p(s, \pi_{F,p}) = \left((1 - a_{1,p}p^{-s})(1 - a_{2,p}p^{-s})(1 - a_{3,p}p^{-s})(1 - a_{4,p}p^{-s}) \right)^{-1}$$

For unramified primes, i.e for $p \nmid N$, these are the Satake-*p*-parameters associated to F. Hence

$$L_p(s,\pi_p) = \left((1 - a_{0,p}^{(F)}p^{-s})(1 - a_{0,p}^{(F)}a_{1,p}^{(F)}p^{-s})(1 - a_{0,p}^{(F)}a_{2,p}^{(F)}p^{-s})(1 - a_{0,p}^{(F)}a_{1,p}^{(F)}a_{2,p}^{(F)}p^{-s}) \right)^{-1}$$

for all $p \nmid N$. At ramified primes, the local L-factor is still an inverse of a polynomial in p^{-s} but the degree can be less than 4. Hence, we can write the local factor with 4 constants $a_{i,p}$ but these can be zero as well. Since we are interested in signs of $\mu_F(p)$, it is enough to study normalized eigenvalues $\lambda_F(p) = \frac{\mu_F(p)}{p^{2k-\frac{3}{2}}}$. From Lemma 2.1, we conclude that $\lambda_F(p) = a_{1,p} + a_{2,p} + a_{3,p} + a_{4,p}$. If the local factors are written as Dirichlet series, say $L_p(s, \pi_F) = \sum_{r=1}^{\infty} a_{\pi_F}(p^r) p^{-rs}$, then at unramified primes $\lambda_F(p) = a_{\pi_F}(p)$.

For a Hecke eigenform in class \mathbf{G} , there exists an irreducible cuspidal automorphic representation π of $\operatorname{GL}_4(\mathbb{A}_{\mathbb{O}})$ such that $\lambda_F(p) = a_{\pi}(p)$ for all unramified primes. Similarly, for a cusp form in class Y, there exists two irreducible cuspidal automorphic representations π and τ of $\operatorname{GL}_2(\mathbb{A}_{\mathbb{O}})$ such that for all unramified primes $p, \lambda_F(p) = a_{\pi}(p) + a_{\tau}(p)$.

In the remainder of the section we prove a few technical results that will be used for the main theorem. For any two real valued functions f(x) and g(x) we use the following notations.

1. f(x) = O(g(x)) if there exists a constant c such that $|f(x)| \le c|g(x)|$ for sufficiently large x.

2.
$$f(x) = o(g(x) \text{ if } \lim_{x \to \infty} \frac{f(x)}{g(x)} = 0.$$

Lemma 2.2. Let π be a self dual, unitary, cuspidal automorphic representation of $GL_m(\mathbb{A}_{\mathbb{Q}})$ for $m \leq 4$. If $L_p(s,\pi) = \sum_{r=1}^{\infty} a_{\pi}(p^r) p^{-rs}$ and $a_{\pi}(p)$ is bounded for all but finitely many primes then

$$\sum_{p \le x} a_{\pi}(p)^2 = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$$

Proof. Say S is the finite set of primes such that a_{π} is bounded for all $p \notin S$. Applying [14, Theorem 3] for π with $\tau_0 = 0$, we get

$$\sum_{p \le x} (\log p) a_{\pi}(p)^2 = x + O(x e^{-c\sqrt{\log x}}).$$

This can be written as,

$$\lim_{x \to \infty} \frac{\left|\sum_{p \le x} (\log p) a_{\pi}(p)^2 - x\right|}{x e^{-c\sqrt{\log x}}} < \infty$$

Since $\lim_{x \to \infty} e^{c\sqrt{\log x}} = \infty$,

$$\lim_{x \to \infty} \frac{\sum_{p \le x} (\log p) a_{\pi}(p)^2 - x}{x} = 0.$$

Hence,

$$\sum_{p \le x} (\log p) a_{\pi}(p)^2 = x + o(x).$$

It implies that

$$\sum_{p \le x} (\log p) a_{\pi}(p)^2 - \sum_{p \le x} (\log x) a_{\pi}(p)^2 + \sum_{p \le x} (\log x) a_{\pi}(p)^2 = x + o(x)$$

We note that,

$$\lim_{x \to \infty} \frac{\sum_{p \le x} (\log x) a_{\pi}(p)^2 - \sum_{p \le x} (\log p) a_{\pi}(p)^2}{x} = \lim_{x \to \infty} \sum_{p \le x} \frac{\log x - \log p}{x} a_{\pi}(p)^2$$
$$= \lim_{x \to \infty} \sum_{p \le x, \ p \in S} \frac{\log x - \log p}{x} a_{\pi}(p)^2 + \lim_{x \to \infty} \sum_{p \le x, \ p \notin S} \frac{\log x - \log p}{x} a_{\pi}(p)^2.$$

Since the first limit has finite summation and $\lim_{x\to\infty} \frac{\log x - \log p}{x} = 0$, the first limit is 0. Say $|a_{\pi}(p)| \leq M$ for $p \notin S$ then

$$\lim_{x \to \infty} \sum_{p \le x, \ p \notin S} \frac{\log x - \log p}{x} a_{\pi}(p)^2 \le M^2 \lim_{x \to \infty} \sum_{p \le x} \frac{\log x - \log p}{x}.$$

 $\sum_{p \leq x} \log p \text{ is called first Chebyshev's function and it is denoted by } \vartheta(x).$

$$\lim_{x \to \infty} \sum_{p \le x} \frac{\log x - \log p}{x} = \lim_{x \to \infty} \sum_{p \le x} \frac{\log x}{x} - \lim_{x \to \infty} \frac{\vartheta(x)}{x} = \lim_{x \to \infty} \frac{\pi(x) \log x}{x} - \lim_{x \to \infty} \frac{\vartheta(x)}{x}.$$

Prime number theorem states that, $\lim_{x \to \infty} \frac{\pi(x) \log x}{x} = \lim_{x \to \infty} \frac{\vartheta(x)}{x} = 1$. Hence

$$\lim_{x \to \infty} \sum_{p \le x, \ p \notin S} \frac{\log x - \log p}{x} = 0$$

And we conclude that,

$$\lim_{x \to \infty} \frac{\sum_{p \le x} (\log x) a_{\pi}(p)^2 - \sum_{p \le x} (\log p) a_{\pi}(p)^2}{x} = 0.$$

Hence, $\sum_{p \le x} (\log p) a_{\pi}(p)^2 - \sum_{p \le x} (\log x) a_{\pi}(p)^2 = o(x)$ and
 $\sum_{p \le x} (\log x) a_{\pi}(p)^2 = x + o(x)$

Dividing the above equation by $\log x$ on both sides proves the lemma.

Lemma 2.3. Let π_1 and π_2 be cuspidal automorphic representations of $\operatorname{GL}_m(\mathbb{A}_{\mathbb{Q}})$ for $m \leq 4$. Assume that they have trivial central character, $\pi_1 \not\cong \pi_2$ and there exists a finite set of primes S such that $|a_{\pi_1}(p) a_{\pi_2}(p)| \leq M$ for some positive constant M and for all $p \notin S$. Then

$$\sum_{p \le x} a_{\pi_1}(p) a_{\pi_2}(p) = o\left(\frac{x}{\log x}\right).$$

Proof. Apply [14, Theorem 3] for π_1 and π_2 with $\tau_0 = 0$, we get

$$\sum_{p \le x} (\log p) a_{\pi_1}(p) a_{\pi_2}(p) = O(x e^{-c\sqrt{\log x}}).$$

Since $\lim_{x \to \infty} e^{c\sqrt{\log x}} = \infty$, similar to previous lemma we conclude that

$$\sum_{p \le x} (\log p) a_{\pi_1}(p) a_{\pi_2}(p) = o(x),$$

which can be written as

$$\sum_{p \le x} (\log p) a_{\pi_1}(p) a_{\pi_2}(p) - \sum_{p \le x} (\log x) a_{\pi_1}(p) a_{\pi_2}(p) + \sum_{p \le x} (\log x) a_{\pi_1}(p) a_{\pi_2}(p) = o(x).$$

We note that,

$$\lim_{x \to \infty} \frac{\sum_{p \le x} (\log x) a_{\pi_1}(p) a_{\pi_2}(p) - \sum_{p \le x} (\log p) a_{\pi_1}(p) a_{\pi_2}(p)}{x} = \lim_{x \to \infty} \sum_{p \le x} \frac{\log x - \log p}{x} a_{\pi_1}(p) a_{\pi_2}(p)$$
$$= \lim_{x \to \infty} \sum_{p \le x, \ p \in S} \frac{\log x - \log p}{x} a_{\pi_1}(p) a_{\pi_2}(p) + \lim_{x \to \infty} \sum_{p \le x, \ p \notin S} \frac{\log x - \log p}{x} a_{\pi_1}(p) a_{\pi_2}(p).$$

Using the bound $|a_{\pi_1}(p)a_{\pi_2}(p)| \leq M$ for $p \notin S$, the above summation is

$$\leq \lim_{x \to \infty} \sum_{p \leq x, \ p \in S} \frac{\log x - \log p}{x} a_{\pi_1}(p) a_{\pi_2}(p) + M \lim_{x \to \infty} \sum_{p \leq x} \frac{\log x - \log p}{x} = 0$$

Similar to the previous lemma, the first limit is zero since the summation is a finite sum and the second limit is zero using the Prime number theorem. Hence,

$$\sum_{p \le x} (\log p) a_{\pi_1}(p) a_{\pi_2}(p) - \sum_{p \le x} (\log x) a_{\pi_1}(p) a_{\pi_2}(p) = o(x)$$
$$\sum_{p \le x} (\log x) a_{\pi_1}(p) a_{\pi_2}(p) = o(x).$$

Corollary 2.3.1. Let $F \in S_k(\Gamma^{(2)}(N))$ be a normalized Hecke eigenform for all primes $p \nmid N$ and satisfies the Ramanujan conjecture. Let $\lambda_F(p)$ represent the eigenvalue for the operator T(p)for all $p \nmid N$. Then

$$\sum_{p \le x, \ p \nmid N} \lambda_F(p)^2 = m \frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$$

where

and

$$m = \begin{cases} 1 & \text{if } F \in \mathbf{G} \\ 2 & \text{if } F \in \mathbf{Y}. \end{cases}$$

Proof. Let π be a self dual unitary cuspidal automorphic representation over $\operatorname{GL}_m(\mathbb{Q})$ for $m \leq 4$. Lemma 2.2 can be written as

$$\sum_{p \le x, \ p \mid N} a_{\pi}(p)^2 + \sum_{p \le x, \ p \nmid N} a_{\pi}(p)^2 = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

The finite sum can be absorbed into $o\left(\frac{x}{\log x}\right)$ and we conclude

$$\sum_{p \le x, \ p \nmid N} a_{\pi}(p)^2 = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

If F is in class **G** then there exists a self dual, unitary cuspidal automorphic representation π of $\operatorname{GL}_4(\mathbb{Q})$ such that $\lambda_F(p) = a_{\pi}(p)$ for all $p \nmid N$. Hence, the corollary follows from Lemma 2.2. If F is in class **Y** then $\lambda_F(p) = a_{\pi_1}(p) + a_{\pi_2}(p)$ for all $p \nmid N$ where π_1, π_2 are distinct self dual, unitary cuspidal automorphic representations of $\operatorname{GL}_2(\mathbb{Q})$. In this case,

$$\lambda_F(p)^2 = a_{\pi_1}(p)^2 + a_{\pi_2}(p)^2 + 2a_{\pi_1}a_{\pi_2}.$$

Applying Lemma 2.2 and 2.3 we get,

$$\sum_{p \le x, \ p \nmid N} \lambda_F(p)^2 = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right) + \frac{x}{\log x} + o\left(\frac{x}{\log x}\right) + o\left(\frac{x}{\log x}\right) = 2\frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

3 Main result

The main result of this article is on comparing the signs of eigenvalues of two distinct modular forms satisfying the Ramanujan conjecture.

Lemma 3.1. Let $F \in S_{k_1}(\Gamma^{(2)}(N_1))$ and $G \in S_{k_2}(\Gamma^{(2)}(N_2))$ be two Hecke eigenforms satisfying the Ramanujan conjecture. Assume that, for some $c \in (0, 4)$ and $\alpha > \frac{15}{16}$, $\#\{p \le x : |\lambda_G(p)| > c\} \ge \alpha \frac{x}{\log x}$ for sufficiently large x. Let S contain all the primes p dividing N_1 and N_2 . Then,

$$\sum_{p \le x, \ p \notin S} \lambda_F(p)^2 \lambda_G(p)^2 \ge \beta \frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$$

for some $\beta \in (0, 1)$.

Proof. Prime Number Theorem states that $\#\{p \le x : p \text{ is prime}\} = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$. Since removing finitely many primes would not effect the asymptotic behavior, we conclude that

$$\#\{p \le x : p \text{ is prime and } p \notin S\} = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$$

Say $S_g(x) = \{p \le x : |\lambda_G(p)| > c\}$. The above equation can be written as,

$$\#\{p \le x : p \notin S_g(x) \text{ and } p \notin S\} = \frac{x}{\log x} + o\left(\frac{x}{\log x}\right) - \#\{p \le x : p \in S_g(x) \text{ and } p \notin S\}.$$

Under the assumption, $\#S_g(x) \ge \alpha \frac{x}{\log x}$ we get,

$$\#\{p \le x : p \notin S_g \text{ and } p \notin S\} \le (1-\alpha)\frac{x}{\log x} + o\left(\frac{x}{\log x}\right)$$

for sufficiently large x. By Weissauer's bound proved in [13], $|\lambda_F(p)| \leq 4$ for all primes $p \notin S$.

$$\sum_{p \le x, \ p \notin S_g(x)} \lambda_F(p)^2 \le 4^2 \# \{ p \le x : p \notin S \cup S_g(x) \} \le 16(1-\alpha) \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

Combining it with Corollary 2.3.1 we get,

$$\sum_{p \le x, \ p \notin S} \lambda_F(p)^2 \ge (16\alpha + m - 16) \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$
$$\sum_{p \le x, \ p \notin S} \lambda_F(p)^2 \lambda_G(p)^2 \ge \sum_{p \le x, \ p \in S_g(x)} \lambda_F(p)^2 \lambda_G(p)^2 \ge c^2 (16\alpha + m - 16) \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

This proves that

$$\sum_{p \le x, \ p \notin S} \lambda_F(p)^2 \lambda_G(p)^2 \ge c^2 (16\alpha + m - 16) \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$
(2)

Lemma 3.2. Let $F \in S_{k_1}(\Gamma^{(2)}(N_1))$ and $G \in S_{k_2}(\Gamma^{(2)}(N_2))$ be two normalised Hecke eigenforms satisfying the Ramanujan conjecture. Assume that if both F, G lift to class Y then

$$L(s, \pi_F, \text{spin}) = L(s, \pi_1)L(s, \pi_2) \text{ and } L(s, \pi_G, \text{spin}) = L(s, \tau_1)L(s, \tau_2)$$

where π_1, π_2, τ_1 and τ_2 are all distinct automorphic representation over $\operatorname{GL}_2(\mathbb{A}_{\mathbb{Q}})$. Under these assumptions, if $\lambda_F(p)$ and $\lambda_G(p)$ are eigenvalues of F and G respectively then

$$\sum_{p \le x, \ p \nmid N} \lambda_F(p) \lambda_G(p) = o\left(\frac{x}{\log x}\right).$$
(3)

Proof. Let π_F , π_G be automorphic representations associated with F and G respectively. There are 4 different possibilities for their L-functions depending on the class of lifts.

- 1. $L(s, \pi_F, \text{spin}) = L(s, \pi_1)$ and $L(s, \pi_G, \text{spin}) = L(s, \pi_2)$ such that π_1, π_2 are distinct self dual, unitary cuspidal automorphic representations of $\text{GL}_4(\mathbb{A}_{\mathbb{Q}})$.
- 2. $L(s, \pi_F, \text{spin}) = L(s, \pi)$ such that π is self dual, unitary cuspidal automorphic representations of $\text{GL}_4(\mathbb{A}_{\mathbb{Q}})$. $L(s, \pi_G, \text{spin}) = L(s, \tau_1)L(s, \tau_2)$ where τ_1, τ_2 are distinct self dual cuspidal automorphic representations of $\text{GL}_2(\mathbb{A}_{\mathbb{Q}})$.
- 3. $L(s, \pi_F, \text{spin}) = L(s, \pi_1)L(s, \pi_2)$ and $L(s, \pi_G, \text{spin}) = L(s, \tau_1)L(s, \tau_2)$ where π_1, π_2, τ_1 and τ_2 are distinct self dual cuspidal automorphic representations of $\text{GL}_2(\mathbb{A}_{\mathbb{Q}})$.
- 4. $L(s, \pi_F, \text{spin}) = L(s, \pi)L(s, \tau_1)$ and $L(s, \pi_G, \text{spin}) = L(s, \pi)L(s, \tau_2)$ where π is self dual cuspidal automorphic representations of $\text{GL}_2(\mathbb{A}_{\mathbb{Q}})$. τ_1 and τ_2 are distinct self dual cuspidal automorphic representations of $\text{GL}_2(\mathbb{A}_{\mathbb{Q}})$.

Assumptions imply that the fourth case is not possible. Hence,

1. $\lambda_F(p) = a_{\pi_1}(p)$ and $\lambda_G(p) = a_{\pi_2}(p)$. Observe that a_{π_1} and a_{π_2} satisfy the conditions for Lemma 2.3. Hence,

$$\sum_{p \nmid N, p \le x} \lambda_F(p) \lambda_G(p) = \sum_{p \nmid N, p \le x} a_{\pi_1}(p) a_{\pi_2}(p) = o\left(\frac{x}{\log x}\right).$$

2. If $\lambda_F(p) = a_{\pi}(p)$ and $\lambda_G(p) = a_{\tau_1}(p) + a_{\tau_2}(p)$, then

$$\sum_{p \nmid N, p \leq x} \lambda_F(p) \lambda_G(p) = \sum_{p \nmid N, p \leq x} a_\pi(p) a_{\tau_1}(p) + \sum_{p \nmid N, p \leq x} a_\pi(p) a_{\tau_2}(p) = o\left(\frac{x}{\log x}\right).$$

3. If $\lambda_F(p) = a_{\pi_1}(p) + a_{\pi_2}(p)$. and $\lambda_G(p) = a_{\tau_1}(p) + a_{\tau_2}(p)$, then $\sum_{p \nmid N, p \leq x} \lambda_F(p) \lambda_G(p)$

$$= \sum_{p \nmid N, p \le x} a_{\pi_1}(p) a_{\tau_1}(p) + \sum_{p \nmid N, p \le x} a_{\pi_1}(p) a_{\tau_2}(p) + \sum_{p \nmid N, p \le x} a_{\pi_2}(p) a_{\tau_1}(p) + \sum_{p \nmid N, p \le x} a_{\pi_2}(p) a_{\tau_2}(p) = o\left(\frac{x}{\log x}\right).$$

Theorem 3.3. Let F, G be two cusp forms satisfying the conditions of Lemma 3.2. Assume there exists a $c \in (0, 4)$ and $\alpha > \frac{15}{16}$ such that

$$\#\{p \le x : |\lambda_G(p)| > c\} \ge \alpha \frac{x}{\log x}$$

for sufficiently large x. Then, the set of primes $\{p : \lambda_F(p)\lambda_G(p) < 0\}$, has positive density. Proof. Let $S = \{p : p \nmid N\}$. Consider the sum

$$S^{-}(x) = \sum_{p \le x, \ p \notin S} (\lambda_F(p)^2 \lambda_G(p)^2 - 16\lambda_F(p)\lambda_G(p)) = \sum_{p \le x, \ p \notin S} \lambda_F(p)\lambda_G(p)[\lambda_F(p)\lambda_G(p) - 16].$$

For $p \notin S$, $|\lambda_F(p)\lambda_G(p)| \leq 16$. Hence, for p such that $\lambda_F(p)\lambda_G(p) > 0$, $\lambda_F(p)\lambda_G(p) - 16 < 0$. Therefore,

$$S^{-}(x) \leq \sum_{p \leq x, \notin S} \sum_{\lambda_F(p)\lambda_G(p) < 0} (\lambda_F(p)^2 \lambda_G(p)^2 - 16\lambda_F(p)\lambda_G(p))$$

$$\leq 512.\#\{p \leq x : p \notin S \text{ and } \lambda_F(p)\lambda_G(p) < 0\}.$$

From Lemma 3.1 and 3.2, we conclude that

$$S^{-}(x) = \sum_{p \le x, \ p \notin S} (\lambda_F(p)^2 \lambda_G(p)^2 - 16\lambda_F(p)\lambda_G(p)) \ge c^2 (16\alpha + m - 16) \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

Combining the inequalities,

$$\#\{p \le x : p \notin S \text{ and } \lambda_F(p)\lambda_G(p) < 0\} \ge \frac{c^2(16\alpha + m - 16)}{512} \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

Since S contains finitely many primes, we can add them to the set to conclude

$$\#\{p \le x : \lambda_F(p)\lambda_G(p) < 0\} \ge \frac{c^2(16\alpha + m - 16)}{512} \frac{x}{\log x} + o\left(\frac{x}{\log x}\right).$$

Hence for $\alpha > \frac{15}{16}$ the set of primes $\{p : \lambda_F(p)\lambda_G(p) < 0\}$ has positive density.

References

- Nagarjuna Chary Addanki. On signs of Hecke eigenvalues of Ikeda lifts. arXiv 2401.08855, 2024.
- [2] Anatoli Andrianov. Introduction to Siegel modular forms and Dirichlet series. Universitext. Springer, New York, 2009.
- [3] James Arthur. The endoscopic classification of representations, volume 61 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 2013. Orthogonal and symplectic groups.
- [4] Mahdi Asgari and Ralf Schmidt. Siegel modular forms and representations. Manuscripta Math., 104(2):173–200, 2001.
- [5] Stefan Breulmann. On Hecke eigenforms in the Maaßspace. Math. Z., 232(3):527–530, 1999.
- [6] Sanoli Gun, Winfried Kohnen, and Biplab Paul. Arithmetic behaviour of Hecke eigenvalues of Siegel cusp forms of degree two. *Ramanujan J.*, 54(1):43–62, 2021.
- [7] Winfried Kohnen. Sign changes of Hecke eigenvalues of Siegel cusp forms of genus two. Proc. Amer. Math. Soc., 135(4):997–999, 2007.
- [8] E. Kowalski, Y.-K. Lau, K. Soundararajan, and J. Wu. On modular signs. Math. Proc. Cambridge Philos. Soc., 149(3):389–411, 2010.
- [9] Hans Maass. über eine Spezialschar von Modulformen zweiten Grades. III. Invent. Math., 53(3):255–265, 1979.
- [10] Ameya Pitale and Ralf Schmidt. Sign changes of Hecke eigenvalues of Siegel cusp forms of degree 2. Proc. Amer. Math. Soc., 136(11):3831–3838, 2008.
- [11] Ralf Schmidt. Iwahori-spherical representations of GSp(4) and Siegel modular forms of degree 2 with square-free level. J. Math. Soc. Japan, 57(1):259–293, 2005.
- [12] Ralf Schmidt. Packet structure and paramodular forms. Trans. Amer. Math. Soc., 370(5):3085–3112, 2018.
- [13] Rainer Weissauer. Endoscopy for GSp(4) and the cohomology of Siegel modular threefolds, volume 1968 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2009.
- [14] Jie Wu and Yangbo Ye. Hypothesis H and the prime number theorem for automorphic representations. Funct. Approx. Comment. Math., 37:461–471, 2007.