CRITICAL THRESHOLD FOR WEAKLY INTERACTING LOG-CORRELATED FOCUSING GIBBS MEASURES

DAMIANO GRECO, TADAHIRO OH, LIYING TAO, AND LEONARDO TOLOMEO

Abstract. We study log-correlated Gibbs measures on the d-dimensional torus with weakly interacting focusing quartic potentials whose coupling constants tend to 0 as we remove regularization. In particular, we exhibit a phase transition for this model by identifying a critical threshold, separating the weakly and strongly coupling regimes; in the weakly coupling regime, we show that the frequency-truncated measures converge to the base Gaussian measure (possibly with a renormalized L^2 -cutoff), whereas, in the strongly coupling regime, we prove nonconvergence of the frequency-truncated measures, even up to a subsequence. Our result answers an open question posed by Brydges and Slade (1996).

CONTENTS

1. Log-correlated Gibbs measures

In this paper, we study the Gibbs measure ρ on the d-dimensional torus on $\mathbb{T}^d = (\mathbb{R}/2\pi\mathbb{Z})^d$, formally given by

$$
d\rho(u) = Z^{-1} \exp\left(\lambda \int_{\mathbb{T}^d} u^4 dx\right) d\mu(u),\tag{1.1}
$$

where the coupling constant $\lambda > 0$ denotes the strength of focusing (i.e. attractive) interaction and μ denotes the log-correlated Gaussian free field on \mathbb{T}^d , formally given by

$$
d\mu = Z^{-1} e^{-\frac{1}{2} ||u||_{H^{d/2}}^2} du.
$$
\n(1.2)

In particular, our interest is to study the *weakly interacting* case whose meaning we will make precise in the following; see [\(1.9\)](#page-1-0).

2020 Mathematics Subject Classification. 60H30, 81T08, 82B26, 35Q55.

Key words and phrases. Gibbs measure; log-correlated Gaussian field; phase transition.

Let us first introduce some notations. Recall that the Gaussian measure μ in [\(1.2\)](#page-0-1) corresponds to the induced probability measure under the map:^{[1](#page-1-1)}

$$
\omega \in \Omega \longmapsto u(\omega) = \sum_{n \in \mathbb{Z}^d} \frac{g_n(\omega)}{\langle n \rangle^{\frac{d}{2}}} e_n,
$$
\n(1.3)

where $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$, $e_n = e^{in \cdot x}$, and $\{g_n\}_{n \in \mathbb{Z}^d}$ is a sequence of mutually independent standard complex-valued Gaussian random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ conditioned that $g_{-n} = \overline{g_n}$, $n \in \mathbb{Z}^d$. It is easy to see from [\(1.3\)](#page-1-2) that a typical element under μ is merely a distribution, thus requiring a renormalization on the interaction potential $\lambda \int_{\mathbb{T}^d} u^4 dx$. Given $N \in \mathbb{N}$, we define the frequency projector π_N onto the frequencies $\{|n| \leq N\}$ by setting

$$
\pi_N f = \sum_{|n| \le N} \hat{f}(n) e_n.
$$
\n(1.4)

Note that, for each fixed $x \in \mathbb{T}^d$, $\pi_N u(x)$ is a mean-zero real-valued Gaussian random variable with variance:

$$
\sigma_N = \mathbb{E}\big[(\pi_N u)^2(x)\big] = \sum_{|n| \le N} \frac{1}{\langle n \rangle^d} \sim \log N \longrightarrow \infty,
$$
\n(1.5)

as $N \to \infty$. We then define the Wick renormalized power $\cdot (\pi_N u)^k$: by setting

$$
:(\pi_N u)^k(x) : \stackrel{\text{def}}{=} H_k(\pi_N u(x); \sigma_N), \tag{1.6}
$$

where $H_k(x;\sigma)$ is the Hermite polynomial of degree k with a variance parameter σ ; see [\[9,](#page-13-1) [15\]](#page-13-2) for further discussions. By setting

$$
R_N(u) = \int_{\mathbb{T}^d} \, : (\pi_N u)^4 : dx,\tag{1.7}
$$

we define the truncated Gibbs measure ρ_N by

$$
d\rho_N(u) = Z_N^{-1} e^{\lambda_N R_N(u)} d\mu(u) = Z_N^{-1} \exp\left(\lambda_N \int_{\mathbb{T}^d} \, : (\pi_N u)^4 : dx\right) d\mu(u). \tag{1.8}
$$

A standard argument shows that $R_N(u)$ is Cauchy in $L^p(\mu)$ for any finite $p \ge 1$; see Lemma [2.4.](#page-4-1) When $\lambda_N \equiv \lambda < 0$ (i.e. the defocusing case),^{[2](#page-1-3)} Nelson's estimate allows us to define the defocusing log-correlated Gibbs measure ρ in [\(1.1\)](#page-0-2) as the unique limit^{[3](#page-1-4)} of the truncated Gibbs measures ρ_N in [\(1.8\)](#page-1-5); see [\[11,](#page-13-3) [17\]](#page-13-4). On the other hand, when $\lambda_N \equiv \lambda > 0$ (i.e. the focusing case), it is known that the Gibbs measure ρ is not normalizable to be a probability measure even with a renormalized L^2 -cutoff; see [\[5,](#page-13-5) [15\]](#page-13-2).^{[4](#page-1-6)}

In [\[5,](#page-13-5) p. 489], Brydges and Slade proposed to study the limiting behavior of the following weakly interacting truncated Gibbs measure ρ_N with a renormalized L^2 -cutoff:

$$
d\rho_N(u) = Z_N^{-1} \mathbf{1}_{\{|\int_{\mathbb{T}^d} : (\pi_N u)^2 : dx| \le K_N\}} e^{\lambda_N R_N(u)} d\mu(u), \tag{1.9}
$$

¹We endow \mathbb{T}^d with the normalized Lebesgue measure $dx_{\mathbb{T}^d} = (2\pi)^{-d} dx$. With a slight abuse of notation, we still use dx to denote the normalized Lebesgue measure.

²Here, the notation " $\lambda_N \equiv \lambda$ " means that the sequence $\{\lambda_N\}_{N\in\mathbb{N}}$ is constant, taking the value λ .

³When $d = 2$, this corresponds to the so-called Φ_2^4 -measure.

⁴We note that the regularization used in [\[5\]](#page-13-5) is different from the frequency truncation and is based on the approximation $(1 - \Delta + \varepsilon \Delta^2)^{-1}$ (as $\varepsilon \to 0$) of the covariance operator $(1 - \Delta)^{-1}$ of the Gaussian free field on \mathbb{T}^2 .

where $R_N(u)$ is as in [\(1.7\)](#page-1-7), by taking $\lambda_N \to 0$ and $K_N \to \infty$ as $N \to \infty$, and investigate existence of a "critical point, separating the weak and strong coupling regimes". Our main result answers this question posed by Brydges and Slade.

Theorem 1.1. Let $\{\lambda_N\}_{N\in\mathbb{N}}$ be a non-increasing sequence of positive numbers tending to 0 as $N \to \infty$, and let $\{K_N\}_{N\in\mathbb{N}}$ be a non-decreasing sequence of positive numbers. Then, there exist $\lambda^* \geq \lambda_* > 0$ such that the following statements hold:

(i) (weakly coupling regime). Suppose that

$$
\lambda_N \le \lambda_*(K_N + \log N)^{-1} \tag{1.10}
$$

for any $N \in \mathbb{N}$. Then, given any $p \geq 1$ we have

$$
\sup_{N \in \mathbb{N}} Z_N := \sup_{N \in \mathbb{N}} \left\| \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (\pi_N u)^2 : dx | \le K_N \}} e^{\lambda_N R_N(u)} \right\|_{L^p(\mu)} < \infty.
$$
 (1.11)

In particular, we have

$$
\lim_{N \to \infty} \mathbf{1}_{\{|\int_{\mathbb{T}^d} : (\pi_N u)^2 : dx| \le K_N\}} e^{\lambda_N R_N(u)} = \mathbf{1}_{\{|\int_{\mathbb{T}^d} : u^2 : dx| \le K\}} \quad in \ L^p(\mu), \tag{1.12}
$$

where $K = \lim_{N \to \infty} K_N \in (0, \infty]$. Here,

$$
\int_{\mathbb{T}^d} :u^2\colon dx=\lim_{N\to\infty}\int_{\mathbb{T}^d}:(\pi_Nu)^2\colon dx,
$$

where the limit is understood in $L^p(\mu)$ and μ -almost surely as in Lemma [2.4.](#page-4-1) As a consequence, we have

- (i.a) If $K = \lim_{N \to \infty} K_N = \infty$, then the truncated Gibbs measure ρ_N in [\(1.9\)](#page-1-0) converges in total variation to the base Gaussian measure μ in [\(1.2\)](#page-0-1) as $N \to \infty$.
- (i.b) If $K = \lim_{N \to \infty} K_N < \infty$, then the truncated Gibbs measure ρ_N in [\(1.9\)](#page-1-0) converges in total variation to the base Gaussian measure with a renormalized L^2 -cutoff:

$$
\mathbf{1}_{\{|\int_{\mathbb{T}^d} : u^2 : dx| \le K\}} d\mu,
$$

as $N \to \infty$.

(ii) (strongly coupling regime). Suppose that

$$
\lambda_N \ge \lambda^* (K_N + \log N)^{-1} \tag{1.13}
$$

for any sufficiently large $N \gg 1$. Then, we have

$$
\sup_{N \in \mathbb{N}} Z_N = \sup_{N \in \mathbb{N}} \mathbb{E}_{\mu} \Big[\mathbf{1}_{\{ |\int_{\mathbb{T}^d} : (\pi_N u)^2 : dx | \le K_N \}} e^{\lambda_N R_N(u)} \Big] = \infty.
$$
 (1.14)

As a consequence, the truncated Gibbs measure ρ_N in [\(1.9\)](#page-1-0) does not converge to any limit in total variation, even up to a subsequence.

Theorem [1.1](#page-2-0) in particular states that the weakly interacting focusing log-correlated Gibbs measure is trivial in the sense that, as we remove regularization, we either obtain the base Gaussian measure (possibly with a renormalized L^2 -cutoff) or non-normalizability / non-convergence.

In a seminal work [\[10\]](#page-13-6), Lebowitz, Rose, and Speer initiated the study of focusing Gibbs measures;^{[5](#page-2-1)} see also [\[3,](#page-13-7) [5,](#page-13-5) [6\]](#page-13-8). In a series of recent works [\[16,](#page-13-9) [13,](#page-13-10) [15,](#page-13-2) [14\]](#page-13-11), the second and fourth authors with their collaborators completed this research program on the (non-)construction of focusing Gibbs measures for any dimension and any power, in particular, by treating critical

⁵More precisely, focusing Φ_d^k -measures, where the base Gaussian measure has covariance $(1 - \Delta)^{-1}$.

models, exhibiting delicate phase transitions, when $d = 1$ and $d = 3$. When $d = 2$, there is no such phase transition in the case $\lambda_N \equiv \lambda > 0$ in [\(1.9\)](#page-1-0). Theorem [1.1](#page-2-0) shows that the weakly interacting model [\(1.9\)](#page-1-0) (proposed by Brydges and Slade [\[5\]](#page-13-5)) is critical when $\lambda_N \sim$ $(K_N + \log N)^{-1}$, nicely complementing the critical models in $d = 1, 3$ studied in [\[16,](#page-13-9) [13,](#page-13-10) [14\]](#page-13-11).

Remark 1.2. (i) While we stated our result in the real-valued setting, a similar result holds in the complex-valued setting by replacing the Hermite polynomials with the Laguerre polynomial; see [\[17\]](#page-13-4) for a further discussion.

(ii) Let $d = 2$. Consider the following weakly interacting truncated nonlinear Schrödinger equation on \mathbb{T}^2 :

$$
i\partial_t u_N + (1 - \Delta)u_N - 4\lambda_N \pi_N(|\pi_N u_N|^2 \pi_N u_N) = 0 \qquad (1.15)
$$

with the initial data distributed by ρ_N in [\(1.9\)](#page-1-0). A standard argument shows that ρ_N is an invariant measure for [\(1.15\)](#page-3-3). Moreover, as a dynamical consequence of Theorem [1.1](#page-2-0) and [\[4\]](#page-13-12), we see that, as $N \to \infty$, the solution u_N to [\(1.15\)](#page-3-3)

- converges to the solution u to the linear Schrödinger equation $i\partial_t u + (1 \Delta)u = 0$, with the initial data distributed by the log-correlated Gaussian measure μ in [\(1.2\)](#page-0-1) (which is an invariant measure for the dynamics), if [\(1.10\)](#page-2-2) holds,
- does not converge to any meaningful limit if (1.13) holds.

See [\[15,](#page-13-2) Subsection 1.2] for other dynamical models related to the log-correlated Gibbs measures.

2. Preliminary lemmas

2.1. Deterministic estimates. We first recall Young's inequality in the general setting; see [\[8,](#page-13-13) Theorem 156 on p. 111].

Lemma 2.1. Let f be a strictly increasing function on \mathbb{R}_+ such that $f(0) = 0$ and its inverse f^{-1} is also strictly increasing. Then, for any $a, b \ge 0$, we have

$$
ab \le \int_0^a f(x)dx + \int_0^b f^{-1}(x)dx
$$
\n(2.1)

with equality if and only if $b = f(a)$. In particular, applying [\(2.1\)](#page-3-4) to $f(x) = e^x - 1$ and $f^{-1}(x) = \log(1+x)$ (with b replaced by $b-1$), we have

$$
ab \le e^a + b \log b - b \tag{2.2}
$$

for any $a > 0$ and $b > 1$.

2.2. Tools from stochastic analysis. In this subsection, we state several useful lemmas from stochastic analysis. We first state the Wiener chaos estimate ([\[19,](#page-13-14) Theorem I.22]); see [\[18,](#page-13-15) Lemma 3.2] for the following particular version.

Lemma 2.2. Let ${g_n}_{n\in\mathbb{Z}^d}$ be a sequence of independent standard real-valued Gaussian random variables. Given $k \in \mathbb{N}$, let $\{P_j\}_{j \in \mathbb{N}}$ be a sequence of polynomials in $\bar{g} = \{g_n\}_{n \in \mathbb{Z}^d}$ of degree at most k. Then, for finite $p \geq 1$, we have

$$
\bigg\|\sum_{j\in\mathbb{N}}P_j(\bar{g})\bigg\|_{L^p(\Omega)}\leq (p-1)^{\frac{k}{2}}\bigg\|\sum_{j\in\mathbb{N}}P_j(\bar{g})\bigg\|_{L^2(\Omega)}.
$$

Next, we recall the following orthogonality result [\[12,](#page-13-16) Lemma 1.1.1].

Lemma 2.3. let f and g be jointly Gaussian random variables with mean zero and variances σ_f and σ_g . Then, we have

$$
\mathbb{E}\big[H_k(f;\sigma_f)H_\ell(g;\sigma_g)\big] = \delta_{k\ell}k!\big\{\mathbb{E}[fg]\big\}^k,
$$

where $H_k(x, \sigma)$ denotes the Hermite polynomial of degree k with a variance parameter σ .

The following convergence result follows from a standard computation, using Lemmas [2.2](#page-3-5) and [2.3;](#page-4-2) see, for example, [\[17\]](#page-13-4) for the proof when $d = 2$, which can be easily generalized to any dimension $d \in \mathbb{N}$.

Lemma 2.4. Let $k \in \mathbb{N}$. Then, given any finite $p \geq 1$, $\int_{\mathbb{T}^d} : (\pi_N u)^k : dx$ converges to a unique limit, denoted by $\int_{\mathbb{T}^d} u^k \colon dx$, in $L^p(\mu)$ and μ -almost surely, as $N \to \infty$, where μ is as in [\(1.2\)](#page-0-1). In particular, given a sequence $\{\lambda_N\}_{N\in\mathbb{N}}$ of positive numbers tending to 0 as $N\to\infty$, $\lambda_N R_N(u)$ converges to 0 in $L^p(\mu)$ and μ -almost surely, as $N \to \infty$, where $R_N(u)$ is as in [\(1.7\)](#page-1-7).

The next lemma plays an important role in studying convergence of the indicator function $1_{\{|\int_{\mathbb{T}^d} :u^2_N: dx|\leq K_N\}}$; see [\[15,](#page-13-2) Lemma 2.4] for the proof. See also [\[7,](#page-13-17) Remark 5.12].

Lemma 2.5. Let μ be as in [\(1.2\)](#page-0-1). Then, we have

$$
\mu\bigg(\int_{\mathbb{T}^d} :u^2\colon dx=K\bigg)=0
$$

for any $K \in \mathbb{R}$, where $\int_{\mathbb{T}^d} u^2 dx$ is the limit of $\int_{\mathbb{T}^d} u(x_n u)^2 dx$ as $N \to \infty$.

2.3. Variational formulation. We prove Theorem [1.1,](#page-2-0) using a variational formula for the partition function, recently popularized in a seminal work [\[1\]](#page-13-18) by Barashkov and Gubinelli; see also [\[13,](#page-13-10) [15,](#page-13-2) [14,](#page-13-11) [20\]](#page-14-0). First, we introduce some notations. Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $W(t)$ be a cylindrical Brownian motion in $L^2(\mathbb{T}^d)$. Namely, we have

$$
W(t) = \sum_{n \in \mathbb{Z}^d} B_n(t)e_n,
$$

where ${B_n}_{n \in \mathbb{Z}^d}$ is a sequence of mutually independent complex-valued^{[6](#page-4-3)} Brownian motions conditioned that $\overline{B_n} = B_{-n}$, $n \in \mathbb{Z}^d$. Then, we define a centered Gaussian process $Y(t)$ by

$$
Y(t) = \langle \nabla \rangle^{-\frac{d}{2}} W(t). \tag{2.3}
$$

In the following, we use the shorthand notation: $Y = Y(1)$. Then, we have Law $(Y) = \mu$, where μ is the log-correlated Gaussian measure defined in [\(1.2\)](#page-0-1). Given $N \in \mathbb{N}$, we set $Y_N = \pi_N Y$. such that Law(Y_N) = $(\pi_N)_*\mu$, i.e. the pushforward of μ under the frequency projector π_N in [\(1.4\)](#page-1-8).

Next, let \mathbb{H}_a denote the space of drifts, which are progressively measurable processes belonging to $L^2([0,1];L^2(\mathbb{T}^d))$, P-almost surely. Then, the Boué-Dupuis variational formula [\[2,](#page-13-19) [23\]](#page-14-1) reads as follow; see [\[24\]](#page-14-2) and [\[20,](#page-14-0) Appendix A] for infinite-dimensional versions.

Lemma 2.6. Given $N \in \mathbb{N}$, let Y_N be as above. Suppose that $F: C^\infty(\mathbb{T}^d) \to \mathbb{R}$ is measurable such that $\mathbb{E}[|F(Y_N)|^p] < \infty$ and $\mathbb{E}[|e^{F(Y_N)}|^q] < \infty$ for some $1 < p, q < \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$. Then, we have

$$
\log \mathbb{E}\Big[e^{F(Y_N)}\Big] = \sup_{\theta \in \mathbb{H}_a} \mathbb{E}\bigg[F(Y_N + \Theta_N) - \frac{1}{2} \int_0^1 \|\theta(t)\|_{L_x^2}^2 dt\bigg],
$$

 6 By convention, we normalize B_n such that $Var(B_n(t)) = t$. In particular, B_0 is a standard real-valued Brownian motion.

where the expectation $\mathbb{E} = \mathbb{E}_{\mathbb{P}}$ is taken with respect to the underlying probability measure \mathbb{P} . Here, $\Theta_N = \pi_N \Theta$, where the process Θ is defined by

$$
\Theta(t) = \int_0^t \langle \nabla \rangle^{-\frac{d}{2}} \theta(t') dt'. \tag{2.4}
$$

We conclude this section by stating basic lemmas in applying the variational formula (Lemma [2.6\)](#page-4-4); see [\[15,](#page-13-2) Lemmas 3.2 and 3.5].

Lemma 2.7. (i) Let $\varepsilon > 0$. Then, given any finite $p \ge 1$, we have

$$
\mathbb{E}\Big[\|Y_N\|_{W^{-\varepsilon,\infty}}^p+\|:Y_N^2\colon\|_{W^{-\varepsilon,\infty}}^p+\|:Y_N^3\colon\|_{W^{-\varepsilon,\infty}}^p\Big]\leq C_{\varepsilon,p}<\infty,
$$

uniformly in $N \in \mathbb{N}$.

(ii) For any $\theta \in \mathbb{H}_a$, we have

$$
\|\Theta\|_{H^{\frac{d}{2}}}^2 \le \int_0^1 \|\theta(t)\|_{L^2}^2 dt.
$$

Lemma 2.8. Given $N \in \mathbb{N}$, let $Y_N = \pi_N Y(1)$, where Y is as in [\(2.3\)](#page-4-5). Then, there exist small $\varepsilon > 0$ and a constant $c_0 = c_0(\varepsilon) > 0$ such that for any $\delta > 0$, we have

$$
\left| \int_{\mathbb{T}^d} :Y_N^3 \colon \Theta dx \right| \leq C(\delta) \| :Y_N^3 \colon \|_{W^{-\varepsilon,\infty}}^2 + \delta \|\Theta\|_{H^{\frac{d}{2}}}^2,
$$

$$
\left| \int_{\mathbb{T}^d} :Y_N^2 \colon \Theta^2 dx \right| \leq C(\delta) \| :Y_N^2 \colon \|_{W^{-\varepsilon,\infty}}^4 + \delta \left(\|\Theta\|_{H^{\frac{d}{2}}}^2 + \|\Theta\|_{L^4}^4 \right),
$$

$$
\left| \int_{\mathbb{T}^d} Y_N \Theta^3 dx \right| \leq C(\delta) \|Y_N\|_{W^{-\varepsilon,\infty}}^{c_0} + \delta \left(\|\Theta\|_{H^{\frac{d}{2}}}^2 + \|\Theta\|_{L^4}^4 \right),
$$

uniformly in $N \in \mathbb{N}$.

3. Weakly coupling regime: normalizability

In this section, we present a proof of Theorem [1.1](#page-2-0) (i). Let $K = \lim_{N \to \infty} K_N \in (0, \infty]$. Then, it follows from Lemmas [2.4](#page-4-1) and [2.5](#page-4-6) that $1_{\{| \int_{\mathbb{T}^d} : (\pi_N u)^2 : dx | \le K_N\}} e^{\lambda_N R_N(u)}$ converges to $\mathbf{1}_{\{| \int_{\mathbb{T}^d} : u^2 : dx | \leq K\}}$ in probability (with respect to μ) as $N \to \infty$. Then, the desired $L^p(\mu)$ convergence [\(1.12\)](#page-2-4) follows from the uniform integrability bound [\(1.11\)](#page-2-5); see [\[21,](#page-14-3) Remark 3.8]. See also the discussion at the end of Section 2 in [\[17\]](#page-13-4).

Given $N \in \mathbb{N}$, define $\mathcal{W}_N(\theta)$ by

$$
\mathcal{W}_N(\theta) = \mathbb{E}\bigg[\lambda_N R_N(Y + \Theta) \cdot \mathbf{1}_{\{\|f_{\mathbb{T}^d} : (Y_N + \Theta_N)^2 : dx\| \le K_N\}} - \frac{1}{2} \int_0^1 \|\theta(t)\|_{L_x^2}^2 dt\bigg],
$$

where Θ is as in [\(2.4\)](#page-5-1). Then, from Lemma [2.6](#page-4-4) with [\(2.3\)](#page-4-5), we see that [\(1.11\)](#page-2-5) follows once we prove

$$
\sup_{N \in \mathbb{N}} \log \mathbb{E}_{\mu} \bigg[\exp \Big(\lambda_N R_N(u) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (\pi_N u)^2 : dx | \le K_N \}} \Big) \bigg] = \sup_{N \in \mathbb{N}} \sup_{\theta \in \mathbb{H}_a} \mathcal{W}_N(\theta) < \infty. \tag{3.1}
$$

From (1.6) and (1.7) we have

$$
\lambda_N R_N(Y + \Theta) = \lambda_N \int_{\mathbb{T}^d} :Y_N^4 \colon dx + 4\lambda_N \int_{\mathbb{T}^d} :Y_N^3 \colon \Theta_N dx + 6\lambda_N \int_{\mathbb{T}^d} :Y_N^2 \colon \Theta_N^2 dx
$$

+ 4\lambda_N \int_{\mathbb{T}^d} Y_N \Theta_N^3 dx + \lambda_N \int_{\mathbb{T}^d} \Theta_N^4 dx. (3.2)

By applying Lemma [2.8](#page-5-2) and Lemma [2.7](#page-5-3) to [\(3.2\)](#page-5-4), we have

$$
\mathcal{W}_N(\theta) \le C_0 + \mathbb{E}\bigg[2\lambda_N \|\Theta_N\|_{L^4}^4 \cdot \mathbf{1}_{\{\|\int_{\mathbb{T}^d} : (Y_N^2 + \Theta_N)^2 : dx\le K\}} - \frac{1}{4} \int_0^1 \|\theta(t)\|_{L^2_x}^2 dt\bigg],\tag{3.3}
$$

uniformly in $N \in \mathbb{N}$ and $\theta \in \mathbb{H}_a$. Hence, [\(3.1\)](#page-5-5) (and thus [\(1.11\)](#page-2-5)) follows from [\(3.3\)](#page-6-0) once we prove the following proposition.

Proposition 3.1. There exists small $\lambda_* > 0$ such that if [\(1.10\)](#page-2-2) holds, then we have

$$
\sup_{\theta \in \mathbb{H}_a} \mathbb{E} \left[\lambda_N \| \Theta_N \|_{L^4}^4 \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (Y_N^2 + \Theta_N)^2 : dx | \le K_N \}} - \frac{1}{10} \int_0^1 \| \theta(t) \|_{L^2_x}^2 dt \right] \lesssim 1,
$$
 (3.4)

uniformly in $N \in \mathbb{N}$.

Suppose that λ_N decays at a rate of the form $\lambda_N \lesssim N^{-\kappa}$ for some $\kappa > 0$. Then, by Sobolev's inequality, interpolation, and (standard) Young's inequality, we have

$$
\lambda_N \|\Theta_N\|_{L^4}^4 \le C \|\Theta_N\|_{H^{\frac{d-\kappa}{4}}}^4 \le C' \|\Theta_N\|_{H^{\frac{d}{2}}}^{\frac{2(d-\kappa)}{d}} \|\Theta_N\|_{L^2}^{\frac{2(d+\kappa)}{d}} \le C' \|\Theta_N\|_{L^2}^{\frac{2(d+\kappa)}{d}} \tag{3.5}
$$
\n
$$
\le \frac{1}{10} \|\Theta_N\|_{H^{\frac{d}{2}}}^2 + C'' \|\Theta_N\|_{L^2}^{\frac{2(d+\kappa)}{\kappa}},
$$

where the first term on the right-hand side is then controlled by Lemma [2.7](#page-5-3) (ii). When $\lambda_N \sim$ $(\log N)^{-1}$ (essentially corresponding to $\kappa = 0$), such an argument does not work, exhibiting the critical nature of our problem. While our argument is motivated by those in [\[13,](#page-13-10) Subsection 5.6] and [\[14,](#page-13-11) Subsection 3.2], there is an additional difficulty in our current problem as compared to those in [\[13,](#page-13-10) [14\]](#page-13-11) in the following sense. In [\[13,](#page-13-10) [14\]](#page-13-11), the essential part in estimating the potential energies was reduced to estimating $\|\Theta_N\|_{L^2}^6$; see [\[13,](#page-13-10) (5.76)] and [\[14,](#page-13-11) (3.21)]. On the other hand, when $\kappa = 0$, [\(3.5\)](#page-6-1) would give us an infinite power of the L^2 -norm of Θ_N ; see the first term on the right-hand side of [\(3.10\)](#page-7-0).

Proof of Proposition [3.1.](#page-6-2) On $A_N := \{ | \int_{\mathbb{T}^d} : (Y_N^2 + \Theta_N)^2 : dx | \le K_N \}$, we have

$$
\|\Theta_N\|_{L^2}^2 \le K_N + \sigma_N + 2 \bigg| \int_{\mathbb{T}^d} Y_N \Theta_N dx \bigg|, \tag{3.6}
$$

where $\sigma_N \sim \log N$ is as in [\(1.5\)](#page-1-10). First, suppose that we have

$$
\|\Theta_N\|_{L^2}^2 \lesssim K_N + \sigma_N. \tag{3.7}
$$

Then, from Sobolev's inequality, interpolation, (3.6) , and (3.7) with (1.5) and (1.10) followed by Lemma 2.7 (ii), we obtain

$$
\lambda_N \|\Theta_N\|_{L^4}^4 \cdot \mathbf{1}_{\{\|\int_{\mathbb{T}^d} : (Y_N^2 + \Theta_N)^2 : dx| \le K_N\}}\n\le C\lambda_N \|\Theta_N\|_{L^2}^2 \|\Theta_N\|_{H^{\frac{d}{2}}}^2 \cdot \mathbf{1}_{\{\|\int_{\mathbb{T}^d} : (Y_N^2 + \Theta_N)^2 : dx| \le K_N\}}\n\le C'\lambda_* \|\Theta_N\|_{H^{\frac{d}{2}}}^2 \le \frac{1}{10} \int_0^1 \|\theta(t)\|_{L^2}^2 dt,
$$

provided that λ_* is sufficiently small. This yields [\(3.4\)](#page-6-5) under the condition [\(3.7\)](#page-6-4).

In view of [\(3.6\)](#page-6-3), it remains to consider the case:

$$
\|\Theta_N\|_{L^2}^2 \lesssim \left| \int_{\mathbb{T}} Y_N \Theta_N dx \right|.
$$
 (3.8)

We note that the following argument holds under a weaker assumption:

$$
\lambda_N \le \lambda_* (\log N)^{-1}, \quad N \in \mathbb{N}.\tag{3.9}
$$

By applying [\(2.2\)](#page-3-6) in Lemma [2.1,](#page-3-7) [\(3.9\)](#page-7-1), and Bernstein's inequality (recall that supp $\widehat{\Theta}_N \subset \{|n| \leq$ $N\},$ we have

$$
\lambda_N \|\Theta_N\|_{L^4}^4 \lesssim \lambda_*^{\frac{1}{2}} (\log N)^{-1} \|\Theta_N\|_{L^2}^2 \left(\lambda_*^{\frac{1}{2}} \|\Theta_N\|_{L^2}^2 \frac{\|\Theta_N\|_{H^{\frac{d}{2}}}^2}{\|\Theta_N\|_{L^2}^2} \right)
$$

\n
$$
\leq \lambda_*^{\frac{1}{2}} (\log N)^{-1} \|\Theta_N\|_{L^2}^2 e^{\lambda_*^{\frac{1}{2}} \|\Theta_N\|_{L^2}^2}
$$

\n
$$
+ \lambda_*^{\frac{1}{2}} (\log N)^{-1} \|\Theta_N\|_{H^{\frac{d}{2}}}^2 \log \frac{\|\Theta_N\|_{H^{\frac{d}{2}}}^2}{\|\Theta_N\|_{L^2}^2}
$$

\n
$$
\lesssim \lambda_*^{\frac{1}{2}} (\log N)^{-1} e^{2\lambda_*^{\frac{1}{2}} \|\Theta_N\|_{L^2}^2} + \lambda_*^{\frac{1}{2}} \|\Theta_N\|_{H^{\frac{d}{2}}}^2.
$$

\n(3.10)

We now claim that there exists a non-negative random variable $X_N(\omega)$ with

$$
\sup_{N \in \mathbb{N}} \mathbb{E}[X_N] < \infty \tag{3.11}
$$

such that

$$
e^{2\lambda_*^{\frac{1}{2}} \|\Theta_N\|_{L^2}^2} \lesssim 1 + \|\Theta_N\|_{H^{\frac{d}{2}}}^2 + X_N(\omega). \tag{3.12}
$$

Then, [\(3.4\)](#page-6-5) follows from [\(3.10\)](#page-7-0), [\(3.12\)](#page-7-2), and Lemma [2.7](#page-5-3) (ii), provided that λ_* is sufficiently small.

The remaining part of the proof is devoted to proving [\(3.12\)](#page-7-2). We proceed as in the proof of [\[14,](#page-13-11) Lemma 3.6] (see also [\[13,](#page-13-10) Subsection 5.6]). Define the sharp frequency projections ${\{\Pi_j\}}_{j\in\mathbb{N}}$ by setting $\Pi_1 = \pi_2$ and $\Pi_j = \pi_{2^j} - \pi_{2^{j-1}}$. We also set $\Pi_{\leq j} = \sum_{k=1}^j \Pi_k$ and $\Pi_{>j} = \text{Id} - \Pi_{\leq j}$. Then, write Θ_N as

$$
\Theta_N = \sum_{j=1}^{\infty} (\alpha_j \Pi_j Y_N + w_j),
$$

where

$$
\alpha_j := \begin{cases} \frac{\langle \Theta_N, \Pi_j Y_N \rangle}{\|\Pi_j Y_N\|_{L^2}^2}, & \text{if } \|\Pi_j Y_N\|_{L^2} \neq 0, \\ 0, & \text{otherwise}, \end{cases} \quad \quad \text{and} \quad \quad w_j := \Pi_j \Theta_N - \alpha_j \Pi_j Y_N.
$$

Noting that w_j is orthogonal to $\Pi_j Y_N$ and Y_N in $L^2(\mathbb{T}^d)$, we have

$$
\|\Theta_N\|_{L^2}^2 = \sum_{j=1}^{\infty} \left(\alpha_j^2 \|\Pi_j Y_N\|_{L^2}^2 + \|w_j\|_{L^2}^2 \right),
$$

$$
\int_{\mathbb{T}^d} Y_N \Theta_N dx = \sum_{j=1}^{\infty} \alpha_j \|\Pi_j Y_N\|_{L^2}^2.
$$
 (3.13)

Given small $N \in \mathbb{N}$, fix a random number $j_0 \in \mathbb{N}$ (to be chosen later). Then, arguing as in the proof of Lemma 3.6 in [\[14\]](#page-13-11) (see [\[14,](#page-13-11) $(3.49)-(3.51)$]), we obtain

$$
\left| \sum_{j=1}^{\infty} \alpha_j \|\Pi_j Y_N\|_{L^2}^2 \right| \lesssim \|\Theta_N\|_{H^{\frac{d}{2}}} \|\Pi_{>j_0} Y_N\|_{H^{-\frac{d}{2}}} + \|\Pi_{\leq j_0} Y_N\|_{L^2}^2.
$$
 (3.14)

Since Y_N is spatially homogeneous, we have

$$
\|\Pi_{>j_0} Y_N\|_{H^{-\frac{d}{2}}}^2 = \int_{\mathbb{T}^d} \left(\langle \nabla \rangle^{-\frac{d}{2}} \Pi_{>j_0} Y_N \right)^2 : dx + \mathbb{E} \left[\left(\langle \nabla \rangle^{-\frac{d}{2}} \Pi_{>j_0} Y_N \right)^2 \right],\tag{3.15}
$$

where the last term is independent of $x \in \mathbb{T}^d$ (and hence we suppressed its x-dependence). From (2.3) , we have

$$
\widetilde{\sigma}_{j_0} := \mathbb{E}\left[(\langle \nabla \rangle^{-\frac{d}{2}} \Pi_{>j_0} Y_N)^2 \right] = \sum_{|n|>2^{j_0}} \frac{1}{\langle n \rangle^{2d}} \sim 2^{-dj_0}.
$$
\n(3.16)

Proceeding as in the proof of Lemma 2.5 in [\[17\]](#page-13-4) with Lemma [2.3,](#page-4-2) we have

$$
\mathbb{E}\left[\left(\int_{\mathbb{T}^d} :(\langle\nabla\rangle^{-\frac{d}{2}}\Pi_{>j_0}Y_N)^2:dx\right)^2\right] \n= \int_{\mathbb{T}^d_x\times\mathbb{T}^d_y} \mathbb{E}\left[H_2(\langle\nabla\rangle^{-\frac{d}{2}}\Pi_{>j_0}Y_N(x);\tilde{\sigma}_{j_0})H_2(\langle\nabla\rangle^{-\frac{d}{2}}\Pi_{>j_0}Y_N(y);\tilde{\sigma}_{j_0})\right]dxdy \n= 2\int_{\mathbb{T}^d_x\times\mathbb{T}^d_y} \left\{\mathbb{E}\left[\langle\nabla\rangle^{-\frac{d}{2}}\Pi_{>j_0}Y_N(x)\cdot\langle\nabla\rangle^{-\frac{d}{2}}\Pi_{>j_0}Y_N(y)\right]\right\}^2 dxdy \n= 2\sum_{\substack{n_1,n_2\in\mathbb{Z}^d\\2^{j_0} <|n_j| \le N}} \frac{1}{\langle n_1\rangle^{2d}\langle n_2\rangle^{2d}} \int_{\mathbb{T}^3_x\times\mathbb{T}^3_y} e_{n_1+n_2}(x-y)dxdy \n\lesssim \sum_{\substack{n_1,n_2\in\mathbb{Z}^d\\|n|>2^{j_0}}}\frac{1}{\langle n\rangle^{4d}} \sim 2^{-3dj_0}.
$$
\n(3.17)

Now, define a non-negative random variable $B_{1,N}(\omega)$ by setting

$$
B_{1,N}(\omega) = \left(\sum_{k=1}^{\infty} 2^{\frac{5}{2}dk} \left(\int_{\mathbb{T}^d} \left(\langle \nabla \rangle^{-\frac{d}{2}} \Pi_{>k} Y_N \right)^2 : dx\right)^2\right)^{\frac{1}{2}}.
$$
 (3.18)

From [\(3.15\)](#page-8-0), [\(3.16\)](#page-8-1), and [\(3.18\)](#page-8-2), we obtain

$$
\|\Pi_{>j_0} Y_N\|_{H^{-\frac{d}{2}}}^2 \lesssim 2^{-\frac{5d}{4}j_0} B_{1,N}(\omega) + 2^{-dj_0}.
$$
 (3.19)

Let us now consider the second term on the right-hand side of [\(3.14\)](#page-7-3). As in [\(3.15\)](#page-8-0), write

$$
\|\Pi_{\leq j_0} Y_N\|_{L^2}^2 = \int_{\mathbb{T}^d} : (\Pi_{\leq j_0} Y_N)^2 : dx + \mathbb{E}\big[(\Pi_{\leq j_0} Y_N)^2 \big]. \tag{3.20}
$$

We have

$$
\mathbb{E}\left[\left(\Pi_{\leq j_0} Y_N\right)^2\right] = \sum_{|n| \leq 2^{j_0}} \frac{1}{\langle n \rangle^d} \sim j_0. \tag{3.21}
$$

Proceeding as in [\(3.17\)](#page-8-3), we have

$$
\mathbb{E}\left[\left(\int_{\mathbb{T}^d} : (\Pi_k Y_N)^2 : dx\right)^2\right] \lesssim 2^{-dk}.\tag{3.22}
$$

As in [\(3.18\)](#page-8-2), define a non-negative random variable $B_{2,N}(\omega)$ by setting

$$
B_{2,N}(\omega) = \sum_{k=1}^{\infty} \left| \int_{\mathbb{T}^d} \left| \left(\Pi_k Y_N \right)^2 \right| \, dx \right|.
$$
 (3.23)

Thus, from [\(3.20\)](#page-8-4), [\(3.21\)](#page-8-5), and [\(3.22\)](#page-8-6), we have

$$
\|\Pi_{\leq j_0} Y_N\|_{L^2}^2 \lesssim B_{2,N}(\omega) + j_0. \tag{3.24}
$$

We now choose $j_0 = j_0(\omega)$ by

$$
2^{\frac{d}{2}j_0(\omega)} \sim 2 + ||\Theta_N(\omega)||_{H^{\frac{d}{2}}}.
$$
\n(3.25)

Then, putting [\(3.8\)](#page-6-6), [\(3.13\)](#page-7-4), [\(3.14\)](#page-7-3), [\(3.19\)](#page-8-7), and [\(3.24\)](#page-9-1) together with [\(3.25\)](#page-9-2), we have

$$
\|\Theta_N\|_{L^2}^2 \le \left(2^{-\frac{d}{2}j_0} + 2^{-\frac{5d}{8}j_0} B_{1,N}^{\frac{1}{2}}(\omega)\right) \|\Theta_N\|_{H^{\frac{d}{2}}} + B_{2,N}(\omega) + j_0
$$

\$\lesssim \log\left(2 + \|\Theta_N\|_{H^{\frac{d}{2}}}\right) + B_{1,N}^{\frac{1}{2}}(\omega) + B_{2,N}(\omega).

Hence, by Young's inequality and choosing $\lambda_* > 0$ sufficiently small, we obtain

$$
e^{2\lambda_*^{\frac{1}{2}}\|\Theta_N\|_{L^2}^2} \lesssim 1 + \|\Theta_N\|_{H^{\frac{d}{2}}}^2 + X_N,
$$

yielding [\(3.12\)](#page-7-2), where $X_N = X_N(\omega)$ is defined by

$$
X_N(\omega) = e^{\gamma B_{1,N}^{\frac{1}{2}}(\omega) + \gamma B_{2,N}(\omega)}
$$
\n(3.26)

with some small constant $\gamma > 0$, which we now choose to guarantee the uniform bound [\(3.11\)](#page-7-5). From [\(3.18\)](#page-8-2), Minkowski's integral inequality, the Wiener chaos estimate (Lemma [2.2\)](#page-3-5), and [\(3.17\)](#page-8-3) (with j_0 replaced by k), we see that

$$
\mathbb{E}\big[B_{1,N}^p\big]\lesssim p
$$

for any finite $p \ge 1$, uniformly in $N \in \mathbb{N}$. Then, it follows from [\[22,](#page-14-4) Lemma 4.5] that there exists $\gamma > 0$ such that

$$
\mathbb{E}\left[e^{\gamma B_{1,N}^{\frac{1}{2}}}\right] \lesssim 1,\tag{3.27}
$$

uniformly in $N \in \mathbb{N}$. A similar computation applied to [\(3.23\)](#page-8-8) yields

$$
\mathbb{E}\left[e^{\gamma B_{2,N}}\right] \lesssim 1,\tag{3.28}
$$

uniformly in $N \in \mathbb{N}$. Hence, the bound [\(3.11\)](#page-7-5) follows from [\(3.26\)](#page-9-3), [\(3.27\)](#page-9-4), and [\(3.28\)](#page-9-5).

4. Strongly coupling regime: non-normalizability

In this section, we present a proof of Theorem [1.1](#page-2-0) (ii) by following closely the argument in [\[15,](#page-13-2) Section 3], which was in turn is inspired by the recent works by the fourth author with Weber [\[20\]](#page-14-0) and by the second and fourth authors with Okamoto [\[13,](#page-13-10) [14\]](#page-13-11). For this purpose, we first recall notations and preliminary results from [\[15\]](#page-13-2).

Let $f: \mathbb{R}^d \to \mathbb{R}$ be a real-valued Schwartz function with $||f||_{L^2(\mathbb{R}^d, \frac{dx}{(2\pi)^d})} = 1$ such that its Fourier transform \hat{f} is supported on $\{\xi \in \mathbb{R}^d : |\xi| \leq 1\}$ with $\hat{f}(0) = 0$. Define a function f_M on \mathbb{T}^d by

$$
f_M = M^{-\frac{d}{2}} \sum_{|n| \le M} \widehat{f}\left(\frac{n}{M}\right) e_n,\tag{4.1}
$$

where $\widehat{f} = \mathcal{F}_{\mathbb{R}^d}(f)$ denotes the Fourier transform on \mathbb{R}^d .

Lemma 4.1 (Lemma 3.3 in [\[15\]](#page-13-2)). Let $\alpha > 0$. Then, we have

$$
\int_{\mathbb{T}^d} f_M^2 dx = 1 + O(M^{-\alpha}),
$$
\n(4.2)

$$
\int_{\mathbb{T}^d} f_M^4 dx \sim M^d,\tag{4.3}
$$

$$
\int_{\mathbb{T}^d} (\langle \nabla \rangle^{-\frac{d}{2}} f_M)^2 dx \sim M^{-d} \tag{4.4}
$$

for any $M \gg 1$.

We also define $Q(u)$ by

$$
Q(u) = \int_{\mathbb{T}^d} u^4 dx.
$$
\n(4.5)

As in [\[15\]](#page-13-2), the divergence of $Q(f_M)$ (see [\(4.3\)](#page-10-0)) is what allows us to prove [\(1.14\)](#page-2-6).

In the following, we split the proof of Theorem [1.1](#page-2-0) (ii) into two cases, depending on whether $\sup_{N\in\mathbb{N}}(\log N)^{-1}K_N<\infty$ or not. In the former case, we use exactly the same drift θ^0 as in [\[15\]](#page-13-2), which we recall now. In [\[15,](#page-13-2) Lemma 3.4], given $M \gg 1$, we constructed the approximation process ζ_M to Y in [\(2.3\)](#page-4-5) by solving the stochastic differential equation [\[15,](#page-13-2) (3.18)] on low frequencies $\{|n| \leq M\}$ and setting $\widehat{\zeta}_M(n, t) \equiv 0$ on high frequencies $\{|n| > M\}$. In [\[15,](#page-13-2) (3.27)], we then defined a drift θ^0 by setting

$$
\theta^{0}(t) = \langle \nabla \rangle^{\frac{d}{2}} \bigg(-\frac{d}{dt} \zeta_{M}(t) + \sqrt{\alpha_{M,N}} f_{M} \bigg),
$$

where $\alpha_{M,N}$ is as in [\[15,](#page-13-2) (3.25)]. As in [\(2.4\)](#page-5-1), we then set

$$
\Theta^0 = \int_0^1 \langle \nabla \rangle^{-\frac{d}{2}} \theta^0(t) dt = -\zeta_M + \sqrt{\alpha_{M,N}} f_M.
$$

From [\[15,](#page-13-2) (3.26) and (3.34)] and the frequency supports of f_M and ζ_M , we have

$$
\alpha_{M,N} = \sigma_M(1 + o(1)) \sim \log M,\tag{4.6}
$$

$$
\mathbb{E}\left[\|\Theta^0\|_{H^{\frac{d}{2}}}^2\right] \le \mathbb{E}\left[\int_0^1 \|\theta^0(t)\|_{L_x^2}^2 dt\right] \lesssim M^d \log M,
$$
\n
$$
\pi_N \Theta^0 = \Theta_N^0 = \Theta^0
$$
\n(4.7)

for any $N \geq M \gg 1$.

The following lemma follows from a slight modification of the proof of [\[15,](#page-13-2) (3.42)].

Lemma 4.2. Let $\{K_N\}_{N\in\mathbb{N}}$ be a non-decreasing sequence of positive numbers. There exists $M_0 = M_0(K_1) \in \mathbb{N}$ such that

$$
\mathbb{P}\bigg(\Big|\int_{\mathbb{T}^d} (:Y_N^2 : +2Y_N\Theta^0 + (\Theta^0)^2) dx\Big| \le K_N\bigg) \ge \frac{1}{2}
$$

for any $N \geq M \geq M_0(K_1)$.

We now present a proof of Theorem [1.1](#page-2-0) (ii).

Proof of Theorem [1.1](#page-2-0) (ii). We only prove the divergence in (1.14) , since the claimed nonconvergence then follows from Lemma [2.4.](#page-4-1) Noting that

$$
\mathbb{E}_{\mu}\left[\exp\left(\min(\lambda_{N}R_{N}(u),L)\right)\cdot\mathbf{1}_{\{|\int_{\mathbb{T}^d}:(\pi_Nu)^2:dx|\leq K_N\}}\right]
$$

$$
\geq \mathbb{E}_{\mu}\left[\exp\left(\min(\lambda_{N}R_{N}(u),L)\cdot\mathbf{1}_{\{|\int_{\mathbb{T}^d}:(\pi_Nu)^2:dx|\leq K_N\}}\right)\right]-1,
$$

it suffices to prove

$$
\liminf_{N \to \infty} \lim_{L \to \infty} \mathbb{E}_{\mu} \Big[\exp \big(\min \big(\lambda_N R_N(u), L \big) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (\pi_N u)^2 : dx | \le K_N \}} \big) \Big] = \infty. \tag{4.8}
$$

We split the proof into two cases.

• Case 1: $\sup_{N \in \mathbb{N}} (\log N)^{-1} K_N < \infty$. In this case, from [\(1.13\)](#page-2-3), we have

$$
\lambda_N \ge C\lambda^* (\log N)^{-1}, \quad N \in \mathbb{N}, \tag{4.9}
$$

for some $C > 0$. In the following, we take $N \geq M \gg 1$.

From [\(3.2\)](#page-5-4) (with Θ_N replaced by Θ^0) and Lemma [2.8,](#page-5-2) we have

$$
\lambda_N R_N(Y + \Theta^0) \ge (1 - \delta)\lambda_N Q(\Theta^0)
$$

- $c(\delta)\lambda_N \left(||\cdot Y_N^3 \cdot ||_{W^{-\varepsilon,\infty}}^2 + ||\cdot Y_N^2 \cdot ||_{W^{-\varepsilon,\infty}}^4 + ||Y_N||_{W^{-\varepsilon,\infty}}^{c_0} \right)$
- $\lambda_N |R_N(Y)| - \delta\lambda_N ||\Theta^0||_{H^{\frac{d}{2}}}^2.$ (4.10)

See [\[15,](#page-13-2) (3.41)]. Proceeding as in [\[15,](#page-13-2) (3.31)-(3.33)] with [\(4.6\)](#page-10-1), Lemma [4.1,](#page-10-2) and Lemma [4.2,](#page-10-3) we have

$$
\mathbb{E}\Big[\min((1-\delta)\lambda_N Q(\Theta^0), L) \cdot \mathbf{1}_{\{\mid \int_{\mathbb{T}^d} : (Y_N + \Theta^0)^2 : dx \mid \le K_N\}}\Big] \ge C\lambda_N \alpha_{M,N}^2 M^d - C(\delta)\lambda_N \alpha_{M,N} \mathbb{E}[\zeta_M^2] \|f_M\|_{L^2}^2 \ge C'\lambda_N M^d (\log M)^2 - C''(\delta)\lambda_N (\log M)^2.
$$
\n(4.11)

for any small $\delta > 0$ and $N \ge M \gg 1$, provided that $L \gg \lambda_N \alpha_{M,N}^2 Q(f_M) \sim \lambda_N \alpha_{M,N}^2 M^d$. Thus, it follows from the variational formula (Lemma [2.6\)](#page-4-4), [\(4.10\)](#page-11-0), [\(4.11\)](#page-11-1), Lemmas [2.7](#page-5-3) and [2.4,](#page-4-1) and [\(4.7\)](#page-10-4) that

$$
\log \mathbb{E}_{\mu} \Big[\exp \Big(\min \big(\lambda_N R_N(u), L \big) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : u_N^2 : dx | \le K_N \}} \Big) \Big] \ge \mathbb{E} \Big[\min \big(\lambda_N R_N(Y + \Theta^0), L \big) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (Y_N + \Theta^0)^2 : dx | \le K_N \}} - \frac{1}{2} \int_0^1 \| \theta^0(t) \|_{L_x^2}^2 dt \Big] \qquad (4.12) \ge \lambda_N M^d (\log M)^2 - C(\delta) \lambda_N (\log M)^2 - C'M^d \log M - C''(\delta)
$$

for any small $\delta > 0$ and $N \ge M \gg 1$, provided $L \gg \lambda_N \alpha_{M,N}^2 M^d$. In view of [\(4.9\)](#page-11-2), by setting $M = N$ in [\(4.12\)](#page-11-3), taking $L \to \infty$, and then $N \to \infty$, [\(4.8\)](#page-11-4) follows from (4.12), provided that λ^* is sufficiently large.

• Case 2: $\sup_{N \in \mathbb{N}} (\log N)^{-1} K_N = \infty$. In this case, from (1.13) , we have

$$
\lambda_N \ge C\lambda^* K_N^{-1}, \quad N \in \mathbb{N}, \tag{4.13}
$$

for some $C > 0$. Under the condition (4.13) , a deterministic drift suffices for our purpose. Given $M \gg 1$, let f_M be as in [\(4.1\)](#page-9-6) and we define a drift θ_γ by

$$
\theta_{\gamma}(t) = \sqrt{\gamma K_M} \langle \nabla \rangle^{\frac{d}{2}} f_M,
$$
\n(4.14)

for some small $\gamma > 0$ is independent of M (to be chosen later). We then set

$$
\Theta_{\gamma} = \int_0^1 \langle \nabla \rangle^{-\frac{d}{2}} \theta_{\gamma}(t) dt = \sqrt{\gamma K_M} f_M.
$$
\n(4.15)

From the frequency support of f_M , we have

 $\pi_N \Theta_\gamma = \Theta_\gamma$

for any $N \geq M \gg 1$. From [\(4.15\)](#page-12-0), Cauchy-Schwarz's inequality (in time), [\(4.14\)](#page-12-1), Bernstein's inequality (with supp $\widehat{f}_M \subset \{|n| \le M\}$), and Lemma [4.1,](#page-10-2) we have

$$
\|\Theta_{\gamma}\|_{H^{\frac{d}{2}}}^{2} \leq \int_{0}^{1} \|\theta_{\gamma}(t)\|_{L_{x}^{2}}^{2} dt \lesssim \gamma K_{M} M^{d}
$$
\n(4.16)

for any $M \gg 1$. From [\(4.5\)](#page-10-5) and Lemma [4.1,](#page-10-2) we also have

$$
Q(\Theta_{\gamma}) \sim \gamma^2 K_M^2 M^d \tag{4.17}
$$

for any $M \gg 1$.

We claim that by choosing $\gamma > 0$ sufficiently small, we have

$$
\mathbb{P}\bigg(\Big|\int_{\mathbb{T}^d} :Y_N^2: +2Y_N\Theta_\gamma + \Theta_\gamma^2 dx\Big| \le K_N\bigg) \ge \frac{1}{2} \tag{4.18}
$$

for any $N = M \gg 1$. Then, from the variational formula (Lemma [2.6\)](#page-4-4), [\(4.10\)](#page-11-0) (with Θ^0 replaced with Θ_{γ} , [\(4.16\)](#page-12-2), [\(4.17\)](#page-12-3), and Lemmas [2.7](#page-5-3) and [2.4,](#page-4-1) we have

$$
\log \mathbb{E}_{\mu} \Big[\exp \Big(\min \big(\lambda_N R_N(u), L \big) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : u_N^2 : dx | \le K_N \}} \Big) \Big] \ge \mathbb{E} \Big[\min \big(\lambda_N R_N(Y + \Theta_\gamma), L \big) \cdot \mathbf{1}_{\{ | \int_{\mathbb{T}^d} : (Y_N + \Theta_\gamma)^2 : dx | \le K_N \}} - \frac{1}{2} \int_0^1 \| \theta_\gamma(t) \|_{L_x^2}^2 dt \Big] \qquad (4.19) \ge \lambda_N \gamma^2 K_M^2 M^d - C \gamma K_M M^d - C'(\delta)
$$

for any small $\delta > 0$ and $N = M \gg 1$, provided $L \gg \lambda_N \gamma^2 K_M^2 M^d$. In view of [\(4.13\)](#page-11-5), by setting $M = N$ in [\(4.19\)](#page-12-4), taking $L \to \infty$, and then $N \to \infty$, [\(4.8\)](#page-11-4) follows from (4.19), provided that $\lambda^* = \lambda^*(\gamma)$ is sufficiently large.

It remains to prove [\(4.18\)](#page-12-5). From Lemma [2.7,](#page-5-3) we have

$$
\mathbb{E}\left[\left|\int_{\mathbb{T}^d} :Y_N^2: +2Y_N\Theta_\gamma + (\Theta_\gamma)^2 dx\right|^2\right] \n\lesssim 1 + \mathbb{E}\left[\left|\int_{\mathbb{T}^d} Y_N\Theta_\gamma dx\right|^2\right] + \mathbb{E}\left[\left(\int_{\mathbb{T}^d} \Theta_\gamma^2 dx\right)^2\right]
$$
\n(4.20)

From (4.15) , (2.3) , and (4.4) in Lemma [4.1,](#page-10-2) we have

$$
\mathbb{E}\left[\left|\int_{\mathbb{T}^d} Y_N \Theta_\gamma dx\right|^2\right] = \gamma K_M \mathbb{E}\left[\left|\int_{\mathbb{T}^d} Y_N f_M dx\right|^2\right] = \gamma K_M \sum_{|n| \le M} \langle n \rangle^{-d} |\widehat{f}_M(n)|^2
$$
\n
$$
\lesssim \gamma K_M M^{-d}.
$$
\n(4.21)

From (4.15) and (4.2) , we have

$$
\mathbb{E}\left[\left(\int_{\mathbb{T}^d} \Theta_\gamma^2 dx\right)^2\right] = \gamma^2 K_M^2 \|f_M\|_{L^2}^4 \sim \gamma^2 K_M^2 \tag{4.22}
$$

for any $M \gg 1$. Hence, by applying Chebyshev's inequality, (4.20) , (4.21) , and (4.22) , we obtain

$$
\mathbb{P}\left(\left|\int_{\mathbb{T}^d} (:Y_N^2 : +2Y_N \Theta_\gamma + (\Theta_\gamma)^2) dx\right| \ge K_N\right) \n\lesssim \frac{1}{K_N^2} + \gamma \frac{K_M}{M^d K_N^2} + \gamma^2 \frac{K_M^2}{K_N^2}.
$$
\n(4.23)

Finally, recalling that $K_N \gtrsim \log N \to \infty$ as $N \to \infty$, the bound [\(4.18\)](#page-12-5) follows from setting $M = N$ sufficiently large and choosing $\gamma > 0$ sufficiently small in (4.23). $M = N$ sufficiently large and choosing $\gamma > 0$ sufficiently small in [\(4.23\)](#page-13-21).

Acknowledgements. D.G. and T.O. were supported by the European Research Council (grant no. 864138 "SingStochDispDyn"). T.O. was also supported by the EPSRC Mathematical Sciences Small Grant (grant no. EP/Y033507/1).

REFERENCES

- [1] N. Barashkov, M. Gubinelli, A variational method for Φ_3^4 , Duke Math. J. 169 (2020), no. 17, 3339-3415.
- [2] M. Boué, P. Dupuis, A variational representation for certain functionals of Brownian motion, Ann. Probab. 26 (1998), no. 4, 1641–1659.
- [3] J. Bourgain, *Periodic nonlinear Schrödinger equation and invariant measures*, Comm. Math. Phys. 166 (1994), no. 1, 1–26.
- [4] J. Bourgain, *Invariant measures for the 2D-defocusing nonlinear Schrödinger equation*, Comm. Math. Phys. 176 (1996), no. 2, 421–445.
- [5] D.C. Brydges, G. Slade, *Statistical mechanics of the 2-dimensional focusing nonlinear Schrödinger equation*, Comm.Math. Phys. 182, (1996), 485–504.
- [6] E. Carlen, J. Fröhlich, J. Lebowitz, Exponential relaxation to equilibrium for a one-dimensional focusing non-linear Schrödinger equation with noise, Comm. Math. Phys. 342 (2016), no. 1, 303–332.
- [7] A. Chapouto, G. Li, T. Oh, Deep-water and shallow-water limits of statistical equilibria for the intermediate long wave equation, arXiv:2409.06905 [math.AP].
- [8] G.H. Hardy, J.E. Littlewood, G. Pólya, *Inequalities*, Reprint of the 1952 edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1988. xii+324 pp.
- [9] H. Kuo, Introduction to stochastic integration, Universitext. Springer, New York, 2006. xiv+278 pp.
- [10] J. Lebowitz, H. Rose, E. Speer, *Statistical mechanics of the nonlinear Schrödinger equation*, J. Statist. Phys. 50 (1988), no. 3-4, 657–687.
- [11] E. Nelson, A quartic interaction in two dimensions, 1966 Mathematical Theory of Elementary Particles (Proc. Conf., Dedham, Mass., 1965), pp. 69–73, M.I.T. Press, Cambridge, Mass.
- [12] D. Nualart, The Malliavin calculus and related topics, Second edition. Probability and its Applications (New York). Springer-Verlag, Berlin, 2006. xiv+382 pp.
- [13] T. Oh, M. Okamoto, L. Tolomeo, *Focusing* Φ_3^4 -model with a Hartree-type nonlinearity, to appear in Mem. Amer. Math. Soc.
- [14] T. Oh, M. Okamoto, L. Tolomeo, Stochastic quantization of the Φ_3^3 -model, to appear in Mem. Eur. Math. Soc.
- [15] T. Oh, K. Seong, L. Tolomeo, A remark on Gibbs measures with log-correlated Gaussian fields, Forum Math. Sigma 12 (2024), Paper No. e50.
- [16] T. Oh, P. Sosoe, L. Tolomeo, Optimal integrability threshold for Gibbs measures associated with focusing NLS on the torus, Invent. Math. 227 (2022), no. 3, 1323–1429.
- [17] T. Oh, L. Thomann, A pedestrian approach to the invariant Gibbs measures for the 2-d defocusing nonlinear Schrödinger equations, Stoch. Partial Differ. Equ. Anal. Comput. 6 (2018), 397-445.
- [18] T. Oh, N. Tzvetkov, Quasi-invariant Gaussian measures for the two-dimensional defocusing cubic nonlinear wave equation, J. Eur. Math. Soc. 22 (2020), no. 6, 1785–1826.
- [19] B. Simon, The $P(\varphi)$ ₂ Euclidean (quantum) field theory, Princeton Series in Physics. Princeton University Press, Princeton, N.J., 1974. xx+392 pp.
- [20] L. Tolomeo, H. Weber, Phase transition for invariant measures of the focusing Schrödinger equation, arXiv:2306.07697 [math.AP].
- [21] N. Tzvetkov, *Invariant measures for the defocusing nonlinear Schrödinger equation*, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 7, 2543–2604.
- [22] N. Tzvetkov, Construction of a Gibbs measure associated to the periodic Benjamin-Ono equation, Probab. Theory Related Fields 146 (2010), no. 3-4, 481–514.
- [23] A. Ustünel, Variational calculation of Laplace transforms via entropy on Wiener space and applications, J. Funct. Anal. 267 (2014), no. 8, 3058–3083.
- [24] X. Zhang, A variational representation for random functionals on abstract Wiener spaces, J. Math. Kyoto Univ. 49 (2009), no. 3, 475–490.

Damiano Greco, School of Mathematics, The University of Edinburgh, and The Maxwell Institute for the Mathematical Sciences, James Clerk Maxwell Building, The King's Buildings, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom Email address: dgreco@ed.ac.uk

Tadahiro Oh, School of Mathematics, The University of Edinburgh, and The Maxwell Institute for the Mathematical Sciences, James Clerk Maxwell Building, The King's Buildings, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom, and School of Mathematics and Statistics, Beijing Institute of Technology, Beijing 100081, China

Email address: hiro.oh@ed.ac.uk

Liying Tao, Graduate School of China Academy of Engineering Physics, Beijing, 100088, China Email address: taoliying20@gscaep.ac.cn

Leonardo Tolomeo, School of Mathematics, The University of Edinburgh, and The Maxwell Institute for the Mathematical Sciences, James Clerk Maxwell Building, The King's Buildings, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom

Email address: l.tolomeo@ed.ac.uk