

Bounding the photon mass with gravitationally lensed fast radio bursts

Chen-Ming Chang,^{1,2,*} Jun-Jie Wei,^{1,2,†} Ke-Lai Meng,^{1,‡} Song-Bo Zhang,^{1,3} Hao-Xuan Gao,¹ Jin-Jun Geng,^{1,2} and Xue-Feng Wu^{1,2,§}

¹Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210023, China

²School of Astronomy and Space Sciences, University of Science and Technology of China, Hefei 230026, China

³CSIRO Space and Astronomy, Australia Telescope National Facility, PO Box 76, Epping, NSW 1710, Australia

(Dated: December 16, 2024)

The gravitational time delays of macro-lenses can be used to constrain the rest mass of the photon with high accuracy. Assuming a point-mass + external shear lens model, we prove that an upper limit of the photon mass can be derived directly from two observables—the time delay Δt and the leading-to-trailing flux ratio R of strongly lensed fast radio bursts (FRBs). Using the observed values of Δt and R of a lensed FRB candidate, i.e., FRB 20190308C, as a reference, we obtain a strict upper limit of the photon mass between $m_\gamma < 5.3 \times 10^{-42}$ kg, for a given external shear strength of $\gamma' = 0.01$, and $m_\gamma < 2.1 \times 10^{-41} - 2.4 \times 10^{-42}$ kg, within the external shear range of $0 < \gamma' < 1$. This provides the most stringent limit to date on the photon mass through gravitational lensing time delays, improving by 1 to 2 orders of magnitude the previous results obtained from lensed active galactic nuclei.

I. INTRODUCTION

As one of the fundamental postulates of Maxwell's electromagnetism and Einstein's special relativity, the principle of invariance of light speed implies that the rest mass of the photon should be exactly zero. Nevertheless, there exist some theories involving a finite photon rest mass, such as the famous de Broglie-Proca theory [1, 2], the model of the nonvanishing photon mass as an explanation of dark energy [3], and other new ideas in the Standard-Model Extension with effectively massive photons [4]. Despite the great success of the postulate of the constancy of light speed, those new theories with massive photons are interesting and worthy to explore, whereas the ultimate word on the photon mass (m_γ) stems from empirical facts.

Over the last few decades, various kinds of experimental approaches have been performed to push the empirical boundary on the masslessness of photons (see [5–11] for reviews). These experiments include measurements of the frequency dependence of the speed of light ($m_\gamma \leq 3.8 \times 10^{-51}$ kg; [12–28]), tests of Coulomb's inverse square law ($m_\gamma \leq 1.6 \times 10^{-50}$ kg; [29]), measurement of Jupiter's magnetic field ($m_\gamma \leq 8 \times 10^{-52}$ kg; [30]), analysis of the mechanical stability of magnetized gas in galaxies ($m_\gamma \leq 3 \times 10^{-63}$ kg; [31]), tests of Ampère's law ($m_\gamma \leq 8.4 \times 10^{-48}$ kg; [32]), magnetohydrodynamics of the solar wind ($m_\gamma \leq 1.4 \times 10^{-49} - 3.4 \times 10^{-51}$ kg; [33–35]), Cavendish torsion balance methods ($m_\gamma \leq 1.2 \times 10^{-54}$ kg; [36, 37]), estimates of supermassive black-hole spin ($m_\gamma \leq 7 \times 10^{-56}$ kg; [38]), analysis of pulsar spindown ($m_\gamma \leq 6.3 \times 10^{-53}$ kg; [39]), gravitational deflection of massive photons ($m_\gamma \leq 1.7 \times 10^{-40} - 4.1 \times 10^{-45}$ kg; [40–44]), and so on. Among these experiments on photon mass, the resulting constraints obtained from the gravitational deflection

of light are not the tightest ones; however, in view of model dependence of many experimental methods (see e.g. [6, 9]), tests of the photon mass using different independent methods (such as gravitational deflection) are always interesting and important.

The semi-classical gravity predicts that the deflection of massive photons in an external gravitational field would be energy-dependent [40, 41, 45, 46]. Therefore, an upper bound on the photon mass can be obtained by comparing the difference between the measured deflection angle and the calculated deflection angle for massless photons [40]. Exploiting the gravitational deflection of radio waves by the Sun, Accioly and Paszko [41] obtained an upper limit of $m_\gamma \leq 10^{-43}$ kg. Based on the astrometry of several strong gravitational lensing systems, Qian [42] investigated the photon mass limit at a cosmological scale, yielding $m_\gamma \leq 8.7 \times 10^{-42}$ kg. With the precise astrometry of the gravitationally lensed quasar MG J2016+112, Egorov *et al.* [43] further improved the limit to be $m_\gamma \leq 4.1 \times 10^{-45}$ kg. However, these astrometric limits do not use lens models and simply assume that the angular separation of lensed images is equivalent to the deflection angle of light. Glicenstein [44] argued that this is a strong assumption. Modeling lens galaxies with central supermassive black holes by a singular isothermal model, Glicenstein [44] used the time delays between compact images from three lensed active galactic nuclei (AGNs) to derive a photon mass limit of $m_\gamma \leq 1.7 \times 10^{-40}$ kg.

Fast radio bursts (FRBs) are bright millisecond-long radio flashes originating at cosmological distances [47–51]. Their cosmological origin, energetic nature, and high all-sky rate make them ideal for probing cosmology (e.g., [52]). With tens of thousands of signals that will be guaranteed in the future, FRBs have gained attention as potential targets for lensing studies [53–59]. Very recently, Chang *et al.* [60] employed the autocorrelation algorithm to search for potential lensed FRBs in the first Canadian Hydrogen Intensity Mapping Experiment (CHIME) FRB catalogue, and identified FRB 20190308C as a lensed candidate with a significance of 3.4σ . The information about the time delay and flux ratio between the two substructures of FRB 20190308C can be easily extracted. Inspired by

* changcm@pmo.ac.cn

† jjwei@pmo.ac.cn

‡ mkl@pmo.ac.cn

§ xfwu@pmo.ac.cn

Glicenstein [44], a natural question arises: is it possible to improve the photon mass lensing limits by using the gravitational time delays of lensed FRBs?

In this work, we propose a new method to place an upper limit on the photon mass by applying the time delay information from lensed FRBs. The rest of this paper is arranged as follows. In Section II, we discuss the photon-mass dependence of the time delay in the Chang-Refsdal lens model. The constraints on the photon mass from a lensed FRB candidate are presented in Section III. Finally, a brief summary and discussion are provided in Section IV.

II. PHOTON-MASS DEPENDENCE OF THE TIME DELAY IN THE CHANG-REFSDAL LENS MODEL

The Chang-Refsdal lens model describes the lensing effect of a star, which can be considered as a point-mass lens under the gravitational perturbation of a background galaxy. The lens equation of the Chang-Refsdal lens model is given by [61–65]

$$\begin{aligned}\beta_1 - \theta_1 &= \gamma\theta_1 - \theta_E^2 \frac{\theta_1}{|\theta|^2}, \\ \beta_2 - \theta_2 &= -\gamma\theta_2 - \theta_E^2 \frac{\theta_2}{|\theta|^2},\end{aligned}\quad (1)$$

where β and θ represent the positions in the source and deflector planes, respectively, θ_E is the Einstein angle, and γ is the external shear strength. Note that $\alpha = \beta - \theta$ stands for the deflection angle of light.

Assuming a weak gravitational field, the basic formulas for the time delay and position of lensed images of a massive photon source were derived by Lowenthal [40] and Glicenstein [44]. These studies show that the deflection angle for massive photons is similar as that of massless photons, except for the $(1 + \frac{1}{2}\mu^2)$ multiplicative factor. Here $\mu^2 = \frac{m_\gamma^2 c^2}{P_0^2}$, where m_γ is the rest mass of the photon and P_0 is the time component of the four-momentum. Therefore, for the scenario of massive photons, the lens equation (Equation 1) can be simply rewritten by replacing γ and θ_E^2 with $\gamma' = (1 + \frac{1}{2}\mu^2)\gamma$ and $\theta_E'^2 = (1 + \frac{1}{2}\mu^2)\theta_E^2$. Scaling the angular coordinates with θ_E' : $y' = \beta/\theta_E'$ and $x' = \theta/\theta_E'$, the dimensionless lens equation reads as

$$\begin{aligned}y'_1 &= (1 + \gamma')x'_1 - \frac{x'_1}{x_1'^2 + x_2'^2}, \\ y'_2 &= (1 - \gamma')x'_2 - \frac{x'_2}{x_1'^2 + x_2'^2}.\end{aligned}\quad (2)$$

This lens equation can have up to four solutions, resulting in multiple imaging scenarios. It is difficult to obtain the general analytical solutions to the lens equation, so do the expressions for the time delay Δt and flux ratio R between lensed images. Fortunately, the “permitted region” in the R - Δt space can be determined with the three boundary conditions that the source is on the symmetry axis of the lensing system (i.e., $y'_1 = 0$ or

$y'_2 = 0$) or at the tips of the inner caustics, thereby proving bounds on the photon mass.

Chen *et al.* [64] focused on two-image configurations with $\gamma \ll 1$. For the case of $\gamma \ll 1$, the size of caustic is much less than θ_E' and the cross section of four-image configurations can be ignored. Their analysis can be extended to the case of $\gamma < 1$ if we only consider two-image configurations.

For $y'_1 = 0$, the lower boundary of the permitted region in the R - Δt space can be determined. The solutions for Equation (2) are

$$\begin{aligned}x'_1 &= 0, \\ x'_2 &= \frac{y'_2 \pm \sqrt{y_2'^2 + 4(1 - \gamma')}}{2(1 - \gamma')};\end{aligned}\quad (3)$$

and

$$\begin{aligned}x'_1 &= \pm \sqrt{\frac{1}{1 + \gamma'} - \frac{y_2'^2}{4\gamma'^2}}, \\ x'_2 &= -\frac{y'_2}{2\gamma'}.\end{aligned}\quad (4)$$

Using Equation (3) and the magnification of each image (see Chen *et al.* [64] for the detailed derivation), we obtain the time delay

$$\Delta t = \frac{4GM}{c^3} (1 + z_l) \left[\frac{y_2' s_2'}{2(1 - \gamma')} + \ln \left(\frac{s_2' + y_2'}{s_2' - y_2'} \right) \right], \quad (5)$$

and the flux ratio

$$R = \frac{(y_2'^2 + 2 + y_2' s_2')(y_2' s_2' - 4\gamma') + 8\gamma'^2 + 2\gamma' y_2'^2}{(y_2'^2 + 2 - y_2' s_2')(y_2' s_2' + 4\gamma') - 8\gamma'^2 - 2\gamma' y_2'^2}, \quad (6)$$

where M and z_l are the point mass and redshift of the lens, respectively, and $s_2' = \sqrt{y_2'^2 + 4(1 - \gamma')}$.

For $y'_2 = 0$, the solutions for Equation (2) are

$$\begin{aligned}x'_1 &= \frac{y'_1 \pm \sqrt{y_1'^2 + 4(1 + \gamma')}}{2(1 + \gamma')}, \\ x'_2 &= 0;\end{aligned}\quad (7)$$

and

$$\begin{aligned}x'_1 &= \frac{y'_1}{2\gamma'}, \\ x'_2 &= \pm \sqrt{\frac{1}{1 - \gamma'} - \frac{y_1'^2}{4\gamma'^2}}.\end{aligned}\quad (8)$$

When $y'_2 = 0$, the corresponding formulas for the time delay and flux ratio can be treated as the upper boundary of the permitted region, i.e.,

$$\Delta t = \frac{4GM}{c^3} (1 + z_l) \left[\frac{y_1' s_1'}{2(1 + \gamma')} + \ln \left(\frac{s_1' + y_1'}{s_1' - y_1'} \right) \right], \quad (9)$$

and

$$R = \frac{(y_1'^2 + 2 + y_1' s_1')(y_1' s_1' + 4\gamma') + 8\gamma'^2 - 2\gamma' y_1'^2}{(y_1'^2 + 2 - y_1' s_1')(y_1' s_1' - 4\gamma') - 8\gamma'^2 + 2\gamma' y_1'^2}, \quad (10)$$

where $s_1' = \sqrt{y_1'^2 + 4(1 + \gamma')}$.

When the source is located at the tips of the inner caustics, i.e., $y_1' = 0$ and $y_2' = \pm 2\gamma'/\sqrt{1 + \gamma'}$ (or $y_1' = \pm 2\gamma'/\sqrt{1 - \gamma'}$ and $y_2' = 0$), the left boundary of the permitted region in the R - Δt space can be determined. The lower limit on Δt can be written as

$$\Delta t_{\min} = \frac{4GM}{c^3} (1 + z_l) \left[\frac{2\gamma'}{1 - \gamma'^2} + \ln \left(\frac{1 + \gamma'}{1 - \gamma'} \right) \right]. \quad (11)$$

The corresponding leading-to-trailing flux ratio R is 0 or $+\infty$.

These boundary conditions can still provide some information about the mass of the lens and the photon-mass-dependent time delay between lensed images in the absence of the general analytical solutions. Therefore, they can be used for further study on photon mass limits.

III. PHOTON MASS LIMIT FROM A LENSED FRB CANDIDATE

The permitted region of all possible R - Δt pairs for a point-mass + external shear lens model with the lens mass $M(1 + z_l) = 4277 M_\odot$ and the equivalent external shear strength $\gamma' = 0.01$ is shown in Figure 1. One can see from this plot that all possible R - Δt pairs between two lensed images are actually bracketed by three boundary lines. The lower boundary of the permitted region (solid curve on bottom) corresponds to the R - Δt relation along the y_2' -axis (i.e., $y_1' = 0$), which is determined by Equations (5) and (6). The upper boundary (solid curve on top) corresponds to the R - Δt relation along the y_1' -axis (i.e., $y_2' = 0$), which is determined by Equations (9) and (10). This $y_2' = 0$ curve reaches its minimum at $y_1' = \sqrt{\frac{2\gamma'(1+2\gamma')}{1-\gamma'}}$ with

$$R_{\min} = \frac{(4\gamma'^2 + 2 + \Delta')(\Delta' + 4\gamma' - 4\gamma'^2) + 16\gamma'^4 - 20\gamma'^3 + 4\gamma'^2}{(4\gamma'^2 + 2 - \Delta')(\Delta' - 4\gamma' + 4\gamma'^2) - 16\gamma'^4 + 20\gamma'^3 - 4\gamma'^2}, \quad (12)$$

where $\Delta' = \sqrt{8\gamma'^3 + 20\gamma'^2 + 8\gamma'}$. It is obvious that R_{\min} is larger than 1 when $\gamma' > 0$. The left boundary (vertical dashed line) corresponds to the lower limit of Δt , which is determined by Equation (11). As shown in Figure 1, with the fixed R , the observed time delay Δt_{obs} between the lensed images should always be larger than the lower limit Δt_{\min} (vertical dashed line). With Equation (11), it is thus easy to obtain

$$\frac{4GM}{c^3} (1 + z_l) \cdot 4 \left(1 + \frac{1}{2}\mu^2 \right) \gamma < \Delta t_{\min} < \Delta t_{\text{obs}}. \quad (13)$$




FIG. 1. Distributions of the leading-to-trailing flux ratio R and time delay Δt for a point-mass + external shear lens model. The red shaded area represents the permitted region of all possible R - Δt pairs with the lens mass $M(1 + z_l) = 4277 M_\odot$ and the equivalent external shear strength $\gamma' = 0.01$. The vertical dashed line and two black solid curves correspond to the left, upper, and lower boundaries of all possible R - Δt pairs.

So the photon mass can be constrained as

$$m_\gamma < \frac{P_0}{c} \sqrt{\frac{c^3 \Delta t_{\text{obs}}}{8GM(1 + z_l) \gamma}} - 2. \quad (14)$$

Very recently, Chang *et al.* [60] searched for potential lensed FRBs within the first CHIME/FRB catalogue using the autocorrelation algorithm and verification through signal simulations. Only FRB 20190308C was identified as a plausible candidate for gravitational lensing. The observed time delay and flux ratio between the two substructures of FRB 20190308C are $\Delta t_{\text{obs}} = 8.85$ ms and $R_{\text{obs}} = 0.5$. As an example, we now use the time delay information from FRB 20190308C to demonstrate how to obtain the constraints on the lens mass $M(1 + z_l)$, thereby placing constraints on the photon mass m_γ . For the doubly lensed FRB 20190308C with $R_{\text{obs}} = 0.5$ and $\Delta t_{\text{obs}} = 8.85$ ms, the upper boundary does not provide any useful information because 0.5 will always be smaller than R_{\min} for any γ' . Therefore, only two $M(1 + z_l)$ - γ' relations derived from the lower and left boundaries are seen in Figure 2. The $M(1 + z_l)$ - γ' relation corresponding to the lower boundary (blue line) is derived from Equations (5) and (6). The $M(1 + z_l)$ - γ' relation corresponding to the left boundary (orange line) is derived from Equation (11). For a moderate shear of $\gamma' = 0.01$, the lower limit on the lens mass is about $M(1 + z_l) = 4277 M_\odot$. Since the photon mass term $\mu \ll 1$, it is reasonable to assume that $\gamma \simeq \gamma' = 0.01$. With the observed time delay $\Delta t_{\text{obs}} = 8.85$ ms, the lowest observed frequency $\nu = \frac{P_0 c}{h} = 400$ MHz, and the lower lens mass limit $M(1 + z_l) = 4277 M_\odot$ corresponding to $\gamma' = 0.01$, a stringent upper limit on the photon mass from Equation (14) is

$$m_\gamma < 5.3 \times 10^{-42} \text{ kg} \quad (15)$$




FIG. 2. $M(1+z_l)$ - γ' relations that correspond to the situation of $R_{\text{obs}} = 0.5$ and $\Delta t_{\text{obs}} = 8.85$ ms. The orange curve represents the upper limit of $M(1+z_l)$ obtained with Equation (11). The blue curve represents the lower limit of $M(1+z_l)$ obtained with Equations (5) and (6).

for FRB 20190308C.

In our above analysis, the external shear strength is set to be $\gamma' = 0.01$. To explore the effect of different γ' values, we estimate the sensitivity as we vary γ' . For $\gamma' < 1$, the solutions (i.e., Equation 3) for the dimensionless lens equation always exist when $y'_1 = 0$. Therefore, $|y'_2|$ needs to be greater than $\frac{2\gamma'}{\sqrt{1+\gamma'}}$ to satisfy the two-image condition and ensure that the boundary conditions discussed in Section II remain applicable. As shown in Figure 2, with a fixed γ' , a lower limit on the lens mass limit $M(1+z_l)$ can be obtained, leading to the establishment of an upper limit on the photon mass m_γ . Figure 3 shows that as γ' increases, m_γ decreases first and then increases. The upper limit of the photon mass has a minimum value of $m_\gamma = 2.4 \times 10^{-42}$ kg, corresponding to $\gamma' = 0.38$ and a minimum lens mass of $M(1+z_l) = 221.7 M_\odot$. The maximum value of the upper photon mass limit is difficult to determine because, as γ' approaches 1, the wave properties of light become significant, making the geometric approximation of the lensing equation invalid. Therefore, we calculate the upper limit of the photon mass to be $m_\gamma < 2.1 \times 10^{-41}$ kg when $\gamma' = 0.99$, corresponding to a minimum lens mass of $M(1+z_l) = 4.3 M_\odot$. That is, within the range of $0 < \gamma' < 1$, the photon mass can be constrained to be

$$m_\gamma < 2.1 \times 10^{-41} - 2.4 \times 10^{-42} \text{ kg} \quad (16)$$

for FRB 20190308C, which is almost 10-100 times tighter than the constraints from the time delays of lensed AGNs [44].

For $\gamma' > 1$, there are two caustics, each with one cusp on the y'_2 axis and two cusps off-axis. The boundary conditions for the two-image scenario are not clear. When $y'_1 = 0$, the two-image scenario has two possibilities: either $|y'_2| \geq \frac{2\gamma'}{\sqrt{1+\gamma'}}$ or $|y'_2| \leq 2\sqrt{\gamma'-1}$. For $y'_1 = 0$ and $|y'_2| \geq \frac{2\gamma'}{\sqrt{1+\gamma'}}$, the time




FIG. 3. Sensitivity of the upper photon mass limit m_γ to the equivalent external shear strength γ' .

delay and flux ratio become:

$$\Delta t = \frac{4GM}{c^3} (1+z_l) \left[\frac{y'_2 s'_2}{2(\gamma'-1)} - \ln \left(\frac{y'_2 + s'_2}{y'_2 - s'_2} \right) \right], \quad (17)$$

and

$$R = \frac{-(y'^2_2 + 2 - y'_2 s'_2)(y'_2 s'_2 + 4\gamma') + 8\gamma'^2 + 2\gamma' y'^2_2}{(y'^2_2 + 2 + y'_2 s'_2)(y'_2 s'_2 - 4\gamma') + 8\gamma'^2 + 2\gamma' y'^2_2}. \quad (18)$$

For $y'_1 = 0$ and $|y'_2| \leq 2\sqrt{\gamma'-1}$, $\Delta t = 0$ and $R = 1$. Numerical calculations suggest that it may still be possible to impose certain constraints on the combined lens mass $M(1+z_l)$ using Equations (17) and (18) by adding some restrictions, such as limiting the leading-to-trailing flux ratio $R < 1$. However, the information obtained from the dynamic spectrum may not be sufficient to constrain the photon mass m_γ due to the lack of an analytical relationship between $M(1+z_l)$ and m_γ .

IV. SUMMARY AND DISCUSSION

It has been suggested that the strong lensing effect of a point mass + external shear lens model on a single-peak FRB can produce double peaks (i.e., lensed images). Based on this lens model, here we proposed a method of using the two observables of the time delay Δt and the leading-to-trailing flux ratio R from lensed FRBs to set a stringent upper limit on the photon mass. In particular, we showed the process of constraining photon mass using the observed values of Δt and R from a lensed FRB candidate, i.e., FRB 20190308C, as a reference.

For a point mass + external shear lens model, there is no one-to-one correspondence between the upper photon mass limit m_γ and Δt and R due to the extra freedom of the external shear. Nevertheless, we showed that an upper limit on m_γ can still be derived from Δt and R for a given external shear strength of γ' (Section III). For FRB 20190308C with $\Delta t =$

8.85 ms and $R = 0.5$, we obtained a strict constraint on the photon mass $m_\gamma < 5.3 \times 10^{-42}$ kg for a fixed external shear strength of $\gamma' = 0.01$. We also inspected the influences of different γ' values, finding that this effect has a modest impact on the photon mass limits. That is, within the range of $0 < \gamma' < 1$, one can derive $m_\gamma < 2.1 \times 10^{-41} - 2.4 \times 10^{-42}$ kg.

Previously, by analyzing the gravitational time delays from lensed AGNs, Glicenstein [44] set a severe limit on the photon mass of $m_\gamma \leq 1.7 \times 10^{-40}$ kg. In the present Letter, using the sharp features of the lensed FRB signals, we have obtained the most stringent limit to date on the photon mass through gravitational lensing time delays, namely $\sim 2.1 \times 10^{-41} - 2.4 \times 10^{-42}$ kg, which represents an improvement of 1 to 2 orders of magnitude over the results previously obtained from lensed AGNs.

So far, only a lensed FRB candidate with a significance of 3.4σ has been identified [60]. Nevertheless, given the high all-sky event rate and sustained efforts in FRB searches, more FRB signals lensed by point-mass lenses with higher significance are expected to be identified in the near future. Due to

the short-lived nature and unpredictability of FRBs, it may be hard to perform a full lens modeling with the observed data. The method presented in this work offers an alternative, straightforward way of constraining the photon mass from easily obtained observables of Δt and R .

ACKNOWLEDGMENTS

This work is supported by the National SKA Program of China (2022SKA0130100), the National Natural Science Foundation of China (grant Nos. 12422307, 12373053, 12321003, and 12041306), the Key Research Program of Frontier Sciences (grant No. ZDBS-LY-7014) of Chinese Academy of Sciences, International Partnership Program of Chinese Academy of Sciences for Grand Challenges (114332KYBS20210018), the CAS Project for Young Scientists in Basic Research (grant No. YSBR-063), and the Natural Science Foundation of Jiangsu Province (grant No. BK20221562).

-
- [1] L. De Broglie, *J. Phys. Radium* **3**, 422 (1922).
 [2] A. Proca, *J. Phys. Radium* **7** 7, 347 (1936).
 [3] S. Kouwn, P. Oh, and C.-G. Park, *Phys. Rev. D* **93**, 083012 (2016), arXiv:1512.00541 [astro-ph.CO].
 [4] A. D. A. M. Spallicci, J. A. Helayël-Neto, M. López-Corredoira, and S. Capozziello, *European Physical Journal C* **81**, 4 (2021), arXiv:2011.12608 [astro-ph.CO].
 [5] A. S. Goldhaber and M. M. Nieto, *Reviews of Modern Physics* **43**, 277 (1971).
 [6] L.-C. Tu, J. Luo, and G. T. Gillies, *Reports on Progress in Physics* **68**, 77 (2005).
 [7] Y. Z. Zhang, “Special Relativity And Its Experimental Foundation,” Special Relativity And Its Experimental Foundation. Series: Advanced Series on Theoretical Physical Science, ISBN: 978-981-02-2749-4. WORLD SCIENTIFIC, Edited by Yuan Zhong Zhang, vol. 4 (1997).
 [8] L. B. Okun, *Acta Physica Polonica B* **37**, 565 (2006), arXiv:hep-ph/0602036 [hep-ph].
 [9] A. S. Goldhaber and M. M. Nieto, *Reviews of Modern Physics* **82**, 939 (2010), arXiv:0809.1003 [hep-ph].
 [10] G. Spavieri, J. Quintero, G. T. Gillies, and M. Rodríguez, *European Physical Journal D* **61**, 531 (2011).
 [11] J.-J. Wei and X.-F. Wu, *Frontiers of Physics* **16**, 44300 (2021), arXiv:2102.03724 [astro-ph.HE].
 [12] B. Lovell, F. L. Whipple, and L. H. Solomon, *Nature (London)* **202**, 377 (1964).
 [13] B. Warner and R. E. Nather, *Nature (London)* **222**, 157 (1969).
 [14] B. E. Schaefer, *Phys. Rev. Lett.* **82**, 4964 (1999), arXiv:astro-ph/9810479 [astro-ph].
 [15] X.-F. Wu, *et al.*, *Astrophys. J. Lett.* **822**, L15 (2016), arXiv:1602.07835 [astro-ph.HE].
 [16] L. Bonetti, *et al.*, *Physics Letters B* **757**, 548 (2016), arXiv:1602.09135 [astro-ph.HE].
 [17] L. Bonetti, *et al.*, *Physics Letters B* **768**, 326 (2017), arXiv:1701.03097 [astro-ph.HE].
 [18] B. Zhang, Y.-T. Chai, Y.-C. Zou, and X.-F. Wu, *Journal of High Energy Astrophysics* **11**, 20 (2016), arXiv:1607.03225 [astro-ph.HE].
 [19] L. Shao and B. Zhang, *Phys. Rev. D* **95**, 123010 (2017), arXiv:1705.01278 [hep-ph].
 [20] J.-J. Wei, E.-K. Zhang, S.-B. Zhang, and X.-F. Wu, *Research in Astronomy and Astrophysics* **17**, 13 (2017), arXiv:1608.07675 [astro-ph.HE].
 [21] J.-J. Wei and X.-F. Wu, *JCAP* **2018**, 045 (2018), arXiv:1803.07298 [astro-ph.HE].
 [22] J.-J. Wei and X.-F. Wu, *Research in Astronomy and Astrophysics* **20**, 206 (2020), arXiv:2006.09680 [astro-ph.HE].
 [23] N. Xing, *et al.*, *Astrophys. J. Lett.* **882**, L13 (2019), arXiv:1907.00583 [astro-ph.HE].
 [24] H. Wang, X. Miao, and L. Shao, *Physics Letters B* **820**, 136596 (2021), arXiv:2103.15299 [astro-ph.HE].
 [25] C.-M. Chang, J.-J. Wei, S.-b. Zhang, and X.-F. Wu, *JCAP* **2023**, 010 (2023), arXiv:2207.00950 [astro-ph.HE].
 [26] H.-N. Lin, L. Tang, and R. Zou, *Mon. Not. R. Astron. Soc.* **520**, 1324 (2023), arXiv:2301.12103 [gr-qc].
 [27] B. Wang, J.-J. Wei, X.-F. Wu, and M. López-Corredoira, *JCAP* **2023**, 025 (2023), arXiv:2304.14784 [astro-ph.HE].
 [28] J.-Y. Ran, B. Wang, and J.-J. Wei, *Chinese Physics Letters* **41**, 059501 (2024), arXiv:2404.17154 [astro-ph.CO].
 [29] E. R. Williams, J. E. Faller, and H. A. Hill, *Phys. Rev. Lett.* **26**, 721 (1971).
 [30] J. Davis, L., A. S. Goldhaber, and M. M. Nieto, *Phys. Rev. Lett.* **35**, 1402 (1975).
 [31] G. V. Chibisov, *Soviet Physics Uspekhi* **19**, 624 (1976).
 [32] M. A. Chernikov, C. J. Gerber, H. R. Ott, and H. J. Gerber, *Phys. Rev. Lett.* **68**, 3383 (1992).
 [33] D. D. Ryutov, *Plasma Physics and Controlled Fusion* **39**, A73 (1997).
 [34] D. D. Ryutov, *Plasma Physics and Controlled Fusion* **49**, B429 (2007).
 [35] A. Retinò, A. D. A. M. Spallicci, and A. Vaivads, *Astroparticle Physics* **82**, 49 (2016), arXiv:1302.6168 [hep-ph].
 [36] R. Lakes, *Phys. Rev. Lett.* **80**, 1826 (1998).
 [37] J. Luo, L.-C. Tu, Z.-K. Hu, and E.-J. Luan, *Phys. Rev. Lett.* **90**,

- 081801 (2003).
- [38] P. Pani, V. Cardoso, L. Gualtieri, E. Berti, and A. Ishibashi, *Phys. Rev. Lett.* **109**, 131102 (2012), arXiv:1209.0465 [gr-qc].
- [39] Y.-P. Yang and B. Zhang, *Astrophys. J.* **842**, 23 (2017), arXiv:1701.03034 [astro-ph.HE].
- [40] D. D. Lowenthal, *Phys. Rev. D* **8**, 2349 (1973).
- [41] A. Accioly and R. Paszko, *Phys. Rev. D* **69**, 107501 (2004).
- [42] L. Qian, *Science China Physics, Mechanics, and Astronomy* **55**, 523 (2012), arXiv:1406.1592 [astro-ph.CO].
- [43] P. Egorov, M. Guzinin, H. Hakobyan, and S. Troitsky, *Mon. Not. R. Astron. Soc.* **437**, L90 (2014), arXiv:1306.2716 [astro-ph.CO].
- [44] J. F. Glicenstein, *Astrophys. J.* **850**, 102 (2017), arXiv:1710.11587 [astro-ph.HE].
- [45] A. Accioly, A. Azeredo, H. Mukai, and E. D. R. Neto, *Progress of Theoretical Physics* **104**, 103 (2000).
- [46] A. Accioly and S. Ragusa, *Classical and Quantum Gravity* **19**, 5429 (2002).
- [47] D. R. Lorimer, M. Bailes, M. A. McLaughlin, D. J. Narkevic, and F. Crawford, *Science* **318**, 777 (2007), arXiv:0709.4301 [astro-ph].
- [48] J. M. Cordes and S. Chatterjee, *Ann. Rev. Astron. Astrophys.* **57**, 417 (2019), arXiv:1906.05878 [astro-ph.HE].
- [49] E. Petroff, J. W. T. Hessels, and D. R. Lorimer, *Astron. Astrophys. Rev.* **27**, 4 (2019), arXiv:1904.07947 [astro-ph.HE].
- [50] E. Petroff, J. W. T. Hessels, and D. R. Lorimer, *Astron. Astrophys. Rev.* **30**, 2 (2022), arXiv:2107.10113 [astro-ph.HE].
- [51] B. Zhang, *Reviews of Modern Physics* **95**, 035005 (2023), arXiv:2212.03972 [astro-ph.HE].
- [52] W. Deng and B. Zhang, *Astrophys. J. Lett.* **783**, L35 (2014), arXiv:1401.0059 [astro-ph.HE].
- [53] J. B. Muñoz, E. D. Kovetz, L. Dai, and M. Kamionkowski, *Phys. Rev. Lett.* **117**, 091301 (2016), arXiv:1605.00008 [astro-ph.CO].
- [54] Y. K. Wang and F. Y. Wang, *Astron. & Astrophys.* **614**, A50 (2018), arXiv:1801.07360 [astro-ph.CO].
- [55] K. Liao, S. B. Zhang, Z. Li, and H. Gao, *Astrophys. J. Lett.* **896**, L11 (2020), arXiv:2003.13349 [astro-ph.CO].
- [56] M. W. Sammons, *et al.*, *Astrophys. J.* **900**, 122 (2020), arXiv:2002.12533 [astro-ph.CO].
- [57] K. Krochek and E. D. Kovetz, *Phys. Rev. D* **105**, 103528 (2022), arXiv:2112.03721 [astro-ph.CO].
- [58] C. Leung, *et al.*, *Phys. Rev. D* **106**, 043017 (2022), arXiv:2204.06001 [astro-ph.HE].
- [59] H. Zhou, *et al.*, *Astrophys. J.* **928**, 124 (2022), arXiv:2109.09251 [astro-ph.CO].
- [60] C. Chang, *et al.*, arXiv e-prints, arXiv:2406.19654 (2024), arXiv:2406.19654 [astro-ph.HE].
- [61] K. Chang and S. Refsdal, *Nature (London)* **282**, 561 (1979).
- [62] K. Chang and S. Refsdal, *Astron. & Astrophys.* **132**, 168 (1984).
- [63] J. H. An and N. W. Evans, *Mon. Not. R. Astron. Soc.* **369**, 317 (2006), arXiv:astro-ph/0601457 [astro-ph].
- [64] X. Chen, Y. Shu, W. Zheng, and G. Li, *Astrophys. J.* **912**, 134 (2021), arXiv:2105.05868 [astro-ph.GA].
- [65] H.-X. Gao, *et al.*, *Mon. Not. R. Astron. Soc.* **516**, 453 (2022), arXiv:2204.03823 [astro-ph.HE].