Sharp L^1 -convergence rates to the Barenblatt solutions for the compressible Euler equations with time-varying damping *

Jun-Ren Luo^a, Ti-Jun Xiao^{b†}

^a College of Science, University of Shanghai for Science and Technology, Shanghai 200093, China

^b Shanghai Key Laboratory for Contemporary Applied Mathematics

School of Mathematical Sciences, Fudan University, Shanghai 200433, China

Abstract: We study the asymptotic behavior of compressible isentropic flow when the initial mass is finite and the friction varies with time, which is modeled by the compressible Euler equation with time-dependent damping. In this paper, we obtain the best L^1 -convergence rates to date, for any $\gamma \in (1, +\infty)$ and $\nu \in [0, 1)$. Here, γ is the adiabatic gas exponent, and ν is the physical parameter in the damping term. The key to the analysis lies in a new perspective on the relationship between the density function and the Barenblatt solution of the porous medium equation, and finding the relevant lower bound for the case of $\gamma < 2$ is a tricky problem. Specialized to $\nu = 0$, these convergence rates also show an essential improvement over the original rates. Moreover, for all $\gamma \in (1, +\infty)$, the results in this work are the first to present a unified form of L^1 -convergence rates. Indeed, even for $\nu = 0$, as noted in 2011, "the current rate is difficult to improve with the current method". Our results are therefore an encouraging advancement.

Keywords: Convergence rates; Compressible Euler equations; Barenblatt solutions; Time-dependent damping.

2020 AMS Subject Classification: 35L65; 76S05; 35K65

^{*}The work was supported partly by the NSF of China (12171094, 11831011), and the Shanghai Key Laboratory for Contemporary Applied Mathematics (08DZ2271900).

[†]Corresponding author. E-mail: tjxiao@fudan.edu.cn

1 Introduction

In this paper, we are concerned with the compressible Euler equations with frictional time-dependent damping in Euler coordinates as follows

$$\begin{cases} \rho_t + (\rho u)_x = 0, \\ (\rho u)_t + (\rho u^2 + p(\rho))_x = -\frac{\alpha}{(1+t)^{\nu}}\rho u, \end{cases}$$
(1.1)

with finite initial mass

$$(\rho, u)(x, 0) = (\rho_0, u_0)(x), \quad \rho_0(x) \ge 0, \quad \int_{-\infty}^{+\infty} \rho_0(x) dx = M > 0,$$
 (1.2)

where ρ , u, and p, denote the density, velocity, and pressure respectively. Since we study a polytropic perfect gas, the state equation of the fluid is given by

$$p(\rho) = \kappa \rho^{\gamma},$$

with $\kappa = \frac{(\gamma-1)^2}{4\gamma}$, and $\gamma > 1$ is the adiabatic gas exponent. The damping term $-\frac{\alpha}{(1+t)^{\nu}}\rho u$ with physical parameter $0 \le \nu < 1$ has the time-dependent frictional effect, and $\alpha = \kappa = \frac{(\gamma-1)^2}{4\gamma}$, which will simplify the form of the entropy functions used below (cf. [14, 17]). Also, we use momentum $m =: \rho u$ in what follows for simplicity.

When $\nu = 0$, the system (1.1) is equivalent to the following decoupled system if taking time asymptotically,

$$\begin{cases} \bar{\rho}_t = (\bar{\rho}^{\gamma})_{xx}, \\ \bar{m} = -(\bar{\rho}^{\gamma})_x, \end{cases}$$
(1.3)

where $(1.3)_1$ is the famous porous medium equation(PME), and $(1.3)_2$ is the Darcy's law, that is, the inertial terms in the momentum equation decay to zero faster than the other terms caused by the damping effect of the frictional force term (see Hsiao and Liu [10]). Later, in the pioneering work of [21], Nishihara improved the decay rates by utilizing detailed energy estimates when the initial data are small perturbation around $\bar{\rho}$ and thus away from vacuum. For more works with small initial data, we refer to [22, 25, 31]. By using the vanishing viscosity method, Zhu [32] studied the asymptotic behavior of weak entropy solution. As for the properties of the solutions near the vacuum, we refer the reader to [15, 18, 19]. Then, Huang, Marcati and Pan [13] obtained the decay rates of the weak entropy solutions of the system (1.1) to the corresponding Barenblatt solutions (1.3) in L^2 or L^{γ} space for any $\gamma > 1$. Later, in [14], Huang, Pan and Wang studied the asymptotic behavior in L^1 space with $1 < \gamma < 3$ and obtained

$$\|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} \le C(1+t)^{-\frac{1}{4(\gamma+1)}+\varepsilon}.$$

Then, the convergence rates were improved to

$$\|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} \le C(1+t)^{-\frac{1}{(\gamma+1)^2}+\varepsilon},$$

for $\gamma \geq 2$ in [8]. For Euler equation or Euler-Poisson equation with friction, we refer to [1, 7]. For compressible Euler equations with physical vacuum free boundaries, see [29, 30],

When $0 < \nu < 1$, the system (1.1) becomes the compressible Euler equations with degenerate linear time-dependent damping. In view of Darcy's law, the solutions of (1.1) should be time-asymptotically equivalent to the Barenblatt solutions of the following system

$$\begin{cases} \bar{\rho}_t = (1+t)^{\nu} \, (\bar{\rho}^{\gamma})_{xx} \,, \\ \bar{m} = -(1+t)^{\nu} \, (\bar{\rho}^{\gamma})_x \,, \end{cases}$$
(1.4)

where $(1.4)_1$ is the PME with time-dependent diffusion and $(1.4)_2$ is the Dracy's law. Recently, there have been various works for the compressible Euler equations with timedependent damping in the literature; see [5, 12, 27, 33]. Pan [23, 24] proved that $\nu =$ $1, \alpha = 2$ is the critical threshold to separate the global existence and non-existence of C^1 solutions in one dimension. For more blow-up phenomena, we refer to [3, 11, 26]. Also, Cui-Yin-Zhang-Zhu [4] and Li-Li-Mei-Zhang [16] considered independently the convergence of smooth solutions with $0 \leq \nu < 1$ and $\alpha > 0$.

In [9], for $1 < \gamma < 3$, the authors proved that any L^{∞} weak entropy solution of problem (1.1)-(1.2) converges to the Barenblatt's solution of equation (1.4) with the same mass in L^1 , and at the rate

$$||(\rho - \bar{\rho}(\cdot, t))||_{L^1} \le C(1+t)^{-\alpha(\varepsilon)},$$

with

$$\alpha(\varepsilon) = \begin{cases} \frac{\nu+1}{4(\gamma+1)} - \varepsilon, & \nu \in \left(0, \frac{\gamma}{\gamma+2}\right], \\ \frac{1-\nu}{4} - \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma+2}, 1\right). \end{cases}$$

At the same time, the authors in [6] obtained

$$||(\rho - \bar{\rho}(\cdot, t))||_{L^1} \le C(1+t)^{k(\epsilon)},$$

with

$$k(\epsilon) = \begin{cases} -\frac{\gamma^2 - \gamma - 1 - \nu \left(\gamma^2 + \gamma - 1\right)}{(\gamma + 1)(2\gamma - 1)} + \epsilon, & \text{if } \frac{1 + \sqrt{5}}{2} < \gamma \le 2, \quad 0 \le \nu < \frac{\gamma^2 - \gamma - 1}{\gamma^2 + \gamma - 1}, \\ -\frac{1 - \nu (2\gamma + 1)}{(\gamma + 1)^2} + \epsilon, & \text{if } \gamma \ge 2, \qquad 0 \le \nu < \frac{1}{2\gamma + 1}. \end{cases}$$

In this paper, we intend to study the asymptotic behavior of L^{∞} weak entropy solutions for the compressible Euler equations with time-dependent damping. By using elaborate energy estimates and relative weak entropies, we will obtain better convergence results for weak entropy solutions of (1.1) to the corresponding Barenblatt solutions of (1.4) with the same finite initial mass. We will establish L^{γ} estimate for any $\gamma \geq 2$ and $0 \leq \nu < 1$ and L^1 estimate for any $\gamma > 1$ and $0 \leq \nu < 1$, which are better compared with those in [6, 8, 9, 14]. Furthermore, it is worth noting that our results present for the first time a uniform form of L^1 estimate over the entire $(1, +\infty)$ range of γ . Indeed, even for $\nu = 0$ it is not easy to make some improvements, and as pointed out in [14], "the current rate is difficult to improve with the current method". Our results are therefore an encouraging advancement.

The outline of the paper is as follows. In section 2, we give a quick review of some information on Barenblatt's solution, and state our main result, explaining the basic proof ideas. Then, in section 3, we recall or derive some important lemmas. Section 4 is devoted to extending the related results of [9] about some key intermediate estimates. Finally, we prove our main result in section 5.

2 Preliminaries and main result

Throughout this paper, $\|\cdot\|_{L^p}$ stands for $L^p(\mathbf{R})(1 \le p \le +\infty)$. In particular, we use $\|\cdot\|$ instead of $\|\cdot\|_2$ and denote $\int_{-\infty}^{+\infty} f(x,t)dx = \int f(x,t)dx$.

Since we study the large time behavior of global solutions to the system (1.1) with vacuum, it is suitable to consider L^{∞} weak entropy solutions. Next, we state the definition of L^{∞} weak entropy solutions for the system (1.1)-(1.2) (cf. [6, 9]).

Definition 2.1. The functions $(\rho, m)(x, t) \in L^{\infty}$ are called an entropy solution of (1.1)-(1.2), if for any nonnegative test function $\phi \in \mathcal{D}(\mathbf{R} \times \mathbf{R}_+)$, it holds that

$$\begin{cases} \int_0^t \int_{-\infty}^{+\infty} \left(\rho\phi_t + m\phi_x\right) dx ds + \int_{-\infty}^{+\infty} \rho_0(x)\phi(x,0) dx = 0,\\ \int_0^t \int_{-\infty}^{+\infty} \left\{m\phi_t + \left(\frac{m^2}{\rho} + \rho^\gamma\right)\phi_x - \frac{m}{(1+s)^\lambda}\phi\right\} dx ds + \int_{-\infty}^{+\infty} m_0(x)\phi(x,0) dx = 0,\\ \int_0^t \int_{-\infty}^{+\infty} \left(\eta\phi_t + q\phi_x - \frac{m}{(1+s)^\lambda}\eta\phi\right) dx ds + \int_{-\infty}^{+\infty} \eta(x,0)\phi(x,0) dx \ge 0, \end{cases}$$

where (η, q) is any weak convex entropy-flux pair $(\eta(\rho, m), q(\rho, m))$ satisfying

$$\nabla q = \nabla \eta \nabla f, \ f = \left(m, \frac{m^2}{\rho} + \kappa \rho^\gamma\right)^t, \ \eta(0, 0) = 0.$$

Next, we recall some results on the entropies available for (1.1). According to [17], all

weak entropies of (1.1) are given by the following formula

$$\eta(\rho,m) = \int g(\xi)\chi(\xi;\rho,u)\mathrm{d}\xi = \rho \int_{-1}^{1} g\left(u+z\rho^{\theta}\right) \left(1-z^{2}\right)^{\lambda}\mathrm{d}z,$$

$$q(\rho,m) = \int g(\xi)(\theta\xi+(1-\theta)u)\chi(\xi;\rho,u)\mathrm{d}\xi$$

$$= \rho \int_{-1}^{1} g\left(u+z\rho^{\theta}\right) \left(u+\theta z\rho^{\theta}\right) \left(1-z^{2}\right)^{\lambda}\mathrm{d}z,$$
(2.1)

where $m = \rho u, \theta = \frac{\gamma - 1}{2}, \lambda = \frac{3 - \gamma}{2(\gamma - 1)}, g(\xi)$ is any smooth function of ξ , and

$$\chi(\xi;\rho,u) = \left(\rho^{\gamma-1} - (\xi-u)^2\right)_+^{\lambda}.$$

The two common choices of $g(\xi)$ are $g(\xi) = \frac{1}{2}|\xi|^2$ and $g(\xi) = |\xi|^{\frac{2\gamma}{\gamma-1}}$.

Then, we give some background information on the Barenblatt solutions and consider

$$\begin{cases} \bar{\rho}_t = (1+t)^{\nu} \left(\bar{\rho}^{\gamma} \right)_{xx}, \\ \bar{\rho}(x,-1) = M \delta(x), \quad M > 0, \end{cases}$$

$$(2.2)$$

with $0 \le \nu < 1$ and $\gamma > 1$. It is shown in [6, 9] that (2.2) admits a unique solution given by

$$\bar{\rho}(x,t) = (1+t)^{-\frac{1+\nu}{\gamma+1}} \left(A_0 - B_0 x^2 (1+t)^{-\frac{2(1+\nu)}{\gamma+1}} \right)_+^{\frac{1}{\gamma-1}}, \qquad (2.3)$$

where $(f)_{+} = \max\{0, f\},\$

$$B_0 = \frac{(\gamma - 1)(1 + \nu)}{2\gamma(\gamma + 1)}, \quad \text{and} \quad A_0^{\frac{\gamma + 1}{2(\gamma - 1)}} = M\sqrt{B_0} \left(\int_{-1}^1 \left(1 - y^2\right)^{1/(\gamma - 1)} dy\right)^{-1}.$$

As a result, the velocity is given by

$$\bar{u}(x,t) = \frac{1+\nu}{(\gamma+1)(1+t)}x, \ |x| < \sqrt{\frac{A_0}{B_0}}(1+t)^{\frac{1+\nu}{\gamma+1}}.$$
(2.4)

Now, we state our main result.

Theorem 2.2. Suppose that $\rho_0(x) \in L^1(\mathbf{R}) \cap L^{\infty}(\mathbf{R}), u_0(x) \in L^{\infty}(\mathbf{R})$ and

$$M = \int_{-\infty}^{+\infty} \rho_0(x) dx > 0, \quad \rho_0(x) \ge 0.$$

Let $1 < \gamma < +\infty$, $0 \le \nu < 1$, and (ρ, m) be an L^{∞} entropy solution of the Cauchy problem (1.1)-(1.2). Let $\bar{\rho}$ be the Barenblatt's solution of porous medium equation (1.3) with mass M and $\bar{m} = -(\bar{\rho}^{\gamma})_x$. Define

$$y(x,t) = -\int_{-\infty}^{x} (\rho - \bar{\rho})(r,t) dr,$$

and let $y(x,0) \in L^2(\mathbf{R})$. If $2 \leq \gamma < +\infty$, then for any $\varepsilon > 0$ and t > 0,

$$\|(\rho - \bar{\rho})(\cdot, t)\|_{L^{\gamma}}^{\gamma} \le C_{\varepsilon}(1+t)^{-\mu(\varepsilon)}, \qquad (2.5)$$

where

$$\mu(\varepsilon) := \begin{cases} \frac{\gamma^2 + \gamma - 1}{(\gamma + 1)^2} (1 + \nu) + \varepsilon, & \nu \in \left[0, \frac{\gamma}{\gamma + 2}\right], \\ \frac{2\gamma - 1 - \nu}{\gamma + 1} + \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma + 2}, 1\right). \end{cases}$$
(2.6)

Furthermore, for any $\gamma > 1$, it holds that

$$\|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} \le C_{\varepsilon} (1+t)^{-k(\varepsilon)}, \qquad (2.7)$$

with

$$k(\varepsilon) := \begin{cases} \frac{\gamma}{2(\gamma+1)^2}(1+\nu) - \varepsilon, & \nu \in \left[0, \frac{\gamma}{\gamma+2}\right], \\ \frac{\gamma}{2(\gamma+1)}(1-\nu) - \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma+2}, 1\right]. \end{cases}$$
(2.8)

Remark 2.3. It is shown in [6] that

$$||(\rho - \bar{\rho}(\cdot, t))||_{L^1} \le C(1+t)^{k_1(\epsilon)},$$

with

$$k_1(\epsilon) := \begin{cases} -\frac{\gamma^2 - \gamma - 1 - \nu \left(\gamma^2 + \gamma - 1\right)}{(\gamma + 1)(2\gamma - 1)} + \epsilon, & \text{if } \frac{1 + \sqrt{5}}{2} < \gamma \le 2, & 0 \le \nu < \frac{\gamma^2 - \gamma - 1}{\gamma^2 + \gamma - 1}, \\ -\frac{1 - \nu (2\gamma + 1)}{(\gamma + 1)^2} + \epsilon, & \text{if } \gamma \ge 2, & 0 \le \nu < \frac{1}{2\gamma + 1}, \end{cases}$$

and in [9] for $1 < \gamma < 3$ that

$$||(\rho - \bar{\rho}(\cdot, t))||_{L^1} \le C(1+t)^{-\alpha_1(\varepsilon)},$$

with

$$\alpha_1(\varepsilon) = \begin{cases} \frac{\nu+1}{4(\gamma+1)} - \varepsilon, & \nu \in \left(0, \frac{\gamma}{\gamma+2}\right], \\ \frac{1-\nu}{4} - \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma+2}, 1\right). \end{cases}$$

It is easy to check that $\alpha_1(\varepsilon) < k_1(\varepsilon)$,

$$\frac{\gamma^2 - \gamma - 1}{\gamma^2 + \gamma - 1} < \frac{\gamma}{\gamma + 2}$$
, and $\frac{1}{2\gamma + 1} < \frac{\gamma}{\gamma + 2}$,

which gives

$$k_1(\epsilon) < \frac{\gamma}{2(\gamma+1)^2}(1+\nu) + \varepsilon$$

This means that the L^1 convergence rates in Theorem 2.2 are better. Furthermore, it is worth mentioning that the L^1 convergence estimate in Theorem 2.2 is valid for any $1 < \gamma < +\infty$. **Remark 2.4.** When $\nu = 0$, it is shown in [8] for $\gamma \ge 2$ and [14] for $1 < \gamma < 3$ that $\|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} \le C(1+t)^{-\frac{1}{(\gamma+1)^2}+\varepsilon}$ and $\|(\rho - \bar{\rho})(\cdot, t)\|_{L^1} \le C(1+t)^{-\frac{1}{4(\gamma+1)}+\varepsilon}$ respectively. It is easy to check that $\frac{\gamma}{2(\gamma+1)^2} > \frac{1}{4(\gamma+1)}$ when $\gamma > 1$ and $\frac{\gamma}{2(\gamma+1)^2} > \frac{1}{(\gamma+1)^2}$ when $\gamma \ge 2$, which means that the decay rates in Theorem 2.2 are better than the previous ones.

Now, we explain the basic proof ideas. We will follow the strategy in [6, 9], employing the entropy inequalities and time-weighted entropy estimates with specific index. An important part of the proof process is to extend key intermediate estimates from [9, Lemmas 3.8 and 3.9] (with $1 < \gamma < 3$) to $1 < \gamma < +\infty$ (Theorem 4.6). This will be realized by Taylor's formula, the energy inequality and in-depth analysis of η (defined in Lemma 4.1).

For $\gamma \geq 2$, we choose the mechanical energy(see [6, 9]),

$$\eta = \frac{1}{2}\frac{m^2}{\rho} + \frac{\kappa}{\gamma - 1}\rho^{\gamma},$$

which measures the L^{γ} norm while the entropy inequality measures the $L^{\gamma+1}$ norm in density. To overcome the mismatch between the exponents, we find the relationship between

$$\rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma+1)\bar{\rho}^{\gamma}(\rho-\bar{\rho}) \text{ and } \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma\bar{\rho}^{\gamma-1}(\rho-\bar{\rho});$$

see Lemma 3.1. As a result, we can obtain the decay rates of the relative weak entropy η_* (Theorem 5.2), by making use of the relative entropy inequality and time-weighted entropy estimates.

For the L^1 estimate with $1 < \gamma < +\infty$, the case $\gamma \ge 2$ can be handled by Lemma 3.2. However, for $1 < \gamma < 2$,

$$C|\rho - \bar{\rho}|^2 \le \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1}(\rho - \bar{\rho}) \le C'|\rho - \bar{\rho}|^{\gamma},$$

which means that Lemma 3.2 is not valid, and the index 2 of $|\rho - \bar{\rho}|^2$ is not suitable. Hence, we should seek suitable lower bound of

$$\rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}),$$

which is quite difficult. By a careful decomposition of the region, we are able to overcome this difficulty; see Lemma 3.3 below.

3 Some lemmas and priori estimates

In this section, we state or prove some auxiliary results, which are important for proving our main result. **Lemma 3.1.** Let $1 < \gamma < +\infty$, and $0 \le \rho$, $\overline{\rho} \le C$. Then, there exists a constant c > 0, such that

$$\rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma+1)\bar{\rho}^{\gamma}(\rho-\bar{\rho}) \ge c(\rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma\bar{\rho}^{\gamma-1}(\rho-\bar{\rho}))^{\frac{\gamma+1}{\gamma}}.$$
(3.1)

Proof. When $\bar{\rho} = 0$, (3.1) is true for any $0 < c \leq 1$. For $\rho = 0$, (3.1) becomes $\gamma \bar{\rho}^{\gamma+1} \geq c(\gamma-1)^{\frac{\gamma+1}{\gamma}} \bar{\rho}^{\gamma+1}$, and it is equivalent to proving that $\frac{\gamma}{(\gamma-1)^{\frac{\gamma+1}{\gamma}}}$, has a positive lower bound, which is guaranteed by $\gamma < +\infty$.

Now, assume $\bar{\rho} \neq 0, \rho \neq 0$. Then $\bar{\rho} > 0$. Notice that (3.1) is equivalent to

$$\left(\frac{\rho}{\bar{\rho}}\right)^{\gamma+1} - 1 - (\gamma+1)\left(\frac{\rho}{\bar{\rho}} - 1\right) \ge c \left[\left(\frac{\rho}{\bar{\rho}}\right)^{\gamma} - 1 - \gamma\left(\frac{\rho}{\bar{\rho}} - 1\right)\right]^{\frac{\gamma+1}{\gamma}}$$

Set $\frac{\rho}{\bar{\rho}} = x$ and

$$f(x) = x^{\gamma+1} - 1 - (\gamma+1)(x-1) - c \left[x^{\gamma} - 1 - \gamma(x-1)\right]^{\frac{\gamma+1}{\gamma}};$$

it is obvious that f(1) = 0 and

$$\frac{f'(x)}{\gamma+1} = x^{\gamma} - 1 - \frac{c}{\gamma} [x^{\gamma} - 1 - \gamma(x-1)]^{\frac{1}{\gamma}} (\gamma x^{\gamma-1} - \gamma)$$
$$= x^{\gamma} - 1 - c [x^{\gamma} - 1 - \gamma(x-1)]^{\frac{1}{\gamma}} (x^{\gamma-1} - 1).$$

Next, we claim that there exists $0 < c \leq 1$, such that $f'(x) \leq 0, \forall x \in (0,1)$, and $f'(x) \geq 0, \forall x \in (1, +\infty)$. That is,

$$f(x) \ge f(1) = 0, \quad \forall x \in [0, +\infty),$$

which will conclude the proof. To this end, we set

$$h(x) = \frac{x^{\gamma} - 1}{[x^{\gamma} - 1 - \gamma(x - 1)]^{\frac{1}{\gamma}} (x^{\gamma - 1} - 1)}$$

and divide the proof into two cases.

Case
$$A: x \in (1, +\infty)$$
.
Observing $[(1+t)^k - 1] \sim kt \ (t \to 0)$, where $k > 0$ is a constant, we find that

$$\lim_{x \to 1^+} \frac{x^{\gamma} - 1}{\left[x^{\gamma} - 1 - \gamma(x - 1)\right]^{\frac{1}{\gamma}} (x^{\gamma - 1} - 1)} = \lim_{x \to 1^+} \frac{\gamma x}{\left[x^{\gamma} - 1 - \gamma(x - 1)\right]^{\frac{1}{\gamma}} (\gamma - 1)x} = +\infty,$$

due to

$$[x^{\gamma} - 1 - \gamma(x - 1)]^{\frac{1}{\gamma}} \to 0$$
, as $x \to 1^+$.

In addition,

$$\lim_{x \to +\infty} h(x) = \lim_{x \to +\infty} \frac{1 - \frac{1}{x^{\gamma}}}{\left[1 - \frac{1}{x^{\gamma}} - \gamma \left(\frac{1}{x^{\gamma-1}} - \frac{1}{x^{\gamma}}\right)\right]^{\frac{1}{\gamma}} \left(1 - \frac{1}{x^{\gamma-1}}\right)} = 1.$$

Hence, there exist $0 < \delta < 1$ and X > 2 such that, when $x \in (1, 1 + \delta)$ or x > X,

$$h(x) \ge \frac{1}{2}.$$

When $x \in [1 + \delta, X]$, $\min_{x \in [1+\delta, X]} h(x) = h(x_0) > 0$, since h(x) is a continuous function. In conclusion, for any $x \in (1, +\infty)$, $h(x) \ge \min\left\{h(x_0), \frac{1}{2}\right\}$.

Case $B: x \in (0, 1)$.

In this case, $x^{\gamma-1} - 1 < 0$,

$$\lim_{x \to 1^{-}} \frac{x^{\gamma} - 1}{\left[x^{\gamma} - 1 - \gamma(x - 1)\right]^{\frac{1}{\gamma}} (x^{\gamma - 1} - 1)} = \lim_{x \to 1^{-}} \frac{\gamma x}{\left[x^{\gamma} - 1 - \gamma(x - 1)\right]^{\frac{1}{\gamma}} (\gamma - 1)x} = +\infty,$$

and

$$\lim_{x \to 0^+} \frac{x^{\gamma} - 1}{\left[x^{\gamma} - 1 - \gamma(x - 1)\right]^{\frac{1}{\gamma}} (x^{\gamma - 1} - 1)} = (\gamma - 1)^{-\frac{1}{\gamma}}.$$

Hence, there exists $0 < \delta_1 < \frac{1}{4}$, such that when $x \in (1 - \delta_1, 1)$ or $x \in (0, \delta_1)$,

$$h(x) \ge \frac{1}{2}(\gamma - 1)^{-\frac{1}{\gamma}}.$$

If $x \in [\delta_1, 1 - \delta_1]$, we have, similarly to *Case A*,

$$\min_{x \in [\delta_1, 1-\delta_1]} h(x) = h(x_1) > 0,$$

and

$$h(x) \ge \min\left\{h(x_1), \frac{1}{2}(\gamma - 1)^{-\frac{1}{\gamma}}\right\}.$$

Consequently, we deduce

$$h(x) \ge \min\left\{h(x_0), h(x_1), \frac{1}{2}, \frac{1}{2}(\gamma - 1)^{-\frac{1}{\gamma}}\right\} > 0.$$

Hence, noting $x^{\gamma} - 1 - \gamma(x - 1) > 0$, $\forall x \neq 1$, and letting

$$c = \frac{1}{2} \min\left\{\frac{\gamma}{(\gamma - 1)^{\frac{\gamma + 1}{\gamma}}}, h(x_0), h(x_1), \frac{1}{2}, \frac{1}{2}(\gamma - 1)^{-\frac{1}{\gamma}}\right\},\$$

justifies the claim and so completes the proof.

Lemma 3.2. [13, 14] Assume $0 \le \rho, \bar{\rho} \le C$ and $\gamma > 1$. Then there exist two positive constants d_1 and d_2 such that

$$\begin{cases} (\rho^{\gamma} - \bar{\rho}^{\gamma}) (\rho - \bar{\rho}) \geq |\rho - \bar{\rho}|^{\gamma+1}, \\ d_1 \left(\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}\right) (\rho - \bar{\rho})^2 \leq \rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma+1)\bar{\rho}^{\gamma}(\rho - \bar{\rho}) \\ \leq d_2 \left(\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}\right) (\rho - \bar{\rho})^2, \\ d_1 \left(\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}\right) (\rho - \bar{\rho})^2 \leq (\rho^{\gamma} - \bar{\rho}^{\gamma}) (\rho - \bar{\rho}) \\ \leq d_2 \left(\rho^{\gamma-1} + \bar{\rho}^{\gamma-1}\right) (\rho - \bar{\rho})^2. \end{cases}$$

From the above lemma, we see

$$\rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) \ge C |\rho - \bar{\rho}|^{\gamma},$$

only holds for $\gamma \geq 2$. Hence, we give a lemma to roughly describe the lower bound of $\rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1}(\rho - \bar{\rho})$ for $\gamma \in (1, 2)$, which is essentially important in the proof of (2.7). **Lemma 3.3.** Let $1 < \gamma < 2$, and $0 \leq \rho$, $\bar{\rho} \leq C$. Then, there exist two positive constants c_1 and c_2 , such that

(1). If $\rho = 0$, or $\bar{\rho} = 0$, or $|\rho - \bar{\rho}| \ge \frac{1}{2}\bar{\rho}$, then

$$c_1|\rho - \bar{\rho}|^{\gamma} \le \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1}(\rho - \bar{\rho}).$$
(3.2)

(2). If $\rho \neq 0$, $\bar{\rho} \neq 0$ and $|\rho - \bar{\rho}| < \frac{1}{2}\bar{\rho}$, then

$$c_2\bar{\rho}^{\gamma-2}|\rho-\bar{\rho}|^{\gamma} \le \rho^{\gamma}-\bar{\rho}^{\gamma}-\gamma\bar{\rho}^{\gamma-1}(\rho-\bar{\rho}).$$
(3.3)

Proof. To begin with, we prove (3.2). When $\rho = 0$, (3.2) is true for any $0 < c_1 \leq \gamma - 1$. When $\bar{\rho} = 0$, (3.2) is true for any $0 < c_1 \leq 1$. Now, assume $\rho \neq 0$, $\bar{\rho} \neq 0$. Hence, $\bar{\rho} > 0$. Notice (3.2) is equivalent to

$$c_1 \left| \frac{\rho}{\bar{\rho}} - 1 \right|^{\gamma} \le \left(\frac{\rho}{\bar{\rho}} \right)^{\gamma} - 1 - \gamma \left(\frac{\rho}{\bar{\rho}} - 1 \right).$$

Set $\frac{\rho}{\bar{\rho}} = x$ and

$$f(x) = \frac{x^{\gamma} - 1 - \gamma(x - 1)}{|x - 1|^{\gamma}}$$

It is obvious that $x \ge \frac{3}{2}$ or $0 < x \le \frac{1}{2}$ by $|\rho - \bar{\rho}| \ge \frac{1}{2}\bar{\rho}$ and $\rho \ne 0$, $\bar{\rho} \ne 0$. Then, we divide into two cases.

Case $I: x \geq \frac{3}{2}$. Since

$$\lim_{x \to +\infty} f(x) = \lim_{x \to +\infty} \frac{x^{\gamma} - 1 - \gamma(x - 1)}{|x - 1|^{\gamma}} = 1,$$

there exists X > 2 such that $f(x) \ge \frac{1}{2}$ for x > X. When $x \in \left[\frac{3}{2}, X\right]$,

$$\min_{x \in [3/2, X]} f(x) = f(x_0) > 0,$$

since f(x) is a continuous function.

As a result, for any $x \ge \frac{3}{2}$, $f(x) \ge \min\left\{f(x_0), \frac{1}{2}\right\}$.

 $Case~II: 0 < x \leq \frac{1}{2}.$ In this case,

$$\lim_{x \to 0^+} f(x) = \lim_{x \to 0^+} \frac{x^{\gamma} - 1 - \gamma(x - 1)}{|x - 1|^{\gamma}} = \gamma - 1.$$

Hence, there exists $0 < \delta_1 < \frac{1}{4}$ such that

$$f(x) \ge \frac{\gamma - 1}{2}, \quad \forall x \in (0, \delta_1).$$

If $x \in [\delta_1, \frac{1}{2}]$, similarly as in *Case I*,

$$\min_{x \in [\delta_1, 1/2]} f(x) = f(x_1) > 0,$$

and

$$f(x) \ge \min\left\{f(x_1), \frac{\gamma - 1}{2}\right\}.$$

Consequently, we deduce that

$$f(x) \ge \min\left\{f(x_0), f(x_1), \frac{\gamma - 1}{2}\right\} > 0,$$

due to $1 < \gamma < 2$. Hence, letting

$$c_1 = \frac{1}{2} \min\left\{f(x_0), f(x_1), \frac{\gamma - 1}{2}\right\}$$

completes the proof of (3.2).

As for (3.3), we note

$$0 < \frac{1}{2}\bar{\rho} < \rho < \frac{3}{2}\bar{\rho}.$$

Then, it follows from Taylor's formula that, with $\delta_2 \in (0, 1)$,

$$\rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) = \frac{\gamma(\gamma - 1)}{2} (\rho + \delta_2 \bar{\rho})^{\gamma-2} |\rho - \bar{\rho}|^2$$
$$\geq \frac{\gamma(\gamma - 1)}{2} \left(\frac{3}{2}\bar{\rho} + \delta_2 \bar{\rho}\right)^{\gamma-2} |\rho - \bar{\rho}|^2$$
$$\geq \frac{\gamma(\gamma - 1)}{2} \left(\frac{3}{2}\bar{\rho} + \bar{\rho}\right)^{\gamma-2} |\rho - \bar{\rho}|^2.$$

Hence, (3.3) holds with

$$c_2 = \frac{\gamma(\gamma - 1)}{2} \left(\frac{2}{5}\right)^{2-\gamma} > 0$$

Next, we recall the invariant region theory for L^{∞} weak entropy solution to system (1.1)-(1.2), which was proved in [9].

Theorem 3.4. Suppose $0 \le \nu < 1$ and $(\rho_0, u_0)(x) \in L^{\infty}(\mathbf{R})$ satisfies

$$0 \le \rho_0(x) \le C$$
, $|m_0(x)| \le C\rho_0(x)$.

Let $(\rho, u) \in L^{\infty}(\mathbf{R} \times [0, T])$ be an L^{∞} weak entropy solution of the system (1.1)-(1.2) with $\gamma > 1$. Then (ρ, m) satisfies

$$0 \le \rho(x,t) \le C, \quad |m(x,t)| \le C\rho(x,t),$$

where the constant C depends solely on the initial data.

Let (ρ, m) be a weak entropy solution of system (1.1)-(1.2) satisfying conditions in Theorem 2.2, with $m = \rho u$, then

$$\begin{cases} \rho_t + m_x = 0, \\ m_t + \left(\frac{m^2}{\rho} + p(\rho)\right)_x = -\frac{\alpha}{(1+t)^{\nu}}m. \end{cases}$$
(3.4)

Next, suppose that $\bar{\rho}$ is the Barenblatt's solution of the porous medium equation with the same total mass M as ρ , and $\bar{m} = -(\bar{\rho}^{\gamma})_x$. In order to maintain consistency in form with the above equation, setting $\bar{R} = \bar{m}_t + \left(\frac{\bar{m}^2}{\bar{\rho}}\right)_x$, one has

$$\begin{cases} \bar{\rho}_t + \bar{m}_x = 0, \\ \bar{m}_t + \left(\frac{\bar{m}^2}{\bar{\rho}} + p(\bar{\rho})\right)_x = -\frac{\alpha}{(1+t)^{\nu}}\bar{m} + \bar{R}. \end{cases}$$
(3.5)

Then, define

$$\begin{cases} w = \rho - \bar{\rho}, \\ z = m - \bar{m}. \end{cases}$$

It follows from (3.4) and (3.5) that

$$\begin{cases} w_t + z_x = 0, \\ z_t + \left(\frac{m^2}{\rho}\right)_x + (p(\rho) - p(\bar{\rho}))_x + \frac{\alpha}{(1+t)^{\nu}}z = -\bar{m}_t. \end{cases}$$
(3.6)

Set $y = -\int_{-\infty}^{x} w(r,t) dr$. It holds that

$$y_x = -w, \quad z = y_t.$$

Hence, the equation (3.6) turns into a nonlinear wave equation with source terms, degenerate at vacuum:

$$y_{tt} + \left(\frac{m^2}{\rho}\right)_x + (p(\rho) - p(\bar{\rho}))_x + \frac{\alpha}{(1+t)^{\nu}}y_t = -\bar{m}_t.$$
(3.7)

We now derive the estimate of $\bar{\rho}^{\beta_1} \bar{u}^{\beta_2}$ and $\bar{\rho}^{\delta} \left(\frac{\bar{R}}{\bar{\rho}}\right)$ in the following lemma.

Lemma 3.5. Let $1 \le p < +\infty$,

$$\beta_1 \ge -\frac{\gamma - 1}{p},\tag{3.8}$$

 $\beta_2 \geq 0$, and $\delta \geq 0$. Then, it holds that

$$\left\|\bar{\rho}^{\beta_1}\bar{u}^{\beta_2}\right\|_{L^p} \le C(1+t)^{-\frac{p[(1+\nu)\beta_1 + (\gamma-\nu)\beta_2] - (1+\nu)}{p(r+1)}},\tag{3.9}$$

$$\left\|\bar{\rho}^{\beta_1}\bar{u}^{\beta_2}\right\|_{L^{\infty}} \le C(1+t)^{-\frac{(1+\nu)\beta_1+(\gamma-\nu)\beta_2}{r+1}},\tag{3.10}$$

and

$$\left\|\bar{\rho}^{\delta}\left(\frac{\bar{R}}{\bar{\rho}}\right)\right\|_{L^{p}} \leq C(1+t)^{-\frac{p[\delta(1+\nu)+2\gamma+1-\nu]-(1+\nu)}{p(\gamma+1)}}.$$
(3.11)

Proof. On one hand, using (2.3), (2.4) and $\xi = x(t+1)^{-\frac{1+\nu}{\gamma+1}}$, we have

$$\begin{split} & \left\|\bar{\rho}^{\beta_{1}}\bar{u}^{\beta_{2}}\right\|_{L^{p}}^{p} \\ &= \int \left\{ (1+t)^{-\frac{1+\nu}{\gamma+1}} \left[\left(A_{0} - B_{0}(1+t)^{-\frac{2(1+\nu)}{\gamma+1}}x^{2}\right)_{+} \right]^{\frac{1}{\gamma-1}} \right\}^{\beta_{1}p} \left[\frac{x(1+\nu)}{(1+t)(\gamma+1)} \right]^{p\beta_{2}} dx \\ &\leq C \int_{-\sqrt{\frac{A_{0}}{B_{0}}}}^{\sqrt{\frac{A_{0}}{B_{0}}}} (1+t)^{-\frac{1+\nu}{\gamma+1}\beta_{1}p}(1+t)^{-\frac{\gamma-\nu}{\gamma+1}p\beta_{2}} \left[\left(A_{0} - B_{0}\xi^{2}\right)_{+} \right]^{\frac{\beta_{1}p}{\gamma-1}} \xi^{p\beta_{2}}(1+t)^{\frac{1+\nu}{\gamma+1}} d\xi \\ &\leq C(1+t)^{-\frac{p[(1+\nu)\beta_{1}+(\gamma-\nu)\beta_{2}]-(1+\nu)}{r+1}}, \end{split}$$

where the last inequality is from

$$\int_{-\sqrt{\frac{A_0}{B_0}}}^{\sqrt{\frac{A_0}{B_0}}} \left[\left(A_0 - B_0 \xi^2 \right)_+ \right]^{\frac{\beta_1 p}{\gamma - 1}} \xi^{p\beta_2} d\xi \le C \int_{-1}^1 \left[\left(A_0 - A_0 \tau^2 \right)_+ \right]^{\frac{\beta_1 p}{\gamma - 1}} \sqrt{\frac{A_0}{B_0}} d\tau$$
$$= C \int_0^1 (1 - \tau)^{\frac{\beta_1 p}{\gamma - 1}} (1 + \tau)^{\frac{\beta_1 p}{\gamma - 1}} d\tau$$
$$\le C \int_0^1 (1 - \tau)^{\frac{\beta_1 p}{\gamma - 1}} d\tau \le C,$$

by $p\beta_2 \ge 0$, $\sqrt{\frac{A_0}{B_0}}\tau = \xi$, and (3.8). Hence, we can obtain (3.9). Particularly,

$$\left\|\bar{\rho}^{\beta_1}\bar{u}^{\beta_2}\right\|_{L^{\infty}} = ess \sup_{x \in R} \left(\bar{\rho}^{\beta_1}\bar{u}^{\beta_2}\right) \le C(1+t)^{-\frac{(\gamma-\nu)\beta_2 + (1+\nu)\beta_1}{r+1}},$$

by $x = C(1+t)^{\frac{1+\nu}{\gamma+1}}$, which gives (3.10).

On the other hand, since $\bar{R} = \bar{m}_t + \left(\frac{\bar{m}^2}{\bar{\rho}}\right)_x$, and $\bar{m} = \bar{\rho}\bar{u}$, we have

$$\bar{R} = (\bar{\rho}\bar{u})_t + (\bar{\rho}\bar{u}^2)_x = \bar{\rho}_t\bar{u} + \bar{\rho}\bar{u}_t + \bar{\rho}_x\bar{u}^2 + 2\bar{\rho}\bar{u}\bar{u}_x = \bar{\rho}\bar{u}_t + \bar{\rho}\bar{u}\bar{u}_x$$

by $\bar{\rho}_t + (\bar{\rho}\bar{u})_x = 0$. Combining the above equality and (2.4), we derive

$$\frac{\bar{R}}{\bar{\rho}} = \bar{u}_t + \bar{u}\bar{u}_x = -\frac{(1+\nu)(\gamma-\nu)}{(1+t)^2(\gamma+1)^2}x_t$$

which implies that with $\xi = x(t+1)^{-\frac{1+\nu}{\gamma+1}}$,

$$\begin{split} & \left\| \bar{\rho}^{\delta} \left(\frac{\bar{R}}{\bar{\rho}} \right) \right\|_{L^{p}}^{p} \\ &= \int \left\{ (1+t)^{-\frac{1+\nu}{\gamma+1}} \left[\left(A_{0} - B_{0}(1+t)^{-\frac{2(1+\nu)}{\gamma+1}} x^{2} \right)_{+} \right]^{\frac{1}{\gamma-1}} \right\}^{\delta p} \left[\frac{(1+\nu)(\gamma-\nu)x}{(1+t)^{2}(\gamma+1)^{2}} \right]^{p} dx \\ &\leq C \int_{-\sqrt{\frac{A_{0}}{B_{0}}}}^{\sqrt{\frac{A_{0}}{B_{0}}}} (1+t)^{-\frac{(1+\nu)\delta p}{\gamma+1}} (1+t)^{-\frac{p(2\gamma+1-\nu)}{\gamma+1}} \left[(A_{0} - B_{0}\xi)_{+} \right]^{\frac{\delta p}{\gamma-1}} \xi^{p} (1+t)^{\frac{1+\nu}{\gamma+1}} d\xi \\ &\leq C (1+t)^{-\frac{p[\delta(1+\nu)+2\gamma+1-\nu]-(1+\nu)}{\gamma+1}}, \end{split}$$

from which we complete the proof of (3.11).

4 Extension of key intermediate estimates

In this section, we are going to extend the results in [9, Lemmas 3.8 and 3.9] (with $\gamma \in (1,3)$) to $\gamma \in (1,+\infty)$. In [9], the authors used the theory of divergence-measure fields (see Chen and Frid [2]), (3.7) and the following two entropy inequalities

$$\eta_{et} + q_{ex} + \frac{m^2}{(1+t)^{\nu}\rho} \le 0, \tag{4.1}$$

and

$$\tilde{\eta}_t + \tilde{q}_x + \frac{2C_2m^2}{(1+t)^{\nu}} + \frac{A_mm}{(1+t)^{\nu}} \le 0,$$
(4.2)

where

$$\eta_e = \frac{\rho}{2} \int_{-1}^{1} \left| u + z\rho^{\theta} \right|^2 \left(1 - z^2 \right)^{\lambda} dz = \frac{1}{2} \frac{m^2}{\rho} + \frac{\kappa}{\gamma - 1} \rho^{\gamma},$$

and

$$\tilde{\eta} = \rho \int_{-1}^{1} \left| u + z\rho^{\theta} \right|^{\frac{2\gamma}{\gamma+2}} (1-z^2)^{\lambda} dz = C_1 \rho^{\gamma+1} + C_2 m^2 + A(\rho,m),$$

with

$$C_{1} = \int_{-1}^{1} |z|^{\frac{2\gamma}{\gamma-1}} (1-z^{2})^{\lambda} dz,$$

$$C_{2} = \frac{\gamma(\gamma+1)}{(\gamma-1)^{2}} \int_{-1}^{1} |z|^{\frac{2}{\gamma-1}} (1-z^{2})^{\lambda} dz = \frac{2\gamma(\gamma+1)}{(\gamma-1)^{2}} C_{1},$$
(4.3)

and

$$A_m m \ge 3A(\rho, m) \ge 0. \tag{4.4}$$

In addition, they make use of

$$\eta^* = \tilde{\eta} - C_1 \bar{\rho}^{\gamma+1} - C_1 (\gamma + 1) \bar{\rho}^{\gamma} (\rho - \bar{\rho}),$$

and obtain

$$\eta_t^* + \{\cdots\}_x + \frac{2C_2}{(1+t)^{\lambda}} (m-\bar{m})^2 + \frac{A_m m}{(1+t)^{\lambda}} \le C_1(\gamma+1) \left(\bar{\rho}_x^{\gamma} y\right)_t + 2C_1(\gamma+1)\bar{\rho}_t^{\gamma} y_x, \quad (4.5)$$

where C_1 and C_2 are defined in (4.3), and $\{\cdots\}_x$ denotes terms which vanish after integrating over **R**.

According to the theory of divergence-measure fields, (3.7) and (4.1) are the same in both cases, and hence we only need to consider (4.2), (4.4) and (4.5). For this, we state and give two lemmas. The first one is the classical Taylor's formula.

Lemma 4.1. Let $f(\xi) = |\xi|^k$ with $k \in \mathbf{R}_+$. Then, for any $0 \le n \le k, n \in \mathbf{N}$, it holds that

$$f(u+z) - f(z) - f'(z)u - \dots - \frac{f^{(n)}(z)}{n!}u^n = u^{n+1} \int_0^1 \frac{(1-s)^n}{n!} f^{(n+1)}(su+z)ds.$$

The second one is in the following.

Lemma 4.2. For $1 < \gamma < +\infty$ and $g(\xi) = |\xi|^{\frac{2\gamma}{\gamma-1}}$, it holds that

$$\eta = \rho \int_{-1}^{1} g\left(u + z\rho^{\theta}\right) \left(1 - z^{2}\right)^{\lambda} dz = C_{1}\rho^{\gamma+1} + C_{2}m^{2} + B(\rho, m),$$

where C_1 , C_2 are defined in (4.3), and

$$B_m m \ge 2B(\rho, m) \ge 0.$$

Proof. According to (2.1) and Lemma 3.2, for $g(\xi) = |\xi|^{\frac{2\gamma}{\gamma-1}}$, we obtain

$$\eta = \rho \int_{-1}^{1} g\left(u + z\rho^{\theta}\right) \left(1 - z^{2}\right)^{\lambda} dz$$

$$= \rho \int_{-1}^{1} \left[g(z\rho^{\theta}) + g'(z\rho^{\theta})u + u^{2} \int_{0}^{1} (1 - s)g''(su + z\rho^{\theta})ds\right] \left(1 - z^{2}\right)^{\lambda} dz$$

$$= C_{1}\rho^{\gamma+1} + \frac{2\gamma(\gamma+1)}{(\gamma-1)^{2}}\rho u^{2} \int_{-1}^{1} \int_{0}^{1} (1 - s)|su + z\rho^{\theta}|^{\frac{2}{\gamma-1}} \left(1 - z^{2}\right)^{\lambda} dsdz$$

$$= C_{1}\rho^{\gamma+1} + \frac{2\gamma(\gamma+1)}{(\gamma-1)^{2}}\rho^{2}u^{2} \int_{-1}^{1} \int_{0}^{1} (1 - s) \left|s\frac{u}{\rho^{\theta}} + z\right|^{\frac{2}{\gamma-1}} \left(1 - z^{2}\right)^{\lambda} dsdz,$$
(4.6)

due to the fact that

$$\int_{-1}^{1} g'(z\rho^{\theta}) \left(1 - z^{2}\right)^{\lambda} dz = 0.$$

Then, with $a = \frac{u}{\rho^{\theta}}$, we set

$$h(a) = \int_{-1}^{1} |sa + z|^{\frac{2}{\gamma - 1}} (1 - z^{2})^{\lambda} dz$$

= $\int_{0}^{1} |sa + z|^{\frac{2}{\gamma - 1}} (1 - z^{2})^{\lambda} dz + \int_{-1}^{0} |sa + z|^{\frac{2}{\gamma - 1}} (1 - z^{2})^{\lambda} dz$
= $\int_{0}^{1} |sa + z|^{\frac{2}{\gamma - 1}} (1 - z^{2})^{\lambda} dz + \int_{0}^{1} |sa - z|^{\frac{2}{\gamma - 1}} (1 - z^{2})^{\lambda} dz,$

which implies that

$$h(a) = h(-a).$$

It is obvious that

$$h(0) = \int_{-1}^{1} |z|^{\frac{2}{\gamma-1}} \left(1 - z^2\right)^{\lambda} dz = \frac{(\gamma-1)^2}{\gamma(\gamma+1)} C_2,$$

and

$$\begin{aligned} h'(a) \\ &= \frac{2s}{\gamma - 1} \int_{-1}^{1} \frac{|sa + z|^{\frac{2}{\gamma - 1}}}{sa + z} (1 - z^{2})^{\lambda} dz \\ &= \frac{2s}{\gamma - 1} \left[\int_{-1}^{-sa} \frac{|sa + z|^{\frac{2}{\gamma - 1}}}{sa + z} (1 - z^{2})^{\lambda} dz + \int_{-sa}^{1} (sa + z)^{\frac{3 - \gamma}{\gamma - 1}} (1 - z^{2})^{\lambda} dz \right] \\ &= \frac{2s}{\gamma - 1} \left[\int_{sa - 1}^{0} \frac{|v|^{\frac{2}{\gamma - 1}}}{v} (1 - (v - sa)^{2})^{\lambda} dv + \int_{0}^{1 + sa} v^{\frac{3 - \gamma}{\gamma - 1}} (1 - (v - sa)^{2})^{\lambda} dv \right] \\ &= \frac{2s}{\gamma - 1} \left[-\int_{0}^{1 - sa} v^{\frac{3 - \gamma}{\gamma - 1}} (1 - (v + sa)^{2})^{\lambda} dv + \int_{0}^{1 + sa} v^{\frac{3 - \gamma}{\gamma - 1}} (1 - (v - sa)^{2})^{\lambda} dv \right] \\ &\geq 0, \end{aligned}$$

$$(4.7)$$

where the last inequality is from

$$1 - (v - sa)^2 \ge 1 - (v + sa)^2$$
,

for any $v \in [0, 1 - sa]$ and $s \ge 0$. Hence, $h(a) \ge h(0) = \frac{(\gamma - 1)^2}{\gamma(\gamma + 1)}C_2$.

Combining the above arguments and (4.6), we deduce that

$$\begin{split} \eta &= C_1 \rho^{\gamma+1} + \frac{2\gamma(\gamma+1)}{(\gamma-1)^2} \rho^2 u^2 \int_{-1}^1 \int_0^1 (1-s) \left| sa + z \right|^{\frac{2}{\gamma-1}} \left(1 - z^2 \right)^{\lambda} ds dz \\ &= C_1 \rho^{\gamma+1} + \frac{2\gamma(\gamma+1)}{(\gamma-1)^2} m^2 \int_0^1 (1-s) h(a) ds \\ &= C_1 \rho^{\gamma+1} + C_2 m^2 + m^2 \left[\frac{2\gamma(\gamma+1)}{(\gamma-1)^2} \int_0^1 (1-s) h(a) ds - C_2 \right], \end{split}$$

and

$$B(\rho,m) = m^2 \left[\frac{2\gamma(\gamma+1)}{(\gamma-1)^2} \int_0^1 (1-s)h(a)ds - C_2 \right]$$

$$\geq C_2 m^2 \left(2\int_0^1 (1-s)ds - 1 \right) = 0.$$

To prove $B_m m \ge 2B(\rho, m)$, we rewrite $B(\rho, m)$ as follows

$$B(\rho,m) = \frac{2\gamma(\gamma+1)}{(\gamma-1)^2} \frac{m^2}{\rho} \int_{-1}^{1} \int_{0}^{1} (1-s) |su+z\rho^{\theta}|^{\frac{2}{\gamma-1}} \left(1-z^2\right)^{\lambda} dsdz - C_2 m^2.$$
(4.8)

Consequently,

$$B_{m} = \frac{4\gamma(\gamma+1)}{(\gamma-1)^{2}} \frac{m}{\rho} \int_{-1}^{1} \int_{0}^{1} (1-s) |su+z\rho^{\theta}|^{\frac{2}{\gamma-1}} (1-z^{2})^{\lambda} dsdz - 2C_{2}m + \frac{4\gamma(\gamma+1)}{(\gamma-1)^{3}} u^{2} \int_{-1}^{1} \int_{0}^{1} (1-s)s \frac{|su+z\rho^{\theta}|^{\frac{2}{\gamma-1}}}{su+z\rho^{\theta}} (1-z^{2})^{\lambda} dsdz,$$

and

$$B_m m = \frac{4\gamma(\gamma+1)}{(\gamma-1)^2} \frac{m^2}{\rho} \int_{-1}^{1} \int_{0}^{1} (1-s) |su+z\rho^{\theta}|^{\frac{2}{\gamma-1}} (1-z^2)^{\lambda} dsdz - 2C_2 m^2 + \frac{4\gamma(\gamma+1)}{(\gamma-1)^3} u^2 m \int_{-1}^{1} \int_{0}^{1} (1-s) s \frac{|su+z\rho^{\theta}|^{\frac{2}{\gamma-1}}}{su+z\rho^{\theta}} (1-z^2)^{\lambda} dsdz.$$

$$(4.9)$$

Comparing this with (4.8) gives $B_m m \ge 2B(\rho, m)$. Hence, we only need to show the nonnegativity of the last term on the right-side of (4.9).

To this end, we set

$$h_1(a) = u \int_{-1}^{1} \frac{|su + z\rho^{\theta}|^{\frac{2}{\gamma - 1}}}{su + z\rho^{\theta}} (1 - z^2)^{\lambda} dz$$

= $\rho a \int_{-1}^{1} \frac{|sa + z|^{\frac{2}{\gamma - 1}}}{sa + z} (1 - z^2)^{\lambda} dz$
= $\rho a \int_{0}^{1} \frac{|sa + z|^{\frac{2}{\gamma - 1}}}{sa + z} (1 - z^2)^{\lambda} dz + \rho a \int_{0}^{1} \frac{|sa - z|^{\frac{2}{\gamma - 1}}}{sa - z} (1 - z^2)^{\lambda} dz,$

which implies that

$$h_1(-a) = -\rho a \int_0^1 \frac{|-sa+z|^{\frac{2}{\gamma-1}}}{-sa+z} (1-z^2)^\lambda dz - \rho a \int_0^1 \frac{|-sa-z|^{\frac{2}{\gamma-1}}}{-sa-z} (1-z^2)^\lambda dz$$
$$= \rho a \int_0^1 \frac{|sa-z|^{\frac{2}{\gamma-1}}}{sa-z} (1-z^2)^\lambda dz + \rho a \int_0^1 \frac{|sa+z|^{\frac{2}{\gamma-1}}}{sa+z} (1-z^2)^\lambda dz$$
$$= h_1(a).$$

In addition, for $\rho \geq 0$ and $a \geq 0$, we have

$$h_1(a) = \rho a \int_{-1}^1 \frac{|sa+z|^{\frac{2}{\gamma-1}}}{sa+z} \left(1-z^2\right)^{\lambda} dz = \frac{\gamma-1}{2s} \rho a h'(a) \ge 0.$$

where the last inequality is from (4.7). Consequently, $h_1(a) \ge 0, \forall a \in R$ by $h_1(-a) = h_1(a)$.

Hence, we have the estimate of the last term on the right-side of (4.9) as follows

$$u^{2}m \int_{-1}^{1} \int_{0}^{1} (1-s)s \frac{|su+z\rho^{\theta}|^{\frac{2}{\gamma-1}}}{su+z\rho^{\theta}} (1-z^{2})^{\lambda} dsdz$$
$$= um \int_{0}^{1} (1-s)sh_{1}(a)ds$$
$$= \rho u^{2} \int_{0}^{1} (1-s)sh_{1}(a)ds \ge 0,$$

and so complete the proof.

Remark 4.3. In Lemma 4.2, we only obtain $B_m m \ge 2B(\rho, m) \ge 0$. Actually, it does not matter whether the coefficient is 2 or 3. It just needs to be larger than 1. In addition, we can obtain (4.5) with $A_m m$ replaced by $B_m m$, since C_1 and C_2 are the same in both cases.

Before making the extension, we recall Lemma 3.7 of [9].

Lemma 4.4. [9] For $\nu \in (0,1)$, $1 < \gamma < 3$ and any $\beta_0 > 0$, it holds that

$$(1+t)^{\omega} \int \eta_e dx + \left(1 - \frac{\omega}{2}\right) \int_0^t \int (1+\tau)^{\omega-\lambda} \frac{m^2}{\rho} dx d\tau$$

$$\leq C + C \int_0^t \int (1+\tau)^{\beta_0} |\rho - \bar{\rho}|^{\gamma+1} dx d\tau,$$

$$(4.10)$$

where η_e is defined in (4.1) and $\omega < \min\left\{\frac{(\gamma-1)(\nu+1)}{\gamma+1}, \frac{(\gamma-1)(1+\beta_0)}{\gamma}\right\}$.

Next, we recall Lemmas 3.6,3.8 and 3.9 of [9] as follows.

Lemma 4.5. [9] For $\nu \in (0, 1)$, $1 < \gamma < 3$, it holds that

$$(1+t)^{\mu_1(\varepsilon)} \int |\rho - \bar{\rho}|^{\gamma+1} dx + (1+t)^{\mu_1(\varepsilon)} \int m^2 dx + \int_0^t \int (1+\tau)^{\mu_1(\varepsilon)-\nu} y_\tau^2 dx d\tau + \int_0^t \int (1+\tau)^{\mu_1(\varepsilon)-\nu} A dx d\tau \le C$$

and

$$(1+t)^{\theta_{1}(\varepsilon)-\nu} \int y^{2} dx + \int_{0}^{t} \int (1+\tau)^{\theta_{1}(\varepsilon)} \left(\rho^{\gamma} - \bar{\rho}^{\gamma}\right) \left(\rho - \bar{\rho}\right) dx d\tau + \int_{0}^{t} \int (1+\tau)^{\theta_{1}(\varepsilon)} m^{2} dx d\tau + \left(\nu - \theta_{1}(\varepsilon)\right) \int_{0}^{t} \int (1+\tau)^{\theta_{1}(\varepsilon)-\nu-1} y^{2} dx d\tau \leq C,$$

$$(4.11)$$

where for any small $\varepsilon > 0$,

$$\mu_1(\varepsilon) = \min\left\{1, 1 + \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)}\right\} - \varepsilon := \tilde{\mu}_1 - \varepsilon,$$

and

$$\theta_1(\varepsilon) = \min\left\{\tilde{\mu}_1 - \nu, \nu, \frac{\gamma - \nu}{\gamma + 1}\right\} - \varepsilon =: \tilde{\theta}_1 - \varepsilon.$$

Furthermore, it holds that

$$(1+t)^{\mu_{k+1}(\varepsilon)} \int |\rho - \bar{\rho}|^{\gamma+1} dx + (1+t)^{\mu_{k+1}(\varepsilon)} \int m^2 dx + \int_0^t \int (1+\tau)^{\mu_{k+1}(\varepsilon)-\nu} y_\tau^2 dx d\tau + \int_0^t \int (1+\tau)^{\mu_{k+1}(\varepsilon)-\nu} A dx d\tau \le C,$$

$$(4.12)$$

where

$$\mu_{k+1}(\varepsilon) = \tilde{\mu}_{k+1} - \varepsilon = \min\left\{1 + \tilde{\theta}_k, 1 + \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)} + \frac{\tilde{\theta}_k}{2}\right\} - \varepsilon, \quad \forall k \in \mathbf{N},$$
(4.13)

and

$$\tilde{\theta}_k = \min\left\{\tilde{\mu}_k - \nu, \nu, \frac{\gamma - \nu}{\gamma + 1}\right\},\tag{4.14}$$

are increasing sequences with $\tilde{\theta}_0 = 0$.

Now, we are ready to extend the related results in [9] to $\gamma \in (1, +\infty)$.

Theorem 4.6. Under the conditions of Theorem 2.2, let $\nu \in [0,1)$ and $\gamma > 1$. Then it holds that

$$(1+t)^{\mu^{*}(\varepsilon)} \int |\rho - \bar{\rho}|^{\gamma+1} dx + \int_{0}^{t} \int (1+\tau)^{\mu^{*}(\varepsilon)-\nu} y_{\tau}^{2} dx d\tau + \int_{0}^{t} \int (1+\tau)^{\theta^{*}} (\rho^{\gamma} - \bar{\rho}^{\gamma}) (\rho - \bar{\rho}) dx d\tau \leq C,$$

$$(4.15)$$

where

$$\mu^*(\varepsilon) := \begin{cases} 1+\nu - \frac{\nu+1}{2(\gamma+1)} - \varepsilon, & \nu \in \left[0, \frac{\gamma}{\gamma+2}\right], \\ \frac{3}{2} + \frac{\nu}{2} - \frac{\nu+1}{\gamma+1} - \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma+2}, 1\right], \end{cases}$$
(4.16)

and

$$\theta^* := \begin{cases} \nu, & \nu \in \left[0, \frac{\gamma}{\gamma+2}\right], \\ \frac{\gamma-\nu}{\gamma+1}, & \nu \in \left[\frac{\gamma}{\gamma+2}, 1\right). \end{cases}$$
(4.17)

Proof. We pay attention to the case $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$. The other case $\nu \in \left(0, \frac{\gamma}{\gamma+2}\right]$ has been dealt with in [9]; For case $\nu = 0$, see [14](with $1 < \gamma < 3$) and [8](with $\gamma \ge 2$).

It follows from (4.13) and (4.14) that

$$\tilde{\mu}_{k+1} = \begin{cases} 1 + \tilde{\theta}_k, & \tilde{\theta}_k \in \left(-\infty, \frac{\nu\gamma - 1}{\gamma + 1}\right], \\ 1 + \frac{\nu}{2} - \frac{\nu + 1}{2(\gamma + 1)} + \frac{\tilde{\theta}_k}{2}, & \tilde{\theta}_k \in \left(\frac{\nu\gamma - 1}{\gamma + 1}, \infty\right), \end{cases}$$
(4.18)

and

$$\tilde{\theta}_k = \min\left\{\tilde{\mu}_k - \nu, \frac{\gamma - \nu}{\gamma + 1}\right\}, \quad \nu \in \left[, \frac{\gamma}{\gamma + 2}, 1\right), \tag{4.19}$$

since $\nu \ge \frac{\gamma - \nu}{\gamma + 1}$ provided that $\nu \in \left[\frac{\gamma}{\gamma + 2}, 1\right)$.

First, we claim that there exists $k_1 \in \mathbf{N}$ such that for any $k \geq k_1$,

$$\tilde{\theta}_k \ge \tilde{\theta}_{k_1} > \frac{\nu\gamma - 1}{\gamma + 1}.\tag{4.20}$$

If not, it follows from (4.19) that

$$0 < \tilde{\theta}_k = \tilde{\mu}_k - \nu \le \frac{\nu\gamma - 1}{\gamma + 1} < \frac{\gamma - \nu}{\gamma + 1}, \quad \forall k \in \mathbf{N},$$
(4.21)

Combining this and (4.18) shows

$$\tilde{\mu}_{k+1} = 1 + \tilde{\theta}_k = 1 + \tilde{\mu}_k - \nu = k(1-\nu) + \tilde{\mu}_1 \to +\infty, \text{ as } k \to +\infty,$$

which contradicts (4.21), and hence (4.20) holds.

Next, we claim that there exists $k_2 \ge k_1$ such that

$$\tilde{\mu}_{k_2} - \nu \ge \frac{\gamma - \nu}{\gamma + 1}.\tag{4.22}$$

If not, then

$$\tilde{\theta}_k = \tilde{\mu}_k - \nu < \frac{\gamma - \nu}{\gamma + 1}, \quad \forall k \ge k_1.$$
(4.23)

Noting the above inequality, (4.18) and (4.20), we deduce that

$$\begin{split} \tilde{\mu}_{k+1} - \nu &= 1 + \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)} + \frac{\tilde{\mu}_k - \nu}{2} - \nu \\ &= \left(1 + \frac{1}{2}\right) \left(1 + \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)} - \nu\right) + \frac{1}{2^2} (\tilde{\mu}_{k-1} - \nu) \\ &= \left(1 + \frac{1}{2} + \dots + \frac{1}{2^{k-k_1}}\right) \left(1 + \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)} - \nu\right) + \frac{1}{2^{k+1-k_1}} (\tilde{\mu}_{k_1} - \nu) \\ &\geq \left(1 + \frac{1}{2} + \dots + \frac{1}{2^{k-k_1}}\right) \left(1 - \frac{\nu}{2} - \frac{\nu+1}{2(\gamma+1)}\right). \end{split}$$

Since

$$1 - \frac{\nu}{2} - \frac{\nu + 1}{2(\gamma + 1)} \ge \frac{1}{2} - \frac{1}{\gamma + 1} = \frac{\gamma - 1}{2(\gamma + 1)} > 0,$$

due to $\gamma > 1$, we know that for k_3 sufficiently large, it holds that

$$\tilde{\theta}_{k_3} = \tilde{\mu}_{k_3+1} - \nu > \frac{\gamma - \nu}{\gamma + 1},$$

which contradicts (4.23) and consequently (4.22) holds.

Hence, from (4.19) and (4.22), it follows that

$$\lim_{k \to \infty} \tilde{\theta}_k = \frac{\gamma - \nu}{\gamma + 1},$$

which, together with (4.18) and (4.20), justifies the conclusion.

Remark 4.7. Based on (4.11) and (4.12), we added the last term on the left-side of (4.15), which is essential in estimating I_1 and I_3 (\hat{I}_1 and \hat{I}_3) defined in the next section, since

$$\int_0^t \int (1+\tau)^{\theta^*} |\rho - \bar{\rho}|^{\gamma+1} dx d\tau \le C \int_0^t \int (1+\tau)^{\theta^*} |\rho^\gamma - \bar{\rho}^\gamma| |\rho - \bar{\rho}| dx d\tau \le C$$

is better than

$$(1+t)^{\mu^*(\varepsilon)} \int |\rho - \bar{\rho}|^{\gamma+1} dx \le C,$$

by

$$\mu^*(\varepsilon) < \theta^* + 1.$$

Observing (4.10) and (4.15), and choosing $\beta_0 = \theta^*$, we obtain

$$(1+t)^{\omega} \int \eta_e dx + \left(1 - \frac{\omega}{2}\right) \int_0^t \int (1+\tau)^{\omega-\nu} \frac{m^2}{\rho} dx d\tau \le C, \tag{4.24}$$

with $0 < \omega < \min\left\{\frac{(\gamma-1)(\lambda+1)}{\gamma+1}, \frac{(\gamma-1)(1+\theta^*)}{\gamma}\right\}$. Next, we will show $\frac{(\gamma-1)(\nu+1)}{\gamma+1} < \frac{(\gamma-1)(1+\theta^*)}{\gamma},$

for any $\nu \in [0, 1)$. In the case $\nu \in \left[0, \frac{\gamma}{\gamma+2}\right]$, the above inequality is obvious, since $\theta^* = \nu$. For $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$,

$$\frac{1+\theta^*}{\gamma} - \frac{1+\nu}{\gamma+1} = \frac{2\gamma+1-\nu}{\gamma(\gamma+1)} - \frac{1+\nu}{\gamma+1} = \frac{1-\nu}{\gamma} > 0,$$

which implies that (4.25) still holds for $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$. Hence, for any $\varepsilon > 0$, we can obtain (4.24) with

$$\omega = \frac{(\gamma - 1)(\nu + 1)}{\gamma + 1} - \varepsilon, \qquad (4.26)$$

which is helpful in the estimate of $I_4(\hat{I}_4)$ defined below.

(4.25)

5 Proof of the main result

In this section, we detail the proof of Theorem 2.2. First, to obtain the L^{γ} estimates, we set $v = (\rho, m)^t$, $\bar{v} = (\bar{\rho}, \bar{m})^t$, where $(\cdot, \cdot)^t$ means the transpose of the vector (\cdot, \cdot) , and define

$$\eta_* = \eta(v) - \eta(\bar{v}) - \nabla \eta(\bar{v})(v - \bar{v}) = \frac{1}{2}Q_* + \frac{\kappa}{\gamma - 1}P_*,$$

$$q_* = q(v) - q(\bar{v}) - \nabla \eta(\bar{v})(f(v) - f(\bar{v})),$$
(5.1)

where η and q are weak convex entropy-flux pair defined in (2.1) with $g(\xi) = \frac{1}{2}|\xi|^2$,

$$f(v) = \left(m, \frac{m^2}{\rho}\right)^t, f(\bar{v}) = \left(\bar{m}, \frac{\bar{m}^2}{\bar{\rho}}\right)^t,$$

and

$$P_{*} = \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) \ge 0,$$

$$Q_{*} = \frac{m^{2}}{\rho} - \frac{\bar{m}^{2}}{\bar{\rho}} + \frac{\bar{m}^{2}}{\bar{\rho}^{2}} (\rho - \bar{\rho}) - \frac{2\bar{m}}{\bar{\rho}} (m - \bar{m}) \ge 0,$$
(5.2)

due to the convexity of ρ^{γ} and $\frac{m^2}{\rho}$.

According to (3.8) of [6], we have the following lemma describing the properties of η_*, q_* and P_*, Q_*

Lemma 5.1. Under the conditions of Theorem 2.2, let $g(\xi) = \frac{1}{2}|\xi|^2$. Then, it holds for any t > 0 and $1 < \gamma < +\infty$ that

$$\eta_{*t} + q_{*x} + \frac{Q_*}{(1+t)^{\nu}} \le \frac{\bar{m}\,\bar{R}}{\bar{\rho}}\,(\rho - \bar{\rho}) - \frac{\bar{R}}{\bar{\rho}}\,(m - \bar{m}) + \frac{\bar{m}}{\bar{\rho}}(P_* + Q_*)_x.$$
(5.3)

Next, we give the estimate for the relative entropy η_* .

Theorem 5.2. Under the conditions of Theorem 2.2, it holds for any small positive constant ε , t > 0 and $1 < \gamma < +\infty$ that

$$(1+t)^{\phi(\varepsilon)} \int \eta_* dx + \int_0^t \int (1+\tau)^{\phi(\varepsilon)-\nu} Q_* dx d\tau \le C,$$
(5.4)

where η_* is defined in (5.1), Q_* is defined in (5.2) and

$$\phi(\varepsilon) = \begin{cases} \frac{\gamma^2 + \gamma - 1}{(1 + \gamma)^2} (1 + \nu) - \varepsilon, & \nu \in \left[0, \frac{\gamma}{\gamma + 2}\right], \\ \frac{2\gamma - 1 - \nu}{\gamma + 1} - \varepsilon, & \nu \in \left[\frac{\gamma}{\gamma + 2}, 1\right]. \end{cases}$$
(5.5)

Proof. Since the estimates of $|\rho - \bar{\rho}|^{\gamma+1}$ and $|y_t|^2 = |m - \bar{m}|^2$ are different due to the different value of ν according to Theorem 4.6, we divide the proof into two cases.

Case $I: \nu \in \left[0, \frac{\nu}{\nu+2}\right)$.

Setting

$$\mu(\varepsilon) = \frac{\gamma^2 + \gamma - 1}{(\gamma + 1)^2} (1 + \nu) - \varepsilon, \qquad (5.6)$$

multiplying (5.3) with $(1+t)^{\mu(\varepsilon)}$ and integrating over $(0,t) \times \mathbf{R}$, we have

$$(1+t)^{\mu(\varepsilon)} \int \eta_* dx + \int_0^t \int (1+\tau)^{\mu(\varepsilon)-\nu} Q_* dx d\tau$$

$$\leq \int_0^t \int (1+\tau)^{\mu(\varepsilon)} \frac{\bar{m}}{\bar{\rho}} \frac{\bar{R}}{\bar{\rho}} \left(\rho - \bar{\rho}\right) dx d\tau + \int_0^t \int (1+\tau)^{\mu(\varepsilon)} \frac{\bar{R}}{\bar{\rho}} \left(m - \bar{m}\right) dx d\tau$$

$$+ \int_0^t \int \mu(\varepsilon) (1+\tau)^{\mu(\varepsilon)-1} P_* dx d\tau + \int_0^t \int \mu(\varepsilon) (1+\tau)^{\mu(\varepsilon)-1} Q_* dx d\tau + C$$

$$=: I_1 + I_2 + I_3 + I_4 + C,$$
(5.7)

where we have used

$$\int_{-\infty}^{+\infty} \frac{\bar{m}}{\bar{\rho}} (P_* + Q_*)_x = -\int_{-\infty}^{+\infty} \left(\frac{\bar{m}}{\bar{\rho}}\right)_x (P_* + Q_*),$$

and

$$\left(\frac{\bar{m}}{\bar{\rho}}\right)_x = \bar{u}_x = \frac{1+\nu}{(1+t)(1+\gamma)} > 0.$$

To estimate I_1 , we see from (3.9) and (3.11) that

$$\begin{split} \int \frac{\bar{m}}{\bar{\rho}} \frac{\bar{R}}{\bar{\rho}} \left(\rho - \bar{\rho}\right) dx &\leq \left| \left| \frac{\bar{m}}{\bar{\rho}} \right| \right|_{L^{\frac{2(\gamma+1)}{\gamma}}} \left| \left| \frac{\bar{R}}{\bar{\rho}} \right| \right|_{L^{\frac{2(\gamma+1)}{\gamma}}} \left| \left| \rho - \bar{\rho} \right| \right|_{L^{\gamma+1}} \\ &\leq C(1+t)^{-\frac{3\gamma^2 + 3\gamma + 1 - \nu(3\gamma+2)}{(\gamma+1)^2}} \left| \left| \rho - \bar{\rho} \right| \right|_{L^{\gamma+1}}, \end{split}$$

~

which gives

$$I_{1} \leq \int_{0}^{t} (1+\tau)^{\mu(\varepsilon) - \frac{3\gamma^{2} + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^{2}}} ||\rho - \bar{\rho}||_{L^{\gamma + 1}} d\tau$$

$$\leq \int_{0}^{t} \int (1+\tau)^{\theta^{*}} |\rho - \bar{\rho}|^{\gamma + 1} dx d\tau$$

$$+ C \int_{0}^{t} (1+\tau)^{\left[\mu(\varepsilon) - \frac{3\gamma^{2} + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^{2}} - \frac{\theta^{*}}{\gamma + 1}\right] \frac{\gamma + 1}{\gamma}} d\tau$$

$$\leq C,$$
(5.8)

due to (4.15) and

$$\begin{split} \mu(\varepsilon) &- \frac{3\gamma^2 + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^2} - \frac{\theta^*}{\gamma + 1} + \frac{\gamma}{\gamma + 1} \\ &= \frac{\gamma^2 + \gamma - 1}{(\gamma + 1)^2} (1 + \nu) - \varepsilon - \frac{2\gamma^2 + 2\gamma + 1 - \nu(2\gamma + 1)}{(\gamma + 1)^2} \\ &= -\frac{\gamma^2 + \gamma + 2 + \nu(\gamma^2 - \gamma - 2)}{(\gamma + 1)^2} - \varepsilon < 0, \end{split}$$

by (4.17) and $\nu \in \left[0, \frac{\gamma}{\gamma+2}\right]$. As for I_2 , since $m - \bar{m} = y_t$, we deduce by (3.11) that

$$\int \frac{\bar{R}}{\bar{\rho}} (m - \bar{m}) \, dx \le \left| \left| \frac{\bar{R}}{\bar{\rho}} \right| \right| ||m - \bar{m}|| \le C(1 + t)^{-\frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)}} ||y_t||,$$

which implies

$$I_{2} \leq \int_{0}^{t} (1+\tau)^{\mu(\varepsilon) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)}} ||y_{\tau}|| d\tau$$

$$\leq \int_{0}^{t} \int (1+\tau)^{\mu^{*}(\varepsilon) - \nu} y_{\tau}^{2} dx d\tau$$

$$+ C \int_{0}^{t} (1+\tau)^{2 \left[\mu(\varepsilon) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)} - \frac{\mu^{*}(\varepsilon) - \nu}{2} \right]} d\tau$$

$$\leq C,$$

(5.9)

due to (4.15) and

$$\begin{split} \mu(\varepsilon) &- \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)} - \frac{\mu^*(\varepsilon) - \nu}{2} + \frac{1}{2} \\ &= \frac{\gamma^2 + \gamma - 1}{(\gamma + 1)^2} (1 + \nu) - \frac{\varepsilon}{2} - \frac{8\gamma + 1 - 7\nu}{4(\gamma + 1)} \\ &= -\frac{4\gamma^2 + 5\gamma + 5 - \nu(4\gamma^2 + 11\gamma + 6)}{4(\gamma + 1)^2} - \frac{\varepsilon}{2} < 0, \end{split}$$

by (4.16) and $\nu \in \left[0, \frac{\gamma}{\gamma+2}\right]$. Then, to estimate I_3 , using Lemma 3.1 and 3.2, we first get

$$P_* = \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho})$$

$$\leq C \left[\rho^{\gamma+1} - \bar{\rho}^{\gamma+1} - (\gamma + 1) \bar{\rho}^{\gamma} (\rho - \bar{\rho}) \right]^{\frac{\gamma}{\gamma+1}}$$

$$\leq C \left[(\rho^{\gamma} - \bar{\rho}^{\gamma}) (\rho - \bar{\rho}) \right]^{\frac{\gamma}{\gamma+1}}.$$

Hence, it follows from (3.9) that

$$\begin{split} \int \bar{\rho}^{\delta} P_* \bar{\rho}^{-\delta} dx &\leq C \int \bar{\rho}^{\delta} \left[\left(\rho^{\gamma} - \bar{\rho}^{\gamma} \right) \left(\rho - \bar{\rho} \right) \right]^{\frac{\gamma}{\gamma+1}} \bar{\rho}^{-\delta} dx \\ &\leq C \left(\int \bar{\rho}^{\frac{\gamma+1}{\gamma}\delta} \left[\left(\rho^{\gamma} - \bar{\rho}^{\gamma} \right) \left(\rho - \bar{\rho} \right) \right] dx \right)^{\frac{\gamma}{\gamma+1}} \left(\int \bar{\rho}^{-\delta(\gamma+1)} dx \right)^{\frac{1}{\gamma+1}} \\ &\leq C || \bar{\rho}^{\frac{\gamma+1}{\gamma}\delta} ||_{L^{\infty}}^{\frac{\gamma}{\gamma+1}} \left(\int \left(\rho^{\gamma} - \bar{\rho}^{\gamma} \right) \left(\rho - \bar{\rho} \right) dx \right)^{\frac{\gamma}{\gamma+1}} \cdot (1+t)^{\frac{(\gamma+1)\delta+1}{(\gamma+1)^2}(1+\nu)} \\ &\leq C (1+t)^{\frac{1+\nu}{(\gamma+1)^2}} \left(\int \left(\rho^{\gamma} - \bar{\rho}^{\gamma} \right) \left(\rho - \bar{\rho} \right) dx \right)^{\frac{\gamma}{\gamma+1}}, \end{split}$$

where the last inequality is from (3.10). Consequently, we have the estimate of I_3 as follows

$$I_{3} \leq C \int_{0}^{t} (1+\tau)^{\mu(\varepsilon)-1+\frac{1+\nu}{(\gamma+1)^{2}}} \left(\int \left(\rho^{\gamma}-\bar{\rho}^{\gamma}\right) \left(\rho-\bar{\rho}\right) dx \right)^{\frac{\gamma}{\gamma+1}} d\tau$$

$$\leq \int_{0}^{t} \int (1+\tau)^{\theta^{*}} \left(\rho^{\gamma}-\bar{\rho}^{\gamma}\right) \left(\rho-\bar{\rho}\right) dx d\tau$$

$$+ C \int_{0}^{t} (1+\tau)^{\left[\mu(\varepsilon)-1+\frac{1+\nu}{(\gamma+1)^{2}}-\frac{\gamma\theta^{*}}{\gamma+1}\right](\gamma+1)} d\tau$$

$$\leq C,$$
(5.10)

due to (4.15), $\theta^* = \nu$ and

$$\mu(\varepsilon) - 1 + \frac{1+\nu}{(\gamma+1)^2} - \frac{\gamma\theta^*}{\gamma+1} + \frac{1}{\gamma+1} = -\varepsilon < 0.$$

Now, we are left to estimate I_4 . Rewrite I_4 as follows

$$I_{4} = \int_{0}^{t} \int \mu(\varepsilon)(1+\tau)^{\mu(\varepsilon)-1} Q_{*} dx d\tau$$

$$\leq C \int_{0}^{t} \int (1+\tau)^{\mu(\varepsilon)-1} \left[\frac{m^{2}}{\rho} - \frac{\bar{m}^{2}}{\bar{\rho}} + \frac{\bar{m}^{2}}{\bar{\rho}^{2}} (\rho - \bar{\rho}) - \frac{2\bar{m}}{\bar{\rho}} (m - \bar{m}) \right] dx d\tau \qquad (5.11)$$

$$=: I_{41} + I_{42} + I_{43} + I_{44}.$$

First, using (4.24) and (4.26), we have the estimation of I_{41} as follows,

$$I_{41} = \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1} \frac{m^2}{\rho} dx d\tau \le C,$$
 (5.12)

where the last inequality is from

$$\begin{split} \mu(\varepsilon) &-1 - \left[\frac{(\gamma-1)(\nu+1)}{\gamma+1} - \nu - \varepsilon\right] \\ = &\frac{\gamma^2 + \gamma - 1}{(\gamma+1)^2}(1+\nu) - \frac{(\gamma-1)(\nu+1)}{\gamma+1} + \nu - 1 \\ = &\frac{\gamma}{(\gamma+1)^2} + \nu - 1 < 0, \end{split}$$

by $\nu < \frac{\gamma}{\gamma+2}$. As for I_{42} , it follows from (3.9) that

$$I_{42} = \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1} \frac{\bar{m}^2}{\bar{\rho}} dx d\tau \le \int_0^t (1+\tau)^{\mu(\varepsilon)-1-\frac{2\gamma-2\nu}{\gamma+1}} d\tau \le C,$$
(5.13)

due to $\nu < \frac{\gamma}{\gamma+2}$ and

$$\begin{split} \mu(\varepsilon) &- \frac{2\gamma - 2\nu}{\gamma + 1} \le &\frac{\gamma^2 + \gamma - 1}{(\gamma + 1)^2} (1 + \nu) - \frac{2\gamma - 2\nu}{\gamma + 1} \\ &= &- \frac{\gamma^2 + \gamma + 1 - \nu(\gamma^2 + 3\gamma + 1)}{(\gamma + 1)^2} < 0. \end{split}$$

Then, to estimate I_{43} and I_{44} , using (3.9), we first have

$$\int \frac{\bar{m}^2}{\bar{\rho}^2} (\rho - \bar{\rho}) dx \le \left| \left| \bar{u}^2 \right| \right|_{L^{\frac{\gamma+1}{\gamma}}} \left| \left| \rho - \bar{\rho} \right| \right|_{L^{\gamma+1}} \le C(1+t)^{-\frac{2\gamma^2 + \gamma - \nu(2\gamma+3)}{(\gamma+1)^2}} \left| \left| \rho - \bar{\rho} \right| \right|_{L^{\gamma+1}},$$

and

$$\int \frac{2\bar{m}}{\bar{\rho}} (m-\bar{m}) dx \le C ||\bar{u}|| ||m-\bar{m}|| \le C(1+t)^{-\frac{2\gamma-1-3\nu}{2(\gamma+1)}} ||y_t||,$$

which infers that

$$I_{43} = \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1} \frac{\bar{m}^2}{\bar{\rho}^2} (\rho-\bar{\rho}) dx d\tau$$

$$\leq \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1-\frac{2\gamma^2+\gamma-\nu(2\gamma+3)}{(\gamma+1)^2}} ||\rho-\bar{\rho}||_{L^{\gamma+1}} dx d\tau$$

$$\leq \int_0^t (1+\tau)^{\left[\mu(\varepsilon)-\frac{3\gamma^2+3\gamma+1-\nu(2\gamma+3)}{(\gamma+1)^2}-\frac{\theta^*}{\gamma+1}\right]\frac{\gamma+1}{\gamma}} d\tau$$

$$+ C \int_0^t \int (1+\tau)^{\theta^*} |\rho-\bar{\rho}|^{\gamma+1} dx d\tau$$

$$\leq C,$$
(5.14)

and

$$I_{44} = \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1} \frac{2\bar{m}}{\bar{\rho}} (m-\bar{m}) dx d\tau$$

$$\leq C \int_0^t \int (1+\tau)^{\mu(\varepsilon)-1-\frac{2\gamma-1-3\nu}{2(\gamma+1)}} ||y_\tau|| dx d\tau$$

$$\leq \int_0^t (1+\tau)^2 \left[\frac{\mu(\varepsilon)-\frac{4\gamma+1-3\nu}{2(\gamma+1)} - \frac{\mu^*(\varepsilon)-\nu}{2}}{2} \right] d\tau$$

$$+ C \int_0^t \int (1+\tau)^{\mu^*(\varepsilon)-\nu} y_\tau^2 dx d\tau$$

$$\leq C,$$

(5.15)

where we have used (4.15), (5.8) and (5.9).

Thus, in view of (5.11)-(5.15), we deduce that

$$I_4 \leq C.$$

Plugging this and (5.8)-(5.10) into (5.7), we immediately obtain (5.4) for $\nu \in \left[0, \frac{\gamma}{\gamma+2}\right)$. *Case II*: $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$. In this case, setting $2\gamma - 1 - \nu$

$$\hat{\mu}(\varepsilon) = \frac{2\gamma - 1 - \nu}{\gamma + 1} - \varepsilon, \qquad (5.16)$$

multiplying (5.3) with $(1+t)^{\hat{\mu}(\varepsilon)}$ and integrating over $(0,t) \times \mathbf{R}$, we have

$$(1+t)^{\hat{\mu}(\varepsilon)} \int \eta_* dx + \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)-\nu} Q_* dx d\tau$$

$$\leq \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)} \frac{\bar{m}}{\bar{\rho}} \frac{\bar{R}}{\bar{\rho}} (\rho-\bar{\rho}) dx d\tau + \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)} \frac{\bar{R}}{\bar{\rho}} (m-\bar{m}) dx d\tau$$

$$+ \int_0^t \int \hat{\mu}(\varepsilon) (1+\tau)^{\hat{\mu}(\varepsilon)-1} P_* dx d\tau + \int_0^t \int \hat{\mu}(\varepsilon) (1+\tau)^{\hat{\mu}(\varepsilon)-1} Q_* dx d\tau + C$$

$$=: \hat{I}_1 + \hat{I}_2 + \hat{I}_3 + \hat{I}_4 + C.$$
(5.17)

First, we estimate \hat{I}_1 , \hat{I}_2 and \hat{I}_3 . Since the estimates of $\frac{\bar{R}}{\bar{\rho}}$, $\bar{\rho}$ and $\frac{\bar{m}}{\bar{\rho}}$ are the same in the two cases, it follows from (4.15), (5.8), (5.9) and (5.10) that

$$\hat{I}_{1} \leq \int_{0}^{t} (1+\tau)^{\hat{\mu}(\varepsilon) - \frac{3\gamma^{2} + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^{2}}} ||\rho - \bar{\rho}||_{L^{\gamma + 1}} d\tau \\
\leq \int_{0}^{t} \int (1+\tau)^{\theta^{*}} |\rho - \bar{\rho}|^{\gamma + 1} dx d\tau \\
+ C \int_{0}^{t} (1+\tau)^{\left[\hat{\mu}(\varepsilon) - \frac{3\gamma^{2} + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^{2}} - \frac{\theta^{*}}{\gamma + 1}\right] \frac{\gamma + 1}{\gamma}} d\tau \\
\leq C,$$
(5.18)

$$\hat{I}_{2} \leq \int_{0}^{t} (1+\tau)^{\hat{\mu}(\varepsilon) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)}} ||y_{\tau}|| d\tau
\leq \int_{0}^{t} \int (1+\tau)^{\mu^{*}(\varepsilon) - \nu} y_{\tau}^{2} dx d\tau
+ C \int_{0}^{t} (1+\tau)^{2 \left[\hat{\mu}(\varepsilon) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)} - \frac{\mu^{*}(\varepsilon) - \nu}{2} \right]} d\tau
\leq C,$$
(5.19)

and

$$\hat{I}_{3} \leq C \int_{0}^{t} (1+\tau)^{\hat{\mu}(\varepsilon)-1+\frac{1+\nu}{(\gamma+1)^{2}}} \left(\int \left(\rho^{\gamma}-\bar{\rho}^{\gamma}\right) \left(\rho-\bar{\rho}\right) dx \right)^{\frac{\gamma}{\gamma+1}} d\tau$$

$$\leq \int_{0}^{t} \int (1+\tau)^{\theta^{*}} \left(\rho^{\gamma}-\bar{\rho}^{\gamma}\right) \left(\rho-\bar{\rho}\right) dx d\tau$$

$$+ C \int_{0}^{t} (1+\tau)^{\left[\mu(\varepsilon)-1+\frac{1+\nu}{(\gamma+1)^{2}}-\frac{\gamma\theta^{*}}{\gamma+1}\right](\gamma+1)} d\tau$$

$$\leq C$$
(5.20)

due to

$$\hat{\mu}(\varepsilon) - \frac{3\gamma^2 + 3\gamma + 1 - \nu(3\gamma + 2)}{(\gamma + 1)^2} - \frac{\theta^*}{\gamma + 1} + \frac{\gamma}{\gamma + 1} \\ = \frac{2\gamma - 1 - \nu}{\gamma + 1} - \varepsilon - \frac{2\gamma^2 + 3\gamma + 1 - \nu(3\gamma + 3)}{(\gamma + 1)^2}$$

$$\begin{split} &= \frac{2\nu - 2}{\gamma + 1} - \varepsilon < 0, \\ &\hat{\mu}(\varepsilon) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)} - \frac{\mu^*(\varepsilon) - \nu}{2} + \frac{1}{2} \\ &= \frac{2\gamma - 1 - \nu}{\gamma + 1} - \frac{\varepsilon}{2} - \frac{1}{4} + \frac{1}{2} \left(\frac{\nu}{2} + \frac{\nu + 1}{\gamma + 1}\right) - \frac{4\gamma + 1 - 3\nu}{2(\gamma + 1)} \\ &= \frac{\nu - 1}{\gamma + 1} + \frac{\nu - 1}{4} - \frac{\varepsilon}{2} < 0, \end{split}$$

and

$$\hat{\mu}(\varepsilon) - 1 + \frac{1+\nu}{(\gamma+1)^2} - \frac{\gamma\theta^*}{\gamma+1} + \frac{1}{\gamma+1} = -\varepsilon < 0.$$

by (4.16), (4.17) and $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$. As for \hat{I}_4 , we rewrite it as follows

$$\hat{I}_{4} = \int_{0}^{t} \int \hat{\mu}(\varepsilon)(1+\tau)^{\hat{\mu}(\varepsilon)-1} Q_{*} dx d\tau
\leq C \int_{0}^{t} \int (1+\tau)^{\hat{\mu}(\varepsilon)-1} \left[\frac{m^{2}}{\rho} - \frac{\bar{m}^{2}}{\bar{\rho}} + \frac{\bar{m}^{2}}{\bar{\rho}^{2}} (\rho - \bar{\rho}) - \frac{2\bar{m}}{\bar{\rho}} (m - \bar{m}) \right] dx d\tau$$

$$=: \hat{I}_{41} + \hat{I}_{42} + \hat{I}_{43} + \hat{I}_{44}.$$
(5.21)

First, we estimate \hat{I}_{41} and \hat{I}_{42} . It follows from $\nu \in \left[\frac{\gamma}{\gamma+2}, 1\right)$, (4.24) and (4.26) that

$$\hat{I}_{41} = \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)-1} \frac{m^2}{\rho} dx d\tau \le C,$$
(5.22)

where the last inequality is from

$$\begin{split} \hat{\mu}(\varepsilon) &-1 - \left[\frac{(\gamma-1)(\nu+1)}{\gamma+1} - \nu - \varepsilon\right] \\ = & \frac{2\gamma - 1 - \nu}{\gamma+1} - \frac{(\gamma-1)(\nu+1)}{\gamma+1} + \nu - 1 \\ = & \frac{\nu-1}{\gamma+1} - \varepsilon < 0, \end{split}$$

and

$$\hat{I}_{42} = \int_{0}^{t} \int (1+\tau)^{\hat{\mu}(\varepsilon)-1} \frac{\bar{m}^{2}}{\bar{\rho}} dx d\tau \le \int_{0}^{t} (1+\tau)^{\hat{\mu}(\varepsilon)-1-\frac{2\gamma-2\nu}{\gamma+1}} d\tau \le C,$$
(5.23)

due to

$$\hat{\mu}(\varepsilon) - \frac{2\gamma - 2\nu}{\gamma + 1} = \frac{2\gamma - 1 - \nu}{\gamma + 1} - \frac{2\gamma - 2\nu}{\gamma + 1} - \varepsilon < \frac{\nu - 1}{\gamma + 1} < 0.$$

Now, we estimate \hat{I}_{43} and \hat{I}_{44} . Similar to (5.14) and (5.15), we deduce that

$$\hat{I}_{43} = \int_{0}^{t} \int (1+\tau)^{\hat{\mu}(\varepsilon)-1} \frac{\bar{m}^{2}}{\bar{\rho}^{2}} (\rho-\bar{\rho}) dx d\tau
\leq \int_{0}^{t} \int (1+\tau)^{\hat{\mu}(\varepsilon)-1-\frac{2\gamma^{2}+\gamma-\nu(2\gamma+3)}{(\gamma+1)^{2}}} ||\rho-\bar{\rho}||_{L^{\gamma+1}} dx d\tau
\leq \int_{0}^{t} (1+\tau)^{\left[\hat{\mu}(\varepsilon)-\frac{3\gamma^{2}+3\gamma+1-\nu(2\gamma+3)}{(\gamma+1)^{2}}-\frac{\theta^{*}}{\gamma+1}\right]\frac{\gamma+1}{\gamma}} d\tau
+ C \int_{0}^{t} \int (1+\tau)^{\theta^{*}} |\rho-\bar{\rho}|^{\gamma+1} dx d\tau
\leq C,$$
(5.24)

and

$$\hat{I}_{44} = \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)-1} \frac{2\bar{m}}{\bar{\rho}} (m-\bar{m}) dx d\tau
\leq C \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)-1-\frac{2\gamma-1-3\nu}{2(\gamma+1)}} ||y_\tau|| dx d\tau
\leq \int_0^t (1+\tau)^2 \left[\hat{\mu}(\varepsilon) - \frac{4\gamma+1-3\nu}{2(\gamma+1)} - \frac{\mu^*(\varepsilon)-\nu}{2} \right] d\tau
+ C \int_0^t \int (1+\tau)^{\mu^*(\varepsilon)-\nu} y_\tau^2 dx d\tau
\leq C,$$
(5.25)

by (4.15), (5.18) and (5.19).

Substituting (5.22)-(5.25) into (5.21), we have $\hat{I}_4 \leq C$, which together with (5.17)-(5.20) yields that

$$(1+t)^{\hat{\mu}(\varepsilon)} \int \eta_* dx + \int_0^t \int (1+\tau)^{\hat{\mu}(\varepsilon)-\nu} Q_* dx d\tau \le C,$$

and so justifies (5.4) by (5.16) for $\nu \in \left[\frac{\gamma}{\gamma+2}1\right)$.

Now, we are ready to prove Theorem 2.2. From (5.1), (5.2) and Theorem 5.2, we have

$$(1+t)^{\phi(\varepsilon)} \int P_* dx \le C.$$

Combining this, (5.5) and Lemma 3.2, we obtain (2.5) and (2.6) with $\gamma \geq 2$.

As for (2.7) and (2.8), we distinguish two cases.

 $Case \ A: \gamma \geq 2.$

Using Lemma 3.2, (3.9) and (2.5), we obtain the L^1 convergence rate on density as follows

$$\int_{-\infty}^{+\infty} |\rho - \bar{\rho}| dx \le \left(\int_{-\infty}^{+\infty} \bar{\rho}^{\gamma - 2} |\rho - \bar{\rho}|^2 dx\right)^{\frac{1}{2}} \left(\int_{-\infty}^{+\infty} \bar{\rho}^{2 - \gamma} dx\right)^{\frac{1}{2}}$$

$$\leq C(1+t)^{-\frac{\phi(\varepsilon)}{2}}(1+t)^{\frac{\gamma-1}{2(\gamma+1)}(1+\nu)}$$

This and (5.5) together justify (2.7) for $\gamma \geq 2$.

 $Case \ B: 1 < \gamma < 2.$

In this case, let

$$\Omega_1 = \left\{ x : \rho = 0, \, \bar{\rho} = 0, \, |\rho - \bar{\rho}| \ge \frac{1}{2}\bar{\rho} \right\},\\ \Omega_2 = \left\{ x : \rho \neq 0, \, \bar{\rho} \neq 0, \, |\rho - \bar{\rho}| < \frac{1}{2}\bar{\rho} \right\}.$$

In Ω_1 , from Lemma 3.5, (3.2), (5.1), Theorem 5.2 and the Hölder inequality, we see that for any $\delta < \frac{(\gamma-1)^2}{2\gamma}$,

$$\int_{\Omega_{1}} |\rho - \bar{\rho}| dx \leq \left(\int_{\Omega_{1}} |\rho - \bar{\rho}|^{\gamma} dx \right)^{\frac{1}{\gamma}} \left| \left| \bar{\rho}^{\delta} \right| \right|_{L^{\frac{2\gamma}{\gamma-1}}(\Omega_{1})} \left| \left| \bar{\rho}^{-\delta} \right| \right|_{L^{\frac{2\gamma}{\gamma-1}}(\Omega_{1})} \\
\leq C \left(\int_{\Omega_{1}} \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma-1} (\rho - \bar{\rho}) dx \right)^{\frac{1}{\gamma}} \left| \left| \bar{\rho}^{\delta} \right| \right|_{L^{\frac{2\gamma}{\gamma-1}}} \left| \left| \bar{\rho}^{-\delta} \right| \right|_{L^{\frac{2\gamma}{\gamma-1}}} \\
\leq C \left(\int \eta_{*} dx \right)^{\frac{1}{\gamma}} (1 + t)^{\frac{\gamma-1}{\gamma(\gamma+1)}(1+\nu)} \\
\leq C (1 + t)^{-\frac{\phi(\varepsilon)}{\gamma}} (1 + t)^{\frac{\gamma-1}{\gamma(\gamma+1)}(1+\nu)}.$$
(5.26)

On the other hand, in Ω_2 , from Lemma 3.5, (3.3), (5.1), Theorem 5.2 and the Hölder inequality, we deduce that

$$\int_{\Omega_2} |\rho - \bar{\rho}| dx \leq \left(\int_{\Omega_2} \bar{\rho}^{\gamma - 2} |\rho - \bar{\rho}|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega_2} \bar{\rho}^{2 - \gamma} dx \right)^{\frac{1}{2}} \\
\leq C \left(\int_{\Omega_2} \rho^{\gamma} - \bar{\rho}^{\gamma} - \gamma \bar{\rho}^{\gamma - 1} (\rho - \bar{\rho}) dx \right)^{\frac{1}{2}} \left(\int \bar{\rho}^{2 - \gamma} dx \right)^{\frac{1}{2}} \\
\leq C \left(\int \eta_* dx \right)^{\frac{1}{2}} (1 + t)^{\frac{\gamma - 1}{2(\gamma + 1)}(1 + \nu)} \\
\leq C (1 + t)^{-\frac{\phi(\varepsilon)}{2}} (1 + t)^{\frac{\gamma - 1}{2(\gamma + 1)}(1 + \nu)}.$$
(5.27)

Using (5.26) and (5.27) yields

$$\begin{split} \int |\rho - \bar{\rho}| dx &= \int_{\Omega_1} |\rho - \bar{\rho}| dx + \int_{\Omega_2} |\rho - \bar{\rho}| dx \\ &\leq C(1+t)^{-\frac{1}{\gamma} \left[\phi(\varepsilon) - \frac{\gamma - 1}{\gamma + 1}(1+\nu) \right]} + C(1+t)^{-\frac{1}{2} \left[\phi(\varepsilon) - \frac{\gamma - 1}{\gamma + 1}(1+\nu) \right]} \\ &\leq C(1+t)^{-\frac{1}{2} \left[\phi(\varepsilon) - \frac{\gamma - 1}{\gamma + 1}(1+\nu) \right]}, \end{split}$$

where the last inequality is from $1 < \gamma < 2$ and $\phi(\varepsilon) - \frac{\gamma - 1}{\gamma + 1}(1 + \nu) > 0$. Thus, combining the above inequality and (5.5), we obtain (2.7) with $1 < \gamma < 2$.

Hence, we complete the proof of Theorem 2.2.

Declarations

Data availability statement: My manuscript has no associated data.

Conflict of interest: The authors declare no conflict of interest.

References

- J. Carrillo, Y.P. Choi, O. Tse, Convergence to equilibrium in Wasserstein distance for damped Euler equations with interaction forces, Comm. Math. Phys. 365(2019) 329-361.
- [2] G.Q. Chen, H. Frid, Divergence-measure fields and hyperbolic conservation laws, Arch. Ration. Mech. Anal. 147 (1999) 89-118.
- [3] S.H. Chen, H.T. Li, J.Y. Li, M. Mei, K.J. Zhang, Global and blow-up solutions for compressible Euler equations with time-dependent damping, J. Differential Equations 268 (2020) 5035-5077.
- [4] H.B. Cui, H.Y. Yin, J.S. Zhang, C.J. Zhu, Convergence to nonlinear diffusion waves for solutions of Euler equations with time-depending damping, J. Differential Equations 264 (2018) 4564-4602.
- [5] H.B. Cui, H.Y. Yin, C.J. Zhu, L.M. Zhu, Convergence to diffusion waves for solutions of Euler equations with time-depending damping on quadrant, Sci. China Math. 62 (2019) 33-62.
- [6] H.B. Cui, H.Y. Yin, C.J. Zhu, Convergence rates to the Barenblatt solutions for the compressible Euler equations with time-dependent damping, J. Differential Equations 374 (2023) 761-788.
- [7] D. Gallenmüller, P. Gwiazda, A. Świerczewska-Gwiazda, J. Woźnicki, Cahn-Hillard and Keller-Segel systems as high-friction limits of Euler-Korteweg and Euler-Poisson equations, Calc. Var. Partial Differential Equations 63 (2024) 47.
- [8] S.F. Geng, F.M. Huang, L¹-convergence rates to the Barenblatt solution for the damped compressible Euler equations, J. Differential Equations 266 (2019) 7890-7908.
- S.H. Geng, F.M. Huang, X.C. Wu, L¹ -convergence to generalized Barenblatt solution for compressible Euler equations with time-dependent damping, SIAM J. Math. Anal. 53 (2021) 6048-6072.
- [10] L. Hsiao, T.P. Liu, Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping, Commun. Math. Phys. 143 (1992) 599-605.
- [11] F. Hou, I. Witt, H.C. Yin, Global existence and blowup of smooth solutions of 3-D potential equations with time-dependent damping, Pac. J. Math. 292 (2018) 389-426.
- [12] F. Hou, H.C. Yin, On the global existence and blowup of smooth solutions to the multi-dimensional compressible Euler equations with time-depending damping, Nonlinearity 30 (2017) 2485-2517.

- [13] F.M. Huang, P. Marcati, R.H. Pan, Convergence to the Barenblatt solution for the compressible Euler equations with damping and vacuum, Arch. Ration. Mech. Anal. 176 (2005) 1-24.
- [14] F.M. Huang, R.H. Pan, Z. Wang, L1 convergence to the Barenblatt solution for compressible Euler equations with damping, Arch. Ration. Mech. Anal. 200 (2011) 665-689.
- [15] M.N. Jiang, C.J. Zhu, Convergence to strong nonlinear diffusion waves for solutions to p-system with damping on quadrant, J. Differential Equations 246 (2009) 50-77.
- [16] H.T. Li, J.Y. Li, M. Mei, K.J. Zhang, Convergence to nonlinear diffusion waves for solutions of p-system with time-dependent damping, J. Math. Anal. Appl. 456 (2017) 849-871.
- [17] P.L. Lions, B. Perthame, E. Tadmor, Kinetic formulation of the isentropic gas dynamics and p-systems, Commun. Math. Phys. 163 (1994) 169-172.
- [18] T.P. Liu, Compressible flow with damping and vacuum, Jpn. J. Ind. Appl. Math. 13 (1996) 25-32.
- [19] T.P. Liu, T. Yang, Compressible Euler equations with vacuum, J. Differential Equations 140 (1997) 223-237.
- [20] T. Luo, H.H. Zeng, Global existence of smooth solutions and convergence to Barenblatt solutions for the physical vacuum free boundary problem of compressible Euler equations with damping, Commun. Pure Appl. Math. 69 (2016) 1354-1396.
- [21] K. Nishihara, Convergence rates to nonlinear diffusion waves for solutions of system of hyperbolic conservation laws with damping, J. Differential Equations 131 (1996) 171-188.
- [22] K. Nishihara, W.K. Wang, T. Yang, Lp-convergence rate to nonlinear diffusion waves for p-system with damping, J. Differential Equations 161 (2000) 191-218.
- [23] X.H. Pan, Global existence of solutions to 1-d Euler equations with time-dependent damping, Nonlinear Anal. 132 (2016) 327-336.
- [24] X.H. Pan, Blow up of solutions to 1-d Euler equations with time-dependent damping, J. Math. Anal. Appl. 442 (2016) 435-445.
- [25] T.C. Sideris, B. Thomases, D.H. Wang, Long time behavior of solutions to the 3D compressible Euler equations with damping, Commun. Partial Differ. Equ. 28 (2003) 795-816.
- [26] Y. Sugiyama, Singularity formation for the 1D compressible Euler equations with variable damping coefficient, Nonlinear Anal. 170 (2018) 70-87.
- [27] Y. Sugiyama, Remark on global existence of solutions to the 1D compressible Euler equation with time-dependent damping, Adv. Stud. Pure Math. 85 (2020) 379-389.
- [28] J.L. Vázquez, Barenblatt solutions and asymptotic behaviour for a nonlinear fractional heat equation of porous medium type, J. Eur. Math. Soc. 16 (2014) 769-803.
- [29] H.H. Zeng, Time-asymptotics of physical vacuum free boundaries for compressible inviscid flows with damping, Calc. Var. Partial Differential Equations 61(2022), 59.

- [30] H.H. Zeng, Almost global solutions to the three-dimensional isentropic inviscid flows with damping in a physical vacuum around Barenlatt solutions, Arch. Ration. Mech. Anal. 239 (2021) 553-597.
- [31] H.J. Zhao, Convergence to strong nonlinear diffusion waves for solutions of p-system with damping, J. Differential Equations 174 (2001) 200-236.
- [32] C.J. Zhu, Convergence rates to nonlinear diffusion waves for weak entropy solutions to p-system with damping, Sci. China Ser. A 46 (2003) 562-575.
- [33] X.S. Zhu, W.K. Wang, The regular solutions of the isentropic Euler equations with degenerate linear damping, Chin. Ann. Math. Ser. B 26 (2005) 583-598.