# CRITICAL NORM BLOW-UP RATES FOR THE ENERGY SUPERCRITICAL NONLINEAR HEAT EQUATION

## TOBIAS BARKER, HIDEYUKI MIURA, AND JIN TAKAHASHI

Abstract. We prove the first classification of blow-up rates of the critical norm for solutions of the energy supercritical nonlinear heat equation, without any assumptions such as radial symmetry or sign conditions. Moreover, the blow-up rates we obtain are optimal, for solutions that blow-up with bounded  $L^{n(p-1)/2,\infty}(\mathbf{R}^n)$ -norm up to the blow-up time.

We establish these results by proving quantitative estimates for the energy supercritical nonlinear heat equation with a robust new strategy based on quantitative  $\varepsilon$ -regularity criterion averaged over certain comparable time scales. With this in hand, we then produce the quantitative estimates using arguments inspired by Palasek [\[31\]](#page-55-0) and Tao [\[38\]](#page-56-0) involving quantitative Carleman inequalities applied to the Navier-Stokes equations.

Our work shows that energy structure is not essential for establishing blowup rates of the critical norm for parabolic problems with a scaling symmetry. This paves the way for establishing such critical norm blow-up rates for other nonlinear parabolic equations.

#### CONTENTS



<sup>2020</sup> Mathematics Subject Classification. Primary 35K58; Secondary 35B33, 35B44, 35B65. Key words and phrases. Nonlinear heat equation, critical norm, blow-up rate,  $\varepsilon$ -regularity.

#### <span id="page-1-2"></span>1. Introduction

<span id="page-1-1"></span><span id="page-1-0"></span>1.1. Background. We consider the following nonlinear heat equation:

(1.1) 
$$
\begin{cases} u_t = \Delta u + |u|^{p-1}u & \text{in } \mathbf{R}^n \times (0,T), \\ u(\cdot,0) = u_0 & \text{on } \mathbf{R}^n, \end{cases}
$$

where  $n \geq 1$  and  $p > 1$ . The equation enjoys the invariance under the scaling  $u(x,t) \mapsto \lambda^{2/(p-1)}u(\lambda x, \lambda^2 t)$  for  $\lambda > 0$ . The invariance defines the critical Lebesgue space  $L^{q_c}(\mathbf{R}^n)$ , where

$$
q_c := \frac{n(p-1)}{2}.
$$

By local-existence theory, [\(1.1\)](#page-1-2) has a unique local-in-time classical solution (see Subsection [1.7](#page-8-0) for a definition) if  $u_0 \in L^{q_c}(\mathbf{R}^n)$  with  $q_c > 1$ . The solution can be continued as a classical solution up to the maximal time of existence  $T \leq \infty$ . If  $T < \infty$ , the blow-up occurs in the sense of  $L^{\infty}$ . In the pioneering work of Giga and Kohn [\[13\]](#page-55-1), they studied the blow-up rate of the  $L^{\infty}$ -norm, and then they proposed the following question in [\[13,](#page-55-1) Section 7].

<span id="page-1-3"></span>**Question 1.** But it is natural to ask about other norms as well, for example  $||u||_{L^q}$ : do they blow up as  $t \to T$ , and if so, at what rate?

For  $q > q_c$ , local-existence theory provides an answer. Indeed,

$$
||u(\cdot,t)||_{L^q(\mathbf{R}^n)} \geq C(n,p,q)(T-t)^{-\frac{n}{2}(\frac{1}{q_c}-\frac{1}{q})}
$$

holds for finite time blow-up solutions u, see [\[40,](#page-56-1) Section 6] and [\[34,](#page-55-2) Remark 16.2] (iii)]. However, in the critical case  $q = q_c$ , even the first part of Question [1](#page-1-3) does not follow from local-existence theory and has been asked in Brezis-Cazenave [\[5,](#page-54-2) Open problem 7] and Quittner-Souplet [\[34,](#page-55-2) OP2.1, Section 55].

For  $q = q_c$ , the subtlety of Question [1](#page-1-3) is shown by the fact [\[7,](#page-54-3) [8,](#page-55-3) [35\]](#page-55-4) that there exist type II blow-up solutions satisfying  $\sup_{0 \leq t \leq T} ||u(\cdot, t)||_{L^{q_c}(\mathbf{R}^n)} < \infty$  for the energy critical nonlinearity  $p = p_S$  with  $3 \le n \le 5$ , where

$$
p_S := \frac{n+2}{n-2} \quad \text{ for } n \ge 3.
$$

Nevertheless, under the type I assumption, Mizoguchi and Souplet [\[27\]](#page-55-5) showed that  $\lim_{t\to T} ||u(\cdot, t)||_{L^{q_c}(\mathbf{R}^n)} = \infty$  holds for each  $p > 1$ . Here we note that the blow-up is of type I if  $\limsup_{t\to T} (T-t)^{1/(p-1)}||u(\cdot,t)||_{L^{\infty}(\mathbf{R}^n)} < \infty$  and type II if it is not of type I. In addition, the proof in  $[27]$  is inspired by Escauriaza-Seregin-Sverák's seminal result showing the regularity of solutions  $v$  of the 3-dimensional Navier-Stokes equations under the assumption that  $\sup_{0 \le t \le T} ||v(\cdot, t)||_{L^3(\mathbf{R}^n)} < \infty$ .

For the energy supercritical nonlinearity  $p > p<sub>S</sub>$ , the second and third author [\[23,](#page-55-6) [24\]](#page-55-7) removed the type I assumption to show  $\limsup_{t\to T} ||u(\cdot, t)||_{L^{q_c}(\mathbf{R}^n)} = \infty$ , and subsequently

$$
\lim_{t\to T}||u(\cdot,t)||_{L^{q_c}(\mathbf{R}^n)}=\infty.
$$

Thus, for  $p > p<sub>S</sub>$ , the first part of Question [1](#page-1-3) is completely resolved. In this paper, we are interested in the second part, which we label Question [2.](#page-1-4)

<span id="page-1-4"></span>Question 2. For  $p > p_s$ , at what rate does  $||u(\cdot, t)||_{L^{q_c}(\mathbf{R}^n)}$  blow up as  $t \to T$ ?

Such an open problem has also been stated in [\[24,](#page-55-7) Remark 1.6]. The main challenge for resolving Question [2](#page-1-4) is that the qualitative results [\[23,](#page-55-6) [24\]](#page-55-7) (see also [\[25\]](#page-55-8)) crucially use the Giga-Kohn monotonicity formula to classify the blow-up limit as backward self-similar solutions. Such qualitative features are challenging to use in a quantitative way. We are able to overcome this obstacle to obtain our main results below by using quantitative Carleman inequalities established by Palasek [\[31\]](#page-55-0) and Tao [\[38\]](#page-56-0), which in turn use Carleman inequalities established by Escauriaza-Seregin-Sverák  $[9]$ . As a by-product, our method also can be used to estimate the blow-up rate of infinite-time blow-up solutions. In the next subsection, we list our main results.

<span id="page-2-0"></span>1.2. Statement of results. Our first main result addresses Question [2](#page-1-4) by means of a quadruple logarithmic rate in the general setting.

<span id="page-2-1"></span>**Theorem 1.** Let  $n \geq 3$ ,  $p > p_S$  and u be a classical solution of [\(1.1\)](#page-1-2) with  $u_0 \in$  $L^{q_c}(\mathbf{R}^n)$ . If the maximal time of existence  $T > 0$  is finite, then

$$
\limsup_{t \to T} \frac{\|u(\cdot, t)\|_{L^{q_c}(\mathbf{R}^n)}}{\left(\log \log \log \log \left(\frac{1}{(T-t)^{\frac{1}{2(p-1)}}}\right)\right)^c} = \infty,
$$

where  $c > 0$  is a constant depending only on n and p.

While Theorem [1](#page-2-1) provides a rate for Question [2,](#page-1-4) we do not know if the rate is optimal. To the best of our knowledge, for constructed blow-up solutions to the nonlinear heat equation, the  $L^{q_c}$ -norm typically diverges at a logarithmic rate even when the blow-up is type II, see for example [\[36,](#page-56-2) Corollary 1.5] for  $p = p_{JL}$  :=  $(n-2\sqrt{n-1})/(n-4-2\sqrt{n-1})(>ps)$  with  $n \ge 11$ , [\[28,](#page-55-10) Corollary 3] for  $p > p_{JL}$ and [\[16,](#page-55-11) Subsection 8.2.1] for  $p = p_S$  with  $n = 6$ . In particular, the solutions constructed by Seki [\[36\]](#page-56-2) and Mukai-Seki [\[28\]](#page-55-10) satisfy

<span id="page-2-2"></span>(1.2) 
$$
\sup_{0
$$

<span id="page-2-3"></span>(1.3) 
$$
C' \log \left( \frac{1}{T-t} \right) \leq \int_{\mathbf{R}^n} |u(x,t)|^{q_c} dx \leq C'' \log \left( \frac{1}{T-t} \right),
$$

where we also refer to [\[19,](#page-55-12) Subsection 4.1] and [\[20,](#page-55-13) Proposition C.1] for deriving [\(1.2\)](#page-2-2).

Our second main result shows that for solutions satisfying [\(1.2\)](#page-2-2), the lower bound of the blow-up rate in [\(1.3\)](#page-2-3) is generic.

<span id="page-2-4"></span>**Theorem 2.** Let  $n \geq 3$ ,  $p > p_S$  and u be a classical solution of [\(1.1\)](#page-1-2) with  $u_0 \in$  $L^{q_c}(\mathbf{R}^n)$ . There exist constants  $M_0, C > 0$  depending only on n and p such that if maximal time of existence  $T > 0$  is finite and u satisfies

$$
\sup_{0
$$

with a constant  $M \geq M_0$ , then

$$
\int_{\mathbf{R}^n} |u(x,t)|^{q_c} dx \ge e^{-e^{e^{M^C}}} \log\left(\frac{t}{T-t}\right)
$$

for all  $(1 + M^{-2(2p^2 + 3p + 2)})^{-1}T < t < T$ .

Our third result gives the first classification of infinite-time blow-up solutions in terms of the critical norm, which answers a question raised in [\[23,](#page-55-6) Remark 1.9]. In what follows, we write  $B(x, r) := \{y \in \mathbb{R}^n; |x - y| < r\}$  for  $x \in \mathbb{R}^n$  and  $r > 0$ .

<span id="page-3-0"></span>**Theorem 3.** Let  $n \geq 3$ ,  $p > p_S$  and u be a global-in-time classical solution of [\(1.1\)](#page-1-2). There exist constants  $C, M_0 > 0$  depending only on n and p such that if  $\limsup_{t\to\infty} |u(0, t)| = \infty$  and

<span id="page-3-1"></span>
$$
\sup_{0
$$

with a constant  $M \geq M_0$ , then we conclude that there exists a sequence  $t_n \to \infty$ such that

(1.4) 
$$
\int_{B(0,t_n^{\frac{1}{2}}e^{e^{M^C}})} |u(x,t_n)|^{q_c} dx \geq e^{-e^{e^{M^C}}}\log t_n.
$$

Theorem [3](#page-3-0) can be applied to the infinite-time blow-up solutions constructed by Poláčik-Yanagida's works [\[32,](#page-55-14) Theorem 1.3] for  $p \geq p_{JL}$  and [\[33,](#page-55-15) Theorem 1] for  $p_S < p < p_{JL}$ . We note that the blow-up rate of  $L^{\infty}$ -norm is studied in [\[10,](#page-55-16) [11,](#page-55-17) [26\]](#page-55-18) for  $p \geq p_{JL}$  and that Theorem [3](#page-3-0) is also applicable to the solutions in these papers.

Remark 4. In order for the assumptions in Theorem [3](#page-3-0) to apply to the known solutions, it seems necessary that the integral in [\(1.4\)](#page-3-1) is over a finite ball. In particular, we expect that there is no infinite-time blow-up solutions with  $u_0 \in$  $L^{q_c}(\mathbf{R}^n)$ . As evidence for this, under  $p > p_S$  and  $\nabla u_0 \in L^{q_*}(\mathbf{R}^n)$  with  $q_* :=$  $n(p-1)/(p+1)$  (this implies  $u_0 \in L^{q_c}(\mathbf{R}^n)$  by the Sobolev inequality), we can prove that any associated global-in-time classical solution  $u$  satisfies

$$
\sup_{1
$$

by applying the argument of Souplet [\[37,](#page-56-3) Proof of Theorem 2, (7.1)] with the aid of  $[37,$  Proposition 5.1(ii)], the convolution inequalities and the approximation  $v_0 \in C_0^{\infty}(\mathbf{R}^n)$  satisfying  $\|\nabla(u_0 - v_0)\|_{L^{q_*}(\mathbf{R}^n)} < \delta$  for a small constant  $\delta > 0$ .

The above theorems hinge on quantitative estimates for solutions of

(1.5) 
$$
u_t = \Delta u + |u|^{p-1}u \quad \text{in } \mathbb{R}^n \times (-1,0],
$$

which are classical up to (and including)  $t = 0$ . Our main quantitative estimate is as follows. It will be applied to the rescaled function

<span id="page-3-4"></span><span id="page-3-2"></span>
$$
u_t(x,s) := (t^{\frac{1}{2}})^{\frac{2}{p-1}} u(t^{\frac{1}{2}}x, ts+t)
$$

<span id="page-3-3"></span>with  $t \in (0, T)$  fixed.

**Proposition 5** (main quantitative estimate). Let  $n \geq 3$ ,  $p > p_S$  and u be a classical solution of [\(1.5\)](#page-3-2). Then there exist constants  $C_0, C_1, M_0 > 0$  depending only on n and p such that the following statement holds true. Suppose that u satisfies

(1.6) 
$$
\sup_{-1 < t < 0} ||u(\cdot, t)||_{L^{q_c, \infty}(\mathbf{R}^n)} \leq M
$$

with a constant  $M \geq M_0$ . Let  $x_0 \in \mathbb{R}^n$  and set

<span id="page-3-5"></span>(1.7) 
$$
N := \int_{B(x_0, e^{e^{M C_0}})} |u(x, 0)|^{q_c} dx,
$$

<span id="page-3-6"></span>(1.8) 
$$
t_* := -M^{-p(2p+3)-1} \exp\left(-Ne^{e^{e^{M^{C_0}}}}\right).
$$

Then,

$$
(1.9) \t\t ||u||_{L^{\infty}(B(x_0, \frac{1}{4}(-t_*)^{\frac{1}{2}}) \times (\frac{1}{16}t_*, 0))} \leq C_1(-t_*)^{-\frac{1}{p-1}}.
$$

<span id="page-4-4"></span>In the next subsection, we compare our method with the previous literature.

<span id="page-4-0"></span>1.3. Comparison with previous literature. In the cases  $q_c = 1, 2$ , from [\[41\]](#page-56-4) and [\[34,](#page-55-2) Propositions 16.3, 16.3a], it is possible to obtain a single logarithmic blow-up rate of the critical norm by elementary direct arguments. Otherwise, there are very few results in the literature regarding quantitative blow-up rates of critical norms for evolution equations with a scaling symmetry. To the best of our knowledge, the first quantitative blow-up rate for a critical norm for more general nonlinearities was obtained by Merle and Raphäel [\[21\]](#page-55-19) for radial solutions of the  $L^2$  supercritical nonlinear Schrödinger equation.

In the parabolic setting, a breakthrough work of Tao [\[38\]](#page-56-0) established that if a solution v of the 3-dimensional Navier-Stokes equations in  $\mathbb{R}^n \times (-1,0)$  first loses smoothness at  $t = 0$ , then the critical  $L^3$ -norm becomes unbounded with a quantitative estimate

<span id="page-4-2"></span>(1.10) 
$$
\limsup_{t \to 0} \frac{\|v(\cdot, t)\|_{L^3(\mathbf{R}^3)}}{\left(\log \log \log \left(\frac{1}{(-t)^{c_0}}\right)\right)^{c_1}} = \infty.
$$

To show this, Tao's aim is the following:

<span id="page-4-3"></span>Tao's objective. Assume

(1.11) 
$$
A := \sup_{-1 \leq t < 0} ||v(\cdot, t)||_{L^3(\mathbf{R}^n)} < \infty.
$$

If the following statement

<span id="page-4-1"></span>(1.12) 
$$
N^{-1} \sup_{-1/2 < t < 0} \|P_N v\|_{L^3(\mathbf{R}^n)} < A^{-c} \quad \text{for all } N \ge N_*
$$

fails, find a quantitative upper bound  $N_*(A)$  for N, where  $P_N$  is a Littlewood-Paley projection on the frequency N.

Once [\(1.12\)](#page-4-1) is established for all  $N \geq N_*(A)$ , this produces quantitative estimates of  $||v||_{L^\infty(\mathbf{R}^3 \times (-1/4,0))}$  in terms of A, which can then be used to prove [\(1.10\)](#page-4-2). See the survey [\[3\]](#page-54-4) for more details.

In [\[38\]](#page-56-0), Tao achieves the above objective by

- (1) Backward propagation mechanism based on the contraposition of [\(1.12\)](#page-4-1).
- (2) Establishing regions of quantitative regularity.
- (3) Applying quantitative Carleman inequalities based on [\[9\]](#page-55-9) to the quantity  $\nabla \times v.$
- (4) Summing of scales of the  $L^3$ -norm of v at the final time over disjoint annuli to produce the upper bound  $N_*(A)$ .

For obtaining our main quantitative estimate (Proposition [5\)](#page-3-3), which in turn implies Theorems [1,](#page-2-1) [2](#page-2-4) and [3,](#page-3-0) we encounter several challenges in the above (1), (2) and (3) in the context of the energy supercritical heat equation. Let us describe these now in more details. The novelty of the results will also be described.

First, we discuss the difficulties regarding the quantitative backward propagation. For 3-dimensional Navier-Stokes equations, Tao's objective (the contraposition of [\(1.12\)](#page-4-1)) produces a sequence of the following 'frequency bubbles of concentration': For all  $n \in \mathbb{N}$ , there exists a frequency  $N_n \in (0, \infty)$  and  $(x_n, t_n) \in$  $\mathbb{R}^3 \times (-1, t_{n-1}) \subset \mathbb{R}^3 \times (-1, 0)$  such that

$$
N_n^{-1} |P_{N_n} u(x_n, t_n)| > A^{-c}
$$

with

$$
x_n = x_0 + A^C(-t_n)^{\frac{1}{2}}, \quad N_n \sim A^C(-t_n)^{-\frac{1}{2}},
$$

see [\[3\]](#page-54-4) for more details. The frequency bubbles of concentration utilizes the Duhamel formula, localized estimates of Fourier multipliers and paraproduct decompositions. It seems non-trivial to obtain such a frequency based backward propagation mechanism for non-smooth nonlinearities, such as those we are faced with for the energy supercritical nonlinear heat equations in the general case.

The first author and Prange [\[2\]](#page-54-5) extended Tao's results and proved an analogue of Theorem [2](#page-2-4) for the 3-dimensional Navier-Stokes equations by using a different objective. For brevity, we state this objective in a less general setting than in [\[2\]](#page-54-5) (same setting as Tao [\[38\]](#page-56-0)).

Barker and Prange's objective. Assume  $A := \sup_{-1 \le t \le 0} ||v(\cdot, t)||_{L^3(\mathbf{R}^n)} < \infty$ . If the following statement

<span id="page-5-0"></span>(1.13) 
$$
(-t)^{\frac{1}{2}} \int_{B(x_0, A^c(-t)^{\frac{1}{2}})} |\nabla \times v(x, t)|^2 dx \leq A^{-c}
$$

fails, find a quantitative upper bound  $t_*(A) \in (-1,0)$  for t such that necessarily  $t \le t_*(A).$ 

Once [\(1.13\)](#page-5-0) is established for all  $t_*(A) < t < 0$ , this produces quantitative estimates of  $||v||_{L^{\infty}(\mathbf{R}^3 \times (-1/4,0))}$  in terms of A. A key part in achieving Barker and Prange's objective in [\[2\]](#page-54-5) is showing 'backward propagation of vorticity concentration'. Namely, the initial concentration

<span id="page-5-1"></span>(1.14) 
$$
(-t)^{\frac{1}{2}} \int_{B(x_0, (-t)^{\frac{1}{2}})} |\nabla \times v(x, t)|^2 dx > A^{-c}
$$

propagates backwards in time and holds true for all times that are sufficiently in the past of t. Both Barker and Prange's objective and backward propagation of vorticity concentration crucially hinge on 'local-in-space smoothing' for the 3-dimensional Navier-Stokes equations. Namely, if the initial data  $v_0$  satisfies  $||v_0||_{L^3(\mathbf{R}^n)} \leq M$ and  $||v_0||_{L^6(B(0,1))} \leq N$ , then the solution v is quantitatively bounded on  $B(0, 1/2) \times$  $(0, T(M, N))$ . The proof of local-in-space smoothing crucially uses the local energy structure of the 3-dimensional Navier-Stokes equations. In particular, it is not obviously clear that it holds for other nonlinear parabolic equations such as the energy supercritical nonlinear heat equation.

To overcome these difficulties, we pursue a different strategy for producing quantitative estimates for the energy supercritical nonlinear heat equation (Proposition [5\)](#page-3-3). Our strategy is based on quantitative partial regularity over comparable time scales under the assumption [\(1.6\)](#page-3-4).

New objective. Assume that  $u$  is a solution of the energy supercritical nonlinear heat equation on  $\mathbb{R}^n \times (-1,0)$  satisfying the Lorentz norm bound [\(1.6\)](#page-3-4). Let N be defined by [\(1.7\)](#page-3-5). If the following statement

<span id="page-6-0"></span>
$$
(1.15) \qquad \qquad (-t)^{\frac{2}{p-1}-\frac{n}{2}} \int_{t}^{t/2} \int_{B(x_0,A(M,p)(-t)^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt \le M^{-c(p)}
$$

fails, find a quantitative upper bound  $t_*(M, N) \in (-1, 0)$  for t such that necessarily  $t \leq t_*(M,N).$ 

Once [\(1.15\)](#page-6-0) is established for  $t_*(M, N) < t < 0$ , this will then imply the quantitative bounds in Proposition [5.](#page-3-3) To pursue this objective, we must first quantify the  $\varepsilon$ -regularity criterion (Proposition [7\)](#page-8-2) concerning [\(1.15\)](#page-6-0) with [\(1.6\)](#page-3-4), which is accomplished using the Giga-Kohn monotonicity formula [\[13,](#page-55-1) Proposition 2.1] and Blatt-Struwe's  $\varepsilon$ -regularity criterion [\[4,](#page-54-6) Proposition 4.1]. With this in hand, we can show that the initial concentration

$$
(-t)^{\frac{2}{p-1}-\frac{n}{2}}\int_{t}^{t/2}\int_{B(x_0,A(M,p)(-t)^{\frac{1}{2}})}|u(x,t)|^{p+1}dxdt > M^{-c(p)}
$$

propagates backwards in time and holds for times that are (quantifiably) sufficiently in the past of t. This forms a crucial part of our proof of Proposition [5.](#page-3-3)

Next, we discuss the obstacle on the quantitative regions of regularity. In the works [\[1,](#page-54-7) [2,](#page-54-5) [17,](#page-55-20) [31,](#page-55-0) [38\]](#page-56-0) on the Navier-Stokes equations, a key part in the production of the quantitative estimates is the use of quantitative Carleman inequalities in Tao [\[38,](#page-56-0) Section 4]. Applying such Carleman inequalities requires the vorticity to satisfy certain differential inequalities, which requires determining regions of quantitative regularity of the velocity. In [\[38\]](#page-56-0), to apply quantitative unique continuation Carleman inequality, it is used that a solution to 3-dimensional Navier-Stokes equations satisfying

$$
\sup_{t \in I} \|v(\cdot, t)\|_{L^3(\mathbf{R}^n)} \le M
$$

for some interval I, there exists a quantifiable sub-interval  $I' \subset I$  (epoch of regularity) such that v is quantitatively bounded on  $\mathbb{R}^3 \times I'$ . This crucially uses the energy structure of the Navier-Stokes equations and the Sobolev embedding theorem in  $n = 3$ , which imply that solutions in  $\mathbb{R}^3$  with bounded energy are in subcritical spaces on many time slices. Here we refer to a Lebesgue space as 'subcritical' for a parabolic evolution equation with scaling symmetry, if it has higher integrability than the (scale-invariant) critical Lebesgue spaces. For the higher-dimensional Navier-Stokes equations  $(n \geq 4)$  with

$$
\sup_{-1
$$

it is not known if solutions possess quantitative epoch of regularity. This is also the case for solutions of the energy supercritical nonlinear heat equation with [\(1.6\)](#page-3-4).

To overcome this obstacle, we utilize ideas from Palasek [\[31\]](#page-55-0) for the higherdimensional Navier-Stokes equations. In particular, we use the quantitative spacetime partial regularity [\(1.15\)](#page-6-0) with [\(1.6\)](#page-3-4) to find quantitative space-time 'slices of regularity' for the solution. In doing this, we face an additional obstacle compared to [\[31\]](#page-55-0) in that the norm  $L^{q_c,\infty}(\mathbf{R}^n)$  in [\(1.6\)](#page-3-4) can have an equal presence at many disjoint scales. We overcome this by performing a Calderón splitting of the solution [\[6\]](#page-54-8), together with the fact that the integrability exponent  $p + 1$  in the quantitative partial regularity [\(1.15\)](#page-6-0) with [\(1.6\)](#page-3-4) is lower than the critical exponent  $q_c$  due to the supercriticality  $p > p<sub>S</sub>$ .

Once we have established quantitative backward propagation and regions of regularity, we then utilize quantitative Carleman inequalities in [\[31\]](#page-55-0) to obtain our main quantitative estimate (Proposition [5\)](#page-3-3). Due to cases involving a non-smooth nonlinearity, we are required to implement such Carleman inequalities in a lower regularity setting compared to previous works on the Navier-Stokes equations.

Remark 6. The quadruple logarithmic blow-up rate in Theorem [1](#page-2-1) is due to the quadruple exponential quantitative estimate in Proposition [5.](#page-3-3) Tao's triple logarithmic blow-up rate [\(1.10\)](#page-4-2) for the 3-dimensional Navier-Stokes equations is due to a triple exponential quantitative estimate, with the reason for the triple exponential estimate outlined in [\[38,](#page-56-0) Remark 1.5]. By comparison, in Palasek's work [\[31\]](#page-55-0) on the higher-dimensional Navier-Stokes equations and this paper, the quantitative estimates involve a further exponential loss for the following reason. For the 3-dimensional Navier-Stokes equations satisfying [\(1.11\)](#page-4-3) and [\(1.14\)](#page-5-1), quantitative unique continuation implies that for certain temporal scales  $T_1$  and sufficiently large R,

$$
\int_{-T_1}^{-T_1/2} \int_{R/2 \leq |x| \leq 2R} |\nabla \times v(x,t)|^2 dx dt \geq T_1^{\frac{1}{2}} e^{-\frac{A^c R^2}{T_1}}
$$

(see [\[3,](#page-54-4) (99)], for example). Yet in [\[31\]](#page-55-0) and our setting, the use of iterated quantitative unique continuation produces a much smaller lower bound than the above Gaussian lower bound for the  $L^2$ -norm of the solution. In both settings, this necessitates the subsequent use of far larger spatial scales for applying quantitative backward uniqueness than those used for the 3-dimensional Navier-Stokes equations. This requirement, together with pigeonhole arguments, produces an extra (quadruple) exponential in the quantitative estimates compared to the 3 dimensional Navier-Stokes equations.

<span id="page-7-0"></span>1.4. Final remarks. Our work shows that energy structure is not an essential feature when proving quantitative blow-up rates of critical norms of nonlinear parabolic equations with a scaling symmetry. To prove our results, we only required the quantitative partial regularity at comparable time scales [\(1.15\)](#page-6-0) for solutions satisfying [\(1.6\)](#page-3-4), along with the fact that regular solutions satisfy the correct differential inequality to apply quantitative Carleman inequalities. This opens the door for obtaining quantitative blow-up rates of critical norms for other nonlinear parabolic equations.

<span id="page-7-1"></span>1.5. Organization of the paper. The rest of this paper is organized as follows. In Section [2,](#page-8-1) we prove the quantitative  $\varepsilon$ -regularity theorem. As applications, in Section [3,](#page-17-0) we prepare regularity estimates concerning propagation of concentration and regions of regularity. In Section [4,](#page-28-0) we give Carleman inequalities regarding quantitative backward uniqueness and unique continuation. By combining these ingredients, we prove our main quantitative estimate (Proposition [5\)](#page-3-3) in Section [5.](#page-38-0) In Section [6,](#page-52-0) we show Theorems [1,](#page-2-1) [2](#page-2-4) and [3.](#page-3-0)

<span id="page-7-2"></span>1.6. Notation. For  $(x, t) \in \mathbb{R}^n \times \mathbb{R}$  and  $r > 0$ , we write  $B(x, r) := \{y \in \mathbb{R}^n; |x |y| < r$  and  $B(r) := B(0,r)$ . We denote by  $Q((x,t), r) := B(x,r) \times (t - r^2, t)$ and  $Q(r) := Q((0,0), r)$  the backward parabolic cylinders. Throughout this paper,  $C$  denotes positive constants depending only on  $n$  and  $p$  unless otherwise stated. To stress the dependence, we also write  $C(a, b, \ldots)$  when it depends only on a, b, . . . . Each of the constants may change from line to line during the proofs. We set  $q_c := n(p-1)/2$  and  $q_* := n(p-1)/(p+1)$ , where  $L^{q_*}$  is the scaling invariant critical space for the gradient of solutions. We denote  $C^{2,1}$  to be the space of functions which are twice continuously differentiable in the space variable and once in the time variable.

<span id="page-8-0"></span>1.7. Notions of solutions. We say that 'u is a classical solution of  $(1.1)$  with  $u_0 \in L^q(\mathbf{R}^n)$   $(q \ge 1)$  if u belongs to  $C([0,T); L^q(\mathbf{R}^n)) \cap L^{\infty}_{loc}((0,T); L^{\infty}(\mathbf{R}^n)) \cap$  $C^{2,1}(\mathbf{R}^n \times (0,T))$  and satisfies [\(1.1\)](#page-1-2). Such a solution exists uniquely for any  $u_0 \in$  $L^{q_c}(\mathbf{R}^n)$  with  $q_c > 1$ , see [\[5,](#page-54-2) [39\]](#page-56-5) and [\[34,](#page-55-2) Remark 15.4 (i)]. We note that Theorems [1](#page-2-1) and [2](#page-2-4) can be applied to classical solutions of [\(1.1\)](#page-1-2) with  $u_0 \in L^{q_c}(\mathbf{R}^n)$ .

By a classical solution of  $(1.1)$  (resp.  $(1.5)$ ) without mentioning the initial data, we mean a function in  $C^{2,1}(\mathbf{R}^n \times (0,T))$  (resp.  $C^{2,1}(\mathbf{R}^n \times (-1,0]))$  without specifying the initial data. In particular, we do not impose  $C([0,T); L^q(\mathbf{R}^n))$   $(q \ge 1)$ or  $L^{\infty}_{loc}((0,T);L^{\infty}(\mathbf{R}^n))$ . We can apply Theorem [3](#page-3-0) and Proposition [5](#page-3-3) to classical solutions without mentioning the initial data.

## 2. QUANTITATIVE  $\varepsilon$ -REGULARITY

<span id="page-8-1"></span>Let u be a classical solution of  $(1.5)$  satisfying the Lorentz norm bound  $(1.6)$ . The goal of this section is to show the following quantitative  $\varepsilon$ -regularity result.

<span id="page-8-2"></span>**Proposition 7** (Quantitative  $\varepsilon$ -regularity). Assume  $p > p<sub>S</sub>$  and [\(1.6\)](#page-3-4) with a constant  $M > 1$ . Then there exist constants  $C > 0$  and  $0 < \varepsilon_1 < 1$  depending only on n and p such that the following holds for any  $0 < \varepsilon < \varepsilon_0 := M^{-2(p+1)^2} \varepsilon_1^{2(p+1)}$ : If

<span id="page-8-3"></span>
$$
(2.1) \qquad \begin{cases} \delta^{\frac{4}{p-1}-n} \int_{t_0-\delta^2}^{t_0-\delta^2/2} \int_{B(x_0,A\delta)} |u(x,t)|^{p+1} dx dt \leq \varepsilon \quad \text{for some } x_0 \in \mathbf{R}^n \\ -1/16 < t_0 \leq 0, 0 < \delta < 1/4 \text{ and } A > (24\log(M^{p+1}/\varepsilon))^{1/2}, \end{cases}
$$

then

$$
||u||_{L^{\infty}(Q((x_0,t_0),\delta/4))} \leq CM \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{2}{p-1}},
$$
  

$$
||\nabla u||_{L^{\infty}(Q((x_0,t_0),\delta/4))} \leq CM \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{p+1}{p-1}}.
$$

We prove Proposition [7](#page-8-2) based on the analysis of the Giga–Kohn weighted energy. For  $\tilde{x} \in \mathbb{R}^n$ ,  $-1 < t < t \leq 0$ , we define the energy  $E_{(\tilde{x}, \tilde{t})}$  by

$$
E_{(\tilde{x},\tilde{t})}(t) := (\tilde{t} - t)^{\frac{p+1}{p-1}} \int_{\mathbf{R}^n} \left( \frac{1}{2} |\nabla u(x,t)|^2 - \frac{1}{p+1} |u(x,t)|^{p+1} + \frac{1}{2(p-1)(\tilde{t}-t)} |u(x,t)|^2 \right) K_{(\tilde{x},\tilde{t})}(x,t) dx,
$$

where  $K$  is the backward heat kernel given by

$$
K_{(\tilde{x},\tilde{t})}(x,t) := (\tilde{t} - t)^{-\frac{n}{2}} e^{-\frac{|x - \tilde{x}|^2}{4(\tilde{t} - t)}}.
$$

We recall the backward similarity variables  $(\eta, \tau)$  by

$$
\eta := \frac{x - \tilde{x}}{(\tilde{t} - t)^{1/2}}, \quad \tau := -\log(\tilde{t} - t).
$$

,

Then the backward rescaled solution  $w_{(\tilde{x}, \tilde{t})}$  is defined by

$$
w_{(\tilde{x},\tilde{t})}(\eta,\tau) := e^{-\frac{1}{p-1}\tau}u(\tilde{x} + e^{-\frac{1}{2}\tau}\eta,\tilde{t} - e^{-\tau}) = (\tilde{t} - t)^{\frac{1}{p-1}}u(x,t).
$$

The corresponding Giga–Kohn energy  $\mathcal{E}_{(\tilde{x},\tilde{t})}$  is given by

<span id="page-9-1"></span>(2.2) 
$$
\mathcal{E}_{(\tilde{x},\tilde{t})}(\tau) := \int_{\mathbf{R}^n} \left( \frac{1}{2} |\nabla w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^2 - \frac{1}{p+1} |w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^{p+1} + \frac{1}{2(p-1)} |w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^2 \right) \rho(\eta) d\eta,
$$

where  $\rho(\eta) := e^{-|\eta|^2/4}$ . We note that

<span id="page-9-7"></span>(2.3) 
$$
E_{(\tilde{x},\tilde{t})}(t) = \mathcal{E}_{(\tilde{x},\tilde{t})}(\tau) \text{ with } \tau = -\log(\tilde{t}-t).
$$

Moreover, by setting  $\tau_0 := -\log(\tilde{t} + 1)$ , we see that w satisfies

<span id="page-9-0"></span>(2.4) 
$$
\rho w_{\tau} = \nabla \cdot (\rho \nabla w) - \frac{1}{p-1} w \rho + |w|^{p-1} w \rho \quad \text{in } \mathbb{R}^n \times (\tau_0, \infty).
$$

Here and below, we often suppress the subscript  $(\tilde{x}, \tilde{t})$  if it is clear from the context.

<span id="page-9-2"></span>We recall the Giga–Kohn monotonicity formula [\[12,](#page-55-21) [13\]](#page-55-1).

Lemma 8. For any  $\tilde{x} \in \mathbb{R}^n$ ,  $-1 < t' < t < \tilde{t} \leq 0$  and  $\tau = -\log(\tilde{t} - t)$ ,

<span id="page-9-6"></span>
$$
(2.5) \quad \frac{d\mathcal{E}_{(\tilde{x},\tilde{t})}}{d\tau}(\tau) = -\int_{\mathbf{R}^n} |\partial_\tau w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^2 \rho(\eta) d\eta,
$$

<span id="page-9-3"></span>
$$
(2.6) \quad E_{(\tilde{x},\tilde{t})}(t) = E_{(\tilde{x},\tilde{t})}(t') - \int_{t'}^{t} \int_{\mathbf{R}^n} (\tilde{t} - s)^{\frac{2}{p-1} - 1} |S_{(\tilde{x},\tilde{t})}(x,s)|^2 K_{(\tilde{x},\tilde{t})}(x,s) dx ds,
$$

$$
S_{(\tilde{x},\tilde{t})}(x,t) := \frac{1}{p-1} u(x,t) + \frac{1}{2} (x - \tilde{x}) \cdot \nabla u(x,t) - (\tilde{t} - t) u_t(x,t).
$$

In particular,  $\mathcal{E}_{(\tilde{x},\tilde{t})}(\tau)$  and  $E_{(\tilde{x},\tilde{t})}(t)$  are nonincreasing in  $\tau$  and t, respectively.

Proof. See [\[13,](#page-55-1) Proposition 2.1].

$$
\Box
$$

The monotonicity guarantees the nonnegativity of  $E_{(\tilde{x},\tilde{t})}$ . Moreover, the critical norm bound [\(1.6\)](#page-3-4) implies the uniform boundedness of  $E_{(\tilde{x},\tilde{t})}$ .

<span id="page-9-4"></span>**Lemma 9.** There exists  $C > 0$  depending only on n and p such that for any  $\tilde{x} \in \mathbb{R}^n$ and  $-1/4 \leq t < \tilde{t} \leq 0$ ,

$$
0 \le E_{(\tilde{x}, \tilde{t})}(t) \le CM^{2p}.
$$

Proof. From [\(2.4\)](#page-9-0) and the integration by parts, it follows that

$$
\frac{1}{2}\frac{d}{d\tau}\int_{\mathbf{R}^n}|w|^2\rho d\eta = \int_{\mathbf{R}^n}\left(-|\nabla w|^2 - \frac{1}{p-1}|w|^2 + |w|^{p+1}\right)\rho d\eta.
$$

Then, [\(2.2\)](#page-9-1) shows that

(2.7) 
$$
\frac{1}{2}\frac{d}{d\tau}\int_{\mathbf{R}^n}|w|^2\rho d\eta = -2\mathcal{E}(\tau) + \frac{p-1}{p+1}\int_{\mathbf{R}^n}|w|^{p+1}\rho d\eta.
$$

This together with Lemma [8](#page-9-2) and the Hölder inequality yields

<span id="page-9-5"></span>
$$
\frac{1}{2}\frac{d}{d\tau}\int_{\mathbf{R}^n}|w(\eta,\tau)|^2\rho(\eta)d\eta\geq-2\mathcal{E}(\tau')+C\left(\int_{\mathbf{R}^n}|w(\eta,\tau)|^2\rho(\eta)d\eta\right)^{\frac{p+1}{2}}
$$

for all  $\tau_0 < \tau' < \tau < \infty$  with  $\tau_0 := -\log(\tilde{t} + 1)$ . This inequality and a contradiction argument show that  $E_{(\tilde{x},\tilde{t})}(t) = \mathcal{E}_{(\tilde{x},\tilde{t})}(\tau) \ge 0$  for any  $\tilde{x} \in \mathbb{R}^n$  and  $-1 \le t < \tilde{t} \le 0$ . For details, see [\[13,](#page-55-1) Propositions 2.1, 2.2] and [\[34,](#page-55-2) Proposition 23.8].

We show the upper bound, which hinges on the strategy in [\[24,](#page-55-7) Lemma 2.4]. From  $(2.6)$  and  $(1.6)$ , it follows that

<span id="page-10-0"></span>
$$
E_{(\tilde{x},\tilde{t})}(t) \le E_{(\tilde{x},\tilde{t})}(-1/4)
$$
  
(2.8) 
$$
\le C(\tilde{t} + 1/4)^{\frac{p+1}{p-1}} \int_{\mathbf{R}^n} \left( |\nabla u(x, -1/4)|^2 + \frac{|u(x, -1/4)|^2}{\tilde{t} + 1/4} \right) K_{(\tilde{x},\tilde{t})}(x, -1/4) dx
$$
  

$$
\le CM^2 + C(\tilde{t} + 1/4)^{\frac{p+1}{p-1}} \int_{\mathbf{R}^n} |\nabla u(x, -1/4)|^2 K_{(\tilde{x},\tilde{t})}(x, -1/4) dx
$$

for  $-1/4 \le t < \tilde{t} \le 0$ . By using the Duhamel formula for solutions of the nonlinear heat equation, we have an integral equation for  $u(x, -1/4)$ . Differentiating the integral equation and using  $|\nabla G| \leq CK_1$  with the heat kernel  $G(x,t) :=$  $(4\pi t)^{-n/2}e^{-|x|^2/(4t)}$  and  $K_1(x,t) := t^{-(n+1)/2}e^{-|x|^2/(8t)}$ , we see that

$$
|\nabla u(x, -1/4)| \le C \int_{\mathbf{R}^n} K_1(x - y, 1/4)|u(y, -1/2)| dy
$$
  
+  $C \int_{-1/2}^{-1/4} \int_{\mathbf{R}^n} K_1(x - y, -1/4 - s)|u(y, s)|^p dy ds$   
=:  $CU_1(x) + CU_2(x)$ .

The Hölder inequality in the Lorentz spaces (see [\[29,](#page-55-22) Section IV]) gives

$$
U_1(x) \le C \left\| |u(\cdot, -1/2)| e^{-\frac{|x-1|^2}{2}} \right\|_{L^1(\mathbf{R}^n)}
$$
  
 
$$
\le C \|u(\cdot, -1/2)\|_{L^{q_c,\infty}(\mathbf{R}^n)} \left\| e^{-\frac{|x-1|^2}{2}} \right\|_{L^{\frac{q_c}{q_c-1},1}(\mathbf{R}^n)} \le CM.
$$

Again by the Hölder inequality with  $q_* := n(p-1)/(p+1) > 2$  for  $p > p<sub>S</sub>$ , we have

$$
\int_{\mathbf{R}^n} |\nabla u(x, -1/4)|^2 K_{(\tilde{x}, \tilde{t})}(x, -1/4) dx \le C \int_{\mathbf{R}^n} (|U_1|^2 + |U_2|^2) K_{(\tilde{x}, \tilde{t})}(x, -1/4) dx
$$
\n
$$
\le CM^2 + C(\tilde{t} + 1/4)^{-\frac{n}{2}} \left\| |U_2|^2 e^{-\frac{|1-\tilde{x}|^2}{4(\tilde{t}+1/4)}} \right\|_{L^1(\mathbf{R}^n)}
$$
\n
$$
\le CM^2 + C(\tilde{t} + 1/4)^{-\frac{n}{2}} \| |U_2|^2 \|_{L^{\frac{q_*}{2}, \infty}(\mathbf{R}^n)} \left\| e^{-\frac{|1-\tilde{x}|^2}{4(\tilde{t}+1/4)}} \right\|_{L^{\frac{q_*}{q_*-2}, 1}(\mathbf{R}^n)}
$$
\n
$$
\le CM^2 + C(\tilde{t} + 1/4)^{-\frac{p+1}{p-1}} \| U_2 \|_{L^{q_*,\infty}(\mathbf{R}^n)}^2 \le C(\tilde{t} + 1/4)^{-\frac{p+1}{p-1}} (M^2 + \| U_2 \|_{L^{q_*,\infty}(\mathbf{R}^n)}^2).
$$

Recall [\(2.8\)](#page-10-0). Then, for  $-1/4 \le t < \tilde{t} \le 0$ , we also have

<span id="page-10-1"></span>
$$
\begin{aligned} (2.9) \qquad E_{(\tilde{x},\tilde{t})}(t) &\leq CM^2 + C(\tilde{t} + 1/4)^{\frac{p+1}{p-1}} \left( M^2 + (\tilde{t} + 1/4)^{-\frac{p+1}{p-1}} \| U_2 \|_{L^{q_*,\infty}(\mathbf{R}^n)}^2 \right) \\ &\leq CM^2 + C \| U_2 \|_{L^{q_*,\infty}(\mathbf{R}^n)}^2. \end{aligned}
$$

We estimate  $U_2$  by a modification of [\[22,](#page-55-23) Theorem 18.1]. By the change of variables in the definition of  $U_2$ , we set

$$
U_2(x) = \int_0^\infty S(x, s)ds,
$$
  
\n
$$
S(x, s) := \chi_{(0, 1/4)}(s) \int_{\mathbf{R}^n} K_1(x - y, s) |u(y, -1/4 - s)|^p dy.
$$

For  $\lambda > 0$  and  $\tau > 0$ , define  $D_{\lambda} := \{x \in \mathbb{R}^n : U_2(x) > \lambda\}$  and

$$
U_2(x) = \left(\int_0^{\tau} + \int_{\tau}^{\infty}\right)S(x,s)ds =: V_{\tau}(x) + W_{\tau}(x).
$$

Let us estimate the Lebesgue measure  $|D_\lambda|$ . By the Hölder inequality, we have

$$
S(x,s) \leq Cs^{-\frac{n}{2}-\frac{1}{2}}\chi_{(0,1/4)}(s) \left\| |u(\cdot, -1/4-s)|^p e^{-\frac{|x-1|^2}{8s}} \right\|_{L^1(\mathbf{R}^n)}
$$
  
\n
$$
\leq Cs^{-\frac{n}{2}-\frac{1}{2}}\chi_{(0,1/4)}(s) \| |u(\cdot, -1/4-s)|^p \|_{L^{\frac{q_c}{p},\infty}(\mathbf{R}^n)} \left\| e^{-\frac{|x-1|^2}{8s}} \right\|_{L^{\frac{q_c}{q_c-p},1}(\mathbf{R}^n)}
$$
  
\n
$$
\leq Cs^{-\frac{np}{2q_c}-\frac{1}{2}}\chi_{(0,1/4)}(s) \| u(\cdot, -1/4-s) \|_{L^{q_c,\infty}(\mathbf{R}^n)}^p \leq CM^p s^{-\frac{p}{p-1}-\frac{1}{2}}
$$

for any  $s > 0$ . Then,

<span id="page-11-0"></span>
$$
W_{\tau}(x) \leq CM^p \int_{\tau}^{\infty} s^{-\frac{p}{p-1} - \frac{1}{2}} ds = C'M^p \tau^{-\frac{p+1}{2(p-1)}},
$$

where  $C' > 0$  is independent of t. For  $\lambda > 0$ , we choose  $\tau$  such that

(2.10) 
$$
C'M^p \tau^{-\frac{p+1}{2(p-1)}} = \frac{\lambda}{2}.
$$

Then  $W_{\tau} \leq \lambda/2$ , and so  $\tilde{D}_{\lambda} := \{x \in \mathbb{R}^n; V_{\tau}(x) > \lambda/2\}$  satisfies  $D_{\lambda} \subset \tilde{D}_{\lambda}$ . From the change of variables, it follows that

$$
S(x,s) = \int_{\mathbf{R}^n} K_1(z,s) \chi_{(0,1/4)}(s) |u(z+x,-1/4-s)|^p dz
$$
  
 
$$
\leq C s^{-\frac{n}{2}-\frac{1}{2}} \int_{\mathbf{R}^n} \chi_{(0,1/4)}(s) |u(z+x,-1/4-s)|^p e^{-\frac{|z|^2}{8s}} dz
$$

and so by O'Neil's convolution inequality [\[29,](#page-55-22) Theorem 2.6]

$$
||S(\cdot,s)||_{L^{\frac{qc}{p},\infty}(\mathbf{R}^n)} \leq Cs^{-\frac{n}{2}-\frac{1}{2}} \int_{\mathbf{R}^n} \chi_{(0,1/4)}(s) ||u(\cdot,-1/4-s)|^p ||_{L^{\frac{qc}{p},\infty}(\mathbf{R}^n)} e^{-\frac{|z|^2}{8s}} dz
$$
  
 
$$
\leq CM^p s^{-\frac{n}{2}-\frac{1}{2}} \int_{\mathbf{R}^n} e^{-\frac{|z|^2}{8s}} dz \leq CM^p s^{-\frac{1}{2}}
$$

for any  $s > 0$ . Thus, by Minkowski's inequality for Lorentz spaces [\[18,](#page-55-24) Lemma 1],

$$
\|V_{\tau}\|_{L^{\frac{q_c}{p},\infty}(\mathbf{R}^n)}\leq \int_0^\tau \|S(\cdot,s)\|_{L^{\frac{q_c}{p},\infty}(\mathbf{R}^n)}ds\leq CM^p\tau^{\frac{1}{2}}.
$$

This together with the Hölder inequality for the Lorentz spaces shows that  $\,$ 

$$
\int_{\tilde{D}_{\lambda}} V_{\tau}(x) dx \leq C \|\chi_{\tilde{D}_{\lambda}}\|_{L^{\frac{qc}{qc-p},1}(\mathbf{R}^n)} \|V_{\tau}\|_{L^{\frac{qc}{p},\infty}(\mathbf{R}^n)} \leq CM^p |\tilde{D}_{\lambda}|^{1-\frac{p}{qc}} \tau^{\frac{1}{2}}.
$$

On the other hand,  $\int_{\tilde{D}_{\lambda}} V_{\tau}(x)dx \geq (\lambda/2)|\tilde{D}_{\lambda}|$ . By  $D_{\lambda} \subset \tilde{D}_{\lambda}$  and  $(2.10)$ , we obtain

$$
|D_\lambda| \leq |\tilde{D}_\lambda| \leq C \lambda^{-\frac{q_c}{p}} M^{q_c} \tau^{\frac{q_c}{2p}} = C M^{pq_*} \lambda^{-q_*},
$$

and so  $\lambda |\{x \in \mathbf{R}^n; U_2(x) > \lambda\}|^{1/q_*} \leq CM^p$  for  $\lambda > 0$ . Hence we obtain

$$
||U_2||_{L^{q_*,\infty}(\mathbf{R}^n)} \le ||U||_{L^{q_*,\infty}(\mathbf{R}^n)} \le CM^p.
$$

This together with [\(2.9\)](#page-10-1) and  $M > 1$  gives the desired inequality.

We prove the following lemma based on Blatt-Struwe's  $\varepsilon$ -regularity criterion [\[4,](#page-54-6) Proposition 4.1].

<span id="page-12-2"></span>**Lemma 10.** There exist  $0 < \tilde{\varepsilon}_1 < 1$  and  $C > 0$  depending only on n and p such that the following holds for any  $0 < \tilde{\varepsilon} \leq \tilde{\varepsilon}_1$ : If there exists  $0 < \delta < 1/2$  such that

$$
(2.11) \tI_r(\tilde{x}, \tilde{t}) := (r/2)^{\frac{4}{p-1} - n} \int_{\tilde{t} - r^2/4}^{\tilde{t} - r^2/16} \int_{B(\tilde{x}, r/2)} |u(x, t)|^{p+1} dx dt \le \tilde{\varepsilon}
$$

for any  $r > 0$ ,  $\tilde{x} \in \mathbb{R}^n$  and  $-1 < \tilde{t} \leq 0$  satisfying  $Q((\tilde{x}, \tilde{t}), r) \subset Q(\delta/2)$ , then

<span id="page-12-0"></span>
$$
||u||_{L^{\infty}(Q(\delta/4))} \leq C\tilde{\varepsilon}^{\frac{1}{p+1}}\delta^{-\frac{2}{p-1}}, \quad ||\nabla u||_{L^{\infty}(Q(\delta/4))} \leq C\tilde{\varepsilon}^{\frac{1}{p+1}}\delta^{-\frac{p+1}{p-1}}.
$$

*Proof.* Let  $\tilde{\varepsilon}_1 > 0$  be a constant chosen later and let  $0 < \tilde{\varepsilon} < \tilde{\varepsilon}_1$ . Set  $v(y, s) :=$  $\delta^{2/(p-1)}u(\delta y,\delta^2 s)$ . Note that  $v_t = \Delta v + |v|^{p-1}v$  in  $\mathbb{R}^n \times (-4,0)$ , since  $-4\delta^2 > -1$ . If  $Q((\tilde{x}, \tilde{t}), r) \subset Q(1/2)$  and  $(r/4)^2 \leq -\tilde{t}$ , then  $Q((\tilde{x}, \tilde{t} + (r/4)^2), r) \subset Q(1/2)$ , and so the change of variables and [\(2.11\)](#page-12-0) show that

<span id="page-12-1"></span>
$$
(r/2)^{\frac{4}{p-1}-n} \int_{\tilde{t}-r^2/16}^{\tilde{t}} \int_{B(\tilde{x},r/2)} |v(y,s)|^{p+1} dyds
$$
  

$$
\leq (r/2)^{\frac{4}{p-1}-n} \int_{(\tilde{t}+r^2/16)-r^2/4}^{(\tilde{t}+r^2/16)-r^2/16} \int_{B(\tilde{x},r/2)} |v(y,s)|^{p+1} dyds \leq \tilde{\varepsilon}.
$$

Let  $(x', t') \in Q(1/3)$  and  $\lambda := (-t')^{1/2}$ . Then  $\tilde{v}(x, t) := \lambda^{2/(p-1)} v(\lambda x + x', \lambda^2 t + t')$ satisfies  $\tilde{v}_t = \Delta \tilde{v} + |\tilde{v}|^{p-1}\tilde{v}$  in  $\mathbb{R}^n \times (-4/\lambda^2, 0)$ . If  $B((\tilde{x} - x')/\lambda, r/\lambda) \subset B(1/2)$  and  $((\tilde{t}-r^2)/\lambda^2, \tilde{t}/\lambda^2) \subset (-1/4, 0)$ , then  $Q((\tilde{x}, \tilde{t}+t'), r) \subset Q(1/2)$  and  $(r/4)^2 \leq -(\tilde{t}+t')$ . Therefore, [\(2.12\)](#page-12-1) shows that

$$
(r/4\lambda)^{\frac{4}{p-1}-n} \int_{(\tilde{t}-(r/4)^2)/\lambda^2}^{\tilde{t}/\lambda^2} \int_{B((\tilde{x}-x')/\lambda, r/4\lambda)} |\tilde{v}(x,t)|^{p+1} dx dt
$$
  
=  $(r/4)^{\frac{4}{p-1}-n} \int_{\tilde{t}+t'-(r/4)^2}^{\tilde{t}+t'} \int_{B(\tilde{x},r/4)} |v(y,s)|^{p+1} dy ds \le C\tilde{\varepsilon}$ 

if  $Q(((\tilde{x} - x')/\lambda, \tilde{t}/\lambda^2), r/\lambda) \subset Q(1/2)$ . Replacing  $(\tilde{x}, \tilde{t})$  and r with  $(\lambda \tilde{x} + x', \lambda^2 \tilde{t})$ and  $\lambda r$ , respectively, we see that

$$
(r/4)^{\frac{4}{p-1}-n} \iint_{Q((\tilde{x},\tilde{t}),r/4)} |\tilde{v}(x,t)|^{p+1} dx dt \le C\tilde{\varepsilon}
$$

if  $Q((\tilde{x},\tilde{t}),r) \subset Q(1/2)$ . Hence  $\|\tilde{v}\|_{M^{p+1,2(p+1)/(p-1)}(Q(1/3))} \leq C\tilde{\varepsilon}^{1/(p+1)}$ , where  $\|\cdot\|$  $\|M^{p+1,2(p+1)/(p-1)}(Q(1/3))$  is the parabolic Morrey norm on the backward parabolic cylinder  $Q(1/3)$ , see [\[4,](#page-54-6) Section 2] for the definition. Taking  $\tilde{\varepsilon}_1$  (>  $\tilde{\varepsilon}$ ) sufficiently small depending only on  $n$  and  $p$ , we apply [\[4,](#page-54-6) Proposition 4.1] to see that

$$
\|\tilde{v}\|_{L^{\infty}(Q(1/4))} + \|\nabla \tilde{v}\|_{L^{\infty}(Q(1/5))} \leq C \|\tilde{v}\|_{M^{p+1, \frac{2(p+1)}{p-1}}(Q(1/3))} \leq C \tilde{\varepsilon}^{\frac{1}{p+1}},
$$

and so

$$
\lambda^{\frac{2}{p-1}}|v(\lambda x + x', \lambda^2 t + t')| + \lambda^{\frac{p+1}{p-1}}|\nabla v(\lambda x + x', \lambda^2 t + t')| \le C\tilde{\varepsilon}^{\frac{1}{p+1}}
$$

for  $(x, t) \in Q(1/5)$ . Letting  $(x, t) \to (0, 0)$  and using  $\lambda = (-t')^{1/2}$  yield  $|v(x', t')| \le$  $C\tilde{\epsilon}^{1/(p+1)}(-t')^{-1/(p-1)}$  and  $|\nabla v(x',t')| \leq C\tilde{\epsilon}^{1/(p+1)}(-t')^{-(p+1)/(2(p-1))}$ . Recall that  $(x', t') \in Q(1/3)$ . Hence we obtain

<span id="page-13-0"></span> $|v(y, s)| \leq C \tilde{\varepsilon}^{\frac{1}{p+1}}(-s)^{-\frac{1}{p-1}}$  for  $y \in B(1/3), -1/9 \leq s < 0$ ,

<span id="page-13-2"></span>
$$
(2.14) \qquad |\nabla v(y,s)| \le C\tilde{\varepsilon}^{\frac{1}{p+1}}(-s)^{-\frac{p+1}{2(p-1)}} \qquad \text{for } y \in B(1/3), \ -1/9 \le s < 0.
$$

We give an  $L^{\infty}$  bound of v based on the argument of [\[14,](#page-55-25) Theorem 2.1]. Let  $\phi \in C_0^{\infty}(\mathbf{R}^n)$  satisfy  $\phi = 1$  in  $B(7/24)$ ,  $\phi = 0$  in  $\mathbf{R}^n \setminus B(1/3)$  and  $0 \le \phi \le 1$ . Set  $w := \phi v$ . Then w satisfies

$$
w_t - \Delta w = |v|^{p-1}w - 2\nabla \cdot (v\nabla \phi) + v\Delta \phi.
$$

From the Duhamel formula and convolution inequalities on the heat kernel G and  $\nabla G$ , together with [\(2.13\)](#page-13-0) and  $(t-s)^{-1/2} > 1$  for  $-1/9 < s < t < 0$ , it follows that

<span id="page-13-1"></span>
$$
||w(\cdot,t)||_{L^{\infty}(\mathbf{R}^{n})}
$$
  
\n
$$
\leq ||w(\cdot,-1/9)||_{L^{\infty}(\mathbf{R}^{n})} + \int_{-1/9}^{t} ||v(\cdot,s)||_{L^{\infty}(B(1/3))}^{p-1} ||w(\cdot,s)||_{L^{\infty}(\mathbf{R}^{n})} ds
$$
  
\n
$$
+ C \int_{-1/9}^{t} (t-s)^{-\frac{1}{2}} ||v(\cdot,s)||_{L^{\infty}(B(1/3))} ds
$$
  
\n
$$
\leq C \tilde{\varepsilon}^{\frac{1}{p+1}} \left(1 + \int_{-1/9}^{t} (t-s)^{-\frac{1}{2}}(-s)^{-\frac{1}{p-1}} ds\right)
$$
  
\n
$$
+ C \tilde{\varepsilon}^{\frac{p-1}{p+1}} \int_{-1/9}^{t} (-s)^{-1} ||w(\cdot,s)||_{L^{\infty}(\mathbf{R}^{n})} ds.
$$

Similar computations in [\[14,](#page-55-25) Lemma 2.2] show that

<span id="page-13-3"></span>
$$
(2.16) \qquad \int_{-1/9}^{t} (t-s)^{-\frac{1}{2}}(-s)^{-\frac{1}{p-1}} ds \le \begin{cases} C\left(1 + \log \frac{1}{-t}\right) & \text{if } \frac{1}{2} \ge \frac{1}{p-1}, \\ C(-t)^{-(\frac{1}{p-1}-\frac{1}{2})} & \text{if } \frac{1}{2} < \frac{1}{p-1}. \end{cases}
$$

By setting

$$
\alpha := \begin{cases} \frac{1}{4(p-1)} & \text{if } \frac{1}{2} \ge \frac{1}{p-1}, \\ \frac{1}{p-1} - \frac{1}{2} & \text{if } \frac{1}{2} < \frac{1}{p-1}, \end{cases}
$$

we see that

$$
\int_{-1/9}^{t} (t-s)^{-\frac{1}{2}}(-s)^{-\frac{1}{p-1}} ds \le C(-t)^{-\alpha}.
$$

Then there exist  $C_1, C_2 > 0$  depending only on n and p such that

$$
||w(\cdot,t)||_{L^{\infty}(\mathbf{R}^n)} \leq C_1 \tilde{\varepsilon}^{\frac{1}{p+1}}(-t)^{-\alpha} + C_2 \tilde{\varepsilon}^{\frac{p-1}{p+1}} \int_{-1/9}^t (-s)^{-1} ||w(\cdot,s)||_{L^{\infty}(\mathbf{R}^n)} ds.
$$

We fix  $\tilde{\varepsilon}_1$  such that  $C_2 \tilde{\varepsilon}_1^{(p-1)/(p+1)} \le \min((4(p-1))^{-1}, 4^{-1}).$ 

Since  $t \mapsto (-t)^{-\alpha}$  is nondecreasing, Gronwall's inequality yields

$$
||w(\cdot,t)||_{L^{\infty}(\mathbf{R}^n)} \leq C_1 \tilde{\varepsilon}^{\frac{1}{p+1}} (-t)^{-\alpha} \exp\left(C_2 \tilde{\varepsilon}^{\frac{p-1}{p+1}} \int_{-1/9}^t (-s)^{-1} ds\right)
$$
  

$$
\leq C_1 \tilde{\varepsilon}^{\frac{1}{p+1}} (-t)^{-\alpha} \exp\left(C_2 \tilde{\varepsilon}^{\frac{p-1}{p+1}} \log \frac{1}{-t}\right)
$$
  

$$
= C_1 \tilde{\varepsilon}^{\frac{1}{p+1}} (-t)^{-\alpha - C_2 \tilde{\varepsilon}^{\frac{p-1}{p+1}}}_{1} \leq \begin{cases} C_1 \tilde{\varepsilon}^{\frac{1}{p+1}} (-t)^{-\frac{1}{2(p-1)}} & \text{if } \frac{1}{2} \geq \frac{1}{p-1}, \\ C_1 \tilde{\varepsilon}^{\frac{1}{p+1}} (-t)^{-\frac{1}{p-1}+\frac{1}{4}} & \text{if } \frac{1}{2} < \frac{1}{p-1}. \end{cases}
$$

We observe that the above estimate is an improvement of  $(2.13)$ . In the first case  $(1/2 \ge 1/(p-1))$ , we can bootstrap the above argument one more time to get the required boundedness of v on  $Q(1/4)$ . In the second case  $(1/2 < 1/(p-1))$ , we can apply finitely many bootstraps to enter a similar scenario to the first case. In all scenarios, we get

$$
||v||_{L^{\infty}(Q(1/4))} \leq ||w||_{L^{\infty}(\mathbf{R}^n \times (-1/9,0))} \leq C \tilde{\varepsilon}^{\frac{1}{p+1}},
$$

and so  $||u||_{L^{\infty}(Q(\delta/4))} \leq C\tilde{\varepsilon}^{1/(p+1)}\delta^{-2/(p-1)}$ .

Let us next consider an  $L^{\infty}$  bound of  $\nabla v$ . We note that *i*-th derivative  $v_{y_i}$ satisfies  $\partial_t v_{y_i} = \Delta v_{y_i} + p|v|^{p-1}v_{y_i}$ . Setting  $\tilde{w} := \phi v_{y_i}$  gives

$$
\tilde{w}_t - \Delta \tilde{w} = p|v|^{p-1}\tilde{w} - 2\nabla \cdot (v_{y_i}\nabla \phi) + v_{y_i}\Delta \phi.
$$

Similar computations to [\(2.15\)](#page-13-1) with [\(2.14\)](#page-13-2) show that

$$
\begin{split} &\|\tilde{w}(\cdot,t)\|_{L^{\infty}(\mathbf{R}^n)}\\ &\leq \|\tilde{w}(\cdot,-1/9)\|_{L^{\infty}(\mathbf{R}^n)} + p \int_{-1/9}^t \|v(\cdot,s)\|_{L^{\infty}(B(1/3))}^{p-1} \|\tilde{w}(\cdot,s)\|_{L^{\infty}(\mathbf{R}^n)} ds\\ &+ C \int_{-1/9}^t (t-s)^{-\frac{1}{2}} \|v_{y_i}(\cdot,s)\|_{L^{\infty}(B(1/3))} ds\\ &\leq C \tilde{\varepsilon}^{\frac{1}{p+1}} \left(1+\int_{-1/9}^t (t-s)^{-\frac{1}{2}}(-s)^{-\frac{p+1}{2(p-1)}} ds\right)\\ &+ C \tilde{\varepsilon}^{\frac{p-1}{p+1}} \int_{-1/9}^t (-s)^{-1} \|\tilde{w}(\cdot,s)\|_{L^{\infty}(\mathbf{R}^n)} ds. \end{split}
$$

Then by an analog of [\(2.16\)](#page-13-3), we have

$$
\|\tilde{w}(\cdot,t)\|_{L^{\infty}(\mathbf{R}^n)} \leq C\tilde{\varepsilon}^{\frac{1}{p+1}}(-t)^{-\frac{1}{p-1}} + C\tilde{\varepsilon}^{\frac{p-1}{p+1}} \int_{-1/9}^t (-s)^{-1} \|\tilde{w}(\cdot,s)\|_{L^{\infty}(\mathbf{R}^n)} ds.
$$

From similar computations as for w with  $\tilde{\varepsilon}_1$  ( $>\tilde{\varepsilon}$ ) replaced by a smaller constant depending only on n and p if necessary, it follows that  $||v_{y_i}||_{L^{\infty}(Q(1/4))} \leq C\tilde{\varepsilon}^{1/(p+1)}$ and that  $\|\nabla u\|_{L^{\infty}(Q(\delta/4))} \leq C\tilde{\varepsilon}^{1/(p+1)}\delta^{-(p+1)/(p-1)}$ . The proof is complete.  $\Box$ 

**Remark 11.** Note that with a slight adjustment to the test function  $\phi$  and the use of another test function  $\tilde{\phi}$  in defining  $\tilde{w} := \tilde{\phi}v_{y_i}$ , we could get that v is bounded in  $Q(1/3)$  and could then infer that for  $-1/9 < t < 0$ ,

$$
\|\tilde{w}(\cdot,t)\|_{L^{\infty}(\mathbf{R}^n)} \leq C\tilde{\varepsilon}^{\frac{1}{p+1}}(-t)^{-\frac{1}{p-1}}.
$$

We could subsequently bootstrap this to get boundedness of  $\tilde{w}$  in  $Q(1/4)$ . We do not pursue this for notational convenience.

We prove the desired quantitative  $\varepsilon$ -regularity. This is achieved by using the Giga-Kohn monotonicity formula (Lemma [8\)](#page-9-2), uniform bounds on the Giga-Kohn energy (Lemma [9\)](#page-9-4) and the previous lemma. See also related arguments in [\[23,](#page-55-6) [24,](#page-55-7) [25\]](#page-55-8).

Proof of Proposition [7.](#page-8-2) Assume [\(2.1\)](#page-8-3). By translation invariance and scaling, it suffices to prove the case  $(x_0, t_0) = (0, 0)$ . For  $r > 0$ ,  $\tilde{x} \in \mathbb{R}^n$  and  $-1 < \tilde{t} \leq 0$  with  $Q((\tilde{x}, \tilde{t}), r) \subset Q(\delta/2)$ , we note that  $|\tilde{x}| \leq \delta/2$ ,  $-\delta^2/4 \leq \tilde{t} \leq 0$  and  $r \leq \delta/2$  hold. Define  $I_r(\tilde{x},\tilde{t})$  as in [\(2.11\)](#page-12-0). By using the backward similarity variables,

$$
I_r(\tilde{x}, \tilde{t}) \le C \int_{\tilde{t}-r^2/4}^{\tilde{t}-r^2/16} (\tilde{t}-t)^{\frac{2}{p-1}} \int_{B(\tilde{x},r/2)} |u(x,t)|^{p+1} (\tilde{t}-t)^{-\frac{n}{2}} e^{-\frac{|x-\tilde{x}|^2}{4(\tilde{t}-t)}} dx dt
$$
  

$$
\le C \int_{-\log(r^2/4)}^{-\log(r^2/16)} \int_{\mathbf{R}^n} |w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^{p+1} \rho(\eta) d\eta d\tau =: J_r(\tilde{x}, \tilde{t}).
$$

From [\(2.7\)](#page-9-5) and [\(2.5\)](#page-9-6), it follows that for  $w = w_{(\tilde{x}, \tilde{t})}$ ,

$$
J_r = \frac{p+1}{p-1} \int_{-\log(r^2/16)}^{-\log(r^2/16)} \int_{\mathbf{R}^n} w w_{\tau} \rho d\eta d\tau + \frac{2(p+1)}{p-1} \int_{-\log(r^2/4)}^{-\log(r^2/16)} \mathcal{E}_{(\tilde{x},\tilde{t})} d\tau
$$
  
\n
$$
\leq C \left( \iint |w|^2 \rho \right)^{\frac{1}{2}} \left( \int_{-\log(r^2/4)}^{-\log(r^2/16)} \int_{\mathbf{R}^n} |w_{\tau}|^2 \rho \right)^{\frac{1}{2}} + C \int_{-\log(r^2/4)}^{-\log(r^2/16)} \mathcal{E}_{(\tilde{x},\tilde{t})} d\tau
$$
  
\n
$$
\leq C(J_r)^{\frac{1}{p+1}} \left( \int_{-\log(r^2/4)}^{-\log(r^2/16)} \int_{\mathbf{R}^n} \rho \right)^{\frac{1}{2} - \frac{1}{p+1}} (\mathcal{E}(-\log(r^2/4)) - \mathcal{E}(-\log(r^2/16)))^{\frac{1}{2}}
$$
  
\n
$$
+ C\mathcal{E}(-\log(r^2/4)) \int_{-\log(r^2/4)}^{-\log(r^2/16)} d\tau.
$$

Here and below, we often suppress the subscript  $(\tilde{x},\tilde{t})$ . On the other hand, straightforward computations with [\(1.6\)](#page-3-4) yield

$$
J_r = \int_{\tilde{t}-r^2/4}^{\tilde{t}-r^2/16} (\tilde{t}-t)^{\frac{2}{p-1}-\frac{n}{2}} \left\| |u(\cdot,t)|^{p+1} e^{-\frac{|.\cdot-\tilde{x}|^2}{4(\tilde{t}-t)}} \right\|_{L^1(\mathbf{R}^n)} dt
$$
  
\n
$$
\leq C \int_{\tilde{t}-r^2/4}^{\tilde{t}-r^2/16} (\tilde{t}-t)^{\frac{2}{p-1}-\frac{n}{2}} \|u(\cdot,t)\|_{L^{q_c,\infty}(\mathbf{R}^n)}^{p+1} \left\| e^{-\frac{|.\cdot-\tilde{x}|^2}{4(\tilde{t}-t)}} \right\|_{L^{\frac{q_c}{q_c-(p+1)},1}(\mathbf{R}^n)} dt
$$
  
\n
$$
\leq C M^{p+1} \int_{\tilde{t}-r^2/4}^{\tilde{t}-r^2/16} (\tilde{t}-t)^{\frac{2}{p-1}-\frac{n}{2}+\frac{n}{2}(1-\frac{p+1}{q_c})} dt = C M^{p+1} \log \frac{r^2/4}{r^2/16} \leq C M^{p+1}.
$$

This together with  $\mathcal{E}_{(\tilde{x},\tilde{t})}(-\log(r^2/16)) \ge 0, M > 1, (2.3)$  $\mathcal{E}_{(\tilde{x},\tilde{t})}(-\log(r^2/16)) \ge 0, M > 1, (2.3)$  and  $(2.6)$  gives

$$
J_r \le CM \left(\mathcal{E}(-\log(r^2/4))\right)^{\frac{1}{2}} + C\mathcal{E}(-\log(r^2/4)) \le CMf(E(\tilde{t} - r^2/4))
$$
  

$$
\le CMf(E(-\delta^2/2)) \le CMf\left(\frac{1}{\delta^2} \int_{-\delta^2}^{-\delta^2/2} E(t)dt\right),
$$

where  $f(s) := s + s^{1/2}$  for  $s \ge 0$ .

Recall [\(2.3\)](#page-9-7). Then,

$$
\frac{1}{\delta^2} \int_{-\delta^2}^{-\delta^2/2} E(t)dt = \int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \frac{e^{-\tau}}{\delta^2} \mathcal{E}(\tau) d\tau \leq C \int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \mathcal{E}(\tau) d\tau.
$$

By  $(2.7)$ , we can also see that

$$
\int_{-\log(\tilde{t}+\delta^2/2)}^{\log(\tilde{t}+\delta^2/2)} \mathcal{E}(\tau)d\tau = -\frac{1}{2} \iint w w_{\tau} \rho + \frac{p-1}{2(p+1)} \iint |w|^{p+1} \rho
$$
  
\n
$$
\leq C \left( \iint |w|^2 \rho \right)^{\frac{1}{2}} \left( \int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \int_{\mathbf{R}^n} |w_{\tau}|^2 \rho \right)^{\frac{1}{2}} + C \iint |w|^{p+1} \rho
$$
  
\n
$$
\leq C \left( \iint |w|^{p+1} \rho \right)^{\frac{1}{p+1}} \left( \int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \int_{\mathbf{R}^n} \rho \right)^{\frac{1}{2} - \frac{1}{p+1}}
$$
  
\n
$$
\times \left( \mathcal{E}(-\log(\tilde{t}+\delta^2)) - \mathcal{E}(-\log(\tilde{t}+\delta^2/2)) \right)^{\frac{1}{2}} + C \iint |w|^{p+1} \rho.
$$

Since  $0 \leq \mathcal{E}_{(\tilde{x},\tilde{t})}(-\log(\tilde{t}+\delta^2)) = E_{(\tilde{x},\tilde{t})}(-\delta^2) \leq CM^{2p}$  by Lemma [9,](#page-9-4) we obtain

$$
\int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \mathcal{E}_{(\tilde{x},\tilde{t})}(\tau) d\tau \le CM^p \tilde{f}\left(\int_{-\log(\tilde{t}+\delta^2)}^{-\log(\tilde{t}+\delta^2/2)} \int_{\mathbf{R}^n} |w_{(\tilde{x},\tilde{t})}(\eta,\tau)|^{p+1} \rho(\eta) d\eta d\tau\right)
$$
  

$$
\le CM^p \tilde{f}\left(\int_{-\delta^2}^{-\delta^2/2} (\tilde{t}-t)^{\frac{2}{p-1}} \int_{\mathbf{R}^n} |u(x,t)|^{p+1} K_{(\tilde{x},\tilde{t})}(x,t) dxdt\right),
$$

where  $\tilde{f}(s) := s + s^{1/(p+1)}$  for  $s \geq 0$ .

Let  $0 < \varepsilon_0 < 1$  be a constant chosen later and let  $0 < \varepsilon < \varepsilon_0$ . Then [\(2.1\)](#page-8-3) yields

$$
I_r(\tilde{x},\tilde{t}) \leq CM^{p+1}(f \circ \tilde{f}) \left( \varepsilon + \delta^{\frac{4}{p-1}} \int_{-\delta^2}^{-\delta^2/2} \int_{\mathbf{R}^n \setminus B(A\delta)} |u(x,t)|^{p+1} K_{(\tilde{x},\tilde{t})}(x,t) dxdt \right).
$$

We use the notation  $K_{(\tilde{y},\tilde{s})}$  even for  $\tilde{s} > 0$  and observe that if  $Q((\tilde{x},\tilde{t}), r) \subset Q(\delta/2)$ , then

$$
K_{(\tilde{x},\tilde{t})}(x,t) \le CK_{(0,\delta^2/2)}(x,t) \le Ce^{-\frac{A^2}{24}}K_{(0,\delta^2)}(x,t)
$$

for  $x \in \mathbb{R}^n \setminus B(A\delta)$  and  $-\delta^2 < t < -\delta^2/2$ , and thus

$$
I_r \leq CM^{p+1}(f \circ \tilde{f}) \left( \varepsilon + \delta^{\frac{4}{p-1}} e^{-\frac{A^2}{24}} \int_{-\delta^2}^{-\delta^2/2} \int_{\mathbf{R}^n} |u|^{p+1} K_{(0,\delta^2)} dx dt \right).
$$

By the Hölder inequality and  $(1.6)$ , we have

$$
\int_{-\delta^2}^{-\delta^2/2} \int_{\mathbf{R}^n} |u|^{p+1} K_{(0,\delta^2)} dx dt
$$
\n
$$
\leq C \int_{-\delta^2}^{-\delta^2/2} (\delta^2 - t)^{-\frac{n}{2}} |||u|^{p+1} ||_{L^{\frac{qc}{p+1},\infty}(\mathbf{R}^n)} \Bigg|| e^{-\frac{|v|^2}{4(\delta^2 - t)}} \Bigg||_{L^{\frac{qc}{qc - (p+1)},1}(\mathbf{R}^n)} dt
$$
\n
$$
\leq CM^{p+1} \int_{-\delta^2}^{-\delta^2/2} (\delta^2 - t)^{-\frac{p+1}{p-1}} dt = CM^{p+1} \delta^{-\frac{4}{p-1}}.
$$

Therefore, if  $0 < \varepsilon < \varepsilon_0$ ,  $Q((\tilde{x}, \tilde{t}), r) \subset Q(\delta/2)$  and A is chosen as in [\(2.1\)](#page-8-3), then

$$
I_r \leq CM^{p+1}(f \circ \tilde{f})(\varepsilon + CM^{p+1}e^{-\frac{A^2}{24}}) \leq CM^{p+1}(f \circ \tilde{f})(\varepsilon + C\varepsilon)
$$
  

$$
\leq CM^{p+1}(f \circ \tilde{f})(\varepsilon) \leq \tilde{C}M^{p+1}\varepsilon^{\frac{1}{2(p+1)}} \leq \tilde{C}M^{p+1}\varepsilon^{\frac{1}{2(p+1)}}_0 = \tilde{\varepsilon}_1
$$

with some constants  $\tilde{C} > 0$  depending only on n and p. Here,  $\tilde{\varepsilon}_1$  is given in Lemma [10](#page-12-2) and  $\varepsilon_0$  is chosen such that  $\varepsilon_0 := \tilde{C}^{-2(p+1)} M^{-2(p+1)^2} \tilde{\varepsilon}_1^{2(p+1)}$ . Thus, we can apply Lemma [10](#page-12-2) with  $\tilde{\varepsilon} = \tilde{C} M^{p+1} \varepsilon^{1/(2(p+1))}$  to obtain

$$
||u||_{L^{\infty}(Q(\delta/4))} \leq CM \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{2}{p-1}}, \quad ||\nabla u||_{L^{\infty}(Q(\delta/4))} \leq CM \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{p+1}{p-1}}.
$$

<span id="page-17-0"></span>The proof is complete, with  $\varepsilon_1$  in Proposition [7](#page-8-2) defined as  $\varepsilon_1 := \tilde{\varepsilon}_1/\tilde{C}$ .

# <span id="page-17-2"></span>3. ESTIMATES VIA  $\varepsilon$ -REGULARITY

Let  $u$  be a classical solution of  $(1.5)$  satisfying the Lorentz norm bound  $(1.6)$ . We give several estimates based on the above quantitative  $\varepsilon$ -regularity. We set  $\varepsilon_0 := M^{-2(p+1)^2} \varepsilon_1^{2(p+1)}$  as in Proposition [7,](#page-8-2) where  $\varepsilon_1$  depends only on n and p. In the rest of this paper, we fix

(3.1) 
$$
\varepsilon := \frac{\varepsilon_0}{2}, \quad A := 48 \log \frac{M^{p+1}}{\varepsilon},
$$

<span id="page-17-1"></span>unless otherwise stated.

3.1. Propagation of concentration. We prove the backward-in-time propagation of concentration in the form of its contraposition. After that, we give estimates on intersecting regions of concentration and quantitative regularity.

<span id="page-17-7"></span>**Proposition 12** (Backward propagation). Let  $p > p<sub>S</sub>$  and assume [\(3.1\)](#page-17-2). There exists  $M_* > 1$  depending only on n and p such that the following statement holds true. Assume that u satisfies [\(1.6\)](#page-3-4) with  $M \geq M_*$ . If  $t', t'' \in (-1,0)$  and u satisfy

<span id="page-17-3"></span>(3.2) 
$$
-\frac{1}{16} \le t'' \le M^{(p-1)(2p+4)} A^{\frac{n(p-1)}{p+1}} t',
$$

<span id="page-17-4"></span>
$$
(3.3) \qquad \qquad (-t'')^{\frac{2}{p-1}-\frac{n}{2}} \int_{t''}^{t''/2} \int_{B(x_0,A(-t'')^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt \le \varepsilon,
$$

then we conclude that

(3.4) 
$$
(-t')^{\frac{2}{p-1}-\frac{n}{2}} \int_{t'}^{t'/2} \int_{B(x_0,A(-t')^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt \leq \varepsilon.
$$

*Proof.* Without loss of generality set  $x_0 = 0$ . The assumptions [\(3.2\)](#page-17-3) and [\(3.3\)](#page-17-4) allow us to apply Proposition [7,](#page-8-2) which gives that

<span id="page-17-6"></span><span id="page-17-5"></span>
$$
||u||_{L^{\infty}(Q((-t'')^{\frac{1}{2}}/4))} \leq CM \varepsilon^{\frac{1}{2(p+1)^2}} (-t'')^{-\frac{1}{p-1}},
$$

and so

(3.5) 
$$
\sup_{Q(\frac{1}{4}(-t'')^{\frac{1}{2}})} |u(x,t)|^{p+1} \leq C^{p+1} M^{p+1} \varepsilon^{\frac{1}{2(p+1)}} (-t'')^{-\frac{p+1}{p-1}}.
$$

From  $(3.2)$ , we have that for M sufficiently large,

$$
B(A(-t')^{\frac{1}{2}}) \times (t', t'/2) \subset Q((-t'')^{\frac{1}{2}}/4).
$$

Thus,  $(3.2)$  and  $(3.5)$  imply that

$$
\begin{aligned} &(-t')^{\frac{2}{p-1}-\frac{n}{2}}\int_{t'}^{t'/2}\int_{B(A(-t')^{\frac{1}{2}})}|u(x,t)|^{p+1}dxdt\leq C(M^{p+1}A^{n})\left(\frac{-t'}{-t''}\right)^{\frac{p+1}{p-1}}\\ &\leq CM^{-(2p+3)(p+1)}\leq \varepsilon,\end{aligned}
$$

as required.  $\hfill \square$ 

From now on we will use the terminology  $M$  being sufficiently large' to mean that  $M \geq M_0$ , where  $M_0 > 1$  is a sufficiently large constant depending only on n and p. We will also utilize the notation  $M_k := M^{c_k(n,p)}$  for  $k \in \mathbb{N}$  with M being sufficiently large and with  $c_k(n, p) > 0$  being sufficiently larger than  $c_{k-1}(n, p)$  etc.

The rest of this subsection is devoted to estimates on intersecting regions of concentration and quantitative regularity. A related statement for the higherdimensional Navier-Stokes equations was previously proven in [\[31,](#page-55-0) Proposition 5.1].

<span id="page-18-1"></span>**Proposition 13** (Intersecting regions). Let  $p > p<sub>S</sub>$  and assume [\(3.1\)](#page-17-2). Then there exist  $C, C > 0$  depending only on n and p such that the following statement holds for all M sufficiently large: Assume that u satisfies [\(1.6\)](#page-3-4) and that  $-1/64 < t'' < 0$ satisfies

<span id="page-18-2"></span>(3.6) 
$$
(-t'')^{\frac{2}{p-1}-\frac{n}{2}}\int_{t''}^{t''/2}\int_{B(0,A(-t'')^{\frac{1}{2}})}|u(x,t)|^{p+1}dxdt > \varepsilon.
$$

Then we conclude that there exist backward parabolic cylinders  $Q((x_*,t_*-r^2/8),\hat{\delta}r)$ and  $Q(z_*, r)$  with  $z_* = (x_*, t_*)$  satisfying

<span id="page-18-3"></span>
$$
(3.7) \qquad Q((x_*,t_*-r^2/8),\hat{\delta}r) \subset Q(z_*,r) \subset B(0,20M_1(-t'')^{\frac{1}{2}}) \times (2t'',t''/4)
$$

with  $r := 8M_2^{-(6n+2)}(-t'')^{1/2}$  and  $\hat{\delta} := M_2^{-1}$  such that

<span id="page-18-4"></span>
$$
(3.8) \t\t ||u||_{L^{\infty}(Q(z_*,r/2))} \le CM^{-2}r^{-\frac{2}{p-1}}, \t ||\nabla u||_{L^{\infty}(Q(z_*,r/2))} \le CM^{-2}r^{-\frac{p+1}{p-1}},
$$

<span id="page-18-5"></span>
$$
(3.9) \qquad \int_{Q((x_*,t_*-r^2/8),\hat{\delta}r)} |u|^2 dx dt \ge M_2^{-\bar{C}} (\hat{\delta}r)^{n+2-\frac{4}{p-1}}.
$$

To prove this proposition, we prepare several estimates based on the following interpolation inequality on a domain  $\Omega \subset \mathbf{R}^n$ .

<span id="page-18-0"></span>**Lemma 14.** Let  $1 \leq \tilde{p} < \tilde{r} < \tilde{q} \leq \infty$ ,  $f \in L^{\tilde{p},\infty}(\Omega) \cap L^{\tilde{q},\infty}(\Omega)$  and  $0 < \tilde{\theta} < 1$  be such that

$$
\frac{1}{\tilde{r}} = \frac{\tilde{\theta}}{\tilde{p}} + \frac{1 - \tilde{\theta}}{\tilde{q}} \quad \left(\text{equivalently, } \theta = \frac{\frac{1}{\tilde{r}} - \frac{1}{\tilde{q}}}{\frac{1}{\tilde{p}} - \frac{1}{\tilde{q}}}\right).
$$

Then,  $f \in L^{\tilde{r}}(\Omega)$  and

$$
||f||_{L^{\tilde{r}}(\Omega)} \leq \left(\frac{\tilde{r}}{\tilde{r}-\tilde{p}}+\frac{\tilde{r}}{\tilde{q}-\tilde{r}}\right)^{\frac{1}{\tilde{r}}}||f||_{L^{\tilde{p},\infty}(\Omega)}^{\tilde{\theta}}||f||_{L^{\tilde{q},\infty}(\Omega)}^{1-\tilde{\theta}}.
$$

*Proof.* See [\[15,](#page-55-26) Proposition 1.1.14] for example.  $\Box$ 

By using Lemma [14,](#page-18-0) we also prepare estimates for  $f : \Omega \times I \to \mathbf{R}$  with  $\Omega \times I \subset$  $\mathbf{R}^n \times \mathbf{R}$  bounded.

<span id="page-19-4"></span>**Lemma 15.** Let  $p > p_S$  and define  $\phi_i = \phi_i(p, n)$   $(i = 1, 2, 3)$  by

<span id="page-19-2"></span>(3.10) 
$$
\phi_1 := \frac{\frac{1}{p+1} - \frac{1}{q_c}}{\frac{1}{p-1} - \frac{1}{q_c}}, \quad \phi_2 := \frac{\frac{1}{p-1} - \frac{1}{q_c}}{\frac{1}{2} - \frac{1}{q_c}}, \quad \phi_3 := \frac{\frac{1}{p+1} - \frac{1}{q_c}}{\frac{1}{3q_c/4} - \frac{1}{q_c}}
$$

Suppose that

$$
\sup_{t\in I} ||f(\cdot,t)||_{L^{q_c,\infty}(\Omega)} \leq M.
$$

.

Then there exists  $C > 0$  depending only on n and p such that the following inequalities hold with  $\iint = \iint_{\Omega \times I}$ :

(i) 
$$
\iint |f|^{p+1} \le CM^{(1-\phi_1)(p+1)} \left( \iint |f|^{p-1} \right)^{\frac{(p+1)\phi_1}{p-1}} |I|^{1-\frac{(p+1)\phi_1}{p-1}}.
$$
  
\n(ii) 
$$
\iint |f|^{p-1} \le \begin{cases} \left( \iint |f|^2 \right)^{\frac{p-1}{2}} |\Omega \times I|^{1-\frac{p-1}{2}} & \text{if } p-1 \le 2, \\ CM^{(1-\phi_2)(p-1)} \left( \iint |f|^2 \right)^{\frac{(p-1)\phi_2}{2}} |I|^{1-\frac{(p-1)\phi_2}{2}} & \text{if } p-1 > 2. \end{cases}
$$
  
\n(iii) 
$$
\iint |f|^{p+1} \le CM^{(1-\phi_3)(p+1)} \left( \iint |f|^{\frac{3q_c}{4}} \right)^{\frac{4(p+1)\phi_3}{3q_c}} |I|^{1-\frac{4(p+1)\phi_3}{3q_c}} \text{ if } \frac{3q_c}{4} < p+1.
$$

*Proof.* (i) We apply Lemma [14](#page-18-0) with  $\tilde{p} = p - 1$ ,  $\tilde{q} = q_c$  and  $\tilde{r} = p + 1$  for each  $t \in I$ . Raising this to the power  $p + 1$  and integrating with respect to time gives that

<span id="page-19-0"></span>
$$
(3.11) \qquad \int_{I} \|f(\cdot,t)\|_{L^{p+1}(\Omega)}^{p+1} dt \le CM^{(1-\phi_1)(p+1)} \int_{I} \|f(\cdot,t)\|_{L^{p-1}(\Omega)}^{(p+1)\phi_1} dt.
$$

Noting that

$$
\frac{1}{p+1} = \frac{\phi_1}{p-1} + \frac{1-\phi_1}{q_c} > \frac{\phi_1}{p-1},
$$

we have that  $(p+1)\phi_1 < p-1$ . This allows us to apply Hölder's inequality to the right-hand side of [\(3.11\)](#page-19-0) to obtain the desired conclusion.

(ii) The case  $p - 1 \leq 2$  follows immediately from Hölder's inequality. As for the case  $p - 1 > 2$ , we apply Lemma [14](#page-18-0) with  $\tilde{p} = 2$ ,  $\tilde{q} = q_c$  and  $\tilde{r} = p - 1$ . The rest of the proof is the same as (i).

(iii) This follows from Lemma [14](#page-18-0) with  $\tilde{p} = 3q_c/4$ ,  $\tilde{q} = q_c$  and  $\tilde{r} = p + 1$ .

The following corollary of Proposition [7](#page-8-2) will be convenient in several places.

<span id="page-19-3"></span>Corollary 16. Let  $p > p_S$  and  $C^* > 0$ . Then there exists  $M_0 > 1$  depending only on n, p and  $C^*$  such that the following statement holds for any  $M \geq M_0$  and  $0 < \delta < (2M)^{-1}$ : If u satisfies [\(1.6\)](#page-3-4) and

<span id="page-19-1"></span>
$$
(3.12) \quad \delta^{\frac{4}{p-1}-n} \int_{t_0-\delta^2}^{t_0} \int_{B(x_0,\delta)} |u|^{p+1} dx dt \le C^* M^{-(n+3-\frac{2(p+1)}{p-1}+6(p+1)^2+\frac{2(p+1)^3}{p-1})},
$$

then there exists  $C > 0$  depending only on n and p such that

$$
||u||_{L^{\infty}(Q((x_0,t_0),\delta/2))} \leq CM^{-3}\delta^{-\frac{2}{p-1}}, \quad ||\nabla u||_{L^{\infty}(Q((x_0,t_0),\delta/2))} \leq CM^{-2}\delta^{-\frac{p+1}{p-1}}.
$$

Proof. In this proof, let

$$
\varepsilon := M^{-(6(p+1)^2 + \frac{2(p+1)^3}{p-1})}
$$
,  $A := 48 \log \frac{M^{p+1}}{\varepsilon}$ .

Clearly, we have that for  $M$  sufficiently large,

$$
(3.13) \t\t 0 < \varepsilon < \varepsilon_0,
$$

where  $\varepsilon_0$  is as in Proposition [7.](#page-8-2) Without loss of generality, let  $x_0 = 0$ . Note that for M sufficiently large

$$
(3.14) \t\t A \le M/2.
$$

Let us rescale

<span id="page-20-1"></span><span id="page-20-0"></span>
$$
u_{\lambda}(x,t) := \lambda^{\frac{2}{p-1}} u(\lambda x, \lambda^2 t + t_0) \quad \text{with } \lambda = M^{-1}.
$$

Then,  $u_{\lambda}$  satisfies the scale-invariant bound  $(1.6)$  on the rescaled time interval. Let  $\tilde{z}_0 = (\tilde{t}_0, \tilde{x}_0) \in Q((0, t_0), \delta/2)$ . Using  $0 < \delta < (2M)^{-1}$  and  $(3.14)$ , we observe that

$$
\begin{aligned} \hat{z}_0 &= (\hat{x}_0, \hat{t}_0) := (\tilde{x}_0/\lambda, (\tilde{t}_0 - t_0)/\lambda^2) \in Q(M\delta/2), \quad \hat{t}_0 > -1/16, \\ B(\hat{x}_0, A\delta) &< (\hat{t}_0 - \delta^2/2, \hat{t}_0 - \delta^2) \subset Q(M\delta). \end{aligned}
$$

These together with  $(3.12)$ ,  $(3.13)$  and  $(3.14)$  show that

$$
\delta^{\frac{4}{p-1}-n} \int_{\hat{t}_0-\delta^2}^{\hat{t}_0-\delta^2/2} \int_{B(\hat{x}_0,A\delta)} |u_\lambda(x,t)|^{p+1} dx dt \leq \delta^{\frac{4}{p-1}-n} \int_{Q(M\delta)} |u_\lambda(x,t)|^{p+1} dx dt
$$
  
=  $M^{n+2-\frac{2(p+1)}{p-1}} \left( \delta^{\frac{4}{p-1}-n} \int_{t_0-\delta^2}^{t_0} \int_{B(\delta)} |u(x,t)|^{p+1} dx dt \right) \leq C^* M^{-1} \varepsilon.$ 

By [\(3.13\)](#page-20-1) and  $t_0 > -1/16$ , we can apply Proposition [7](#page-8-2) to  $u_\lambda$  to see that

$$
|u(\tilde{x}_0, \tilde{t}_0)| \le CM^{\frac{p+1}{p-1}} \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{2}{p-1}} = CM^{-3} \delta^{-\frac{2}{p-1}},
$$
  

$$
|\nabla u(\tilde{x}_0, \tilde{t}_0)| \le CM^{\frac{p+1}{p-1}} \varepsilon^{\frac{1}{2(p+1)^2}} \delta^{-\frac{p+1}{p-1}} = CM^{-2} \delta^{-\frac{p+1}{p-1}}.
$$

Since  $\tilde{z}_0 = (\tilde{x}_0, \tilde{t}_0) \in Q((0, t_0), \delta/2)$  was taken arbitrarily, this implies that

$$
||u||_{L^{\infty}(Q((0,t_0),\delta/2))} \le CM^{-3}\delta^{-\frac{2}{p-1}}, \quad ||\nabla u||_{L^{\infty}(Q((0,t_0),\delta/2))} \le CM^{-2}\delta^{-\frac{p+1}{p-1}},
$$
as required.

We state a useful version of  $\varepsilon$ -regularity, which will be used in the proof of Proposition [13.](#page-18-1)

<span id="page-20-2"></span>**Lemma 17.** Let  $p > p_S$  and  $C^* > 0$ . Define  $\alpha(n, p) := \max(a_1(n, p), a_2(n, p))$  with

$$
a_1(n,p) := \frac{3q_c}{4} \left( \frac{n+3}{p+1} - \frac{2}{p-1} + 6(p+1) + \frac{2(p+1)^2}{p-1} \right),
$$
  

$$
a_2(n,p) := \frac{3q_c}{4\phi_3} \left( \frac{n+3}{p+1} - \frac{2}{p-1} + 6(p+1) + \frac{2(p+1)^2}{p-1} + (1-\phi_3) \right),
$$

where  $\phi_3$  is defined by [\(3.10\)](#page-19-2). Then there exists  $M_0 > 1$  depending only on n, p and  $C^*$  such that the following statement holds for any  $M \geq M_0$  and  $0 < \delta < (2M)^{-1}$ : If u satisfies [\(1.6\)](#page-3-4) and

$$
\delta^{-(\frac{n}{4}+2)} \int_{t_0-\delta^2}^{t_0} \int_{B(x_0,\delta)} |u(x,t)|^{\frac{3q_c}{4}} dx dt \le C^* M^{-\alpha(n,p)},
$$

then there exists  $C > 0$  depending only on n and p such that

$$
||u||_{L^{\infty}(Q((x_0,t_0),\delta/2))} \leq CM^{-3}\delta^{-\frac{2}{p-1}}, \quad ||\nabla u||_{L^{\infty}(Q((x_0,t_0),\delta/2))} \leq CM^{-2}\delta^{-\frac{p+1}{p-1}}.
$$

*Proof.* In the case that  $3q_c/4 \geq p+1$ , we obtain the conclusion by applying Hölder's inequality and Corollary [16.](#page-19-3) In the case that  $3q_c/4 < p+1$ , we apply Lemma [15](#page-19-4) (iii) and Corollary [16.](#page-19-3)

We are now in a position to prove Proposition [13](#page-18-1) by using Lemma [15,](#page-19-4) Lemma [17](#page-20-2) and performing a counting argument inspired by [\[31,](#page-55-0) Proposition 5.1]. The scaleinvariant bound [\(1.6\)](#page-3-4) is not countably additive over disjoint spatial scales, which requires a further judicious choice of indices in the counting argument.

*Proof of Proposition [13.](#page-18-1)* In the proof,  $C_n^*$ ,  $C_n^{**}$  and  $C_n^{***}$  will be used to denote certain constants depending only on  $n$ . We divide the proof into 4 steps. Step 1: rescaling and derived functions

Let  $-1/64 < t'' < 0$  and let u satisfy [\(3.6\)](#page-18-2). Set *I* := (-1/64, -1/128). Define the rescaled function  $u_{\lambda}: \mathbf{R}^n \times (1/(64t''),0) \to \mathbf{R}$  by

<span id="page-21-8"></span>(3.15) 
$$
u_{\lambda}(x,t) := \lambda^{\frac{2}{p-1}} u(\lambda x, \lambda^2 t) \quad \text{with } \lambda := 8(-t'')^{\frac{1}{2}}.
$$

As u satisfies [\(1.6\)](#page-3-4) and [\(3.6\)](#page-18-2), we get that  $u_{\lambda}$  satisfies

<span id="page-21-0"></span>(3.16) 
$$
\sup_{-1 \leq t < 0} \|u_{\lambda}(\cdot, t)\|_{L^{q_c, \infty}(\mathbf{R}^n)} \leq M,
$$

<span id="page-21-1"></span>(3.17) 
$$
\iint_{B(A/8)\times I} |u_{\lambda}(x,t)|^{p+1} dx dt \ge C(p)\varepsilon.
$$

Using  $(3.16)$ ,  $(3.17)$  and Lemma [15](#page-19-4) (i), we infer that

<span id="page-21-2"></span>
$$
(3.18)\qquad \qquad \iint_{B(A/8)\times I} |u_{\lambda}(x,t)|^{p-1} dx dt \ge C(p,n) M^{-\frac{(1-\phi_1)(p-1)}{\phi_1}} \varepsilon^{\frac{p-1}{(p+1)\phi_1}},
$$

where  $\phi_1$  is as in [\(3.10\)](#page-19-2). Define  $U: \mathbb{R}^n \times (1/(64t''),0) \to \mathbb{R}$  by

<span id="page-21-7"></span>(3.19) 
$$
U(x,t) := |u_\lambda(x,t)|^{\frac{p-1}{2}}.
$$

By [\(3.16\)](#page-21-0) and [\(3.18\)](#page-21-2), we get that, for M sufficiently large and appropriate  $M_1$ ,

<span id="page-21-3"></span>(3.20) 
$$
\sup_{-1 < t < 0} \|U(\cdot, t)\|_{L^{n,\infty}(\mathbf{R}^n)} \le M^{\frac{p-1}{2}},
$$

<span id="page-21-5"></span>(3.21) 
$$
\iint_{B(M_1)\times I} |U(x,t)|^2 dx dt \ge M_1^{-1}.
$$

Step 2: families of cubes and the regularity of  $u_{\lambda}$ 

For appropriately chosen  $M_2 := M^{c_2(n,p)}$ , define

(3.22) 
$$
\hat{\delta} := M_2^{-1}, \quad l := M_2^{-(6n+2)}
$$

With this, we define the family of cubes (backward parabolic cylinders)

$$
C_0 := \{Q(z_0, l); z_0 = (x_0, t_0) \in ((\hat{\delta}l)^n \mathbf{Z}^n \times (\hat{\delta}l)^2 \mathbf{Z}) \cap (B(2M_1) \times 2I)\}.
$$

.

Here, for  $0 < r \leq 3$ , we write

<span id="page-21-6"></span><span id="page-21-4"></span>
$$
rI := \left(-\frac{3}{256} - \frac{r}{256}, -\frac{3}{256} + \frac{r}{256}\right) \subset [-1, 0].
$$

Then there exists  $C(n) > 1$  such that

(3.23) 
$$
\frac{1}{C(n)} M_1^n(\hat{\delta}l)^{-(n+2)} \leq #\mathcal{C}_0 \leq C(n) M_1^n(\hat{\delta}l)^{-(n+2)},
$$

where  $\#\mathcal{C}_0$  is the cardinality of  $\mathcal{C}_0$ . Now, let us define the subfamily of cubes

$$
\mathcal{C}_1 := \left\{ Q \in \mathcal{C}_0; \; \|U\|_{L^{\frac{3n}{4}}(Q)}^{\frac{3n}{4}} > M_1^{-1} l^n \right\}.
$$

Here  $M_2 = M^{c_2(p,n)}$  is chosen appropriately to ensure that if

<span id="page-22-1"></span>
$$
\frac{1}{l^{\frac{n}{4}+2}}\|U\|_{L^{\frac{3n}{4}}(Q)}^{\frac{3n}{4}} \leq M_1^{-1}l^{\frac{3n}{4}-2},
$$

then  $u_{\lambda}$  satisfies the hypothesis in Lemma [17](#page-20-2) with  $\delta = l$ , which in turn implies that

<span id="page-22-0"></span>
$$
(3.24) \t\t ||u_{\lambda}||_{L^{\infty}(Q(z_0,l/2))} \leq Cl^{-\frac{2}{p-1}}M^{-2} \t with Q = Q(z_0,l).
$$

Hence, to show that there exists a cube  $Q \in \mathcal{C}_0$  satisfying [\(3.24\)](#page-22-0), it suffices to show that  $C_0 \setminus C_1 \neq \emptyset$ . To show this, it is sufficient to show that

$$
\# \mathcal{C}_1 < \# \mathcal{C}_0.
$$

We prove [\(3.25\)](#page-22-1). Using the definition of  $C_1$ , the fact that each of the cubes in  $C_1$ overlaps with at most  $C(n)l^{n+2}(\hat{\delta}l)^{-2-n}$  (=  $C(n)\hat{\delta}^{-2-n}$ ) other cubes in  $\mathcal{C}_1$ , [\(3.20\)](#page-21-3) and Hölder's inequality for Lorentz spaces, we have

$$
M_1^{-1}l^n(\#\mathcal{C}_1) \le \sum_{Q \in \mathcal{C}_0} ||U||_{L^{\frac{3n}{4}}(Q)}^{\frac{3n}{4}} \le C(n)\hat{\delta}^{-2-n} \iint_{B(3M_1)\times 3I} |U|^{\frac{3n}{4}} dxdt
$$
  

$$
\le C(n)M_1^{\frac{n}{4}}M^{\frac{3n(p-1)}{8}}\hat{\delta}^{-2-n}.
$$

Thus,

<span id="page-22-5"></span>(3.26) 
$$
\#\mathcal{C}_1 \leq C(n)M_1^{\frac{n}{4}+1}M^{\frac{3n(p-1)}{8}}l^2(\hat{\delta}l)^{-(n+2)}.
$$

Recalling [\(3.23\)](#page-21-4), we see that for  $M_2 = M^{c_2(p,n)}$  chosen appropriately, we have that  $\#\mathcal{C}_1 < \#\mathcal{C}_0$ . This implies that there exists a cube  $Q \in \mathcal{C}_0$  such that [\(3.24\)](#page-22-0) holds. Step 3: concentration of u on a cube descending from  $C_0 \setminus C_1$ 

For  $Q = Q((x_0, t_0), l) \in C_0$ , we define the descendant  $Q' := Q((x_0, t_0 - l^2/8), \delta l)$ . These cubes are such that  $B(M_1) \times I \subset \{Q'; Q \in C_0\}$ . Using this and [\(3.21\)](#page-21-5) gives

.

<span id="page-22-6"></span>(3.27) 
$$
\sum_{Q \in C_0 \setminus C_1} \iint_{Q'} |U|^2 dx dt + \sum_{Q \in C_1} \iint_{Q'} |U|^2 dx dt \ge M_1^{-1}
$$

By  $(3.20)$  and Hölder's inequality in the Lorentz spaces, we have

(3.28) 
$$
||U||_{L^{2}(Q')}^{2} \leq C(n)M^{p-1}(\hat{\delta}l)^{n}
$$

for every  $Q \in \mathcal{C}_0$ . Now we are going to show that if  $\mathcal{C}_1 \neq \emptyset$ , then

(3.29) 
$$
\sum_{Q \in \mathcal{C}_1} \iint_{Q'} |U|^2 dx dt \ge \frac{1}{2} M_1^{-1}
$$

cannot occur. Assume for contradiction that [\(3.29\)](#page-22-2) holds. We define

<span id="page-22-4"></span><span id="page-22-3"></span><span id="page-22-2"></span>
$$
C_2 := \{ Q \in C_1; ||U||_{L^2(Q')}^2 > M_1^{-5} \hat{\delta}^{2+n} l^{n+1-\frac{8}{3n}} \}.
$$

By [\(3.29\)](#page-22-2), we get

(3.30) 
$$
\sum_{Q \in \mathcal{C}_1 \setminus \mathcal{C}_2} \iint_{Q'} |U|^2 dx dt + \sum_{Q \in \mathcal{C}_2} \iint_{Q'} |U|^2 dx dt \ge \frac{1}{2} M_1^{-1}.
$$

Now for  $Q \in C_1 \setminus C_2$ , we have that

$$
\iint_{Q'}|U|^2dxdt\leq M_1^{-5}\hat{\delta}^{2+n}l^{n+1-\frac{8}{3n}}.
$$

Using this, [\(3.28\)](#page-22-3) and [\(3.30\)](#page-22-4), we get that

$$
(3.31) \qquad \frac{1}{2}M_1^{-1} \leq \sum_{Q \in \mathcal{C}_1 \setminus \mathcal{C}_2} \iint_{Q'} |U|^2 dxdt + \sum_{Q \in \mathcal{C}_2} \iint_{Q'} |U|^2 dxdt
$$
  
 
$$
\leq (\#\mathcal{C}_1 \setminus \mathcal{C}_2)M_1^{-5} \hat{\delta}^{2+n} l^{n+1-\frac{8}{3n}} + C(n)(\#\mathcal{C}_2)M^{p-1}(\hat{\delta}l)^n.
$$

Using [\(3.26\)](#page-22-5) gives that, for appropriately chosen  $M_2 = M^{c_2(p,n)}$ ,

<span id="page-23-1"></span><span id="page-23-0"></span>
$$
(\#\mathcal{C}_1 \setminus \mathcal{C}_2)M_1^{-5}\hat{\delta}^{2+n}l^{n+1-\frac{8}{3n}} \le M_1^{\frac{n}{4}-4}M^{\frac{3n}{8}(p-1)}l^{1-\frac{8}{3n}}
$$
  
=  $M_1^{\frac{n}{4}-4}M^{\frac{3n}{8}(p-1)}M_2^{-(6n+2)(1-\frac{8}{3n})} \le \frac{M_1^{-1}}{4}.$ 

Substituting this into [\(3.31\)](#page-23-0) then gives that

(3.32) 
$$
\#\mathcal{C}_2 \ge \frac{M_1^{-1}M^{1-p}}{4C(n)(\hat{\delta}l)^n}.
$$

Next, note that from the definition of  $C_2$  and Hölder's inequality, we have that for  $Q \in \mathcal{C}_2$ ,

<span id="page-23-2"></span>(3.33) 
$$
\iint_{Q'} |U|^{\frac{3n}{4}} dx dt \geq C_n^* \hat{\delta}^{n+2} l^{\frac{8+5n}{8}} M_1^{-\frac{15n}{8}}.
$$

Noting that  $\bigcup_{Q \in \mathcal{C}_2} Q' \subset B(3M_1) \times 3I$  and that each descendant  $Q'$  from  $\mathcal{C}_2$  intersects with at most  $C_n^{**}$  descendants, we get

<span id="page-23-3"></span>
$$
(3.34) \quad \sum_{Q \in \mathcal{C}_2} \iint_{Q'} |U|^{\frac{3n}{4}} dx dt \leq C_n^{**} \iint_{B(3M_1) \times 3I} |U|^{\frac{3n}{4}} dx dt \leq C_n^{***} M^{\frac{3n(p-1)}{8}} M_1^{\frac{n}{4}},
$$

where the last inequality follows from  $(3.20)$  and Hölder's inequality for Lorentz spaces. Combining  $(3.32)$ ,  $(3.33)$  and  $(3.34)$  gives

$$
\begin{split} & \sum_{Q\in \mathcal{C}_2} \iint_{Q'} |U|^{\frac{3n}{4}} dx dt \geq (\#\mathcal{C}_2) C_n^* \hat{\delta}^{n+2} l^{\frac{8+5n}{8}} M_1^{-\frac{15n}{8}} \geq \frac{C_n^* \hat{\delta}^{n+2} l^{\frac{8+5n}{8}} M_1^{-\frac{15n}{8}-1} M^{1-p}}{4C(n) (\hat{\delta} l)^n}, \\ & \sum_{Q\in \mathcal{C}_2} \iint_{Q'} |U|^{\frac{3n}{4}} dx dt \leq C_n^{***} M^{\frac{3n(p-1)}{8}} M_1^{\frac{n}{4}}. \end{split}
$$

This, together with [\(3.22\)](#page-21-6), implies that

<span id="page-23-4"></span>
$$
(3.35) \tC_n^*(4C(n))^{-1}M_1^{-\frac{15n}{8}-1}M^{1-p}M_2^{\frac{(3n-8)(6n+2)}{8}-2} \leq C_n^{***}M^{\frac{3n(p-1)}{8}}M_1^{\frac{n}{4}}.
$$

Noting that  $((3n-8)(6n+2)/8)-2\geq 1/2$  for  $n\geq 3$  and that  $M_2\geq 1$ , we see that [\(3.35\)](#page-23-4) implies that

<span id="page-23-5"></span>
$$
(3.36) \tC_n^*(4C(n))^{-1}M_1^{-\frac{15n}{8}-1}M^{1-p}M_2^{\frac{1}{2}} \leq C_n^{***}M^{\frac{3n(p-1)}{8}}M_1^{\frac{n}{4}}.
$$

Thus, for  $M_2 = M^{c_2(p,n)}$  chosen appropriately, [\(3.36\)](#page-23-5) gives a contradiction. Hence [\(3.29\)](#page-22-2) cannot occur.

Step 4: conclusion

As [\(3.29\)](#page-22-2) cannot occur, by [\(3.27\)](#page-22-6), we must have

(3.37) 
$$
\sum_{Q \in C_0 \setminus C_1} \iint_{Q'} |U|^2 dx dt \ge \frac{1}{2} M_1^{-1}.
$$

By [\(3.23\)](#page-21-4), we have that

<span id="page-24-1"></span>
$$
\#\mathcal{C}_0 \le \frac{C_n M_1^n}{(\hat{\delta}l)^{n+2}}.
$$

From this,  $(3.37)$ , the pigeonhole principle and the definition of U in  $(3.19)$ , we see that there exists  $Q \in C_0 \setminus C_1$  such that

<span id="page-24-2"></span>
$$
\iint_{Q'} |u_{\lambda}|^{p-1} dx dt \ge \frac{M_1^{-n-1}(\hat{\delta}l)^{n+2}}{2C_n}.
$$

From this, [\(3.16\)](#page-21-0), Lemma [15](#page-19-4) (ii) and [\(3.22\)](#page-21-6), it follows that there exists  $\bar{C} > 0$ depending only on  $n$  and  $p$  and satisfying

(3.38) 
$$
\iint_{Q'} |u_{\lambda}|^2 dx dt \ge M^{-\bar{C}} \quad \text{with } Q \in \mathcal{C}_0 \setminus \mathcal{C}_1
$$

for  $M$  sufficiently large. We also get from  $Q \in C_0 \setminus C_1$  and  $(3.24)$  that

<span id="page-24-3"></span>(3.39) 
$$
||u_{\lambda}||_{L^{\infty}(Q(z_0, l/2))} \leq Cl^{-\frac{2}{p-1}}M^{-2} \text{ with } Q = Q(z_0, l).
$$

Moreover, from the definition of  $C_0$ , we see that for M sufficiently large,

<span id="page-24-4"></span>
$$
(3.40) \tQ = Q(z_0, l) \subset B(5M_1/2) \times (-1/32, -1/256).
$$

Using  $(3.38)$ ,  $(3.39)$  and  $(3.40)$ , we can undo the rescaling given by  $(3.15)$  to get the desired conclusion for the original function  $u$ .

<span id="page-24-0"></span>3.2. Annuli and slices of regularity. We give annuli of quantitative regularity by using Proposition [7](#page-8-2) and arguments in [\[2,](#page-54-5) Section 6]. Compared to the arguments for the 3-dimensional Navier-Stokes equations in [\[2\]](#page-54-5), the lack of analogous energy structure for the energy supercritical nonlinear heat equation prevents us transferring [\(3.41\)](#page-24-5) to a countably additive supercritical bound. We overcome this by applying a Calderón type splitting [\[6\]](#page-54-8). This enables us to apply the  $\varepsilon$ -regularity criterion below the critical exponent (Proposition [7\)](#page-8-2), together with the pigeonhole principle.

<span id="page-24-6"></span>**Proposition 18** (Annuli of regularity). Let  $p > p<sub>S</sub>$ ,  $T<sub>1</sub> > 0$  and  $\lambda > 2$ . Define  $\beta(n,p) := (4(p+1))^{-1} \max(a_3(n,p), a_4(n,p))$  with

$$
a_3(n,p) := 4n(6p^3 + 6p^2 - 6p - 8) + 4n^2(p-1),
$$
  
\n
$$
a_4(n,p) := n^2(p-1) + n(6p^3 + 6p^2 - 7p - 9) + 2(6p^3 + 18p^2 + 19p + 7).
$$

Suppose that u is a classical solution to [\(1.5\)](#page-3-2) on  $\mathbb{R}^n \times [-T_1, 0]$  with

(3.41) 
$$
\sup_{-T_1 < t < 0} \|u(\cdot, t)\|_{L^{q_c, \infty}(\mathbf{R}^n)} \leq M
$$

and M sufficiently large. Then, for any  $R \geq 2M$ , there exists

<span id="page-24-5"></span>
$$
\hat{R} \in [R, R^{\lambda^{M^{q_c+1+\beta(n,p)}}}]
$$

such that

$$
||u||_{L^{\infty}} \left( \left\{ x \in \mathbf{R}^n : 2T_1^{\frac{1}{2}} \hat{R} < |x| < \frac{1}{2} T_1^{\frac{1}{2}}(\hat{R})^{\lambda} \right\} \times \left( -\frac{1}{2} T_1, 0 \right) \right) \leq M^{-1} T_1^{-\frac{1}{p-1}},
$$
\n
$$
||\nabla u||_{L^{\infty}} \left( \left\{ x \in \mathbf{R}^n : 2T_1^{\frac{1}{2}} \hat{R} < |x| < \frac{1}{2} T_1^{\frac{1}{2}}(\hat{R})^{\lambda} \right\} \times \left( -\frac{1}{2} T_1, 0 \right) \right) \leq M^{-1} T_1^{-\frac{p+1}{2(p-1)}}.
$$

Proof. By means of the rescaling

$$
u_{T_1}(x,t) := T_1^{\frac{1}{p-1}} u(T_1^{\frac{1}{2}}x, T_1t),
$$

we can assume without loss of generality that  $T_1 = 1$ .

Let us decompose

(3.42) 
$$
u = u_- + u_+
$$
 with  $u_-(x,t) := \chi_{\{(x,t); |u(x,t)| \leq 1\}}(x,t)u(x,t).$   
Then by  $(q_c + p + 1)/2 < q_c < 2q_c$  and [15, p.22], we get that

<span id="page-25-1"></span>
$$
||u_{-}(\cdot,t)||_{L^{2q_c}(\mathbf{R}^n)}^{2q_c} + ||u_{+}(\cdot,t)||_{L^{\frac{q_c+p+1}{2}}(\mathbf{R}^n)}^{\frac{q_c+p+1}{2}} \le CM^{q_c}
$$

for  $t \in [-1,0]$ . Thus,

$$
\sum_{k=0}^{\infty} \int_{-1}^{0} \int_{R^{\lambda^k} < |x| < R^{\lambda^{k+1}}} |u_-(x,t)|^{2q_c} + |u_+(x,t)|^{\frac{q_c+p+1}{2}} dx dt \le CM^{q_c}.
$$

By the pigeonhole principle, there exists  $k_0 \in \{0, 1, \ldots, \lceil M^{q_c+1+\beta(n,p)} \rceil\}$  such that for  $M$  sufficiently large,

<span id="page-25-0"></span>
$$
(3.43) \qquad \int_{-1}^{0} \int_{R^{\lambda^{k_0}} < |x| < R^{\lambda^{k_0+1}}} |u - (x, t)|^{2q_c} + |u + (x, t)|^{\frac{q_c + p + 1}{2}} dx dt \le M^{-\beta(n, p)}.
$$

Here  $[c]$  is the least integer with  $[c] \geq c$ . Now fix

$$
y_0 \in \{2R^{\lambda^{k_0}} < |x| < (1/2)R^{\lambda^{k_0+1}}\}, \quad s_0 \in (-1/2, 0).
$$

As M is sufficiently large,  $\lambda > 2$  and  $R \ge 2M$ , we see that

$$
B(y_0, M) \subset \{R^{\lambda^{k_0}} < |x| < R^{\lambda^{k_0+1}}\} \quad \text{for } y_0 \in \{2R^{\lambda^{k_0}} < |x| < (1/2)R^{\lambda^{k_0+1}}\}.
$$
\nUsing this and (3.43), we get that

$$
\int_{-1}^{0} \int_{B(y_0,M)} |u_-(x,t)|^{2q_c} + |u_+(x,t)|^{\frac{q_c+p+1}{2}} dx dt \le M^{-\beta(n,p)}.
$$

Applying Hölder's inequality and using  $(3.42)$  gives that

$$
\int_{-1}^{0} \int_{B(y_0,M)} |u|^{p+1} dx dt \le C \left( M^{n-(1+\frac{\beta(n,p)}{n})\frac{p+1}{p-1}} + M^{-\frac{4\beta(n,p)(p+1)+n(n(p-1)-2(p+1))}{n(p-1)+2(p+1)}} \right)
$$

.

Define the rescaled function  $\bar{u}: \mathbf{R}^n \times [-1,0] \to \mathbf{R}$  by

(3.44) 
$$
\bar{u}(x,t) := 2^{-\frac{1}{p-1}} u(2^{-\frac{1}{2}}x + y_0, 2^{-1}t + s_0).
$$

Then  $\bar{u}$  satisfies [\(1.6\)](#page-3-4) and

<span id="page-25-2"></span>
$$
5^{n-\frac{4}{p-1}}\int_{-1/25}^{-1/50}\int_{B(M)}|\bar{u}|^{p+1}dxdt
$$
  
\$\leq C\left(M^{n-(1+\frac{\beta(n,p)}{n})\frac{p+1}{p-1}}+M^{-\frac{4\beta(n,p)(p+1)+n(n(p-1)-2(p+1))}{n(p-1)+2(p+1)}}\right).

Using this and the definition of  $\beta(n, p)$  in Proposition [18,](#page-24-6) we get that for M sufficiently large

$$
5^{n-\frac{4}{p-1}} \int_{-1/25}^{-1/50} \int_{B(M)} |\bar{u}|^{p+1} dx dt \le M^{-6(p+1)^2}.
$$

For M sufficiently large this enables us to apply Proposition [7](#page-8-2) with  $(x_0, t_0) = (0, 0)$ ,  $\delta = 1/5, \, \varepsilon = M^{-6(p+1)^2}$  and  $A = 2(24 \log(M^{p+1}/\varepsilon))^{1/2}$ . This then gives that for M sufficiently large,

$$
|\bar{u}(0,0)| \le CM^{-2}, \quad |\nabla \bar{u}(0,0)| \le CM^{-2}.
$$

Bearing in mind the definition of  $\bar{u}$  in [\(3.44\)](#page-25-2), this gives for M sufficiently large,

$$
|u(y_0, s_0)| \le M^{-1}
$$
,  $|\nabla u(y_0, s_0)| \le M^{-1}$ .

As y<sub>0</sub> and s<sub>0</sub> were taken arbitrarily in  $\{2R^{\lambda^{k_0}} < |x| < (1/2)R^{\lambda^{k_0+1}}\} \times (-1/2,0),$ we then obtain the desired conclusion with  $\hat{R} := R^{\lambda^{k_0}}$ .

We give a slice of regularity based on Corollary [16,](#page-19-3) which is inspired by [\[31,](#page-55-0) Proposition 3.5]. Compared to [\[31\]](#page-55-0), we cannot use countable additivity of the scaleinvariant bound to obtain a quantity to which we can obtain a slice of regularity via the pigeonhole principle and  $\varepsilon$ -regularity. As was the case in Proposition [18,](#page-24-6) we overcome this by applying a Calderón type splitting [\[6\]](#page-54-8).

<span id="page-26-2"></span>**Proposition 19** (Slice of regularity). Let  $p > p<sub>S</sub>$ ,  $T<sub>1</sub> > 0$ ,  $z<sub>0</sub> = (x<sub>0</sub>, t<sub>0</sub>) \in \mathbb{R}^n \times$  $[-T_1/2, 0]$  and  $R \le (T_1/4)^{1/2}$ . Define  $\gamma := \max(2, a_5(n, p), a_6(n, p))$  with

$$
a_5(n,p) := \frac{p-1}{p+3} \left( q_c + n + 7 + 12p + 6p^2 + \frac{2(p+1)^3 - 4}{p-1} \right),
$$
  

$$
a_6(n,p) := \frac{\frac{(p+1)q_c}{n(p-1) - (p+1)} + n + 7 + 12p + 6p^2 + \frac{2(p+1)^3 - 4}{p-1}}{\frac{2(p+1)}{p-1} - \frac{p+1}{n(p-1) - (p+1)}}.
$$

Suppose that u is a classical solution to [\(1.5\)](#page-3-2) on  $\mathbb{R}^n \times [-T_1, 0]$  satisfying [\(3.41\)](#page-24-5) and with M sufficiently large. Then there exist a direction  $\theta \in \mathbf{S}^{n-1}$  and a time interval  $I \subset [t_0 - R^2, t_0]$  with  $|I| = (R/M^{\gamma})^2$  such that within the slice

$$
S = \left\{ x \in \mathbf{R}^n; \begin{aligned} &\text{dist}(x, x_0 + \mathbf{R}_+ \theta) \le 10M^{-\gamma} |(x - x_0) \cdot \theta|, \\ &|x - x_0| \ge 20R \end{aligned} \right\} \times I
$$
\n
$$
\subset \mathbf{R}^n \times [-T_1, 0],
$$

the following estimates holds:

<span id="page-26-1"></span>
$$
(3.45) \t\t ||u||_{L^{\infty}(S)} \leq M^{-1} \left(\frac{R}{M^{\gamma}}\right)^{-\frac{2}{p-1}}, \t ||\nabla u||_{L^{\infty}(S)} \leq M^{-1} \left(\frac{R}{M^{\gamma}}\right)^{-\frac{p+1}{p-1}}.
$$

Proof. By means of the rescaling

<span id="page-26-0"></span>
$$
u_R(y,t) = R^{\frac{2}{p-1}} u(Ry + x_0, R^2t + t_0),
$$

we can take  $z_0 = (x_0, t_0) = (0, 0)$  and  $R = 1$  without loss of generality. Let us decompose

(3.46) 
$$
u = u_{-} + u_{+} \quad \text{with } u_{-}(x,t) := \chi_{\{(x,t); |u(x,t)| \le 1\}}(x,t)u(x,t).
$$
  
Then by  $p + 1 < q_c < n(p-1) - (p+1)$  and [15, p.22], we get that for  $t \in [-1,0]$ ,  

$$
||u_{-}(\cdot,t)||_{L^{n}(p-1)-(p+1)}^{n(p-1)-(p+1)}(\mathbf{R}^n) + ||u_{+}(\cdot,t)||_{L^{p+1}(\mathbf{R}^n)}^{p+1} \le CM^{q_c}.
$$

Thus,

<span id="page-27-1"></span>
$$
(3.47) \qquad \int_{-1}^{0} \int_{\mathbf{R}^n} (|u_-(x,t)|^{n(p-1)-(p+1)} + |u_+(x,t)|^{p+1}) dx dt \le CM^{q_c}.
$$

Consider the collection of space-time slices of the form

<span id="page-27-0"></span>(3.48) 
$$
S^* = \{x \in \mathbb{R}^n; \text{ dist}(x, \mathbb{R}_+ \theta) \le 20M^{-\gamma} |x \cdot \theta|, |x| \ge 10\}
$$

$$
\times [-10M^{-2\gamma}k, -10M^{-2\gamma}(k-1)]
$$

with  $\theta \in \mathbf{S}^{n-1}$  and for  $k \in \{1, ..., \lfloor (1/10)M^{2\gamma} \rfloor\}$ , where  $\lfloor c \rfloor$  is the greatest integer with  $|c| \leq c$ . We may find a collection S of disjoint slices of the form [\(3.48\)](#page-27-0) within  $\mathbf{R}^n \times [-1,0]$  such that

$$
\#\mathcal{S} \ge \frac{M^{2\gamma} M^{\gamma(n-1)}}{C(n)} = \frac{M^{\gamma(n+1)}}{C(n)}.
$$

Thus [\(3.47\)](#page-27-1) and the pigeonhole principle implies that there exists a slice  $S^* \in \mathcal{S}$  of the form

$$
S^* = S_x^* \times [-10M^{-2\gamma}k, -10M^{-2\gamma}(k-1)],
$$
  
\n
$$
S_x^* := \{ x \in \mathbf{R}^n; \text{ dist}(x, \mathbf{R}_+\theta) \le 20M^{-\gamma}|x \cdot \theta|, |x| \ge 10 \}
$$

such that

$$
\int_{-10M^{-2\gamma}k}^{-10M^{-2\gamma}(k-1)} \int_{S_x^*} (|u_-|^{n(p-1)-(p+1)} + |u_+|^{p+1}) dx dt \le CM^{q_c - (n+1)\gamma}.
$$

Now fix

(3.49) 
$$
y \in \{x \in \mathbb{R}^n; \text{ dist}(x, \mathbb{R}_+ \theta) \le 10M^{-\gamma} |x \cdot \theta|, |x| \ge 20\},\
$$

$$
s \in (-10M^{-2\gamma}(k-1) - M^{-2\gamma}, -10M^{-2\gamma}(k-1)).
$$

For such  $(y, s)$ , we have

<span id="page-27-3"></span>
$$
Q((y,s),M^{-\gamma}) \subset S_x^* \times (-10M^{-2\gamma}(k-1) - M^{-2\gamma}, -10M^{-2\gamma}(k-1)).
$$

Thus,

$$
\iint_{Q((y,s),M^{-\gamma})} (|u_-|^{n(p-1)-(p+1)} + |u_+|^{p+1}) dx dt \le CM^{q_c-(n+1)\gamma}.
$$

Using Hölder's inequality, [\(3.46\)](#page-26-0) and the definition of  $\gamma$  in the statement of the proposition gives that for  $M$  sufficiently large,

$$
(3.50)
$$
\n
$$
M^{(n-\frac{4}{p-1})\gamma} \iint_{Q((y,s),M^{-\gamma})} |u|^{p+1} dx dt
$$
\n
$$
\leq C \left( M^{q_c - \frac{(p+1)(q_c + \gamma)}{p-1}} + M^{\frac{(p+1)(q_c + \gamma)}{n(p-1) - (p+1)}} - \frac{2(p+1)\gamma}{p-1} \right)
$$
\n
$$
\leq CM^{-(n+7+12p+6p^2 + \frac{2(p+1)^3 - 4}{p-1})}.
$$

Noting that, as  $\gamma \geq 2$ , taking  $M^{-\gamma} < (2M)^{-1}$  for M sufficiently large, Thus, for M sufficiently large, [\(3.50\)](#page-27-2) allows us to apply Corollary [16](#page-19-3) with  $\delta = M^{-\gamma}$  giving

<span id="page-27-2"></span>
$$
|u(y,s)| \le M^{-1} \left(\frac{1}{M^{\gamma}}\right)^{-\frac{2}{p-1}}, \quad |\nabla u(y,s)| \le M^{-1} \left(\frac{1}{M^{\gamma}}\right)^{-\frac{p+1}{p-1}}
$$

.

As  $(y, s)$  are chosen to be any space-time points satisfying  $(3.49)$ , we obtain the desired conclusion [\(3.45\)](#page-26-1) for  $z_0 = (x_0, t_0) = (0, 0)$  and  $R = 1$ .

## 4. Carleman inequalities

<span id="page-28-0"></span>The quantitative unique continuation Carleman inequality we use is a higherdimensional analogue of that given by Tao [\[38,](#page-56-0) Proposition 4.3], in a lower regularity setting. The quantitative backward uniqueness Carleman inequality that we use is from Palasek [\[30,](#page-55-27) Proposition 9], in a lower regularity setting (and with a certain adjustment of indices). We require a lower regularity setting  $(C^{2,1})$  compared to previous works, since bounded solutions to the nonlinear heat equation with nonsmooth nonlinearity are not necessarily  $C^{\infty}$ , in contrast with the Navier-Stokes equations, for example.

The proof of these Carleman inequalities (in both the smooth and lower regularity settings) hinges on the following general inequality proven by Tao [\[38,](#page-56-0) Lemma 4.1]. Throughout this section, let L be the backward heat operator

<span id="page-28-1"></span>
$$
L:=\partial_t+\Delta.
$$

<span id="page-28-3"></span>**Lemma 20.** (General Carleman inequality, [\[38,](#page-56-0) Lemma 4.1]) Let  $g: \mathbb{R}^n \times [t_1, t_2] \rightarrow$ **R** be smooth and let  $D^2g$  be the bilinear form expressed in coordinates (with usual summation convention) as

(4.1) 
$$
D^2g(a,b) := (\partial_i \partial_j g) a_{ik} b_{jk}.
$$

Let  $F: \mathbf{R}^n \times [t_1, t_2] \to \mathbf{R}$  denote the function

(4.2) 
$$
F := \partial_t g - \Delta g - |\nabla g|^2.
$$

Then, for any vector-valued function  $v \in C_0^{\infty}(\mathbf{R}^n \times [t_1, t_2]; \mathbf{R}^m)$ ,

<span id="page-28-2"></span>
$$
\frac{d}{dt} \int_{\mathbf{R}^n} \left( |\nabla v|^2 + \frac{1}{2} F|v|^2 \right) e^g dx
$$
\n
$$
\geq \int_{\mathbf{R}^n} \left( \frac{1}{2} (LF)|v|^2 + 2D^2 g(\nabla v, \nabla v) - \frac{1}{2} |Lv|^2 \right) e^g dx
$$

for all  $t \in [t_1, t_2]$ . Moreover, v also satisfies

$$
\int_{t_1}^{t_2} \int_{\mathbf{R}^n} \left( \frac{1}{2} (LF) |v|^2 + 2D^2 g(\nabla v, \nabla v) \right) e^g dx dt
$$
  

$$
\leq \frac{1}{2} \int_{t_1}^{t_2} \int_{\mathbf{R}^n} |Lv|^2 e^g dx dt + \int_{\mathbf{R}^n} \left( |\nabla v|^2 + \frac{1}{2} F |v|^2 \right) e^g dx \Big|_{t=t_1}^{t=t_2}.
$$

We state the quantitative backward uniqueness and unique continuation Carleman inequalities, and then we show an iterated quantitative unique continuation.

<span id="page-28-6"></span>**Proposition 21** (Backward uniqueness Carleman inequality). Let  $0 < r_- < r_+ <$  $\infty$  and  $0 < T_1 < \infty$ . Define a space-time annulus A by

<span id="page-28-5"></span><span id="page-28-4"></span>
$$
\mathcal{A} := \{ (x, t) \in \mathbf{R}^n \times \mathbf{R}; \ r_- \le |x| \le r_+, \ t \in [-T_1, 0] \}.
$$

Let  $w \in C^{2,1}(\mathcal{A})$  satisfy the differential inequality

(4.3) 
$$
|\partial_t w - \Delta w| \le \frac{|w(x,t)|}{C_{\text{Carl}}T_1} + \frac{|\nabla w(x,t)|}{(C_{\text{Carl}}T_1)^{1/2}} \quad on \mathcal{A}
$$

with a constant  $C_{\text{Carl}} \geq 3n$ . Assume

(4.4) r 2 <sup>−</sup> ≥ 4CCarlT1.

Then there exists  $C(n) > 0$  depending only on n such that

$$
\int_{-T_1/4}^{0} \int_{10r - \leq |x| \leq r+/2} \left( \frac{|w(x,t)|^2}{T_1} + |\nabla w(x,t)|^2 \right) dx dt
$$
  
 
$$
\leq C(n) C_{\text{Carl}} e^{-\frac{r_- \cdot r_+}{4C_{\text{Carl}} T_1}} X + C(n) C_{\text{Carl}} e^{-\frac{r_- \cdot r_+}{4C_{\text{Carl}} T_1} + \frac{2r_+^2}{C_{\text{Carl}} T_1} Y},
$$

where

$$
\begin{split} X &:= \iint_{\mathcal{A}} e^{\frac{2|x|^2}{C_{\text{Carl}}T_1}} \left( \frac{|w(x,t)|^2}{T_1} + |\nabla w(x,t)|^2 \right) dxdt, \\ Y &:= \int_{r_- \leq |x| \leq r_+} |w(x,0)|^2 dx. \end{split}
$$

Proof. Let us briefly outline how the corresponding proof in [\[30,](#page-55-27) Proposition 9] also applies in our less regular setting. Without loss of generality, suppose  $20r-\leq r_+$ . By the change of variables, we have

$$
X = \int_0^{T_1} \int_{r_- \le |x| \le r_+} e^{\frac{2|x|^2}{C_{\text{Carl}} T_1}} \left( \frac{|w(x, -t)|^2}{T_1} + |\nabla w(x, -t)|^2 \right) dx dt.
$$

Then by the pigeonhole principle, there exists  $T_0 \in [T_1/2, 3T_1/4]$  such that

<span id="page-29-1"></span>(4.5) 
$$
\int_{r_{-} \leq |x| \leq r_{+}} e^{\frac{2|x|^{2}}{C_{\text{Carl}}T_{1}}}\left(\frac{|w(x, -T_{0})|^{2}}{T_{1}} + |\nabla w(x, -T_{0})|^{2}\right)dx \leq 4T_{1}^{-1}X.
$$

Let

<span id="page-29-2"></span>(4.6) 
$$
g(x,t) := \frac{r_+(T_0-t)|x|}{2C_{\text{Carl}}(T_1)^2} + \frac{|x|^2}{C_{\text{Carl}}T_1}.
$$

By using the definitions [\(4.1\)](#page-28-1) and [\(4.2\)](#page-28-2), together with  $r^2 \geq 4C_{\text{Carl}}T_1$  and  $C_{\text{Carl}} \geq$ 3n, direct computations readily show that for  $(x, t) \in \{r_{-} \leq |x| \leq r_{+}\} \times [0, T_{0}],$ 

<span id="page-29-0"></span>(4.7) 
$$
F(x,t) \le 0
$$
,  $D^2 g(x,t) \ge \frac{2}{C_{\text{Carl}} T_1} Id$ ,  $LF(x,t) \ge \frac{4}{C_{\text{Carl}}(T_1)^2}$ ,

where Id is the identity matrix. Let  $\psi$  be a smooth spatial cut-off compactly supported in  $\{r_-\leq |x|\leq r_+\}$  that equals 1 in  $\{2r_-\leq |x|\leq r_+/2\}$  and obeys

<span id="page-29-3"></span>(4.8) 
$$
|\nabla^j \psi(x)| \le C(n,j) r_{-}^{-j} \quad \text{for } j = 0, 1, 2.
$$

The main difference in the less regular setting compared with [\[30,](#page-55-27) Proposition 9], is that we do not have sufficient regularity to directly apply Lemma [20](#page-28-3) (with the weight g as above) to  $\psi(x)w(x, -t)$  on  $\mathbb{R}^n \times [0, T_0]$ . Instead, we apply Lemma [20](#page-28-3) with the weight g to  $\psi(x)v_{1/M,1/N}(x,t)$  on  $\mathbb{R}^n \times [\varepsilon, T_0]$ . Here,

$$
v_{\frac{1}{M},\frac{1}{N}}(x,t) := \int_0^{T_1} \omega_{\mathbf{R},\frac{1}{M}}(t-s) \int_{r_{-\leq}|y| \leq r_{+}} \omega_{\mathbf{R}^n,\frac{1}{N}}(x-y) w(y,-s) dy ds
$$

and  $\omega_{\mathbf{R},1/M}$  (resp.  $\omega_{\mathbf{R}^n,1/N}$ ) denotes a standard mollifier on  $\mathbf{R}$  (resp.  $\mathbf{R}^n$ ) supported in  $[-1/M, 1/M]$  (resp.  $\{|x| \le 1/N\}$ ).

Applying Lemma [20,](#page-28-3) together with using [\(4.7\)](#page-29-0), gives

$$
\int_{\varepsilon}^{T_0} \int_{\mathbf{R}^n} \left( \frac{2}{C_{\text{Carl}}(T_1)^2} |\psi v_{\frac{1}{M},\frac{1}{N}}|^2 + \frac{4}{C_{\text{Carl}}T_1} |\nabla(\psi v_{\frac{1}{M},\frac{1}{N}})|^2 \right) e^g dxdt
$$
  
\n
$$
\leq \frac{1}{2} \int_{\varepsilon}^{T_0} \int_{\mathbf{R}^n} |L(\psi v_{\frac{1}{M},\frac{1}{N}})|^2 e^g dxdt + \int_{\mathbf{R}^n} |\nabla(\psi v_{\frac{1}{M},\frac{1}{N}})(x,T_0)|^2 e^{g(x,T_0)} dx
$$
  
\n
$$
+ \frac{1}{2} \int_{\mathbf{R}^n} |F(x,\varepsilon)| |\psi v_{\frac{1}{M},\frac{1}{N}}(x,\varepsilon)|^2 e^{g(x,\varepsilon)} dx.
$$

Now we take the limit  $N \to \infty$ , then  $M \to \infty$  and finally  $\varepsilon \to 0$ . Using  $w \in C^{2,1}(\mathcal{A}),$  $T_0 \in [T_1/2, 3T_1/4]$  and standard properties of mollification yield that

$$
\int_0^{T_0} \int_{\mathbf{R}^n} \left( \frac{2}{C_{\text{Carl}}(T_1)^2} |\psi w(x, -t)|^2 + \frac{4}{C_{\text{Carl}}(T_1)} |\nabla(\psi w(x, -t))|^2 \right) e^g dx dt
$$
  
\n
$$
\leq \frac{1}{2} \int_0^{T_0} \int_{\mathbf{R}^n} |L(\psi w(x, -t))|^2 e^g dx dt + \int_{\mathbf{R}^n} |\nabla(\psi w)(x, -T_0)|^2 e^{g(x, T_0)} dx
$$
  
\n
$$
+ \int_{\mathbf{R}^n} |F(x, 0)| |\psi w(x, 0)|^2 e^{g(x, 0)} dx.
$$

This together with [\(4.3\)](#page-28-4) (with  $C_{\text{Carl}} \geq 3n$ ), [\(4.4\)](#page-28-5), [\(4.5\)](#page-29-1), [\(4.6\)](#page-29-2) and [\(4.8\)](#page-29-3) allows us to apply the same arguments used in [\[30,](#page-55-27) Proposition 9] (particularly, [\[30,](#page-55-27) pp. 1521–1523]), which then yields the desired conclusion. In particular, the reader can readily check that  $w \in C^{2,1}(\mathcal{A})$  is sufficient regularity for this final step.

We remark that the setting in [\[30,](#page-55-27) Proposition 9] concerns  $x \in \mathbb{R}^{d_1+d_2}$ , with the (anisotropic) annuli in the quantitative backward uniqueness Carleman inequality being in  $\mathbb{R}^{d_1}$ . This is more general than we require and we can consider the reasoning in [\[30,](#page-55-27) Proposition 9] with just  $d_1 = n$  and  $d_2 = 0$ . Moreover, we just require that the time derivative and second spatial derivatives are square integrable to perform the final step, along with  $(4.3)$ .

<span id="page-30-2"></span>**Proposition 22** (Unique continuation Carleman inequality). Let  $r > 0$ ,  $0 < T_1$  $\infty$  and  $1 \leq C_0 < \infty$ . Define a space-time cylinder C by

<span id="page-30-0"></span>
$$
\mathcal{C} := \{ (x, t) \in \mathbf{R}^n \times \mathbf{R}; \ |x| \le r, \ t \in [-T_1, 0] \}.
$$

Let  $w \in C^{2,1}(\mathcal{C})$  satisfy the differential inequality

(4.9) 
$$
|\partial_t w - \Delta w| \le \frac{|w(x,t)|}{C_0 T_1} + \frac{|\nabla w(x,t)|}{(C_0 T_1)^{1/2}} \quad on \ \mathcal{C}.
$$

Assume

$$
(4.10) \t\t\t r^2 \ge 16000T_1.
$$

Then there exists  $C(n) > 0$  depending only on n such that

<span id="page-30-1"></span>
$$
\int_{-2\overline{s}}^{-\overline{s}} \int_{|x| \le r/2} \left( \frac{|w(x,t)|^2}{T_1} + |\nabla w(x,t)|^2 \right) e^{\frac{|x|^2}{4t}} dx dt
$$
  

$$
\le C(n) e^{-\frac{r^2}{500\overline{s}}} X + C(n) (\overline{s})^{\frac{n}{2}} \left( \frac{3e^{\overline{s}}}{\underline{s}} \right)^{\frac{r^2}{200\overline{s}}} Y
$$

for all  $0 < \underline{s} \leq \overline{s} < T_1/(10000n)$ , where

$$
X := \int_{-T_1}^{0} \int_{|x| \le r} \left( \frac{|w(x,t)|^2}{T_1} + |\nabla w(x,t)|^2 \right) dxdt,
$$
  

$$
Y := \int_{|x| \le r} |w(x,0)|^2 \, (s)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4s}} dx.
$$

Proof. Let us briefly outline the modification required to the proof of [\[38,](#page-56-0) Proposition 4.3] to our setting with general dimension and lower regularity. Clearly,

$$
X = \int_0^{T_1} \int_{|x| \le r} \left( \frac{|w(x, -t)|^2}{T_1} + |\nabla w(x, -t)|^2 \right) dx dt
$$

and by the pigeonhole principle there exists  $S_0 \in [T_1/200, T_1/100]$  such that

(4.11) 
$$
\int_{|x| \le r} \left( \frac{|w(x, -S_0)|^2}{T_1} + |\nabla w(x, -S_0)|^2 \right) dx \le 200 T_1^{-1} X.
$$

Let  $\alpha := r^2/(400\bar{s})$  and

<span id="page-31-2"></span>
$$
g(x,t) := \frac{-|x|^2}{4(t+1)} - \frac{n}{2}\log(t+1) - \alpha\log\left(\frac{t+1}{S_0 + 1}\right) + \alpha\frac{t+1}{S_0 + 1}.
$$

where we note that in [\[38,](#page-56-0) Proposition 4.3], this weight is considered in the case  $n = 3$ . Using the definitions [\(4.1\)](#page-28-1) and [\(4.2\)](#page-28-2), direct computations readily give that

<span id="page-31-0"></span>(4.12) 
$$
F(x,t) = \frac{\alpha}{S_0 + s} - \frac{\alpha}{t + s}, \quad D^2 g(x,t) = -\frac{1}{2(t + s)} Id,
$$

$$
LF(x,t) = \frac{\alpha}{(t + s)^2}.
$$

Let  $\psi \in C_0^{\infty}(B(r))$  be a smooth spatial cut-off such that  $\psi = 1$  on  $B(0, r/2)$  and

 $|\nabla^j \psi(x)| \le C(n,j)r^{-j}$  for  $j = 0, 1, 2$ .

The main difference in the less regular setting compared with [\[38,](#page-56-0) Proposition 4.3], is that we do not have sufficient regularity to directly apply Lemma [20](#page-28-3) (with the weight g as above) to  $\psi(x)w(x, -t)$  on  $\mathbb{R}^n \times [0, S_0]$ . Instead, we apply Lemma [20](#page-28-3) with the weight g to  $\psi v_{1/M,1/N}(x,t)$  on  $\mathbb{R}^n \times [0, S_0]$ . Here,

$$
v_{\frac{1}{M},\frac{1}{N}}(x,t):=\int_0^{T_1}\omega_{\mathbf{R},\frac{1}{M}}(t-s)\int_{|y|\leq r}\omega_{\mathbf{R}^n,\frac{1}{N}}(x-y)w(y,-s)dyds.
$$

Applying Lemma [20,](#page-28-3) together with using [\(4.12\)](#page-31-0), gives that for  $t \in [0, S_0]$ ,

<span id="page-31-1"></span>
$$
(4.13) \quad \frac{dE_{M,N}}{dt}(t) \ge \int_{\mathbf{R}^n} \left( \frac{\alpha |\psi v_{\frac{1}{M},\frac{1}{N}}|^2}{2(t+\underline{s})^2} - \frac{|\nabla(\psi v_{\frac{1}{M},\frac{1}{N}})|^2}{t+\underline{s}} - \frac{|L(\psi v_{\frac{1}{M},\frac{1}{N}})|^2}{2} \right) e^g dx,
$$

$$
E_{M,N}(t) := \int_{\mathbf{R}^n} \left( |\nabla(\psi v_{\frac{1}{M},\frac{1}{N}})|^2 + \frac{1}{2} \left( \frac{\alpha}{S_0 + \underline{s}} - \frac{\alpha}{t+\underline{s}} \right) |\psi v_{\frac{1}{M},\frac{1}{N}}|^2 \right) e^g dx.
$$

Using [\(4.13\)](#page-31-1),  $S_0 \in [T_1/200, T_1/100]$  and  $0 < s \leq \overline{s} < T_1/(10000n)$ , we get that for  $t \in [0, S_0],$ 

<span id="page-32-0"></span>
$$
(4.14) \frac{d}{dt} \left( \left( (t+s) + \frac{(t+s)2}{10S_0} \right) E_{M,N}(t) \right)
$$
  

$$
\geq \int_{\mathbf{R}^n} \left( \frac{\alpha |\psi v_{\frac{1}{M},\frac{1}{N}}|^2}{10(S_0+\underline{s})} + \frac{(t+s)|\nabla(\psi v_{\frac{1}{M},\frac{1}{N}})|^2}{10(S_0+\underline{s})} - (t+s)|L(\psi v_{\frac{1}{M},\frac{1}{N}})|^2 \right) e^g dx.
$$

Now take  $\varepsilon \in (0, S_0]$ . Using this, [\(4.14\)](#page-32-0) and the fundamental theorem of calculus, one can deduce that

$$
\int_{\varepsilon}^{S_{0}} \int_{\mathbf{R}^{n}} \left( \frac{\alpha |\psi v_{\frac{1}{M}, \frac{1}{N}}|^{2}}{10(S_{0} + \underline{s})} + \frac{(t + \underline{s})|\nabla(\psi v_{\frac{1}{M}, \frac{1}{N}})|^{2}}{10(S_{0} + \underline{s})} \right) e^{g} dx dt
$$
\n
$$
\leq \int_{\varepsilon}^{S_{0}} \int_{\mathbf{R}^{n}} (t + \underline{s}) |L(\psi v_{\frac{1}{M}, \frac{1}{N}})|^{2} e^{g} dx dt + \left( \left( (t + \underline{s}) + \frac{(t + \underline{s})^{2}}{10S_{0}} \right) E_{M, N}(t) \right) \Big|_{t = \varepsilon}^{t = S_{0}}
$$
\n
$$
\leq \int_{\varepsilon}^{S_{0}} \int_{\mathbf{R}^{n}} (t + \underline{s}) |L(\psi v_{\frac{1}{M}, \frac{1}{N}})|^{2} e^{g} dx + C(n) S_{0} \int_{\mathbf{R}^{n}} |\nabla(\psi v_{\frac{1}{M}, \frac{1}{N}})(x, S_{0})|^{2} e^{g(x, S_{0})} dx
$$
\n
$$
+ C(n) \alpha \int_{\mathbf{R}^{n}} |\psi v_{\frac{1}{M}, \frac{1}{N}}(x, \varepsilon)|^{2} e^{g(x, \varepsilon)} dx.
$$

We now take the limit of the above inequality as  $N \to \infty$ ,  $M \to \infty$  and finally  $\varepsilon \to 0$ . Using that  $w \in C^{2,1}(\mathcal{C}), S_0 \in [T_1/200, T_1/100]$  and standard properties of mollification yields that

$$
\int_{0}^{S_{0}} \int_{\mathbf{R}^{n}} \left( \frac{\alpha}{10(S_{0}+s)} |\psi w(x,-t)|^{2} + \frac{t+s}{10(S_{0}+s)} |\nabla(\psi w(x,-t))|^{2} \right) e^{g} dx dt
$$
  
\n
$$
\leq \int_{0}^{S_{0}} \int_{\mathbf{R}^{n}} (t+s) |L(\psi w(x,-t))|^{2} e^{g} dx + C(n) S_{0} \int_{\mathbf{R}^{n}} |\nabla(\psi w)(x,-S_{0})|^{2} e^{g(x,S_{0})} dx
$$
  
\n
$$
+ C(n) \alpha \int_{\mathbf{R}^{n}} |\psi w(x,0)|^{2} e^{g(x,0)} dx.
$$

This together with [\(4.9\)](#page-30-0) (with  $C_0 \ge 1$ ), [\(4.10\)](#page-30-1),  $0 < s \le \bar{s} < T_1/(10000n)$  and [\(4.11\)](#page-31-2) allows us to apply similar arguments used in [\[38,](#page-56-0) Proposition 4.3] (particularly, from [\[38,](#page-56-0) (4.17)]), albeit in the setting with general dimension. Then we obtain the desired conclusion. In particular, the reader can readily check that  $w \in C^{2,1}(\mathcal{C})$  is sufficient regularity for this final step.  $\Box$ 

We will use the terminology ' $\eta$  being sufficiently small' to mean that  $0 < \eta \leq$  $\eta_0(n, p)$ , where  $\eta_0$  is positive and sufficiently small. Let us prove the iterated quantitative unique continuation, which is based upon [\[31,](#page-55-0) Proposition 4.2] that corresponds to the case  $p = 2$ .

<span id="page-32-1"></span>**Proposition 23** (Iterated unique continuation). Let  $p > 1$ ,  $0 < \eta \leq 1$ ,  $0 < T_1$  $\infty$ ,  $1 \leq C_0 < \infty$  and  $\theta \in \mathbf{S}^{n-1}$ . Define

<span id="page-32-2"></span>(4.15) 
$$
S := \{x \in \mathbf{R}^n : |x| > 10T_1^{\frac{1}{2}}, \text{ dist}(x, \mathbf{R}_+ \theta) \le \eta |x \cdot \theta| \} \times [-\eta T_1, 0].
$$

Let  $w \in C^{2,1}(S)$  satisfy

(4.16) 
$$
\|\nabla^j w\|_{L^{\infty}(S)} \le (\eta T_1)^{-\frac{1}{p-1} - \frac{j}{2}} \quad \text{for } j = 0, 1,
$$

$$
|\partial_t w - \Delta w| \le \frac{|w(x, t)|}{C_0 \eta T_1} + \frac{|\nabla w(x, t)|}{(C_0 \eta T_1)^{1/2}} \quad \text{on } S.
$$

Assume that for every  $t \in [-\eta T_1, 0],$ 

<span id="page-33-3"></span><span id="page-33-0"></span>
$$
\int_{B(R_0 \theta, \eta^5 R_0)} |w(x, t)|^2 dx \ge \epsilon T_1^{\frac{n}{2} - \frac{2}{p-1}},
$$

where  $20T_1^{1/2} \le R_0 \le \eta^{-2}T_1^{1/2}$  and  $0 < \epsilon \le \eta^8$ . Then, for  $\eta$  sufficiently small,

(4.17) 
$$
\int_{B(R\theta,\eta^2 R)} |w(x,t)|^2 dx \geq \epsilon \left(\frac{R}{R_0}\right)^{\eta^{-4}} T_1^{\frac{n}{2} - \frac{2}{p-1}}
$$

for all  $t \in [-\eta T_1/2, 0]$  and  $R > R_0$ .

<span id="page-33-1"></span>To prove Proposition [23,](#page-32-1) we prepare the following lemma:

Lemma 24. Let  $p > 1, 0 < p \le 1, 0 < T_1 < \infty, 1 \le C_0 < \infty, \theta \in S^{n-1}$  and  $a \in (\eta/2, 3\eta/4]$ . Define S by [\(4.15\)](#page-32-2) and let  $w \in C^{2,1}(S)$  satisfy [\(4.16\)](#page-33-0). Assume that

<span id="page-33-2"></span>
$$
\int_{B(R\theta,\eta^5 R)} |w(x,t)|^2 dx \ge \epsilon_0 T_1^{\frac{n}{2} - \frac{2}{p-1}} \quad \text{for every } t \in [-aT_1, 0],
$$
\n
$$
\text{where } R \ge 20T_1^{\frac{1}{2}} \text{ and } 0 < \epsilon_0 \le \min\left(\eta^8, \left(\frac{R^2}{T_1}\right)^{-5000n\eta}, e^{-\frac{40000n\eta^4 R^2}{T_1}}\right)
$$

Then, for  $\eta$  sufficiently small and  $R' = (1 + \eta^3)R$ ,

$$
\int_{B(R'\theta,\eta^5 R')} |w(x,t)|^2 dx \ge \epsilon_0^{\eta^{-2}} T_1^{\frac{n}{2} - \frac{2}{p-1}}
$$

.

for all  $t \in [-aT_1 + 2\eta^5 R^2 (\log(1/\epsilon_0))^{-1}, 0].$ 

Proof. Lemma [24](#page-33-1) is based on [\[31,](#page-55-0) Lemma 4.3]. However, we required minor differences on the condition for a and  $\epsilon_0$  for the statement and proof to hold. For the readers' convenience, we include the proof (following [\[31,](#page-55-0) Lemma 4.3]) with these adjustments.

First note that by means of the rescaling

$$
w_{T_1}(x,t) := T_1^{\frac{1}{p-1}} w(T_1^{\frac{1}{2}}x, T_1t),
$$

we can take  $T_1 = 1$  without loss of generality. Let R be given in [\(4.18\)](#page-33-2). Note also that for  $0 < \eta \le 1$ , the fact that  $0 < \epsilon_0 \le \min(\eta^8, e^{-40000n\eta^4 R^2})$  and  $a \in (\eta/2, 3\eta/4]$ ensures that

$$
-\frac{3}{4}\eta \le -a \le -a + \frac{2\eta^5 R^2}{\log(1/\epsilon_0)} < 0.
$$

Now fix  $t' \in [-a + 2\eta^5 R^2 (\log(1/\epsilon_0))^{-1}, 0]$  and define

$$
r := \eta^2 R
$$
,  $T_c := \min(\eta/4, \eta^5 R^2)$ ,  $t_0 := \frac{\eta^5 R^2}{\log(1/\epsilon_0)}$ ,  $t_1 := \frac{\eta^{15} R^2}{\log(1/\epsilon_0)}$ .

As  $0 < \epsilon_0 \le \eta^8$ ,  $0 < a \le 3\eta/4$  and  $R' = (1 + \eta^3)R$ , for  $\eta$  sufficiently small,

<span id="page-34-0"></span>(4.19) 
$$
B(R'\theta, r) \times [t'-T_c, t'] \subset S,
$$

<span id="page-34-3"></span>
$$
(4.20) \t B(R'\theta, r/2) \supset B(R\theta, \eta^5 R), \t [t'-2t_0, t'-t_0] \subset [-a, 0],
$$

 $|x - R'\theta| \leq 2\eta^3 R$  for  $x \in B(R\theta, \eta^5 R)$ .

As  $0 < \epsilon_0 \le \min(\eta^8, e^{-40000n\eta^4 R^2})$ , we also get that for  $\eta$  sufficiently small,

<span id="page-34-1"></span>(4.21) 
$$
r^2 \ge 16000T_c, \quad 0 < t_1 < t_0 \le \frac{T_c}{10000n}.
$$

Furthermore, from [\(4.16\)](#page-33-0), [\(4.19\)](#page-34-0) and  $T_c \leq \eta/4$ , it follows that

<span id="page-34-2"></span>(4.22) 
$$
|\partial_t w - \Delta w| \le \frac{|w(x,t)|}{4C_0T_c} + \frac{|\nabla w(x,t)|}{(4C_0T_c)^{1/2}} \quad \text{on } B(R'\theta,r) \times [t'-T_c,t'].
$$

Then [\(4.21\)](#page-34-1) and [\(4.22\)](#page-34-2), along with the fact that  $1 \leq C_0 < \infty$ , allows us to apply Proposition [22](#page-30-2) with  $T_c$ ,  $\bar{s} = t_0$  and  $\underline{s} = t_1$  to

$$
(x,t) \mapsto w(x + R'\theta, t + t')
$$
 for  $(x,t) \in B(r) \times [-T_c, 0].$ 

This yields

(4.23) 
$$
Z \leq C(n)e^{-\frac{r^2}{500t_0}}X + C(n)(t_0)^{\frac{n}{2}} \left(\frac{3et_0}{t_1}\right)^{\frac{r^2}{200t_0}} Y,
$$

where

<span id="page-34-5"></span>
$$
X := \int_{t'-T_c}^{t'} \int_{B(R'\theta,r)} (T_c^{-1}|w|^2 + |\nabla w|^2) dx dt,
$$
  
\n
$$
Y := \int_{B(R'\theta,r)} |w(x,t')|^2 (t_1)^{-\frac{n}{2}} e^{-\frac{|x-R'\theta|^2}{4t_1}} dx,
$$
  
\n
$$
Z := \int_{t'-2t_0}^{t'-t_0} \int_{B(R'\theta,r/2)} (T_c^{-1}|w|^2 + |\nabla w|^2) e^{-\frac{|x-R'\theta|^2}{4(t'-t)}} dx dt.
$$

Using [\(4.18\)](#page-33-2) and [\(4.20\)](#page-34-3), recalling that  $t_0 := \eta^5 R^2 (\log(1/\epsilon_0))^{-1}$  and  $T_c \le \eta^5 R^2$ , we see that

$$
Z \ge t_0 T_c^{-1} e^{-\frac{\eta^6 R^2}{t_0}} \epsilon_0 \ge t_0 (\eta^5 R^2)^{-1} e^{-\frac{\eta^6 R^2}{t_0}} \epsilon_0 = \frac{\epsilon_0^2}{\epsilon_0^{1-\eta} \log(1/\epsilon_0)}.
$$

Then, for  $0 < \eta < 1/2$  and  $0 < \epsilon_0 \leq \eta^8$ , we have

<span id="page-34-4"></span>
$$
Z \ge \frac{\epsilon_0^2}{\epsilon_0^{1/2} \log(1/\epsilon_0)}.
$$

As  $\lim_{x\to 0} x^{1/2} \log x = 0$  and  $0 < \epsilon_0 \leq \eta^8$ , we infer that for  $\eta$  sufficiently small,

$$
(4.24) \t\t Z \ge \epsilon_0^2.
$$

Now let us show that  $e^{-r^2/(500t_0)}X$  is much smaller than the lower bound of Z. Using [\(4.16\)](#page-33-0), [\(4.19\)](#page-34-0) and  $T_c \in [0, \eta/4]$ , we get that

$$
e^{-\frac{r^2}{500t_0}}X \leq C(n)\epsilon_0^{\frac{1}{500\eta}}\eta^{2n}R^n\eta^{-\frac{2}{p-1}}.
$$

As  $\epsilon_0 \le \min(\eta^8, R^{-10000n\eta})$  and  $R \ge 20$ , we see that for  $\eta$  sufficiently small,

<span id="page-35-0"></span>
$$
(4.25) \quad e^{-\frac{r^2}{500t_0}} X \leq C(n)\epsilon_0^{\frac{1}{500\eta}} \eta^{2n} R^n \eta^{-\frac{2}{p-1}} = C(n)\epsilon_0^{\frac{1}{1000\eta}} (\epsilon_0^{\frac{1}{2000\eta}} \eta^{2n-\frac{2}{p-1}}) \epsilon_0^{\frac{1}{2000\eta}} R^n
$$
  

$$
\leq C(n)\epsilon_0^{\frac{1}{1000\eta}} (\eta^{\frac{1}{250\eta}+2n-\frac{2}{p-1}}) R^{-4n} \leq C(n)\epsilon_0^{\frac{1}{1000\eta}} (\eta^{\frac{1}{250\eta}+2n-\frac{2}{p-1}}).
$$

As  $\epsilon_0 \leq \eta^8$ , for  $\eta$  sufficiently small, we have that

$$
\epsilon_0 \in (0, 1), \quad \frac{1}{1000\eta} \ge 2, \quad \frac{1}{250\eta} + 2n - \frac{2}{p-1} \ge 3.
$$

Using these and  $(4.25)$ , we get for  $\eta$  sufficiently small that

$$
e^{-\frac{r^2}{500t_0}}X \leq C(n)\epsilon_0^{\frac{1}{1000\eta}}\left(\eta^{\frac{1}{250\eta}+2n-\frac{2}{p-1}}\right) \leq C(n)\epsilon_0^2\eta^3 \leq \epsilon_0^2\eta^2.
$$

From this and [\(4.24\)](#page-34-4), we see that for  $\eta$  sufficiently small,  $e^{-r^2/500t_0}X$  is negligible compared to  $Z$ . Thus  $(4.23)$  gives that

$$
\epsilon_0^2 \le C(n) (3e\eta^{-10})^{\frac{n}{2} + \frac{\log(1/\epsilon_0)}{200\eta}} \int_{B(R'\theta, r)} |w(x, t')|^2 e^{-\frac{|x - R'\theta|^2}{4t_1}} dx.
$$

<span id="page-35-1"></span>Using  $\epsilon_0 \leq \eta^8$  gives for  $\eta$  sufficiently small,

$$
\epsilon_0^2 \le C(n)\eta^{-8n - \frac{8\log(1/\epsilon_0)}{50\eta}} \int_{B(R'\theta,r)} |w(x,t')|^2 e^{-\frac{|x-R'\theta|^2}{4t_1}} dx
$$
\n
$$
\le C(n) \left(\frac{1}{\epsilon_0}\right)^{n + \frac{1}{50\eta} \log(\frac{1}{\eta^8})} \int_{B(R'\theta,r)} |w(x,t')|^2 e^{-\frac{|x-R'\theta|^2}{4t_1}} dx.
$$

For  $\eta$  sufficiently small we have that  $n + (50\eta)^{-1} \log(1/\eta^8) \le 1/(2\eta^2)$ . From this,  $(4.26)$  and  $r = \eta^2 R$ , we get that

(4.27) 
$$
\epsilon_0^2 \leq C(n)\epsilon_0^{-\frac{1}{2\eta^2}}Y_1 + C(n)\epsilon_0^{-\frac{1}{2\eta^2}}Y_2,
$$

where

<span id="page-35-2"></span>
$$
Y_1 := \int_{\eta^5 R' \le |x - R'\theta| < \eta^2 R} |w(x, t')|^2 e^{-\frac{|x - R'\theta|^2}{4t_1}} dx,
$$
\n
$$
Y_2 := \int_{|x - R'\theta| < \eta^5 R'} |w(x, t')|^2 e^{-\frac{|x - R'\theta|^2}{4t_1}} dx.
$$

Using [\(4.16\)](#page-33-0) and  $\epsilon_0 \leq \min(\eta^8, R^{-10000n\eta})$ , we get that

$$
C(n)\epsilon_0^{-\frac{1}{2\eta^2}}Y_1 \leq C(n)\epsilon_0^{-\frac{1}{2\eta^2} + \frac{(1+\eta^3)^2}{4\eta^5}}\eta^{2n-\frac{2}{p-1}}R^n \leq C(n)\epsilon_0^{-\frac{1}{2\eta^2} + \frac{(1+\eta^3)^2}{4\eta^5} - \frac{1}{10000\eta}}\eta^{2n-\frac{2}{p-1}}
$$
  
=  $C(n)\epsilon_0^{-\frac{1}{2\eta^2} + \frac{(1+\eta^3)^2}{8\eta^5} - \frac{1}{10000\eta}}\epsilon_0^{\frac{(1+\eta^3)^2}{8\eta^5}}\eta^{2n-\frac{2}{p-1}}$   
 $\leq C(n)\epsilon_0^{-\frac{1}{2\eta^2} + \frac{(1+\eta^3)^2}{8\eta^5} - \frac{1}{10000\eta}}\eta^{2n-\frac{2}{p-1} + \frac{(1+\eta^3)^2}{\eta^5}}.$ 

Using this and the fact that for  $\eta$  sufficiently small,

$$
-\frac{1}{2\eta^2} + \frac{(1+\eta^3)^2}{8\eta^5} - \frac{1}{10000\eta} \ge 2, \quad 2n - \frac{2}{p-1} + \frac{(1+\eta^3)^2}{\eta^5} \ge 3,
$$

we infer that

$$
C(n)\epsilon_0^{-\frac{1}{2\eta^2}}Y_1 \leq C(n)\epsilon_0^2\eta^3 \leq \epsilon_0^2\eta^2.
$$

Thus, for  $\eta$  sufficiently small,  $C(n)\epsilon_0^{-1/(2\eta^2)}Y_1$  can be absorbed into the left-handside of [\(4.27\)](#page-35-2). Then,

<span id="page-36-0"></span>
$$
(4.28) \qquad \int_{|x-R'\theta|<\eta^5 R'} |w(x,t')|^2 dx \ge \frac{1}{C(n)} \epsilon_0^{2+\frac{1}{2\eta^2}} = \frac{1}{C(n)\epsilon_0^{(1/(2\eta^2))-2}} \epsilon_0^{\eta^{-2}}.
$$

As  $\epsilon_0 \leq \eta^8$ , we see that for  $\eta$  sufficiently small,

$$
\frac{1}{C(n)\epsilon_0^{(1/(2\eta^2))-2}} \ge \frac{1}{C(n)\epsilon_0} \ge 1.
$$

Using this and  $(4.28)$ , we obtain that for  $\eta$  sufficiently small,

$$
\int_{|x-R'\theta|<\eta^5 R'} |w(x,t')|^2 dx \geq \epsilon_0^{\eta^{-2}}.
$$

As  $t' \in [-a + 2\eta^5 R^2 (\log(1/\epsilon_0))^{-1}, 0]$  was taken arbitrarily, we obtain the desired conclusion for  $T_1 = 1$ .

Let us prove the iterated unique continuation.

Proof of Proposition [23.](#page-32-1) Proposition [23](#page-32-1) is based on [\[31,](#page-55-0) Proposition 4.2] however we required the final conclusion [\(4.17\)](#page-33-3) to be over a larger domain of integration for the statement and proof to hold. The different condition on  $a$  in Lemma [24](#page-33-1) also necessitates adjustments in the iterated parameters [\(4.29\)](#page-36-1). For the readers' convenience, we include the proof (following [\[31,](#page-55-0) Proposition 4.2]) with these adjustments.

By means of the rescaling

<span id="page-36-1"></span>
$$
w_{T_1}(x,t) := T_1^{\frac{1}{p-1}} w(T_1^{\frac{1}{2}}x, T_1t),
$$

we can assume without loss of generality that  $T_1 = 1$ . For  $k \in \mathbb{N}$ , define

(4.29) 
$$
\epsilon_k := \epsilon^{\eta^{-2k}}, \qquad R_k := R_0 (1 + \eta^3)^k,
$$

$$
a_0 = \frac{3\eta}{4}, \qquad a_k := \frac{3\eta}{4} - 2 \sum_{i=0}^{k-1} \frac{\eta^5 R_i^2}{\log(1/\epsilon_i)},
$$

where  $R_0$  is given in the assumption of Proposition [23.](#page-32-1) Fix  $R \geq R_0$ . Then there exists  $N\in\mathbf{N}$  such that

(4.30) 
$$
R_N \le R < R_{N+1} \quad \text{with } N = \lfloor \log_{1+\eta^3} \frac{R}{R_0} \rfloor.
$$

Let  $x \in B(R_N \theta, \eta^5 R_N)$ . Then, from [\(4.29\)](#page-36-1) and [\(4.30\)](#page-36-2), it follows that

<span id="page-36-2"></span>
$$
|x - R\theta| \le |x - R_N \theta| + R - R_N < \eta^5 R_N + R_{N+1} - R_N
$$
  
=  $(\eta^5 + \eta^3) R_N \le (\eta^5 + \eta^3) R.$ 

Then, for  $\eta$  sufficiently small,

(4.31) 
$$
B(R\theta, \eta^2 R) \supseteq B(R_N\theta, \eta^5 R_N).
$$

By L'Hôpital's rule

<span id="page-36-3"></span>
$$
\lim_{x \to 0} \frac{x^{\frac{4}{3}} \log x}{\log(1+x)} = 0.
$$

Thus, for all  $\eta$  sufficiently small,

$$
\frac{2\eta^4 \log(\eta^3)}{3\log(1+\eta^3)} \ge -1,
$$

and so

<span id="page-37-0"></span>
$$
\frac{2}{\log_{\eta}(1+\eta^3)} = \frac{2\log(\eta^3)}{3\log(1+\eta^3)} \ge -\eta^{-4}.
$$

From this and  $R \ge R_0$ , we infer that for  $\eta$  sufficiently small,

$$
\eta^{-2N} \leq \eta^{-2\log_{1+\eta^3} \frac{R}{R_0}} = \eta^{-2\frac{\log_{\eta}(R/R_0)}{\log_{\eta}(1+\eta^3)}} = \left(\frac{R}{R_0}\right)^{-\frac{2}{\log_{\eta}(1+\eta^3)}} \leq \left(\frac{R}{R_0}\right)^{\eta^{-4}}.
$$

As  $\epsilon \leq \eta^8$ , we infer from this that for  $\eta$  sufficiently small,

(4.32) 
$$
\epsilon_N := \epsilon^{\eta^{-2N}} \geq \epsilon^{\left(\frac{R}{R_0}\right)^{\eta^{-4}}}
$$

Note that as  $\epsilon \leq \eta^8$  and  $R_0 \leq \eta^{-2}$ , we get for  $\eta$  sufficiently small,

<span id="page-37-1"></span>
$$
0 \le 2\eta^5 \sum_{i=0}^{k-1} \frac{R_i^2}{\log(1/\epsilon_i)} \le \frac{2\eta}{\log(1/\eta^8)} \sum_{i=0}^{k-1} (\eta + \eta^4)^{2i} \le \frac{2\eta(\log(1/\eta^8))^{-1}}{1 - (\eta + \eta^4)^2} < \frac{\eta}{4}
$$

.

for  $k = 1, \ldots, N$ . Thus, for  $\eta$  sufficiently small,  $a_k$  given by [\(4.29\)](#page-36-1) satisfies (4.33)  $a_k \in (\eta/2, 3\eta/4]$  for all  $k = 0, ..., N$ .

<span id="page-37-5"></span>From  $(4.31)$ ,  $(4.32)$  and  $(4.33)$ , we see that to show  $(4.17)$ , it suffices to show

(4.34) 
$$
\int_{B(R_N\theta,\eta^5 R_N)} |w(x,t)|^2 dx \ge \epsilon_N = \epsilon^{\eta^{-2N}} \quad \text{for all } t \in [-a_N,0].
$$

This is achieved by iteratively applying Lemma [24](#page-33-1) for  $k = 0, \ldots, N$ , with the parameters defined in [\(4.29\)](#page-36-1). To show that this can be done, it suffices to show that  $\epsilon_k$  and  $R_k$  satisfy the hypothesis in Lemma [24](#page-33-1) for each  $k = 0, \ldots, N$ , where we note that  $(4.33)$  shows that  $a_k$  satisfies the hypothesis in Lemma [24.](#page-33-1) First it is clear that as  $0 < \epsilon \leq \eta^8$ , for  $\eta$  sufficiently small,

(4.35) 
$$
\epsilon_k := \epsilon^{\eta^{-2k}} \leq \eta^8 \quad \text{for all } k = 0, \dots, N.
$$

Next, let us verify that

(4.36) 
$$
\epsilon_k \le e^{-40000n\eta^4 R_k^2} \quad \text{for all } k = 0, \dots, N.
$$

To see this, first note that for  $\eta$  sufficiently small (independent of k), we have that

<span id="page-37-3"></span><span id="page-37-2"></span>
$$
40000n(\eta + \eta^4)^{2k} \le 8\log(1/\eta) \quad \text{ for all } k = 0, \dots, N.
$$

Using this,  $R_k := R_0(1+\eta^3)^k$ ,  $R_0 \leq \eta^{-2}$  and  $\epsilon \leq \eta^8$ , we get that for all  $k = 0, ..., N$ ,

$$
40000n\eta^{4}R_{k}^{2} \le 40000n(1+\eta^{3})^{2k} \le 8\eta^{-2k}\log(1/\eta) \le \eta^{-2k}\log(1/\epsilon).
$$

Thus, for all  $k = 0, \ldots, N$ , we obtain

$$
\epsilon_k := \epsilon^{\eta^{-2k}} = e^{-\eta^{-2k} \log(1/\epsilon)} \le e^{-40000n\eta^4 R_k^2}.
$$

Finally, let us verify that

(4.37) 
$$
\epsilon_k \le (R_k^2)^{-5000n\eta} \quad \forall k = 0, \dots, N.
$$

First note that for  $\eta$  sufficiently small (independent of k), for all  $k = 0, \ldots, N$ ,

<span id="page-37-4"></span>
$$
(1 + \eta^3)^{2k} \le \frac{1}{\eta^{2k}}, \quad \log\left(\frac{1}{\eta^{2k}}\right) \le \frac{1}{\eta^{2k}}.
$$

Using this and  $\epsilon \leq \eta^8$ , we infer that for  $\eta$  sufficiently small (independent of k) and for all  $k = 0, \ldots, N$ ,

$$
5000n\eta^{2k+1}\log(\eta^{-4}(1+\eta^3)^{2k})
$$
  
= 20000n\eta^{2k+1}\log(1/\eta) + 5000n\eta^{2k+1}\log((1+\eta^3)^{2k})  

$$
\leq 20000n\eta^{2k+1}\log(1/\eta) + 5000n\eta^{2k+1}\log(1/(\eta^{2k}))
$$
\leq 20000n\eta\log(1/\eta) + 5000n\eta \leq 8\log(1/\eta) \leq \log(1/\epsilon).
$$
$$

Using this,  $R_k := R_0(1 + \eta^3)^k$  and the fact that  $R_0 \leq \eta^{-2}$ , for  $\eta$  sufficiently small (independent of k) we have that for all  $k = 0, \ldots, N$ ,

$$
5000n\eta \log(R_k^2) \le 5000n\eta \log(\eta^{-4}(1+\eta^3)^{2k}) \le \eta^{-2k} \log(1/\epsilon).
$$

Thus, for  $\eta$  sufficiently small (independent of k), we have that for all  $k = 0, \ldots, N$ 

$$
\epsilon_k := \epsilon^{\eta^{-2k}} = e^{-\eta^{-2k} \log(1/\epsilon)} \le e^{-5000n\eta \log(R_k^2)} = (R_k^2)^{-5000n\eta}
$$

as required. Combining  $(4.33)$  with  $(4.35)$ ,  $(4.36)$  and  $(4.37)$  means that we can iteratively apply Lemma [24](#page-33-1) for  $k = 0, \ldots, N$ , with the parameters defined in [\(4.29\)](#page-36-1). This gives [\(4.34\)](#page-37-5), which implies [\(4.17\)](#page-33-3) with  $T_1 = 1$  by the aforementioned reasoning. The proof of Proposition [23](#page-32-1) is complete.

# 5. Proof of main quantitative estimate

<span id="page-38-0"></span>To prove our main quantitative estimate (Proposition [5\)](#page-3-3), we show the following proposition by applying the propositions in Sections [3](#page-17-0) and [4.](#page-28-0)

<span id="page-38-3"></span>**Proposition 25.** Let  $n \geq 3$ ,  $p > p_S$  and u be a classical solution of [\(1.5\)](#page-3-2). Then there exist constants  $C_0$ ,  $M_0 > 0$  depending only on n and p such that the following statement holds true. If there exist  $M \geq M_0$ ,  $x_0 \in \mathbb{R}^n$  and  $t' \in (-M^{-p(2p+3)}, 0)$ such that u satisfies [\(1.6\)](#page-3-4) and

<span id="page-38-4"></span>(5.1) 
$$
(-t')^{\frac{2}{p-1}-\frac{n}{2}}\int_{t'}^{t'/2}\int_{B(x_0,A(-t')^{\frac{1}{2}})}|u(x,t)|^{p+1}dxdt > \varepsilon,
$$

where  $\varepsilon$  and A are constants given by [\(3.1\)](#page-17-2), then we conclude that

<span id="page-38-1"></span>(5.2) 
$$
N := \int_{B(x_0, e^{e^{M C_0}})} |u(x,0)|^{q_c} dx \ge Me^{-e^{e^{M C_0}}},
$$

<span id="page-38-2"></span>(5.3) 
$$
-t' \ge \exp\left(-Ne^{e^{e^{M^{C_0}}}}\right).
$$

**Remark 26.** [\(5.2\)](#page-38-1) is equivalent to  $\exp(-Ne^{e^{e^{M C_0}}}) \leq e^{-M}$ , and so the lower bound [\(5.3\)](#page-38-2) in the conclusion is consistent with the upper bound  $-t' < M^{-p(2p+3)}$  in the assumption of Proposition [25.](#page-38-3)

*Proof of Proposition [25.](#page-38-3)* Without loss of generality, we can assume  $x_0 = 0$ . Let  $t' \in (-M^{-p(2p+3)}, 0)$  satisfy [\(5.1\)](#page-38-4), where M is sufficiently large. For the meaning of the terminology 'M being sufficiently large', see just before Proposition [13.](#page-18-1) We divide the proof into 12 steps.

Step 1: backward propagation

<span id="page-38-5"></span>Let  $t'' \in (-1/64, 0)$  satisfy

(5.4) 
$$
-M^{-3} \le t'' \le M^{(p-1)(2p+4)} A^{\frac{n(p-1)}{p+1}} t'.
$$

,

We note that  $(3.2)$  is satisfied. Now we have  $(5.1)$  with  $x_0 = 0$ , the opposite inequality of [\(3.4\)](#page-17-6) with  $x_0 = 0$ . Then the contraposition of Proposition [12](#page-17-7) shows that the opposite inequality of  $(3.3)$  with  $x_0 = 0$  also holds, that is, the estimate

<span id="page-39-0"></span>(5.5) 
$$
(-t'')^{\frac{2}{p-1}-\frac{n}{2}} \int_{t''}^{t''/2} \int_{B(0,A(-t'')^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt > \varepsilon
$$

holds, where  $(5.5)$  is the same as  $(3.6)$ .

Step 2: preparation of slice of regularity

By  $(5.5)$  (or  $(3.6)$ ) and Proposition [13,](#page-18-1) there exist a point

$$
z_* = (x_*, t_*) \in B(0, 20M_1(-t'')^{\frac{1}{2}}) \times (2t'', t''/4)
$$

and backward parabolic cylinders  $Q((x_*, t_* - r^2/8), \hat{\delta}r)$  and  $Q(z_*, r)$  satisfying

<span id="page-39-1"></span>
$$
(5.6) \qquad Q((x_*, t_* - r^2/8), \hat{\delta}r) \subset Q(z_*, r) \subset B(0, 20M_1(-t'')^{\frac{1}{2}}) \times (2t'', t''/4)
$$

with  $r := 8M_2^{-(6n+2)}(-t'')^{1/2}$  and  $\hat{\delta} := M_2^{-1}$  such that

<span id="page-39-2"></span>
$$
(5.7) \t\t ||u||_{L^{\infty}(Q(z_*,r/2))} \le CM^{-2}r^{-\frac{2}{p-1}}, \t ||\nabla u||_{L^{\infty}(Q(z_*,r/2))} \le CM^{-2}r^{-\frac{p+1}{p-1}},
$$
  

$$
(5.8) \t \int_{\Omega} |u|^2 dx dt \ge M_2^{-\bar{C}}(\hat{\delta}r)^{n+2-\frac{4}{p-1}}.
$$

<span id="page-39-3"></span>
$$
.8) \qquad \int_{Q((x_*,t_*-r^2/8),\hat{\delta}r)} |u|^2 dxdt \geq 0
$$

We note that  $(5.6)$ ,  $(5.7)$  and  $(5.8)$  are the same as  $(3.7)$ ,  $(3.8)$  and  $(3.9)$ .

In steps 2-4 we follow the arguments of [\[31\]](#page-55-0) for the higher-dimensional Navier-Stokes equations. This enables the transfer of the lower bound [\(5.8\)](#page-39-3) to large scales in a slice of regularity by using  $(5.7)$ , quantitative unique continuation and its iterated counterpart.

By Proposition [19](#page-26-2) with  $R = \hat{\delta}r$ ,  $T_1 = 1$ ,  $x_0 = x_*$ ,  $t_0 = t_* - (r^2/8) + 100(\hat{\delta}r)^2$ , there exists a direction  $\theta \in \mathbf{S}^{n-1}$  and a time interval  $I \subset [t_* - (r^2/8) + 99(\hat{\delta}r)^2, t_* (r^2/8) + 100(\hat{\delta}r)^2$  with  $|I| = (\hat{\delta}r/M^{\gamma})^2$  such that within the slice

$$
S = S_* \times I \subset \mathbf{R}^n \times [-1, 0],
$$

$$
S_* := \{ x \in \mathbf{R}^n; \text{dist}(x, x_* + \mathbf{R}_+ \theta) \le 10M^{-\gamma} |(x - x_*) \cdot \theta|, \ |x - x_*| \ge 20 \hat{\delta} r \},
$$

we have

<span id="page-39-4"></span>
$$
(5.9) \t\t ||u||_{L^{\infty}(S)} \leq M^{-1} \left(\frac{\hat{\delta}r}{M^{\gamma}}\right)^{-\frac{2}{p-1}}, \t ||\nabla u||_{L^{\infty}(S)} \leq M^{-1} \left(\frac{\hat{\delta}r}{M^{\gamma}}\right)^{-\frac{p+1}{p-1}}.
$$

Here,  $\gamma = \gamma(n, p)$  is defined in Proposition [19.](#page-26-2)

Step 3: unique continuation Carleman inequality

Fix  $\tilde{t} \in I$ . Let us apply Proposition [22](#page-30-2) for

 $u_1(x,t) := u(x + x_* + 30\hat{\delta}r\theta, t + \tilde{t}), \quad x \in \overline{B(r/4)}, t \in [-1500000n(\hat{\delta}r)^2, 0],$ 

where  $B(r/4) = B(0, r/4)$ . Since

$$
x + x_{*} + 30\hat{\delta}r\theta \in B(x_{*}, r/2) \quad \text{for } x \in \overline{B(r/4)},
$$
  

$$
t + \tilde{t} \in (t_{*} - (r/2)^{2}, t_{*}) \qquad \text{for } t \in [-1500000n(\hat{\delta}r)^{2}, 0],
$$

we can see from [\(5.7\)](#page-39-2) that

$$
|\partial_t u_1 - \Delta u_1| \le \frac{C_1}{M^{2(p-1)}r^2}|u_1| \quad \text{in } \overline{B(r/4)} \times [-1500000n(\hat{\delta}r)^2, 0]
$$

for some  $C_1 = C_1(n,p) > 0$ . Notice that  $(r/4)^2 \ge 16000 \times 1500000n(\hat{\delta}r)^2$ . By Proposition [22](#page-30-2) with  $T_1 = 1500000n(\hat{\delta}r)^2$ ,  $C_0 = M^{2(p-1)}/(1500000C_1n\hat{\delta}^2) \ge 1$  and

$$
\underline{s} = \left(\frac{\hat{\delta}^4}{M_3^4}\right)(\hat{\delta}r)^2 \le \overline{s} = 75(\hat{\delta}r)^2 < \frac{1500000n(\hat{\delta}r)^2}{10000n},
$$

we have  $Z_1 \leq X_1 + Y_1$ , where

$$
X_1 := C(n)e^{-\frac{(r/4)^2}{500 \times 75(\tilde{\delta}r)^2}} \int_{-1500000n(\hat{\delta}r)^2}^{0} \int_{B(r/4)} \left( \frac{|u_1|^2}{1500000n(\hat{\delta}r)^2} + |\nabla u_1|^2 \right) dx dt,
$$
  
\n
$$
Y_1 := C(n) \left( 75(\hat{\delta}r)^2 \right)^{\frac{n}{2}} \left( \frac{3e \times 75(\hat{\delta}r)^2}{(\hat{\delta}^4/M_3^4)(\hat{\delta}r)^2} \right)^{\frac{(r/4)^2}{200 \times 75(\hat{\delta}r)^2}}
$$
  
\n
$$
\times \int_{B(r/4)} |u_1(x,0)|^2 \left( \frac{\hat{\delta}^4}{M_3^4} (\hat{\delta}r)^2 \right)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4(\hat{\delta}^4/M_3^4)(\hat{\delta}r)^2}} dx,
$$
  
\n
$$
Z_1 := \int_{-2 \times 75(\hat{\delta}r)^2}^{-75(\hat{\delta}r)^2} \int_{B(r/8)} \left( \frac{|u_1|^2}{1500000n(\hat{\delta}r)^2} + |\nabla u_1|^2 \right) e^{-\frac{|x|^2}{4t}} dx dt.
$$

From the change of variables and [\(5.8\)](#page-39-3), it follows that

$$
Z_{1} \geq \frac{1}{C(n)(\hat{\delta}r)^{2}} \int_{-150(\hat{\delta}r)^{2}}^{-75(\hat{\delta}r)^{2}} \int_{B(r/8)} |u(x+x_{*}+30\hat{\delta}r\theta, t+\tilde{t})|^{2} e^{-\frac{|x|^{2}}{4t}} dxdt
$$
  
\n
$$
\geq \frac{1}{C(n)(\hat{\delta}r)^{2}} e^{-\frac{31^{2}(\hat{\delta}r)^{2}}{4(-150(\hat{\delta}r)^{2}}}\int_{-150(\hat{\delta}r)^{2}}^{-75(\hat{\delta}r)^{2}} \int_{B(31\hat{\delta}r)} |u(x+x_{*}+30\hat{\delta}r\theta, t+\tilde{t})|^{2} dxdt
$$
  
\n
$$
= \frac{e^{\frac{31^{2}}{600}}}{C(n)(\hat{\delta}r)^{2}} \int_{\tilde{t}-150(\hat{\delta}r)^{2}}^{t-75(\hat{\delta}r)^{2}} \int_{B(x_{*}+30\hat{\delta}r\theta, 31\hat{\delta}r)} |u(y,s)|^{2} dyds
$$
  
\n
$$
\geq \frac{1}{C(n)(\hat{\delta}r)^{2}} \int_{t_{*}-(r^{2}/8)}^{t_{*}-(r^{2}/8)} \int_{B(x_{*},\hat{\delta}r)} |u(y,s)|^{2} dyds
$$
  
\n
$$
\geq \frac{1}{C(n)(\hat{\delta}r)^{2}} M_{2}^{-\overline{C}(n,p)}(\hat{\delta}r)^{n+2-\frac{4}{p-1}}.
$$

Then by  $\hat{\delta} = M_2^{-1}$ , we obtain

<span id="page-40-0"></span>(5.10) 
$$
Z_1 \ge \frac{1}{C(n)} M_2^{\frac{4}{p-1} - n - \overline{C}(n, p)} r^{n - \frac{4}{p-1}} \ge e^{-M_2} r^{n - \frac{4}{p-1}}.
$$

By [\(5.7\)](#page-39-2) and  $M^4 \geq \hat{\delta}$ , we see that

$$
X_1 \leq Ce^{-\frac{1}{600000\delta^2}} \int_{-1500000n(\hat{\delta}r)^2}^{0} \int_{B(r/4)} \left( \frac{r^{-\frac{4}{p-1}} M^{-4}}{1500000n(\hat{\delta}r)^2} + r^{-\frac{2(p+1)}{p-1}} M^{-4} \right) dx dt
$$
  

$$
\leq \frac{C}{M^4} \left( 1 + \hat{\delta}^2 \right) r^{n - \frac{4}{p-1}} e^{-\frac{M_2^2}{600000}} \leq e^{-\frac{M_2^2}{600000}} r^{n - \frac{4}{p-1}}.
$$

Then,  $X_1$  can be absorbed in the right-hand side of  $(5.10)$ , and so

<span id="page-40-1"></span>(5.11) 
$$
Y_1 \ge e^{-2M_2} r^{n - \frac{4}{p-1}}.
$$

On the other hand, by splitting

$$
Y_1 \leq C(n) \left(\frac{M_3^4}{\hat{\delta}^4}\right)^{\frac{n}{2} + \frac{1}{240000\hat{\delta}^2}} \int_{B(\hat{\delta}r/M_3)} |u_1(x,0)|^2 e^{-\frac{M_3^4 |x|^2}{4\hat{\delta}^4(\hat{\delta}r)^2}} dx
$$
  
+ 
$$
C(n) \left(\frac{M_3^4}{\hat{\delta}^4}\right)^{\frac{n}{2} + \frac{1}{240000\hat{\delta}^2}} \int_{B(r/4) \backslash B(\hat{\delta}r/M_3)} |u_1(x,0)|^2 e^{-\frac{M_3^4 |x|^2}{4\hat{\delta}^4(\hat{\delta}r)^2}} dx =: Y_1' + Y_1''
$$

and by using  $(5.7)$ ,  $(n/2) + (M_2^2/240000) \le M_2^2/5$  and  $M_3^{M_2^2} \le e^{M_3^2}$ , we see that

$$
Y_{1}^{\prime\prime} \leq C(n)M^{-4} \left(\frac{225eM_{3}^{4}}{\hat{\delta}^{4}}\right)^{\frac{n}{2}+\frac{1}{240000\hat{\delta}^{2}}} r^{-\frac{4}{p-1}} \int_{B(r/4)\backslash B(\hat{\delta}r/M_{3})} e^{-\frac{M_{3}^{4}|x|^{2}}{4\hat{\delta}^{4}(\hat{\delta}r)^{2}}} dx
$$
  

$$
\leq \frac{C(n)}{M^{4}} \left(225eM_{2}^{4}M_{3}^{4}\right)^{\frac{n}{2}+\frac{M_{2}^{2}}{240000}} r^{n-\frac{4}{p-1}} e^{-\frac{M_{3}^{2}}{4\hat{\delta}^{4}}}
$$
  

$$
\leq (M_{3}^{5})^{\frac{M_{2}^{2}}{5}} e^{-\frac{M_{2}^{4}M_{3}^{2}}{8}} e^{-\frac{M_{3}^{4}M_{3}^{2}}{8}} r^{n-\frac{4}{p-1}} \leq \frac{M_{3}^{M_{2}^{2}}}{e^{M_{3}^{2}}} e^{-M_{3}^{2}} r^{n-\frac{4}{p-1}} \leq e^{-M_{3}^{2}} r^{n-\frac{4}{p-1}}.
$$

Thus,  $Y_1''$  can be absorbed in the right-hand side of  $(5.11)$ . Then we have

$$
e^{-3M_2}r^{n-\frac{4}{p-1}} \le Y'_1 \le C(M_2^4M_3^4)^{\frac{n}{2}+\frac{1}{240000\delta^2}} \int_{B(\hat{\delta}r/M_3)} |u_1(x,0)|^2 dx
$$
  

$$
\le M_3^{M_2^2} \int_{B(x_*+30\hat{\delta}r\theta,\hat{\delta}r/M_3)} |u(y,\tilde{t})|^2 dy.
$$

Since  $e^{-3M_2} M_3^{-M_2^2} \ge e^{-M_3}$ , we obtain

<span id="page-41-0"></span>(5.12) 
$$
\int_{B(x_*+30\hat{\delta}r\theta,\hat{\delta}r/M_3)} |u(y,\tilde{t})|^2 dy \geq e^{-M_3}r^{n-\frac{4}{p-1}} \text{ for any } \tilde{t} \in I.
$$

Step 4: iterated unique continuation on thinner slice

Let us apply Proposition [23](#page-32-1) to

$$
u_2(x,t) := u(x + x_*, t + \tau), \quad (x,t) \in \tilde{S}.
$$

Here we set  $\tau := \sup I \leq 0$  and

$$
\tilde{S} := \tilde{S}_0 \times [-M_2^{-3}(\hat{\delta}r)^2, 0] \subset \mathbf{R}^n \times [-1, 0],
$$
  

$$
\tilde{S}_0 := \{ x \in \mathbf{R}^n; \text{ dist}(x, \mathbf{R}_+ \theta) \le M_2^{-3} |x \cdot \theta|, |x| \ge 30 \hat{\delta} r \}.
$$

For  $x \in \tilde{S}_0$  and  $t \in [-M_2^{-3}(\hat{\delta}r)^2, 0]$ , we can check that  $x + x_* \in S_*$  and  $t + \tau \in I$ . Particularly,  $[\tau - M_2^{-3}(\hat{\delta}r)^2, \tau] \subset I$ . Indeed,

dist
$$
(x + x_*, x_* + \mathbf{R}_+ \theta)
$$
 = dist $(x, \mathbf{R}_+ \theta) \le M_2^{-3} |x \cdot \theta| \le 10 M^{-\gamma} |(x + x_* - x_*) \cdot \theta|$ ,  
\n $|x + x_* - x_*| = |x| \ge 30 \hat{\delta} r \ge 20 \hat{\delta} r$ ,  $t + \tau \le \tau = \sup I$ ,  
\n $t + \tau \ge \sup I - M_2^{-3} (\hat{\delta} r)^2 \ge \sup I - (\hat{\delta} r / M^{\gamma})^2 = \sup I - |I| = \inf I$ .

Then by [\(5.9\)](#page-39-4), we observe that

$$
||u_2||_{L^{\infty}(\tilde{S})} \le ||u||_{L^{\infty}(S)} \le M^{\frac{2\gamma}{p-1}-1}(\hat{\delta}r)^{-\frac{2}{p-1}} \le \left(M^{-3} \times (\hat{\delta}r)^2\right)^{-\frac{1}{p-1}},
$$
  

$$
||\nabla u_2||_{L^{\infty}(\tilde{S})} \le ||\nabla u||_{L^{\infty}(S)} \le M^{\frac{2\gamma}{p-1}-1}(\hat{\delta}r)^{-\frac{2}{p-1}-1} \le \left(M^{-3} \times (\hat{\delta}r)^2\right)^{-\frac{1}{p-1}-\frac{1}{2}},
$$

and that the condition on the first line in [\(4.16\)](#page-33-0) is satisfied with  $T_1 = (\hat{\delta}r)^2$  and  $\eta = M_2^{-3}$ . Moreover, since

$$
|\partial_t u_2 - \Delta u_2| \le |u_2|^{p-1} |u_2| \le \frac{|u_2|}{M^{-(2\gamma - (p-1))} (\hat{\delta}r)^2} \le \frac{|u_2|}{M_2^{-3} \times (\hat{\delta}r)^2}
$$

in  $\tilde{S}$ , the condition on the second line in [\(4.16\)](#page-33-0) is also satisfied with  $C_0 = 1$ . Notice that  $0 < \eta \le \min(C_0^{-1}, 1) = 1$ . From  $(5.12)$  and  $[\tau - M_2^{-3}(\hat{\delta}r)^2, \tau] \subset I$ , it follows that

$$
\int_{B(30\hat{\delta}r\theta, M_3^{-1}\hat{\delta}r)} |u(x+x_*, t+\tau)|^2 dx \geq e^{-M_3}r^{n-\frac{4}{p-1}}
$$

for  $t \in [-M_2^{-3}(\hat{\delta}r)^2, 0]$ . In particular,

$$
\int_{B(30\hat{\delta}r\theta,30M_2^{-15}\hat{\delta}r)} |u_2(x,t)|^2 dx \geq e^{-M_3}r^{n-\frac{4}{p-1}} = \hat{\delta}^{\frac{4}{p-1}-n}e^{-M_3}(\hat{\delta}r)^{n-\frac{4}{p-1}}
$$

with  $20(\hat{\delta}r) \leq 30\hat{\delta}r \leq M_2^6(\hat{\delta}r)$  and  $0 < \hat{\delta}^{(4/(p-1))-n}e^{-M_3} \leq M_2^{-24}$ . Then, Proposi-tion [23](#page-32-1) with  $R_0 = 30\hat{\delta}r$ ,  $\epsilon = \hat{\delta}^{(4/(p-1)) - n}e^{-M_3}$  and  $4/(p-1) < n$  yields

$$
\int_{B(\tilde{R}_0 \theta, M_2^{-6}\tilde{R}_0)} |u_2(x,t)|^2 dx \ge \left(\hat{\delta}^{\frac{4}{p-1}-n} e^{-M_3}\right)^{\left(\frac{\tilde{R}_0}{30\tilde{\delta}r}\right)^{M_2^{12}}} (\hat{\delta}^r)^{n-\frac{4}{p-1}}
$$
  

$$
\ge e^{-M_3 \left(\frac{\tilde{R}_0}{30\tilde{\delta}r}\right)^{M_2^{12}}} (\hat{\delta}^r)^{n-\frac{4}{p-1}}
$$

for all  $t \in [-M_2^{-3}(\hat{\delta}r)^2/2, 0]$  and  $\tilde{R}_0 \geq 30\hat{\delta}r$ . Since  $r = 8M_2^{-(6n+2)}(-t'')^{1/2}$  and  $\hat{\delta} = M_2^{-1}$ , we have

<span id="page-42-0"></span>
$$
\int_{B(\tilde{R}_0 \theta, M_2^{-6} \tilde{R}_0)} |u(x + x_*, t + \tau)|^2 dx
$$
\n
$$
\geq e^{-M_3 \left(\frac{M_2^{6n+3} \tilde{R}_0}{240(-t'')^{1/2}}\right)^{M_2^{12}}} (8M_2^{-(6n+3)})^{n - \frac{4}{p-1}} (-t'')^{\frac{n}{2} - \frac{2}{p-1}}
$$
\n
$$
\geq e^{-2M_3 \left(\frac{M_2^{6n+3} \tilde{R}_0}{240(-t'')^{1/2}}\right)^{M_2^{12}}} (-t'')^{\frac{n}{2} - \frac{2}{p-1}}
$$

for all  $t \in [32M_2^{-(12n+9)}t'$ , 0] and  $\tilde{R}_0 \ge 240M_2^{-(6n+3)}(-t'')^{1/2}$ . Step 5: preparation of annulus of regularity

<span id="page-42-2"></span>By Proposition [18](#page-24-6) with  $\lambda = M_4$ ,  $R = M_4$  and  $T_1 = M^3(-t'')$ , there exists  $Mq_c+1+\beta$ 

$$
(5.14) \t\t M_4 \le R_1 \le M_4^{M_4^{M_3}}.
$$

such that

<span id="page-42-1"></span>
$$
(5.15) \t\t ||u||_{L^{\infty}(\mathcal{A})} \leq T_1^{-\frac{1}{p-1}} M^{-1}, \t ||\nabla u||_{L^{\infty}(\mathcal{A})} \leq T_1^{-\frac{p+1}{2(p-1)}} M^{-1},
$$
  

$$
\mathcal{A} := \left\{ (x, t); \ 2T_1^{\frac{1}{2}} R_1 < |x| < \frac{1}{2} T_1^{\frac{1}{2}} R_1^{M_4}, \ t \in \left[ -\frac{1}{2} T_1, 0 \right] \right\}.
$$

Based on [\[31\]](#page-55-0) for the higher-dimensional Navier-Stokes equations, we apply quantitative backward uniqueness in a sub-annulus of  $A$  and use the lower bound [\(5.13\)](#page-42-0). However, in [\[31\]](#page-55-0), the annulus used for backward uniqueness is similar to the annulus of regularity, which seems to create a gap in the arguments involving subsequent uses of unique continuation remaining inside the annulus of regularity. As in [\[38\]](#page-56-0) and subsequently [\[2\]](#page-54-5), we instead apply backward uniqueness Carleman inequalities in a much smaller annulus than the annulus of regularity.

Let  $T_2 := M^{-2}T_1 = M(-t'')$ . We define a smaller annulus  $\tilde{A}$  by

$$
\tilde{\mathcal{A}} := \left\{ (x, t); \ M_3 T_2^{\frac{1}{2}} R_1 < |x| < M_3^{-1} T_2^{\frac{1}{2}} R_1^{M_4}, \ t \in [-T_2, 0] \right\}
$$
\n
$$
\subset \left\{ (x, t); \ 2MT_2^{\frac{1}{2}} R_1 < |x| < \frac{1}{2} MT_2^{\frac{1}{2}} R_1^{M_4}, \ t \in \left[ -\frac{1}{2} M^2 T_2, 0 \right] \right\} = \mathcal{A}.
$$

Note that the estimates in [\(5.15\)](#page-42-1) also hold in  $\tilde{\mathcal{A}}$ . We substitute  $20M_3T_2^{1/2}R_1(=$  $20M^{1/2}M_3R_1(-t'')^{1/2}$  into  $\tilde{R}_0$  in [\(5.13\)](#page-42-0). Then the change of variables yields

$$
Z_2 := \int_{B(x_*+20M_3T_2^{1/2}R_1\theta, 20M_2^{-6}M_3T_2^{1/2}R_1)} |u(y, s)|^2 dy
$$
  
\n
$$
\geq e^{-2M_3 \left(\frac{M_2^{6n+3}20M^{1/2}M_3R_1(-t'')^{1/2}}{240(-t'')^{1/2}}\right)^{M_2^{12}} (-t'')^{\frac{n}{2} - \frac{2}{p-1}}
$$
  
\n
$$
\geq e^{-2M_3 \left(\frac{M_2^{6n+3}M^{1/2}M_3}{12}R_1\right)^{M_2^{12}} (-t'')^{\frac{n}{2} - \frac{2}{p-1}}
$$

for all  $t \in [\tau + 32M_2^{-(12n+9)}t'', \tau] = [\tau - 32M^{-1}M_2^{-(12n+9)}T_2, \tau]$ . Since  $R_1 \ge M_4$ , we see that

$$
Z_2 \ge e^{-R_1^{4M_2^{12}}} (-t'')^{\frac{n}{2} - \frac{2}{p-1}} \ge e^{-R_1^{4M_2^{12}}} (M^{-1}T_2)^{\frac{n}{2} - \frac{2}{p-1}} \ge e^{-R_1^{M_3}} T_2^{\frac{n}{2} - \frac{2}{p-1}}.
$$

Hence by integrating  $Z_2$  over  $s \in [\tau - 32M^{-1}M_2^{-(12n+9)}T_2, \tau]$ , we obtain

<span id="page-43-0"></span>
$$
(5.16) \qquad \int_{\tau-\frac{32T_2}{MM_2^{12n+9}}}^{\tau} \int_{B(x_*+20M_3T_2^{\frac{1}{2}}R_1\theta, 20M_3M_2^{-6}T_2^{\frac{1}{2}}R_1)} |u(x,t)|^2 dxdt
$$
  

$$
\geq \frac{32}{MM_2^{12n+9}} e^{-R_1^{M_3}} T_2^{\frac{n}{2}-\frac{2}{p-1}+1} \geq e^{-2R_1^{M_3}} T_2^{\frac{n}{2}-\frac{2}{p-1}+1}.
$$

## Step 6: backward uniqueness Carleman inequality—two cases

Let us apply Proposition [21](#page-28-6) to u. By [\(5.15\)](#page-42-1),  $M > 1$  and  $T_2 = M^{-2}T_1$ , we have

$$
|\partial_t u - \Delta u| \le \frac{|u|}{T_1 M^{p-1}} \le \frac{|u|}{T_1} = \frac{|u|}{M^2 \times T_2} \quad \text{in } \tilde{\mathcal{A}}.
$$

We apply Proposition [21](#page-28-6) on  $\tilde{A}$  with  $r_- = M_3 T_2^{1/2} R_1$ ,  $r_+ = M_3^{-1} T_2^{1/2} R_1^{M_4}$  and  $C_{\text{Carl}} = M^2$ . Here we note that  $C_{\text{Carl}} \geq 3n$  and  $r^2 \geq 4C_{\text{Carl}}T_2$  hold by taking M so large that  $M^2 \geq 3n$  and by using  $R_1 \geq M_4$ , respectively. Then, we have  $Z_3 \leq X_3 + Y_3$ , where

$$
X_3 := C(n)M^2 e^{-\frac{R_1^{1+M_4}}{4M^2}} \iint_{\tilde{\mathcal{A}}} e^{\frac{2|x|^2}{T_1}} \left( \frac{|u|^2}{T_2} + |\nabla u|^2 \right) dxdt,
$$
  
\n
$$
Y_3 := C(n)M^2 e^{-\frac{R_1^{1+M_4}}{4M^2}} e^{\frac{2R_1^{2M_4}}{M^2 M_3^2}} \int_{M_3 T_2^{1/2} R_1 \le |x| \le \frac{T_2^{1/2} R_1^{M_4}}{M_3}} |u(x,0)|^2 dx,
$$
  
\n
$$
Z_3 := \int_{-\frac{T_2}{4}}^0 \int_{10M_3 T_2^{1/2} R_1 \le |x| \le \frac{T_2^{1/2}}{2M_3} R_1^{M_4}} \left( \frac{|u|^2}{T_2} + |\nabla u|^2 \right) dxdt.
$$

We estimate  $Z_3$  by using [\(5.16\)](#page-43-0). Recall that  $x_* \in B(0, 20M_1(-t'')^{1/2})$  in the first part of this section, and so  $|x_*| \leq 20M^{-1/2}M_1T_2^{1/2}$  since  $T_2 = M(-t'')$ . For

$$
x \in B(x_{*} + 20M_{3}T_{2}^{1/2}R_{1}\theta, 20M_{2}^{-6}M_{3}T_{2}^{1/2}R_{1}), \text{ we observe that}
$$
\n
$$
|x| \leq |x - x_{*} - 20M_{3}T_{2}^{\frac{1}{2}}R_{1}\theta| + |x_{*}| + 20M_{3}T_{2}^{\frac{1}{2}}R_{1} \leq 22M_{3}T_{2}^{\frac{1}{2}}R_{1} \leq \frac{T_{2}^{\frac{1}{2}}}{2M_{3}}R_{1}^{M_{4}},
$$
\n
$$
|x| \geq |20M_{3}T_{2}^{\frac{1}{2}}R_{1}\theta + x_{*}| - |x - x_{*} - 20M_{3}T_{2}^{\frac{1}{2}}R_{1}\theta|
$$
\n
$$
\geq 20M_{3}T_{2}^{\frac{1}{2}}R_{1} - 20M^{-\frac{1}{2}}M_{1}T_{2}^{\frac{1}{2}} - 20M_{2}^{-6}M_{3}T_{2}^{\frac{1}{2}}R_{1} \geq 10M_{3}T_{2}^{\frac{1}{2}}R_{1}.
$$
\nThus

Thus,

$$
B(x_*+20M_3T_2^{\frac{1}{2}}R_1\theta,20M_2^{-6}M_3T_2^{\frac{1}{2}}R_1) \subset \left\{ 10M_3T_2^{\frac{1}{2}}R_1 \leq |x| \leq \frac{T_2^{\frac{1}{2}}}{2M_3}R_1^{M_4} \right\}.
$$

Recall Step 2 that

$$
r = 8M_2^{-(6n+2)}(-t'')^{\frac{1}{2}} = 8M^{-1/2}M_2^{-(6n+2)}T_2^{\frac{1}{2}}, \quad \hat{\delta} = M_2^{-1},
$$
  

$$
\tau = \sup I \ge t_* - (r^2/8) + 99(\hat{\delta}r)^2.
$$

Moreover, the inclusions in [\(5.6\)](#page-39-1) particularly show that  $t_* - (r^2/8) - (\hat{\delta}r)^2 \geq 2t''$ . Then we see that

$$
0 \ge \tau \ge \tau - \frac{32T_2}{MM_2^{12n+9}} \ge 2t'' + 100(\hat{\delta}r)^2 - \frac{32T_2}{MM_2^{12n+9}}
$$
  
=  $2t'' + \frac{6400T_2}{MM_2^{12n+6}} - \frac{32T_2}{MM_2^{12n+9}} \ge 2t'' = -\frac{2T_2}{M} \ge -\frac{T_2}{4}.$ 

Then,  $[\tau - 32M^{-1}M_2^{-(12n+9)}T_2, \tau] \subset [-T_2/4, 0]$ . By [\(5.16\)](#page-43-0), we see that

$$
Z_3 \ge \int_{\tau-\frac{32T_2}{MM_2^{12n+9}}}^{\tau} \int_{B(x_*+20M_3T_2^{\frac{1}{2}}R_1\theta,20\frac{M_3}{M_2^6}T_2^{\frac{1}{2}}R_1)} \frac{|u|^2}{T_2} dx dt
$$
  

$$
\ge \frac{1}{T_2} e^{-2R_1^{M_3}} T_2^{\frac{n}{2}-\frac{2}{p-1}+1} \ge e^{-2R_1^{M_3}} T_2^{\frac{n}{2}-\frac{2}{p-1}}.
$$

Hence, at least, one of the following holds:

<span id="page-44-1"></span>(5.17) 
$$
Y_3 \ge e^{-3R_1^{M_3}} T_2^{\frac{n}{2} - \frac{2}{p-1}},
$$

<span id="page-44-2"></span>(5.18) 
$$
X_3 \ge e^{-3R_1^{M_3}} T_2^{\frac{n}{2} - \frac{2}{p-1}}.
$$

Eventually, in both cases, we can derive the same estimate as can be seen in [\(5.19\)](#page-44-0) and [\(5.27\)](#page-49-0) below. However, the second case requires more involved arguments than the first case.

# Step 7: estimate for first case

If [\(5.17\)](#page-44-1) is satisfied, we see that

$$
\begin{aligned} e^{-3R_1^{M_3}}T_2^{\frac{n}{2}-\frac{2}{p-1}} &\le Y_3 \le C(n)M^2 e^{\frac{2R_1^{2M_4}}{M^2M_3^2}}\int_{M_3T_2^{1/2}R_1 \le |x| \le \frac{T_2^{1/2}R_1^{M_4}}{M_3}}|u(x,0)|^2 dx \\ &\le e^{R_1^{3M_4}}\int_{M_3T_2^{1/2}R_1 \le |x| \le \frac{T_2^{1/2}R_1^{M_4}}{M_3}}|u(x,0)|^2 dx. \end{aligned}
$$

Hence by expanding the integral domain, we obtain

<span id="page-44-0"></span>(5.19) 
$$
\int_{T_2^{\frac{1}{2}} R_1 \leq |x| \leq T_2^{\frac{1}{2}} R_1^{M_4}} |u(x,0)|^2 dx \geq e^{-R_1^{4M_4}} T_2^{\frac{n}{2} - \frac{2}{p-1}}.
$$

Step 8: estimate for second case by triple pigeonhole—first pigeonhole in space variable

In steps  $8-11$ , we show that  $(5.18)$  also implies  $(5.19)$  by utilizing arguments in [\[38\]](#page-56-0) involving the pigeonhole principle and a further use of quantitative unique continuation.

We examine the case [\(5.18\)](#page-44-2) and deduce the same inequality as [\(5.19\)](#page-44-0). By  $T_2 =$  $M^{-2}T_1$  and  $R_1 \geq M_4$ , we have

$$
\begin{split} &X_3\leq C(n)M^2e^{-\frac{R_1^{1+M_4}}{4M^2}}\int_{-T_2}^0\int_{M_3T_2^{\frac{1}{2}}R_1\leq |x|\leq \frac{T_2^{\frac{1}{2}}}{M_3}R_1^{M_4}}e^{\frac{2|x|^2}{T_1}}\left(\frac{|u|^2}{T_2}+|\nabla u|^2\right)dxdt\\ &\leq e^{-\frac{R_1^{1+M_4}}{5M^2}}\int_{-\frac{T_1}{M^2}}^0\int_{\frac{M_3}{M}T_1^{\frac{1}{2}}R_1\leq |x|\leq \frac{T_1^{\frac{1}{2}}}{M M_3}R_1^{M_4}}e^{\frac{2|x|^2}{T_1}}\left(\frac{M^2|u|^2}{T_1}+|\nabla u|^2\right)dxdt\\ &\leq e^{-\frac{R_1^{1+M_4}}{6M^2}}\int_{-\frac{T_1}{M^2}}^0\int_{M_3^{\frac{1}{2}}T_1^{\frac{1}{2}}R_1\leq |x|\leq M_3^{-\frac{1}{2}}T_1^{\frac{1}{2}}R_1^{M_4}}e^{\frac{2|x|^2}{T_1}}\left(\frac{|u|^2}{T_1}+|\nabla u|^2\right)dxdt.\end{split}
$$

This together with [\(5.18\)](#page-44-2) gives

$$
X'_3 := \int_{-\frac{T_1}{M^2}}^0 \int_{M_3^{\frac{1}{2}} T_1^{\frac{1}{2}} R_1 \le |x| \le M_3^{-\frac{1}{2}} T_1^{\frac{1}{2}} R_1^{M_4}} e^{\frac{2|x|^2}{T_1}} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt
$$
  
\n
$$
\ge e^{\frac{R_1^{1+M_4}}{6M^2}} e^{-3R_1^{M_3}} M^{\frac{4}{p-1} - n} T_1^{\frac{n}{2} - \frac{2}{p-1}} \ge e^{\frac{R_1^{\frac{1}{2} + M_4}}{6M^2}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

We claim that there exists  $M_3^{1/2}T_1^{1/2}R_1 \leq R_3 \leq M_3^{-1/2}T_1^{1/2}R_1^{M_4}$  satisfying

$$
\int_{-\frac{T_1}{M^2}}^0 \int_{R_3 \leq |x| \leq 2R_3} e^{\frac{2|x|^2}{T_1}} \left(\frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt \geq e^{\frac{R_1^{M_4}}{6M^2}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

To obtain a contradiction, we suppose that

<span id="page-45-0"></span>
$$
(5.20) \qquad \qquad \int_{-\frac{T_1}{M^2}}^0 \int_{R_3 \le |x| \le 2R_3} e^{\frac{2|x|^2}{T_1}} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt < e^{\frac{R_1^{M_4}}{6M^2}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

for any  $M_3^{1/2}T_1^{1/2}R_1 \leq R_3 \leq M_3^{-1/2}T_1^{1/2}R_1^{M_4}$ . We take the positive integer k' satisfying  $2^{k'} \le R_1^{M_4-1}/M_3 \le 2^{k'+1}$ . Note that

$$
M_3^{-\frac{1}{2}}T_1^{\frac{1}{2}}R_1^{M_4} \le 2^{k'+1}M_3^{\frac{1}{2}}T_1^{\frac{1}{2}}R_1 \le 2M_3^{-\frac{1}{2}}T_1^{\frac{1}{2}}R_1^{M_4}.
$$

Then from [\(5.20\)](#page-45-0), it follows that

$$
X_3' \leq \sum_{k=0}^{k'} \int_{-\frac{T_1}{M^2}}^0 \int_{2^k M_3^{\frac{1}{2}} T_1^{\frac{1}{2}} R_1 \leq |x| \leq 2^{k+1} M_3^{\frac{1}{2}} T_1^{\frac{1}{2}} R_1} e^{\frac{2|x|^2}{T_1}} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt
$$
  
<  $(k' + 1) e^{\frac{R_1^{M_4}}{6M^2}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.$ 

Since  $k' \leq (\log(R_1^{M_4-1}/M_3)/\log 2) \leq 2M_4 \log R_1 \leq R_1^2$  by  $R_1 \geq M_4$ , we obtain

$$
X'_3< (R_1^2+1) e^{\frac{R_1^{M_4}}{6M^2}}T_1^{\frac{n}{2}-\frac{2}{p-1}}\leq e^{\frac{R_1^{\frac{1}{2}+M_4}}{6M^2}}T_1^{\frac{n}{2}-\frac{2}{p-1}}\leq X'_3,
$$

a contradiction. Hence the claim follows. In particular, by  $e^{2|x|^2/T_1} \leq e^{8R_3^2/T_1}$  for  $|x| \leq 2R_3$  and by  $e^{R_1^{M_4}/(6M^2)} \geq 1$ , we see that

<span id="page-46-0"></span>(5.21) 
$$
\tilde{X}_3 := \int_{-\frac{T_1}{M^2}}^0 \int_{R_3 \le |x| \le 2R_3} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt
$$

$$
\ge e^{\frac{R_1^{M_4}}{6M^2} - \frac{8R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \ge e^{-\frac{8R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

Step 9: estimate for second case by triple pigeonhole—second pigeonhole in time variable

We decompose

$$
\tilde{X}_3 = \int_{-\frac{T_1}{M^2}}^{-T_1 e^{-\frac{10R_3^2}{T_1}}} \int_{R_3 \leq |x| \leq 2R_3} (\cdots) + \int_{-T_1 e^{-\frac{10R_3^2}{T_1}}}^{0} \int_{R_3 \leq |x| \leq 2R_3} (\cdots) =: \tilde{X}'_3 + \tilde{X}''_3.
$$

Note that

$$
\{R_3 \le |x| \le 2R_3\} \subset \left\{2T_1^{\frac{1}{2}}R_1 < |x| < \frac{1}{2}T_1^{\frac{1}{2}}R_1^{M_4}\right\}.
$$

Thus, we can apply [\(5.15\)](#page-42-1) to see that

$$
\tilde X''_3 \leq \frac{C(n)}{M^2} R_3^n T_1^{1-\frac{p+1}{p-1}} e^{-\frac{10R_3^2}{T_1}} = \frac{C(n)}{M^2} \left(\frac{R_3^2}{T_1}\right)^{\frac{n}{2}} e^{-\frac{10R_3^2}{T_1}} T_1^{\frac{n}{2}-\frac{2}{p-1}} \leq e^{-\frac{9R_3^2}{T_1}} T_1^{\frac{n}{2}-\frac{2}{p-1}}.
$$

Hence  $\tilde{X}_3''$  can be absorbed by the right-hand side of [\(5.21\)](#page-46-0), and then

$$
\tilde{X}'_3 = \int_{-\frac{T_1}{M^2}}^{-T_1 e^{-\frac{10R_3^2}{T_1}}} \int_{R_3 \leq |x| \leq 2R_3} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt \geq e^{-\frac{9R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

We claim that there exists  $T_1e^{-10R_3^2/T_1} \le -t_3 \le T_1/M^2$  satisfying

$$
(5.22) \t\t \hat{X}_3 := \int_{t_3}^{\frac{t_3}{2}} \int_{R_3 \leq |x| \leq 2R_3} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt \geq e^{-\frac{10R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

To obtain a contradiction by the same argument as in Step 8, we suppose that

<span id="page-46-1"></span>
$$
\int_{t_3}^{\frac{t_3}{2}} \int_{R_3 \leq |x| \leq 2R_3} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt < e^{-\frac{10R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

for any  $T_1e^{-10R_3^2/T_1} \leq -t_3 \leq T_1/M^2$ . We take the positive integer  $\tilde{k}'$  satisfying

$$
2^{-\tilde{k}'-1} \le M^2 e^{-\frac{10R_3^2}{T_1}} \le 2^{-\tilde{k}'}
$$

.

Then, we can observe that

$$
-T_1e^{-\frac{10R_3^2}{T_1}} \le -2^{-\tilde{k}'-1}M^{-2}T_1 \le -2^{-1}T_1e^{-\frac{10R_3^2}{T_1}}.
$$

In addition,  $\tilde{k}' \log 2 \leq 10R_3^2/T_1 - \log M^2$ . Thus,

$$
\tilde{X}'_3 \le \sum_{k=0}^{\tilde{k}'} \int_{-2^{-k} \frac{T_1}{M^2}}^{-2^{-k-1} \frac{T_1}{M^2}} \int_{R_3 \le |x| \le 2R_3} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt < (\tilde{k}' + 1) e^{-\frac{10R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \n\le \left( \frac{10}{\log 2} \left( \frac{R_3^2}{T_1} \right) + 1 \right) e^{-\frac{10R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \le e^{-\frac{9R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \le \tilde{X}'_3,
$$

a contradiction. The claim follows.

# Step 10: estimate for second case by triple pigeonhole—third pigeonhole by covering

We take a covering  $\{B(\tilde{x}_i, (-t_3)^{1/2})\}_{i=1,\dots,N_{\text{cov}}}$  of  $\{R_3 \leq |x| \leq 2R_3\}$  such that  $R_3 \leq |\tilde{x}_i| \leq 2R_3$ , where  $N_{\text{cov}} \leq C_{\text{cov}} (R_3/(-t_3)^{1/2})^n$  with some constant  $C_{\text{cov}} > 0$ depending only on *n*. We claim that there exists  $x_3 := \tilde{x}_{i'}$   $(1 \leq i' \leq N_{\text{cov}})$  satisfying

$$
(5.23) \qquad \int_{t_3}^{\frac{t_3}{2}} \int_{B(x_3, (-t_3)^{\frac{1}{2}})} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt \geq e^{-\frac{10nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

To obtain a contradiction, suppose that

<span id="page-47-0"></span>
$$
\int_{t_3}^{\frac{t_3}{2}} \int_{B(\tilde{x}_i, (-t_3)^{\frac{1}{2}})} \left( \frac{|u|^2}{T_1} + |\nabla u|^2 \right) dx dt < e^{-\frac{10nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

for any  $1 \le i \le N_{\text{cov}}$ . By  $-t_3 \ge T_1 e^{-10R_3^2/T_1}$ ,  $n \ge 3$  and  $(5.22)$ ,

$$
\hat{X}_3 \le \sum_{i=1}^{N_{\text{cov}}} \int_{t_3}^{\frac{t_3}{2}} \int_{B(\tilde{x}_i, (-t_3)^{\frac{1}{2}})} \left( \frac{|u|^2}{r^2} + |\nabla u|^2 \right) dx dt < N_{\text{cov}} e^{-\frac{10nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \n\le C_{\text{cov}} \left( \frac{R_3^2}{T_1} \right)^{\frac{n}{2}} e^{-\frac{5nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \le e^{-\frac{4nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \le e^{-\frac{12R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \le \hat{X}_3,
$$

a contradiction. The claim follows.

Step 11: final estimate for second case by unique continuation Carleman inequality again

We apply Proposition [22](#page-30-2) to

$$
u_4(x,t) := u(x+x_3,t), \quad x \in B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}}), \ t \in [10000nt_3,0].
$$

We recall that

<span id="page-47-1"></span>(5.24) 
$$
R_3 \le |x_3| \le 2R_3, \qquad M_3^{\frac{1}{2}} T_1^{\frac{1}{2}} R_1 \le R_3 \le M_3^{-\frac{1}{2}} T_1^{\frac{1}{2}} R_1^{M_4},
$$

$$
T_1 e^{-\frac{10R_3^2}{T_1}} \le -t_3 \le M^{-2} T_1, \quad R_1 \ge M_4.
$$

Then, for  $x \in B(0, 100nT_1^{-1/2}R_3(-t_3)^{1/2}),$ 

$$
|x+x_3| \le \left(2 + 100n\left(\frac{-t_3}{T_1}\right)^{1/2}\right) R_3 \le \left(\frac{2}{M_3^{\frac{1}{2}}} + \frac{100n}{MM_3^{\frac{1}{2}}}\right) T_1^{\frac{1}{2}} R_1^{M_4} \le \frac{1}{2} T_1^{\frac{1}{2}} R_1^{M_4},
$$
  

$$
|x+x_3| \ge \left(1 - 100n\left(\frac{-t_3}{T_1}\right)^{1/2}\right) R_3 \ge \left(M_3^{\frac{1}{2}} - \frac{100nM_3^{\frac{1}{2}}}{M}\right) T_1^{\frac{1}{2}} R_1 \ge 2T_1^{\frac{1}{2}} R_1.
$$

Moreover, for  $t \in [10000nt_3, 0]$ ,

$$
0 \ge t \ge 10000nt_3 \ge -10000nM^{-2}T_1 \ge -\frac{1}{2}T_1.
$$

Thus,  $(x+x_3,t) \in \mathcal{A}$  for  $(x,t) \in B(0,100nT_1^{-1/2}R_3(-t_3)^{1/2}) \times [10000nt_3,0]$ , where  $\mathcal A$  is defined in [\(5.15\)](#page-42-1). By (5.15), we see that

$$
|\partial_t u_4 - \Delta u_4| \le \frac{|u_4|}{T_1 M^{p-1}} \le \frac{|u_4|}{T_1} = \frac{|u_4|}{(10000n)^{-1} (T_1/(-t_3)) \times 10000n(-t_3)}
$$

in 
$$
B(0; 100nT_1^{-1/2}R_3(-t_3)^{1/2}) \times [10000nt_3, 0]
$$
. Since

$$
\frac{T_1/(-t_3)}{10000n} \ge \frac{M^2}{10000n} \ge 1,
$$
  
\n
$$
(100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})^2 \ge 10000n^2M_3R_1^2(-t_3) \ge 16000 \times 10000n(-t_3),
$$

we can apply Proposition [22](#page-30-2) to  $u_4$  on  $C = B(0, 100nT_1^{-1/2}R_3(-t_3)^{1/2}) \times [10000nt_3, 0]$ with  $\underline{s} = \overline{s} = (-t_3)/2$ . Then we obtain  $Z_4 \le X_4 + Y_4$ , where

$$
X_4 := C(n)e^{-\frac{(100nT_1^{-1/2}R_3(-t_3)^{1/2})^2}{500(-t_3/2)}}\times \int_{10000nt_3}^{0} \int_{B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} \left(\frac{|u_4|^2}{10000n(-t_3)} + |\nabla u_4|^2\right) dxdt,
$$
  
\n
$$
Y_4 := C(n)(-t_3/2)^{\frac{n}{2}} (3e)^{\frac{(100nT_1^{-1/2}R_3(-t_3)^{1/2})^2}{200(-t_3/2)}}\times \int_{B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} |u_4(x,0)|^2(-t_3/2)^{-\frac{n}{2}}e^{-\frac{|x|^2}{4(-t_3/2)}} dx,
$$
  
\n
$$
Z_4 := \int_{t_3}^{\frac{t_3}{2}} \int_{B(0,50nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} \left(\frac{|u_4|^2}{10000n(-t_3)} + |\nabla u_4|^2\right) e^{\frac{|x|^2}{4t}} dxdt.
$$

From the change of variables, the relation

$$
50nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}} \ge 50nM_3^{\frac{1}{2}}(-t_3)^{\frac{1}{2}} \ge (-t_3)^{\frac{1}{2}}
$$

and [\(5.23\)](#page-47-0) with  $-t_3 \leq M^{-2}T_1$ , it follows that

<span id="page-48-0"></span>
$$
Z_{4} \geq \int_{t_{3}}^{\frac{t_{3}}{2}} \int_{B(0,50nT_{1}^{-\frac{1}{2}}R_{3}(-t_{3})^{\frac{1}{2}})} \frac{|u_{4}|^{2}}{10000n(-t_{3})} e^{\frac{|x|^{2}}{4t}} dxdt
$$
  
\n
$$
\geq \frac{T_{1}}{10000n(-t_{3})} e^{\frac{-t_{3}}{4(t_{3}/2)}} \int_{t_{3}}^{\frac{t_{3}}{2}} \int_{B(x_{3}, (-t_{3})^{\frac{1}{2}})} \frac{|u|^{2}}{T_{1}} dxdt
$$
  
\n
$$
\geq \frac{T_{1}}{10000n e^{1/2}(-t_{3})} e^{-\frac{10nR_{3}^{2}}{T_{1}}} T_{1}^{\frac{n}{2} - \frac{2}{p-1}} \geq \frac{M^{2}}{10000n e^{1/2}} e^{-\frac{10nR_{3}^{2}}{T_{1}}} T_{1}^{\frac{n}{2} - \frac{2}{p-1}}
$$
  
\n
$$
\geq e^{-\frac{10nR_{3}^{2}}{T_{1}}} T_{1}^{\frac{n}{2} - \frac{2}{p-1}}.
$$

On the other hand, by [\(5.15\)](#page-42-1),  $T_1 e^{-10R_3^2/T_1} \le -t_3 \le M^{-2}T_1$  and  $n \ge 3$ , we see that

$$
X_4 \leq C(n)e^{-\frac{10000n^2R_3^2}{250T_1}}(10000n(-t_3))(100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})^n
$$
  
\$\times \left( \frac{(T\_1^{-\frac{1}{p-1}}M^{-1})^2}{10000n(-t\_3)} + (T\_1^{-\frac{p+1}{2(p-1)}}M^{-1})^2 \right)\$  
\$\leq C(n)M^{-4}e^{-\frac{40n^2R\_3^2}{T\_1}}\left(\frac{R\_3^2}{T\_1}\right)^{\frac{n}{2}}\left(e^{\frac{10R\_3^2}{T\_1}}+1\right)T\_1^{-\frac{2}{p-1}}(M^{-2}T\_1)^{\frac{n}{2}}\$  
\$\leq C(n)M^{-n-4}e^{-\frac{20n^2R\_3^2}{T\_1}}T\_1^{\frac{n}{2}-\frac{2}{p-1}}\$.

From  $R_3^2/T_1 \geq M_3 R_1^2 \geq M_3 M_4^2$ , it follows that

$$
X_4 \le C(n)M^{-n-4}e^{-\frac{10n^2R_3^2}{T_1}}e^{-\frac{10n^2R_3^2}{T_1}}T_1^{\frac{n}{2}-\frac{2}{p-1}}
$$
  
 
$$
\le C(n)M^{-n-4}e^{-10n^2M_3M_4^2}e^{-\frac{10n^2R_3^2}{T_1}}T_1^{\frac{n}{2}-\frac{2}{p-1}} \le e^{-\frac{10n^2R_3^2}{T_1}}T_1^{\frac{n}{2}-\frac{2}{p-1}}.
$$

Therefore,  $X_4$  can be absorbed by the right-hand side of  $(5.25)$ , and so

$$
(5.26) \tY_4 \ge e^{-\frac{20nR_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}} \ge e^{-\frac{20n^2R_3^2}{T_1}} T_1^{\frac{n}{2} - \frac{2}{p-1}}.
$$

For  $Y_4$ , using  $R_3^2/T_1 \geq M_3 M_4^2$ , we compute that

<span id="page-49-1"></span>
$$
Y_4 \leq C(n)(3e)^{\frac{100n^2R_3^2}{T_1}} \int_{B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} |u_4(x,0)|^2 dx
$$
  
\n
$$
\leq C(n)(3e)^{-\frac{100n^2R_3^2}{T_1}} (e^3)^{\frac{200n^2R_3^2}{T_1}} \int_{B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} |u_4(x,0)|^2 dx
$$
  
\n
$$
\leq e^{\frac{600n^2R_3^2}{T_1}} \int_{B(0,100nT_1^{-\frac{1}{2}}R_3(-t_3)^{\frac{1}{2}})} |u_4(x,0)|^2 dx.
$$

From this and [\(5.26\)](#page-49-1) (with a change of variables), we infer that

$$
Y_4' := \int_{B(x_3, 100nT_1^{-\frac{1}{2}} R_3(-t_3)^{\frac{1}{2}})} |u(x,0)|^2 dx \ge e^{-\frac{620n^2 R_3^2}{T_1} T_1^{\frac{n}{2} - \frac{2}{p-1}}}.
$$

Recall [\(5.24\)](#page-47-1) and  $T_2 = M^{-2}T_1$ . Then, for  $x \in B(x_3, 100nT_1^{-1/2}R_3(-t_3)^{1/2})$ ,

$$
|x| \le |x_3| + |x - x_3| \le \left(2 + 100n\left(\frac{-t_3}{T_1}\right)^{1/2}\right) R_3 \le \left(2 + \frac{100n}{M}\right) R_3
$$
  

$$
\le 3M_3^{-\frac{1}{2}} T_1^{\frac{1}{2}} R_1^{M_4} = 3M_3^{-\frac{1}{2}} M T_2^{\frac{1}{2}} R_1^{M_4} \le T_2^{\frac{1}{2}} R_1^{M_4},
$$
  

$$
|x| \ge |x_3| - |x - x_3| \ge \left(1 - 100nT_1^{-\frac{1}{2}}(-t_3)^{\frac{1}{2}}\right) R_3 \ge \left(1 - \frac{100n}{M}\right) R_3
$$
  

$$
\ge \frac{1}{2} M_3^{\frac{1}{2}} T_1^{\frac{1}{2}} R_1 = \frac{1}{2} M_3^{\frac{1}{2}} M T_2^{\frac{1}{2}} R_1 \ge T_2^{\frac{1}{2}} R_1.
$$

and so  $B(x_3, 100nT_1^{-1/2}R_3(-t_3)^{1/2}) \subset \{T_2^{1/2}R_1 \leq |x| \leq T_2^{1/2}R_1^{M_4}\}.$  From these together with  $R_3 \leq M_3^{-1/2} T_1^{1/2} R_1^{M_4}$  and  $R_1 \geq M_4$ , it follows that

<span id="page-49-0"></span>
$$
(5.27)
$$
\n
$$
\int_{T_2^{\frac{1}{2}} R_1 \leq |x| \leq T_2^{\frac{1}{2}} R_1^{M_4}} |u(x,0)|^2 dx \geq Y_4' \geq e^{-\frac{620n^2 R_3^2}{T_1} T_1^{\frac{n}{2} - \frac{2}{p-1}}} = M^{n - \frac{4}{p-1}} e^{-\frac{620n^2 R_3^2}{M_3} T_1^{\frac{n}{2} - \frac{2}{p-1}}} \geq M^{n - \frac{4}{p-1}} e^{-\frac{620n^2}{M_3} R_1^{2M_4}} T_2^{\frac{n}{2} - \frac{2}{p-1}} \geq M^{n - \frac{4}{p-1}} e^{-R_1^{2M_4} T_2^{\frac{n}{2} - \frac{2}{p-1}}}.
$$

Therefore, [\(5.19\)](#page-44-0) also holds in this case. In particular, [\(5.19\)](#page-44-0) holds in both cases  $(5.17)$  and  $(5.18)$ . By applying the Hölder inequality, we obtain

$$
e^{-R_1^{4M_4}} T_2^{\frac{n}{2} - \frac{2}{p-1}} \leq \int_{T_2^{\frac{1}{2}} R_1 \leq |x| \leq T_2^{\frac{1}{2}} R_1^{M_4}} |u(x,0)|^2 dx
$$
  

$$
\leq C \left( \int_{T_2^{\frac{1}{2}} R_1 \leq |x| \leq T_2^{\frac{1}{2}} R_1^{M_4}} |u(x,0)|^{q_c} dx \right)^{\frac{2}{q_c}} \left( T_2^{\frac{1}{2}} R_1^{M_4} \right)^{n(1-\frac{2}{q_c})},
$$

and so

$$
\begin{aligned} &\int_{T_2^{\frac{1}{2}} R_1 \leq |x| \leq T_2^{\frac{1}{2}} R_1^{M_4}} |u(x,0)|^{q_c} dx \geq \frac{1}{C} R_1^{-n(\frac{q_c}{2}-1)M_4} e^{-\frac{q_c}{2} R_1^{4M_4}} \\ &\geq \frac{1}{C} e^{-R_1^{M_4}} e^{-\frac{q_c}{2} R_1^{4M_4}} \geq e^{-R_1^{5M_4}}. \end{aligned}
$$

## Step 12: final summation

In analogy to [\[38\]](#page-56-0) for the Navier-Stokes equations, we use the conclusion of Step 11 and a summing of disjoint spatial scales to conclude. This then implies the lower bound [\(5.3\)](#page-38-2).

From  $(5.14)$  and  $(5.19)$ , it follows that

$$
\begin{split} &\int_{T_2^\frac{1}{2}R_1\leq |x|\leq T_2^\frac{1}{2}R_1^{M_4}}|u(x,0)|^{q_c}dx\geq e^{-R_1^{5M_4}}\geq e^{-(M_4^{M_4^{M^{q_c+1+\beta}}})^{5M_4}}\\ &=e^{-M_4^{5M_4\times M_4^{M^{q_c+1+\beta}}}}=e^{-M_4^{5M_4^{1+M^{q_c+1+\beta}}}}\geq e^{-M_4^{e^{M_4}}}\geq e^{-e^{e^{M_5}}}. \end{split}
$$

Similarly, we have  $1 \leq M_4 \leq R_1 \leq R_1^{M_4} \leq e^{e^{M_5}}$ . Thus,

(5.28) 
$$
\int_{T_2^{\frac{1}{2}} \leq |x| \leq e^{e^{M_5}} T_2^{\frac{1}{2}} } |u(x,0)|^{q_c} dx \geq e^{-e^{e^{M_5}}}
$$

Recall that  $T_2 = M^{-2}T_1 = M(-t'')$  and  $t''$  is arbitrarily chosen as in [\(5.4\)](#page-38-5), that is,

.

<span id="page-50-0"></span>
$$
M^{(p-1)(2p+4)}A^{\frac{n(p-1)}{p+1}}(-t') \le -t'' \le M^{-3}.
$$

Thus, we can take any  $T_2 \in [T_{\min}, T_{\max}]$ , where

<span id="page-50-2"></span>
$$
T_{\min} := M^{(p-1)(2p+4)+1} A^{\frac{n(p-1)}{p+1}}(-t'), \quad T_{\max} := M^{-2}.
$$

This together with [\(5.28\)](#page-50-0) implies that

$$
(5.29) \qquad \int_{(e^{e^{M_5}})^{k-1}T_{\min}^2 \leq |x| \leq (e^{e^{M_5}})^k T_{\min}^{\frac{1}{2}}}|u(x,0)|^{q_c} dx \geq e^{-e^{e^{M_5}}}
$$

holds for  $k = 1, \ldots, k'' - 1$ , where  $k''$  is the integer satisfying

<span id="page-50-3"></span>
$$
(5.30) \qquad (e^{e^{M_5}})^{k''-1} T_{\min}^{\frac{1}{2}} \leq e^{e^{M_5}} T_{\max}^{\frac{1}{2}} \leq (e^{e^{M_5}})^{k''} T_{\min}^{\frac{1}{2}} \left(= e^{k'' e^{M_5}} T_{\min}^{\frac{1}{2}}\right).
$$

Since  $A$  in the definition of  $T_{\min}$  is determined in  $(3.1)$  (see also Proposition [7](#page-8-2) for  $\varepsilon = \varepsilon_0/2$  in the definition of A), we have

$$
k'' \ge e^{-M_5} \log \left( e^{e^{M_5} \frac{T_{\text{max}}^{1/2}}{T_{\text{min}}^{1/2}}} \right) = e^{-M_5} \log \left( e^{e^{M_5} \frac{1}{M(p-1)(p+2) + \frac{3}{2} \int_{4}^{\frac{n(p-1)}{2(p+1)}} (-t')^{\frac{1}{2}}} \right)
$$
  
= 1 + e^{-M\_5} \log \left( \frac{1}{M^{(p-1)(p+2) + \frac{3}{2}} (48 \log(\frac{M^{p+1}}{\varepsilon\_0/2}))^{\frac{n(p-1)}{2(p+1)}} (-t')^{\frac{1}{2}}} \right)  

$$
\ge 1 + e^{-M_5} \log \left( \frac{1}{M^{(p-1)(p+2) + \frac{3}{2}} (48 \log(\frac{CM^{p+1}}{M^{-2(p+1)^2}}))^{\frac{n(p-1)}{2(p+1)}} (-t')^{\frac{1}{2}}} \right).
$$

For M sufficiently large, we note that

<span id="page-50-1"></span>
$$
(5.31) \qquad \left(48\log\frac{CM^{p+1}}{M^{-2(p+1)^2}}\right)^{\frac{n(p-1)}{2(p+1)}} \le \left(M^{\frac{(p+1)}{n(p-1)}}\right)^{\frac{n(p-1)}{2(p+1)}} \le M^{\frac{1}{2}}.
$$

Now choose  $0 < \sigma_p < 1/2$  depending only on p and satisfying

$$
(p-1)(p+2) + 1 - p(2p+3)\left(\frac{1}{2} - \sigma_p\right) = -1 - \frac{1}{2}p + p(2p+3)\sigma_p < 0.
$$

Then, from this choice of  $\sigma_p$  and [\(5.31\)](#page-50-1), we get

$$
k'' \ge 1 + e^{-M_5} \log \left( \frac{1}{M^{(p-1)(p+2)+1}(-t')^{\frac{1}{2}}} \right)
$$
  
 
$$
\ge 1 + e^{-M_5} \log \left( \frac{1}{M^{(p-1)(p+2)+1}(-t')^{\frac{1}{2}-\sigma_p}(-t')^{\sigma_p}} \right)
$$
  
 
$$
\ge 1 + e^{-M_5} \log \left( \frac{1}{M^{(p-1)(p+2)+1}(-t')^{\frac{1}{2}-\sigma_p}} \right) + \sigma_p e^{-M_5} \log \frac{1}{-t'}.
$$

Recall from our standing assumptions in Proposition [25](#page-38-3) that  $-t' < M^{-p(2p+3)}$ . Then we note that  $M^{(p-1)(p+2)+1}(-t')^{(1/2)-\sigma_p} < 1$  and obtain

$$
k'' \ge 1 + \sigma_p e^{-M_5} \log \frac{1}{-t'} \ge 1 + e^{-2M_5} \log \frac{1}{-t'}.
$$

Summing up the lower estimates [\(5.29\)](#page-50-2) for  $k = 1, ..., k'' - 1$ , we have

$$
\int_{T_{\min}^{\frac{1}{2}} \leq |x| \leq (e^{e^{M_5}})^{k''-1} T_{\min}^{\frac{1}{2}} |u(x,0)|^{q_c} dx \geq (k''-1)e^{-e^{e^{M_5}}}
$$
  

$$
\geq e^{-2M_5} e^{-e^{e^{M_5}}} \log \frac{1}{-t'} \geq e^{-e^{e^{M_6}}} \log \frac{1}{-t'}.
$$

By [\(5.30\)](#page-50-3) and  $T_{\text{max}} = M^{-2}$ , we also have

$$
\{T^{\frac{1}{2}}_{\min} \leq |x| \leq (e^{e^{M_5}})^{k''-1} T^{\frac{1}{2}}_{\min} \} \subset \{ |x| \leq e^{e^{M_5}} T^{\frac{1}{2}}_{\max} \} \subset \{ |x| \leq e^{e^{M_6}} \}.
$$

Thus,

$$
N := \int_{|x| \le e^{e^{M_6}}} |u(x,0)|^{q_c} dx \ge e^{-e^{e^{M_6}}} \log \frac{1}{-t'}.
$$

This is equivalent to  $-t' \ge \exp(-Ne^{e^{tM_6}})$ . Then, [\(5.3\)](#page-38-2) with  $x_0 = 0$  follows. On the other hand, from [\(5.28\)](#page-50-0) and  $T_2 \le T_{\text{max}} \le M^{-2}$ , it follows that

$$
N \ge \int_{T_2^{\frac{1}{2}} \le |x| \le e^{e^{M_5}} T_2^{\frac{1}{2}} } |u(x,0)|^{q_c} dx \ge e^{-e^{e^{M_5}}} \ge Me^{-e^{e^{M_6}}}
$$

This shows  $(5.2)$ . The proof of Proposition [25](#page-38-3) is complete.

.

By using Proposition [25,](#page-38-3) we show our main quantitative estimate.

*Proof of Proposition [5.](#page-3-3)* Assume that u satisfies the Lorentz norm bound  $(1.6)$  and define  $N$  by  $(1.7)$ . We consider 2 cases.  $\overline{M}^C$ <sub>0</sub>

Case 1: 
$$
N < Me^{-e^{e^2}}
$$

<span id="page-51-0"></span>Since [\(5.2\)](#page-38-1) does not hold, the contraposition of Proposition [25](#page-38-3) shows that

$$
(5.32) \qquad \qquad (-t')^{\frac{2}{p-1}-\frac{n}{2}} \int_{t'}^{t'/2} \int_{B(x_0, A(-t')^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt \le \varepsilon
$$

for all  $t' \in (-M^{-p(2p+3)}, 0)$ . Here  $\varepsilon = \varepsilon_0/2$ ,  $\varepsilon_0$  is defined in Proposition [7](#page-8-2) and A is defined in  $(3.1)$ . Now  $t_*$  defined by  $(1.8)$  satisfies

<span id="page-52-1"></span>
$$
(5.33) \t -t_* = M^{-p(2p+3)-1} \exp\left(-Ne^{e^{e^{M^{C_0}}}}\right) \le M^{-p(2p+3)-1} < M^{-p(2p+3)}.
$$

By [\(5.32\)](#page-51-0) with  $t' = t_*$ , we get

<span id="page-52-2"></span>
$$
(5.34) \qquad \qquad (-t_*)^{\frac{2}{p-1}-\frac{n}{2}} \int_{t_*}^{t_*/2} \int_{B(x_0,A(-t_*)^{\frac{1}{2}})} |u(x,t)|^{p+1} dx dt \le \varepsilon.
$$

Then by Proposition [7](#page-8-2) with  $t_0 = 0$  and  $\delta = (-t_*)^{\frac{1}{2}}$  and by  $\varepsilon < M^{-2(p+1)^2} \varepsilon_1^{2(p+1)}$ with the fact that  $\varepsilon_1$  depends only on n and p, we get

$$
||u||_{L^{\infty}(B(x_0, \frac{1}{4}(-t_*))^{\frac{1}{2}})\times(\frac{1}{16}t_*,0))} \leq CM\varepsilon^{\frac{1}{2(p+1)^2}}(-t_*)^{-\frac{1}{p-1}} \leq C\varepsilon_1^{\frac{1}{p+1}}(-t_*)^{-\frac{1}{p-1}} \leq C(n,p)(-t_*)^{-\frac{1}{p-1}}.
$$

Hence we obtain [\(1.9\)](#page-4-4).

**Case 2:**  $N \ge Me^{-e^{e^{M^{C_0}}}}$ By [\(5.33\)](#page-52-1), we have  $t_* \in (-M^{-p(2p+3)}, 0)$ . Since

$$
-t_* = M^{-p(2p+3)-1} \exp\left(-Ne^{e^{e^{M^{C_0}}}}\right) < \exp\left(-Ne^{e^{e^{M^{C_0}}}}\right),
$$

<span id="page-52-0"></span>we observe that [\(5.3\)](#page-38-2) does not hold. Then by the contraposition of Proposition [25,](#page-38-3) we must have  $(5.34)$ . By the same reasoning as Case 1, we get  $(1.9)$ .

## 6. Proof of main theorems

We are now in a position to prove our main theorems (Theorems [1,](#page-2-1) [2](#page-2-4) and [3\)](#page-3-0).

Proof of Theorem [1.](#page-2-1) By applying Proposition [5](#page-3-3) (specifically [\(1.9\)](#page-4-4)) to

<span id="page-52-4"></span>
$$
u_t(x,s) := (t^{\frac{1}{2}})^{\frac{2}{p-1}} u(t^{\frac{1}{2}}x, ts+t),
$$

the following statement holds true. If  $M_0$  is sufficiently large and if u is a classical solution on  $\mathbf{R}^n \times (0, t]$  with

(6.1) 
$$
\max\left(M_0, \sup_{s\in(0,t)}\|u(\cdot,s)\|_{L^{q_c}(\mathbf{R}^n)}\right) \leq M_t
$$

for some  $M_t > 0$ , then there exists  $C_* > 0$  depending only on n and p such that

(6.2) 
$$
t^{\frac{1}{p-1}} \|u(\cdot,t)\|_{L^{\infty}(\mathbf{R}^n)} \leq e^{e^{e^{M_t^{C*}}}}
$$

With this in hand, we proceed with proving Theorem [1.](#page-2-1)

We suppose for contradiction that

<span id="page-52-5"></span>
$$
\limsup_{t\to T}\frac{\|u(\cdot,t)\|_{L^{q_c}(\mathbf{R}^n)}}{\left(\log\log\log\log\left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/(2C_*)}}<\infty.
$$

.

This implies that there exists  $N_* > 0$  such that for all  $t \in (0, T)$ ,

<span id="page-52-3"></span>
$$
(6.3) \quad \sup_{s\in(0,t)} \|u(\cdot,s)\|_{L^{q_c}(\mathbf{R}^n)} \le N_* \left(\log\log\log\log\left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/(2C_*)}.
$$

Since  $\lim_{t\to T} \log \log \log (1/(T-t)^{1/(2(p-1))}) = \infty$ , there exists  $0 < \alpha < T$  depending only on p,  $C_*, T, M_0$  and  $N_*$  such that for all  $t \in (T - \alpha, T)$ ,

<span id="page-53-0"></span>(6.4) 
$$
\max(N_*^2, M_0) \le \left(\log \log \log \left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/C_*},
$$

<span id="page-53-1"></span>(6.5) 
$$
t^{-\frac{1}{p-1}}(T-t)^{\frac{1}{2(p-1)}} < \frac{1}{2}(p-1)^{-\frac{1}{p-1}}.
$$

Using [\(6.3\)](#page-52-3) and [\(6.4\)](#page-53-0) gives that for all  $t \in (T - \alpha, T)$ ,

$$
\sup_{s \in (0,t)} \|u(\cdot,s)\|_{L^{q_c}(\mathbf{R}^n)} \le N_* \left(\log \log \log \log \left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/(2C_*)}
$$
  

$$
\le \left(\log \log \log \log \left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/C_*},
$$
  

$$
M_0 \le \left(\log \log \log \log \left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/C_*}
$$

This shows that [\(6.1\)](#page-52-4) is satisfied with

<span id="page-53-2"></span>
$$
M_t := \left(\log \log \log \left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/C_*}
$$

.

By substituting this into [\(6.2\)](#page-52-5) and using [\(6.5\)](#page-53-1), we have for all  $t \in (T - \alpha, T)$ ,

(6.6) 
$$
||u(\cdot,t)||_{L^{\infty}(\mathbf{R}^n)} \leq t^{-\frac{1}{p-1}}(T-t)^{-\frac{1}{2(p-1)}} < \frac{(p-1)^{-\frac{1}{p-1}}}{2(T-t)^{\frac{1}{p-1}}}.
$$

But from [\[34,](#page-55-2) Proposition 23.1], we have that for all  $t \in (0, T)$ ,

(6.7) 
$$
||u(\cdot,t)||_{L^{\infty}(\mathbf{R}^n)} \geq \frac{(p-1)^{-\frac{1}{p-1}}}{(T-t)^{\frac{1}{p-1}}}.
$$

This contradicts [\(6.6\)](#page-53-2). Hence, we must have

<span id="page-53-3"></span>
$$
\limsup_{t\to T}\frac{\|u(\cdot,t)\|_{L^{q_c}(\mathbf{R}^n)}}{\left(\log\log\log\log\left(\frac{1}{(T-t)^{1/(2(p-1))}}\right)\right)^{1/(2C_*)}}=\infty.
$$

This shows the desired conclusion with  $c := 1/(2C_*)$ . The proof is complete.  $\square$ 

Proof of Theorem [2.](#page-2-4) Let

(6.8) 
$$
(1 + M^{-2(2p^2 + 3p + 2)})^{-1}T < t < T.
$$

We see that the assumptions of Theorem [2](#page-2-4) allow us to apply Proposition [5](#page-3-3) (specifically  $(1.9)$  to

<span id="page-53-4"></span>
$$
u_t(x,s) := (t^{\frac{1}{2}})^{\frac{2}{p-1}} u(t^{\frac{1}{2}}x, ts+t).
$$

Then we can check that

$$
t^{\frac{1}{p-1}} \|u(\cdot,t)\|_{L^{\infty}(\mathbf{R}^n)} \leq C(n,p) M^{\frac{p(2p+3)+1}{p-1}} \exp\left(\frac{1}{p-1} e^{e^{e^{M^{C_0}}}} \int_{\mathbf{R}^n} |u(x,t)|^{q_c} dx\right)
$$
  

$$
\leq M^{\frac{p(2p+3)+2}{p-1}} \exp\left(\frac{1}{p-1} e^{e^{e^{M^{C_0}}}} \int_{\mathbf{R}^n} |u(x,t)|^{q_c} dx\right)
$$

for M sufficiently large. This together with  $(6.7)$  gives

$$
(p-1)^{-\frac{1}{p-1}}\left(\frac{t}{T-t}\right)^{\frac{1}{p-1}}M^{-\frac{2p^2+3p+2}{p-1}} \leq \exp\left(\frac{1}{p-1}e^{e^{tM^{C_0}}}\int_{\mathbf{R}^n}|u(x,t)|^{q_c}dx\right).
$$

From [\(6.8\)](#page-53-4), we see that  $1 \le (t/(T-t))^{1/(2(p-1))} M^{-(2p^2+3p+2)/(p-1)}$ . Thus,

$$
(p-1)^{-\frac{1}{p-1}}\left(\frac{t}{T-t}\right)^{\frac{1}{2(p-1)}} \leq \exp\left(\frac{1}{p-1}e^{e^{e^{M^{C_0}}}}\int_{\mathbf{R}^n}|u(x,t)|^{q_c}dx\right),
$$

and so

$$
\int_{\mathbf{R}^n} |u(x,t)|^{q_c} dx \ge \frac{1}{2} e^{-e^{e^{M^C 0}}} \log \left( \frac{t}{T-t} \right) - e^{-e^{e^{M^C 0}}} \log(p-1).
$$

This together with [\(6.8\)](#page-53-4) gives the desired conclusion with  $C := 2C_0$ , for M sufficiently large.  $\Box$ 

Proof of Theorem [3.](#page-3-0) We see that the assumptions of Theorem [3](#page-3-0) allow us to apply Proposition [5](#page-3-3) (specifically [\(1.9\)](#page-4-4)) to

$$
u_t(x,s) := (t^{\frac{1}{2}})^{\frac{2}{p-1}} u(t^{\frac{1}{2}}x, ts+t).
$$

This gives

<span id="page-54-9"></span>
$$
(6.9) \quad M^{-\frac{2p^2+3p+2}{p-1}}|u(0,t)|t^{\frac{1}{p-1}} \leq \exp\left(\frac{1}{p-1}e^{e^{uC_0}}\int_{|x|\leq t^{\frac{1}{2}}e^{e^{M C_0}}}|u(x,t)|^{q_c}dx\right).
$$

As (by assumption) lim  $\sup_{t\to\infty} |u(0,t)| = \infty$ , there exists  $t_n \to \infty$  such that

$$
M^{-\frac{2p^2+3p+2}{p-1}}|u(0,t_n)| \ge 1 \quad \text{ for all } n \in \mathbf{N}.
$$

Substituting this into  $(6.9)$  then readily gives the desired conclusion.

## <span id="page-54-0"></span>**ACKNOWLEDGMENTS**

The first author thanks Philippe Souplet for references and the Institute of Science Tokyo for its hospitality. The second author was supported in part by JSPS KAKENHI Grant Numbers 17K05312, 21H00991 and 21H04433. The third author was supported in part by JSPS KAKENHI Grant Numbers 22H01131, 22KK0035 and 23K12998.

### <span id="page-54-1"></span>**REFERENCES**

- <span id="page-54-7"></span>[1] T. Barker, Localized quantitative estimates and potential blow-up rates for the Navier-Stokes equations. SIAM J. Math. Anal. 55 (2023), no. 5, 5221–5259.
- <span id="page-54-5"></span>[2] T. Barker, C. Prange, Quantitative regularity for the Navier-Stokes equations via spatial concentration. Comm. Math. Phys. 385 (2021), no. 2, 717–792.
- <span id="page-54-4"></span>[3] T. Barker, C. Prange, From concentration to quantitative regularity: a short survey of recent developments for the Navier-Stokes equations. Vietnam J. Math. 52 (2024), no. 3, 707–734.
- <span id="page-54-6"></span>[4] S. Blatt, M. Struwe, An analytic framework for the supercritical Lane-Emden equation and its gradient flow. Int. Math. Res. Not. IMRN 2015, no. 9, 2342–2385.
- <span id="page-54-2"></span>[5] H. Brezis, T. Cazenave, A nonlinear heat equation with singular initial data. J. Anal. Math. 68 (1996), 277–304.
- <span id="page-54-8"></span>[6] C. P. Calderón, Existence of weak solutions for the Navier-Stokes equations with initial data in  $L^p$ . Trans. Amer. Math. Soc. 318 (1990), no. 1, 179-200.
- <span id="page-54-3"></span>[7] M. del Pino, M. Musso, J. Wei, Type II blow-up in the 5-dimensional energy critical heat equation. Acta Math. Sin. (Engl. Ser.) 35 (2019), no. 6, 1027–1042.
- <span id="page-55-3"></span>[8] M. del Pino, M. Musso, J. Wei, Q. Zhang, Y. Zhou, Type II Finite time blow-up for the three dimensional energy critical heat equation, preprint, [arXiv:2002.05765.](http://arxiv.org/abs/2002.05765)
- <span id="page-55-9"></span>[9] L. Escauriaza, G. A. Seregin, V. Šverák, Backward uniqueness for parabolic equations. Arch. Ration. Mech. Anal. 169 (2003), no. 2, 147–157.
- <span id="page-55-16"></span>[10] M. Fila, J. R. King, M. Winkler, E. Yanagida, Optimal lower bound of the grow-up rate for a supercritical parabolic equation. J. Differential Equations 228 (2006), no. 1, 339–356.
- <span id="page-55-17"></span>[11] M. Fila, M. Winkler, E. Yanagida, Grow-up rate of solutions for a supercritical semilinear diffusion equation. J. Differential Equations 205 (2004), no. 2, 365–389.
- <span id="page-55-21"></span>[12] Y. Giga, R. V. Kohn, Asymptotically self-similar blow-up of semilinear heat equations. Comm. Pure Appl. Math. 38 (1985), no. 3, 297–319.
- <span id="page-55-1"></span>[13] Y. Giga, R. V. Kohn, Characterizing blowup using similarity variables. Indiana Univ. Math. J. 36 (1987), no. 1, 1–40.
- <span id="page-55-25"></span>[14] Y. Giga, R. V. Kohn, Nondegeneracy of blowup for semilinear heat equations. Comm. Pure Appl. Math. 42 (1989), no. 6, 845–884.
- <span id="page-55-26"></span>[15] L. Grafakos, Classical Fourier analysis. Second edition. Graduate Texts in Mathematics, 249. Springer, New York, 2008.
- <span id="page-55-11"></span>[16] J. Harada, A type II blowup for the six dimensional energy critical heat equation. Ann. PDE 6 (2020), no. 2, Paper No. 13, 63 pp.
- <span id="page-55-20"></span>[17] R. Hu, P. T. Nguyen, Q. H. Nguyen, P. Zhang. Quantitative bounds for bounded solutions to the Navier-Stokes equations in endpoint critical Besov spaces, preprint, [arXiv:2411.06483.](http://arxiv.org/abs/2411.06483)
- <span id="page-55-24"></span>[18] R. Mandel, Real interpolation for mixed Lorentz spaces and Minkowski's inequality. Journal of Analysis & Its Applications/Zeitschrift für Analysis & ihre Anwendungen. 2023 Jul 1;42.
- <span id="page-55-12"></span>[19] H. Matano, F. Merle, Classification of type I and type II behaviors for a supercritical nonlinear heat equation. J. Funct. Anal. 256 (2009), no. 4, 992–1064.
- <span id="page-55-13"></span>[20] H. Matano, M. Merle, Threshold and generic type I behaviors for a supercritical nonlinear heat equation. J. Funct. Anal. 261 (2011), no. 3, 716–748.
- <span id="page-55-19"></span>[21] F. Merle, P. Raphël, Blow up of the critical norm for some radial  $L^2$  super critical nonlinear Schrödinger equations. Amer. J. Math. 130 (2008), no. 4, 945–978.
- <span id="page-55-23"></span>[22] Y. Meyer, Wavelets, paraproducts, and Navier-Stokes equations. Current developments in mathematics, 1996 (Cambridge, MA), 105–212, Int. Press, Boston, MA, 1997.
- <span id="page-55-6"></span>[23] H. Miura, J. Takahashi, Blow-up of the critical norm for a supercritical semilinear heat equation, to appear in J. Eur. Math. Soc. (JEMS), arXiv: 2206.10790.
- <span id="page-55-7"></span>[24] H. Miura, J. Takahashi, Critical norm blow-up for the energy supercritical nonlinear heat equation, preprint, arXiv: 2310.09750.
- <span id="page-55-8"></span>[25] H. Miura, J. Takahashi, On critical norm blow-up for a nonlinear heat equation, to appear in Proceedings of the conference "Critical Phenomena in Nonlinear Partial Differential Equations, Harmonic Analysis, and Functional Inequalities."
- <span id="page-55-18"></span>[26] N. Mizoguchi, Growup of solutions for a semilinear heat equation with supercritical nonlinearity. J. Differential Equations 227 (2006), no. 2, 652–669.
- <span id="page-55-5"></span>[27] N. Mizoguchi, Ph. Souplet, Optimal condition for blow-up of the critical  $L<sup>q</sup>$  norm for the semilinear heat equation. Adv. Math. 355 (2019), 106763, 24 pp.
- <span id="page-55-10"></span>[28] A. Mukai, Y. Seki, Refined construction of type II blow-up solution for semilinear heat equations with Joseph-Lundgren supercritical nonlinearity. Discrete Contin. Dyn. Syst. 41 (2021), no. 10, 4847–4885.
- <span id="page-55-27"></span><span id="page-55-22"></span>[29] R. O'Neil, Convolution operators and  $L(p, q)$  spaces. Duke Math. J. 30 (1963), 129–142.
- [30] S. Palasek, Improved quantitative regularity for the Navier-Stokes equations in a scale of critical spaces. Arch. Ration. Mech. Anal. 242 (2021), no. 3, 1479–1531.
- <span id="page-55-0"></span>[31] S. Palasek, A minimum critical blowup rate for the high-dimensional Navier-Stokes equations. J. Math. Fluid Mech. 24 (2022), no. 4, Paper No. 108, 28 pp.
- <span id="page-55-14"></span>[32] P. Poláčik, E. Yanagida, On bounded and unbounded global solutions of a supercritical semilinear heat equation. Math. Ann. 327 (2003), no. 4, 745–771.
- <span id="page-55-15"></span>[33] P. Poláčik, E. Yanagida, Global unbounded solutions of the Fujita equation in the intermediate range. Math. Ann. 360 (2014), no. 1-2, 255–266.
- <span id="page-55-2"></span>[34] P. Quittner, Ph. Souplet, Superlinear parabolic problems. Blow-up, global existence and steady states. Second edition. Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser/Springer, Cham, 2019.
- <span id="page-55-4"></span>[35] R. Schweyer, Type II blow-up for the four dimensional energy critical semi linear heat equation. J. Funct. Anal. 263 (2012), no. 12, 3922–3983.
- <span id="page-56-3"></span><span id="page-56-2"></span>[36] Y. Seki, Type II blow-up mechanisms in a semilinear heat equation with critical Joseph-Lundgren exponent. J. Funct. Anal. 275 (2018), no. 12, 3380–3456.
- [37] Ph. Souplet, Morrey spaces and classification of global solutions for a supercritical semilinear heat equation in  $\mathbb{R}^n$ . J. Funct. Anal. 272 (2017), no. 5, 2005–2037.
- <span id="page-56-0"></span>[38] T. Tao, Quantitative bounds for critically bounded solutions to the Navier-Stokes equations. Nine mathematical challenges—an elucidation, 149–193, Proc. Sympos. Pure Math., 104, Amer. Math. Soc., Providence, RI, 2021.
- <span id="page-56-5"></span>[39] F. B. Weissler, Local existence and nonexistence for semilinear parabolic equations in  $L^p$ . Indiana Univ. Math. J. 29 (1980), no. 1, 79–102.
- <span id="page-56-1"></span>[40] F. B. Weissler, Existence and nonexistence of global solutions for a semilinear heat equation. Israel J. Math. 38 (1981), no. 1-2, 29–40.
- <span id="page-56-4"></span>[41] F. B. Weissler,  $L^p$ -energy and blow-up for a semilinear heat equation. Nonlinear functional analysis and its applications, Part 2 (Berkeley, Calif., 1983), 545–551, Proc. Sympos. Pure Math., 45, Part 2, Amer. Math. Soc., Providence, RI, 1986.

Department of Mathematical Sciences, University of Bath, Bath BA2 7AY. UK Email address, Corresponding author: tb2130@bath.ac.uk

Department of Mathematics, Institute of Science Tokyo, Tokyo 152-8551, Japan Email address: hideyuki@math.titech.ac.jp

Department of Mathematical and Computing Science, Institute of Science Tokyo, Tokyo 152-8552, Japan

Email address: takahashi@c.titech.ac.jp