Twice Epi-Differentiability of Orthogonally Invariant Matrix Functions and Application

Jiahuan He

School of Science, Harbin University of Science and Technology, Harbin, 150080, P.R. China

Chao Kan and Wen Song[∗]

School of Mathematical and Sciences, Harbin Normal University,

Harbin, 150025, P.R. China

Abstract. In this paper, our focus lies on the study of the second-order variational analysis of orthogonally invariant matrix functions. It is well-known that an orthogonally invariant matrix function is an extended-real-value function defined on $\mathbb{M}_{m,n}$ $(n \leq m)$ of the form $f \circ \sigma$ for an absolutely symmetric function $f: \mathbb{R}^n \to [-\infty, +\infty]$ and the singular values $\sigma: \mathbb{M}_{m,n} \to \mathbb{R}^n$. We establish several second-order properties of orthogonally invariant matrix functions, such as parabolic epi-differentiability, parabolic regularity, and twice epi-differentiability when their associated absolutely symmetric functions enjoy some properties. Specifically, we show that the nuclear norm of a real $m \times n$ matrix is twice epi-differentiable and we derive an explicit expression of its second-order epi-derivative. Moreover, for a convex orthogonally invariant matrix function, we calculate its second subderivative and present sufficient conditions for twice epi-differentiability. This enables us to establish second-order optimality conditions for a class of matrix optimization problems.

Keywords: orthogonally invariant matrix functions; subderivatives; second subderivatives; twice epi-differentiability; second-order optimality conditions

Mathematics Subject classification: 15A18; 49J52; 49J53; 94A11

1 Introduction

Consider convex matrix optimization problem

$$
\min_{X \in \mathbb{M}_{m,n}} \psi(X) + \theta(X),
$$

where $\mathbb{M}_{m,n}$ stands for the Euclidean space of $m \times n$ real matrices, with inner product $\langle X, Y \rangle =$ $tr(X^TY)$, $\psi: \mathbb{M}_{m,n} \to \mathbb{R}$ is a twice continuously differentiable convex function, and $\theta: \mathbb{M}_{m,n} \to$ [−∞, +∞] is a proper closed convex function. The problems described above represent a broad and significant class of convex matrix optimization problems with extensive applications in many fields, including matrix completion, rank minimization, graph theory, and machine learning. The examples of θ include the indicator function over the positive semidefinite cone [\[21\]](#page-27-0), the nuclear norm (defined as the sum of all singular values of a matrix) [\[22\]](#page-28-0), the spectral norm (which represents the largest singular value of a matrix) [\[12\]](#page-27-1), and the matrix Ky Fan k-norm $(1 \leq k \leq n)$ function (which corresponds to the sum of k largest singular values of a matrix) [\[4\]](#page-27-2). The functions θ in the aforementioned examples belong to a distinguished class, which can be expressed either in the form of

$$
\theta(X) = (g \circ \lambda)(X), \ X \in \mathbb{S}^n
$$

with the function $g: \mathbb{R}^n \to [-\infty, +\infty]$ being proper closed convex and symmetric, and λ is a function, which assigns to each matrix $X \in \mathbb{S}^n$ its eigenvalue vector $(\lambda_1(X), \ldots, \lambda_n(X))$ arranged in nonincreasing order, referred to as spectral functions [\[8\]](#page-27-3), or in the form of

$$
\theta(X) = (f \circ \sigma)(X), \ X \in \mathbb{M}_{m,n} \tag{1.1}
$$

with the function $f: \mathbb{R}^n \to [-\infty, +\infty]$ being proper closed convex and absolutely symmetric, and σ is a function, which assigns to each matrix $X \in M_{m,n}$ its singular value vector $(\sigma_1(X), \ldots, \sigma_n(X))$

[∗]Corresponding author, E-mail address: wsong@hrbnu.edu.cn

arranged in nonincreasing order, known as orthogonally invariant matrix functions [\[9\]](#page-27-4) or singular value functions [\[10\]](#page-27-5). Recall that $F: \mathbb{M}_{m,n} \to [-\infty, +\infty]$ is an orthogonally invariant matrix function if for any matrix X in $\mathbb{M}_{m,n}$, any $m \times m$ orthogonal matrix U_m and any $n \times n$ orthogonal matrix U_n , one has

$$
F(U_m^T X U_n) = F(X).
$$

The work of Lewis and Sendov [\[10,](#page-27-5) [11\]](#page-27-6) serves as a foundational contribution to the variational analysis of this class of functions. Lewis [\[9\]](#page-27-4) showed that convexity, lower semi-continuity, differentiability, and essentially smoothness of the absolutely symmetric function f in (1.1) are inherited by the orthogonally invariant matrix function θ . A similar observation was made by Lewis and Sendov [\[10\]](#page-27-5) about Fréchet differentiability, regularity, and strictly differentiability, and by Cui et al. [\[2\]](#page-27-7) about the \mathcal{C}^2 -cone reducibility and the metric subregularity of their subdifferentials. The calculation of various notions of subdifferentials for orthogonally invariant matrix functions, including the limiting subdifferential, the Clarke subdifferential, and the proximal subdifferential, which play an important role in second-order variational analysis, was demonstrated in Lewis and Sendov [\[10,](#page-27-5) [11\]](#page-27-6). The central question addressed in this paper is whether the remarkable pattern observed can be extended to other significant second-order variational properties.

It was proved in Mohammadi and Sarabi [\[15\]](#page-27-8) that important second-order variational properties of a composite function $g \circ \varphi$, where $g: \mathbb{Y} \to [-\infty, +\infty]$ is convex and $\varphi: \mathbb{X} \to \mathbb{Y}$ is twice differentiable with X and Y being finite-dimensional Hilbert space, can be established at any $\bar{x} \in \text{dom}(q \circ \varphi)$, provided that g is parabolically regular and that the following metric subregularity constraint qualification is satisfied: There exists a constant $\kappa \geq 0$, such that the estimate

$$
dist(x, dom(g \circ \varphi)) \leq \kappa dist(\varphi(x), dom g),
$$

holds for all x in a neighborhood of \bar{x} . It was observed in [\[3\]](#page-27-9) that the metric subregularity constraint qualification, when applied to spectral functions, is inherently satisfied. In addition, because a quadratic expansion of parabolic nature suffices to substitute for the twice differentiability of the inner function, Mohammadi and Sarabi [\[14\]](#page-27-10) characterized parabolic regularity for spectral functions and calculated their second subderivative when the symmetric functions associated with them are convex. In the spirit of Mohammadi and Sarabi [\[14,](#page-27-10)[15\]](#page-27-8), we study second-order variational properties, including parabolic epi-differentiability and twice epi-differentiability, of orthogonally invariant matrix functions.

The first-order directional derivatives of the singular values, as developed in [\[5\]](#page-27-11), serve as a powerful tool for characterizing the first-order necessary optimality conditions in matrix cone optimization problems. To derive the second-order necessary or sufficient conditions for the matrix optimization problem, it is essential to examine the second-order directional derivatives of the singular values. Zhang et al. [\[23\]](#page-28-1) proposed a direct method to derive the formula for the secondorder directional derivative of any eigenvalue of a symmetric matrix, as presented in Torki [\[20\]](#page-27-12). From this, they established a corresponding formula for the second-order directional derivative of any singular value of a matrix.

The structure of the paper is organized as follows. In Section 2, we review key concepts from variational analysis, along with important notions related to eigenvalues and singular values, which are central to the discussions throughout the paper. Section 3 presents explicit expressions for the first- and second-order epi-derivatives of the nuclear norm. In Section 4, we establish a chain rule for the subderivatives of orthogonally invariant matrix functions in [\(1.1\)](#page-0-0). Additionally, we derive the tangent cone and second-order tangent sets associated with orthogonally invariant matrix sets. It is further demonstrated that the subderivative is a symmetric function with respect to a subset of the space of signed permutation matrices. Finally, in Section 5, we compute the second subderivative of orthogonally invariant matrix functions under the assumption that the corresponding absolutely symmetric functions are convex. We also present sufficient conditions for the twice epi-differentiability of orthogonally invariant matrix functions. As an application, we derive second-order optimality conditions for a class of matrix optimization problems.

2 Preliminaries

2.1 Notation

Throughout the whole paper, we will assume that m and n are natural numbers and $n \leq m$. The notations and concepts of convex analysis that we employ are standard $[6, 16, 18]$ $[6, 16, 18]$ $[6, 16, 18]$ $[6, 16, 18]$. Given a nonempty set $C \subset \mathbb{R}^n$, the indicator function of the set C, denoted by δ_C , is defined by $\delta_C(x) = 0$ if $x \in C$ and $+\infty$ otherwise. The support function of the set C is defined by $\sigma_C(\cdot) := \sup_{s \in C} \langle s, \cdot \rangle$.

The distance function from a point x to the set C, denoted by $dist(x, C)$ or $d_C(x)$, is defined by $dist(x, C) = d_C(x) := \inf\{|x - y\| : y \in C\}.$ The tangent cone to C at \bar{x} is defined by

$$
T_C(\bar{x}) = \{ w \in \mathbb{R}^n \colon \exists t_k \downarrow 0, w_k \to w \text{ as } k \to \infty \text{ with } \bar{x} + t_k w_k \in C \}.
$$

The second-order tangent set to $C \subset \mathbb{R}^n$ at $\bar{x} \in C$ for a tangent vector $w \in T_C(\bar{x})$ is given by

$$
T_C^2(\bar{x}, w) = \{u \in \mathbb{R}^n : \exists t_k \downarrow 0, u_k \to u \text{ as } k \to \infty \text{ with } \bar{x} + t_k w + \frac{1}{2} t_k^2 u_k \in C\}.
$$

A set $C \subset \mathbb{R}^n$ is called parabolically derivable at \bar{x} for w if $T_C^2(\bar{x}, w)$ is nonempty, and for each $u \in T_C^2(\bar{x}, w)$, there exists $\epsilon > 0$ and an arc $\xi : [0, \epsilon] \to C$ with $\xi(0) = \bar{x}, \xi'_{+}(0) = w$, and $\xi''_{+}(0) = u$, where $\xi''_{+}(0) = \lim_{t \downarrow 0} [\xi(t) - \xi(0) - t\xi'_{+}(0)] / \frac{1}{2}t^2$.

Let $g: \mathbb{R}^n \to \mathbb{R} \cup \{\pm \infty\}$ be a function. The domain and epigraph of g are defined as dom $g :=$ $\{x \in \mathbb{R}^n : g(x) < \infty\}$ and epig := $\{(x, r) \in \mathbb{R}^n \times \mathbb{R} : g(x) \leq r\}$. g is proper if domg $\neq \emptyset$ and $g(x) > -\infty$ for all $x \in \text{dom}g$. g is convex and lower semi-continuous (lsc) if epig is convex and closed, respectively. The function g is called locally Lipschitz continuous around \bar{x} relative to $C \subset \text{dom} g$ with constant $l \geq 0$ if $\bar{x} \in C$ with $g(\bar{x})$ finite, and there exists a neighborhood U of \bar{x} such that

$$
|g(x) - g(y)| \le l ||x - y||
$$
 for all $x, y \in U \cap C$.

We say that g is locally Lipschitz continuous relative to C if it is locally Lipschitz continuous around every $\bar{x} \in C$ relative to C. The directional derivative of g at \bar{x} in the direction w is defined as

$$
g'(\bar{x}; w) := \lim_{t \downarrow 0} \frac{g(\bar{x} + tw) - g(\bar{x})}{t},
$$

and the subderivative of q at \bar{x} in the direction w is defined as

$$
dg(\bar{x})(w) := \liminf_{\substack{t \downarrow 0 \\ w' \to w}} \frac{g(\bar{x} + tw') - g(\bar{x})}{t}.
$$

It is known that if q is locally Lipschitz continuous around \bar{x} relative to its domain, then dom $dq(\bar{x}) =$ $T_{\text{dom}g}(\bar{x})$ (cf. Mohammadi and Sarabi [\[15,](#page-27-8) Proposition 2.2]).

In what follows, we'll review the classical notion of second-order variational analysis. Let's first recall the concepts of the second subderivative and parabolic regularity for functions, respectively. Given a function $g: \mathbb{R}^n \to \mathbb{R} \cup {\{\pm \infty\}}$ and a point $\bar{x} \in \mathbb{R}^n$ with $g(\bar{x})$ finite, we say g is (parabolic) second order directionally differentiable at \bar{x} if g is directionally differentiable at \bar{x} and for any $w, z \in \mathbb{R}^n$,

$$
\lim_{\tau \downarrow 0} \frac{g(\bar{x} + \tau w + \frac{1}{2}\tau^2 z) - g(\bar{x}) - \tau g'(\bar{x}; w)}{\frac{1}{2}\tau^2}
$$
 exists;

and the above limit is said to be the (parabolic) second order directional derivative of g at \bar{x} along the directions w and z, denoted by $g''(x; w, z)$.

Define the parametric family of second-order difference quotients for g at \bar{x} for $v \in \mathbb{R}^n$ by

$$
\triangle_{\tau}^2 g(\bar{x} \mid v)(w) := \frac{g(\bar{x} + \tau w) - g(\bar{x}) - \tau \langle v, w \rangle}{\frac{1}{2}\tau^2} \text{ with } w \in \mathbb{R}^n, \tau > 0.
$$

The second subderivative of g at \bar{x} for $v \in \mathbb{R}^n$ is defined by

$$
d^2 g(\bar{x} | v)(w) = \liminf_{\substack{\tau \downarrow 0 \\ w' \to w}} \triangle^2_{\tau} g(\bar{x} | v)(w')
$$
 with $w \in \mathbb{R}^n$.

We say that g is twice epi-differentiable at \bar{x} for $v \in \mathbb{R}^n$ if the functions $\triangle^2_{\tau} g(\bar{x} \mid v)$ epi-converge to $d^2g(\bar{x} \mid v)$ as $\tau \downarrow 0$, see [\[18,](#page-27-15) Definition 13.6].

Now consider another kind of second-order difference quotient, which is called a parabolic difference quotient, defined by

$$
\triangle_{\tau}^{2} g(\bar{x})(w \mid z) := \frac{g(\bar{x} + \tau w + \frac{1}{2}\tau^{2}z) - g(\bar{x}) - \tau dg(\bar{x})(w)}{\frac{1}{2}\tau^{2}} \text{ with } w, z \in \mathbb{R}^{n}, \tau > 0.
$$

The parabolic subderivative of g at \bar{x} for w with $dq(\bar{x})(w)$ finite with respect to z is defined by

$$
d^2g(\bar{x})(w \mid z) := \liminf_{\substack{\tau \downarrow 0 \\ z' \to z}} \triangle^2_{\tau}g(\bar{x})(w \mid z').
$$

g is said to be parabolically epi-differentiable at \bar{x} for w if dom $d^2g(\bar{x})(w \mid \cdot) \neq \emptyset$ and for every $z \in \mathbb{R}^n$ and every sequence $\tau_k \downarrow 0$ there exist sequences $z_k \rightarrow z$ such that

$$
d^2 g(\bar{x})(w \mid z) = \lim_{k \to \infty} \frac{g(\bar{x} + \tau_k w + \frac{1}{2}\tau_k^2 z_k) - g(\bar{x}) - \tau_k dg(\bar{x})(w)}{\frac{1}{2}\tau_k^2}.
$$

We say that q is parabolically epi-differentiable at \bar{x} if it satisfies the above condition at \bar{x} for any w with $dq(\bar{x})(w)$ finite. It has been demonstrated in Mohammadi and Sarabi [\[15,](#page-27-8) Proposition] 4.1] that if q is locally Lipschitz continuous around \bar{x} relative to its domain and parabolically epi-differentiable at \bar{x} for $w \in T_{\text{dom}g}(\bar{x})$, then $\text{dom} d^2 g(\bar{x})(w \mid \cdot) = T_{\text{dom}g}^2(\bar{x}, w)$ and $\text{dom} g$ is parabolically derivable at \bar{x} for w. Below, we record an important relationship between the second subderivative and the parabolic subderivative of functions, used extensively in our paper (see [\[18,](#page-27-15) Proposition 13.64]).

Proposition 2.1. For $g: \mathbb{R}^n \to \mathbb{R} \cup {\{\pm \infty\}}$, any point $\bar{x} \in \mathbb{R}^n$ with $g(\bar{x})$ finite and any vector w with $dg(\bar{x})(w)$ finite, let v be such that $dg(\bar{x})(w) = \langle v, w \rangle$. Then

$$
d^2 g(\bar{x} \mid v)(w) \le \inf_{z} \{d^2 g(\bar{x})(w \mid z) - \langle v, z \rangle\}.
$$
 (2.1)

We say that a function $g: \mathbb{R}^n \to \mathbb{R} \cup {\pm \infty}$ is parabolically regular at a point \bar{x} for a vector v if $g(\bar{x})$ is finite and the inequality in [\(2.1\)](#page-3-0) holds with equality for every w satisfying $dg(\bar{x})(w) = \langle v, w \rangle$. The critical cone of a function $g: \mathbb{R}^n \to \mathbb{R} \cup \{\pm \infty\}$ at a point \bar{x} for a vector v is defined by

$$
K_g(\bar{x}, v) := \{w \in \mathbb{R}^n : dg(\bar{x})(w) = \langle v, w \rangle\}.
$$

2.2 Eigenvalues and Singular Values

Given an $m \times n$ matrix X and index sets $I \subset \{1, \ldots, m\}$, $J \subset \{1, \ldots, n\}$, denote by X_{ij} the (i, j) th entry of X and denote by X_{IJ} the submatrix of X obtained by removing all the rows of X not in I and all the columns of X not in J. The matrix X_I is the submatrix of X with columns specified by I, unless otherwise specified. For any vector $x \in \mathbb{R}^n$, let $diag(x)$ denote the matrix where $(\text{diag}(x))_{ii} = x_i$ for all i, and $(\text{diag}(x))_{ij} = 0$ for $i \neq j$. It's worth noting that $\text{diag}(x)$ may represent an $m \times n$, $n \times n$, or $m \times m$ matrix (the latter when $x \in \mathbb{R}^m$). However, the context will always clarify which dimension is applicable. Let \mathbb{S}^n be the space of all real $n \times n$ symmetric matrices. Let \mathcal{O}^n be the set of all $n \times n$ orthogonal matrices and $\mathcal{O}^{m,n}$ denote the Cartesian product $\mathcal{O}^m \times \mathcal{O}^n$. The induced Frobenius norm of $X \in \mathbb{M}_{m,n}$ is defined via the trace inner product by $||X|| = \sqrt{\text{tr}(X^TX)}.$

If $A \in \mathbb{S}^n$, then we can arrange its n real eigenvalues in the decreasing order:

$$
\lambda_1(A) \geq \lambda_2(A) \geq \cdots \geq \lambda_s(A) \geq \cdots \geq \lambda_n(A),
$$

where $\lambda_s(A)$ is the sth largest eigenvalue of A (counting multiplicity of each of them). For any $A \in \mathbb{S}^n$, there exists $U \in \mathcal{O}^n$ for which we have

$$
A = U(\text{diag}(\lambda(A)))U^T \text{ with } \lambda(A) = (\lambda_1(A), \dots, \lambda_n(A)).
$$
\n(2.2)

For a given matrix $A \in \mathbb{S}^n$, the set of such orthogonal matrices U in [\(2.2\)](#page-3-1) is denoted by $\mathcal{O}^n(A)$. We say that two matrices A and B in \mathbb{S}^n have a simultaneous ordered spectral decomposition if there exists $U \in \mathcal{O}^n$ such that $A = U(\text{diag}(\lambda(A)))U^T$ and $B = U(\text{diag}(\lambda(B)))U^T$. It is well-known that any two matrices A and B in \mathbb{S}^n satisfy the inequality

$$
\langle A, B \rangle \le \langle \lambda(A), \lambda(B) \rangle,\tag{2.3}
$$

which is known as Fan's inequality. Moreover, equality in (2.3) amounts to A and B admitting a simultaneous ordered spectral decomposition. We denote by l_s^{λ} the number of eigenvalues, ranking before s, which are equal to $\lambda_s(A)$ (including $\lambda_s(A)$) and j_s^{λ} the number of eigenvalues, ranking strictly after s, which are equal to $\lambda_s(A)$. According to [\[20,](#page-27-12) Proposition 1.4, Lemma 1.1], we have the following result.

Lemma 2.2. Let $A \in \mathbb{S}^n$ and $U = [u_1, \ldots, u_n] \in \mathcal{O}^n$ such that

$$
U^T A U = \text{diag}(\lambda_1(A), \dots, \lambda_n(A)).
$$

If we set

$$
U_s = [u_{s-l_s^{\lambda}+1}, \dots, u_{s+j_s^{\lambda}}],
$$

\n
$$
U_s^c = [u_1, \dots, u_{s-l_s^{\lambda}}, u_{s+j_s^{\lambda}+1}, \dots, u_n],
$$

then for a small perturbation matrix $E \in \mathbb{S}^n$,

$$
\lambda_s(A+E) = \lambda_s(A) + \lambda_{l_s^{\lambda}} \left(U_s^T E U_s + U_s^T E U_s^c (\lambda_s(A) I - \Lambda_s)^{-1} U_s^{cT} E U_s \right) + O(||E||^3),
$$

where $\Lambda_s = \text{diag}(\lambda_1(A), \ldots, \lambda_{s-l_s^{\lambda}}(A), \lambda_{s+j_s^{\lambda}+1}(A), \ldots, \lambda_n(A)).$

For any $X \in \mathbb{M}_{m,n}$ with rank r, there exists $(U, V) \in \mathcal{O}^{m,n}$ for which we have a singular value decomposition:

$$
X = U\Sigma(X)V^T,\t\t(2.4)
$$

where $\Sigma(X) := \text{diag}(\sigma(X)) \in \mathbb{M}_{m,n}$ is a diagonal matrix with singular values in descending order:

$$
\sigma_1(X) \geq \sigma_2(X) \geq \ldots \geq \sigma_r(X) > 0 = \sigma_{r+1}(X) = \ldots = \sigma_n(X)
$$

on the diagonal. For a given matrix $X \in M_{m,n}$, the set of such orthogonal matrices U and V in [\(2.4\)](#page-4-0) is denoted by $\mathcal{O}^{m,n}(X)$. We say that two matrices X and Y in $\mathbb{M}_{m,n}$ have a simultaneous ordered singular value decomposition if there exists $(U, V) \in \mathcal{O}^{m,n}$ such that $X = U\Sigma(X)V^T$ and $Y = U\Sigma(Y)V^T$. The next lemma, known as Von Neumann's trace theorem, shows precisely when two matrices X and Y admit simultaneous ordered singular value decomposition (see [\[10,](#page-27-5) Theorem 4.6]).

Lemma 2.3. Any matrices X and Y in $\mathbb{M}_{m,n}$ satisfy the inequality

$$
\langle X, Y \rangle \le \langle \sigma(X), \sigma(Y) \rangle. \tag{2.5}
$$

Equality holds if and only if X and Y have a simultaneous ordered singular value decomposition.

It is not hard to see that any two matrices X and Y in $\mathbb{M}_{m,n}$, the estimate

$$
\|\sigma(X) - \sigma(Y)\| \le \|X - Y\|
$$
\n(2.6)

always holds. Denote the three index sets α , β , β ₀ by

$$
\alpha := \{1, \dots, r\}, \beta := \{r+1, \dots, n\}, \text{ and } \beta_0 = \{n+1, \dots, m\}.
$$

Assume that $\mu_1 > \cdots > \mu_t > \mu_{t+1} = 0$ are the distinct singular values of $X \in M_{m,n}$, and define the index sets

$$
\alpha_i := \{ s \in \alpha \colon \sigma_s(X) = \mu_i \} \text{ for all } i = 1, \dots, t. \tag{2.7}
$$

Obviously, $\alpha = \bigcup_{i=1}^{t} \alpha_i$. Define $\hat{\beta} := \beta \cup \beta_0$. Partition $U \in \mathcal{O}^m$ as

$$
U = \begin{bmatrix} U_{\alpha_1} & U_{\alpha_2} & \cdots & U_{\alpha_t} & U_{\hat{\beta}} \end{bmatrix},
$$

where $U_{\alpha_i} \in \mathbb{R}^{m \times |\alpha_i|}$ for $i = 1, ..., t$, and $U_{\hat{\beta}} \in \mathbb{R}^{m \times |\hat{\beta}|}$. Similarly,

$$
V = \begin{bmatrix} V_{\alpha_1} & V_{\alpha_2} & \cdots & V_{\alpha_t} & V_{\alpha_{t+1}} \end{bmatrix} \in \mathcal{O}^n,
$$

where $V_{\alpha_i} \in \mathbb{R}^{n \times |\alpha_i|}$ for $i = 1, \ldots, t, t + 1$ with $\alpha_{t+1} := \beta$.

For a matrix $X \in M_{m,n}$, its nuclear norm is given by

$$
||X||_* := \sum_{i=1}^n \sigma_i(X).
$$

Denote $\mathbb{B} := \{X : ||X||_* \leq 1\}$ as the unit sublevel set of the nuclear norm, and then its polar is the closed convex set

$$
\mathbb{B}^o = \Big\{ Z \colon \sigma_1(Z) := \max_{1 \le i \le n} \sigma_i(Z) \le 1 \Big\}.
$$

From $[22, Example 2]$, we know that the subdifferential of the nuclear norm at X is

$$
\partial ||X||_* = \left\{ U \text{diag}(s) V^T : X = U \Sigma(X) V^T, s \in \partial ||\sigma(X)||_1 \right\}
$$

=
$$
\left\{ U \text{diag}(s) V^T : X = U \Sigma(X) V^T, s_i = 1, i \in \alpha; |s_i| \leq 1, i \in \beta \right\}
$$

=
$$
\left\{ U_{\alpha} V_{\alpha}^T + U_{\beta} Z V_{\beta}^T : X = \begin{bmatrix} U_{\alpha} & U_{\beta} \end{bmatrix} \Sigma(X) \begin{bmatrix} V_{\alpha} & V_{\beta} \end{bmatrix}^T, \sigma_1(Z) \leq 1 \right\},
$$

where $diag(s)$ is an $m \times n$ diagonal matrix with s on its diagonal and $Z \in \mathbb{R}^{(m-r)\times(n-r)}$.

Define the linear operator $\mathcal{B}(\cdot) : \mathbb{M}_{m,n} \to \mathbb{S}^{m+n}$ by

$$
\mathcal{B}(X) := \begin{bmatrix} 0 & X \\ X^T & 0 \end{bmatrix}, \ X \in \mathbb{M}_{m,n}.
$$

It follows from [\[7,](#page-27-16) Theorem 7.3.3] that

$$
P^T \mathcal{B}(X) P = \begin{bmatrix} \Sigma_n(X) & 0 & 0 \\ 0 & 0_{m-n} & 0 \\ 0 & 0 & -\Sigma_n(X) \end{bmatrix},
$$
 (2.8)

where the orthogonal matrix $P \in \mathcal{O}^{m+n}$ is given by

$$
P:=\frac{1}{\sqrt{2}}\begin{bmatrix}U_\alpha&U_\beta&\sqrt{2}U_{\beta_0}&-U_\alpha&-U_\beta\\V_\alpha&V_\beta&0&V_\alpha&V_\beta\end{bmatrix}.
$$

Obviously, for each $s \in \alpha \cup \beta$, we have that $\sigma_s(X) = \lambda_s(\mathcal{B}(X))$. Therefore, by applying the definition of directional derivative, we derive that $\sigma'_s(X;H) = \lambda'_s(\mathcal{B}(X); \mathcal{B}(H))$ for any $H \in M_{m,n}$.

Similar to the symmetric case, for matrix $X_0 \in M_{m,n}$, we denote by $l_s(X_0)$ the number of singular values, ranking before s, which are equal to $\sigma_s(X_0)$ (including $\sigma_s(X_0)$) and $j_s(X_0)$ the number of singular values, ranking strictly after s, which are equal to $\sigma_s(X_0)$, respectively, i.e., we define $l_s(X_0)$ and $j_s(X_0)$ such that

$$
\sigma_1(X_0) \geq \cdots \geq \sigma_{s-l_s(X_0)}(X_0) > \sigma_{s-l_s(X_0)+1}(X_0) = \cdots = \sigma_s(X_0) = \cdots = \sigma_{s+j_s(X_0)}(X_0) > \sigma_{s+j_s(X_0)+1}(X_0) \geq \cdots \geq \sigma_n(X_0).
$$

We use $r_s(X_0)$ to denote the multiplicity of $\sigma_s(X_0)$. In later discussions, when the dependence of l_s , j_s and r_s , $s \in \alpha \cup \beta$, on X_0 can be seen clearly from the context, we often drop X_0 from these notations. The following proposition on the directional derivatives of the singular value of a matrix was explored in $[11, Section 5.1]$.

Proposition 2.4. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r be given and have the singular value decomposition [\(2.4\)](#page-4-0). Let $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. Then for each $s \in \alpha_i, i = 1, \ldots, t$,

$$
\sigma'_s(X_0; H) = \frac{1}{2}\lambda_{l_s} \left(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \right).
$$

For each $s \in \beta$, one has that

$$
\sigma'_s(X_0; H) = \sigma_{l_s}\big(U_{\hat{\beta}}^T H V_{\beta}\big).
$$

Assume that $\eta_1^i > \cdots > \eta_{N_i}^i$ are the distinct eigenvalues of $\frac{1}{2} (U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i}),$ and define the index sets

$$
\beta_j^i := \left\{ s \in \alpha_i \colon \lambda_s \left(\frac{1}{2} \left(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \right) \right) = \eta_j^i \right\} \text{ for all } i = 1, \dots, t, j = 1, \dots, N_i. \tag{2.9}
$$

Denote the distinct singular values of $U_{\hat{\beta}}^T H V_{\beta}$ by $\eta_1^{t+1} > \cdots > \eta_{N_{t+1}}^{t+1} > \eta_{N_{t+1}+1}^{t+1} = 0$, and define

$$
\beta_k^{t+1} := \left\{ s \in \beta \colon \sigma_s \left(U_{\hat{\beta}}^T H V_{\beta} \right) = \eta_k^{t+1} \right\} \text{ for all } k = 1, \dots, N_{t+1}, N_{t+1} + 1. \tag{2.10}
$$

For each $s \in \{1, \ldots, n\}$, there exists $i \in \{1, \ldots, t, t+1\}$ such that $s \in \alpha_i$ and $l_s(X_0) \in \{1, \ldots, |\alpha_i|\}$. Furthermore, we can find $j \in \{1, ..., N_i, N_{t+1} + 1\}$ such that $l_s(X_0) \in \beta_j^i$. Define now the integer $l_s(X_0, H)$ by

$$
\tilde{l}_s(X_0, H) := \begin{cases} l_{l_s(X_0)} \Big(\frac{1}{2} \big(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \big) \Big) & \text{if } s \in \alpha, \\ l_{l_s(X_0)} \big(U_{\hat{\beta}}^T H V_{\beta} \big) & \text{if } s \in \beta, \end{cases}
$$

which, in fact, signifies the number of eigenvalues of $\frac{1}{2} (U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i})$ that are equal to $\lambda_{l_s(X_0)}\Big(\frac{1}{2}\big(U_{\alpha_i}^THV_{\alpha_i}+V_{\alpha_i}^TH^TU_{\alpha_i}\big)\Big)$, but are ranked before $\lambda_{l_s(X_0)}\Big(\frac{1}{2}\big(U_{\alpha_i}^THV_{\alpha_i}+V_{\alpha_i}^TH^TU_{\alpha_i}\big)\Big)$, including $\lambda_{l_s(X_0)}\Big(\frac{1}{2}\big(U_{\alpha_i}^THV_{\alpha_i}+V_{\alpha_i}^TH^TU_{\alpha_i}\big)\Big)$ for $s \in \alpha$, and the number of singular values of

 $U_{\hat{\beta}}^T H V_{\beta}$ that are equal to $\sigma_{l_s(X_0)}(U_{\hat{\beta}}^T H V_{\beta})$, but are ranked before $\sigma_{l_s(X_0)}(U_{\hat{\beta}}^T H V_{\beta})$, including $\sigma_{l_s(X_0)}(U_{\hat{\beta}}^THV_{\beta})$ for $s \in \beta$. As before, we often drop X_0 and H from $\tilde{l}_s(X_0, H)$ when the dependence of \tilde{l}_s on X_0 and H can be seen clearly from the context.

Similarly, [\[23,](#page-28-1) Theorem 3.1] derives the following explicit formulas of the (parabolic) second order directional derivatives of the singular values.

Proposition 2.5. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r be given and have the singular value decomposition [\(2.4\)](#page-4-0). Suppose that the direction $H, W \in M_{m,n}$. Then the following assertions hold:

(i) For each $s \in \alpha$, there exists $i \in \{1, ..., t\}$ and $j \in \{1, ..., N_i\}$ such that $s \in \alpha_i$ and $l_s(X_0) \in \beta_j^i$, and there exists $Q^i \in \mathcal{O}^{|\alpha_i|} \Big(\frac{1}{2} \big(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \big) \Big)$ such that

$$
\sigma_s''(X_0; H, W) = \lambda_{\tilde{l}_s} \left(\left(Q^i_{\beta^i_j} \right)^T P^T_{\alpha_i} \left[\mathcal{B}(W) - 2\mathcal{B}(H) P^c_{\alpha_i} (A_s - \mu_i I)^{-1} P^{c \, T}_{\alpha_i} \mathcal{B}(H) \right] P_{\alpha_i} Q^i_{\beta^i_j} \right), \quad (2.11)
$$

where $\Lambda_s \in \mathbb{R}^{(m+n-r_s)\times(m+n-r_s)}$ is a diagonal matrix whose diagonal elements are the eigenvalues of $P^T \mathcal{B}(X_0) P$ that are not equal to $\sigma_s(X_0)$.

(ii) For each $s \in \beta$, if $l_s(X_0) \in \beta_k^{t+1}$ for some $k \in \{1, ..., N_{t+1}\}$, then there exists $(Q_{\hat{\beta}\hat{\beta}}, \widehat{Q}_{\beta\beta}) \in$ $\mathcal{O}^{|\hat{\beta}|,|\beta|}\big(U_{\hat{\beta}}^THV_{\beta}\big)$ such that

$$
\sigma_s''(X_0; H, W) = \frac{1}{2} \lambda_{\tilde{l}_s} \left(\left(Q_{\hat{\beta}\beta_k^{t+1}} \right)^T \left[U_{\hat{\beta}}^T W V_{\beta} - 2U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H V_{\beta} \right] \hat{Q}_{\beta \beta_k^{t+1}} + \left(\hat{Q}_{\beta \beta_k^{t+1}} \right)^T \left[U_{\hat{\beta}}^T W V_{\beta} - 2U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H V_{\beta} \right]^T Q_{\hat{\beta}\beta_k^{t+1}} \right), \tag{2.12}
$$

and if $l_s(X_0) \in \beta^{t+1}_{N_{t+1}+1}$, then there exists $(Q_{\hat{\beta}\hat{\beta}}, \widehat{Q}_{\beta\beta}) \in \mathcal{O}^{|\hat{\beta}|, |\beta|}(U_{\hat{\beta}}^T H V_{\beta})$ such that

$$
\sigma_s''(X_0; H, W) = \sigma_{\tilde{l}_s} \left(\begin{bmatrix} Q_{\hat{\beta}\beta_{N_{t+1}+1}}^{t+1} & Q_{\hat{\beta}\beta_0} \end{bmatrix}^T \begin{bmatrix} U_{\hat{\beta}}^T W V_{\beta} - 2U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \end{bmatrix} \hat{Q}_{\beta\beta_{N_{t+1}+1}^{t+1}} \right). \tag{2.13}
$$

Combining Proposition [2.4](#page-5-0) with Proposition [2.5,](#page-6-0) we obtain the following result, which is important for our development in this paper.

Corollary 2.6. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r be given and have the singular value decomposition [\(2.4\)](#page-4-0). Suppose that the direction $H, W \in \mathbb{M}_{m,n}$. Then for any $t > 0$ sufficiently small, we have

$$
\sigma\left(X_0 + tH + \frac{1}{2}t^2W + o(t^2)\right) = \sigma(X_0) + t\sigma'(X_0; H) + \frac{1}{2}t^2\sigma''(X_0; H, W) + o(t^2). \tag{2.14}
$$

3 Twice Epi-Differentiability of the Nuclear Norm

In this section, we study the first- and second-order epi-differentiability of the nuclear norm. The subsequent proposition is a direct consequence of Lemma [2.2.](#page-3-3)

Proposition 3.1. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r be given and have the singular value decomposition [\(2.4\)](#page-4-0). Let $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. Then for $s \in \alpha \cup \beta$, we have

$$
\sigma_s(X_0+H) = \sigma_s(X_0) + \lambda_{l_s}\left(P_s^T\mathcal{B}(H)P_s + P_s^T\mathcal{B}(H)P_s^c(\sigma_s(X_0)I - A_s)^{-1}P_s^{cT}\mathcal{B}(H)P_s\right) + O\left(\|\mathcal{B}(H)\|^3\right),
$$

where the columns of P_s form an orthonormal basis of eigenvectors of $\mathcal{B}(X_0)$ associated with $\sigma_s(X_0)$, P_s^c is the submatrix of P obtained by removing all the columns of P_s , and $\Lambda_s \in \mathbb{R}^{(m+n-r_s)\times (m+n-r_s)}$ is a diagonal matrix whose diagonal elements are the eigenvalues of $P^T\mathcal{B}(X_0)P$ that are not equal to $\sigma_s(X_0)$.

Moreover, for any $\tau > 0$ we have the following results:

(i) For any $s \in \alpha$ there exists $i \in \{1, \ldots, t\}$ and $j \in \{1, \ldots, N_i\}$ such that $s \in \alpha_i$ and $l_s(X_0) \in$ β_j^i , and we have

$$
\sigma_s(X_0 + \tau H) = \sigma_s(X_0) + \tau \lambda_{l_s} \left(P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i} \right) +
$$

$$
\frac{\tau^2}{2} \lambda_{\tilde{l}_s} \left(2 \left(Q_{\beta_j^i}^i \right)^T P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c (\sigma_s(X_0)I - A_s)^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H) P_{\alpha_i} Q_{\beta_j^i}^i \right) + O(\tau^3),
$$

(3.1)

where $Q^i \in \mathcal{O}^{|\alpha_i|} \left(P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i} \right)$. (ii) If $n > r$, then for $s \in \beta$ we have

$$
\sigma_s(X_0+H)=\sigma_s(X_0)+\sigma_{l_s}\big(U_{\hat{\beta}}^THV_{\beta}-U_{\hat{\beta}}^THV_{\alpha}\Sigma_{\alpha}^{-1}(X_0)U_{\alpha}^THV_{\beta}\big)+O(\|\mathcal{B}(H)\|^3).
$$

Furthermore, if $l_s(X_0) \in \beta_k^{t+1}$ for some $k \in \{1, ..., N_{t+1}\}$, then there exists $(Q_{\hat{\beta}\hat{\beta}}, \hat{Q}_{\beta\beta}) \in$ $\mathcal{O}^{|\hat{\beta}|,|\beta|}\big(U_{\hat{\beta}}^THV_{\beta}\big)$ such that

$$
\sigma_s(X_0 + \tau H) = \sigma_s(X_0) + \tau \sigma_{l_s} \left(U_{\hat{\beta}}^T H V_{\beta} \right) + \frac{\tau^2}{4} \lambda_{\tilde{l}_s} \left(\left(Q_{\hat{\beta}\beta_k^{t+1}} \right)^T \left[-U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right] \hat{Q}_{\beta \beta_k^{t+1}} + \left(\hat{Q}_{\beta \beta_k^{t+1}} \right)^T \left[-U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right]^T Q_{\hat{\beta}\beta_k^{t+1}} \right) + O(\tau^3),
$$

and if $l_s(X_0) \in \beta^{t+1}_{N_{t+1}+1}$, then there exists $(Q_{\hat{\beta}\hat{\beta}}, \widehat{Q}_{\beta\beta}) \in \mathcal{O}^{|\hat{\beta}|, |\beta|}(U_{\hat{\beta}}^T H V_{\beta})$ such that

$$
\sigma_s(X_0 + \tau H) = \sigma_s(X_0) + \tau \sigma_{l_s} (U_{\hat{\beta}}^T H V_{\beta})
$$

+
$$
\frac{\tau^2}{2} \sigma_{\tilde{l}_s} \left(\left[Q_{\hat{\beta} \beta^{t+1}_{N_{t+1}+1}} Q_{\hat{\beta} \beta_0} \right]^T \left[-U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right] \widehat{Q}_{\beta \beta^{t+1}_{N_{t+1}+1}} \right) + O(\tau^3).
$$

Proof It follows from Lemma [2.2](#page-3-3) and (2.8) that

$$
\sigma_s(X_0 + H) = \lambda_s(\mathcal{B}(X_0 + H)) = \lambda_s(\mathcal{B}(X_0) + \mathcal{B}(H))
$$

= $\lambda_s(\mathcal{B}(X_0)) + \lambda_{l_s}(P_s^T\mathcal{B}(H)P_s + P_s^T\mathcal{B}(H)P_s^c(\lambda_s(\mathcal{B}(X_0)))I - A_s)^{-1}P_s^{cT}\mathcal{B}(H)P_s) + O(||\mathcal{B}(H)||^3)$
= $\sigma_s(X_0) + \lambda_{l_s}(P_s^T\mathcal{B}(H)P_s + P_s^T\mathcal{B}(H)P_s^c(\sigma_s(X_0)I - A_s)^{-1}P_s^{cT}\mathcal{B}(H)P_s) + O(||\mathcal{B}(H)||^3).$

For any $s \in \alpha$, there exists $i \in \{1, ..., t\}$ and $j \in \{1, ..., N_i\}$ such that $s \in \alpha_i$ and $l_s(X_0) \in \beta_j^i$. By using Lemma [2.2](#page-3-3) with $A = P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}$ and $E = P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c (\sigma_s(X_0)I - A_s)^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H) P_{\alpha_i}$, we have

$$
\sigma_{s}(X_{0} + \tau H) = \sigma_{s}(X_{0}) + \lambda_{l_{s}}(\tau P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}} + \tau^{2} P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\sigma_{s}(X_{0})I - A_{s})^{-1} P_{\alpha_{i}}^{c} \mathcal{B}(H) P_{\alpha_{i}}) + O(\tau^{3})
$$
\n
$$
= \sigma_{s}(X_{0}) + \tau \lambda_{l_{s}}(P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}} + \tau P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\sigma_{s}(X_{0})I - A_{s})^{-1} P_{\alpha_{i}}^{c} \mathcal{B}(H) P_{\alpha_{i}}) + O(\tau^{3})
$$
\n
$$
= \sigma_{s}(X_{0}) + \tau \left[\lambda_{l_{s}}(P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}) + \right.
$$
\n
$$
\tau \lambda_{\tilde{l}_{s}}\left(\left(Q_{\beta_{j}^{i}}^{i}\right)^{T} P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\sigma_{s}(X_{0})I - A_{s})^{-1} P_{\alpha_{i}}^{c} \mathcal{B}(H) P_{\alpha_{i}} Q_{\beta_{j}^{i}}^{i} \right) + O(\tau^{3})
$$
\n
$$
= \sigma_{s}(X_{0}) + \tau \lambda_{l_{s}}(P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}) +
$$
\n
$$
\frac{\tau^{2}}{2} \lambda_{\tilde{l}_{s}}\left(2\left(Q_{\beta_{j}^{i}}^{i}\right)^{T} P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\sigma_{s}(X_{0})I - A_{s})^{-1} P_{\alpha_{i}}^{c} \mathcal{B}(H) P_{\alpha_{i}} Q_{\beta_{j}^{i}}^{i} \right) + O(\tau^{3}).
$$

In particular, we have

$$
\sigma'_s(X_0;H) = \lambda_{l_s}\big(P_{\alpha_i}^T \mathcal{B}(H)P_{\alpha_i}\big) = \frac{1}{2}\lambda_{l_s}\big(U_{\alpha_i}^T HV_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i}\big).
$$

If $n > r$, then $\beta \neq \emptyset$. For any $s \in \beta$, we have $\sigma_s(X_0) = 0$,

$$
P_s = \frac{1}{\sqrt{2}} \begin{bmatrix} U_\beta & \sqrt{2}U_{\beta_0} & -U_\beta \\ V_\beta & 0 & V_\beta \end{bmatrix}, \text{ and } \Lambda_s = \begin{bmatrix} \Sigma_\alpha(X_0) & 0 \\ 0 & -\Sigma_\alpha(X_0) \end{bmatrix}.
$$

By some elementary calculations, we can obtain that

$$
P_s^T \mathcal{B}(H) P_s = S \mathcal{B}(V_\beta^T H^T U_{\hat{\beta}}) S^T,
$$

$$
P_s^T \mathcal{B}(H) P_s^c (\sigma_s(X_0)I - A_s)^{-1} P_s^{cT} \mathcal{B}(H) P_s = -S \mathcal{B}(V_\beta^T H^T U_\alpha \Sigma_\alpha^{-1}(X_0) V_\alpha^T H^T U_{\hat{\beta}}) S^T,
$$

where

$$
S = \frac{1}{2} \begin{bmatrix} \sqrt{2}I_{|\beta|} & 0 & \sqrt{2}I_{|\beta|} \\ 0 & 2I_{|\beta_0|} & 0 \\ \sqrt{2}I_{|\beta|} & 0 & -\sqrt{2}I_{|\beta|} \end{bmatrix} \in \mathcal{O}^{2|\beta|+|\beta_0|}.
$$

Therefore, we have

$$
\lambda_{l_s} (P_s^T \mathcal{B}(H) P_s + P_s^T \mathcal{B}(H) P_s^c (\sigma_s(X_0)I - A_s)^{-1} P_s^{cT} \mathcal{B}(H) P_s)
$$
\n
$$
= \lambda_{l_s} \left(S \mathcal{B} (V_\beta^T H^T U_{\hat{\beta}}) S^T - S \mathcal{B} (V_\beta^T H^T U_\alpha \Sigma_\alpha^{-1}(X_0) V_\alpha^T H^T U_{\hat{\beta}}) S^T \right)
$$
\n
$$
= \lambda_{l_s} \left(S \mathcal{B} (V_\beta^T H^T U_{\hat{\beta}} - V_\beta^T H^T U_\alpha \Sigma_\alpha^{-1}(X_0) V_\alpha^T H^T U_{\hat{\beta}}) S^T \right)
$$
\n
$$
= \lambda_{l_s} \left(\mathcal{B} (V_\beta^T H^T U_{\hat{\beta}} - V_\beta^T H^T U_\alpha \Sigma_\alpha^{-1}(X_0) V_\alpha^T H^T U_{\hat{\beta}}) \right)
$$
\n
$$
= \sigma_{l_s} \left(V_\beta^T H^T U_{\hat{\beta}} - V_\beta^T H^T U_\alpha \Sigma_\alpha^{-1}(X_0) V_\alpha^T H^T U_{\hat{\beta}} \right)
$$
\n
$$
= \sigma_{l_s} \left(U_\beta^T H V_\beta - U_\beta^T H V_\alpha \Sigma_\alpha^{-1}(X_0) U_\alpha^T H V_\beta \right),
$$

and then $\sigma_s(X_0+H) = \sigma_s(X_0) + \sigma_{l_s}(U_{\hat{\beta}}^THV_{\beta}) + O(||\mathcal{B}(H)||^2)$. In particular, we have $\sigma'_s(X_0; H) =$ $\sigma_{l_s}(U_{\hat{\beta}}^THV_{\beta})$. Similar to the discussions of first-order directional derivatives of singular values, we can obtain their second-order directional derivatives. \Box

Define the sum of all the singular values that are equal to $\sigma_s(X_0)$ but are ranked before $\sigma_s(X_0)$ by

$$
\Psi_s(X_0)=\sigma_{s-l_s+1}(X_0)+\cdots+\sigma_s(X_0),
$$

and the sum of the first s-largest singular values of X_0 by

$$
\Phi_s(X_0)=\sigma_1(X_0)+\cdots+\sigma_s(X_0).
$$

Lemma 3.2. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r satisfy $n > r$ and $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. Then, we have

$$
\Psi_n(X_0 + H) = \sigma_{\mathbb{B}^o} \left(U_{\hat{\beta}}^T H V_{\beta} - U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right) + O\big(\| \mathcal{B}(H) \|^3 \big), \tag{3.2}
$$

where $\sigma_{\mathbb{B}^o}$ is the support function of the set $\mathbb{B}^o \subset \mathbb{R}^{|\beta| \times |\beta|}$. As a consequence,

$$
\Psi_n(X_0 + H) = \sigma_{\mathbb{B}^\circ} \left(U_\beta^T H V_\beta \right) + O\big(\| \mathcal{B}(H) \|^2 \big). \tag{3.3}
$$

 \Box

Proof From Proposition [3.1,](#page-6-1) we obtain

$$
\Psi_n(X_0 + H) = \sigma_{r+1}(X_0 + H) + \cdots + \sigma_n(X_0 + H)
$$
\n
$$
= \sigma_{r+1}(X_0) + \sigma_1 \left(U_{\hat{\beta}}^T H V_{\beta} - U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right) + \cdots +
$$
\n
$$
\sigma_n(X_0) + \sigma_{l_n} \left(U_{\hat{\beta}}^T H V_{\beta} - U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right) + O\left(\| \mathcal{B}(H) \|^3 \right)
$$
\n
$$
= (\sigma_1 + \cdots + \sigma_{l_n}) \left(U_{\hat{\beta}}^T H V_{\beta} - U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right) + O\left(\| \mathcal{B}(H) \|^3 \right)
$$
\n
$$
= \sigma_{\mathbb{B}^\circ} \left(U_{\hat{\beta}}^T H V_{\beta} - U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \right) + O\left(\| \mathcal{B}(H) \|^3 \right).
$$

Lemma 3.3. The sum of the first r-largest singular values Φ_r is \mathcal{C}^2 on the set of $m \times n$ matrices of rank r. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r be given and $H \in \mathbb{M}_{m,n}$ be a matrix, the second-order differential of Φ_r at X_0 in the direction H is given by

$$
D^{2}\Phi_{r}(X_{0})(H,H) = \sum_{k_{l} \leq r} \text{tr}\Big(2P_{k_{l}}^{T}\mathcal{B}(H)P_{k_{l}}^{c}(\sigma_{k_{l}}(X_{0})I - A_{k_{l}})^{-1}P_{k_{l}}^{c}^{T}\mathcal{B}(H)P_{k_{l}}\Big).
$$

Proof It follows from [\[5,](#page-27-11) Proposition 8] that Φ_r is \mathcal{C}^2 on the set of $m \times n$ matrices of rank r. For any $s \in \alpha$, it follows from [\(3.1\)](#page-6-2) that the second-order directional derivative of σ_s at X_0 in the direction H is

$$
\sigma_s''(X_0; H) = \lim_{\tau \downarrow 0} \frac{\sigma_s(X_0 + \tau H) - \sigma_s(X_0) - \tau \sigma_s'(X_0; H)}{\frac{\tau^2}{2}}
$$

=
$$
\lambda_{\tilde{l}_s} \left(2Q_{l_s}^T P_s^T \mathcal{B}(H) P_s^c(\sigma_s(X_0)I - \Lambda_s)^{-1} P_s^{cT} \mathcal{B}(H) P_s Q_{l_s} \right).
$$

Denote by k_1, \ldots, k_q the set of the indices of all the leading singular values of X_0 , i.e.

$$
\sigma_{k_1}(X_0) > \sigma_{k_2}(X_0) > \cdots > \sigma_{k_q}(X_0).
$$

Assume that there exists $k_l \leq r$ such that

$$
\sigma_{k_l-1}(X_0) > \sigma_{k_l}(X_0) = \cdots = \sigma_{k_l+j_{k_l}}(X_0) > \sigma_{k_{l+1}}(X_0).
$$

Set $G_{k_l} := P_{k_l}^T \mathcal{B}(H) P_{k_l}^c(\sigma_{k_l}(X_0)I - A_{k_l})^{-1} P_{k_l}^{c}{}^T \mathcal{B}(H) P_{k_l}$. Then for any $q \in \{k_l, k_l + 1, \ldots, k_l + j_{k_l}\},$ we have $\sqrt{2}$ T $\overline{ }$ $\sqrt{2}$ $\mathcal I$ $\overline{ }$

$$
\left(2Q_{l_q}^T G_{k_l} Q_{l_q}\right) v_{l_q} = \lambda_{\tilde{l}_q} \left(2Q_{l_q}^T G_{k_l} Q_{l_q}\right) v_{l_q},
$$

where v_{l_q} is a unit eigenvector of $2Q_{l_q}^T G_{k_l} Q_{l_q}$ associated with eigenvalue $\lambda_{\tilde{l}_q} (2Q_{l_q}^T G_{k_l} Q_{l_q})$. Therefore, by some elementary calculations, we obtain that

$$
D^{2}\Phi_{r}(X_{0})(H,H) = \sum_{k_{l}\leq r} (\sigma_{k_{l}} + \sigma_{k_{l}+1} + \ldots + \sigma_{k_{l}+j_{k_{l}}})''(X_{0};H)
$$

=
$$
\sum_{k_{l}\leq r} \lambda_{\tilde{l}_{k_{l}}} \left(2Q_{l_{k_{l}}}^{T} G_{k_{l}} Q_{l_{k_{l}}}\right) + \lambda_{\tilde{l}_{k_{l}+1}} \left(2Q_{l_{k_{l}+1}}^{T} G_{k_{l}} Q_{l_{k_{l}+1}}\right) + \cdots + \lambda_{\tilde{l}_{k_{l}+j_{k_{l}}}} \left(2Q_{l_{k_{l}+j_{k_{l}}}}^{T} G_{k_{l}} Q_{l_{k_{l}+j_{k_{l}}}}\right)
$$

=
$$
\sum_{k_{l}\leq r} tr\left(2Q^{T} G_{k_{l}} Q\right) = \sum_{k_{l}\leq r} tr\left(2G_{k_{l}} \right),
$$

and the proof is complete.

Recall from Rockafellar and Wets [\[18,](#page-27-15) Definition 7.20] that a function $g: \mathbb{R}^n \to \mathbb{R} \cup {\pm \infty}$ is said to be semidifferentiable at $\bar{x} \in \mathbb{R}^n$ with $g(\bar{x})$ finite for $w \in \mathbb{R}^n$, if the (possibly infinite) limit

 \Box

 \Box

$$
\lim_{\substack{t \downarrow 0 \\ w' \to w}} \frac{g(\bar{x} + tw') - g(\bar{x})}{t}
$$

exists. This limit is the semiderivative of g at \bar{x} for w.

Lemma 3.4. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r satisfy $n > r$ and $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. The function Ψ_n is semidifferentiable at X_0 for H and its semiderivative coincides with subderivative, i.e. $\Psi'_n(X_0; H) = d\Psi_n(X_0)(H)$. The regular subdifferential of Ψ_n at X_0 is given by

$$
\hat{\partial}\Psi_n(X_0) = \left\{ U_{\hat{\beta}} Z V_{\beta}^T \colon X_0 = \begin{bmatrix} U_{\alpha} & U_{\hat{\beta}} \end{bmatrix} \Sigma(X_0) \begin{bmatrix} V_{\alpha} & V_{\beta} \end{bmatrix}^T, Z \in \mathbb{B}^o \subset \mathbb{R}^{|\hat{\beta}| \times |\beta|} \right\}.
$$

Proof Note that $\Psi_n(X_0) = 0$ and $\sigma_{\mathbb{B}^{\circ}}(U_{\hat{\beta}}^T H V_{\beta})$ is continuous in H. From [\(3.3\)](#page-8-0), we have

$$
\Psi'_n(X_0;H) = \lim_{\substack{\tau \downarrow 0 \\ H' \to H}} \frac{\Psi_n(X_0 + \tau H') - \Psi_n(X_0)}{\tau} = \sigma_{\mathbb{B}^o} (U_{\hat{\beta}}^T H V_{\beta}).
$$

Then Ψ_n is semidifferentiable at X_0 for H and also epi-differentiable. Furthermore, we have $\Psi'_n(X_0; H) = d\Psi_n(X_0)(H)$ and

$$
\hat{\partial}\Psi_n(X_0) = \left\{ U_{\hat{\beta}} Z V_{\beta}^T : X_0 = \begin{bmatrix} U_{\alpha} & U_{\hat{\beta}} \end{bmatrix} \Sigma(X_0) \begin{bmatrix} V_{\alpha} & V_{\beta} \end{bmatrix}^T, \sigma_1(Z) \le 1 \right\},
$$

$$
= \left\{ U_{\hat{\beta}} Z V_{\beta}^T : X_0 = \begin{bmatrix} U_{\alpha} & U_{\hat{\beta}} \end{bmatrix} \Sigma(X_0) \begin{bmatrix} V_{\alpha} & V_{\beta} \end{bmatrix}^T, Z \in \mathbb{B}^o \subset \mathbb{R}^{|\hat{\beta}| \times |\beta|} \right\}.
$$

Next, we present the main conclusion of this section. From this, the second-order epi-derivative of the nuclear norm can be derived.

Theorem 3.5. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r satisfy $n > r$ and $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. The function Ψ_n is twice epi-differentiable at X_0 . The second-order epi-derivative of Ψ_n at X_0 relative to any $\Omega \in \partial \Psi_n(X_0)$ is given explicitly by

$$
d^2\Psi_n(X_0 \mid \Omega)(H) = \begin{cases} -2\langle \Omega, HV_\alpha \Sigma_\alpha^{-1} U_\alpha^T H \rangle & \text{if } \Psi_n'(X_0; H) = \langle \Omega, H \rangle, \\ +\infty & \text{if } \Psi_n'(X_0; H) > \langle \Omega, H \rangle. \end{cases}
$$

Proof Let $\Omega \in \hat{\partial}\Psi_n(X_0)$. Given $X_0, H \in \mathbb{M}_{m,n}$, $H_k \to H$, and $t_k \to 0^+$, we focus our attention on the second-order difference quotient

$$
\frac{\Psi_n(X_0 + t_k H_k) - \Psi_n(X_0) - t_k \langle \Omega, H_k \rangle}{\frac{1}{2}t_k^2}.
$$
\n(3.4)

To begin with, assume that $\Psi'_n(X_0; H) > \langle \Omega, H \rangle$. By using the relation [\(3.3\)](#page-8-0) with $H = t_k H_k$, [\(3.4\)](#page-10-0) becomes

$$
\frac{\Psi_n'(X_0; H_k) - \langle \Omega, H_k \rangle}{\frac{1}{2}t_k} + O(||\mathcal{B}(H_k)||^2),
$$

which goes to $+\infty$ as $k \to +\infty$.

Assume now that $\Psi'_n(X_0; H) = \langle \Omega, H \rangle$. Let $H_k \to H$ and $t_k \to 0^+$. Since $\Omega \in \hat{\partial} \Psi_n(X_0)$, it is possible to write $\Omega = U_{\hat{\beta}} Z V_{\beta}^T$ for some $Z \in \mathbb{B}^o$. By using the development [\(3.2\)](#page-8-1), the difference quotient [\(3.4\)](#page-10-0) can be written as

$$
\sigma_{\mathbb{B}^o} \left(U_{\hat{\beta}}^T H_k V_{\beta} - t_k U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k V_{\beta} \right) - \left\langle Z, U_{\hat{\beta}}^T H_k V_{\beta} \right\rangle + O(t_k). \tag{3.5}
$$

Moreover, since $Z \in \mathbb{B}^o$, it is possible to bound from below the previous quantity by

$$
\frac{\left\langle Z, U_{\hat{\beta}}^T H_k V_{\beta} - t_k U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k V_{\beta} \right\rangle - \left\langle Z, U_{\hat{\beta}}^T H_k V_{\beta} \right\rangle}{\frac{1}{2} t_k} + O(t_k)
$$

= $- 2 \left\langle Z, U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k V_{\beta} \right\rangle + O(t_k)$
= $- 2 \left\langle U_{\hat{\beta}} Z V_{\beta}^T, H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k \right\rangle + O(t_k)$

which converges to $-2\langle \Omega, HV_{\alpha} \Sigma_{\alpha}^{-1}(X_0)U_{\alpha}^TH \rangle$.

Let $t_k \to 0^+$. We have to exhibit a sequence $\{H_k\}_k$ which converges to H such that the limit of [\(3.4\)](#page-10-0) equals $-2\langle \Omega, HV_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H \rangle$. Let us consider $H_k = H + t_k H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H_k$. The difference quotient (3.4) , which is equal to (3.5) , therefore becomes

$$
\frac{\sigma_{\mathbb{B}^o} (U_{\hat{\beta}}^T H V_{\beta}) - \langle \Omega, H \rangle - t_k \langle \Omega, H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k \rangle}{\frac{1}{2} t_k} + O(t_k)
$$

=
$$
\frac{\Psi_n'(X_0; H) - \langle \Omega, H \rangle}{\frac{1}{2} t_k} - 2 \langle \Omega, H_k V_{\alpha} \Sigma_{\alpha}^{-1} (X_0) U_{\alpha}^T H_k \rangle + O(t_k),
$$

which converges to $-2\langle \Omega, HV_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H \rangle$, as required.

This, combined with Lemma [3.3](#page-8-2) and [\[17,](#page-27-17) Proposition 2.10], implies the following corollary.

Corollary 3.6. Let $X_0 \in \mathbb{M}_{m,n}$ of rank r satisfy $n > r$ and $H \in \mathbb{M}_{m,n}$ be a small perturbation matrix. The nuclear norm is twice epi-differentiable at X_0 . The second-order epi-derivative of the nuclear norm at X_0 relative to any $\Omega \in \partial \|X_0\|_*$ is given explicitly by

$$
d^{2} \|\cdot\|_{*}(X_{0} \mid \Omega)(H) = D^{2} \Phi_{r}(X_{0})(H, H) + d^{2} \Psi_{n}(X_{0} \mid \Omega)(H).
$$

4 Subderivatives of Orthogonally Invariant Matrix Functions

In this section, we present a chain rule for the subderivative of orthogonally invariant matrix functions in Theorem [4.3.](#page-12-0) Similarly, we also give that of absolutely symmetric function in Theorem [4.8.](#page-16-0) These important results are central to our developments in this paper. Recall that Q_{\pm} is a signed permutation matrix if all its components are either 0 or ± 1 and each row and each column has exactly one nonzero element. Let \mathbf{P}_{\pm}^n denote the set of all $n \times n$ signed permutation matrices, and \mathbf{P}^n denote the set of all $n \times n$ permutation matrices. Recall also that a function $f: \mathbb{R}^n \to [-\infty, +\infty]$ is called absolutely symmetric if for every $x \in \mathbb{R}^n$ and every $n \times n$ signed permutation matrix Q_{\pm} , we have $f(Q_{\pm}x) = f(x)$. It is well-known (cf. Lewis [\[10,](#page-27-5) Proposition 5.1) that for any orthogonally invariant matrix function $F: \mathbb{M}_{m,n} \to [-\infty, +\infty]$, there exists an absolutely symmetric function $f: \mathbb{R}^n \to [-\infty, +\infty]$ satisfying

$$
F = f \circ \sigma. \tag{4.1}
$$

 \Box

Indeed, f can be chosen as a composite function of F and the linear mapping $x \mapsto diag(x)$ with $x \in \mathbb{R}^n$, namely,

$$
f = F \circ \text{diag.} \tag{4.2}
$$

A set $\Gamma \subset M_{m,n}$ is called an orthogonally invariant matrix set if δ_{Γ} is an orthogonally invariant matrix function. Similarly, $\Delta \subset \mathbb{R}^n$ is called an absolutely symmetric set if δ_{Δ} is an absolutely symmetric function. Therefore, it is easy to see that for any orthogonally invariant matrix set $\Gamma \subset \mathbb{M}_{m,n},$ there exists an absolutely symmetric set $\Delta \subset \mathbb{R}^n$ such that

$$
\Gamma = \{ X \in \mathbb{M}_{m,n} \colon \sigma(X) \in \Delta \},\tag{4.3}
$$

where Δ can be chosen as

$$
\Delta = \{x \in \mathbb{R}^n : \text{diag}(x) \in \Gamma\}. \tag{4.4}
$$

Next, we are going to justify a similar result as [\[3,](#page-27-9) Proposition 2.3] for orthogonally invariant matrix sets, which allows us to obtain a chain rule for the subderivative of orthogonally invariant matrix functions via the established theory for composite functions in [\[13,](#page-27-18) Theorem 3.4].

Proposition 4.1. Let K be an absolutely symmetric subset of \mathbb{R}^n . Then the distance function $d_{\sigma^{-1}(K)}$ to the orthogonally invariant matrix set $\sigma^{-1}(K)$ satisfies:

$$
d_{\sigma^{-1}(K)} = d_K \circ \sigma. \tag{4.5}
$$

In particular, if F is an orthogonally invariant matrix function, then for any $X \in M_{m,n}$, we have

$$
dist(X, domF) = dist(\sigma(X), domf), \qquad (4.6)
$$

where f is taken by (4.1) .

Proof To see that $d_{\sigma^{-1}(K)}$ is an orthogonally invariant matrix function, we fix $X \in M_{m,n}$ and $(U, V) \in \mathcal{O}^{m,n}$ such that $X = U \text{diag}(\sigma(X)) V^T$. Then we have

$$
d_K(\sigma(X)) = \inf_{y \in K} \|\sigma(X) - y\|
$$

\n
$$
= \inf_{y \in K} \|\text{diag}(\sigma(X)) - \text{diag}(y)\|
$$

\n
$$
\geq \inf_{Y \in \sigma^{-1}(K)} \|\text{diag}(\sigma(X)) - Y\|
$$

\n
$$
= \inf_{Y \in \sigma^{-1}(K)} \|U \text{diag}(\sigma(X))V^T - UYV^T\|
$$

\n
$$
= \inf_{U^T \bar{Y} V \in \sigma^{-1}(K)} \|X - \bar{Y}\|
$$

\n
$$
= \inf_{\bar{Y} \in \sigma^{-1}(K)} \|X - \bar{Y}\|
$$

\n
$$
= d_{\sigma^{-1}(K)}(X).
$$

On the other hand, it follows from [\(2.6\)](#page-4-1) that

$$
d_{\sigma^{-1}(K)}(X) = \inf_{Y \in \sigma^{-1}(K)} ||X - Y||
$$

\n
$$
\geq \inf_{Y \in \sigma^{-1}(K)} ||\sigma(X) - \sigma(Y)||
$$

\n
$$
= \inf_{y \in K} ||\sigma(X) - y||
$$

\n
$$
= d_K(\sigma(X)).
$$

In particular, if F is an orthogonally invariant matrix function, there exists an absolutely symmetric function $f: \mathbb{R}^n \to [-\infty, +\infty]$ satisfying [\(4.1\)](#page-10-2). Combining this with (4.1), we obtain that

$$
\text{dom}F = \{ X \in \mathbb{M}_{m,n} \colon \sigma(X) \in \text{dom}f \} = \sigma^{-1}(\text{dom}f) \tag{4.7}
$$

is an orthogonally invariant matrix set and dom f is an absolutely symmetric set. From (4.5) , we can deduce that $dist(X, domF) = dist(\sigma(X), domf)$ for any $X \in M_{m,n}$. \Box

To prove a chain rule for subderivatives of orthogonally invariant matrix functions, let's first recall a useful characterization of the subdifferential of orthogonally invariant matrix functions.

Proposition 4.2. Assume that $f: \mathbb{R}^n \to [-\infty, +\infty]$ is a proper, convex, lsc, and absolutely symmetric function. Then the following properties are equivalent:

(i) $Y \in \partial (f \circ \sigma)(X);$

(ii) $\sigma(Y) \in \partial f(\sigma(X))$ and the matrices X and Y have a simultaneous ordered singular value decomposition.

Proof According to Lewis [\[9,](#page-27-4) Theorem 2.4, Corollary 2.6], $f \circ \sigma$ is convex and lsc if and only if s convex and lsc. The claimed equivalence then follows from Lewis [9, Corollary 2.5]. f is convex and lsc. The claimed equivalence then follows from Lewis $[9,$ Corollary 2.5].

Theorem 4.3. (Subderivatives of Orthogonally Invariant Matrix Functions). Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be an absolutely symmetric function and let $X \in \mathbb{M}_{m,n}$ with $(f \circ \sigma)(X)$ finite. If f satisfies one of the following conditions:

(a) f is lsc and convex with $\partial f(\sigma(X)) \neq \emptyset;$

(b) f is locally Lipschitz continuous around $\sigma(X)$ relative to its domain.

Then for all $H \in \mathbb{M}_{m,n}$, we have

$$
d(f \circ \sigma)(X)(H) = df(\sigma(X))(\sigma'(X; H)).
$$
\n(4.8)

Proof Choose any $H \in \mathbb{M}_{m,n}$. According to Proposition [2.4,](#page-5-0) $\sigma'(X; \cdot)$ is a Lipschitz-continuous and positively homogeneous function. Moreover, $\sigma'(X;E) + O(t^2 ||E||^2)/t \to \sigma'(X;H)$ as $t \downarrow 0$ and $E \to H$. Combining this with the definition of subderivative, we derive

$$
d(f \circ \sigma)(X)(H) = \liminf_{\substack{t \downarrow 0 \\ E \to H}} \frac{f(\sigma(X + tE)) - f(\sigma(X))}{t}
$$

\n
$$
= \liminf_{\substack{t \downarrow 0 \\ E \to H}} \frac{f(\sigma(X) + t\sigma'(X; E) + O(t^2 \|E\|^2)) - f(\sigma(X))}{t}
$$

\n
$$
= \liminf_{\substack{t \downarrow 0 \\ E \to H}} \frac{f(\sigma(X) + t(\sigma'(X; E) + \frac{O(t^2 \|E\|^2)}{t})) - f(\sigma(X))}{t}
$$

\n
$$
\geq df(\sigma(X))(\sigma'(X; H)). \tag{4.9}
$$

This establishes the inequality " \geq " in [\(4.8\)](#page-12-1). For the opposite inequality, let's consider two cases. First, if $df(\sigma(X))(\sigma'(X;H)) = \infty$, the latter inequality clearly holds. So, we'll assume $df(\sigma(X))(\sigma'(X;H)) < \infty$. If f is convex and lsc, $f \circ \sigma$ is also convex and lsc according to [\[9,](#page-27-4) Corol-lary 2.6]. Furthermore, Lewis and Sendov [\[10,](#page-27-5) Theorem 7.1] and the non-emptiness of $\partial f(\sigma(X))$ imply that $\partial(f \circ \sigma)(X) \neq \emptyset$. Thus, it follows from Bonnans and Shapiro [\[1,](#page-27-19) Proposition 2.126]
that $d(f \circ \sigma)(X)(H) = \sup \langle Y, H \rangle$. Let $\epsilon > 0$ and choose $Y \in \partial(f \circ \sigma)(X)$ such that that $d(f \circ \sigma)(X)(H) =$ Sup
 $Y \in \partial (f \circ \sigma)(X)$ $\langle Y, H \rangle$. Let $\epsilon > 0$ and choose $Y \in \partial (f \circ \sigma)(X)$ such that $d(f \circ \sigma)(X)(H) \leq \langle Y, H \rangle + \epsilon$. Since $Y \in \partial (f \circ \sigma)(X)$, it follows from Proposition [4.2](#page-12-2) that there exists $(U, V) \in \mathcal{O}^{m,n}$ such that $\sigma(Y) \in \partial f(\sigma(X))$. Setting $\Sigma(Y) := U^T Y V$ and applying Proposition

[2.4](#page-5-0) and the fact that $\left\langle \Sigma(Y)_{\alpha_i \alpha_i}, \frac{1}{2} (V_{\alpha_i}^T H^T U_{\alpha_i} - U_{\alpha_i}^T H V_{\alpha_i}) \right\rangle = 0$ for any $i = 1, ..., t$, we get

$$
d(f \circ \sigma)(X)(H) \leq \langle Y, H \rangle + \epsilon = \langle \Sigma(Y), U^{T}HV \rangle + \epsilon
$$
\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, U_{\alpha_{i}}^{T}HV_{\alpha_{i}} \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle + \epsilon
$$
\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, U_{\alpha_{i}}^{T}HV_{\alpha_{i}} \rangle + \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, \frac{1}{2}(V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}} - U_{\alpha_{i}}^{T}HV_{\alpha_{i}}) \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle + \epsilon
$$
\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, \frac{1}{2}(U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle + \epsilon
$$
\n
$$
\leq \sum_{i=1}^{t} \langle \lambda(\Sigma(Y)_{\alpha_{i}\alpha_{i}}), \frac{1}{2}\lambda(U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \langle \sigma(\Sigma(Y)_{\beta\beta}), \sigma(U_{\beta}^{T}HV_{\beta}) \rangle + \epsilon
$$
\n
$$
= \sum_{i=1}^{t} \sum_{s \in \alpha_{i}} \langle \sigma_{s}(Y), \frac{1}{2}\lambda_{l_{s}}(U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \sum_{s \in \beta} \langle \sigma_{s}(Y), \sigma_{l_{s}}(U_{\beta}^{T}HV_{\beta}) \rangle + \epsilon
$$
\n
$$
= \sum_{i=1}^{t} \sum_{s \in \alpha_{i}} \langle \sigma_{s}(Y), \sigma'_{s}(X;H) \rangle + \sum_{s \in \beta} \langle \sigma_{s}(Y), \sigma'_{s}(X;H) \
$$

This second inequality follows from Fan's inequality [\(2.3\)](#page-3-2) and von Neumann's trace inequality [\(2.5\)](#page-4-2) and the last inequality stems from f being lsc and convex and $\sigma(Y) \in \partial f(\sigma(X))$. As $\epsilon \downarrow 0$, we obtain the opposite inequality in (4.9) , thus proving (4.8) in this case. Assume that f is locally Lipschitz continuous around $\sigma(X)$ relative to its domain. To prove the opposite inequality in [\(4.9\)](#page-12-3), we need only consider the case where $df(\sigma(X))(\sigma'(X;H)) < \infty$. By the definition of subderivatives, there exist sequences $t_k \downarrow 0$ and $v_k \rightarrow \sigma'(X; H)$ such that

$$
df(\sigma(X))(\sigma'(X;H)) = \lim_{k \to \infty} \frac{f(\sigma(X) + t_k v_k) - f(\sigma(X))}{t_k}.
$$
\n(4.10)

Without loss of generality, we assume that $\sigma(X) + t_k v_k \in \text{dom } f$ for all $k \in \mathbb{N}$. From [\(4.6\)](#page-11-1), we get

$$
dist(X + t_k H, domF) = dist(\sigma(X + t_k H), domf)
$$
 for all $k \in \mathbb{N}$,

which, together with Proposition [3.1,](#page-6-1) implies that for all $k \in \mathbb{N}$,

$$
\operatorname{dist}\left(H, \frac{\operatorname{dom} F - X}{t_k}\right) = \frac{1}{t_k} \operatorname{dist}(\sigma(X) + t_k \sigma'(X; H) + O(t_k^2), \operatorname{dom} f)
$$

$$
= \frac{1}{t_k} ||\sigma(X) + t_k \sigma'(X; H) + O(t_k^2) - (\sigma(X) + t_k v_k) ||
$$

$$
= \left\|\sigma'(X; H) - v_k + \frac{O(t_k^2)}{t_k}\right\|.
$$

Then for each $k \in \mathbb{N}$, we can find a matrix $H_k \in \frac{\text{dom} F - X}{t_k}$ satisfying

$$
||H - H_k|| < \left\| \sigma'(X;H) - v_k + \frac{O(t_k^2)}{t_k} \right\| + \frac{1}{k}.
$$

This allows us to have $X + t_k H_k \in \text{dom} F$ for all k and $H_k \to H$ as $k \to \infty$. This, coupled with (4.10) , Proposition [3.1](#page-6-1) and the imposed assumption on f, leads us to obtain

$$
df(\sigma(X))(\sigma'(X;H)) = \lim_{k \to \infty} \left[\frac{f(\sigma(X + t_k H_k)) - f(\sigma(X))}{t_k} + \frac{f(\sigma(X) + t_k v_k) - f(\sigma(X + t_k H_k))}{t_k} \right]
$$

\n
$$
\geq \liminf_{k \to \infty} \frac{f(\sigma(X + t_k H_k)) - f(\sigma(X))}{t_k} - l \lim_{k \to \infty} \left\| \frac{\sigma(X + t_k H_k) - \sigma(X)}{t_k} - v_k \right\|
$$

\n
$$
\geq d(f \circ \sigma)(X)(H) - l \lim_{k \to \infty} \left\| \sigma'(X;H_k) + \frac{O(t_k^2)}{t_k} - v_k \right\|
$$

\n
$$
= d(f \circ \sigma)(X)(H),
$$

where $l \geq 0$ is a Lipschitz constant of f around $\sigma(X)$ relative to its domain. This verifies the inequality "<" in (4.8), and ends the proof. inequality " \leq " in [\(4.8\)](#page-12-1), and ends the proof.

As a direct result of Theorem [4.3,](#page-12-0) we derive a straightforward representation of tangent cones for orthogonally invariant matrix sets. Note that reducing (4.6) , which was stated for the orthogonally invariant matrix function in (4.1) , to the orthogonally invariant matrix set Γ in (4.3) leads us the estimate

$$
dist(X, \Gamma) = dist(\sigma(X), \Delta) \text{ for any } X \in M_{m,n},
$$
\n(4.11)

where Δ is taken from [\(4.4\)](#page-11-3).

Corollary 4.4. (Tangent Cone to the Orthogonally Invariant Matrix Sets). Let Γ be an orthogo-nally invariant matrix set represented by [\(4.3\)](#page-11-2). Then for any $X \in \Gamma$, we have

$$
T_{\Gamma}(X) = \{ H \in \mathbb{M}_{m,n} \colon \sigma'(X;H) \in T_{\Delta}(\sigma(X)) \}.
$$

Proof Using the absolutely symmetric set Δ from [\(4.3\)](#page-11-2), we apply Theorem [4.3](#page-12-0) to the absolutely symmetric function δ_{Δ} and then

$$
d\delta_{\Gamma}(X)(H) = d\delta_{\Delta}(\sigma(X))(\sigma'(X;H)).
$$

The representation of the tangent cone to Γ at $X \in \Gamma$ follows from two key facts: $d\delta_{\Gamma}(X) = \delta_{T_{\Gamma}(X)}$
and $d\delta_{\Delta}(\sigma(X)) = \delta_{T_{\Gamma}}(\Gamma(X))$ and $d\delta_{\Delta}(\sigma(X)) = \delta_{T_{\Delta}(\sigma(X))}$.

In what follows, we are going to present the second-order tangent sets of orthogonally invariant matrix sets. To this end, we begin by demonstrating that certain second-order approximations of orthogonally invariant matrix sets possess an outer Lipschitzian property. Define the set-valued mapping $G_H: \mathbb{R}^n \rightrightarrows \mathbb{M}_{m,n}$ via the second-order tangent set to the absolutely symmetric set Δ in (4.4) by

$$
G_H(b) = \{ W \in \mathbb{M}_{m,n} \colon \sigma''(X; H, W) + b \in T^2_{\Delta}(\sigma(X), \sigma'(X; H)) \}. \tag{4.12}
$$

Proposition 4.5. Assume that Γ is an orthogonally invariant matrix set represented by [\(4.3\)](#page-11-2) and that $X \in \Gamma$ and $H \in T_{\Gamma}(X)$. Then, the mapping G_H in [\(4.12\)](#page-13-1) enjoys the following uniform outer Lipschitzian property at the origin:

$$
G_H(b) \subset G_H(0) + ||b|| \mathbb{B} \text{ for any } b \in \mathbb{R}^n. \tag{4.13}
$$

Proof Let $b \in \mathbb{R}^n$ and pick any $W \in G_H(b)$. It follows from (4.12) that $\sigma''(X; H, W) + b \in$ $T_{\Delta}^2(\sigma(X), \sigma'(X; H))$. By the definition of second-order tangent set, there exists a sequence $t_k \downarrow 0$ such that

$$
\sigma(X) + t_k \sigma'(X; H) + \frac{1}{2} t_k^2 \sigma''(X; H, W) + \frac{1}{2} t_k^2 b + o(t_k^2) \in \Delta \text{ for any } k \in \mathbb{N}.
$$

For any k sufficiently large, we conclude from (2.14) that

$$
\sigma\left(X + t_k H + \frac{1}{2}t_k^2 W\right) = \sigma(X) + t_k \sigma'(X; H) + \frac{1}{2}t_k^2 \sigma''(X; H, W) + o(t_k^2),
$$

which, coupled with [\(4.11\)](#page-13-2), leads us to

$$
\text{dist}\left(X + t_k H + \frac{1}{2}t_k^2 W, \Gamma\right) = \text{dist}\left(\sigma(X + t_k H + \frac{1}{2}t_k^2 W), \Delta\right) \le \frac{1}{2}t_k^2 \|b\| + o(t_k^2).
$$

Thus, there exists $Y_k \in \Gamma$ such that

$$
||E_k|| \leq \frac{1}{2}||b|| + \frac{o(t_k^2)}{t_k^2},
$$

where $E_k := \frac{X+t_kH+\frac{1}{2}t_k^2W-Y_k}{t_k^2}$. Passing to a subsequence, if necessary, implies that there exists $E \in \mathbb{M}_{m,n}$ such that $\tilde{E_k} \to E$ as $k \to \infty$. This brings us to

$$
||E|| \le \frac{1}{2}||b||. \tag{4.14}
$$

Since $X + t_k H + \frac{1}{2} t_k^2 W - t_k^2 E_k = Y_k \in \Gamma$, it follows from [\(4.3\)](#page-11-2) and [\(2.14\)](#page-6-3) that

$$
\sigma(X) + t_k \sigma'(X;H) + \frac{1}{2} t_k^2 \sigma''(X;H,W - 2E) + o(t_k^2) = \sigma(X + t_k H + \frac{1}{2} t_k^2 W - t_k^2 E_k) \in \Delta.
$$

By the definition of the second-order tangent set, we get

$$
\sigma''(X; H, W - 2E) \in T_\Delta^2(\sigma(X), \sigma'(X; H)),
$$

which leads us to $W - 2E \in G_H(0)$. This, combined with [\(4.14\)](#page-14-0), implies [\(4.13\)](#page-14-1) and thus ends the proof. proof.

Proposition 4.6. (Second-Order Tangent Sets of Orthogonally Invariant Matrix Sets). Assume that Γ is an orthogonally invariant matrix set represented by [\(4.3\)](#page-11-2) and that $X \in \Gamma$ and $H \in T_{\Gamma}(X)$. Then, we have

$$
T_{\Gamma}^{2}(X,H) = \{ W \in \mathbb{M}_{m,n} \colon \sigma''(X;H,W) \in T_{\Delta}^{2}(\sigma(X), \sigma'(X;H)) \},
$$
\n(4.15)

where Δ is taken from [\(4.4\)](#page-11-3). Moreover, if the absolutely symmetric set Δ is parabolically derivable at $\sigma(X)$ for $\sigma'(X; H)$, then Γ is parabolically derivable at X for H.

Proof Pick any $W \in T_{\Gamma}^2(X, H)$. By the definition of second-order tangent set, there exists a sequence $t_k \downarrow 0$ such that

$$
X + t_k H + \frac{1}{2}t_k^2 W + o(t_k^2) \in \Gamma.
$$

It follows from (4.3) and (2.14) that

$$
\sigma(X) + t_k \sigma'(X;H) + \frac{1}{2} t_k^2 \sigma''(X;H,W) + o(t_k^2) = \sigma(X + t_k H + \frac{1}{2} t_k^2 W + o(t_k^2)) \in \Delta.
$$

Thus we get $\sigma''(X; H, W) \in T^2_{\Delta}(\sigma(X), \sigma'(X; H))$ and prove the inclusion "⊂" in [\(4.15\)](#page-14-2). To prove the opposite inclusion in (4.15) , we take any $W \in M_{m,n}$ satisfying $\sigma''(X; H, W) \in T^2_{\Delta}(\sigma(X), \sigma'(X; H)).$ By the definition of second-order tangent set, there exists a sequence $t_k \downarrow 0$ such that

$$
\sigma(X) + t_k \sigma'(X;H) + \frac{1}{2} t_k^2 \sigma''(X;H,W) + o(t_k^2) \in \Delta.
$$

Combining (2.14) with (4.11) , we obtain that for any $t > 0$ sufficiently small,

$$
\begin{split} \text{dist}\left(X+tH+\frac{1}{2}t^2W,\Gamma\right) &= \text{dist}\left(\sigma\left(X+tH+\frac{1}{2}t^2W\right),\Delta\right) \\ &= \text{dist}\left(\sigma(X)+t\sigma'(X;H)+\frac{1}{2}t^2\sigma''(X;H,W)+o(t^2),\Delta\right). \end{split}
$$

Thus we get that $X + t_k H + \frac{1}{2} t_k^2 W \in \Gamma$, hence that $W \in T_{\Gamma}^2(X, H)$. This verifies the inclusion " \supset " in (4.15) , and then (4.15) is proved.

To prove the parabolic derivability of Γ at X for H, we first show that $T_{\Gamma}^2(X, H)$ is a non-empty set. Suppose that the absolutely symmetric set Δ is parabolically derivable at $\sigma(X)$ for $\sigma'(X;H)$, it follows from the definition of parabolically derivable that $T^2_{\Delta}(\sigma(X), \sigma'(X; H))$ is nonempty. Thus there exists $p \in T^2_{\Delta}(\sigma(X), \sigma'(X; H))$ and $\widetilde{W} \in M_{m,n}$ such that

$$
\sigma''(X; H, \widetilde{W}) + b \in T_\Delta^2(\sigma(X), \sigma'(X; H)),
$$

where $b := p - \sigma''(X; H, \overline{W})$. It is easy to obtain $\overline{W} \in G_H(b)$ by [\(4.12\)](#page-13-1). According to the outer Lipschitzian property of the set-valued mapping G_H , there exists $W \in G_H(0)$ such that $\|W - \tilde{W}\| \le$ b. This leads us to $\sigma''(X; H, W) \in T^2_{\Delta}(\sigma(X), \sigma'(X; H))$. By [\(4.15\)](#page-14-2), we have $W \in T^2_{\Gamma}(X, H)$, which justifies $T_{\Gamma}^2(X, H) \neq \emptyset$. It remains to prove that for each $W \in T_{\Gamma}^2(X, H)$, there exists $\epsilon > 0$ and an arc $\xi : [0, \epsilon] \to \Gamma$ with $\xi(0) = X$, $\xi'_{+}(0) = H$, and $\xi''_{+}(0) = W$. It suffices to show that for all $t \in [0, \epsilon],$

$$
X + tH + \frac{1}{2}t^2W + o(t^2) \in \Gamma.
$$
 (4.16)

Pick any $W \in T_{\Gamma}^2(X, H)$, we have $\sigma''(X; H, W) \in T_{\Delta}^2(\sigma(X), \sigma'(X; H))$ by [\(4.15\)](#page-14-2). Since the absolutely symmetric set Δ is parabolically derivable at $\sigma(X)$ for $\sigma'(X;H)$, there exists $\epsilon' > 0$ and an arc $\zeta: [0, \epsilon'] \to \Delta$ with $\zeta(0) = \sigma(X), \, \zeta'_+(0) = \sigma'(X; H),$ and $\zeta''_+(0) = \sigma''(X; H, W)$. Thus we get $\zeta(t) = \sigma(X) + t\sigma'(X;H) + \frac{1}{2}t^2\sigma''(X;H,W) + o(t^2) \in \Delta$ for all $t \in [0,\epsilon']$. This, combined with [\(2.14\)](#page-6-3) and [\(4.3\)](#page-11-2), implies [\(4.16\)](#page-15-0) for some $\epsilon \leq \epsilon'$. This justifies that Γ is parabolically derivable at X for H , and completes the proof. \Box

We proceed by proving a chain rule for subderivatives of absolutely symmetric function. We begin with presenting a counterpart of the estimate in [\(4.6\)](#page-11-1) for domains of absolutely symmetric functions.

Proposition 4.7. Let $F: \mathbb{M}_{m,n} \to [-\infty, +\infty]$ be an orthogonally invariant matrix function, rep-resented by [\(4.1\)](#page-10-2). Then for any $x \in \mathbb{R}^n$, we have

$$
dist(x, dom f) = dist(diag(x), dom F),
$$
\n(4.17)

where f is taken by (4.1) .

Proof For any $x \in \mathbb{R}^n$, there exists a signed permutation matrix $Q_{\pm} \in \mathbf{P}_{\pm}^n$ such that $\sigma(\text{diag}(x)) =$ $Q_{+}x$. As pointed out before, the absolutely symmetric function f can be represented as a composite function of the orthogonally invariant matrix function F and the linear mapping $x \mapsto diag(x)$ with $x \in \mathbb{R}^n$. Thus, it follows from (4.2) that

$$
\text{dom} f = \{ x \in \mathbb{R}^n \colon \text{diag}(x) \in \text{dom} F \} \tag{4.18}
$$

is an absolutely symmetric set. This, combined with (4.7) , implies that $Q_{\pm}^T \sigma(X) \in \text{dom} f$ for any $X \in \text{dom} F$. Then by von Neumann's trace inequality (2.5) , we obtain

dist
$$
(x, \text{dom } f) \le ||x - Q_{\pm}^T \sigma(X)|| = ||Q_{\pm} x - \sigma(X)|| = ||\sigma(\text{diag}(x)) - \sigma(X)|| \le ||\text{diag}(x) - X||,
$$

for all $X \in \text{dom} F$, which, in turn, leads us to

$$
dist(x, dom f) \leq dist(diag(x), dom F). \tag{4.19}
$$

 \Box

To prove the opposite inequality, pick any $y \in \text{dom } f$. By [\(4.18\)](#page-15-1), we have $\text{diag}(y) \in \text{dom } F$ and then

$$
dist(diag(x), domF) \le ||diag(x) - diag(y)|| = ||x - y||,
$$

which implies the opposite inequality in (4.19) , and completes the proof.

Note that the linear mapping $x \mapsto diag(x)$ with $x \in \mathbb{R}^n$ is twice continuously differentiable. This, coupled with [\(4.17\)](#page-15-3) and [\[13,](#page-27-18) Theorem 3.4], brings us to the following theorem.

Theorem 4.8. (Subderivatives of Absolutely Symmetric Functions). Let $F: \mathbb{M}_{m,n} \to [-\infty, +\infty]$ be an orthogonally invariant matrix function, represented by (4.1) , and let its associated absolutely symmetric function f be locally Lipschitz continuous relative to its domain. Then, for any $X \in$ $\mathbb{M}_{m,n}$ with $F(X)$ finite and any $z \in \mathbb{R}^n$, we have

$$
df(\sigma(X))(z) = dF(\text{diag}(\sigma(X)))(\text{diag}(z)).\tag{4.20}
$$

Proof Since f is locally Lipschitz continuous relative to its domain, by (2.6) we obtain that F is locally Lipschitz continuous relative to its domain. According to [\[13,](#page-27-18) Theorem 3.4], we justify $(4.20).$ $(4.20).$ \Box

Similar to Corollary [4.4,](#page-13-3) we proceed to derive a refined representation of tangent cones for absolutely symmetric sets according to Theorem [4.8.](#page-16-0)

Corollary 4.9. (Tangent Cone to the Absolutely Symmetric Sets). Let Δ be an absolutely sym-metric set represented by [\(4.4\)](#page-11-3). Then for any $X \in \Gamma$, we have

$$
T_{\Delta}(\sigma(X)) = \{ z \in \mathbb{R}^n \colon \text{diag}(z) \in T_{\Gamma}(\text{diag}(\sigma(X))) \}.
$$

We close this section by revealing that the subderivative of f at $\sigma(X)$ is a symmetric function with respect to $\mathbf{P}_{\pm}^n(X)$, which is a subset of \mathbf{P}_{\pm}^n consisting of all $n \times n$ block diagonal matrices in the form $Q_{\pm} = \text{diag}(Q_1, \ldots, Q_t, Q_{t+1})$, where $Q_i \in \mathbb{R}^{|\alpha_i| \times |\alpha_i|}$ is a signed permutation matrix for any $i = 1, \ldots, t$ with α_i taken from [\(2.7\)](#page-4-3), t being the number of distinct nonzero singular values of X and $Q_{t+1} \in \mathbb{R}^{|\beta| \times |\beta|}$ is also a signed permutation matrix. Obviously, we have that $\mathbf{P}_{\pm}^n(X) \subset \mathbf{P}_{\pm}^n$ and that if $Q_{\pm} \in \mathbf{P}_{\pm}^n(X)$, then $Q_{\pm} \sigma(X) = \sigma(X)$.

Proposition 4.10. Assume that $f: \mathbb{R}^n \to [-\infty, +\infty]$ is an absolutely symmetric function and $X \in \mathbb{M}_{m,n}$ with $f(\sigma(X))$ finite. Then, for any $v \in \mathbb{R}^n$ and any signed permutation matrix $Q_{\pm} \in$ $\mathbf{P}_{\pm}^n(X)$, we have

$$
df(\sigma(X))(Q_{\pm}v) = df(\sigma(X))(v),
$$

which means that the subderivative $v \mapsto df(\sigma(X))(v)$ is symmetric with respect to $\mathbf{P}_{\pm}^n(X)$.

Proof For any $v \in \mathbb{R}^n$ and $Q_{\pm} \in \mathbf{P}_{\pm}^n(X)$, it follows from the absolutely symmetric property of f that

$$
df(\sigma(X))(v) = \liminf_{\substack{v \to v \\ v \to v}} \frac{f(\sigma(X) + \tau v') - f(\sigma(X))}{\tau}
$$

$$
= \liminf_{\substack{v \to v \\ v \to v}} \frac{f(\sigma(X) + \tau Q_{\pm} v') - f(\sigma(X))}{\tau}
$$

$$
\geq \liminf_{\substack{v \to v \\ v \to Q_{\pm}v}} \frac{f(\sigma(X) + \tau w) - f(\sigma(X))}{\tau}
$$

$$
= df(\sigma(X))(Q_{\pm}v).
$$

According to $Q^{-1}_{\pm} = \text{diag}(Q^{-1}_1, \ldots, Q^{-1}_t, Q^{-1}_{t+1}) \in \mathbf{P}^n_{\pm}(X)$, we can similarly show the opposite inequality above, and the proof is complete. \Box

5 Second Subderivatives of Orthogonally Invariant Matrix Functions

This section is dedicated to the study of second order variational analysis of orthogonally invariant matrix functions. Our objective is to compute the second subderivatives of such functions when the corresponding absolutely symmetric functions are convex. Additionally, we derive second-order optimality conditions for a class of matrix optimization problems. We begin our investigation of the second subderivative of orthogonally invariant matrix functions by establishing a lower bound for it.

Proposition 5.1. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be lsc, convex, and absolutely symmetric. Assume that $\mu_1 > \cdots > \mu_t$ are the distinct nonzero singular values of $X \in \mathbb{M}_{m,n}$, and that $(U, V) \in$

 $\mathcal{O}^{m,n}(X) \cap \mathcal{O}^{m,n}(Y)$. Then, for any $H \in \mathbb{M}_{m,n}$, we have

$$
d^{2}(f \circ \sigma)(X \mid Y)(H) \geq d^{2}f(\sigma(X) \mid \sigma(Y))(\sigma'(X; H))
$$

+
$$
2\sum_{i=1}^{t} \left\langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T}B(H)P_{\alpha_{i}}^{c}(\mu_{i}I - \Lambda_{\alpha_{i}})^{-1}P_{\alpha_{i}}^{c}{}^{T}B(H)P_{\alpha_{i}} \right\rangle
$$

+
$$
2\left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^{T}HV_{\alpha}\Sigma_{\alpha}^{-1}(X)U_{\alpha}^{T}HV_{\beta} \right\rangle,
$$
 (5.1)

where the columns of P_{α_i} form an orthonormal basis of eigenvectors of $\mathcal{B}(X)$ associated with μ_i , $P_{\alpha_i}^c$ is the submatrix of P obtained by removing all the columns of P_{α_i} , and $\Lambda_{\alpha_i} \in \mathbb{R}^{(m+n-|\alpha_i|)\times(m+n-|\alpha_i|)}$ is a diagonal matrix whose diagonal elements are the eigenvalues of $P^T\mathcal{B}(X)P$ that are not equal to μ_i for all $i \in \{1, \ldots, t\}.$

Proof Let $H \in \mathbb{M}_{m,n}$ and pick sequences $H_K \to H$ and $t_k \downarrow 0$. Setting $\Delta_{t_k} \sigma(X)(H_k) := \sigma(X + t_k H_k) - \sigma(X)$ we obtain $\frac{H_k - \sigma(\Lambda)}{t_k}$, we obtain

$$
\Delta_{t_k}^2 (f \circ \sigma)(X \mid Y)(H_k) = \frac{f(\sigma(X + t_k H_k)) - f(\sigma(X)) - t_k \langle Y, H_k \rangle}{\frac{1}{2}t_k^2}
$$

=
$$
\frac{f(\sigma(X) + t_k \Delta_{t_k} \sigma(X)(H_k)) - f(\sigma(X)) - t_k \langle \sigma(Y), \Delta_{t_k} \sigma(X)(H_k) \rangle}{\frac{1}{2}t_k^2} + \frac{\langle \sigma(Y), \Delta_{t_k} \sigma(X)(H_k) \rangle - \langle Y, H_k \rangle}{\frac{1}{2}t_k}
$$

=
$$
\Delta_{t_k}^2 f(\sigma(X) \mid \sigma(Y))(\Delta_{t_k} \sigma(X)(H_k)) + \frac{\langle \sigma(Y), \Delta_{t_k} \sigma(X)(H_k) \rangle - \langle Y, H_k \rangle}{\frac{1}{2}t_k}.
$$

It follows from $Y = U\Sigma(Y)V^T$ that

$$
\langle Y, H_k \rangle = \langle U \Sigma(Y) V^T, H_k \rangle = \langle \Sigma(Y), U^T H_k V \rangle = \sum_{i=1}^t \langle \Sigma(Y)_{\alpha_i \alpha_i}, U_{\alpha_i}^T H_k V_{\alpha_i} \rangle + \langle \Sigma(Y)_{\hat{\beta} \beta}, U_{\hat{\beta}}^T H_k V_{\beta} \rangle.
$$

Moreover, it results from Fan's inequality [\(2.3\)](#page-3-2), von Neumann's trace inequality [\(2.5\)](#page-4-2), and Proposition [3.1](#page-6-1) that

$$
\langle \sigma(Y), \Delta_{t_k} \sigma(X)(H_k) \rangle
$$
\n
$$
= \sum_{i=1}^t \sum_{s \in \alpha_i} \frac{\sigma_s(Y)(\sigma_s(X + t_k H_k) - \sigma_s(X))}{t_k} + \sum_{s \in \beta} \frac{\sigma_s(Y)(\sigma_s(X + t_k H_k) - \sigma_s(X))}{t_k}
$$
\n
$$
= \sum_{i=1}^t \sum_{s \in \alpha_i} \sigma_s(Y)\lambda_{l_s} (P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i} + t_k P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^c {}^T \mathcal{B}(H_k) P_{\alpha_i}) + O(t_k^2)
$$
\n
$$
+ \sum_{s \in \beta} \sigma_s(Y)\sigma_{l_s} (U_{\beta}^T H_k V_{\beta} - t_k U_{\beta}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H_k V_{\beta}) + O(t_k^2)
$$
\n
$$
\geq \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i} + t_k P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^c {}^T \mathcal{B}(H_k) P_{\alpha_i} \right\rangle
$$
\n
$$
+ \left\langle \Sigma(Y)_{\beta \beta}, U_{\beta}^T H_k V_{\beta} - t_k U_{\beta}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H_k V_{\beta} \right\rangle + O(t_k^2).
$$

Thus, we have

$$
\frac{\langle \sigma(Y), \Delta_{t_k}\sigma(X)(H_k) \rangle - \langle Y, H_k \rangle}{\frac{1}{2}t_k}
$$
\n
$$
\geq 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i\alpha_i}, \frac{1}{2} (V_{\alpha_i}^T H_k^T U_{\alpha_i} - U_{\alpha_i}^T H_k V_{\alpha_i}) + P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H_k) P_{\alpha_i} \right\rangle
$$
\n
$$
+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H_k V_{\beta} \right\rangle + O(t_k^2)
$$
\n
$$
= 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i\alpha_i}, P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H_k) P_{\alpha_i} \right\rangle
$$
\n
$$
+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H_k V_{\beta} \right\rangle + O(t_k^2),
$$

and further

$$
\Delta_{t_k}^2(f \circ \sigma)(X \mid Y)(H_k) \geq \Delta_{t_k}^2 f(\sigma(X) \mid \sigma(Y))(\Delta_{t_k} \sigma(X)(H_k))
$$

+
$$
2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H_k) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c} {}^T \mathcal{B}(H_k) P_{\alpha_i} \right\rangle
$$

+
$$
2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H_k V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H_k V_{\beta} \right\rangle + O(t_k^2),
$$

which, in turn, brings us to the lower estimate in [\(5.1\)](#page-17-0) for the second subderivative of $f \circ \sigma$ at X
for Y because $\triangle_{t} \sigma(X)(H_b) \to \sigma'(X \cdot H)$ as $k \to \infty$ for Y because $\triangle_{t_k} \sigma(X)(H_k) \to \sigma'(X; H)$ as $k \to \infty$.

We now advance to a discussion concerning the critical cone of orthogonally invariant matrix functions.

Proposition 5.2. (Critical Cone of Orthogonally Invariant Matrix Functions). Let $f: \mathbb{R}^n \to$ $[-\infty, +\infty]$ be lsc, convex, and absolutely symmetric. Assume that $\mu_1 > \cdots > \mu_t$ are the distinct nonzero singular values of $X \in \mathbb{M}_{m,n}$, and that $Y \in \partial(f \circ \sigma)(X)$. Then, we have $H \in K_{f \circ \sigma}(X, Y)$ if and only if $\sigma'(X;H) \in K_f(\sigma(X), \sigma(Y))$, the matrices $\Sigma(Y)_{\alpha_i \alpha_i}$ and $\frac{1}{2}(U_{\alpha_i}^THV_{\alpha_i} + V_{\alpha_i}^TH^TU_{\alpha_i})$ have a simultaneous ordered spectral decomposition, and the matrices $\Sigma(\bar{Y})_{\hat{\beta}\beta}^{}$ and $U_{\hat{\beta}}^TH\bar{V}_{\beta}^{}$ have a simultaneous ordered singular value decomposition for any $i = 1, \ldots, t$ and $(U, V) \in \mathcal{O}^{m,n}(X) \cap$ $\mathcal{O}^{m,n}(Y)$.

Proof It follows from $Y \in \partial (f \circ \sigma)(X)$ and Proposition [4.2](#page-12-2) that $\sigma(Y) \in \partial f(\sigma(X))$ and we can find $(U, V) \in \mathcal{O}^{m,n}(X) \cap \mathcal{O}^{m,n}(Y)$. From Theorem [4.3,](#page-12-0) we can conclude that

$$
\langle Y, H \rangle = \langle \Sigma(Y), U^{T}HV \rangle
$$

\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, U_{\alpha_{i}}^{T}HV_{\alpha_{i}} \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle
$$

\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, U_{\alpha_{i}}^{T}HV_{\alpha_{i}} \rangle + \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, \frac{1}{2} (V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}} - U_{\alpha_{i}}^{T}HV_{\alpha_{i}}) \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle
$$

\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, \frac{1}{2} (U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \langle \Sigma(Y)_{\beta\beta}, U_{\beta}^{T}HV_{\beta} \rangle
$$

\n
$$
\leq \sum_{i=1}^{t} \langle \lambda(\Sigma(Y)_{\alpha_{i}\alpha_{i}}), \frac{1}{2} \lambda(U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \langle \sigma(\Sigma(Y)_{\beta\beta}), \sigma(U_{\beta}^{T}HV_{\beta}) \rangle
$$

\n
$$
= \sum_{i=1}^{t} \sum_{s \in \alpha_{i}} \langle \sigma_{s}(Y), \frac{1}{2} \lambda_{l_{s}} (U_{\alpha_{i}}^{T}HV_{\alpha_{i}} + V_{\alpha_{i}}^{T}H^{T}U_{\alpha_{i}}) \rangle + \sum_{s \in \beta} \langle \sigma_{s}(Y), \sigma_{s}(Y, \sigma_{s}(Y, \beta_{i}W_{\beta})) \rangle
$$

\n
$$
= \sum_{i=1}^{t} \sum_{s \in \alpha_{i}} \langle \sigma_{s}(Y), \sigma_{s}'(X;H) \rangle + \sum_{s \in \beta} \langle \sigma_{s}(Y), \sigma_{s}'(X;H) \rangle
$$

\n
$$
= \langle \
$$

The condition $H \in K_{f \circ \sigma}(X, Y)$ is equivalent to equality between the left and right hand sides (and hence throughout), and the claimed equivalence then results from $\sigma'(X; H) \in K_f(\sigma(X), \sigma(Y)),$ Fan's inequality [\(2.3\)](#page-3-2), and von Neumann's trace theorem (Lemma [2.3\)](#page-4-4).

The following result provides sufficient conditions for the parabolic epi-differentiability of orthogonally invariant matrix functions. Furthermore, it derives a valuable formula for the parabolic subderivatives of this class of functions.

Theorem 5.3. (Parabolic Subderivatives of Orthogonally Invariant Matrix Functions). Let $f: \mathbb{R}^n \to$ $[-\infty, +\infty]$ be an absolutely symmetric function and let $X \in \mathbb{M}_{m,n}$ with $(f \circ \sigma)(X)$ finite. Assume that $H \in T_{dom(f \circ \sigma)}(X)$ and that f is locally Lipschitz continuous relative to its domain and parabolically epi-differentiable at $\sigma(X)$ for $\sigma'(X;H)$. Then the parabolic subderivative of orthogonally invariant matrix function $f \circ \sigma$ at X for H with respect to $W \in M_{m,n}$ is

$$
d^{2}(f \circ \sigma)(X)(H \mid W) = d^{2}f(\sigma(X))(\sigma'(X;H) \mid \sigma''(X;H,W)),
$$
\n(5.2)

and $f \circ \sigma$ is parabolically epi-differentiable at X for H.

Proof Let $F := f \circ \sigma$ and pick $W \in M_{m,n}$. Since f is parabolically epi-differentiable at $\sigma(X)$ for $\sigma'(X;H)$, domf is parabolically derivable at $\sigma(X)$ for $\sigma'(X;H)$. We conclude from [\(4.7\)](#page-11-5), Proposition [4.6,](#page-14-3) Corollary [4.4,](#page-13-3) and [\(4.15\)](#page-14-2) that dom F is parabolically derivable at X for H and the tangent cone of domF at X and second-order tangent set of domF at X for $H \in T_{\text{dom}F}(X)$ are, respectively,

$$
T_{\text{dom}F}(X) = \{ H \in \mathbb{M}_{m,n} \colon \sigma'(X;H) \in T_{\text{dom}f}(\sigma(X)) \}
$$

and

$$
T^{2}_{\text{dom}F}(X,H) = \{ W \in \mathbb{M}_{m,n} \colon \sigma''(X;H,W) \in T^{2}_{\text{dom}f}(\sigma(X),\sigma'(X;H)) \} \neq \emptyset. \tag{5.3}
$$

Since $H \in T_{\text{dom}F}(X)$ amounts to $\sigma'(X; H) \in T_{\text{dom}f}(\sigma(X))$, we deduce from the imposed assumption on f that

$$
\operatorname{dom} \mathrm{d}^2 f(\sigma(X))(\sigma'(X;H) \mid \cdot) = T^2_{\operatorname{dom} f}(\sigma(X), \sigma'(X;H)). \tag{5.4}
$$

The proof falls naturally into two cases. Let us first assume $W \notin T^2_{\text{dom} F}(X, H)$. Combining [\(5.3\)](#page-19-0) with (5.4) , we obtain that

$$
d^2 f(\sigma(X))(\sigma'(X;H) | \sigma''(X;H,W)) = \infty.
$$

On the other hand, by definition, it is not hard to see that the inclusion dom $d^2F(X)(H | \cdot) \subset$ $T^2_{\text{dom}F}(X, H)$ always holds for any $H \in T_{\text{dom}F}(X)$. Thus we have $d^2F(X)(H \mid W) = \infty$. This implies [\(5.2\)](#page-18-0) for every $W \notin T^2_{\text{dom} F}(X, H)$. Now consider an arbitrary sequence $t_k \downarrow 0$, set $W_k := W$ for all $k \in \mathbb{N}$, and obtain that

$$
d^{2}F(X)(H \mid W) \leq \liminf_{k \to \infty} \frac{F(X + t_{k}H + \frac{1}{2}t_{k}^{2}W_{k}) - F(X) - t_{k}dF(X)(H)}{\frac{1}{2}t_{k}^{2}} \leq \limsup_{k \to \infty} \frac{F(X + t_{k}H + \frac{1}{2}t_{k}^{2}W_{k}) - F(X) - t_{k}dF(X)(H)}{\frac{1}{2}t_{k}^{2}}.
$$

Since $d^2F(X)(H | W) = \infty$ for all $W \notin T^2_{dom F}(X, H)$, the inequality in the above expression can take equality.

Turn to the case $W \in T^2_{\text{dom} F}(X, H)$. Combining [\(5.3\)](#page-19-0) with [\(5.4\)](#page-19-1), we obtain that

$$
d^2 f(\sigma(X))(\sigma'(X;H) \mid \sigma''(X;H,W)) < \infty. \tag{5.5}
$$

Since domF is parabolically derivable at X for H, for arbitrary sequence $t_k \downarrow 0$, there exists a sequence $W_k \to W$ as $k \to \infty$ such that

$$
X_k := X + t_k H + \frac{1}{2} t_k^2 W_k = X + t_k H + \frac{1}{2} t_k^2 W + o(t_k^2) \in \text{dom} F.
$$

Because f is parabolically epi-differentiable at $\sigma(X)$ for $\sigma'(X; H)$, for the vector $\sigma''(X; H, W) \in \mathbb{R}^n$, corresponding to the above t_k , we find a sequence $w_k \to \sigma''(X; H, W)$ such that

$$
d^2 f(\sigma(X))(\sigma'(X;H) \mid \sigma''(X;H,W)) = \lim_{k \to \infty} \frac{f(y_k) - f(\sigma(X)) - t_k df(\sigma(X)) (\sigma'(X;H))}{\frac{1}{2}t_k^2},
$$

where $y_k := \sigma(X) + t_k \sigma'(X; H) + \frac{1}{2} t_k^2 w_k$. It follows from [\(5.5\)](#page-19-2) that $y_k \in \text{dom} f$ for all k sufficiently large. We deduce from (4.8) and (2.14) that

$$
d^{2}F(X)(H|W) \leq \liminf_{k \to \infty} \frac{F(X + t_{k}H + \frac{1}{2}t_{k}^{2}W_{k}) - F(X) - t_{k}dF(X)(H)}{\frac{1}{2}t_{k}^{2}}
$$

\n
$$
\leq \limsup_{k \to \infty} \frac{F(X_{k}) - F(X) - t_{k}dF(X)(H)}{\frac{1}{2}t_{k}^{2}}
$$

\n
$$
= \limsup_{k \to \infty} \frac{f(\sigma(X_{k})) - f(\sigma(X)) - t_{k}df(\sigma(X))(\sigma'(X;H))}{\frac{1}{2}t_{k}^{2}}
$$

\n
$$
\leq \limsup_{k \to \infty} \frac{f(y_{k}) - f(\sigma(X)) - t_{k}df(\sigma(X))(\sigma'(X;H))}{\frac{1}{2}t_{k}^{2}} + \limsup_{k \to \infty} \frac{f(\sigma(X_{k})) - f(y_{k})}{\frac{1}{2}t_{k}^{2}}
$$

\n
$$
\leq d^{2}f(\sigma(X))(\sigma'(X;H) | \sigma''(X;H,W)) + c \limsup_{k \to \infty} ||\sigma''(X;H,W) + \frac{\sigma(t_{k}^{2})}{\frac{1}{2}t_{k}^{2}} - w_{k}||
$$

\n
$$
= d^{2}f(\sigma(X))(\sigma'(X;H) | \sigma''(X;H,W)),
$$

where $c \geq 0$ is a Lipschitz constant of f around $\sigma(X)$ relative to its domain. On the other hand, we conclude from the definition of parabolic subderivative that for any sequence $t_k \downarrow 0$ and any sequence $W_k \to W$ as $k \to \infty$, one has

$$
\liminf_{k \to \infty} \frac{F(X + t_k H + \frac{1}{2} t_k^2 W_k) - F(X) - t_k dF(X)(H)}{\frac{1}{2} t_k^2}
$$
\n
$$
= \liminf_{k \to \infty} \frac{f(\sigma(X) + t_k \sigma'(X; H) + \frac{1}{2} t_k^2 \sigma''(X; H, W) + o(t_k^2)) - f(\sigma(X)) - t_k d f(\sigma(X)) (\sigma'(X; H))}{\frac{1}{2} t_k^2}
$$
\n
$$
\geq \liminf_{t \to 0, t \to 0,
$$

which verifies the inequality " \geq " in [\(5.2\)](#page-18-0). In summary, for any $W \in T^2_{\text{dom} F}(X, H)$ and any sequence $t_k \downarrow 0$, there exists a sequence $W_k \rightarrow W$ such that

$$
d^{2}F(X)(H \mid W) = \lim_{k \to \infty} \frac{F(X + t_{k}H + \frac{1}{2}t_{k}^{2}W_{k}) - F(X) - t_{k}dF(X)(H)}{\frac{1}{2}t_{k}^{2}},
$$

and

$$
d^{2}F(X)(H \mid W) = d^{2}f(\sigma(X))(\sigma'(X;H) \mid \sigma''(X;H,W)),
$$

which coupled with [\(5.5\)](#page-19-2), leads us to $T^2_{\text{dom} F}(X, H) \subset \text{dom} d^2 F(X)(H | \cdot)$. We conclude from $T^2_{\text{dom}F}(X, H) \neq \emptyset$ that dom $d^2F(X)(H \mid \cdot) \neq \emptyset$, and thus F is parabolically epi-differentiable at X for H.

Recall from Proposition [4.10](#page-16-2) that the subderivative of f at $\sigma(X)$ is a symmetric function with respect to a subset of \mathbf{P}_{\pm}^n . The following result shows the parabolic subderivative of f at $\sigma(X)$ for $\sigma'(X;H)$ is also a symmetric function with respect to a subset of \mathbf{P}_{\pm}^n . Now suppose that N_i is the number of distinct eigenvalues of $\frac{1}{2} \left(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \right)$ for $i = 1, \ldots, t$ and that N_{t+1} is the number of distinct nonzero singular values of $U_{\hat{\beta}}^T H V_{\beta}$. Pick the index sets β_j^i for $i = 1, \ldots, t$, $j = 1, ..., N_i$ from [\(2.9\)](#page-5-2) and β_k^{t+1} for $k = 1, ..., N_{t+1}, N_{t+1} + 1$ from [\(2.10\)](#page-5-3). Denote by $\mathbf{P}_{\pm}^n(X, H)$ a subset of \mathbf{P}_{\pm}^n consisting of all $n \times n$ block diagonal matrices in the form $Q_{\pm} = \text{diag}(Q_1, \ldots, Q_t, Q_{t+1})$ such that for each $i = 1, ..., t$, the $|\alpha_i| \times |\alpha_i|$ signed permutation matrix Q_i has a block diagonal representation

$$
Q_i = \text{diag}\left(B_1^i, \ldots, B_{N_i}^i\right),
$$

and $|\beta| \times |\beta|$ signed permutation matrix Q_{t+1} has a block diagonal representation

$$
Q_{t+1} = \text{diag}\left(B_1^{t+1}, \ldots, B_{N_{t+1}}^{t+1}, B_{N_{t+1}+1}^{t+1}\right),
$$

where $B_j^i \in \mathbb{R}^{|\beta_j^i| \times |\beta_j^i|}$ is a signed permutation matrix for any $i = 1, \ldots, t, j = 1, \ldots, N_i$ and $B_j^{t+1} \in \mathbb{R}^{|\beta_k^{t+1}| \times |\beta_k^{t+1}|}$ is also a signed permutation matrix for any $k = 1, \ldots, N_{t+1}, N_{t+1} + 1$. It is obvious that $\mathbf{P}_{\pm}^n(X, H) \subset \mathbf{P}_{\pm}^n(X) \subset \mathbf{P}_{\pm}^n$ and that if $Q_{\pm} \in \mathbf{P}_{\pm}^n(X, H)$, then $Q_{\pm} \sigma(X) = \sigma(X)$ and $Q_{\pm} \sigma'(X;H) = \sigma'(X;H).$

Proposition 5.4. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be an absolutely symmetric function and $X, H \in \mathbb{M}_{m,n}$ with $f(\sigma(X))$ and $df(\sigma(X))(\sigma'(X;H))$ finite. Then, for any $u \in \mathbb{R}^n$ and any signed permutation $matrix Q_{\pm} \in \mathbf{P}_{\pm}^n(X;H)$, we have

$$
d^2 f(\sigma(X))(\sigma'(X;H) \mid Q_{\pm}u) = d^2 f(\sigma(X))(\sigma'(X;H) \mid u),
$$

which means that the parabolic subderivative $u \mapsto d^2 f(\sigma(X))(\sigma'(X;H) \mid u)$ is symmetric with respect to $\mathbf{P}^n_{\pm}(X;H)$.

Proof This follows by the same method as in Proposition [4.10.](#page-16-2)

As mentioned before, we can partition any vector $z \in \mathbb{R}^n$ into $z = (z_{\alpha_1}, \ldots, z_{\alpha_t}, z_{\beta})$ with $\alpha_i, i = 1, \ldots, t$, taken from (2.7) . Pick $i \in \{1, \ldots, t\}$ and recall from (2.9) that the index set $\alpha_i = \bigcup_{j=1}^{N_i} \beta_j^i$. This allows us to partition further z_{α_i} and z_{β} , respectively, into $z_{\alpha_i} = (z_{\beta_1^i}, \dots, z_{\beta_{N_i}^i})$ and $z_{\beta} = (z_{\beta_1^{t+1}}, \ldots, z_{\beta_{N_{t+1}}^{t+1}}, z_{\beta_{N_{t+1}+1}^{t+1}}),$ where $z_{\beta_j^i} \in \mathbb{R}^{|\beta_j^i|}$ for $i \in \{1, \ldots, t\}, j \in \{1, \ldots, N_i\},$ and $z_{\beta_k^{t+1}} \in \mathbb{R}^{|\beta_k^{t+1}|}$ for $k \in \{1, ..., N_{t+1}, N_{t+1} + 1\}$. In short, we can equivalently write z as

$$
\left(z_{\beta_1^1}, \ldots, z_{\beta_{N_1}^1}, \ldots, z_{\beta_1^t}, \ldots, z_{\beta_{N_t}^t}, z_{\beta_1^{t+1}}, \ldots, z_{\beta_{N_{t+1}}^{t+1}}, z_{\beta_{N_{t+1}+1}^{t+1}}\right),\tag{5.6}
$$

 \Box

where t, N_{t+1} , taken from (2.7) and (2.10) , and N_i , $i = 1, \ldots, t$, taken from (2.9) , stand for the number of distinct nonzero singular values of $X \in M_{m,n}$ and $U_{\hat{\beta}}^T H V_{\beta}$, and the number of distinct eigenvalues of $\frac{1}{2}(U_{\alpha_i}^THV_{\alpha_i} + V_{\alpha_i}^TH^TU_{\alpha_i}),$ respectively. Thus, the representation of z in [\(5.6\)](#page-20-0) is associated with the signed permutation matrices with representation

diag
$$
(B_1^1, \ldots, B_{N_1}^1, \ldots, B_1^t, \ldots, B_{N_t}^t, B_1^{t+1}, \ldots, B_{N_{t+1}}^{t+1}, B_{N_{t+1}+1}^{t+1})
$$

in $\mathbf{P}_{\pm}^n(X, H)$, where $B_j^i \in \mathbb{R}^{|\beta_j^i| \times |\beta_j^i|}$ is an signed permutation matrix for any $i = 1, \ldots, t, j =$ $1, \ldots, N_i$ and $B_k^{t+1} \in \mathbb{R}^{|\beta_k^{t+1}| \times |\beta_k^{t+1}|}$ is also an signed permutation matrix for any $k = 1, \ldots, N_{t+1}, N_{t+1}+1$ 1. Denote by $\mathbb{R}_{\downarrow}^{n}$ the set of all vectors (v_1, \ldots, v_n) such that $v_1 \geq \ldots \geq v_n$.

Proposition 5.5. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be lsc, convex, and absolutely symmetric. Assume that $Y \in \partial (f \circ \sigma)(X)$ and $H \in K_{f \circ \sigma}(X, Y)$. If f is parabolically regular at $\sigma(X)$ for $\sigma(Y)$, then the following assertions hold:

(i) There exists $\bar{z} \in \mathbb{R}^n$ with representation [\(5.6\)](#page-20-0), where $z_{\beta_j^i} \in \mathbb{R}_{\downarrow}^{|\beta_j^i|}$ \downarrow^{p_j} for $i \in \{1, \ldots, t\}, j \in$ $\{1,\ldots,N_i\}$ and $z_{\beta_k^{t+1}} \in \mathbb{R}_{\downarrow}^{|\beta_k^{t+1}|}$ $\bigcup_{k}^{\mu_{k}}$ \quad for $k \in \{1, ..., N_{t+1}, N_{t+1} + 1\}$, satisfying $d^2 f(\sigma(X) | \sigma(Y))(\sigma'(X;Y)) = d^2 f(\sigma(X))(\sigma'(X;Y) | \bar{z}) - \langle \sigma(Y), \bar{z} \rangle.$ (5.7)

(ii) There exists a matrix $\widehat{W} \in M_{m,n}$ such that $\sigma''(X; H, \widehat{W}) = \overline{z}$, where \overline{z} satisfies the above condition (i).

Proof Because $Y \in \partial(f \circ \sigma)(X)$ and $H \in K_{f \circ \sigma}(X, Y), \sigma(Y) \in \partial f(\sigma(X))$ and $\sigma'(X; H) \in$ $K_f(\sigma(X), \sigma(Y))$ by Proposition [4.2](#page-12-2) and Proposition [5.2,](#page-18-1) respectively. It follows from the parabolic regularity of f at $\sigma(X)$ for $\sigma(Y) \in \partial f(\sigma(X))$ that

$$
d^2 f(\sigma(X), \sigma(Y))(\sigma'(X; H)) = \inf_{z} \{d^2 f(\sigma(X))(\sigma'(X; H) \mid z) - \langle \sigma(Y), z \rangle\}.
$$

As explained above, there exists $\hat{z} \in \mathbb{R}^n$ with representation [\(5.6\)](#page-20-0) satisfying [\(5.7\)](#page-21-0). It is sufficient to show that the components of each $z_{\beta_i^i} \in \mathbb{R}^{|\beta_j^i|}$ for $i \in \{1, \ldots, t\}$, $j \in \{1, \ldots, N_i\}$ and $z_{\beta_k^{t+1}} \in \mathbb{R}^{|\beta_k^{t+1}|}$ for $k \in \{1, ..., N_{t+1}, N_{t+1}+1\}$ have nonincreasing order. We choose $|\beta_j^i| \times |\beta_j^i|$ signed permutation matrix B_j^i for any $i = 1, ..., t, j = 1, ..., N_i$ and $|\beta_k^{t+1}| \times |\beta_k^{t+1}|$ signed permutation matrix B_k^{t+1} for $k = 1, ..., N_{t+1}, N_{t+1} + 1$ such that $\bar{z}_{\beta_j^i} = B_j^i z_{\beta_j^i} \in \mathbb{R}_{\downarrow}^{|\beta_j^i|}$ $\frac{|\beta_j^i|}{\downarrow}$ and $\bar{z}_{\beta_k^{t+1}} = B_k^{t+1} z_{\beta_k^{t+1}} \in \mathbb{R}_{\downarrow}^{|\beta_k^{t+1}|}$ $↓^{\nu_k}$ '. Let

$$
Q_{\pm} := \text{diag}\left(B_1^1, \ldots, B_{N_1}^1, \ldots, B_1^t, \ldots, B_{N_t}^t, B_1^{t+1}, \ldots, B_{N_{t+1}}^{t+1}, B_{N_{t+1}+1}^{t+1}\right),\,
$$

then $Q_{\pm} \in \mathbf{P}_{\pm}^{n}(X, H)$. Set

$$
\bar{z} = \left(\bar{z}_{\beta_1^1}, \dots, \bar{z}_{\beta_{N_1}^1}, \dots, \bar{z}_{\beta_1^t}, \dots, \bar{z}_{\beta_{N_t}^t}, \bar{z}_{\beta_1^{t+1}}, \dots, \bar{z}_{\beta_{N_{t+1}}^{t+1}}, \bar{z}_{\beta_{N_{t+1}+1}^{t+1}}\right)
$$
(5.8)

and observe that $\bar{z} = Q_{\pm} \hat{z}$. We deduce from Proposition [5.4](#page-20-1) that

$$
d^2 f(\sigma(X))(\sigma'(X;H) \mid \bar{z}) = d^2 f(\sigma(X))(\sigma'(X;H) \mid \hat{z}). \tag{5.9}
$$

On the other hand, suppose that

$$
\left(\sigma(Y)_{\beta_1^1},\ldots,\sigma(Y)_{\beta_{N_1}^1},\ldots,\sigma(Y)_{\beta_1^t},\ldots,\sigma(Y)_{\beta_{N_t}^t},\sigma(Y)_{\beta_1^{t+1}},\ldots,\sigma(Y)_{\beta_{N_{t+1}}^{t+1}},\sigma(Y)_{\beta_{N_{t+1}}^{t+1}+1}\right)
$$

is a partition of the vector $\sigma(Y)$ corresponding to [\(5.6\)](#page-20-0) and observe that $\sigma(Y)_{\beta_j^i} \in \mathbb{R}_{\downarrow}^{|\beta_j^i|}$ $\downarrow^{\rho_{j+}}$ for $i \in \{1, ..., t\}, j \in \{1, ..., N_i\}$ and $\sigma(Y)_{\beta_k^{t+1}} \in \mathbb{R}_{\downarrow}^{|\beta_k^{t+1}|}$ $\bigcup_{k=1}^{\infty}$ for $k \in \{1, \ldots, N_{t+1}, N_{t+1} + 1\}$. It is not hard to get that

$$
\langle \sigma(Y), \hat{z} \rangle = \sum_{i=1}^{t} \sum_{j=1}^{N_i} \langle \sigma(Y)_{\beta_j^i}, z_{\beta_j^i} \rangle + \sum_{k=1}^{N_{t+1}+1} \langle \sigma(Y)_{\beta_k^{t+1}}, z_{\beta_k^{t+1}} \rangle
$$

$$
\leq \sum_{i=1}^{t} \sum_{j=1}^{N_i} \langle \sigma(Y)_{\beta_j^i}, \bar{z}_{\beta_j^i} \rangle + \sum_{k=1}^{N_{t+1}+1} \langle \sigma(Y)_{\beta_k^{t+1}}, \bar{z}_{\beta_k^{t+1}} \rangle
$$

$$
= \langle \sigma(Y), \bar{z} \rangle.
$$

This, together with [\(5.9\)](#page-21-1) and \hat{z} with representation [\(5.6\)](#page-20-0) satisfying [\(5.7\)](#page-21-0), justifies

$$
d^2 f(\sigma(X) | \sigma(Y))(\sigma'(X;Y)) \ge d^2 f(\sigma(X))(\sigma'(X;Y) | \bar{z}) - \langle \sigma(Y), \bar{z} \rangle.
$$

We conclude from the inequality [\(2.1\)](#page-3-0) that [\(5.7\)](#page-21-0) holds for \bar{z} .

Turning now to proof (ii), pick the vector $\bar{z} \in \mathbb{R}^n$ from [\(5.8\)](#page-21-2). We can equivalently write via the index sets α_i , $i = 1, \ldots, t$, from (2.7) that

$$
\bar{z} = (\bar{z}_{\alpha_1}, \dots, \bar{z}_{\alpha_t}, \bar{z}_{\beta}) \text{ with } \bar{z}_{\alpha_i} = \left(\bar{z}_{\beta_1^i}, \dots, \bar{z}_{\beta_{N_i}^i}\right) \text{ and } \bar{z}_{\beta} = \left(\bar{z}_{\beta_1^{t+1}}, \dots, \bar{z}_{\beta_{N_{t+1}}^{t+1}}, \bar{z}_{\beta_{N_{t+1}+1}^{t+1}}\right),
$$

where $\bar{z}_{\beta_j^i} \in \mathbb{R}^{|\beta_j^i|}_{\downarrow}$ $\downarrow^{\beta_j^i}$ for $i \in \{1, ..., t\}, j \in \{1, ..., N_i\}$, and $\bar{z}_{\beta_k^{t+1}} \in \mathbb{R}^{\vert \beta_k^{t+1} \vert}$ 1}. It suffices to prove that there exists a matrix $\widehat{W} \in M_{m,n}$ satisfying $\sigma''(X; H, \widehat{W}) = \overline{z}$. Pick the $\big\downarrow^{p_k}$ for $k \in \{1, \ldots, N_{t+1}, N_{t+1}+\}$ $|\alpha_i| \times |\alpha_i|$ matrix $Q^i \in \mathcal{O}^{|\alpha_i|} \Big(\frac{1}{2} \big(U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i} \big) \Big), i = 1, \ldots, t$, and take the two matrices $(Q_{\hat{\beta}\hat{\beta}}, \widehat{Q}_{\beta\beta}) \in \mathcal{O}^{|\hat{\beta}|, |\beta|} (U_{\hat{\beta}}^T H V_{\beta}).$ Consider the $|\alpha| \times |\alpha|$ block diagonal matrix

$$
A_{\alpha\alpha} = \text{diag}\left(Q^1 \text{diag}(\bar{z}_{\alpha_1})(Q^1)^T, \dots, Q^t \text{diag}(\bar{z}_{\alpha_t})(Q^t)^T\right),\tag{5.10}
$$

and the $|\hat{\beta}| \times |\beta|$ matrix

$$
A_{\hat{\beta}\beta} = Q_{\hat{\beta}\hat{\beta}} \left(\text{diag}(\bar{z}_{\beta}) \right)_{\hat{\beta}\beta} \hat{Q}_{\beta\beta}^T. \tag{5.11}
$$

It is clear that $A_{\hat{\beta}\beta} = Q_{\hat{\beta}\beta} (\text{diag}(\bar{z}_{\beta}))_{\beta\beta} \hat{Q}_{\beta\beta}^T$. We claim that there exists a matrix $\widehat{W} \in M_{m,n}$ satisfying

$$
P_{\alpha_i}^T \mathcal{B}(\widehat{W}) P_{\alpha_i} = P_{\alpha_i}^T \Big[2\mathcal{B}(H) P_{\alpha_i}^c (A_s - \mu_i I)^{-1} P_{\alpha_i}^{c} {}^T \mathcal{B}(H) + P_{\alpha} A_{\alpha \alpha} P_{\alpha}^T \Big] P_{\alpha_i} \tag{5.12}
$$

and

$$
U_{\hat{\beta}}^T \widehat{W} V_{\beta} = U_{\hat{\beta}}^T \Big[2H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H + U_{\hat{\beta}} A_{\hat{\beta}\beta} V_{\beta}^T \Big] V_{\beta}.
$$
 (5.13)

Indeed, let $W = diag(W_{\alpha_1 \alpha_1}, \ldots, W_{\alpha_t \alpha_t}, W_{\hat{\beta}\beta}) \in M_{m,n}$ satisfy that

$$
W_{\alpha_i\alpha_i} + W_{\alpha_i\alpha_i}^T = P_{\alpha_i}^T \Big[2\mathcal{B}(H) P_{\alpha_i}^c (A_s - \mu_i I)^{-1} P_{\alpha_i}^c{}^T \mathcal{B}(H) + P_{\alpha} A_{\alpha\alpha} P_{\alpha}^T \Big] P_{\alpha_i},\tag{5.14}
$$

for all $i = 1, \ldots, t$, and

$$
W_{\hat{\beta}\beta} = U_{\hat{\beta}}^T \left[2HV_{\alpha} \Sigma_{\alpha}^{-1}(X)U_{\alpha}^T H + U_{\hat{\beta}} A_{\hat{\beta}\beta} V_{\beta}^T \right] V_{\beta}.
$$
 (5.15)

Set $\widehat{W} = UWV^{T}$. Then by some elementary calculations, we can obtain that

$$
\mathcal{B}(\widehat{W}) = P_{\alpha} [U_{\alpha}^T \widehat{W} V_{\alpha} + V_{\alpha}^T \widehat{W}^T U_{\alpha}] P_{\alpha}^T = P_{\alpha} [W_{\alpha \alpha} + W_{\alpha \alpha}^T] P_{\alpha}^T
$$

and

$$
W_{\hat{\beta}\beta} = U_{\hat{\beta}}^T U W V^T V_{\beta} = U_{\hat{\beta}}^T \widehat{W} V_{\beta}.
$$

This, together with (5.14) and (5.15) , justifies that (5.12) and (5.13) hold. Next we show that $\sigma''(X; H, W) = \overline{z}$. For any $s \in \{1, \ldots, n\}$, we first consider the case $s \in \alpha$. There exists $i \in$ $\{1, \ldots, t\}$ and $j \in \{1, \ldots, N_i\}$ such that $s \in \alpha_i$ and $l_s \in \beta_j^i$. It follows from (2.11) that

$$
\sigma_s''(X; H, \widehat{W}) = \lambda_{\tilde{l}_s} \left(\left(Q_{\beta_j^i}^i \right)^T P_{\alpha_i}^T \left[\mathcal{B}(\widehat{W}) - 2\mathcal{B}(H) P_{\alpha_i}^c (A_s - \mu_i I)^{-1} P_{\alpha_i}^c{}^T \mathcal{B}(H) \right] P_{\alpha_i} Q_{\beta_j^i}^i \right)
$$

\n
$$
= \lambda_{\tilde{l}_s} \left(\left(Q_{\beta_j^i}^i \right)^T \left[P_{\alpha_i}^T P_{\alpha} A_{\alpha \alpha} P_{\alpha}^T P_{\alpha_i} \right] Q_{\beta_j^i}^i \right)
$$

\n
$$
= \lambda_{\tilde{l}_s} \left(\left(Q_{\beta_j^i}^i \right)^T Q^i \text{diag}(\bar{z}_{\alpha_i}) (Q^i)^T Q_{\beta_j^i}^i \right)
$$

\n
$$
= \lambda_{\tilde{l}_s} \left(\text{diag} \left(\bar{z}_{\beta_j^i} \right) \right) = \left(\bar{z}_{\beta_j^i} \right)_{\tilde{l}_s} = \bar{z}_s.
$$

We now turn to the case $s \in \beta$. If $l_s(X) \in \beta_k^{t+1}$ for some $k \in \{1, ..., N_{t+1}\}$, we deduce from (2.12) that

$$
\sigma_s''(X;H,\widehat{W}) = \frac{1}{2}\lambda_{\tilde{l}_s}\left(\left(Q_{\hat{\beta}\beta_k^{t+1}}\right)^T \left[U_{\hat{\beta}}^T\widehat{W}V_{\beta} - 2U_{\hat{\beta}}^THV_{\alpha}\Sigma_{\alpha}^{-1}(X_0)U_{\alpha}^THV_{\beta}\right]\widehat{Q}_{\beta\beta_k^{t+1}} \n+ \left(\widehat{Q}_{\beta\beta_k^{t+1}}\right)^T \left[U_{\hat{\beta}}^T\widehat{W}V_{\beta} - 2U_{\hat{\beta}}^THV_{\alpha}\Sigma_{\alpha}^{-1}(X_0)U_{\alpha}^THV_{\beta}\right]^T Q_{\hat{\beta}\beta_k^{t+1}}\right) \n= \frac{1}{2}\lambda_{\tilde{l}_s}\left(\left(Q_{\hat{\beta}\beta_k^{t+1}}\right)^T \left[U_{\hat{\beta}}^TU_{\hat{\beta}}A_{\hat{\beta}\beta}V_{\beta}^TV_{\beta}\right]\widehat{Q}_{\beta\beta_k^{t+1}} + \left(\widehat{Q}_{\beta\beta_k^{t+1}}\right)^T \left[U_{\hat{\beta}}^TU_{\hat{\beta}}A_{\hat{\beta}\beta}V_{\beta}^TV_{\beta}\right]^T Q_{\hat{\beta}\beta_k^{t+1}}\right) \n= \frac{1}{2}\lambda_{\tilde{l}_s}\left(\left(Q_{\hat{\beta}\beta_k^{t+1}}\right)^T Q_{\hat{\beta}\beta}(\text{diag}(\bar{z}_{\beta}))_{\beta\beta}\widehat{Q}_{\beta\beta_k^{t+1}}^T + \left(\widehat{Q}_{\beta\beta_k^{t+1}}\right)^T \widehat{Q}_{\beta\beta}(\text{diag}(\bar{z}_{\beta}))_{\beta\beta}Q_{\hat{\beta}\beta}^T Q_{\hat{\beta}\beta_k^{t+1}}\right) \n= \lambda_{\tilde{l}_s}\left(\text{diag}\left(\bar{z}_{\beta_k^{t+1}}\right)\right) = \left(\bar{z}_{\beta_k^{t+1}}\right)_{\tilde{l}_s} = \bar{z}_s.
$$

On the other hand, if $l_s(X) \in \beta^{t+1}_{N_{t+1}+1}$, we conclude from [\(2.13\)](#page-6-6) that

$$
\sigma_s''(X;H,\widehat{W}) = \sigma_{\tilde{l}_s} \left(\begin{bmatrix} Q_{\hat{\beta}\beta_{N_{t+1}+1}}^{t+1} & Q_{\hat{\beta}\beta_0} \end{bmatrix}^T \begin{bmatrix} U_{\hat{\beta}}^T \widehat{W} V_{\beta} - 2U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X_0) U_{\alpha}^T H V_{\beta} \end{bmatrix} \widehat{Q}_{\beta\beta_{N_{t+1}+1}} \right)
$$

\n
$$
= \sigma_{\tilde{l}_s} \left(\begin{bmatrix} Q_{\hat{\beta}\beta_{N_{t+1}+1}}^{t+1} & Q_{\hat{\beta}\beta_0} \end{bmatrix}^T \begin{bmatrix} U_{\hat{\beta}}^T U_{\hat{\beta}} A_{\hat{\beta}\beta} V_{\beta}^T V_{\beta} \end{bmatrix} \widehat{Q}_{\beta\beta_{N_{t+1}+1}} \right)
$$

\n
$$
= \sigma_{\tilde{l}_s} \left(\begin{bmatrix} Q_{\hat{\beta}\beta_{N_{t+1}+1}}^{t+1} & Q_{\hat{\beta}\beta_0} \end{bmatrix}^T Q_{\hat{\beta}\hat{\beta}} (\text{diag}(\bar{z}_{\beta}))_{\hat{\beta}\beta} \widehat{Q}_{\beta\beta}^T \widehat{Q}_{\beta\beta_{N_{t+1}+1}} \right)
$$

\n
$$
= \sigma_{\tilde{l}_s} \left(\left(\text{diag} \left(\bar{z}_{\beta_{N_{t+1}+1}}^{t+1} \right) \right)_{(\beta_{N_{t+1}+1}^{t+1})} \widehat{Q}_{\beta\beta_{N_{t+1}+1}}^{t+1} \right) = \left(\bar{z}_{\beta_{N_{t+1}+1}^{t+1}}^{t+1} \right)_{\tilde{l}_s} = \bar{z}_s.
$$

In summary, we can find a matrix $\widehat{W} \in M_{m,n}$ satisfying simultaneously [\(5.12\)](#page-22-2) and [\(5.13\)](#page-22-3) such that $\sigma''(X; H, \widehat{W}) = \overline{z}$, where \overline{z} comes from (i). \Box

Finally, we present a exact formula for second subderivative of orthogonally invariant matrix functions.

Theorem 5.6. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be an absolutely symmetric function, and be lsc, convex, and locally Lipschitz continuous relative to its domain. Assume that $\mu_1 > \cdots > \mu_t$ are the distinct nonzero singular values of $X \in \mathbb{M}_{m,n}$ and $Y \in \partial(f \circ \sigma)(X)$, and that f is parabolically epidifferentiable at $\sigma(X)$ and parabolically regular at $\sigma(X)$ for $\sigma(Y)$. Then $f \circ \sigma$ is parabolically regular at X for Y, and for any $H \in K_{f \circ \sigma}(X, Y)$ we have

$$
d^{2}(f \circ \sigma)(X \mid Y)(H) = d^{2}f(\sigma(X) \mid \sigma(Y))(\sigma'(X; H))
$$

+
$$
2 \sum_{i=1}^{t} \left\langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T}B(H)P_{\alpha_{i}}^{c}(\mu_{i}I - \Lambda_{\alpha_{i}})^{-1}P_{\alpha_{i}}^{c}{}^{T}B(H)P_{\alpha_{i}} \right\rangle
$$

+
$$
2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^{T}HV_{\alpha}\Sigma_{\alpha}^{-1}(X)U_{\alpha}^{T}HV_{\beta} \right\rangle,
$$

where $(U, V) \in \mathcal{O}^{m,n}(X) \cap \mathcal{O}^{m,n}(Y)$, the columns of P_{α_i} form an orthonormal basis of eigenvectors of $\mathcal{B}(X)$ associated with μ_i , $P^c_{\alpha_i}$ is the submatrix of P obtained by removing all the columns of P_{α_i} , and $\Lambda_{\alpha_i} \in \mathbb{R}^{(m+n-|\alpha_i|)\times(m+n-|\alpha_i|)}$ is a diagonal matrix whose diagonal elements are the eigenvalues of $P^T \mathcal{B}(X)P$ that are not equal to μ_i for all $i \in \{1, \ldots, t\}.$

Proof Set $F := f \circ \sigma$. Since $H \in K_F(X, Y)$, it follows from Proposition [5.2](#page-18-1) that there exists

$$
\bar{Q}^i \in \mathcal{O}^{|\alpha_i|}(\Sigma(Y)_{\alpha_i \alpha_i}) \bigcap \mathcal{O}^{|\alpha_i|}(\frac{1}{2}(U_{\alpha_i}^THV_{\alpha_i} + V_{\alpha_i}^TH^TU_{\alpha_i}))\bigg),
$$

and

$$
(\bar{Q}_{\hat{\beta}\hat{\beta}}, \hat{\bar{Q}}_{\beta\beta}) \in \mathcal{O}^{|\hat{\beta}|, |\beta|}(\Sigma(Y)_{\hat{\beta}\beta}) \bigcap \mathcal{O}^{|\hat{\beta}|, |\beta|}(U_{\hat{\beta}}^T H V_{\beta})
$$

 \overline{a}

for any $i \in \{1, ..., t\}$ and $(U, V) \in \mathcal{O}^{m,n}(X) \cap \mathcal{O}^{m,n}(Y)$ such that

$$
\Sigma(Y)_{\alpha_i \alpha_i} = \overline{Q}^i \Sigma(Y)_{\alpha_i \alpha_i} (\overline{Q}^i)^T,
$$

$$
\frac{1}{2} (U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i}) = \overline{Q}^i \text{diag}\left(\lambda \left(\frac{1}{2} (U_{\alpha_i}^T H V_{\alpha_i} + V_{\alpha_i}^T H^T U_{\alpha_i})\right)\right) (\overline{Q}^i)^T,
$$

and

$$
\Sigma(Y)_{\hat{\beta}\beta} = \bar{Q}_{\hat{\beta}\hat{\beta}} \Sigma(Y)_{\hat{\beta}\beta} (\hat{\bar{Q}}_{\beta\beta})^T,
$$

$$
U_{\hat{\beta}}^T H V_{\beta} = \bar{Q}_{\hat{\beta}\hat{\beta}} \Sigma (U_{\hat{\beta}}^T H V_{\beta}) (\widehat{\bar{Q}}_{\beta\beta})^T.
$$

Consider the $m \times n$ block diagonal matrix

$$
\bar{A} = \text{diag}\left(\bar{Q}^1 \text{diag}(\bar{z}_{\alpha_1})(\bar{Q}^1)^T, \dots, \bar{Q}^t \text{diag}(\bar{z}_{\alpha_t})(\bar{Q}^t)^T, \bar{Q}_{\hat{\beta}\hat{\beta}} (\text{diag}(\bar{z}_{\beta}))_{\hat{\beta}\beta} \widehat{\bar{Q}}_{\beta\beta}^T\right),\right)
$$

where \bar{Q}^i replace the matrices Q^i in the definition of the matrix $A_{\alpha\alpha}$ in [\(5.10\)](#page-22-4) for $i = 1, \ldots, t$, and $\bar{Q}_{\hat{\beta}\hat{\beta}}, \bar{Q}_{\beta\beta}$ replace the matrices $Q_{\hat{\beta}\hat{\beta}}, \widehat{Q}_{\beta\beta}$ in the definition of the matrix $A_{\hat{\beta}\beta}$ in [\(5.11\)](#page-22-5), respectively. It is not hard to see that the same conclusion can be achieved as the one in Proposition [5.5](#page-21-3) for the above \overline{A} . We can deduce from Proposition [5.5](#page-21-3) that there exists $\overline{z} \in \mathbb{R}^n$ with representation [\(5.6\)](#page-20-0) satisfying [\(5.7\)](#page-21-0), where $z_{\beta_j^i} \in \mathbb{R}_{\downarrow}^{|\beta_j^i|}$ $\downarrow^{\beta_j^i}$ for $i \in \{1, ..., t\}, j \in \{1, ..., N_i\}$ and $z_{\beta_k^{t+1}} \in \mathbb{R}_{\downarrow}^{|\beta_k^{t+1}|}$ $\downarrow^{\scriptscriptstyle{P_k} \cdots \scriptscriptstyle{P_k}}$ for $k \in$ $\{1,\ldots,N_{t+1},N_{t+1}+1\}$, and further there exists a matrix $\widehat{W} \in \mathbb{M}_{m,n}$ such that $\sigma''(X;H,\widehat{W})=\overline{z}$. This, together with $Y = U \Sigma(Y) V^T$ and the fact that $\left\langle \Sigma(Y)_{\alpha_i \alpha_i}, \frac{1}{2} (V_{\alpha_i}^T \widehat{W}^T U_{\alpha_i} - U_{\alpha_i}^T \widehat{W} V_{\alpha_i}) \right\rangle = 0$ for any $i = 1, \ldots, t$, we obtain that

$$
d^{2} f(\sigma(X))(\sigma'(X;H) | \sigma''(X;H,\widehat{W})) - \langle Y, \widehat{W} \rangle
$$

\n
$$
= d^{2} f(\sigma(X))(\sigma'(X;H) | \bar{z}) - \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, U_{\alpha_{i}}^{T} \widehat{W} V_{\alpha_{i}} \rangle - \langle \Sigma(Y)_{\hat{\beta}\beta}, U_{\hat{\beta}}^{T} \widehat{W} V_{\beta} \rangle
$$

\n
$$
= d^{2} f(\sigma(X) | \sigma(Y))(\sigma'(X;Y)) + \langle \sigma(Y), \bar{z} \rangle - \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} B(\widehat{W}) P_{\alpha_{i}} \rangle - \langle \Sigma(Y)_{\hat{\beta}\beta}, U_{\hat{\beta}}^{T} \widehat{W} V_{\beta} \rangle
$$

\n
$$
= d^{2} f(\sigma(X) | \sigma(Y))(\sigma'(X;Y)) + \langle \sigma(Y), \bar{z} \rangle
$$

\n
$$
- \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} [2B(H) P_{\alpha_{i}}^{c}(A_{s} - \mu_{i} I)^{-1} P_{\alpha_{i}}^{c} {}^{T} B(H) + P_{\alpha} \bar{A}_{\alpha\alpha} P_{\alpha}^{T} P_{\alpha_{i}} \rangle
$$

\n
$$
- \langle \Sigma(Y)_{\hat{\beta}\beta}, U_{\hat{\beta}}^{T} [2H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^{T} H + U_{\hat{\beta}} \bar{A}_{\hat{\beta}\beta} V_{\beta}^{T} P_{\beta} \rangle
$$

\n
$$
= d^{2} f(\sigma(X) | \sigma(Y))(\sigma'(X;H)) + \langle \sigma(Y), \bar{z} \rangle
$$

\n
$$
+ 2 \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} B(H) P_{\alpha_{i}}^{c} (\mu_{i} I - A_{\alpha_{i}})^{-1} P_{\alpha_{i}}^{c} {}^{T} B(H) P_{\alpha_{i}} \rangle + 2 \langle \Sigma(Y)_{\hat{\beta}\beta}, -U
$$

From the definition of \overline{A} and the representation of \overline{z} , we deduce that

$$
\langle \sigma(Y), \bar{z} \rangle = \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_i \alpha_i}, \text{diag}(\bar{z}_{\alpha_i}) \rangle + \langle \Sigma(Y)_{\hat{\beta}\beta}, \left(\text{diag}(\bar{z}_{\beta}) \right)_{\hat{\beta}\beta} \rangle
$$

\n
$$
= \sum_{i=1}^{t} \langle \bar{Q}^i \Sigma(Y)_{\alpha_i \alpha_i} (\bar{Q}^i)^T, \bar{Q}^i \text{diag}(\bar{z}_{\alpha_i}) (\bar{Q}^i)^T \rangle + \langle \bar{Q}_{\hat{\beta}\hat{\beta}} \Sigma(Y)_{\hat{\beta}\beta} \hat{\bar{Q}}^T_{\beta\beta}, \bar{Q}_{\hat{\beta}\hat{\beta}} \left(\text{diag}(\bar{z}_{\beta}) \right)_{\hat{\beta}\beta} \hat{\bar{Q}}^T_{\beta\beta} \rangle
$$

\n
$$
= \sum_{i=1}^{t} \langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \Big[P_{\alpha} \bar{A}_{\alpha \alpha} P_{\alpha}^T \Big] P_{\alpha_i} \rangle + \langle \Sigma(Y)_{\hat{\beta}\beta}, U_{\hat{\beta}}^T \Big[U_{\hat{\beta}} \bar{A}_{\hat{\beta}\beta} V_{\beta}^T \Big] V_{\beta} \rangle.
$$

Thus, from (5.1) , Proposition [2.1](#page-3-4) and (5.2) we get

$$
d^{2} f(\sigma(X) | \sigma(Y))(\sigma'(X;H)) + 2 \sum_{i=1}^{t} \left\langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\mu_{i}I - \Lambda_{\alpha_{i}})^{-1} P_{\alpha_{i}}^{c}^{T} \mathcal{B}(H) P_{\alpha_{i}} \right\rangle
$$

+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^{T} H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^{T} H V_{\beta} \right\rangle
\n
$$
\leq d^{2} F(X | Y) (H)
$$

\n
$$
\leq \inf_{W \in \mathbb{M}_{m,n}} \left\{ d^{2} F(X) (H | W) - \langle Y, W \rangle \right\}
$$

\n
$$
= \inf_{W \in \mathbb{M}_{m,n}} \left\{ d^{2} f(\sigma(X)) (\sigma'(X;H) | \sigma''(X;H,W)) - \langle Y, W \rangle \right\}
$$

\n
$$
\leq d^{2} f(\sigma(X)) (\sigma'(X;H) | \sigma''(X;H,\widehat{W})) - \langle Y, \widehat{W} \rangle
$$

\n
$$
= d^{2} f(\sigma(X) | \sigma(Y)) (\sigma'(X;H)) + 2 \sum_{i=1}^{t} \left\langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\mu_{i}I - \Lambda_{\alpha_{i}})^{-1} P_{\alpha_{i}}^{c}^{T} \mathcal{B}(H) P_{\alpha_{i}} \right\rangle
$$

\n
$$
+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^{T} H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^{T} H V_{\beta} \right\rangle.
$$

It follows immediately that $d^2F(X | Y)(H) = \inf_{W \in \mathbb{M}_{m,n}} \{d^2F(X)(H | W) - \langle Y, W \rangle\}$ for any $H \in$ $K_F(X, Y)$, which implies that F is parabolically regular at X for Y. Moreover, this verifies the proposed formula for the second subderivative of F at X for Y for any $H \in K_F(X, Y)$, thereby completing the proof. completing the proof.

Combining Theorem [5.6](#page-23-0) with [\[15,](#page-27-8) Theorem 3.8], we provide sufficient conditions for twice epidifferentiability of orthogonally invariant matrix function.

Proposition 5.7. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be an absolutely symmetric function, and be lsc, convex, and locally Lipschitz continuous relative to its domain. Assume that $Y \in \partial (f \circ \sigma)(X)$ and f is parabolically epi-differentiable at $\sigma(X)$ and parabolically regular at $\sigma(X)$ for $\sigma(Y)$. Then $f \circ \sigma$ is twice epi-differentiable at X for Y .

As an application, we present second-order optimality conditions for the following optimization problem according to [\[18,](#page-27-15) Theorem 13.24]. Given a twice differentiable function $\psi: \mathbb{M}_{m,n} \to \mathbb{R}$ and an absolutely symmetric function $f: \mathbb{R}^n \to [-\infty, +\infty]$, consider the optimization problem

$$
\min_{X \in \mathbb{M}_{m,n}} \psi(X) + (f \circ \sigma)(X).
$$

From [\[18,](#page-27-15) Exercise 13.18], we can derive the following conclusion.

Theorem 5.8. Let $f: \mathbb{R}^n \to [-\infty, +\infty]$ be an absolutely symmetric function, and be lsc, convex, and locally Lipschitz continuous relative to its domain. Assume that X_0 is a feasible solution to (P) with f being parabolically epi-differentiable at $\sigma(X_0)$ and parabolically regular at $\sigma(X_0)$ for $\sigma(Y)$ and $Y \in \partial (f \circ \sigma)(X_0)$. If $-\nabla \psi(X_0) \in \partial (f \circ \sigma)(X_0)$, then the following second-order optimality conditions for (P) hold.

(i) If X_0 is a local minimizer of (P) , then the second-order necessary condition

$$
\nabla^2 \psi(X_0)(H, H) + d^2 f(\sigma(X) | \sigma(Y)) (\sigma'(X; H)) + 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c(\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H) P_{\alpha_i} \right\rangle
$$

+
$$
2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H V_{\beta} \right\rangle \geq 0
$$

holds for all $H \in K_{f \circ \sigma}(X_0, -\nabla \psi(X_0)).$

(ii) Having the second-order sufficient condition

$$
\nabla^2 \psi(X_0)(H, H) + d^2 f(\sigma(X) | \sigma(Y)) (\sigma'(X; H)) + 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c(\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}{}^T \mathcal{B}(H) P_{\alpha_i} \right\rangle
$$

+
$$
2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H V_{\beta} \right\rangle > 0
$$

holds for all $H \in K_{f \circ \sigma}(X_0, -\nabla \psi(X_0))$ is equivalent to having the existence $\epsilon > 0$ and $c > 0$ such that

 $\psi(X) + (f \circ \sigma)(X) \geq \psi(X_0) + (f \circ \sigma)(X_0) + c||X - X_0||^2$ when $X \in \mathbb{B}_{\epsilon}(X_0)$.

In particular, when absolutely symmetric function f is a polyhedral function, whose epigraph is a polyhedral convex set, we obtain from $[18,$ Exercise 13.61 and $[15,$ Example 3.2 that f is parabolically epi-differentiable and parabolically regular at $\sigma(X)$. Moreover, it follows from Theorem [5.6](#page-23-0) and [\[15,](#page-27-8) Proposition 3.4] that dom $d^2(f \circ \sigma)(X | Y) = K_{f \circ \sigma}(X, Y)$. Further, we can conclude the following result from [\[18,](#page-27-15) Proposition 13.9].

Corollary 5.9. Let an absolutely symmetric function $f: \mathbb{R}^n \to [-\infty, +\infty]$ be polyhedral. Assume that $\mu_1 > \cdots > \mu_t$ are the distinct nonzero singular values of $X \in \mathbb{M}_{m,n}$ and that $Y \in \partial (f \circ \sigma)(X)$. Then for any $H \in K_{f \circ \sigma}(X, Y)$, we have

$$
d^{2}(f \circ \sigma)(X \mid Y)(H) = \delta_{K_{f \circ \sigma}(X,Y)}(H) + 2\sum_{i=1}^{t} \left\langle \Sigma(Y)_{\alpha_{i}\alpha_{i}}, P_{\alpha_{i}}^{T} \mathcal{B}(H) P_{\alpha_{i}}^{c} (\mu_{i}I - \Lambda_{\alpha_{i}})^{-1} P_{\alpha_{i}}^{c}{}^{T} \mathcal{B}(H) P_{\alpha_{i}} \right\rangle
$$

+
$$
2\left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^{T} H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^{T} H V_{\beta} \right\rangle,
$$

where $(U, V) \in \mathcal{O}^{m,n}(X) \cap \mathcal{O}^{m,n}(Y)$, the columns of P_{α_i} form an orthonormal basis of eigenvectors of $\mathcal{B}(X)$ associated with μ_i , $P^c_{\alpha_i}$ is the submatrix of P obtained by removing all the columns of P_{α_i} , and $\Lambda_{\alpha_i} \in \mathbb{R}^{(m+n-|\alpha_i|)\times(m+n-|\alpha_i|)}$ is a diagonal matrix whose diagonal elements are the eigenvalues of $P^T \mathcal{B}(X)P$ that are not equal to μ_i for all $i \in \{1, \ldots, t\}.$

Remark 5.10. If the absolutely symmetric function f is ℓ_1 -norm, then orthogonally invariant matrix function $f \circ \sigma$ is nuclear norm. The above corollary gives the second subderivative of nuclear norm as that in Corollary [3.6](#page-10-3) using a different approach.

In what follows, we present second-order optimality conditions for the optimization problem (P) with f being polyhedral.

Corollary 5.11. Assume that X_0 is a feasible solution to (P) with f being polyhedral. If $-\nabla \psi(X_0) \in$ $\partial (f \circ \sigma)(X_0)$, then the following second-order optimality conditions for (P) hold.

(i) If X_0 is a local minimizer of (P) , then the second-order necessary condition

$$
\nabla^2 \psi(X_0)(H, H) + 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}^T \mathcal{B}(H) P_{\alpha_i} \right\rangle
$$

+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H V_{\beta} \right\rangle \ge 0

holds for all $H \in K_{f \circ \sigma}(X_0, -\nabla \psi(X_0)).$

(ii) Having the second-order sufficient condition

$$
\nabla^2 \psi(X_0)(H, H) + 2 \sum_{i=1}^t \left\langle \Sigma(Y)_{\alpha_i \alpha_i}, P_{\alpha_i}^T \mathcal{B}(H) P_{\alpha_i}^c (\mu_i I - \Lambda_{\alpha_i})^{-1} P_{\alpha_i}^{c}^T \mathcal{B}(H) P_{\alpha_i} \right\rangle
$$

+ 2 \left\langle \Sigma(Y)_{\hat{\beta}\beta}, -U_{\hat{\beta}}^T H V_{\alpha} \Sigma_{\alpha}^{-1}(X) U_{\alpha}^T H V_{\beta} \right\rangle > 0 (5.16)

holds for all $H \in K_{f \circ \sigma}(X_0, -\nabla \psi(X_0))$ is equivalent to having the existence $\epsilon > 0$ and $c > 0$ such that

$$
\psi(X) + (f \circ \sigma)(X) \ge \psi(X_0) + (f \circ \sigma)(X_0) + c||X - X_0||^2 \text{ when } X \in \mathbb{B}_{\epsilon}(X_0).
$$

In the case of f being polyhedral, the second-order sufficient condition (5.16) is consistent with the "no-gap" second order sufficient condition in [\[2,](#page-27-7) Theorem 3.5], and we have provided the specific expression for the second term in the latter.

6 Conclusion

The primary objective of this work was to explore several second-order properties of orthogonally invariant matrix functions, with a focus on parabolic epi-differentiability, parabolic regularity, and twice epi-differentiability. Our results are based on the concept of metric subregularity constraint qualification, which, in this context, is automatically satisfied. For a convex orthogonally invariant matrix function, we derive the exact formula for its second subderivatives and establish sufficient conditions for twice epi-differentiability. Furthermore, we present second-order optimality conditions for a class of matrix optimization problems. In particular, inspired by Torki [\[20\]](#page-27-12), we adopt a different approach to calculate the first- and second-order epi-derivatives of the nuclear norm.

Acknowledgements

The research of the second author was supported by the National Natural Sciences Grant of China (No.11701126). The research of the third author was supported by the National Natural Sciences Grant of China (No. 11871182).

References

- [1] Bonnans, J.F., Shapiro, A.: Perturbation Analysis of Optimization Problems. Springer, New York (2000)
- [2] Cui, Y., Ding, C. and Zhao, X.: Quadratic Growth Conditions for Convex Matrix Optimization Problems Associated with Spectral Functions. SIAM J. Optim. 27(4): 2332-2355 (2017)
- [3] Daniilidis, A., Lewis, A., Malick, J. and Sendov, H.: Prox-Regularity of Spectral Functions and Spectral Sets. J. Convex Anal. 15(3): 547-560 (2008)
- [4] Ding, C.: Variational Analysis of the Ky Fan k-norm. Set-Valued Var. Anal. 25: 265-296 (2017)
- [5] Ding, C., Sun, D. and Toh, K.C.: An Introduction to a Class of Matrix Cone Programming. Math. Program. 144(1-2): 141-179 (2014)
- [6] Hiriart-Urruty, J.-B., Lemaréchal, C.: Fundamentals of Convex Analysis. Springer, Berlin (2001)
- [7] Horn, R.A., Johnson, C.R.: Matrix Analysis (2nd ed.). Cambridge university press, Cambridge (2012)
- [8] Lewis, A.S.: Nonsmooth Analysis of Eigenvalues. Math. Program. 84(1): 1-24 (1999)
- [9] Lewis, A.S.: The Convex Analysis of Unitarily Invariant Matrix Functions. J. Convex Anal. 2(1–2): 173-183 (1995)
- [10] Lewis, A.S., Sendov, H.S.: Nonsmooth Analysis of Singular Values. Part I: Theory. Set-Valued Anal. 13: 213-241 (2005)
- [11] Lewis, A.S., Sendov, H.S.: Nonsmooth Analysis of Singular Values. Part II: Applications. Set-Valued Anal. 13: 243-264 (2005)
- [12] Mathias, R.: The Spectral Norm of a Nonnegative Matrix. Linear Algebra Appl. 139: 269-284 (1990)
- [13] Mohammadi, A., Mordukhovich, B.S. and Sarabi, M.E.: Variational Analysis of Composite Models with Applications to Continuous Optimization. Math. Oper. Res. 47(1): 397–426 (2021)
- [14] Mohammadi, A., Sarabi, E.: Parabolic Regularity of Spectral Functions. Math. Oper. Res. (2024) https://doi.org/10.1287/moor.2023.0010
- [15] Mohammadi, A., Sarabi, M.E.: Twice Epi-Differentiability of Extended-Real-Valued Functions with Applications in Composite Optimization. SIAM J. Optim. 30: 2379–2409 (2020)
- [16] Rockafellar, R.T.: Convex Analysis. Princeton University Press, Princeton (1970)
- [17] Rockafellar, R.T.: First- and Second-Order Epi-Differentiability in Nonlinear Programming. Trans. Amer. Math. Soc. 307(1): 75-108 (1988)
- [18] Rockafellar, R.T., Wets, R.J.-B.: Variational Analysis. Springer, Berlin (1998)
- [19] Torki, M.: First-and Second-Order Epi-Differentiability in Eigenvalue Optimization. J. Math. Anal. Appl. 234(2): 391-416 (1999)
- [20] Torki, M.: Second-Order Directional Derivatives of All Eigenvalues of a Symmetric Matrix. Nonlinear Anal. 46(8): 1133-1150 (2001)
- [21] Vandenberghe, L., Boyd, S.: Semidefinite Programming. SIAM Rev. 38(1): 49-95 (1996)
- [22] Watson, G.A.: Characterization of the Subdifferential of Some Matrix Norms. Linear Algebra Appl. 170(1): 33-45 (1992)
- [23] Zhang, L., Zhang, N. and Xiao, X.: On the Second-order Directional Derivatives of Singular Values of Matrices and Symmetric Matrix-valued Functions. Set-Valued Var. Anal. 21(3): 557-586 (2013)