Quantum chromatic numbers of some graphs in Hamming schemes

 $Xiwang Cao^{a,1}$, Keqin Feng^b, Ying-Ying Tan^c

 ${}^{a}S$ chool of Mathematics, Nanjing University of Aeronautics and Astronautics, China b Department of Mathematics, Tsinghua University, China c Anhui Jianzhu University, China

Abstract

The study of quantum chromatic numbers of graphs is a hot research topic in recent years. However, the infinite family of graphs with known quantum chromatic numbers are rare, as far as we know, the only known such graphs (except for complete graphs, cycles, bipartite graphs and some trivial cases) are the Hadamard graphs H_n with 2^n vertices and n being a multiple of 4. In this paper, we define a class of graphs named as generalized Hadamard graphs, we determined the quantum chromatic numbers of one class of such graphs. Notably, this is the second known family of graphs with the quantum chromatic numbers are explicitly determined except for some cases aforementioned. We also provide some bounds for the quantum chromatic numbers of some other generalized Hadamard graphs. Consequently, we can obtain the quantum chromatic numbers of products of some graphs.

Keywords: chromatic number, quantum chromatic number, colouring, quantum colouring *2000 MSC:* 05C15, 05E30, 94B25, 97K30

1. Introduction

In recent years, combinatorial designs and graph theory have become useful tools in the study of quantum communications and quantum information processing, mainly reflected in the following aspects:

- Quantum states constructed from graphs and hypergraphs are used to study entanglement phenomena and construct high-performance quantum error correction codes [\[1,](#page-14-0) [2](#page-14-1)];
- Spherical designs are used to construct various types of unbiased bases for quantum measurement [\[5](#page-14-2)];
- Classical combinatorial designs are extended to quantum versions (quantum orthogonal Latin squares, etc.) to study and construct various maximally entangled quantum states[\[4](#page-14-3)];
- Quantum state transfer are employed for transmitting quantum information using quantum networks, for example, the so-called perfect state transfer, uniform mixing. etc.[\[3](#page-14-4)];

¹This research is supported by National Natural Science Foundation of China, Grant No. 12171241

• Extend some parameters and concepts of the classical graphs to that of the quantum version, such as quantum homomorphism, quantum chromatic numbers, quantum independent numbers, etc.[\[2](#page-14-1)]

Let Γ be a simple graph whose vertex set is V and edge set E. A colouring on Γ is an assignment of colors to vertices of the graph such that the two vertices of each edge have different colours. Graph colouring is of particular interesting since it finds applications in quantum information theory and communication as seen in [\[1\]](#page-14-0). Classical graph colouring can be interpreted as a so-called non-local game, where two players Alice and Bob collaborate to answer pairs of questions without communication using some prior agreed-upon strategy. Quantum coloring of graphs is a modification of the classical graph coloring where the players may use "quantum" strategies meaning a shared entangled resource is available. In the colouring games, we are interested in the minimum number of the colours needed to win. For a classical graph Γ , this minimum number is denoted by $\chi(\Gamma)$, and termed as the chromatic number. For quantum graphs, by $\chi_q(\Gamma)$ and called the quantum chromatic number. The mathematical definition of these notations will be given in the next section. Notably, it was shown the tremendous quantum advantage in [\[1](#page-14-0)] that Hadamard graphs can provide examples of an exponential gap between the quantum and classical chromatic numbers using a result in [\[6\]](#page-14-5). However, in general, it is difficult to determine the chromatic number of a given graph, even more harder to evaluate or estimate the quantum chromatic number of the graph. In [\[9\]](#page-14-6), Ji proved that determining this number is NP hard. Up to date, except for complete graphs, cycles, bipartite graphs and some trivial cases, only a few lower bounds on the chromatic numbers are known for some sporadic graphs, as far as we know, the only non-trivial family of graphs whose quantum chromatic numbers are explicitly determined is the class of Hadamard graphs which were defined by Ito [\[8\]](#page-14-7) in the year 1985. Using the representations of certain groups and extensive computations, Ito obtained the spectra of the Hadamaed graphs. Very recently, Menamara [\[12\]](#page-14-8) determined the quantum chromatic numbers of the Hadamard graphs by using character sums over finite fields. Menamara computed the exact value of the quantum chromatic numbers of Hadamard graphs of order $n = 2^N$ for N a multiple of 4 using the upper bound derived by Avis etal [\[1\]](#page-14-0), as well as an application of the Hoffman-like lower bound of Elphick and Wocjan [\[4](#page-14-3)] that was generalized by Ganesan [\[7\]](#page-14-9) for quantum graphs. The main result in $[12]$ is as follows:

Theorem 1.1. [\[12](#page-14-8)] (Exact quantum chromatic number of Hadamard graphs). Let H_N be the *Hadamard graph on* 2 ^N *vertices,* N *a multiple of 4. Then,*

$$
\chi_q(H_N) = N. \tag{1}
$$

In this paper, we give a new method for calculating the spectrum of the Hadamard graph in Theorem [1.1](#page-1-0) by using some properties of Krawchouk polynomials, we also determined the quantum chromatic numbers of some other graphs in Hamming schemes, some bounds on the quantum chromatic numbers of some graphs in Hamming schemes are provided.

The organization of the paper is as follows: In Sect. [2,](#page-2-0) we give some backgrounds on quantum information and quantum measurements, as well as some basic concepts of graph theory. In Sect. [3,](#page-6-0) we define a class of generalized Hadamrd graphs, and determined their spectra. Using some spectral bounds on the quantum chromatic numbers, we successfully obtained the quantum chromatic numbers of one class of such graphs, see Theorem [3.3.](#page-9-0) Some bounds of quantum chromatic numbers of other class of generalized Hadamard graphs are provided as well (Theorem [3.5](#page-11-0) and Proposition [3.6\)](#page-12-0). By utilizing the products of graphs, we can also get the quantum chromatic numbers of some graphs (Theorem [3.11\)](#page-13-0).

2. Preliminaries

2.1. Some basic concepts of quantum communication

2.1.1. Quantum state

 \sum^{K} , where \sum is a q-ary set. In most cases, \sum is chosen as the finite field \mathbb{F}_q with q elements, In digital communications, information is represented as an K-tuple $c = (c_0, c_1, \dots, c_{K-1}) \in$ where q is a prime power. Then c can be viewed as a vector in the space $\mathbb{F}_q^{\hat{K}}$.

In quantum communication, each qubit, denoted by $|v\rangle = (v_0, v_1, \dots, v_{K-1})^T$, is a unite vector in the K-dimensional vector space \mathbb{C}^K . For every $|v\rangle = (v_0, v_1, \dots, v_{K-1})^T$, $|u\rangle =$ $(u_0, u_1, \dots, u_{K-1})^T \in \mathbb{C}^K$, define the inner product

$$
\langle u|v\rangle = \sum_{i=0}^{K-1} u_i v_i^*.
$$

If $\langle u|v \rangle = 0$, we say that $|u\rangle$ and $|v\rangle$ are separable. A quantum state is a vector in the space $\mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2} \otimes \cdots \mathbb{C}^{d_K}$ which is the tensor product of complex spaces. Take $V_i = \mathbb{C}^{d_i}$, $1 \leq i \leq K$, and choose an orthnormal basis of V_i as $|0\rangle, |1\rangle, \cdots, |d_i - 1\rangle$. Then

$$
\{|e_1 \otimes \cdots \otimes |e_K\rangle : 0 \le e_i \le d_i - 1, (1 \le i \le K)\}
$$

forms an orthnormal basis of $\mathfrak{V} := \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2} \otimes \cdots \mathbb{C}^{d_K}$. Thus each quantum state in \mathfrak{V} can be uniquely represented as

$$
|v\rangle = \sum_{0 \leq e_i \leq d_i-1, 1 \leq i \leq K} a_{e_1, \dots, e_K} |e_1\rangle \otimes \dots \otimes |e_K\rangle, a_{e_1, \dots, e_K} \in \mathbb{C}.
$$

2.1.2. Quantum measurement

Let $H = (h_{ij})_{0 \le i,j \le K-1}$ be a Hermite matrix. Then the quantum measurement of H on $|v\rangle \in$ \mathbb{C}^K is defined by $H|v\rangle$. In quantum communication, H can be written as $H = \sum_{i,j=0}^{K-1} h_{ij} |i\rangle\langle j|.$

Generally speaking, it is not easy to devise a measurement procedure which uniquely identifies the given quantum state from the statistical date produced by the measurements. For example, if the state of the quantum system is given by an $K \times K$ density matrix, the complete measurement statistics of one fixed *von Neumann* measurement is not sufficient to reconstruct the state, see e.g. [\[10\]](#page-14-10). However, it is possible to perform a somewhat general measurement procedure on a quantum system, namely a positive operator-valued measurement (or POVM for short), see [\[15](#page-14-11)]. Mathematically, a POVM is a collection of some semi-positive operators $E_i \geq 0$, each E_i is a K by K matrix, called POVM elements, satisfying the summation of all these operators is equal to I_K the identity matrix. POVMs constitute a basic ingredient in many applications of quantum information processing: quantum tomography, quantum key distribution required in cryptography, discrete Wigner function, quantum teleportation, quantum error correction codes, dense coding, teleportation, entanglement swapping, covariant cloning and so on, see for example [\[14](#page-14-12)].

2.1.3. Projective measurement

In a quantum measurement, people usually use some projective matrices $P = (p_{ij})_{0 \le i,j \le K}$: $\mathbb{C}^K \to \mathbb{C}^K$. A Hermite matrix P is called projective if $P^2 = P = P^*$. Suppose that $|v\rangle$ is contained in the image of P, that is, there is a vector $|a\rangle \in \mathbb{C}^K$ such that $P|a\rangle = |v\rangle$. Then

$$
P|v\rangle = P^2|a\rangle = P|a\rangle = |v\rangle.
$$

Thus $P|_{\text{Im}(P)} = id$. Then there exists a unitary matrix U such that $U^*PU = \text{diag}(I_r, 0)$, where $r = \text{rank}(P)$.

Finally, a set of projective operators $\{P_1, P_2, \cdots, P_K\}$ in $\mathbb{C}^{K \times K}$ is called a complete POVM if $P_i P_j = 0_K$ for every $1 \leq i \neq j \leq K$, and $\sum_{i=1}^K P_i = I_K$. In this case, it can be proved that there exists a unitary matrix U such that

$$
U^* P_i U = \text{diag}(0, 0, \cdots, 1, 0, \cdots, 0), 1 \le i \le K,
$$

where 1 is in the *i*-th entry. Moreover, $\mathbb{C}^K = \text{Im}(P_1) \oplus \text{Im}(P_2) \oplus \cdots \oplus \text{Im}(P_K)$.

2.2. Quantum homomorphism of graphs and graph colouring

Let $\Gamma = (V, E)$ be a simple graph with $n = |V|$ vertices and $m = |E|$ edges. A homomorphism φ from a graph $\Gamma_1 = (V_1, E_1)$ to a graph $\Gamma_2 = (V_2, E_2)$ is a mapping $\varphi : \Gamma_1 \to \Gamma_2$ satisfying $(\varphi(u), \varphi(v)) \in E_2$ if $(u, v) \in E_1$. For example, if $\Gamma_2 = K_c$ is a complete graph on c vertices, then $\varphi : \Gamma = (V, E) \to K_c$ is a homomorphism means that if $(u, v) \in E$, then $\varphi(u) \neq \varphi(v)$. We name the minimum number c such that there exists a homomorphism from Γ to K_c the chromatic number of Γ and denote it by $\chi(\Gamma)$. The maximum number c such that there is a homomorphism from K_c to Γ is called the clique number of Γ and denoted by $\omega(\Gamma)$. Let $\bar{\Gamma}$ be the complement graph of Γ . Then $\alpha(\Gamma) := \omega(\bar{\Gamma})$ is called the independent number of Γ.

Definition 2.1. *A quantum homomorphism from a graph* $\Gamma_1 = (V_1, E_1)$ *to a graph* $\Gamma_2 =$ (V_2, E_2) *means that there is a positive integer d such that for every* $x \in V_1$ *, there exists a complete orthogonal projective system* $\mathfrak{F}_x = \{P_{x,y} : y \in V_2\}$ *satisfying the following two conditions:*

- 1. (Completeness) For every $x \in V_1$, \mathfrak{F}_x *is a complete orthogonal system, namely,* $P_{x,y}^2 =$ $P_{x,y} = P_{x,y}^*$ and, when $y \neq y'$, we have $P_{x,y}P_{x,y'} = 0_d$. Moreover, $\sum_{y \in V_2} P_{x,y} = I_d$.
- 2. (Orthogonality) For every $x, x' \in V_1$, $y, y' \in V_2$, we have $P_{x,y}P_{x',y'} = 0$ whence $(x, x') \in V_1$ $E_1, (y, y') \in E_2.$

It is easy to see that a classical graph homomorphism is actually a quantum homomorphism.

Definition 2.2. *The quantum chromatic number of a graph* Γ *, denoted by* $\chi_q(\Gamma)$ *, is the minimum number* c *such that there exists a quantum homomorphism from* Γ *to the complete graph* K_c .

By definition, we see that for every graph Γ,

$$
\chi_q(\Gamma) \le \chi(\Gamma). \tag{2}
$$

It seems that how to determine the quantum chromatic number of a given graph is very hard. Up to date, except for some sporadic graphs with small size of vertices and some trivial cases, the only known class of graphs whose quantum chromatic numbers are determined are the Hadamard graphs H_N with N a multiple of 4. This situation motivates the study of quantum chromatic numbers of graphs. The following questions are of particular interesting.

- For a specific class of graphs, determine their chromatic numbers;
- Separable problem: find graphs such that their chromatic numbers are strictly less than their quantum chromatic numbers, note that $\chi_q(H_N) = N$ when $4|N$, however, $\chi(H_N) \ge$ $2^{N/2}$ when N is bigger enough;
- Find some lower or upper bounds for the chromatic numbers of some class of graphs.

For more information about quantum chromatic numbers, we refer the reader to [\[2,](#page-14-1) [4](#page-14-3), [5](#page-14-2)].

2.3. Spectra of Cayley graphs and bounds on quantum chromatic numbers

Let $\Gamma = (V, E)$ be a simple graph with $|V| = n, |E| = m$, A be its adjacency matrix. The spectrum of A is also termed the spectrum of Γ. For every $x \in V$, the number of its neighborhoods is defined as its valency (or degree). If we label the vertices of Γ as x_1, \dots, x_n , and denote the valency of x_i by k_i . Then $D := diag(k_1, \dots, k_n)$ is the degree matrix of Γ . $L = D - A$ (resp. $L^+ = D + A$) is the Laplace (resp. signless Laplace) matrix of Γ. Suppose that the eigenvalues of A, L and L^+ are $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$, $\theta_1 \geq \theta_2 \geq \cdots \geq \theta_n (= 0)$, and $\delta_1 \geq \delta_2 \geq \cdots \delta_n$, respectively. The following result is known, see for example, [\[4](#page-14-3)] and the references therein.

Theorem 2.3. *Let notations be defined as above. Then*

$$
\chi(\Gamma) \ge 1 + \max\left\{\frac{\lambda_1}{|\lambda_n|}, \frac{2m}{2m - n\delta_n}, \frac{\lambda_1}{\lambda_1 - \delta_1 + \theta_1}, \frac{n^+}{n^-}, \frac{n^-}{n^+}, \frac{S^+}{S^-}, \frac{S^-}{S^+}\right\},\tag{3}
$$

where n^+ (resp. n^-) is the number of positive (resp. negative) eigenvalues of Γ , and S^+ (resp. S [−]*) is the summation of the squares of positive (resp. negative) eigenvalues of* Γ*.*

A spectral lower bound on the quantum chromatic numbers is provided in $|4|$.

Lemma 2.4. [\[4\]](#page-14-3) For any graph Γ with eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$, we have

$$
\chi_q(\Gamma) \ge 1 + \frac{\lambda_1}{|\lambda_n|}.\tag{4}
$$

Let G be a finite group. A representation of G is a homomorphism $\rho: G \to GL(U)$ for a (finite-dimensional) non-zero vector space U. The dimension of U is called the degree of ρ . Two representations $\rho: G \to GL(U)$ and $\varrho: G \to GL(W)$ are *equivalent*, denoted by $\rho \sim \varrho$, if there exists an isomorphism $T: U \to W$ such that $\rho_g = T \varrho_g T^{-1}$ for all $g \in G$. For every representation $\rho: G \to GL(U)$ of G, the character of χ_{ρ} is defined by:

$$
\chi_{\rho}: G \to \mathbb{C}, \chi_{\rho}(g) = \text{tr}(\rho(g)) \text{ for all } g \in G,
$$

where $tr(\rho(g))$ is the trace of the representation matrix with respect to a basis of U.

A subspace W of U is said to be G-invariant if $\rho(q)\omega \in W$ for every $q \in G$ and $\omega \in$ W. Obviously, $\{0\}$ and U are G-invariant subspaces, called trivial subspaces. If U has no non-trivial G-invariant subspaces, then ρ is called an irreducible representation and χ_{ρ} and irreducible character of G.

Let S be a subset of G with $S^{-1} := \{s^{-1} : s \in S\} = S$ and $G = \langle S \rangle$. A Cayley graph over G with connection set S is defined by $\Gamma := \text{Cav}(G, S)$ where the vertex set is G and two elements $x, y \in G$ are adjacent if and only if $xy^{-1} \in S$. If S is conjugation closed meaning that for every $x \in G$ and $s \in S$, we have $x^{-1}sx \in S$. In this case, the Cayley graph Cay (G, S) is called normal. For normal Cayley graphs, the following result is well-known.

Lemma 2.5. [\[17,](#page-14-13) pp. 69-70] Let $G = \{g_1, \dots, g_n\}$ be a finite group of order n and let $\rho^{(1)}, \cdots, \rho^{(s)}$ be a complete set of unitary representatives of the equivalent classes of irreducible *representations of* G. Let χ_i be the character of $\rho^{(i)}$ and d_i be the degree of χ_i . Let S be a *symmetric set and further that* $gSq^{-1} = S$ *for all* $g \in G$ *. Then the eigenvalues of the adjacency matrix* A *of the Cayley graph* Cay(G, S) *with respect to* S *are given by*

$$
\lambda_k = \frac{1}{d_k} \sum_{g \in S} \chi_k(g), 1 \le k \le s,
$$

each λ_k has multiplicity d_k^2 . Moreover, the vectors

$$
v_{ij}^{(k)} = \frac{\sqrt{d_k}}{|G|} \left(\rho_{ij}^{(k)}(g_1), \cdots, \rho_{ij}^{(k)}(g_n) \right)^T, 1 \le i, j \le d_k
$$

 ${\it form \ an \ orthonormal \ basis \ for \ the \ eigenspace } {\it eq}$ ${\it V}_{\lambda_k}.$

Note that a proof of Lemma [2.5](#page-5-0) can also be found in [\[13](#page-14-14), Theorem 9].

As a consequence, if G is a finite abelian group, we assume that G is decomposed as a direct sum of cyclic groups, $G = \mathbb{Z}_{n_1} \oplus \cdots \oplus \mathbb{Z}_{n_r}$, then the spectrum of the Cayley graph $\Gamma = \text{Cay}(G, S)$ is given by

$$
\lambda_g = \sum_{s \in S} \chi_g(s),\tag{5}
$$

where $\chi_g(s) = \prod_{j=1}^s \xi_{n_j}^{g_js_j}, \forall g = (g_1, \dots, g_r) \in G, s = (s_1, \dots, s_r) \in S$, and ξ_{n_j} is a primitive n_j -th root of unity in \mathbb{C} .

2.4. Krawtchouk polynomials

For positive integers n, ℓ , and q, the Krawchouk polynomial in variable x is defined by

$$
K_{\ell}^{n,q}(x) = \sum_{j=0}^{\ell} (-1)^{j} (q-1)^{\ell-j} {x \choose j} {n-x \choose \ell-j}.
$$
 (6)

Krawchouk polynomials are a kind of orthogonal polynomials and have many important applications in many fields such as coding theory, function analysis and approximation etc. For our purpose, we list some of the properties of such polynomials as follows.

Theorem 2.6. *[\[11\]](#page-14-15) The Krawchouk polynomials have the following properties.*

1. (Orthogonality Relations): For every $i, j, (i, j = 0, 1, \dots, n)$

$$
\sum_{d=0}^{n} K_i^n(d) K_j^n(d) (q-1)^d \binom{n}{d} = q^n (q-1)^i \binom{n}{i} \delta_{i,j}.
$$
 (7)

2. (Recursive Relation): For any $k = 1, \dots, n$ and any real x

$$
K_k^n(x) = K_k^{n-1}(x-1) - K_{k-1}^{n-1}(x-1)
$$
\n
$$
K_k^n(x) = K_k^{n-1}(x) + (a-1)K_k^{n-1}(x)
$$
\n(8)

$$
K_k^n(x) = K_k^{n-1}(x) + (q-1)K_{k-1}^{n-1}(x)
$$
\n(9)

$$
K_k^{n-1}(x) = \sum_{j=0}^k K_j^n(x)(1-q)^{k-j}.
$$
 (10)

3. (Reciprocal Law):

$$
(q-1)^{i} \binom{n}{i} K_d^n(i) = (q-1)^d \binom{n}{d} K_i^n(d). \tag{11}
$$

4. (Generating Function):

$$
\sum_{k=0}^{n} K_k^n(d) z^k = (1-z)^d (1 + (q-1)z)^{n-d}.
$$
 (12)

5. (Inversion Formula):

$$
f(x) = \sum_{j=0}^{n} f_j K_j^n(x)
$$
 (13)

if and only if for every $i = 0, 1, \cdots, n$,

$$
f_i = q^{-n} \sum_{j=0}^{n} f(j) K_j^n(i).
$$
 (14)

3. Main results

Let $V_n = \{(x_0, x_1, \dots, x_{n-1}) : x_i \in \mathbb{F}_2\}$, where \mathbb{F}_2 is the binary field. V_n is a *n*-dimensional vector space over \mathbb{F}_2 . For $x = (x_0, x_1, \dots, x_{n-1}) \in V_n$, the Hamming weight of x, denoted by $wt(x)$, is the nonzero coordinates of x, the support of x is supp $(x) := \{i : 0 \le i \le n-1, x_i = 1\}.$ For $x, y \in V_n$, the Hamming distance between x and y is $d(x, y) := wt(x - y)$. The following defined Hadamard graph is isomorphic to that defined by Ito [\[8](#page-14-7)].

Definition 3.1. Let n be a positive integer with $4|n$. The Hadamard graph $H_n = (V_n, E_n)$, *where* V_n *is the n*-dimensional vector space over \mathbb{F}_2 . Two vectors $x, y \in V_n$ *are adjacent if and only if* $d(x, y) = n/2$.

In what follows, we give a generalization of Hadamard graphs.

Definition 3.2. Let n be a positive integer with $2|n, \ell$ be a positive integer. We define the *generalized Hadamard graph* $GH_{n,\ell} = (V_n, E_n)$ *as the* V_n *is the n*-dimensional vector space over \mathbb{F}_2 *, two vectors* $x, y \in V_n$ are adjacent if and only if $d(x, y) = \ell$.

Note that if ℓ is odd, then $GH_{n,\ell}$ is a bipartite graph and then its quantum chromatic is 2. Thus in next sequel, we assume that ℓ is even.

In this section, we first give a new method to calculate the spectrum of H_n and then prove that $\chi_q(H_n) = n$. Then for the generalized Hadamard graphs, we present some new results on the quantum chromatic numbers of such graphs.

3.1. Proof of Theorem [1.1](#page-1-0)

Firstly, it is easy to see that $H_n = \text{Cay}(V_n, S)$, where $S = \{x \in V_n : wt(x) = n/2\}$. The character group of V_n (as an elementary commutative 2-group of rank n) is $\widehat{V}_n = \{\phi_a : a \in$ V_n , where $\phi_a(x) = (-1)^{x \cdot a}$, $x \cdot a$ is the inner product of x and a, i.e., $x \cdot a = \sum_{i=0}^{n-1} x_i a_i$, $a = (a_0, \dots, a_{n-1})$. By [\(5\)](#page-5-1), the eigenvalues of H_n are

$$
\lambda_a = \sum_{s \in S} (-1)^{s \cdot a}, a \in V_n. \tag{15}
$$

Obviously, $\lambda_0 = |S| = \binom{n}{n}$ $n \choose n/2$. Take $a = 1_n := (1, 1, \dots, 1)$. Then

$$
\lambda_{1_n} = \sum_{s \in S} (-1)^{s \cdot 1_n} = \sum_{s \in S} (-1)^{wt(s)} = \sum_{s \in S} (-1)^{n/2} = \sum_{s \in S} 1 = |S| = \binom{n}{n/2} = \lambda_0.
$$

And for every $a \in V_n$, $a \neq 0_n, 1_n$, then $\lambda_a < \lambda_0$. Thus $\lambda_{\max} = \binom{n}{n}$ $\binom{n}{n/2}$ with multiplicity 2. H_n has two isomorphic components. Below, we proceed to find λ_{\min} .

For $a \neq 0_n, 1_n \in V_n$,

$$
\lambda_a = \sum_{s \in S} (-1)^{s \cdot a} = \sum_{x \in V_n : wt(x) = n/2} (-1)^{x \cdot a}.
$$

Suppose that $a = (a_0, \dots, a_{n-1}) \in V_n$, $wt(a) = r$, $1 \le wt(a) < n$. Assume that supp $(a) =$ $\{i_1, i_2, \dots, i_r\}$. Let x run through V_n with weight $n/2$. If $|\text{supp}(x) \cup \text{supp}(a)| = j$, then $x \cdot a = j$. A simple argument shows that

$$
\lambda_a = \sum_{s \in S} (-1)^{s \cdot a} = \sum_{x \in V_n : wt(x) = n/2} (-1)^{x \cdot a} = \sum_{j=0}^{n/2} (-1)^j {r \choose j} {n-r \choose n/2 - j} = K_{n/2}^n(r).
$$

Thus we build a link between the eigenvalues of H_n and the Krawchouk polynomials. By using the Reciprocal Law of the Krawchouk polynomials (see Theorem [2.6\)](#page-5-2), we have

$$
K_{n/2}^n(r) = \frac{\binom{n}{n/2}}{\binom{n}{r}} K_r^n(n/2).
$$

Since $K_r^n(n/2)$ is the coefficient of x^r in $(1-x)^{n/2}(1+x)^{n-n/2} = (1-x^2)^{n/2}$. Thus, if $r = 2j+1$ is odd, then $\lambda_a = K_{n/2}^n(2j + 1) = 0$; if $r = 2j$ for some j, then

$$
\lambda_a = (-1)^j \frac{\binom{n}{n/2} \binom{n/2}{j}}{\binom{n}{2j}}.
$$
\n(16)

Now, it is easy to see that

$$
\lambda_{\min} = (-1)^j \frac{\binom{n}{n/2} \binom{n/2}{1}}{\binom{n}{2}} = -\frac{\binom{n}{n/2}}{n-1} = -\frac{\lambda_{\max}}{n-1}.
$$
\n(17)

Then, by the spectral bounds in (4) , we obtain

$$
\chi_q(H_n) \ge 1 + \frac{\lambda_{\max}}{|\lambda_{\min}|} = n. \tag{18}
$$

Next, we show that $\chi_q(H_n) \leq n$. To this end, we need to find a quantum homomorphism of H_n . Very recently, Menamara [\[12](#page-14-8)] found such a homomorphism. We provide his result for completeness.

For every $x = (x_0, x_1, \dots, x_{n-1}) \in V_n$, and $0 \le \alpha \le n-1$, we define the following operators:

$$
P_x^{\alpha} = (a_x^{\alpha}(i,j))_{0 \le i,j \le n-1}, a_x^{\alpha}(i,j) = \frac{1}{n} \xi_n^{(j-i)\alpha} (-1)^{x_i + x_j},
$$
\n(19)

where $\xi_n = e^{\frac{2\pi\sqrt{-1}}{n}}$ is a *n*-th root of unity in C. Then it is obvious that P_x^{α} is a Hermite matrix, moreover, let $(P_x^{\alpha})^2 = (b(i,j))_{0 \le i,j \le n-1}$. Then

$$
b(i,j) = \sum_{k=0}^{n-1} a_x^{\alpha}(i,k) a_x^{\alpha}(k,j)
$$

=
$$
\frac{1}{n^2} \sum_{k=0}^{n-1} \xi_n^{(k-i)\alpha} (-1)^{x_i + x_k} \xi_n^{(j-k)\alpha} (-1)^{x_k + x_k}
$$

=
$$
\frac{1}{n^2} \xi_n^{(j-i)\alpha} (-1)^{x_i + x_j} \sum_{k=0}^{n-1} 1
$$

=
$$
a_x^{\alpha}(i,j).
$$

Thus $(P_x^{\alpha})^2 = P_x^{\alpha}$. That is, P_x^{α} is a projection.

For every $x \in V_n$, let $\Delta_x = \{P_x^{\alpha} : 0 \leq \alpha \leq n-1\}$. We aim to prove Δ_x is a complete orthogonal system of $\mathbb{C}^{n \times n}$. Indeed, for every $0 \leq \alpha \neq \alpha' \leq n-1$, denote $P_x^{\alpha} P_x^{\alpha'} = (c(i,j)).$ Then

$$
c(i,j) = \sum_{k=0}^{n-1} \frac{1}{n^2} \xi_n^{(k-i)\alpha} (-1)^{x_i + x_k} \xi_n^{(j-k)\alpha'} (-1)^{x_j + x_k}
$$

=
$$
\frac{1}{n^2} \xi_n^{j\alpha' - i\alpha} (-1)^{x_i + x_j} \sum_{k=0}^{n-1} \xi_n^{k(\alpha - \alpha')}
$$

= 0.

Therefore, $P_x^{\alpha} P_x^{\alpha'} = 0$. Furthermore, we can prove that for every $x \in V_n$, the above defined

 \triangle_x is complete, i.e., $\sum_{\alpha=0}^{n-1} P_x^{\alpha} = I_n$. Let $\sum_{\alpha=0}^{n-1} P_x^{\alpha} = (u(i,j))_{0 \le i,j \le n-1}$. Then

$$
u(i,j) = \sum_{\alpha=0}^{n-1} a_x^{\alpha}(i,j)
$$

=
$$
\frac{1}{n} \sum_{\alpha=0}^{n-1} \xi_n^{(j-i)\alpha} (-1)^{x_i + x_j}
$$

=
$$
\delta_{i,j} (-1)^{x_i + x_j}
$$

=
$$
\delta_{i,j}.
$$

Thus $\sum_{\alpha=0}^{n-1} P_x^{\alpha} = I_n$.

Finally, let $x, y \in V_n$ with $(x, y) \in E$ be an edge of H_n , that is $d(x, y) = 2t$, then

$$
(P_x^{\alpha} P_y^{\alpha})(i, j) = \frac{1}{n^2} \sum_{k=0}^{n-1} \xi_n^{(i-k)\alpha} (-1)^{x_i + x_k} \xi_n^{(k-j)\alpha} (-1)^{y_j + y_k}
$$

$$
= \frac{1}{n^2} \xi_n^{(i-j)\alpha} (-1)^{x_i + y_j} \sum_{k=0}^{n-1} (-1)^{x_k + y_k}
$$

$$
= \frac{1}{n^2} \xi_n^{(i-j)\alpha} (-1)^{x_i + y_j} (-2t + 4t - 2t)
$$

$$
= 0.
$$

Thus the set $\mathfrak F$ provides a quantum colouring of H_n . Therefore, by the definition of quantum chromatic numbers, we know that

$$
\chi_q(H_n) \le n. \tag{20}
$$

Combining [\(18\)](#page-8-0) and [\(20\)](#page-9-1), we have $\chi_q(H_n) = n$ as required.

3.2. Some new results

3.2.1. Quantum chromatic number of a kind of generalized Hadamard graphs

Firstly, we have the following result:

Theorem 3.3. Let $V_n = \mathbb{F}_2^n$ be the *n*-dimensional vector space over \mathbb{F}_2 , $S = \{x \in V_n : wt(s) = \mathbb{F}_2^n\}$ $ℓ$ *}. Define a graph by* $GH_{n,ℓ} := Cay(V_n, S)$ *. If* $n = 4t − 1$ *and* $ℓ = 2t$ *for some positive integer t*, then the spectrum of $GH_{n,\ell}$ is

$$
\lambda_a = \begin{cases}\n(-1)^j \frac{\binom{4t-1}{2t} \binom{2t-1}{j}}{\binom{4t-1}{2t-1}} & \text{if } wt(a) = r = 2j, \ 0 \le j \le 2t - 1, \\
(-1)^{j+1} \frac{\binom{4t-1}{2t} \binom{2t-1}{j}}{\binom{4t-1}{2j+1}} & \text{if } wt(a) = r = 2j + 1, \ 0 \le j \le 2t - 1.\n\end{cases} (21)
$$

Moreover,

$$
\chi_q(GH_{n,\ell}) = n+1. \tag{22}
$$

Proof. For every $a = (a_0, a_1, \dots, a_{n-1}) \in V_n$, if $wt(a) = r$, the corresponding eigenvalue of $GH_{n,\ell}$ is

$$
\lambda_a = \sum_{s \in S} (-1)^{x \cdot a}.
$$

It is readily seen that the maximum eigenvalue of $GH_{n,\ell}$ is $\lambda_{\max} = \binom{n}{\ell}$ λ_{ℓ}^{n} = λ_{1n} since ℓ is even.

For $a \neq 0_n, 1_n \in V_n$,

$$
\lambda_a = \sum_{s \in S} (-1)^{s \cdot a} = \sum_{x \in V_n : wt(x) = \ell} (-1)^{x \cdot a}.
$$

A similar analysis as in Section [3.1](#page-7-0) leads to

$$
\lambda_a = \sum_{s \in S} (-1)^{s \cdot a} = \sum_{x \in V_n : wt(x) = \ell} (-1)^{x \cdot a} = K^n_{\ell}(r). \tag{23}
$$

By the Reciprocal Law of the Krawchouk polynomials (see Theorem [2.6\)](#page-5-2), we have

$$
K_{\ell}^{n}(r) = \frac{\binom{n}{\ell}}{\binom{n}{r}} K_{r}^{n}(\ell).
$$

Since $K_r^n(\ell)$ is the coefficient of x^r in the expansion of

$$
(1-x)^{\ell}(1+x)^{n-\ell} = (1-x^2)^{2t-1}(1-x) = \sum_{j=0}^{2t-1} (-1)^j {\binom{2t-1}{j}} (x^{2j} - x^{2j+1}).
$$

Therefore, if $r = 2j$ for some j, then

$$
\lambda_a = (-1)^j \frac{\binom{4t-1}{2t} \binom{2t-1}{j}}{\binom{4t-1}{2j}}.
$$
\n(24)

If $r = 2j + 1$ is odd, then

$$
\lambda_a = (-1)^{j+1} \frac{\binom{4t-1}{2t} \binom{2t-1}{j}}{\binom{4t-1}{2j+1}}.
$$
\n(25)

By $(24)-(25)$ $(24)-(25)$, one can check that

$$
\lambda_{\min} = -\frac{\binom{4t-1}{2t}}{4t-1} = -\frac{\lambda_{\max}}{4t-1}.\tag{26}
$$

Thanks to Lemma [2.4,](#page-4-1) we have

$$
\chi_q(GH_{4t-1,2t}) \ge 4t = n+1. \tag{27}
$$

In order to prove $\chi_q(GH_{4t-1,2t}) \leq 4t$, we need to find a proper quantum colouring of such a graph. To this end, for every $x = (x_0, x_1, \dots, x_{n-1}) \in V_n$, we extend it to V_{n+1} by adding a coordinate at the end of x as $\tilde{x} = (x_0, x_1, \dots, x_{n-1}, 0)$, i.e., $x_n = 0$. Define the following set of operators on \mathbb{C}^{4t} :

$$
\mathfrak{F} := \{ P_x^{\alpha} : x \in V_n, 0 \le \alpha \le 4t - 1 \},\
$$

where

$$
P_x^{\alpha} = (p_x^{\alpha}(i,j))_{0 \le i,j \le 4t-1}, p_x^{\alpha}(i,j) = \frac{1}{4t} \xi_{4t}^{(j-i)\alpha} (-1)^{x_i + x_j}.
$$

It is obvious that each operator is Hermitian, and the (i, j) -th entry of $(P_x^{\alpha})^2$ is

$$
((P_x^{\alpha})^2)(i,j) = \sum_{k=0}^{4t-1} p_x^{\alpha}(i,k) p_x^{\alpha}(k,j) = \frac{1}{(4t)^2} \xi_{4t}^{(j-i)\alpha} (-1)^{x_i+x_j} \sum_{k=0}^{4t-1} 1 = p_x^{\alpha}(i,j).
$$

Thus P_x^{α} is a projection. Similarly, one can prove that

$$
P_x^{\alpha} P_x^{\alpha'} = 0 \text{ if } \alpha \neq \alpha',
$$

and
$$
\sum_{0 \leq \alpha \leq 4t-1} P_x^{\alpha} = I_{4t}.
$$

Now, we assume that $(x, y) \in E_{4t-1,2t}$ is an edge of $GH_{4t-1,2t}$, then we need to prove that $P_x^{\alpha}P_y^{\alpha} = 0$, where $x = (x_0, x_1, \cdots, x_{4t-2}), y = (y_0, y_1, \cdots, y_{4t-2})$ and $x_{4t-1} = y_{4t-1} = 0$. Moreover, $d(x, y) = 2t$. Since

$$
\sum_{k=0}^{4t-1} (-1)^{x_k+y_k} = 1 + \sum_{k=0}^{4t-2} (-1)^{x_k+y_k} = 1 - 2t + 2t - 1 = 0,
$$

it follows that

$$
(P_x^{\alpha} P_y^{\alpha})_{i,j} = \sum_{k=0}^{4t-1} p_x^{\alpha}(i,k) p_y^{\alpha}(k,j) = \frac{1}{(4t)^2} \xi_4^{(j-i)\alpha} (-1)^{x_i+y_j} \sum_{k=0}^{4t-1} (-1)^{x_k+y_k} = 0.
$$

Thus $P_x^{\alpha} P_y^{\alpha} = 0$ if $(x, y) \in E_{4t-1,2t}$.

Therefore, \mathfrak{F} gives a quantum colouring of $GH_{4t-1,2t}$, and then

$$
\chi_q(GH_{4t-1,2t}) \le 4t = n+1. \tag{28}
$$

Combining [\(27\)](#page-10-2) and [\(28\)](#page-11-1) yields $\chi_q(GH_{4t-1,2t}) = 4t = n + 1$. This completes the proof. \Box

Remark 3.4. *We note that, except for some trivial cases, the family of graphs in Theorem [3.3](#page-9-0) is the second known family of graphs whose quantum chromatic numbers are exactly determined, up to our knowledge.*

3.2.2. A upper bound for quantum chromatic numbers of generalized Hadamard graphs

Let ℓ be a positive integer. $GH_{n,\ell} = \text{Cay}(V_n, S)$, where $S = \{x \in V_n : wt(x) = \ell\}$. If ℓ is odd, then $GH_{n,\ell}$ is a bipartite graph and then $\chi_q(GH_{n,\ell})=2$. Thus we assume that $\ell=2t$ for some integer t . In this case, we have the following result:

Theorem 3.5. Let notations be defined as above, if $\ell \geq n/2$, then

$$
\chi_q(GH_{n,\ell}) \le 2\ell. \tag{29}
$$

Proof. Denote $d = 2\ell$. For every $x = (x_0, x_1, \dots, x_{n-1} \in V_n$, we let $x_n = \dots x_{d-1} = 0$. Define a set of operators as follows:

$$
\mathfrak{F} = \{P_x^{\alpha} : x \in V_n, 0 \le \alpha \le d - 1\},\
$$

where $P_x^{\alpha}(i, j) = \frac{1}{d} \xi_d^{(i-j)\alpha}$ $\int_{d}^{(i-j)\alpha} (-1)^{x_i+x_j}.$

Using a similar method as in the previous section, we can prove that $\mathfrak F$ forms a complete orthogonal system of \mathbb{C}^d .

When $0 \le \alpha \le d-1$, we claim that if $x, y \in V_n$ with $d(x, y) = \ell$, it holds that $P_x^{\alpha} P_y^{\alpha} = 0$. Note that for $y \in V_n$, we define $y_n = \cdots = y_{d-1} = 0$. Indeed, we have

$$
P_x^{\alpha} P_y^{\alpha}(i,j) = \frac{1}{d^2} \sum_{k=0}^{d-1} \xi_d^{(i-k)\alpha} (-1)^{x_i + x_k} \xi_d^{(j-k)\alpha} (-1)^{y_k + y_j} = \frac{1}{d^2} \xi_d^{(j-i)\alpha} (-1)^{x_i + x_j} \sum_{k=0}^{d-1} (-1)^{x_k + y_k}.
$$

Now,

$$
\sum_{k=0}^{d-1} (-1)^{x_k+y_k} = \sum_{k=0}^{n-1} (-1)^{x_k+y_k} + \sum_{k=n}^{d-1} (-1)^{x_k+y_k} = (-\ell+n-\ell) + (d-n) = 0.
$$

Therefore, $\mathfrak F$ gives a proper quantum colouring of $GH_{n,\ell}$, and then the desired result follows.

 \Box

Particularly, if $n = 4t + 2$, $\ell = 2t + 2$, we have

Proposition 3.6. Let $GH_{n,\ell} = \text{Cay}(V_n, S)$, where $S = \{x \in V_n : wt(x) = \ell\}$. If $n = 4t+2, \ell =$ $2t + 2$, then the spectrum of $GH_{n,\ell}$ is given by

$$
\lambda_a = \begin{cases}\n(-1)^j \frac{\left[\binom{2t}{j} - \binom{2t}{j-1}\right]\binom{n}{\ell}}{\binom{n}{2j}} & \text{if } wt(a) = r = 2j, \\
(-1)^j \frac{2\binom{2t}{j}\binom{n}{\ell}}{\binom{n}{2j+1}} & \text{if } wt(a) = r = 2j+1.\n\end{cases}
$$
\n(30)

Moreover,

$$
\ell \le \chi_q(GH_{n,\ell}) \le 2\ell. \tag{31}
$$

Proof. Since $\ell > n/2$, the upper bound $\chi_q(GH_{n,\ell}) \leq 2\ell$ follows from Theorem [3.5.](#page-11-0) Now, we prove the lower bound. In order to prove it, we compute the eigenvalues of $GH_{n,\ell}$. For every $a \in V_n$, if $wt(a) = r$, the eigenvalue of $GH_{n,\ell}$ corresponding to a is

$$
\lambda_a = \sum_{x \in V_n, wt(x) = \ell} (-1)^{x \cdot a} = K^n_{\ell}(r) = \frac{\binom{n}{\ell}}{\binom{n}{r}} K^n_r(\ell).
$$

When $n = 4t + 2$, $t = 2t + 2$, $K_r^n(\ell)$ is the coefficient of x^r in the expansion of

$$
(1-x)^{2t+2}(1+x)^{2t} = 1 + x^{4t+2} + \sum_{j=1}^{2t} (-1)^j \left[\binom{2t}{j} - \binom{2t}{j-1} \right] x^{2j} - 2 \sum_{j=1}^{2t} \binom{2t}{j} x^{2t+1}.
$$

Thus, we have

$$
\lambda_a = \begin{cases}\n(-1)^j \frac{\left[\binom{2t}{j} - \binom{2t}{j-1}\right]\binom{n}{\ell}}{\binom{n}{2j}} & \text{if } wt(a) = r = 2j, \\
(-1)^j \frac{2\binom{2t}{j}\binom{n}{\ell}}{\binom{n}{2j+1}} & \text{if } wt(a) = r = 2j+1.\n\end{cases}
$$
\n(32)

It is easy to check that $\lambda_{\min} = -\binom{n}{\ell}$ $\binom{n}{\ell}/(2t+1)$. By the spectral bounds on the quantum chromatic numbers, see Lemma [2.4,](#page-4-1) we have

$$
\chi_q(GH_{4t+2,2t+2}) \ge 2t+2 = \ell.
$$

This completes the proof.

 \Box

Remark 3.7. *Form the discussions before this, we can see that for the generalized Hadamrd graphs* $GH_{n,\ell}$ *, we can use the properties of Krawchouk polnomials to evaluate the spectra of these graphs, and then by the spectral bounds on quantum chromatic numbers, we can get a lower bound on such numbers, especially, when* $n - 2\ell$ *is small. If* $\ell > n/2$ *, we have a upper bound for the quantum chromatic numbers. But for* $\ell < n/2$, we don't know how to find such *a upper bound. The following question is still open.*

Open question: Find a upper bound on $\chi_q(GH_{n,\ell})$ when $2\ell < n$.

3.3. Quantum chromatic numbers of products of graphs

Let Γ_1 and Γ_2 be two graphs. The product of Γ_1 and Γ_2 , denoted by $\Gamma_1 \times \Gamma_2$, is defined by $\Gamma_1 \times \Gamma_2 = (V, E)$, where $V = V(\Gamma_1) \times V(\Gamma_2) = \{(x_1, x_2) : x_1 \in V(\Gamma_1), x_2 \in V(\Gamma_2) \text{ and }$ $E = \{((x_1, x_2), (y_1, y_2)) : (x_1, y_1) \in E(\Gamma_1) \text{ and } (x_2, y_2) \in E(\Gamma_2)\}\.$ Let the eigenvalues of Γ_1 be $\lambda_1 \geq \cdots \geq \lambda_n$ and the eigenvalues of Γ_2 be $\mu_1 \geq \cdots \geq \mu_m$. Then the eigenvalues of $\Gamma_1 \times \Gamma_2$ are $\{\lambda_i \mu_j : 1 \le i \le n, 1 \le j \le m\}$. Thus $\lambda_{\max}(\Gamma_1 \times \Gamma_2) = \lambda_1 \mu_1$, and $\lambda_{\min}(\Gamma_1 \times \Gamma_2) =$ $\min\{\lambda_1\mu_m, \lambda_n\mu_1\}.$ In [\[16\]](#page-14-16), a upper bound on the quantum chromatic numbers of products of graphs is given as follows:

Lemma 3.8. *[\[16\]](#page-14-16) Let notations be defined as above. Then*

$$
\chi_q(\Gamma_1 \times \Gamma_2) \le \min\{\chi_q(\Gamma_1), \chi_q(\Gamma_2)\}.
$$
\n(33)

By Lemma [3.8,](#page-13-1) Elphick and Wocjan [\[4\]](#page-14-3) proved the following result.

Theorem 3.9 ([\[4\]](#page-14-3), Prop. 3.2). *Suppose that the eigenvalues of* Γ_1 *are* $\lambda_1 \geq \cdots \geq \lambda_n$ *and the eigenvalues of* Γ_2 *are* $\mu_1 \geq \cdots \geq \mu_m$. If $\frac{\lambda_1}{|\lambda_n|} \geq \frac{\mu_1}{|\mu_m|}$ $\frac{\mu_1}{|\mu_m|}$, then

$$
\chi_q(\Gamma_1 \times \Gamma_2) \ge 1 + \frac{\mu_1}{|\mu_n|}.\tag{34}
$$

As a consequence, we have

Corollary 3.10. *If* $\chi_q(\Gamma_1) = 1 + \frac{\lambda_1}{|\lambda_n|}$ and $\chi_q(\Gamma_2) = 1 + \frac{\mu_1}{|\mu_m|}$, then $\chi_a(\Gamma_1 \times \Gamma_2) = \min{\chi_a(\Gamma_1), \chi_a(\Gamma_2)}.$ (35)

By Corollary [3.10,](#page-13-2) we get the following result for generalized Hadamard graphs.

Theorem 3.11. Let n_i, ℓ_i $(i = 1, 2)$ and $s \geq t$ be positive integers. Then

- *1.* $\chi_q(GH_{4t,2t} \times GH_{4s,2s}) = 4t;$
- 2. $\chi_a(GH_{4t-1,2t} \times GH_{4s,2s}) = 4t;$
- *3.* $\chi_q(GH_{4t,2t} \times GH_{4s-1,2s}) = 4t$;
- χ_a (*GH*_{4t−1,2t} × *GH*_{4s−1,2s}) = 4*t*.

References

References

- [1] David Avis, Jun Hasegawa, Yosuke Kikuchi, and Yuuya Sasaki, A quantum protocol to win the graph colouring game on all hadamard graphs, IEICE Trans. Fundam. Electron. Commun. Comput. Sci. E89-A (2006), no. 5, 1378-1381.
- [2] Peter J. Cameron, Ashley Montanaro, Michael W. Newman, Simone Severini, and Andreas Winter, On the quantum chromatic number of a graph, Electron. J. Combin. 14 (2007), No. 1, Research Paper 81, 15 pages.
- [3] Ada Chan and William J. Martin, Quantum isomorphism of graphs from association schemes, Journal of Combinatorial Theory, Series B, 164 (2024), 340-363.
- [4] Clive Elphick and Pawel Wocjan, Spectral lower bounds for the quantum chromatic number of a graph, Journal of Combinatorial Theory, Series A, 168 (2019), 338-347.
- [5] Keqin Feng, Quantum chromatic numbers: A survey (in Chinese), Journal of Hebei Normal University (Natural Science), 17(5)(2023), 433-446.
- [6] Peter Frankl and Vojtech R¨odl, Forbidden intersections, Trans. Amer. Math. Soc. 300(1)(1987), 259-286.
- [7] Priyanga Ganesan, Spectral bounds for the quantum chromatic number of quantum graphs, Linear Algebra Appl. 674 (2023), 351-376.
- [8] Noboru Ito, Hadamard graphs. I, Graphs Combin. 1 (1985), 57-64.
- [9] Zhengfeng Ji, Binary constraint system games and locally commutative reductions, ArXiv abs/1310.3794 (2013).
- [10] A. Klappenecker, M. Rötteler, I. E. Spharlinski, A Winterhof, On approximately symmetric informationally complete positive operator-valued measures and related systems of quantum states. J. Math Phys. A, 46: 082104(2005).
- [11] Vladimir I. Levenshtein, Krawtchouk polynomials and universal bounds for codes and designs in Hamming spaces, IEEE Trans. Inform. Theory, 41(1995), 1303-1321.
- [12] A.M. Menamara, ArXiv: 2410. 0042 v2 [math. OA], 14, Oct. 2024.
- [13] M.R. Murty, Ramanujan graphs, J. Ramanujan Math. Sco. 18, No. 1(2003), 1-20.
- [14] M.A. Nilsen and I.L. Chuang, Quantum computation and quantum information, Cambridge University Press, 2000.
- [15] A. Peres, Quantum Theory: Concepts and Methods. Kluwer Academic Publishers, 1993.
- [16] R.Santiago, A.M. Mcnamara, Quantum chromatic numbers of products of graphs, ArXiv:2408.11911v1 [math.OA], 21 Aug 2024.
- [17] B. Steinberg, Representation Theory of Finite Groups: An Introductory Approach. Universitext. DOI: 10.1007/978-1-4614-0776-8-7.

4. Appendix

For readers convenience, we provide some tables on the spectra of some generalized Hadamard graphs. Note that if $wt(a) = r$, we denote by $\rho_{\ell}^{n}(r)$ the eigenvalue of $GH_{n,\ell}$ corresponding to a.

Table 2. $\rho_{\ell}^4(r)$

$n =$	$=$			2	
$r=0$			6		
		$\overline{2}$		-2	
		$\left(\right)$	-2		
Q		-2		$\mathcal{D}_{\mathcal{L}}$	

Table 3. $\rho_{\ell}^5(r)$

$n=5$	$\ell=0$		$\mathcal{D}_{\mathcal{A}}$			5
$r=0$		5	10	10	5	
		२	$\mathcal{D}_{\mathcal{L}}$	- 2	\mathbf{Q}	
2			-2	-2		
3			-2	$\overline{2}$		
		-3	2	2	-3	
ь					5	

Table 4. $\rho_{\ell}^6(r)$

Table 5. $\rho_{\ell}^7(r)$

Table 6. $\rho_{\ell}^{8}(r)$

$n=8$	ι ι \prime $\ell=0$		$\overline{2}$	3	4	5			8
$r=0$		8	28	56	70	56	28	8	
		6	14	14	$\overline{0}$	-14	-14	-6	-1
$\overline{2}$		$\overline{4}$	4	-4	-10	-4	4	4	1
3		$\overline{2}$	-2	-6	$\overline{0}$	6	$\overline{2}$	-2	-1
4		$\overline{0}$	-4	θ	6	$\boldsymbol{0}$	-4	$\boldsymbol{0}$	1
$\bf 5$		-2	-2	6	$\overline{0}$	-6	$\overline{2}$	$\overline{2}$	-1
6		-4	4	4	-10	4	4	-4	
		-6	14	-14	$\overline{0}$	14	-14	6	-1
8		-8	28	-56	-70	-56	28	-8	1

Table 7. $\rho_{\ell}^9(r)$

$n=9$	\cdot \cdot \cdot \cdot $\ell=0$		2	3	4	5	6		8	9
$r=0$		9	36	84	126	126	84	36	9	
		7	20	28	14	-14	-28	-20	-7	
$\overline{2}$		5	8	$\overline{0}$	-14	-14	$\overline{0}$	8	5	
3		3	θ	-8	-6	6	8	θ	-3	
4		1	-4	-4	6	6	-4	-4		
5		-1	-4	4	6	-6	-4	4	1	-1
6		-3	θ	8	-6	-6	8	Ω	-3	
		-5	8	θ	-14	14	$\overline{0}$	-8	5	
8		-7	20	-28	14	14	-28	20	-7	
9		-9	36	-84	126	-126	84	-36	9	

Table 8. $\rho_{\ell}^{10}(r)$

