

Exploring the anisotropic gravitational wave background from all-sky mock gravitational wave event catalogues

ZHENCHENG LI,^{1,2} ZHEN JIANG,¹ YUN LIU,^{1,2} XI-LONG FAN,³ LIANG GAO,^{1,2,4,5} YUN CHEN,^{1,2} AND TENGPENG XU^{1,2}

¹*National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, China*

²*School of Astronomy and Space Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China*

³*School of Physics and Technology, Wuhan University, Wuhan 430072, China*

⁴*School of Physics and Astronomy, Beijing Normal University, Beijing, 100875, China*

⁵*School of Physics and Microelectronics, Zhengzhou University, Zhengzhou 450001, China*

ABSTRACT

Anisotropic stochastic gravitational wave background (SGWB) serves as a potential probe of the large-scale structure (LSS) of the universe. In this work, we explore the anisotropic SGWB from local ($z < \sim 0.085$) merging stellar mass compact binaries, specifically focusing on merging stellar binary black holes, merging neutron-star–black-hole binaries, and merging binary neutron stars. The analysis employs seven all-sky mock lightcone gravitational wave event catalogues, which are derived from the Millennium simulation combined with a semi-analytic model of galaxy formation and a binary population synthesis model. We calculate the angular power spectra C_ℓ at multipole moments ℓ , expressed as $\log_{10}[\ell(\ell + 1)C_\ell/(2\pi)]$, based on the skymaps of the overdensity δ_{GW} in the anisotropic SGWB. The spectra for all three source types exhibit an approximately linear increase with $\log_{10}\ell$ at higher ℓ (e.g., $\ell > \sim 30 - 300$) in seven catalogues, with a characteristic slope of ~ 2 . The spectra of seven catalogues exhibit considerable variations, arising from fluctuations in spatial distribution, primarily in the radial distribution, of nearby sources (e.g., < 50 Mpc/h). After subtracting these nearby sources, the variations become much smaller and the spectra for the three source types become closely aligned (within discrepancies of a factor of ~ 2 across $\ell = 1 - 1000$ for all catalogues). We also find that including further sources results in a rapid decrease in the anisotropy.

Keywords: gravitational wave, anisotropy

1. INTRODUCTION

Various gravitational wave (GW) sources, such as merging supermassive ($M > 10^7 M_\odot$) binary black holes (SMBBHs), merging stellar binary black holes (BBHs), merging neutron-star–black-hole binaries (NS-BHs), merging binary neutron stars (BNSs), contribute to the total stochastic GW background (SGWB). For the summary of the sources, see recent reviews, e.g. Christensen (2019) and Renzini et al. (2022). Recently, pulsar timing array (PTA) collaborations, including EPTA (Antoniadis et al. 2023), NANOGrav (Agazie et al. 2023a), PPTA (Reardon et al. 2023), and CPTA (Xu et al. 2023), have provided strong ev-

idence for the existence of the SGWB. Furthermore, the anticipated detection of the SGWB by both ground-based detectors, such as the advanced Laser Interferometer Gravitational-Wave Observatory (aLIGO) (Harry & LIGO Scientific Collaboration 2010) and the Einstein Telescope (ET) (Punturo et al. 2010), and space-based detectors, such as the Laser Interferometer Space Antenna (LISA) (Amaro-Seoane et al. 2017), is forthcoming, promising significant advancements in our understanding of the universe.

The SGWB can be divided into isotropic and anisotropic parts. Beyond the extensively studied isotropic part (e.g. Abbott et al. 2021; Bavera et al. 2022; Agazie et al. 2023b), the anisotropic part provides additional insights, particularly concerning the spatial distribution of GW sources. Several studies have explored the anisotropic SGWB, for example, the anisotropy from merging SMBBHs (Mingarelli et al. 2017; Bécsy et al.

2022; Agazie et al. 2023c; Gardiner et al. 2024), and the anisotropy from merging BBHs, NSBHs, BNSs (Cusin et al. 2018; Jenkins et al. 2018, 2019; Capurri et al. 2021). Jenkins et al. (2018) and Jenkins et al. (2019) first use the mock lightcone GW event catalogue to investigate the anisotropy of merging stellar mass compact binaries. Specifically, they employ a post-processing analytical approach to compute the GW events within the galaxies contained in the all-sky mock lightcone galaxy catalogue sourced from Blaizot et al. (2005) and De Lucia & Blaizot (2007). However, their results suggest a significantly higher anisotropy level compared to the prediction of Cusin et al. (2018), particularly at very small scales. The discrepancies among these studies remain under discussion.

The GW event catalogues could be obtained from a semi-analytic model of galaxy formation, Galaxy Assembly with Binary Evolution (GABE) (Jiang et al. 2019), combined with a rapid binary population synthesis model COSMIC v3.3.0 (Breivik et al. 2020). Using the binary population data generated by COSMIC, merging BNSs, NSBHs and BBHs can be explored within the cosmological framework provided by GABE. For example, Li et al. (2024) calculates the total SGWB from these sources using this GABE-COSMIC methodology. However, the original event catalogue lacks information on the source positions within the celestial sphere. In this work, we use the original event catalogue to construct seven all-sky mock lightcone GW event catalogues, which incorporate source position information across the celestial sphere, to investigate the anisotropic SGWB from the local ($z < \sim 0.085$) merging BBHs, NSBHs, and BNSs.

The structure of this paper is as follows: In Section 2, we present the fundamental equations used to calculate the anisotropic SGWB and the angular power spectrum. Section 3 briefly outlines the GABE-COSMIC methodology to produce the GW event catalogues and details the methodology employed for generating mock lightcone GW event catalogues. The results, including the anisotropic SGWB and angular power spectra, are discussed in Section 4. Finally, the primary conclusions of the study are summarized in Section 5.

2. FORMALISM

2.1. Anisotropic SGWB from ground-based detector compact binaries

The SGWB from the observed direction \hat{e} is defined by the dimensionless energy density (Cusin et al. 2017; Jenkins et al. 2018):

$$\Omega_{\text{GW}}(f, \hat{e}) = \frac{1}{\rho_c} \frac{d^3 \rho_{\text{GW}}(f, \hat{e})}{d \ln(f) d^2 \Omega}, \quad (1)$$

where $\rho_c = 3H_0^2 c^2 / (8\pi G)$ is the critical energy density, $d^3 \rho_{\text{GW}}$ is the present-day energy density of GWs from the observed frequency interval $\{f, f + df\}$ and the solid angle element $d^2 \Omega$ centered on \hat{e} . The Ω_{GW} could be calculated using the Phinney formula (Phinney 2001) in the homogeneous and isotropic universe:

$$\Omega_{\text{GW}}(f, \hat{e}) = \frac{1}{4\pi \rho_c} \int \mathcal{N}(z, \hat{e}) \frac{1}{1+z} \left[\frac{dE(f_r)}{d \ln(f_r)} \right]_{f_r=(1+z)f} dz, \quad (2)$$

where $\mathcal{N}(z, \hat{e}) = dN(z, \hat{e}) / (dV_{c,e} dz)$ is the number of GW sources per comoving volume per redshift at observed direction \hat{e} ; dN is the number of GW sources at comoving volume element dV_c , redshift element dz , at observed direction \hat{e} ; $dE(f_r) / d \ln(f_r)$ is the energy spectrum of a single GW source in logarithmic interval, f is the observed frequency and $f_r = (1+z)f$ is the frequency in the cosmic rest frame of the GW source. In this work, we adopt the same energy spectrum as used in Zhu et al. (2011) and Ajith et al. (2008), assuming that this spectrum remains applicable for merging BNSs and NSBHs. By dividing the full-direction comoving element presented in Phinney (2001) by 4π , we can derive the comoving volume element at the observed direction \hat{e} as follows:

$$dV_{c,e} = \frac{cd_c^2}{H(z)} dz, \quad (3)$$

where $H(z) = H_0[\Omega_m(1+z)^3 + \Omega_\Lambda]^{0.5}$ and d_c is the comoving distance:

$$d_c(z) = \int_0^z \frac{cdz'}{H_0 \sqrt{\Omega_m(1+z')^3 + \Omega_\Lambda}}. \quad (4)$$

Using $dz/dt = (1+z)H(z)$, $\mathcal{N}(z, \hat{e})$ could be calculated using the merger rate of GW sources per cosmic time $R(z, \hat{e}) = dN(z, \hat{e})/dt$, at observed direction \hat{e} :

$$\mathcal{N}(z, \hat{e}) = \frac{dN(z, \hat{e})}{c(1+z)d_c^2 dt dz} = \frac{R(z, \hat{e})}{c(1+z)d_c^2 dz}, \quad (5)$$

We now extend the applicability of the above formulas to an inhomogeneous and anisotropic universe, assuming that all GW events occur within galaxies. Consequently, $\mathcal{N}(z, \hat{e})$ and $\Omega_{\text{GW}}(f, \hat{e})$ might vary across different directions in accordance with the distribution of galaxies. By substituting Eq. 5 into Eq. 2, we compute the direction-dependent SGWB by summing the contributions from

all galaxies within the galaxy catalogue:

$$\begin{aligned}\Omega_{\text{GW}}(f, \hat{e}) &= \frac{f}{4\pi c \rho_c} \sum_{k=1}^{k_{\text{tot}}} \sum_{m_1, m_2}^{m \times n} \frac{R_k(z_k, m_1, m_2)}{(1+z_k)d_{c,k}^2} \delta_K(\hat{e}, \hat{e}_k) \\ &\times \left[\frac{dE(f_r, m_1, m_2)}{df_r} \right]_{f_r=(1+z_k)f} \\ &= \sum_{k=1}^{k_{\text{tot}}} \Omega_{\text{GW,gal},k}(f) \delta_K(\hat{e}, \hat{e}_k),\end{aligned}\quad (6)$$

where m and n are the bin numbers of primary star mass m_1 and secondary star mass m_2 , respectively, which are determined by the division of the $m_1 \times m_2$ space; k is the index of the galaxies; k_{tot} is the total number of the galaxies in the catalogue; R_k is the merger rate of merging stellar compact binaries in the galaxy k at z_k in the primary mass-secondary mass space m_1, m_2 ; δ_K is the Kronecker delta function; \hat{e}_k is the direction of the galaxy k , and see Sec. 3.2 for details on the positions of our galaxies; $\Omega_{\text{GW,gal},k}$ is the contribution of the galaxy k to the SGWB. In this work, the R_k is approximated as the mean merger rate of galaxy k , denoted \bar{R}_k . Details regarding the calculation of \bar{R}_k can be found in Sec. 3. This study focuses exclusively on the anisotropy arising from the spatial distribution of GW sources. For a more comprehensive formalism that incorporates additional effects such as Doppler, Kaiser, and gravitational potential influences, see Cusin et al. (2017) and Bertacca et al. (2020). Note that some of these effects have a negligible impact on anisotropy. For example, the kinematic dipole resulting from the observer's peculiar motion has been reported as minor (Jenkins et al. 2018; Jenkins 2022).

In this work, the skymaps are constructed as HEALPix maps (Górski et al. 2005) composed of N_{pix} equal area pixels, where $N_{\text{pix}} = 12 \times N_{\text{side}}^2$ is the number of pixels of the HEALPix skymap, with N_{side} denoting the parameter utilized in the HEALPix framework. A pixel may correspond to a range of directions, specifically $\hat{e}_{i,1}, \hat{e}_{i,2}, \dots, \hat{e}_{i,n_i}$. In practice, $\Omega_{\text{GW}}(f, \hat{e})$ is computed as follows:

$$\Omega_{\text{GW}}(f, \hat{e}) = \frac{1}{\Delta\Omega} \sum_{n=1}^{n=n_i} \sum_{k=1}^{k_{\text{tot}}} \Omega_{\text{GW,gal},k}(f) \delta_K(\hat{e}_{i,n}, \hat{e}_k), \quad (7)$$

where i denotes the index of the pixel corresponding to the observed direction \hat{e} , n_i is the total number of the directions within the pixel and $\Delta\Omega$ is the pixel area.

The overdensity δ_{GW} in the anisotropic SGWB from ground-based detector compact binaries is defined as follows:

$$\delta_{\text{GW}}(f, \hat{e}) = \frac{\Omega_{\text{GW}}(f, \hat{e}) - \bar{\Omega}_{\text{GW}}(f)}{\bar{\Omega}_{\text{GW}}(f)}, \quad (8)$$

where $\bar{\Omega}_{\text{GW}}(f) = \frac{1}{4\pi} \int_{\mathcal{S}^2} \Omega_{\text{GW}}(f, \hat{e}) d^2\Omega$ is the mean SGWB over the sky, \mathcal{S}^2 is the surface of the celestial sphere.

2.2. Angular power spectrum

To quantify the anisotropy of the SGWB at various scales, the angular power spectrum C_ℓ is adopted in this work. The angular power spectrum C_ℓ quantifies the amplitude of statistical fluctuations in the angular power of the SGWB at scales $\theta \sim 180^\circ/\ell$, where ℓ denotes the multipole moment. The C_ℓ is calculated using the healpy package (Górski et al. 2005; Zonca et al. 2019). The over-density δ_{GW} at each frequency on the celestial sphere can be expanded as:

$$\delta_{\text{GW}}(\hat{e}) = \sum_{\ell, m} a_{\ell m} Y_{\ell m}(\hat{e}), \quad (9)$$

where $Y_{\ell m}$ are spherical harmonics and:

$$a_{\ell m} = \frac{4\pi}{N_{\text{pix}}} \sum_{i=0}^{N_{\text{pix}}-1} Y_{\ell m}^*(\hat{e}_i) \delta_{\text{GW}}(\hat{e}_i), \quad (10)$$

where i denotes the index of the pixel, and the superscript star is complex conjugation. Consequently, the angular power spectrum can be expressed as:

$$C_\ell = \frac{1}{2\ell+1} \sum_{m=-\ell}^{m=+\ell} |a_{\ell m}|^2. \quad (11)$$

We calculate C_ℓ up to $\ell_{\text{max}} = 3N_{\text{side}} - 1$. To maintain consistency with the previous literature, we multiply C_ℓ by $\ell(\ell+1)/(2\pi)$.

3. SIMULATION OF SOURCES

3.1. The GW events in the GABE-COSMIC methodology

GABE is a semi-analytic model of galaxy formation, specifically addressing the evolution of binaries. This model is based on the Millennium simulation (Springel et al. 2005), which has a simulation box volume of $(500/h)^3 \text{Mpc}^3$. In this simulation, a fiducial cosmological model with $\Omega_m = 0.25$, $\Omega_\Lambda = 0.75$ and $H_0 = 73 \text{km/s/Mpc}$ (WMAP1, Spergel et al. 2003) is adopted. Further details on the generation of merging compact binaries using the GABE-COSMIC methodology can be found in Li et al. (2024), and we utilize datasets identical to those described therein. Here, we provide a concise overview.

¹ See https://healpix.sourceforge.io/html/fac_anafast.htm

We begin by discussing the formation of merging stellar mass compact binaries. Within the GABE framework, the total stellar populations in galaxies result from star formation events. Each star formation event triggers the birth of a simple stellar population (SSP) under conditions such as an unstable gaseous disk or galaxy mergers. The number, mass distribution, and delay time distribution of merging stellar mass compact binaries originating from each SSP are determined using the rapid binary population synthesis model COSMIC. The final population of merging stellar mass compact binaries within galaxies is a composite of these individual SSP contributions.

Upon executing GABE, we record for each galaxy the redshift, evolution history, spatial coordinates within the simulation box (x_0, x_1, x_2) , mean merger rates $\bar{R}_k(z_k, m_1, m_2)$, and the mass distribution of the merging BNSs, NSBHs, and BBHs contained within each galaxy, across 64 snapshots at discrete redshifts. The selection of the discrete redshifts is aligned with the Millennium simulation. The mean merger rate $\bar{R}_k = \Delta N_k / \Delta t$, where Δt represents the time interval between the current snapshot of galaxy k and its preceding snapshot, and ΔN_k denotes the number of merging sources of interest during Δt within the galaxy k .

3.2. All-sky mock lightcone GW event catalogue construction

Using the GW event catalogue derived from the GABE-COSMIC methodology, we create mock lightcone GW event catalogues restricted to the local universe, specifically for $d_c < 250 \text{ Mpc/h}$ ($z < \sim 0.085$). Our methodology is detailed below. The initial simulation box is replicated eight times, with the observer positioned at the center of the resultant new box, as illustrated in Fig. 1. The coordinates of objects within the new box are derived from the relation $(X_0, X_1, X_2) = (x_0 - aL, x_1 - bL, x_2 - cL)$, where a, b, c can take values of 0 or 1 and $L = 500 \text{ Mpc/h}$. The surface of the past-lightcone of the observer, representing the observable mock universe, at $z = z_i$, corresponds to the spherical surface with the radius equal to the comoving distance $d_c(z_i)$.

A galaxy crosses the surface of the past-lightcone of the observer when its distance to the observer, given by $r = \sqrt{X_0^2 + X_1^2 + X_2^2}$, matches the comoving distance that light can travel from $z = 0$ to the redshift of the target galaxy, denoted as $d_c(z)$. As galaxies in the original catalogue from the GABE-COSMIC methodology are stored in discrete snapshots, the galaxy, along with its associated GW events, is included in the mock lightcone catalogue if it meets two criteria: 1) in the initial




Figure 1. The new box, generated by replicating the initial simulation box eight times. The observer is positioned at the center, specifically at coordinates $(0, 0, 0)$ Mpc/h, with the surface of the past-lightcone at redshift $z = z_i$ presented for reference.

snapshot at z_{ori} , $r_{\text{ori}} < d_c(z_{\text{ori}})$, and 2) in the subsequent snapshot at z_{des} , $r_{\text{des}} > d_c(z_{\text{des}})$.

To determine the exact time $t(z_{\text{obs}})$ of the galaxy as it crosses the surface of the past-lightcone between z_{ori} and z_{des} , two assumptions are employed: 1) the galaxy follows a straight-line trajectory between two adjacent snapshots, and 2) the comoving distance could be approximated to increase linearly with cosmic time $t(z)$ during the interval. These assumptions yield the following two equations:

$$\begin{cases} \frac{t(z_{\text{obs}}) - t(z_{\text{ori}})}{t(z_{\text{des}}) - t(z_{\text{ori}})} = \frac{r_{\text{obs}} - r_{\text{ori}}}{r_{\text{des}} - r_{\text{ori}}} \\ \frac{t(z_{\text{obs}}) - t(z_{\text{ori}})}{t(z_{\text{des}}) - t(z_{\text{ori}})} = \frac{d_c(z_{\text{obs}}) - d_c(z_{\text{ori}})}{d_c(z_{\text{des}}) - d_c(z_{\text{ori}})} \end{cases} \quad (12)$$

Consequently, $t(z_{\text{obs}})$ corresponds to the time at which the two equations intersect, with the condition that at this intersection $r_{\text{obs}} = d_c(z_{\text{obs}})$. The position of the galaxy and the mean merger rates of merging stellar mass compact binaries within the galaxy could be determined through linear interpolation:

$$X_{i,\text{obs}} = X_{i,\text{ori}} + \frac{t(z_{\text{obs}}) - t(z_{\text{ori}})}{t(z_{\text{des}}) - t(z_{\text{ori}})} \times (X_{i,\text{des}} - X_{i,\text{ori}}), \quad (13)$$

Table 1. Observer coordinates in the new box for the models r2' through r7', in addition to the observer coordinate (0, 0, 0) Mpc/h for the model 'r1'.

Model	r2	r3	r4
Pos/(Mpc/h)	(125, 0, 0)	(250, 0, 0)	(250, 125, 0)
Model	r5	r6	r7
Pos/(Mpc/h)	(250, 250, 0)	(250, 250, 125)	(250, 250, 250)

$$\begin{aligned}
 \bar{R}_{\text{obs}}(z_{\text{obs}}, m_1, m_2) &= \bar{R}_{\text{ori}}(z_{\text{ori}}, m_1, m_2) \\
 &+ \frac{t(z_{\text{obs}}) - t(z_{\text{ori}})}{t(z_{\text{des}}) - t(z_{\text{ori}})} \times (\bar{R}_{\text{des}}(z_{\text{des}}, m_1, m_2) \\
 &- \bar{R}_{\text{ori}}(z_{\text{ori}}, m_1, m_2)),
 \end{aligned}
 \tag{14}$$

where $i = 0, 1, 2$. The right ascension (RA) and declination (DEC) of the galaxy, which denote its directional orientation, are determined under the assumption that the coordinate system of the observer aligns with the equatorial coordinate system. By applying the aforementioned procedure to all galaxies with distances to the observer less than 250Mpc/h in the original catalogue from the GABE-COSMIC methodology, we construct the all-sky mock lightcone GW event catalogue. We designate the all-sky mock lightcone GW event catalogue, constructed with the observer positioned at coordinates (0, 0, 0) Mpc/h as described above, as the 'r1' model. In addition to the r1 model, we change the observer's location to generate six additional mock catalogues, labeled from 'r2' to 'r7' models. The coordinates of the observers for these models are detailed in Table. 1. We select the r2 model as our fiducial model, as it demonstrates the most significant anisotropy in the merging BBHs among the seven models (see results in Sec. 4.1).

4. RESULTS

4.1. The angular power spectra

The HEALPix skymap of the overdensity δ_{GW} in the anisotropic SGWB from the merging BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz with $N_{\text{side}} = 512$ for the r2 model is shown in Fig. 2. The overdensity distinctly traces the large-scale structure (LSS) of the universe, highlighting the filaments, nodes, and voids of the cosmic web. The skymaps for merging BNSs and NSBHs display similar characteristics, which are not presented here.

The upper panel of Fig. 3 presents the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BNSs, NSBHs and BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz respectively, with $N_{\text{side}} = 512$, for the seven models, ranging from r1 to r7. All the spectra are contained

within the colored regions, with the spectra for models r2 and r4 distinctly highlighted. The spectra for all three source types exhibit relatively flat shapes below $\ell \sim 10$, followed by steep upward trends. For higher ℓ (e.g., $\ell > \sim 30 - 300$), the trends are especially typical, with slopes $\beta \approx 2$ in logarithmic space. The spectra across our seven catalogues exhibit considerable variations. For example, the spectra for merging BBHs exhibit a variation spanning approximately a factor of 2 to one order of magnitude for $\ell = 1$ to ~ 80 and about 1–2 orders of magnitude for $\ell = \sim 80$ to 1000. Our results are not directly comparable to those of Jenkins et al. (2018) and Jenkins et al. (2019) for several reasons. First, the mock catalogue used by Jenkins et al. (2018) extends to $z \sim 0.78$, significantly surpassing the redshift range of the catalogue in our study. Consequently, the spectra derived from closer sources may be higher, as discussed later. Additionally, their catalogue includes only galaxies with apparent AB magnitudes limited to 18 in the r filter from SDSS, whereas our catalogue encompasses all galaxies. However, we present a comparison here to highlight certain commonalities. Our spectra exhibit shapes similar to those reported by Jenkins et al. (2018) and Jenkins et al. (2019) for higher ℓ (e.g., $\ell > \sim 30 - 300$). Their spectra for merging BBHs² are generally situated near the lower boundary of the spectrum range for merging BBHs we obtained. We also conduct a brief comparison with the findings of Cusin et al. (2018), although our results are not directly comparable for several reasons. In addition to the limited redshift range explored in our study, differences in the GW energy spectrum and the models of merging stellar mass compact binaries might also contribute to the disparity. The lower boundary of our spectra for merging BBHs exhibits significantly higher amplitudes (> 2 orders of magnitude for the majority of ℓ) compared to the spectrum in Cusin et al. (2018)³.

The distribution of several nearby sources might contribute significantly to the fluctuations in anisotropy. To illustrate this effect, the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BNSs, NSBHs and BBHs between 50 and 250 Mpc/h at 60 Hz are shown respectively in the lower panel of Fig. 3. After the subtraction of the nearby sources, the growth of the spectra relatively slows down within the range

² The spectrum reported by Jenkins et al. (2018) includes all three types of merging stellar-mass compact binaries but introduces only a $\mathcal{O}(1)$ difference, as noted by the authors.

³ In this study, the comparison is made with the reproduced spectrum of Cusin et al. (2018), presented in Fig. 2 of Jenkins et al. (2019) as the 'CMBquick + Halofit' model at 65.75 Hz.




Figure 2. The HEALPix skymap of the overdensity δ_{GW} in the anisotropic SGWB from the merging BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz with $N_{\text{side}} = 512$ for the r2 model.

$\sim 10 < \ell < \sim 600$, yet they continue to exhibit the typical sharp shapes for $\ell > \sim 600$. The deviation in the spectra for the r1 to r7 models is significantly reduced across $\ell = 1 - 1000$. The spectra for all three merging types exhibit a variation of less than a factor of ~ 4 for $\ell = 1$ to ~ 20 , and less than a factor of ~ 2 for $\ell = \sim 20$ to 1000. The sharp reduction in deviation after the subtraction indicates that the significant discrepancies observed in the spectra of the upper panel of Fig. 3 are primarily attributable to the nearby sources. For each of seven mock catalogues, approximately 10^5 galaxies are situated within 50 Mpc/h, accounting for roughly 0.5–1% of the total galaxy population. However, the SGWB scales as d_c^{-2} , as indicated by Eq.6. Consequently, nearer sources contribute more significantly to the SGWB. The deviation regions of the spectra after the subtraction of nearby sources exhibit greater similarity to the error regions described in Jenkins et al. (2018) and Jenkins et al. (2019) for higher ℓ (e.g., $\ell > \sim 20$). However, the shapes of the spectra more closely resemble the results reported by Cusin et al. (2018) within the range $\sim 10 < \ell < \sim 100$. Note that, to enable a more reasonable comparison with the exist-

ing literature in the future work, in addition to using a larger simulation box, several other factors should be addressed. For example, when comparing with Jenkins et al. (2019), it is essential to apply a magnitude filter to our galaxy catalogues. Furthermore, understanding the influence of the GW energy spectra and the models of merging stellar mass compact binaries on anisotropy is essential for comparisons with Cusin et al. (2018).

From the observational perspective, the contribution to the total SGWB from nearby sources would be the first to be subtracted by future GW detectors, as these discrete events are the most readily detectable. This process alleviates the challenge of accurately predicting the angular power spectra from mock GW event catalogues. After the subtraction of the nearby sources, the spectra for merging BNSs, NSBHs and BBHs at 60 Hz display notable similarities (within discrepancies of a factor of ~ 2 across $\ell = 1 - 1000$ for all catalogues). This result suggests that the anisotropy is primarily influenced by the distribution of GW source hosts and might serve as a valuable probe of the LSS of the universe.




Figure 3. Upper panel: The angular power spectra of the overdensity δ_{GW} in the SGWB from merging BNSs, NSBHs and BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz respectively, with $N_{\text{side}} = 512$, for the seven models, ranging from r1 to r7. All the spectra are contained within the colored regions, with the spectra for models r2 and r4 distinctly highlighted. Lower panel: Similar to the upper panel, but with nearby ($< 50\text{Mpc}/h$) sources subtracted and exclusively highlights the spectrum for the r2 model. The approximate scale of ℓ is indicated on the upper axis, along with reference lines (grey dashed lines) exhibiting slopes of $\beta = 2$ (note that the intercepts α are variable) in both panels.

4.2. The effects of the radial distribution

The spatial distribution of the sources could be decomposed into two parts: the angular distribution on the celestial sphere, referred to as ‘azimuthal distribution’, and the distribution at different comoving distances, referred to as ‘radial distribution’. The influence of the radial distribution to the anisotropy is assessed through the implementation of the following smoothing

process in our work. We partition the comoving distances of galaxies, along with the merging stellar mass compact binaries they contain, into m equally spaced bins within the range $[a, b]$ Mpc/h, referred to as ‘ m -bin approximation’. Here, a and b denote the lower and upper boundaries of the comoving distances of the considered sources, respectively. For example, in the 5-bin approximation, the bin boundaries are defined as $[a, a + \frac{(b-a)}{5}, a + \frac{2(b-a)}{5}, a + \frac{3(b-a)}{5}, a + \frac{4(b-a)}{5}, b]$ Mpc/h. The comoving distances of the galaxies within each bin, denoted as $d_{c,i,1}, d_{c,i,2} \dots d_{c,i,n_i}$ are then replaced by the mean comoving distance for that bin, given by $\bar{d}_{c,i} = \sum_0^{n_i} d_{c,i}/n_i$, where i represents the bin index and n_i is the total number of the galaxies within the bin. The variation in the spectra between those calculated directly and those obtained using the ‘ m -bin approximation’ is interpreted as the anisotropy potentially influenced by the radial distribution. We anticipate that the extreme case of $m = 1$, corresponding to the 1-bin approximation, captures the anisotropy approximately arising from the azimuthal distribution. In this work, we use the approximation model to study sources up to $d_c = 50\text{Mpc}/h$ ($a = 0, b = 50$), between $50 - 250\text{Mpc}/h$ ($a = 50, b = 250$) and up to $d_c = 250\text{Mpc}/h$ ($a = 0, b = 250$).

To evaluate the impact of the radial distribution of both nearby and further sources on the anisotropy, we calculate the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs up to $d_c = 50\text{Mpc}/h$ and between $50 - 250\text{Mpc}/h$, derived from direct calculation, the 5-bin approximation model, and the 1-bin approximation model, respectively, with $N_{\text{side}} = 512$ at 60 Hz in the r2 model. The results are presented in Fig. 4. The colored regions illustrate the influence of the radial distribution. The variance of the spectra for the sources within $d_c = 50\text{Mpc}/h$ is within a factor of ~ 3 for $\ell = 1$ to ~ 10 , a factor of ~ 2 to approximately one order of magnitude for $\ell = \sim 10$ to ~ 50 , and $\sim 1-2$ orders of magnitude for $\ell = \sim 50$ to 1000. For the sources situated between $50 - 250\text{Mpc}/h$, the variance is within a factor of ~ 3 for $\ell = 1$ to ~ 10 , and within about 30% for $\ell = \sim 10$ to 1000. Consequently, when modeling the anisotropy with the subtraction of nearby sources, the radial distribution of sources becomes less significant for higher ℓ (e.g., $\ell > \sim 10$). The directly calculated spectrum for sources up to $d_c = 50\text{Mpc}/h$ exhibits the typical upward shape for $\ell > \sim 30$, as shown by the ‘BBHs r2’ spectrum in the upper panel of Fig. 3. However, the directly calculated spectrum for sources within the range of $50 - 250\text{Mpc}/h$ do not display this shape until $\ell > \sim 600$, as shown by the spectra in the lower panel of Fig. 3.




Figure 4. Upper panel: The angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs within $d_c = 50\text{Mpc}/h$ and between $d_c = 50 - 250\text{Mpc}/h$, derived from direct calculation, the 5-bin approximation model, and the 1-bin approximation model, respectively, with $N_{\text{side}} = 512$ at 60 Hz in the r2 model. The colored regions illustrate the influence of the radial distribution. Lower panel: The angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz, derived from the 5-bin approximation model, with $N_{\text{side}} = 512$, for the seven models, ranging from r1 to r7. All the spectra are contained within the colored regions, with the spectra for the r2 model distinctly highlighted. The approximate scale of ℓ is indicated on the upper axis, along with reference lines (grey dashed lines) exhibiting slopes of $\beta = 2$.

Fig.4 also shows the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs up to $d_c = 250\text{Mpc}/h$ at 60 Hz, derived from the 5-bin approximation, with $N_{\text{side}} = 512$, for the seven models, ranging from r1 to r7. All the spectra are contained within the blue regions, with the spectra for the r2 model distinctly

highlighted. The spectra exhibit minimal variation (less than a factor of ~ 3.5) across our seven mock catalogues for $\ell > \sim 10$. This observation also suggests that the significant deviations in the spectra presented in the upper panel of Fig. 3 for $\ell > \sim 10$ might primarily arise from fluctuations in the radial distribution of the nearby sources. The fluctuations might arise from Poisson noise associated with the radial distribution. The primary term contributing to the radial dependence in Eq. 6, d_c^{-2} , originates from the calculation of the comoving number density \mathcal{N} in Eq. 5. Consequently, the radial distribution of sources represents the number density of galaxies within each comoving volume element $dV_{c,e}$ at various redshifts, which inherently includes Poisson noise. When the number of galaxies is small (e.g. $\sim 10^5$ within 50 Mpc/h in our models), this noise might predominate in the estimation of the anisotropic SGWB.

We note that the spectra for both merging BBHs up to $d_c = 50\text{Mpc}/h$ and between 50 – 250Mpc/h exhibit typical shapes for higher ℓ (e.g. $\ell > \sim 300 - 600$) even under the extreme 1-bin approximation. Consequently, other factors might contribute to the Poisson noise. The typical shapes of our spectra exhibit slopes similar to those of the modeled Poisson noise spectra presented in Cusin et al. (2019). Additionally, the spectral behavior aligns with the predictions made in Cusin et al. (2019), which suggest that the high ℓ (e.g., $\ell > \sim 50$ to ~ 600 , based on the different choice of integral cut-off, shown in their Fig. 9.) portion of the angular power spectrum is predominantly influenced by Poisson noise arising from the azimuthal distribution of mock catalogues. This might indicate that the Poisson noise from the azimuthal distribution is entangled within our spectra.

4.3. The effects of including farther sources

Fig. 5 presents the ratio of the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs up to various comoving distances, relative to the angular power spectrum of the overdensity δ_{GW} in the SGWB from merging BBHs up to 50 Mpc/h, at 60 Hz with $N_{\text{side}} = 512$ in the r2 model. As sources at higher redshifts are included, the ratio decreases across $\ell = 1 - 1000$. For instance, the contribution from sources beyond 100 Mpc to the total anisotropy is expected to be less than $\sim 30 - 40\%$. Meanwhile, the rate of decrease seems to slow. In comparison to the spectrum that incorporates sources up to 100 Mpc/h ($C_{\ell,100\text{Mpc}/h}$), the spectrum that includes sources up to 150 Mpc/h ($C_{\ell,150\text{Mpc}/h}$) shows a reduction of approximately 1.7 – 2.5 times across the range $\ell = 1 - 1000$. However, when comparing the spectrum with sources up to 200 Mpc/h ($C_{\ell,200\text{Mpc}/h}$) to that with sources up to




Figure 5. The ratio of the angular power spectra of the overdensity δ_{GW} in the SGWB from merging BBHs up to various comoving distances, relative to the angular power spectrum of the overdensity δ_{GW} in the SGWB from merging BBHs up to 50 Mpc/h, at 60 Hz with $N_{\text{side}} = 512$ in the r2 model. The approximate scale of ℓ is indicated on the upper axis.

250 Mpc/h ($C_{\ell, 250\text{Mpc/h}}$), the latter exhibits a decrease of roughly 1.3 – 1.6 times across the range $\ell = 1 - 1000$. Although a larger simulation box is necessary for a comprehensive exploration of the universe, our findings establish an upper limit for the anisotropy.

5. SUMMARY AND DISCUSSION

Anisotropic SGWB could serve as a probe of the spatial distribution of GW sources. We explore the anisotropic SGWB from local merging BBHs, NSBHs, and BNSs ($d_c < 250\text{Mpc/h}$) by constructing seven all-sky mock lightcone GW event catalogues. The HEALPix skymaps of the overdensity δ_{GW} in the anisotropic SGWB were generated using these mock catalogues. The HEALPix skymap of the r2 model for merging BBHs shown in Fig. 2, serves as an example to illustrate the anisotropic distribution of the GW sources.

Based on the HEALPix skymaps, the angular power spectra in term of $\log_{10}[\ell(\ell+1)C_{\ell}/(2\pi)]$, are calculated to quantify the anisotropy level of the SGWB from merging stellar mass compact binaries. As shown in the upper panel of Fig. 3, the spectra for all three source types exhibit a typical shape at higher ℓ (e.g., $\ell > \sim 30 - 300$) across all catalogues, featuring an approximately linear increase with $\log_{10}\ell$: $\log_{10}[\ell(\ell+1)C_{\ell}/(2\pi)] \propto \beta(\log_{10}\ell)$, with a characteristic slope of $\beta \sim 2$. The spectra for our seven catalogues exhibit considerable variations. For example, the spectra for merging BBHs in different cata-

logues vary by approximately a factor of 2 to one order of magnitude for $\ell = 1$ to ~ 80 , and by about 1–2 orders of magnitude for $\ell = \sim 80$ to 1000. The fluctuation in the distribution of the nearby (e.g. < 50 Mpc/h) sources should contribute to the variations of the spectra for merging BBHs, NSBHs and BNSs. Subtracting these nearby sources results in significant reductions of the variations of the spectra. Shown in the lower panel of Fig. 3, after the subtraction of the nearby sources, all spectra vary within a factor of ~ 4 for $\ell = 1$ to ~ 20 , and within a factor of ~ 2 for $\ell = \sim 20$ to 1000. The spectra for the three source types then become closely aligned (within discrepancies of a factor of ~ 2 across $\ell = 1 - 1000$ for all catalogues). This result indicates that the SGWB anisotropy is primarily influenced by the distribution of the hosts of GW sources and could be used as a prob of LSS of the universe.

We further examine the contribution of the radial distribution of GW sources to the anisotropy of the SGWB. The results for merging BBHs are shown in Figs. 4 and 5. The random distance of the nearby sources makes the main contribution to the anisotropy of the SGWB (the red region in the upper panel of Figs. 4). After the subtraction of the nearby sources, the radial distribution has an insignificant effect on the spectra (implied by the green region in the upper panel of Fig.4), and the anisotropy from the azimuthal distribution (approximately estimated using the 5-bin approximation) of sources varies within ~ 3.5 times across our seven mock catalogues (the blue region in the lower panel of Fig.4). Therefore, the considerable spectral variations for $\ell > \sim 30$ shown in the upper panel of the Fig. 3 could be attributed to the radial distribution of the nearby sources. This random radial distribution of nearby sources might be the main Poisson noise, and contribute to the typical spectral shape as shown in Cusin et al. (2019). However, even with the extreme 1-bin approximation, namely the anisotropy approximately originating from the azimuthal distribution, the spectra continue to exhibit typical shapes for higher ℓ (e.g. $\ell > \sim 300 - 600$). This indicate that the variation of spectra for the seven mock catalogues also include the Poisson noises from the azimuthal distribution. We also find that including further sources results in a rapid decrease in the anisotropy. However, the rates of decrease seem to diminish as sources at higher z are included.

Besides of above results from the spectrum for the merging BBHs shown in Fig.4 and Fig.5, similar conclusions could also be obtained for both merging NSBHs and BNSs results. Furthermore, we investigate the effects of the observation frequency and the angular resolution of skymap on the spectra. We find that dur-

ing the most sensitive observation frequency for LIGO, ranging from 20 to 100 Hz, the spectra exhibit non negligible variation. The spectra calculated from the HEALPix skymaps at different resolutions, specifically with $N_{\text{side}} = 32, 64, 128, 256, 512$, are also consistent across the various configurations.

ACKNOWLEDGMENTS

This work has been supported by the National Natural Science Foundation of China (Nos.11988101, 11922303, 12033008 and 11673031), the K. C. Wong Education Foundation, the National Key Research and Development Program of China (Grant No.2020YFC2201400, SQ2021YFC220045-03).

REFERENCES

- Abbott, R., Abbott, T. D., Abraham, S., et al. 2021, *PhRvD*, 104, 022004, doi: [10.1103/PhysRevD.104.022004](https://doi.org/10.1103/PhysRevD.104.022004)
- Agazie, G., Anumarlapudi, A., Archibald, A. M., et al. 2023a, *ApJL*, 951, L8, doi: [10.3847/2041-8213/acdac6](https://doi.org/10.3847/2041-8213/acdac6)
- . 2023b, *ApJL*, 952, L37, doi: [10.3847/2041-8213/ace18b](https://doi.org/10.3847/2041-8213/ace18b)
- . 2023c, *ApJL*, 956, L3, doi: [10.3847/2041-8213/acf4fd](https://doi.org/10.3847/2041-8213/acf4fd)
- Ajith, P., Babak, S., Chen, Y., et al. 2008, *PhRvD*, 77, 104017, doi: [10.1103/PhysRevD.77.104017](https://doi.org/10.1103/PhysRevD.77.104017)
- Amaro-Seoane, P., Audley, H., Babak, S., et al. 2017, arXiv e-prints, arXiv:1702.00786, doi: [10.48550/arXiv.1702.00786](https://doi.org/10.48550/arXiv.1702.00786)
- Antoniadis, J., Arumugam, P., Arumugam, S., et al. 2023, arXiv e-prints, arXiv:2306.16214, doi: [10.48550/arXiv.2306.16214](https://doi.org/10.48550/arXiv.2306.16214)
- Bavera, S. S., Franciolini, G., Cusin, G., et al. 2022, *A&A*, 660, A26, doi: [10.1051/0004-6361/202142208](https://doi.org/10.1051/0004-6361/202142208)
- Bécsy, B., Cornish, N. J., & Kelley, L. Z. 2022, *ApJ*, 941, 119, doi: [10.3847/1538-4357/aca1b2](https://doi.org/10.3847/1538-4357/aca1b2)
- Bertacca, D., Ricciardone, A., Bellomo, N., et al. 2020, *PhRvD*, 101, 103513, doi: [10.1103/PhysRevD.101.103513](https://doi.org/10.1103/PhysRevD.101.103513)
- Blaizot, J., Wadadekar, Y., Guiderdoni, B., et al. 2005, *MNRAS*, 360, 159, doi: [10.1111/j.1365-2966.2005.09019.x](https://doi.org/10.1111/j.1365-2966.2005.09019.x)
- Breivik, K., Coughlin, S., Zevin, M., et al. 2020, *ApJ*, 898, 71, doi: [10.3847/1538-4357/ab9d85](https://doi.org/10.3847/1538-4357/ab9d85)
- Capurri, G., Lapi, A., Baccigalupi, C., et al. 2021, *JCAP*, 2021, 032, doi: [10.1088/1475-7516/2021/11/032](https://doi.org/10.1088/1475-7516/2021/11/032)
- Christensen, N. 2019, *Reports on Progress in Physics*, 82, 016903, doi: [10.1088/1361-6633/aae6b5](https://doi.org/10.1088/1361-6633/aae6b5)
- Cusin, G., Dvorkin, I., Pitrou, C., & Uzan, J.-P. 2018, *PhRvL*, 120, 231101, doi: [10.1103/PhysRevLett.120.231101](https://doi.org/10.1103/PhysRevLett.120.231101)
- . 2019, *PhRvD*, 100, 063004, doi: [10.1103/PhysRevD.100.063004](https://doi.org/10.1103/PhysRevD.100.063004)
- Cusin, G., Pitrou, C., & Uzan, J.-P. 2017, *PhRvD*, 96, 103019, doi: [10.1103/PhysRevD.96.103019](https://doi.org/10.1103/PhysRevD.96.103019)
- De Lucia, G., & Blaizot, J. 2007, *MNRAS*, 375, 2, doi: [10.1111/j.1365-2966.2006.11287.x](https://doi.org/10.1111/j.1365-2966.2006.11287.x)
- Gardiner, E. C., Kelley, L. Z., Lemke, A.-M., & Mitridate, A. 2024, *ApJ*, 965, 164, doi: [10.3847/1538-4357/ad2be8](https://doi.org/10.3847/1538-4357/ad2be8)
- Górski, K. M., Hivon, E., Banday, A. J., et al. 2005, *ApJ*, 622, 759, doi: [10.1086/427976](https://doi.org/10.1086/427976)
- Harry, G. M., & LIGO Scientific Collaboration. 2010, *Classical and Quantum Gravity*, 27, 084006, doi: [10.1088/0264-9381/27/8/084006](https://doi.org/10.1088/0264-9381/27/8/084006)
- Jenkins, A. C. 2022, arXiv e-prints, arXiv:2202.05105, doi: [10.48550/arXiv.2202.05105](https://doi.org/10.48550/arXiv.2202.05105)
- Jenkins, A. C., O’Shaughnessy, R., Sakellariadou, M., & Wysocki, D. 2019, *PhRvL*, 122, 111101, doi: [10.1103/PhysRevLett.122.111101](https://doi.org/10.1103/PhysRevLett.122.111101)
- Jenkins, A. C., Sakellariadou, M., Regimbau, T., & Slezak, E. 2018, *PhRvD*, 98, 063501, doi: [10.1103/PhysRevD.98.063501](https://doi.org/10.1103/PhysRevD.98.063501)
- Jiang, Z., Wang, J., Gao, L., et al. 2019, *Research in Astronomy and Astrophysics*, 19, 151, doi: [10.1088/1674-4527/19/10/151](https://doi.org/10.1088/1674-4527/19/10/151)
- Li, Z., Jiang, Z., Fan, X.-L., et al. 2024, *MNRAS*, 527, 5616, doi: [10.1093/mnras/stad3576](https://doi.org/10.1093/mnras/stad3576)
- Mingarelli, C. M. F., Lazio, T. J. W., Sesana, A., et al. 2017, *Nature Astronomy*, 1, 886, doi: [10.1038/s41550-017-0299-6](https://doi.org/10.1038/s41550-017-0299-6)
- Phinney, E. S. 2001, arXiv e-prints, astro, doi: [10.48550/arXiv.astro-ph/0108028](https://doi.org/10.48550/arXiv.astro-ph/0108028)
- Punturo, M., Abernathy, M., Acernese, F., et al. 2010, *Classical and Quantum Gravity*, 27, 194002, doi: [10.1088/0264-9381/27/19/194002](https://doi.org/10.1088/0264-9381/27/19/194002)
- Reardon, D. J., Zic, A., Shannon, R. M., et al. 2023, *ApJL*, 951, L6, doi: [10.3847/2041-8213/acdd02](https://doi.org/10.3847/2041-8213/acdd02)
- Renzini, A. I., Goncharov, B., Jenkins, A. C., & Meyers, P. M. 2022, *Galaxies*, 10, 34, doi: [10.3390/galaxies10010034](https://doi.org/10.3390/galaxies10010034)
- Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175, doi: [10.1086/377226](https://doi.org/10.1086/377226)
- Springel, V., White, S. D. M., Jenkins, A., et al. 2005, *Nature*, 435, 629, doi: [10.1038/nature03597](https://doi.org/10.1038/nature03597)
- Xu, H., Chen, S., Guo, Y., et al. 2023, *Research in Astronomy and Astrophysics*, 23, 075024, doi: [10.1088/1674-4527/acdfa5](https://doi.org/10.1088/1674-4527/acdfa5)

Zhu, X.-J., Howell, E., Regimbau, T., Blair, D., & Zhu,
Z.-H. 2011, *ApJ*, 739, 86,

doi: [10.1088/0004-637X/739/2/86](https://doi.org/10.1088/0004-637X/739/2/86)

Zonca, A., Singer, L., Lenz, D., et al. 2019, *Journal of Open
Source Software*, 4, 1298, doi: [10.21105/joss.01298](https://doi.org/10.21105/joss.01298)