

HIGHER DIRECT IMAGES OF THE STRUCTURE SHEAF VIA THE HILBERT-CHOW MORPHISM.

YAO YUAN

ABSTRACT. Let X be a projective smooth surface over \mathbb{C} with $H^2(\mathcal{O}_X) = 0$. Let $M = M(L, \chi)$ be the moduli space of 1-dimensional semistable sheaves with determinant $\mathcal{O}_X(L)$ and Euler characteristic χ . We have the Hilbert-Chow morphism $\pi : M \rightarrow |L|$. We give explicit forms of the higher direct images $R^i\pi_*\mathcal{O}_M$ under some mild conditions on M and $|L|$. Our result shows that $R^i\pi_*\mathcal{O}_M$ are direct sums of line bundles. In particular, using our result one gets explicit formulas for the Euler characteristic of $\pi^*\mathcal{O}_{|L|}(m)$, which in $X = \mathbb{P}^2$ case was once conjectured by Chung-Moon.

Keywords: Moduli spaces of 1-dimensional sheaves on a surface, Hilbert-Chow morphism, abelian fibration, compactified Jacobian, higher direct image.

CONTENTS

1. Introduction.	2
1.1. The main result.	2
1.2. Acknowledgements.	3
2. Proof of the main result.	3
2.1. The relative compactified Jacobians.	3
2.2. Moduli spaces of 1-dimensional sheaves on a surface.	5
2.3. Proof of Theorem 1.1.	5
2.4. An equality in binomial coefficients.	7
References	8

1. INTRODUCTION.

1.1. The main result. We work over the complex number \mathbb{C} .

Let X be a projective smooth surface with K_X the canonical line bundle. Let L be an effective divisor class on X . Let $M(L, \chi)$ be the moduli space of 1-dimensional semistable sheaves with determinant $\mathcal{O}_X(L)$ and Euler characteristic χ . We have the Hilbert-Chow morphism

$$\pi : M(L, \chi) \rightarrow |L|, \mathcal{F} \mapsto \text{supp}(\mathcal{F})$$

sending each sheaf \mathcal{F} to its Fitting support. We write $M = M(L, \chi)$ for simplicity if there is no confusion on L and χ .

The main purpose of the paper is to study the higher direct image $R^i \pi_* \mathcal{O}_M$ on $|L|$. Denote by M^s the subset of M consisting of stable sheaves. Denote by $|L|^{int}$ ($|L|^{sm}$, resp.) the subset of $|L|$ consisting of integral curves (smooth curves, resp.). Let g_L be the arithmetic genus of curves in $|L|$ and $\ell := \dim |L|$. Our main theorem is as follows.

Theorem 1.1. *Let X be a surface with $h^1(\mathcal{O}_X) = q$ and $h^2(\mathcal{O}_X) = 0$. Let L be an effective divisor class with $L.K_X < 0$ and $h^1(\mathcal{O}_X(-L)) = 0$. Assume M has rational singularities with $M \setminus \pi^{-1}(|L|^{int})$ of codimension ≥ 2 and assume moreover $|L| \setminus |L|^{int}$ has codimension ≥ 2 and $|L|^{sm} \neq \emptyset$. Then for all i we have*

$$(1.1) \quad R^i \pi_* \mathcal{O}_M \cong \wedge^i \left(\mathcal{O}_{|L|}^{\oplus q} \oplus \mathcal{O}_{|L|}(-1)^{\oplus g_L - q} \right).$$

We write the following theorem as a special case for Theorem 1.1 with X rational.

Theorem 1.2. *Let X be a rational surface with $L.K_X < 0$. Assume M has rational singularities with $M \setminus \pi^{-1}(|L|^{int})$ of codimension ≥ 2 and assume moreover $|L| \setminus |L|^{int}$ has codimension ≥ 2 and $|L|^{sm} \neq \emptyset$. Then for all i we have*

$$(1.2) \quad R^i \pi_* \mathcal{O}_M \cong \mathcal{O}_{|L|}(-i)^{\oplus \binom{g_L}{i}}.$$

Corollary 1.3. *Let X, M, L, π be as in Theorem 1.2, then we have $H^i(M, \mathcal{O}_M) = 0$ for all $i > 0$ and*

$$\chi(M, \pi^* \mathcal{O}_{|L|}(m)) = \binom{m + \ell - g_L}{m} = \binom{m - L.K_X - 1}{m}.$$

Proof. By Theorem 1.2 we have $H^i(R^j\pi_*\mathcal{O}_M) = 0$ except for $i = j = 0$ and

$$\begin{aligned} \chi(M, \pi^*\mathcal{O}_{|L|}(m)) &= \sum_{i=0}^{g_L} (-1)^i \chi(R^i\pi_*\mathcal{O}_M(m)) \\ (1.3) \qquad \qquad \qquad &= \sum_{i=0}^{g_L} (-1)^i \binom{g_L}{i} \binom{m-i+\ell}{\ell} = P_{\ell-g_L, \ell-g_L}(m, g_L), \end{aligned}$$

where $P_{k,l}(m, n)$ is defined in (2.7). By Lemma 2.6 we have

$$\chi(M, \pi^*\mathcal{O}_{|L|}(m)) = \binom{m+\ell-g_L}{\ell-g_L} = \binom{m+\ell-g_L}{m}.$$

The last thing left to show is $\ell - g_L = -L.K_X - 1$. By assumption $\dim M = \dim \pi^{-1}(|L|^{int}) = g_L + \ell$. On the other hand since $L.K_X < 0$, $\pi^{-1}(|L|^{int})$ is smooth of the expected dimension $L^2 + 1$. We also have $g_L = \chi(\mathcal{O}_X(-L)) - \chi(\mathcal{O}_X) + 1 = \frac{L^2 + L.K_X}{2} + 1$. Therefore $\ell = \frac{L^2 - L.K_X}{2}$ and we are done. \square

Remark 1.4. *If X is a del Pezzo surface, then M always has rational singularities since it is a quotient of a smooth scheme. Moreover π is flat over M^s by Corollary 1.3 in [12]. Therefore $\text{codim}(M^s \setminus \pi^{-1}(|L|^{int})) = \text{codim}(|L| \setminus |L|^{int})$.*

Remark 1.5. *Corollary 1.3 applies to $X = \mathbb{P}^2$, $L = dH$ and $(\chi, d) = 1$ and proves the conjecture by Chung-Moon ([4, Conjecture 8.2] or [7, Conjecture 6.13]).*

Remark 1.6. *Let X be a K3 surface. If $M = M^s$, then it is smooth and π is a Lagrangian fibration. By Matsushita's result ([8, Theorem 1.3]) we have*

$$(1.4) \qquad \qquad \qquad R^i\pi_*\mathcal{O}_M \cong \Omega_{|L|}^i,$$

where $\Omega_{|L|}^i = \wedge^i \Omega_{|L|}^1$ with $\Omega_{|L|}^1$ the sheaf of differentials over $|L|$. We will give an explanation in Remark 2.5 on why there is a difference between (1.1) and (1.4).

We will prove our main result in the next section.

1.2. Acknowledgements. I would like to thank Qizheng Yin for introducing to me the work of Arinkin on compactified Jacobians. I thank the referees.

2. PROOF OF THE MAIN RESULT.

2.1. The relative compactified Jacobians. In this subsection we prove a property on the higher direct images of the structure sheaf of relative compactified Jacobians to the base.

Let $f : \mathcal{C} \rightarrow \mathcal{B}$ be a flat proper family of integral curves with planar singularities of genus $g \geq 1$. Let $\mathcal{C}^{sm} \subset \mathcal{C}$ be the open subscheme where f is smooth. Let $h^d : \overline{\mathcal{J}}^d \rightarrow \mathcal{B}$ be the relative compactified Jacobian (of degree d) and let $\mathcal{J}^d \subset \overline{\mathcal{J}}^d$ be the relative Jacobian, i.e. for each $b \in \mathcal{B}$, \mathcal{J}_b^d parametrizes line bundles with degree d on \mathcal{C}_b . We assume that \mathcal{C}, \mathcal{B} and $\overline{\mathcal{J}}^d$ are smooth.

Proposition 2.1. *With notations as above, we have*

$$(2.1) \quad h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong \mathcal{O}_{\mathcal{B}};$$

$$(2.2) \quad R^1 h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong R^1 f_* \mathcal{O}_{\mathcal{C}};$$

and

$$(2.3) \quad R^i h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong \wedge^i R^1 h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d}, \text{ for any } i = 1, \dots, g.$$

Proof. By Altman-Iarrobino-Kleiman's result in [1], every fiber of h^d is projective and irreducible and hence $h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong \mathcal{O}_{\mathcal{B}}$.

To prove (2.2), let us first assume that there is a section $s : \mathcal{B} \rightarrow \mathcal{C}^{sm}$. Hence $\overline{\mathcal{J}}^d$ are naturally isomorphic to each other. We may let $d = 0$ and denote $\overline{\mathcal{J}} = \overline{\mathcal{J}}^0, h = h^0$. By [3, Theorem B] we see that $\overline{\mathcal{J}} \cong \overline{\text{Pic}}^0(\overline{\mathcal{J}})$. Therefore we have a natural embedding $\iota : \mathcal{C} \hookrightarrow \overline{\text{Pic}}^0(\overline{\mathcal{J}}) \cong \overline{\mathcal{J}}$ as schemes over \mathcal{B} . The surjective morphism $\mathcal{O}_{\overline{\mathcal{J}}} \rightarrow \mathcal{O}_{\mathcal{C}}$ induces the following morphism

$$R^1 h_* \mathcal{O}_{\overline{\mathcal{J}}} \rightarrow R^1 f_* \mathcal{O}_{\mathcal{C}},$$

which is an isomorphism restricted to all fibers by [2, Theorem 1.2 (ii), Proposition 6.1]. Hence $R^1 h_* \mathcal{O}_{\overline{\mathcal{J}}} \cong R^1 f_* \mathcal{O}_{\mathcal{C}}$.

If there is no section $s : \mathcal{B} \rightarrow \mathcal{C}^{sm}$, we may do an étale surjective base change $\sigma : \mathcal{B}' \rightarrow \mathcal{B}$ such that $f' : \mathcal{C}^{sm'} \rightarrow \mathcal{B}'$ has a section. Then

$$R^1 h_*^{d'} \mathcal{O}_{\overline{\mathcal{J}}^{d'}} \cong \sigma^* R^1 h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong R^1 f'_* \mathcal{O}_{\mathcal{C}'} \cong \sigma^* R^1 f_* \mathcal{O}_{\mathcal{C}}.$$

Therefore $R^1 h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d} \cong R^1 f_* \mathcal{O}_{\mathcal{C}}$ because σ is étale and surjective.

(2.3) is just a relative version of Arinkin's result ([2, Theorem 1.2 (ii)]). Firstly by [2, Theorem 1.2 (ii)] we have both $R^i h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d}$ and $\wedge^i R^1 h_*^d \mathcal{O}_{\overline{\mathcal{J}}^d}$ are locally free of the same rank, hence it is enough to show (2.3) holds in codimension 1. Therefore we may assume \mathcal{B} to be a curve.

Let $D \subset \mathcal{B}$ consist of finite reduced points such that \mathcal{C}_x is smooth for all $x \notin D$. Take a resolution $\eta : \tilde{\mathcal{J}}^d \rightarrow \overline{\mathcal{J}}^d$ such that η is an isomorphism restricted to $(h^d)^{-1}(\mathcal{B} \setminus D)$ and $(\eta \circ h^d)^{-1}D$ is a normal crossing divisor on $\tilde{\mathcal{J}}$. Since $(h^d)^{-1}(D)$ is reduced, we have that $(\eta \circ h^d)^{-1}D$ is reduced. We also have $R^i \eta_* \mathcal{O}_{\tilde{\mathcal{J}}^d} = 0$ for $i > 0$ and $\eta_* \mathcal{O}_{\tilde{\mathcal{J}}^d} \cong \mathcal{O}_{\overline{\mathcal{J}}^d}$ because $\overline{\mathcal{J}}^d$ is smooth. Therefore

it is enough to show (2.3) with h^d replaced by $\eta \circ h^d$. (2.3) holds over $\mathcal{B} \setminus D$ by the functoriality of variations of Hodge structures. Since $(\eta \circ h^d)^{-1}D$ is a reduced normal crossing divisor, the monodromy of the variations of the Hodge structures are unipotent. Therefore there are functorial canonical extensions of Hodge bundles (see [10, Theorem 6.16] or [9, Corollary 11.18]) and hence (2.3) extends to the whole \mathcal{B} (see also [8, Lemma 4.9]).

□

2.2. Moduli spaces of 1-dimensional sheaves on a surface. Now let $M = M(L, \chi)$ parametrize 1-dimensional semistable sheaves with determinant $\mathcal{O}_X(L)$ and Euler characteristic χ on a projective surface X . Let $\pi : M \rightarrow |L|$ be the Hilbert-Chow morphism. Let g_L be the arithmetic genus of curves in $|L|$. We have the following proposition on the dualizing sheaf ω_M of M .

Proposition 2.2. *Let $L.K_X < 0$ and $|L|^{sm} \neq \emptyset$, then the canonical line bundle over $\pi^{-1}(|L|^{int})$ is $\pi^*\mathcal{O}_{|L|}(1)^{\otimes L.K_X}|_{\pi^{-1}(|L|^{int})}$. In particular if M is normal and Cohen-Macaulay and $M \setminus \pi^{-1}(|L|^{int})$ has codimension ≥ 2 , then the dualizing sheaf ω_M of M is $\pi^*\mathcal{O}_{|L|}(1)^{\otimes L.K_X}$.*

Proof. The proof of the first statement is analogous to [11, Proposition C.0.15]. Notice that $\text{Ext}^2(\mathcal{F}, \mathcal{F}) = 0$ for any $\mathcal{F} \in \pi^{-1}(|L|^{int})$ by $L.K_X < 0$, and hence $\pi^{-1}(|L|^{int})$ is smooth.

Now if M is normal and Cohen-Macaulay, then ω_M is the push forward of the canonical line bundle of its smooth locus. Hence the proposition. □

Proposition 2.3. *Let $L.K_X < 0$ and $|L|^{sm} \neq \emptyset$. If M has rational singularities and $M \setminus \pi^{-1}(|L|^{int})$ has codimension ≥ 2 , then $R^i\pi_*\mathcal{O}_M = 0$ except for $i = 0, 1, \dots, g_L$, and $R^i\pi_*\mathcal{O}_M$ is locally free for all $i = 0, 1, \dots, g_L$.*

Proof. Since M has rational singularities, we can take a resolution $\xi : \widetilde{M} \rightarrow M$ such that ξ is birational, \widetilde{M} is smooth and $R^i\xi_*\omega_{\widetilde{M}} = R^i\xi_*\mathcal{O}_{\widetilde{M}} = 0$ for all $i > 0$ and $\xi_*\mathcal{O}_{\widetilde{M}} \cong \mathcal{O}_M$, $\xi_*\omega_{\widetilde{M}} \cong \omega_M$ where $\omega_{\widetilde{M}}$ is the canonical line bundle on \widetilde{M} . Therefore we may replace M by \widetilde{M} and assume that M is smooth.

By Kollár's result ([5, Theorem 2.1 (i) (ii)]), we have $R^i\pi_*\omega_M = 0$ except for $i = 0, 1, \dots, g_L$, and $R^i\pi_*\omega_M$ is torsion free for all $i = 0, 1, \dots, g_L$. On the other hand, $\omega_M \cong \pi^*\mathcal{O}_{|L|}(1)^{\otimes L.K_X}$ by Proposition 2.2. Hence $R^i\pi_*\mathcal{O}_M$ are torsion-free. Then by [6, Proposition 3.12], $R^i\pi_*\mathcal{O}_M$ are Cohen-Macaulay sheaves on $|L|$ hence locally free since $|L|$ is smooth. □

2.3. Proof of Theorem 1.1. In this subsection, we let X be a surface with $h^1(\mathcal{O}_X) = q$ and $h^2(\mathcal{O}_X) = 0$. Let L be an effective divisor class with

$h^1(\mathcal{O}_X(-L)) = 0$. Denote by $\mathcal{C}_L \subset X \times |L|$ the universal curve.

$$(2.4) \quad \begin{array}{ccc} \mathcal{C}_L & \hookrightarrow & X \times |L| \\ & \searrow q & \downarrow p \\ & X & |L| \end{array}$$

Lemma 2.4. $R^1 p_* \mathcal{O}_{\mathcal{C}_L} \cong \mathcal{O}_{|L|}^{\oplus q} \oplus \mathcal{O}_{|L|}(-1)^{\oplus gL-q}$.

Proof. We have an exact sequence on $X \times |L|$

$$(2.5) \quad 0 \rightarrow q^* \mathcal{O}_X(-L) \otimes p^* \mathcal{O}_{|L|}(-1) \rightarrow \mathcal{O}_{X \times |L|} \rightarrow \mathcal{O}_{\mathcal{C}} \rightarrow 0.$$

Push (2.5) forward to $|L|$ and we have

$$0 \rightarrow R^1 p_* \mathcal{O}_{X \times |L|} \rightarrow R^1 p_* \mathcal{O}_{\mathcal{C}_L} \rightarrow R^2 p_*(q^* \mathcal{O}_X(-L) \otimes p^* \mathcal{O}_{|L|}(-1)) \rightarrow 0.$$

Hence

$$0 \rightarrow \mathcal{O}_{|L|}^{\oplus q} \rightarrow R^1 p_* \mathcal{O}_{\mathcal{C}_L} \rightarrow \mathcal{O}_{|L|}(-1)^{\oplus gL-q} \rightarrow 0.$$

and

$$R^1 p_* \mathcal{O}_{\mathcal{C}_L} \cong \mathcal{O}_{|L|}^{\oplus q} \oplus \mathcal{O}_{|L|}(-1)^{\oplus gL-q}.$$

The lemma is proved. \square

Proof of Theorem 1.1. Since $\pi^{-1}(|L|^{int})$ and $|L|^{int}$ are smooth, π restricted to $\pi^{-1}(|L|^{int})$ is flat. Combine Proposition 2.1 and Lemma 2.4 and we have over $|L|^{int}$

$$(2.6) \quad R^i \pi_* \mathcal{O}_{\pi^{-1}(|L|^{int})} \cong R^i \pi_* \mathcal{O}_M|_{|L|^{int}} \cong \wedge^i \left(\mathcal{O}_{|L|}^{\oplus q} \oplus \mathcal{O}_{|L|}(-1)^{\oplus gL-q} \right) |_{|L|^{int}}.$$

Hence the theorem since $R^i \pi_* \mathcal{O}_M$ are locally free and $|L| \setminus |L|^{int}$ has codimension ≥ 2 . \square

Remark 2.5. If X is a K3 surface with L an effective divisor class. Denote by $\mathcal{C}_L \subset X \times |L|$ the universal curve. If $|L|^{int} \neq \emptyset$, then $|L| \cong \mathbb{P}^{gL}$ and we have the following exact sequence on $|L|$

$$0 \rightarrow R^1 p_* \mathcal{O}_{\mathcal{C}_L} \rightarrow R^2 p_*(q^* \mathcal{O}_X(-L) \otimes p^* \mathcal{O}_{|L|}(-1)) \rightarrow R^2 p_* \mathcal{O}_{X \times |L|} \rightarrow 0.$$

Therefore

$$0 \rightarrow R^1 p_* \mathcal{O}_{\mathcal{C}_L} \rightarrow \mathcal{O}_{|L|}(-1)^{gL+1} \rightarrow \mathcal{O}_{|L|} \rightarrow 0,$$

which is compatible with the fact that $R^1 \pi_* \mathcal{O}_M \cong \Omega_{|L|}^1$ for M smooth. So the main reason to cause the difference between (1.1) and (1.4) is whether the cohomology group $H^2(\mathcal{O}_X)$ vanishes.

2.4. An equality in binomial coefficients. In this subsection we prove an equality in binomial coefficients used in the proof of Corollary 1.3. Lemma 2.6 below maybe obvious to experts, but we still give a proof here. For $k, l, m, n \in \mathbb{Z}_{\geq 0}$ define

$$(2.7) \quad P_{k,l}(m, n) := \sum_{i=0}^n (-1)^i \binom{n}{i} \binom{m+n-i+k}{n+l}.$$

Lemma 2.6. *For any $k, l, m, n \in \mathbb{Z}_{\geq 0}$ we have*

$$P_{k,l}(m, n) = \binom{m+k}{l}.$$

Proof. We prove the lemma in several steps.

Step 1: We show that $P_{0,0}(m, n) = 1$ for all $m, n \in \mathbb{Z}_{\geq 0}$.

It is easy to see that $P_{0,0}(m, n) = P_{0,0}(n, m)$ and $P_{0,0}(0, n) = P_{0,0}(n, 0) = 1$ for all $n \in \mathbb{Z}_{\geq 0}$. Now we assume $P_{0,0}(m, n) = 1$ if either $n < n_0$ or $m < m_0$, it is enough to show $P_{0,0}(m_0, n_0) = 1$. Use the equality $\binom{m+1}{k} = \binom{m}{k} + \binom{m}{k-1}$ and we get

$$P_{0,0}(m_0, n_0) = P_{0,0}(m_0 - 1, n_0) - P_{0,0}(m_0 - 1, n_0 - 1) + P_{0,0}(m_0, n_0 - 1).$$

Hence by induction assumption we get $P_{0,0}(m_0, n_0) = 1$. We are done.

Step 2: We show that $P_{0,l}(m, n) = \binom{m}{l}$ for all $m, n \in \mathbb{Z}_{\geq 0}$.

It is easy to see $P_{0,l}(m, n) = 0$ for $m < l$ and $P_{0,l}(l, n) = 1 = \binom{l}{l}$. We assume $P_{0,l}(m, n) = \binom{m}{l}$ if either $l < l_0$ or $m < m_0$, it is enough to show $P_{0,l_0}(m_0, n) = \binom{m_0}{l_0}$. We have

$$P_{0,l_0}(m_0, n) = P_{0,l_0-1}(m_0 - 1, n - 1) + P_{0,l_0}(m_0 - 1, n).$$

Hence $P_{0,l_0}(m_0, n) = \binom{m_0-1}{l_0-1} + \binom{m_0-1}{l_0} = \binom{m_0}{l_0}$. We are done.

Step 3 (the last step): We show that $P_{k,l}(m, n) = \binom{m+k}{l}$ for all $m, n \in \mathbb{Z}_{\geq 0}$.

We do induction on k . Assume $P_{k,l}(m, n) = \binom{m+k}{l}$ for $k < k_0$, it is enough to show $P_{k_0,l}(m, n) = \binom{m+k_0}{l}$. We have

$$P_{k_0,l}(m, n) = P_{k_0-1,l}(m, n) + P_{k_0-1,l-1}(m, n).$$

Hence $P_{k_0,l}(m, n) = \binom{m+k_0-1}{l} + \binom{m+k_0-1}{l-1} = \binom{m+k_0}{l}$. The lemma is proved. \square

REFERENCES

- [1] A. B. Altman, A. Iarrobino, and S. L. Kleiman, *Irreducibility of the compactified Jacobian*, in: Real and complex singularities (Oslo, 1976), Sijthoff and Noordhoff International Publishers, Alphen aan den Rijn (1977), 1-12.
- [2] D. Arinkin, *Cohomology of line bundles on compactified Jacobians*, Math. Res. Lett. 18 (2011), no. 6, 1215-1226.
- [3] D. Arinkin, *Autoduality of compactified Jacobians for curves with plane singularities*, J. Algebraic Geom. 22 (2013), no. 2, 363-388.
- [4] K. Chung and H.-B. Moon, *Chow ring of the moduli space of stable sheaves supported on quartic curves*, Q. J. Math., 68 (2017), pp. 851-887.
- [5] J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. Math., 123(1), 11-42 (1986).
- [6] J. Kollár, *Higher direct images of dualizing sheaves II*, Ann. Math., 124(1), 171-202 (1986).
- [7] Y. Kononov, W. Lim, M. Moreira, and W. Pi, *Cohomology rings of the moduli of one-dimensional sheaves on the projective plane*, arXiv: 2403.06277.
- [8] D. Matsushita, *Higher direct images of dualizing sheaves of Lagrangian fibrations*, Am. J. Math., 127(2), 243-259 (2005).
- [9] Chris A. M. Peters and Joseph H. M. Steenbrink, *Mixed Hodge structures*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge/A Series of Modern Surveys in Mathematics, Volume 52. Springer-Verlag Berlin Heidelberg.
- [10] W. Schmid, *Variation of Hodge structure: the singularities of the period mapping*, Invent. Math. 22, 211-319 (1973).
- [11] Y. Yuan, *Determinant line bundles on Moduli spaces of pure sheaves on rational surfaces and Strange Duality*, Asian J. Math. 16 (2012), no. 3, 451-478.
- [12] Y. Yuan, *Sheaves on non-reduced curves in a projective surface*, Sci. China Math. Vol. 66, No. 2: 237-250.

Yao Yuan

Beijing National Center for Applied Mathematics,
Academy for Multidisciplinary Studies,
Capital Normal University, 100048, Beijing, China
E-mail: 6891@cnu.edu.cn.