Stochastic Variational Inference for Structured Additive Distributional Regression

Gianmarco Callegher* \mathbf{D}^1 \mathbf{D}^1 , Thomas Kneib \mathbf{D}^1 , Johannes Söding \mathbf{D}^2 \mathbf{D}^2 , and Paul Wiemann \mathbb{B}^3 \mathbb{B}^3

> 1 University of Göttingen ²Max Planck Institute for Multidisciplinary Sciences ³The Ohio State University

> > December 16, 2024

Abstract

In structured additive distributional regression, the conditional distribution of the response variables given the covariate information and the vector of model parameters is modelled using a P-parametric probability density function where each parameter is modelled through a linear predictor and a bijective response function that maps the domain of the predictor into the domain of the parameter. We present a method to perform inference in structured additive distributional regression using stochastic variational inference. We propose two strategies for constructing a multivariate Gaussian variational distribution to estimate the posterior distribution of the regression coefficients. The first strategy leverages covariate information and hyperparameters to learn both the location vector and the precision matrix. The second strategy tackles the complexity challenges of the first by initially assuming independence among all smooth terms and then introducing correlations through an additional set of variational parameters. Furthermore, we present two approaches for estimating the smoothing parameters. The first treats them as free parameters and provides point estimates, while the second accounts for uncertainty by applying a variational approximation to the posterior distribution. Our model was benchmarked against state-of-the-art competitors in logistic and gamma regression simulation studies. Finally, we validated our approach by comparing its posterior estimates to those obtained using Markov Chain Monte Carlo on a dataset of patents from the biotechnology/pharmaceutics and semiconductor/computer sectors.

[∗]Corresponding author: gianmarco-callegher@uni-goettingen.de

Acronyms

EMD Earth Mover's Distance INLA Integrated Nested Laplace Approximation KL Kullback-Leibler MCMC Markov chain Monte Carlo NUTS No U-Turn sampler SVI Stochastic Variational Inference VI Variational Inference WD Wasserstein Distance

1 Introduction

In recent years, interest has been increasing in flexible forms of regression models that overcome some of the limitations of classical forms of statistical modelling such as (generalized) linear models. Roughly, these developments can be grouped into (i) flexibility with respect to the chosen response distribution such that one can focus on other aspects than the conditional mean of the response variable [\[17,](#page-24-0) [18\]](#page-25-0), (ii) flexibility with respect to the predictor specification going beyond linear combinations of covariate effects in semiparametric regression [\[27,](#page-25-1) [28\]](#page-25-2), (iii) allowing for dependence and unobserved heterogeneity between observations by including random or spatial effects [\[2,](#page-24-1) [4,](#page-24-2) [5\]](#page-24-3). Many of these developments can be cast into the framework of structured additive distributional regression models [\[17\]](#page-24-0), where the response distribution is specified based on a set of latent variables (e.g. the regression predictors or the regression coefficients) equipped with Gaussian (prior) distributions.

The complexity of such flexible forms of models requires well-suited inferential procedures. Bayesian inference based on Markov chain Monte Carlo (MCMC) simulation techniques has proven very valuable in this regard (as can be seen from the abundance of Bayesian inferential procedures in the references above). This is owed on the one hand to the possibility of providing additional information to the model by including suitable forms of regularisation priors that enforce desirable model properties such as shrinkage, smoothness or sparsity, and on the other hand to the implementation of divide-and-conquer strategies in MCMC, where large and complex models can be decomposed into smaller building blocks.

The main drawback of MCMC-based inference, however, is its high demand for computation time, especially when learning models with thousands or millions of parameters. Other drawbacks include potentially slow mixing of the Markov chain, e.g. in case of highly correlated parameters, which further adds to the computation time, and the need to carefully monitor convergence in distribution over a high-dimensional parameter space. Therefore, fast and efficient approximate Bayesian inference methods are growing in popularity.

A very successful example is conducting inference using the *Integrated Nested Laplace Approxima*tion (INLA) [\[25\]](#page-25-3). INLA has become extremely widely used in many application areas of statistics due to the availability of fast, free software and the wide applicability of Latent Gaussian Models. In particular, INLA offers a wide range of modelling options for the (univariate) predictor of a regression model (relying on a structured additive predictor decomposition) and for considering dependence and unobserved heterogeneity with random and spatio-temporal effects. INLA gains much speed by adding the structured additive predictors to the set of Gaussian latent variables, which results in a sparse precision matrix of the joint prior to the latent variables. Consequently, this trick allows it to use fast sparse matrix operations and reduce its time complexity such that it can be applied to tasks with tens of thousands of latent variables.

Variational Inference (VI) is another class of approximate Bayesian inference methods where the posterior distribution is approximated by the best approximator from a class of pre-specified distributions [\[3,](#page-24-4) [29\]](#page-25-4). More precisely, let z denote the set of latent variables, y are the observed response values, and λ are some additional hyperparameters in the model specification that are not part of the latent variables. The posterior $p(\mathbf{z}|\mathbf{y},\boldsymbol{\lambda})$ of the latent variables z given the data y and the hyperparameters λ shall be approximated by a *variational distribution* $q(\mathbf{z}|\mathbf{y}, \boldsymbol{\phi})$ characterized by the parameter vector ϕ (which typically depends on the hyperparameters λ). In mean-field VI, the distribution q is assumed to be the product of distributions over subsets of latent variables such that, in some special cases, when the factorization is chosen carefully, the variational distribution that best approximates the posterior can be found analytically. However, the factorization in mean-field VI requires strong assumptions on posterior independence which limit its applicability and, when violated, can lead to severe underestimation of posterior uncertainty due to a bad approximation [\[3\]](#page-24-4).

Stochastic Variational Inference (SVI) has been developed with the goal to be applicable to a much broader range of models, overcoming the strong limitations of mean-field VI [\[29,](#page-25-4) [8\]](#page-24-5). Here, the variational distribution $q(\mathbf{z}|\mathbf{y}, \boldsymbol{\phi})$ is taken to be from a family of distributions that – given the right parameters ϕ – can approximate a broad range of posteriors well. The model and variational parameters are found by maximizing the *evidence lower bound (ELBO)* with respect to λ and ϕ . Since the model likelihood (here a.k.a. evidence) is equal to the ELBO minus the Kullback-Leibler (KL) divergence between the posterior $p(\mathbf{z}|\mathbf{y},\boldsymbol{\lambda})$ and its variational approximation $q(\mathbf{z}|\mathbf{y},\boldsymbol{\lambda},\boldsymbol{\phi})$, maximizing the ELBO is equivalent to minimizing their KL divergence and thereby approximating p by q . Stochastic gradient descent minimization relies on a Monte Carlo estimate of the gradient of the ELBO with respect to $\bm{\lambda}$ and $\bm{\phi}$, with samples \mathbf{z}^s drawn from $q(\mathbf{z}|\mathbf{y},\bm{\lambda},\bm{\phi})$. Since q is chosen so that we can independently draw samples from it (without the need for a Markov chain), mixing is instantaneous, removing a frequent complication of MCMC.

In this project, we propose to develop stochastic VI methods for structured additive distributional regression models. We use multivariate Gaussians as variational distribution, as proposed in [\[16\]](#page-24-6), for the latent variables, which has been shown to be more precise than the Laplace approximation [\[7,](#page-24-7) [21\]](#page-25-5). This is due to the fact that the latter is limited by approximating the integrand only at a single point, its mode, while in variational approaches the entire distribution is approximated. In particular, our variational distribution q is composed of two multiplicative components. The first component approximates the likelihood $p(y_i|\boldsymbol{\eta}_i=\mathbf{X}_i\boldsymbol{\beta})$ over $\boldsymbol{\beta}$ and is represented by the product of N P -dimensional Gaussian distributions $\mathcal{N}\big(\bm\eta_i|\bm\mu_i,(\mathbf{L}_i\mathbf{L}_i^{\sf T})^{-1}\big)$, where P is the number of response parameters. This choice is supported by the Bernstein–von Mises theorem. The second component is the unnormalized prior. This allows us to exploit the information given by the covariates X and the hyper-parameters λ for a faster estimation. We propose two methods for estimating the variance parameters λ . The first method involves computing the gradient of the ELBO with respect to λ , yielding a point estimate of the variance parameters. The second method accounts for uncertainty by utilizing a variational distribution q_{λ} , parameterized by a vector ϕ_{λ} . We learn ϕ_{λ} through stochastic maximization of a two-stage ELBO.

The structure of this article is as follows: In [Section 2,](#page-4-0) we briefly introduce the necessary background on structured additive distributional regression. [Section 3](#page-5-0) introduces the main ideas of VI and then describes how we specifically exploit (stochastic) VI inference for structured additive distributional regression. In [Section 4,](#page-13-0) presents a simulation study in which we compare our models with MCMC and INLA. [Section 5](#page-20-0) evaluates our model by contrasting its posterior estimate with results obtained through MCMC on a dataset containing patents spanning the fields of biotechnology/pharmaceutics and semiconductor/computer. In [Section 6,](#page-23-0) we summarize our results and briefly discuss limitations and potential directions for future research.

2 Structured additive distributional regression

2.1 Observation model

Given observational data $(y_i, \mathbf{x}_i),\; i=1,\dots,N$ of a response variable y and covariates \mathbf{x}_i we assume that the conditional distribution of the responses given the covariates is modelled using a P-parametric probability density function (pdf) as

$$
p(y_i|\mathbf{x}_i,\boldsymbol{\beta})=p(y_i|\boldsymbol{\nu}_i),
$$

where $\bm{\nu}_i=(\nu_1(\mathbf{x}_i,\bm{\beta}_1),\ldots,\nu_P(\mathbf{x}_i,\bm{\beta}_P))^{\sf T}$ is the P -dimensional vector of parameters that characterizes the distribution (for example, in terms of location, scale and skewness) and β = $(\beta_1^{\mathsf{T}}$ $_{1}^{\mathsf{T}},\ldots,\bm{\beta}_{I}^{\mathsf{T}}$ $P^\mathsf{T}_P)^\mathsf{T} \, \in \, \mathbb{R}^{Q}$ is the vector of regression coefficients. Each parameter $\nu_p(\mathbf{x}_i,\bm{\beta}_p),\ p=0$ $1,\ldots,P$, with $\bm{\beta}_p\,\in\,\mathbb{R}^{Q_p}$, is then related to a regression predictor $\eta_p(\mathbf{x}_i,\bm{\beta}_p)$ via a bijective response function h_n as

$$
\nu_{i,p} \equiv \nu_p(\mathbf{x}_i, \boldsymbol{\beta}_p) = h_p(\eta_p(\mathbf{x}_{i,p}, \boldsymbol{\beta}_p)),
$$

to ensure appropriate restrictions on the parameter space of $\nu_p(\mathbf{x}_{i,p},\bm{\beta}_p)$ (such as positivity for scale parameters).

The regression predictor $\eta_p(\mathbf{x}_{i,p},\bm{\beta}_p)$ is assumed to be of structured additive form, i.e. it is composed of J_p functions as

$$
\eta_{i,p}\equiv\eta_p(\mathbf{x}_{i,p},\boldsymbol{\beta}_p)=\sum_{j=1}^{J_p}f_{p,j}(\mathbf{x}_{i,p,j},\boldsymbol{\beta}_{p,j}),
$$

where $f_{p,j}(\mathbf{x}_i,\bm{\beta}_{p,j})$ are effects of different types such as nonlinear effects of continuous covariates, spatial effects, random effects, etc. The different effect types can be represented through basis function expansion as

$$
f_{i,p,j} \equiv f_{p,j}(\mathbf{x}_i, \boldsymbol{\beta}_{p,j}) = \sum_{l=1}^{Q_{p,j}} \beta_{p,j,l} B_{p,j,l}(\mathbf{x}_i), \qquad (1)
$$

where $B_{p,j,l}$ are known basis functions.

As a major advantage, this setup offers considerable flexibility in choosing from a variety of response distributions and effect types which are all linear in the basis coefficients $\beta_{p,j}$ such that each term $f_{p,j}$ of the structured additive predictor η_p is associated with a design matrix $\mathbf{B}_{p,j}$. This design matrix is obtained by evaluating the basis functions at the observed covariates, i.e.

$$
\mathbf{X}_{p,j} = \begin{pmatrix} B_{p,j,1}(\mathbf{x}_1) & \dots & B_{p,j,Q_{p,j}}(\mathbf{x}_1) \\ \vdots & & \vdots \\ B_{p,j,1}(\mathbf{x}_N) & \dots & B_{p,j,Q_{p,j}}(\mathbf{x}_N) \end{pmatrix} \in \mathbb{R}^{N \times Q_{p,j}}.
$$

The vector of function evaluations $\mathbf{f}_{p,j}=(f_{p,j}(\mathbf{x}_1,\bm{\beta}_{p,j}),\ldots,f_{p,j}(\mathbf{x}_N,\bm{\beta}_{p,j}))^{\sf T}$ can then be written as

$$
\mathbf{f}_{p,j} = \mathbf{X}_{p,j} \boldsymbol{\beta}_{p,j},
$$

where $\bm{\beta}_{p,j} = (\beta_{p,j,1}, \ldots, \beta_{p,j,Q_{p,j}})^{\mathsf{T}} \in \mathbb{R}^{Q_{p,j}}$.

2.2 Prior specifications

To regularize estimation, we assume informative priors for the basis coefficients that implement prior assumptions such as smoothness or shrinkage. This can be achieved by (potentially improper) normal priors for the coefficient vectors $\beta_{p,j}$, where

$$
p(\boldsymbol{\beta}_{p,j}|\lambda_{p,j}^2) \propto \exp\left\{-\frac{\lambda_{p,j}^2}{2}\boldsymbol{\beta}_{p,j}^{\mathsf{T}}\mathbf{K}_{p,j}\boldsymbol{\beta}_{p,j}\right\},\,
$$

with (potentially rank-deficient) precision matrix ${\bf K}_{p,j}\in\mathbb{R}^{Q_{p,j}\times Q_{p,j}}$ and precision parameter $\lambda_{p,j}^2$ governing the impact of the prior, similar to a smoothing parameter in penalized likelihood estimation. The model specification is completed by assuming an additional hyperprior for $\lambda_{p,j}^2$ which depends on the chosen effect type and whether it is subject to effect selection or regularization.

We define a prior for our latent variables $\boldsymbol{\beta} = (\boldsymbol{\beta}_1^T)$ $_{1}^{T},\ldots,\bm{\beta}_{I}^{T}$ $\left(T\overline{P}\right)^T$ by a joint normal distribution,

$$
p(\boldsymbol{\beta}|\boldsymbol{\lambda}) = \prod_{p=1}^P \mathcal{N}(\boldsymbol{\beta}_p | \boldsymbol{0}, \mathbf{K}_{p,\boldsymbol{\lambda}_p}) = \mathcal{N}(\boldsymbol{\beta} | \boldsymbol{0}, \mathbf{K}_{\boldsymbol{\lambda}}),
$$

with precision matrix

$$
\mathbf{K}_{\pmb{\lambda}} \equiv \mathsf{block\text{-}diag}\big(\mathbf{K}_{1,\pmb{\lambda}_1},\ldots,\mathbf{K}_{P,\pmb{\lambda}_P}\big),
$$

where

$$
\mathbf{K}_{p, \boldsymbol{\lambda}_p} \equiv \mathsf{block\text{-}diag}\big(\lambda_{p,1}^2 \mathbf{K}_{p,1}, \ldots, \lambda_{p,J_p}^2 \mathbf{K}_{p,J_p}\big).
$$

2.3 Posterior

Assuming conditional independence of the observations given the model parameters, the logposterior distribution for all model parameters is given by

$$
\ln p(\boldsymbol{\beta}, \boldsymbol{\lambda} | \mathbf{y}, \mathbf{X}) = \ln p(\mathbf{y} | \boldsymbol{\beta}, \mathbf{X}) + \ln p(\boldsymbol{\beta} | \boldsymbol{\lambda}) + \ln p(\boldsymbol{\lambda}) + \text{const}
$$

=
$$
\sum_{i=1}^{N} \log p(\mathbf{y}_i | \boldsymbol{\eta}_i = \mathbf{X}_i \boldsymbol{\beta}) + \ln p(\boldsymbol{\beta} | \boldsymbol{\lambda}) + \ln p(\boldsymbol{\lambda}) + \text{const.}
$$
 (2)

Since distributional regression allows for very flexible types of response distributions and usually deals with a large dimension of regression coefficients β , the posterior is not analytically tractable. In the introduction we discussed the limitations of MCMC methods in high-dimensional parameter spaces. We therefore derive an approximate approach based on stochastic variational inference in the following section.

3 Stochastic variational inference for structured additive distributional regression

Variational Inference (VI) is a powerful technique in the context of probabilistic modelling and Bayesian inference. It offers a flexible and scalable approach for approximating complex probability distributions, especially in scenarios where exact inference is computationally intractable. In this section, we introduce the main ideas of VI and then describe how to specifically develop (stochastic) VI inference for structured additive distributional regression.

3.1 Background on variational inference

To introduce ideas for VI in a generic setup, let $y = y_{1:N}$ be a set of observed variables (e.g. the responses of a regression model) and assumed to be drawn from a distribution

$$
p(\mathbf{y}|\boldsymbol{\theta}) = \int p(\mathbf{y}|\mathbf{z}, \boldsymbol{\theta}) p(\mathbf{z}|\boldsymbol{\theta}) \, d\mathbf{z},
$$

where $\bm{\theta}$ are the model's hyperparameters (e.g. $\bm{\lambda}$ in [Equation 2\)](#page-5-1) and $\mathbf{z}=z_{1:M}$ is a set of latent parameters (e.g. regression coefficients β). Besides finding the best hyperparameters θ , a central goal of variational inference is to approximate the conditional density $p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta})$ of the latent variables given y, by framing the problem as an optimization task. The idea is to find the so-called variational distribution $q(\mathbf{z})$ from a predefined family of distributions Q that best approximates the true posterior $p(\mathbf{z}|\mathbf{y}, \theta)$ among all members of the variational family. We use a family of densities $q(\mathbf{z}|\boldsymbol{\phi})$ over the latent variables z, parameterized by the free variational parameter $\boldsymbol{\phi}$. The optimization finds the member of this family $q(\mathbf{z}|\boldsymbol{\phi}^{\star})$ that is closest in Kullback-Leibler (KL) divergence to the conditional of interest, i.e.

$$
\phi^* = \underset{\phi}{\arg\min} \mathcal{D}_{\mathsf{KL}}(q(\mathbf{z}|\phi) || p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta}))
$$

=
$$
\underset{\phi}{\arg\min} \mathbb{E}_{q(\mathbf{z}|\phi)} [\ln q(\mathbf{z}|\boldsymbol{\phi})] - \mathbb{E}_{q(\mathbf{z}|\phi)} [\ln p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta})] + \ln p(\mathbf{y}|\boldsymbol{\theta}).
$$

where $\mathcal{D}_{\mathsf{KL}}$ denotes the Kullback-Leibler divergence.

Variational inference offers several advantages over Markov chain Monte Carlo (MCMC) methods. It is generally faster and more scalable, making it suitable for large datasets and complex models. Moreover, VI offers a global sampling scheme, since we already deal with a proper distribution which serves as an approximation of the posterior distribution, while in MCMC, the posterior is retrieved by aggregating individual posterior samples. Despite its advantages, VI comes with its own set of challenges. One of the primary challenges is the choice of the variational family Q , which directly impacts the quality of the approximation. Additionally, finding the optimal variational parameters often involves solving complex optimization problems, which may require careful tuning.

3.1.1 Evidence lower bound

Using $\int q(\mathbf{z}|\boldsymbol{\phi})d\mathbf{z} = 1$, we can rewrite the log evidence in the following way,

$$
\ln p(\mathbf{y}|\boldsymbol{\theta}) = \int q(\mathbf{z}|\boldsymbol{\phi}) \ln p(\mathbf{y}|\boldsymbol{\theta}) d\mathbf{z}
$$

\n
$$
= \int q(\mathbf{z}|\boldsymbol{\phi}) \ln \frac{p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta})}{p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta})} d\mathbf{z}
$$

\n
$$
= \int q(\mathbf{z}|\boldsymbol{\phi}) \ln \left(\frac{p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta})}{q(\mathbf{z}|\boldsymbol{\phi})} \frac{q(\mathbf{z}|\boldsymbol{\phi})}{p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta})}\right) d\mathbf{z}
$$

\n
$$
= \int q(\mathbf{z}|\boldsymbol{\phi}) \ln \frac{p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta})}{q(\mathbf{z}|\boldsymbol{\phi})} d\mathbf{z} + \int q(\mathbf{z}|\boldsymbol{\phi}) \ln \frac{q(\mathbf{z}|\boldsymbol{\phi})}{p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta})} d\mathbf{z}
$$

\n
$$
= \text{ELBO}(\boldsymbol{\phi}, \boldsymbol{\theta}) + \mathcal{D}_{\text{KL}}(q(\mathbf{z}|\boldsymbol{\phi}) || p(\mathbf{z}|\mathbf{y}, \boldsymbol{\theta}))
$$

\n
$$
\geq \text{ELBO}(\boldsymbol{\phi}, \boldsymbol{\theta}),
$$

where we have used the fact that the KL divergence $\mathcal{D}_{KL}(q||p)$ is nonnegative for any probability distributions p and q , and we defined the evidence lower bound (ELBO) as

$$
\text{ELBO}(\boldsymbol{\phi}, \boldsymbol{\theta}) = \int q(\mathbf{z}|\boldsymbol{\phi}) \ln \frac{p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta})}{q(\mathbf{z}|\boldsymbol{\phi})} d\mathbf{z} = \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[\ln p(\mathbf{z}, \mathbf{y}|\boldsymbol{\theta}) \right] - \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[\ln q(\mathbf{z}|\boldsymbol{\phi}) \right],
$$

which is a lower bound of the log evidence for any $q(\mathbf{z})$. Because ELBO $(\boldsymbol{\phi}, \boldsymbol{\theta}) = \ln p(\mathbf{y}|\boldsymbol{\theta})$ – $\mathcal{D}_{\text{KL}}(q(\mathbf{z}|\boldsymbol{\phi})||p(\mathbf{z}|\mathbf{y}))$ is equal to a constant (w.r.t $\boldsymbol{\phi}$) minus the KL divergence, maximization of the ELBO with respect to ϕ is equivalent to minimizing the KL divergence with respect to ϕ . Also, because the ELBO is a lower bound of the log evidence, maximization of the ELBO with respect to θ approximates the maximization of the log evidence when the KL divergence is small.

To get some intuition, we can write the ELBO as

$$
\text{ELBO}(\boldsymbol{\phi}, \boldsymbol{\theta}) = \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[p(\mathbf{y}|\mathbf{z}, \boldsymbol{\theta}) \right] + \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[p(\mathbf{z}|\boldsymbol{\theta}) \right] - \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[q(\mathbf{z}|\boldsymbol{\phi}) \right] = \mathrm{E}_{q(\mathbf{z}|\boldsymbol{\phi})} \left[p(\mathbf{y}|\mathbf{z}, \boldsymbol{\theta}) \right] - \mathcal{D}_{\mathsf{KL}} \left(q(\mathbf{z}|\boldsymbol{\phi}) || p(\mathbf{z}|\boldsymbol{\theta}) \right).
$$

The first term is an expected likelihood which encourages densities $q(z|\phi)$ that place their mass on latent variables that explain the observed data. The second term, the negative KL divergence between the variational density and the prior, encourages densities to be similar to the prior. Thus, the variational objective mirrors the usual balance between likelihood and prior in Bayesian statistics.

3.1.2 Stochastic variational inference

Stochastic Variational Inference $(\mathsf{SVl})^1$ $(\mathsf{SVl})^1$ is an extension of VI that introduces efficient stochastic estimates of the gradient of the ELBO with respect to model and variational parameters and uses stochastic gradient descent to maximize the ELBO.

 1 The word "Stochastic" frequently denotes the use of stochastic optimization techniques to manage large datasets, which typically involves the use of stochastic minibatches. It is imperative to explicitly clarify that we are discussing a stochastic approximation of the ELBO with respect to samples drawn from the variational distribution.

To obtain these efficient estimators of the ELBO gradients, we write the gradients as expectation values $E_q(z|\phi)[f(z)]$, since in that case we can approximate the expectation values by sample means $S^{-1}\sum_{s=1}^S f(\mathbf{z}^s)$ using samples $\mathbf{z}^s\sim q(\mathbf{z}|\boldsymbol{\phi}).$ To compute an estimate of the gradient with respect to the model parameters θ , we can simply take the sample mean of the gradient of the terms in the sum:

$$
\nabla_{\theta} \text{ELBO}(\phi, \theta) = \nabla_{\theta} \int q(\mathbf{z}|\phi) \ln \frac{p(\mathbf{y}, \mathbf{z}|\theta)}{q(\mathbf{z}|\phi)} d\mathbf{z}
$$

\n
$$
= \int q(\mathbf{z}|\phi) \nabla_{\theta} \ln \frac{p(\mathbf{y}, \mathbf{z}|\theta)}{q(\mathbf{z}|\phi)} d\mathbf{z}
$$

\n
$$
= \mathbf{E}_{q(\mathbf{z}|\phi)} \left[\nabla_{\theta} \ln \frac{p(\mathbf{y}, \mathbf{z}|\theta)}{q(\mathbf{z}|\phi)} \right]
$$

\n
$$
\approx \frac{1}{S} \sum_{s=1}^{S} \nabla_{\theta} \ln \frac{p(\mathbf{y}, \mathbf{z}^{s}|\theta)}{q(\mathbf{z}^{s}|\phi)}, \text{ with } \mathbf{z}^{s} \sim q(\mathbf{z}|\phi).
$$

The gradient in the sum is usually computed using automatic differentiation.

For the gradient with respect ϕ , however, we cannot move the term $q(\mathbf{z}|\phi)$ out of the gradient as done above, since $q(\mathbf{z}|\boldsymbol{\phi})$ depends on $\boldsymbol{\phi}$. To take this dependence into account, we use the so-called reparameterization trick [\[12\]](#page-24-8). We find a transformation ${\rm g}_{\rm z}(\phi,\epsilon):\mathbb{R}^Q\to\mathbb{R}^Q$ such that sampling $\bm{\epsilon}^s\sim\mathcal{N}(\bm{0},\bf{I})$ ensures that the transformed variables $\mathbf{z}^s=\mathbf{g}_{\mathbf{z}}(\bm{\phi},\bm{\epsilon}^s)$ are sampled from $q(\mathbf{z}|\boldsymbol{\phi})$. For Gaussian $q(\mathbf{z}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}, \boldsymbol{\Sigma})$, for instance, this can be achieved simply by the linear transformation $g_z(\mu, C, \epsilon) = \mu + C\epsilon$ for any matrix C satisfying $\Sigma = CC^{\top}$.

With this reparameterization of stochastic samples $\mathbf{z}^s = \mathbf{g}_{\mathbf{z}}(\boldsymbol{\phi}, \boldsymbol{\epsilon}^s)$, we can write

$$
\nabla_{\phi} \text{ELBO}(\phi, \theta) = \nabla_{\phi} E_{q(\mathbf{z}|\phi)} \left[\ln \frac{p(\mathbf{y}, \mathbf{z}|\theta)}{q(\mathbf{z}|\phi)} \right]
$$

\n
$$
= \nabla_{\phi} E_{\mathcal{N}(\epsilon|\mathbf{0}, \mathbf{I})} \left[\ln \frac{p(\mathbf{y}, \mathbf{g}_{\mathbf{z}}(\phi, \epsilon)|\theta)}{q(\mathbf{g}_{\mathbf{z}}(\phi, \epsilon)|\phi)} \right]
$$

\n
$$
= E_{\mathcal{N}(\epsilon|\mathbf{0}, \mathbf{I})} \left[\nabla_{\phi} \ln \frac{p(\mathbf{y}, \mathbf{g}_{\mathbf{z}}(\phi, \epsilon)|\theta)}{q(\mathbf{g}_{\mathbf{z}}(\phi, \epsilon)|\phi)} \right]
$$

\n
$$
\approx \frac{1}{S} \sum_{s=1}^{S} \nabla_{\phi} \ln \frac{p(\mathbf{y}, \mathbf{g}_{\mathbf{z}}(\phi, \epsilon^{s})|\theta)}{q(\mathbf{g}_{\mathbf{z}}(\phi, \epsilon^{s})|\phi)}, \text{ with } \epsilon^{s} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}).
$$

Hence, the gradient $\nabla_\phi\text{ELBO}(\phi,\bm\theta)$ can be estimated as a simple sample mean of $\nabla_\phi\ln\frac{p(\mathbf{y},\mathbf{g}_{\mathbf{z}}(\phi,\epsilon^s))}{q(\mathbf{y},\epsilon(\phi,\epsilon^s))d\delta}$ $\frac{p(\mathbf{y}, \mathbf{g}_{\mathbf{z}}(\boldsymbol{\phi}, \boldsymbol{\epsilon}^{\circ}))}{q(\mathbf{g}_{\mathbf{z}}(\boldsymbol{\phi}, \boldsymbol{\epsilon}^{\circ})|\boldsymbol{\phi})},$ and these gradient terms are usually computed using automatic differentiation methods.

3.2 Local variational inference for the regression coefficients

In this section, we describe our methodology for constructing a multivariate Gaussian variational distribution to estimate the posterior distribution $p(\beta|\lambda, y, X)$ in the context of structured additive distributional regression. Specifically, we propose two distinct approaches for building the variational distribution q . The first approach utilizes the information provided by the covariates X and the hyper-parameters λ to learn both the location vector and the precision matrix. The second approach addresses the asymptotic complexity issues of the first by initially assuming independence among all smooth terms $f_{i,p,j}$ and then reintroducing correlation through an additional set of variational parameters. We then explain how to estimate the variational parameters. Lastly, we present two methods for estimating the smoothing parameters λ . The first method treats λ as free parameters and provides point estimates, whereas the second method accounts for uncertainty by employing a variational approximation to the posterior distribution.

The multivariate normal distribution is favoured for the variational distribution of β , supported by the Bernstein–von Mises theorem and its conjugacy properties. The Bernstein-von Mises theorem states that, under regularity conditions such as smoothness of the likelihood function and the prior not being too informative, the posterior distribution $p(\beta|\lambda, y, X)$ is approximately normal for large sample size N . Conversely, the variational distribution for the smoothing parameters can differ based on the chosen parametrization and the feasibility of obtaining an analytical posterior solution.

Unlike traditional variational inference, which seeks a global approximation for all latent variables simultaneously, local variational inference focuses on approximating the posterior for a subset of latent variables associated with individual data points. This approach is particularly advantageous in large-scale data settings, as it allows for parallel computation and reduces the complexity of the inference process. By optimizing a local variational objective for each data point, this method facilitates more flexible and fine-grained approximations, enhancing the overall accuracy and computational efficiency of the inference in complex models.

In this section, we outline the construction of a variational distribution for estimating the regression coefficients β , designed for a fast approximation of the joint distribution. Our variational distribution q is composed of two multiplicative components. The first component approximates the likelihood $p(y_i|\bm{\eta}_i=\mathbf{X}_i\bm{\beta})$ over $\bm{\beta}$ and is represented by the product of N P -dimensional Gaussian distributions $\mathcal{N}\big(\bm{\eta}_i|\bm{\mu}_i, (\mathbf{L}_i\mathbf{L}_i^{\sf T})^{-1}\big)$. The Bernstein–von Mises theorem supports this choice. The second component is the unnormalized prior. Consequently, the unnormalized variational distribution is given by:

$$
\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}) = -\frac{1}{2} \sum_{i=1}^{N} (\mathbf{X}_i \boldsymbol{\beta} - \boldsymbol{\mu}_i)^{\mathsf{T}} \mathbf{L}_i \mathbf{L}_i^{\mathsf{T}} (\mathbf{X}_i \boldsymbol{\beta} - \boldsymbol{\mu}_i) - \frac{1}{2} \boldsymbol{\beta}^{\mathsf{T}} \mathbf{K}_{\boldsymbol{\lambda}} \boldsymbol{\beta} + \text{const},
$$
 (3)

where

$$
\mathbf{X}_i := \begin{pmatrix} \mathbf{x}_{i,1}^{\mathsf{T}} & \mathbf{0}^{\mathsf{T}} & \cdots & \mathbf{0}^{\mathsf{T}} \\ \mathbf{0}^{\mathsf{T}} & \mathbf{x}_{i,2}^{\mathsf{T}} & \cdots & \mathbf{0}^{\mathsf{T}} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0}^{\mathsf{T}} & \mathbf{0}^{\mathsf{T}} & \cdots & \mathbf{x}_{i,P}^{\mathsf{T}} \end{pmatrix} \in \mathbb{R}^{P \times Q},
$$

are the covariates for data sample i. The set of variational parameters ϕ is represented by all latent variables' local variational distribution parameters $(\bm{\mu}_1^{\sf T},\bm{l}_1^T,\ldots,\bm{\mu}_N^{\sf T},\bm{l}_N^{\sf T})^{\sf T}$, where

$$
\mathbf{l}_i = \psi(\mathbf{L}_i), \quad \text{for } i = 1, \dots, N
$$

is the log-Cholesky parametrization [\[22\]](#page-25-6) of the matrix \mathbf{L}_i , *i.e.*:

$$
\psi^{-1}(\mathbf{l}_i) = \begin{pmatrix} \exp{(l_{i,1})} & 0 & \cdots & \mathbf{0} \\ l_{i,2} & \exp{(l_{i,3})} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ l_{i,P(P+1)/2-P+1} & l_{i,P(P+1)/2-P+2} & \cdots & \exp{(l_{i,P(P+1)/2})} \end{pmatrix} \in \mathbb{R}^{P \times P}.
$$

The normalized variational distribution over the latent variables can be written as

$$
\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}) = -\frac{1}{2} \sum_{i=1}^{N} (\boldsymbol{\beta}^{\mathsf{T}} \mathbf{X}_{i}^{\mathsf{T}} \mathbf{L}_{i} \mathbf{L}_{i}^{\mathsf{T}} \mathbf{X}_{i} \boldsymbol{\beta} - 2 \boldsymbol{\beta}^{\mathsf{T}} \mathbf{X}_{i}^{\mathsf{T}} \mathbf{L}_{i} \mathbf{L}_{i}^{\mathsf{T}} \boldsymbol{\mu}_{i}) - \frac{1}{2} \boldsymbol{\beta}^{\mathsf{T}} \mathbf{K}_{\boldsymbol{\lambda}} \boldsymbol{\beta} + \text{const} \n= \ln \mathcal{N}(\boldsymbol{\beta} | \tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda}, \boldsymbol{\phi}}, \boldsymbol{\Lambda}_{\boldsymbol{\lambda}, \boldsymbol{\phi}}^{-1}),
$$
\n(4)

with

$$
\Lambda_{\lambda,\phi} = \sum_{i=1}^{N} \mathbf{X}_{i}^{\mathsf{T}} \mathbf{L}_{i} \mathbf{L}_{i}^{\mathsf{T}} \mathbf{X}_{i} + \mathbf{K}_{\lambda},
$$
\n
$$
\tilde{\boldsymbol{\beta}}_{\lambda,\phi} = \Lambda_{\lambda,\phi}^{-1} \sum_{i=1}^{N} \mathbf{X}_{i}^{\mathsf{T}} \mathbf{L}_{i} \mathbf{L}_{i}^{\mathsf{T}} \boldsymbol{\mu}_{i}.
$$
\n(5)

In order to ensure that the matrix $\Lambda_{\lambda,\phi}$ has full-rank, we need to ensure the same property for each matrix $\mathbf{X}_i.$ To do so, we leverage the QR-reparametrization presented in [Section A.1.](#page-26-0)

3.3 A block-diagonal variational distribution

The model proposed in [Equation 5](#page-10-0) necessitates computing the Cholesky decomposition of the matrix $\Lambda_{\lambda,\phi}$ to solve the linear system needed to determine $\tilde{\boldsymbol{\beta}}_{\lambda,\phi}$ and to sample from $\mathcal{N}(\boldsymbol{\beta}|\tilde{\boldsymbol{\beta}}_{\lambda,\phi},\Lambda_{\lambda,\phi}^{-1})$ $\frac{-1}{\lambda, \phi}$). This process has a complexity of $O(Q^3)$, rendering the model impractical for large numbers of regression coefficients. In this section, we introduce an alternative variational distribution. This distribution assumes independence between each smooth term $f_{i,p,j}$ [\(Equation 1\)](#page-4-1) initially, then reintroduces dependence using a new set of variational parameters. We employ a Gaussian distribution $\mathcal{N}\big(f_{i,p,j}|\mu_{i,p,j},\sigma^2_{i,p,j}\big)$ to approximate each smooth term $f_{i,p,j}.$ Consequently, the variational distribution of $\beta_{p,j}$ takes the following form:

$$
\ln q(\boldsymbol{\beta}_{p,j}|\lambda_{p,j}, \mathbf{X}_{p,j}, \boldsymbol{\phi}_{p,j}) = \ln \mathcal{N}(\boldsymbol{\beta}_{p,j}|\tilde{\boldsymbol{\beta}}_{p,j}(\lambda_{p,j}, \boldsymbol{\phi}_{p,j}), \boldsymbol{\Lambda}_{p,j}^{-1}(\lambda_{p,j}, \boldsymbol{\phi}_{p,j}))
$$

with

$$
\Lambda_{p,j}(\lambda_{p,j}, \phi_{p,j}) = \sum_{i=1}^{N} \frac{\mathbf{x}_{i,p,j}^{T} \mathbf{x}_{i,p,j}}{\sigma_{i,p,j}^{2}} + \mathbf{K}_{p,j}(\lambda_{p,j}),
$$
\n
$$
\tilde{\boldsymbol{\beta}}_{p,j}(\lambda_{p,j}, \phi_{p,j}) = \Lambda_{p,j}^{-1}(\lambda_{p,j}, \phi_{p,j}) \sum_{i=1}^{N} \frac{\mu_{i,p,j} \mathbf{x}_{i,p,j}^{T}}{\sigma_{i,p,j}^{2}},
$$
\n(6)

where $\mathbf{x}_{i,p,j} \, \in \, \mathbb{R}^{Q_{p,j}}$ and μ_i and $\log \sigma_i$ are the latent variables' local variational distribution parameters.

The full variational distribution $q(\beta|\lambda, \mathbf{X}, \phi)$ can be written as:

$$
\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}) = \ln \mathcal{N}(\boldsymbol{\beta}|\tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda}, \boldsymbol{\phi}}, \boldsymbol{\Lambda}_{\boldsymbol{\lambda}, \boldsymbol{\phi}}^{-1}),
$$

where $\tilde{\bm{\beta}}_{\bm{\lambda},\bm{\phi}}$ is the vector obtained by stacking all $\tilde{\bm{\beta}}_{p,j}(\lambda_{p,j},\bm{\phi})$ and $\bm{\Lambda}_{\bm{\lambda},\bm{c}}^{-1}$ $_{\boldsymbol{\lambda},\boldsymbol{\phi}}^{-1}$ is a block-diagonal matrix obtained by stacking all $\Lambda_{p,j}^{-1}(\lambda_{p,j},\boldsymbol{\phi}_{p,j})$ along the diagonal.

To account for correlation, we can learn an additional vector $\tilde{{\boldsymbol \beta}}^{(c)}_{\boldsymbol \phi}$ and a full lower triangular matrix $\mathbf{L}_{\boldsymbol\phi}^{(c)}$ $\stackrel{(c)}{\phi}$. The resulting full variational distribution is:

$$
\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}) = \mathcal{N}\left(\tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda}, \boldsymbol{\phi}} + \tilde{\boldsymbol{\beta}}_{\boldsymbol{\phi}}^{(c)}, \left(\left(\mathbf{L}_{\boldsymbol{\lambda}, \boldsymbol{\phi}} + \mathbf{L}_{\boldsymbol{\phi}}^{(c)}\right) \left(\mathbf{L}_{\boldsymbol{\lambda}, \boldsymbol{\phi}} + \mathbf{L}_{\boldsymbol{\phi}}^{(c)}\right)^{\mathsf{T}}\right)^{-1}\right)
$$
(7)

where ${\rm L}_{\phi,\lambda}$ is the Cholesky factor of $\Lambda_{\lambda,\phi}$, *i.e.* $\Lambda_{\lambda,\phi}={\rm L}_{\phi,\lambda}{\rm L}_{\phi,\lambda}^{\rm T}$. The complexity for obtaining the Cholesky decomposition of the precision matrix is then reduced to $O\big(Q^2+\sum_p\sum_jQ^3_{p,j}\big).$

3.4 Estimation of the variational parameters

We optimize the variational parameters ϕ by maximizing the evidence lower bound:

$$
ELBO(\boldsymbol{\phi}) = E_{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})} \left[\ln \frac{p(\mathbf{y}, \boldsymbol{\beta}|\boldsymbol{\lambda})}{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})} \right],
$$
(8)

where $q(\bm{\beta}|\bm{\phi},\bm{\lambda},\mathbf{X})$ can be both [Equation 4](#page-10-1) or [Equation 7.](#page-11-0) To do that, we draw S samples $\bm{\beta}^s\sim$ $\mathcal{N} \big(\tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda},\boldsymbol{\phi}}, \boldsymbol{\Lambda}_{\boldsymbol{\lambda},\boldsymbol{\epsilon}}^{-1}$ $^{-1}_{\bm{\lambda},\bm{\phi}})$ from $q(\bm{\beta}|\bm{\phi},\bm{\lambda},\mathbf{X})$. This can be done by computing the Cholesky decomposition $\Lambda_{\pmb{\lambda},\pmb{\phi}} = \mathbf{L}_{\pmb{\phi},\pmb{\lambda}} \mathbf{L}_{\pmb{\phi},\pmb{\lambda}}^\mathsf{T}$, sampling $\pmb{\epsilon}^s \sim \mathcal{N}(\pmb{0},\mathbf{I}_Q)$ and setting:

$$
\boldsymbol{\beta}^s = \mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\epsilon}^s, \boldsymbol{\phi}) := \tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda}, \boldsymbol{\phi}} + \mathbf{L}_{\boldsymbol{\phi}, \boldsymbol{\lambda}}^{-T} \boldsymbol{\epsilon}^s.
$$
\n(9)

We can now proceed by maximizing the ELBO with respect to ϕ using the reparameterization trick, which consists of using the function $g_\beta(\lambda, X, \epsilon, \phi)$, which depends on ϕ , to replace the expectation over β by an expectation over ϵ :

$$
\mathrm{E}_{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})} \left[\ln \frac{p(\mathbf{y}, \boldsymbol{\beta}|\boldsymbol{\lambda})}{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})} \right] = \mathrm{E}_{\boldsymbol{\epsilon}} \left[\ln \frac{p(\mathbf{y}, \mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\epsilon}, \boldsymbol{\phi})|\boldsymbol{\lambda})}{q(\mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\epsilon}, \boldsymbol{\phi})|\boldsymbol{\lambda}, \boldsymbol{\phi})} \right]
$$

To optimize our parameters ϕ , we need to compute the gradient w.r.t. ϕ of this expression:

$$
\nabla_{\phi} \mathcal{E}_{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \phi)} \left[\ln \frac{p(\mathbf{y}, \boldsymbol{\beta}|\boldsymbol{\lambda})}{q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})} \right] = \mathcal{E}_{\epsilon} \left[\nabla_{\phi} \ln \frac{p(\mathbf{y}, \mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\epsilon}, \boldsymbol{\phi}) | \boldsymbol{\lambda})}{q(\mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\epsilon}, \boldsymbol{\phi}) | \boldsymbol{\lambda}, \boldsymbol{\phi})} \right] \approx \frac{1}{S} \sum_{s=1}^{S} \nabla_{\phi} \left[\ln p(\mathbf{y}, \boldsymbol{\beta}^{s} | \boldsymbol{\lambda}) - \ln q(\boldsymbol{\beta}^{s} | \boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}) \right]. \tag{10}
$$

A sketch of the training algorithm is described in [Algorithm 1.](#page-11-1)

Algorithm 1 Train SVI-DR by stochastic variational inference.

Input: Training data: responses (y_1, \ldots, y_N) , covariates $(\mathbf{X}_1, \ldots, \mathbf{X}_P)$ and step-size α

1: Initialize ϕ 2: for $t = 1, ..., T$ do 3: $\phi^{t+1} \leftarrow \phi^t + \alpha \nabla_\phi \mathsf{ELBO}(\phi^t)$ using [Equation 10](#page-11-2) 4: end for 5: $\mathsf{return}~ \tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda},\phi}$ and $\Lambda^{-1}_{\boldsymbol{\lambda},\phi}$ $\frac{-1}{\lambda, \phi}$.

3.5 Estimation of the variance parameters

We propose two methods for estimating the variance parameters λ . The first method treats λ as hyperparameters and provides point estimates, whereas the second method accounts for uncertainty by employing a variational approximation to the posterior distribution.

3.5.1 A point estimate of the variance parameters

We optimize the hyper-parameters λ by maximizing the evidence lower bound, [Equation 8.](#page-11-3) To do so, we extend [Equation 10](#page-11-2) by computing the gradient w.r.t. to λ as well:

$$
\nabla_{\phi,\lambda} E_{q(\beta|\lambda,\mathbf{X},\phi)} \left[\ln \frac{p(\mathbf{y},\beta|\lambda)}{q(\beta|\lambda,\mathbf{X},\phi)} \right] = E_{\epsilon} \left[\nabla_{\phi,\lambda} \ln \frac{p(\mathbf{y},\mathbf{g}_{\beta}(\lambda,\mathbf{X},\epsilon,\phi)|\lambda)}{q(\mathbf{g}_{\beta}(\lambda,\mathbf{X},\epsilon,\phi)|\lambda,\mathbf{X},\phi)} \right]
$$

$$
\approx \frac{1}{S} \sum_{s=1}^{S} \nabla_{\phi,\lambda} \left[\ln p(\mathbf{y},\beta^{s}|\lambda) - \ln q(\beta^{s}|\lambda,\mathbf{X},\phi) \right]. \tag{11}
$$

[Algorithm 2](#page-12-0) shows an updated version of [Algorithm 1](#page-11-1) accordingly.

Algorithm 2 Train SVI-DR by stochastic variational inference.

Input: Training data: responses (y_1, \ldots, y_N) , covariates $(\mathbf{X}_1, \ldots, \mathbf{X}_P)$ and step-size α

1: Initialize ϕ 2: for $t = 1, ..., T$ do 3: $\phi^{t+1} \leftarrow \phi^t + \alpha \nabla_\phi \mathsf{ELBO}(\phi^t, \boldsymbol{\lambda}^t)$ using [Equation 11](#page-12-1) 4: $\bm{\lambda}^{t+1} \leftarrow \bm{\lambda}^t + \alpha \nabla_{\bm{\lambda}} \mathsf{ELBO}(\bm{\phi}^t,\bm{\lambda}^t)$ using [Equation 11](#page-12-1) 5: end for 6: $\mathsf{return}~ \tilde{\boldsymbol{\beta}}_{\boldsymbol{\lambda},\phi}$ and $\Lambda^{-1}_{\boldsymbol{\lambda},\phi}$ $\frac{-1}{\lambda, \phi}$.

3.5.2 A variational distribution for the variance parameters

Instead of treating λ as hyperparameters to optimize, we can include it into the set of latent variables and sample from it. We use [Equation 3,](#page-9-0) including the variance-parameters, to approximate [Equation 2.](#page-5-1) This results in the following variational distribution:

$$
\ln q(\boldsymbol{\beta}, \boldsymbol{\lambda} | \mathbf{X}, \boldsymbol{\phi}) = \ln q(\boldsymbol{\beta} | \boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi}_{\beta}) + \ln q(\boldsymbol{\lambda} | \boldsymbol{\phi}_{\lambda})
$$

The variational distribution $q(\bm{\lambda}|\bm{\phi}_{\lambda})$ assumes independence among all smoothing parameters $\lambda_{p,j}$, i.e.

$$
q(\boldsymbol{\lambda}|\boldsymbol{\phi}_{\lambda}) = \prod_{p=1}^{P} \prod_{j=1}^{J_p} q(\lambda_{p,j}|\boldsymbol{\phi}_{\lambda_{p,j}}).
$$

For example, we can choose a Log-normal distribution for each $q(\lambda_{p,j}|\boldsymbol{\phi}_{\lambda_{p,j}})$. In that case, the vector of variational parameters ϕ_λ contains all the log-location and log-scale parameters that characterize each hyper-variational normal distribution $q(\lambda_{p,j} | \phi_{p,j})$.

We optimize the variational parameters $\bm{\phi}=(\bm{\phi}_{\beta},\bm{\phi}_{\lambda})$ by maximizing the evidence lower bound

$$
ELBO(\boldsymbol{\phi}) = E_{q(\boldsymbol{\beta},\boldsymbol{\lambda}|\mathbf{X},\boldsymbol{\phi})} [\ln p(\mathbf{y},\boldsymbol{\beta},\boldsymbol{\lambda})] - E_{q(\boldsymbol{\beta},\boldsymbol{\lambda}|\mathbf{X},\boldsymbol{\phi})} [\ln q(\boldsymbol{\beta},\boldsymbol{\lambda}|\mathbf{X},\boldsymbol{\phi})]. \tag{12}
$$

To do that, we once again use the reparametrization trick. First, we draw S_λ samples $\bm{\lambda}^{s_\lambda}$ from $q(\bm{\lambda}|\bm{\phi}_{\lambda})$. Each $\bm{\lambda}^{s_\lambda}$ maps to a different variational distribution $\mathcal{N}(\tilde{\bm{\beta}}_{\bm{\lambda}^{s_\lambda},\bm{\phi}},\bm{\Lambda}_{\bm{\lambda}^{s_\lambda}}^{-1}$ $(\overline{\lambda^{s_{\lambda}}}, \phi)$ from which we draw S_β samples of β using [Equation 9:](#page-11-4)

$$
\boldsymbol{\beta}^s = \mathbf{g}_{\boldsymbol{\beta}}(\boldsymbol{\lambda}^{s_{\lambda}}, \mathbf{X}, \boldsymbol{\epsilon}, \boldsymbol{\phi}_{\beta}).
$$

To optimize our parameters ϕ , we need to compute the gradient w.r.t. ϕ of [Equation 12:](#page-13-1)

$$
\nabla_{\boldsymbol{\phi}} \mathcal{E}_{q(\boldsymbol{\beta},\boldsymbol{\lambda}|\mathbf{X},\boldsymbol{\phi})} \left[\ln \frac{p(\mathbf{y},\boldsymbol{\beta},\boldsymbol{\lambda})}{q(\boldsymbol{\beta},\boldsymbol{\lambda}|\mathbf{X},\boldsymbol{\phi})} \right] = \approx \frac{1}{S_{\lambda} \cdot S_{\beta}} \sum_{S_{\lambda}=1}^{S_{\lambda}} \sum_{s_{\beta}=1}^{S_{\beta}} \nabla_{\boldsymbol{\phi}} \left[\ln p(\mathbf{y},\boldsymbol{\beta}^{s_{\beta}},\boldsymbol{\lambda}^{s_{\lambda}}) - \ln q(\boldsymbol{\beta}^{s_{\beta}},\boldsymbol{\lambda}^{s_{\lambda}}|\mathbf{X},\boldsymbol{\phi}) \right].
$$
\n(13)

The estimation algorithm is similar to [Algorithm 1,](#page-11-1) where, instead of updating the variational parameters using [Equation 10](#page-11-2) (line 3), one has to use [Equation 13.](#page-13-2)

3.6 Asymptotic complexity

[Table 3](#page-27-0) in the Appendix provides a comparison of the asymptotic complexity for a single iteration of various models focused solely on estimating the regression coefficients β , including: svi local, the model presented in [Section 3.2;](#page-8-0) svi local bd, a the block-diagonal variant designed for faster Cholesky decomposition of the precision matrix presented in [Section 3.3;](#page-10-2) svi classic, a more flexible version of the model from [\[16\]](#page-24-6) that learns a full precision matrix for the coefficients β rather than using a factor structure; as well as INLA and the No U-Turn sampler (NUTS) [\[9\]](#page-24-9). Due to the QR-reparametrization applied to each design matrix, X_i the precision matrix in INLA becomes dense, leading to a cubic complexity for its factorization. We employ one NUTS for each block $\beta_{p,i}$ of the regression coefficients. Assuming that the number of observations N is lower than the number of regression coefficients β , *i.e.* $N < Q$, the asymptotic complexity of the model discussed in [Section 3.2](#page-8-0) and the model in [Section 3.3](#page-10-2) is primarily driven by the Cholesky decomposition of the matrix $\Lambda_{\lambda,\phi}$. Adopting the block diagonal model notably reduces this complexity. The "classical" model introduced by [\[16\]](#page-24-6) achieves the optimal asymptotic complexity of $\mathcal{O}(Q^2)$. While comparing INLA with SVI in terms of asymptotic complexity is challenging, it is important to note that INLA's efficiency can be significantly enhanced by exploiting the sparsity of the precision matrix, reducing factorization to $\mathcal{O}(Q^{\frac{3}{2}})$ [\[26\]](#page-25-7). Conversely, the bottleneck of NUTS lies in the number of Leapfrog steps, which is non-deterministic and can lead to an exponential increase in the algorithm's complexity.

4 Simulations

4.1 Goals of the simulation study

We aim to evaluate the ability of our SVI approach to approximate the posterior in structured additive distributional regression accurately. We have considered two different situations: (i) Binary response whose likelihood terms are notoriously hard to approximate by normal distributions [\[1\]](#page-24-10) and (ii) gamma responses for which both the location and the scale parameter depend on covariates as an example of distributional regression with skewed response distribution. While the skewness makes the likelihood terms hard to fit using our variational normal distributions, the complex dependence between location and scale of the gamma distribution also challenges the SVI variant with block-diagonal covariance structure introduced in [Section 3.3.](#page-10-2)

In both cases, we consider MCMC with a large number of posterior samples as the gold standard against which we benchmark the approximation to the posterior provided by SVI. To quantify the deviation between the approximate posterior and the posterior derived from MCMC, we rely on the Wasserstein Distance (WD), also known as the Earth Mover's Distance (EMD) or the Mallows distance. The Wasserstein distance is a metric that can be used to quantify the difference between two probability distributions in a metric space. Let p and q be two probability distributions defined on a metric space (M, d) , where M is a set and d is the metric. Let $\Omega(p, q)$ denote the set of all joint distributions on $M \times M$ with marginals p and q. The r-th Wasserstein distance between p and q for $r \in [1, +\infty]$ is defined as

$$
W_r(p,q) = \left(\inf_{\omega \in \Omega(p,q)} \mathcal{E}_{(\theta_p,\theta_q)\sim \omega} \left[d(\theta_p,\theta_q)^r\right]\right)^{1/r}.
$$

4.2 Model specification

We conduct tests on various implementations of our approach as well as two different competitors. The first one is the popular framework INLA [\[26\]](#page-25-7), for Integrated Nested Laplace Approximation. The second one belongs to the realm of SVI and is inspired by [\[16\]](#page-24-6). For our approach, we focus on two distinct aspects: (i) Determination of the smoothing parameters by direct estimation via a point estimate (indicated with the suffix point in the following) or employing a variational distribution also for the smoothing parameters (suffix hyper). In all approaches, we use a log-precision parametrization of the hyper-parameters

$$
\boldsymbol{\tau} = (\log \lambda_{1,1}, \ldots, \log \lambda_{1,J_p}, \ldots, \log \lambda_{P,1}, \ldots, \log \lambda_{P,J_P})^{\mathsf{T}}
$$

in accordance with the specifications in INLA. (ii) The construction of the variational family via a local variational distribution for β (explained in [Section 3.2](#page-8-0) and [Section 3.3\)](#page-10-2) versus employing an amortized variational distribution. In the latter, instead of individual variational parameters $\boldsymbol{\mu}_i$ and \mathbf{l}_i for each data point i , we introduce a parameterized function represented by a neural network with weights and bias parameters $\bm{\phi}$. This function takes $y_i, \mathbf{X}_i, \bm{\tau}$ as inputs and yields the respective parameters $\boldsymbol{\mu}_i$ and \mathbf{l}_i :

$$
flatten(\boldsymbol{\mu}_i, l_i) = NN_{\phi}(y_i, \mathbf{X}_i, \boldsymbol{\tau}) \in \mathbb{R}^{P+P(P+1)/2}.
$$
\n(14)

This leads us to the following eight competing approaches:

- 1. svi local SVI with a local variational distribution as presented in [Section 3.2.](#page-8-0)
- 2. svi local bd SVI with a local variational distribution and block-diagonal dependence structure as presented in [Section 3.3.](#page-10-2)
- 3. svi amortized SVI with amortized variational distribution as presented in [Section 3.2](#page-8-0) where the local variational parameters $\boldsymbol{\mu}_i$ and \mathbf{l}_i are modelled using a neural network with weights and bias parameters $\boldsymbol{\phi}$ and input y_i , \mathbf{X}_i and $\boldsymbol{\tau}.$
- 4. svi amortized bd SVI with amortized variational distribution and block-diagonal de-pendence structure as presented in [Section 3.3,](#page-10-2) where the local variational parameters $\mu_{i,p,j}$ and $\sigma^2_{i,p,j}$ are modelled using a neural network with weights and bias parameters $\bm{\phi}_{p,j}$ and input y_i , $\mathbf{x}_{i,p,j}$ and $\tau_{p,j}$ for $p=1,\ldots,P$ and $j=1,\ldots,J_p.$
- 5. svi_classic SVI where the variational distributions for the regression coefficients β and the smoothing parameters τ are assumed to be independent. This model is inspired by [\[16\]](#page-24-6) but, instead of employing a factor structure for the precision matrix for the coefficients β, we learn a full matrix. The variational distribution for β is a full Multivariate Normal Distribution parametrized by the location vector β and the Cholesky factor L:

$$
q(\boldsymbol{\beta}) = \mathcal{N}\left(\tilde{\boldsymbol{\beta}}, \left(\mathbf{L}\mathbf{L}^{\mathsf{T}}\right)^{-1}\right).
$$

The variational parameters ϕ are simply $\tilde{\beta}$ and l, where $I = \psi(L)$ is the log-Cholesky parametrization of the matrix L.

6. svi classic joint – SVI overcoming the independence limitations imposed by svi classic using a single multivariate normal distribution, parametrized by the location vector $\tilde{\gamma}$ and the Cholesky factor ${\bf L}$, to model the vector of all model parameters $\pmb{\gamma}=(\pmb{\beta}^{\sf T},\pmb{\tau}^{\sf T})^{\sf T}$:

$$
q(\boldsymbol{\gamma}) = \mathcal{N}\left(\tilde{\boldsymbol{\gamma}}, \left(\mathbf{C} \mathbf{C}^{\mathsf{T}}\right)^{-1}\right)
$$

.

The variational parameters ϕ are simply $\tilde{\gamma}$ and c, such that $\mathbf{c} = \psi(\mathbf{C})$.

- 7. inla Integrated Nested Laplace Approximation (INLA) [\[26\]](#page-25-7) is a statistical method used for Bayesian inference. It provides an efficient alternative to MCMC techniques for approximating the posterior distributions of model parameters leveraging on the nested Laplace approximations. INLA has become extremely widely used in many application areas of statistics due to the availability of fast, free software [\[20\]](#page-25-8).
- 8. mcmc Markov Chain Monte Carlo. We used the software Liesel [\[24\]](#page-25-9) with one NUTS [\[9\]](#page-24-9) for each block of the regression coefficients β and another one for all smoothing parameters τ .

All models are estimated using a Log-Gamma (see [Equation 16](#page-28-0) for the Log-Gamma pdf) prior on τ with concentration 1.0 and rate 0.01. In each simulation scenario, we use NUTS to generate 8000 warmup samples, followed by 4000 posterior samples and 4000 terminal samples. Each SVI model with a point estimate for the smoothing parameters undergoes training for 8000 epochs, with the sample size S for the approximation of the ELBO set to 64. When using a variational distribution for the hyper-parameters τ , we split the estimation into two stages: first, we estimate τ with a point estimate for 6000 epochs and then, we consider normal distribution as the variational approximation for each $\tau_{p,j}$ where we keep the point estimates as the location parameters and only estimate the variance parameters of the variational distributions for 2000 additional epochs. This allows us for a faster estimation, reducing the number of epochs and the number of samples S_{τ} , and consequently the number of Cholesky decomposition of the matrix $\Lambda_{\tau,\epsilon}^{-1}$ $_{\bm{\tau},\bm{\phi}}^{-1}$, for the estimation of all variational parameters. In this case, we set S_β to 32 and S_τ to 2. All neural networks adopted for the amortized model have a single hidden layer of 16 neurons. All SVI models are trained using the Adam optimizer [\[11\]](#page-24-11) with a learning rate set to 0.01.

The tests and simulations were conducted on a 2021 MacBook Pro equipped with an Apple M1 Pro CPU, 32GB of memory, running macOS Sequoia 15.0.1, and a 512GB SSD.

4.3 Binary regression

In this section, we consider logistic regression models with binary responses and correlated sparseregion covariates, which can lead to biased estimates and high uncertainty [\[6\]](#page-24-12), providing a challenging case for the normal distributions that we utilize in the construction of the variational family. For each of the 100 simulation replications, we generate $N = 200$ observations according to the following scheme:

- Generate $x_{i,1}$ from a three-component mixture of $\mathcal{U}(0, \frac{\pi}{3})$ $\frac{\pi}{3}$), $\mathcal{U}(\frac{\pi}{3})$ $\frac{\pi}{3}, \frac{2\pi}{3}$ $(\frac{2\pi}{3})$, and $\mathcal{U}(\frac{2\pi}{3})$ $(\frac{2\pi}{3}, \pi)$ with mixture weights $9/20$, $2/20$, and $9/20$.
- Generate $z_{i,2}$ from a two-component mixture of $\mathcal{U}(-\pi,-\frac{\pi}{3})$ $\frac{\pi}{3}$) and $\mathcal{U}(-\frac{\pi}{3})$ $(\frac{\pi}{3},0)$ with mixture weights 18/20 and 2/20.
- \bullet Generate $x_{i,2} = (\rho \ast x_{i,1}) + \sqrt{1-\rho^2} \ast z_{i,2}$ where $\rho = -0.7$ governs the correlation between x_1 and x_2 .
- Generate $y_i \sim \text{Bernoulli}(\sin(1.75 * x_{i,1}) + \cos(-1.75 * x_{i,2})).$

We model the nonlinear effects of x_{i1} and x_{i2} based on cubic P-splines with 10 knots and secondorder random walk prior. For each simulation replication, we change the data and initialize all SVI models and MCMC with a different seed.

Figure 1: Logistic model - Boxenplots of the WD between the MCMC posterior for β and the estimated posterior for each of the SVI approaches. Each simulation replicate produces one sample of WD from which the boxenplots are calculated. The panels show the distances after 100, 500, 1000 and 2000, 4000 and 8000 epochs, respectively. Here, we use a point-estimate for τ .

[Figure 1](#page-16-0) illustrates the convergence of the optimisation for determining the variational distribution based on the WD for the regression coefficients β in each SVI model after 100, 500, 1000, 2000, 4000 and 8000 epochs. The figure reveals that all variations of our model converge to a lower distance in fewer iterations compared to the classic or joint model. Despite the strong correlation between the two covariates, the difference between the block-diagonal model and the full model is negligible. Additionally, there is no observable difference between the local and amortized variational distributions. [Figure 8](#page-28-1) in the appendix shows similar information for the posterior of the precision parameters τ . As mentioned earlier, we initially estimated each model with a point estimate $\hat{\tau}$ for the hyper-parameters and then constructed a Normal variational distribution centred at $\hat{\tau}$. Consequently, there are minimal to no improvements in the WD across epochs for each model variation.

Figure 2: Logistic model - WD between the posterior for β (left) and τ (right) determined from MCMC and the approximate posterior by each of the SVI approaches and INLA. Each simulation replicate produces one sample of WD from which the boxenplots are calculated. As a baseline, we include the WD to a second MCMC posterior estimate.

[Figure 2](#page-17-0) evaluates the quality of the variational families for approximating the posterior at the end of the estimation process and compares them with INLA as well as the results from a separate MCMC chain. The latter provides us with a graphical reference for convergence, given the lack of an absolute scale for the WD. For the regression coefficients (left panel), the variations of our model exhibit similar behaviour, although using a variational distribution for the precision parameter τ tends to result in a slightly lower WD. For the precision parameters τ , our model shows consistent results with both MCMC, while INLA exhibits a higher median value and, also in this case, higher variance.

[Table 1](#page-18-0) shows the execution time for each model, where simulation replicates of each model were run in parallel. It is evident that each SVI model surpasses the performance of the MCMC by a factor of 3 to 5 and, is comparable to INLA when we treat the smoothing parameter as free parameters. As expected, all hyper models require more time to complete the estimation process than their point counterparts, with a time difference ranging from 24 to 39 seconds. Interestingly, the block diagonal models do not provide any improvements over their full counterparts.

4.4 Gamma regression

In this second simulation scenario, we consider a distribution regression setup with gamma distributed responses obtained from the following scenario:

- Generate $x_i \sim \mathcal{U}(0, \pi)$.
- Compute $\mu_i = 3 + \exp{(\sin(1.75 * x_i))}$ and $\sigma_i^2 = \exp{(\cos(-2 * x_i))^2}$.

Table 1: Logistic model - Execution time for each model

 \bullet Generate $y_i \sim \mathsf{Gamma}(\mu_i, \sigma_i^2)$ where μ_i is the mean of the Gamma distribution and σ_i^2 is the variance.

The effect of the covariate x_i on both the location and the variance are modelled with a cubic P-spline with 10 knots and second-order random walk prior. For each of the 100 simulation replications, we generate $N = 500$ observations and initialize all SVI models and MCMC with a different seed. Note that in this scenario, we could not use the software INLA, since there are no APIs for estimating both moments of a Gamma response variable.

[Figure 9](#page-28-2) depicts the WD for the regression coefficients β between each SVI model using a point estimate of the hyper-parameters τ and MCMC at epochs 100, 500, 1000, 2000, 4000 and 8000. Also in this scenario, all variations of our model converge more quickly than the leading competitors. However, unlike the logistic regression, the learning process is significantly influenced by the non-orthogonality between the mean and variance of a Gamma distribution, favouring therefore the full model, which can exploit the information given by the covariates X to learn the correlation between the two response parameters, over the block-diagonal approach. Unlike the logistic regression results, when looking at the WD for the precision parameters τ in [Figure 10,](#page-29-0) the svi_classic_hyper model does not show any improvement, in contrast to all other models, even after extensive training of 2000 epochs. The difference between the posterior distribution estimated by MCMC and the one estimated by svi_classic_hyper for τ may be influenced by the model's independence assumption between the regression coefficients and the smoothing parameters.

[Figure 3](#page-19-0) displays the WD between the estimated posterior of all models and MCMC posterior at the conclusion of the estimation process. Similar to the logistic regression case, all SVI models demonstrate comparable performance in both median and variance. However, using a variational distribution for τ appears to result in a slightly improved WD. As for the smoothing parameters τ , the results reflect what was shown in [Figure 10,](#page-29-0) with our model emulating the posterior

Figure 3: Gamma model - WD between the posterior for β (left) and τ (right) determined from MCMC and the approximate posterior by each of the SVI approaches and from a separate MCMC chain. Each simulation replicate produces one sample of WD from which the boxenplots are calculated.

distribution of the precision parameters estimated by MCMC better than svi_classic_hyper. Notice that, in contrast to the logistic regression simulations, in this case, no model's posterior distribution is comparable to MCMC.

As for the logistic model, the execution times for each model are summarized in [Table 2.](#page-19-1) Once again each SVI model outperforms MCMC, in this case by a factor ranging from 4 to 6. In contrast to the logistic regression, the block diagonal models brings an improvement in terms of performance compared to their full parts, with an execution time that reduces by roughly 40 seconds.

5 Applications

We now evaluate our model by contrasting its posterior estimate with results obtained through MCMC on a dataset containing patents spanning the fields of biotechnology/pharmaceutics and semiconductor/computer. The variables encompassed in this dataset are detailed in [Table 4.](#page-29-1) Our objective in this analysis is to develop a model that predicts the number of citations, leveraging a range of explanatory variables for the count response variable "number of claims". Following the prior research in $[10]$, we excluded about 1% of observations with extreme values in certain variables, resulting in a dataset of 4,804 patents.

We assume a Negative Binomial response and model the mean $\mu > 0$ and the dispersion $\delta > 0$ parameter with an additive model. In particular, we assume that both parameters are a function of all the available covariates:

 $\mu_i=\beta_{\mu,0}+f({\sf year}_i)+f({\sf{ncountry}}_i)+f({\sf{nclaim}}{s_i})+{\sf opp}_i+{\sf biopharm}_i+{\sf ustwin}_i+{\sf patus}_i+{\sf pattsgr}_i$ $\delta_i=\beta_{\delta,0}+f({\sf year}_i)+f({\sf{ncountry}}_i)+f({\sf{nclains}}_i)+{\sf opp}_i+{\sf biopharm}_i+{\sf ustwin}_i+{\sf patus}_i+{\sf patsgrr}_i,$ where all functions f are cubic P-splines with 10 knots.

As already analyzed in [\[15\]](#page-24-14), this study faces challenges in estimating the regression coefficients β , where the marginal posterior distributions may deviate slightly from normality. This deviation can lead to biased results if a Gaussian approximation of the posterior distribution, as outlined in [Section 3,](#page-5-0) is used. The authors highlighted that the number of citations is highly overdispersed, with minimal evidence for zero-inflation, which resulted in a "flat likelihood," leading to nonstationary MCMC paths when estimating the model's parameters. The authors in [\[13\]](#page-24-15) tackled the problem using scale-dependent priors, instead of classical Inverse Gamma, mitigating convergence issues and providing more reliable results without extensive manual intervention.

We compare the posterior distribution estimated by our model presented in [Section 3.2](#page-8-0) and [Section 3.3](#page-10-2) with the one estimated by MCMC and by svi_classic_joint by plotting the Kernel density estimate of the marginal distributions of a subset of regression coefficients β and the hyper-parameters τ . As for the simulation studies, we used the software Liesel [\[24\]](#page-25-9) with one NUTS for each block of the regression coefficients β and another one for all smoothing parameters τ .

[Figure 4](#page-21-0) presents the Kernel density estimates for six regression coefficients β , with mean μ displayed in the first row and dispersion δ in the second row. Notably, the intercept is represented by the first plot in each row. The results indicate that all models exhibit similar distributional behaviour, with the SVI models encompassing the posterior mass of the MCMC models, despite the latter demonstrating greater estimation uncertainty. Only the intercept for the dispersion parameters shows some bias.

[Figure 5](#page-21-1) displays the Kernel density estimates for all hyperparameters τ on a log scale. The marginal posterior distributions from all SVI models align with those estimated by MCMC. Although the posterior distribution of the precision parameter for the P-spline modelling the number of countries for dispersion appears biased, the log scale of the x-axis reveals that this hyperparameter takes on a high value (> 54) , indicating minimal penalization. Therefore, it is difficult to assert that the model is biased, as the posterior mode lacks a practical interpretation.

[Figure 6](#page-22-0) and [Figure 7](#page-22-1) are particularly helpful in this regard. They demonstrate that, although there are minor variations in the estimated regression coefficients β and smoothing parameters

Figure 4: Patents citations dataset - Kernel density for six regression coefficients β for the loglocation (first row) and for the log-dispersion (second row).

Figure 5: Patents citations dataset - Kernel density for all smoothing parameters τ . The first three belong to the log-location while the last three to the log-dispersion.

 τ , all models produced similar non-linear effects for each covariate in both the log-location and log-dispersion, with comparable credible intervals.

Figure 6: Patents citations dataset - Estimated splines nonlinear effects of continuous covariates year, ncountry and nclaims on the log-location.

Figure 7: Patents citations dataset - Estimated splines nonlinear effects of continuous covariates year, ncountry and nclaims on the log-dispersion.

In conclusion, despite the challenges highlighted by the authors in [\[13\]](#page-24-15) when estimating the posterior distribution of the regression coefficients β and smoothing parameters τ , where the use of an Inverse-Gamma hyper-prior led to non-stationary MCMC chains, a Gaussian variational distribution, supported by the Bernstein von-Mises theorem for the regression parameters, yields results comparable to those obtained by MCMC in terms of both posterior mode and uncertainty.

6 Conclusion

In this paper, we introduced an SVI model designed for posterior estimation within the framework of structured additive distributional regression. We proposed two distinct strategies for constructing the variational distribution q, leveraging information from the covariates X , potentially the observations y, and the hyperparameters λ to expedite the estimation process. The first approach utilizes both the covariates X and the hyperparameters λ to simultaneously learn the location vector and the precision matrix. The second approach addresses the computational complexity of the first by initially assuming independence among the smooth terms $f_{i,p,j}$ and later reintroducing correlation through a set of additional variational parameters. We also presented two methods for estimating the smoothing parameters λ . The first treats λ as free parameters and provides point estimates, while the second incorporates uncertainty by using a variational approximation to the posterior distribution. We evaluated our approach against two alternative methods: the widely-used INLA framework [\[26\]](#page-25-7) for Integrated Nested Laplace Approximation and a more advanced version of the variational inference model from [\[16\]](#page-24-6), which directly estimates the Cholesky factor of the full precision matrix for greater modelling flexibility. In the simulation study detailed in [Section 4,](#page-13-0) we assessed the ability of our SVI model to accurately approximate the posterior in structured additive distributional regression. Two scenarios were considered: (i) binary response models, which pose challenges for normal distribution approximations of likelihood terms [\[1\]](#page-24-10), and (ii) gamma response models, where both the location and scale parameters depend on covariates, illustrating a case of distributional regression with a skewed response distribution. Across both tasks, our model variations demonstrated faster convergence to a lower WD compared to the classic and joint models. In the logistic regression case, the posterior estimates for both the regression coefficients β and the model parameters λ were similar to those produced by MCMC, though INLA exhibited greater variance in terms of WD for λ . In the gamma regression scenario, our model more accurately replicated the MCMC-estimated posterior distribution of the smoothing parameters than svi_classic_hyper. Finally, in [Section 5,](#page-20-0) we validated our model by comparing its posterior estimates with those obtained via MCMC on a dataset of patents related to biotechnology/pharmaceutics and semiconductor/computer fields. Again, the kernel density estimates for a subset of the regression coefficients β and smoothing parameters λ , along with the plots of the estimated non-linear effects on both response parameters of the Negative Binomial likelihood, revealed comparable results across all models. We plan to extend our framework in two key directions. First, we intend to explore the use of Normalizing Flows, a class of deep generative models that transform a simple probability distribution, often Gaussian, into a more complex one through a series of invertible mappings [\[23\]](#page-25-10), as an alternative for the variational distribution. Incorporating Normalizing Flows within structured additive distributional regression could provide significant advantages, as they offer a flexible approach to modelling complex, high-dimensional probability distributions commonly found in real-world data. Unlike traditional SVI methods that rely on fixed distributional assumptions, Normalizing Flows can capture intricate dependencies and non-linear relationships between covariates and the response distribution, potentially leading to more accurate posterior estimates and better uncertainty quantification. Second, we aim to expand the model to handle multivariate responses by employing copulas, building on the work of [\[15,](#page-24-14) [14,](#page-24-16) [19\]](#page-25-11). This extension would allow for the modelling of dependencies between multiple response variables, further enhancing the applicability of the framework to more complex data scenarios.

References

- [1] J. H. Albert and S. Chib. Bayesian analysis of binary and polychotomous response data. Journal of the American statistical Association, 88(422):669–679, 1993.
- [2] S. Banerjee, B. P. Carlin, and A. E. Gelfand. Hierarchical Modelling and Analysis for Spatial Data (second edition). Chapman & Hall / CRC, 2014.
- [3] D. M. Blei, A. Kucukelbir, and J. D. McAuliffe. Variational inference: A review for statisticians. J Am Stat Assoc, 112(518):859–877, 2017.
- [4] L. Fahrmeir and T. Kneib. Bayesian Smoothing and Regression for Longitudinal, Spatial and Event History Data. Oxford University Press, 2011.
- [5] L. Fahrmeir, T. Kneib, and S. Lang. Penalized structured additive regression for space-time data: a bayesian perspective. Statistica Sinica, pages 731–761, 2004.
- [6] J. Friedman. The elements of statistical learning: Data mining, inference, and prediction. (No Title), 2009.
- [7] P. Hall, J. T. Ormerod, and M. P. Wand. Theory of Gaussian variational approximation for a Poisson mixed model. Statistica Sinica, pages 369–389, 2011.
- [8] M. D. Hoffman, D. M. Blei, C. Wang, and J. Paisley. Stochastic variational inference. J Machine Learning Res, 14(1):1303–1347, 2013.
- [9] M. D. Hoffman, A. Gelman, et al. The no-u-turn sampler: adaptively setting path lengths in hamiltonian monte carlo. J. Mach. Learn. Res., 15(1):1593–1623, 2014.
- [10] A. Jerak and S. Wagner. Modeling probabilities of patent oppositions in a bayesian semiparametric regression framework. Empirical Economics, 31:513–533, 2006.
- [11] D. P. Kingma and J. Ba. Adam: A method for stochastic optimization. arXiv:1412.6980, 2014.
- [12] D. P. Kingma and M. Welling. Auto-encoding variational Bayes. arXiv:1312.6114, 2013.
- [13] N. Klein and T. Kneib. Scale-dependent priors for variance parameters in structured additive distributional regression. Bayesian Analysis, 11:1071–1106, 2016. doi: 10.1214/15-BA983.
- [14] N. Klein and T. Kneib. Simultaneous inference in structured additive conditional copula regression models: a unifying Bayesian approach. Statistics and Computing, 26:841–860, 2016. ISSN 0960-3174. doi: 10.1007/s11222-015-9573-6.
- [15] N. Klein, T. Kneib, S. Klasen, and S. Lang. Bayesian structured additive distributional regression for multivariate responses. J Royal Stat Soc C, 64:569–591, 2015. ISSN 00359254. doi: 10.1111/rssc.12090.
- [16] J. Kleinemeier and N. Klein. Scalable estimation for structured additive distributional regression through variational inference. arXiv preprint arXiv:2311.07371, 2023.
- [17] T. Kneib. Beyond mean regression. Statistical Modelling, 13:275–303, 2013. ISSN 1471- 082X. doi: 10.1177/1471082X13494159.
- [18] T. Kneib, A. Silbersdorff, and B. Säfken. Rage Against the Mean - A review of distributional regression approaches. Econometrics Stat, 2021.
- [19] L. Kock and N. Klein. Truly multivariate structured additive distributional regression. $arXiv$ preprint arXiv:2306.02711, 2023.
- [20] F. Lindgren and H. Rue. Bayesian spatial modelling with r-inla. Journal of statistical software, 63(19), 2015.
- [21] J. T. Ormerod and M. P. Wand. Gaussian variational approximate inference for generalized linear mixed models. J Comput Graph Stat, 21(1):2-17, 2012.
- [22] J. C. Pinheiro and D. M. Bates. Unconstrained parametrizations for variance-covariance matrices. Statistics and computing, 6:289–296, 1996.
- [23] D. J. Rezende and S. Mohamed. Variational inference with normalizing flows. arXiv:1505.05770, 2015.
- [24] H. Riebl, P. F. V. Wiemann, and T. Kneib. Liesel: A probabilistic programming framework for developing semi-parametric regression models and custom bayesian inference algorithms, 2022. URL <https://arxiv.org/abs/2209.10975>.
- [25] H. Rue, S. Martino, and N. Chopin. Approximate bayesian inference for latent gaussian models by using integrated nested laplace approximations. Journal of the royal statistical society: Series b (statistical methodology), 71(2):319–392, 2009.
- [26] H. Rue, A. Riebler, S. H. Sørbye, J. B. Illian, D. P. Simpson, and F. K. Lindgren. Bayesian computing with inla: a review. Annual Review of Statistics and Its Application, 4:395–421, 2017.
- [27] D. Ruppert, M. P. Wand, and R. J. Carroll. Semiparametric Regression. Cambridge University Press, 2003.
- [28] S. N. Wood. Generalized Additive Models: An Introduction with R (second edition). Chapman & Hall / CRC, 2017.
- [29] C. Zhang, J. Bütepage, H. Kjellström, and S. Mandt. Advances in variational inference. IEEE Transactions on Pattern Analysis and Machine Intelligence, 41(8):2008–2026, 2018.

A Appendix

A.1 Identifiability

In additive regression models, the additive effects have to be constrained to ensure identifiability since

$$
f_{p,1}(\mathbf{x}_i, \boldsymbol{\beta}_{p,1}) + f_{p,2}(\mathbf{x}_i, \boldsymbol{\beta}_{p,2}) = f_{p,1}(\mathbf{x}_i, \boldsymbol{\beta}_{p,1}) + c + f_{p,2}(\mathbf{x}_i, \boldsymbol{\beta}_{p,2}) - c
$$

This is usually achieved by

$$
\sum_i^N f_{p,j}^{\text{new}}(x_{i,j},\beta_{p,j})=0,
$$

with

$$
f_{p,j}(x_{i,j}, \beta_{p,j}) = f_{p,j}^{\text{new}}(x_{i,j}, \beta_{p,j}) + \beta_{0,p,j}.
$$

To do so, we calculate the vector $\mathbf{C}_{p,j}$ of columns means of $\mathbf{X}_{p,j}$ and we build the QRdecomposition of

$$
\mathbf{C}_{p,j} = \mathbf{Q}_{p,j} \mathbf{R}_{p,j},
$$

where $\mathbf{Q}_{p,j}$ is an orthogonal matrix and

$$
\mathbf{R}_{p,j} = (\tilde{R}_{p,j}, 0, \ldots, 0)^{\mathsf{T}},
$$

where $\tilde{R}_{p,j}$ is the upper triangular matrix of dimension $1\times1.$ Then, we split

$$
\mathbf{Q}_{p,j}=\left[\mathbf{Z}_{p,j}^{(a)},\mathbf{Z}_{p,j}^{(b)}\right],
$$

where $\mathbf{Z}_{p,j}^{(a)}$ is a $Q_{p,j}\times 1$ column vector. Finally, we write the new design as

$$
\mathbf{X}_{p,j}^{\mathsf{new}} = \mathbf{X}_{p,j} \mathbf{Z}_{p,j}^{(b)}
$$

and the new penalty matrix as:

$$
\mathbf{K}^{\mathsf{new}}_{p,j}(\lambda_{p,j}) = \left(\mathbf{Z}^{(b)}_{p,j}\right)^{\mathsf{T}} \mathbf{K}_{p,j} \mathbf{Z}^{(b)}_{p,j}.
$$

The design matrix for one parameter p can be written as:

$$
\mathbf{X}_{p} = \left[\mathbf{1}, \mathbf{X}_{p,1}^{\mathsf{new}}, \ldots, \mathbf{X}_{p,J_{p}}^{\mathsf{new}}\right] \in \mathbb{R}^{N \times Q_{p}}
$$
(15)

where $Q_p \coloneqq (Q_{p,1} + \cdots + Q_{p,J_p})$ and the penalty matrix as:

$$
\mathbf{K}_{p}(\boldsymbol{\lambda}_{p}) = \begin{pmatrix} 0 & \dots & 0 \\ 0 & \lambda_{p,1}^{2} \mathbf{K}_{p,1}^{\text{new}} \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_{p,J_{p}}^{2} \mathbf{K}_{p,J_{p}}^{\text{new}} \end{pmatrix}.
$$

Model Model **Asymptotic complexity** svi_local $3 + NPQ^2 + SNQ + S\sum$ Construction of q $\ln p(\mathbf{y}|\mathbf{\beta}, \mathbf{X})$ $\ln p(\mathbf{\beta}|\mathbf{\lambda})$ $\ln q(\mathbf{\beta}|\mathbf{\lambda}, \mathbf{X}, \mathbf{\phi})$ P $p=1$ \sum J_p $j=1$ $Q_{p,j}^2 + SQ^2$) svi_local_bd $2+\sum$ P $p=1$ \sum J_p $j=1$ $Q_{p,j}^3+N\sum$ P $p=1$ \sum J_p $j=1$ $Q_{p,j}$ $\overline{}$ Construction of q Construction of q + SNQ $\ln p(\mathbf{y}|\boldsymbol{\beta}, \mathbf{X})$ $+ S \sum$ P $p=1$ \sum J_p $j=1$ $Q_{p,j}^2$ $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$ $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$ + SQ^2 $\widetilde{\beta(\lambda,2)}$ $\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})$) svi _cclassic $\mathcal{O}($ 2 Construction of q $+$ SNQ $\ln p(\mathbf{y}|\boldsymbol{\beta}, \mathbf{X})$ $+ S \sum$ P $_{p=1}$ \sum J_p $j=1$ $Q_{p,j}^2$ $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$ $+$ SQ^2 $\ln q(\boldsymbol{\beta}|\boldsymbol{\lambda}, \mathbf{X}, \boldsymbol{\phi})$) INLA $\mathcal{O}(\hspace{1cm}Q^3$ Factorization of the precision matrix $\ln p(\mathbf{y}|\boldsymbol{\beta}, \mathbf{X})$ $+ NQ + \sum$ P Jp $p=1$ \sum $j=1$ $Q_{p,j}^2)$ 2 $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$ **NUTS** \sum P $p=1$ \sum J_p $j=1$ 2 $D_{p,j}$ Leapfrog steps Gradient of $\ln p(\mathbf{y}, \boldsymbol{\beta}_{p,j} | \boldsymbol{\lambda}_{p,j}, \mathbf{X}_{p,j})$ $Q_{p,j}$ $+ NQ + \sum$ $\ln p(\mathbf{y}|\boldsymbol{\beta}, \mathbf{X})$ P $p=1$ \sum J_p $j=1$ $Q_{p,j}^2$ $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$ $\ln p(\boldsymbol{\beta}|\boldsymbol{\lambda})$

A.2 Asymptotic complexity

Table 3: Asymptotic complexity of different SVI models compared with INLA and the No-U-Turn Sampler (NUTS). Due to the QR-reparametrization applied to each design matrix X_i the precision matrix in INLA becomes dense, leading to a cubic complexity for its factorization. We employ one No-U-turn sampler (NUTS) for each block $\beta_{p,j}$ of the regression coefficients.

A.3 Log-Gamma distribution

The probability density function of the Log-Gamma distribution with concentration parameter $\alpha > 0$ and rate parameter $\beta > 0$ is:

$$
p(x) = \frac{e^{x^{\alpha}}e^{-e^{x}\beta}}{\Gamma(\alpha)\beta^{-\alpha}}.
$$
\n(16)

A.4 Additional figures and tables

Figure 8: Logistic model - WD between the posterior for τ determined from MCMC and the approximate posterior by each of the SVI approaches after 100, 500, 1000 and 2000 epochs. Each simulation replicate produces one sample of WD from which the boxenplots are calculated.

Figure 9: Gamma model - WD after 100, 500, 1000, 2000, 4000, and 8000 epochs between each VI model with a point estimate for τ and MCMC for β . Each simulation replicate produces one sample of WD from which the boxenplots are calculated.

Figure 10: Gamma model - WD after 100, 500, 1000 and 2000 epochs between each VI model and MCMC for τ . Each simulation replicate produces one sample of WD from which the boxenplots are calculated.

Table 4: Patent citations: description of variables including summary statistics