

Microcanonical phase space and entropy in curved spacetime

Avinandan Mondal*, Dawood Kothawala†

*Department of Physics, Indian Institute of Technology Madras,
Chennai 600036.*

Abstract

We discuss the structure of microcanonical ensembles in inertial and non-inertial frames attached to a confined system of positive energy particles in curved spacetime. Under certain physically reasonable assumptions that ensure the existence of such ensembles, we derive two generic characteristics of the leading curvature corrections: (1) they are characterized by Ricci and Einstein tensors, and (2) their contribution is proportional to the boundary area of the system. We also obtain exact analytical results for microcanonical ensembles in de Sitter spacetime, which reproduce the perturbative results in the limit $\Lambda \rightarrow 0$, but have interesting limits when the size of the system is comparable to $\Lambda^{-1/2}$. We give a general argument to highlight two distinct sources of divergences in the phase space volume, coming from redshift and spatial geometry, and illustrate this by comparing and contrasting the results for (i) geodesic box in de Sitter, (ii) geodesic box in Schwarzschild, and (iii) uniformly accelerated box in Minkowski. Finally, we extend these results to N particle systems in the massless (ultra-relativistic) limit for certain restricted class of spacetimes.

*Email: ph20b005@smail.iitm.ac.in

†Email: dawood@iitm.ac.in

Contents

1	Introduction	2
2	Constant energy hypersurface: a covariant description	3
2.1	One particle phase space volume	5
2.2	Multi-particle systems	5
2.2.1	Massless (ultra-relativistic) particles in Minkowski	6
2.2.2	Massless (ultra-relativistic) particles in metrics with $g^{0\mu} = 0$	7
3	Single particle system: Exact results for specific spacetimes	9
3.1	Uniform Acceleration in Flat Spacetime	9
3.2	Static spherically symmetric black holes	10
3.3	de Sitter (dS) spacetime	12
4	Leading curvature corrections in arbitrary curved spacetimes	14
4.1	One particle phase space volume	14
4.2	Microcanonical entropy and temperature	18
4.3	Multi-particle systems	19
5	Discussion	21
A	Derivation of the Formula for $\Gamma(E)$	22

1 Introduction

Statistical mechanics of particle systems in relativistic settings and self-gravitating systems has been a long-standing topic of interest from both a purely academic perspective and for its relevance in astrophysical situations. Extensive discussions on this topic exist in the literature, and some notable contributions include works on isothermal spheres in Newtonian gravity [1, 2], relativistic kinetic theory [3], and self-gravitating quantum gases [4], among others. The unique issues posed by gravity due to its long range nature and, in the general relativistic framework, due to self-gravitating particles back reacting on the metric, are well acknowledged in the community, and it is expected that a satisfactory clarification of some of these issues might be of fundamental significance. For instance, negative specific heat capacities associated with self-gravitating systems like isothermal spheres [2] and black holes [5] highlight the inequivalence of statistical ensembles and it has been pointed out in [5] that microcanonical ensemble must be used to discuss

thermodynamics of systems where gravitational interactions are important. Extreme cases when such considerations become crucial are quantum black holes which display thermodynamic properties.

In this work, we use microcanonical ensemble to describe confined system of particles in a given background geometry. The main aim is to bring out the purely geometric aspects of phase space volume in curved spacetime, in the hope that these will help understand peculiar thermodynamic properties of systems in a gravitational field. While we do not include self-gravity in our analysis, in the sense that no back reaction on the underlying geometry due to the particles is taken into account, we expect this to be possible based on the results we derives. A large part of the work is devoted to single particle system which, although it has very little significance in statistical mechanics, exhibits non-trivial and interesting results like divergence of phase space volume as the system approaches event horizon of a static black hole or as the system size approaches cosmological horizon of a de Sitter spacetime. Another justification for doing single particle phase space calculations lie in the result that at least for massless systems in some restricted class of spacetimes, we have the phase volume of a massless N -particle system related to that of a one particle system by a simple relation.

Conventions: We work with $(-, +, +, +)$ metric signature. The Greek indices run over spatial components, whereas Latin indices run over both space and time components. Also, we shall be explicitly dealing with $(3+1)$ dimensional spacetimes although generalizations to $(D + 1)$ dimensional spacetimes will be obvious from our analysis. Finally, the matrix version of a rank-2 tensor U written in some coordinate system will be denoted by $[U]$ and the matrix components as $[U]_{ab}$ where the indices a, b are matrix indices.

2 Constant energy hypersurface: a covariant description

Let $\{x^i | i = 0, 1, 2, 3\}$ be a coordinate chart on the region of interest of some arbitrary spacetime (we are allowing coordinate singularities for this chart). $\{x^\alpha | \alpha = 1, 2, 3\}$ are then a chart on each $x^0 = \text{constant}$ submanifolds. Also, we assume that our spacetime has a Killing vector field ξ^i which is timelike at the "nice regions of spacetime" (by "nice" we refer to regions such as exterior of event horizon of a black hole, interior of cosmological horizon of de Sitter, etc.).

Suppose we have a container with N free particles in this spacetime. Let $p_{(a)}^i$ be the momentum vector field defined on the world line of the a -th particle. The Hamiltonian of the system is given by¹ :

$$H = - \sum_{a=1}^N \xi^i p_{(a)i} \quad (1)$$

Then, if the system has at most energy E , then phase space volume accessible to the system is given by:

$$\Gamma(E) = \frac{1}{N!} \int d^3\mathbf{x} d^3\mathbf{p} \Theta(E - H) \quad (2)$$

where we assumed indistinguishable particles. Θ is the Heaviside step function and integral is over entire phase space.²

Then, accessible phase space volume for the system with energy E is given by:

$$\Omega(E) = \frac{1}{N!} \int d^3\mathbf{x} d^3\mathbf{p} \delta(E - H) = \frac{d\Gamma(E)}{dE} \quad (3)$$

Note that the number of accessible microstates is actually defined as:

$$\Omega(E) = \frac{1}{N! h^{3N}} \int d^3\mathbf{x} d^3\mathbf{p} \delta(E - H) \quad (4)$$

where we have N no. of particles moving in a spacetime with three spatial dimensions and h is Planck's constant. The factor of h^{3N} makes $\Omega(E)$ dimensionless and serves as a quantity denoting denoting "smallest phase space volume that can be occupied by a microstate because of Heisenberg's uncertainty". However, we shall put $h = 1$ and assume that things have been made dimensionless.

Then, the microcanonical entropy is defined as:

$$S = \log \Omega(E) \quad (5)$$

¹We assume $H > 0$ always.

²The choice behind the volume form on the phase space to be $d^3\mathbf{x}d^3\mathbf{p} = dx^1 \wedge dx^2 \wedge dx^3 \wedge dp_1 \wedge dp_2 \wedge dp_3$ is actually motivated by the following facts: it is invariant under coordinate transformations in constant x^0 foliations and for Minkowski spacetime, this volume form is invariant under Lorentz transformations. Of course in a general spacetime, under general coordinate transformations that mix space and time it is not an invariant volume form. Also note that simply $d^3\mathbf{x}$ would refer to $dx^1 \wedge dx^2 \wedge dx^3$ and just $d^3\mathbf{p}$ would refer to $dp_1 \wedge dp_2 \wedge dp_3$.

and other thermodynamic quantities like temperature, specific heat, etc. can be derived from this.

2.1 One particle phase space volume

One can obtain explicit geometric expressions for the above quantities for the phase space of a single particle assuming that the metric g_{ij} can be described in an approximately time independent manner, through (1). This is a rather lengthy computation that follows the route similar to the one sketched in [7], but generalized to arbitrary metrics; we refer the reader to Appendix-(A) for details.

One obtains the following expression for the phase space volume of a single particle system of energy at most E :

$$\Gamma(E) = \int_V d^3\mathbf{x} \sqrt{\text{Det}([h])} \left(\frac{4\pi}{3}\right) \left[\lambda + \sum_{\mu} \frac{\gamma_{\mu}^2}{4}\right]^{3/2} \quad (6)$$

where V is the spatial extent of the box in which the particle resides; the matrix $[h]$ is the matrix form of the metric on spatial foliations defined as:

$$h_{\alpha\beta} = g_{\alpha\beta} - \frac{g_{0\alpha}g_{0\beta}}{g_{00}} \quad (7)$$

and $\gamma_{\nu} = 2E(g^{0\mu}[\omega]_{\mu\nu})$ and $\lambda = (-g^{00})E^2 - m^2$; further where $[\omega]$ is the matrix that diagonalizes the matrix $[g]$ composed of spatial components of inverse metric to identity matrix.

2.2 Multi-particle systems

For a system with N particles, from (2) and (1), it is obvious that we have to evaluate the integral:

$$\Gamma(E) = \frac{1}{N!} \int d^3\mathbf{x} d^3\mathbf{p} \Theta\left(E - \sum_{k=1}^N H_k\right) \quad (8)$$

where:

$$H_k = -\xi^i p_{(k)i} \quad (9)$$

Now, we recall that $\Theta(x)$ has an integral representation given by:

$$\Theta(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{ixs}}{s - i0^+} \quad (10)$$

Hence,

$$\Theta\left(E - \sum_{k=1}^N H_k\right) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} \prod_{k=1}^N e^{-iH_k s} \quad (11)$$

Computing this for arbitrary curved spacetime and with non-zero mass is a difficult job. Hence, we focus on $m = 0$ case (ultra-relativistic particles or simply massless particles). In the below subsection we compute it for Minkowski spacetime and we shall see that our computation shall give us a direct hint for computing it for some quite general class of spacetimes.

2.2.1 Massless (ultra-relativistic) particles in Minkowski

For Minkowski spacetime, (11) can be written as:

$$\Theta\left(E - \sum_{k=1}^N H_k\right) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} \prod_{k=1}^N e^{-i(|\mathbf{p}_k|^2 + m^2)^{\frac{1}{2}} s} \quad (12)$$

where \mathbf{p}_k is the 3-momentum of k -th particle. Thus,

$$\begin{aligned} & \int \prod_{k=1}^N d^3 \mathbf{p}_k \Theta\left(E - \sum_{k=1}^N H_k\right) \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} \prod_{k=1}^N \left[d^3 \mathbf{p}_k e^{-i(|\mathbf{p}_k|^2 + m^2)^{\frac{1}{2}} s} \right] \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} (4\pi)^N \left[\int_0^{E - Nm} dp p^2 e^{-i(p^2 + m^2)^{\frac{1}{2}} s} \right]^N \end{aligned} \quad (13)$$

Note that the limit of integral over p is from 0 to $E - Nm$ in the last expression as the magnitude of momentum of any particle can never exceed the total energy of the system subtracted by rest mass. In fact any bound larger than or equal to $E - Nm$ shall work. Now, we put $m = 0$ and continue the computation of (13):

$$\begin{aligned} & \int \prod_{k=1}^N d^3 \mathbf{p}_k \Theta\left(E - \sum_{k=1}^N H_k\right) \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} (4\pi)^N \left[\int_0^E dp p^2 e^{-ips} \right]^N \\ &= \frac{(4\pi)^N}{2\pi i} \int_{-\infty}^{\infty} ds \frac{e^{iEs}}{s - i0^+} \left[\frac{2i + e^{-iEs}(-2i + Et(2 + iEs))}{s^3} \right]^N \end{aligned} \quad (14)$$

Now, denoting $f(s) = \frac{2i+e^{-iEs}(-2i+Et(2+iEs))}{s^3}$, it can be easily shown by doing appropriate expansions that $f(s)$ has no pole at $s = 0$ in the complex s -plane. It's singularity at $s = 0$ is a removable one. In fact one obtains:

$$\lim_{s \rightarrow 0} f(s) = \frac{E^3}{3} \quad (15)$$

Now, performing the last integral in (14) by the method of contour integration and closing the contour in the upper half plane (as $E > 0$) and noting (15), we obtain:

$$\int \prod_{k=1}^N d^3 \mathbf{p}_k \Theta \left(E - \sum_{k=1}^N H_k \right) = \left(\frac{4\pi}{3} \right)^N E^{3N} \quad (16)$$

Considering the system is in a container of volume V , we further conclude:

$$\Gamma(E) = \frac{1}{N!} \left(\frac{4\pi V}{3} \right)^N E^{3N} \quad (17)$$

Now, note that the integral (8) for massless particles in Minkowski can be written as:

$$\begin{aligned} \Gamma(E) &= \frac{1}{N!} \int dx^\alpha dp_\alpha \Theta \left(E - \sum_{k=1}^N H_k \right) \\ &= V^N \int \prod_{k=1}^N d^3 \mathbf{p}_k \Theta \left(E - \sum_{k=1}^N |\mathbf{p}_k| \right) \\ &= \frac{1}{N!} (4\pi V)^N \int_0^\infty dp_1 \dots \int_0^\infty dp_N \left(\prod_{i=1}^N p_i^2 \right) \Theta \left(E - \sum_{k=1}^N p_k \right) \end{aligned} \quad (18)$$

Thus, comparing this with (17), we can say that:

$$\int_0^\infty dp_1 \dots \int_0^\infty dp_N \left(\prod_{i=1}^N p_i^2 \right) \Theta \left(E - \sum_{k=1}^N p_k \right) = \left(\frac{E^3}{3} \right)^N \quad (19)$$

We shall be needing this result (19) in the next subsection.

2.2.2 Massless (ultra-relativistic) particles in metrics with $g^{0\mu} = 0$

For $m = 0$ and $g^{0\mu} = 0$, the single particle Hamiltonian in (81) reduces to:

$$H = \frac{\sqrt{g^{\mu\nu} p_\mu p_\nu}}{(-g^{00})^{\frac{1}{2}}} \quad (20)$$

Now, using same definition of \tilde{p}_μ as in (83), we have:

$$H = \frac{|\tilde{\mathbf{p}}|}{(-g^{00})^{\frac{1}{2}}} \quad (21)$$

Hence,

$$\Gamma(E) = \frac{1}{N!} \int \prod_{k=1}^N d^3 \mathbf{x}_k \int \prod_{k=1}^N d^3 \mathbf{p}_k \left(\prod_{i=1}^N \text{Det}([\omega(\mathbf{x}_i)]) \right) \Theta \left[E - \left(\sum_{i=1}^N \frac{|\tilde{\mathbf{p}}_i|}{\sqrt{-g^{00}(\mathbf{x}_i)}} \right) \right] \quad (22)$$

where as we had defined earlier while doing single particle computation:

$$[\omega(\mathbf{x}_i)]^T [g(\mathbf{x}_i)] [\omega(\mathbf{x}_i)] = [\mathbb{1}] \quad (23)$$

So, doing angular integrals in momentum space, we have:

$$\begin{aligned} \Gamma(E) &= \frac{(4\pi)^N}{N!} \int \prod_{k=1}^N d^3 \mathbf{x}_k \left(\prod_{i=1}^N \text{Det}([\omega(\mathbf{x}_i)]) \right) \int_0^\infty dp_1 \dots \int_0^\infty dp_N \\ &\quad \left(\prod_{i=1}^N p_i^2 \right) \Theta \left[E - \left(\sum_{i=1}^N \frac{p_i}{\sqrt{-g^{00}(\mathbf{x}_i)}} \right) \right] \\ &= \frac{(4\pi)^N}{N!} \int \prod_{k=1}^N d^3 \mathbf{x}_k \left(\prod_{i=1}^N \text{Det}([\omega(\mathbf{x}_i)](-g^{00}(\mathbf{x}_i))^{\frac{3}{2}}) \right) \int_0^\infty dq_1 \dots \int_0^\infty dq_N \\ &\quad \left(\prod_{i=1}^N q_i^2 \right) \Theta \left[E - \sum_{i=1}^N q_i \right] \end{aligned} \quad (24)$$

where $q_i = \frac{p_i}{\sqrt{-g^{00}(\mathbf{x}_i)}}$

Now, using (19) in (24), we arrive at:

$$\Gamma(E) = \frac{1}{N!} \left(\frac{4\pi E^3}{3} \right)^N \int \prod_{i=1}^N \left(d^3 \mathbf{x}_i \text{Det}([\omega(\mathbf{x}_i)])(-g^{00}(\mathbf{x}_i))^{3/2} \right) \quad (25)$$

Further recall that $\text{Det}([\omega(\mathbf{x}_i)]) = \sqrt{\text{Det}([h(\mathbf{x}_i)])}$ where in this case $h_{\mu\nu} = g_{\mu\nu} - \frac{g_{0\mu}g_{0\nu}}{g_{00}} = g_{\mu\nu}$. So, denoting the matrix with entries as covariant components of metric ($g_{\mu\nu}$) as $[g_1]$, we have:

$$\Gamma(E) = \frac{1}{N!} \left(\frac{4\pi}{3} \right)^N E^{3N} \left[\int_V d^3 \mathbf{x} \left(\text{Det}([g_1])(-g^{00})^3 \right)^{1/2} \right]^N \quad (26)$$

Labelling single particle phase volume for energy E as $\Gamma_1(E)$ and fetching its expression from (6) and then comparing with (26), we conclude that:

$$\Gamma(E) = \frac{\Gamma_1(E)^N}{N!} \quad (27)$$

The result essentially states that for massless particles in spacetimes described by coordinate chart where metric has no spatio-temporal cross components, studying single particle phase volume is enough. We devote our next sections for studying single particle phase volumes, but however we allow massive particles and most general spacetimes as we have an exact Eq. (6) for single particle system in arbitrary backgrounds.

3 Single particle system: Exact results for specific spacetimes

3.1 Uniform Acceleration in Flat Spacetime

The spacetime metric as described by a particle uniformly accelerated in Cartesian-like coordinate system in a background Minkowski spacetime is given by:

$$g_{00} = -(1 + a_\mu y^\mu)^2 \quad (28)$$

$$g_{\mu\nu} = \delta_{\mu\nu} \quad (29)$$

where $a^i = u^k \nabla_k u^i$ is the acceleration of the centroid of the box.

So, for a single free particle in a box, we can obtain $\Gamma(E)$ by performing the integration in the expression (6). Hence, we have:

$$\Gamma(E) = \int_V d^3\mathbf{x} \left(\frac{4\pi}{3}\right) \left[\frac{E^2}{(1 + a_\mu x^\mu)^2} - m^2 \right]^{3/2} \quad (30)$$

Now, we take the box to be spherical and denoting magnitude of the 3-acceleration as $|\mathbf{a}|$, we have:

$$\Gamma(E) = \left(\frac{4\pi}{3}\right) \int_0^R dr r^2 \int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi \left[\frac{E^2}{(1 + |\mathbf{a}|r \cos\theta)^2} - m^2 \right]^{3/2} \quad (31)$$

Now, the angular integrals for massive case gives hypergeometric functions and then the radial integral is not exactly solvable. So, we rather concentrate on the massless limit. In the massless case, although the integral is exactly solvable, it is still quite complicated. So, we focus on two regimes:

firstly, when the box is small ($R \ll 1/|\mathbf{a}|$) and when the box approaches the Rindler horizon ($R \rightarrow (1/|\mathbf{a}|)^-$)

Case I: Box radius is small compared to Rindler horizon ($R \ll 1/|\mathbf{a}|$)

In this regime, we obtain:

$$\begin{aligned}\Gamma(E) &= \left(\frac{8\pi^2}{3}\right) \left[\frac{2}{3} E^3 R^3 + \frac{4}{5} E^3 |\mathbf{a}|^2 R^5 + \mathcal{O}(R^6) \right] \\ &= \frac{4\pi}{3} E^3 \cdot V \left[1 + \frac{3}{10\pi} A |\mathbf{a}|^2 + \mathcal{O}(R^3) \right]\end{aligned}\quad (32)$$

where V and A denote volume and area of the spherical box containing the particle.

Case II: Box radius approaches Rindler horizon ($R \rightarrow (1/|\mathbf{a}|)^-$)

In this regime, the leading order divergent behaviour of $\Gamma(E)$ is captured by the following expression:

$$\Gamma(E)_{sing} \sim \left(\frac{8\pi^2}{3}\right) E^3 \left[\frac{1}{2|\mathbf{a}|^4 \left(\frac{1}{|\mathbf{a}} - R\right)} - \frac{1}{2} \log \left(\frac{1}{|\mathbf{a}} - R\right) \right]\quad (33)$$

A similar analysis for accelerated box of classical ideal gas in some symmetric spacetimes with Killing horizons has been done in [8]. In this context, it is worth noting that the leading order divergent behaviour of both phase space volume and entropy, as computed from expression (33), will be different from the one mentioned in [8], for two reasons: the difference in spatial geometry of the box (spherical for our case, cubical for [8]), and the ensemble being used (microcanonical vs. canonical).

3.2 Static spherically symmetric black holes

For a static spherically symmetric black hole, in Schwarzschild coordinates have the line element given by:

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 d\Omega^2\quad (34)$$

As $g_{0\alpha} = 0$, so $[h] = [g]$ and $\gamma_\mu = 0$. Thus,

$$\begin{aligned}\Gamma(E) &= \int_V d^3\mathbf{x} \sqrt{\text{Det}([g])} \left(\frac{4\pi}{3}\right) \lambda^{3/2} \\ &= \int_V dr d\theta d\phi \frac{r^2 \sin(\theta)}{\sqrt{f(r)}} \left(\frac{4\pi}{3}\right) \left(\frac{E^2}{f(r)} - m^2\right)^{3/2}\end{aligned}\quad (35)$$

From the expression in (35) it is readily observed that the phase volume diverges when $f(r) = 0$, i.e. for a static black hole when $r = 2GM$ (i.e. at the event horizon). So, as the particle container approaches the event horizon, the phase volume that can be occupied by the particle diverges for a given energy E .³ This has also been observed in [7].

Computation in Painleve-Gullstrand coordinates and covariance of the phase space volume

Since Schwarzschild coordinate chart is not well-behaved at the event horizon, we now redo the computation using Painleve-Gullstrand coordinates which are well-behaved at the horizon. Then in these coordinates, we have the line element given as:

$$ds^2 = -f(r)dT^2 + 2\sqrt{1 - pf(r)}dTdr + pdr^2 + r^2d\Omega^2 \quad (36)$$

where $p \in (0, 1]$.

It can be obtained that, $g^{00} = -K = -p$. So, $\lambda = KE^2 - m^2 = pE^2 - m^2$. The $[g]$ matrix for black hole in Painleve coordinates is given by:

$$[g] = \text{Diag} \left(pf(r), \frac{1}{r^2}, \frac{1}{r^2 \sin^2 \theta} \right) \quad (37)$$

Clearly $[g]$ is already diagonal and hence:

$$[\omega] = \text{Diag} \left(\frac{1}{\sqrt{pf(r)}}, r, r \sin \theta \right) \quad (38)$$

Now as $n^\mu = g^{0\mu}$, so:

$$[n]^T = \left(\sqrt{1 - pf(r)}, 0, 0 \right) \quad (39)$$

and as $\gamma_\nu = 2E(n^\mu[\omega]_{\mu\nu})$, so:

$$[\gamma]^T = 2E[n]^T[\omega] = \frac{2E}{\sqrt{pf(r)}} \left(\sqrt{1 - pf(r)}, 0, 0 \right) \quad (40)$$

³Actually, E is the energy of the particle measured at infinity and it is conserved along the geodesic. Also, another thing to note is that the divergence of phase space volume doesn't come only because the locally measured energy is infinitely redshifted $E_{\text{loc}} = E/\sqrt{f}$ at $f = 0$, but there is a divergence coming from the volume form itself.

Hence, putting all together:

$$\begin{aligned}\Gamma(E) &= \int_V dr d\theta d\phi \frac{r^2 \sin(\theta)}{\sqrt{pf(r)}} \left(\frac{4\pi}{3}\right) \left(pE^2 - m^2 + \frac{E^2(1 - pf(r))}{pf(r)}\right)^{3/2} \\ &= \int_V dr d\theta d\phi \frac{r^2 \sin(\theta)}{\sqrt{pf(r)}} \left(\frac{4\pi}{3}\right) \left(\frac{E^2(1 - pf(r) + p^2 f(r))}{pf(r)} - m^2\right)^{3/2} \quad (41)\end{aligned}$$

When $p = 1$, it can be readily seen that (41) matches with (35). Also, note that for any $p \in (0, 1]$, (41) diverges when $r = 2GM$ is achieved.

3.3 de Sitter (dS) spacetime

The dS metric as described in locally inertial coordinates tied to a particle moving along a geodesic is given by [9]:

$$ds^2 = -\cos^2(Hr)dt^2 + dr^2 + \frac{\sin^2(Hr)}{H^2}d\Omega^2 \quad (42)$$

with $H = \frac{\pi}{2}\sqrt{\Lambda}$ with Λ being the cosmological constant. The curve $r = 0$ is the worldline of the particle (or, in our case, the centroid of the box).⁴ It can be easily obtained that:

$$\Gamma(E) = \int_V dr d\Omega \left(\frac{4\pi}{3}\right) \left(\frac{E^2}{\cos^2(Hr)} - m^2\right)^{\frac{3}{2}} \frac{\sin^2(Hr)}{H^2} \quad (43)$$

We assume the box to be a spherical box of radius R . So, then after performing angular integrals in (43), we have:

$$\begin{aligned}\Gamma(E) &= \frac{16\pi^2}{3} \int_0^R dr \frac{\sin^2(Hr)}{H^2} \left(\frac{E^2}{\cos^2(Hr)} - m^2\right)^{\frac{3}{2}} \\ &= \frac{16\pi^2}{3H^3} \int_0^\Psi dz \sin^2 z \left(\frac{E^2}{\cos^2 z} - m^2\right)^{3/2} \quad (44)\end{aligned}$$

where in the second line we performed change of variables by defining $z = Hr$ and have defined $\Psi = HR$.

The integral in (44) cannot be exactly solved for general R . So, we will check two cases depending on relative value of R w.r.t. cosmological horizon

⁴Obviously, $\Lambda = \frac{4H^2}{\pi^2}$ is the cosmological constant and $\frac{1}{\sqrt{\Lambda}} = \frac{\pi}{2H}$ is the radius of cosmological horizon

radius $\frac{\pi}{2H}$: firstly, when $R \rightarrow 0^+$ (or $R \ll \frac{\pi}{2H}$) which is equivalently $\Psi \rightarrow 0^+$ and secondly, when $R \rightarrow \frac{\pi}{2H}^-$ which is equivalently $\Psi \rightarrow \frac{\pi}{2}^-$

Case I : $\Psi \rightarrow 0^+$; $R \ll \frac{\pi}{2H}$

By performing small z expansion of the integrand of second line in (44) one can obtain:

$$\begin{aligned}
\Gamma(E) &= \frac{16\pi^2}{3H^3}(E^2 - m^2)^{3/2} \int_0^\Psi dz \left(z^2 - \frac{z^4}{3} + \mathcal{O}(z^6) \right) \\
&\quad \left[1 + \frac{3}{2} \left(\frac{E^2}{E^2 - m^2} \right) z^2 + \mathcal{O}(z^4) \right] \\
&= \frac{16\pi^2}{3H^3}(E^2 - m^2)^{3/2} \int_0^\Psi dz \left[z^2 + \left\{ \frac{3}{2} \left(\frac{E^2}{E^2 - m^2} \right) - \frac{1}{3} \right\} z^4 + \mathcal{O}(z^6) \right] \\
&= \frac{16\pi^2}{3H^3}(E^2 - m^2)^{3/2} \left[\frac{\Psi^3}{3} + \left\{ \frac{3}{2} \left(\frac{E^2}{E^2 - m^2} \right) - \frac{1}{3} \right\} \frac{\Psi^5}{5} + \mathcal{O}(\Psi^7) \right] \quad (45)
\end{aligned}$$

Now putting $\psi = HR$ in last expression in (45), we obtain:

$$\begin{aligned}
\Gamma(E) &= \frac{16\pi^2}{9}(E^2 - m^2)^{3/2} \left[R^3 + \left\{ \frac{9}{2} \left(\frac{E^2}{E^2 - m^2} \right) - 1 \right\} \frac{H^2 R^5}{5} + \right. \\
&\quad \left. \frac{1}{H^3} \mathcal{O}(H^7 R^7) \right] \quad (46)
\end{aligned}$$

Case II: $\Psi \rightarrow \frac{\pi}{2}^-$; $R \rightarrow \left(\frac{\pi}{2H} \right)^-$

It is obvious that the integral (44) diverges for $\Psi \rightarrow \frac{\pi}{2}$ and unlike the case of static black hole, the divergence is solely from the infinite redshift of energy at the cosmological horizon. The singular part of the integral which diverges at the cosmological horizon can be easily obtained by writing $\sin z = \cos(\frac{\pi}{2} - z)$ and $\cos z = \sin(\frac{\pi}{2} - z)$ and then performing change of variables from z to $x = \frac{\pi}{2} - z$ and then performing a small parameter expansion in x followed by integrating the terms that will result in singular behaviour as $x \rightarrow 0$. Hence, one can obtain the singular behaviour of $\Gamma(E)$ as $R \rightarrow \frac{\pi}{2H}$ as:

$$\Gamma(E)_{\text{sing}} = \frac{16\pi^2}{3H^3} \left\{ \frac{E^3}{\left(\frac{\pi}{2} - RH \right)^2} \right\} - \left\{ \frac{1}{2} + \frac{3}{2} \left(\frac{m^2}{E^2} \right) \right\} E^3 \ln \left(\frac{\pi}{2} - RH \right) \quad (47)$$

We note that the spatial volume element doesn't contribute to divergence of $\Gamma(E)$ at cosmological horizon for de Sitter and at Rindler horizon for accelerated particle in flat spacetime and hence this divergence is somewhat different from the divergence of $\Gamma(E)$ at event horizon for Schwarzschild black hole where both the spatial volume element and the λ factor ($-g^{00}$ term) with energy (E) diverges at event horizon.

4 Leading curvature corrections in arbitrary curved spacetimes

4.1 One particle phase space volume

In the earlier sections, we performed analysis in some special spacetimes. Now, we do a perturbative analysis on any general spacetime for particles confined in a box small enough compared to curvature scale of the spacetime. Hence, around the worldline of centroid of the box, we can perform a perturbative analysis using Fermi normal coordinates. To this end, consider a perturbed metric over Minkowski where the metric tensor is given by:

$$g_{ab} = \begin{bmatrix} -1 - 2\epsilon q & \epsilon l_1 & \epsilon l_2 & \epsilon l_3 \\ \epsilon l_1 & 1 + \epsilon h_{11} & \epsilon h_{12} & \epsilon h_{13} \\ \epsilon l_2 & \epsilon h_{12} & 1 + \epsilon h_{22} & \epsilon h_{23} \\ \epsilon l_3 & \epsilon h_{13} & \epsilon h_{23} & 1 + \epsilon h_{33} \end{bmatrix} \quad (48)$$

From (94), we obtain:

$$h_{\alpha\beta} = g_{\alpha\beta} + \mathcal{O}(\epsilon^2) \quad (49)$$

Hence,

$$\begin{aligned} \sqrt{\text{Det}([h])} &= [1 + \epsilon(h_{11} + h_{22} + h_{33}) + \mathcal{O}(\epsilon^2)]^{1/2} \\ &= 1 + \epsilon \left(\frac{h_{11} + h_{22} + h_{33}}{2} \right) + \mathcal{O}(\epsilon^2) \end{aligned} \quad (50)$$

Also, we have:

$$-g^{00} = K = 1 - 2\epsilon q + \mathcal{O}(\epsilon^2) \quad (51)$$

Now, note that as $\gamma_\nu = n^\mu \omega_{\mu\nu}$ and as $n^\mu = g^{0\mu} = \epsilon l^\mu$, so, $\gamma_\nu^2 \sim \mathcal{O}(\epsilon^2)$. Thus volume occupied by constant energy hypersurface in 3-momentum space is

given using (87):

$$\int d^3\mathbf{p} \Theta(E + \xi^i p_i) = \left(\frac{4\pi}{3}\right) \left[(1 - 2\epsilon q) E^2 - m^2 + \mathcal{O}(\epsilon^2) \right]^{3/2} \left[1 + \frac{\epsilon}{2} (h_{11} + h_{22} + h_{33}) + \mathcal{O}(\epsilon^2) \right] \quad (52)$$

This can be simplified to finally obtain:

$$\int d^3\mathbf{p} \Theta(E + \xi^i p_i) = \left(\frac{4\pi}{3}\right) (E^2 - m^2)^{3/2} \left[1 + \epsilon \left\{ \left(\frac{h_{11} + h_{22} + h_{33}}{2} \right) - 3\epsilon q \left(\frac{E^2}{E^2 - m^2} \right) \right\} + \mathcal{O}(\epsilon^2) \right] \quad (53)$$

We can now proceed to evaluate the curvature corrections by inserting the explicit forms for q and $h_{\mu\nu}$ in Fermi normal coordinates (FNC)⁵. The FNC constitute a coordinate system on spacetime which is constructed around a timelike curve in spacetime such that the coordinate chart is flat (metric is Minkowski and Christoffel symbols vanish) on the curve. The coordinate chart is given by $\{\tau, y^1, y^2, y^3\}$ where τ is proper time along the curve. The metric components in FNC is given in [10] to be:

$$g_{00} = -(1 + a_\mu y^\mu)^2 + R_{0\mu 0\nu} y^\mu y^\nu + \mathcal{O}(y^3) \quad (54)$$

$$g_{0\mu} = -\frac{2}{3} R_{0\rho\mu\nu} y^\rho y^\nu + \mathcal{O}(y^3) \quad (55)$$

$$g_{\mu\nu} = \delta_{\mu\nu} - \frac{1}{3} R_{\mu\rho\nu\sigma} y^\rho y^\sigma + \mathcal{O}(y^3) \quad (56)$$

where $a^i = u^k \nabla_k u^i$ is the acceleration of the centroid of the box.

Now, we shall assume that our box is a spherical box.⁶ This will require a coordinate change as FNC as described in (54) are adapted to a locally Cartesian-like coordinate chart. We shall not write FNC in locally spherical coordinate chart, rather we shall do change of variables in the integration.

Now, to proceed, we shall first consider curvature as the small parameter (note that $\mathcal{O}(y^3)$ terms in FNC also contain higher order curvature terms,

⁵Our convention will be as follows: Repeated indices when one is in upstairs and other is in downstairs will be automatically summed unless otherwise stated. But for summing over repeated indices placed at the same level, an explicit summation symbol will be used.

⁶This is motivated by the fact that we shall be using our calculations mostly for spherically symmetric spacetimes and a spherical box is a natural choice for it

so, we can equivalently treat the box radius as well as curvature as small parameters) and then invert the metric. Then, we will consider acceleration to be small and keep only terms till quadratic order in acceleration. Note that dimensionally, acceleration is $[L]^{-1}$, whereas curvature is $[L]^{-2}$ and hence we keep linear order terms in curvature and quadratic order in acceleration. So, essentially, the dimensionless small parameters in our perturbative calculations are essentially R^2 .(curvature) and $R^2|\mathbf{a}|^2$ and we consider both to be of same order. So, we obtain the following till first order in curvature and quadratic order in acceleration.

$$K = -g^{00} = 1 - 2a_\mu y^\mu + (R_{0\mu 0\nu} + 3a_\mu a_\nu)y^\mu y^\nu \quad (57)$$

$$h_{\mu\nu} = g_{\mu\nu} = \delta_{\mu\nu} - \frac{1}{3}R_{\mu\rho\nu\sigma}y^\rho y^\sigma \quad (58)$$

where $h_{\alpha\beta}$ is the metric on spatial foliations as appearing in (6). Now, using (50), we say that:

$$\sqrt{\text{Det}([h])} = 1 - \frac{1}{6} \sum_{\mu=1}^3 R_{\mu\alpha\mu\beta} y^\alpha y^\beta \quad (59)$$

Also, spatio-temporal cross terms of the metric appear only in quadratic order in (6) and hence quadratic in curvature. So, we ignore them in our computation.

So, putting all of them in (6) retaining only the order of terms we are interested in (first order in curvature and second order in acceleration) and also ignoring all acceleration-curvature cross terms appearing in $\mathcal{O}(y^3)$, the phase volume is:

$$\begin{aligned} \Gamma(E) &= \int_V d^3\mathbf{x} \left(\frac{4\pi}{3}\right) \left(1 - \frac{1}{6} \sum_{\mu=1}^3 R_{\mu\alpha\mu\beta} y^\alpha y^\beta\right) \left[(1 - 2a_\mu y^\mu + (R_{0\mu 0\nu} + 3a_\mu a_\nu)y^\mu y^\nu) E^2 - m^2 \right]^{3/2} \\ &= \int_V d^3\mathbf{x} \left(\frac{4\pi}{3}\right) (E^2 - m^2)^{3/2} \left[1 - \frac{1}{6} \sum_{\mu=1}^3 R_{\mu\alpha\mu\beta} y^\alpha y^\beta + \left(\frac{E^2}{E^2 - m^2}\right) \left\{ -3a_\mu y^\mu + \frac{3}{2}(R_{0\mu 0\nu} + 3a_\mu a_\nu)y^\mu y^\nu \right\} + \frac{3}{2} \left(\frac{E^2}{E^2 - m^2}\right)^2 a_\mu a_\nu y^\mu y^\nu \right] \end{aligned} \quad (60)$$

where the spatial integrations are to be performed inside a spherical box of radius R .

Replacing Cartesian $\{y^1, y^2, y^3\}$ with the usual spherical polar $\{r, \theta, \phi\}$ by the usual coordinate transformation, one can write: ⁷

$$\begin{aligned}
& \int_V d^3\mathbf{y} \left(-\frac{1}{6} \sum_{\mu=1}^3 R_{\mu\alpha\mu\beta} y^\alpha y^\beta \right) \\
&= -\frac{1}{6} \sum_{\mu=1}^3 \int_V dr d\Omega r^2 R_{\mu\alpha\mu\beta} y^\alpha y^\beta \\
&= \frac{4\pi R^5}{45} G^0_0
\end{aligned} \tag{61}$$

where V denotes a spherical box at centered at origin with radius R , and $d\Omega = \sin\theta d\theta d\phi$. Note that in evaluating the above integral (61), we have used the fact that:

$$\int_V dr r^2 d\Omega_2 x^2 = \int_V dr r^2 d\Omega_2 y^2 = \int_V dr r^2 d\Omega_2 z^2 = \left(\frac{4\pi}{3} \right) \frac{R^5}{5} \tag{62}$$

and ⁸

$$\sum_{\mu\nu} R_{\mu\nu\mu\nu} = -2G^0_0 \tag{63}$$

Further we note that:

$$\int_V d^3\mathbf{y} y^\mu = 0 \tag{64}$$

and

$$\int_V d^3\mathbf{y} R_{0\mu 0\nu} y^\mu y^\nu = \left(\frac{4\pi}{3} \right) \frac{R^5}{5} \sum_{\mu} R_{0\mu 0\mu} = \left(\frac{4\pi}{3} \right) \frac{R^5}{5} R^0_0 \tag{65}$$

and hence similarly,

$$\int_V d^3\mathbf{y} a_\mu a_\nu y^\mu y^\nu = \left(\frac{4\pi}{3} \right) \frac{R^5}{5} \sum_{\mu} (a_\mu)^2 = \left(\frac{4\pi}{3} \right) \frac{R^5}{5} |\mathbf{a}|^2 \tag{66}$$

⁷And also note the facts that $\int dr r^2 d\Omega_2 x = \int dr r^2 d\Omega_2 y = \int dr r^2 d\Omega_2 z = 0$ and also $\int dr r^2 d\Omega_2 xy = \int dr r^2 d\Omega_2 yz = \int dr r^2 d\Omega_2 zx = 0$

⁸To prove the following (63), first note that $R_{\mu\nu\mu\nu} = R^{\mu\nu}{}_{\mu\nu}$ (no summation over any index) as the metric is Minkowski along the curve and as these are Riemann tensors defined on the curve, so the indices can be freely raised or lowered. Then, $\sum_{\mu\nu} R_{\mu\nu\mu\nu} = R^{\mu\nu}{}_{\mu\nu}$. Finally note that $R^{12}{}_{12} + R^{23}{}_{23} + R^{13}{}_{13} = -G^0_0$

Putting all of them in (60), we obtain:

$$\Gamma(E) = \frac{4\pi}{3}(E^2 - m^2)^{3/2} \left[\frac{4\pi}{3}R^3 + \frac{4\pi R^5}{45}G^0_0 + \left(\frac{E^2}{E^2 - m^2} \right) \left(\frac{2\pi R^5}{5}R^0_0 + \frac{6\pi R^5}{5}|\mathbf{a}|^2 \right) + \left(\frac{E^2}{E^2 - m^2} \right)^2 \left(\frac{2\pi R^5}{5}|\mathbf{a}|^2 \right) \right] \quad (67)$$

Now, writing in terms of volume and area of the box and clubbing curvature dependent terms and acceleration dependent terms separately together, we obtain:

$$\Gamma(E) = \frac{4\pi}{3}(E^2 - m^2)^{3/2}.V \left[1 + \frac{A}{60\pi}G^0_0 + \left(\frac{E^2}{E^2 - m^2} \right) \frac{3A}{40\pi}R^0_0 + \left\{ \left(\frac{E^2}{E^2 - m^2} \right) \frac{9}{40\pi} + \left(\frac{E^2}{E^2 - m^2} \right)^2 \frac{3}{40\pi} \right\} A|\mathbf{a}|^2 \right] \quad (68)$$

We again emphasize that (68) is obtained by taking first order in curvature and quadratic order in acceleration terms. Also, we emphasize the point that Eq. (68) is only valid when V and A are computed for a spherical box and it is not valid for boxes of arbitrary geometry.

We can perform two important consistency checks:

- By putting dS values of $R^0_0 = 3H^2$ and $G^0_0 = -3H^2$ and putting $a^i = 0$ in (67), one gets back the small R expression (46) for $\Gamma(E)$ in dS.
- By putting mass and curvature terms to zero, we get back Rindler result (32) when box radius is very small compared to Rindler horizon.

4.2 Microcanonical entropy and temperature

Although there isn't much point of doing statistical mechanics with single particle systems, nevertheless in this subsection we shall find expressions for entropy and temperature for one particle systems as they might give insights to statistical behaviour of many particle systems in curved backgrounds.

The number of accessible microstates for a given value of energy E can be

obtained from (68) as:

$$\begin{aligned}
\Omega(E) &= \frac{d\Gamma(E)}{dE} \\
&= 4\pi(E^2 - m^2)^{\frac{1}{2}} E V \left[1 + \frac{A}{60\pi} G^0_0 + \frac{A}{40\pi} \left(\frac{3E^2 - 2m^2}{E^2 - m^2} \right) R^0_0 + \right. \\
&\quad \left. \frac{A}{40\pi} \left(\frac{12E^4 - 19E^2m^2 + 6m^4}{(E^2 - m^2)^2} \right) |\mathbf{a}|^2 \right] \quad (69)
\end{aligned}$$

Now, taking its logarithm and using the fact that $R^2|\mathbf{a}|^2 \sim R^2(\text{curvature}) \ll 1$, the microcanonical entropy is given by:

$$\begin{aligned}
S(E) &= \log \Omega(E) \\
&= \log\{4\pi E(E^2 - m^2)^{1/2}\} + \log V + A \left[\frac{1}{60\pi} G^0_0 + \right. \\
&\quad \left. \frac{1}{40\pi} \left(\frac{3E^2 - 2m^2}{E^2 - m^2} \right) R^0_0 + \frac{1}{40\pi} \left(\frac{12E^4 - 19E^2m^2 + 6m^4}{(E^2 - m^2)^2} \right) |\mathbf{a}|^2 \right] \quad (70)
\end{aligned}$$

Now, the inverse temperature β is given by:

$$\begin{aligned}
\beta &= \frac{\partial S(E)}{\partial E} \\
&\approx \frac{1}{E} + \frac{E}{E^2 - m^2} - A \left[\frac{1}{20\pi} \left(\frac{Em^2}{(E^2 - m^2)^2} \right) R^0_0 + \right. \\
&\quad \left. \frac{1}{20\pi} \left(\frac{Em^2(5E^2 - 7m^2)}{(E^2 - m^2)^3} \right) |\mathbf{a}|^2 \right] \quad (71)
\end{aligned}$$

From Eq. (70) and (71) it is obvious that leading order curvature correction terms in entropy and inverse temperature are proportional to surface area of the system.

4.3 Multi-particle systems

In Section-(2.2), we discussed about phase volume occupied by a restricted class of multiparticle systems, essentially particles with $m = 0$ and background spacetimes with $g_{0\mu} = 0$. In this restricted case, we had obtained (26) and hence subsequently (27). Now, we can employ them and use the one particle computations in Section-(4.2) to derive microcanonical entropy and temperature expressions for these N -particle systems.

From (27), we have:

$$\begin{aligned}\Omega(E) &= \frac{1}{(N-1)!} \Gamma_1(E)^{N-1} \frac{d\Gamma_1(E)}{dE} \\ &= \frac{1}{(N-1)!} \Gamma_1(E)^{N-1} \Omega_1(E)\end{aligned}\quad (72)$$

where $\Omega_1(E)$ is single particle constant energy phase volume. Hence, entropy will be:

$$\begin{aligned}S(E) &= \log \Omega(E) \\ &= -\log((N-1)!) + (N-1) \log(\Gamma_1(E)) + \log \Omega_1(E)\end{aligned}\quad (73)$$

Now, employing Stirling's approximation and taking thermodynamic limit (N very large), we have:

$$S(E) = N - N \log N + N \log(\Gamma_1(E)) + \log(\Omega_1(E))\quad (74)$$

Now, we can employ this to evaluate entropy of massless limits of many systems of interest. However, we focus our attention to an accelerated system in curved spacetime with the assumption $R^2(\text{curvature}) \sim R^2|\mathbf{a}|^2 \ll 1$.

Taking massless cases of Eq. (70), (68) and plugging them in Eq. (74), we get:

$$\begin{aligned}S(E) = N - N \log N + N \left[\log \left(\frac{4\pi}{3} E^3 \right) + \log V + A \left(\frac{1}{60\pi} G^0_0 + \right. \right. \\ \left. \left. \frac{3}{40\pi} R^0_0 + \frac{3}{10\pi} |\mathbf{a}|^2 \right) \right] + \log\{4\pi E^2\} + \log V + A \left[\frac{1}{60\pi} G^0_0 + \right. \\ \left. \frac{3}{40\pi} R^0_0 + \frac{3}{10\pi} |\mathbf{a}|^2 \right]\end{aligned}\quad (75)$$

Now, clubbing all N dependent (and E, V, A , curvature and acceleration independent) constants as C_N and taking thermodynamic limit (large N limit), we can further simplify the expression (75) to obtain:

$$S(E) \approx C_N + 3N \log E + N \log V + N A \left[\frac{1}{60\pi} G^0_0 + \left(\frac{3}{40\pi} \right) (R^0_0 + 4|\mathbf{a}|^2) \right]\quad (76)$$

Furthermore, we note that the microcanonical inverse temperature β is given by:

$$\beta = \frac{3N}{E}\quad (77)$$

So, for massless particles in the restricted class of spacetimes considered, entropy has leading order curvature correction terms appearing as proportional to area in Eq. (76), but the inverse temperature has no first order curvature (or quadratic order acceleration) correction terms as is evident from Eq. (77)⁹. In fact, the expression in (77) is precisely the result we obtain by applying the general equipartition theorem as described in [11] to an ultra-relativistic gas in flat spacetime.

5 Discussion

To summarize, in this work we have explicitly computed correction terms to microcanonical phase volume, entropy and temperature of inertial and non-inertial systems in curved spacetimes with a timelike Killing field (with the field being timelike in some “nice” patch of the manifold, so that we have a notion of a “conserved energy”). We have further observed that these corrections to thermodynamic quantities lead to quite non-trivial consequences such as divergent behavior near null surfaces as well as area dependent curvature and acceleration terms. We highlight some of the major qualitative inferences of this work in the following paragraphs.

The divergent behavior of $\Gamma(E)$ (and hence other thermodynamic quantities, including entropy) at event horizon of Schwarzschild black hole (see (41)) is due to infinite redshift of energy (i.e. divergence of $(-g^{00})$ term) as well as due to divergence of spatial volume element. However, the divergence of $\Gamma(E)$ at cosmological horizon of de Sitter (see (47)) and Rindler horizon for uniform acceleration in flat spacetime (see (33)) is only due to infinite redshift of energy.

Apart from that, it is also worth pointing out that although quite appealing, the area dependent curvature correction terms appearing in entropy expressions of (70) and (76) have no direct relationship with the Bekenstein-Hawking entropy of black holes. This is because we are not considering self-gravitation here and hence the system does not describe a gravitational collapse or black hole. Nevertheless, we expect that the above analysis would still hold provided one uses, for G_{ab} and R_{ab} , their expressions as obtained from Einstein equations with the system itself as a source. This work is under progress.

⁹Even, one particle β also has no first order curvature correction terms if particles are massless even in a completely arbitrary spacetime as is evident by putting $m = 0$ in (71)

We also note that entropy of massless N -particle systems for accelerated motion in curved spacetime¹⁰ has area dependent first order correction terms. But, microcanonical inverse temperature has no correction term at this order and just reproduces the standard equi-partition result for ultra-relativistic ideal gas in flat background.

As already emphasized, the most natural and relevant extension of this work is to include self-gravitation of the system itself into account (through the metric), and compute the various thermodynamic quantities. This would be the first step towards understanding the purely (quantum) statistical mechanical contribution to thermodynamic properties of black holes.

A Derivation of the Formula for $\Gamma(E)$

Consider a metric g_{ij} which do not depend on time coordinate x^0 . Hence, this spacetime has a Killing field $\xi^i = (1, 0, 0, 0)$. So,

$$H = -\xi^i p_i = -p_0 \quad (78)$$

and it is conserved if the particle follows a geodesic trajectory. Now,

$$\begin{aligned} p_i p^i &= -m^2 \\ \implies g^{00}(p_0)^2 + 2g^{0\mu} p_\mu p_0 + g^{\mu\nu} p_\mu p_\nu &= -m^2 \\ \implies K(p_0)^2 - (2n^\mu p_\mu)p_0 - (g^{\mu\nu} p_\mu p_\nu + m^2) &= 0 \end{aligned} \quad (79)$$

where we have put $g^{00} = -K$ and $g^{0\mu} = n^\mu$.

Now, the quadratic equation (79) has solutions:

$$p_0^\pm = \frac{n^\mu p_\mu \pm \sqrt{(n^\mu p_\mu)^2 + K(g^{\mu\nu} p_\mu p_\nu + m^2)}}{K} \quad (80)$$

Since we need $H = -p_0 > 0$, hence we take $p_0 < 0$ solution out of (80) and assuming that $K > 0$ and $\mathbf{p} \cdot \mathbf{p} = g^{\mu\nu} p_\mu p_\nu > 0$ we take the p_0^- solution. Thus,

$$H = \frac{-n^\mu p_\mu + \sqrt{(n^\mu p_\mu)^2 + K(g^{\mu\nu} p_\mu p_\nu + m^2)}}{K} \quad (81)$$

To evaluate the integral (2), we first evaluate $\int dp_\alpha \Theta(E + \xi^i p_i)$ which essentially corresponds to the volume occupied by the surface $E = H$ in the

¹⁰for the restricted class of metrics where $g^{0\mu} = 0$

3-momentum space.

The surface $H = E$ in the 3-momentum space is given by:

$$\begin{aligned} (KE + n^\mu p_\mu)^2 &= (n^\mu p_\mu)^2 + K(g^{\mu\nu} p_\mu p_\nu + m^2) \\ \implies g^{\mu\nu} p_\mu p_\nu - 2En^\mu p_\mu &= KE^2 - m^2 \end{aligned} \quad (82)$$

Now for the following part of the computation of volume occupied by this surface, objects with square brackets are matrices on the 3-momentum space (assumed as a vector space) and their indices are not tensor indices but matrix indices and always placed downstairs. However Einstein summation convention will still be in force in the sense that repeated indices will be summed but placement of indices won't matter. Now, define:

$$p_\mu = [\omega]_{\mu\nu} \tilde{p}_\nu \quad \text{s.t.} \quad g^{\mu\nu} p_\mu p_\nu = [\mathbb{1}]_{\mu\nu} \tilde{p}_\mu \tilde{p}_\nu \quad (83)$$

So, the surface (82) can be written in p coordinate system in momentum space as:

$$\sum_\mu (\tilde{p}_\mu)^2 - 2E(n^\mu [\omega]_{\mu\nu}) \tilde{p}_\nu = KE^2 - m^2 \quad (84)$$

$$\implies \sum_\mu (\tilde{p}_\mu^2 - \gamma_\mu \tilde{p}_\mu) = \lambda \quad (85)$$

where $\gamma_\nu = 2E(n^\mu [\omega]_{\mu\nu})$ and $\lambda = KE^2 - m^2$.

Now, completing squares in (85), we arrive at the equation of the surface to be:

$$\sum_\mu \left(\tilde{p}_\mu - \frac{\gamma_\mu}{2} \right)^2 = \lambda + \sum_\mu \frac{\gamma_\mu^2}{4} \quad (86)$$

Now, clearly the volume occupied by surface (86) in 3-momentum space (which essentially a sphere in \tilde{p} coordinate system) and hence the momentum integral that we set forth to evaluate first is given by:

$$\int d^3\mathbf{p} \Theta(E + \xi^i p_i) = \frac{4}{3}\pi \left[\lambda + \sum_\mu \frac{\gamma_\mu^2}{4} \right]^{3/2} \text{Det}([\omega]) \quad (87)$$

where the factor of $\text{Det}([\omega])$ comes because of change of variables in integration.

Now, the change of basis condition (83) can be written in matrix notation as:

$$[p] = [\omega][\tilde{p}] \quad \text{s.t.} \quad [p]^T [g][p] = [\tilde{p}]^T [\tilde{g}] \quad (88)$$

where $[p]$ is column vector corresponding to the 3-momentum vector p^α and $[g]$ is the matrix corresponding to $g^{\mu\nu}$. So, (88) implies that:

$$[\tilde{p}]^T [\omega]^T [g] [\omega] [\tilde{p}] = [\tilde{p}]^T [\tilde{p}] \quad (89)$$

$$\implies [\omega]^T [g] [\omega] = [1] \quad (90)$$

So, we have:

$$[\omega] = [D][\chi] \quad (91)$$

where $[D]$ is the matrix with eigenvectors of $[g]$ as columns ¹¹ and

$$[\chi] = \text{Diag}\left(\frac{1}{\sqrt{\lambda_1}}, \frac{1}{\sqrt{\lambda_2}}, \frac{1}{\sqrt{\lambda_3}}\right) \quad (92)$$

where λ_i 's are eigenvalues of $[g]$.

In [6], it has been shown that the spatial line element on constant time foliations is given by:

$$dl^2 = h_{\alpha\beta} dx^\alpha dx^\beta \quad (93)$$

where

$$h_{\alpha\beta} = g_{\alpha\beta} - \frac{g_{0\alpha}g_{0\beta}}{g_{00}} \quad (94)$$

and also

$$g^{\mu\rho} h_{\rho\nu} = \delta_\nu^\mu \implies [g]^{-1} = [h] \quad (95)$$

Thus, from (91) we can write:

$$\text{Det}([\omega]) = \text{Det}([D]) \frac{1}{\sqrt{\lambda_1 \lambda_2 \lambda_3}} \quad (96)$$

Now,

$$[D]^T [g] [D] = \text{Diag}(\lambda_1, \lambda_2, \lambda_3) \quad (97)$$

$$\implies \text{Det}([D]) = \frac{\sqrt{\lambda_1 \lambda_2 \lambda_3}}{\sqrt{\text{Det}([g])}} \quad (98)$$

Putting $\text{Det}([D])$ from (98) in (96) and finally using (95), we have:

$$\text{Det}([\omega]) = \frac{1}{\sqrt{\text{Det}[g]}} = \sqrt{\text{Det}([h])} \quad (99)$$

¹¹ $[g]$ being symmetric, it is orthogonally diagonalizable and hence $[D]^T = [D]^{-1}$

Hence, combining (99) with (87), we can write:

$$\Gamma(E) = \int_V d^3\mathbf{x} \sqrt{\text{Det}([h])} \left(\frac{4\pi}{3}\right) \left[\lambda + \sum_{\mu} \frac{\gamma_{\mu}^2}{4}\right]^{3/2} \quad (100)$$

where V is the spatial extent of the box in which the particle resides. This is precisely the expression (6).

References

- [1] T. Padmanabhan, *Statistical mechanics of gravitating systems*, Physics Reports, **188**, No. 5(1990), pp. 285–362.
- [2] D. Lynden-Bell, Roger Wood, Astronomer Royal, *The Gravo-Thermal Catastrophe in Isothermal Spheres and the Onset of Red-Giant Structure for Stellar Systems*, Monthly Notices of the Royal Astronomical Society, Vol. 138, Issue 4, Feb 1968, pp. 495–525.
- [3] D. Alba, H.W. Crater, L. Lusanna, *On the Relativistic Micro-Canonical Ensemble and Relativistic Kinetic Theory for N Relativistic Particles in Inertial and Non-Inertial Rest Frames*, arXiv:1202.4667v2.
- [4] Pierre-Henri Chavanis, *Statistical mechanics of self-gravitating systems in general relativity: I. The quantum Fermi gas*, Eur. Phys. J. Plus (2020) **135**, 290, arXiv:1908.10806.
- [5] S.W. Hawking, *Black holes and thermodynamics*, Physical Review D, Vol. 13 No. 2, Jan 1976, pp. 191-197.
- [6] L.D. Landau & E.M. Lifshitz, *The Classical Theory of Fields*, 3rd Ed., Sec. 84.
- [7] T.Padmanabhan, *Phase volume occupied by a test particle around an incipient black hole*, Phys. Letts. A. (1989) **136**, pp. 203-205.
- [8] S. Kolekar & T. Padmanabhan, *Ideal Gas in a strong Gravitational field: Area dependence of Entropy*, Phys. Rev. D **83**, 064034 (2011), arXiv:1012.5421v2; S. Bhattacharya, S. Chakraborty & T. Padmanabhan, *Entropy of a box of gas in an external gravitational field — revisited*, Phys. Rev. D **96**, 084030 (2017), arXiv:1702.08723v2.
- [9] C. Chicone & B. Mashhoon, *Explicit Fermi Coordinates and Tidal Dynamics in de Sitter and Godel Spacetimes*, arXiv:gr-qc/0511129v2.

- [10] D. Kothawala, *Box of Ideal Gas in Free Fall*, arXiv:gr-qc/1108.6301v2.
- [11] R.C. Tolman, *A General Theory of Energy Partition with Applications to Quantum Theory*, Physical Review (1918). 11 (4), pp. 261–275.