AN EQUIDISTRIBUTION RESULT FOR DIFFERENCES ASSOCIATED WITH SQUARE PYRAMIDAL NUMBERS

ANJI DONG, KATERINA SAETTONE, KENDRA SONG AND ALEXANDRU ZAHARESCU

ABSTRACT. We provide an asymptotic formula for the average value of the sequence <u>A351830</u>: $a_n = |P_n - y_n^2|$ for $1 \le n \le x$, where P_n is the *n*-th square pyramidal number and y_n^2 is the closest square to P_n . Moreover, we supply asymptotic formulas for the *k*-th moment of the same sequence, for any fixed natural number k.

1. INTRODUCTION

In the 1500s, Sir Walter Raleigh proposed the problem of counting the cannonballs in a square pyramid during a sea voyage, which is now known as the Cannonball problem. Later, the problem was officially proposed by Lucas [10], which studies what integers are both a square and a square pyramidal number. It was eventually proved by Watson [11] that the only solutions are 0, 1, and 4900.

There are many integer sequences associated with the Cannonball problem, for example, see <u>A001032</u>, <u>A351830</u>, <u>A350886</u>, <u>A350887</u>, and <u>A350888</u> in the On-Line Encyclopedia of Integer Sequences (OEIS) [9]. In this paper, we establish additional properties of the sequence <u>A351830</u>. This is given by

(1.1)
$$a_n = |P_n - y_n^2|,$$

where P_n is the *n*-th square pyramidal number defined as

(1.2)
$$P_n = \sum_{i=1}^n i^2 = \frac{2n^3 + 3n^2 + n}{6},$$

and y_n^2 is the closest square to P_n . In fact, Paolo Xausa [12] computed the first ten thousand elements of the sequence.

Note that the only values of n such that $a_n = 0$ are 0, 1, and 24, which correspond to the solutions to the Cannonball problem. Moreover, noted by Conway and Sloane [3, 4], the solution corresponding to n = 24 allows for the Lorentzian construction of the Leech lattice. Further applications and generalizations of the Cannonball problem are discussed by Laub [7], Beeckmans [1], and Bennett [2].

For real $x \ge 1$, let A(x) represent the average value of (a_n) for $1 \le n \le x$. More precisely,

(1.3)
$$A(x) := \frac{1}{x} \sum_{1 \le n \le x} a_n.$$

Hence, since a_n represents the distance between the *n*-th square pyramidal number and the closest square, A(x) represents the average distance from a square pyramidal number to its closest square up to the $\lfloor x \rfloor$ -th square pyramidal number. Our primary goal is to provide an asymptotic formula for A(x).

²⁰²⁰ Mathematics Subject Classification. Primary: 11L07. Secondary: 11B99, 11K38, 11K06.

Key words and phrases. Cannonball problem, exponential sums, discrepancy, equidistribution.

Theorem 1.1. For real $x \ge 1$, the function A(x), given by (1.3), satisfies the asymptotic formula

(1.4)
$$A(x) = \frac{1}{5\sqrt{3}}x^{3/2} + O(x^{4/3}),$$

where the constant implied by the big O symbol is absolute.

Next, we generalize Theorem 1.1 by considering higher moments of the sequence (a_n) . For each $k \in \mathbb{N}$, let $M_k(x)$ represent the k-th moment of $(a_n)_{n \leq x}$, that is,

$$(1.5) M_k(x) := \sum_{1 \le n \le x} a_n^k.$$

By Theorem 1.1,

$$M_1(x) = xA(x) = \frac{1}{5\sqrt{3}}x^{5/2} + O(x^{7/3}).$$

More generally, we have the following result.

Theorem 1.2. For any positive integer k,

(1.6)
$$M_k(x) = \frac{x^{\frac{3}{2}k+1}}{3^{k/2}(\frac{3}{2}k+1)(k+1)} + O_k\left(x^{\frac{3}{2}k+\frac{5}{6}}\right)$$

where M_k is defined in (1.5), and the constant implied by the O_k symbol depends at most on k.

Structure of the Paper. The paper is organized as follows. We begin by introducing some standard notation in Section 2. In Section 3, we provide some auxiliary results needed in the proofs of the main theorems. Lastly, in Section 4, we complete the proofs of Theorems 1.1 and 1.2.

2. GENERAL NOTATION

We employ some standard notation used throughout the article.

- The expressions f(X) = O(g(X)), f(X) ≪ g(X), and g(X) ≫ f(X) are equivalent to the statement that |f(X)| ≤ C|g(X)| for all sufficiently large X, where C > 0 is an absolute constant. A subscript of the form ≪_α means that the implied constant may depend on the parameter α. Dependence on several parameters is indicated similarly, as in ≪_{α,λ}.
- Given $\alpha \in \mathbb{R}$, the notation $\|\alpha\|$ denotes the smallest distance of α to an integer.
- Given a set S, the notation |S| and #S both stand for the cardinality of S.
- For x ∈ ℝ, the notation [x] denotes the ceiling of x, which is the smallest integer larger than x. The notation [x] denotes the floor of x, which is the largest integer smaller than x.
- For real $x \in \mathbb{R}$, $\{x\}$ denotes the fractional part of x, that is, x |x|.
- The notation e(x) stands for $\exp(2\pi i x)$.

3. AUXILIARY PROPOSITIONS

In this section, we aim to show that (a_n) , defined in (1.1), is equidistributed in $[0, \frac{1}{2}]$. We begin with the following lemma by Kuipers and Niederreiter [6].

Lemma 3.1 ([6, Thm. 2.7, p. 17]). Let a and b be integers with a < b. Assume there exists some $\rho \in \mathbb{R}^+$ such that f is twice differentiable on [a, b] with $f''(x) \ge \rho > 0$ or $f''(x) \le -\rho < 0$ for all $x \in [a, b]$. Then,

(3.1)
$$\left|\sum_{n=a}^{b} e(f(n))\right| \leq \left(\left|f'(b) - f'(a)\right| + 2\right) \left(\frac{4}{\sqrt{\rho}} + 3\right).$$

Lemma 3.2 ([8, Weyl's Criterion, p. 1]). A sequence (a_n) is uniformly distributed if and only if for each integer $k \neq 0$,

$$\lim_{N \to \infty} \sum_{n=1}^{N} e(ka_n) = 0.$$

Proposition 3.3. The sequence $\{\sqrt{P_n}\}$ is uniformly distributed in the unit interval [0, 1].

Proof. Let
$$h(x) = \sqrt{P_x}$$
. It is clear that

$$h'(x) = \frac{2x(x+1) + (2x+1)(x+1) + x(2x+1)}{2\sqrt{6}\sqrt{x(x+1)(2x+1)}},$$

$$h''(x) = \frac{8x + 4(x+1) + 2}{2\sqrt{6}\sqrt{x(x+1)(2x+1)}} - \frac{(2x(x+1) + (2x+1)(x+1) + x(2x+1))^2}{4\sqrt{6}(x(x+1)(2x+1))^{3/2}}$$

Let $a \in \mathbb{Z} \setminus \{0\}$. Since h'' is monotonically decreasing, take $\rho = h''(N)$. Then Lemma 3.1 gives us

$$\lim_{N \to \infty} \frac{1}{N} \left| \sum_{n=1}^{N} e(ah(n)) \right| \le \lim_{N \to \infty} \frac{1}{N} (|ah'(N) - ah'(1)| + 2) \left(\frac{4}{\sqrt{ah''(N)}} + 3 \right) = 0.$$

Upon using Lemma 3.2, we see that $\sqrt{P_n}$ is uniformly distributed. Therefore, the sequence $(\sqrt{P_n})$ is uniformly distributed mod 1, that is, the sequence of the fractional parts, $\{\sqrt{P_n}\}$, is uniformly distributed in the interval [0, 1].

Proposition 3.4. The sequence (a_n) , defined as $|\sqrt{P_n} - y_n|$, is uniformly distributed in the interval [0, 1/2].

Proof. Note that $(\lfloor \sqrt{P_n} \rfloor)^2 \leq P_n \leq (\lceil \sqrt{P_n} \rceil)^2$, and since $\lceil \sqrt{P_n} \rceil - \lfloor \sqrt{P_n} \rfloor \leq 1$, it follows that y_n is either $\lfloor \sqrt{P_n} \rfloor$ or $\lceil \sqrt{P_n} \rceil$. Therefore, $|\sqrt{P_n} - y_n|$ is at most 1/2.

Consider the following set

(3.2)
$$E(x) = \left\{ n \colon 1 \le n \le x, |\sqrt{P_n} - y_n| \ne ||\sqrt{P_n}|| \right\}$$

which we refer to as the exceptional set depending on x. For all $n \in E(x)$, the number n must satisfy one of the two following cases:

Case 1:
$$y_n = \lfloor \sqrt{P_n} \rfloor$$
 and $\|\sqrt{P_n}\| = |\sqrt{P_n} - (y_n + 1)| < |\sqrt{P_n} - y_n|$.
By definition of y_n in (1.1), we have

(3.3)
$$|P_n - y_n^2| < |P_n - (y_n + 1)^2|.$$

Since $y_n = \lfloor \sqrt{P_n} \rfloor$, (3.3) is equivalent to

$$P_n - y_n^2 < (y_n + 1)^2 - P_n$$

so $\sqrt{P_n} < \sqrt{y_n^2 + y_n + 1/2}$. Moreover, under the conditions of Case 1, we also have $\sqrt{P_n} > y_n + \frac{1}{2}$. These two inequalities yield

$$y_n + \frac{1}{2} < \sqrt{P_n} < \sqrt{y_n^2 + y_n + 1/2}$$
$$< \left(y_n + \frac{1}{2}\right) \left(1 + \frac{1}{8(y_n + \frac{1}{2})^2}\right)$$

Therefore,

(3.4)
$$\frac{1}{2} < \{\sqrt{P_n}\} < \frac{1}{2} + \frac{1}{\sqrt{P_n}}$$

Case 2: $y_n = \lceil \sqrt{P_n} \rceil$, $\|\sqrt{P_n}\| = |\sqrt{P_n} - (y_n - 1)| < |\sqrt{P_n} - y_n|$. Through arguments analogous to those in Case 1, we obtain the bounds:

$$\sqrt{y_n^2 - y_n + 1/2} < \sqrt{P_n} < y_n - \frac{1}{2},$$

which again leads to

(3.5)
$$\frac{1}{2} - \frac{1}{\sqrt{P_n}} < \{\sqrt{P_n}\} < \frac{1}{2}$$

To bound |E(x)|, we first break the interval [1, x] into dyadic pieces, and bound the number of $n \text{ in } [1, \sqrt{x}]$ trivially by $O(\sqrt{x})$. Now, for $\sqrt{x} < n < x$, using the closed form of P_n , we obtain

(3.6)
$$\sqrt{P_n} = 2\sqrt{\frac{n(n+1)(2n+1)}{6}}$$
$$= \frac{1}{\sqrt{3}}n^{3/2}\sqrt{\left(1+\frac{1}{n}\right)\left(1+\frac{1}{2n}\right)}$$
$$= \frac{1}{\sqrt{3}}n^{3/2} + \frac{\sqrt{3}}{4}n^{1/2} + O(n^{-1/2})$$

Combining (3.5) and (3.6), we achieve that

$$\left|\left\{\sqrt{P_n}\right\} - \frac{1}{2}\right| \le x^{-3/4},$$

for sufficiently large x. Thus,

(3.7)
$$|E(x)| = \#\left\{n: 1 \le n \le x, \left|\{\sqrt{P_n}\} - \frac{1}{2}\right| \le x^{-3/4}\right\} + O(\sqrt{x}).$$

Taking into account all of the above, for all $0 \le \alpha < \beta \le 1/2$, we have

$$\lim_{x \to \infty} \frac{\#\{n \colon \|\sqrt{P_n}\| \in [\alpha, \beta]\}}{x} - \lim_{x \to \infty} \frac{|E(x)|}{x} \le \lim_{x \to \infty} \frac{\#\{n \colon 1 \le n \le x, |\sqrt{P_n} - y_n| \in [\alpha, \beta]\}}{x}$$

and

$$\lim_{x \to \infty} \frac{\#\{n \colon 1 \le n \le x, |\sqrt{P_n} - y_n| \in [\alpha, \beta]\}}{x} \le \lim_{x \to \infty} \frac{\#\{n \colon \|\sqrt{P_n}\| \in [\alpha, \beta]\}}{x} + \lim_{x \to \infty} \frac{|E(x)|}{x}.$$

Applying Proposition 3.3,

$$\lim_{x \to \infty} \frac{\#\{n \colon \|\sqrt{P_n}\| \in [\alpha, \beta]\}}{x} = \lim_{x \to \infty} \frac{\#\{n \colon \{\sqrt{P_n}\} \in [\alpha, \beta] \cup [1 - \beta, 1 - \alpha]\}}{x}$$
$$= 2(\beta - \alpha).$$

Moreover,

$$\lim_{x \to \infty} \frac{|E(x)|}{x} = \lim_{x \to \infty} \frac{\#\{n \colon \{\sqrt{P_n}\} \in [\frac{1}{2} - x^{-3/4}, \frac{1}{2} + x^{-3/4}]\}}{x} + x^{-1/2}$$
$$= \lim_{x \to \infty} 2x^{-3/4} + x^{-1/2} = 0.$$

Thus,

$$\lim_{x \to \infty} \frac{\#\{n \colon 1 \le n \le x, |\sqrt{P_n} - y_n| \in [\alpha, \beta]\}}{x} = 2(\beta - \alpha)$$

which, by the definition of equidistribution, completes the proof of Proposition 3.4.

4. PROOF OF THEOREMS 1.1 AND 1.2

In this section, we provide proofs for Theorems 1.1 and 1.2. Before proving the theorems, we first introduce two useful lemmas. The following lemma is a modified version of an auxiliary lemma by Das, Robles, Dirk, and the fourth author [5]. The proof is essentially the same as the original one, so we omit it here.

Lemma 4.1 ([5, Lem. 5.1]). Let F and G be continuous, real-valued functions on \mathbb{R}_+ such that F is strictly decreasing and G is increasing. Further, suppose that

(4.1)
$$\lim_{x \to \infty} F(x) = \lim_{x \to 0} G(x) = 0 \text{ and } \lim_{x \to 0} F(x) = \lim_{x \to \infty} G(x) = \infty$$

Set $H(x) := \max\{F(x), G(x)\}$. We then have

(4.2)
$$\min_{x \in (0,\infty)} H(x) = F(U) = G(U),$$

where U is the solution of the equation F(U) = G(U).

Definition 4.2. Let (u_n) be a sequence of points in the circle group $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. For $0 \le \alpha \le 1$,

$$Z(N;\alpha) = \#\{n \in \mathbb{Z} : 1 \le n \le N, \ 0 \le u_n \le \alpha \bmod 1\}$$

Let $D(N; \alpha) = Z(N; \alpha) - N\alpha$. Then the discrepancy of the sequence is

$$D(N) = \sup_{\alpha \in [0,1]} |D(N;\alpha)|.$$

Lemma 4.3 (Erdös-Turán inequality, [8, Coro. 1.1, p. 8]). Let $(x_n)_{n \leq N}$ be a finite sequence with discrepancy D(N). Then, for any positive integer K,

$$D(N) \le \frac{N}{K+1} + 3\sum_{m=1}^{K} \frac{1}{m} \bigg| \sum_{n \le N} e(mx_n) \bigg|.$$

Since the result of Theorem 1.1 follows immediately from Theorem 1.2 by taking k = 1, we begin with the proof of Theorem 1.2.

Proof of Theorem 1.2. We write

(4.3)
$$a_n = \left|\sqrt{P_n} - y_n\right| \left|\sqrt{P_n} + y_n\right|$$

Then,

$$M_k(x) = \sum_{1 \le n \le x} a_n^k = \sum_{1 \le n \le x} |\sqrt{P_n} - y_n|^k |\sqrt{P_n} + y_n|^k.$$

By the discussion in the proof of Proposition 3.4, y_n is either $\lfloor \sqrt{P_n} \rfloor$ or $\lceil \sqrt{P_n} \rceil$, so

$$(4.4) y_n = \sqrt{P_n} + O(1),$$

and thus $y_n + \sqrt{P_n} = 2\sqrt{P_n} + O(1)$. Moreover,

(4.5)

$$\left(\sqrt{P_n} + y_n\right)^k = \left(2\sqrt{P_n} + O(1)\right)^k$$

$$= \binom{k}{0} \left(2\sqrt{P_n}\right)^k + O\left(\binom{k}{1} \left(2\sqrt{P_n}\right)^{k-1}\right)$$

$$= \left(2\sqrt{P_n}\right)^k + O_k\left(n^{\frac{3}{2}(k-1)}\right).$$

Upon using the closed form of $\sqrt{P_n}$ in (3.6), we have

$$\left(\sqrt{P_n} + y_n\right)^k = \left(2\sqrt{\frac{n(n+1)(2n+1)}{6}}\right)^k + O_k\left(n^{\frac{3}{2}(k-1)}\right)$$
$$= \left(\frac{2}{\sqrt{3}}n^{3/2}\sqrt{\left(1+\frac{1}{n}\right)\left(1+\frac{1}{2n}\right)}\right)^k + O_k\left(n^{\frac{3}{2}(k-1)}\right)$$
$$= \left(\frac{2}{\sqrt{3}}\right)^k n^{\frac{3}{2}k} \left(1+O\left(\frac{1}{n}\right)\right)^k + O_k\left(n^{\frac{3}{2}(k-1)}\right)$$
$$= \left(\frac{2}{\sqrt{3}}\right)^k n^{\frac{3}{2}k} + O_k\left(n^{\frac{3}{2}k-1}\right).$$

Using (4.6), upon trivially bounding $|\sqrt{P_n} - y_n| \le 1$ in the error term, we now have that

$$M_k(x) = \sum_{1 \le n \le x} |\sqrt{P_n} - y_n|^k \left(\frac{2}{\sqrt{3}}\right)^k n^{\frac{3}{2}k} + O_k(x^{\frac{3}{2}k}).$$

By Proposition 3.4, choose an even $L \in \mathbb{N}$, which will be optimized later, we have

$$M_k(x) = \left(\frac{2}{\sqrt{3}}\right)^k \sum_{j=1}^{L/2} \sum_{\substack{1 \le n \le x\\ \frac{j-1}{L} < |\sqrt{P_n} - y_n| \le \frac{j}{L}}} |\sqrt{P_n} - y_n|^k n^{\frac{3}{2}k} + O_k(x^{\frac{3}{2}k}).$$

Note that

(4.7)
$$U_k(x,L) \le M_k(x) \le V_k(x,L),$$

where

(4.8)
$$U_k(x,L) = \left(\frac{2}{\sqrt{3}}\right)^k \sum_{j=1}^{L/2} \left(\frac{j-1}{L}\right)^k \sum_{\substack{1 \le n \le x \\ \frac{j-1}{L} < |\sqrt{P_n} - y_n| \le \frac{j}{L}} n^{\frac{3}{2}k} + O_k\left(x^{\frac{3}{2}k}\right),$$

and

(4.9)
$$V_k(x,L) = \left(\frac{2}{\sqrt{3}}\right)^k \sum_{j=1}^{L/2} \left(\frac{j}{L}\right)^k \sum_{\substack{1 \le n \le x \\ \frac{j-1}{L} < |\sqrt{P_n} - y_n| \le \frac{j}{L}}} n^{\frac{3}{2}k} + O_k(x^{\frac{3}{2}k}).$$

We remark that one can obtain the main term of $U_k(x, L)$ and $V_k(x, L)$ at this point by use of the fact that $|\sqrt{P_n} - y_n|$ is equidistributed in [0, 1/2] from Proposition 3.4. However, we want a more accurate error term, so we proceed as follows. To evaluate $V_k(x, L)$, we focus on the inner sum on the right of (4.9). Let $M \leq x$ be in \mathbb{N} , and define

(4.10)
$$S_j := \sum_{\substack{1 \le n \le x \\ \frac{j-1}{L} < |\sqrt{P_n} - y_n| \le \frac{j}{L}}} n^{\frac{3}{2}k} = \sum_{0 \le \ell \le M-1} S_{j,\ell,M}$$

where

(4.11)
$$S_{j,\ell,M} := \sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ \frac{j-1}{L} \le |\sqrt{P_n} - y_n| < \frac{j}{L}}} n^{\frac{3}{2}k}.$$

(4.6)

We therefore have the bounds

$$\left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \le n^{\frac{3k}{2}} \le \left(\frac{(\ell+1)x}{M}\right)^{\frac{3k}{2}}.$$

Note that

$$\left(\frac{(\ell+1)x}{M}\right)^{\frac{3k}{2}} = \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \left(\frac{\ell+1}{\ell}\right)^{\frac{3k}{2}}$$
$$= \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \left(1 + O_k\left(\frac{1}{\ell}\right)\right)$$
$$= \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} + O_k\left(\left(\frac{\ell x}{M}\right)^{\frac{3k}{2}}\frac{1}{\ell}\right).$$

Hence,

(4.12)
$$n^{\frac{3k}{2}} = \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} + O_k\left(\frac{\ell^{\frac{3k}{2}-1}x^{\frac{3k}{2}}}{M^{\frac{3k}{2}}}\right).$$

In summary, (4.11) can be rewritten as

$$S_{j,\ell,M} = \sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ \frac{j-1}{L} < |\sqrt{P_n} - y_n| \le \frac{j}{L}}} \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} + O\left(\frac{\ell^{\frac{3k}{2}-1}x^{\frac{3k}{2}}}{M^{\frac{3k}{2}}}\right)$$

$$= \sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ \frac{j-1}{L} < \{\sqrt{P_n}\} \le \frac{j}{L}}} \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} + \sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ 1-\frac{j}{L} < \{\sqrt{P_n}\} \le 1-\frac{j-1}{L}}} \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}}$$

$$(4.13) \qquad + O\left(\sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ \frac{1}{2} - x^{-\frac{3}{4}} < \{\sqrt{P_n}\} \le \frac{1}{2} - x^{-\frac{3}{4}}}} \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} + \frac{\ell^{\frac{3k}{2}-1}x^{\frac{3k}{2}}}{M^{\frac{3k}{2}}}\right),$$

where the second equality follows from (3.7). We now focus on the first sum in (4.13).

$$\sum_{\substack{\frac{\ell x}{M} \le n < \frac{(\ell+1)x}{M} \\ \frac{j-1}{L} < \sqrt{P_n} \le \frac{j}{L}}} \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} = \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \# \left\{ n \in \left[\frac{\ell x}{M}, \frac{(\ell+1)x}{M}\right] : \left\{\sqrt{P_n}\right\} \in \left(\frac{j-1}{L}, \frac{j}{L}\right) \right\}$$
$$= \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \left(\frac{x}{LM} + O(D(\mathcal{U}_{\ell,M}))\right),$$

where

(4.14)
$$\mathcal{U}_{\ell,M} = \left\{ \{\sqrt{P_n}\} \colon n = \left\lfloor \frac{\ell x}{M} \right\rfloor + i, \ 1 \le i \le \left\lfloor \frac{x}{M} \right\rfloor \right\}$$

and $D(\mathcal{U}_{\ell,M})$ represents the discrepancy of the family $\mathcal{U}_{\ell,M}$. Applying the same arguments to the second sum and error term in (4.13), we have

(4.15)
$$S_{j,\ell,M} = \left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} \cdot \frac{2x}{LM} + O\left(\left(\frac{\ell x}{M}\right)^{\frac{3k}{2}} D(\mathcal{U}_{\ell,M}) + \frac{\ell^{\frac{3k}{2}} x^{\frac{3k}{2} + \frac{1}{4}}}{M^{\frac{3k}{2} + 1}}\right).$$

To bound $D(\mathcal{U}_{\ell,M})$, we apply Lemma 4.3. Then, for any $K \geq 1$,

(4.16)
$$D(\mathcal{U}_{\ell,M}) \leq \left\lfloor \frac{x}{M} \right\rfloor \frac{1}{K+1} + 3\sum_{m=1}^{K} \frac{1}{m} \left\lfloor \sum_{1 \leq i \leq \left\lfloor \frac{x}{M} \right\rfloor} e(m\sqrt{P_i}) \right\rfloor.$$

Now our focus is to obtain an upper bound for the exponential sum in (4.16), where

(4.17)
$$\sum_{1 \le i \le \left\lfloor \frac{x}{M} \right\rfloor} e(m\sqrt{P_i}) = \sum_{\left\lfloor \frac{\ell x}{M} \right\rfloor \le n \le \left\lfloor \frac{(\ell+1)x}{M} \right\rfloor} e(m\sqrt{P_n}).$$

By Proposition 3.3, we obtain

$$\left| \sum_{\lfloor \frac{\ell x}{M} \rfloor \le n \le \lfloor \frac{(\ell+1)x}{M} \rfloor} e(m\sqrt{P_n}) \right| \\ \le \left(m \left| h' \left(\left\lfloor \frac{(\ell+1)x}{M} \right\rfloor \right) - h' \left(\left\lfloor \frac{\ell x}{M} \right\rfloor \right) \right| + 2 \right) \left(4 \left(mh'' \left(\left\lfloor \frac{(\ell+1)x}{M} \right\rfloor \right) \right)^{-1/2} + 3 \right).$$

Using Taylor series expansion, we achieve that

$$\left|h'\left(\left\lfloor\frac{(\ell+1)x}{M}\right\rfloor\right) - h'\left(\left\lfloor\frac{\ell x}{M}\right\rfloor\right)\right| \ll \frac{x}{M}h''\left(\frac{\ell x}{M}\right).$$

Thus,

$$\left|\sum_{\lfloor\frac{\ell x}{M}\rfloor \leq n \leq \lfloor\frac{(\ell+1)x}{M}\rfloor} e(m\sqrt{P_n})\right| \ll \left(\frac{mx}{M}h''\left(\frac{\ell x}{M}\right) + 2\right) \left(4\left(mh''\left(\lfloor\frac{(\ell+1)x}{M}\rfloor\right)\right)^{-1/2} + 3\right)$$
$$\ll \left(\frac{mx}{M} \cdot \frac{\sqrt{M}}{\sqrt{\ell x}} + 2\right) \left(\frac{\ell^{1/4}x^{1/4}}{m^{1/2}M^{1/4}} + 3\right)$$
$$\ll \frac{m^{\frac{1}{2}}x^{\frac{3}{4}}}{\ell^{\frac{1}{4}}M^{\frac{3}{4}}} + \frac{m\sqrt{x}}{\sqrt{M\ell}} + \frac{\ell^{\frac{1}{4}}x^{\frac{1}{4}}}{m^{\frac{1}{2}}M^{\frac{1}{4}}}.$$

$$(4.18)$$

Substituting (4.18) into (4.16), we obtain

(4.19)
$$D(\mathcal{U}_{\ell,M}) \ll \frac{x}{KM} + \sum_{m=1}^{K} \frac{x^{\frac{3}{4}}}{m^{\frac{1}{2}}\ell^{\frac{1}{4}}M^{\frac{3}{4}}} + \frac{\sqrt{x}}{\sqrt{M\ell}} + \frac{\ell^{\frac{1}{4}}x^{\frac{1}{4}}}{m^{\frac{3}{2}}M^{\frac{1}{4}}} \\ \ll \frac{x}{KM} + \frac{K^{\frac{1}{2}}x^{\frac{3}{4}}}{\ell^{\frac{1}{4}}M^{\frac{3}{4}}} + \frac{K\sqrt{x}}{\sqrt{M\ell}} + \frac{\ell^{\frac{1}{4}}x^{\frac{1}{4}}}{K^{\frac{1}{2}}M^{\frac{1}{4}}}.$$

Combining (4.15) and (4.19), we see that

$$S_{j,\ell,M} = \frac{2\ell^{\frac{3k}{2}}x^{\frac{3k}{2}+1}}{LM^{\frac{3k}{2}+1}} + O\left(\frac{\ell^{\frac{3k}{2}}x^{\frac{3k}{2}+1}}{KM^{\frac{3k}{2}+1}} + \frac{K^{\frac{1}{2}}\ell^{\frac{3k}{2}-\frac{1}{4}}x^{\frac{3k}{2}+\frac{3}{4}}}{M^{\frac{3k}{2}+\frac{3}{4}}} + \frac{K\ell^{\frac{3k}{2}-\frac{1}{2}}x^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{3k}{2}+\frac{1}{4}}}\right) + \frac{\ell^{\frac{3k}{2}-1}x^{\frac{3k}{2}}}{M^{\frac{3k}{2}}} + \frac{\ell^{\frac{3k}{2}}x^{\frac{3k}{2}+\frac{1}{4}}}{M^{\frac{3k}{2}+1}}\right).$$

$$(4.20)$$

Upon substitution of (4.20) back into (4.10), we show that

$$S_{j} = \sum_{0 \leq \ell \leq M-1} \left(\frac{2\ell^{\frac{3k}{2}} x^{\frac{3k}{2}+1}}{LM^{\frac{3k}{2}+1}} + O\left(\frac{\ell^{\frac{3k}{2}} x^{\frac{3k}{2}+1}}{KM^{\frac{3k}{2}+1}} + \frac{K^{\frac{1}{2}}\ell^{\frac{3k}{2}-\frac{1}{4}} x^{\frac{3k}{2}+\frac{3}{4}}}{M^{\frac{3k}{2}+\frac{3}{4}}} + \frac{K\ell^{\frac{3k}{2}-\frac{1}{2}} x^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{3k+1}{2}}\ell^{\frac{1}{2}}} + \frac{\ell^{\frac{3k}{2}+\frac{1}{4}} x^{\frac{3k}{2}+\frac{1}{4}}}{K^{\frac{1}{2}}M^{\frac{3k}{2}+\frac{1}{4}}} \right) \\ + \frac{\ell^{\frac{3k}{2}-1} x^{\frac{3k}{2}}}{M^{\frac{3k}{2}}} + \frac{\ell^{\frac{3k}{2}} x^{\frac{3k}{2}+\frac{1}{4}}}{M^{\frac{3k}{2}+1}} \right) \right) \\ = \frac{2x^{\frac{3k}{2}+1}}{L(\frac{3}{2}k+1)} + O_k \left(\frac{x^{\frac{3k}{2}+1}}{LM} + \frac{x^{\frac{3k}{2}+1}}{K} + K^{\frac{1}{2}} x^{\frac{3k}{2}+\frac{3}{4}} + \frac{Kx^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{1}{2}}} + \frac{Mx^{\frac{3k}{2}+\frac{1}{4}}}{K^{\frac{1}{2}}} + x^{\frac{3k}{2}+\frac{1}{4}} \right).$$

Therefore, substituting the above back into the expression for $V_k(x, L)$ in (4.9), and applying Euler-Maclaurin summation to the sum over j, we obtain

$$\begin{aligned} V_{k}(x,L) &= \left(\frac{2}{L\sqrt{3}}\right)^{k} \sum_{j=1}^{L/2} j^{k} \left(\frac{2x^{\frac{3k}{2}+1}}{L(\frac{3}{2}k+1)} + O_{k} \left(\frac{x^{\frac{3k}{2}+1}}{LM} + \frac{x^{\frac{3k}{2}+1}}{K} + K^{\frac{1}{2}}x^{\frac{3k}{2}+\frac{3}{4}} \right. \\ &+ \frac{Kx^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{1}{2}}} + \frac{Mx^{\frac{3k}{2}+\frac{1}{4}}}{K^{\frac{1}{2}}} + x^{\frac{3k}{2}+\frac{1}{4}} \right) \\ &= \frac{2^{k}}{3^{k/2}L^{k}} \left(\frac{(\frac{L}{2})^{k+1}}{k+1} + \frac{(\frac{L}{2})^{k}}{2} + O(L^{k-1})\right) \left(\frac{2x^{\frac{3k}{2}+1}}{L(\frac{3}{2}k+1)} + O_{k} \left(\frac{x^{\frac{3k}{2}+1}}{LM} + \frac{x^{\frac{3k}{2}+1}}{K} \right) \\ &+ K^{\frac{1}{2}}x^{\frac{3k}{2}+\frac{3}{4}} + \frac{Kx^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{1}{2}}} + \frac{Mx^{\frac{3k}{2}+\frac{1}{4}}}{K^{\frac{1}{2}}} + x^{\frac{3k}{2}+\frac{1}{4}} \right) \\ &= \frac{x^{\frac{3k}{2}+1}}{3^{k/2}(k+1)(\frac{3}{2}k+1)} + O_{k} \left(\frac{x^{\frac{3k}{2}+1}}{L} + \frac{x^{\frac{3k}{2}+1}}{K} + K^{\frac{1}{2}}x^{\frac{3k}{2}+\frac{3}{4}} + \frac{Kx^{\frac{3k}{2}+\frac{1}{2}}}{M^{\frac{1}{2}}} \right) \\ (4.21) &\qquad + \frac{Mx^{\frac{3k}{2}+\frac{1}{4}}}{K^{\frac{1}{2}}} + x^{\frac{3k}{2}+\frac{1}{4}} \right). \end{aligned}$$

Finally, we need to optimize M, K, and L to minimize the error term in (4.21). To do this, apply Lemma 4.1 with

$$F(M) = \frac{Kx^{\frac{3k}{2} + \frac{1}{2}}}{M^{\frac{1}{2}}}, G(M) = \frac{Mx^{\frac{3k}{2} + \frac{1}{4}}}{K^{\frac{1}{2}}}.$$

We obtain $M = Kx^{1/6}$. Substituting M back into (4.21), and applying Lemma 4.1 with

$$F(K) = \frac{x^{\frac{3k}{2}+1}}{K}, G(K) = K^{\frac{1}{2}} x^{\frac{3k}{2}+\frac{3}{4}},$$

we have $K = x^{1/6}$. Plugging the optimal values of M and K back into (4.21), and taking L arbitrarily large, we eventually get

$$V_k(x,L) = \frac{x^{\frac{3k}{2}+1}}{3^{k/2}(k+1)\left(\frac{3}{2}k+1\right)} + O_k\left(x^{\frac{3k}{2}+\frac{5}{6}}\right).$$

A similar calculation yields

$$U_k(x,L) = \frac{x^{\frac{3}{2}k+1}}{3^{k/2}(\frac{3}{2}k+1)(k+1)} + O_k\left(x^{\frac{3k}{2}+\frac{5}{6}}\right).$$

Thus, using (4.7), we have, for any $k \in \mathbb{N}$,

$$M_k(x) = \frac{x^{\frac{3}{2}k+1}}{3^{k/2}(\frac{3}{2}k+1)(k+1)} + O_k\left(x^{\frac{3k}{2}+\frac{5}{6}}\right).$$

This finishes the proof of Theorem 1.2.

Theorem 1.1 now follows as a corollary of Theorem 1.2 by taking k = 1.

REFERENCES

- [1] L. Beeckmans, Squares expressible as sum of consecutive squares, *The American Mathematical Monthly*, **25** (1994), 437–442.
- [2] M. A. Bennett, Lucas' square pyramid problem revisited, Acta Arithmetica, 105 (2002), 341–347.
- [3] J. H. Conway and N. J. A. Sloane, Lorentzian forms for the Leech lattice, *Bulletin (New Series) of the American Mathematical Society*, **6** (1982), 215–217.
- [4] J. H. Conway and N. J. A. Sloane, Sphere Packings, Lattices and Groups. Springer, (1999), pp. 524–528.
- [5] M. Das, N. Robles, A. Zaharescu, and D. Zeindler, Partitions into semiprimes, arxiv preprint arXiv: 2212.12489 [math.NT], 2022. Available at https://arxiv.org/abs/2212.12489.
- [6] I. Kuipers and H. Niederreiter, Uniform Distribution of Sequences, Wiley-Interscience, (1974).
- [7] M. Laub, Squares expressible as a sum of *n* consecutive squares, *The American Mathematical Monthly*, **97** (1990), 622–625.
- [8] H. L. Montgomery, *Ten Lectures on the Interface Between Analytic Number Theory and Harmonic Analysis.* American Mathematical Society, (1994).
- [9] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, Published electronically at https://oeis.org, 2024.
- [10] Questions, Nouvelles annales de mathématiques : journal des candidats aux écoles polytechnique et normale, 14 (1875), 336. Available at http://www.numdam.org/item/NAM_1875_2_14_336_0/.
- [11] G. N. Watson, The problem of the square pyramid, Messenger of Mathematics, 48 (1918), 1–22.
- [12] P. Xausa, The On-Line Encyclopedia of Integer Sequences, 2024. Available at https://oeis.org/A351830/b351830.txt.

ANJI DONG: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, ALT-GELD HALL, 1409 WEST GREEN STREET, URBANA, IL, 61801, USA *Email address*: anjid2@illinois.edu

KATERINA SAETTONE: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, ALTGELD HALL, 1409 WEST GREEN STREET, URBANA, IL, 61801, USA *Email address*: kas18@illinois.edu

KENDRA SONG: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, ALTGELD HALL, 1409 WEST GREEN STREET, URBANA, IL, 61801, USA *Email address*: kendras4@illinois.edu

ALEXANDRU ZAHARESCU: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, ALTGELD HALL, 1409 WEST GREEN STREET, URBANA, IL, 61801, USA AND SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P. O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA *Email address*: zaharesc@illinois.edu

10