Tensor Product CFTs and One-Character Extensions

Chethan N. Gowdigere^{1 ab}, Sachin Kala^{2 ab}, Jagannath Santara^{3 c}

^aNational Institute of Science Education and Research Bhubaneshwar, P.O. Jatni, Khurdha, 752050, Odisha, INDIA

 b Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400094, INDIA

> c Department of Physics, Indian Institute of Technology Madras, Chennai 600036, INDIA

E-mail: ¹chethan.gowdigere@niser.ac.in, ²sachin.kala@niser.ac.in, ³jagannath.santara@physics.iitm.ac.in

Abstract

We study one-character CFTs obtained as one-character extensions of the tensor products of a single CFT \mathcal{C} . The motivation comes from the fact that 28 of the 71 CFTs in the Schelleken's list of $c = 24$ CFTs are such CFTs. We study for C : (i) any two-character WZW CFT with vanishing Wronskian index, (ii) the Ising CFT, (iii) the infinite class of $D_{r,1}$ CFTs and the $A_{4,1}$ CFT. The characters being S-invariant homogenous polynomials of the characters of \mathcal{C} , when organised in terms of a S-invariant basis, take compact forms allowing for closed form answers for high central charges. We find a S-invariant basis for each of the CFTs studied. As an example, one can find an explicit expression for the character of the monster CFT as a degree-48 polynomial of the characters of the Ising CFT. In some CFTs, some of the S-invariant polynomials of characters compute, after using the q-series of the characters, to a constant value. Hence, the characters of one-character extensions are more properly elements of the quotient ring of polynomials (of characters) with the ideal needed for the quotient, generated by S-invariant polynomials that compute to a constant.

1 Introduction

Two dimensional conformal field theories (CFTs) are a subject of great interest for a whole range of theoretical scientists [\[1](#page-41-0)[–6\]](#page-42-0). They are studied intensely by theoretical physicists, of both the high-energy and condensed matter varieties, as well as mathematical physicists and physical mathematicians. They are studied by high-energy theorists, especially string theorists, because of it's relevance to the world-sheet description of string theory and string perturbation theory. They are studied by condensed matter physicists because they describe two dimensional critical systems and the fixed point of the renormalisation group. They are also relevant for the Kondo problem, describe universality classes of quantum Hall systems and find applications in entanglement entropy as well as quantum computing. Two dimensional CFTs have contributed to many areas of pure

[6 Conclusions and Future Directions](#page-40-0) 41

mathematics viz. infinite dimensional Lie algebras, theory of modular forms, representation theory, etc.

A classification of such an object, which seems to possess such a wide range of interest and application, should be desirable. But it seems that classifying all CFTs may be quite hard and perhaps is at some distance in the future. A subclass of CFTs that one may attempt a classification for, are the so-called rational conformal field theories (RCFTs). These are two dimensional CFTs with some rationality properties : the central charge and the conformal dimensions of fields are restricted to be rational numbers. The rationality results in a finiteness property [\[7\]](#page-42-1) : the partition function can be written in terms of a finite number of characters.

A classification scheme for two dimensional rational conformal field theories was given by Mathur, Mukhi and Sen (MMS) in 1988. The scheme is based on properties of the torus partition function of a RCFT. The torus partition function of a RCFT can be written in terms of a finite number of characters. One denotes this by \bf{n} and is one of the two positive integers that forms a label for every RCFT. The other positive integer (actually, non-negative) comes about as follows. The n characters of a RCFT, it turns out, constitute a set of linearly independent solutions for a certain n'th order ordinary differential equation, known as the modular linear differential equation (MLDE) [\[8\]](#page-42-2). Thus there is a MLDE associated to every RCFT. Now, one considers the Wronskian of the characters, which turns out to be modular of weight $n(n - 1)$. From the Riemann-Roch theorem a.k.a the valence formula for modular functions/forms, one obtains information about the location of the zeroes of the Wronskian. This zeroes' information is encoded in what is termed as the Wronskian index or the Mukhi index^{[1](#page-2-0)} of a RCFT, denoted here by l. of a RCFT, denoted here by l. According to the MMS scheme, one should identify any RCFT by a label [n, l] consisting of the number of characters and the Wronskian/Mukhi index of that RCFT.

A second crucial feature of the MMS classification program of RCFT is the fact that the $[n, l]$ label is also a label for MLDEs. Given a $[n, l]$, one can construct a unique MLDE, unique upto the presence of a finite number of parameters. A given $[n, l]$ CFT solves the $[n, l]$ MLDE for some values of it's parameters. Another $[n, l]$ CFT solves the same $[n, l]$ MLDE but for different values of the parameters. Thus to obtain all $[n, l]$ CFTs, one only needs to solve one MLDE, over it's entire parameter space. In this way, the MMS classification program reduces the infinitely hard problem of classifying all RCFTs to an infinite number of (still hard but hopefully only) finitely hard problems of classifying RCFTs with a given $[n, l]$. Historically, [2, 0] RCFTs were the first to be classified [\[9,](#page-42-3)10]. The classification of $[3, 0]$ CFTs was begun in [\[10\]](#page-42-4) and was completed in [\[26\]](#page-43-0); this was achieved after the classification of $\mathbf{3, 0}$ characters $\mathbf{22}, \mathbf{23}, \mathbf{24}.$

This paper largely concerns CFTs with $n = 1$. They have a single character which can be

¹This important detail about a RCFT, which is part of the identification label of a RCFT in the MMS scheme, had not been named for most of it's existence. It was given the name "Wronskian index" by Prof. Sunil Mukhi in 2016. The name "Mukhi index" has also been used by some workers in this field.

thought of as an identity character, with a q-series : $\chi(q) = q^{-\frac{c}{24}}(1 + a_1q + \ldots)$. They have extremely simple modular transformation properties [\[19\]](#page-43-4). The identity is self-conjugate, and hence the charge conjugation matrix C must be equal to 1. Therefore $S = \pm 1$ and the character satisfies $\chi(-\frac{1}{\tau}) = \pm \chi(\tau)$. Choosing $\tau = i$ and from $\chi(i) \neq 0$, we conclude that $S = 1$. Thus the character of a one-character CFT is S-invariant. Furthermore, since $(ST)^3 = C$, we find that $T = e^{\frac{2k\pi i}{3}}$, $k \in \mathbb{Z}$. On the other hand we have that the action of T on a character is given in terms of the central charge as $T = e^{-\frac{c}{24}2\pi i}$. It follows that the central charge c must be a multiple of 8. When c is a multiple of 24, we have $S = 1$ as well as $T = 1$ and hence the character itself is modular invariant. In this case, the character itself can serve as a modular invariant partition function (as in [\[20\]](#page-43-5)). For one-character CFTs with central charges given by other multiples of 8, we have $S = 1$ (S-invariance), but $T \neq 1$. The partition function of such CFTs is simply $\chi \overline{\chi}$.

To put one-character CFTs within the purview of the MMS classification scheme : their Wronskian/Mukhi indices, being $\frac{c}{4}$, are even numbers. One can readily write down the corresponding first-order MLDEs. But MLDE methods have not been that effective in studying one-character CFTs as they have been for multiple-character CFTs. Even in this paper, we will develop methods that avoid MLDEs and are very algebraic, but at the end of the say, produce solutions of the MLDEs.

The classification of one-character CFTs, available in the literature, is as follows. First, $c = 8$ or $\mathbf{[1,2]}$ theories: there is exactly one such CFT viz. the $E_{8,1}$ CFT. Then for $c = 16$ or $\mathbf{[1,4]}$ theories : there are exactly two CFTs viz. $E_{8,1} \otimes E_{8,1}$ and the one-character extension of $D_{16,1}$, which we denote by $\mathcal{E}_1[D_{16,1}]^2$ $\mathcal{E}_1[D_{16,1}]^2$. For $c = 24$ or $[1,6]$ CFTs, there is the famous classification due to Schellekens [\[19\]](#page-43-4) where there are a total of 71 one-character CFTs. For $c > 24$, we are very far from a classification; some relevant papers are [\[21\]](#page-43-6) and [\[25\]](#page-43-7).

We make the following observation about the one-character CFTs in the Schellekens classification. 28 of the 71 CFTs are one-character extensions of tensor products of a single WZW CFT and we reproduce them here in table [1.](#page-4-0) The second column contains the numbering of the CFT in the original list of [\[19\]](#page-43-4) and the third column contains the one-character CFT. Inspired by this observation, we are now motivated to search for one-character CFTs by considering tensor products of a single mutliple-character CFT. In this paper, as an initial study, we consider tensor products of $[2, 0]$ and $[3, 0]$ CFTs. We first have to take the tensor products followed by searching for one-character extensions.

Let us denote the single CFT that we wish to tensor by $\mathcal C$ and let us take the tensor product of

²Given a multiple-character CFT \mathcal{C} , if one can find a combination of characters which has the modular properties of a one-character CFT, one would have found a one-character extension of C. The one-character CFT thus constructed is denoted by $\mathcal{E}_1[\mathcal{C}]$. Here $D_{16,1}$ is a three-character CFT. It has three characters χ_0, χ_1, χ_2 where the subscripts are the conformal dimensions. It turns out that $\chi_0 + \chi_2$ is S-invariant and $T = e^{\frac{4\pi i}{3}}$ and hence defines a one-character CFT.

#	Schellekens No.	One-character CFT	#	Schellekens No.	One-character CFT
1.	Sch. 1	$\mathcal{E}_1[U(1)^{\otimes 24}]$	15.	Sch. 38	$\mathcal{E}_1[C_{4,1}^{\otimes 4}]$
2.	Sch. 2	$\mathcal{E}_1[A_{1,4}^{\otimes 12}]$	16.	Sch. 41	$\mathcal{E}_1[B_{6.2}^{\otimes 2}]$
3.	Sch. 4	$\mathcal{E}_1[C_{4,10}]$	17.	Sch. 42	$\mathcal{E}_1[D_{4.1}^{\otimes 6}]$
4.	Sch. 5	$\mathcal{E}_1[A_{1,2}^{\otimes 16}]$	18.	Sch. 46	$\mathcal{E}_1[A_{6.1}^{\otimes 4}]$
5.	Sch. 6	$\mathcal{E}_1[A_{2.3}^{\otimes 6}]$	19.	Sch. 51	$\mathcal{E}_1[A_{8.1}^{\otimes 3}]$
6.	Sch. 9	$\mathcal{E}_1[A_{4.5}^{\otimes 2}]$	20.	Sch. 54	$\mathcal{E}_1[D_{6.1}^{\otimes 4}]$
7.	Sch. 11	$\mathcal{E}_1[A_{6,7}]$	21.	Sch. 57	$\mathcal{E}_1[B_{12,2}]$
8.	Sch. 12	$\mathcal{E}_1[C_{2.2}^{\otimes 6}]$	22.	Sch. 58	$\mathcal{E}_1[E_{6.1}^{\otimes 4}]$
9.	Sch. 15	$\mathcal{E}_1[A_{1,1}^{\otimes 24}]$	23.	Sch. 60	$\mathcal{E}_1[A_{12.1}^{\otimes 2}]$
10.	Sch. 23	$\mathcal{E}_1[B_{3.2}^{\otimes 4}]$	24.	Sch. 61	$\mathcal{E}_1[D_{8.1}^{\otimes 3}]$
11.	Sch. 24	$\mathcal{E}_1[A_{2.1}^{\otimes 12}]$	25.	Sch. 66	$\mathcal{E}_1[D_{12,1}^{\otimes 2}]$
12.	Sch. 29	$\mathcal{E}_1[B_{4.2}^{\otimes 3}]$	26.	Sch. 67	$\mathcal{E}_1[A_{24,1}]$
13.	Sch. 30	$\mathcal{E}_1[A_{3.1}^{\otimes 8}]$	27.	Sch. 68	$\mathcal{E}_1[E_{8,1}^{\otimes 3}]$
14.	Sch. 37	$\mathcal{E}_1[A_{4.1}^{\otimes 6}]$	28.	Sch. 70	$\mathcal{E}_1[D_{24.1}]$

Table 1: One-character extensions of the tensor product of a single CFT, that are in the Schellekens list of $c = 24$ CFTs [\[19\]](#page-43-4).

N copies of C. We will denote the tensor product CFT by $\mathcal{C}^{\otimes N}$. The N will depend on the central charge of $\mathcal C$ as well as the central charge of the one-character extension we wish to construct. We will give details pertaining to N in section [2.](#page-7-0) $\mathcal{C}^{\otimes N}$ is a multiple-character CFT. We will first learn how to construct it's characters. When C has two characters, say χ_0, χ_1 , then the characters of $\mathcal{C}^{\otimes N}$ are the monomials of degree N that can obtained from the homogenous polynomial $(\chi_0 + \chi_1)^N$. Thus, in this case, $\mathcal{C}^{\otimes N}$ has $N+1$ characters. Similarly, when $\mathcal C$ has three characters, say χ_0, χ_1, χ_2 , then the characters of $\mathcal{C}^{\otimes N}$ are the monomials of degree N that can obtained from the homogenous polynomial $(\chi_0 + \chi_1 + \chi_2)^N$. Thus in this case, $\mathcal{C}^{\otimes N}$ has $\frac{(N+1)(N=2)}{2}$ characters. The identity character of $\mathcal{C}^{\otimes N}$ is χ_0^N and hence $\mathcal{C}^{\otimes N}$ has a central charge equal to N times the central charge of $\mathcal{C}.$

Now that we understand how to construct tensor product CFTs, we can now proceed to the next step of the process in this paper viz. constructing one-character extensions. The character of a one-character extension CFT, $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$ is a certain (homogenous) polynomial of the characters of the tensor product CFT. This polynomial includes the identity character χ_0^N and other characters that have a commensurate q-series with the identity character^{[3](#page-4-1)}. Let us denote by b the number of such characters. The candidate character of the one-character extension is a polynomial with

³Every character of $\mathcal{C}^{\otimes N}$ has a q-series $q^{-\frac{cN}{24}+h}(a_0 + a_1q + a_2q^2 + ...)$ where h is it's conformal dimension; having a q-series commensurate with the identity character translates to h being a positive integer.

 $b+1$ terms (the $+1$ is for the identity character) and there are b number of arbitrary co-efficients in it. This polynomial is a homogenous polynomial of degree N . Let us denote this polynomial by $P(\chi_0, \chi_1, \ldots)$. Now we need to impose the modular data of a one-character extension, which we reviewed above, on $P(\chi_0, \chi_1, \ldots)$. The choice of monomials already ensures the correct action of T viz. by multiplication by a cube root of 1. Thus we have to only impose S-invariance :

$$
SP(\chi_0, \chi_1, \ldots) \equiv P(S(\chi_0), S(\chi_1), \ldots) = P(\chi_0, \chi_1, \ldots) \tag{1}
$$

In the above, the first \equiv is simply the definition : the S-transformation of $P(\chi_0, \chi_1, \ldots)$ is derived from the S-transformation of the χ_i 's, that is the modular S-matrix of C. The second equality is the imposition of S-invariance. This equality is an equality of two homogenous polynomials of degree N . Thus it results in as many equations as there are monomials of degree N , which are way larger than the number of undetermined coefficients in $P(\chi_0, \chi_1, \ldots)$. Thus we have an overdetermined linear system of equations for the b undetermined coefficients in $P(\chi_0, \chi_1, \ldots)$. This is our procedure. We solve simultaneously some finite number of linear equations and we determine the polynomial $P(\chi_0, \chi_1, \ldots)$. Our procedure is thus very simple technically : algebraic manipulations of homogenous polynomials in polynomial rings; we do not solve any MLDEs here.

All these considerations are implemented in the subsequent sections. In section two, we begin by working out the systematics of the tensor product and the extension. Given a CFT \mathcal{C} , it is not always possible to obtain a one-character CFT with every central charge. For example, it is impossible to obtain a one-character CFT with central charge 8 when $C = G_{2,1}$. Hence, we begin in the next section by deriving, given \mathcal{C} , what are the allowed values of N and what are the values of the central charges 8k of the one-character extension that can be achieved.

Then, in section three, we apply the above delineated procedure when $\mathcal C$ is a [2,0] WZW CFT. Recall that the list of [2,0] WZW CFTs is in one-to-one correspondence with the Deligne series of exceptional Lie algebras viz. $\{A_{1,1}, A_{2,1}, G_{2,1}, D_{4,1}, F_{4,1}, E_{6,1}, E_{7,1}\}$. We describe in full detail the computations for $A_{1,1}$. We are able to obtain one-character extensions $\mathcal{E}_1[A_{1,1}]$ with central charges upto 120. Then we give the results for $A_{2,1}$, $G_{2,1}$ and $D_{4,1}$. The other WZW CFTs in the [2,0] classification are $F_{4,1}$, $E_{6,1}$ and $E_{7,1}$. It is well-known that they can be obtained as a coset CFT : coseting $E_{8,1}$ by $G_{2,1}$, $A_{2,1}$, and $A_{1,1}$ respectively. Hence the modular S-matrices of $F_{4,1}$, $E_{6,1}$ and $E_{7,1}$ are identical to those of $G_{2,1}$, $A_{2,1}$, and $A_{1,1}$ respectively. We can adapt the computations for the latter to obtain results for the former.

In section [4,](#page-27-0) we apply the above described procedure for a few (including an infinite class) of [3, 0] WZW CFTs. We study the $A_{4,1}$ CFT first. Then we study the Ising CFT. FInally we study the infinite set of $D_{r,1}$ CFTs for all r. It turns out there are four categories (i) r is an odd number, (ii) r is twice of an odd number, (iii) r is four times and odd number, (iv) r is a multiple of eight; and there is a common formulation for tall he solutions in any category.

Before we proceed further in summarizing the contents of this paper, we should dwell on a major caveat. MLDE studies of CFTs always proceed in two waves. In the first wave, one works only with characters and the emphasis is on scanning all of the MLDE parameter space and obtaining solutions with non-negative coefficients in the q -series. These solutions are referred to as admissible characters. One has to note that these solutions are far from being characters of CFTs. For example, every multiple of a non-identity character is also a solution, but clearly all multiples of all characters will not have the correct modular properties. In the second wave of analysis, one worries about which of the admissible character solutions are characters of CFTs. For example, for $[2, 0]$ CFTs, the first wave of analysis was done in the seminal paper [\[9\]](#page-42-3), and the second wave in [\[10\]](#page-42-4). It is useful to note that except one, all admissible character solutions of the $[2, 0]$ MLDE resulted in CFTs. For [3, 0] CFTs, the first wave of analysis was done in the papers [\[22\]](#page-43-1), [\[23\]](#page-43-2), [\[24\]](#page-43-3) and the second wave in [\[26\]](#page-43-0). Here the dichotomy between admissible character solutions (around 300 in number) and actual CFTs (around 45 in number) is a little more severe. This dichotomy between admissible characters and actual CFTs is most severe for one-character CFTs. For example, for $c = 24$, there are an infinite number of admissible characters, but only 71 actual CFTs, the celebrated Schellekens list of CFTs [\[19\]](#page-43-4). For $c > 24$, the first wave of analysis is readily performed and admissible characters are found to be infinite in number, for each central charge. But the second wave of analysis has not been done yet and a list of actual CFTs is not available. Now comes the caveat : in this paper, we are very much within the first wave of analysis. We are working with admissible characters only. Imposing modular data as we do via [\(1\)](#page-5-0) results in only admissible characters. We may use the expression "one-character extension CFT" but we are always only working with a "character of a possible one-character extension CFT".

In this paper, the way we have setup the problem of finding characters of possible one-character CFTs is algebraic. All our computations are within a polynomial ring. Remember that our answers are homogenous polynomials that are invariant under the S-transformation. To start with we work with the natural basis of generators in which the problem is defined viz. the characters of the $\mathcal C$ CFT. One of the major results of this paper is that this is a very inefficient basis of generators for the homogenous polynomial algebra. We find an alternate basis of generators, where the basis elements themselves are S-invariant. We discovered this basis from the answers of the first few cases. For computations of every two-character CFT that we perform in section [2,](#page-7-0) we find two S-invariant polynomial basis $\{P_1, P_2\}$. We express all our answers in terms of this basis and the expressions become compact. Similarly for the three-characer CFTs we study in section [3,](#page-12-0) we find three S-invariant polynomial basis $\{P_1, P_2, P_3\}$. Because of the efficient computations and compact forms, we are able to find explicit answers, for example to questions such as: what is the degree 48 polynomial of characters of the Ising CFT that is the character of the monster CFT?

At the outset, it seems that the character of the one-character CFT $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$ is an element

of the polynomial ring of characters of \mathcal{C} . But surprisingly we find that (for some CFTs) it is properly an element of a quotient ring of this polynomial ring. We find that for some CFTs there are polynomial relations that exist between the characters. And then these polynomial relations generate ideals which should be used to obtain the quotient ring. This is the content of section [5.](#page-38-0)

2 Systematics of Tensor Products and One-character Extensions

In this section, we will describe the general theory and working method of this paper. We will first discuss the tensor product procedure and the characters of a tensor product. Then we discuss the procedure of obtaining one-character extensions. We will discuss how to obtain an ansatz for the character of a one-character extension and impose the modular properties of one-character CFTs on the ansatz. We will learn to impose the correct T -transformation and then the methodology to impose S-invariance. We will also elucidate on the relationship between the number of tensor product factors, the central charge of the one-character extension and explicitly display this in tables for the CFTs we consider in this paper.

The CFT which is being tensored is denoted by \mathcal{C} . Let us denote it's central charge by $c(\mathcal{C})$ and let n_c denote the number of characters of C. In this paper we have $n_c = 2, 3$ and C is either a $[2, 0]$ or a $[3, 0]$ WZW CFT. Let us denote the characters of C by

$$
\chi_0, \chi_1, \dots, \chi_{(n_c-1)}.\tag{2}
$$

Here, χ_0 is the identity character. The non-identity characters χ_i have conformal dimensions h_i , , $i = 1, 2, \ldots (n_c - 1)$. The numbering of characters is such that as i increases, h_i increases. For an unitary CFT, this is just a matter of naming the characters. For non-unitary CFTs (which we do not deal with, for the most part, in this paper), this can be done when one works with the unitary presentation of the non-unitary CFT. The q-series of the characters take the form :

$$
\chi_i(q) = q^{-\frac{c(C)}{24} + h_i} \left(p_0^{(i)} + p_1^{(i)} q + p_2^{(i)} q^2 + \ldots \right)
$$
 (3)

with the Fourier coefficients $p_k^{(i)}$ $k_k^{(i)}$, $k = 0, 1...$ all non-negative integers and $p_0^{(0)} = 1$. The action of S-transformation on the characters can be written as a $n_c \times n_c$ and the this matrix can be extracted from the modular S-matrix of the CFT (which can be bigger in size, when multiple "primary fields" have the same character). We will denote this $n_c \times n_c$ matrix by

$$
S_{ij}, \qquad i, j = 0, 1, \dots (n_c - 1). \tag{4}
$$

Let us denote by N the number of copies of the C CFT that we take in the tensor product : $\mathcal{C}^{\otimes N}$. The characters of this tensor product CFT are nothing but the monomials of the homogenous $degree-N$ polynomial

$$
(\chi_0 + \chi_0 + \dots \chi_{(n_c - 1)})^N
$$
 (5)

and they can be written explicitly :

$$
\Pi_{i=0}^{(n_C-1)} \chi_i^{d_i}, \qquad \sum_{i=0}^{(n_C-1)} d_i = N. \tag{6}
$$

For $n_c = 2$, the number characters of $C^{\otimes N}$ is $(N + 1)$ and for $n_c = 3$, the number characters of $\mathcal{C}^{\otimes N}$ is $\frac{(N+1)(N+2)}{2}$. The q-series of the character [\(6\)](#page-8-0) begins with $q^{-\frac{c(C)N}{24}+\sum_{i=0}^{(n_C-1)}h_id_i}$. F or the identity character χ_0^N , the q-series is $q^{-\frac{c(C)N}{24}}(1+\ldots)$. Hence, we find that the central charge of tensor product CFT, $C^{\otimes N}$, is thus $Nc(C)$.

A one-character extension of a CFT D is a one-character CFT constructed from the CFT D . The notation for this is $\mathcal{E}_1[\mathcal{D}]$. In this paper, we mostly study characters of one-character CFTs and not the CFTs themselves. We will use the same notation $(\mathcal{E}_1[\mathcal{D}])$ for the character as well. The character of the one-character extension $\mathcal{E}_1[\mathcal{D}]$ has the extremely simple modular properties of one-character CFTs, which we have already detailed in the introduction. There is a S-invariance and T acts on the single character by mutliplication of a cube root of unity, which results in the central charge of a one-character CFT to be a multiple of 8.

Our ultimate goal is to construct CFTs which are one-character extensions of the tensor product CFT $\mathcal{C}^{\otimes N}$ with $\mathcal C$ either a $[\mathbf{2},\mathbf{0}]$ or $[\mathbf{3},\mathbf{0}]$ WZW CFTs. But in this paper, we will only address a smaller goal and achieve this to a large extent for the cases we consider. The smaller goal is the following : using the characters of the tensor product CFT $\mathcal{C}^{\otimes N}$ and the accompanying modular $data⁴$ $data⁴$ $data⁴$ as ingredients, construct a character with the modular properties of a one-character CFT. A subsequent study (not in this paper) should then establish which of the characters of this paper will result in a genuine one-character CFT and this genuine one-character CFT should have the notation $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$. For example, in table [1,](#page-4-0) every $\mathcal{E}_1[\ldots]$ refers to a genuine one-character CFT (and not just a character with the required modular properties). But, with a slight abuse of notation, when we write $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$, we may mean only (the character for) a possibility for a one-character CFT.

The first step in constructing the character for a possible one-character CFT is to identify all characters of $\mathcal{C}^{\otimes N}$ that have a q-series commensurate with the identity character. This happens for the character in (6) when the d_i 's are such that

$$
\sum_{i=1}^{(n_C-1)} h_i d_i = M, \qquad M \in \mathbf{Z}^+. \tag{7}
$$

We should remember that the d_i 's need to satisfy the homogeneity constraint also:

$$
\sum_{i=0}^{(n_{\mathcal{C}}-1)} d_i = N \tag{8}
$$

⁴The modular data for the $\mathcal{C}^{\otimes N}$ CFT can be derived from the modular data for \mathcal{C} .

Thus, to find all characters of $\mathcal{C}^{\otimes N}$ that have a q-series commensurate with the identity character χ_0^N , we need to simultaneously solve, for the d_i 's, the two equations [\(7\)](#page-8-2) and [\(8\)](#page-8-3). Note that d_0 does not occur in (7) . Solutions for (7) and (8) exist for a finite number of values for m, only for $m < Nh_{(n,-1)}$. Let us denote the number of solutions by b and the solutions themselves by

$$
d_i^{(I)}, \t I = 1, 2, \dots b.
$$
\t(9)

Consider the following sum of characters :

$$
\chi_0^N + \sum_{I=1}^b p_I \left(\Pi_{i=0}^{(n_C-1)} \chi_i^{d_i^{(I)}} \right) \equiv P(\chi_0, \chi_1, \dots, \chi_{(n_C-1)}), \tag{10}
$$

where the p_1 's are arbitrary constants at this stage. Later, they will be fixed (partially, if not completely) by requiring $P(\chi_0, \chi_1, \ldots, \chi_{(n_C-1)})$ to satisfy the modular properties of a one-character CFT.

The q-series of each of the summands in [\(10\)](#page-9-0) begins^{[5](#page-9-1)} with $q^{-\frac{c(C)N}{24}}$. This means that the qseries of $P(\chi_0, \chi_1, \ldots \chi_{(n_{\mathcal{C}}-1)})$ begins with $q^{-\frac{c(\mathcal{C})N}{24}}$. Let us denote by $c_{(1)}$ the central charge of the one-character CFT which we want to construct, the one whose character is [\(10\)](#page-9-0). We can thus conclude

$$
c_{(1)} = Nc(\mathcal{C}).\tag{11}
$$

Now we impose the modular properties of a one-character CFT. First we impose the Ttransformation: to be a cube root of unity, which means $c_{(1)}$ is multiple of 8. - hence we have

$$
c_{(1)} = 8k, \t\t(12)
$$

$$
Nc(\mathcal{C}) = 8k, \qquad k \in \mathbf{Z}^+ \tag{13}
$$

Imposing the S transformation forms the bulk of the paper, in sections two and three. Here, in this section, we will note the consequences of the T-transformation. (13) is the main equation for this section, which will help us answer the questions :

(1) Given a C, what all $c_{(1)}$'s can be achieved? In other words what all central-charges of onecharacter CFTs can be achieved by the process of taking tensor products and one-character extensions?

(2) And for each of central charges of one-character CFT that is possible, what is the number of tensor product factors of $\mathcal C$ i.e. N needed?

⁵After factoring out this beginning, the rest of the q-series, the integral powers, starts with $q^{h_i d_i^{(I)}}$ for the summand in [\(10\)](#page-9-0) with coefficient p_I . For the identity character χ_0^N , this rest of the q-series start with q^0 . Thus the integral power part of the q-series of each of the summands of [\(10\)](#page-9-0) can be added like usual power series. This is the meaning of commensurate q-series.

Since, C is a RCFT, it's central charge $c(C)$ is a rational number. - Let it be $\frac{m}{n}$ with m and n being co-prime positive integers. We then have, following [\(13\)](#page-9-2),

$$
N = \frac{8k}{m}n\tag{14}
$$

N is a whole number. All the factors of m need to be cancelled by factors of the numerator. One of the factors of m viz. $gcd(m, 8)$ is cancelled by the 8 of the numerator. The other factor of m viz. $\frac{m}{\gcd(m,8)}$ needs to be cancelled by k in the numerator because n, being co-prime with m cannot cancel it. Thus all possible values of k are given by

$$
k = \text{multiples of } \frac{m}{\text{gcd}(m, 8)}\tag{15}
$$

and the number of tensor product factors N follows from (14) and is given by

$$
N = \text{multiples of } \frac{8n}{\text{gcd}(m, 8)}.\tag{16}
$$

The above answers the two questions we asked earlier. The answers are summarised for tensor products of $[2, 0]$ CFTs in table [2](#page-10-1) and for $[3, 0]$ CFTs in [3.](#page-11-0) For $A_{1,1}$, following entry 1. of table [2,](#page-10-1) we find that the allowed tensor powers are all multiples of 8 and the corresponding $c_{(1)}$'s are (the same) multiples of 8. For, $A_{1,1}$, $A_{2,1}$ and $D_{4,1}$ CFTs, every possible $c_{(1)}$ can be achieved by some tensor power. For the other five $[2, 0]$ CFTs not all $c_{(1)}$'s can be achieved. We find that for $G_{2,1}, c_{(1)} = 56, 112, 168, 224, \ldots$ can be achieved by taking tensor powers $N = 20, 40, 60, 80 \ldots$ respectively. For $F_{4,1}$, the one-character CFT with the smallest central charge that one can hope to obtain is 104. From table [3,](#page-11-0) entry 1. we see that one can hope to obtain one-character CFTs for every possible central charge by taking some number of tensor powers of the Ising CFT viz. $\mathcal{M}(4,3)$. An interesting case is the monster CFT which can be obtained by taking tensor products of 48 copies of the Ising CFT. We will see this in section [4](#page-27-0) where we can give explicitly the character of the monster CFT in terms of the Ising characters.

	N	$20 \mid 40 \mid 60 \mid 80 \mid 100 \mid 120 \mid 140 \mid \ldots$			
	6. $E_{6,1}$ $c_{(1)} = 8k$ 24 48 72 96 120 144 168				
		$4 \mid 8 \mid 12 \mid 16 \mid 20 \mid 24 \mid 28 \mid $			
	7. $E_{7,1}$ $c_{(1)} = 8k$ 56 112 168 224 280 336 392				
		$8 \mid 16 \mid 24 \mid 32 \mid 40 \mid 48 \mid 56 \mid \ldots$			

Table 2: Allowed values of number of tensor product factors N and central charge $c_{(1)}$ of onecharacter extensions for [2, 0] CFTs.

$\#$	\mathcal{C}									
1.	$\mathcal{M}(4,3)$	8k	8	16	24	32	40	48	56	\ldots .
		\boldsymbol{N}	16	32	48	64	80	96	112	\ldots
2.	$A_{4,1}$	8k	8	16	24	32	40	48	56	\sim \sim \sim
		N	2	$\overline{4}$	$6\,$	8	10	12	14	\ldots
3.	$D_{r,1}$	8k	$8(2p+1)$	$16(2p + 1)$		$24(2p+1)$ $32(2p+1)$		$40(2p+1)$ $48(2p+1)$ $56(2p+1)$		\sim \sim \sim
	$r = 2p + 1$	\boldsymbol{N}	8	16	24	32	40	48	56	\ldots .
	$p\in {\bf Z}^+$									\ldots
4.	$D_{r,1}$	8k	$8(2p + 1)$	$16(2p + 1)$		$24(2p+1)$ $32(2p+1)$		$40(2p+1)$ $48(2p+1)$ $56(2p+1)$		\ldots
	$r = 2(2p + 1)$	N	$\overline{4}$	8	12	16	20	24	28	\ldots
	$p \in \mathbf{Z}^+$									\ldots
5.	$D_{r,1}$	8k	$8(2p + 1)$	$16(2p + 1)$		$24(2p+1)$ $32(2p+1)$		$40(2p+1)$ $48(2p+1)$ $56(2p+1)$		\ldots .
	$r = 4(2p + 1)$	\boldsymbol{N}	2	$\overline{4}$	6	8	10	12	14	\ldots .
	$p \in \mathbf{Z}^+$									\ldots
6.	$D_{r,1}$	8k	8p'	16p'	24p'	32p'	40p'	48p'	56p'	\ldots .
	$r=8p'$	\boldsymbol{N}	$\mathbf{1}$	$\overline{2}$	3	$\overline{4}$	5	6	$\overline{7}$	\cdots
	$p'\in{\bf Z}^+$									\cdots

Table 3: Allowed values of number of tensor product factors N and central charge $c_{(1)}$ of onecharacter extensions for [3, 0] CFTs.

In [\(10\)](#page-9-0), what we have is an ansatz for the character of the one-character extension $P(\chi_0, \ldots \chi_{(n_c-1)})$. This ansatz together with (13) incorporates the correct T-transformation property. What is left to do is to impose S-invariance. First we learn how to S-transform the ansatz character P : the S-transformation of P is derived from the S-transformation of the characters χ_i , from [\(4\)](#page-7-1).

$$
S(P(\chi_0, \chi_1, \dots, \chi_{(n_c-1)})) \equiv P(S(\chi_0), S(\chi_1), \dots, S(\chi_{(n_c-1)})), \qquad S(\chi_i) = \sum_j S_{ij} \chi_j.
$$
 (17)

Now we can impose S-invariance via

$$
P(S(\chi_0), S(\chi_1), \dots S(\chi_{(n_c-1)})) = P(\chi_0, \chi_1, \dots \chi_{(n_c-1)}).
$$
\n(18)

We know that the right hand side is a certain homogenous polynomial of degree N with a b monomials (and b is smaller than the total number of monomials of degree N). But the left hand side, which is also a homogenous polynomial of degree N , may contain all the monomials. In anycase, we can describe (18) as the vanishing of a homogenous polynomial of degree N. Hence, we have as many equations as there are monomials of degree N . Each of these equations is a linear equation for the b unknown coefficients p_I of the ansatz character P. We thus have an over-determined linear system of equations. Thus, to make the ansatz character P S-invariant, we need to chose the p_1 's to be from a solution of these linear system of equations. We will use the notation $P_{c_{(1)}=8k}^{\mathcal{C}}(\chi_0,\ldots)$ to denote the S-invariant character $(P(\chi_0,\ldots,\chi_{(n_{\mathcal{C}}-1)})$ is used to denote the ansatz character before imposing S-invariance).

This is what we do in this paper, we pick a CFT $\mathcal C$ and a N and a k , in accordance with the tables [2,](#page-10-1) [3.](#page-11-0) We set up the linear system of equations and solve them. It turns out that we always obtain solutions. Only very rarely, we get the rank of the coefficient matrix of the linear system of equations to be exactly b, giving unique solutions. For most of the cases, we the rank of the coefficient matrix is less than b giving families of solutions.

3 [2, 0] CFTs : Tensor Products and One-character Extensions

In this section, we study [2, 0] CFTs. We implement the procedure described in section [2](#page-7-0) for each of the $[2, 0]$ WZW CFTs. We start with the smallest central charge viz. $A_{1,1}$ and go up the sequence in increasing central charges. We will solve the linear system of equations associated with S-invariance for $A_{1,1}$, $A_{2,1}$, $G_{2,1}$ and $D_{4,1}$. The solutions of the first three are also solutions of $E_{7,1}$, $E_{6,1}$ and $F_{4,1}$ respectively. We will describe the $A_{1,1}$ case in some detail and be somewhat brief for the others.

3.1 $C = A_{1,1}$

 $A_{1,1}$ is a [2,0] CFT with central charge 1. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{1}{4}$ $\frac{1}{4}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{1}{24}}(1 + 3q + 4q^2 + 7q^3 + 13q^4 + 19q^5 + 29q^6 + 43q^7 + 62q^8 + 90q^9 + 126q^{10} + \dots)
$$

\n
$$
\chi_1 = q^{\frac{5}{24}}(2 + 2q + 6q^2 + 8q^3 + 14q^4 + 20q^5 + 34q^6 + 46q^7 + 70q^8 + 96q^9 + 138q^{10} + \dots)
$$
\n(19)

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + \chi_1}{\sqrt{2}}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_1}{\sqrt{2}}.
$$
 (20)

The allowed values of k and N can be read from entry 1. of table [2.](#page-10-1) Here, every possible $c_{(1)} = 8k$ is possible and the corresponding number of tensor product factors is $N = 8k$.

We begin with $N = 8, c_{(1)} = 8$. $A_{1,1}^{\otimes 8}$ has nine characters : $\chi_0^m \chi_1^n$, $m + n = 8$ with m, n nonnegative integers. The conformal dimension of $\chi_0^m \chi_1^n$ is $\frac{n}{4}$. Solving [\(7\)](#page-8-2) and [\(8\)](#page-8-3) we obtain the characters which have q-series commensurate with the identity : $\{\chi_0^8, \chi_0^4 \chi_1^4, \chi_1^8\}$. Hence, the ansatz character (10) which incorporates the relevant T-transformation is

$$
P(\chi_0, \chi_1) = \chi_0^8 + p_1 \chi_0^4 \chi_1^4 + p_2 \chi_1^8,\tag{21}
$$

which has two arbitrary coefficients.

Now, we impose S-invariance, (18) , (20) . We get a system of 9 linear equations for the two variables p_1 and p_2 . Remarkably, there is a unique solution. We thus have the following unique character for $N = 8, c_{(1)} = 8$:

$$
P_{c_{(1)}=8}^{A_{1,1}}(\chi_0,\chi_1) = \chi_0^8 + 14\,\chi_0^4\,\chi_1^4 + \chi_1^8. \tag{22}
$$

We can now compute the q-series of $P_{c_{(1)}=8}^{A_{1,1}}(\chi_0, \chi_1)$ using [\(19\)](#page-12-4) and we get $P_{c_{(1)}=8}^{A_{1,1}}(\chi_0, \chi_1)(q) = j^{\frac{1}{3}}$. This is the unique character possible at $c_{(1)} = 8$. But we have obtained this well-known result by a few algebraic manipulations. Only for $c_{(1)} = 8$, we have a unique character possible and that unique characters corresponds to a unique CFT viz. $E_{8,1}$. Hence, what we have derived is the fact that

$$
\mathcal{E}_1[A_{1,1}^{\otimes 8}] = E_{8,1} \tag{23}
$$

We then study the second entry of row 1. of table [2](#page-10-1) viz. $N = 16, c_{(1)} = 16$. $A_{1,1}^{\otimes 16}$ has 17 characters : $\chi_0^m \chi_1^n$, $m + n = 16$ with m, n non-negative integers. The conformal dimension of $\chi_0^m \chi_1^n$ is $\frac{n}{4}$. Solving [\(7\)](#page-8-2) and [\(8\)](#page-8-3) we obtain the characters which have q-series commensurate with the identity : $\{\chi_0^{16}, \chi_0^{12} \chi_1^4, \chi_0^8 \chi_1^8, \chi_0^4 \chi_1^{12}, \chi_1^{16}\}\$. Hence, the ansatz character [\(10\)](#page-9-0) which incorporates the relevant T-transformation is

$$
P(\chi_0, \chi_1) = \chi_0^{16} + p_1 \chi_0^{12} \chi_1^4 + p_2 \chi_0^8 \chi_1^8 + p_3 \chi_0^4 \chi_1^{12} + p_4 \chi_1^{16},
$$
 (24)

which has four arbitrary coefficients.

.

Now, we impose S-invariance, (18) , (20) . We get a system of 17 linear equations for the 4 variables p_1, p_2, p_3, p_4 . Remarkably, again there is a unique solution. We thus have the following unique character for $N = 16$, $c_{(1)} = 16$:

$$
P_{c_{(1)}=16}^{A_{1,1}}(\chi_0,\chi_1) = \chi_0^{16} + 28\chi_0^{12}\chi_1^4 + 198\chi_0^8\chi_1^8 + 28\chi_0^4\chi_1^{12} + \chi_1^{16}
$$
 (25)

We can now compute the q-series of $P_{c_{(1)}=16}^{A_{1,1}}(\chi_0,\chi_1)$ using [\(19\)](#page-12-4) and we get $P_{c_{(1)}=16}^{A_{1,1}}(\chi_0,\chi_1)(q) = j^{\frac{2}{3}}$. This is the unique character possible at $c_{(1)} = 16$. But we have obtained this well-known result by a few algebraic manipulations. Only for $c_{(1)} = 16$, we have a unique character possible but that unique characters corresponds to two different CFT viz. $E_{8,1}^{\otimes 2}$ and $\mathcal{E}_1[D_{16,1}]$. Hence, what we have derived is the fact that

$$
\mathcal{E}_1[A_{1,1}^{\otimes 16}] = E_{8,1}^{\otimes 2} \quad \text{or} \quad \mathcal{E}_1[D_{16,1}]. \tag{26}
$$

We further note that the S-invariant polynomial of degree 16 in (25) is nothing but the square of the S-invariant polynomial of degree 8 in [\(22\)](#page-13-1). Let us denote the latter by

$$
P_1(\chi_0, \chi_1) \equiv \chi_0^8 + 14 \chi_0^4 \chi_1^4 + \chi_1^8 \tag{27}
$$

so that we can write both the results for $c_{(1)} = 8, 16$ in terms of this P_1 :

$$
P_{c_{(1)}=8}^{A_{1,1}}(\chi_0,\chi_1) = P_1(\chi_0,\chi_1), \qquad P_{c_{(1)}=16}^{A_{1,1}}(\chi_0,\chi_1) = P_1(\chi_0,\chi_1)^2
$$
\n(28)

One of the major themes of this paper is the existence of a few S-invariant polynomials for every CFT C. Every character of any one-character extension $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$ can be written as a polynomial of these S-invariant polynomials. What we have discovered here in [\(27\)](#page-14-0) is the first S-invariant polynomial for $A_{1,1}$. And in [\(28\)](#page-14-1), we have written the characters for $\mathcal{E}_1[\mathcal{A}_{1,1}^{\otimes 8}]$ and $\mathcal{E}_1[\mathcal{A}_{1,1}^{\otimes 16}]$ as polynomials of P_1 . Note that as polynomials of the characters, they are of degree 8 and 16 respectively, but as polynomials of P_1 , they are of degree 1 and 2 respectively. This reduction in degrees when one goes from the description in terms of characters and the description in terms of S-invariant polynomials makes for compact expressions and reporting results for large N. We will see that there is another S-invariant polynomial $P_2(\chi_0, \chi_1)$ for $A_{1,1}$. And all results for $A_{1,1}$ can be written as polynomials for P_1 and P_2 . P_2 shows up first in the next case.

We then study the third entry of row 1. of table [2](#page-10-1) viz. $N = 24$, $c_{(1)} = 24$. $A_{1,1}^{\otimes 24}$ has 25 characters: $\chi_0^m \chi_1^n$, $m + n = 24$ with m, n non-negative integers. The conformal dimension of $\chi_0^m \chi_1^n$ is $\frac{n}{4}$. Solving (7) and (8) we obtain the characters which have q-series commensurate with the identity $\{\chi_0^{24-4r}\chi_1^{4r} \mid r=0,1,\ldots 6\}$. Hence, the ansatz character [\(10\)](#page-9-0) which incorporates the relevant T-transformation is

$$
P(\chi_0, \chi_1) = \sum_{r=0}^{6} p_r \ \chi_0^{24-4r} \chi_1^{4r}, \text{ where } p_0 = 1 \tag{29}
$$

which has 6 arbitrary coefficients.

.

Now, we impose S-invariance, (18) , (20) . We get a system of 25 equations for 6 variables p_1, \ldots, p_6 . This time we do not get a unique solution. The co-efficient matrix of the set of linear equations has rank 5. One of the coefficients is undetermined and the rest are given in terms of this coefficient :

$$
P_{c_{(1)}=24}^{A_{1,1}}(\chi_0, \chi_1) = \chi_0^{24} + p_1 \chi_0^{20} \chi_1^4 + (759 - 4p_1) \chi_0^{16} \chi_1^8 + (2576 + 6p_1) \chi_0^{12} \chi_1^{12} + (759 - 4p_1) \chi_0^8 \chi_1^{16} + p_1 \chi_0^4 \chi_1^{20} + \chi_1^{24} \chi_1^{20} + \chi_1^{24} \chi_1^{20} + \chi_1^{25} \chi_1^{20} + \chi_1^{26} \chi_1^{20} + \chi_1^{27} \chi_1^{20} + \chi_1^{28} \chi_1^{20} + \chi_1^{29} \chi_1^{20} + \chi_1^{20} \
$$

The polynomial in [\(30\)](#page-14-2) is S-invariant for all values of p_1 . In particular setting $p_1 = 0$ gives one of the summands of $P_{c_{(1)}=24}^{A_{1,1}}(\chi_0,\chi_1)$ which is separately S-invariant. This then implies that the coefficient of p_1 is separately S-invariant. Examining the coefficient of p_1 , we discover that it is the fourth power of the following S-invariant polynomial

$$
P_2(\chi_0, \chi_1) \equiv \chi_0^5 \chi_1 - \chi_0 \chi_1^5. \tag{31}
$$

The final answer for is $c_{(1)} = 24$ can be written in terms of the two S-invariant polynomials viz. the P_1 of [\(27\)](#page-14-0) and the P_2 of [\(31\)](#page-15-0).

$$
P_{c_{(1)}=24}^{A_{1,1}}(\chi_0, \chi_1) = P_1^3 - 42P_2^4 + p_1 P_2^4 \tag{32}
$$

We can now compute the *q*-series of $P_{c_{(1)}=24}^{A_{1,1}}(\chi_0, \chi_1)$ using [\(19\)](#page-12-4):

$$
P_{c_{(1)}=24}^{A_{1,1}}(\chi_0,\chi_1)(q) = j + 16p_1 - 672
$$
\n(33)

To obtain an admissible character, we need to choose p_1 such that $744+16p_1-672$ is a non-negative integer i.e. $p_1 \geq -4, p_1 \in \mathbb{Z}$.

Let us note that there is a fundamental difference between the two S-invariant polynomials [\(27\)](#page-14-0) and (31) . When we evaluate the q-series for the latter, we obtain the constant 2. We will refer to the latter as a constant S-invariant polynomial and discuss the significance of the same in section [5.](#page-38-0) The former has a non-trivial q-series and we will refer to it a as a non-constant S -invariant polynomial. Let us collect all the information about the two S-invariant polynomials of $A_{1,1}$ in one place :

$$
P_1(\chi_0, \chi_1) = \chi_0^8 + 14 \chi_0^4 \chi_1^4 + \chi_1^8 = j^{\frac{1}{3}}
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^5 \chi_1 - \chi_0 \chi_1^5 = 2
$$
\n(34)

It turns out that the character $P_{\text{cav}}^{A_{1,1}}$ $c_{(1)}=8k(\chi_0,\chi_1)$ for every k can be written as polynomials of P_1 and P_2 . Again let us note the compactness of the expressions. In equation [\(32\)](#page-15-1) we have a degree-4 polynomial and when written in terms of the characters χ_0, χ_1 as in [\(30\)](#page-14-2) we have a degree-24 polynomial.

We can now describe the rest of the results that we are reporting here for $A_{1,1}$: for $32 \le N \le 128$, $4 \leq k \leq 16$. First, we solve [\(7\)](#page-8-2) and [\(8\)](#page-8-3) and find that there are a total of $b = 2k$ monomials commensurate with the identity character. The ansatz character (10) has 2k arbitrary coefficients. Then we impose S-invariance - then we impose S-invariance, (18) , (20) . This amounts to solving $8k+1$ equations for $2k$ variables. It turns out that the rank of the coefficient matrix is $2k-\left[\frac{k}{3}\right]$ $\frac{k}{3}$ where [Q] represents the integer part of the rational number Q. So that the solution has $\left[\frac{k}{3}\right]$ undetermined (linear combinations of) coefficients. This means that S-invariant character $P_{\text{cusp}}^{A_{1,1}}$ $c_{(1)}=8k(\chi_0,\chi_1)$ can be

written as a sum of of $\left[\frac{k}{3}\right]+1$ different summands with each summand being separately S-invariant :

$$
P_{c_{(1)}=8k}^{A_{1,1}}(\chi_0, \chi_1) = \mathcal{Z}_0 + \sum_{j=1}^{\left[\frac{k}{3}\right]} p_j \mathcal{Z}_j \tag{35}
$$

The number of p 's is $\left[\frac{k}{3}\right]$. The number of \mathcal{Z} 's is $\left[\frac{k}{3}\right]+1$.

In table [4,](#page-17-0) we collect all the results for $k = 1, 2, \ldots 16$. The first column contains the number of tensor product factors N (which equals $8k$). In the second column, we give the central charge of the one-character extension $c_{(1)}$. The third column contains the character of the one-character extension $P_{\text{c}}^{A_{1,1}}$ $c_{(1)}=8k(\chi_0,\chi_1)$. The number of summands ranges from 1, for $k=1$, to 6 for $k=16$. For $k = 16$, the last entry of the table [4](#page-17-0) we have six S-invariant summands $\mathcal{Z}_0, \mathcal{Z}_1, \ldots \mathcal{Z}_5$.

The expressions for the Z's when written in terms of the characters χ_0, χ_1 are very voluminous. Each of them. Is a degree N polynomial. For example for $N = 128$, the last entry of table [4,](#page-17-0) we have six homogenous degree 128 polynomials. But, when written in terms of the S-invariant polynomials [\(34\)](#page-15-2), they are in fact polynomials of much smaller degree. For $N = 128$, the highest degree term that occurs is 21.

This is one of the results to report in this paper. Solving the huge linear system of equations that follow from S-invariance, organising in terms of some number of S-invariant summands and writing each of the summands in terms of S-invariant polynomials and thus achieving compact expressions.

7.	$\mathcal{E}_1[A_{1,1}^{\otimes 56}]$	$\mathcal{Z}_0 = P_1^7 - 98P_1^4P_2^4 + 973P_1P_2^8$	$j^{\frac{1}{3}}(j^2+N_1j+N_2)$
		$\mathcal{Z}_1 = P_1^4 P_2^4 - 52P_1 P_2^8$	$N_1 = 16 (p_1 - 98)$
		$Z_2 = P_1 P_2^8$	$N_2 = 256(p_2 - 52p_1 + 973)$
8.	$\mathcal{E}_1[A_{1,1}^{\otimes 64}]$	$\mathcal{Z}_0 = P_1^8 - 112P_1^5P_2^4 + 1896P_1^2P_2^8$	$j^{\frac{2}{3}}(j^2+N_1j+N_2)$
		$\mathcal{Z}_1 = P_1^5 P_2^4 - 66 P_1^2 P_2^8$	$N_1 = 16(p_1 - 112)$
		$Z_2 = P_1^2 P_2^8$	$N_2 = 256(p_2 - 66p_1 + 1896)$
9.	$\mathcal{E}_1[A_1^{\otimes 72}]$	$\mathcal{Z}_0 = P_1^9 - 126P_1^6P_2^4 + 3015P_1^3P_2^8 - 4398P_2^{12}$	$j^3 + N_1 j^2 + N_2 j + N_3$
		$\mathcal{Z}_1 = P_1^6 P_2^4 - 80 P_1^3 P_2^8 + 104 P_2^{12}$	$N_1 = 16(p_1 - 126)$
		$Z_2 = P_1^3 P_2^8 - 34P_2^{12}$	$N_2 = 256(p_2 - 80p_1 + 3015)$
		$\mathcal{Z}_3 = P_2^{12}$	$N_3 = 4096(104p_1 - 34p_2 + p_3 - 4398)$
10.	$\mathcal{E}_1[A_{1,1}^{\otimes 80}]$	$\mathcal{Z}_0 = P_1^{10} - 140P_1^7P_2^4 + 4330P_1^4P_2^8 - 15200P_1P_2^{12}$	$j^{\frac{1}{3}}(j^3+N_1j^2+N_2j+N_3)$
		$\mathcal{Z}_1 = P_1^7 P_2^4 - 94 P_1^4 P_2^8 + 775 P_1 P_2^{12}$	$N_1 = 16(p_1 - 140)$
		$\mathcal{Z}_2 = P_1^4 P_2^8 - 48 P_1 P_2^{12}$	$N_2 = -256(94p_1 - p_2 - 4330)$
		$Z_3 = P_1 P_2^{12}$	$N_3 = 4096(775p_1 - 48p_2 + p_3 - 15200)$
11.	$\mathcal{E}_1[A_{1,1}^{\otimes 88}]$	$\mathcal{Z}_0 = P_1^{11} - 154P_1^8P_2^4 + 5841P_1^5P_2^8 - 38126P_1^2P_2^{12}$	$j^{\frac{2}{3}}(j^3+N_1j^2+N_2j+N_3)$
		$\mathcal{Z}_1 = P_1^8 P_2^4 - 108 P_1^5 P_2^8 + 1642 P_1^2 P_2^{12}$	$N_1 = 16(p_1 - 154)$
		$\mathcal{Z}_2 = P_1^5 P_2^8 - 62P_1^2 P_2^{12}$	$N_2 = -256(108p_1 - p_2 - 5841)$
		$\mathcal{Z}_3 = P_1^2 P_2^{12}$	$N_3 = 4096(1642p_1 - 62p_2 + p_3 - 38126)$
12.	$\mathcal{E}_1[A_{1,1}^{\otimes 96}]$	$\mathcal{Z}_0 = P_1^{12} - 168P_1^9P_2^4 + 7548P_1^6P_2^8 - 75920P_1^3P_2^{12} + 28086P_2^{16}$	$j^4 + N_1 j^3 + N_2 j^2 + N_3 j + N_4$
		$\mathcal{Z}_1 = P_1^9 P_2^4 - 122 P_1^6 P_2^8 + 2705 P_1^3 P_2^{12} - 4302 P_2^{16}$	$N_1 = 16(p_1 - 168)$
		$\mathcal{Z}_2 = P_1^6 P_2^8 - 76P_1^3 P_2^{12} - 22P_2^{16}$	$N_2 = -256(122p_1 - p_2 - 7548)$
		$\mathcal{Z}_3 = P_1^3 P_2^{12} - 30 P_2^{16}$	$N_3 = 4096(2705p_1 - 76p_2 + p_3 - 75920)$
		$\mathcal{Z}_4 = P_2^{16}$	$N_4 = -65536(4302p_1 + 22p_2 + 30p_3 - p_4 - 28086)$
	13. $\mathcal{E}_1[A_{1,1}^{\otimes 104}]$	$\mathcal{Z}_0 = P_1^{13} - 182P_1^{10}P_2^4 + 9451P_1^7P_2^8 - 131326P_1^4P_2^{12} + 203333P_1P_2^{16}$	$j^{\frac{1}{3}}(j^4+N_1j^3+N_2j^2+N_3j+N_4)$
		$\mathcal{Z}_1 = P_1^{10} P_2^4 - 136 P_1^7 P_2^8 + 3964 P_1^4 P_2^{12} - 12560 P_1 P_2^{16}$	$N_1 = 16 (p_1 - 182)$
		$\mathcal{Z}_2 = P_1^7 P_2^8 - 90 P_1^4 P_2^{12} + 593 P_1 P_2^{16}$	$N_2 = -256(136p_1 - p_2 - 9451)$
		$\mathcal{Z}_3 = P_1^4 P_2^{12} - 44 P_1 P_2^{16}$	$N_3 = 4096 (3964p_1 - 90p_2 + p_3 - 131326)$
		$\mathcal{Z}_4 = P_1 P_2^{16}$	$N_4 = -65536 (12560p_1 - 593p_2 + 44p_3 - p_4 - 203333)$
	14. $\mathcal{E}_1[A_{1,1}^{\otimes 112}]$	$\mathcal{Z}_0 = P_1^{14} - 196P_1^{11}P_2^4 + 11550P_1^8P_2^8 - 207088P_1^5P_2^{12} + 652141P_1^2P_2^{16}$	$j^{\frac{2}{3}}(j^4+N_1j^3+N_2j^2+N_3j+N_4)$
		$\mathcal{Z}_1 = P_1^{11} P_2^4 - 150 P_1^8 P_2^8 + 5419 P_1^5 P_2^{12} - 32158 P_1^2 P_2^{16}$	$N_1 = 16 (p_1 - 196)$
		$\mathcal{Z}_2 = P_1^8 P_2^8 - 104 P_1^5 P_2^{12} + 1404 P_1^2 P_2^{16}$	$N_2 = -256(150p_1 - p_2 - 11550)$
		$\mathcal{Z}_3 = P_1^5 P_2^{12} - 58 P_1^2 P_2^{16}$	$N_3 = 4096 (5419p_1 - 104p_2 + p_3 - 207088)$
		$\mathcal{Z}_4 = P_1^2 P_2^{16}$	$N_4 = -65536 (32158p_1 - 1404p_2 + 58p_3 - p_4 - 652141)$
	15. $\mathcal{E}_1[A_{1,1}^{\otimes 120}]$	$\mathcal{Z}_0 = P_1^{15} - 210P_1^{12}P_2^4 + 13845P_1^9P_2^8 - 305950P_1^6P_2^{12} + 1571490P_1^3P_2^{16} - 492372P_2^{20}$	$j^5 + N_1 j^4 + N_2 j^3 + N_3 j^2 + N_4 j + N_5$
		$\mathcal{Z}_1 = P_1^{12} P_2^4 - 164 P_1^9 P_2^8 + 7070 P_1^6 P_2^{12} - 65840 P_1^3 P_2^{16} + 10093 P_2^{20}$	$N_1 = 16 (p_1 - 210)$
		$\mathcal{Z}_2 = P_1^9 P_2^8 - 118 P_1^6 P_2^{12} + 2411 P_1^3 P_2^{16} - 4670 P_2^{20}$	$N_2 = -256(164p_1 - p_2 - 13845)$
		$\mathcal{Z}_3 = P_1^6 P_2^{12} - 72 P_1^3 P_2^{16} - 132 P_2^{20}$	$N_3 = 4096 (7070p_1 - 118p_2 + p_3 - 305950)$
		$\mathcal{Z}_4 = P_1^3 P_2^{16} - 26 P_2^{20}$	$N_4 = -65536 (65840p_1 - 2411p_2 + 72p_3 - p_4 - 1571490)$
		$\mathcal{Z}_5 = P_2^{20}$	$N_5 = 1048576(10093p_1 - 4670p_2 - 132p_3 - 26p_4 + p_5 - 492372)$
16.		$\mathcal{E}_1[A_{1,1}^{\otimes 128}] \quad \mathcal{Z}_0=p_1^{16}-224 P_1^{13} P_2^4+16336 P_1^{10} P_2^8-430656 P_1^7 P_2^{12}+3196776 P_1^4 P_2^{16}-2696256 P_1 P_2^{20}$	$i^{\frac{1}{3}}(i^5+N_1i^4+N_2i^3+N_3i^2+N_4i+N_5)$
		$\mathcal{Z}_1=P_1^{13}P_2^4-178P_1^{10}P_2^8+8917P_1^7P_2^{12}-116350P_1^4P_2^{16}+158178P_1P_2^{20}$	$N_1 = 16 (p_1 - 224)$
		$\mathcal{Z}_2 = P_1^{10} P_2^8 - 132 P_1^7 P_2^{12} + 3614 P_1^4 P_2^{16} - 10608 P_1 P_2^{20}$	$N_2 = -256(178p_1 - p_2 - 16336)$
		$\mathcal{Z}_3 = P_1^7 P_2^{12} - 86 P_1^4 P_2^{16} + 427 P_1 P_2^{20}$	$N_3 = 4096 (8917p_1 - 132p_2 + p_3 - 430656)$
		$\mathcal{Z}_4 = P_1^4 P_2^{16} - 40 P_1 P_2^{20}$	$N_4 = -65536 (116350p_1 - 3614p_2 + 86p_3 - p_4 - 3196776)$
		$\mathcal{Z}_5 = P_1 P_2^{20}$	$N_5 = 1048576(158178p_1 - 10608p_2 + 427p_3 - 40p_4 + p_5 - 2696256)$

Table 4: Characters of one-character extensions of tensor products of $A_{1,1}$ (and $E_{7,1}$) written in a basis of S-invariant polynomials

3.2 $C = A_{2,1}$

 $A_{2,1}$ is a [2,0] CFT with central charge 2. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{1}{3}$ $\frac{1}{3}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{1}{12}} \left(1 + 8q + 17q^2 + 46q^3 + 98q^4 + 198q^5 + 371q^6 + 692q^7 + 1205q^8 + 2082q^9 + 3463q^{10} + \ldots \right)
$$

$$
\chi_1 = q^{\frac{1}{4}} \left(3 + 9q + 27q^2 + 57q^3 + 126q^4 + 243q^5 + 465q^6 + 828q^7 + 1458q^8 + 2463q^9 + 4104q^{10} + \ldots \right) \tag{36}
$$

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + 2\chi_1}{\sqrt{3}}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_1}{\sqrt{3}}.
$$
 (37)

The allowed values of k and N can be read from entry [2.](#page-10-1) of table 2. Here, every possible $c_{(1)} = 8k$ is possible and the corresponding number of tensor product factors is $N = 4k$.

We will be brief in our description of the results here. The analysis proceeds very similar to $A_{1,1}$. We discover the first S-invariant polynomial when we are working with $c_{(1)} = 8$. This is a degree-4 non-constant S-invariant polynomial. Just as for the $A_{1,1}$ case in section [4.1,](#page-27-1) we discover the second S-invariant polynomial when working $c_{(1)} = 24$. This is a constant S-invariant polynomial of degree-4. The two *S*-invariant polynomials for $A_{2,1}$ are :

$$
P_1(\chi_0, \chi_1) = \chi_0^4 + 8\chi_0 \chi_1^3 = j^{\frac{1}{3}}
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^3 \chi_1 - \chi_1^4 = 3
$$
\n(38)

It is a peculiarity of $A_{2,1}$ that both the S-invariant polynomials are of the same degree. This then results in the fact that the characters of $\mathcal{E}_1[A_{2,1}^{\otimes N}]$ are homogenous polynomials of the S-invariant polynomials also.

In table [5,](#page-20-1) we collect all the results for $k = 1, 2, \ldots 16$. For $c_{(1)} = 8k$ the character has $\left[\frac{k}{3}\right]$ arbitrary constants (the *p*'s) and there are $1 + \left[\frac{k}{3}\right]$ *S*-invariant summands the *Z*'s. Each summand, when expressed in terms of the characters χ_0, χ_1 are polynomials of degree 4k. But when expressed in terms of the S-invariant polynomials [\(38\)](#page-18-1), they are polynomials of only degree k. Because of this, we can report in a compact manner, computations upto $c_{(1)} = 128$ for which we have polynomials of degree 16 only.

4.
$$
\mathcal{E}_1[A_2^{\otimes 16}]
$$

\n $\mathcal{Z}_0 = P_1^A - 32P_1P_2^3$
\n $\mathcal{Z}_1 = P_1P_2^3$
\n $\mathcal{Z}_2 = P_1^2 - 40P_1^2P_2^3$
\n $\mathcal{Z}_3 = P_1^2P_2^3$
\n6. $\mathcal{E}_1[A_2^{\otimes 24}]$
\n $\mathcal{Z}_0 = P_1^6 - 48P_1^3P_2^3 + 48P_2^6$
\n $\mathcal{Z}_1 = P_1^2P_2^3$
\n $\mathcal{Z}_2 = P_2^6$
\n $\mathcal{Z}_3 = P_1^5P_2^3 - 21P_2^6$
\n $\mathcal{Z}_4 = P_1^3P_2^3 - 21P_2^6$
\n $\mathcal{Z}_5 = P_2^6$
\n $\mathcal{Z}_6 = P_1^7 - 48P_1^3P_2^3 + 48P_2^6$
\n $\mathcal{Z}_7 = P_2^7$
\n $\mathcal{Z}_8 = P_2^8$
\n $\mathcal{Z}_9 = P_1^7$
\n $\mathcal{Z}_1 = P_1^3P_2^3 - 21P_2^6$
\n $\mathcal{Z}_2 = P_2^6$
\n $\mathcal{Z}_3 = P_1^4P_2^3$
\n $\mathcal{Z}_4 = P_1^4P_2^3 - 29P_1P_2^6$
\n $\mathcal{Z}_5 = P_1P_2^5$
\n $\mathcal{Z}_6 = P_1^7 - 56P_1^4P_2^3 + 576P_1^2P_2^6$
\n $\mathcal{Z}_7 = 729(p_2 - 21p_1 + 48)$
\n7. $\mathcal{E}_1[A_2^{\otimes 24}]$
\n $\mathcal{Z}_6 = P_1^8 - 64P_1^5P_2^3 + 376P_1^2P_2^6$

12. $\mathcal{E}_1[A_2]$ 848	$\mathcal{Z}_0 = P_1^{12} - 96P_1^9P_2^3 + 2400P_1^6P_2^6 - 12704P_1^3P_2^9 + 96P_2^{12}$	$i^4 + N_1 i^3 + N_2 i^2 + N_3 i + N_4$
	$\mathcal{Z}_1 = P_1^9 P_2^3 - 69 P_1^6 P_2^6 + 807 P_1^3 P_2^9 - 1013 P_2^{12}$	$N_1 = 27(p_1 - 96)$
	$\mathcal{Z}_2 = P_1^6 P_2^6 - 42 P_1^3 P_2^9 - 57 P_2^{12}$	$N_2 = -729(69p_1 - p_2 - 2400)$
	$\mathcal{Z}_3 = P_1^3 P_2^9 - 15 P_2^{12}$	$N_3 = 19683(807p_1 - 42p_2 + p_3 - 12704)$
	$\mathcal{Z}_4 = P_2^{12}$	$N_4 = -531441(1013p_1 + 57p_2 + 15p_3 - p_4 - 96)$
13. $\mathcal{E}_1[A_2^{\otimes 52}]$	$\mathcal{Z}_0 = P_1^{13} - 104P_1^{10}P_2^3 + 3016P_1^7P_2^6 - 22360P_1^4P_2^9 + 16328P_1P_2^{12}$	$i^{\frac{1}{3}}(i^4+N_1i^3+N_2i^2+N_3i+N_4)$
	$\mathcal{Z}_1 = P_1^{10} P_2^3 - 77 P_1^7 P_2^6 + 1207 P_1^4 P_2^9 - 2029 P_1 P_2^{12}$	$N_1 = 27 (p_1 - 104)$
	$\mathcal{Z}_2 = P_1^7 P_2^6 - 50 P_1^4 P_2^9 + 127 P_1 P_2^{12}$	$N_2 = -729(77p_1 - p_2 - 3016)$
	$Z_3 = P_1^4 P_2^9 - 23P_1 P_2^{12}$	$N_3 = 19683(1207p_1 - 50p_2 + p_3 - 22360)$
	$\mathcal{Z}_4 = P_1 P_2^{12}$	$N_4 = -531441 (2029p_1 - 127p_2 + 23p_3 - p_4 - 16328)$
14. $\mathcal{E}_1[A_{2,1}^{\otimes 56}]$	$\mathcal{Z}_0 = P_1^{14} - 112 P_1^{11} P_2^3 + 3696 P_1^8 P_2^6 - 35728 P_1^5 P_2^9 + 56560 P_1^2 P_2^{12}$	$i^{\frac{2}{3}}(i^4+N_1i^3+N_2i^2+N_3i+N_4)$
	$\mathcal{Z}_1 = P_1^{11} P_2^3 - 85 P_1^8 P_2^6 + 1671 P_2^5 P_2^9 - 5029 P_1^2 P_2^{12}$	$N_1 = 27 (p_1 - 112)$
	$\mathcal{Z}_2 = P_1^8 P_2^6 - 58P_1^5 P_2^9 + 375P_1^2 P_2^{12}$	$N_2 = -729(85p_1 - p_2 - 3696)$
	$\mathcal{Z}_3 = P_1^5 P_2^9 - 31 P_1^2 P_2^{12}$	$N_3 = 19683(1671p_1 - 58p_2 + p_3 - 35728)$
	$\mathcal{Z}_4 = P_1^2 P_2^{12}$	$N_4 = -531441 (5029p_1 - 375p_2 + 31p_3 - p_4 - 56560)$
15. $\mathcal{E}_1[A_2]_0^{\otimes 60}$	$\mathcal{Z}_0 = P_1^{15} - 120P_1^{12}P_2^3 + 4440P_1^9P_2^6 - 53320P_1^6P_2^9 + 140760P_1^3P_2^{12} - 41184P_2^{15}$	$i^5 + N_1 i^4 + N_2 i^3 + N_3 i^2 + N_4 i + N_5$
	$\mathcal{Z}_1 = P_1^{12} P_2^3 - 93 P_1^9 P_2^6 + 2199 P_1^6 P_2^9 - 10525 P_1^3 P_2^{12} - 3420 P_2^{15}$	$N_1 = 27 (p_1 - 120)$
	$\mathcal{Z}_2 = P_1^9 P_2^6 - 66 P_1^6 P_2^9 + 687 P_1^3 P_2^{12} - 1264 P_2^{15}$	$N_2 = -729(93p_1 - p_2 - 4440)$
	$\mathcal{Z}_3 = P_1^6 P_2^9 - 39 P_1^3 P_2^{12} - 96 P_2^{15}$	$N_3 = 19683(2199p_1 - 66p_2 + p_3 - 53320)$
	$\mathcal{Z}_4 = P_1^3 P_2^{12} - 12 P_2^{15}$	$N_4 = -531441 (10525p_1 - 687p_2 + 39p_3 - p_4 - 140760)$
	$\mathcal{Z}_5 = P_2^{15}$	$N_5 = -14348907 (3420p_1 + 1264p_2 + 96p_3 + 12p_4 - p_5 + 41184)$
	16. $\mathcal{E}_1[A_{2,1}^{\otimes 64}]$ $\mathcal{Z}_0 = P_1^{16} - 128P_1^{13}P_2^3 + 5248P_1^{10}P_2^6 - 75648P_1^7P_2^9 + 292992P_1^4P_2^{12} - 127488P_1P_2^{15}$	$i^{\frac{1}{3}}(i^5 + N_1i^4 + N_2i^3 + N_3i^2 + N_4i + N_5)$
	$\mathcal{Z}_1 = P_1^{13} P_2^3 - 101 P_1^{10} P_2^6 + 2791 P_1^7 P_2^9 - 19029 P_1^4 P_2^{12} + 10788 P_1 P_2^{15}$	$N_1 = 27 (p_1 - 128)$
	$\mathcal{Z}_2 = P_1^{10} P_2^6 - 74 P_1^7 P_2^9 + 1063 P_1^4 P_2^{12} - 1776 P_1 P_2^{15}$	$N_2 = -729(101p_1 - p_2 - 5248)$
	$\mathcal{Z}_3 = P_1^7 P_2^9 - 47 P_1^4 P_2^{12} + 64 P_1 P_2^{15}$	$N_3 = 19683(2791p_1 - 74p_2 + p_3 - 75648)$
	$\mathcal{Z}_4 = P_1^4 P_2^{12} - 20 P_1 P_2^{15}$	$N_4 = -531441 (19029p_1 - 1063p_2 + 47p_3 - p_4 - 292992)$
	$\mathcal{Z}_5 = P_1 P_2^{15}$	$N_5 = 14348907 (10788p_1 - 1776p_2 + 64p_3 - 20p_4 + p_5 - 127488)$

Table 5: Characters of one-character extensions of tensor products of $A_{2,1}$ (and $E_{6,1}$) written in a basis of S-invariant polynomials

3.3 $C = G_{2,1}$

 $G_{2,1}$ is a [2,0] CFT with central charge $\frac{14}{5}$. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{2}{5}$ $\frac{2}{5}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{7}{60}} \left(1 + 14q + 42q^2 + 140q^3 + 350q^4 + 840q^5 + 1827q^6 + 3858q^7 + 7637q^8 + 14756q^9 + \ldots \right)
$$

$$
\chi_1 = q^{\frac{17}{60}} \left(7 + 34q + 119q^2 + 322q^3 + 819q^4 + 1862q^5 + 4025q^6 + 8218q^7 + 16177q^8 + \ldots \right) \tag{39}
$$

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{2\chi_0 + (1 + \sqrt{5})\chi_1}{\sqrt{2(\sqrt{5} + 5)}}, \qquad S(\chi_1) = \frac{(1 + \sqrt{5})\chi_0 - 2\chi_1}{\sqrt{2(\sqrt{5} + 5)}} \tag{40}
$$

The allowed values of k and N can be read from entry 3. of table [2.](#page-10-1) The allowed tensor powers N are multiples of 20: $N = 20, 40, 60, 80...$ and the corresponding $c_{(1)}$'s are multiples of 56: $c_{(1)} = 56, 112, 168, 224 \ldots$

We start the analysis with the smallest number of tensor powers i.e. the $N = 20, c_{(1)} = 56$ case. We discover the first S-invariant polynomial P_1 here. It is a non-constant S-invariant polynomial which we call P_1 . Then we work on the next higher number of tensor powers, the $N = 40, c_{(1)} = 112$ case. For this, we find that the character is P_1^2 . The second S-invariant polynomial, which is also a non-constant S-invariant polynomial is discovered when working on the $N = 60, c_{(1)} = 168$ case. It is a degree-12 polynomial of the characters. We collect both the S-invariant polynomials for $G_{2,1}$ and their q -series here:

$$
P_1(\chi_0, \chi_1) = \chi_0^{20} - 228\chi_0^{15}\chi_1^5 + 494\chi_0^{10}\chi_1^{10} + 228\chi_0^5\chi_1^{15} + \chi_1^{20} = j^{\frac{1}{3}}(j^2 - 1456j - 3670016)
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^{11}\chi_1 + 11\chi_0^6\chi_1^6 - \chi_0\chi_1^{11} = 7j - 4096
$$
\n(41)

There is a similarity between all the three CFTs we have studied so far viz. $A_{1,1}$, $A_{2,1}$ and $G_{2,1}$. In each of them, one discovers P_1 for the smallest $c_{(1)}$ and P_2 while working with three times the smallest $c_{(1)}$. We collect the results for $G_{2,1}$ in table [6.](#page-21-1)

#	N	$c_{(1)}$	$P_{c_{(1)}=8k}^{G_{2,1}}(\chi_0,\chi_1)$	q -series
	1. $\mathcal{E}_1[G_{2,1}^{\otimes 20}]$	56	$\mathcal{Z}_0=P_1$	$j^{\frac{1}{3}}(j^2+N_1j+N_2)$ $N_1=-1456, N_2=-3670016$
	2. $\mathcal{E}_1[G_{2.1}^{\otimes 40}]$	112	$\mathcal{Z}_0 = P_1^2$	$j^{\frac{2}{3}}(j^4+N_1j^3+N_2j^2+N_3j+N_4)$ $N_1=-2912$
				$N_2 = -5220096$, $N_3 = 10687086592$, $N_4 = 13469017440256$
	3. $\mathcal{E}_1[G_{2.1}^{\otimes 60}]$		$168 \mid \mathcal{Z}_0 = P_1^3 + 684 P_2^5$	$j^7 + N_1 j^6 + N_2 j^5 + N_3 j^4 + N_4 j^3 + N_5 j^2 + N_6 j + N_7$
			$\mathcal{Z}_1 = P_2^5$	$N_1 = -4368, N_2 = 7(2401p_1 + 977964), N_3 = -1003520(49p_1 + 4643)$
				$N_4 = 8220835840 (7p_1 + 6864), N_5 = -6734508720128 (5p_1 + 12156)$
				$N_6 = 1970324836974592(5p_1 - 21668)$
				$N_7 = -1152921504606846976 (p_1 + 684)$

Table 6: Characters of one-character extensions of tensor products of $G_{2,1}$ written in a basis of S-invariant polynomials

3.4 $C = D_{4,1}$

 $D_{4,1}$ is a [2,0] CFT with central charge 4. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{1}{2}$. Let us note the q-series for the characters :

$$
\chi_0 = q^{-\frac{1}{6}} \left(1 + 28q + 134q^2 + 568q^3 + 1809q^4 + 5316q^5 + 13990q^6 + 34696q^7 + 80724q^8 + \ldots \right)
$$

\n
$$
\chi_1 = q^{\frac{1}{3}} \left(8 + 64q + 288q^2 + 1024q^3 + 3152q^4 + 8704q^5 + 22208q^6 + 53248q^7 + \ldots \right) \tag{42}
$$

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + 3\chi_1}{2}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_1}{2}.
$$
 (43)

The allowed values of k and N can be read from entry 4. of table [2.](#page-10-1) Here, every possible $c_{(1)} = 8k$ is possible and the corresponding number of tensor product factors is $N = 2k$.

We will be brief in our description of the results here. The analysis proceeds very similar to $A_{1,1}$. We discover the first S-invariant polynomial when we are working with $c_{(1)} = 8$. This is a quadratic non-constant S-invariant polynomial. Just as for the $A_{1,1}$ case in section [4.1,](#page-27-1) we discover the second S-invariant polynomial when working $c_{(1)} = 24$. This is a constant S-invariant polynomial of degree-3. The two S-invariant polynomials for $D_{4,1}$ are:

$$
P_1(\chi_0, \chi_1) = \chi_0^2 + 3\chi_1^2 = j^{\frac{1}{3}}
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^2 \chi_1 - \chi_1^3 = 8
$$
\n(44)

In table [7,](#page-23-0) we collect all the results for $k = 1, 2, \ldots 16$. For $c_{(1)} = 8k$ the character has $\left[\frac{k}{3}\right]$ arbitrary constants (the *p*'s) and there are $1 + \left[\frac{k}{3}\right]$ *S*-invariant summands the *Z*'s. Each summand, when expressed in terms of the characters χ_0, χ_1 are polynomials of degree 2k. But when expressed in terms of the S-invariant polynomials [\(44\)](#page-22-0), they are polynomials of much smaller degree.. Because of this, we can report in a compact manner, computations upto $c_{(1)} = 128$ for which we have polynomials of degree 16 only.

Table 7: Characters of one-character extensions of tensor products of $D_{4,1}$ written in a basis of S-invariant polynomials

3.5 $C = F_{4,1}$

 $F_{4,1}$ is a [2,0] CFT with central charge $\frac{26}{5}$. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{3}{5}$ $\frac{3}{5}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{13}{60}} \left(1 + 52q + 377q^2 + 1976q^3 + 7852q^4 + 27404q^5 + 84981q^6 + 243230q^7 + 649454q^8 + \ldots \right)
$$

$$
\chi_1 = q^{\frac{23}{60}} \left(26 + 299q + 1702q^2 + 7475q^3 + 27300q^4 + 88452q^5 + 260650q^6 + 714727q^7 + \ldots \right) \tag{45}
$$

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{2\chi_0 + (1 + \sqrt{5})\chi_1}{\sqrt{2(\sqrt{5} + 5)}}, \qquad S(\chi_1) = \frac{(1 + \sqrt{5})\chi_0 - 2\chi_1}{\sqrt{2(\sqrt{5} + 5)}} \tag{46}
$$

The allowed values of k and N can be read from entry 5. of table [2.](#page-10-1) The allowed tensor powers N are multiples of 20: $N = 20, 40, 60, 80...$ and the corresponding $c_{(1)}$'s are multiples of 104: $c_{(1)} = 104, 208, 312, 416 \ldots$

Before we try to solve for the characters of one-character extensions of the tensor products of the $F_{4,1}$ CFT, we note from table [2,](#page-10-1) from entries 3. and 5., that the N values for $F_{4,1}$ is identical to those of $G_{2,1}$. So we are looking at the monomials of the same degree. Then we note that the h_1 for both these CFT have 5 in the denominator. What this means is that a monomial which solves the equations [\(7\)](#page-8-2) and [\(8\)](#page-8-3) for $G_{2,1}$ also solves the equations for $F_{2,1}$ (but for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) for $F_{4,1}$ for a given N is exactly the same as the ansatz character for $G_{2,1}$.

Now consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for both the $F_{4,1}$ and $G_{2,1}$ CFTs are identical [\(40\)](#page-20-2), [\(46\)](#page-24-1), we get the same set of linear equations and hence the same solutions as we had obtained for $G_{2,1}$. We have the same two S-invariant polynomials but they compute (using (45)) to different q-series :

$$
P_1(\chi_0, \chi_1) = \chi_0^{20} - 228\chi_0^{15}\chi_1^5 + 494\chi_0^{10}\chi_1^{10} + 228\chi_0^5\chi_1^{15} + \chi_1^{20}
$$

= $j^{\frac{1}{3}}(j^4 - 2184j^3 + 1086696j^2 - 2815351344j + 402462867456)$ (47)

$$
P_2(\chi_0, \chi_1) = \chi_0^{11}\chi_1 + 11\chi_0^6\chi_1^6 - \chi_0\chi_1^{11} = 26j^2 - 23517j - 2985984
$$

We will give the results for $F_{4,1}$ in table [8.](#page-25-1) Note that the first two columns of tables [6](#page-21-1) and [8](#page-25-1) are identical.

	2. $\mathcal{E}_1[F_{4,1}^{\otimes 40}]$ 208 $\mathcal{Z}_0 = P_1^2$	$i^{\frac{2}{3}}(i^8+N_1i^7+N_2i^6+N_3i^5+N_4i^4+N_5i^3+N_6i^2+N_7i+N_8)$ $N_1 = -4368$, $N_2 = 6943248$, $N_3 = -10377390816$
		$N_4 = 14283288601920, \quad N_5 = -7876819893286656, \quad N_6 = 8800912766588537088$
		$N_7 = -2266148749604686921728, \quad N_8 = 161976359680905823911936$
	3. $\mathcal{E}_1[F_4^{860}]$ 312 $\mathcal{Z}_0 = P_1^3 + 684P_2^5$	$N_1^{13} + N_1 i^{12} + N_2 i^{11} + N_3 i^{10} + N_4 i^9 + N_5 i^8 + N_6 i^7 + N_7 i^6 + N_8 i^5$
	$\mathcal{Z}_1 = P_2^5$	$+N_9i^4+N_{10}i^3+N_{11}i^2+N_{12}i+N_{13}$
		$N_1 = -6552$, $N_2 = 17569656$, $N_3 = 208(57122p_1 - 120079917)$
		$N_4 = -28080 (1913587p_1 - 727880976), \quad N_5 = 17740944 (5094505p_1 - 554277348)$
		$N_6 = -79834248 (792100465p_1 - 379310231988)$, $N_7 = 538881174 (14546868115p_1 - 128180655844668)$
		$N_8 = 21824687547 (400892800009p_1 + 2226113564872716)$, $N_9 = -4022726408663040 (223797971p_1 + 7800667315497)$
		$N_{10} = -177952543594745167872 (4686985p_1 - 53539931088),$ $N_{11} = -4541567931053090216607744 (30145p_1 + 321850188)$
		$N_{12} = -27903393368390186290837979136(335p_1 - 2107116)$, $N_{13} = -237376313799769806328950291431424(p_1 + 684)$

Table 8: Characters of one-character extensions of tensor products of $F_{4,1}$ written in a basis of S-invariant polynomials

3.6 $C = E_{6.1}$

 $E_{6,1}$ is a [2,0] CFT with central charge 6. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{2}{3}$ $\frac{2}{3}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{1}{4}} \left(1 + 78q + 729q^2 + 4382q^3 + 19917q^4 + 77274q^5 + 264644q^6 + 827388q^7 + 2400597q^8 + \ldots \right)
$$

\n
$$
\chi_1 = q^{\frac{5}{12}} \left(27 + 378q + 2484q^2 + 12312q^3 + 49896q^4 + 177660q^5 + 570807q^6 + 1695762q^7 + \ldots \right) \tag{48}
$$

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + 2\chi_1}{\sqrt{3}}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_1}{\sqrt{3}}.
$$
 (49)

The allowed values of k and N can be read from entry 6. of table [2.](#page-10-1) Here, only $c_{(1)} = 24k$ is possible and the corresponding number of tensor product factors is $N = 4k$.

Before we try to solve for the characters of one-character extensions of the tensor products of the $E_{6,1}$ CFT, we note from table [2,](#page-10-1) from entries 2. and 6., that the N values for $E_{6,1}$ is identical to those of $A_{2,1}$. This is because they have the same value for $\frac{8n}{\gcd(m,8)}$. Hence we are looking at the monomials of the same degree. Then we note that the h_1 for both these CFT have 3 in the denominator. What this means is that a monomial which solves the equations [\(7\)](#page-8-2) and [\(8\)](#page-8-3) for $A_{2,1}$ also solves the equations for $E_{6,1}$ (but maybe for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) for $E_{6,1}$ for a given N is exactly the same as the ansatz character for $A_{2,1}$.

Now consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for both the $A_{2,1}$ and $E_{6,1}$ CFTs are identical (37) , (49) , we get the same set of linear equations and hence the same solutions as we had obtained for $A_{2,1}$. We have the same two S-invariant polynomials but they compute (using (48)) to different *q*-series :

$$
P_1(\chi_0, \chi_1) = \chi_0^4 + 8\chi_0 \chi_1^3 = j - 432
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^3 \chi_1 - \chi_1^4 = 27j^{\frac{1}{3}}
$$
\n(50)

The results for $E_{6,1}$ can be read off from the first two columns of table [5](#page-20-1) after translating the first column: $A_{2,1}$ is replaced by $E_{6,1}$.

3.7 $C = E_{7,1}$

 $E_{7,1}$ is a [2,0] CFT with central charge 7. Let us denote it's two characters by χ_0, χ_1 , the latter is the non-identity character with conformal dimension $h_1 = \frac{3}{4}$ $\frac{3}{4}$. Let us note the *q*-series for the characters :

$$
\chi_0 = q^{-\frac{7}{24}}(1+133q+1673q^2+11914q^3+63252q^4+278313q^5+1070006q^6+3711557q^7+11862767q^8+\ldots)
$$

\n
$$
\chi_1 = q^{\frac{11}{24}}(56+968q+7504q^2+42616q^3+194768q^4+772576q^5+2742040q^6+8933456q^7+\ldots)
$$
\n(51)

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + \chi_1}{\sqrt{2}}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_1}{\sqrt{2}}.
$$
 (52)

The allowed values of k and N can be read from entry 7. of table [2.](#page-10-1) Here, every possible $c_{(1)} = 56k$ is possible and the corresponding number of tensor product factors is $N = 8k$.

Before we try to solve for the characters of one-character extensions of the tensor products of the $E_{7,1}$ CFT, we note from table [2,](#page-10-1) from entries 1. and 7., that the N values for $E_{7,1}$ is identical to those of $A_{1,1}$. This is because they have the same value for $\frac{8n}{\gcd(m,8)}$. Hence we are looking at the monomials of the same degree. Then we note that the h_1 for both these CFT have 4 in the denominator. What this means is that a monomial which solves the equations [\(7\)](#page-8-2) and [\(8\)](#page-8-3) for $A_{1,1}$ also solves the equations for $E_{7,1}$ (but maybe for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) for $E_{7,1}$ for a given N is exactly the same as the ansatz character for $A_{1,1}$.

Now consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for both the $A_{1,1}$ and $E_{7,1}$ CFTs are identical [\(20\)](#page-12-3), [\(52\)](#page-26-1), we get the same set of linear equations and hence the same solutions as we had obtained for $A_{1,1}$. We have the same two S-invariant polynomials but they compute (using (51)) to different q-series :

$$
P_1(\chi_0, \chi_1) = \chi_0^8 + 14 \chi_0^4 \chi_1^4 + \chi_1^8 = j^{\frac{1}{3}} (j^2 - 672j - 145152)
$$

\n
$$
P_2(\chi_0, \chi_1) = \chi_0^5 \chi_1 - \chi_0 \chi_1^5 = 56j - 3456
$$
\n(53)

The results for $E_{7,1}$ can be read off from the first two columns of table [4](#page-17-0) after translating the first column: $A_{1,1}$ is replaced by $E_{7,1}$.

To conclude this section which only studied two-character WZW CFTs, we note that in every case, we have two S-invariant polynomials $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$ $((34), (38), (41), (44), (47), (50), (53))$. In every case, the existence of these S-invariant polynomials have aided our computations heavily. Take the character of any of the $\mathcal{E}_1[\mathcal{C}^{\otimes N}]$ in any of the tables of this section [2.](#page-7-0) If the S-invariant polynomials did not exist, we would have them as polynomials of the basic characters χ_0, χ_1 , their degree would be much higher, sometimes as high as 8 times.

Apart from contributing to the compactness of expressions, one asks why there are only two Sinvariant polynomials. Perhaps there is an answer from polynomial algebra. Having to do with the fact that the basic characters, which are two in number, form some kind of a basis of generators, not just for the entire polynomial algebra but also for the sub-algebra of S-invariant polynomials. Now, the polynomials P_1 and P_2 , also seem to form a basis of generators for the sub-algebra of S-invariant polynomials. And since any two sets of bases should have the same cardinality, we perhaps have an explanation for the existence of two S-invariant polynomials P_1 and P_2 for two-character theories. This kind of argument will perhaps gain more strength after studying three-character CFTs in section [4,](#page-27-0) where we find three S-invariant polynomials P_1 , P_2 and P_3 for every three-character CFT studied there.

4 [3, 0] CFTs : Tensor Products and One-character Extensions

In this section, we study some of the (but an infinite number of) [3, 0] WZW CFTs. We implement the procedure described in section [2](#page-7-0) for each of the following $[2, 0]$ WZW CFTs. We start with the $A_{4,1}$ CFT in [4.1.](#page-27-1) Then we take on the interesting example of the Ising CFT a.k.a $\mathcal{M}(4,3)$ in section [4.2.](#page-29-0) Then we study the infinite class of CFTs $D_{r,1}$ in section [4.3.](#page-32-0) We are able to solve for an infinite number of CFTs. This is because, the infinite class of $D_{r,1}$ CFTs can be separated into four infinite classes of CFTs. These are (i) when r is a odd number, (ii) r is 2 times an odd number, (iii) r is a 4 times an odd number, (iv) r is a multiple of 8. We study these separately in sections [4.3.1,](#page-32-1) [4.3.2,](#page-34-0) [4.3.3,](#page-35-0) [4.3.4.](#page-37-0) All CFTs in any of these four classes of CFTs have a common answer which we will explain below. We thus are able to solve for all infinite CFTs $D_{r,1}$.

$$
\textbf{4.1} \quad \mathcal{C} = A_{4,1}
$$

 $A_{4,1}$ is a [3,0] CFT with central charge 4. Let us denote it's three characters by χ_0, χ_1, χ_2 , the latter two are the non-identity characters with conformal dimensions $h_1 = \frac{2}{5}$ $\frac{2}{5}$, $h_2 = \frac{3}{5}$ $\frac{3}{5}$ respectively. Let us note the q -series for the characters :

$$
\chi_0 = q^{-\frac{1}{6}}(1 + 24q + 124q^2 + 500q^3 + 1625q^4 + 4752q^5 + 12524q^6 + 31000q^7 + 72250q^8 + 161000q^9 + \ldots)
$$

\n
$$
\chi_1 = q^{\frac{7}{30}}(5 + 50q + 220q^2 + 820q^3 + 2525q^4 + 7070q^5 + 18125q^6 + 43780q^7 + 100030q^8 + 219230q^9 + \ldots)
$$

\n
$$
\chi_2 = q^{\frac{13}{30}}(10 + 65q + 300q^2 + 1025q^3 + 3140q^4 + 8565q^5 + 21760q^6 + 51785q^7 + 117500q^8 + 255110q^9 + \ldots)
$$

\n(54)

and the modular S-transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0}{\sqrt{5}} + \frac{2\chi_1}{\sqrt{5}} + \frac{2\chi_2}{\sqrt{5}}
$$

\n
$$
S(\chi_1) = \frac{\chi_0}{\sqrt{5}} + \frac{(\sqrt{5} - 1)\chi_1}{2\sqrt{5}} - \frac{(\sqrt{5} + 1)\chi_2}{2\sqrt{5}}
$$

\n
$$
S(\chi_2) = \frac{\chi_0}{\sqrt{5}} - \frac{(\sqrt{5} + 1)\chi_1}{2\sqrt{5}} + \frac{(\sqrt{5} - 1)\chi_2}{2\sqrt{5}}
$$
(55)

The allowed values of k and N can be read from entry 2. of table [3.](#page-11-0) Here, every possible $c_{(1)} = 8k$ is possible and the corresponding number of tensor product factors is $N = 2k$.

We follow all the steps laid out in section [2.](#page-7-0) We solve the set of equations [\(7\)](#page-8-2), [\(8\)](#page-8-3) and construct the ansatz character (10) . We cannot come up with a closed formula for b the number of monomials with q series commensurate to the identity character. But we can readily compute them. And we can set up the overdetermined system of linear equations and find solutions. The answers are tabulated in table [9.](#page-29-1)

We start with smallest possible $c_{(1)}$. This is the $N = 2, c_{(1)} = 8$ case, whose solution gives us our first S-invariant polynomial $P_1(\chi_0, \chi_1, \chi_2)$. This is a non-constant quadratic polynomial. The solution for the $N = 4, c_{(1)} = 16$ is simply $P_1(\chi_0, \chi_1, \chi_2)^2$. While solving the $N = 6, c_{(1)} = 24$, we discover the second S-invariant polynomial $P_2(\chi_0, \chi_1, \chi_2)$ which is of degree-6 and computes (using [\(54\)](#page-27-2)) to a constant. Then, while solving the $N = 10c_{(1)} = 40$, we discover the third S-invariant polynomial $P_3(\chi_0, \chi_1, \chi_2)$ which is of degree-10 and computes (using [\(54\)](#page-27-2)) to a constant. We collect all the three S-invariant polynomials for $A_{4,1}$ below :

$$
P_1(\chi_0, \chi_1, \chi_2) = \chi_0^2 + 4\chi_1 \chi_2 = j^{\frac{1}{3}}
$$

\n
$$
P_2(\chi_0, \chi_1, \chi_2) = \chi_0^4 \chi_1 \chi_2 - \chi_0^2 \chi_1^2 \chi_2^2 - \chi_0 \chi_1^5 - \chi_0 \chi_2^5 + 2\chi_1^3 \chi_2^3 = 50
$$

\n
$$
5 P_3(\chi_0, \chi_1, \chi_2) = 5 \chi_0^6 \chi_1^2 \chi_2^2 - 4\chi_0^5 \chi_1^5 - 4\chi_0^5 \chi_2^5 - 10\chi_0^4 \chi_1^3 \chi_2^3 + 10\chi_0^3 \chi_1^6 \chi_2 + 10v\chi_0^3 \chi_1 \chi_2^6 + 5\chi_0^2 \chi_1^4 \chi_2^4
$$

\n
$$
-10\chi_0 \chi_1^7 \chi_2^2 - 10\chi_0 \chi_1^2 \chi_2^7 + \chi_1^{10} + 6\chi_1^5 \chi_2^5 + \chi_2^{10} = 0
$$

\n(56)

We have tabulated the results for $\mathcal{E}_1[A_{4,1}^{\otimes N}]$ for $N \leq 24$ and $c_{(1)} \leq 96$ in table [9.](#page-29-1) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

7. $\mathcal{E}_1[A_4]$ 56	$Z_0 = P_1^7 - 28P_1^4P_2 + 35P_1^2P_3 + 49P_1P_2^2$	$i^{\frac{1}{3}}(i^2+N_1i+N_2)$
	$Z_1 = P_1^4 P_2 - \frac{5P_1^2 P_3}{4} - \frac{55P_1 P_2^2}{4}$	$N_1 = 50(p_1 - 28)$
	$\mathcal{Z}_2 = P_1 P_2^2$	$N_2 = -625(55p_1 - 4p_2 - 5p_3 - 196)$
	$Z_3 = P_1 \left(\frac{5P_2^2}{4} - \frac{5P_1P_3}{4} \right)$	
8. $\mathcal{E}_1[A_4^{\otimes 16}]$ 64	$\mathcal{Z}_0 = P_1^8 - 32P_1^5P_2 + 40P_1^3P_3 + 120P_1^2P_2^2 - 160P_2P_3$	$j^{\frac{2}{3}}(j^2+N_1j+N_2)$
	$Z_1 = P_1^5 P_2 - \frac{5P_1^3 P_3}{4} - \frac{71P_1^2 P_2^2}{4} - 23P_2 P_3$	$N_1 = 50 (p_1 - 32)$
	$Z_2 = P_1^2 P_2^2 - 6P_2 P_3$	$N_2 = -625(71p_1 - 4p_2 - 5p_3 - 480)$
	$Z_3 = -\frac{1}{4}5P_1^3P_3 + \frac{5P_1^2P_2^2}{4} + 5P_2P_3$	
	$\mathcal{Z}_4 = P_2 P_3$	
9. $\mathcal{E}_1[A_{4.1}^{\otimes 18}]$ 72	$\mathcal{Z}_0=P_1^9-36P_1^6P_2+45P_1^4P_3+207P_1^3P_2^2-45P_1P_2P_3-183P_2^3$	$i^3 + N_1 i^2 + N_2 i + N_3$
	$Z_1 = P_1^6 P_2 - \frac{5P_1^4 P_3}{4} - \frac{87P_1^3 P_2^2}{4} + \frac{165P_1 P_2 P_3}{4} - \frac{97P_2^3}{4}$	$N_1 = 50 (p_1 - 36)$
	$Z_2 = P_1^3 P_2^2 - \frac{5P_1P_2P_3}{2} - \frac{15P_2^3}{2}$	$N_2 = -625(87p_1 - 4p_2 - 5p_3 - 828)$
	$Z_3 = -\frac{1}{4}5P_1^4P_3 + \frac{5P_1^3P_2^2}{4} + \frac{55P_1P_2P_3}{4} - \frac{35P_2^3}{4}$	$N_3 = -31250 (97p_1 + 30p_2 + 35p_3 - 4p_5 - 5p_6 + 732)$
	$\mathcal{Z}_4 = P_2^3$	
	$\mathcal{Z}_5 = \frac{5P_2^3}{4} - \frac{5P_1P_2P_3}{4}$	
10. $\mathcal{E}_1[A_4^{\otimes 20}]$ 80	$\mathcal{Z}_0=P_1^{10}-40P_1^7P_2+50P_1^5P_3+310P_1^4P_2^2-250P_1^2P_2P_3-270P_1P_2^3-1280P_3^2$	$i^{\frac{1}{3}}(i^3+N_1i^2+N_2i+N_3)$
	$\mathcal{Z}_1 = P_1^7 P_2 - \frac{5 P_1^5 P_3}{4} - \frac{103 P_1^4 P_2^2}{4} + \frac{205}{4} P_1^2 P_2 P_3 + \frac{87 P_1 P_2^3}{4} - 310 P_3^2$	$N_1 = 50 (p_1 - 40)$
	$Z_2 = P_1^4 P_2^2 - \frac{5}{3} P_1^2 P_2 P_3 - \frac{23 P_1 P_2^3}{3} - 45 P_3^2$	$N_2 = -625(103p_1 - 4p_2 - 5p_3 - 1240)$
	$Z_3 = -\frac{1}{4}5P_1^5P_3 + \frac{5P_1^4P_2^2}{4} + \frac{75}{4}P_1^2P_2P_3 - \frac{55P_1P_2^3}{4} - 15P_3^2$	$N_3 = 31250(87p_1 - 46p_2 - 55p_3 + 4p_4 + 5p_5 - 1080)$
	$Z_4 = P_1 P_2^3 - P_3^2$	
	$\mathcal{Z}_5 = -\frac{5}{4}P_1^2P_2P_3 + \frac{5P_1P_2^3}{4} + 5P_3^2$	
	$\mathcal{Z}_6 = P_3^2$	
11. $\mathcal{E}_1[A_4^{\otimes 22}]$ 88	$\mathcal{Z}_0=P_1^{11}-44P_1^8P_2+55P_1^6P_3+429P_1^5P_2^2-495P_1^3P_2P_3-605P_1^2P_2^3+1760P_1P_3^2-2200P_2^2P_3$	$i^{\frac{2}{3}}(i^3+N_1i^2+N_2i+N_3)$
	$\mathcal{Z}_1 = P_1^8 P_2 - \tfrac{5 P_1^6 P_3}{4} - \tfrac{119 P_1^5 P_2^2}{4} + \tfrac{245}{4} P_1^3 P_2 P_3 + \tfrac{335 P_1^2 P_3^3}{4} + 380 P_1 P_3^2 - 590 P_2^2 P_3$	$N_1 = 50 (p_1 - 44)$
	$Z_2 = P_1^5 P_2^2 - \frac{5}{2} P_1^3 P_2 P_3 - \frac{31 P_1^2 P_2^3}{2} + 60 P_1 P_3^2 - 85 P_2^2 P_3$	$N_2 = -625(119p_1 - 4p_2 - 5p_3 - 1716)$
	$\mathcal{Z}_3 = -\frac{1}{4}5P_1^6P_3 + \frac{5P_1^5P_2^2}{4} + \frac{95}{4}P_1^3P_2P_3 - \frac{75P_1^2P_3^3}{4} + \frac{85P_1P_3^2}{2} - \frac{155P_2^2P_3}{2}$	$N_3 = 31250 (335p_1 - 62p_2 - 75p_3 + 4p_4 + 5p_5 - 2420)$
	$Z_4 = P_1^2 P_2^3 + \frac{5P_1P_3^2}{4} - \frac{25P_2^2P_3}{4}$	
	$\mathcal{Z}_5=-\frac{1}{4}5P_1^3P_2P_3+\frac{5P_1^2P_3^3}{4}+\frac{5P_1P_3^2}{2}+\frac{5P_2^2P_3}{2}$	
	$Z_6 = 2P_2^2P_3 - P_1P_2^2$	
	$Z_7 = \frac{5P_2^2P_3}{4} - \frac{5P_1P_3^2}{4}$	
	12. $\mathcal{E}_1[A_A^{(2)24}]$ 96 $\mathcal{Z}_0 = P_1^{12} - 48P_1^9P_2 + 60P_1^7P_3 + 564P_1^6P_2^2 - 780P_1^4P_2P_3 - 1252P_1^3P_2^3 + 75P_1^2P_3^2 + 1290P_1P_2^2P_3 - 789P_2^4P_3^2$	$i^4 + N_1 i^3 + N_2 i^2 + N_3 i + N_4$
	$\mathcal{Z}_1 = P_1^9 P_2 - \frac{5 P_1^7 P_3}{4} - \frac{135 P_1^6 P_2^2}{4} + \frac{285}{4} P_1^4 P_2 P_3 + \frac{647 P_1^3 P_2^3}{4} - \frac{575 P_1^2 P_3^2}{16} + \frac{335}{8} P_1 P_2^2 P_3 - \frac{4159 P_2^4 P_3^2}{16}$	$N_1 = 50 (p_1 - 48)$
	$\mathcal{Z}_2=P_1^6P_2^2-\tfrac{5}{2}P_1^4P_2P_3-\tfrac{39P_1^3P_2^3}{2}+\tfrac{25P_1^2P_3^2}{16}+\tfrac{375}{8}P_1P_2^2P_3-\tfrac{599P_2^4}{16}$	$N_2 = -625(135p_1 - 4p_2 - 5p_3 - 2256)$
	$Z_3 = -\frac{1}{4}5P_1^7P_3 + \frac{5P_1^6P_2^2}{4} + \frac{115}{4}P_1^4P_2P_3 - \frac{95P_1^3P_3^3}{4} - \frac{75P_1^2P_3^2}{4} + \frac{65}{2}P_1P_2^2P_3 - 50P_2^4$	$N_3 = 31250 (647p_1 - 78p_2 - 95p_3 + 4p_4 + 5p_5 - 5008)$
	$Z_4 = P_1^3 P_2^3 - \frac{15}{4} P_1 P_2^2 P_3 - \frac{21 P_2^4}{4}$	$N_4 = -390625 (4159p_1 + 599p_2 + 800p_3 + 84p_4 + 95p_5 - 16p_6 - 20p_7 - 25p_8 + 12624)$
	$\mathcal{Z}_5 = -\frac{1}{4}5P_1^4P_2P_3 + \frac{5P_1^3P_2^3}{4} + \frac{25P_1^2P_3^2}{16} + \frac{75}{8}P_1P_2^2P_3 - \frac{95P_2^4}{16}$	
	$\mathcal{Z}_6 = P_2^4$	
	$Z_7 = \frac{5P_2^4}{4} - \frac{5}{4}P_1P_2^2P_3$	
	$\mathcal{Z}_8=\frac{25P_1^2P_3^2}{16}-\frac{25}{8}P_1P_2^2P_3+\frac{25P_2^4}{16}$	

Table 9: Characters of one-character extensions of tensor products of $A_{4,1}$ written in a basis of S-invariant polynomials

4.2 $C = \mathcal{M}(4, 3)$ a.k.a the Ising CFT

The Virasoro minimal model $\mathcal{M}(4,3)$ is a $[\mathbf{3},\mathbf{0}]$ CFT with central charge $\frac{1}{2}$. Let us denote it's three characters by χ_0, χ_1, χ_2 , the latter two are the non-identity characters with conformal dimensions $h_1 = \frac{1}{16}, h_2 = \frac{1}{2}$ $\frac{1}{2}$ respectively. Let us note the q-series for the characters :

$$
\chi_0(q) = q^{-\frac{1}{48}}(1+q^2+q^3+2q^4+2q^5+3q^6+3q^7+5q^8+5q^9+7q^{10}+8q^{11}+\ldots)
$$

\n
$$
\chi_1(q) = q^{\frac{1}{24}}(1+q+q^2+2q^3+2q^4+3q^5+4q^6+5q^7+6q^8+8q^9+10q^{10}+\ldots)
$$

\n
$$
\chi_2(q) = q^{23/48}(1+q+q^2+q^3+2q^4+2q^5+3q^6+4q^7+5q^8+6q^9+8q^{10}+\ldots)
$$
\n(57)

and the modular S -transformations for the characters :

$$
S(\chi_0) = \frac{\chi_0 + \sqrt{2}\chi_1 + \chi_2}{2}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_2}{\sqrt{2}}, \qquad S(\chi_2) = \frac{\chi_0 - \sqrt{2}\chi_1 + \chi_2}{2} \tag{58}
$$

The allowed values of k and N can be read from entry 1. of table [3.](#page-11-0) Here, every possible $c_{(1)} = 8k$ is possible and the corresponding number of tensor product factors is $N = 16k$. This CFT is one of the hardest in terms of degree of polynomials required : for the smallest $c_{(1)} = 8$, we need to work with homogenous polynomials of degree 16 in 3 variables, which has 153 monomials.

We follow all the steps laid out in section [2.](#page-7-0) We solve the set of equations [\(7\)](#page-8-2), [\(8\)](#page-8-3) and construct the ansatz character (10) . We cannot come up with a closed formula for b the number of monomials with q series commensurate to the identity character. But we can readily compute them. And we can set up the overdetermined system of linear equations and find solutions. The answers are tabulated in table [10.](#page-31-0)

We start with smallest possible $c_{(1)}$. This is the $N = 16$, $c_{(1)} = 8$ case. Even at this stage, we have to contend with two different S-invariant polynomials (P_2, P_3) and $P_{c_{(1)}=8k}^{\mathcal{M}(4,3)}$ $c_{(1)}=8k(\chi_0,\chi_1,\chi_2)$ is a combination $\mathcal{Z}_0 + p_1 \mathcal{Z}_1$. In the next step $N = 32$, $c_{(1)} = 16$ we meet a third S-invariant polynomial (P_1) . We collect all the three S-invariant polynomials for $\mathcal{M}(4,3)$ below :

$$
P_1 = \chi_0^{23} \chi_2 - \chi_0^{21} \chi_2^3 - 21 \chi_0^{19} \chi_2^5 + 85 \chi_0^{17} \chi_2^7 - 134 \chi_0^{15} \chi_2^9 + 256 \chi_0^{14} \chi_1^8 \chi_2^2 + 70 \chi_0^{13} \chi_2^{11} + 3584 \chi_0^{12} \chi_1^8 \chi_2^4 \tau_0 \chi_0^{11} \chi_2^{13} + 16128 \chi_0^{10} \chi_1^8 \chi_2^6 - 134 \chi_0^9 \chi_2^{15} + 25600 \chi_0^8 \chi_1^8 \chi_2^8 - 256 \chi_0^7 \chi_1^{16} \chi_2 + 85 \chi_0^7 \chi_2^{17} + 16128 \chi_0^6 \chi_1^8 \chi_2^{10} - 1792 \chi_0^5 \chi_1^{16} \chi_2^3 - 21 \chi_0^5 \chi_2^{19} + 3584 \chi_0^4 \chi_1^8 \chi_2^{12} - 1792 \chi_0^3 \chi_1^{16} \chi_2^5 - \chi_0^3 \chi_2^{21} + 256 \chi_0^2 \chi_1^8 \chi_2^{14} - 256 \chi_0 \chi_1^{16} \chi_2^7 + \chi_0 \chi_2^{23} = 1 P_2 = \chi_0^{16} + 140 \chi_0^{12} \chi_2^4 + 448 \chi_0^{10} \chi_2^6 + 870 \chi_0^8 \chi_2^8 + 240 \chi_0^7 \chi_1^8 \chi_2 + 448 \chi_0^6 \chi_2^{10} + 1680 \chi_0^5 \chi_1^8 \chi_2^3 + 140 \chi_0^4 \chi_2^{12} + 1680 \chi_0^3 \chi_1^8 \chi_2^5 + 240 \chi_0 \chi_1^8 \chi_2^7 + 8 \chi_1^{16} + \chi_2^{16} = j^{-\frac{1}{3}} P_3 = \chi_0^7 \chi_2 + 7 \chi_0^5 \chi_2^3 +
$$

 P_1 is a degree-24 S-invariant polynomial which evaluates to a constant (1). P_2 is a degree-16 non-constant S-invariant polynomial. P_3 is a degree-8 S-invariant polynomial which evaluates to zero.

We have tabulated the results for $\mathcal{E}_1[\mathcal{M}(4,3)^{\otimes N}]$ for $N \leq 96$ and $c_{(1)} \leq 48$ in table [10.](#page-31-0) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

It is well-known that the character of the monster CFT is a one-character extension of 48 copies of the Ising model. Here, we can report the character of the monster CFT in terms of the characters of the Ising model. From the third entry of table [10,](#page-31-0) we have

$$
P_{c_{(1)}=24}^{\mathcal{M}(4,3)}(\chi_0,\chi_1,\chi_2) = \mathcal{Z}_0 + p_1 \mathcal{Z}_1 + p_2 \mathcal{Z}_2 + p_3 \mathcal{Z}_3 + p_4 \mathcal{Z}_4 + p_5 \mathcal{Z}_5 + p_6 \mathcal{Z}_6.
$$
 (59)

There are seven summands in [\(59\)](#page-30-0). As a q-series, this evaluates to $j + p_1 + p_4 + p_6 - 744$. This is because \mathcal{Z}_2 , \mathcal{Z}_3 and \mathcal{Z}_5 vanish. Now, it is well-known that the character of the monster CFT is j −744. To obtain the character of the monster CFT from [\(59\)](#page-30-0), we need to choose $p_1 + p_4 + p_6 = 0$. This gives us the character of the monster CFT:

$$
\begin{aligned} \chi_{\text{Monster}} \quad &= P_2^3 - 744 P_1^2 \\ &= P_2^3 - 744 \end{aligned} \tag{60}
$$

The first line is a presentation in which the homogeneity of the Ising characters is clear. The second line is in the simplest form.

Table 10: Characters of one-character extensions of tensor products of $\mathcal{M}(4,3)$ written in a basis of $S\mbox{-}\mathrm{invariant}$ polynomials

4.3 $C = D_{r,1}$

In this subsection, we study the infinite class of CFTs $D_{r,1}$ where r is a positive integer and $r \geq 3$. Each of these CFTs is a [3,0] CFT [\[27\]](#page-43-8). A $D_{r,1}$ CFT has a central charge equalling r. Let us denote it's three characters by χ_0, χ_1, χ_2 , the latter two are the non-identity characters with conformal dimensions $h_1 = \frac{r}{8}$ $\frac{r}{8}$, $h_2 = \frac{1}{2}$ $\frac{1}{2}$ respectively. Let us note the *q*-series for the characters (these formulae are sourced from chapter 14 of [\[3\]](#page-41-1)):

$$
\chi_0(q) = \frac{\theta_3^r(q) + \theta_4^r(q)}{2\eta^r}, \qquad \chi_1(q) = \frac{\theta_2^r(q)}{2\eta^r}, \qquad \chi_2(q) = \frac{\theta_3^r(q) - \theta_4^r(q)}{2\eta^r}
$$
(61)

where are the θ -constants : $\theta_i(\tau) \equiv \theta_i(0|\tau)$ with q-series given by :

$$
\theta_2(q) = 2q^{\frac{1}{8}} \prod_{n=1}^{\infty} (1 - q^n)(1 + q^n)^2, \qquad \theta_3(q) = \prod_{n=1}^{\infty} (1 - q^n)(1 + q^{n - \frac{1}{2}})^2
$$

$$
\theta_4(q) = \prod_{n=1}^{\infty} (1 - q^n)(1 - q^{n - \frac{1}{2}})^2, \qquad \eta(q) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n).
$$

The modular S-transformations for the characters is given by :

$$
S(\chi_0) = \frac{\chi_0 + 2\chi_1 + \chi_2}{2}, \qquad S(\chi_1) = \frac{\chi_0 - \chi_2}{2}, \qquad S(\chi_2) = \frac{\chi_0 - 2\chi_1 + \chi_2}{2}.
$$
 (62)

We note that the S-transformations do not have a r dependence. This means that we can hope that imposing S-invariance can be done independent of r. Before we get to imposing S-invariance, we need to construct the ansatz character (10) . This has a r-dependence. We can see this from examining entries 3, 4, 5 and 6 of table [3.](#page-11-0) We find that the number of tensor powers N depends on r and this dependence makes us divide into four (infinite) sub-classes (i) when r is a odd number, (ii) r is 2 times an odd number, (iii) r is a 4 times an odd number, (iv) r is a multiple of 8. And then it turns out that there is a common solution for all the infinite members of a sub-class. We will take up each sub-class below.

4.3.1 $C = D_{r,1}$, $r = 2p + 1$

From row [3](#page-11-0) of table 3 it is clear that all $D_{r,1}$ CFTs with $r = 2p + 1$ have the same N values. Hence for all such r, we are looking for monomials of the same degree. Then we note that h_1 and h_2 values for all $r = 2p + 1$ have the same structure, i.e. denominators are 2 and 8. What this means is that a monomial which solves the equations (7) and (8) for any odd r also solves the equations for any other odd r (but for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) is common for all D_r with $r = 2p + 1$.

Then we consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for all $D_{r,1}$ CFTs are identical, we get the same set of linear equations and hence the same solutions for all odd r . These solutions are tabulated in table [11.](#page-34-1) The solutions are given not in terms of the basic characters of the $D_{r,1}$ CFT but in terms of S-invariant polynomials. There are three of them :

$$
P_1(\chi_0, \chi_1, \chi_2) = \chi_0^8 + 14 \chi_0^4 \chi_2^4 + 112 \chi_0^3 \chi_1^4 \chi_2 + 112 \chi_0 \chi_1^4 \chi_2^3 + 16 \chi_1^8 + \chi_2^8
$$

\n
$$
P_2(\chi_0, \chi_1, \chi_2) = \chi_0^3 \chi_2 + \chi_0 \chi_2^3 - 2 \chi_1^4
$$

\n
$$
P_3(\chi_0, \chi_1, \chi_2) = \chi_0^2 \chi_1 - \chi_1 \chi_2^2 = 2^{r-1}.
$$
\n(63)

We note that one of the S-invariant polynomials is a constant S-invariant polynomial. The others have a non-trivial q -series that do depend on r .

We have tabulated the results for $\mathcal{E}_1[D_{2p+1,1}^{\otimes N}]$ for $N \leq 48$ in table [11.](#page-34-1) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

	$6.\ \left\vert \mathcal{E}_{1}[D_{(2p+1),1}^{<38}] \right\vert\left. 48(2p+1)\right\vert\left. Z_{0}=125376P_{3}^{16}-41984P_{2}^{3}P_{3}^{12}+263040P_{1}P_{2}P_{3}^{12}-66912P_{2}^{6}P_{3}^{8}+703680P_{1}P_{2}^{4}P_{3}^{8}-1440P_{1}^{3}P_{3}^{8}+88608P_{1}^{2}P_{2}^{2}P_{3}^{8}-78720P_{2}^{9}P_{3}^{4}-$
	$+3936P_1P_2^7P_3^4+28224P_1^2P_2^5P_3^4-3360P_1^3P_2^3P_3^4-672P_1^4P_2P_3^4-13360P_2^{12}-1968P_1P_2^{10}+246P_1^2P_2^8+P_1^6+336P_1^3P_2^6-84P_1^4P_2^4$
	$\mathcal{Z}_1=-2785P_3^{16}+104352P_2^3P_3^{12}-5056P_1P_2P_3^{12}+251312P_2^6P_3^8+35452P_1P_2^4P_3^8+4P_1^3P_3^8-2944P_1^2P_2^2P_3^8-24528P_2^9P_3^4$
	$+18560P_1P_2^7P_3^4-2100P_1^2P_2^5P_3^4-560P_1^3P_2^3P_3^4+4P_1^4P_2P_3^4+4808P_2^{12}-884P_1P_2^{10}+362P_1^2P_2^8-63P_1^3P_2^6-2P_1^4P_2^4+P_1^5P_2^2$
	$\mathcal{Z}_2 = 16P_3^{16} - 10496P_2^3P_3^{12} + 32P_1P_2P_3^{12} + 1696P_2^6P_3^8 - 3568P_1P_2^4P_3^8 + 24P_1^2P_2^2P_3^8 + 10112P_2^9P_3^4$
	$-1120P_1P_2^7P_3^4 - 464P_1^2P_2^5P_3^4 + 8P_1^3P_2^3P_3^4 - 1341P_2^{12} + 304P_1P_2^{10} - 38P_1^2P_2^8 - 4P_1^3P_2^6 + P_1^4P_2^4$
	$\mathcal{Z}_3 = -1384P_3^{16} - 1672P_2^3P_3^{12} - 1564P_1P_2P_3^{12} + 792P_2^6P_3^8 - 3246P_1P_2^4P_3^8 + 2P_1^3P_3^8 - 422P_1^2P_2^2P_3^8$
	$-164P_2^9P_3^4+104P_1P_2^7P_3^4-55P_1^2P_2^5P_3^4+3P_1^3P_2^3P_3^4+P_1^4P_2P_3^4$
	$\mathcal{Z}_4 = 154P_2^{12} - 9P_1P_2^{10} - 372P_3^4P_2^9 - 6P_1^2P_2^8 - 384P_1P_3^4P_2^7 - 3408P_3^8P_2^6 + P_1^3P_2^6 + 12P_1^2P_3^4P_2^5 + 60P_1P_3^8P_2^4 + 160P_3^12P_2^3$
	$\mathcal{Z}_5 = 8P_3^{16} - 1832P_2^3P_3^{12} + 12P_1P_2P_3^{12} - 1992P_2^6P_3^8 - 302P_1P_2^4P_3^8 + 6P_1^2P_2^2P_3^8 + 135P_2^9P_3^4 - 40P_1P_2^7P_3^4 + P_1^2P_2^5P_3^4 + P_1^3P_2^3P_3^4$
	$\mathcal{Z}_6 = -688P_3^{16} - 3184P_2^3P_3^{12} - 304P_1P_2P_3^{12} - 24P_2^6P_3^8 - 22P_1P_2^4P_3^8 + P_1^3P_3^8 + 8P_1^2P_2^2P_3^8$
	$\mathcal{Z}_7 = 24 P_2^{12} - 8 P_1 P_2^{10} - 320 P_3^4 P_2^9 + P_1^2 P_2^8 + 16 P_1 P_3^4 P_2^7 + 112 P_3^8 P_2^6$
	$\mathcal{Z}_8 = 40 P_2^3 P_3^{12} - 198 P_2^6 P_3^8 + 10 P_1 P_2^4 P_3^8 - 21 P_2^9 P_3^4 - P_1 P_2^7 P_3^4 + P_1^2 P_2^5 P_3^4$
	$Z_9 = 4P_3^{16} - 160P_2^3P_3^{12} + 4P_1P_2P_3^{12} - 17P_2^6P_3^8 + 6P_1P_2^4P_3^8 + P_1^2P_2^2P_3^8$
	$\mathcal{Z}_{10} = -10 P^{12}_2 + P_1 P^{10}_2 + 20 P^4_3 P^9_2, \quad \mathcal{Z}_{11} = -3 P^4_3 P^9_2 + P_1 P^4_3 P^7_2 + 14 P^8_3 P^6_2$
	$\mathcal{Z}_{12} = 8P_2^3 P_3^{12} + 4P_2^6 P_3^8 + P_1 P_2^4 P_3^8, \quad \mathcal{Z}_{13} = P_2^{12}$
	$\mathcal{Z}_{14}=2P^{16}_3+11P^{3}_2P^{12}_3+P_1P_2P^{12}_3,\quad \mathcal{Z}_{15}=P^{9}_2P^{4}_3$
	$\mathcal{Z}_{16} = P_2^6 P_3^8$, $\mathcal{Z}_{17} = P_2^3 P_3^{12}$, $\mathcal{Z}_{18} = P_3^{16}$

Table 11: Characters of one-character extensions of tensor products of $D_{2p+1,1}$ written in a basis of S-invariant polynomials

4.3.2 $C = D_{r,1}$, $r = 2(2p + 1)$

From row 4 of table [3](#page-11-0) it is clear that all $D_{r,1}$ CFTs with $r = 2(2p + 1)$ have the same N values. Hence for all such r , we are looking for monomials of the same degree. Then we note that h_1 and h_2 values for all $r = 2(2p + 1)$ have the same structure, i.e. denominators are 2 and 4. What this means is that a monomial which solves the equations (7) and (8) for any r of this class also solves the equations for any other r of this class (but for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) is common for all D_r with $r = 2(2p + 1)$.

Then we consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for all $D_{r,1}$ CFTs are identical, we get the same set of linear equations and hence the same solutions for all $r = 2(2p+1)$. These solutions are tabulated in table [12.](#page-35-1) The solutions are given not in terms of the basic characters of the $D_{r,1}$ CFT but in terms of S-invariant polynomials. There are three of them :

$$
P_1(\chi_0, \chi_1, \chi_2) = \chi_0^4 + 12 \chi_0 \chi_1^2 \chi_2 + 2 \chi_1^4 + \chi_2^4
$$

\n
$$
P_2(\chi_0, \chi_1, \chi_2) = \chi_0^2 \chi_1 - \chi_1 \chi_2^2 = 2^{r-1}
$$

\n
$$
P_3(\chi_0, \chi_1, \chi_2) = \chi_0 \chi_2 - \chi_1^2
$$
\n(64)

We note that one of the S-invariant polynomials is a constant S-invariant polynomial. The others have a non-trivial q -series that do depend on r .

We have tabulated the results for $\mathcal{E}_1[D_{2(2)}^{\otimes N}]$ $\sum_{2(2p+1),1}^{\otimes N}$ for $N \leq 24$ in table [12.](#page-35-1) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

4.3.3
$$
C = D_{r,1}, \quad r = 4(2p+1)
$$

From row 5 of table [3](#page-11-0) it is clear that all $D_{r,1}$ CFTs with $r = 4(2p + 1)$ have the same N values. Hence for all such r , we are looking for monomials of the same degree. Then we note that h_1 and h_2 values for all $r = 2(2p + 1)$ have the same structure, i.e. denominators are 2 and 2 respectively. What this means is that a monomial which solves the equations (7) and (8) for any r of this class also solves the equations for any other r of this class (but for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) is common for all D_r with $r = 4(2p + 1)$.

Then we consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for all $D_{r,1}$ CFTs are identical, we get the same set of linear equations and hence the same solutions for all $r = 4(2p+1)$. These solutions are tabulated in table [13.](#page-37-1) The solutions are given not in terms of the basic characters of the $D_{r,1}$ CFT but in terms of S-invariant polynomials. There are three of them :

$$
P_1(\chi_0, \chi_1, \chi_2) = \chi_0^2 \chi_1 - \chi_1 \chi_2^2 = 2^{r-1}
$$

\n
$$
P_2(\chi_0, \chi_1, \chi_2) = \chi_0^2 + 4 \chi_1 \chi_2 - \chi_2^2
$$

\n
$$
P_3(\chi_0, \chi_1, \chi_2) = \chi_2 - \chi_1
$$
\n(65)

We note that one of the S-invariant polynomials is a constant S-invariant polynomial. The others have a non-trivial q-series that do depend on r .

We have tabulated the results for $\mathcal{E}_1[D_{4(2)}^{\otimes N}]$ $\binom{\otimes N}{4(2p+1),1}$ for $N \leq 12$ in table [13.](#page-37-1) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

#	\boldsymbol{N}	$c_{(1)}$	$P_{c_{(1)}}^{D_{4(2p+1),1}}(\chi_0,\chi_1,\chi_2)$
1.	$\mathcal{E}_1[D_{4(2p+1),1}^{\; \otimes 2}]$	$8(2p + 1)$	$\mathcal{Z}_0 = P_2$
			$\mathcal{Z}_1 = P_3^2$
2.	$\mathcal{E}_1[D_{4(2p+1),1}^{\otimes 4}]$	$16(2p + 1)$	$\mathcal{Z}_0 = 8P_1P_3 + P_2^2 + 8P_2P_3^2$
			$Z_1 = -2P_1P_3 - P_2P_3^2$
			$Z_2 = -P_1P_3 - P_2P_3^2$
			$\mathcal{Z}_3 = P_2^4$
3.	$\mathcal{E}_1[D_{4(2p+1),1}^{\,\otimes 6}]$	$24(2p+1)$	$\mathcal{Z}_0 = P_2^3 - 9P_1^2 - 6P_1P_2P_3 + 48P_1P_3^3 + 3P_2^2P_3^2 + 48P_2P_3^4, \quad \mathcal{Z}_1 = P_1^2$
			$\mathcal{Z}_2 = P_1^2 + P_1 P_2 P_3 - 4P_1 P_3^3 - 4P_2 P_3^4$, $\mathcal{Z}_3 = P_1^2 + 2P_1 P_2 P_3 + P_2^2 P_3^2$
			$\mathcal{Z}_4 = -4P_1P_3^3 - 3P_2P_3^4$, $\mathcal{Z}_5 = -P_1P_3^3 - P_2P_3^4$, $\mathcal{Z}_6 = P_3^6$
4.			$\mathcal{E}_1[D_{4(2p+1),1}^{\otimes 8}]\left[\right.\\ \left.32(2p+1)\left.\right \right.\\ \left. \mathcal{Z}_0=P_2^4-12P_1^2P_2-128P_1^2P_3^2-8P_1P_2^2P_3-192P_1P_2P_3^3+256P_1P_3^5+4P_2^3P_3^2-64P_2^2P_3^4+256P_2P_3^6\right]$
			$\mathcal{Z}_1 = P_1^2 P_2 - 9P_1^2 P_3^2 - 14P_1 P_2 P_3^3 + 16P_1 P_3^5 - 5P_2^2 P_3^4 + 16P_2 P_3^6$
			$\mathcal{Z}_2 = P_1^2 P_2 + 4P_1^2 P_3^2 + P_1 P_2^2 P_3 + 4P_1 P_2 P_3^3 - 16P_1 P_3^5 - 16P_2 P_3^6$
			$\mathcal{Z}_3 = P_1^2 P_2 + 8P_1^2 P_3^2 + 2P_1 P_2^2 P_3 + 16P_1 P_2 P_3^3 + P_2^3 P_3^2 + 8P_2^2 P_3^4$
			$\mathcal{Z}_4 = 6P_1^2P_3^2 + 8P_1P_2P_3^3 - 8P_1P_3^5 + 3P_2^2P_3^4 - 8P_2P_3^6$
			$\mathcal{Z}_5 = 3P_1^2P_3^2 + 5P_1P_2P_3^3 - 4P_1P_3^5 + 2P_2^2P_3^4 - 4P_2P_3^6$
			$\mathcal{Z}_6 = P_1^2 P_3^2 + 2P_1 P_2 P_3^3 + P_2^2 P_3^4$, $\mathcal{Z}_7 = -6P_1 P_3^5 - 5P_2 P_3^6$
			$\mathcal{Z}_8 = -P_1 P_3^5 - P_2 P_3^6, \quad \mathcal{Z}_9 = P_3^8$

Table 13: Characters of one-character extensions of tensor products of $D_{4(2p+1),1}$ written in a basis of S-invariant polynomials

4.3.4 $C = D_{r,1}$, $r = 8 p$

From row 6 of table [3](#page-11-0) it is clear that all $D_{r,1}$ CFTs with $r = 8p$ have the same N values. Hence for all such r, we are looking for monomials of the same degree. Then we note that h_1 and h_2 values for all $r = 8p$ have the same structure, i.e. denominators are 2 and 1 respectively. What this means is that a monomial which solves the equations [\(7\)](#page-8-2) and [\(8\)](#page-8-3) for any r of this class also solves the equations for any other r of this class (but for different values of M). We thus conclude that the ansatz character [\(10\)](#page-9-0) is common for all D_r with $r = 8p$.

Then we consider the S-invariance equations [\(18\)](#page-12-2). Since the S-matrix for all $D_{r,1}$ CFTs are identical, we get the same set of linear equations and hence the same solutions for all $r = 8p$. These solutions are tabulated in table [14.](#page-38-1) The solutions are given not in terms of the basic characters of the $D_{r,1}$ CFT but in terms of S-invariant polynomials. There are three of them :

$$
P_1(\chi_0, \chi_1, \chi_2) = \chi_0^2 \chi_1 - \chi_1 \chi_2^2 = 2^{r-1}
$$

\n
$$
2P_2(\chi_0, \chi_1, \chi_2) = \chi_1^2 + \chi_2^2 - 2\chi_0 \chi_1
$$

\n
$$
P_3(\chi_0, \chi_1, \chi_2) = \chi_0 + \chi_1
$$
\n(66)

We note that one of the S-invariant polynomials is a constant S-invariant polynomial. The others have a non-trivial q-series that do depend on r .

We have tabulated the results for $\mathcal{E}_1[D_{8p,1}^{\otimes N}]$ for $N\leq 6$ in table [14.](#page-38-1) We obtain compact expressions because we have used P_1, P_2, P_3 as a basis rather than χ_0, χ_1, χ_2 .

#	N	$c_{(1)}$	$P_{c_{(1)}}^{D_{8p,1}}(\chi_0,\chi_1,\chi_2)$
1.	$\mathcal{E}_1[D_{8p,1}]$	8p	$\mathcal{Z}_0 = P_3$
2.	$\mathcal{E}_1[D_{8n,1}^{\otimes 2}]$	16p	$\mathcal{Z}_0 = P_3^2 + 2P_2$
			$\mathcal{Z}_1=-P_2$
3.	$\mathcal{E}_1[D_{8p,1}^{\otimes 3}]$	24p	$\mathcal{Z}_0 = P_2^3 + 3P_1 + 6P_2P_3$
			$\mathcal{Z}_1=P_1$
			$\mathcal{Z}_2 = -(2P_1 + 2P_2P_3)$
4.	$\mathcal{E}_1[D_{8p,1}^{\otimes 4}]$	32p	$\mathcal{Z}_0 = P_3^4 - 4P_1P_3 - 2P_2^2$
			$\mathcal{Z}_1 = P_1 P_3 - P_2^2$
			$\mathcal{Z}_2 = P_2^2$
			$\mathcal{Z}_3 = 2P_1P_3 + P_2^2 + 2P_2P_3^2$
5.	$\mathcal{E}_1[D_{8p,1}^{\otimes 5}]$	40p	$\mathcal{Z}_0 = P_3^5 - 10P_1P_2 - 5P_1P_3^2 - 10P_2^2P_3$
			$\mathcal{Z}_1 = -2P_1P_2 + P_1P_3^2 - 4P_2^2P_3$
			$\mathcal{Z}_2 = P_2^2 P_3$
			$\mathcal{Z}_3 = 2P_1P_2 + 2P_1P_3^2 + 3P_2^2P_3 + 2P_2P_3^3$
			$\mathcal{Z}_4 = 2P_1P_2 + 2P_2^2P_3$
6.	$\mathcal{E}_1[D_{8p1}^{\,\otimes 6}]$	48p	$Z_0 = P_3^6 - 69P_1^2 - 132P_1P_2P_3 - 6P_1P_3^3 + 8P_2^3 - 60P_2^2P_3^2$
			$Z_1 = -21P_1^2 - 38P_1P_2P_3 + P_1P_3^3 + 2P_2^3 - 20P_2^2P_3^2$
			$\mathcal{Z}_2 = P_1^2$
			$\mathcal{Z}_3 = 5P_1^2 + 12P_1P_2P_3 + 2P_1P_3^3 + 8P_2^2P_3^2 + 2P_2P_3^4$
			$\mathcal{Z}_4 = 6P_1^2 + 12P_1P_2P_3 - P_2^3 + 6P_2^2P_3^2$
			$\mathcal{Z}_5 = 4P_1^2 + 6P_1P_2P_3 - P_2^3 + 2P_2^2P_3^2$
			$\mathcal{Z}_6 = 4P_1^2 + 8P_1P_2P_3 + 4P_2^2P_3^2$

Table 14: Characters of one-character extensions of tensor products of $D_{8p,1}$ written in a basis of S-invariant polynomials

5 "Constant" S-invariant Polynomials

In this section, we will first collect all the constant S-invariant polynomials for various CFTs that we have discovered in this paper.

*A*_{1,1}:
$$
\chi_0^5 \chi_1 - \chi_0 \chi_1^5 = 2
$$

\n*A*_{2,1}: $\chi_0^3 \chi_1 - \chi_1^4 = 3$
\n*D*_{4,1}: $\chi_0^2 \chi_1 - \chi_1^3 = 8$
\n*A*_{4,1}: (i) $\chi_0^4 \chi_1 \chi_2 - \chi_0^2 \chi_1^2 \chi_2^2 - \chi_0 \chi_1^5 - \chi_0 \chi_2^5 + 2 \chi_1^3 \chi_2^3 = 50$
\n(ii) $5 \chi_0^6 \chi_1^2 \chi_2^2 - 4 \chi_0^5 \chi_1^5 - 4 \chi_0^5 \chi_2^5 - 10 \chi_0^4 \chi_1^3 \chi_2^3 + 10 \chi_0^3 \chi_1^6 \chi_2$
\n $+ 10 \chi_0^3 \chi_1 \chi_2^6 + 5 \chi_0^2 \chi_1^4 \chi_2^4 - 10 \chi_0 \chi_1^7 \chi_2^2 - 10 \chi_0 \chi_1^2 \chi_2^7$
\n $+ \chi_1^{10} + 6 \chi_1^5 \chi_2^5 + \chi_2^{10} = 0$
\n*M*(4,3): (i) $\chi_0^7 \chi_2 + 7 \chi_0^5 \chi_2^3 + 7 \chi_0^3 \chi_2^5 + \chi_0 \chi_2^7 - \chi_1^8 = 0$
\n(ii) $\chi_0^{23} \chi_2 - \chi_0^{21} \chi_2^3 - 21 \chi_0^{19} \chi_2^5 + 85 \chi_0^{17} \chi_2^7 - 134 \chi_0^{15} \chi_2^9$
\n $+ 256 \chi_0^{14} \chi_1^8 \chi_2^2 + 70 \chi_0^{13} \chi_2^{11} + 3584 \chi_0^{12} \chi_1^8 \chi_2^4 + 70 \chi_0^{11} \chi_2^{13}$
\n $+ 16128 \chi_0^{10} \chi_1^8 \chi_2^6 - 134$

Also, consider the following S-invariant polynomial of the $\mathcal{M}(5, 2)$ a.k.a Lee-Yang CFT.

$$
\mathcal{M}(5,2): \qquad \qquad \chi_0^{11} \chi_1 + 11 \chi_0^6 \chi_1^6 - \chi_0 \chi_1^{11} = 1 \tag{68}
$$

In the above [\(68\)](#page-39-0), we have two infinite series which evaluate to the constant 1. This polynomial relation for these particular infinite series was known to Ramanujan and is believed to be at the heart of the Rogers-Ramnujan identities. We can refer to this [\(68\)](#page-39-0) as the Rogers-Ramanujan polynomial relation. Hence, we can think of what we have obtained in this paper, collected in [\(67\)](#page-39-1), as generalizations of Rogers-Ramanujan-like polynomial relations from the Lee-Yang CFT to other CFTs. We expect this kind of generalization to exist for other CFTs as well, which we hope to investigate.

Another lesson to learn from the above polynomial relations among characters [\(67\)](#page-39-1) is the following. When we start our analysis, the character of a one-character extension of tensor product of N copies of a CFT is an element of the polynomial ring generated by the characters of the CFT. In fact it is a homogenous polynomial of degree N . But the existence of these polynomial relations means that the expressions for the character of the one-character extension can be simplified and they may not look homogenous any more. More importantly, the correct way to account for the existence of these polynomial relations is that the ideal generated by them can be used to quotient the polynomial ring of characters and the character of the one-character extension is properly an element of this quotient ring.

6 Conclusions and Future Directions

First, let us summarise our results in this paper. In section [2](#page-7-0) we set up an operational procedure to construct characters of one-character extensions of tensor products of a single CFT \mathcal{C} . The technical elements involved in the procedure was very basic and algebraic : polynomial algebra and solving systems of linear equations. We implemented this procedure in section [3](#page-12-0) for the [2,0] CFTs. For each of $A_{1,1}$, $A_{2,1}$ and $D_{4,1}$ we could construct characters with central charges $c_{(1)} = 8, 16, \ldots 128$. For $G_{2,1}$ we could construct characters with central charges $c_{(1)} = 52, 104, 168$. For $F_{4,1}$ we could construct characters with central charges $c_{(1)} = 104, 208, 312$. For $E_{6,1}$ we could construct characters with central charges $c_{(1)} = 24, 48, \ldots, 384$. For $E_{7,1}$, we could construct characters with central charges $c_{(1)} = 56, 112, \ldots 896$.

The venue of the problem is polynomial rings. The statement of the problem has a natural basis of generators of the ring I.e. the characters of \mathcal{C} . The answers we are seeking I.e. characters of one-character extensions are homogenous polynomials of the generators. But more importantly, as we discovered, they are S-invariant. We were able to discover another basis of generators which have the distinction from the earlier basis used in that they are S-invariant. We discovered such a basis with two S-invariant polynomials for each of the [2, 0] CFTs.

Apart from it being aesthetically pleasing to work with a S-invariant basis to obtain answers that are S-invariant, we are rewarded with brevity. The characters when expressed in terms of the S-invariant polynomial basis gave compact forms and this is one of the reasons we could go to such high central charges in this paper.

In section [3,](#page-12-0) we have studied some [3, 0] WZW CFTs and the Ising CFT which is not a WZW CFT but a [3, 0] CFT. For $A_{4,1}$ we could construct characters with central charges $c_{(1)} = 8, 16, \ldots 96$. For the Ising CFT we could construct characters with central charges $c_{(1)} = 8, 16, \ldots 48$. Then we solved the problem we set for ourselves for an infinite number of CFTs the $D_{r,1}$ CFTs. We had to divide the problem into four categories, each with an infinite number of CFTs. We could give a common solution for all the infinite number of CFTs in each of the four categories.

Here again, for each of the $\mathbf{3,0}$ CFTs, we constructed a new basis of generators for the homogenous polynomials, which contained only S-invariant polynomials. If we had not found this S-invariant basis, we could not probably have written this paper. We could provide, for example, the explicit degree-48 polynomial of characters of the Ising CFT that is the character of the monster CFT.

We ask the question, hoped to be answered in the future : is there a way to set up and solve the problem of finding characters of one-character extensions of tensor product CFTs with the S-invariant basis being incorporated at a much earlier stage of the analysis, perhaps from the very beginning?

Another important result/aspect of our work is the content of section [5.](#page-38-0) Some of the basis S-invariant polynomials of characters, in some (not all) CFTs, evaluate magically (after using the q-series of the characters) to a constant. We use "magically" because each character is an infinite series has an infinite amount of information and there are an infinite number of cancellations that happen when it all reduces to one term, the constant term of the series. We called such S-invariant polynomials that evaluate to a constant as a constant S-invariant polynomial. Historically, the constant S-invariant polynomial associated to the Lee-Yang CFT (reproduced here in (68)), was known to Ramanujan and is considered a basis for the Rogers-Ramanujan identities in number theory. We seem to have stumbled upon such objects for many CFTs. See [\(67\)](#page-39-1). We wonder if there are more such constant S-invariant polynomials of characters for other CFTs. This is an avenue of investigation for the future.

Finally, we noted that the constant S-invariant polynomial relations can generate ideals in the polynomial ring of characters. And one should properly work in the corresponding quotient ring. Also, the character of a one-character extension of a tensor product CFT properly belongs to this quotient ring.

Acknowledgments

CNG thanks Prof. Sunil Mukhi for raising a question during the course of the collaboration [\[26\]](#page-43-0), which triggered the investigations of this paper. CNG thanks him in general for general inspiration to pursue this subject of CFTs, MLDEs etc. CNG also thanks (i) the CERN theory group, (ii) Prof. Sudipta Mukherji and IOP, Bhubaneshwar, for their warm hospitality, during the course of this work. JS would like to acknowledge the warm hospitality of The Abdus Salam International Centre for Theoretical Physics, Trieste where part of the work had been done. JS is indebted to Suresh Govindarajan for helpful discussions. JS would also like to acknowledge the warm hospitality of IIT Ropar where he met one of the authors and discussed this project physically.

References

- [1] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, "Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory," Nucl. Phys. B 241 (1984) 333.
- [2] E. Witten, "Nonabelian Bosonization in Two-Dimensions," Commun. Math. Phys. 92 (1984), 455-472
- [3] P. Di Francesco, P. Mathieu and D. Senechal, "Conformal Field Theory," Springer-Verlag, New York, 1997.
- [4] G. W. Moore and N. Seiberg, "LECTURES ON RCFT," RU-89-32.
- [5] J. Fuchs, I. Runkel and C. Schweigert, "Twenty-five years of two-dimensional rational conformal field theory," J. Math. Phys. 51 (2010), 015210 [\[arXiv:0910.3145](http://arxiv.org/abs/0910.3145) [hep-th]].
- [6] M. R. Gaberdiel, "An Introduction to conformal field theory," Rept. Prog. Phys. 63 (2000), 607-667 [\[arXiv:hep-th/9910156](http://arxiv.org/abs/hep-th/9910156) [hep-th]].
- [7] G. Anderson and G. W. Moore, "Rationality in Conformal Field Theory," Commun. Math. Phys. 117 (1988), 441
- [8] T. Eguchi and H. Ooguri, "Differential Equations for Conformal Characters in Moduli Space," Phys. Lett. B 203 (1988), 44
- [9] S. D. Mathur, S. Mukhi and A. Sen, "On the Classification of Rational Conformal Field Theories," Phys. Lett. B 213 (1988), 303-308
- [10] S. D. Mathur, S. Mukhi and A. Sen, "Reconstruction of Conformal Field Theories From Modular Geometry on the Torus," Nucl. Phys. B 318 (1989), 483-540
- [11] S. G. Naculich, "Differential Equations For Rational Conformal Characters," Nucl. Phys. B 323 (1989), 423-440
- [12] H. R. Hampapura and S. Mukhi, "On 2d Conformal Field Theories with Two Characters," JHEP 01 (2016), 005 [\[arXiv:1510.04478](http://arxiv.org/abs/1510.04478) [hep-th]].
- [13] M. R. Gaberdiel, H. R. Hampapura and S. Mukhi, "Cosets of Meromorphic CFTs and Modular Differential Equations," JHEP 04 (2016), 156 [\[arXiv:1602.01022](http://arxiv.org/abs/1602.01022) [hep-th]].
- [14] H. R. Hampapura and S. Mukhi, "Two-dimensional RCFT's without Kac-Moody symmetry," JHEP 07 (2016), 138 [\[arXiv:1605.03314](http://arxiv.org/abs/1605.03314) [hep-th]].
- [15] S. Mukhi and G. Muralidhara, "Universal RCFT Correlators from the Holomorphic Bootstrap," JHEP 02 (2018), 028 [\[arXiv:1708.06772](http://arxiv.org/abs/1708.06772) [hep-th]].
- [16] A. R. Chandra and S. Mukhi, "Towards a Classification of Two-Character Rational Conformal Field Theories," JHEP 04 (2019), 153 [\[arXiv:1810.09472](http://arxiv.org/abs/1810.09472) [hep-th]].
- [17] A. R. Chandra and S. Mukhi, "Curiosities above $c = 24$," SciPost Phys. 6 (2019) no.5, 053 [\[arXiv:1812.05109](http://arxiv.org/abs/1812.05109) [hep-th]].
- [18] S. Mukhi, "Classification of RCFT from Holomorphic Modular Bootstrap: A Status Report," [\[arXiv:1910.02973](http://arxiv.org/abs/1910.02973) [hep-th]].
- [19] A. N. Schellekens, "Meromorphic C = 24 conformal field theories," Commun. Math. Phys. 153 (1993), 159-186 [\[arXiv:hep-th/9205072](http://arxiv.org/abs/hep-th/9205072) [hep-th]].
- [20] E. Witten, "Three-Dimensional Gravity Revisited," [\[arXiv:0706.3359](http://arxiv.org/abs/0706.3359) [hep-th]].
- [21] O. D. King, "A mass formula for unimodular lattices with no roots," Mathematics of Computation 72 (2003), 839 [\[math/0012231\]](http://arxiv.org/abs/math/0012231).
- [22] J. Kaidi, Y. H. Lin and J. Parra-Martinez, "Holomorphic modular bootstrap revisited," JHEP 12 (2021), 151 [\[arXiv:2107.13557](http://arxiv.org/abs/2107.13557) [hep-th]].
- [23] A. Das, C. N. Gowdigere and J. Santara, "Classifying three-character RCFTs with Wronskian index equalling 0 or 2," JHEP 11 (2021), 195 [\[arXiv:2108.01060](http://arxiv.org/abs/2108.01060) [hep-th]].
- [24] J. B. Bae, Z. Duan, K. Lee, S. Lee and M. Sarkis, "Bootstrapping fermionic rational CFTs with three characters," JHEP 01 (2022), 089 [\[arXiv:2108.01647](http://arxiv.org/abs/2108.01647) [hep-th]].
- [25] A. Das, C. N. Gowdigere and S. Mukhi, "New meromorphic CFTs from cosets," JHEP 07 (2022), 152 [\[arXiv:2207.04061](http://arxiv.org/abs/2207.04061) [hep-th]].
- [26] A. Das, C. N. Gowdigere and S. Mukhi, "Meromorphic cosets and the classification of threecharacter CFT," JHEP 03 (2023), 023 [\[arXiv:2212.03136](http://arxiv.org/abs/2212.03136) [hep-th]].
- [27] A. Das, C. N. Gowdigere and J. Santara, "Wronskian Indices and Rational Conformal Field Theories," JHEP 04 (2021), 294 [\[arXiv:2012.14939](http://arxiv.org/abs/2012.14939) [hep-th]].