Learning payoffs while routing in skill-based queues

Sanne van Kempen, Jaron Sanders, Fiona Sloothaak, Maarten G. Wolf

Abstract

Motivated by applications in service systems, we consider queueing systems where each customer must be handled by a server with the right skill set. We focus on optimizing the routing of customers to servers in order to maximize the total payoff of customer-server matches. In addition, customer-server dependent payoff parameters are assumed to be unknown *a priori*. We construct a machine learning algorithm that adaptively learns the payoff parameters while maximizing the total payoff and prove that it achieves polylogarithmic regret. Moreover, we show that the algorithm is asymptotically optimal up to logarithmic terms by deriving a regret lower bound. The algorithm leverages the basic feasible solutions of a static linear program as the action space. The regret analysis overcomes the complex interplay between queueing and learning by analyzing the convergence of the queue length process to its stationary behavior. We also demonstrate the performance of the algorithm numerically, and have included an experiment with time-varying parameters highlighting the potential of the algorithm in non-static environments.

1 Introduction

Service systems such as contact centers, computer networks, and manufacturing systems are widely used in practice [1, 2, 3]. Achieving the highest possible quality of service in such systems is consequently of general importance. However, actually doing so is typically quite challenging because of complex interactions within the system, between e.g. customers and servers. Moreover, service provisioning is an intrinsically uncertain process.

In this paper, we develop a machine learning algorithm that can attain the highest possible performance in one such service system. Specifically, we focus on a finite skill-based queueing system with customer–server dependent random payoffs. One such system is illustrated in Figure 1. We consider different customers to have different needs, and different servers to be better at helping certain types of customer over others. In fact, some servers might even be unable to help some types of customers. We model these aspects by assuming that there are compatibility relations between customers and servers, and by assuming that whenever a customer is served by a server, a random payoff is generated that depends on the specific customer–server pairing. Since the different servers are shared between the different customer types, and since they are limited in their number, the highest possible average reward can only be achieved by optimizing how one matches customers to servers.



Figure 1: A skill-based queueing system with compatibility lines and customer–server dependent payoffs. Here, the λ_i denote arrival rates, the μ_j denote service rates, and the θ_{ij} indicate the average payoff generated upon service completion of a type-*i* customer at server *j*.

Within queueing literature, a canonical tool for doing server allocation is routing policies. Routing policies decide which customer is served by which server, and at what time. However, routing policies typically do not take into account (i) compatibility relations or (ii) payoffs generated by different customer–server pairs. Furthermore, they usually do not consider (iii) uncertainty in and/or a lack of knowledge on model parameters (such as the average payoffs θ_{ij}).

Our machine learning algorithm, Algorithm 1 in Section 4, does take into account aspects (i)–(iii). Letting $D_{ij}(t)$ refer to the number of departures of type-*i* customers at server *j* up to time *t*, and \mathcal{L} to the set of compatibility lines, Algorithm 1 maximizes the long-term expected reward rate

$$\frac{1}{t} \sum_{(ij)\in\mathcal{L}} \theta_{ij} \mathbb{E}(D_{ij}(t)).$$
(1)

Algorithm 1 does so while (a) honoring the queueing dynamics and compatibility relations; (b) guaranteeing the stability of all queues; (c) being able to hone in on the true system parameters θ_{ij} when either misspecified or unknown *a priori*; and (d) dealing with the exploration–exploitation dilemma efficiently. These aspects (a)-(d) will be discussed in more detail in Section 1.1 below, but may be summarized in one sentence as follows: we establish in Theorem 2 that Algorithm 1 achieves a polylogarithmic regret, and in Theorem 1 that this is (asymptotically), up to logarithmic terms, the best possible within a certain class of stable policies.

The nature of the optimization problem that we study, required us to combine techniques from different fields to develop Algorithm 1 and prove Theorems 1 and 2. Firstly, we were inspired by techniques for Multi-Armed Bandit (MAB) problems from *machine learning* for the learning aspect. Secondly, in the regret analysis of Algorithm 1, we had to rely on *queueing theory* to establish the typical behavior of the queue length process under the dynamic routing policy. Finally, a key idea was to leverage *combinatorial optimization* to come up with 'good' actions for Algorithm 1.

Let us briefly discuss these three facets in more detail:

How queueing dynamics complicate decision making and how MAB techniques can be adapted to cope.

Contrary to the classical MAB setting [4], we have to deal with queueing dynamics and stability constraints that complicate the problem. Specifically, both the decision moments and the set of available routing decisions are subject to the underlying random queueing dynamics: we can only route a customer if there are a compatible customer and server available. Moreover, simply routing each customer to the server with the highest payoff might not result in a stable system in the long run.

To overcome this, we define episodes and only make a decision on the routing strategy at the start of each episode. The estimated reward of the chosen action is updated after each episode based on observed payoff samples. This way, similar to the MAB setting, the algorithm learns the average reward of each action adaptively. To face the exploration–exploitation tradeoff efficiently, we use Upper Confidence Bounds (UCB) for the payoff parameters [5].

How queueing theory is used to analyze Algorithm 1.

Each episode, Algorithm 1 chooses an action that determines a fixed routing policy for the duration of that episode. This means that the queue length process will show different behavior in each episode. Worse yet, at the moment the routing policy changes, the queueing system will likely be far from its new equilibrium behavior. It then takes some time to adjust to the new environment. Still, in order to prove convergence results of Algorithm 1, we must show that sufficient payoff samples of all the different lines are collected. To this end, we utilize known results of the stationary behavior and analyze the typical time of convergence of the queue length process to stationarity.

How combinatorial optimization allows one to identify 'good' actions.

The optimal routing problem inherently poses combinatorial challenges. In particular, the optimal routing problem can be considered as a stochastic variant of the optimal transport problem [6] with a finite set of possible solutions. We formulate an optimal transport linear program (LP) and exploit its structural properties in the construction of our algorithm: first, we characterize the basic feasible solutions of the LP in terms of routing rates. We then consider the different sets of routing rates as candidate routing policies in an MAB setting. In the analysis, we exploit the dual of the LP to decompose the regret.

1.1 Main contributions

Our main contributions can be summarized as follows:

Regret lower bound. We present an asymptotic regret lower bound for a class of routing policies satisfying certain stability constraints in Theorem 1. The stability constraints ensure that the queue backlog does not grow infinitely large. In particular, we prove that any policy of this class must suffer $\Omega(\ln(t))$ regret to learn the payoff parameters.

In an MAB setting, a regret lower bound is often described in terms of a *suboptimality gap* for each suboptimal action. The gap measures the instant loss in reward by choosing this action over the optimal action [7, 5, 4]. A straightforward implementation of this approach proves to be difficult since in our model, some suboptimal routing decision might be necessary to maintain stability. Therefore, it is not trivial to identify the optimal action at each decision moment, nor to quantify the suboptimality gaps.

Our solution to this problem is as follows: we split the lines into an 'optimal' and 'suboptimal' set, based on the optimal solution of the optimal transport LP, and consequently quantify the suboptimality gaps using the dual. [8, 9] use a similar method to obtain a regret lower bound in an MAB setting with cost constraints, although their work is limited to programs with only two equality constraints. The proof of Theorem 1 also uses a change–of–measure argument, a technique that is frequently used in the MAB setting [4, 7, 10, 11].

Adaptive learning & routing algorithm. We present a machine learning algorithm, that utilizes the basic feasible solutions of the optimal transport LP as actions in Algorithm 1—a method that, to the best of our knowledge, has not been explored before in this setting. We consider the related optimal transport LP, where the objective function (that includes the payoff parameter θ) is unknown. However, by the properties of the LP, the set of basic feasible solutions is finite and the optimal solution is attained at one of these solutions [6]. Hence, we identify the set of basic feasible solutions as actions. Each solution uniquely defines a set of routing rates. Moreover, the objective function is approximated using UCB indices of the payoff parameters.

The learning is schematically depicted in Figure 2 and works as follows. At the start of each episode, the algorithm chooses the action with maximal (estimated) reward. During the episode, customers are routed according to the rates of the chosen action. In particular, customers are assigned a label corresponding to one of the compatible servers upon arrival (illustrated by colors in Figure 2). Each server serves customers with matching label in First–Come–First–Served (FCFS) order, and service completions generate payoffs. If, at the start of an episode, the chosen action differs from the action of the previous episode, the algorithm assigns new labels for all queueing customers sampled according to the routing rates of the new action.

For each line, we maintain a UCB index of its payoff parameter. At the end of the episode, the payoff samples are used to update the UCB indices. The reward of an action is then computed as the sum over all lines of the UCB index weighted by the routing rate of the action. We assume Bernoulli payoffs for simplicity.



(b) Routing dynamics and possible queue length behavior.

Figure 2: Schematic overview of Algorithm 1.

Regret upper bound. We show that the asymptotic regret of our policy is $\mathcal{O}(\ln^{2\beta}(t))$ for any $\beta > 1$ in Theorem 2. This regret is only slightly worse than the $\mathcal{O}(\ln(t))$ regret of the benchmark UCB policy [5] in the classical MAB setting, even though our learning problem is more challenging. The regret analysis relies on concentration inequalities that bound the probability that the empirical average of payoff samples deviates far from its mean. However, it differs from the standard MAB techniques since the number of routing decision and therefore the number of samples is subject to queueing dynamics. We deal with this issue by splitting each episode into a transient and a stationary phase and analyzing the convergence time of the queue length process to its stationary measure. We let the episodes grow in length such that the probability that the queue length process does not reach stationarity decreases sufficiently fast.

Numerical experiments. Lastly, we analyze Algorithm 1 and its performance in several numerical experiments. We find that in our experiments, the average reward of Algorithm 1 converges quickly to the optimal reward and chooses the optimal action most of the time, even in a larger system with many different actions. The algorithm outperforms a benchmark greedy policy that myopically optimizes the instantaneous payoff without considering long-term effects. Even though the theoretical analysis of Algorithm 1 is for a static setting, we test its robustness against the change of a parameter value and find that the algorithm adapts adequately. This highlights the potential applicability of our algorithm beyond static environments.

1.2 Related literature

The majority of literature on optimal routing in skill-based queues does not account for customer–server dependent payoffs, and focuses on waiting time minimization [3, 12, 13, 14, 15, 16, 17, 18, 19, 20]. In this scenario, the $c\mu$ (or $c\mu/\theta$ in case of abandonments) routing policy that prioritizes customer types based on a ranking depending on the holding costs, is known to be asymptotically optimal [20]. Therefore, an intuitive approach for an adaptive learning policy is to route according to a variant of the $c\mu$ policy where the unknown parameters are replaced by empirical estimators. In our case however, the objective in (1) is difficult to evaluate in full generality, since the departure process depends on the availability of servers and the state of the queues which are random and fairly intractable. Even for simple routing policies like FCFS Assign–the–Longest–Idle–Server (ALIS), the expected number of type-*i* departures at server *j* is, to the best of our knowledge, unknown for finite time. Only a select few convergence results are known for specific queueing systems [14, 15].

On top of the classical exploration–exploitation tradeoff, learning in queueing systems requires one to deal with limited capacity and the interplay between queueing and learning (queueing degrades learning and *vice versa*). Some works decouple learning and queueing by splitting the time horizon into distinct exploration and exploitation phases [21, 20, 22], while others directly implement a reinforcement learning algorithm [23, 19], or apply Bayesian inference [24, 25]. Our Algorithm 1, however, is an integrated learning and routing policy.

The focus in our work lies on online payoff maximization in skill-based queues. Payoff maximization problems with unknown utility functions are well studied in literature [26, 27, 28, 29, 30, 31]. In the setting of queueing systems, there exists a variety of algorithms based on different approaches. For example, [32, 33, 34] propose utility guided algorithms based on a static LP. This method is an extension of the Lyapunov drift plus penalty reward, where the Lyapunov drift assures stability and the penalty is used for payoff maximization. [32] obtain moment bounds for the maximal queue length in the system and an instance independent regret upper bound of order $\mathcal{O}(\sqrt{t \ln(t)})$. However, the algorithm suffers a linear loss in reward due to stability constraints. In our work, we instead split the regret into a queueing and a learning component.

Other algorithms for online payoff maximization in queueing systems are considered by [35, 36], who use Lyapunov drift analysis. A different method is used by [37], who present a routing algorithm that combines the Join-the-Shortest-Queue (JSQ) routing scheme with a confidence ball learning algorithm (introduced by [38]) in the setting of optimization under bandit feedback. It is shown that, given a fixed horizon, the threshold parameter K in the JSQ-K algorithm can be tuned in such a way that polylogarithmic regret can be attained, but no guarantees are provided on the convergence of the routing rates to their the optimal values. In our work, we consider an asymptotic result and therefore our algorithm does not require the knowledge of the time horizon. Lastly, in [26], a primal-dual method is used for reward maximization in an online advertisement setting. An algorithm is presented where the current queue length is deducted from the estimated server-customer payoffs to penalize congestion and near optimality with respect to an oracle reward is proven.

The analysis of our learning algorithm relies on establishing convergence properties of an episodic queue length process to a stationary probability measure. Similar convergence properties have been used in [22, 39, 40]. We let the episodes grow in duration so that convergence is achieved with increasing probability. [41, 42, 43, 23] use similar concepts in different settings of optimal control in queueing networks.

The methodology of our learning algorithm bears most resemblance with the achievable region approach in [44, 45], where the goal is to solve an optimization problem with an unknown utility function using the feasible region spanned by a set of constraints. [44, 45] consider the optimal routing problem in a static setting with known parameters, while our policy integrates learning and optimization in an online fashion.

1.3 Outline

The remainder of this work is organized as follows. In Section 2, we discuss the model, optimal routing problem, and regret formulation. Next, in Section 3, we study the regret of routing policies and present an asymptotic regret lower bound for a class of routing policies. In Section 4, we present the adaptive learning algorithm and show that its asymptotic regret is of polylogarithmic order. Lastly, in Section 5, we present a numerical implementation of the algorithm.

2 Model description

In this section we describe the queueing system and routing policies in more detail. We analyze a related deterministic optimal transport LP and discuss its properties. Lastly, we present a definition of regret of a routing policy, which is based on the optimal transport LP.

2.1 Arrival and service process

We consider a continuous-time queueing system with a fixed set of customer types $\mathcal{I} = \{1, \ldots, I\}$, servers $\mathcal{J} = \{1, \ldots, J\}$, and a set of lines connecting compatible customer types and servers $\mathcal{L} = \{(ij) : i \in \mathcal{I}, j \in \mathcal{J}\}$. Here, $I, J \in \mathbb{N}_{>0}$. Let $L = |\mathcal{L}| \in \mathbb{N}_{>0}$ denote the number of compatible lines. To exclude trivial cases, we assume that L > I + J - 1. We denote the set of servers that are compatible with customer type i by \mathcal{S}_i , and similarly, we denote the set of customer types that are compatible with server j by \mathcal{C}_j , i.e.,

$$S_i = \{ j \in \mathcal{J} : (ij) \in \mathcal{L} \},$$

$$C_j = \{ i \in \mathcal{I} : (ij) \in \mathcal{L} \}.$$
(2)
(3)

An example of such a queueing network is given in Figure 3.



Figure 3: A queueing model with I = 3 types of customers and J = 3 servers with compatibility lines. Here, e.g., $S_1 = \{1, 2, 3\}$ and $C_1 = \{1, 3\}$.

For $i \in \mathcal{I}$, type-*i* customers are assumed to arrive according to a Poisson process with rate $\lambda_i > 0$. They wait in queue *i* until they are allocated to a compatible (and idle) server. We assume that there are no customers in the system at time t = 0. For $j \in \mathcal{J}$, service times of customers at server *j* are assumed exponentially distributed with rate $\mu_j > 0$. We also assume that service times are independent between servers and across customers. Note that the service time distribution only depends on the server and not the customer type. For convenience, we write $\lambda = (\lambda_1, \ldots, \lambda_I)$ and similarly $\mu = (\mu_1, \ldots, \mu_J)$. For stability, we assume that for any subset of customer types $\mathcal{A} \subseteq \mathcal{I}$,

$$\sum_{i \in \mathcal{A}} \lambda_i < \sum_{j \in \mathcal{S}_{\mathcal{A}}} \mu_j.$$
(4)

Here, $S_{\mathcal{A}} = \bigcup_{i \in \mathcal{A}} S_i$ denotes the set of servers that are compatible with any of the customer types $i \in \mathcal{A}$. By [15, Theorem 2.1], there exists a routing policy discipline such that the joint queue length process under that discipline is ergodic.

Upon the service completion of a type-*i* customer at server *j*, a random payoff Y_{ij} is obtained, sampled from a payoff distribution $P_{\theta_{ij}}$ with support $\mathcal{X} \subseteq \mathbb{N}_{\geq 0}$ and unknown and finite mean $\theta_{ij} \in \mathbb{R}_{\geq 0}$. Note that while all lines share the same payoff distribution, the mean payoff differs per line. We denote the vector of unknown mean payoffs as $\theta = (\theta_{ij})_{(ij)\in\mathcal{L}} \in \mathbb{R}^L_{\geq 0}$. As quantification of the difference between the payoff distribution for two lines (ij) and $(k\ell) \in \mathcal{L}$, we use the Kullback–Leibler (KL) divergence, given by

$$I(\theta_{ij}, \theta_{k\ell}) := \sum_{x \in \mathcal{X}} P_{\theta_{ij}}(x) \ln\left(\frac{P_{\theta_{ij}(x)}}{P_{\theta_{k\ell}}(x)}\right).$$
(5)

We assume that the payoff distribution is such that

$$0 < I(\theta_{ij}, \theta_{k\ell}) < \infty \quad \text{when} \quad \theta_{ij} < \theta_{k\ell}.$$
 (6)

2.2 Routing policies

A routing policy π determines which customer is served by which server at what time. We denote the probability measure and expectation with respect to routing policy π and payoff vector $\theta \in \mathbb{R}^{L}_{\geq 0}$ by $\mathbb{P}^{\theta}_{\pi}$ and $\mathbb{E}^{\theta}_{\pi}$, respectively. Moreover, we denote the total number of service completions, i.e., departures, of type-*i* customers at server *j* up to time $t \geq 0$ by $D_{ij}(t)$. We consider a class of routing policies that satisfy for $\varepsilon \geq 0$ and for any payoff vector $\theta \in \mathbb{R}_{\geq 0}$ the following conditions:

$$\lim_{t \to \infty} t\lambda_i - \sum_{j \in \mathcal{S}_i} \mathbb{E}^{\theta}_{\pi}(D_{ij}(t)) < \infty, \quad \forall i \in \mathcal{I},$$
(7)

$$t(\mu_j - \varepsilon) - \sum_{i \in \mathcal{C}_j} \mathbb{E}^{\theta}_{\pi}(D_{ij}(t)) \ge 0, \quad \forall j \in \mathcal{J}, \quad \forall t \ge 0.$$
(8)

Constraint (7) is a notion of stability: we require that the difference between the expected number of arrivals and departures, i.e., the expected queue length, of any customer type does not diverge. Note that the set of policies satisfying (7) is nonempty by assumption (4). Constraint (8) requires that the expected workload of server j is at most $(\mu_j - \varepsilon)/\mu_j$. If $\varepsilon > 0$, (8) prevents critically loaded servers.

We consider the reward of a policy π with respect to payoff vector $\theta \in \mathbb{R}^{L}_{\geq 0}$ up to time $t \geq 0$ as the total average payoff, i.e.,

$$\sum_{(ij)\in\mathcal{L}}\theta_{ij}\mathbb{E}^{\theta}_{\pi}(D_{ij}(t)).$$
(9)

As discussed previously, (9) can be difficult to compute in full generality. Instead, we consider a static version of the optimal control problem which is discussed next.

2.3 Optimal transport LP

For a payoff vector $\theta \in \mathbb{R}_{\geq 0}^L$ and $\varepsilon \geq 0$, let

$$LP(\theta, \varepsilon): \max_{x} \sum_{(ij)\in\mathcal{L}} \theta_{ij} x_{ij},$$
 (10a)

s.t.
$$\sum_{j \in S_i} x_{ij} = \lambda_i, \quad \forall i \in \mathcal{I},$$
 (10b)

$$\sum_{i \in C_i} x_{ij} \le \mu_j - \varepsilon, \quad \forall j \in \mathcal{J},$$
(10c)

$$x_{ij} \ge 0, \quad \forall (ij) \in \mathcal{L}.$$
 (10d)

The objective of $LP(\theta, \varepsilon)$ is to maximize the average payoff per time unit. Each optimization variable $x_{ij} \in [0, \lambda_i]$ can be interpreted as the long-term rate of type-*i* customers that is routed to server *j* per time unit. Constraint (10b) requires that all customers are routed to some server and constraint (10c) states that the capacity of each server must not be exceeded. Here, $\varepsilon \ge 0$ is a fixed amount of slack at each server. The feasibility of $LP(\theta, \varepsilon)$ depends on the value of ε , which will be discussed later. $LP(\theta, \varepsilon)$ can be regarded as a variant of the assignment problem initially presented in [46], where the resources are divisible. Similar LP formulations of the static planning problem in the context of reward maximization in queueing models are used in [33, 47, 37, 26, 27, 28].

We briefly discuss some fundamental properties of LPs. Consider an LP in standard form: $\max_x c'x$ s.t. $Ax = b, x \ge 0$ with $A \in \mathbb{R}^{m \times n}$. For $B \subseteq \{1, \ldots, n\}$, we denote by $\mathbf{B} = [A_{B(1)}, \ldots, A_{B(m)}] \in \mathbb{R}^{n \times n}$ the matrix formed by the columns $B(1), \ldots, B(m)$. We call $B \subseteq \{1, \ldots, n\}$ a basis if |B| = m and \mathbf{B} is invertible (or equivalently, has full rank); see [6]. In this case, $x_B = \mathbf{B}^{-1}b \in \mathbb{R}^n$ is a basic solution. A basis (or equivalently a basic solution) is *feasible* if $\mathbf{B}^{-1}b \ge 0$, and nondegenerate if $(\mathbf{B}^{-1}b)_k \ne 0$ for all $k \in B$. Lastly, recall that if an LP has an optimal solution, then it has an optimal basic feasible solution [6, Theorem 2.7].

In the remainder of this work, we assume nondegeneracy, i.e., we assume that all basic feasible solutions of $LP(\theta, \varepsilon)$ and its dual are nondegenerate. We note that this assumption implies that each basic feasible solution of the primal (dual) implies a unique basic feasible solution of the dual (primal) (see e.g. [6, Exercise 4.12]).

2.3.1 Basic feasible solutions

We will now characterize the basic feasible solutions of $LP(\theta, \varepsilon)$ in (10). Since $LP(\theta, \varepsilon)$ has a finite number of linear inequality constraints, the set of basic feasible solutions is finite [6, Corollary 2.1].

It is a known result that a basic feasible solution of $LP(\theta, \varepsilon)$ induces a spanning forest on the bipartite graph $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$ [37, Proposition 2]. We extend the result of [37] by providing the explicit form of any basis (see Lemma 1 below) and any basic solution (see Lemma 2 below). The proofs are presented in Appendix C.1 and Appendix C.2, respectively.

The introduction of Lemma 1 and 2 do require some more notation. For $\mathscr{L} \subseteq \mathscr{L}$ and $\mathscr{J} \subseteq \mathscr{J}$ we denote by $\mathscr{G}(\mathscr{L}, \mathscr{J})$ the subgraph of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$ induced by \mathscr{L} and \mathscr{J} . We say that $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is a *spanning forest* of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$ if (i) it is a union of trees, i.e., it contains no cycles, (ii) each $v \in \mathcal{I} \cup \mathcal{J}$ is contained in $\mathscr{G}(\mathscr{L}, \mathscr{J})$, and (iii) each tree of $\mathcal{G}(\mathcal{L}, \mathscr{J})$ contains a unique root node $j \in \mathscr{J}$. In such a tree, parent and child nodes of servers are customer types and *vice versa*. We denote the subtree rooted in v, including v itself, by $\cap(v)$. We let $\mathsf{C}(\cap(v))$ denote the set of customer types contained in the subtree rooted in v, i.e., $\mathsf{C}(\cap(v)) = \cap(v) \cap \mathcal{I}$. Similarly, $\mathsf{S}(\cap(v)) = \cap(v) \cap \mathcal{J}$. An example is shown in Figure 4.



Figure 4: Consider the queueing system in Figure 3 and let $\mathscr{L} = \{(11), (13), (23), (33)\}$ and $\mathscr{J} = \{1, 2\}$. $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is then a spanning forest of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$ consisting of two trees.

We are now in position to state Lemma 1 and 2:

Lemma 1. Let $\varepsilon \geq 0$, $\mathscr{L} \subseteq \mathcal{L}$, and $\mathscr{J} \subseteq \mathcal{J}$. $B = \mathscr{L} \cup \mathscr{J}$ is a basis of $LP(\theta, \varepsilon)$ if and only if $\mathcal{G}(\mathscr{L}, \mathscr{J})$ is a spanning forest of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$.

Lemma 2. Let $\varepsilon \ge 0$, $\mathscr{L} \subseteq \mathscr{L}$, and $\mathscr{J} \subseteq \mathscr{J}$. If $B = \mathscr{L} \cup \mathscr{J}$ is a basis of $LP(\theta, \varepsilon)$, then its corresponding basic solution $x \in \mathbb{R}^L$ satisfies

$$x_{ij} = \begin{cases} \sum_{k \in \mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell \in \mathsf{S}(\cap(i))} (\mu_\ell - \varepsilon), & \text{if } i \text{ is a child of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), \\ \sum_{\ell \in \mathsf{S}(\cap(j))} (\mu_\ell - \varepsilon) - \sum_{k \in \mathsf{C}(\cap(j))} \lambda_k, & \text{if } i \text{ is the parent of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), \\ 0 & \text{if } (ij) \in \mathcal{L} \setminus \mathscr{L}. \end{cases}$$
(11)

2.3.2 Feasibility

Note that by (4), $LP(\theta, 0)$ is feasible, i.e, there exists at least one basic feasible solution. The feasibility region of $LP(\theta, \varepsilon)$ in (10) depends on ε : the larger ε , the smaller the feasible region. Lemma 3 characterizes the range of ε such that the $LP(\theta, \varepsilon)$ has the same set of feasible bases as $LP(\theta, 0)$. The lemma is proven in Appendix C.3.

Lemma 3. Let $B = \mathscr{L} \cup \mathscr{J}$ with $\mathscr{L} \subseteq \mathscr{L}$ and $\mathscr{J} \subseteq \mathscr{J}$ be a nondegenerate feasible basis of $LP(\theta, 0)$. If

$$0 \le \varepsilon < \min_{j \in \mathcal{J}} \frac{\sum_{\ell \in \mathsf{S}(\bigcap(j))} \mu_{\ell} - \sum_{k \in \mathsf{S}(\bigcap(j))} \lambda_{k}}{|\mathsf{S}(\bigcap(j))|},$$
(12)

then B also is a nondegenerate feasible basis of $LP(\theta, \varepsilon)$.

2.4 Regret formulation

We use the value of $LP(\theta, \varepsilon)$ in (10) as the oracle reward to define the regret of a learning policy. Let $\varepsilon > 0$ satisfy (12), $\theta \in \mathbb{R}^{L}_{\geq 0}$ be a payoff vector, and $x^{\theta} \in \mathbb{R}^{L}_{\geq 0}$ be the optimal basic feasible solution of $LP(\theta, \varepsilon)$ in (10). We define the (expected) regret of a routing policy π at time $t \geq 0$ as

$$R^{\theta}_{\pi}(t) := \sum_{(ij)\in\mathcal{L}} \theta_{ij} \left(t x^{\theta}_{ij} - \mathbb{E}^{\theta}_{\pi}(D_{ij}(t)) \right).$$
(13)

Lemma 4 states that the asymptotic regret is nonnegative and is proven in Appendix C.4.

Lemma 4. Let $\varepsilon \geq 0$ and let π satisfy (7) and (8). Then $\lim_{t\to\infty} R^{\theta}_{\pi}(t)/t \geq 0$.

3 Regret lower bound

We provide an asymptotic lower bound for the regret for routing policies satisfying constraints (7) and (8) in Theorem 1. To this end, we first provide a regret decomposition in Lemma 5.

3.1 Regret decomposition

Let $\varepsilon > 0$ satisfy (12), $\theta \in \mathbb{R}_{\geq 0}^{L}$ be a payoff vector, and $x^{\theta} \in \mathbb{R}_{\geq 0}^{L}$ be the unique optimal solution of $LP(\theta, \varepsilon)$ in (10). We define the set of optimal lines $O(\theta)$ as the lines with non-zero load in the optimal solution $x^{\theta} \in \mathbb{R}^{L}$ and the set of suboptimal lines $O^{c}(\theta)$ as its complement:

$$O(\theta) = \{ (ij) \in \mathcal{L} : x_{ij}^{\theta} > 0 \},$$

$$(14)$$

$$O^{c}(\theta) = \mathcal{L} \setminus O(\theta).$$
⁽¹⁵⁾

The dual problem [6] of $LP(\theta, \varepsilon)$ is given by

$$D(\theta,\varepsilon): \min_{v,w} \sum_{i\in\mathcal{I}} \lambda_i v_i + \sum_{j\in\mathcal{J}} (\mu_j - \varepsilon) w_j$$
(16a)

s.t.
$$v_i + w_j - \theta_{ij} \ge 0, \quad \forall (ij) \in \mathcal{L},$$
 (16b)

$$v_i \in \mathbb{R}, \quad \forall i \in \mathcal{I},$$
 (16c)

$$w_j \ge 0, \quad \forall j \in \mathcal{J}.$$
 (16d)

By strong duality, we have that $D(\theta, \varepsilon)$ is feasible since $LP(\theta, \varepsilon)$ is feasible. Moreover, x^{θ} and v^{θ} , w^{θ} are an optimal primal-dual pair if and only if they satisfy the complementary slackness conditions [6, Theorem 4.5], which are given by

$$x_{ij}^{\theta}(v_i^{\theta} + w_j^{\theta} - \theta_{ij}) = 0, \quad \forall (ij) \in \mathcal{L},$$

$$(17)$$

$$\left(\mu_j - \varepsilon - \sum_{i \in \mathcal{C}_j} x_{ij}^{\theta}\right) w_j^{\theta} = 0, \quad \forall j \in \mathcal{J}.$$
(18)

We note that the optimal solutions of $LP(\theta, \varepsilon)$ and $D(\theta, \varepsilon)$ are unique due to the nondegeneracy assumption. Moreover, the nondegeneracy assumption implies strict complementarity (see e.g. Exercises 4.12 and 4.20c in [6]).

For a line $(ij) \in \mathcal{L}$, we define

$$\phi_{ij}^{\theta} = v_i^{\theta} + w_j^{\theta} - \theta_{ij} \tag{19}$$

as its suboptimality gap. Here, $v^{\theta} \in \mathbb{R}^{I}$, $w^{\theta} \in \mathbb{R}^{J}_{\geq 0}$ is the optimal solution of $D(\theta, \varepsilon)$ in (16). Note that $\phi_{ij}^{\theta} = 0$ for any $(ij) \in O(\theta)$ by (17) and $\phi_{k\ell} > 0$ for any $(k\ell) \in O^{c}(\theta)$ by strict complementarity.

Lemma 5 presents a regret decomposition based on the line classification. The proof can be found in Appendix C.5.

Lemma 5. Let $v^{\theta} \in \mathbb{R}^{I}$, $w^{\theta} \in \mathbb{R}_{\geq 0}^{J}$ be the optimal solution of $D(\theta, \varepsilon)$ in (16). Let π satisfy (7) and (8). Then the regret in (13) can be expressed as

$$R_{\pi}^{\theta}(t) = \underbrace{\sum_{(k\ell)\in O^{c}(\theta)} \phi_{k\ell}^{\theta} \mathbb{E}_{\pi}^{\theta}(D_{k\ell}(t))}_{\mathrm{II}} + \underbrace{\sum_{j\in\mathcal{J}} w_{j}^{\theta}\left(t(\mu_{j}-\varepsilon) - \sum_{i\in\mathcal{C}_{j}} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t))\right)}_{\mathrm{II}} + \underbrace{\sum_{i\in\mathcal{I}} v_{i}^{\theta}\left(t\lambda_{i} - \sum_{j\in\mathcal{S}_{i}} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t))\right)}_{\mathrm{III}}.$$
 (20)

In decomposition (20), term I represents the loss in reward due to customers being routed over suboptimal lines. The other two summations represent regret accumulated by deviating from the oracle routing rates. In particular, II measures regret from not utilizing the allowed capacity from servers and III measures regret accumulated from customers that are still waiting in the queue or are in service at time t, since we do not obtain any payoff until a customer finishes service. Note that since $v_i^{\theta} \in \mathbb{R}$, the queue length can actually contribute to negative regret, since it might be beneficial to let customers wait until a high pay-off server is available.

3.2 Regret lower bound

Before we can state the asymptotic regret lower bound, we need to introduce some further notation. Let $\theta \in \mathbb{R}^{L}_{\geq 0}$. We define $A(\theta, ij, a) \in \mathbb{R}^{L}_{\geq 0}$ element-wise by

$$A(\theta, ij, a)_{k\ell} = \begin{cases} a & \text{if } (k\ell) = (ij), \\ \theta_{k\ell} & \text{otherwise.} \end{cases}$$
(21)

In particular, $A(\theta, ij, a)$ represents a payoff vector that is (almost) equal to θ except for index (ij). For $(ij) \in O^{c}(\theta)$, we define

$$\Delta(\theta, ij) = \left\{ a \ge 0 : \ (ij) \in O(A(\theta, ij, a)) \right\}$$

$$(22)$$

as the set of parameter values that make (ij) an optimal line without changing the payoff values of the other lines. Note that by construction, either $\Delta(\theta, ij) = \emptyset$, or $\Delta(\theta, ij) = (a_0, \infty)$ for some $a_0 \in \mathbb{R}_{\geq 0}$. We define

$$K(\theta, ij) = \inf\{I(\theta_{ij}, a) : a \in \Delta(\theta, ij)\}$$
(23)

as the minimum distance in terms of KL divergence between the parameter θ_{ij} and the set of parameter values that make (ij) an optimal line. Here, the KL divergence I is defined in (5).

Lemma 6 states that any suboptimal line could become optimal by changing its payoff value. It is proven in Appendix C.6.

Lemma 6. For any $(ij) \in O^c(\theta)$, we have $\Delta(\theta, ij) \neq \emptyset$.

Theorem 1 provides an asymptotic regret lower bound and is proven in Appendix A.

Theorem 1. Let π satisfy (7) and (8). Moreover, we assume that π is consistent, i.e., that for any deterministic payoff vector $\xi \in \mathbb{R}_{\geq 0}^{L}$,

$$\lim_{t \to \infty} \mathbb{P}^{\xi}_{\pi}(D_{ij}(t) < b \ln(t)) = 0, \qquad \forall b > 0, \qquad \forall (ij) \in O(\xi).$$
(24)

Then, for any $\theta \in \mathbb{R}^{L}_{\geq 0}$,

$$\lim_{t \to \infty} \frac{R_{\pi}^{\theta}(t)}{\ln(t)} \ge \sum_{(k\ell) \in O^c(\theta)} \frac{\phi_{k\ell}^{\theta}}{K(\theta, k\ell)} > 0.$$
⁽²⁵⁾

The lower bound in (25) is a summation over all suboptimal lines of the suboptimality gap divided by the infimum KL distance as defined in (23). This means that a suboptimal line that is nearly optimal, in the sense that its suboptimality gap is small, leads to a large contribution to the regret, since it is difficult for a policy to distinguish between optimal and nearly optimal lines. Note that the lower bound is not redundant as strict positivity is guaranteed.

Assumption (24) is a notion of consistency which is a common assumption in regret analysis in MAB literature [11] as it excludes policies that are specialized on a subset of problem instances. Intuitively speaking, it states that optimal lines are sufficiently explored. In particular, (24) requires that the number of departures of optimal lines eventually grows larger than logarithmic with probability one. In classical bandit literature, an assumption on the concentration of the expectation of the number of optimal decisions is often sufficient, since there is exactly one decision at each time step. In our case however, the number of departures is random even for stationary policies. It remains an open question whether a similar regret lower bound can be proven for a class of policies satisfying a weaker constraint such as $\lim_{t\to\infty} \mathbb{E}^{\theta}_{\pi}(tx_{ij}^{\theta} - D_{ij}(t))/t^{\theta} = 0$ for any b > 0 and $(ij) \in \mathcal{L}$, in combination with assumptions (7) and (8).

Theorem 1 states that the regret is $\Omega(\ln(t))$, which is the same order as the lower bound for classical MABs [4, Theorem 2]. The proof of Theorem 1 in Appendix A is structured in a similar way as those presented in [4, 7, 10, 11] and uses a change–of–measure argument. Since payoffs are random samples, each line is associated with a level of uncertainty that decreases with the number of obtained samples, which in our case corresponds to departures. In order to prove the theorem, we construct a second, 'confusing' system which has the same parameters as the original model with the exception of the average payoff $\theta_{k\ell}$ for some line $(k\ell) \in \mathcal{L}$, which we set to a high value such that line $(k\ell)$ is an optimal line in the second model. We then prove that a consistent policy needs $\Omega(\ln(t))$ samples of payoffs of line $(k\ell)$ in order to distinguish between the two models.

Recall that Theorem 1 assumes that the payoff distribution is discrete (see the end of Section 2.1). We note that the proof can be adapted to include payoff distributions defined by a probability density function $f(\cdot, \theta_{ij})$: in this case, the KL divergence in (5) changes to

$$I(\theta_{ij}, \theta_{k\ell}) := \int_0^\infty f(x; \theta_{ij}) \ln\left(\frac{f(x; \theta_{ij})}{f(x; \theta_{k\ell})}\right) \mathrm{d}x.$$
(26)

Similarly, the Radon–Nikodym (RN) derivative in the proof of Theorem 1 in Appendix A changes to $d\mathbb{P}_{\pi}^{\theta}/d\mathbb{P}_{\pi}^{\theta'} = \prod_{i=1}^{m} f(X_i; \theta_{k\ell})/f(X_i; \theta'_{kl})$ and equations (67), (70), and (71) change accordingly. Extension of our result to general probability measures requires careful construction of the RN derivative. This can be done in line with the discussion following [11, Proposition 4.8].

4 Adaptive queue routing policy

In this section, we introduce a routing algorithm called Adaptive UCB Queue Routing Algorithm (UCB QR) and show that its asymptotic regret is close to the lower bound in Theorem 1. Specifically, Theorem 2 proves that its regret is polylogarithmic as time grows large.

4.1 Description of the algorithm

We consider a fixed network structure $\mathcal{I}, \mathcal{J}, \mathcal{L}, \lambda, \mu$ and $\varepsilon > 0$ satisfying (12). We assume that the payoff distribution of line $(ij) \in \mathcal{L}$ is Bernoulli distributed with parameter $\theta_{ij} \in [0, 1]$.

We consider the basic feasible solutions of $LP(\theta, \varepsilon)$ in (10) as the different actions in an MAB setting. Since the set of basic feasible solutions of $LP(\theta, \varepsilon)$ is finite (see Section 2.3.1), the set of actions is finite and will be denoted by \mathcal{A} . Concretely, each action $a \in \mathcal{A}$ is associated to a basic feasible solution $x^a \in \mathbb{R}^L_{\geq 0} LP(\theta, \varepsilon)$. We call the action corresponding to the optimal basic feasible solution the optimal action, which is unique by the nondegeneracy assumption.

For $a \in \mathcal{A}$, we denote the set of lines with positive load by $\mathcal{L}^a \subseteq \mathcal{L}$, i.e.,

$$\mathcal{L}^a := \{ (ij) \in \mathcal{L} : \ x^a_{ij} > 0 \}.$$

$$\tag{27}$$

The long-term average reward $r^a \in \mathbb{R}_{\geq 0}$ will be denoted by

$$r^a := \sum_{(ij)\in\mathcal{L}^a} x^a_{ij}\theta_{ij},\tag{28}$$

and its associated suboptimality gap by

$$d^a := \max_{\tilde{a}} r^{\tilde{a}} - r^a.$$
⁽²⁹⁾

The UCB QR algorithm is given in Algorithm 1.

Algorithm 1 Adaptive UCB Queue Routing Algorithm (UCB QR).

- 1: Initialize k = 1, for all $(ij) \in \mathcal{L}$ initialize $T_{ij}(0) = 0$, $\hat{\theta}_{ij}(0) = 0$, and $U_{ij}(0) = \infty$ and for all $a \in \mathcal{A}$ initialize $U^a(0) = \infty$.
- 2: for each episode $k = 1, 2, \ldots$ of length H_k do
- 3: Choose action $A_k = \operatorname{argmax}_a U^a(k-1)$, with ties broken arbitrarily.
- 4: **if** $A_k \neq A_{k-1}$ **then**

5: Reallocate customers to virtual queues: for each waiting type-*i* customer, assign it to the virtual queue of server *j* with probability $x_{ij}^{A_k}/\lambda_i$.

- 6: For each $j \in \mathcal{J}$, order the customers in virtual queue j by their arrival time.
- 7: end if
- 8: for time $t = 0, \ldots, H_k$ do
- 9: Serve customers according to Algorithm 2: FCFS $RR(A_k)$.
- 10: end for

11: For each line $(ij) \in \mathcal{L}^{A_k}$, observe $N_{ij}^k \in \mathbb{N}_{\geq 0}$ payoff samples $(Y_{ij}^\ell)_{\ell=1,\ldots,N_{ij}^k}$.

12: For each line $(ij) \in \mathcal{L}^{A_k}$, update

$$T_{ij}(k) = T_{ij}(k-1) + N_{ij}^k, (30)$$

$$\hat{\theta}_{ij}(k) = \frac{\hat{\theta}_{ij}(k-1)T_{ij}(k-1) + \sum_{\ell=1}^{N_{ij}^{\kappa}} Y_{ij}^{\ell}}{T_{ij}(k)},$$
(31)

$$U_{ij}(k) = \begin{cases} \hat{\theta}_{ij}(k) + \sqrt{\frac{\ln(k)}{T_{ij}(k)}} & \text{if } T_{ij}(k) > 0, \\ \infty & \text{if } T_{ij}(k) = 0. \end{cases}$$
(32)

13: Update the UCB index of action A_k ,

$$U^{A_k}(k) = \sum_{(ij)\in\mathcal{L}^{A_k}} x_{ij}^{A_k} U_{ij}(k).$$
(33)

14: end for

The learning algorithm works as follows. We maintain UCB indices for the average payoffs of lines $(ij) \in \mathcal{L}$ and use these to compute UCB indices for each action $a \in \mathcal{A}$. At the start of each episode, we choose the action $a \in \mathcal{A}$ with the highest UCB index. During the episode, we route customers according to the rates $x^a \in \mathbb{R}^L_{\geq 0}$ using the FCFS Random–Routing (FCFS RR) scheme in Algorithm 2. We use virtual queues to implement the FCFS RR scheme. For each departure of a type-*i* customer at server *j*, we obtain a payoff sample from a Bernoulli distribution with parameter $\theta_{ij} \in [0, 1]$. At the end of each episode, we use the obtained payoff

Algorithm 2 FCFS RR(a).

1: for each arrival of a type-*i* customer do

- 2: Assign customer to the virtual queue of server j with probability x_{ij}^a/λ_i .
- 3: If the server is idle, start service.
- 4: **end for**
- 5: for each service completion at server j do
- 6: Obtain a payoff $Y_{ij} \sim \text{Ber}(\theta_{ij})$, where *i* is the departing customer's type.
- 7: If the virtual queue of server j is nonempty, start service of the customer at the head of the queue.
- 8: end for

samples to update the UCB indices. Note that customers may be reallocated to a different virtual queue when a change in action occurs at episode transitions. Episode k has a predetermined length $H_k \in \mathbb{R}_{>0}$ where H_k is a fixed constant that may depend on k.

For each line $(ij) \in \mathcal{L}$, we keep track of $T_{ij}(k)$, the total number of departures of type-*i* customers at server j up to episode number $k \in \mathbb{N}_{\geq 1}$. Similarly, we keep track of the empirical mean $\hat{\theta}_{ij}(k)$ of type-(ij) payoffs and a UCB estimator $U_{ij}(k)$. The counters and empirical means are both initialized at zero while the UCB estimator is initialized at infinity as an incentive for the algorithm to obtain at least one sample from each line. The number of type-(ij) departures in episode k, N_{ij}^k , in line 11 is random due to the queueing dynamics.

4.2 Regret upper bound

Theorem 2 gives an asymptotic upper bound on the regret as defined in (13) of Algorithm 1.

We denote the minimal positive rate on a line across any action as $x_{\min} := \min_{a \in \mathcal{A}} \min_{(ij) \in \mathcal{L}^a} x_{ij}^a$. Moreover, $\tilde{\lambda}_j^a := \sum_{i \in \mathcal{C}_i} x_{ij}^a$ denotes the total arrival rate of customers at server j under action a and

$$\rho_j^a := \tilde{\lambda}_j^a / \mu_j \in (0, 1) \tag{34}$$

denotes the load of server j under action a, for each $j \in \mathcal{J}$ and $a \in \mathcal{A}$.

Theorem 2. Let $\beta \in \mathbb{R}_{>1}$ and let $\alpha \in \mathbb{R}_{>1}$ satisfy

$$\alpha \ge \max_{j \in \mathcal{J}} \max_{a \in \mathcal{A}} \left\{ \frac{3(\tilde{\lambda}_{j}^{a} + \mu_{j}) \ln(\rho_{j}^{a}) - 2\sqrt{(\mu_{j} - \tilde{\lambda}_{j}^{a})^{2} + 9\tilde{\lambda}_{j}^{a}\mu_{j} \ln^{2}(\rho_{j}^{a})}{2(\mu_{j} - \tilde{\lambda}_{j}^{a})^{2} \ln(\rho_{j}^{a})}, 1 \right\}.$$
(35)

Let $H_0 \in \mathbb{R}_{\geq 1}$ and the episode lengths H_k , $k = 1, 2, \ldots$, satisfy

$$H_k = \tau_k + H_0 \quad with \quad \tau_k = \alpha \ln^\beta(2Jk) \quad and \quad H_0 \ge \max\left\{\frac{4}{x_{\min}}, 1\right\}.$$
(36)

Lastly, let $\theta \in \mathbb{R}^{L}_{\geq 0}$ be a payoff vector. Then, Algorithm 1 satisfies (24). Moreover, the regret $\mathbb{R}^{\theta}(t)$ of Algorithm 1 satisfies

$$\lim_{t \to \infty} \frac{R^{\theta}(t)}{\alpha \ln^{2\beta}(t)} \le \sum_{(k\ell) \in O^{c}(\theta)} \phi^{\theta}_{k\ell}(\mu_{\ell} - \varepsilon) |\mathcal{A}| + \sum_{j \in \mathcal{J}} w^{\theta}_{j}(\mu_{j} - \varepsilon) |\mathcal{A}|,$$
(37)

where $w^{\theta} \in \mathbb{R}^{J}_{\geq 0}$ is the optimal dual variable of $D(\theta, \varepsilon)$ in (16) and $\phi^{\theta} \in \mathbb{R}^{L}_{\geq 0}$ is defined in (19).

The regret upper bound in Theorem 2 provides a theoretical guarantee on the performance of Algorithm 1. It shows that the regret scales with rate at most $\alpha \ln^{2\beta}(t)$ as the time horizon t grows large. Note that this matches with the regret lower bound in Theorem 1 up to logarithmic terms. From the right side of (37), we observe that the upper bound increases with the cardinality of the action set $|\mathcal{A}|$, this will be discussed in Section 4.4.2. Moreover, the regret for each suboptimal action is upper bounded by the instant regret of suboptimal lines in the set $O^c(\theta)$ (recall (15)) multiplied by the maximal expected departure rate at server j, which is $\mu_j - \varepsilon$ by construction. For a suboptimal line $(ij) \in O^c(\theta)$, the instant regret is expressed by the suboptimality gap $\phi_{ij} \in \mathbb{R}_{>0}$ in (19). The last term in (37) is related to the queueing regret: it represents the regret obtained at the start of optimal episodes when the queue length process is not necessarily close to its stationary behavior. We outline the proof of Theorem 2 in Section 4.3. The full proof is given in Appendix B.

4.2.1 Episode length

In order to analyze the regret of the algorithm, we provide lower bounds on the number of departures for each line. The analysis is challenging since the routing scheme changes at the start of each episode, hence the starting state of the queueing system is not necessarily close to its stationary behavior, as illustrated in Figure 5. This motivates the introduction of a *warmup time* τ_k in the construction of the episode length H_k in (36). The warmup time is followed by a period H_0 of fixed length.

In the analysis of the algorithm, we provide a lower bound on the probability of convergence of the queue length process to its stationary behavior within the warmup time. Constraint (35) guarantees that this bound is sharp enough. The warmup time τ_k is increasing in k, so that this probability converges to 1 at the right speed in terms of the episode number k.



Figure 5: Possible realization of the queue length process of the virtual queue of server $j \in \mathcal{J}$ under Algorithm 1 for two episodes. Episode k consists of a warmup time τ_k followed by a period of fixed length H_0 . Every episode, the algorithm can choose a different action with its own stationary measure.

4.3 **Proof outline**

We sketch the proof of Theorem 2, highlighting the key contributions. The proof can be split into four steps. We shortly explain these steps, followed by a more in-depth elaboration per step. Without loss of generality, we label the first action as the unique optimal action, i.e., $\operatorname{argmax}_{a \in \mathcal{A}} r^a = 1$.

- Step 1. Analyzing the number of departures in FCFS RR. For the proof of Theorem 2, we need to show that Algorithm 1 learns the payoff parameters sufficiently fast. Since payoff samples are collected upon departures, we need to prove a lower bound on the number of departures in the queueing system. We do so via an intermediate step: we consider the scenario where the system is initially empty at the start of an episode, which represents a 'worst case' scenario. Next, we show that this worst case queue length process reaches stationarity within warmup time τ_m with sufficiently high probability (Lemma 7).
- Step 2. Bounding the probability of choosing a suboptimal action. We note that Algorithm 1 chooses a suboptimal action in Line 3 only if the UCB index of the suboptimal action in (33) is at least as high as the UCB index of the optimal action. Broadly speaking, this can happen if (A) the index of the suboptimal action overestimates its true mean, or (B) the index of the optimal action underestimates its true mean. Compared to the classical MAB analysis in [11], event (A) is more complicated to analyze in our model, since the number of payoff samples we obtain within one episode is stochastic rather than deterministic as in the classical MAB setting. We show that event (A) is unlikely by showing that on the one hand, overestimation based on 'sufficient' samples is unlikely and on the other hand, it is unlikely to obtain 'insufficient' samples. Here, the term 'sufficient' is carefully constructed, as discussed below. Event (B) is unlikely by construction of the UCB estimators. We bound the probability of event (A) in Lemma 8 and 9 and event (B) in Lemma 10.
- Step 3. Bounding the number of suboptimal episodes. We use the bounds from Step 2 to bound the number of episodes where Algorithm 1 chooses a suboptimal action in Lemma 11.
- Step 4. Bounding the regret of Algorithm 1. Finally, we prove Theorem 2 in Appendix B using the regret decomposition (20). Term I is the main contributing factor since it measures the regret accumulated by using suboptimal lines. We bound this term using Lemma 11. Term II is bounded by analyzing the queue length behavior in episodes where the algorithm chooses the optimal action. Lastly, we bound term III using the properties of FCFS RR and the constraints in $LP(\theta, \varepsilon)$.

Let us next describe Steps 1-4 in more detail.

4.3.1 Step 1. Analyzing the number of departures in FCFS RR

We consider the queue length process $Q_j^{mA_m}(t)$, $t \in [0, H_m)$ of the virtual queue of server $j \in \mathcal{J}$ during episode $m \in \mathbb{N}_{\geq 1}$, where A_m denotes the action chosen by Algorithm 1 in Line 3. Within an episode, Algorithm 1 routes customers according to the fixed FCFS RR policy in Algorithm 2. Since arrival processes are independent across customer types, we have by the Poisson split and merge properties [48] that the arrival process of $Q_j^{mA_m}(t)$ is a Poisson process with rate $\sum_{i \in \mathcal{I}} x_{ij}^{A_m}$. This implies that $Q_j^{mA_m}(t)$ behaves as an M/M/1 queueing system independently from other servers. Recall that x^{A_m} is a basic feasible solution of $LP(\theta, \varepsilon)$ in (10), so by constraint (10c), we have $\sum_{i \in \mathcal{C}_j} x_{ij}^{A_m} \leq \mu_j - \varepsilon < \mu_j$. Therefore, $\rho_j^{A_m} = \sum_{i \in \mathcal{I}} x_{ij}^{A_m}/\mu_j < 1$ and hence $Q_j^{mA_m}(t)$ is positive recurrent [49], and its stationary distribution is given by

$$p_j^{A_m}(n) := \lim_{t \to \infty} \mathbb{P}(Q_j^{mA_m}(t) = n) = (1 - \rho_j^{A_m})(\rho_j^{A_m})^n.$$
(38)

For each server, we associate two new queue length processes $\hat{Q}_{j}^{mA_{m}}(t)$ and $\underline{Q}_{j}^{mA_{m}}(t)$ where the initial value $\hat{Q}_{j}^{mA_{m}}(0)$ is sampled from the stationary measure $p_{j}^{A_{m}}$ and $\underline{Q}_{j}^{mA_{m}}(0) = 0$. We couple the processes $Q_{j}^{mA_{m}}(t)$, $\hat{Q}_{j}^{mA_{m}}(t)$, and $\underline{Q}_{j}^{mA_{m}}$ in the sense that the sampled arrival times and service completions (if possible) are the same for all processes, as illustrated in Figure 6.



Figure 6: A possible realization of the queue length processes $Q_j^{mA_m}(t)$, $\hat{Q}_j^{mA_m}(t)$, and $\underline{Q}_j^{mA_m}(t)$. Customer arrivals lead to a unit increase and service completions to a unit decrease (unless the queue is empty). At hitting time x the processes $\hat{Q}_j^{mA_m}(t)$ and $\underline{Q}_j^{mA_m}(t)$ collide and evolve identically afterwards.

Note that if $\hat{Q}_{j}^{mA_{m}}(t)$ and $\underline{Q}_{j}^{mA_{m}}(t)$ collide, then the departure process of $\underline{Q}_{j}^{mA_{m}}(t)$ is stationary afterwards. Lemma 7 provides a bound on the hitting time of $\hat{Q}_{j}^{mA_{m}}(t)$ and $\underline{Q}_{j}^{mA_{m}}(t)$. The proof in Appendix C.7 relies on hitting time analysis of an M/M/1 queueing system. Lemma 7 extends the result of [22, Proposition 4], in the sense that we provide a sharper bound by exploiting the known formula for the moment generating function of hitting times in an M/M/1 system. We also provide exact values for the constants in our result, which are not provided in [22].

Lemma 7. Let $\beta \in \mathbb{R}_{>1}$ and let $m \in \mathbb{N}_{\geq 1}$ satisfy $m \geq C_{\beta}$ with

$$C_{\beta} := \frac{1}{2J} \exp\left(\beta^{\frac{1}{\beta-1}}\right). \tag{39}$$

For any $t \ge \tau_m$, we have $\mathbb{P}(\underline{Q}_j^{mA_m}(t) = \hat{Q}_j^{mA_m}(t), \ \forall j \in \mathcal{J}) \ge 1 - 1/m^{\beta}$.

4.3.2 Step 2. Bounding the probability of choosing a suboptimal action

Bound on event (A). Note that the confidence bound in (33) decreases with the number of obtained samples (departures). This means that overestimation of the true mean is likely when the number of samples is small, but becomes less likely as the number of obtained samples increases. We split the analysis into two parts. First, Lemma 8 bounds the probability that Algorithm 1 has not obtained sufficient samples after $C_{\beta} + \ln^{\beta}(k)$ episodes. Here, we quantify 'sufficient' by σ_{ijk}^{a} in (43) below. To provide the lemma, we introduce some notation. Let

$$\overline{\theta}_{ij}[s] := \frac{1}{s} \sum_{\ell=1}^{s} Y_{ij}^{\ell} \tag{40}$$

denote the empirical average payoff of type-(ij) departures based on s samples. Here, the $(Y_{ij}^{\ell})_{\ell=1}^{s}$ denote i.i.d. random variables with distribution Ber (θ_{ij}) . We let $D_{ij}^{mA_m}$ denote the number of type-(ij) departures within episode $m \in \mathbb{N}_{\geq 1}$ but after warmup time τ_m , where A_m denotes the action chosen by Algorithm 1 in Line 3. Moreover, for $a \in \mathcal{A}$ let

$$\overline{T}^a_{ij}[s] := \sum_{m=1}^s D^{ma}_{ij} \tag{41}$$

denote the total number of type-(ij) departures (after warmup periods) after completing $s \in \mathbb{N}_{\geq 1}$ episodes where action a was chosen. We let $\overline{U}^a[s,m]$ denote the UCB index of action $a \in \mathcal{A}$ at episode $m \in \mathbb{N}_{\geq 1}$ if action a was chosen in $s \leq m$ episodes,

$$\overline{U}^{a}[s,m] := \sum_{(ij)\in\mathcal{L}^{a}} x^{a}_{ij} \Big(\overline{\theta}_{ij} \big[\overline{T}^{a}_{ij}[s]\big] + \sqrt{\frac{\ln(m)}{\overline{T}^{a}_{ij}[s]}}\Big).$$
(42)

We define for $k \in \mathbb{N}_{\geq 1}$, $a \in \mathcal{A}$, and $(ij) \in \mathcal{L}^a$ the quantities

$$u_k = \ln^{\beta}(k), \qquad \eta_k = \ln^{\frac{\beta+1}{2}}(k), \qquad \sigma^a_{ijk} = x^a_{ij}H_0(u_k - 2\eta_k).$$
 (43)

We are interested in the event that the number of type-(*ij*) departures under action a after $\lceil C_{\beta} + u_k \rceil$ episodes is at least $\lceil \sigma_{ijk}^a \rceil$, i.e.,

$$E_k^a = \left\{ \overline{T}_{ij}^a \big[\left[C_\beta + u_k \right] \right] \ge \left[\sigma_{ijk}^a \right], \ \forall (ij) \in \mathcal{L}^a \right\}.$$

$$(44)$$

Lemma 8. Let $\beta \in \mathbb{R}_{>1}$ and $a \in \mathcal{A}$. Then $\lim_{k\to\infty} k\mathbb{P}((E_k^a)^c) = 0$.

The proof of Lemma 8 in Appendix C.8 relies on Lemma 7. We use that the number of departures is at least as high as the number of departures in a coupled system that starts from the empty state at the start of the episode (the 'worst case' scenario). We then apply Lemma 7 to show that the worst case process reaches stationarity with high probability. Next, we use Poisson merging and splitting properties (see e.g. [48]) and Burke's Theorem [49, II.2.4 Theorem 2.1] to obtain that the *stationary* departure process of the FCFS RR(a) policy in Algorithm 2 is a Poisson process with rate x_{ij}^a . Lastly, we use a Poisson tail bound (see Lemma C.3) to obtain a lower bound on the number of departures of the stationary process.

For the second part of event (A), Lemma 9 shows that the probability of overestimating the true reward is small if the number of obtained samples (departures) is sufficiently large. The proof in Appendix C.9 relies on Hoeffding's inequality [50, Theorem 2].

Lemma 9. Let
$$\beta \in \mathbb{R}_{>1}$$
, $a \in \mathcal{A}$, and $\xi_k := \ln^{-\frac{\beta}{4}}(k)$, then $\lim_{k \to \infty} \sum_{s=\lceil C_{\beta}+u_k \rceil}^k \mathbb{P}(\overline{U}^a[s,k]-r^a \ge d^a-\xi_k, E_k^a) = 0$.

The value σ_{ijk}^a in (43) is constructed precisely such that Lemma 8 and Lemma 9 hold simultaneously. The intuition behind this value is as follows: if we have observed u_k episodes with action a, then for approximately $u_k - \eta_k$ of those episodes, the number of departures can be bounded from below by the number of departures of the stationary process, which is Poisson distributed. From these episodes, we lose approximately η_k episodes where the Poisson tail bound fails (see Lemma C.3). Hence, we collect payoff samples (departures) at a rate of approximately $u_k - 2\eta_k$, as reflected in the definition of σ_{ijk}^a .

Bound on event (B). For event (B), recall that the true mean of action $a \in \mathcal{A}$ is r^a as defined in (28). By construction, the UCB index in (33) includes a confidence bound (exploration bonus) which implies that the probability of event (B) is small. This is made precise in Lemma 10, which is proven in Appendix C.10. We let $\|\cdot\|_2$ denote the L^2 -norm.

Lemma 10. Let $\beta \in \mathbb{R}_{>1}$, $a \in \mathcal{A}$, $k \in \mathbb{N}_{\geq 1}$, and $0 < \xi < r^a$. Then, $\sum_{m=1}^k \mathbb{P}(U^a(m) < r^a - \xi) \le \pi^2 \|\lambda\|_2^2 / (12\xi^2)$.

4.3.3 Step 3. Bounding the number of suboptimal episodes

We define $S^{a}(k)$ as the number of episodes where Algorithm 1 chooses action $a \in \mathcal{A}$ in Line 3 up to and including episode number k, i.e.,

$$S^{a}(k) := \sum_{m=1}^{k} \mathbb{1}\{A_{m} = a\}.$$
(45)

From Lemma 11 it follows that the expected number of suboptimal episodes grows asymptotically with rate at most $\ln^{\beta}(k)$ with the number of episodes k. The proof of Lemma 11 is given in Appendix C.11, and relies on Lemma 8, 9, and 10.

Lemma 11. Let $\beta \in \mathbb{R}_{>1}$. For any suboptimal action $a \in \mathcal{A} \setminus \{1\}$ we have

$$\lim_{k \to \infty} \frac{\mathbb{E}(S^a(k))}{\ln^\beta(k)} \le 1.$$
(46)

4.3.4 Step 4. Bounding the regret of Algorithm 1

We bound the terms in the regret decomposition (20) individually. For term I, we use that suboptimal lines are only used in suboptimal episodes by construction. Hence, for $(k\ell) \in O^c(\theta)$, the expected number of type- $(k\ell)$ departures can be upper bounded by the maximal departure rate of server ℓ , which is $\mu_{\ell} - \varepsilon$ by construction, multiplied by the total number of suboptimal episodes. This, we bound using Lemma 11.

For term II, we show that the queue length process in consecutive optimal episodes converges sufficiently fast to its stationary behavior. Specifically, we use the bound in Lemma 7: if $A_k = A_{k+1} = 1$ and the queue length processes reaches stationarily within the warmup period of episode k, then the process is stationary at the start of episode k + 1, since the stationary measure does not change in this case.

Lastly, for term III we use the fact that x^{A_m} satisfies (10c) to show that the virtual queues of all servers are positive recurrent under the FCFS $\operatorname{RR}(A_m)$ policy. In view of (10b), this implies that the queue of any customer type is positive recurrent as well. This means that the expected queue lengths remain finite, and hence the contribution of term III to the regret vanishes on a logarithmic scale.

4.4 Discussion

4.4.1 Avoiding carryover effect between episodes

The reallocation in Lines 5-6 avoids a delay in observations when a new episode starts: if the chosen action in episode k differs from the action that was chosen in the previous episode k-1, we reallocate all waiting customers to the virtual queues according to new rates of the action chosen in episode k. After the new reallocation, we order the customers in each virtual queue in order of their arrival times, so that the service order within the virtual queue is FCFS. This reallocation improves the learning efficiency since there is no 'carryover' effect between episodes, unlike other learning algorithms in literature [37, 22].

4.4.2 Transfer learning

The rewards of the actions are related by the common unknown payoff parameter θ via (28). Algorithm 1 utilizes transfer learning in the sense that knowledge about the payoff parameter is shared between actions: specifically, the UCB index of a reward is computed using the UCB indices of the lines. This effect also becomes evident in the numerical analysis in Section 5.1.

Our regret bound in Theorem 1 is pessimistic: in analyzing the convergence of the UCB index U^a in (33), we only account for departures that are obtained in episodes where action a was chosen. However, we possibly observe type-(ij) departures as well in episodes where another action \tilde{a} with $x_{ij}^{\tilde{a}} > 0$ was chosen. This insight can be used in future work to improve the bound in Lemma 8 and in turn improve the regret bound in Theorem 2. In particular, it might be possible to replace the cardinality of the action $|\mathcal{A}|$ in (37) by the number of lines L, which is typically smaller. Such analysis methods are often used in the setting of linear bandits [11].

4.4.3 Parameter values

The required lower bound on α in (35) can be large when the value of the slack parameter $\varepsilon > 0$ is small. As a consequence, the episode length H_k in (36) increases quickly in the number of episodes k. Since Algorithm 1 only chooses a new action at the start of episodes, this can in turn lead to a slow decrease in the rate of regret accumulation. (35) is constructed in such a way to bound a term in the proof of Lemma 7. However, this bound is quite loose. More careful analysis of the individual terms can potentially give a lower value than (35). We show in Section 5.3 that Algorithm 1 still performs well in finite time, even when α is chosen smaller than the lower bound in (35).

4.4.4 Number of actions

The number of actions grows exponentially with the number of queues I, servers J, and lines L in the queueing system. Concretely, the number of bases of $LP(\theta, \varepsilon)$ in (10) is $\binom{L+J}{I+J}$, hence the number of basic feasible solutions grows exponentially with the system's complexity. Reducing the number of actions using dimension reduction techniques is appealing for future research. For example, we could apply a column generation technique on the utility function that uses the UCB indices as proxy for the true payoff parameters [51], or use a branch– and–bound technique like Bender's decomposition to split the optimization problem into a master problem and subproblems [52].

5 Numerical results

In this section we analyze the properties of Algorithm 1 (UCB QR) numerically. In Section 5.1, we consider a small skill-based queueing system. For this system, we analyze the regret of the UCB QR algorithm and compare its long-term average payoff rate with several benchmark routing policies. In Section 5.2, we analyze the robustness of the learning policy against changes in the true payoff parameter. In Section 5.3, we consider a larger queueing system and compare the performance of UCB QR against benchmark policies in several scenarios.

We consider four different benchmark policies which are described as follows.

- Oracle: Almost the same as the UCB QR algorithm, but with Line 3 replaced by $A_k = \operatorname{argmax}_{\tilde{a}} r^{\tilde{a}}$. Note that this policy relies on the true payoff parameter θ . This policy represents the optimal upper bound for the long-term reward rate of any stabilizing policy.
- FCFS ALIS: If there are nonempty compatible queues at a service completion, assign the customer with maximal waiting time. If there are compatible servers idle upon a customer arrival, assign it to the server with maximal idle time. This policy is widely used in practice since both customers and servers experience a sense of fairness [15].
- Greedy: If there are nonempty compatible queues at a service completion of server j, assign the first-inline customer from the queue $\operatorname{argmax}_{i \in \mathcal{C}_j} \theta_{ij}$. If there are compatible servers idle upon a type-*i* customer arrival, assign it to the server $\operatorname{argmax}_{j \in \mathcal{S}_i} \theta_{ij}$. This policy myopically optimizes the instantaneous payoff without considering long-term effects or stability constraints.
- Random: If there are nonempty compatible queues at a service completion, choose one of these queues uniformly at random and assign the first-in-line customer. If there are compatible servers idle at a customer arrival, assign it to one of these servers uniformly at random. This policy serves as a most naive baseline by making completely uninformed decisions.

5.1 Small example

We consider the queueing system illustrated on the left in Figure 7 with $\varepsilon = 0.5$. It can be verified that (12) is satisfied. LP(θ, ε) in (10) has six nondegenerate basic feasible solutions which we label as actions in Figure 7. The optimality of the actions depends on the payoff vector $\theta = \{\theta_{11}, \theta_{12}, \theta_{21}, \theta_{22}\}$ as illustrated in Figure 8.



Figure 7: A skill-based queueing system with I = 2 customer classes, J = 2 servers, and compatibility lines $\mathcal{L} = \{(11), (12), (21), (22)\}$, along with the six different actions. For each action, the routing rates $\{x_{ij}\}_{(ij)\in\mathcal{L}}$ are illustrated next to the corresponding lines.



Figure 8: Optimality conditions for each of the six feasible solutions of $LP(\theta, \varepsilon)$.

We set the true payoff vector as $\theta = \{0.4, 0.1, 0.3, 0.01\}$. This implies that action 2 is optimal and that the optimal long-term payoff rate is approximately 5.4. Moreover, line (12) is suboptimal since it has zero load in the optimal solution. The suboptimality gap of line (12) (recall (19)) is $\phi_{12} = 0.01$. The suboptimality gaps for each action in (29) are given by

$$d^{1} = 1.305, d^{2} = 0, d^{3} = 1.755, d^{4} = 0.055, d^{5} = 1.84, d^{6} = 1.405.$$
 (47)

Recall that the UCB QR algorithm requires configuration parameters α, β , and H_0 which define the episode length according to (36). We set $\alpha = 364$, $\beta = 1.01$, and $H_0 = 10$. These parameters satisfy (35) and (36). All of our numerical results are based on 50 independent replications and all plots show 95% confidence areas, although these intervals are too small to be visible in Figure 9.

We observe in Figure 9 that, as expected, the payoff rate of the UCB QR algorithm converges to the optimal payoff rate of the Oracle policy. In the initial period up to approximately time 2000, the UCB QR algorithm has not yet collected sufficient samples to distinguish the actions. Therefore, it chooses actions uniformly at random, which corresponds to the first few episodes in Figure 11. In turn, the regret accumulation is high since actions with large suboptimality gaps are chosen. After the initial period, we observe in Figure 11 that the algorithm primarily struggles to differentiate the optimal action 2 and action 4. This leads to a sharp decrease in regret accumulation, since action 4 has a small suboptimality gap.

Moreover, we observe in Figure 11 that actions $a \in \{1, 3, 5, 6\}$ are hardly ever chosen by the UCB QR algorithm. As opposed to a classical MAB, the rewards of the actions are related by the common unknown payoff parameter θ . Therefore, it suffices to only sample a subset of actions, as long as we observe sufficient payoff samples from all lines. By only using actions 2 and 4, the algorithm infers sufficient information about all four payoff parameters. In particular, in accordance with Line 12 in Algorithm 1, the indices U_{11} , U_{21} , and U_{22} are updated after episodes with action 2, while U_{11} , U_{12} , and U_{21} are updated after episodes with action 4 does not exploit this feature of transfer learning in the construction of the regret upper bound. This remains an interesting direction for future research.

The FCFS ALIS, Greedy and Random policies maintain a consistent gap to the optimal payoff of the Oracle policy. The value of this gap depends on the true payoff parameter θ . Since the total load in the system is far from critical, the long-term routing rates under the FCFS ALIS policy are close to those of the Random policy. This leads to the minor difference in total payoff rate with respect to the Random policy, as shown in Figure 9. In particular, the FCFS ALIS policy is indistinguishable from the Random policy in terms of reward. The Greedy policy obtains a slightly higher reward making a greedy decision whenever possible.

To summarize, we have shown for a small queueing system that the payoff rate of the UCB QR algorithm converges to the optimal rate and that it chooses the optimal action most of the time. Thereby, it outperforms our benchmark policies that (a) are agnostic to the payoff parameters, or (b) rely on the payoff parameters but make suboptimal routing decisions.



Figure 9: Running average payoff rate over time for all policies.



Figure 10: Regret accumulation over time of the UCB QR algorithm.



Figure 11: Cumulative count of actions over the first 20 episodes of the UCB QR algorithm.

5.2 Changing parameters

Although our theoretical analysis of the UCB QR algorithm in Section 4 is for a static environment, we analyze its robustness against changes in the payoff parameter θ . In this experiment, we use the same payoff vector as in Section 5.1 initially. At one-third of the total runtime, we update the payoff parameter of line (12) to $\theta'_{12} = 0.5$. As a consequence, the optimal action under the new payoff parameter is action 6. We set $\alpha = H_0 = 10$, and $\beta = 1.01$. Our numerical results are based on 100 independent replications.

Observe in Figure 12 that upon the parameter change (at episode number 60, depicted by a vertical line), the UCB QR algorithm at first prefers action 4. This makes sense since action 4 is closest to the initially optimal action 2, while it also includes line (12) which is an optimal line under the updated parameter θ' . From approximately episode number 100 onwards, the learning algorithm switches more to the optimal action 6. Hence, we see that the UCB QR algorithm suffers from a switch-over period, but eventually changes to the optimal action. The convergence after the parameter change is more slow than the initial convergence. We see this effect since the empirical estimators take into account the complete history. Therefore, after the parameter update, learning is hindered by samples obtained before the update. We expect that robustness against changing parameter values can be improved by decreasing the episode length, or altering the empirical estimators in (31) by decreasing the weight of observations from the past. This remains open for future work.

Hence, the numerical results suggest that the UCB QR algorithm can correctly identify changes in the true system parameters.



Figure 12: Cumulative count of actions per episode of the UCB QR algorithm when changing the true payoff parameter of line (21) to $\theta'_{21} = 0.5$ at one-third of the total runtime.

5.3 Complex queueing system

We consider the queueing system illustrated in Figure 13. This system is inspired by the operation of the call center of a real-world telecommunications company. We let $\varepsilon = 0.05$ such that (12) is satisfied. The number of basic feasible solutions of LP(θ, ε) in (10), i.e., the number of actions, is 88. The lower bound in (35) is 165088. However, since a large α slows down learning, we set $\alpha = 10$. We set $\beta = 1.01$, and $H_0 = 10$. Our numerical results are based on 100 independent replications.



Figure 13: Big queueing system with I = 5 customer classes and J = 5 servers. The parameter values are shown in the figure.

We consider three different scenarios with different levels of difficulty for adaptive learning:

- Initial parameters: The first scenario is the system with initial parameters as shown in Figure 13.
- Minimal payoff discrepancy: In this scenario, all-but-one payoff parameters are equal. In particular, $\theta_{ij} = 0.5$ for all $(ij) \in \mathcal{L}$ except for $\theta_{55} = 0.6$. The learning policy must identify this minimal payoff gap.
- Balanced arrival rates: In the third scenario, the arrival rates are equal for all customers, namely $\lambda_i = 42$ for all $i \in \mathcal{I}$. This value is chosen so that the total load of the system is high, namely approximately 0.98.

We compare the UCB QR algorithm with the other benchmark policies as described above. The average number of customers in the system is provided in Table 1 and the payoff rate is illustrated in Figure 14.

Observe in Table 1 that the UCB QR and Oracle policies have a similar average number of customers in the system in all scenarios. The average queue lengths of the FCFS ALIS policy is much smaller than those of Oracle and the UCB QR algorithm. This is to be expected, since this policy inherently aims to decrease the queue length. However, the nonidling nature of this policy leads to a gap in payoff rate with respect to the Oracle policy, as can be seen in Figure 14. For the Greedy and Random policies, the average number of customers in the system is much higher than for the other policies. Despite the high average queue length, the average payoff of Greedy and Random in Figure 14 is very comparable to those of FCFS ALIS. This is explained as follows: the policies obtain high reward by routing customer types with high average payoffs, while other customer types with overall lower average payoffs are not served at all. We find that for the initial parameters and minimal payoff discrepancy scenarios, especially the queue length of customer type 4 is large for these policies, since this customer type is only compatible with server 5 while the service capacity of server 5 is for a large part used by customer types 3 and 5. For this reason, the Greedy policy obtains an even lower average reward than the Random policy.

Observe also in Figure 14 that the convergence of the payoff rate of the UCB QR algorithm to the optimal payoff of the Oracle policy is much faster than in the small queueing system considered in Section 5.1 (see Figure 9), i.e., the regret is smaller. This can be explained by the choice of α (10 vs. 364). A smaller value of α implies that the episodes are shorter. Hence, per time interval there are more decision moments where the policy can switch between actions, which speeds up learning.

Comparing between the different scenarios, we find that in the minimal payoff discrepancy scenario, the UCB QR algorithm chooses suboptimal actions more frequently than in the other scenarios. However, since the suboptimality gaps of the suboptimal actions are small by construction, the average reward is still close to the optimal, as shown in Figure 14b. Out of the three scenarios, the convergence is slowest in Figure 14c. Our belief is that in this case, the algorithm needs approximately the same number of suboptimal episodes to learn the payoff parameters as in the initial scenario, while the suboptimality gaps are larger, i.e., the cost of exploration is higher.

Lastly, we note that Algorithm 1, although it maintains system stability by construction, it is not incentivized to minimize or balance the queue lengths or server loads. Possible extensions of the algorithm where customer waiting times, server loads or fairness constraints are taken into account are interesting for future research. For example, $LP(\theta, \varepsilon)$ in (10) can be altered to include either waiting time constraints, or a penalty factor proportional to the average queue length in the objective function. This would result in a different action space for the algorithm.

To summarize the findings of this section, we have shown that even in a complex queueing system with 88 different actions, the UCB QR algorithm converges quickly to the oracle reward, while maintaining reasonable queue lengths in different scenarios.

Policy	Initial parameters		Min. payoff discrepancy		Balanced arrival rates	
	E	σ	E	σ	\mathbb{E}	σ
Oracle	329	106	269	80	277	90
UCB QR	315	101	147	31	248	71
FCFS ALIS	9	0.3	12	0.8	12	0.8
Greedy	14069	142	17372	148	14193	207
Random	8491	138	8483	130	9447	130

Table 1: Mean (\mathbb{E}) and standard deviation (σ) of the number of customers in the system over the entire simulation time for all policies and scenarios.



Figure 14: Running average payoff rate over time for all policies and scenarios. The legend is the same for all subplots.

A Proof of Theorem 1

Let π satisfy (7) and (8), and let $\theta \in \mathbb{R}_{\geq 0}^{L}$.

We first prove that the regret lower bound is strictly positive, i.e., the second inequality in (25). We have $K(\theta, k\ell) < \infty$ for any $(k\ell) \in O^c(\theta)$ as a consequence of Lemma 6 and assumption (6). Moreover, $\phi_{k\ell} > 0$ for any $(k\ell) \in O^c(\theta)$ by strict complementarity. It remains to be shown that $O^c(\theta)$ is nonempty. Let $B = \mathscr{L} \cup \mathscr{J}$ be the optimal basis of $LP(\theta, \varepsilon)$, then |B| = I + J and $\mathscr{L} = O(\theta)$ by construction. It follows from Lemma 1 that $|\mathscr{J}| \ge 1$ (since the corresponding spanning forest consists of at least one tree), hence $|O(\theta)| = |\mathscr{L}| \le I + J - 1$. Thus, since $O(\theta) \cup O^c(\theta)$, is a partition of all L lines,

$$|O^{c}(\theta)| = L - |O(\theta)| \ge L - (I + J - 1) > 0,$$
(48)

where the last inequality holds by assumption (see Section 2.1). This means that $O^{c}(\theta) \neq \emptyset$ and thus the regret lower bound is strictly positive.

Next, we prove the first inequality in (25). Using the regret decomposition in (20), it is sufficient to show that simultaneously

$$\lim_{t \to \infty} \frac{1}{\ln(t)} \left(\sum_{j \in \mathcal{J}} w_j^{\theta} \left(t(\mu_j - \varepsilon) - \sum_{i \in \mathcal{C}_j} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t)) \right) + \sum_{i \in \mathcal{I}} v_i^{\theta} \left(t\lambda_i - \sum_{j \in \mathcal{S}_i} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t)) \right) \right) \ge 0,$$
(49)

and

$$\lim_{t \to \infty} \frac{\mathbb{E}_{\pi}^{\theta}(D_{k\ell}(t))}{\ln(t)} \ge \frac{1}{K(\theta, k\ell)}, \quad \forall (k\ell) \in O^{c}(\theta).$$
(50)

Proof of (49).

Since $v^{\theta} \in \mathbb{R}^{I}$, $w^{\theta} \in \mathbb{R}_{\geq 0}^{J}$ is the optimal solution of $D(\theta, \varepsilon)$ in (16), we have for any $j \in \mathcal{J}$ that $w_{j}^{\theta} \geq 0$ by (16d). Moreover, π satisfies (8) by assumption. Therefore

$$\lim_{t \to \infty} \frac{w_j^{\theta} \left(t(\mu_j - \varepsilon) - \sum_{i \in \mathcal{C}_j} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t)) \right)}{\ln(t)} \ge 0, \quad \forall j \in \mathcal{J}.$$
(51)

On the other hand, we have for any $i \in \mathcal{I}$ that $v_i^{\theta} \in \mathbb{R}$. Hence by (7)

$$\lim_{t \to \infty} \frac{v_i^{\theta} \left(t \lambda_i - \sum_{j \in \mathcal{S}_i} \mathbb{E}_{\pi}^{\theta} (D_{ij}(t)) \right)}{\ln(t)} = 0, \quad \forall i \in \mathcal{I}.$$
(52)

Combining (51) with (52) and summing over all servers and customer types completes the proof of (49).

Proof of (50).

Let $(k\ell) \in O^c(\theta)$. By Markov's inequality, for any $t \ge 0$ and for any $0 < \eta < 1$,

$$\frac{\mathbb{E}_{\pi}^{\theta}(D_{k\ell}(t))}{\ln(t)} \ge \frac{1-\eta}{K(\theta,k\ell)} \mathbb{P}_{\pi}^{\theta} \Big(D_{k\ell}(t) \ge \frac{(1-\eta)\ln(t)}{K(\theta,k\ell)} \Big).$$
(53)

Hence, to prove (50) it suffices to show that for $0 < \eta < 1$

$$\lim_{t \to \infty} \mathbb{P}^{\theta}_{\pi} \left(D_{k\ell}(t) < \frac{(1-\eta)\ln(t)}{K(\theta, k\ell)} \right) = 0.$$
(54)

Let $0 < \eta < 1$, and let $a, \delta \in \mathbb{R}$ be such that $0 < \delta < \eta/(2 - \eta)$ and $0 < a < \delta$. We note that $I(\cdot, \cdot)$ is continuous and by (6) finite, and that $0 < K(\theta, k\ell) < \infty$ by Lemma 6. This means that we can find a $\theta'_{k\ell} \in \Delta(\theta, k\ell)$ such that

$$0 < K(\theta, k\ell) \le I(\theta_{k\ell}, \theta'_{k\ell}) < (1+\delta)K(\theta, k\ell) < \infty.$$
(55)

Note that the choice of δ implies $1 - \eta < (1 - \delta)/(1 + \delta)$, and therefore,

$$\mathbb{P}_{\pi}^{\theta}\left(D_{k\ell}(t) < \frac{(1-\eta)\ln(t)}{K(\theta,k\ell)}\right) \leq \mathbb{P}_{\pi}^{\theta}\left(D_{k\ell}(t) < \frac{1-\delta}{1+\delta}\frac{\ln(t)}{K(\theta,k\ell)}\right) \stackrel{(55)}{\leq} \mathbb{P}_{\pi}^{\theta}\left(D_{k\ell}(t) < \frac{(1-\delta)\ln(t)}{I(\theta_{k\ell},\theta_{k\ell}')}\right). \tag{56}$$

We define the following functions,

$$g(t) := \frac{(1-\delta)\ln(t)}{I(\theta_{k\ell}, \theta'_{k\ell})}, \quad c(t) := (1-a)\ln(t).$$
(57)

We next prove that

$$\lim_{t \to \infty} \mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t)) = 0.$$
(58)

Consider the log-likelihood ratio of the parameter values $\theta_{k\ell}$ and $\theta'_{k\ell}$ given a sequence X_1, X_2, \ldots, X_n of observations from the reward distribution belonging to type-k customers being served by server ℓ , which is defined by

$$L_n = \sum_{i=1}^n l_i = \sum_{i=1}^n \ln\left(\frac{P_{\theta_{k\ell}}(X_i)}{P_{\theta_{k\ell'}}(X_i)}\right).$$
(59)

By the law of total probability, we have

$$\mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t)) = \mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), \ L_{D_{k\ell}(t)} > c(t)) + \mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), \ L_{D_{k\ell}(t)} \le c(t)).$$
(60)

To complete the proof, we will show that both terms of (60) are o(1) as $t \to \infty$, i.e., we will prove separately that

$$\lim_{t \to \infty} \mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} > c(t)) = 0$$

$$\tag{61}$$

and

$$\lim_{t \to \infty} \mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} \le c(t)) = 0.$$
(62)

Proof of (61). In order to prove (61), we will use the following result, proven in [10, Lemma 2(ii)]: Let $\{Z_i\}_{i\in\mathbb{N}}$ be an i.i.d. sequence of random variables that satisfies the strong law of large numbers, i.e., $\mathbb{P}\left(\lim_{n\to\infty}\sum_{i=1}^{n}Z_i/n=\mu\right)=1$, for some constant μ . Let h_n be an increasing sequence of positive constants such that $h_n \to \infty$ as $n \to \infty$. Then, for any $\zeta > 0$, $\lim_{n\to\infty} \mathbb{P}\left(\max_{s \le \lfloor h_n \rfloor} sum_{i=1}^s Z_i/h_n > (1+\zeta)\mu\right) = 0$. We leverage the definition of the log-likelihood ratio (59) and the properties of the maximum to write

$$\mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), \ L_{D_{k\ell}(t)} > c(t)) \le \mathbb{P}^{\theta}_{\pi}\left(\max_{s \le \lfloor g(t) \rfloor} L_s > c(t)\right).$$
(63)

Next, we divide by g(t) > 0:

$$\mathbb{P}_{\pi}^{\theta}\left(\max_{s\leq \lfloor g(t)\rfloor} L_{s} > c(t)\right) = \mathbb{P}_{\pi}^{\theta}\left(\max_{s\leq \lfloor g(t)\rfloor} \frac{L_{s}}{g(t)} > \frac{c(t)}{g(t)}\right)$$
(64)

$$\stackrel{57)}{=} \mathbb{P}^{\theta}_{\pi} \left(\max_{s \le \lfloor g(t) \rfloor} \frac{L_s}{g(t)} > \frac{1-a}{1-\delta} I(\theta_{k\ell}, \theta'_{k\ell}) \right)$$
(65)

$$= \mathbb{P}_{\pi}^{\theta} \left(\max_{s \le \lfloor g(t) \rfloor} \frac{L_s}{g(t)} > \left(1 + \frac{\delta - a}{1 - \delta} \right) I(\theta_{k\ell}, \theta'_{k\ell}) \right).$$
(66)

We make a few observations. Firstly, note that $\ln(t) \to \infty$ and therefore $g(t) \to \infty$ as $t \to \infty$. Secondly, $(\delta - a)/(1 - \delta) > 0$, since $0 < a < \delta$ by definition and $\delta < 1$. Thirdly, by definition, for every $i \in [n]$

$$\mathbb{E}^{\theta}_{\pi}(l_i) = \mathbb{E}^{\theta}_{\pi} \left(\ln \left(\frac{P_{\theta_{k\ell}}(X_i)}{P_{\theta_{k\ell'}}(X_i)} \right) \right) = I(\theta_{k\ell}, \theta'_{k\ell}), \tag{67}$$

hence by the strong law of large numbers,

$$\mathbb{P}^{\theta}_{\pi}\left(\lim_{n \to \infty} \frac{L_n}{n} = I(\theta_{k\ell}, \theta'_{k\ell})\right) = 1.$$
(68)

Therefore, we can apply [10, Lemma 2(ii)] to conclude that

$$\lim_{t \to \infty} \mathbb{P}_{\pi}^{\theta} \left(\max_{s \le \lfloor g(t) \rfloor} \frac{L_s}{g(t)} > \left(1 + \frac{\delta - a}{1 - \delta} \right) I(\theta_{k\ell}, \theta'_{k\ell}) \right) = 0.$$
(69)

Combining with (63), this completes the proof of (61).

Proof of (62). Consider the same queueing network structure but with a slightly different payoff vector $\theta' = A(\theta, k\ell, \theta'_{k\ell})$, which will be our 'confusing' system. Let x' be the optimal solution to $LP(\theta', \varepsilon)$. In order to prove (62), we describe a change–of–measure transformation to relate the measure of an event under probability law $\mathbb{P}^{\theta}_{\pi}$ by the measure of that same event under the law $\mathbb{P}_{\pi}^{\theta'}$.

Denote the event $\{D_{k\ell}(t) = m, L_m \leq c(t)\}$ by B. Given event B and noting that θ and θ' differ only at index $k\ell$, we find that the RN derivative of $\mathbb{P}^{\theta}_{\pi}$ with respect to $\mathbb{P}^{\theta'}_{\pi}$ has the form $d\mathbb{P}^{\theta}_{\pi}/d\mathbb{P}^{\theta'}_{\pi} = \prod_{i=1}^{m} P_{\theta_{k\ell}}(X_i)/P_{\theta_{k\ell'}}(X_i)$. Moreover, the rewards are sampled independently from the rest of the system. Hence, we can write

$$\mathbb{P}^{\theta}_{\pi}(B) = \int_{B} \prod_{i=1}^{m} \frac{P_{\theta_{k\ell}}(X_i)}{P_{\theta_{k\ell'}}(X_i)} \mathrm{d}\mathbb{P}^{\theta'}_{\pi}.$$
(70)

We exponentiate and use the definition of L_m to write

$$\mathbb{P}_{\pi}^{\theta}(B) = \int_{B} \exp\left(\sum_{i=1}^{m} \ln\left(\frac{P_{\theta_{k\ell}}(X_i)}{P_{\theta_{k\ell'}}(X_i)}\right)\right) d\mathbb{P}_{\pi}^{\theta'} = \int_{B} e^{L_m} d\mathbb{P}_{\pi}^{\theta'}.$$
(71)

Lastly, the definition of B implies that $L_m \leq c(t)$, hence

$$\mathbb{P}^{\theta}_{\pi}(B) \leq \int_{B} e^{c(t)} \mathrm{d}\mathbb{P}^{\theta'}_{\pi} = e^{c(t)} \mathbb{P}^{\theta'}_{\pi}(B).$$
(72)

We next prove (62). Note that $\{D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} \leq c(t)\}$ is a disjoint union of events of the form $\{D_{k\ell}(t) = m, L_{D_{k\ell}(t)} \leq c(t)\}$ for m < g(t), so we can apply the change–of–measure transformation (72) to write

$$\mathbb{P}^{\theta}_{\pi}(D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} \le c(t)) \le e^{c(t)} \mathbb{P}^{\theta'}_{\pi}(D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} \le c(t)).$$
(73)

Recall (21), (22), and that $\theta'_{k\ell} \in \Delta(\theta, k\ell)$ by assumption, implying $(k\ell) \in O(\theta')$. Therefore, by the definition of g(t) in (57) and assumption (24),

$$\mathbb{P}_{\pi}^{\theta'}(D_{k\ell}(t) < g(t), L_{D_{k\ell}(t)} \le c(t)) \le \mathbb{P}_{\pi}^{\theta'}(D_{k\ell}(t) < g(t)) = \mathbb{P}_{\pi}^{\theta'}\left(D_{k\ell}(t) < \frac{(1-\delta)\ln(t)}{I(\theta_{k\ell}, \theta'_{k\ell})}\right) = o(1) \text{ as } t \to \infty.$$
(74)

This completes the proof.

B Proof of Theorem 2

Let

$$\mathcal{T}_k := \sum_{m=1}^k H_m \tag{75}$$

denote the total duration of the first $k \in \mathbb{N}_{\geq 1}$ episodes. Then, $\mathcal{D}_{ij}(\mathcal{T}_k)$ is the number of type-(ij) departures up to and including episode k. Moreover, let \mathcal{D}_{ij}^m denote the number of type-(ij) departures within episode $m \in \mathbb{N}_{\geq 1}$. Note that this quantity is similar to D_{ij}^m as introduced in Section 4.3, although \mathcal{D}_{ij}^m includes the departures within the warmup time of the episode. Note that $D_{ij}(\mathcal{T}_k) = \sum_{m=1}^k \mathcal{D}_{ij}^m$ for any $(ij) \in \mathcal{L}$ and $k \in \mathbb{N}_{\geq 1}$.

First, we show that Algorithm 1 satisfies the consistency assumption (24). Let $\theta \in \mathbb{R}_{\geq 0}^{L}$, $(ij) \in O(\theta)$, $b \in \mathbb{R}_{>0}$, $k \in \mathbb{N}_{\geq 1}$ and $z_k := \lceil k/2 \rceil$. We use a similar analysis as in the proof of Lemma 8 in Appendix C.8. In particular, consider the stationary queue length process as introduced in Section 4.3.1 (dropping the action *a* from notation) and the event $\underline{\Omega}_{ij}^{m-1}$ as introduced in (184).

Recall also that A_m is the action chosen by Algorithm 1 in Line 3 in episode number m. Recall that 1 is the optimal action by assumption. Event $\underline{\Omega}_{ij}^{m-1}$ implies that the number of departures in episode m-1 after the warmup time is at least as large as the number of departures of the stationary queue length process. Consider the event $A_{m-1} = A_m = 1$. In this case, the stationary measure is the same for episodes m-1 and m, so $\underline{\Omega}_{ij}^{m-1}$ implies that $\mathcal{D}_{ij}^m \geq \hat{\mathcal{D}}_{ij}^m$, where $\hat{\mathcal{D}}_{ij}^m$ is the number of type-(ij) departures of the stationary queue length process during episode m. Lastly, from Burke's theorem [49, II.2.4 Theorem 2.1] and the Poisson split and merge properties [48], it follows that on the event $A_m = 1$, we have $\hat{\mathcal{D}}_{ij}^m \sim \text{Poi}(x_{ij}^1H_m)$.

By the law of total probability,

$$\mathbb{P}^{\theta}(D_{ij}(\mathcal{T}_k) < b \ln(\mathcal{T}_k)) \le \mathbb{P}^{\theta}\left(\sum_{m=2}^k \mathcal{D}_{ij}^m < b \ln(\mathcal{T}_k)\right)$$
(76)

$$\leq \mathbb{P}^{\theta} \left(\sum_{m=2}^{k} \mathcal{D}_{ij}^{m} < b \ln(\mathcal{T}_{k}) \mid \sum_{m=2}^{k} \mathbb{1} \left\{ A_{m} = A_{m-1} = 1, \ \underline{\Omega}_{ij}^{m-1} \right\} > z_{k} \right)$$
(77)

$$+ \mathbb{P}^{\theta} \left(\sum_{m=2}^{k} \mathbb{1} \left\{ A_m = A_{m-1} = 1, \ \underline{\Omega}_{ij}^{m-1} \right\} \le z_k \right).$$

$$(78)$$

We analyze these probabilities individually. Let $X_{ij}^m \sim \text{Poi}\left(x_{ij}^1 H_m\right)$ be independent across $m \in \mathbb{N}_{\geq 1}$. We note that $\mathcal{D}_{ij}^m \geq X_{ij}^m$ whenever $A_m = A_{m-1} = 1$ and $\underline{\Omega}_{ij}^{m-1}$, hence

$$\mathbb{P}^{\theta}\left(\sum_{m=2}^{k} \mathcal{D}_{ij}^{m} < b\ln(\mathcal{T}_{k}) \mid \sum_{m=2}^{k} \mathbb{1}\left\{A_{m} = A_{m-1} = 1, \ \underline{\Omega}_{ij}^{m-1}\right\} > z_{k}\right) \leq \mathbb{P}^{\theta}\left(\sum_{m=2}^{z_{k}} X_{ij}^{m} < b\ln(\mathcal{T}_{k})\right).$$
(79)

Note that $\sum_{m=2}^{z_k} X_{ij}^m$ is again Poisson distributed. Moreover, note that in light of the definition of \mathcal{T}_k in (75), there exists a $k_0 \in \mathbb{N}$ such that $\sum_{m=2}^{z_k} x_{ij}^1 H_m \ge b \ln(\mathcal{T}_k)$ for any $k \ge k_0$. Hence, it follows from Lemma C.3 in Appendix C.8 that for $k \ge k_0$,

<

$$\mathbb{P}^{\theta}\left(\sum_{m=2}^{z_{k}} X_{ij}^{m} < b\ln(\mathcal{T}_{k})\right) = \mathbb{P}^{\theta}\left(\sum_{m=2}^{z_{k}} X_{ij}^{m} - \sum_{m=2}^{z_{k}} x_{ij}^{1}H_{m} \le b\ln(\mathcal{T}_{k}) - \sum_{m=2}^{z_{k}} x_{ij}^{1}H_{m}\right)$$
(80)
$$\left(-(b\ln(\mathcal{T}_{k}) - \sum_{m=2}^{z_{k}} x_{ij}^{1} + U_{m})^{2}\right)$$

$$\leq \exp\left(\frac{-(b\ln(\mathcal{T}_k) - \sum_{m=2}^{z_k} x_{ij}^1 H_m)^2}{2\sum_{m=2}^{z_k} x_{ij}^1 H_m}\right).$$
(81)

It is readily verified that

$$\lim_{k \to \infty} \frac{(b \ln(\mathcal{T}_k) - \sum_{m=2}^{z_k} x_{ij}^1 H_m)^2}{2 \sum_{m=2}^{z_k} x_{ij}^1 H_m} = \infty.$$
(82)

Hence, (79) is o(1) as $k \to \infty$.

For the other probability in (78), note

$$\mathbb{P}^{\theta}\left(\sum_{m=2}^{k}\mathbb{1}\left\{A_{m}=A_{m-1}=1,\ \underline{\Omega}_{ij}^{m-1}\right\} \le z_{k}\right) = \mathbb{P}^{\theta}\left(k-\sum_{m=2}^{k}\mathbb{1}\left\{A_{m}=A_{m-1}=1,\ \underline{\Omega}_{ij}^{m-1}\right\} \ge k-z_{k}\right)$$
(83)

$$\leq \mathbb{P}^{\theta} \left(\sum_{m=2}^{k} \left(\mathbb{1}\left\{ A_m \neq 1 \right\} + \mathbb{1}\left\{ A_{m-1} \neq 1 \right\} + \mathbb{1}\left\{ \left(\underline{\Omega}_{ij}^{m-1}\right)^c \right\} \right) \geq k - z_k \right).$$
(84)

From Boole's inequality, we obtain

$$\mathbb{P}^{\theta}\left(\sum_{m=2}^{k} \left(\mathbb{1}\left\{A_{m} \neq 1\right\} + \mathbb{1}\left\{A_{m-1} \neq 1\right\} + \mathbb{1}\left\{\left(\underline{\Omega}_{ij}^{m-1}\right)^{c}\right\}\right) \ge k - z_{k}\right)$$

$$(85)$$

$$\leq 2\mathbb{P}^{\theta}\left(\sum_{m=1}^{k} \mathbb{1}\left\{A_m \neq 1\right\} \geq \frac{k - z_k}{3}\right) + \mathbb{P}^{\theta}\left(\sum_{m=1}^{k} \mathbb{1}\left\{\left(\underline{\Omega}_{ij}^{m-1}\right)^c\right\} \geq \frac{k - z_k}{3}\right).$$
(86)

In view of the definition of $S^{a}(k)$ in (45), $k - z_{k} = \Theta(k)$, Lemma 11, and Markov's inequality, we have

$$\mathbb{P}^{\theta}\left(\sum_{m=1}^{k} \mathbb{1}\left\{A_m \neq 1\right\} \ge \frac{k - z_k}{3}\right) = \mathbb{P}^{\theta}\left(\sum_{a \neq 1} S^a(k) \ge \frac{k - z_k}{3}\right) \le \frac{3\sum_{a \neq 1} \mathbb{E}(S^a(k))}{k - z_k} = o(1) \text{ as } k \to \infty.$$
(87)

Lastly, note that $\mathbb{P}^{\theta}((\underline{\Omega}_{ij}^{m})^{c}) \leq 1/m^{\beta}$ (see (191)) and that the $\underline{\Omega}_{ij}^{m}$ are mutually independent over all episodes. Since $k - z_{k} = \Theta(k)$ is increasing in k and since $\beta > 1$, we conclude that

$$\mathbb{P}^{\theta}\left(\sum_{m=1}^{k} \mathbb{1}\left\{\left(\underline{\Omega}_{ij}^{m-1}\right)^{c}\right\} \geq \frac{k-z_{k}}{3}\right) = o(1) \text{ as } k \to \infty.$$
(88)

Since $\theta \in \mathbb{R}_{\geq 0}^{L}$ was chosen arbitrarily, Algorithm 1 satisfies (24) for any $\xi \in \mathbb{R}_{\geq 0}^{L}$.

Next, we prove the regret upper bound by analyzing the terms in the regret decomposition (20) individually.

Analysis of term I in (20).

Let $(ij) \in O^{c}(\theta)$ be a suboptimal line. Since $(ij) \in O^{c}(\theta)$, there are no departures of type (ij) in episodes where Algorithm 1 chooses the optimal action 1. Therefore, using (75) and the law of total probability,

$$\mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k)) = \sum_{m=1}^k \mathbb{E}^{\theta}(\mathcal{D}_{ij}^m) = \sum_{m=1}^k \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta} (\mathcal{D}_{ij}^m \mid A_m = a) \mathbb{P}(A_m = a).$$
(89)

Note that the expected workload for server j under the FCFS $\operatorname{RR}(a)$ routing algorithm in Algorithm 2 does not exceed capacity $\mu_j - \varepsilon$, since the routing rates $x_{k\ell}^a$ for all lines $(k\ell) \in \mathcal{L}$ satisfy constraint (10c). Hence, for any $a \in \mathcal{A}$

$$\mathbb{E}^{\theta}\left(\mathcal{D}_{ij}^{m} \mid A_{m} = a\right) \leq H_{m}(\mu_{j} - \varepsilon).$$

$$\tag{90}$$

Applying (90) in (89) gives

$$\mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k)) \leq \sum_{m=1}^k \sum_{a \in \mathcal{A} \setminus \{1\}} H_m(\mu_j - \varepsilon) \mathbb{E}^{\theta}(\mathbb{1}\left\{A_m = a\right\}) \leq H_k(\mu_j - \varepsilon) \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta}(S^a(k)).$$
(91)

where we used in last inequality that the episode length H_m is nondecreasing in m by (36), and definition (45). We use $\mathcal{T}_k \geq k$ and Lemma 11 to obtain that for any suboptimal action $a \in \mathcal{A} \setminus \{1\}$,

$$\lim_{k \to \infty} \frac{\mathbb{E}^{\theta}(S^{a}(k))}{\ln^{\beta}(\mathcal{T}_{k})} \leq \lim_{k \to \infty} \frac{\mathbb{E}^{\theta}(S^{a}(k))}{\ln^{\beta}(k)} \leq 1.$$
(92)

Moreover, $\lim_{k\to\infty} \ln(2Jk)/\ln(k) = 1$ and $\alpha \ge 1$ (recall (35)), so $\lim_{k\to\infty} H_k/(\alpha \ln^\beta(k)) \le 1$. Hence,

$$\lim_{k \to \infty} \frac{\mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(\mathcal{T}_k)} \le \lim_{k \to \infty} \frac{H_k(\mu_j - \varepsilon) \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta}(S^a(k))}{\alpha \ln^{2\beta}(k)} \le (\mu_j - \varepsilon)|\mathcal{A}|.$$
(93)

Analysis of term II in (20).

Recall that $(v_i^{\theta})_{i \in \mathcal{I}}$ and $(w_j^{\theta})_{j \in \mathcal{J}}$ in (20) are the optimal dual variables, i.e., the minimizers of (16). By (16d), we have $w_j^{\theta} \ge 0$ for all $j \in \mathcal{J}$. The contribution of term II in (20) is clearly zero for $j \in \mathcal{J}$ with $w_j^{\theta} = 0$.

Let $j \in \mathcal{J}$ be such that $w_j > 0$. Recall that x^1 and v^{θ} , w^{θ} satisfy the complementary slackness conditions (18), thus $w_j^{\theta} > 0$ implies $\mu_j - \varepsilon = \sum_{i \in \mathcal{C}_j} x_{ij}^1$. Hence,

$$\lim_{k \to \infty} \frac{\mathcal{T}_k(\mu_j - \varepsilon) - \sum_{i \in \mathcal{C}_j} \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(\mathcal{T}_k)} = \lim_{k \to \infty} \sum_{i \in \mathcal{C}_i} \frac{\mathcal{T}_k x_{ij}^1 - \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(\mathcal{T}_k)}.$$
(94)

Let $i \in C_j$. By (75) and the law of total probability,

$$\mathcal{T}_k x_{ij}^1 - \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k)) = \sum_{m=1}^k \sum_{a \in \mathcal{A}} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) \mathbb{1} \left\{ A_m = a \right\} \right).$$
(95)

We consider the optimal action a = 1 and all suboptimal actions $a \neq 1$ separately.

For suboptimal actions, note that \mathcal{D}_{ij}^m is nonnegative, the episode length H_m is nondecreasing in m, and recall (45), hence

$$\sum_{m=1}^{\kappa} \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) \mathbb{1} \left\{ A_m = a \right\} \right) \le \sum_{m=1}^{\kappa} \sum_{a \in \mathcal{A} \setminus \{1\}} H_m x_{ij}^1 \mathbb{E}^{\theta} (\mathbb{1} \left\{ A_m = a \right\})$$
(96)

$$\leq H_k x_{ij}^1 \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta}(S^a(k)).$$
(97)

On the other hand, for the optimal action, we consider for each episode m also the preceding episode m - 1. We let $A_0 := 2$. Then

$$\sum_{m=1}^{k} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) \mathbb{1} \left\{ A_m = 1 \right\} \right) = \sum_{m=1}^{k} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) (\mathbb{1} \left\{ A_m = 1, A_{m-1} \neq 1 \right\} + \mathbb{1} \left\{ A_m = A_{m-1} = 1 \right\}) \right).$$
(98)

Since \mathcal{D}_{ij}^m is nonnegative we have, similar to (97),

$$\sum_{m=1}^{k} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) \mathbb{1} \left\{ A_m = 1, A_{m-1} \neq 1 \right\} \right) \le H_k x_{ij}^1 \sum_{m=1}^{k} \mathbb{E}^{\theta} (\mathbb{1} \left\{ A_{m-1} \neq 1 \right\}) = H_k x_{ij}^1 \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta} (S^a(k)).$$
(99)

For the other term in (98), consider the event $\underline{\Omega}_{ij}^{m-1}$ as introduced in (184). By the law of total probability,

$$\mathbb{E}^{\theta}\left((H_{m}x_{ij}^{1}-\mathcal{D}_{ij}^{m})\mathbb{1}\left\{A_{m}=1=A_{m-1}=1\right\}\right)=\mathbb{E}^{\theta}\left((H_{m}x_{ij}^{1}-\mathcal{D}_{ij}^{m})\mathbb{1}\left\{A_{m}=1=A_{m-1}=1\right\}\mid\underline{\Omega}_{ij}^{m-1}\right)\mathbb{P}^{\theta}(\underline{\Omega}_{ij}^{m-1})\quad(100)\\+\mathbb{E}^{\theta}\left((H_{m}x_{ij}^{1}-\mathcal{D}_{ij}^{m})\mathbb{1}\left\{A_{m}=1=A_{m-1}=1\right\}\mid(\underline{\Omega}_{ij}^{m-1})^{c}\right)\mathbb{P}^{\theta}((\underline{\Omega}_{ij}^{m-1})^{c}).$$
(101)

Since $\underline{\Omega}_{ij}^{m-1}$ implies that $\mathcal{D}_{ij}^m \ge \hat{\mathcal{D}}_{ij}^m$, and $\hat{D}_{ij}^m \mathbb{1}\left\{A_m = 1\right\} \sim \operatorname{Poi}\left(x_{ij}^1 H_m\right)$, we have

$$\mathbb{E}^{\theta}\left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m)\mathbb{1}\left\{A_m = 1 = A_{m-1} = 1\right\} \mid \underline{\Omega}_{ij}^{m-1}\right) \leq \mathbb{E}^{\theta}\left((H_m x_{ij}^1 - \hat{\mathcal{D}}_{ij}^m)\mathbb{1}\left\{A_m = 1 = A_{m-1} = 1\right\} \mid \underline{\Omega}_{ij}^{m-1}\right) = 0.$$
(102)

On the other hand, by (191), $\mathbb{P}^{\theta}((\underline{\Omega}_{ij}^{m-1})^c) \leq 1/m^{\beta}$. This, together with $\mathcal{D}_{ij}^m \geq 0$ gives

$$\mathbb{E}^{\theta}\left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m)\mathbb{1}\left\{A_m = 1 = A_{m-1} = 1\right\} \mid (\underline{\Omega}_{ij}^{m-1})^c\right)\mathbb{P}^{\theta}((\underline{\Omega}_{ij}^{m-1})^c) \le \frac{H_m x_{ij}^*}{m^{\beta}}.$$
(103)

Hence, by (101), (102), and (103) we have

$$\sum_{m=2}^{k} \mathbb{E}^{\theta} \left((H_m x_{ij}^1 - \mathcal{D}_{ij}^m) \mathbb{1} \left\{ A_m = A_{m-1} = 1 \right\} \right) \le \sum_{m=2}^{k} \frac{H_m x_{ij}^1}{m^{\beta}}.$$
 (104)

Substitution of (97), (99), and (104) into (95) yields

$$\mathcal{T}_k x_{ij}^1 - \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k)) \le 2H_k x_{ij}^1 \sum_{a \in \mathcal{A} \setminus \{1\}} \mathbb{E}^{\theta}(S^a(k)) + \sum_{m=2}^k \frac{H_m x_{ij}^1}{m^{\beta}}.$$
(105)

For the first term, we invoke Lemma 11 similarly as in (93). For the other term, we note that $m^{\beta-1} \ge H_m$ for $m \ge 1$ large enough since $\beta > 1$, and $\sum_{m=1}^{k} 1/m = \mathcal{O}(\ln(k))$. Hence,

$$\lim_{k \to \infty} \frac{\mathcal{T}_k x_{ij}^1 - \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(k)} \le x_{ij}^1 |\mathcal{A}|.$$
(106)

Concluding from (94) and (106) and using $\sum_{i \in \mathcal{J}} x_{ij}^1 \leq \mu_j - \varepsilon$, we have for any $j \in \mathcal{J}$ with $w_j^{\theta} > 0$ that

$$\lim_{k \to \infty} \frac{\mathcal{T}_k(\mu_j - \varepsilon) - \sum_{i \in \mathcal{C}_j} \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(\mathcal{T}_k)} \le (\mu_j - \varepsilon)|\mathcal{A}|.$$
(107)

Analysis of term III in (20).

Consider an episode $m \in \mathbb{N}_{\geq 1}$ and a customer type $i \in \mathcal{I}$. Algorithm 1 routes type-*i* customers to servers according to the routing rates $x_{ij}^{A_m}$ using the FCFS $\operatorname{RR}(A_m)$ policy as described in Algorithm 2. Since x^{A_m} is a basic feasible solution of $\operatorname{LP}(\theta, \varepsilon)$ in (10), it satisfies (10b). This means that any type-*i* customer is placed in a virtual queue of a compatible server $j \in S_i$. We next show that all virtual queues are positive recurrent.

Let $j \in \mathcal{J}$. Recall that for any $a \in \mathcal{A}$, x^a satisfies (10c). This implies that the queue length process $Q_j^m(t)$, $t \in [0, H_m)$ of the virtual queue of server j during episode m (as introduced in Section 4.3.1) has negative drift:

$$\lim_{h \to 0} \frac{\mathbb{E}^{\theta} \left(Q_j^m(t+h) \right) - \mathbb{E}^{\theta} \left(Q_j^m(t) \right)}{h} = \sum_{(ij) \in \mathcal{L}} x_{ij}^{Am} - \mu_j \le \mu_j - \varepsilon - \mu_j < 0.$$
(108)

From the Foster-Lyapunov criterion [53, Proposition 4.5] follows that the virtual queue length process is positive recurrent. Since all type-i customers are placed in a virtual queue, and all virtual queues are positive recurrent, (7) is satisfied.

From (7) it follows that

$$\lim_{k \to \infty} \frac{\mathcal{T}_k \lambda_i - \sum_{j \in \mathcal{S}_i} \mathbb{E}^{\theta}(D_{ij}(\mathcal{T}_k))}{\alpha \ln^{2\beta}(\mathcal{T}_k)} = 0.$$
(109)

To conclude the proof, we recall that the dual variables $(v_i^{\theta})_{i \in \mathcal{I}}$ and $(w_j^{\theta})_{j \in \mathcal{J}}$ are finite by strong duality [6, Theorem 4.4]. Hence, applying (93), (107), and (109) to (20) gives

$$\lim_{t \to \infty} \frac{R^{\theta}(t)}{\alpha \ln^{2\beta}(t)} = \lim_{k \to \infty} \frac{R^{\theta}(\mathcal{T}_k)}{\alpha \ln^{2\beta}(\mathcal{T}_k)} \le \sum_{(k\ell) \in O^c(\theta)} \phi^{\theta}_{k\ell}(\mu_{\ell} - \varepsilon)|\mathcal{A}| + \sum_{j \in \mathcal{J}} w^{\theta}_j(\mu_j - \varepsilon)|\mathcal{A}|.$$
(110)

C Remaining proofs

C.1 Proof of Lemma 1

Observe that $LP(\theta, \varepsilon)$ in (10) is of the form

$$LP(\theta,\varepsilon): \max_{x,\sigma} \begin{bmatrix} \theta \\ 0 \end{bmatrix}^{\top} \begin{bmatrix} x \\ \sigma \end{bmatrix}, \qquad (111a)$$

s.t.
$$\begin{bmatrix} A' & 0 \\ A'' & I \end{bmatrix} \begin{bmatrix} x \\ \sigma \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu - \varepsilon \end{bmatrix}, x \in \mathbb{R}^{L}_{\geq 0}, \sigma \in \mathbb{R}^{J}_{\geq 0}, \qquad (111b)$$

where

$$A' \in \mathbb{R}^{I \times L}, \ A'_{i,k\ell} = \begin{cases} 1 & \text{if } i = k \text{ and } \ell \in \mathcal{S}_i, \\ 0 & \text{otherwise,} \end{cases}$$
(112)

$$A'' \in \mathbb{R}^{J \times L}, \ A'_{j,k\ell} = \begin{cases} 1 & \text{if } j = \ell \text{ and } k \in \mathcal{C}_j, \\ 0 & \text{otherwise.} \end{cases}$$
(113)

" \Longrightarrow " Let $B = \mathscr{L} \cup \mathscr{J}$ be a basis of $LP(\theta, \varepsilon)$. Since B is a basis of $LP(\theta, \varepsilon)$, |B| = I + J and the matrix

$$\mathbf{B} = \begin{bmatrix} \mathbf{C}_{\mathscr{L}} & \mathbf{D}_{\mathscr{J}} \end{bmatrix} = \begin{bmatrix} A'_{\mathscr{L}} & 0_{\mathscr{J}} \\ A''_{\mathscr{L}} & I_{\mathscr{J}} \end{bmatrix} \in \mathbb{R}^{(I+J) \times (I+J)}$$
(114)

formed by the columns of $\mathscr{L} \cup \mathscr{J}$ has full rank. Here, $\mathbb{C}_{\mathscr{L}}$ is the node edge incidence matrix of $\mathcal{G}(\mathscr{L}, \mathscr{J})$. We will prove that (i) $\mathcal{G}(\mathscr{L}, \mathscr{J})$ contains no cycles, (ii) each $v \in \mathcal{I} \cup \mathcal{J}$ is contained in $\mathcal{G}(\mathscr{L}, \mathscr{J})$, and (iii) each tree of $\mathcal{G}(\mathscr{L}, \mathscr{J})$ contains a unique node $j \in \mathscr{J}$.

For (i), note that the sum of the entries of a row of the incidence matrix is equal to the degree of the corresponding node in the graph. Suppose there is a cycle $\ell_1, \ldots, \ell_n \in \mathscr{L}$. This implies that each row of the matrix $\mathbf{E} = [\mathbf{C}_{\ell_1}, \ldots, \mathbf{C}_{\ell_n}]$ has sum 2, so that $\mathbf{E}v^T = 0$ for any vector $v \in \mathbb{R}^n$ satisfying $\sum_i v_i = 0$. Then, **B** has linearly dependent columns, contradicting the assumption that **B** has full rank.

For (ii), let $i_0 \in \mathcal{I}$ and suppose $i_0 \notin \mathcal{G}(\mathcal{L}, \mathscr{J})$, then row i_0 of $A'_{\mathscr{L}}$ contains only zeros. Therefore, row i_0 row of **B** contains only zeros, which contradicts rank(**B**) = I + J. Similarly, let $j_0 \in \mathcal{J} \setminus \mathcal{G}(\mathcal{L}, \mathscr{J})$, then row j_0 of $A''_{\mathscr{L}}$ and row j_0 of $I_{\mathscr{J}}$ contain only zeros. Therefore, row j_0 of **B** contains only zeros, which contradicts rank(**B**) = I + J.

Lastly, for (iii), note that by (i) we have that $\mathcal{G}(\mathscr{L}, \mathscr{J})$ is a union of $K \geq 1$ trees $\mathcal{T}_1, \ldots, \mathcal{T}_K$ and by (ii), $\mathcal{G}(\mathscr{L}, \mathscr{J})$ has I + J nodes. By [54, Theorem 6], the rank of the incidence matrix of a graph with n nodes that consists of K connected components is n - K, hence rank $(\mathbf{C}_{\mathscr{L}}) = I + J - K$. Since **B** has full rank by assumption, this implies rank $(\mathbf{D}_{\mathscr{J}}) = K$. On the other hand, rank $(\mathbf{C}_{\mathscr{J}}) = |\mathcal{J}|$ by construction, thus $|\mathscr{J}| = K$, i.e., the number of servers \mathscr{J} equals the number of trees. It remains to be shown that each tree contains exactly one server $j \in \mathscr{J}$. Suppose there is a tree that contains two servers $j_1, j_2 \in \mathscr{J}$, then there is a path $\ell_1, \ldots, \ell_n \in \mathscr{L}$ between j_1 and j_2 . It can be verified that the row sum of the matrix $\mathbf{E} = [\mathbf{C}_{\ell_1}, \ldots, \mathbf{C}_{\ell_n}, \mathbf{D}_{j_1}, \mathbf{D}_{j_2}]$ equals 2 for each row, hence $\mathbf{E}v' = 0$ for any vector $v \in \mathbb{R}^{n+2}$ satisfying $\sum_i v_i = 0$. Then, **B** has linearly dependent columns, contradicting the assumption that **B** has full rank.

" \Leftarrow " Let $\mathscr{L} \subseteq \mathscr{L}$ and $\mathscr{J} \subseteq \mathscr{J}$ be such that $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is a spanning forest of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$. Suppose $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is the union of trees $\mathcal{T}_1, \ldots, \mathcal{T}_K$. Let $B = \mathscr{L} \cup \mathscr{J}$ and **B** as in (114). We will prove that **B** has full rank. Since $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is a graph with K connected components and I + J nodes, we have by [54, Theorem 6] that $\operatorname{rank}(\mathbf{C}_{\mathscr{L}}) = I + J - K$. Moreover, $\operatorname{rank}(\mathbf{D}_{\mathscr{J}}) = |\mathscr{J}|$ by construction. Since $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is a spanning forest, each tree contains exactly one server $j \in \mathscr{J}$, hence $\operatorname{rank}(\mathbf{D}_{\mathscr{J}}) = |\mathscr{J}| = K$.

Next, we show that $\mathbf{C}_{\mathscr{L}}$ and $\mathbf{D}_{\mathscr{J}}$ are linearly independent. Consider a linear combination $\sum_{\ell \in \mathscr{L}'} \alpha_{\ell} \mathbf{C}_{\ell}$ of the columns of $\mathbf{C}_{\mathscr{L}}$ with $\alpha_{\ell} \neq 0$ and $\mathscr{L}' \subseteq \mathscr{L}$. Since $\mathscr{G}(\mathscr{L}, \mathscr{J})$ is a forest, the subgraph induced by \mathscr{L}' contains no cycles and therefore must contains at least two nodes with degree 1. This means that the vector $\sum_{\ell \in \mathscr{L}'} \alpha_{\ell} \mathbf{C}_{\ell}$ must have at least two nonzero entries corresponding to the nodes with degree 1. However, any column $b \in \mathbf{D}_{\mathscr{J}}$ has only one nonzero entry and therefore $b \notin \operatorname{span}(\mathbf{C}_{\mathscr{L}})$. Since any column of $\mathbf{D}_{\mathscr{J}}$ is linearly independent of $\mathbf{C}_{\mathscr{L}}$ and the columns of $\mathbf{D}_{\mathscr{J}}$ are linearly independent.

We have shown that $\mathbf{C}_{\mathscr{L}}$ and $\mathbf{D}_{\mathscr{J}}$ are linearly independent and that $\operatorname{rank}(\mathbf{C}_{\mathscr{L}}) + \operatorname{rank}(\mathbf{D}_{\mathscr{J}}) = I + J$. Together, this implies $\operatorname{rank}(\mathbf{B}) = I + J$ and completes the proof. \Box

C.2 Proof of Lemma 2

Let $B = \mathscr{L} \cup \mathscr{J}$ be a basis of $LP(\theta, \varepsilon)$. From Lemma 1, we obtain that $\mathcal{G}(\mathscr{L}, \mathscr{J})$ is a spanning forest of $(\mathcal{I} \cup \mathcal{J}, \mathcal{L})$. Let $x \in \mathbb{R}^L$ be defined as in (11) and define $\sigma \in \mathbb{R}^J$ by $\sigma_j := \mu_j - \varepsilon - \sum_{i \in \mathcal{C}_j} x_{ij}$. Recall that $LP(\theta, \varepsilon)$ is of the form (111). Hence, we must show that $[x, \sigma]^T \in \mathbb{R}^{L+J}$ satisfies

$$\mathbf{B}\begin{bmatrix} x\\ \sigma \end{bmatrix} = \begin{bmatrix} \lambda\\ \mu - \varepsilon \end{bmatrix},\tag{115}$$

where **B** is defined as in (114).

Let $i \in \mathcal{I}$. Recall that *i* is contained in the spanning forest and that *i* cannot be the root of a tree since $\mathscr{J} \subseteq \mathscr{J}$ are the root nodes.

- If *i* is a leaf node, then *i* has a unique parent node $j_0 \in \mathcal{J}$ and so $\mathbf{B}_{ij_0} = 1$ and $\mathbf{B}_{ij} = 0$ for all $j \neq j_0$. Hence, $(\mathbf{B}[x,\sigma])_i = x_{ij_0}$. In this case, $\cap(i) = \{i\}$, so we obtain from (11) that $x_{ij_0} = \lambda_i$.
- If *i* is not a leaf node, then *i* has a unique parent node $j_0 \in \mathcal{J}$ and child nodes $j_1, \ldots, j_m \in \mathcal{J}$ for some $m \ge 1$. Hence, $(\mathbf{B}[x,\sigma])_i = x_{ij_0} + \sum_{k=1}^m x_{ij_k}$. In this case, $\mathsf{C}(\cap(i)) = \{i\} \cup \bigcup_{k=1}^m \mathsf{C}(\cap(j_k))$ and $\mathsf{S}(\cap(i)) = \bigcup_{k=1}^m \mathsf{S}(\cap(j_k))$. Therefore by (11),

$$(\mathbf{B}[x,\sigma])_{i} = x_{ij_{0}} + \sum_{k=1}^{m} x_{ij_{k}} = \sum_{k \in \mathsf{C}(\cap(i))} \lambda_{k} - \sum_{\ell \in \mathsf{S}(\cap(i))} (\mu_{\ell} - \varepsilon) + \sum_{k=1}^{m} \left(\sum_{\ell \in \mathsf{S}(\cap(j_{k}))} (\mu_{\ell} - \varepsilon) - \sum_{k \in \mathsf{C}(\cap(j_{k}))} \lambda_{k}\right) = \lambda_{i}.$$
(116)

Let now $j \in \mathcal{J}$.

- If j is a root node, then j has no parent node and possibly child nodes $i_1, \ldots, i_m \in \mathcal{C}$ for some $m \ge 1$. Since $x_{\ell j} = 0$ for all $(\ell j) \notin \mathscr{L}$, we have $(\mathbf{B}[x, \sigma])_j = \sum_{k=1}^m x_{i_k j} + \sigma_j = \sum_{i \in \mathcal{C}_j} x_{ij} + \sigma_j = \mu_j \varepsilon$.
- If j is a leaf node, then j has a unique parent node $i_0 \in C$ and so $(\mathbf{B}[x,\sigma])_j = x_{i_0j} + \sigma_j$ by definition of σ_j . In this case, $\cap(j) = \{j\}$, so we obtain from (11) that $x_{i_0j} = \mu_j \varepsilon$. Since $x_{\ell j} = 0$ for all $(\ell j) \notin \mathscr{L}$, it follows that $\sigma_j = \mu_j \varepsilon x_{i_0j} = 0$. Hence $(\mathbf{B}[x,\sigma])_j = \mu_j \varepsilon$.
- If j is not a leaf or a root node, then j has a unique parent node $i_0 \in \mathcal{C}$ and child nodes $i_1, \ldots, i_m \in \mathcal{C}$ for some $m \geq 1$ and so $(\mathbf{B}[x,\sigma])_j = x_{i_0j} + \sum_{k=1}^m x_{i_kj} + \sigma_j$. Since $x_{\ell j} = 0$ for all $(\ell j) \notin \mathscr{L}$, it follows that $\sigma_j = \mu_j \varepsilon x_{i_0j} \sum_{k=1}^m x_{i_km} = 0$. In this case, $\mathsf{S}(\mathsf{m}(j)) = \{j\} \cup \bigcup_{k=1}^m \mathsf{S}(\mathsf{m}(i_k))$ and $\mathsf{C}(\mathsf{m}(j)) = \bigcup_{k=1}^m \mathsf{C}(\mathsf{m}(i_k))$. Therefore by (11),

$$(\mathbf{B}[x,\sigma])_j = x_{i_0j} + \sum_{k=1}^m x_{i_kj}$$
(117)

$$= \sum_{\ell \in \mathsf{S}(\widehat{\square}(j))} (\mu_{\ell} - \varepsilon) - \sum_{k \in \mathsf{C}(\widehat{\square}(j))} \lambda_{k} + \sum_{k=1}^{m} \left(\sum_{k \in \mathsf{C}(\widehat{\square}(i_{k}))} \lambda_{k} - \sum_{\ell \in \mathsf{S}(\widehat{\square}(i_{k}))} (\mu_{\ell} - \varepsilon) \right)$$
(118)

$$=\mu_i - \varepsilon. \tag{119}$$

Hence (115) is satisfied and the proof is concluded. \Box

C.3 Proof of Lemma 3

Let $B = \mathscr{L} \cup \mathscr{J}$ be a nondegenerate feasible basis of $LP(\theta, 0)$ and let $\varepsilon \ge 0$ satisfy (12). From Lemma 2, we obtain that $[x^0, \sigma^0]^T \in \mathbb{R}^{L+J}$ defined by

$$x_{ij}^{0} = \begin{cases} \sum_{k \in \mathsf{C}(\cap(i))} \lambda_{k} - \sum_{\ell \in \mathsf{S}(\cap(i))} \mu_{\ell}, & \text{if } i \text{ is a child of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), \\ \sum_{\ell \in \mathsf{S}(\cap(j))} \mu_{\ell} - \sum_{k \in \mathsf{C}(\cap(j))} \lambda_{k}, & \text{if } i \text{ is the parent of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), & \sigma_{j}^{0} = \mu_{j} - \sum_{i \in \mathcal{C}_{j}} x_{ij}^{0}, \end{cases}$$
(120)
0 & \text{if } (ij) \in \mathcal{L} \setminus \mathscr{L},

is the corresponding basic solution of LP($\theta, 0$). Similarly, we obtain that $[x^{\varepsilon}, \sigma^{\varepsilon}]^T \in \mathbb{R}^{L+J}$ defined by

$$x_{ij}^{\varepsilon} = \begin{cases} \sum_{k \in \mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell \in \mathsf{S}(\cap(i))} (\mu_\ell - \varepsilon), & \text{if } i \text{ is a child of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), \\ \sum_{\ell \in \mathsf{S}(\cap(j))} (\mu_\ell - \varepsilon) - \sum_{k \in \mathsf{C}(\cap(j))} \lambda_k, & \text{if } i \text{ is the parent of } j \text{ in } \mathcal{G}(\mathscr{L}, \mathscr{J}), & \sigma_j^{\varepsilon} = \mu_j - \sum_{i \in \mathcal{C}_j} x_{ij}^{\varepsilon}, \quad (121) \\ 0 & \text{if } (ij) \in \mathcal{L} \setminus \mathscr{L}, \end{cases}$$

is the corresponding basic solution of $LP(\theta, \varepsilon)$. It remains to be shown that $[x^{\varepsilon}, \sigma^{\varepsilon}]^T$ is nondegenerate and feasible for $LP(\theta, \varepsilon)$. To this end, we show that (a) $x_{ij}^{\varepsilon} > 0$ for $(ij) \in \mathscr{L}$ and $x_{ij}^{\varepsilon} = 0$ otherwise, and (b) $\sigma_j^{\varepsilon} > 0$ for $j \in \mathscr{J}$ and $\sigma_j^{\varepsilon} = 0$ otherwise.

Proof of (a). Let $(ij) \in \mathscr{L}$. If i is a child of j in $\mathcal{G}(\mathscr{L}, \mathscr{J})$, then by (121),

$$x_{ij}^{\varepsilon} = \sum_{k \in \mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell \in \mathsf{S}(\cap(i))} (\mu_\ell - \varepsilon) \stackrel{\varepsilon > 0}{>} \sum_{k \in \mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell \in \mathsf{S}(\cap(i))} \mu_\ell \stackrel{(120)}{=} x_{ij}^0 > 0, \tag{122}$$

where the last step follows since x^0 is a nondegenerate basic feasible solution of $LP(\theta, 0)$. On the other hand, if *i* is the parent of *j* in $\mathcal{G}(\mathscr{L}, \mathscr{J})$, then by (121),

$$x_{ij}^{\varepsilon} = \sum_{\ell \in \mathsf{S}(\widehat{\square}(j))} (\mu_{\ell} - \varepsilon) - \sum_{k \in \mathsf{C}(\widehat{\square}(j))} \lambda_{k} = \sum_{\ell \in \mathsf{S}(\widehat{\square}(j))} \mu_{\ell} - \sum_{k \in \mathsf{C}(\widehat{\square}(j))} \lambda_{k} - |\mathsf{S}(\square(j))|\varepsilon \stackrel{(12)}{>} 0.$$
(123)

Lastly, let $(ij) \in \mathcal{L} \setminus \mathscr{L}$, then $x_{ij}^{\varepsilon} = 0$ follows directly from (121). Hence, (a) is satisfied.

Proof of (b). Let $j \in \mathcal{J}$. By construction, j is the root of a tree in the spanning forest $\mathcal{G}(\mathcal{L}, \mathcal{J})$ so by (121),

$$\sum_{i \in \mathcal{C}_j} x_{ij}^{\varepsilon} = \sum_{i:(ij) \in \mathscr{L}} \left(\sum_{k \in \mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell \in \mathsf{S}(\cap(i))} (\mu_\ell - \varepsilon) \right)$$
(124)

$$= (|\mathsf{S}(\cap(j))| - 1)\varepsilon + \sum_{i:(ij)\in\mathscr{L}} \left(\sum_{k\in\mathsf{C}(\cap(i))} \lambda_k - \sum_{\ell\in\mathsf{S}(\cap(i))} \mu_\ell\right).$$
(125)

By (12),

$$|\mathsf{S}(\cap(j))|\varepsilon < \sum_{\ell \in \mathsf{S}(\cap(j))} \mu_{\ell} - \sum_{k \in \mathsf{S}(\cap(j))} \lambda_{k} = \mu_{j} + \sum_{i:(ij) \in \mathscr{L}} \left(\sum_{\ell \in \mathsf{S}(\cap(i))} \mu_{\ell} - \sum_{k \in \mathsf{S}(\cap(i))} \lambda_{k}\right).$$
(126)

Here, we used that j is a root and therefore $S(\cap(j))$ is the union of $\{j\}$ and $\bigcup_{i:(ij)\in\mathscr{L}} S(\cap(i))$. Bounding (125) using (126) gives

$$\sigma_j^{\varepsilon} = \mu_j - \varepsilon - \sum_{i \in \mathcal{C}_j} x_{ij}^{\varepsilon} > 0.$$
(127)

Lastly, let $j \in \mathcal{J} \setminus \mathcal{J}$, then j is not a root node in a tree in $\mathcal{G}(\mathcal{L}, \mathcal{J})$. It can be verified that $\sigma_j^{\varepsilon} = 0$ using a similar approach as in the proof of Lemma 2 (see the last two steps in Appendix C.2). Hence, (b) is satisfied and the proof is concluded.

C.4 Proof of Lemma 4

Since π satisfies (7),

$$\lim_{t \to \infty} \sum_{j \in \mathcal{S}_i} \frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t} = \lambda_i, \quad \forall i \in \mathcal{I}.$$
(128)

Moreover, by (8),

$$\lim_{t \to \infty} \sum_{i \in \mathcal{C}_j} \frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t} \le \mu_j - \varepsilon, \quad \forall j \in \mathcal{J}.$$
(129)

Now, since the summation over all lines $(ij) \in \mathcal{L}$ in (13) is finite, we can interchange the summation and limit to obtain

$$\lim_{t \to \infty} \sum_{(ij) \in \mathcal{L}} \frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t} \theta_{ij} = \sum_{(ij) \in \mathcal{L}} \lim_{t \to \infty} \frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t} \theta_{ij}.$$
 (130)

In light of (128) and (129), the set of values $\{\lim_{t\to\infty} \mathbb{E}^{\theta}_{\pi}(D_{ij}(t))/t\}_{(ij)\in\mathcal{L}}$ satisfies the constraints of $LP(\theta,\varepsilon)$ in (10). Since x^{θ} is the optimal solution of $LP(\theta,\varepsilon)$, we have

$$\sum_{(ij)\in\mathcal{L}}\lim_{t\to\infty}\frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t}\theta_{ij} \le \sum_{(ij)\in\mathcal{L}}x_{ij}\theta_{ij}.$$
(131)

The result follows from (13),

$$\lim_{t \to \infty} \frac{R_{\pi}^{\theta}(t)}{t} = \lim_{t \to \infty} \sum_{(ij) \in \mathcal{L}} x_{ij} \theta_{ij} - \frac{\mathbb{E}_{\pi}^{\theta}(D_{ij}(t))}{t} \ge 0.$$
(132)

C.5 Proof of Lemma 5

Let π satisfy (7) and (8). By (13) and (19),

$$R_{\pi}^{\theta}(t) = \sum_{(ij)\in\mathcal{L}} (v_i^{\theta} + w_j^{\theta} - \phi_{ij}^{\theta}) (tx_{ij}^{\theta} - \mathbb{E}_{\pi}^{\theta}(D_{ij}(t))).$$
(133)

By (17), we have $\phi_{ij}^{\theta} x_{ij}^{\theta} = 0$ for all $(ij) \in \mathcal{L}$. Moreover, $\phi_{ij}^{\theta} = 0$ for $(ij) \in O(\theta)$ and $O(\theta) \cup O^{c}(\theta)$ is a partition of \mathcal{L} . Therefore,

$$R^{\theta}_{\pi}(t) = \sum_{(ij)\in\mathcal{L}} (v^{\theta}_i + w^{\theta}_j) t x^{\theta}_{ij} - \sum_{(k\ell)\in\mathcal{L}} (v^{\theta}_k + w^{\theta}_\ell) \mathbb{E}^{\theta}_{\pi}(D_{k\ell}(t)) + \sum_{(mn)\in O^c(\theta)} \phi^{\theta}_{mn} \mathbb{E}^{\theta}_{\pi}(D_{mn}(t)).$$
(134)

Reordering terms gives

$$R^{\theta}_{\pi}(t) = \sum_{(ij)\in\mathcal{L}} w^{\theta}_{j}(tx^{\theta}_{ij} - \mathbb{E}^{\theta}_{\pi}(D_{ij}(t))) + \sum_{(k\ell)\in\mathcal{L}} v^{\theta}_{k}(tx^{\theta}_{k\ell} - \mathbb{E}^{\theta}_{\pi}(D_{k\ell}(t))) + \sum_{(mn)\in O^{c}(\theta)} \phi^{\theta}_{mn} \mathbb{E}^{\theta}_{\pi}(D_{mn}(t)).$$
(135)

As an intermediate step, note that by definition of the sets S_i in (2) and C_j in (3), the following identities hold for any vector $(\gamma_{ij})_{(ij)\in\mathcal{L}}$,

$$\sum_{(ij)\in\mathcal{L}}\gamma_{ij} = \sum_{i\in\mathcal{I}}\sum_{j\in\mathcal{S}_i}\gamma_{ij} = \sum_{j\in\mathcal{J}}\sum_{i\in\mathcal{C}_j}\gamma_{ij}.$$
(136)

Applying (136) to (135) gives

$$R^{\theta}_{\pi}(t) = \sum_{j \in \mathcal{J}} w^{\theta}_{j} \Big(\sum_{i \in \mathcal{C}_{j}} (tx^{\theta}_{ij} - \mathbb{E}^{\theta}_{\pi}(D_{ij}(t))) \Big) + \sum_{k \in \mathcal{I}} v^{\theta}_{k} \Big(\sum_{\ell \in \mathcal{S}_{k}} (tx^{\theta}_{k\ell} - \mathbb{E}^{\theta}_{\pi}(D_{k\ell}(t))) \Big) + \sum_{(mn) \in O^{c}(\theta)} \phi^{\theta}_{mn} \mathbb{E}^{\theta}_{\pi}(D_{mn}(t)).$$
(137)

Since x^{θ} is the optimal solution of $LP(\theta, \varepsilon)$, it satisfies (10b), i.e., for all $k \in \mathcal{I}$, $\sum_{\ell \in S_k} x_{k\ell}^{\theta} = \lambda_k$. Moreover, since the complementary slackness condition (18) are strict, we have for any $j \in \mathcal{J}$ that either $w_j^{\theta} = 0$, or $\sum_{i \in C_j} x_{ij}^{\theta} = \mu_j - \varepsilon$. Hence

$$R_{\pi}^{\theta}(t) = \sum_{j \in \mathcal{J}} w_{j}^{\theta} \left(t(\mu_{j} - \varepsilon) - \sum_{i \in \mathcal{C}_{j}} \mathbb{E}_{\pi}^{\theta}(D_{ij}(t)) \right) + \sum_{k \in \mathcal{I}} v_{k}^{\theta} \left(t\lambda_{k} - \sum_{\ell \in \mathcal{S}_{k}} \mathbb{E}_{\pi}^{\theta}(D_{k\ell}(t)) \right) + \sum_{(mn) \in O^{c}(\theta)} \phi_{mn}^{\theta} \mathbb{E}_{\pi}^{\theta}(D_{mn}(t)).$$
(138)

This concludes the proof of (20). \Box

C.6 Proof of Lemma 6

Let $x^{\theta} \in \mathbb{R}_{\geq 0}^{L}$ be the optimal solution of $LP(\theta, \varepsilon)$ in (10), $v^{\theta} \in \mathbb{R}^{I}, w^{\theta} \in \mathbb{R}_{\geq 0}^{J}$ be the optimal solution of $D(\theta, \varepsilon)$ in (16), and $(ij) \in O^{c}(\theta)$. Let $a_{0} \in \mathbb{R}_{\geq 0}$ be such that $a_{0} > v_{i}^{\theta} + w_{j}^{\theta} \ge \theta_{ij}$. We consider a new program $LP(A(\theta, ij, a_{0}), \varepsilon)$ with $A(\theta, ij, a_{0})$ as in (21) and denote its optimal solution by $x^{A} \in \mathbb{R}_{\geq 0}^{L}$. Suppose that $(ij) \in O^{c}(A(\theta, ij, a_{0}))$, then $x_{ij}^{A} = x_{ij}^{\theta} = 0$. In this case, we must have $x^{A} = x^{\theta}$, since the programs only differ in the reward value for line (ij). By the nondegeneracy assumption, a primal solution implies a unique dual solution, hence v^{θ}, w^{θ} must be optimal for $D(A(\theta, ij, a_{0}))$. However, v^{θ}, w^{θ} violates the feasibility constraint (16b) since $a_{0} > v_{i}^{\theta} + w_{j}^{\theta}$, leading to a contradiction. This implies $(ij) \in O(A(\theta, ij, a_{0}))$ and concludes the proof that $\Delta(\theta, ij)$ is nonempty. \Box

C.7 Proof of Lemma 7

In order to prove Lemma 7, we present an intermediate result in Lemma C.1. Let

$$R_j^{mA_m} := \min\{t \ge 0 : \ \hat{Q}_j^{mA_m}(t) = 0\}$$
(139)

denote the hitting time of state 0 by the process $\hat{Q}_j^{mA_m}(t)$, which is well-defined by positive recurrence of the process $\hat{Q}_i^a(t)$. Lemma C.1 provides a tail bound on $R_j^{mA_m}$.

Lemma C.1. For any $n \in \mathbb{N}$ and $b \in \mathbb{R}_{>0}$,

$$\mathbb{P}(R_j^{mA_m} > nb \mid \hat{Q}_j^{mA_m}(0) = n) \le (\rho_j^{A_m})^{-n/2} \exp\left(-nb(\tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{b^{-2} + 4\tilde{\lambda}_j^{A_m}\mu_j})\right).$$
(140)

Proof of Lemma C.1. As discussed in Section 4.3.1, $\hat{Q}_{j}^{mA_{m}}(t)$ behaves as an M/M/1 queueing system with arrival rate $\tilde{\lambda}_{j}^{A_{m}}$ independently from other servers. Therefore, from [49, Section II.2.2] we obtain for the moment generating function of $R_{j}^{mA_{m}}$,

$$\mathbb{E}\left(\exp(sR_{j}^{mA_{m}}) \mid \hat{Q}_{j}^{mA_{m}}(0) = n\right) = \left(\frac{\tilde{\lambda}_{j}^{A_{m}} + \mu_{j} - s - \sqrt{(\tilde{\lambda}_{j}^{A_{m}} + \mu_{j} - s)^{2} - 4\tilde{\lambda}_{j}^{A_{m}}\mu_{j}}}{2\tilde{\lambda}_{j}^{A_{m}}}\right)^{n}.$$
(141)

The radius of convergence is determined by

$$(\tilde{\lambda}_j^{A_m} + \mu_j - s)^2 - 4\tilde{\lambda}_j^{A_m}\mu_j > 0 \iff s < \tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{4\tilde{\lambda}_j^{A_m}\mu_j} =: \delta.$$
(142)

Let $b \in \mathbb{R}_{>0}$. By Markov's inequality, we have

$$\mathbb{P}(R_j^{mA_m} > nb \mid \hat{Q}_j^{mA_m}(0) = n) \le \inf_{0 < s < \delta} \mathbb{E}(\exp(sR_j^a) \mid \hat{Q}_j^{mA_m}(0) = n) \exp(-nbs).$$
(143)

It can be verified that the infimum within the radius of convergence is attained at

$$s^* = \tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{b^{-2} + 4\tilde{\lambda}_j^{A_m}\mu_j} < \delta.$$
(144)

Substitution of s^* into (143) gives

$$\mathbb{P}(R_{j}^{mA_{m}} > nb \mid \hat{Q}_{j}^{mA_{m}}(0) = n) \leq \exp\left(-nb(\tilde{\lambda}_{j}^{A_{m}} + \mu_{j} - \sqrt{b^{-2} + 4\tilde{\lambda}_{j}^{A_{m}} \mu_{j}})\right) \left(\frac{-b^{-1} + \sqrt{b^{-2} + 4\tilde{\lambda}_{j}^{A_{m}} \mu_{j}}}{2\tilde{\lambda}_{j}^{A_{m}}}\right)^{n}.$$
 (145)

We note that the term

$$\frac{-b^{-1} + \sqrt{b^{-2} + 4\tilde{\lambda}_j^{A_m}\mu_j}}{2\tilde{\lambda}_j^{A_m}}$$
(146)

is increasing in b, and that

$$\lim_{b \to \infty} \frac{-b^{-1} + \sqrt{b^{-2} + 4\tilde{\lambda}_j^{A_m} \mu_j}}{2\tilde{\lambda}_j^{A_m}} = \frac{\sqrt{4\tilde{\lambda}_j^{A_m} \mu_j}}{2\tilde{\lambda}_j^{A_m}} = (\rho_j^{A_m})^{-1/2}.$$
 (147)

Therefore, we can bound (145) further and complete the proof, i.e.,

$$\mathbb{P}(R_j^{mA_m} > nb \mid \hat{Q}_j^{mA_m}(0) = n) \le (\rho_j^{A_m})^{-n/2} \exp\left(-nb(\tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{b^{-2} + 4\tilde{\lambda}_j^{A_m}}\mu_j)\right).$$
(148)

We next present the proof of Lemma 7.

Proof of Lemma 7. We define

$$s^{mA_m} := \frac{-\ln^{\beta}(2Jm)}{\ln(\rho_i^{A_m})} > 0.$$
(149)

Let $t \geq \tau_m$ and $j \in \mathcal{J}$. Note that by construction of the coupling

$$\mathbb{P}(\underline{Q}_{j}^{mA_{m}}(t) = \hat{Q}_{j}^{mA_{m}}(t)) \ge \mathbb{P}(\underline{Q}_{j}^{mA_{m}}(\tau_{m}) = \hat{Q}_{j}^{mA_{m}}(\tau_{m})).$$
(150)

By the law of total probability, we have

$$\mathbb{P}(\underline{Q}_{j}^{mA_{m}}(\tau_{m}) = \hat{Q}_{j}^{mA_{m}}(\tau_{m})) \ge \mathbb{P}(\underline{Q}_{j}^{mA_{m}}(\tau_{m}) = \hat{Q}_{j}^{mA_{m}}(\tau_{m}) \mid \hat{Q}_{j}^{mA_{m}}(0) \le \lfloor s^{mA_{m}} \rfloor) \mathbb{P}(\hat{Q}_{j}^{mA_{m}}(0) \le \lfloor s^{mA_{m}} \rfloor).$$
(151)

Since the initial queue length $\hat{Q}_{j}^{mA_{m}}(0)$ is sampled from the stationary measure $p_{j}^{A_{m}}$ in (38), we have

$$\mathbb{P}(\hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor) = \sum_{i=0}^{\lfloor s^{mA_{m}} \rfloor} (1 - \rho_{j}^{A_{m}})(\rho_{j}^{A_{m}})^{i} = 1 - (\rho_{j}^{A_{m}})^{1 + \lfloor s^{mA_{m}} \rfloor}.$$
(152)

We use that $b^{\ln(a)/\ln(b)} = a$ for any a, b > 0 and (149) to obtain

$$(\rho_j^{A_m})^{\lfloor s^{mA_m} \rfloor} \ge (\rho_j^{A_m})^{s^{mA_m}} = (\rho_j^{A_m})^{-\ln^{\beta}(2Jm)/\ln(\rho_j^{A_m})} = (2Jm)^{-\ln^{\beta-1}(2Jm)},$$
(153)

and so

$$\mathbb{P}(\hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor) \leq 1 - \rho_{j}^{A_{m}}(2Jm)^{-\ln^{\beta-1}(2Jm)} \stackrel{(i)}{\geq} 1 - (2Jm)^{-\ln^{\beta-1}(2Jm)} \stackrel{(ii)}{\geq} 1 - (2Jm)^{-\beta} \stackrel{(iii)}{\geq} 1 - \frac{1}{2Jm^{\beta}}, \quad (154)$$

where (i) follows from $\rho_j^{A_m} < 1$ and 2Jm > 1, (ii) by the assumption $m \ge C_\beta$ and $2Jm \ge 1$, and (iii) by $J \ge 1$, $m \ge 1$, and $\beta > 1$.

We continue to bound the conditional probability in (151). Observe that if the process $\hat{Q}_j^{mA_m}(t)$ hits state 0 at time x, then $\hat{Q}_j^{mA_m}(t) = \underline{Q}_j^{mA_m}(t)$ for all $t \ge x$, as illustrated in Figure 6. Therefore,

$$\mathbb{P}\left(\underline{Q}_{j}^{mA_{m}}(\tau_{m}) = \hat{Q}_{j}^{mA_{m}}(\tau_{m}) \mid \hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor\right) = \mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor\right).$$
(155)

As an intermediate step, we note that for any A and any pairwise disjoint B_1, \ldots, B_n ,

$$\mathbb{P}\left(A \mid \bigcup_{k=1}^{n} B_{k}\right) = \frac{\sum_{k=1}^{n} \mathbb{P}(A \cap B_{k})}{\sum_{j=1}^{n} \mathbb{P}(B_{j})} = \sum_{k=1}^{n} \frac{\mathbb{P}(B_{k})}{\sum_{j=1}^{n} \mathbb{P}(B_{j})} \mathbb{P}(A \mid B_{k}) \ge \min_{k \in \{1, \dots, n\}} \mathbb{P}(A \mid B_{k}),$$
(156)

where the inequality follows since the weighted average of a set of nonnegative numbers is at least as large as the minimum of that set. Therefore, since $\{\hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor\} = \bigcup_{k=0}^{\lfloor s^{mA_{m}} \rfloor} \{\hat{Q}_{j}^{mA_{m}}(0) = k\}$, we have

$$\mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor\right) \geq \min_{k \in \{0,\dots,\lfloor s^{mA_{m}} \rfloor\}} \mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = k\right).$$
(157)

Moreover, since $\hat{Q}_{i}^{mA_{m}}(t)$ is a birth-death process, we have

$$\mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = k_{1}\right) \geq \mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = k_{2}\right), \ \forall k_{1} \leq k_{2} \in \mathbb{N},$$
(158)

which implies that the minimum in (157) is attained at $\lfloor s^{mA_m} \rfloor$, therefore

$$\mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) \leq \lfloor s^{mA_{m}} \rfloor\right) \geq \mathbb{P}\left(R_{j}^{mA_{m}} \leq \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = \lfloor s^{mA_{m}} \rfloor\right).$$
(159)

We obtain from Lemma C.1 that

$$\mathbb{P}\left(R_{j}^{mA_{m}} > \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = \lfloor s^{mA_{m}} \rfloor\right) \leq (\rho_{j}^{A_{m}})^{-\lfloor s^{mA_{m}} \rfloor/2} \exp\left(-\tau_{m}\left(\tilde{\lambda}_{j}^{A_{m}} + \mu_{j} - \sqrt{\left(\frac{\tau_{m}}{\lfloor s^{mA_{m}} \rfloor}\right)^{-2} + 4\tilde{\lambda}_{j}^{A_{m}} \mu_{j}}\right)\right)$$
(160)

$$\leq (\rho_j^{A_m})^{-s^{mA_m/2}} \exp\left(-\tau_m \left(\tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{\left(\frac{\tau_m}{s^{mA_m}}\right)^{-2} + 4\tilde{\lambda}_j^{A_m} \mu_j}\right)\right).$$
(161)

Now, by substitution of τ_m and s^{mA_m} as defined in (36) and (149), respectively, and noting that $\tau_m/s^{mA_m} = -\alpha \ln(\rho_j^{A_m})$, we have

$$\mathbb{P}\left(R_{j}^{mA_{m}} > \tau_{m} \mid \hat{Q}_{j}^{mA_{m}}(0) = \lfloor s^{mA_{m}} \rfloor\right) \leq (\rho_{j}^{A_{m}})^{\ln^{\beta}(2Jm)/(2\ln(\rho_{j}^{A_{m}}))} \\ \cdot \exp\left(-\alpha \ln^{\beta}(2Jm)\left(\tilde{\lambda}_{j}^{A_{m}} + \mu_{j} - \sqrt{\left(\alpha \ln(\rho_{j}^{A_{m}})\right)^{-2} + 4\tilde{\lambda}_{j}^{A_{m}}\mu_{j}}\right)\right)$$
(162)

$$= (2Jm)^{-\ln^{\beta-1}(2Jm)\left(-1/2 + \alpha(\tilde{\lambda}_{j}^{Am} + \mu_{j} - \sqrt{(\alpha\ln(\rho_{j}^{Am}))^{-2} + 4\tilde{\lambda}_{j}^{Am} \mu_{j})}\right)}.$$
 (163)

It can be verified that $f(\gamma) := -\frac{1}{2} + \gamma \left(\tilde{\lambda}_j^{A_m} + \mu_j - \sqrt{(\gamma \ln(\rho_j^{A_m}))^{-2} + 4\tilde{\lambda}_j^{A_m} \mu_j} \right)$ is strictly increasing on $[0, \infty)$ and that

$$\gamma^* = \frac{3(\tilde{\lambda}_j^{A_m} + \mu_j)\ln(\rho_j^{A_m}) - 2\sqrt{(\mu_j - \tilde{\lambda}_j^{A_m})^2 + 9\tilde{\lambda}_j^{A_m}\mu_j\ln^2(\rho_j^{A_m})}}{2(\mu_j - \tilde{\lambda}_j^{A_m})^2\ln(\rho_j^{A_m})} > 0$$
(164)

satisfies $f(\gamma^*) = 1$. By construction of α in (35), we have $\alpha \geq \gamma^*$, and hence

$$\mathbb{P}\left(R_j^{mA_m} > \tau_m \mid \hat{Q}_j^{mA_m}(0) = \lfloor s^{mA_m} \rfloor\right) \le (2Jm)^{-\ln^{\beta-1}(2Jm)}.$$
(165)

Since $m \ge C_{\beta}$ and hence $\log^{\beta-1}(2Jm) \ge \beta$, we obtain

$$\mathbb{P}\left(R_j^{mA_m} > \tau_m \mid \hat{Q}_j^{mA_m}(0) = \lfloor s^{mA_m} \rfloor\right) \le (2Jm)^{-\beta}.$$
(166)

Substitution of (154) and (155) into (151) and subsequently using (150) and (166) and noting that $\beta > 1$ gives

$$\mathbb{P}(\underline{Q}_{j}^{mA_{m}}(t) = \hat{Q}_{j}^{mA_{m}}(t)) \ge \left(1 - \frac{1}{2Jm^{\beta}}\right)^{2} \ge 1 - \frac{1}{Jm^{\beta}}.$$
(167)

To conclude the proof, we note that by independence of the customer arrivals,

$$1 - \mathbb{P}(\underline{Q}_{j}^{mA_{m}}(t) = \hat{Q}_{j}^{mA_{m}}(t), \ \forall j \in \mathcal{J}) = \mathbb{P}(\exists j \in \mathcal{J}: \ \underline{Q}_{j}^{mA_{m}}(t) \neq \hat{Q}_{j}^{mA_{m}}(t)) \stackrel{(167)}{\leq} \sum_{j=1}^{J} \frac{1}{Jm^{\beta}} = \frac{1}{m^{\beta}}.$$
 (168)

C.8 Proof of Lemma 8

In order to prove Lemma 8, we provide two intermediate results.

Lemma C.2. Let $\gamma \in \mathbb{R}_{\geq 1}$, $u \in \mathbb{N}_{\geq 1}$, and X_1, \ldots, X_u be independent random variables with $X_m \sim Ber(1 - 1/m^{\gamma})$ for all $m = 1, \ldots, u$. Then for any $\eta \in \mathbb{R}_{\geq 1}$ such that $\lceil \sum_{m=1}^u 1/m^{\gamma} \rceil \leq \eta \leq u$, we have

$$\mathbb{P}\left(\sum_{m=1}^{u} X_m < u - \eta\right) \le \frac{u^{1+\eta}}{\eta^{(\gamma+1)\eta}} \exp((\gamma+1)\eta).$$
(169)

Proof of Lemma C.2. Let $Y_m := 1 - X_m$ for $m = 1, \ldots, u, \mu := \sum_{m=1}^u 1/m^{\gamma}$, and $\mu \le \eta \le u$. Then, $Y_m \sim \text{Ber}(1/m^{\gamma})$ and

$$\mathbb{P}\Big(\sum_{m=1}^{u} X_m < u - \eta\Big) = \mathbb{P}\Big(\sum_{m=1}^{u} Y_m > \eta\Big).$$
(170)

We note that $\sum_{m=1}^{u} Y_m$ follows a Poisson binomial distribution. Let $a_k := \mathbb{P}(\sum_{m=1}^{u} Y_m = k)$ and $m^* := \arg \max_{k \in [u]} a_k$ refer to the probability mass function and mode of the Poisson binomial distribution, respectively. It is known that the Poisson binomial distribution is strong Rayleigh [55, Corollary 4.2]. As a consequence, $\{a_k\}_{k=1}^{u}$ is ultra-logconcave, and this in turn implies that the sequence is unimodal (see the discussion following [55, Corollary 4.2]) This proves that if $m^* \leq k \leq l$, then $a_l \leq a_k$. What remains is to relate the mode m^* to the mean μ . By [56, Theorem 4], we have $m^* \in \{\lfloor \mu \rfloor, \lceil \mu \rceil\}$. Consequently, if $\lceil \mu \rceil \leq k \leq l$, then $a_l \leq a_k$.

Therefore, since $\lceil \mu \rceil \leq \eta$, we obtain

$$\mathbb{P}\left(\sum_{m=1}^{u} Y_m > \eta\right) \le \sum_{k=\eta}^{u} \mathbb{P}\left(\sum_{m=1}^{u} Y_m = k\right) \le u \mathbb{P}\left(\sum_{m=1}^{u} Y_m = \eta\right).$$
(171)

Since the success probability of Y_m is decreasing in m, the most likely way to get $\sum_{m=1}^{u} Y_m = \eta$ is for the first η indices to be 1. Moreover, the number of combinations of cardinality η from the set $\{1, \ldots, u\}$ is $\binom{u}{\eta}$, and therefore

$$\mathbb{P}\left(\sum_{m=1}^{u} Y_m > \eta\right) \le u\binom{u}{\eta} \prod_{m=1}^{\eta} \mathbb{P}(Y_m = 1)$$
(172)

$$= u \binom{u}{\eta} \prod_{m=1}^{\eta} \frac{1}{m^{\gamma}}$$
(173)

$$= u \binom{u}{\eta} \frac{1}{(\eta!)^{\gamma}}.$$
(174)

By Stirling's formula [57],

$$\eta! \ge \sqrt{2\pi\eta} \ \eta^{\eta} \exp(-\eta) \ge \eta^{\eta} \exp(-\eta). \tag{175}$$

Moreover, since $\exp(\eta) \ge \eta^{\eta}/\eta!$, we have

$$\binom{u}{\eta} \le \frac{u^{\eta}}{\eta!} \le \left(\frac{u}{\eta}\right)^{\eta} \exp(\eta).$$
(176)

Hence

$$\mathbb{P}\left(\sum_{m=1}^{u} Y_m > \eta\right) \le u\left(\frac{u}{\eta}\right)^{\eta} \exp(\eta)\eta^{-\gamma\eta} \exp(\gamma\eta) = \frac{u^{1+\eta}}{\eta^{(\gamma+1)\eta}} \exp((\gamma+1)\eta).$$
(177)

The result follows by substituting (177) into (170).

Lemma C.3. Let
$$\gamma > 0$$
 and $X \sim Poi(\gamma)$. For any $b \in (0, \gamma)$, $\mathbb{P}(X - \gamma \leq -b) \leq \exp\left(-\frac{b^2}{2\gamma}\right)$

Proof of Lemma C.3. Let $\gamma > 0$ and $0 < b < \gamma$. We have for any t < 0,

$$\mathbb{P}(X \le \gamma - b) = \mathbb{P}\left(e^{tX} \ge e^{t(\gamma - b)}\right).$$
(178)

By applying Markov's inequality, we obtain

$$\mathbb{P}(X \le \gamma - b) \le \inf_{t < 0} \mathbb{E}\left(e^{tX}\right) e^{-t(\gamma - b)}.$$
(179)

Since the moment generating function of X is $\mathbb{E}(e^{tX}) = e^{\gamma e^t - 1}$ for $t \in \mathbb{R}$, we have

$$\mathbb{P}(X \le \gamma - b) \le \inf_{t < 0} \mathrm{e}^{\gamma(\mathrm{e}^t - 1)} \mathrm{e}^{-t(\gamma - b)}.$$
(180)

The infimum is attained for $t = \ln((\gamma - b)/\gamma) < 0$, hence we find after substituting that

$$\mathbb{P}(X \le \gamma - b) \le \exp\left(-b - (\gamma - b)\ln\left(\frac{\gamma - b}{\gamma}\right)\right)$$
(181)

$$= \exp\left(-\frac{b^2}{2\gamma}\left(\frac{2\gamma}{b} + \left(\frac{2\gamma(\gamma-b)}{b^2}\right)\ln\left(\frac{\gamma-b}{\gamma}\right)\right)\right).$$
(182)

Let $x = b/\gamma$. From the assumptions, it follows that 0 < x < 1. Now, it can be verified from a Taylor expansion that

$$h(x) = \frac{2}{x} + \frac{2}{x} \left(\frac{1}{x} - 1\right) \ln(1 - x)$$
(183)

satisfies $h(x) \ge 1$ for any 0 < x < 1. Therefore $\mathbb{P}(X \le \gamma - b) \le \exp\left(-\frac{b^2}{2\gamma}\right)$. \Box We next present the proof of Lemma 8.

Proof of Lemma 8. For $(ij) \in \mathcal{L}^a$ and $m \in \mathbb{N}$, consider the processes \hat{Q}_j^{ma} and \underline{Q}_j^{ma} as defined in Section 4.3.1. Let \hat{D}_{ij}^{ma} denote the number of type-(ij) departures within episode m but after warm up time τ_m of the process \hat{Q}_j^{ma} . We introduce the events

$$\Omega_{ij}^{ma} := \{ D_{ij}^{ma} \ge \hat{D}_{ij}^{ma} \}, \quad \underline{\Omega}_{ij}^{ma} := \{ \hat{Q}_{ij}^{mA_m}(\tau_m) = \underline{Q}_{ij}^{mA_m}(\tau_m) \}.$$
(184)

We note that $\underline{\Omega}_{ij}^{ma}$ implies Ω_{ij}^{ma} , hence

$$\mathbb{P}(\underline{\Omega}_{ij}^{ma}) \le \mathbb{P}(\Omega_{ij}^{ma}),\tag{185}$$

and so for any $k \geq 1$,

$$\mathbb{P}\left(\sum_{m=1}^{k} \mathbb{1}\left\{\underline{\Omega}_{ij}^{ma}\right\} \le \sum_{m=1}^{k} \mathbb{1}\left\{\Omega_{ij}^{ma}\right\}\right) = 1.$$
(186)

Suppose $k > \lceil C_{\beta} \rceil$. It follows from (41) and (44) that

$$E_k^a = \bigg\{ \sum_{m=1}^{\lceil C_\beta + u_k \rceil} D_{ij}^{ma} \ge \lceil \sigma_{ijk}^a \rceil, \ \forall (ij) \in \mathcal{L}^a \bigg\}.$$
(187)

By the law of total probability,

$$\mathbb{P}((E_k^a)^c) \le \mathbb{P}\Big(\sum_{m=1}^{\lceil C_\beta + u_k \rceil} \mathbb{1}\left\{\Omega_{ij}^{ma}\right\} < \lceil u_k - \eta_k \rceil\Big) + \mathbb{P}\Big((E_k^a)^c, \sum_{m=1}^{\lceil C_\beta + u_k \rceil} \mathbb{1}\left\{\Omega_{ij}^{ma}\right\} \ge \lceil u_k - \eta_k \rceil\Big).$$
(188)

We bound these terms individually.

It follows from (186) that

$$\mathbb{P}\left(\sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil}\mathbb{1}\left\{\Omega_{ij}^{ma}\right\} < \lceil u_{k}-\eta_{k}\rceil\right) = \mathbb{P}\left(\sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil}\mathbb{1}\left\{\Omega_{ij}^{ma}\right\} < \lceil u_{k}-\eta_{k}\rceil, \sum_{m=1}^{k}\mathbb{1}\left\{\underline{\Omega}_{ij}^{ma}\right\} \le \sum_{m=1}^{k}\mathbb{1}\left\{\Omega_{ij}^{ma}\right\}\right)$$
(189)

$$\leq \mathbb{P}\bigg(\sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil} \mathbb{1}\big\{\underline{\Omega}_{ij}^{ma}\big\} < \lceil u_{k}-\eta_{k}\rceil\bigg).$$
(190)

We observe that event $\underline{Q}_{j}^{ma}(\tau_{m}) = \hat{Q}_{j}^{ma}(\tau_{m})$ implies that there exists an $0 \leq s \leq \tau_{m}$ such that $\hat{Q}_{j}^{ma}(s) = \underline{Q}_{j}^{ma}(s) = 0$. In this case, we have $\hat{Q}_{ij}^{ma}(s) \leq \hat{Q}_{j}^{ma}(s) = 0 = \underline{Q}_{ij}^{ma}(s)$ by construction, and so $\mathbb{1}\left\{\underline{\Omega}_{ij}^{m}\right\} = 1$. Thus, by invoking Lemma 7 we find that

$$\mathbb{P}(\underline{\Omega}_{ij}^{ma}) \ge \mathbb{P}(\underline{\Omega}_{j}^{ma}) \ge 1 - \frac{1}{m^{\beta}}.$$
(191)

Moreover, the events $\underline{\Omega}_{ij}^{ma}$ are mutually independent over all episodes, since the initial values $\hat{Q}_{j}^{ma}(0)$ are sampled from p_{j}^{a} in (38) and $\underline{Q}_{j}^{ma}(0) = 0$. Lastly, it can be verified that $\eta_{k} \geq \lceil \sum_{m=1}^{\lceil u_{k} \rceil} 1/m^{\beta} \rceil$ for k sufficiently large. Hence, it follows from (191) and Lemma C.2 that for k sufficiently large,

$$\mathbb{P}\left(\sum_{m=1}^{\lceil C_{\beta}+u_k\rceil} \mathbb{1}\left\{\underline{\Omega}_{ij}^{ma}\right\} < \lceil u_k - \eta_k\rceil\right) \le \mathbb{P}\left(\sum_{m=\lceil C_{\beta}\rceil}^{\lceil C_{\beta}+u_k\rceil} \mathbb{1}\left\{\underline{\Omega}_{ij}^{ma}\right\} < \lceil u_k - \eta_k\rceil\right) \le \frac{u_k^{1+\eta_k}}{\eta_k^{(\beta+1)\eta_k}} \exp((\beta+1)\eta_k).$$
(192)

By using $a = \exp(\ln(a))$, we have

$$u_{k}^{1+\eta_{k}} = \exp\left((1+\eta_{k})\ln(u_{k})\right) \stackrel{(43)}{=} \exp\left((1+\eta_{k})\beta\ln\ln(k)\right),$$
(193)

$$\eta_k^{(\beta+1)\eta_k} = \exp\left((\beta+1)\eta_k \ln(\eta_k)\right) \stackrel{(43)}{=} \exp\left(\frac{(\beta+1)^2}{2}\eta_k \ln\ln(k)\right),\tag{194}$$

and so

$$\frac{u_k^{1+\eta_k}}{\eta_k^{(\beta+1)\eta_k}} = \exp\left(\left(\beta - \frac{\beta^2 + 1}{2}\eta_k\right)\ln\ln(k)\right).$$
(195)

Substitution of (195) into (192) gives

$$\mathbb{P}\left(\sum_{m=\lceil C_{\beta}\rceil}^{\lceil C_{\beta}+u_{k}\rceil} \mathbb{1}\left\{\underline{\Omega}_{ij}^{ma}\right\} < \lceil u_{k}-\eta_{k}\rceil\right) \leq \exp\left(\beta\ln\ln(k) + \left(\beta+1-\frac{\beta^{2}+1}{2}\ln\ln(k)\right)\eta_{k}\right) \qquad (196)$$

$$\leq \exp\left(\left[2\beta+1-\frac{\beta^{2}+1}{2}\ln\ln(k)\right]\eta_{k}\right). \qquad (197)$$

where the last inequality follows from $\ln \ln(k) \leq \eta_k$ and $\beta > 1$.

For the rightmost probability in (188) we have by Boole's inequality that

$$\mathbb{P}\left(\left(E_{k}^{a}\right)^{c}, \sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil} \mathbb{1}\left\{\Omega_{ij}^{ma}\right\} \geq \lceil u_{k}-\eta_{k}\rceil\right) \\
\leq \sum_{(ij)\in\mathcal{L}^{a}} \mathbb{P}\left(\sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil} D_{ij}^{ma} < \lceil \sigma_{ijk}^{a}\rceil, \sum_{m=1}^{\lceil C_{\beta}+u_{k}\rceil} \mathbb{1}\left\{\Omega_{ij}^{ma}\right\} \geq \lceil u_{k}-\eta_{k}\rceil\right) \\
\stackrel{(184)}{\leq} \sum_{(ij)\in\mathcal{L}^{a}} \mathbb{P}\left(\sum_{m=1}^{\lceil u_{k}-\eta_{k}\rceil} \hat{D}_{ij}^{ma} < \lceil \sigma_{ijk}^{a}\rceil\right).$$
(198)

Recall from (36) that the length of an episode without the warmup period is H_0 . From Burke's Theorem [49, II.2.4 Theorem 2.1] and Poisson split and merge properties [48], we have that $\hat{D}_{ij}^{ma} \stackrel{\text{i.i.d.}}{\sim} \text{Poi}\left(x_{ij}^a H_0\right)$, and hence $\sum_{m=1}^k \hat{D}_{ij}^{ma} \sim \text{Poi}\left(x_{ij}^a H_0 k\right)$. Moreover, $u_k - \eta_k > \eta_k$ for k sufficiently large. Therefore, we obtain from Lemma C.3 and (43) that for k sufficiently large,

$$\mathbb{P}\left(\sum_{m=1}^{\lceil u_k - \eta_k \rceil} \hat{D}_{ij}^{ma} < \lceil \sigma_{ijk}^a \rceil\right) \le \exp\left(\frac{-(x_{ij}^a H_0 \lceil u_k - \eta_k \rceil - \lceil \sigma_{ijk}^a \rceil)^2}{2x_{ij}^a H_0 \lceil u_k - \eta_k \rceil}\right) \le \exp\left(\frac{-x_{ij}^a H_0 \eta_k^2}{2u_k}\right),\tag{199}$$

where we used $x_{ij}^a H_0 [u_k - \eta_k] - [\sigma_{ijk}^a] \ge x_{ij}^a H_0 \eta_k$ in the last equality. We use (36), the bounds $|\mathcal{L}^a| \le L$, $x_{ij}^a \ge x_{\min}$, together with the definitions of η_k and u_k in (43) to write

$$\sum_{(ij)\in\mathcal{L}^a} \mathbb{P}\left(\sum_{m=1}^{|u_k-\eta_k|} \hat{D}_{ij}^{ma} < \lceil \sigma_{ijk}^a \rceil\right) \le \sum_{(ij)\in\mathcal{L}^a} \exp\left(\frac{-x_{ij}^a H_0 \eta_k^2}{2u_k}\right) \le L \exp\left(-\frac{1}{2} x_{\min} H_0 \ln(k)\right)$$
(200)

$$= Lk^{-x_{\min}H_0/2} \le Lk^{-2}.$$
 (201)

Finally, we obtain from (188), (197), and (201),

$$\lim_{k \to \infty} k \mathbb{P}((E_k^a)^c) \le \lim_{k \to \infty} \left(Lk^{-1} + k \exp\left(\left[2\beta + 1 - \frac{\beta^2 + 1}{2}\ln\ln(k)\right]\eta_k\right)\right) = 0.$$
(202)

Here, the last equality follows since

$$k \exp\left(\left[2\beta + 1 - \frac{\beta^2 + 1}{2}\ln\ln(k)\right]\eta_k\right) \propto k^{1 - \log^{(\beta - 1)/2}(k)\ln\ln(k)},\tag{203}$$

and $\beta > 1$. \Box

C.9 Proof of Lemma 9.

In order to prove Lemma 9, we provide an intermediate result in Lemma C.4.

Lemma C.4. Let $(\sigma_k)_{k \in \mathbb{R}_{>0}} \in \mathbb{R}$ satisfy $\sigma_k = \omega(\ln(k))$ and let $\gamma \in \mathbb{R}_{>0}$. Then

$$\lim_{k \to \infty} k \int_{\sigma_k}^{\infty} \exp\left(-2m\left(\gamma - \sqrt{\frac{\ln(k)}{m}}\right)^2\right) \mathrm{d}m = 0.$$
(204)

Proof of Lemma C.4. Substitution of $\tau = \gamma \sqrt{m} - \sqrt{\ln(k)}$ with $dm/d\tau = 2(\tau + \sqrt{\ln(k)})/\gamma^2$ gives

$$\int_{\sigma_k}^{\infty} \exp\left(-2m\left(\gamma - \sqrt{\frac{\ln(k)}{m}}\right)^2\right) \mathrm{d}m = \int_{\gamma\sqrt{\sigma_k} - \sqrt{\ln(k)}}^{\infty} \exp\left(-2\tau^2\right) \frac{2(\tau + \sqrt{\ln(k)})}{\gamma^2} \mathrm{d}\tau.$$
 (205)

It can be verified that

$$\int_{\gamma\sqrt{\sigma_k}-\sqrt{\ln(k)}}^{\infty} \exp\left(-2\tau^2\right) \frac{2(\tau+\sqrt{\ln(k)})}{\gamma^2} \mathrm{d}\tau$$
(206)

$$=\frac{\exp\left(-2(\gamma\sqrt{\sigma_k}-\sqrt{\ln(k)})^2\right)+\sqrt{2\ln(k)\pi}\operatorname{erfc}\left(\sqrt{2}(\gamma\sqrt{\sigma_k}-\sqrt{\ln(k)})\right)}{2\gamma^2},$$
(207)

where $\operatorname{erfc}(x) := 1 - 2/\sqrt{\pi} \int_0^x \exp(-t^2) dt$ is the complementary error function. Now since $\sigma_k = \omega(\ln(k))$, we have

$$\exp\left(-2(\gamma\sqrt{\sigma_k} - \sqrt{\ln(k)})^2\right) = O\left(\exp(-\sigma_k)\right) = o(1/k).$$
(208)

Moreover, we use the bound $\operatorname{erfc}(x) \leq \exp(-x^2)/(x\sqrt{\pi})$ [58] to obtain

$$\sqrt{2\ln(k)\pi}\operatorname{erfc}\left(\sqrt{2}(\gamma\sqrt{\sigma_k} - \sqrt{\ln(k)})\right) = O\left(\frac{\sqrt{\ln(k)}\exp(-\sigma_k)}{\sqrt{\sigma_k} - \sqrt{\ln(k)}}\right) = o(1/k).$$
(209)

Substitution of (208) and (209) into (207) gives the result. \Box

We next present the proof of Lemma 9.

Proof of Lemma 9. Let $\lceil C_{\beta} + u_k \rceil \leq s \leq k$. Recall from (28) and (42) that

$$\mathbb{P}\left(\overline{U}^{a}[s,k] - r^{a} \ge d^{a} - \xi_{k}, \ E_{k}^{a}\right) = \mathbb{P}\left(\sum_{(ij)\in\mathcal{L}^{a}} x_{ij}^{a}\left(\overline{\theta}_{ij}\left[\overline{T}_{ij}^{a}[s]\right] - \theta_{ij} + \sqrt{\frac{\ln(k)}{\overline{T}_{ij}^{a}[s]}}\right) \ge d^{a} - \xi_{k}, \ E_{k}^{a}\right).$$
(210)

From Boole's inequality,¹ we obtain that for any $n, m, z \ge 0$,

$$\mathbb{P}\Big(\sum_{(ij)\in\mathcal{L}^a} x_{ij}^a \Big(\overline{\theta}_{ij}[n] - \theta_{ij} + \sqrt{\frac{\ln(m)}{n}}\Big) \ge z\Big) \le \sum_{(ij)\in\mathcal{L}^a} \mathbb{P}\Big(\overline{\theta}_{ij}[n] - \theta_{ij} + \sqrt{\frac{\ln(m)}{n}} \ge \frac{z}{x_{ij}^a |\mathcal{L}^a|}\Big).$$
(211)

Since E_k^a can be expressed as a union of disjoint events of the form $\{\overline{T}_{ij}^a[s] = m\}$, it follows from (210) and (211) that

$$\mathbb{P}\left(\overline{U}^{a}[s,k] - r^{a} \ge d^{a} - \xi_{k}, \ E_{k}^{a}\right) \le \sum_{(ij)\in\mathcal{L}^{a}} \mathbb{P}\left(\overline{\theta}_{ij}\left[\overline{T}_{ij}^{a}[s]\right] - \theta_{ij} + \sqrt{\frac{\ln(k)}{\overline{T}_{ij}^{a}[s]}} \ge \frac{d^{a} - \xi_{k}}{x_{ij}^{a}|\mathcal{L}^{a}|}, \ E_{k}^{a}\right).$$
(212)

Note that $\overline{T}_{ij}^{a}[\cdot]$ is non decreasing and that $s \geq \lceil C_{\beta} + u_k \rceil$, so on (44), $\overline{T}_{ij}^{a}[s] \geq \lceil \sigma_{ijk}^{a} \rceil$ for all $(ij) \in \mathcal{L}^a$. Therefore

$$\mathbb{P}\left(\overline{U}^{a}[s,k] - r^{a} \ge d^{a} - \xi_{k}, \ E_{k}^{a}\right) \le \sum_{(ij)\in\mathcal{L}^{a}} \mathbb{P}\left(\overline{\theta}_{ij}\left[\overline{T}_{ij}^{a}[s]\right] - \theta_{ij} + \sqrt{\frac{\ln(k)}{\overline{T}_{ij}^{a}[s]}} \ge \frac{d^{a} - \xi_{k}}{x_{ij}^{a}|\mathcal{L}^{a}|}, \ \overline{T}_{ij}[s] \ge \lceil \sigma_{ijk}^{a} \rceil\right).$$
(213)

By partitioning the event based on the value of $\overline{T}_{ij}^{a}[s]$, we find that

$$\mathbb{P}\left(\overline{U}^{a}[s,k] - r^{a} \ge d^{a} - \xi_{k}, \ E_{k}^{a}\right) \le \sum_{(ij)\in\mathcal{L}^{a}} \sum_{m=\lceil\sigma_{ijk}^{a}\rceil}^{\infty} \mathbb{P}\left(\overline{\theta}_{ij}\left[m\right] - \theta_{ij} + \sqrt{\frac{\ln(k)}{m}} \ge \frac{d^{a} - \xi_{k}}{x_{ij}^{a}|\mathcal{L}^{a}|}\right).$$
(214)

Since the right side of (214) is independent of s, we have

$$\sum_{i=\lceil C_{\beta}+u_{k}\rceil}^{k} \mathbb{P}\left(\overline{U}^{a}[s,k]-r^{a} \geq d^{a}-\xi_{k}, E_{k}^{a}\right) \leq k \sum_{(ij)\in\mathcal{L}^{a}} \sum_{m=\lceil\sigma_{ijk}^{a}\rceil}^{\infty} \mathbb{P}\left(\overline{\theta}_{ij}\left[m\right]-\theta_{ij}+\sqrt{\frac{\ln(k)}{m}} \geq \frac{d^{a}-\xi_{k}}{x_{ij}^{a}|\mathcal{L}^{a}|}\right).$$
(215)

¹For any random variables X, Y and any $z \ge 0$, $\{X + Y \ge z\} \subseteq \{X \ge z/2\} \cup \{Y \ge z/2\}$. Therefore $\mathbb{P}(X + Y \ge z) \le \mathbb{P}(\{X \ge z/2\} \cup \{Y \ge z/2\})$. Applying Boole's inequality gives $\mathbb{P}(X + Y \ge z) \le \mathbb{P}(X \ge z/2) + \mathbb{P}(Y \ge z/2)$.

Note that by construction $\sigma_{ijk}^a = \omega(\ln(k))$ (recall (43)) and $\xi_k = o(1)$, so that $(d^a - \xi_k)/(x_{ij}^a |\mathcal{L}^a|) > \sqrt{\ln(k)/\sigma_{ijk}^a}$ for $k \in \mathbb{N}_{\geq 1}$ sufficiently large. For such k, we have by (i) Hoeffding's inequality [50, Theorem 2] and (ii) Lemma C.4 that

$$\sum_{(ij)\in\mathcal{L}^a}\sum_{m=\lceil\sigma_{ijk}^a\rceil}^{\infty} \mathbb{P}\Big(\overline{\theta}_{ij}\big[m\big] - \theta_{ij} + \sqrt{\frac{\ln(k)}{m}} \ge \frac{d^a - \xi_k}{x_{ij}^a|\mathcal{L}^a|}\Big) \stackrel{(i)}{\le} \sum_{(ij)\in\mathcal{L}^a}\sum_{m=\lceil\sigma_{ijk}^a\rceil}^{\infty} \exp\Big(-2m\Big(\frac{d^a - \xi_k}{x_{ij}^a|\mathcal{L}^a|} - \sqrt{\frac{\ln(k)}{m}}\Big)^2\Big)$$
(216)

$$\leq \sum_{(ij)\in\mathcal{L}^a} \int_{\sigma_{ijk}^a}^{\infty} \exp\left(-2m\left(\frac{d^a-\xi_k}{x_{ij}^a|\mathcal{L}^a|}-\sqrt{\frac{\ln(k)}{m}}\right)^2\right) \mathrm{d}m \qquad (217)$$

$$\stackrel{\text{(ii)}}{=} o(1/k). \tag{218}$$

The result follows from (215) and (218). \Box

C.10 Proof of Lemma 10

By definitions (28), (32), and (33), we have that for any $m = 1, \ldots, k$,

$$\mathbb{P}(U^{a}(m) < r^{a} - \xi) = \mathbb{P}\left(\sum_{(ij)\in\mathcal{L}^{a}} x^{a}_{ij}\left(\hat{\theta}_{ij}(m) - \theta_{ij} + \sqrt{\frac{\ln(m)}{T_{ij}(m)}}\right) < -\xi\right).$$
(219)

We use the inequality $\mathbb{P}(A + B < z) \le \mathbb{P}(A < z) + \mathbb{P}(B < z)$ for any events A, B and any $z \in \mathbb{R}$ to write

$$\mathbb{P}\left(U^{a}(m) < r^{a} - \xi\right) \leq \sum_{(ij)\in\mathcal{L}^{a}} \mathbb{P}\left(\hat{\theta}_{ij}(m) - \theta_{ij} + \sqrt{\frac{\ln(m)}{T_{ij}(m)}} < -\frac{\xi}{x_{ij}^{a}}\right).$$
(220)

By partitioning the event based on the value of $T_{ij}(m)$, we find that

$$\mathbb{P}(U^a(m) < r^a - \xi) \le \sum_{(ij)\in\mathcal{L}^a} \sum_{s=1}^{\infty} \mathbb{P}\left(\hat{\theta}_{ij}(m) - \theta_{ij} + \sqrt{\frac{\ln(m)}{T_{ij}(m)}} < -\frac{\xi}{x_{ij}^a}, \ T_{ij}(m) = s\right)$$
(221)

$$\leq \sum_{(ij)\in\mathcal{L}^a} \sum_{s=1}^{\infty} \mathbb{P}\Big(\overline{\theta}_{ij}[s] - \theta_{ij} + \sqrt{\frac{\ln(m)}{s}} < -\frac{\xi}{x_{ij}^a}\Big).$$
(222)

Since the distribution of the payoff parameters are [0, 1]-bounded and independent, we can apply Hoeffding's inequality [50, Theorem 2] to obtain

$$\mathbb{P}(U^a(m) < r^a - \xi) \le \sum_{(ij)\in\mathcal{L}^a} \sum_{s=1}^{\infty} \exp\left(-2s\left(\frac{\xi}{x_{ij}^a} + \sqrt{\frac{\ln(m)}{s}}\right)^2\right)$$
(223)

$$\leq \sum_{(ij)\in\mathcal{L}^a} \sum_{s=1}^{\infty} \exp\left(-2s\left(\frac{\xi}{x_{ij}^a}\right)^2 - 2\ln(m)\right)$$
(224)

$$= \frac{1}{m^2} \sum_{(ij)\in\mathcal{L}^a} \sum_{s=1}^{\infty} \exp\left(-2s\left(\frac{\xi}{x_{ij}^a}\right)^2\right).$$
(225)

We use that for any y > 0 we have

$$\sum_{s=1}^{\infty} \exp(-sy) = \frac{1}{\exp(y) - 1} \le \frac{1}{y},$$
(226)

to write

$$\mathbb{P}(U^{a}(m) < r^{a} - \xi) \leq \frac{1}{m^{2}} \sum_{(ij)\in\mathcal{L}^{a}} \frac{(x_{ij}^{a})^{2}}{2\xi^{2}} \leq \frac{\|\lambda\|_{2}^{2}}{2m^{2}\xi^{2}},$$
(227)

where we used that $x_{ij}^a \leq \lambda_i$ for all $(ij) \in \mathcal{L}^a$. Taking the sum over all episodes gives

$$\sum_{m=1}^{k} \mathbb{P}(U^{a}(m) < r^{a} - \xi) \le \sum_{m=1}^{\infty} \mathbb{P}(U^{a}(m) < r^{a} - \xi) \le \sum_{m=1}^{\infty} \frac{\|\lambda\|_{2}^{2}}{2m^{2}\xi^{2}} = \frac{\pi^{2}\|\lambda\|_{2}^{2}}{12\xi^{2}}.$$
(228)

C.11 Proof of Lemma 11

Let $k \in \mathbb{N}_{\geq 1}$ be such that $k \geq C_{\beta}$. Let $a \in \mathcal{A} \setminus \{1\}$ be a suboptimal action and x^a its corresponding nondegenerate basic feasible solution of $LP(\theta, \varepsilon)$ in (10). Starting from (45), we split the summation based on whether the UCB index $U^a(m)$ is smaller or larger than the true optimal mean r^1 minus some small error $\xi_k := \ln^{-\frac{\beta}{4}}(k)$:

$$S^{a}(k) = \sum_{m=1}^{k} \mathbb{1}\left\{U^{a}(m) < r^{1} - \xi_{k}, \ A_{m} = a\right\} + \sum_{m=1}^{k} \mathbb{1}\left\{U^{a}(m) \ge r^{1} - \xi_{k}, \ A_{m} = a\right\}.$$
(229)

By construction, action a can be chosen only if the UCB index U^a is as large as the indices of the other actions, including the optimal action U^1 (Line 3 in Algorithm 1). Therefore,

$$\sum_{m=1}^{k} \mathbb{1}\left\{ U^{a}(m) < r^{1} - \xi_{k}, \ U^{1}(m) \le U^{a}(m) \right\} \le \sum_{m=1}^{k} \mathbb{1}\left\{ U^{1}(m) < r^{1} - \xi_{k} \right\}.$$
(230)

Taking the expectation of (229) yields

$$\mathbb{E}(S^{a}(k)) \leq \sum_{m=1}^{k} \mathbb{P}(U^{1}(m) < r^{1} - \xi_{k}) + \sum_{m=1}^{k} \mathbb{P}(U^{a}(m) \geq r^{1} - \xi_{k}, A_{m} = a).$$
(231)

We will next analyze the two terms on the right side of (231) more carefully.

Note that $\xi_k \to 0$ as $k \to \infty$, so $\xi_k < r^1$ for k sufficiently large. From Lemma 10, and using the definition of ξ_k , we find

$$\lim_{k \to \infty} \frac{\sum_{m=1}^{k} \mathbb{P}(U^{1}(m) < r^{1} - \xi_{k})}{\ln^{\beta}(k)} \le \lim_{k \to \infty} \frac{\pi^{2} \|\lambda\|_{2}^{2} \ln^{\beta/2}(k)}{12 \ln^{\beta}(k)} = 0.$$
(232)

For the second term in (231), we note that the summation can be simplified as follows. Suppose that $m_1 \in \mathbb{N}_{\geq 1}$ is the first episode in which Algorithm 1 chooses action a in Line 3. Then $U^a(m_1) = \overline{U}^a[1, m_1]$. Similarly, $U^a(m_2) = \overline{U}^a[2, m_2]$ where m_2 is the second episode in which Algorithm 1 chooses action a; so on and so on. Moreover, for fixed a sample path, $\overline{U}[s, k]$ is nondecreasing in k (recall (42)). Therefore,

$$\sum_{m=1}^{k} \mathbb{P}(U^{a}(m) \ge r^{1} - \xi_{k}, A_{m} = a) \le \sum_{s=1}^{k} \mathbb{P}(\overline{U}^{a}[s, k] \ge r^{1} - \xi_{k}).$$
(233)

Substitution of $r^1 = r^a + d^a$ gives

$$\sum_{m=1}^{k} \mathbb{P}(U^{a}(m) \ge r^{1} - \xi_{k}, A_{m} = a) \le \sum_{s=1}^{k} \mathbb{P}(\overline{U}^{a}[s, k] - r^{a} \ge d^{a} - \xi_{k}).$$
(234)

We bound the probability of the first $C_{\beta} + u_k$ episodes in (234) by one,

$$\sum_{m=1}^{k} \mathbb{P}(U^{a}(m) \ge r^{1} - \xi_{k}, A_{m} = a) \le C_{\beta} + u_{k} + \sum_{s \in \lceil C_{\beta} + u_{k} \rceil}^{k} \mathbb{P}\left(\overline{U}^{a}[s, k] - r^{a} \ge d^{a} - \xi_{k}\right).$$
(235)

By (44) and the law of total probability, we obtain

$$\sum_{s=\lceil C_{\beta}+u_k\rceil}^k \mathbb{P}\left(\overline{U}^a[s,k] - r^a \ge d^a - \xi_k\right) \le \sum_{s=\lceil C_{\beta}+u_k\rceil}^k \mathbb{P}\left(\overline{U}^a[s,k] - r^a \ge d^a - \xi_k, \ E_k^a\right) + \sum_{s=\lceil C_{\beta}+u_k\rceil}^k \mathbb{P}\left((E_k^a)^c\right).$$
(236)

From Lemma 8 it follows that

$$\lim_{k \to \infty} \sum_{s = \lceil C_{\beta} + u_k \rceil}^k \mathbb{P}((E_k^a)^c) \le \lim_{k \to \infty} k \mathbb{P}((E_k^a)^c) = 0.$$
(237)

Moreover, Lemma 9 implies immediately that the middle term in (236) converges to zero, hence

$$\lim_{k \to \infty} \sum_{s = \lceil C_{\beta} + u_k \rceil}^k \mathbb{P}\left(\overline{U}^a[s,k] - r^a \ge d^a - \xi_k\right) = 0.$$
(238)

We continue to bound (229),

$$\lim_{k \to \infty} \frac{\mathbb{E}(S^{a}(k))}{\ln^{\beta}(k)} \stackrel{(231)}{\leq} \lim_{k \to \infty} \frac{\sum_{m=1}^{k} \mathbb{P}(U^{1}(m) < r^{1}) + \sum_{m=1}^{k} \mathbb{P}(U^{a}(m) \ge r^{1}, A_{m} = a)}{\ln^{\beta}(k)}$$
(239)

$$\leq \lim_{k \to \infty} \frac{\sum_{m=1}^{k} \mathbb{P}(U^{1}(m) < r^{1})}{\ln^{\beta}(k)} + \frac{C_{\beta} + u_{k} + \sum_{s=\lceil C_{\beta} + u_{k} \rceil}^{k} \mathbb{P}\left(\overline{U}^{a}[s,k] - r^{a} \geq d^{a} - \xi_{k}\right)}{\ln^{\beta}(k)}.$$
 (240)

Now, by (232),

$$\lim_{k \to \infty} \frac{\sum_{m=1}^{k} \mathbb{P}(U^{1}(m) < r^{1})}{\ln^{\beta}(k)} = 0.$$
(241)

Moreover, C_{β} is a constant and $\lim_{k\to\infty} u_k / \log^{\beta}(k) = 1$ (recall (43)). Lastly, by (238),

$$\lim_{k \to \infty} \frac{\sum_{s=\lceil C_{\beta}+u_k \rceil}^k \mathbb{P}\left(\overline{U}^a[s,k] - r^a \ge d^a - \xi_k\right)}{\log^\beta(k)} = 0.$$
(242)

So we find that

$$\lim_{k \to \infty} \frac{\mathbb{E}(S^a(k))}{\ln^\beta(k)} = 1,$$
(243)

which concludes the proof. \Box

Acknowledgements

The authors would like to thank Thomas van Vuren for his insightful comments and fruitful discussions.

This work is part of *Valuable AI*, a research collaboration between the Eindhoven University of Technology and the Koninklijke KPN N.V. Parts of this research have been funded by the EAISI's IMPULS program, and by Holland High Tech — TKI HSTM via the PPS allowance scheme for public-private partnerships.

References

- M. Harchol-Balter. Performance Modeling and Design of Computer Systems: Queueing Theory in Action. Cambridge University Press, UK, 2013.
- [2] Jinsheng Chen, Jing Dong, and Pengyi Shi. A survey on skill-based routing with applications to service operations management. Queueing Systems, 96(1-2):53–82, 2020.
- [3] Ger Koole and Avishai Mandelbaum. Queueing models of call centers: An introduction. Annals of Operations Research, 113(1-4):41-59, 2002.
- [4] Tze Leung Lai and Herbert Robbins. Asymptotically efficient adaptive allocation rules. Advances in Applied Mathematics, 6(1):4–22, 1985.
- [5] Peter Auer, Nicolò Cesa-Bianchi, and Paul Fischer. Finite-time analysis of the multiarmed bandit problem. Machine Learning, 47(2-3):235-256, 2002.
- [6] Dimitris Bertsimas and John N. Tsitsiklis. Introduction to Linear Optimization. Number 1. Athena scientific, Belmont, MA, 1997.
- [7] Apostolos N. Burnetas and Michael N. Katehakis. Optimal adaptive policies for sequential allocation problems. Advances in Applied Mathematics, 17(2):122–142, 1996.
- [8] Apostolos N. Burnetas, Odysseas Kanavetas, and Michael N. Katehakis. Asymptotically optimal multiarmed bandit policies under a cost constraint. Probability in the Engineering and Informational Sciences, 31(3):284–310, 2017.
- [9] Apostolos N. Burnetas and Odysseas Kanavetas. Adaptive policies for sequential sampling under incomplete information and a cost constraint. *Springer Optimization and its Applications*, 71(January):97–112, 2012.
- [10] Apostolos N. Burnetas and Michael N. Katehakis. Optimal adaptive policies for Markov Decision Processes. Mathematics of Operations Research, 22(1):222–255, 1997.
- [11] Tor Lattimore and Csaba Szepesvári. Bandit Algorithms. 2020.
- [12] Itai Gurvich and Ward Whitt. Service-level differentiation in many-server service systems via queue-ratio routing. Operations Research, 58(2):316–328, 2010.
- [13] Jeremy Visschers, Ivo Adan, and Gideon Weiss. A product form solution to a system with multi-type jobs and multi-type servers. *Queueing Systems*, 70(3):269–298, 2012.
- [14] Ivo Adan and Gideon Weiss. Exact FCFS matching rates for two infinite multitype sequences. Operations Research, 60(2):475–489, 2012.

- [15] Ivo Adan and Gideon Weiss. A skill based parallel service system under FCFS-ALIS steady state, overloads, and abandonments. *Stochastic Systems*, 4(1):250–299, 2014.
- [16] Zixian Yang, R. Srikant, and Lei Ying. Learning while scheduling in multi-server systems with unknown statistics: MaxWeight with Discounted UCB. Proceedings of The 26th International Conference on Artificial Intelligence and Statistics, 206:4275—4312, 2023.
- [17] Subhashini Krishnasamy, Ari Arapostathis, Ramesh Johari, and Sanjay Shakkottai. On learning the cμ rule in single and parallel server networks. 2018 56th Annual Allerton Conference on Communication, Control, and Computing, Allerton 2018, pages 153–154, 2018.
- [18] Dabeen Lee and Milan Vojnovic. Scheduling jobs with stochastic holding costs. Advances in Neural Information Processing Systems, 23:19375–19384, 2021.
- [19] Tuhinangshu Choudhury, Gauri Joshi, Weina Wang, and Sanjay Shakkottai. Job dispatching policies for queueing systems with unknown service rates, volume 1. Association for Computing Machinery, 2021.
- [20] Yueyang Zhong, John R. Birge, and Amy Ward. Learning the scheduling policy in time-varying multiclass many server queues with abandonment. SSRN Electronic Journal, (2012), 2022.
- [21] Ramesh Johari, Vijay Kamble, and Yash Kanoria. Matching while learning. Operations Research, 69(2):655– 681, 2021.
- [22] Huiwen Jia, Cong Shi, and Siqian Shen. Online learning and pricing for service systems with reusable resources. *Operations Research*, 72(3):1203–1241, 2022.
- [23] Bai Liu, Qiaomin Xie, and Eytan Modiano. RL-QN: A reinforcement learning framework for optimal control of queueing systems. 2019 57th Annual Allerton Conference on Communication, Control, and Computing, Allerton 2019, pages 663–670, 2019.
- [24] Virag Shah, Lennart Gulikers, Laurent Massoulié, and Milan Vojnović. Adaptive matching for expert systems with uncertain task types. Operations Research, 68(5):1403–1424, 2020.
- [25] Kostas Bimpikis and Mihalis G. Markakis. Learning and hierarchies in service systems. Management Science, 65(3):1268–1285, 2019.
- [26] Bo Tan and R. Srikant. Online advertisement, optimization and stochastic networks. *IEEE Transactions on Automatic Control*, 57(11):2854–2868, 2012.
- [27] Xiaoqi Tan, Bo Sun, Alberto Leon-Garcia, Yuan Wu, and Danny H.K. Tsang. Mechanism design for online resource allocation: A unified approach. *Performance Evaluation Review*, 48(1):11–12, 2020.
- [28] Alberto Vera and Siddhartha Banerjee. The Bayesian prophet: A low-regret framework for online decision making. *Management Science*, 67(3):1368–1391, 2021.
- [29] Alekh Agarwal, Dean P. Foster, Daniel Hsu, Sham M. Kakade, and Alexander Rakhlin. Stochastic convex optimization with bandit feedback. SIAM Journal on Optimization, 23(1):213–240, 2013.
- [30] Shipra Agrawal, Zizhuo Wang, and Yinyu Ye. A dynamic near-optimal algorithm for online linear programming. Operations Research, 62(4):876–890, 2014.
- [31] Hao Yu, Michael J. Neely, and Xiaohan Wei. Online convex optimization with stochastic constraints. Advances in Neural Information Processing Systems, 2017-Decem:1429–1439, 2017.
- [32] Xu Sun, Jingtong Zhao, and Shiwei Chai. Congestion-aware matching and learning for service platforms. 2023.
- [33] Wei Kang Hsu, Jiaming Xu, Xiaojun Lin, and Mark R. Bell. Integrated online learning and adaptive control in queueing systems with uncertain payoffs. *Operations Research*, 70(2):1166–1181, 2022.
- [34] Kim and Milan Vojnovic. Scheduling servers with stochastic bilinear rewards. pages 1–63, 2021.
- [35] Xin Liu, Bin Li, Pengyi Shi, and Lei Ying. POND: Pessimistic–Optimistic oNline Dispatching. 2020.
- [36] Juaren Steiger, Bin Li, and Ning Lu. Learning from Delayed Semi-Bandit Feedback under Strong Fairness Guarantees. In IEEE INFOCOM 2022 - IEEE Conference on Computer Communications, pages 1379– 1388, 2022.

- [37] Xinzhe Fu and Eytan Modiano. Joint learning and control in stochastic queueing networks with unknown utilities. Proceedings of the ACM on Measurement and Analysis of Computing Systems, 6(3), 2022.
- [38] Varsha Dani, Thomas P. Hayes, and Sham M. Kakade. Stochastic linear optimization under bandit feedback. 21st Annual Conference on Learning Theory, COLT 2008, pages 355–366, 2008.
- [39] Boxiao Chen and Cong Shi. Tailored base-surge policies in dual-sourcing inventory systems with demand learning. SSRN Electronic Journal, (2010), 2019.
- [40] Omar Besbes and Assaf Zeevi. On the (surprising) sufficiency of linear models for dynamic pricing with demand learning. *Management Science*, 61(4):723–739, 2015.
- [41] Jaron Sanders, Sem C. Borst, and Johan S.H. Van Leeuwaarden. Online network optimization using product-form Markov processes. *IEEE Transactions on Automatic Control*, 61(7):1838–1853, 2016.
- [42] Céline Comte, Matthieu Jonckheere, Jaron Sanders, and Albert Senen-Cerda. Score-aware policy-gradient methods and performance guarantees using local Lyapunov conditions: applications to product-form stochastic networks and queueing systems. 2023.
- [43] Libin Jiang and Jean Walrand. A distributed CSMA algorithm for throughput and utility maximization in wireless networks. *IEEE/ACM Transactions on Networking*, 18(3):960–972, 2010.
- [44] Dimitris Bertsimas. The achievable region method in the optimal control of queueing systems; formulations, bounds and policies. *Queueing Systems*, 21(3-4):337–389, 1995.
- [45] Marcus Dacre, Kevin Glazebrook, and José Niño-Mora. The achievable region approach to the optimal control of stochastic systems. Journal of the Royal Statistical Society. Series B: Statistical Methodology, 61(4):747-791, 1999.
- [46] L. S. Shapley and M. Shubik. The assignment game I: The core. International Journal of Game Theory, 1(1):111–130, 1971.
- [47] Xinzhe Fu and Eytan Modiano. Learning-NUM: Network Utility Maximization with unknown utility functions and queueing delay. Proceedings of the International Symposium on Mobile Ad Hoc Networking and Computing (MobiHoc), pages 21–30, 2021.
- [48] E. Çinlar and Robert Agnew. On the Superposition of Point Processes. Journal of the Royal Statistical Society: Series B (Methodological), 30:576–581, 1968.
- [49] Joel E. Cohen and Frank P. Kelly. Reversibility and Stochastic Networks, 1981.
- [50] Wassily Hoeffding. Probability Inequalities for Sums of Bounded Random Variables. Journal of the American Statistical Association, 58(301):13–30, 1963.
- [51] Jacques Desrosiers and Marco E. Lübbecke. A primer in column generation. Column Generation, (March 2006):1–32, 2005.
- [52] Fernando García-Muñoz, Sebastián Dávila, and Franco Quezada. A Benders decomposition approach for solving a two-stage local energy market problem under uncertainty. *Applied Energy*, 329(November 2022):120226, 2023.
- [53] Maury Bramson. Stability of Queueing Networks. Springer-Verlag Berlin Heidelberg, 2006.
- [54] Frank Harary. Graphs and Matrices. SIAM Review, 9(1):83–90, 1967.
- [55] Wenpin Tang and Fengmin Tang. The Poisson Binomial Distribution Old & New. Statistical Science, 38(1):108–119, 2023.
- [56] J. N. Darroch. On the distribution of the number of successes in independent trials. The Annals of Mathematical Statistics, 35(3):1317–1321, 1964.
- [57] Herbert Robbins. A remark on Stirling's Formula. The American Mathematical Monthly, 62(1):26–29, 1955.
- [58] George K. Karagiannidis and Athanaszsios S. Lioumpas. An improved approximation for the Gaussian Q-function. *IEEE Communications Letters*, 11(8):644–646, 2007.