Quasi-orthogonal extension of symmetric matrices

Abderrahim Boussaïri^{*,1}, Brahim Chergui¹, Zaineb Sarir¹, and Mohamed Zouagui²

¹Laboratoire Mathématiques Fondamentales et Appliquées, Faculté des Sciences Aïn Chock, Hassan II University of Casablanca, Morocco ²Laboratoire de Recherche Mathématiques et Sciences de l'Ingénieur MSI, International University of Casablanca, School of Engineering, Casa Green Town, Bouskoura, Morocco

December 16, 2024

Abstract

An $n \times n$ real matrix Q is quasi-orthogonal if $Q^{\top}Q = qI_n$ for some positive real number q. If M is a principal sub-matrix of a quasi-orthogonal matrix Q, we say that Q is a quasi-orthogonal extension of M. In a recent work, the authors have investigated this notion for the class of real skew-symmetric matrices. Using a different approach, this paper addresses the case of symmetric matrices.

Keywords: Symmetric matrix; principal sub-matrix; quasi-orthogonal matrix; Seidel matrix.

MSC Classification: 15A18; 15B10.

1 Introduction

All the matrices considered in this paper are real. The identity matrix of order n and the $n \times m$ all-zeros matrix are respectively denoted by I_n and $O_{n,m}$. We omit the subscript when the order is understood. The characteristic polynomial of an $n \times n$ matrix A is $\phi_A(x) := \det(xI_n - A)$.

An $n \times n$ real matrix Q is *quasi-orthogonal* if $Q^{\top}Q = qI_n$, for some positive real number q or equivalently, the matrix $\frac{1}{\sqrt{q}}Q$ is orthogonal. Special classes of quasi-orthogonal matrices are the set of Hadamard matrices and conference matrices. Recall that a *Hadamard matrix* is a square matrix with entries in $\{-1, 1\}$ and whose columns are mutually orthogonal. The order of such matrices must be 1, 2 or a multiple of 4. A *conference matrix* is an $n \times n$ matrix C with 0 on the diagonal and ± 1 off the diagonal

^{*}Corresponding author, email: aboussairi@hotmail.com

such that $C^{\top}C = (n-1)I_n$. Hadamard and conference matrices have been extensively studied since they are related to many combinatorial problems. We refer to the Handbook of Combinatorial Designs [4] for relevant background. Taussky [10] suggested the following generalization of Hadamard and conference matrices. A *weighing matrix* of weight k and order n is an $n \times n$ {-1,0,1} matrix A such that $AA^{\top} = kI_n$. Recent results on these topics can be found in [9].

Let *M* be a square matrix. If *M* is a principal sub-matrix of a quasi-orthogonal matrix *Q*, we say that *Q* is a *quasi-orthogonal extension* of *M*. Any symmetric (resp. skew-symmetric) matrix with 0 in the diagonal and ± 1 off the diagonal is a principal sub-matrix of a symmetric (resp. skew-symmetric) conference matrix. These facts follow from [2, Theorem 3] and [5]. In [3], the authors proved that every skew-symmetric matrix has a skew-symmetric quasi-orthogonal extension.

This paper deals with the real symmetric matrices. Let M be a symmetric matrix. We prove in Section 2 that M has a symmetric quasi-orthogonal extension. We define the *quasi-orthogonality index* of M as the least integer d, denoted by ind(M), such that M has a symmetric quasi-orthogonal extension of order n + d.

The remainder of the paper is organized as follows. In Section 3, we work out the quasi-orthogonality index using the well-known Cauchy's interlace theorem. Section 4 is devoted to $n \times n$ Seidel matrices with minimum quasi-orthogonality index.

2 Existence of symmetric quasi-orthogonal extension of symmetric matrices

Let *S* be a non-zero square symmetric matrix. For a real number λ , we denote by $\mu_{\lambda}(S)$ (or simply by μ_{λ}) its algebraic multiplicity in *S* if λ is an eigenvalue of *S*. We convene that $\mu_{\lambda} = 0$ whenever λ is not an eigenvalue of *S*.

The next theorem guarantees the existence of symmetric quasi-orthogonal extension of symmetric matrices.

Theorem 2.1. Let *S* be an $n \times n$ non-zero real symmetric matrix with spectral radius ρ . Then *S* has a symmetric quasi-orthogonal extension \hat{S} of order $2n - \mu_{\rho} - \mu_{-\rho}$.

Proof. Since the matrix *S* is real and symmetric, it is diagonalizable, and hence there exists an $n \times n$ orthogonal matrix *P* such that

$$P^{\top}SP = \begin{pmatrix} D & O \\ O & \Sigma \end{pmatrix} \tag{1}$$

where *D* is a diagonal matrix and Σ is a diagonal matrix of order $\mu_{\rho} + \mu_{-\rho}$ with diagonal entries $\pm \rho$. Then we have

$$P^{\top}S^2P = \begin{pmatrix} D^2 & O \\ O & \rho^2I \end{pmatrix}.$$

We partition the matrix P as $P := \begin{pmatrix} N & L \end{pmatrix}$, where N and L are sub-matrices of orders $n \times (n - \mu_{-\rho} - \mu_{\rho})$ and $n \times (\mu_{-\rho} + \mu_{\rho})$ respectively. Note that the diagonal matrix $\rho^2 I - D^2$ has positive diagonal entries. We set $M := N \cdot \sqrt{\rho^2 I - D^2}$. Let us consider the following extension \hat{S} of S

$$\hat{S} := \begin{pmatrix} S & M \\ M^\top & -D \end{pmatrix}.$$

We have

$$\hat{S}^2 = \begin{pmatrix} S^2 + MM^\top & SM - MD \\ M^\top S - DM^\top & M^\top M + D^2 \end{pmatrix}.$$

Equality (1) implies that SN = ND. Right multiplying both sides by $\sqrt{\rho^2 I - D^2}$, we get SM = MD. Hence

$$SM - MD = M^{\top}S - DM^{\top} = O.$$
 (2)

As *P* is an orthogonal matrix and $P^{\top}P = \begin{pmatrix} N^{\top}N & N^{\top}L \\ L^{\top}N & L^{\top}L \end{pmatrix}$, we obtain $N^{\top}N = I$. It follows that $M^{\top}M = \rho^2 I - D^2$. Then

$$M^{\top}M + D^2 = \rho^2 I. \tag{3}$$

Moreover

$$MM^{\top} = N(\rho^2 I - D^2)N^{\top}$$
$$= \rho^2 NN^{\top} - ND^2N^{\top}$$

and

$$S^{2} = P \begin{pmatrix} D^{2} & O \\ O & \rho^{2}I \end{pmatrix} P^{\top}$$
$$= \begin{pmatrix} N & L \end{pmatrix} \begin{pmatrix} D^{2} & O \\ O & \rho^{2}I \end{pmatrix} \begin{pmatrix} N^{\top} \\ L^{\top} \end{pmatrix}$$
$$= ND^{2}N^{\top} + \rho^{2}LL^{\top}.$$

It follows that $S^2 + MM^{\top} = \rho^2(NN^{\top} + LL^{\top})$. Given that $PP^{\top} = NN^{\top} + LL^{\top} = I$, we have

$$S^2 + M^{\top}M = \rho^2 I. \tag{4}$$

From (2), (3) and (4) we deduce that \hat{S} is a symmetric quasi-orthogonal extension of order $2n - \mu_{-\rho} - \mu_{\rho}$.

3 Quasi-orthogonality index of a symmetric matrix

The following theorem gives the quasi-orthogonality index for real symmetric matrices.

Theorem 3.1. Let S be an $n \times n$ symmetric matrix with spectral radius ρ . The quasiorthogonality index of S is equal to $n - \mu_{\rho} - \mu_{-\rho}$.

To prove this theorem, we need the following results.

Theorem 3.2 (Cauchy's interlace theorem). Let A be an $n \times n$ symmetric matrix and let B be a principal sub-matrix of order m < n. Suppose A has eigenvalues $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n$ and B has eigenvalues $\beta_1 \leq \beta_2 \leq \cdots \leq \beta_m$. Then

$$\lambda_k \leq \beta_k \leq \lambda_{k+n-m}$$
 for $k = 1, \dots, m$,

in particular, if m = n - 1, we have

$$\lambda_1 \leq \beta_1 \leq \lambda_2 \leq \beta_2 \leq \cdots \leq \beta_{n-1} \leq \lambda_n.$$

Lemma 3.3. Let *S* be a real symmetric matrix of order *m* with spectral radius ρ . Let \hat{S} be a symmetric quasi-orthogonal extension of order $n \ge 2$, with spectral radius $\hat{\rho}$. If $n - m \le (n-1)/2$, then $\hat{\rho} = \rho$ and $n \ge 2m - (\mu_{\rho}(S) + \mu_{-\rho}(S))$.

Proof. Let $\lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n$ and $\beta_1 \leq \beta_2 \leq \cdots \leq \beta_m$ respectively be the eigenvalues of \hat{S} and S. We set $f(S) := m - (\mu_{\rho}(S) + \mu_{-\rho}(S))$. Without loss of generality we can assume that $\mu_{-\hat{\rho}}(\hat{S}) \geq \mu_{\hat{\rho}}(\hat{S})$. We will distinguish two cases:

• Suppose that $n-m \ge \mu_{\hat{\rho}}(\hat{S})$. As \hat{S} is a symmetric quasi-orthogonal matrix, $\mu_{\hat{\rho}}(\hat{S}) + \mu_{-\hat{\rho}}(\hat{S}) = n$. It follows that

$$2\mu_{-\hat{\rho}}(\hat{S}) \ge n.$$

Hence

$$\mu_{-\hat{\rho}}(\hat{S}) > \frac{n-1}{2} \ge n-m.$$

Let $k \in \{1, \ldots, \mu_{-\hat{\rho}}(\hat{S}) - (n-m)\}$. As $k + n - m \leq \mu_{-\hat{\rho}}(\hat{S})$, we have $\lambda_k = \lambda_{k+n-m} = -\hat{\rho}$. By Theorem 3.2, we get $\beta_k = -\hat{\rho}$. Then $\hat{\rho} = \rho$. Moreover, we have

$$f(S) \le m - (\mu_{-\hat{\rho}}(\hat{S}) - (n-m)).$$

Then

$$f(S) \le n - \mu_{-\hat{\rho}}(\hat{S}).$$

Hence

$$f(S) \le \mu_{\hat{\rho}}(S) \le n - m.$$

Consequently, $n \ge 2m - (\mu_{\rho}(S) + \mu_{-\rho}(S))$.

• Suppose that $n - m < \mu_{\hat{\rho}}(\hat{S})$, then $n - m < \mu_{-\hat{\rho}}(\hat{S})$.

If $k \in \{1, \ldots, \mu_{-\hat{\rho}}(\hat{S}) - (n-m)\}$, then we have $k+n-m \leq \mu_{-\hat{\rho}}(\hat{S})$ and $\lambda_k = \lambda_{k+n-m} = -\hat{\rho}$. It follows from Theorem 3.2 that $\beta_k = -\hat{\rho}$ for $k \in \{1, \ldots, \mu_{-\hat{\rho}}(\hat{S}) - (n-m)\}$.

If $k \in {\mu_{-\hat{\rho}}(\hat{S}) + 1, ..., m}$, then $\lambda_k = \lambda_{k+n-m} = \hat{\rho}$ and hence $\beta_k = \hat{\rho}$. Therefore, $f(S) \leq m - [\mu_{-\hat{\rho}}(\hat{S}) - (n - m) + m - \mu_{-\hat{\rho}}(\hat{S})] = n - m$. Consequently, $\hat{\rho} = \rho$ and $n \geq 2m - (\mu_{\rho}(S) + \mu_{-\rho}(S))$.

Proof of Theorem 3.1. Let \hat{S} be a minimal symmetric quasi-orthogonal extension of the matrix S with order \hat{n} . By Theorem 2.1, S has a symmetric quasi-orthogonal extension of order $2n - \mu_{\rho}(S) - \mu_{-\rho}(S)$. Then

$$\hat{n} \le 2n - \mu_{\rho}(S) - \mu_{-\rho}(S).$$

It follows that

$$\hat{n} - 2n \le -\mu_{\rho}(S) - \mu_{-\rho}(S) \le -1,$$

and hence

$$\hat{n} - n \le \frac{\hat{n} - 1}{2}.$$

By Lemma 3.3, we have $\hat{n} \ge 2n - \mu_{\rho}(S) - \mu_{-\rho}(S)$. Thus $\hat{n} = 2n - \mu_{\rho}(S) - \mu_{-\rho}(S)$.

Example 3.4. We will use the proof of Theorem 2.1 to find a minimal symmetric quasi-orthogonal extension of the matrix

$$S = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

The spectrum of S is $\{[-1]^2, [2]^1\}$ *. By Theorem 3.1, we have* ind(S) = 2*. Let*

$$P = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & \frac{-2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{pmatrix}$$

The matrix P is orthogonal and

$$P^{\top}SP = \begin{pmatrix} -1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & 2 \end{pmatrix}$$

Following the proof of Theorem 2.1, we have $N = \begin{pmatrix} \frac{-1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ 0 & \frac{-2}{\sqrt{6}} \end{pmatrix}$, $L = \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{pmatrix}$, $D = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$

and $M = N \cdot \sqrt{4I - D^2}$. A minimal symmetric quasi-orthogonal extension of S is then

$$\hat{S} = \begin{pmatrix} 0 & 1 & 1 & -\frac{1}{2}\sqrt{6} & \frac{1}{2}\sqrt{2} \\ 1 & 0 & 1 & \frac{1}{2}\sqrt{6} & \frac{1}{2}\sqrt{2} \\ 1 & 1 & 0 & 0 & -\sqrt{2} \\ -\frac{1}{2}\sqrt{6} & \frac{1}{2}\sqrt{6} & 0 & 1 & 0 \\ \frac{1}{2}\sqrt{2} & \frac{1}{2}\sqrt{2} & -\sqrt{2} & 0 & 1 \end{pmatrix}$$

4 Symmetric Seidel matrices with minimum quasi-orthogonality index

Following [6], a Seidel matrix is a $\{0, \pm 1\}$ -matrix S with zero diagonal and all offdiagonal entries nonzero such that $S = \pm S^{\top}$. Symmetric Seidel matrices were introduced in [11] in connection with equiangular lines in Euclidean spaces [8].

Let *S* be a symmetric Seidel matrix of order *n* with spectrum $\{[\theta_1]^{m_1}, \ldots, [\theta_r]^{m_r}\}$. Then

$$\sum_{i=1}^{r} m_i = n,\tag{5}$$

$$\sum_{i=1}^{r} m_i \theta_i = 0, \tag{6}$$

$$\sum_{i=1}^{r} m_i \theta_i^2 = n(n-1).$$
(7)

Two Seidel matrices S_1 and S_2 are said to be *equivalent* if there exists a signed permutation matrix P such that $S_2 = \pm P S_1 P^{\top}$. Two equivalent Seidel matrices have the same quasi-orthogonality index.

For an integer $n \ge 2$, we denote by ind(n) the minimum quasi-orthogonality index among all $n \times n$ Seidel matrices.

The two Seidel matrices of order 2 are orthogonal and then ind(2) = 0. Up to equivalence, there is only one Seidel matrix of order 3, namely

$$S = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

As seen in Example 3.4, ind(S) = 2, hence ind(3) = 2. Table 1 lists all non equivalent $n \times n$ Seidel matrices having the minimum quasi-orthogonality index for $n \in \{4, 5, 6, 7\}$.

Table 1: Non equivalent $n \times n$ Seidel matrices with minimum quasi-orthogonality index for $n \in \{4, 5, 6, 7\}$.

Order n		Seidel matrix	Characteristic polynomial
4	2	$\begin{pmatrix} 0 & 1 & -1 & 1 \\ 1 & 0 & 1 & 1 \\ -1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$	$(x^2 - 1)(x^2 - 5)$
5	1	$\begin{pmatrix} 0 & 1 & -1 & -1 & 1 \\ 1 & 0 & 1 & -1 & 1 \\ -1 & 1 & 0 & 1 & 1 \\ -1 & -1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{pmatrix}$	$x(x^2-5)^2$
6	0	$\begin{pmatrix} 0 & 1 & -1 & -1 & 1 & 1 \\ 1 & 0 & 1 & -1 & -1 & 1 \\ -1 & 1 & 0 & 1 & -1 & 1 \\ -1 & -1 & 1 & 0 & 1 & 1 \\ 1 & -1 & -1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}$	$(x^2 - 5)^3$
7	1	$ \begin{pmatrix} 0 & 1 & 1 & -1 & -1 & 1 & 1 \\ 1 & 0 & 1 & -1 & 1 & -1 & 1 \\ 1 & 1 & 0 & 1 & -1 & -1 & 1 \\ -1 & -1 & 1 & 0 & 1 & 1 & 1 \\ -1 & 1 & -1 & 1 & 0 & 1 & 1 \\ 1 & -1 & -1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix} $	$(x+2)(x-1)^2(x^2-9)^2$

Symmetric quasi-orthogonal Seidel matrices are known as symmetric conference matrices. The order n of a symmetric conference matrix is of the form 4k + 2 for some integer k, moreover, n - 1 is the sum of two squares [1]. The first orders of symmetric conference matrices are n = 2, 6, 10, 14, 18. However, there is no conference matrix of order 22 (21 is not the sum of two squares). Greaves and Suda [6] prove that the existence of a symmetric conference matrix of order 4k + 2 is equivalent to the existence of a symmetric Seidel matrix of order 4k + i having a prescribed spectrum for each $i \in \{1, 0, -1\}$. More precisely, they obtain the following result.

Theorem 4.1 ([6]). *The existence of the following are equivalent:*

i) a symmetric Seidel matrix with characteristic polynomial

 $(x^2 - 4k - 1)^{2k+1};$

ii) a symmetric Seidel matrix with characteristic polynomial

$$x(x^2 - 4k - 1)^{2k}$$

iii) a symmetric Seidel matrix with characteristic polynomial

$$(x^2 - 1)(x^2 - 4k - 1)^{2k-1};$$

iv) a symmetric Seidel matrix with characteristic polynomial

$$(x-2)(x+1)^{2}(x^{2}-4k-1)^{2(k-1)}.$$

It follows from Theorem 3.1 that the quasi-orthogonality index of the symmetric Seidel matrices mentioned in assertions i, ii, iii, iii and iv of the above theorem are respectively equal to 0, 1, 2 and 3.

Seidel matrices with quasi-orthogonality index 0 are conference matrices. The following result gives the characteristic polynomials of symmetric Seidel matrices with quasi-orthogonality index 1.

Proposition 4.2. Let *S* be an $n \times n$ Seidel matrix with quasi-orthogonality index equals 1. Then *n* is odd and its characteristic polynomial $\phi_S = x(x^2 - n)^{\frac{n-1}{2}}$.

Proof. Let ρ be the spectral radius of S. As $\operatorname{ind}(S) = 1$ then S has a simple eigenvalue α such that $|\alpha| < \rho$. If $\mu_{-\rho} = 0$ then by (6) we have $(n-1)\rho + \alpha = 0$, which is a contradiction because $|\alpha| < \rho$. We obtain a similar contradiction if $\mu_{\rho} = 0$. Now suppose that $\mu_{-\rho} \neq \mu_{\rho}$. By (6) we get $(\mu_{\rho} - \mu_{-\rho})\rho + \alpha = 0$, which implies that $|\mu_{\rho} - \mu_{-\rho}|\rho = |\alpha|$. This is a contradiction. Then $\mu_{-\rho} = \mu_{\rho}$ and hence $\alpha = 0$ and $n = 2\mu_{\rho} + 1$. By (7) we have $\rho^2 = n$. Then the characteristic polynomial $\phi_S(x) = x(x^2 - n)^{\frac{n-1}{2}}$.

For a given $n \ge 4$, the following theorem provides a spectral characterization of $n \times n$ Seidel matrices with minimum quasi-orthogonality index.

Theorem 4.3. Let S be a Seidel matrix of order $n \ge 4$. Then, we have the following statements

- i) If n = 4k + 2, then $ind(S) \ge 0$. Equality holds if and only if $\phi_S(x) = (x^2 4k 1)^{2k+1}$.
- ii) If n = 4k + 1, then $ind(S) \ge 1$. Equality holds if and only if $\phi_S(x) = x(x^2 4k 1)^{2k}$.
- *iii)* If n = 4k, then $ind(S) \ge 2$. Equality holds if and only if $\phi_S(x) = (x^2 1)(x^2 4k 1)^{2k-1}$.
- iv) If n = 4k-1, then $\operatorname{ind}(S) \ge 3$. Equality holds if and only if $\phi_S(x) = (x-2)(x+1)^2(x^2-4k-1)^{2(k-1)}$ or $\phi_S(x) = (x+2)(x-1)^2(x^2-4k-1)^{2(k-1)}$.

To prove this theorem, we need the following results.

Lemma 4.4. Let $P(x) = (x - a_1)^{\alpha_1} (x - a_2)^{\alpha_2} \dots (x - a_r)^{\alpha_r}$ be an integral polynomial where a_1, \dots, a_r are distinct real numbers and $\alpha_1 \ge \alpha_2 \ge \dots \ge \alpha_r$. If $\alpha_1 = \alpha_2 = \dots = \alpha_k > \alpha_{k+1}$ then $(x - a_1)(x - a_2) \dots (x - a_k)$ is an integral polynomial.

Proof. The result follows from the fact that if a_i and a_j share the same minimal polynomial then $\alpha_i = \alpha_j$.

The next proposition gives a useful property about the determinant of symmetric Seidel matrices.

Proposition 4.5. [7, Corollary 3.6] Let S be a symmetric Seidel matrix of order n. Then $det(S) \equiv 1 - n \pmod{4}$.

Proof of Theorem 4.3. For $n \in \{4, 5, 6, 7\}$, the proof follows from Table 1. Now, we assume that $n \ge 8$.

- i) If ind(S) = 0, then S is quasi-orthogonal and hence $\phi_S(x) = (x^2 4k 1)^{2k+1}$.
- ii) If ind(S) = 0, then S is quasi-orthogonal which implies that n is even, a contradiction. Hence $ind(S) \ge 1$. If ind(S) = 1, then by Proposition 4.2, we have $\phi_S(x) = x(x^2 4k 1)^{2k}$.
- iii) If ind(S) = 0, then S is quasi-orthogonal. Hence S is a symmetric conference matrix. This is a contradiction because $n \not\equiv 2 \pmod{4}$. Consequently $ind(S) \ge 1$.

If ind(S) = 1, then by Proposition 4.2, n is odd which is a contradiction. Then ind(S) > 1.

If $\operatorname{ind}(S) = 2$ then $\phi_S(x) = (x - \alpha)(x - \beta)(x - \rho)^{\mu_{\rho}}(x + \rho)^{\mu_{-\rho}}$ where $|\alpha| < \rho$ and $|\beta| < \rho$. By (6) we have $\alpha + \beta + (\mu_{\rho} - \mu_{-\rho})\rho = 0$. It follows that $|\mu_{\rho} - \mu_{-\rho}| < 2$. Moreover, $\mu_{\rho} + \mu_{-\rho} = 4k - 2$ then $\mu_{-\rho}$ and μ_{ρ} have the same parity. Hence $\mu_{\rho} = \mu_{-\rho}$ and $\beta = -\alpha$. By (7), we get

$$\alpha^2 + (2k-1)\rho^2 = 2k(4k-1) \tag{8}$$

Then

$$(2k-1)\rho^2 \le 2k(4k-1).$$

Hence

$$\rho^2 \le 4k + 1 + \frac{1}{2k - 1}.$$

By Lemma 4.4, we have α^2 and ρ^2 are integers. As $n \ge 8$, we get $k \ge 2$ and hence

$$\rho^2 \le 4k + 1. \tag{9}$$

Since $|\alpha| < \rho$, equality (8) implies that

$$2k\rho^2 > 2k(4k-1).$$

Then

$$\rho^2 > 4k - 1$$

We cannot have $\rho^2 = 4k$, otherwise $\alpha^2 = 2k$ and then $det(S) = 2k(4k)^{2k-1}$ which contradicts Proposition 4.5. Therefore, $\rho^2 = 4k + 1$ and consequently $\alpha^2 = 1$. Hence, $\phi_S(x) = (x^2 - 1)(x^2 - 4k - 1)^{2k-1}$. iv) As *n* is odd, *S* cannot be quasi-orthogonal. Therefore, $ind(S) \ge 1$.

If $\operatorname{ind}(S) = 1$, then by Proposition 4.2, $\det(S) = 0$. However, Proposition 4.5 implies that $\det(S) \equiv 2 \pmod{4}$. This is a contradiction. Consequently, $\operatorname{ind}(S) \ge 2$. If $\operatorname{ind}(S) = 2$ then $\phi_S(x) = (x - \alpha)(x - \beta)(x - \rho)^{\mu_{\rho}}(x + \rho)^{\mu_{-\rho}}$ where $|\alpha| < \rho$ and $|\beta| < \rho$. By (7), we have

$$\alpha^2 + \beta^2 + (4k - 3)\rho^2 = (4k - 1)(4k - 2).$$
(10)

Then

$$(4k-1)\rho^2 > (4k-1)(4k-2)$$

Hence

$$\rho^2 > 4k - 2.$$

Moreover

$$(4k-3)\rho^2 \le (4k-1)(4k-2).$$

Thus

$$\rho^2 \le 4k + \frac{2}{4k - 3}.$$

Moreover, $\mu_{-\rho}$ and μ_{ρ} have different parity because $\mu_{\rho} + \mu_{-\rho} = 4k - 3$. It follows from Lemma 4.4 that ρ is an integer. Since $n \ge 8$, we have $k \ge 3$ and hence

$$4k - 1 \le \rho^2 \le 4k. \tag{11}$$

By (6) we have $\alpha + \beta + (\mu_{\rho} - \mu_{-\rho})\rho = 0$. Thus $|\mu_{\rho} - \mu_{-\rho}| < 2$. Hence $|\mu_{\rho} - \mu_{-\rho}| = 1$ and $\alpha + \beta = \pm \rho$. Then $(\alpha + \beta)^2 = \rho^2$. Using (10), we have $\alpha\beta = (2k - 1)(\rho^2 - 4k + 1)$. Consequently $\alpha\beta$ is an integer. From Proposition 4.5, we have $\det(S) = \pm \alpha\beta\rho^{4k-3} \equiv 2 \pmod{4}$. Then ρ must be odd, otherwise $\rho^2 \equiv 0 \pmod{4}$ and $\det(S) \equiv 0 \pmod{4}$. By (11) we get $\rho^2 = 4k - 1$. Then $\alpha\beta = 0$ and $\det(S) = 0$, a contradiction. Hence $\operatorname{ind}(S) \geq 3$.

If $\operatorname{ind}(S) = 3$, then $\phi_S(x) = (x - \alpha)(x - \beta)(x - \gamma)(x - \rho)^{\mu_{\rho}}(x + \rho)^{\mu_{-\rho}}$ where $|\alpha| < \rho$, $|\beta| < \rho$ and $|\gamma| < \rho$. By (7), we get

$$\alpha^{2} + \beta^{2} + \gamma^{2} + (4k - 4)\rho^{2} = (4k - 1)(4k - 2).$$
(12)

This implies

$$4k - 2 < \rho^2 \le \frac{(4k - 1)(4k - 2)}{4k - 4}.$$

Then

$$4k - 1 \le \rho^2 \le 4k + 1 + \frac{6}{4k - 4}.$$
(13)

By (6) we have $\alpha + \beta + \gamma + (\mu_{\rho} - \mu_{-\rho})\rho = 0$. It follows that $|\mu_{\rho} - \mu_{-\rho}| < 3$. In addition, $\mu_{\rho} + \mu_{-\rho} = 4k - 4$ then $\mu_{-\rho}$ and μ_{ρ} have the same parity. Hence $|\mu_{\rho} - \mu_{-\rho}| \in \{0, 2\}$. We will prove that $\mu_{\rho} = \mu_{-\rho}$. Suppose, for contradiction, that $|\mu_{\rho} - \mu_{-\rho}| = 2$. By Lemma 4.4, ρ is an integer. Moreover, $\phi_S(x)$ is an integral polynomial. Then $(x - \alpha)(x - \beta)(x - \gamma)$ is an integral polynomial. Hence, $\alpha\beta\gamma$ is an integer. Since $n \ge 8$, we have $k \ge 3$. By (13), we get $\rho^2 \in \{4k - 1, 4k, 4k + 1\}$. From Proposition 4.5, det $(S) = \pm \alpha \beta \gamma \rho^{4k-4} \equiv 2 \pmod{4}$. Thus ρ is odd. Hence $\rho^2 = 4k + 1$. By (12), we get

$$\alpha^2 + \beta^2 + \gamma^2 = 6. \tag{14}$$

By Cauchy inequality applied to $\begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ we get $|\alpha + \beta + \gamma| \le 3\sqrt{2}$. But

from (6), we have $|\alpha + \beta + \gamma| = 2\rho = 2\sqrt{4k+1}$. This is a contradiction because $k \ge 3$. Hence $\mu_{-\rho} = \mu_{\rho}$.

By 6, we have

$$\alpha + \beta + \gamma = 0. \tag{15}$$

By Lemma 4.4, ρ^2 is an integer. Moreover, $\phi_S(x)$ is an integral polynomial. Then $(x - \alpha)(x - \beta)(x - \gamma)$ is an integral polynomial. Hence, $\alpha\beta\gamma$ is an integer. From Proposition 4.5, $\det(S) = \alpha\beta\gamma\rho^{4k-4} \equiv 2 \pmod{4}$. Thus ρ^2 is odd. Then, by (13), we have $\rho^2 \in \{4k - 1, 4k + 1\}$.

We will prove that $\rho^2 = 4k + 1$. Suppose, for contradiction, that $\rho^2 = 4k - 1$. Let $S := [s_{ij}]$ and $B := (4k - 1)I - S^2 = [b_{ij}]$. The matrix B is symmetric and its eigenvalues are $(4k - 1) - \alpha^2$, $(4k - 1) - \beta^2$, $(4k - 1) - \gamma^2$ and 0. The first three eigenvalues are positive and the multiplicity of 0 is 4k - 4. Thus B is a positive semi-definite matrix with rank equal to 3. Moreover, for $i \in \{1, 2, ..., n\}$ we have

$$b_{ii} = 4k - 1 - (s_{i1}^2 + \ldots + s_{in}^2) = 1.$$

Since *B* is positive semi-definite, for $i \neq j \in \{1, 2, ..., n\}$ we have

$$\det \begin{pmatrix} 1 & b_{ij} \\ b_{ij} & 1 \end{pmatrix} \ge 0$$

Then, $b_{ij} \in \{0, 1, -1\}$. We also have

$$b_{ij} = -\sum_{k=1}^{n} s_{ik} s_{kj} = -\sum_{k \neq i, k \neq j} s_{ik} s_{kj}.$$

As the off-diagonal entries of *S* are ± 1 , $b_{ij} \equiv 4k - 3 \equiv 1 \pmod{2}$ and hence $b_{ij} \in \{1, -1\}$. It is easy to check that the determinant of a 3×3 symmetric matrix with 1's on the diagonal and ± 1 off-diagonal is equal to 0 or -4. Moreover, the matrix *B* is positive semi-definite, then its principal minors of order 3 must equal to 0. Then, the rank of *B* is at most 2, which leads to a contradiction. Hence $\rho^2 = 4k + 1$. By (12), we get

$$\alpha^2 + \beta^2 + \gamma^2 = 6. \tag{16}$$

Moreover, we have $\alpha^2 \beta^2 \gamma^2 \leq (\frac{\alpha^2 + \beta^2 + \gamma^2}{3})^3$. Then $|\alpha\beta\gamma| \leq 2$. Since $\alpha\beta\gamma$ is an integer and $\det(S) = \alpha\beta\gamma\rho^{4k-4} \equiv 2 \pmod{4}$, then $\alpha\beta\gamma \in \{-2, 2\}$.

case 1 : $\alpha\beta\gamma = 2$, together with the above equations (16) and (15), we find that $\{\alpha, \beta, \gamma\} = \{[2]^1, [-1]^2\}$. Hence, $\phi_S(x) = (x-2)(x+1)^2(x^2-4k-1)^{2(k-1)}$.

case 2 :
$$\alpha\beta\gamma = -2$$
, we get $\{\alpha, \beta, \gamma\} = \{[-2]^1, [1]^2\}$. Hence, $\phi_S(x) = (x+2)(x-1)^2(x^2-4k-1)^{2(k-1)}$.

The converse of the four assertions follows from Theorem 3.1.

Analyzing the proof of Theorem 4.1, one can see that the matrices in Theorem 4.3 are principal sub-matrices of a conference matrix.

References

- [1] V. Belevitch. Theorem of 2*n*-terminal networks with application to conference telephony. *Electrical Communication*, 27:231–244, 1950.
- [2] Béla Bollobás and Andrew Thomason. Graphs which contain all small graphs. *European Journal of Combinatorics*, 2(1):13–15, 1981.
- [3] Abderrahim Boussaïri, Brahim Chergui, Zaineb Sarir, and Mohamed Zouagui. Quasi-orthogonal extension of skew-symmetric matrices. *arXiv preprint arXiv:2410.19594*, 2024.
- [4] Colbourn Ch and J Dinitz. *Handbook of Combinatorial Designs, (Discrete mathematics and its applications, ser. ed. K. Rosen).* CRC Press, Boca Raton, FL, 2007.
- [5] Ronald L Graham and Joel H Spencer. A constructive solution to a tournament problem. *Canadian Mathematical Bulletin*, 14(1):45–48, 1971.
- [6] G. Greaves and S. Suda. Symmetric and skew-symmetric $\{0, \pm 1\}$ -matrices with large determinants. *Journal of Combinatorial Designs*, 25(11):507–522, 2017.
- [7] Gary Greaves, Jacobus H Koolen, Akihiro Munemasa, and Ferenc Szöllősi. Equiangular lines in euclidean spaces. *Journal of Combinatorial Theory, Series A*, 138:208–235, 2016.
- [8] Johannes Haantjes. Equilateral point-sets in elliptic two-and three-dimensional spaces. *Nieuw Arch. Wiskunde* (2), 22:355–362, 1948.
- [9] Christos Koukouvinos and Jennifer Seberry. New weighing matrices and orthogonal designs constructed using two sequences with zero autocorrelation function–a review. *Journal of statistical planning and inference*, 81(1):153–182, 1999.
- [10] Olga Taussky. sums of squares and hadamard matrices. *Combinatorics (Proc. Sympos. Pure Math., Vol. XIX, Univ. California, Los Angeles, Calif., 1968)*, pages 229–233, 1971.
- [11] Jacobus H van Lint and Johan J Seidel. Equilateral point sets in elliptic geometry. In *Geometry and Combinatorics*, pages 3–16. Elsevier, 1991.