ON DIFFEOMORPHISMS OF 4–DIMENSIONAL 1–HANDLEBODIES

DELPHINE MOUSSARD

ABSTRACT. We give a new proof of Laudenbach and Poénaru's theorem, which states that any diffeomorphism of the boundary of a 4–dimensional 1–handlebody extends to the whole handlebody. Our proof is based on the cassification of Heegaard splittings of double handlebodies and a result of Cerf on diffeomorphisms of the 3–ball. Further, we extend this theorem to 4–dimensional compression bodies, namely cobordisms between 3–manifolds constructed using only 1–handles: when the negative boundary is a product of a compact surface by interval, we show that every diffeomorphism of the positive boundary extends to the whole compression body. This invlolves a strong Haken theorem for sutured Heegaard splittings and a classification of sutured Heegaard splittings of double compression bodies. Finally, we show how this applies to the study of relative trisection diagrams for compact 4–manifolds.

1. INTRODUCTION

A famous theorem of Laudenbach and Poénaru asserts that every diffeomorphism of the boundary of a 4–dimensional 1–handlebody extends to a diffeomorphism of the whole handlebody [\[LP72\]](#page-9-0). This result is of great importance in the theory of smooth 4–manifolds because it implies that, given a handle decomposition of a closed 4–manifold X, the attaching information for the $1-$ and $2-$ handles contained in a Kirby diagram is sufficient to determine X up to diffeomorphism. Likewise, in the theory of trisections, it implies that a trisection diagram determines a unique closed 4–manifold up to isotopy. We give here an alternative proof of Laudenbach–Poénaru's theorem, based on two main ingredients. The first one is the uniqueness of the minimal genus Heegaard splitting of the boundary of a 4–dimensional 1–handlebody, that is a double handlebody, due to Carvalho and Oertel [\[CO05\]](#page-9-1). The second ingredient is the fact that every diffeomorphism of a 3–dimensional handlebody, that restricts to the identity on the boundary, is isotopic to the identity; this is based on a result of Cerf [\[Cer68\]](#page-9-2). From that point, our proof of Laudenbach–Poénaru's theorem is very short.

Carvalho and Oertel actually used much of the same machinery that Laudenbach and Poénaru used in their original proof. In particular, both papers relied on Laudenbach's results from [\[Lau73\]](#page-9-3). So at first glance this new proof might be regarded as a repackaging of Laudenbach and Poénaru's original proof. However, in [\[HS21\]](#page-9-4), Hensel and Schultens reprove Carvalho–Oertel's result using brief cut and paste arguments. Thus, we believe that this proof of Laudenbach–Poénaru's theorem represents a true simplification of the original.

We then extend the setting and consider 4–dimensional compression bodies. Such a compression body is a cobordism between two 3–manifolds, its negative boundary and its positive boundary, constructed using only 1–handles. We further require that the negative boundary is a product of a compact surface and an interval. These compression bodies are the building blocks of the so-called relative trisections of compact 4–manifolds with boundary. We generalize Laudenbach–Poénaru's theorem to compression bodies.

Theorem (Theorem [4.4\)](#page-7-0). Let V be a 4-dimensional compression body. Assume the negative boundary of V is a product $P \times I$, where P is a compact oriented surface which contains no 2–sphere. Then every diffeomorphism of the positive boundary of V extends to a diffeomorphism of V .

From this result, we recover the statement, due to Castro, Gay and Pinzón-Caicedo [\[CGPC18\]](#page-9-5), that every relative trisection diagram determines a unique compact 4–manifold up to diffeomorphism, and we extend it to the case when the page of the trisection (the surface P) is allowed to contain closed components. This fails when P is allowed to contain 2–spheres, as we show with an example which was communicated to us by David Gay.

The proof of Theorem [4.4](#page-7-0) follows the lines of our proof of Laudenbach–Poénaru's theorem. The second ingredient is easily adapted to the relative case: given a 3-dimensional compression body C , namely a cobordism between two compact surfaces constructed using only 1–handles, we show that every diffeomorphism of C, which restricts to the identity on its positive boundary, is isotopic to the identity. The first ingredient requires more work. The boundary of a 4–dimensional compression body is diffeomorphic to the connected sum of some products $F \times I$, where F is a compact surface, and some copies of $S^1 \times S^2$. We need to understand the sutured Heegaard splittings of these so-called double compression bodies.

Theorem (Theorem [3.17\)](#page-6-0). Any two sutured Heegaard splittings of a double compression body with the same genus are isotopic.

This will essentially follow from the strong Haken theorem due to Scharlemann [\[Sch24\]](#page-9-6): every 2–sphere embedded in a 3–manifold equipped with a Heegaard splitting is isotopic to a 2–sphere which intersects the Heegaard surface along a single circle. Another proof of this result was given by Hensel and Schultens in [\[HS21\]](#page-9-4), which appears surprisingly simple to us. We apply their technique to check that the strong Haken theorem remains true in the setting of sutured Heegaard splittings.

To prove our relative Laudenbach–Poénaru's theorem, we only need the uniqueness of the minimal genus Heegaard splittings of double compression bodies. However, the general classification result is also useful in the theory of relative trisections. In the original definition of a relative trisection, one has to precisely describe a decomposition of the boundary of the building blocks, namely the 4–dimensional compression bodies, in order to prescribe the way the different pieces should meet. We give a simpler definition of a relative trisection, and Theorem [3.17](#page-6-0) shows that the two definitions are equivalent.

Plan of the paper. In Section [2,](#page-1-0) we reprove Laudenbach–Poénaru's theorem. In Section [3,](#page-2-0) we define sutured Heegaard splittings, we give a proof of the strong Haken theorem in this setting, and we classify the sutured Heegaard splittings of double compression bodies. In Section [4,](#page-6-1) we prove Theorem [4.4,](#page-7-0) we apply it to the study of relative trisection diagrams, and we discuss its failure when there is a 2–sphere in the page P.

Conventions. The boundary of an oriented manifold with boundary is oriented using the outward normal first convention. If M is an oriented manifold, $-M$ represents M with the opposite orientation. If M is a compact manifold and N is a submanifold, we denote by $M \setminus N$ the manifold "M cut along N", which comes with a surjective map $\pi : M \setminus N \to M$ such that π is a diffeomorphism from $\pi^{-1}(M \setminus N)$ to $M \setminus N$ and a double cover from $\pi^{-1}(N)$ to N.

Acknowledgements. I warmly thank Trenton Schirmer for many helpful conversations and for valuable comments on the first version of the paper. I am also grateful to David Gay for helpful conversations.

2. PROOF OF LAUDENBACH-POÉNARU'S THEOREM VIA HEEGAARD SPLITTINGS

A genus–g handlebody is a 3–manifold diffeomorphic to a 3–ball with g 1–handles glued on its boundary. A Heegaard splitting of a closed 3–manifold M is a decomposition $M = H_1 \cup_{\Sigma} H_2$, where H_1 and H_2 are handlebodies and $\Sigma = \partial H_1 = \partial H_2$. A genus–g double handlebody is a 3-manifold diffeomorphic to $\sharp_{i=1}^g (S^1 \times S^2)$.

The first preliminary result we need to reprove Laudenbach–Poénaru's theorem is the classification of minimal genus Heegaard splittings of double handlebodies. This result is recovered in Theorem [3.17](#page-6-0) within the more general setting of double compression bodies.

Theorem 2.1 (Carvalho–Oertel). A genus–g double handlebody admits a unique genus–g Heegaard splitting, up to isotopy.

The second preliminary result is based on the following theorem of Cerf [\[Cer68\]](#page-9-2).

Theorem 2.2 (Cerf). Every diffeomorphism of a 3–ball, which is the identity on the boundary, is isotopic to the identity.

We shall generalize this fact to handlebodies of positive genus. A defining disk system for a 3dimensional handlebody H is a union D of properly embedded disks such that $H \setminus \mathcal{D}$ is a 3-ball.

Lemma 2.3. Let H be a 3-dimensional handlebody. Every two defining disk systems for H which coincide on the boundary are isotopic.

Proof. This follows from a standard innermost disk argument, using the fact that handlebodies are irreducible.

Lemma 2.4. Let H be a 3-dimensional handlebody. Let φ be a diffeomorphism of H. If φ is the identity on ∂H , then φ is isotopic to the identity.

Proof. Pick a defining disk system $\mathcal D$ for H. Then $\varphi(\mathcal D)$ is another defining disk system with the same boundary, thus isotopic to D . By the isotopy extension theorem, there is an ambient isotopy sending $\varphi(\mathcal{D})$ to D, keeping ∂H fixed. Hence, up to isotoping φ , we can assume that $\varphi(\mathcal{D}) = \mathcal{D}$. Now, every diffeomorphism of a 2–disk, which is the identity on the boundary, is isotopic to the identity. Hence we can even assume that φ is the identity on $\partial H \cup \mathcal{D}$. We are led to a diffeomorphism of a 3-ball which is the identity on the boundary, and we apply Cerf's result.

A 4–dimensional 1–handlebody is a compact oriented smooth 4–manifold obtained from a 4–ball by adding a finite number of 1–handles. The number of 1–handles glued is the genus of the handlebody. Note that the boundary of a 4-dimensional 1-handlebody of genus q is a genus–q double handlebody.

Theorem 2.5 (Laudenbach–Poénaru). Let Z be a 4-dimensional 1-handlebody. Then every diffeomorphism of ∂Z extends to a diffeomorphism of Z.

Proof. Fix an identification of Z with a product $H \times I$, with H a 3-dimensional genus–q handlebody, where the vertical boundary $\partial H \times I$ has been collapsed along the I–factor. This induces a foliation of Z by 3–dimensional handlebodies H_t , $t \in [0, 1]$, which meet exactly along their common boundary $\Sigma = \partial H_t$. This surface Σ defines a minimal genus Heegaard splitting of ∂Z .

Now take a diffeomorphism φ of ∂Z . It sends the Heegaard splitting $\partial Z = H_0 \cup_{\Sigma} H_1$ onto another splitting $\partial Z = \varphi(H_0) \cup_{\varphi(\Sigma)} \varphi(H_1)$, with the same genus. By Theorem [2.1,](#page-1-1) they are isotopic. Hence, realizing the isotopy in a collar neighborhood of ∂Z , we can assume that $\varphi(\Sigma) = \Sigma$ and $\varphi(H_t) = H_t$ for each $t = 0, 1$.

Now $\varphi_{|H_0}$ and $\varphi_{|H_1}$ define two diffeomorphisms of H which coincide on the boundary. By Lemma [2.4,](#page-2-1) there is an isotopy φ_t of diffeomorphisms of H from $\varphi_{|H_0}$ to $\varphi_{|H_1}$. The map $\phi: Z \to Z$ induced by the diffeomorphism $(x, t) \mapsto (\varphi_t(x), t)$ of $H \times I$ is the desired diffeomorphism.

3. Sutured Heegaard splittings

3.1. Compression bodies.

Definition 3.1. A compression body C is a cobordism from a connected compact oriented surface $\partial_{+}C$ to a compact oriented surface ∂−C which is constructed using only 2–handles and 3–handles, where enough 3–handles are glued to avoid any S^2 –component in $\partial_{-}C$. A lensed compression body is then obtained by collapsing the vertical boundary of the cobordism so that the boundary of $\partial_{+}C$ becomes identified with the boundary of $\partial_{-}C$.

Note that the definition includes the possibility that ∂ −C be empty. Note also that a compression body can alternatively be constructed by adding 1–handles to a thickening of the negative boundary, a compact oriented surface containing no 2–sphere, or to a 3–ball. In what follows, compression bodies are supposed to be lensed.

Definition 3.2. Let C be a compression body. A defining disk system for C is a collection D of disks properly embedded in C such that $C \setminus \mathcal{D}$ is a thickening of $\partial_{-}C$, or a 3-ball if $\partial_{-}C$ is empty. The boundary $\partial \mathcal{D} \subset \partial_{+}C$ is a *cut-system* for C.

Note that defining disk systems do exist: take for instance the core disks of the 2–handles in the definition (with a minimal number of 2– and 3–handles).

Lemma 3.3. Every compression body is irreducible.

Proof. Let S be a 2–sphere embedded in a compression body C. Let $\mathcal D$ be a defining collection of disks for C . By a standard innermost disk argument, we see that we can isotope S so that it gets disjoint from D. Then S lies in a product $P \times I$ where P is a component of $\partial_{-}C$, that is a compact connected oriented surface different from S^2 (or a disk if ∂ - $C = \emptyset$). We embed this product in \mathbb{R}^3 , which is irreducible. Now S bounds a 3-ball B in \mathbb{R}^3 . If ∂P is non-empty, then $\partial(P \times I)$ is connected, so that it is contained in $\mathbb{R}^3 \setminus B$, and $B \subset (P \times I)$. Now assume P is closed. If S separate the two boundary components of $P \times I$, then the retraction of $P \times I$ onto $P \times \{0\}$ provides a map $f : S \to P$ which induces an isomorphism $f_* : H_2(S) \to H_2(P)$. But f lifts to a map $\tilde{f} : S \to \tilde{P}$, where \tilde{P} is the universal cover of P, which is non-compact since $q(P) > 0$. It follows that f_* factors through $H_2(\tilde{P}) = 0$. We get a contradiction and conclude that S does not separate the boundary of $P \times I$, so that $B \subset (P \times I)$. \Box

Given a compact oriented 3–manifold M, a sutured decomposition of ∂M is a decomposition ∂M = $\partial_0 M \cup_s \partial_1 M$, where $\partial_0 M$ and $\partial_1 M$ are two compact surfaces oriented like M, and s is their common boundary oriented as $\partial(\partial_1M)$. The curve s is called the *suture*. Note that, if ∂_0M and ∂_1M have no closed component, the datum of the oriented suture fully determines the sutured decomposition. A sutured 3–manifold is a manifold whose boundary is equipped with a sutured decomposition.

Definition 3.4. Let M be a sutured 3–manifold. A *sutured Heegaard splitting* of M is a decomposition $M = C_1 \cup_{\Sigma} C_2$ where C_1 and C_2 are compression bodies, $\Sigma = \partial_+ C_1 = -\partial_+ C_2$ is a compact connected oriented surface, $\partial_{-}C_1 = \partial_0 M$ and $\partial_{-}C_2 = \partial_1 M$.

Every sutured 3–manifold admits a sutured Heegaard splitting, unique up to stabilization (see Juhasz [\[Juh06,](#page-9-7) Section 2] or Dissler [\[Dis23,](#page-9-8) Section 3]).

3.2. Strong Haken's theorem for sutured Heegaard splittings.

Definition 3.5. An essential embedded 2–sphere in a 3–manifold with a Heegaard splitting is Haken if it intersects the Heegaard surface transversely along a connected simple closed curve.

Definition 3.6. An embedded 2–sphere S in a compact 3–manifold M is surviving if S is essential in the manifold obtained from M by filling every S^2 -component in ∂M with a 3-ball.

The following theorem is due to Haken in the closed case [\[Hak68\]](#page-9-9). The proof we give here is taken from Jaco [\[Jac80\]](#page-9-10). We reproduce it in order to take care of the small additionnal argument needed in the case of sutured Heegaard splittings.

Theorem 3.7. Let $M = C_1 \cup_{\Sigma} C_2$ be a sutured Heegaard splitting. If there is an essential (resp surviving) embedded 2-sphere $S \subset M$, then there is an essential (resp surviving) embedded 2-sphere $S' \subset M$ which is Haken.

This result will follow from two lemmas. A *spanning annulus* in a compression body C is a properly embedded annulus with an end on each of $\partial_{\pm}C$.

Lemma 3.8. Let C be a compression body. Let S be a planar compact surface with no closed component. Assume $(S, \partial S)$ is embedded in $(C, \partial_{+}C)$. If S is incompressible and boundary-incompressible, then S is a union of disks.

Proof. We construct a family $\mathcal F$ of disks and annuli properly embedded in C , as follows. First take a defining disk system for C. Then, for each component F of $\partial_{-}C$:

- if F is closed, choose a collection of simple closed curves $(\alpha_i, \beta_i)_{i\in I}$ on F which are pairwise disjoint, except that for all i, α_i intersects β_i transversely once, and which cut F into a disk; take spanning annuli $A_i = \alpha_i \times I$ and $B_i = \beta_i \times I$,
- if F has a non-empty boundary, choose a collection of properly embedded arcs $(\gamma_i)_{i\in J}$ which cut F into a disk; take spanning annuli $C_i = \gamma_i \times I$.

We assume that all these disks and annuli are pairwise disjoint, except each pair (A_i, B_i) which intersects along an arc. Note that cutting C along all the disks and annuli in $\mathcal F$ gives a 3-ball.

Suppose S meets $F \in \mathcal{F}$ along a simple closed curve γ that bounds a disk in F. Since S is incompressible, γ also bounds a disk in S. It then follows from the irreducibility of C that F can be isotoped in order to remove this intersection. By a standard innermost disk argument, we see that all such intersections can be removed while keeping the disjointness properties of \mathcal{F} .

Similarly, by ∂ –incompressibility of S and irreducibility of C, we can assume that S never meets an $F \in \mathcal{F}$ along a properly embedded arc which is non-essential in F.

The last possibility is that S intersects an annulus A_i (or equivalently B_i) along an essential curve ξ . It implies that S also meets the annulus B_i along an essential curve ζ (we have removed yet non-essential intersection curves). But then a tubular neighborhood of $\xi \cup \zeta$ in S is a punctured torus. This is impossible since S is a genus–0 surface.

Finally, we can assume that S is disjoint from all F. Cutting C along all disks and annuli in F gives a 3–ball B. Take a boundary component σ of S which innermost in ∂B . The curve σ bounds a disk in B, so, since S is incompressible, this component of S is a disk. Iterating the argument, we see that S is a disjoint union of disks.

Lemma 3.9 (Johnson [\[Joh,](#page-9-11) Lemma 5.24]). Let S be a connected planar compact surface which is not a disk. Let $\alpha_1, \ldots, \alpha_n$ be disjoint properly embedded arcs in S such that:

- $S \setminus \cup_{i=1}^n \alpha_i$ is a union of disks,
- for all i, α_i is essential in $S \setminus \bigcup_{j=i+1}^n \alpha_j$.

Then $S \setminus \cup_{i=1}^n \alpha_i$ has strictly fewer components than ∂S .

Proof of Theorem [3.7.](#page-3-0) Start with $S_1 = S \cap C_1$. Perform on S_1 as many compressions as possible. At each compression, S is surgered into two spheres, at least one of which is essential (resp surviving); keep that one. Note that the number of components of S_1 cannot increase during this process. Then perform on S_1 as many ∂ –compressions as possible. Thanks to Lemma [3.8,](#page-3-1) S_1 is now a union of disks. By Lemma [3.9,](#page-4-0) the number of components of S₁ has strictly decreased. Then apply the same process to $S_2 = S \cap C_2$, and iterate it until $S \cap \Sigma$ is connected.

We now prove a strong Haken's theorem for sutured Heegaard splittings, following Hensel and Schul-tens [\[HS21\]](#page-9-4). We need a condition on the splitting. We say that a sutured Heegaard splitting $M = C_1 \cup_{\Sigma} C_2$ is *admissible* if every 2–sphere in ∂M meets Σ along a connected curve.

Theorem 3.10. Let $M = C_1 \cup_{\Sigma} C_2$ be an admissible sutured Heegaard splitting. Every essential embedded 2–sphere in M is isotopic to a Haken sphere.

Lemma 3.11. In a sutured Heegaard splitting, every non-surviving sphere is isotopic to a Haken sphere.

Proof. The proof of [\[HS21,](#page-9-4) Lemma 3.7] applies verbatim. \square

The proof of Theorem [3.10](#page-4-1) involves the graph of surviving spheres $\mathcal{S}(M)$, defined as follows.

- The vertices are the isotopy classes of surviving spheres in M .
- There is an edge between two vertices if the corresponding isotopy classes can be realized by disjoint spheres.

Lemma 3.12. The graph $S(M)$ is connected.

Proof. Assume there are non-isotopic surviving spheres S and S' in M . Assume they minimize their number of intersection curves within their isotopy classes. If they do intersect, choose an intersection curve c which is innermost in S, thus bounds a disk $\delta \subset S$ whose interior is disjoint from S'. Then surger S' along δ: this provides two embedded spheres in M, having fewer intersection curves with S, at least one of which is surviving. Iterating, we get a path between the isotopy classes of S and S' in $\mathcal{S}(M)$. \Box

Proof of Theorem [3.10.](#page-4-1) We proceed by induction on $(q, n) = (q(\Sigma), |\partial \Sigma|)$ with lexicographic order.

If $g = 0$, then M is a punctured S^3 , hence it contains no surviving sphere and Lemma [3.11](#page-4-2) concludes. Now fix $(q, n) \succ (0, 3)$.

We first prove the following claim: if $S \subset M$ is a surviving Haken sphere and $T \subset M$ is a surviving sphere disjoint from S, then T is isotopic to a Haken sphere. Indeed, let N be the component of $M \setminus S$ (over 1 or 2 components) which contains T. The Heegaard splitting induced on N by that of M has either a strictly lower genus, or a Heegaard surface with strictly less boundary components. Hence we can apply the inductive hypotesis.

Now, the result is given by Lemma [3.11](#page-4-2) if M contains no surviving sphere. Otherwise, M contains a Haken sphere S_0 by Theorem [3.7.](#page-3-0) If S is an essential 2–sphere in M non-isotopic to S_0 , either it is non-surviving and we apply Lemma [3.11,](#page-4-2) or it is surviving and there is a path of surviving spheres from S_0 to S in which successive spheres are disjoint, then we apply the above claim.

3.3. Sutured Heegaard splittings of double compression bodies. We are interested in understanding the Heegaard splittings of double compression bodies, namely compact 3–manifolds obtained by gluing two copies of a given compression body along their positive boundaries via the identity map. Such a double compression body can be written as $(\sharp(\overrightarrow{P} \times I)) \sharp (\sharp^k(S^1 \times S^2))$, where P is a compact surface, connected or not, possibly empty, and $\sharp(P \times I)$ is the connected sum of all the components of $(P \times I)$. We define a sutured decomposition of ∂M as follows: $P \times \{0\} \subset \partial_0 M$, $P \times \{1\} \subset \partial_1 M$, and the suture is $s = \partial P \times {\frac{1}{2}}$. Thanks to Theorem [3.10,](#page-4-1) we essentially need to understand the splittings of products $P \times I$ with P connected, possibly with punctures.

Proposition 3.13. Let P be a compact connected oriented surface of genus $g \geq 0$. Let B_1, \ldots, B_k be disjoint closed 3-balls embedded in the interior of $P \times I$, each of which intersects $P \times \{\frac{1}{2}\}\$ transversely along a 2-disk, where $k \geq 0$. Set $\mathcal{B} = \bigcup_{i=1}^{k} B_i$, $M = (P \times I) \setminus \text{Int}(\mathcal{B})$, and $\Sigma_0 = M \cap (P \times {\frac{1}{2}})$. Define a sutured decomposition of ∂M with the suture $s = \partial \Sigma_0$, $P \times \{0\} \subset \partial_0 M$, and $P \times \{1\} \subset \partial_1 M$. Then the Heegaard splitting of M defined by the Heegaard surface Σ_0 is the unique genus–g Heegaard splitting of M up to isotopy. It is the minimal genus Heegaard splitting of M.

Proof. Let $M = C_1 \cup_{\Sigma} C_2$ be a Heegaard splitting of M. Then Σ is the positive boundary of C_1 , whose negative boundary is the union of a copy of P and some 2–disks. It follows that the genus of Σ is at least that of P. Hence Σ_0 defines a minimal genus Heegaard splitting of M. We now assume that $g(\Sigma) = g(P)$.

For $i = 1, \ldots, k$, choose a properly embedded arc γ_i in C_1 with an end on $\partial B_i \cap C_1$ and the other end on $P \times \{0\}$ (hence the ends of γ_i lie on different components of $\partial_{-}C_1$). Since C_1 is obtained from a thickening of ∂ −C₁ by gluing k 1–handles (which make C₁ connected), the arcs γ_i can be chosen so that C_1 is a regular neighborhood of $\partial_{-}C_1 \cup (\bigcup_{i=1}^{k} \gamma_i)$. Hence the uniqueness of the Heegaard splitting will follow from the uniqueness of the collection of arcs (γ_i) up to isotopy. We consider here isotopies within similar collections of arcs, which means that the ends are not fixed, but they must remain in ∂ −C₁. Such collections of arcs are clearly homotopic, so we need to show that homotopic collections are isotopic. We shall see that we can allow a strand to cross another. Indeed, any point of an arc γ_i is the center of a properly embedded disk in C_1 whose boundary c is parallel to the component of the suture s which lies on ∂B_i . This curve c also bounds a disk δ in C_2 . Hence we can slide part of any γ_j (including γ_i) along this disk δ in order to realize a crossing with γ_i . .

Theorem 3.14. Consider a double compression body $M = (\sharp(P \times I)) \sharp (\sharp^k(S^1 \times S^2))$, where P is any compact surface and $k \geq 0$, with the sutured decomposition of ∂M defined above. The sutured manifold M admits a unique minimal genus Heegaard splitting up to isotopy, and this minimal genus is $g(P) + k$.

Proof. If F is a component of P, the minimal genus Heegaard splitting of $F \times I$ is given in Proposition [3.13.](#page-5-0) The components $S^1 \times S^2$ have a standard genus–1 splitting. The connected sum of all these splittings gives a Heegaard splitting of M of genus $g(P) + k$.

Now let $M = C_1 \cup_{\Sigma} C_2$ be any Heegaard splitting of M.

First assume that $k > 0$. Then, thanks to Theorem [3.10,](#page-4-1) there is a non-separating essential sphere $S \subset M$ which is Haken. Cutting along this sphere and adding $S \cap \Sigma$ to the suture produces a Heegaard splitting of $(\sharp (P' \times I)) \sharp (\sharp^{k-1} (\check{S}^1 \times \check{S}^2))$, where P' is the disjoint union of P and two 2-disks, and the genus has decreased by one. Hence, by an inductive argument, we are led to the case $k = 0$.

We proceed by induction on $(g, n) = (g(\Sigma), |\partial\Sigma|)$ with lexicographic order. The case $(g, n) = (0, 0)$ is reduced to $M = S³$, which has a unique genus–0 splitting. For $(g, n) \succ (0, 0)$, first assume that M

contains an essential 2–sphere S which is non peripheral (*ie* non-parallel to a boundary component). By Theorem [3.10,](#page-4-1) we can assume that S is Haken. Hence, cutting along S provides two double compression bodies, with Heegaard splittings whose genera add up to give $g(\Sigma)$, and for both the value of (g, n) has strictly decreased. We are led to the case when M contains no essential non-peripheral 2–sphere, which gives three possibilities: P is connected, P is the disjoint union of a connected surface and a disk, or P is the disjoint union of three disks. All three cases are covered by Proposition [3.13.](#page-5-0)

In [\[Wal68\]](#page-9-12), Waldhausen classified the Heegaard splittings of the 3–sphere. In [\[ST93\]](#page-9-13), Scharlemann and Thompson classified the Heegaard splittings of a product $S \times I$ where S is a closed surface. Together with the strong Haken theorem and the above result, it provides a full classification of Heegaard splittings of double compression bodies, with the sutured decomposition we have fixed.

Theorem 3.15 (Waldhausen). Every Heegaard splitting of positive genus of $S³$ is stabilized.

This result and the Haken theorem provide a classification of Heegaard splittings of $\sharp^g(S^1 \times S^2)$ up to diffeomorphisms. As explained in [\[HS21\]](#page-9-4), one can get a classification up to isotopy, using the strong Haken theorem and the classification of genus–0 splittings of a punctured 3–ball ([\[HS21,](#page-9-4) Theorem 3.3], also a particular case of Proposition [3.13\)](#page-5-0). This is recovered in Theorem [3.17](#page-6-0) below, for P empty.

Theorem 3.16 (Scharlemann–Thompson). Let $M = P \times I$, where P is a compact connected oriented surface of genus g, with the sutured decomposition of ∂M defined above. Every Heegaard splitting of genus $n > q$ of M is stabilized.

Proof. This is proven in [\[ST93\]](#page-9-13) for P closed. The general case easily follows. For each component c of ∂P , we fill in $c \times I$ with a solid tube and we cap off the Heegaard surface with a disk. The closed case tells us that the splitting we obtain is stabilized. The solid tubes we added do not interact with the stabilization, so the original Heegaard splitting was already stabilized. \square

Theorem 3.17. Consider a double compression body $M = (\sharp(P \times I)) \sharp (\sharp^k(S^1 \times S^2))$, where P is any compact surface and $k \geq 0$, with the sutured decomposition of ∂M defined above. For all $n \geq q(P) + k$, the sutured manifold M admits a unique Heegaard splitting of genus n up to isotopy.

Proof. Assume $n > q(P) + k$. We need to prove that, in this case, the Heegaard splitting is stabilized. As in the proof of Theorem [3.14,](#page-5-1) cutting along non-separating Haken spheres, we reduce to problem to the case $k = 0$.

Thanks to Theorem [3.10,](#page-4-1) there are Haken spheres which realize M as the connected sum of some products $F \times I$ with F a connected surface. We cut along these Haken spheres and then fill in each created boundary with a 3–ball containing a properly embedded 2–disk which caps the Heegaard surface. We obtain a Heegaard splitting on each $F \times I$, one of which has a genus greater than $g(F)$. By Theorem [3.16,](#page-6-2) this splitting is stabilized, so that our initial splitting of M was already stabilized. \square

4. Diffeomorphisms of 4–dimensional compression bodies and relative trisections

4.1. A relative Laudenbach–Poénaru's theorem. In this section, we adapt our proof of Laudenbach– Poénaru's theorem to a relative setting. We had two preliminaries for this proof. The first one is the classification of minimal genus splittings of double handlebodies. We generalized it in Theorem [3.14](#page-5-1) to double compression bodies. The second one is the fact that every diffeomorphism of a 3–dimensional handlebody which is the identity on the boundary is isotopic to the identity. We now generalize it to compression bodies.

Lemma 4.1. Let C be a compression body. Every two defining disk systems for C which coincide on the boundary are isotopic.

Proof. We have proved in Lemma [3.3](#page-3-2) that compression bodies are irreducible. Hence a standard innermost disk argument concludes.

Lemma 4.2. Let C be a compression body. Let φ be a diffeomorphism of C. If φ is the identity on $\partial_{+}C$, then φ is isotopic to the identity.

Proof. Pick a defining disk system D for C . By the same reasoning as in the proof of Lemma [2.4,](#page-2-1) we can assume that φ is the identity on $\partial_+ C \cup \mathcal{D}$. We are led to a diffeomorphism of a product $\partial_- C \times I$ which is the identity on ∂ -C × {1}. Interpolating with the identity provides the required isotopy.

Definition 4.3. A *(lensed) hyper compression body* V is a smooth connected manifold constructed as follows:

- start with $M \times I$ where M is a compact oriented 3-manifold,
- glue 4–dimensional 1–handles along $M \times \{1\}$,
- collapse the vertical boundary along the I–factor.

The negative boundary ∂ −V is defined as $M \times \{0\}$ and the positive boundary is $M \times \{1\}$, so that $\partial V = \partial_{-} V \cup_{\partial} \partial_{+} V.$

We will consider hyper compression bodies with a specific condition on the negative boundary. We say that a hyper compression body V is P–based if ∂ −V is a trivial compression body P × I, where P is a compact oriented surface. Note that it comes with a sutured decomposition of its boundary, defined as in Section [3.3.](#page-5-2) Further, the positive boundary is diffeomorphic to a double compression body $(\sharp (P \times I)) \sharp (\sharp^k (S^1 \times S^2)),$ again with a sutured structure.

Theorem 4.4. Let V be a P-based hyper compression body. Assume that P contains no 2 -sphere. Then every diffeomorphism of $\partial_+ V$ extends to a diffeomorphism of V.

Proof. Like in the proof of Theorem [2.5,](#page-2-2) we see that there is a foliation of V by compression bodies C_t , $t \in [0, 1]$, which intersect along their positive boundary $\Sigma = \partial_+ C_t$, such that $C_0 = \partial_0(\partial_+ V)$ and $C_1 =$ $\partial_1(\partial_+V)$. We conclude with the very same argument, using Lemma [4.2](#page-6-3) instead of Lemma [2.4.](#page-2-1)

4.2. Diagrams of 4–dimensional multisections. In this section, we apply Theorem [4.4](#page-7-0) to the problematic of diagrams in the setting of relative multisections in the sense of Islambouli–Naylor [\[IN24\]](#page-9-14), a generalization of Gay and Kirby's trisections [\[GK16\]](#page-9-15).

Definition 4.5. A multisection of a compact oriented 4-manifold X is a decomposition $X = \bigcup_{i=1}^{n} X_i$ with the following properties (all arithmetic involving indices is mod n):

- (1) $\Sigma = \bigcap_{i=1}^{n} X_i$ is a compact connected oriented surface,
- (2) when $|i j| > 1$, $X_i \cap X_j = \Sigma$.
- (3) $C_i = X_i \cap X_{i+1}$ is a 3-dimensional compression body satisfying $\partial_+ C_i = \Sigma$ and $\partial_- C_i = C_i \cap \partial X$,
- (4) there is a compact oriented surface P, which contains no S^2 , such that each X_i is a P-based hyper compression body with ∂ − $X_i = X_i \cap \partial X$, and the natural sutured decomposition of $\partial(\partial - X_i)$ coincides with the decomposition $\partial(\partial_- X_i) = \partial_- C_{i-1} \cup \partial_- C_i$.

A multisection is called a *trisection* when $n = 3$.

Figure 1. Schematic of a multisection

A diagram of such a multisection is a tuple $(\Sigma; \alpha_1, \ldots, \alpha_n)$ where Σ is the central surface of the multisection and α_i is a cut-system for C_i . Note that the positive boundaries of the X_i are double

compression bodies, so that Theorem [3.17](#page-6-0) implies that each subdiagram $(\Sigma; \alpha_{i-1}, \alpha_i)$ is handleslide diffeomorphic to a diagram as represented in Figure [2.](#page-8-0) Note that any abstract diagram satisfying this property is a diagram of some multisected 4–manifold. When the surface P has no closed component, Castro, Gay and Pinzón-Caicedo proved that a multisection diagram determines a unique 4–manifold up to isotopy [\[CGPC18\]](#page-9-5). Theorem [4.4](#page-7-0) allows us to give a simple proof of this fact and to extend it to any surface P containing no S^2 -component. We will see below that this is optimal.

FIGURE 2. Heegaard diagram for $C_{i-1} \cup C_i$

Proposition 4.6. A multisection diagram determines a unique compact 4–manifold up to diffeomorphism.

Proof. Let X and X' be multisected 4–manifolds with diffeomorphic multisection diagrams $(\Sigma; \alpha_1, \ldots, \alpha_n)$ and $(\Sigma'; \alpha'_1, \ldots, \alpha'_n)$, meaning that there is a diffeomorphism $\varphi : \Sigma \to \Sigma'$ such that $\varphi(\alpha_i) = \alpha'_i$. First extend φ along defining disk systems for the 3–dimensional compression bodies of the multisection. Then extend it to the whole compression bodies (which amounts to extending a diffeomorphism from S^2 to B^3 or from $P \times \{1\}$ to $P \times I$). Finally extend φ to the 4-dimensional compression bodies using Theorem [4.4.](#page-7-0)

4.3. The bad case. In this section, we discuss the failure of Proposition [4.6](#page-8-1) in the case when the surface P is allowed to contain some 2–spheres. Accordingly, we allow 2–spheres in the negative boundary of a 3–dimensional compression body. We will analyse an example which was explained to us by David Gay.

Start with a torus $\Sigma = S^1 \times S^1$ and three parallel essential simple closed curves $\alpha_i \subset \Sigma$. Form the product $\Sigma \times \Delta$ where Δ is a 2–disk. Choose three distinct points $p_i \in \partial \Delta$, $i = 1, 2, 3$, and glue a 4– dimensional 2-handle $D_i \times D^2$ along $\alpha_i \times \{p_i\}$ for each i. More precisely glue the handle along $A_i \times [x_i, y_i]$ where A_i is a closed tubular neighborhood of α_i in Σ and $[x_i, y_i]$ an interval around p_i in $\partial \Delta$, so that $D_i \times \{p_i\}$ is the core of the handle, see Figure [3](#page-9-16) for the order of the points on the circle. It remains to glue some 3–handles.

Define a foliation of Σ by simple closed curves α_t , $t \in [0,3]$, such that, for $i = 1,2,3$, α_i is the curve previously defined, and $\alpha_0 = \alpha_3$. From now on, the indices i are considered modulo 3. For each i, let $t \in [0,1] \mapsto q_i(t) \in \partial \Delta$ be an injective path from y_{i-1} to x_i , chosen so that q_1 , q_2 and q_3 are pairwise disjoint. Set $\Sigma_i = \bigcup_{t \in [0,1]} \{q_i(t)\} \times \alpha_{t+i-1}$ (Figure [3](#page-9-16) represents the projection $\widetilde{\Sigma}_i$ of Σ_i on Σ). Now let D_i^x be a disk parallel to D_i on the boundary of the 2-handle and attached to $\alpha_i \times \{x_i\}$. Similarly define D_{i-1}^y . For $i = 1, 2$, glue a 3-handle B_i along $D_{i-1}^y - D_i^x + \Sigma_i$.

In the gluing of B_3 , we shall give more flexibility in order to produce a family of distinct manifolds. Fix an integer $n \geq 0$. Define a path $q_3^n(t) \in \partial \Delta$ from y_2 to x_3 as a concatenation of q_3 and $n-1$ full turns around Δ . Set $\Sigma_3^n = \bigcup_{t \in [0,1]} \{q_3^n(t)\} \times \alpha_{t+2}$, and glue a 3-handle B_3 along $D_2^y - D_3^x + \Sigma_3^n$.

We claim that this defines trisected manifolds with boundary X_n sharing a common trisection diagram, namely that of Figure [3.](#page-9-16) Since they are constructed by gluing handles, it is easy to compute their homology. We get $H_2(X_n) \cong \mathbb{Z}/n\mathbb{Z}$, so that the X_n are non-homeomorphic manifolds.

FIGURE 3. Decomposition of the surface Σ and schematic of the construction of the manifolds X_n

This failure of Proposition [4.6](#page-8-1) can be analysed as follows. The 3–dimensional pieces of the trisections we constructed are punctured solid tori, say C_i . These are non-irreducible, and each curve α_i on their positive boundary bounds a family of pairwise non-isotopic properly embedded disks indexed by Z. This implies that they admit diffeomorphisms that restrict to the identity on the positive boundary, but are non-isotopic to the identity. The non-existence of such diffeomorphisms was a key point in our proof of the relative Laudenbach–Poénaru theorem. One can check that the X_n are related by the following move. Cut X_n along one of the $C_i \setminus \text{Int}(\partial_+ C_i)$ and reglue via a diffeomorphism of C_i that restricts to the identity on the positive boundary, but is non-isotopic to the identity.

REFERENCES

- [Cer68] Jean Cerf, Sur les difféomorphismes de la sphère de dimension trois $(\Gamma_4 = 0)$, Lecture Notes in Mathematics, vol. No. 53, Springer-Verlag, Berlin-New York, 1968.
- [CGPC18] Nickolas Andres Castro, David T Gay, and Juanita Pinzón-Caicedo, Trisections of 4–manifolds with boundary, Proceedings of the National Academy of Sciences 115 (2018), no. 43, 10861–10868.
- [CO05] Leonardo N Carvalho and Ulrich Oertel, A classification of automorphisms of compact 3–manifolds, arXiv: [math/0510610,](http://arxiv.org/abs/math/0510610) 2005.
- [Dis23] Rudy Dissler, Relative trisections of fiber bundles over the circle, [arXiv:2304.09300,](http://arxiv.org/abs/2304.09300) 2023.
- [GK16] David T Gay and Robion Kirby, Trisecting 4–manifolds, Geometry & Topology 20 (2016), no. 6, 3097–3132.
- [Hak68] Wolfgang Haken, Some results on surfaces in 3–manifolds, Studies in Modern Topology/Prentice-Hall (1968).
- [HS21] Sebastian Hensel and Jennifer Schultens, Strong Haken via sphere complexes, [arXiv:2102.09831.](http://arxiv.org/abs/2102.09831)
- [IN24] Gabriel Islambouli and Patrick Naylor, Multisections of 4–manifolds, Transactions of the American Mathematical Society 377 (2024), no. 2, 1033–1068.
- [Jac80] William Jaco, Lectures on three-manifold topology, CBMS Regional Conference Series in Mathematics, vol. 43, American Mathematical Society, Providence, RI, 1980.
- [Joh] Jesse Johnson, Notes on Heegaard splittings, available on the web.
- [Juh06] András Juhász, Holomorphic discs and sutured manifolds, Algebraic & Geometric Topology 6 (2006), no. 3, 1429–1457.
- [Lau73] François Laudenbach, Sur les 2-sphères d'une variété de dimension 3, Annals of Mathematics 97 (1973), no. 1, 57–81.
- [LP72] François Laudenbach and Valentin Poénaru, A note on 4-dimensional handlebodies, Bulletin de la Société Mathématique de France 100 (1972), 337–344.
- [Sch24] Martin Scharlemann, A strong Haken theorem, Algebraic & Geometric Topology 24 (2024), no. 2, 717–753.
- [ST93] Martin Scharlemann and Abigail Thompson, *Heegaard splittings of (surface)×I are standard*, Mathematische Annalen 295 (1993), no. 1, 549–564.
- [Wal68] Friedhelm Waldhausen, Heegaard-Zerlegungen der 3-sphäre, Topology 7 (1968), 195-203.