Nonequilibrium Fluctuation-Response Relations for State Observables

Krzysztof Ptaszyński,^{1,*} Timur Aslyamov,^{2,†} and Massimiliano Esposito^{2,‡}

¹Institute of Molecular Physics, Polish Academy of Sciences, Mariana Smoluchowskiego 17, 60-179 Poznań, Poland

²Department of Physics and Materials Science, University of Luxembourg, L-1511 Luxembourg City, Luxembourg

(Dated: December 16, 2024)

Time-integrated state observables, which quantify the fraction of time spent in a specific pool of states, are important in many fields, such as chemical sensing or theory of fluorescence spectroscopy. We derive exact identities, called Fluctuation-Response Relations (FRRs), that connect the fluctuations of such observables to their response to external perturbations in nonequilibrium steady state of Markov jump processes. Using these results, we derive novel upper and lower bounds for fluctuations. We further demonstrate their applicability for simplifying calculations of fluctuations in large Markov networks, use them to explain the physical origin of positive and negative correlations of occupation times in a double quantum dot device, and discuss their relevance for model inference.

Introduction-Both the ordered responses to external perturbations and the stochastic fluctuations are among the central topics of statistical physics. Close to equilibrium, the responses and fluctuations are strictly related via a seminal fluctuation-dissipation theorem [1–5]. Far from equilibrium, such universal relations are no longer valid. However, in recent years, significant progress has been made in describing the universal properties of static responses to external perturbations in Markov processes or chemical reaction networks [6-20]. Similarly, research in recent decades has produced a wealth of universal laws describing the properties of fluctuations, such as fluctuation theorems [21-24], generalizations of fluctuation-dissipation theorem [25-33], or Thermodynamic and Kinetic Uncertainty Relations (TURs and KURs), lower bounding the current fluctuations in terms of average current and entropy production or traffic (also called activity) [34–44].

Very recently, significant developments have also been made in connecting the responses and fluctuations, in the spirit of the original fluctuation-dissipation theorem, in systems arbitrarily far from equilibrium [45-47]. In particular, Refs. [48, 49] used information theory to obtain an inequality bounding the precision of response (i.e., its ratio to fluctuations) of an arbitrary trajectory observable in Markov jump processes by the traffic. This result was later generalized to open quantum systems [50, 51]. Complementing those results, Ref. [52] obtained a similar Response-TUR (R-TUR), bounding the precision of response to kinetic perturbations (i.e., perturbations affecting the system kinetics, but not thermodynamic forces) in a Nonequilibrium Steady State (NESS) by the entropy production. This result was later proven in Ref. [33], which further derived exact identities, called Fluctuation-Response Relations (FRRs), relating the current fluctuations in NESS to static responses to kinetic or thermodynamic perturbations. Subsequently, R-TUR has been generalized to transient regime using information theory [50].

The kinetic and thermodynamic bounds derived in Refs. [48–50] were also applicable to precision of response of

time-integrated state observables, which describe the fraction of time spent in a given state. Such observables and their fluctuations have attracted significant attention in many fields, including diffusion [53–60], motion of active particles [61, 62], theory of fluorescence microscopy [63, 64], optical systems [65, 66], or chemical sensing [67–76]. In particular, in the latter context, they enter a famous Berg-Purcell bound for the precision of concentration measurements [77]. Other types of inequalities bounding the fluctuations of state observables have also been derived in Ref. [78] for the timedependent scenario and in Ref. [79] for the stationary case. However, the exact identities relating fluctuations and responses of state observables, analogous to FRRs obtained for current [33], have not been known so far.

In our Letter we close this gap by deriving exactly such identities. Interestingly, we found that they are identical in structure to FRRs proven for currents. The physical meaning of our identities is illustrated by applying them to explain the origin of positive and negative correlations of occupation times in a double-quantum-dot device. We also discuss the relation of our result to recently proven inequalities [48–50, 79], derive the upper bound for fluctuations and the lower bound for the precision of equilibrium response to state energies, and discuss potential applications of our framework for simplifying calculation of fluctuations or inferring the structure of Markov networks using observational data.

Setup—We consider a continuous-time Markov jump process among N discrete states. It is described by the directed graph whose nodes correspond to the system states and the edges e to transitions between states. NESS of the system is defined as

$$d_t \boldsymbol{\pi} = \boldsymbol{\mathbb{W}} \cdot \boldsymbol{\pi} = 0, \qquad (1)$$

where $\pi = (..., \pi_n, ...)^{\top}$ is the vector of state probabilities π_n with $\sum_n \pi_n = 1$. The matrix \mathbb{W} is the rate matrix with off-diagonal elements $W_{nm} = \sum_e [W_{+e}\delta_{s(+e)m}\delta_{t(+e)n} + W_{-e}\delta_{s(-e)m}\delta_{t(-e)n}]$ and the diagonal elements $W_{nn} = -\sum_{m\neq n} W_{mn}$. Here, to each edge *e* we assign a forward (+*e*) and reverse (-*e*) direction, so that the source of the edge $\pm e$, labeled $s(\pm e)$, is a target of edge $\mp e$, labeled $t(\mp e)$. Every transition rate associated with the edge $\pm e$ can be generically

^{*} krzysztof.ptaszynski@ifmpan.poznan.pl

[†] timur.aslyamov@uni.lu

[‡] massimiliano.esposito@uni.lu

parameterized as [10, 13]

$$W_{\pm e} = \exp(X_{\pm e})$$
 with $X_{\pm e} = \pm V_{s(\pm e)} + B_e \pm S_e/2$, (2)

where B_e and S_e are called symmetric and asymmetric edge parameters, respectively, and V_n are called vertex parameters. We note that this parameterization is not unique for a given Markov network, but can acquire a clear physical interpretation within specific physical contexts. For example, taking $V_n = 0$, B_e characterize the kinetic barriers, while S_e is the change in entropy in the reservoir due to a transition *e* that includes changes in thermodynamic forces and energy landscape [10, 80, 81]. However, the energy landscape may be alternatively parameterized by vertex parameters V_n , which is particularly useful at equilibrium; we discuss that later.

State observables—Our object of interest are the state observables that are time integrated along the stochastic trajectory of the system,

$$\hat{o}(t) \equiv \frac{1}{t} \sum_{n} x_n \int_0^t \phi_n(t') dt', \qquad (3)$$

where $\mathbf{x} \equiv (..., x_n, ...)^{\mathsf{T}}$ is an arbitrary vector and $\phi_n(t)$ is the random variable taking value 1 when the state *n* is occupied and 0 otherwise. The average value of this observable is defined as

$$O(t) \equiv \langle \hat{o}(t) \rangle, \tag{4}$$

where $\langle \cdot \rangle$ denotes the average over the ensemble of stochastic trajectories. The covariance of two observables is in turn defined as

$$\langle \langle O(t), O'(t) \rangle \rangle \equiv t \langle \Delta \hat{o}(t) \Delta \hat{o}'(t) \rangle, \qquad (5)$$

where $\Delta \hat{o}(t) \equiv \hat{o}(t) - \langle \hat{o}(t) \rangle$.

We further mostly focus on the long-time version of the quantities considered,

$$O \equiv \lim_{t \to \infty} O(t) = \sum_{n} x_n \pi_n, \qquad (6a)$$

$$\langle\langle O, O' \rangle\rangle \equiv \lim_{t \to \infty} \langle\langle O(t), O'(t) \rangle\rangle.$$
 (6b)

We emphasize that while the average observable O is determined only by the stationary state π , the covariances $\langle \langle O, O' \rangle \rangle$ are dynamical quantities, dependent on the ensemble of stochastic trajectories. They are given by the algebraic expression

$$\langle \langle O, O' \rangle \rangle = x^{\mathsf{T}} \mathbb{C} x', \qquad (7)$$

where $\mathbf{x}' \equiv (..., x'_n, ...)^{\mathsf{T}}$ and $\mathbb{C} = [C_{mn}]$ is the covariance matrix of occupation times of different states. It is defined as

$$C_{mn} \equiv \lim_{t \to \infty} \frac{1}{t} \langle \theta_m(t) \theta_n(t) \rangle, \qquad (8)$$

where $\theta_n(t) \equiv \int_0^t [\phi_n(t') - \pi_n] dt'$. It can be calculated using

the formula (see Appendix A) [58-60, 79]

$$\mathbb{C} = -\mathbb{W}^D \cdot \operatorname{diag}(\boldsymbol{\pi}) - [\mathbb{W}^D \cdot \operatorname{diag}(\boldsymbol{\pi})]^{\mathsf{T}}, \qquad (9)$$

where \mathbb{W}^D is the Drazin inverse of the rate matrix (also called resolvent [79]), see Refs. [82, 83] for a definition and the discussion of its properties.

We also consider the static responses of average observables to perturbations of transition rates. Throughout our Letter, we focus on a situation where the vector x does not depend on the perturbed parameters. For such a case, we have

$$d_{\xi}O = \mathbf{x}^{\mathsf{T}}d_{\xi}\boldsymbol{\pi}, \quad \xi \in \{X_e, B_e, S_e, V_n\},$$
(10)

where $d_{\xi}\pi$ is the static response of the stationary probability vector to the perturbation of transition rates. It can be calculated as [84]

$$d_{\xi}\boldsymbol{\pi} = -\mathbb{W}^D(d_{\xi}\mathbb{W})\boldsymbol{\pi} \,. \tag{11}$$

This expression is rederived in Appendix B. We notice that the Drazin inverse form of Eq. (11) is an alternative to the method from Refs. [13, 14].

Fluctuation-response relations—Let us now present our main result, namely, the exact identities relating the covariance matrix elements to static responses of state probabilities. They are proven in the Appendix C. They read as

$$C_{mn} = \sum_{\pm e} \frac{1}{W_{\pm e} \pi_{s(\pm e)}} d_{X_{\pm e}} \pi_m d_{X_{\pm e}} \pi_n, \qquad (12a)$$

$$C_{mn} = \sum_{e} \frac{\tau_e}{j_e^2} d_{B_e} \pi_m d_{B_e} \pi_n \,, \tag{12b}$$

$$C_{mn} = \sum_{e} \frac{4}{\tau_e} d_{S_e} \pi_m d_{S_e} \pi_n , \qquad (12c)$$

where $\sum_{\pm e}$ denotes the summation over forward and reverse edge directions, \sum_e denotes the summation over forward directions, $j_e = W_{+e}\pi_{s(+e)} - W_{-e}\pi_{t(+e)}$ is the oriented current through the edge e, and $\tau_e = W_{+e}\pi_{s(+e)} + W_{-e}\pi_{t(+e)}$ is the unoriented traffic through that edge. We note that Eq. (12b) can be directly applied only when no edge is detailed balanced, i.e., when all edge currents j_e are nonzero; in the opposite case, one needs to consider a limit of small perturbation from equilibrium. In contrast, the other identities can be applied directly also to equilibrium systems.

The above identities produce analogous fluctuationresponse relation (FRRs) for generic state observables,

$$\langle \langle O, O' \rangle \rangle = \sum_{\pm e} \frac{1}{W_{\pm e} \pi_{s(\pm e)}} d_{X_{\pm e}} O d_{X_{\pm e}} O', \qquad (13a)$$

$$\langle \langle O, O' \rangle \rangle = \sum_{e} \frac{\tau_e}{j_e^2} d_{B_e} O d_{B_e} O', \qquad (13b)$$

$$\langle\langle O, O' \rangle\rangle = \sum_{e} \frac{4}{\tau_e} d_{S_e} O d_{S_e} O' ,$$
 (13c)

which are identical in structure to FRRs for currents derived

in Ref. [33]. The identities (12) and (13) are the main result of our Letter.

Upper bound for fluctuations of state observables—We now show that our result further produces upper bounds for the variance of state observables $\langle \langle O \rangle \rangle \equiv \langle \langle O, O \rangle \rangle$ expressed solely in terms of the rate matrix W and the stationary state π . To derive them, we use the result of Ref. [10], bounding the response to symmetric and asymmetric perturbations as

$$\left| d_{B_e} \pi_n \right| \le \pi_n (1 - \pi_n) \tanh \left(\mathcal{F}_{\max} / 4 \right), \tag{14a}$$

$$\left| d_{S_e} \pi_n \right| \le \pi_n (1 - \pi_n) \,, \tag{14b}$$

where $\mathcal{F}_{max} = \max_c |\mathcal{F}_c|$ is the maximum of the cycle affinities \mathcal{F}_c . Using Eq. (14) with Eq. (12), we obtain upper bounds for the diagonal covariance matrix elements,

$$C_{nn} \le \pi_n^2 (1 - \pi_n)^2 \tanh^2 \left(\mathcal{F}_{\max}/4 \right) \sum_e \frac{\tau_e}{j_e^2},$$
 (15a)

$$C_{nn} \le \pi_n^2 (1 - \pi_n)^2 \sum_e \frac{4}{\tau_e}$$
 (15b)

As proven in the Appendix D, the above relations produce an upper bound for the variance of a generic state observable,

$$\langle \langle O \rangle \rangle \le \mathbf{x}^{\mathsf{T}} \mathbf{x} \tanh^2 \left(\mathcal{F}_{\max}/4 \right) \sum_{n,e} \pi_n^2 (1 - \pi_n)^2 \frac{\tau_e}{j_e^2}, \quad (16a)$$

$$\langle \langle O \rangle \rangle \le \mathbf{x}^{\mathsf{T}} \mathbf{x} \sum_{n,e} \pi_n^2 (1 - \pi_n)^2 \frac{4}{\tau_e} \,.$$
 (16b)

We notice that this result is different from the spectral upper bound for fluctuations of time-integrated fluxes presented in Ref. [85].

Relation to fluctuation-response inequalities—Let us now discuss the relation of our identities to previous results. First, they are strictly related to fluctuation-response inequalities, derived in Refs. [48–50] for the time-dependent variance $\langle \langle O(t) \rangle \rangle \equiv \langle \langle O(t), O(t) \rangle \rangle$:

$$\langle \langle O(t) \rangle \rangle \ge \sum_{\pm e} \frac{1}{W_{\pm e} \pi_{s(\pm e)}} [d_{X_{\pm e}} O(t)]^2,$$
 (17a)

$$\langle \langle O(t) \rangle \rangle \ge \sum_{e} \frac{\tau_e}{j_e^2} [d_{B_e} O(t)]^2,$$
 (17b)

$$\langle \langle O(t) \rangle \rangle \ge \sum_{e} \frac{4}{\tau_{e}} [d_{S_{e}}O(t)]^{2}$$
. (17c)

In that context, $d_{\xi}O(t)$ is the dynamic response to the perturbations introduced at t = 0, for a system initialized in the stationary state of an unperturbed rate matrix. In the longtime limit $t \to \infty$, it converges to the static response $d_{\xi}O$ considered in our paper. As further conjectured numerically in Ref. [50], the above inequalities converge to equalities for $t \to \infty$. This is proven by our analytic result (13).

We further note that the above relations have also been used to derive TURs and KURs, which bound the response to kinetic or thermodynamic perturbations by the stationary entropy production rate $\dot{\sigma} = \sum_{e} j_{e} \ln(W_{+e}/W_{-e})$ or total traffic $\mathcal{T} = \sum_{e} \tau_{e}$. To that end, one introduces single parameters κ , ε and η that parameterize the edge parameters as $X_{\pm e} = X_{\pm e}(\kappa)$, $B_e = B_e(\varepsilon)$ and $S_e = S_e(\eta)$. After applying the Cauchy-Schwarz inequality, one obtains [48–50]

$$\frac{[d_{\kappa}O(t)]^2}{\langle\langle O(t)\rangle\rangle} \le k_{\max}^2 \mathcal{T}, \qquad (18a)$$

$$\frac{[d_{\varepsilon}O(t)]^2}{\langle\langle O(t)\rangle\rangle} \le b_{\max}^2 \min(\dot{\sigma}/2, \mathcal{T}), \qquad (18b)$$

$$\frac{[d_{\eta}O(t)]^2}{\langle\langle O(t)\rangle\rangle} \le \frac{z_{\max}^2\mathcal{T}}{4}, \qquad (18c)$$

where $k_{\max} = \max_{\pm e} |d_{\kappa}X_{\pm e}|$, $b_{\max} = \max_{e} |d_{\varepsilon}B_{e}|$, and $z_{\max} = \max_{e} |d_{\eta}S_{e}|$.

Response to vertex perturbations—Here we complement Eqs. (17)–(18) by deriving similar inequalities, bounding response to perturbations of vertex parameters V_n . Such perturbations have a particularly simple interpretation in the longtime limit $t \rightarrow \infty$ at thermodynamic equilibrium, $\pi_n \propto e^{-\beta E_n}$, where E_n are the state energies and $\beta = 1/(k_B T)$ is the inverse temperature. Then, independent of the particular parameterization of the transition rates, the response to vertex perturbations corresponds to the response to perturbations of state energies: $d_{V_n}O = \beta^{-1}d_{E_n}O$ (see Appendix E).

To derive them, we first rewrite Eq. (17a) as

$$\begin{split} \langle \langle O(t) \rangle \rangle &\geq \sum_{n} \sum_{\pm e} \frac{[\pm \delta_{ns(\pm e)} d_{X_{\pm e}} O(t)]^2}{W_{\pm e} \pi_{s(\pm e)}} \\ &\geq \sum_{n} \frac{[\sum_{\pm e} \pm \delta_{ns(\pm e)} d_{X_{\pm e}} O(t)]^2}{\sum_{\pm e} \delta_{ns(\pm e)} W_{\pm e} \pi_{s(\pm e)}} \,. \end{split}$$
(19)

Here, in the second step, we applied Sedrakyan's inequality, $\sum_i a_i^2/b_i \ge (\sum_i a_i)^2/(\sum_i b_i)$ that holds for real a_i and positive b_i , to each sum with a given n. We than notice $d_{V_n}O = \sum_{\pm e} \pm \delta_{ns(\pm e)} d_{X_{\pm e}}O$ and $\sum_{\pm e} \delta_{ns(\pm e)} W_{\pm e} \pi_{s(\pm e)} = -W_{nn}\pi_n$. As a result, we obtain an inequality bounding the responses of state observables to vertex parameters,

$$\langle \langle O(t) \rangle \rangle \ge \sum_{n} \frac{[d_{V_n} O(t)]^2}{\pi_n |W_{nn}|} .$$
 (20)

In particular, precision of response to perturbation associated with each vertex *n* is bounded by the probability of that state π_n and the escape rate from that state $|W_{nn}|$:

$$\forall_n : \quad \frac{[d_{V_n} O(t)]^2}{\langle \langle O(t) \rangle \rangle} \le \pi_n |W_{nn}| \,. \tag{21}$$

Parameterizing the vertex parameters using a single control parameter v, $V_n = V_n(v)$, and using Sedrykyan's inequality, we further get

$$\frac{[d_{\nu}O(t)]^2}{\langle\langle O(t)\rangle\rangle} \le v_{\max}^2 \mathcal{T}, \qquad (22)$$

where $v_{\text{max}} = \max_n |d_v V_n|$.

Relation to "occupation uncertainty relation"-Let us now



FIG. 1. Scheme of one-dimensional Markov network with each state $n \in \{0, N\}$ apart from n = N being a source of a single edge n + 1 pointing to a tip n + 1.

show that the long-time version of Eq. (20) is equivalent to the so-called occupation uncertainty relation, recently derived in Ref. [79]. To this end, we rewrite Eq. (20) as

$$\langle \langle O \rangle \rangle \ge \sum_{n} \frac{(\boldsymbol{x}^{\intercal} d_{V_n} \boldsymbol{\pi})^2}{\pi_n |W_{nn}|} .$$
 (23)

We then use the exact relation for the static response to vertex perturbation, $d_{V_n}\pi_m = \pi_n(\pi_m - \delta_{nm})$ [10, 13]. In this way we reproduce the result of Ref. [79] (see Appendix F),

$$\langle \langle O \rangle \rangle = \mathbf{x}^{\mathsf{T}} \mathbb{C} \mathbf{x} \ge \mathbf{x}^{\mathsf{T}} \mathbb{Q} \cdot \Lambda^{-1} \cdot \operatorname{diag}(\mathbf{\pi}) \cdot \mathbb{Q}^{\mathsf{T}} \mathbf{x}, \quad (24)$$

where $(\mathbb{Q})_{mn} \equiv \delta_{nm} - \pi_m$ and $\Lambda \equiv \text{diag}(\dots, |W_{nn}|, \dots)$. This inequality, which lower bounds the fluctuations solely in terms of \mathbb{W} and π , complements the analogous upper bounds (16).

Application for calculating fluctuations—We now show that our approach can be used to analytically calculate the fluctuations of state observables in certain models admitting an exact solution, which is otherwise very difficult. We illustrate that on one-dimensional Markov networks [86] shown in Fig. 1. Such models have been applied in many contexts, e.g., to describe chemical bistability (Schlögl model) [87–89], bistable electric circuits [90, 91], magnetic systems (Curie-Weiss model) [84, 92–94], coupled heat engines [95, 96], or disease spread [97]. NESS of such systems is detailed balanced (even out of equilibrium), so that the state probabilities read

$$\pi_n = \pi_0 \prod_{m=1}^n \exp(S_m), \qquad (25)$$

where $S_n = \ln(W_{+n}/W_{-n})$, and π_0 is determined by the normalization condition $\sum_{n=0}^{N} \pi_n = 1$. As a result, one finds

$$d_{S_n}\pi_m = \pi_m \left[\Theta(m-n) - \sum_{k=n}^N \pi_k\right],$$
 (26)

where $\Theta(x)$ is the Heaviside theta with $\Theta(0) = 1$. Inserting this into Eq. (12c), we find an explicit expression for the covariance matrix elements

$$C_{mn} = \sum_{k=1}^{N} \frac{4d_{S_k} \pi_m d_{S_k} \pi_n}{\tau_k} , \qquad (27)$$

with $\tau_k = W_{+k}\pi_{k-1} + W_{-k}\pi_k$, and π_n , $d_{S_k}\pi_n$ as calculated in Eqs. (25)–(26). Consequently, the covariance matrix can be expressed explicitly in terms of transition rates $W_{\pm n}$.



FIG. 2. (a) Scheme of a system consisting of two capacitively coupled quantum dots *L* and *R* (left and right) with the electronic level energies E_L and E_R . Each quantum dot is coupled to a separate reservoir *L* and *R* with chemical potential $\mu = 0$ and inverse temperature $\beta = 1/(k_B T)$. We assume strong capacitative coupling $\beta E_C \gg 1$, so that the dots cannot be simultaneously occupied by an electron. Thus, the system can reside in three charge states $\{\emptyset, L, R\}$, which denote the empty state, and the state with the dot *L* or *R* occupied by a single electron, respectively. The electron transitions to (+) or from (-) the dot $\alpha \in \{L, R\}$ occur at the rate $\Gamma_{\pm\alpha} = \Gamma_{\alpha}/[1 + \exp(\pm\beta E_{\alpha})]$, where Γ_{α} is the tunneling rate. (b) The system dynamics is described by one-dimensional Markov network (see Fig. 1) with $\{L, \emptyset, R\} \rightarrow \{0, 1, 2\}, S_1 = \beta E_L, S_2 = -\beta E_R, W_{\pm 1} = \Gamma_{\mp L}, W_{\pm 2} = \Gamma_{\pm R}$.

Example—Finally, let us illustrate the physical meaning of our identities on the double quantum dot model shown in Fig. 2, which is similar to the experimental setup in Ref. [98]. We note that in such systems the charge states can be monitored in real time using charge counting techniques [99, 100]. This could enable the experimental verification of our result. Since the system can be described using one-dimensional Markov network [Fig. 2(b)], the covariance between times spent in states \emptyset and L can be expressed using Eq. (27) as

$$C_{\mathcal{O}L} = -\frac{4|d_{S_1}\pi_{\mathcal{O}}d_{S_1}\pi_L|}{\tau_1} + \frac{4|d_{S_2}\pi_{\mathcal{O}}d_{S_2}\pi_L|}{\tau_2}$$
$$= -\frac{4\pi_{\mathcal{O}}\pi_L^2(1-\pi_L)}{\Gamma_{+L}\pi_{\mathcal{O}} + \Gamma_{-L}\pi_L} + \frac{4\pi_{\mathcal{O}}\pi_L\pi_R^2}{\Gamma_{+R}\pi_{\mathcal{O}} + \Gamma_{-R}\pi_R}, \qquad (28)$$

with the first (second) term having a negative (positive) sign because of $d_{S_1}\pi_{\emptyset}d_{S_1}\pi_L < 0$ ($d_{S_2}\pi_{\emptyset}d_{S_2}\pi_L > 0$). This expression depends on both the state responses, described solely in terms of S_e [Eqs. (25)–(26)], and the edge traffics τ_e , which depend also on the system's kinetics determined by the tunneling rates. The above equation can be interpreted as follows. On the one hand, the system spends less time in the state \emptyset when it spends more time in L, which leads to the negative covariance (first term of the equation). This effect becomes less pronounced when the kinetics of transition $\emptyset \leftrightarrow L$ accelerates, i.e., when τ_1 increases, because state populations equilibrate faster to their average value. Similarly, the system spends less time in both states \emptyset and *L* if it spends more time in the state R. This leads to their positive covariance (the second term of the equation). Analogously, this effect becomes suppressed when the kinetics of transitions $\emptyset \leftrightarrow R$ accelerates (that is, when the traffic τ_2 increases), which reduces the probability that the system spends an unusually long or short time in the state R. Interestingly, our identity

demonstrates that the magnitude of those competing effects is governed by the response of state populations to asymmetric parameters S_e , which is a nontrivial result. Since the system is at equilibrium, these perturbations directly correspond to perturbations of experimentally controllable state energies: $d_{S_1} = \beta^{-1} d_{E_L}$, $d_{S_2} = -\beta^{-1} d_{E_R}$ (note that this generally does not hold out of equilibrium as E_{α} also affect B_e).

Concluding remarks—As illustrated by our example, identity (13) has a somewhat intuitive interpretation: It means that two state observables can be positively (negatively) correlated only when they respond with the same (opposite) sign to at least one symmetric and asymmetric edge perturbation. Beyond its fundamental importance, this result has practical relevance as it can serve to infer Markov models of physical systems that are consistent with measured data. On the methodological side, as illustrated by the Appendix, our Letter demonstrates that the recently developed algebraic approach to responses [13, 14, 33] and fluctuations [58– 60, 83] can be a powerful tool to determine the universal properties of these quantities. Examining the tightness of the bounds (15)–(16) and (20)–(22) for physically relevant setups is an interesting perspective.

ACKNOWLEDGMENTS

K.P., T.A. and M.E. acknowledge the financial support from, respectively, project No. 2023/51/D/ST3/01203 funded by the National Science Centre, Poland, project ThermoElectroChem (C23/MS/18060819) from Fonds National de la Recherche-FNR, Luxembourg, project TheCirco (IN-TER/FNRS/20/15074473) funded by FRS-FNRS (Belgium) and FNR (Luxembourg).

Appendix A: Equivalence of Eq. (9) and Refs. [58-60]

To demonstrate that Eq. (9) is equivalent to the result of Refs. [58–60], we express the latter using our notation,

$$C_{mn} = -\sum_{k} \frac{1}{\lambda_{k}} \left(\mathbf{1}^{\mathsf{T}} \mathbb{1}^{(m)} r_{k} l_{k}^{\mathsf{T}} \mathbb{1}^{(n)} \boldsymbol{\pi} + \mathbf{1}^{\mathsf{T}} \mathbb{1}^{(n)} r_{k} l_{k}^{\mathsf{T}} \mathbb{1}^{(m)} \boldsymbol{\pi} \right)$$

= $-\mathbf{1}^{\mathsf{T}} \mathbb{1}^{(m)} \mathbb{W}^{D} \mathbb{1}^{(n)} \boldsymbol{\pi} - \mathbf{1}^{\mathsf{T}} \mathbb{1}^{(n)} \mathbb{W}^{D} \mathbb{1}^{(m)} \boldsymbol{\pi}, \qquad (A1)$

where $\mathbf{1} \equiv (1, 1, ...)^{\mathsf{T}}, \mathbb{1}^{(n)} = \operatorname{diag}(\delta_{1n}, \delta_{2n}, ...), \lambda_k$ are nonzero eigenvalues of \mathbb{W} , and $\mathbf{l}_k^{\mathsf{T}}, \mathbf{r}_k$ are the associated left and right eigenvectors, $\mathbf{l}_k^{\mathsf{T}} \mathbb{W} = \lambda_k \mathbf{l}_k^{\mathsf{T}}, \mathbb{W} \mathbf{r}_k = \lambda_k \mathbf{r}_k$. In the second equality we use the spectral decomposition of the Drazin inverse, $\mathbb{W}^D = \sum_k \mathbf{r}_k \mathbf{l}_k^{\mathsf{T}} / \lambda_k$ [83]. Consequently

$$C_{mn} = -(\mathbb{W}^D)_{mn}\pi_n - (\mathbb{W}^D)_{nm}\pi_m, \qquad (A2)$$

which is equivalent to Eq. (9).

Appendix B: Derivation of Eq. (11)

Equation (11) can be derived by acting on Eq. (1) with derivative d_{ξ} . This yields

$$\mathbb{W}d_{\xi}\boldsymbol{\pi} = -(d_{\xi}\mathbb{W})\boldsymbol{\pi} . \tag{B1}$$

Acting on both sides of the above expression with the Drazin inverse \mathbb{W}^D we get

$$\mathbb{W}^{D}\mathbb{W}d_{\xi}\boldsymbol{\pi} = d_{\xi}\boldsymbol{\pi} - \boldsymbol{\pi}(\mathbf{1}^{\mathsf{T}}d_{\xi}\boldsymbol{\pi}) = -\mathbb{W}^{D}(d_{\xi}\mathbb{W})\boldsymbol{\pi}.$$
 (B2)

where we used $\mathbb{W}^{D}\mathbb{W} = \mathbb{1} - \pi \mathbf{1}^{\mathsf{T}}$ [83]. Since the response vector $d_{\xi}\pi$ is traceless, $\mathbf{1}^{\mathsf{T}}d_{\xi}\pi = 0$, which yields Eq. (11).

Appendix C: Proof of fluctuation-response relations (12)

Here we prove our main result. Let us first explicitly express the responses of state probabilities included in the right-hand side of Eqs. (12a)-(12c) in terms of Eq. (11). For generic perturbations, we get

$$d_{X_{\pm e}} \mathbb{W} = \begin{array}{c} \vdots \\ t(\pm e) \\ \vdots \\ s(\pm e) \\ \vdots \end{array} \begin{pmatrix} W_{\pm e} \\ W_{\pm e} \\ -W_{\pm e} \\ \vdots \end{pmatrix}, \quad (C1)$$

and thus $d_{X_{\pm e}}\pi_m = W_{\pm e}\pi_{s(\pm e)}[(\mathbb{W}^D)_{ms(\pm e)} - (\mathbb{W}^D)_{mt(\pm e)}].$ Consequently, using $t(\pm e) = s(\mp e)$, we have $d_{B_e}\pi_m = d_{X_{+e}}\pi_m + d_{X_{-e}}\pi_m = j_e[(\mathbb{W}^D)_{ms(+e)} - (\mathbb{W}^D)_{mt(+e)}]$ and $d_{S_e}\pi_m = (d_{X_{+e}}\pi_m - d_{X_{-e}}\pi_m)/2 = \tau_e[(\mathbb{W}^D)_{ms(+e)} - (\mathbb{W}^D)_{mt(+e)}]/2$. As a result, the right-hand sides of Eqs. (12a)–(12c) are the same, and can be expanded as

$$\sum_{\pm e} \frac{1}{W_{\pm e} \pi_{s(\pm e)}} d_{X_{\pm e}} \pi_m d_{X_{\pm e}} \pi_n$$

$$= \sum_e \frac{\tau_e}{j_e^2} d_{B_e} \pi_m d_{B_e} \pi_n = \sum_e \frac{4}{\tau_e} d_{S_e} \pi_m d_{S_e} \pi_n$$

$$= \sum_e \tau_e [(\mathbb{W}^D)_{ms(+e)} - (\mathbb{W}^D)_{mt(+e)}]$$

$$\times [(\mathbb{W}^D)_{ns(+e)} - (\mathbb{W}^D)_{nt(+e)}]. \quad (C2)$$

We now replace the summation over the edges *e* with the summation over the pairs of states *k* and *l*. To that end, we define the sum of all traffics at the edges connecting states *k* and *l*, $\tau_{kl} \equiv \sum_{e} [\delta_{ks(e)} \delta_{lt(e)} + \delta_{kt(e)} \delta_{ls(e)}] \tau_e = W_{kl} \pi_l + W_{lk} \pi_k$.

The above expression then becomes equal to

$$\sum_{k,l < k} \tau_{kl}(\mathbb{W}^D)_{mk}[(\mathbb{W}^D)_{nk} - (\mathbb{W}^D)_{nl}]$$

$$- \sum_{k,l < k} \tau_{kl}(\mathbb{W}^D)_{ml}[(\mathbb{W}^D)_{nk} - (\mathbb{W}^D)_{nl}]$$

$$= \sum_{k,l < k} \tau_{kl}(\mathbb{W}^D)_{mk}[(\mathbb{W}^D)_{nk} - (\mathbb{W}^D)_{nl}]$$

$$- \sum_{l,k < l} \tau_{lk}(\mathbb{W}^D)_{mk}[(\mathbb{W}^D)_{nl} - (\mathbb{W}^D)_{nk}]$$

$$= \sum_{k,l \neq k} \tau_{kl}(\mathbb{W}^D)_{mk}[(\mathbb{W}^D)_{nk} - (\mathbb{W}^D)_{nl}], \qquad (C3)$$

where in the first step we flipped the indices *k* and *l* in the second sum, while in the second step we used $\tau_{lk} = \tau_{kl}$ and noted $\sum_{k,l < k} (\cdot) + \sum_{l,k < l} (\cdot) = \sum_{k,l < k} (\cdot) + \sum_{k,l > k} (\cdot) = \sum_{k,l \neq k} (\cdot)$.

We now show that the covariance matrix elements C_{mn} are equal to the result of Eq. (C3). First, we note

$$\mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi}) = \mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi}) \cdot (\mathbb{W}^{D} \mathbb{W})^{\mathsf{T}}, \qquad (C4)$$

which results from the identities $(\mathbb{W}^D \mathbb{W})^{\mathsf{T}} = \mathbb{1} - \mathbf{1}\pi^{\mathsf{T}}$ and $\mathbb{W}^D \cdot \operatorname{diag}(\pi) \cdot \mathbf{1} = \mathbb{W}^D \pi = 0$ [83]. As a result

$$[\mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi})]_{mn} = [\mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi}) \cdot \mathbb{W}^{\mathsf{T}} (\mathbb{W}^{D})^{\mathsf{T}}]_{mn}$$
$$= \sum_{k,l} (\mathbb{W}^{D})_{mk} \pi_{k} W_{lk} (\mathbb{W}^{D})_{nl} .$$
(C5)

Analogously,

$$[\mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi})]_{mn}^{\mathsf{T}} = [\mathbb{W}^{D} \cdot \operatorname{diag}(\boldsymbol{\pi})]_{nm}$$
$$= \sum_{k,l} (\mathbb{W}^{D})_{nk} \pi_{k} W_{lk} (\mathbb{W}^{D})_{ml} = \sum_{k,l} (\mathbb{W}^{D})_{mk} \pi_{l} W_{kl} (\mathbb{W}^{D})_{nl},$$
(C6)

where in the last step we flipped indices k and l. Consequently,

$$C_{mn} = -[\mathbb{W}^D \cdot \operatorname{diag}(\boldsymbol{\pi})]_{mn} - [\mathbb{W}^D \cdot \operatorname{diag}(\boldsymbol{\pi})]_{mn}^{\mathsf{T}}$$
$$= -\sum_{k,l} (W_{lk}\pi_k + W_{kl}\pi_l)(\mathbb{W}^D)_{mk}(\mathbb{W}^D)_{nl}.$$
(C7)

Let us now focus on the term $(W_{lk}\pi_k + W_{kl}\pi_l)$. For $k \neq l$, it is equal to τ_{kl} defined below Eq. (C2). For k = l, we note that $-W_{kk}\pi_k = \sum_{l\neq k} W_{lk}\pi_k$ is the probability flux out of the state k. Due to the Kirchhoff's law, it is equal to the flux to that state, $\sum_{l\neq k} W_{kl}\pi_l$. Consequently, $(W_{kk}\pi_k + W_{kk}\pi_k) =$ $-\sum_{l\neq k} (W_{lk}\pi_k + W_{kl}\pi_l) = -\sum_{l\neq k} \tau_{kl}$. As a result, Eq. (C7) becomes

$$C_{mn} = \sum_{k,l \neq k} \tau_{kl} (\mathbb{W}^D)_{mk} [(\mathbb{W}^D)_{nk} - (\mathbb{W}^D)_{nl}], \qquad (C8)$$

which is equal to the result of Eq. (C3). This concludes the proof.

Appendix D: Proof of Eq. (16)

Using properties of matrix and vector norms, the variance of a generic state observable is bounded as

$$\langle \langle O \rangle \rangle = \mathbf{x}^{\mathsf{T}} \mathbb{C} \mathbf{x} \le ||\mathbf{x}||_2 ||\mathbb{C} \mathbf{x}||_2 \le ||\mathbf{x}||_2 (||\mathbb{C}||_2 ||\mathbf{x}||_2)$$
$$= ||\mathbf{x}||_2^2 ||\mathbb{C}||_2 = \mathbf{x}^{\mathsf{T}} \mathbf{x} ||\mathbb{C}||_2.$$
(D1)

Here, $||\mathbf{x}||_2 = \sqrt{\mathbf{x}^{\top}\mathbf{x}}$ is the Euclidean norm of \mathbf{x} and $||\mathbb{C}||_2$ is the spectral norm of \mathbb{C} , which (since \mathbb{C} is positive semidefinite) corresponds to its maximum eigenvalue. As \mathbb{C} is positive semidefinite, its maximum eigenvalue $||\mathbb{C}||_2$ is upper bounded by tr(\mathbb{C}) = $\sum_n C_{nn}$, which corresponds to the sum of all eigenvalues. Using Eqs. (15), we obtain Eq. (16).

Appendix E: Proof of relation $d_{V_n}O = \beta^{-1}d_{E_n}O$

At thermodynamic equilibrium, the state probabilities are expressed as $\pi_n = e^{-\beta E_n}/Z$, where $Z = \sum_m e^{-\beta E_m}$ is the partition function. This yields $d_{E_n}\pi_m = \beta \pi_n(\pi_m - \delta_{nm})$. At the same time, for a generic Markov network, $d_{V_n}\pi_m =$ $\pi_n(\pi_m - \delta_{nm})$ [10, 13]. Thus, at equilibrium, $d_{V_n}O = \beta^{-1}d_{E_n}O$.

Appendix F: Equivalence of Eqs. (23) and (24)

To demonstrate the equivalence of Eqs. (23) and (24), we use the relation $d_{V_n}\pi_m = \pi_n(\pi_m - \delta_{nm})$ to expand r.h.s. of Eq. (23) as

$$\sum_{n} \frac{(\mathbf{x}^{\mathsf{T}} d_{V_{n}} \pi)^{2}}{\pi_{n} |W_{nn}|} = \sum_{n} \frac{\left[\sum_{m} x_{m} \pi_{n} (\pi_{m} - \delta_{nm})\right]^{2}}{\pi_{n} |W_{nn}|}$$
$$= \sum_{n} \frac{\pi_{n} \left[\sum_{m} x_{m} (\pi_{m} - \delta_{nm})\right]^{2}}{|W_{nn}|} .$$
(F1)

Analogously, we expand r.h.s. of Eq. (24) using $[\mathbf{x}^{\mathsf{T}}\mathbb{Q}]_n = [\mathbb{Q}^{\mathsf{T}}\mathbf{x}]_n = \sum_m x_m(\delta_{nm} - \pi_m)$. This yields

$$\boldsymbol{x}^{\mathsf{T}} \mathbb{Q} \cdot \Lambda^{-1} \cdot \operatorname{diag}(\boldsymbol{\pi}) \cdot \mathbb{Q}^{\mathsf{T}} \boldsymbol{x} = \sum_{n} \frac{\pi_{n} [\sum_{m} x_{m} (\pi_{m} - \delta_{nm})]^{2}}{|W_{nn}|},$$
(F2)

which is equal to the result of Eq. (F1).

^[1] R. Kubo, Rep. Prog. Phys. 29, 255 (1966).

^[2] R. Kubo, M. Toda, and N. Hashitsume, Statistical physics II: nonequilibrium statistical mechanics, Vol. 31 (Springer Science)

& Business Media, 2012).

- [3] R. L. Stratonovich, Nonlinear nonequilibrium thermodynamics I: linear and nonlinear fluctuation-dissipation theorems, Vol. 57 (Springer Science & Business Media, 2012).
- [4] U. M. B. Marconi, A. Puglisi, L. Rondoni, and A. Vulpiani, Physics reports 461, 111 (2008).
- [5] D. Forastiere, R. Rao, and M. Esposito, New Journal of Physics 24, 083021 (2022).
- [6] V. Lucarini, Journal of Statistical Physics 162, 312 (2016).
- [7] M. Santos Gutiérrez and V. Lucarini, Journal of Statistical Physics 179, 1572 (2020).
- [8] G. Falasco, T. Cossetto, E. Penocchio, and M. Esposito, New Journal of Physics 21, 073005 (2019).
- [9] J. D. Mallory, A. B. Kolomeisky, and O. A. Igoshin, Proceedings of the National Academy of Sciences 117, 8884 (2020).
- [10] J. A. Owen, T. R. Gingrich, and J. M. Horowitz, Phys. Rev. X 10, 011066 (2020).
- [11] J. A. Owen and J. M. Horowitz, Nature Communications 14, 1280 (2023).
- [12] G. Fernandes Martins and J. M. Horowitz, Phys. Rev. E 108, 044113 (2023).
- [13] T. Aslyamov and M. Esposito, Phys. Rev. Lett. 132, 037101 (2024).
- [14] T. Aslyamov and M. Esposito, Phys. Rev. Lett. 133, 107103 (2024).
- [15] P. E. Harunari, S. Dal Cengio, V. Lecomte, and M. Polettini, Phys. Rev. Lett. 133, 047401 (2024).
- [16] F. Khodabandehlou, C. Maes, and K. Netočný, Affine relationships between steady currents (2024), arXiv:2412.05019 [cond-mat.stat-mech].
- [17] C. Floyd, A. R. Dinner, and S. Vaikuntanathan, arXiv preprint 10.48550/arXiv.2404.03798 (2024).
- [18] D. Frezzato, The Journal of Chemical Physics 160, 234111 (2024).
- [19] C. Floyd, A. R. Dinner, A. Murugan, and S. Vaikuntanathan, arXiv preprint 10.48550/arXiv.2409.05827 (2024).
- [20] Q. Gao, H.-M. Chun, and J. M. Horowitz, Phys. Rev. E 105, L012102 (2022).
- [21] C. Jarzynski, Phys. Rev. Lett. 78, 2690 (1997).
- [22] G. E. Crooks, Phys. Rev. E 60, 2721 (1999).
- [23] M. Esposito, U. Harbola, and S. Mukamel, Rev. Mod. Phys. 81, 1665 (2009).
- [24] U. Seifert, Rep. Prog. Phys. 75, 126001 (2012).
- [25] G. S. Agarwal, Zeitschrift f
 ür Physik A Hadrons and nuclei 252, 25 (1972).
- [26] U. Seifert and T. Speck, Europhys. Lett. 89, 10007 (2010).
- [27] J. Prost, J.-F. Joanny, and J. M. Parrondo, Phys. Rev. Lett. 103, 090601 (2009).
- [28] B. Altaner, M. Polettini, and M. Esposito, Phys. Rev. Lett. 117, 180601 (2016).
- [29] H.-M. Chun and J. M. Horowitz, The Journal of Chemical Physics 158, 174115 (2023).
- [30] N. Shiraishi, Fundamental Theories of Physics. Springer, Singapore (2023).
- [31] Q. Gao, H.-M. Chun, and J. M. Horowitz, Europhys. Lett. 146, 31001 (2024).
- [32] L. Tesser and J. Splettstoesser, Phys. Rev. Lett. 132, 186304 (2024).
- [33] T. Aslyamov, K. Ptaszyński, and M. Esposito, Nonequilibrium fluctuation-response relations: From identities to bounds (2024), arXiv:2410.17140 [cond-mat.stat-mech].
- [34] A. C. Barato and U. Seifert, Phys. Rev. Lett. 114, 158101 (2015).
- [35] T. R. Gingrich, J. M. Horowitz, N. Perunov, and J. L. England, Phys. Rev. Lett. 116, 120601 (2016).

- [36] P. Pietzonka, A. C. Barato, and U. Seifert, Phys. Rev. E 93, 052145 (2016).
- [37] P. Pietzonka, F. Ritort, and U. Seifert, Phys. Rev. E 96, 012101 (2017).
- [38] J. M. Horowitz and T. R. Gingrich, Phys. Rev. E 96, 020103 (2017).
- [39] G. Falasco, M. Esposito, and J.-C. Delvenne, New Journal of Physics 22, 053046 (2020).
- [40] J. M. Horowitz and T. R. Gingrich, Nature Physics 16, 15 (2020).
- [41] T. Van Vu, V. T. Vo, and Y. Hasegawa, Phys. Rev. E 101, 042138 (2020).
- [42] T. Van Vu and K. Saito, Physical Review X 13, 011013 (2023).
- [43] I. Di Terlizzi and M. Baiesi, Journal of Physics A: Mathematical and Theoretical 52, 02LT03 (2018).
- [44] N. Shiraishi, Journal of Statistical Physics 185, 19 (2021).
- [45] A. Dechant and S. ichi Sasa, arXiv preprint (2019).
- [46] A. Dechant and S.-i. Sasa, Proceedings of the National Academy of Sciences 117, 6430 (2020).
- [47] G. Falasco, M. Esposito, and J.-C. Delvenne, Journal of Physics A: Mathematical and Theoretical 55, 124002 (2022).
- [48] J. Zheng and Z. Lu, arXiv preprint 10.48550/arXiv.2403.10952 (2024).
- [49] K. Liu and J. Gu, arXiv preprint arXiv:2410.20800 10.48550/arXiv.2410.20800 (2024).
- [50] E. Kwon, H.-M. Chun, H. Park, and J. S. Lee, arXiv preprint arXiv:2411.18108 10.48550/arXiv.2411.18108 (2024).
- [51] T. Van Vu, arXiv preprint arXiv:2411.19546 10.48550/arXiv.2411.19546 (2024).
- [52] K. Ptaszyński, T. Aslyamov, and M. Esposito, Phys. Rev. Lett. 133, 227101 (2024).
- [53] P. Lévy, Compositio Mathematica 7, 283 (1940).
- [54] A. Dhar and S. N. Majumdar, Phys. Rev. E 59, 6413 (1999).
- [55] C. Godreche and J. Luck, Journal of Statistical Physics 104, 489 (2001).
- [56] S. N. Majumdar and A. Comtet, Phys. Rev. Lett. 89, 060601 (2002).
- [57] S. N. Majumdar, in *The legacy of Albert Einstein: A collection of essays in celebration of the year of physics* (World Scientific, 2007) pp. 93–129.
- [58] A. Lapolla and A. Godec, New Journal of Physics 20, 113021 (2018).
- [59] A. Lapolla and A. Godec, Frontiers in Physics 7, 182 (2019).
- [60] A. Lapolla, D. Hartich, and A. Godec, Phys. Rev. Res. 2, 043084 (2020).
- [61] P. Singh and A. Kundu, Journal of Statistical Mechanics: Theory and Experiment 2019, 083205 (2019).
- [62] P. C. Bressloff, Phys. Rev. E 102, 042135 (2020).
- [63] K. Palo, Ü. Mets, V. Loorits, and P. Kask, Biophysical J. 90, 2179 (2006).
- [64] I. V. Gopich and A. Szabo, Proceedings of the National Academy of Sciences 109, 7747 (2012).
- [65] G. Margolin and E. Barkai, Phys. Rev. Lett. 94, 080601 (2005).
- [66] V. G. Ramesh, K. J. H. Peters, and S. R. K. Rodriguez, Phys. Rev. Lett. 132, 133801 (2024).
- [67] W. Bialek and S. Setayeshgar, Proceedings of the National Academy of Sciences 102, 10040 (2005).
- [68] R. G. Endres and N. S. Wingreen, Proceedings of the National Academy of Sciences 105, 15749 (2008).
- [69] T. Mora and N. S. Wingreen, Physical Review Letters 104, 248101 (2010).
- [70] C. C. Govern and P. R. ten Wolde, Phys. Rev. Lett. 109, 218103 (2012).

- [71] C. C. Govern and P. R. ten Wolde, Phys. Rev. Lett. 113, 258102 (2014).
- [72] K. Kaizu, W. De Ronde, J. Paijmans, K. Takahashi, F. Tostevin, and P. R. Ten Wolde, Biophysical J. 106, 976 (2014).
- [73] A. H. Lang, C. K. Fisher, T. Mora, and P. Mehta, Phys. Rev. Lett. 113, 148103 (2014).
- [74] A. C. Barato and U. Seifert, Phys. Rev. E 92, 032127 (2015).
- [75] S. Ray and A. C. Barato, Journal of Physics A: Mathematical and Theoretical 50, 355001 (2017).
- [76] S. E. Harvey, S. Lahiri, and S. Ganguli, Physical Review E 108, 014403 (2023).
- [77] H. C. Berg and E. M. Purcell, Biophysical J. 20, 193 (1977).
- [78] T. Koyuk and U. Seifert, Phys. Rev. Lett. 125, 260604 (2020).
- [79] K. Macieszczak, Occupation uncertainty relations (2024), arXiv:2411.15118 [cond-mat.stat-mech].
- [80] R. Rao and M. Esposito, New Journal of Physics 20, 023007 (2018).
- [81] G. Falasco and M. Esposito, Macroscopic stochastic thermodynamics (2023), arXiv:2307.12406 [cond-mat.stat-mech].
- [82] G. E. Crooks, On the Drazin inverse of the rate matrix (2018), technical note.
- [83] G. T. Landi, M. J. Kewming, M. T. Mitchison, and P. P. Potts, PRX Quantum 5, 020201 (2024).
- [84] K. Ptaszynski and M. Esposito, Critical heat current fluctuations in Curie-Weiss model in and out of equilibrium (2024), arXiv:2411.19643 [cond-mat.stat-mech].
- [85] G. Bakewell-Smith, F. Girotti, M. Guţă, and J. P. Garrahan, Phys. Rev. Lett. 131, 197101 (2023).
- [86] W. Ledermann and G. E. H. Reuter, Philos. Trans. R. Soc. A 246, 321 (1954).

- [87] F. Schlögl, Z. Physik 253, 147 (1972).
- [88] M. Vellela and H. Qian, Journal of The Royal Society Interface 6, 925 (2009).
- [89] H. Ge and H. Qian, Phys. Rev. Lett. 103, 148103 (2009).
- [90] R. Landauer, J. App. Phys. 33, 2209 (1962).
- [91] P. Hänggi and H. Thomas, Phys. Rep. 88, 207 (1982).
- [92] T. Herpich, T. Cossetto, G. Falasco, and M. Esposito, New J. Phys. 22, 063005 (2020).
- [93] J. Meibohm and M. Esposito, Phys. Rev. Lett. 128, 110603 (2022).
- [94] J. Meibohm and M. Esposito, New J. Phys. 25, 023034 (2023).
- [95] H. Vroylandt, M. Esposito, and G. Verley, Europhys. Lett. 120, 30009 (2017).
- [96] H. Vroylandt, M. Esposito, and G. Verley, Phys. Rev. Lett. 124, 250603 (2020).
- [97] G. T. Nieddu, E. Forgoston, and L. Billings, Journal of the Royal Society Interface 19, 20220253 (2022).
- [98] H. Thierschmann, R. Sánchez, B. Sothmann, F. Arnold, C. Heyn, W. Hansen, H. Buhmann, and L. W. Molenkamp, Nat. Nanotechnol. 10, 854 (2015).
- [99] S. Gustavsson, R. Leturcq, B. Simovič, R. Schleser, T. Ihn, P. Studerus, K. Ensslin, D. C. Driscoll, and A. C. Gossard, Phys. Rev. Lett. 96, 076605 (2006).
- [100] S. Gustavsson, R. Leturcq, M. Studer, I. Shorubalko, T. Ihn, K. Ensslin, D. Driscoll, and A. Gossard, Surface Science Reports 64, 191 (2009).