


Regression trees for nonparametric diagnostics of sequential positivity violations in longitudinal causal inference

Arthur Chatton  *¹, Michael Schomaker^{2,3}, Miquel-Angel Luque-Fernandez^{4,5}, Robert W. Platt^{6,7,8}, and Mireille E. Schnitzer^{1,8,9}

¹Faculté de Pharmacie, Université de Montréal, Montréal, QC, Canada

²Department of Statistics, Ludwig-Maximilians University Munich, Munich, Germany

³Centre for Infectious Disease Epidemiology and Research, University of Cape Town, Cape Town, South Africa

⁴Department of Statistics and Operations Research, University of Granada, Granada, Spain

⁵Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK

⁶Center for Clinical Epidemiology, Lady Davis Institute, Jewish General Hospital, Montréal, QC, Canada

⁷Department of Pediatrics, McGill University, Montréal, QC, Canada

⁸Department of Epidemiology, Biostatistics, Occupational Health, McGill University, Montréal, QC, Canada

⁹Département de Médecine Sociale et Préventive, Université de Montréal, Montréal, QC, Canada

Abstract

Sequential positivity is often a necessary assumption for drawing causal inferences, such as through marginal structural modeling. Unfortunately, verification of this assumption can be challenging because it usually relies on multiple parametric propensity score models, unlikely all correctly specified. Therefore, we propose a new algorithm, called *sequential Positivity Regression Tree* (sPoRT), to check this assumption with greater ease under either static or dynamic treatment strategies. This algorithm also identifies the subgroups found to be violating this assumption, allowing for insights about the nature of the violations and potential solutions. We first present different versions of sPoRT based on either stratifying or pooling over time. Finally, we illustrate its use in a real-life application of HIV-positive children in Southern Africa with and without pooling over time. An R notebook showing how to use sPoRT is available at github.com/ArthurChatton/sPoRT-notebook.

Keywords: Causal inference, Data support, Dynamic treatment regime, Longitudinal

*Contact: arthur.chatton@umontreal.ca

study, Positivity, Regression tree.

Introduction

Causal inference is a two-step process.¹ First, we define a causal parameter of interest in a given population based on counterfactual data. For instance, a marginal structural model (MSM) is a popular choice to represent longitudinal treatment effects. Based on a causal model representing knowledge and assumptions about the data-generating process, one can determine whether the counterfactual scientific question of interest (represented by the causal parameter) can be written as a statistical parameter of only the observed data. Next comes the estimation: one uses an estimator of the statistical parameter defined in the first step. Numerous estimators were developed for parameters defined to represent treatment effects in longitudinal settings, notably based on the g-formula,^{2,3} inverse probability weighting,⁴ and doubly robust estimators, such as the longitudinal targeted maximum likelihood estimator.^{5,6} Tutorials of various levels of technicality on these estimators are available.⁷⁻¹⁰

While causal effects in longitudinal settings were defined and estimators proposed by Robins in the mid-80s,² later works investigated how to deal with common statistical challenges in practice. One major issue in longitudinal causal inference is violations (or near-violations) of the empirical positivity assumption, which requires that all possible confounder patterns are observed following all relevant treatments at all time points.¹¹ Model pooling and smoothing methods allow for a relaxation of this requirement.¹² Statistical covariate reduction methods can also be helpful.¹³ To diagnose the presence of sequential positivity (near-)violations, one may identify covariate-defined subgroups with null (or nearly null) estimated probabilities of some intervention trajectory's level. Thus, the diagnosis of positivity violations is complicated because positivity must be checked across all time points and may rely on many propensity score models, unlikely all correctly specified.^{8,14} This is a direct extension of the positivity assumption in single time-point treatment settings.¹⁵

The *Positivity Regression Trees* (PoRT) algorithm¹⁶ is a regression tree-based explanatory tool to check positivity in single time-point treatment settings that require neither assumptions about the data-generating process nor specification of parametric models. In this study, we propose an extension of the PoRT method to longitudinal settings – sequential PoRT or “sPoRT” – to identify the time-updated subgroups of individuals (defined by baseline and time-dependent

covariate patterns), which lack data support for some category of treatment. We demonstrate the potential of this approach by reanalyzing a study investigating the effect of delaying HIV antiretroviral therapy (ART) among children in Southern Africa.⁸

The PoRT algorithm for single time-point treatment settings

First, consider the following data structure for a single time-point treatment where Y is the outcome, A is the binary treatment, and W is the confounder set. The PoRT algorithm runs a succession of regression trees, where the treatment is regressed on one or more confounders.¹⁶ An overview of the algorithm is as follows. First, fit individual trees by regressing the treatment on each confounder (*i.e.* one tree per confounder). Second, search through each tree for nodes (*i.e.*, subgroups of individuals) with an extreme probability of receiving a treatment level according to two user-defined hyperparameters (α and β , see Table 1). Third, record the problematic subgroups identified in the second step and remove the related covariate(s) from further consideration. After the third step, repeat the above three steps, except this time, increase the number of covariates in each tree by one. The algorithm stops once the user-supplied value of the subgroups’ allowed complexity (hyperparameter γ , Table 1) is exceeded.

Table 1: PoRT main hyperparameters

Hyperparameter	Definition	Suggested default values ¹
α	Minimal proportion of the whole sample size to consider a problematic subgroup	0, 0.05, 0.1
β	Threshold defining positivity violations: $P(A = a W) > \beta$, for all relevant a	0.01, 0.05, $5/(\sqrt{n} \ln n)$
γ	Maximal number of covariates in a tree	2, 3, all

Notation: A , Treatment; n , sample size; W , covariates. ¹ Must be based on domain expertise.

The hyperparameter $\alpha \geq 0$ controls the size of identified subgroups by excluding the smallest ones. Positivity violations in very small subgroups may not impact the estimation when the model smooths over the subpopulation.^{15,17} The hyperparameter β defines the desired level of positivity, such that $P(A = a|W) \leq \beta$ is considered a violation. It represents a threshold where the treatment probability becomes too extreme for the treatment effect in a subgroup to be supported by the data.¹⁸ Recently, Gruber *et al.* proposed a new optimal bound of $5/(\sqrt{n} \ln n)$ to truncate the propensity score,¹⁹ which can be applied as the default value of β . The last hyperparameter, γ , represents the maximum number of covariates used to build a regression tree. For instance, when γ is set to two, the resulting subgroups are defined by a maximum of two covariates (*e.g.*, male patients older than 70 years). Its optimal value depends on the treatment allocation mechanism’s complexity. According to the positivity assumption,

W should include all variables (confounders or not) that are part of the adjustment set used for future estimation.²⁰ The hyperparameter γ allows us to reduce the allowed interactions to better interpret the subgroups. Furthermore, clinically meaningful violations may be missed when the full set W is used to build a tree.¹⁶

PoRT's strength relies on its usage of regression trees. Regression trees are one of the most interpretable data-driven methods,²¹ making their output ideal for obtaining clinically interpretable subgroups defined by covariate values. They also can adapt to non-linear functional forms and incorporate higher-order interactions between covariates.²² Trees' weaknesses are their tendency to overfit and the greedy categorization of continuous variables. PoRT overcomes the first issue by fitting the simplest regression trees first and increasing their complexity sequentially through γ . When applying PoRT, the authors suggested manually categorizing the continuous confounders using meaningful thresholds since the interpretability of the subgroups is key,¹⁶ which also circumvents the second issue.

Motivating example in longitudinal settings

ART is effective for reducing infant mortality,^{23,24} causing the WHO to recommend immediate ART initiation in all HIV-positive children as of 2015.²⁵

Schomaker *et al.* investigated the effect of different ART initiation rules on the child's growth measured by height-for-age z-scores (HAZ) in West and Southern Africa using the observational International epidemiological Databases to Evaluate AIDS (IeDEA) in sub-Saharan Africa.^{8,26-29} They highlight the severity of positivity violations encountered in their analysis. We reanalyse a component of this study to demonstrate how PoRT can be used to identify sequential positivity violations.

Our reanalysis includes data from South Africa, Malawi, and Zimbabwe. HIV-positive children aged 12-59 months at the first measured contact with health care facility (which is the baseline) who acquired the virus before or at birth or during breastfeeding were included. To be eligible for the analysis, children had at least one visit before ART initiation, complete data at enrolment (baseline), and at least one follow-up clinic visit. In total, 2352 children were included in the analysis. Data were recorded at 0, 1, 3, 6, 9, and 12 months after enrolment ($t = 0, 1, 2, 3, 4, 5$ respectively). At nine months, 1893 children were still followed. The most relevant parts of the

posited causal structure from the initial study are depicted in Figure 1. More details regarding censoring and relevant baseline unmeasured confounders are given in the initial study.⁸ The outcome Y_t is HAZ measured at time t ; the treatment A_t refers to whether ART was taken at time t ; the time-varying confounders set L_t includes CD4 cell count, CD4% (*i.e.*, CD4/white blood cell counts $\times 100$), and clinical stage, approximated by weight for age z-score (WAZ);²⁷ the time-fixed confounders set W includes biological sex, age, and year of treatment initiation. Y_0 is the baseline HAZ, and it is included in the adjustment set.




Figure 1: Simplified posited data-generating process for the first time points. Double-headed arrows indicated an effect on all downstream nodes. $A_t; t = 1, 2, 3, 4$ are the treatment status at each time point, $L_t; t = 0, 1, 2, 3, 4$ are the time-varying confounder sets, $Y_t; t = 0, 2, 3, 4, 5$ are the repeatedly measured outcomes. W is the baseline confounder set, and U are potential unmeasured confounders.

Treatment and treatment strategy

Time-varying treatment complicates the estimand definition. Furthermore, data support may vanish over time due to the potentially large number of possible treatment trajectories. Focusing on MSM parameters,³⁰ if treatment can change at any of T time-points, there can be as many as 2^T trajectories. In our example, the treatment, which represents ART initiation, is monotone such that $A_t = 1$ when $A_{t-1} = 1$ (see Figure 2, magenta flow).

Treatment strategies can be used in longitudinal settings to define relevant causal contrasts flexibly. A treatment strategy is a rule that assigns the treatment at each time t of follow-up.³¹




Figure 2: Switches between antiretroviral therapy (ART) usage. Immediate and no initiation are highlighted in magenta and green, respectively. Initiation is highlighted in blue.

To avoid confusion with observed treatment, we will now use the term *intervention rule* for treatment strategy. A *dynamic* rule depends on the value of one or more covariates. Otherwise, the rule is *static*. Common examples of static rules are “always treat” and “never treat”. Dynamic rules are sometimes more realistic or even essential in practice. For instance, when some patients develop a contraindication to a specific treatment, they can no longer receive it and must switch to another one, which any realistic intervention rule must respect.³² We denote an intervention rule at time t by d_t .

We considered two static rules: immediate ART initiation and continued usage (rule 1, $d_t = 1$ for $t = 1, 2, 3, 4$) or no ART initiation at any time (rule 2, $d_t = 0$ for $t = 1, 2, 3, 4$). We also considered two dynamic intervention rules consisting of deferring ART initiation until a particular CD4 threshold is reached (similar to WHO’s 2006 and 2010 guidelines²⁹). The rules are to start treatment ($d_t = 1$) when (i) CD4 count < 750 or CD4% $< 25\%$ (rule 3), or (ii) CD4 count < 350 or CD4% $< 15\%$ (rule 4). For these dynamic rules, we can write $d_t(L_t)$ since the rules are a function of the time-updated covariates. Of note, ART is always continued once initiated to respect the observed treatment monotonicity. In this demonstration, we do not diagnose sparsity in the censoring process, although this would be possible.

sPoRT for interventions at multiple time points

Sequential positivity implies that all individuals in the population have a non-zero probability of continuing to follow a given assigned intervention strategy at each time point, given that they have followed it thus far.⁸ However, the way one may wish to check sequential positivity may vary according to data pooling or smoothing decisions used in the estimation. Rudolph *et al.* illustrated the advantages and pitfalls of different levels of smoothing with respect to positivity.¹⁴ Briefly, stratifying over time and/or treatment patterns improves the flexibility of the estimation when the model parameters vary significantly over time or across treatment histories. However, this stratification increases the number of models to fit, while reducing the amount of data available for their fit. Random positivity violations (or data sparsity) are more likely to occur, yielding either estimation issues or the strong assumption of a correct extrapolation over missing strata.¹⁷ Pooling over time-points and/or treatment histories increases the stability of the estimation by positing a smoothed model that shares its parameters over time or across different treatment histories. However, smoothing over heterogeneous parameters may cause bias, even if the precision increases.³³

Our sPoRT algorithm can be used to check the sequential positivity assumption under either stratification or pooling over time. If stratification is desired, the PoRT process described in Section is applied to each time point independently of each other, using a data subset depending on the intervention rule being investigated. The analyst must specify the time-varying confounders, the time-varying rules indicators, the hyperparameters, and any additional subsetting used for the estimation. The sPoRT algorithm can estimate the relevant treatment probabilities according to the observed treatment pattern (monotone or non-monotone) and the intervention rule type (Table 2). Briefly, observed treatment monotonicity requires subsetting of individuals who have not initiated (or ceased) the treatment at the previous time. Dynamic intervention rules imply checking positivity of continuing to follow the rule, i.e., checking data support for $A_t = d_t(L_t)$ where $d_t(L_t)$ can be 0 or 1 depending on covariates, among individuals who previously complied with the rule. However, a common occurrence is that the data does not support such a stratification over treatment history due to decreasing sample sizes over time. In that case, one can smooth over treatment history, assuming the model and the covariates can correct for the lack of subsetting.

Table 2: Illustration of typical probabilities modeled post-baseline in sPoRT using two time-points

Rule	Pattern of A	Probability modeled
Static	Monotone	$P(A_2 = 1 A_1 = 0, L_2, W, d_2 = 1) > \beta$ $P(A_2 = 0 A_1 = 0, L_2, W, d_2 = 0) > \beta$
	Non-monotone	$P(A_2 = 1 A_1, L_2, W, d_2 = 1) > \beta$ $P(A_2 = 0 A_1, L_2, W, d_2 = 0) > \beta$
Dynamic	Monotone	$P(A_2 = 1 A_1 = 0, L_2, W, d_2(L_2) = 1) > \beta$ $P(A_2 = 0 A_1 = 0, L_2, W, d_2(L_2) = 0) > \beta$
	Non-monotone	$P(A_2 = 1 A_1, L_2, W, d_2(L_2) = 1) > \beta$ $P(A_2 = 0 A_1, L_2, W, d_2(L_2) = 0) > \beta$

Notation: A_t , treatment; $d_t(L_t)$, indicator of whether the rule recommends treatment; L_t , time-varying confounders; W , baseline confounders; β , positivity bound. Subscript indicates time-point. Note: Smoothing over treatment history is applied in the above. Different subsetting can be used depending on the specific context. Monotone refers to a non-decreasing sequence, without loss of generality.

Application results

We ran sPoRT with stratification on time with the following hyperparameters: $\alpha = 0.05$, $\beta = 5/(\sqrt{n_t} \ln n_t)$, where n_t is the sample size of the subset used to fit the trees at time t , and $\gamma = 2$. A detailed explanation of its use with R code is provided in Supplementary material (see also the online notebook that allows for updates in the future). Stratification and smoothing over treatment history were allowed for static and dynamic rules, respectively. It did not find any positivity violation at $t = 1$. We did not check positivity further for the first static rule of immediate initiation because the observed treatment is monotone (ART is not stopped once initiated, as illustrated in Figure 2). No violation was identified for the never initiate ART static rule.

Positivity was challenged for initiating treatment under the two dynamic rules starting from six months (Tables 3 and 4). The third rule of deferring ART with higher thresholds presented two violations at six months and thirteen violations at nine months. The fourth rule, deferring ART with lower thresholds, presented only one and seven violations at months 6 and 9, respectively. In contrast, no positivity violation was identified when pooling over time, regardless of the rule. Many of the detected violations highlight the fact that it was unlikely for very healthy children to initiate ART immediately after becoming formally eligible according to older treatment

rules.

These positivity violations may occur for three reasons, each with a different implication for the analysis. First, sparsity is more frequent in longitudinal settings due to censoring. Furthermore, the monotone treatment pattern reduces the number of untreated individuals over time. For instance, there were only 12 untreated individuals eligible for the fourth rule at nine months with baseline CD4% lower than 15% and a CD4 count between 100 and 225. In the case of sparsity, smoothing should be considered by pooling over time or using estimators able to extrapolate in the missing strata. Second, physicians may have deferred ART initiation for healthier children. Positivity violations will exist when the definition of “healthier” differs from those used in the intervention rule. In our data, ART was initiated according to the WHO 2010 guideline (i.e., the third rule),³⁴ which can explain some other rules’ violations. Careful thinking about the definition of the intervention rules, and making sure that they fall within the variability of clinical practice at the time of data collection, will help to avoid them. Last, structural positivity violations may be present. For instance, HIV-positive pediatric ART coverage was estimated at around 33% in Southern Africa in 2013³⁵ and some unmeasured factors, such as socioeconomic status or nutrition, can lead to or be related to very low rates of ART initiation in some subgroups. Quality of data and expert knowledge are crucial for this last case.

Discussion

In this paper, we described sPoRT, a new algorithm for checking sequential positivity by identifying possibly overlapping subgroups of individuals without support for the defined intervention strategies. The algorithm can incorporate different types of data pooling, matching the desired level of smoothing in the analysis. We implemented the algorithm in the R package PoRT to facilitate its use and provide an R notebook demonstrating its application.

The standard way to check sequential positivity is to compute several propensity scores and use the cumulative product of their resulting weights as a measure of data support, a tiny cumulative weight being a red flag for non-positivity.^{5,8} However, this approach (i) is based on the assumption that all the propensity scores usually fitted using parametric regressions are correctly specified and (ii) does not directly identify problematic subgroups of individuals. Indeed, although individuals with near-zero predicted probabilities of following strategies are easily identifiable, their shared characteristics differing from the remaining individuals may not

be obvious. In contrast, sPoRT provides clear subgroups defined by the values of one or more covariates, possibly at each time point when stratification on time is desired. Using regression trees avoids the parametric assumptions of modeling the propensity scores, adding robustness to the sequential positivity checking. We recommend routine usage of both approaches.

However, we stress that both the fitting of MSMs and usage of sPoRT require the correct a priori identification of the confounders and time-ordering of all variables. Only domain knowledge can inform this. Furthermore, sPoRT depends on its tuning parameters α , β and γ (and those specific to the regression tree to a lesser extent). A grid of pragmatically informed values (as suggested in Table 1) can be considered for these parameters.¹⁶ Unlike Danelian *et al.*,¹⁶ we adapted β automatically to the sample size at a given time using Gruber’s bound.¹⁹ Since the bound is a function of the sample size, it naturally accounts for the decreasing sample size over time in longitudinal contexts. However, this bound was initially defined for cross-sectional studies; thereby, the optimal bound function could slightly differ in longitudinal settings.

Finally, what should be done with the sPoRT output? We recommend checking first whether some patterns regularly appear over time and the subgroups from the first time point to consider potential structural positivity violations. If some subgroups are likely due to structural violations at baseline, they must be excluded by adding exclusion criteria to the study, and then positivity should be rechecked to see if other violations on the variables involved in these subgroups were hidden. Structural positivity violations occurring over time can also be dealt with by adapting the intervention strategy to account for these violations.³² Once the structural violations have been ruled out, sparsity can be investigated through the remaining subgroups. The violations and subgroup sizes may influence the choice of the estimator. For instance, inverse probability weighting is highly sensitive to sparsity.³⁶ In contrast, estimators able to extrapolate over missing strata could be considered, such as the g-formula.³⁷ When the sparsity is high, alternative estimands could also be considered; see Hoffman *et al.* for an overview.³⁸

Competing interests

No competing interest is declared.

Author contributions statement

AC: Conceptualization, Methodology, Software, Writing - Original Draft, and Writing - Review & Editing. MS: Data curation, Methodology, and Writing - Review & Editing. MALF: Writing - Review & Editing. RWP: Supervision and Writing - Review & Editing. MES: Methodology, Supervision, and Writing - Review & Editing.

Acknowledgments

A.C. is supported by an IVADO postdoctoral fellowship (#2022-7820036733). MES holds a tier 2 Canada Research Chair in Causal Inference and Machine Learning in Health Science and a Discovery Grant (#RGPIN-2021-03019) from the Natural Sciences and Engineering Research Council of Canada. This project was approved by the Université de Montréal clinical ethics committee (#2021-359). We also thank Shobna Sawry, Karl-Günter Technau, Frank Tanser, Haby Sam Phiri, Vivian Cox, Cleophas Chimbe, Janet Giddy, and Robin Wood for sharing their data with us. We would also like to highlight the support of the International epidemiologic Databases to Evaluate AIDS Southern Africa (IeDEA-SA), led by Matthias Egger and Mary-Ann Davies; the NIH has supported the collaboration, grant numbers #U01AI069924.

References

- ¹ Petersen ML, van der Laan MJ. Causal Models and Learning from Data: Integrating Causal Modeling and Statistical Estimation. *Epidemiology*. 2014;25(3):418–426.
- ² Robins JM. A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Mathematical Modelling*. 1986;7(9):1393–1512.
- ³ Bang H, Robins JM. Doubly robust estimation in missing data and causal inference models. *Biometrics*. 2005;61(4):962–973.
- ⁴ Robins JM, Hernán MA, Brumback B. Marginal structural models and causal inference in epidemiology. *Epidemiology*. 2000;11(5):550–560.

- ⁵ Petersen M, Schwab J, Gruber S, Blaser N, Schomaker M, van der Laan MJ. Targeted Maximum Likelihood Estimation for Dynamic and Static Longitudinal Marginal Structural Working Models. *Journal of Causal Inference*. 2014;2(2):147–185.
- ⁶ Schnitzer ME, van der Laan MJ, Moodie EEM, Platt RW. Effect of breastfeeding on gastrointestinal infection in infants: a targeted maximum likelihood approach for clustered longitudinal data. *The Annals of Applied Statistics*. 2014;8(2):703–725.
- ⁷ Daniel RM, Cousens SN, De Stavola BL, Kenward MG, Sterne JAC. Methods for dealing with time-dependent confounding. *Statistics in Medicine*. 2013;32(9):1584–1618.
- ⁸ Schomaker M, Luque-Fernandez MA, Leroy V, Davies MA. Using longitudinal targeted maximum likelihood estimation in complex settings with dynamic interventions. *Statistics in Medicine*. 2019;38(24):4888–4911.
- ⁹ Naimi AI, Cole SR, Kennedy EH. An Introduction to G Methods. *International Journal of Epidemiology*. 2016;46(2):756–762.
- ¹⁰ Hernán M, Robins JM. *Causal Inference: What if?* Boca Raton: Chapman & Hall/CRC; 2020.
- ¹¹ Orellana L, Rotnitzky A, Robins JM. Dynamic regime marginal structural mean models for estimation of optimal dynamic treatment regimes, Part II: proofs of results. *The International Journal of Biostatistics*. 2010;6(2):Article 9.
- ¹² Cole SR, Hernán MA. Constructing Inverse Probability Weights for Marginal Structural Models. *American Journal of Epidemiology*. 2008;168(6):656–664.
- ¹³ Schnitzer ME, Sango J, Ferreira Guerra S, van der Laan MJ. Data-adaptive longitudinal model selection in causal inference with collaborative targeted minimum loss-based estimation. *Biometrics*. 2020;76(1):145–157.
- ¹⁴ Rudolph JE, Benkeser D, Kennedy EH, Schisterman EF, Naimi AI. Estimation of the Average Causal Effect in Longitudinal Data with Time-Varying Exposures: The Challenge of Nonpositivity and the Impact of Model Flexibility. *American Journal of Epidemiology*. 2022;191(11):1962–1969.

- ¹⁵ Petersen ML, Porter KE, Gruber S, Wang Y, van der Laan MJ. Diagnosing and responding to violations in the positivity assumption. *Statistical Methods in Medical Research*. 2012;21(1):31–54.
- ¹⁶ Danelian G, Foucher Y, Léger M, Le Borgne F, Chatton A. Identification of in-sample positivity violations using regression trees: The PoRT algorithm. *Journal of Causal Inference*. 2023;11(1):20220032.
- ¹⁷ Léger M, Chatton A, Le Borgne F, Pirracchio R, Lasocki S, Foucher Y. Causal inference in case of near-violation of positivity: comparison of methods. *Biometrical Journal*. 2022;64(8):1389–1403.
- ¹⁸ D’Amour A, Ding P, Feller A, Lei L, Sekhon J. Overlap in observational studies with high-dimensional covariates. *Journal of Econometrics*. 2021;221(2):644–654.
- ¹⁹ Gruber S, Phillips RV, Lee H, van der Laan MJ. Data-Adaptive Selection of the Propensity Score Truncation Level for Inverse-Probability-Weighted and Targeted Maximum Likelihood Estimators of Marginal Point Treatment Effects. *American Journal of Epidemiology*. 2022;191(9):1640–1651.
- ²⁰ Westreich D. *Epidemiology by design: a causal approach to the health sciences*. Oxford University Press; 2020.
- ²¹ Breiman L. Statistical Modeling: The Two Cultures (with comments and a rejoinder by the author). *Statistical Science*. 2001;16(3):199–231.
- ²² Bi Q, Goodman KE, Kaminsky J, Lessler J. What is Machine Learning? A Primer for the Epidemiologist. *American Journal of Epidemiology*. 2019;188(12):2222–2239.
- ²³ Violari A, Cotton MF, Gibb DM, Babiker AG, Steyn J, Madhi SA, et al. Early antiretroviral therapy and mortality among HIV-infected infants. *The New England Journal of Medicine*. 2008;359(21):2233–2244.
- ²⁴ Edmonds A, Yotebieng M, Lusiana J, Matumona Y, Kitetele F, Napravnik S, et al. The Effect of Highly Active Antiretroviral Therapy on the Survival of HIV-Infected Children in a Resource-Deprived Setting: A Cohort Study. *PLoS Medicine*. 2011;8(6):e1001044.
- ²⁵ World Health Organization. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: recommendations for a public health approach.

Geneva; 2016. Available from: <https://www.who.int/publications-detail-redirect/9789241549684>.

- ²⁶ Egger M, Ekouevi DK, Williams C, Lyamuya RE, Mukumbi H, Braitstein P, et al. Cohort Profile: The international epidemiological databases to evaluate AIDS (IeDEA) in sub-Saharan Africa. *International Journal of Epidemiology*. 2012;41(5):1256–1264.
- ²⁷ Schomaker M, Egger M, Ndirangu J, Phiri S, Moultrie H, Technau K, et al. When to Start Antiretroviral Therapy in Children Aged 2–5 Years: A Collaborative Causal Modelling Analysis of Cohort Studies from Southern Africa. *PLOS Medicine*. 2013;10(11):e1001555.
- ²⁸ Schomaker M, Davies MA, Malateste K, Renner L, Sawry S, N’Gbeche S, et al. Growth and Mortality Outcomes for Different Antiretroviral Therapy Initiation Criteria in Children Ages 1–5 Years: A Causal Modeling Analysis. *Epidemiology*. 2016;27(2):237–246.
- ²⁹ Schomaker M, Leroy V, Wolfs T, Technau KG, Renner L, Judd A, et al. Optimal timing of antiretroviral treatment initiation in HIV-positive children and adolescents: a multiregional analysis from Southern Africa, West Africa and Europe. *International Journal of Epidemiology*. 2017;46(2):453–465.
- ³⁰ Petersen ML. Applying a Causal Road Map in Settings with Time-dependent Confounding. *Epidemiology*. 2014;25(6):898–901.
- ³¹ Smith LH, García-Albéniz X, Chan JM, Zhao S, Cowan JE, Broering JM, et al. Emulation of a target trial with sustained treatment strategies: an application to prostate cancer using both inverse probability weighting and the g-formula. *European Journal of Epidemiology*. 2022;37(12):1205–1213.
- ³² Schnitzer ME, Platt RW, Durand M. A tutorial on dealing with time-varying eligibility for treatment: Comparing the risk of major bleeding with direct-acting oral anticoagulants vs warfarin. *Statistics in Medicine*. 2020;39(29):4538–4550.
- ³³ Rudolph JE, Cole SR, Edwards JK. Parametric assumptions equate to hidden observations: comparing the efficiency of nonparametric and parametric models for estimating time to AIDS or death in a cohort of HIV-positive women. *BMC Medical Research Methodology*. 2018;18(1):142.

- ³⁴ World Health Organization. Antiretroviral therapy of HIV infection in infants and children: towards universal access: recommendations for a public health approach - 2010 revision. Geneva; 2010. Available from: <https://iris.who.int/handle/10665/164255>.
- ³⁵ Rosen JG, Muraleetharan O, Walker A, Srivastava M. Pediatric Antiretroviral Therapy Coverage and AIDS Deaths in the “Treat All” Era. *Pediatrics*. 2023;151(6):e2022059013.
- ³⁶ Spreafico M. Positivity violations in marginal structural survival models with time-dependent confounding: a simulation study on IPTW-estimator performance. arXiv (not peer-reviewed). 2024;2403.19606. Available from: <http://arxiv.org/abs/2403.19606>.
- ³⁷ Robins J, Sued M, Lei-Gomez Q, Rotnitzky A. Comment: Performance of Double-Robust Estimators When “Inverse Probability” Weights Are Highly Variable. *Statistical Science*. 2007;22(4):544–559.
- ³⁸ Hoffman KL, Salazar-Barreto D, Williams NT, Rudolph KE, Díaz I. Studying Continuous, Time-varying, and/or Complex Exposures Using Longitudinal Modified Treatment Policies. *Epidemiology*. 2024;35(5):667.

Table 3: Positivity violations identified with the stratified sPoRT for the third rule (dynamic): deferring ART with higher thresholds

Subgroup	Prob.	n^* (%)
$P(A_t = 1 A_{t-1} = 0, L_t, W, d_t(L_t) = 1)$		
One month ($n_1 = 2060, \beta = 0.014$)		
No violation was found	-	-
Three months ($n_2 = 1346, \beta = 0.019$)		
No violation was found	-	-
Six months ($n_3 = 794, \beta = 0.027$)		
CD4 _{c0} < 1600 & CD4% ₀ ≥ 30	0.023	43 (5.4)
CD4 _{c3} < 2025 & CD4% ₀ ≥ 30	0.025	40 (5.0)
Nine months ($n_4 = 565, \beta = 0.033$)		
CD4% ₄ ≥ 25 & age = 4	0.029	34 (6.0)
CD4% ₄ ∈ [25, 30[& age < 3	0.031	32 (5.7)
CD4% ₄ ≥ 25 & CD4 _{c0} ∈ [625, 900[0.020	50 (8.8)
CD4% ₄ ∈ [20, 25[& CD4 _{c0} ∈ [900, 1225[0.000	35 (6.2)
HAZ ₀ ≥ -1 & CD4% ₀ ≥ 25	0.032	31 (5.5)
CD4 _{c4} ∈ [625, 900[& CD4% ₀ ∈ [25, 35[0.024	42 (7.4)
CD4 _{c4} ∈ [1225, 1600[& CD4% ₀ ∈ [10, 25[0.030	33 (5.8)
HAZ ₀ ∈ [-4, -1[& WAZ ₀ ≥ 0	0.024	41 (7.3)
CD4% ₄ ∈ [25, 30[& CD4% ₀ ≥ 25	0.024	42 (7.4)
CD4 _{c4} ∈ [1225, 1600[& WAZ ₀ ≥ -2	0.030	33 (5.8)
CD4% ₄ ≥ 25 & WAZ ₀ ∈ [-2, 0[0.032	63 (11.2)
CD4% ₄ ∈ [20, 25[& WAZ ₀ ≥ 0	0.000	29 (5.1)
CD4% ₄ ≥ 25 & HAZ ₀ ∈ [-2, 1[0.000	33 (5.8)
$P(A_t = 0 A_{t-1} = 0, L_t, W, d_t(L_t) = 0)$		
One month ($n_1 = 292, \beta = 0.052$)		
No violation was found	-	-
Three months ($n_2 = 206, \beta = 0.065$)		
No violation was found	-	-
Six months ($n_3 = 137, \beta = 0.087$)		
No violation was found	-	-
Nine months ($n_4 = 112, \beta = 0.100$)		
No violation was found	-	-

Abbreviations: β , Gruber bound; CD4c, CD4 count; HAZ, height for age z-score; n_t , sample size at time t ; n^* , subgroup size; Prob., probability of being treated; WAZ, weight for age z-score. Subscript indicates time-point. Smoothing over treatment history was considered.

Table 4: Positivity violations identified with the stratified sPoRT for the fourth rule (dynamic): deferring ART with lower thresholds

Subgroup	Prob.	n^* (%)
$P(A_t = 1 A_{t-1} = 0, L_t, W, d_t(L_t) = 1)$		
One month ($n_1 = 1164, \beta = 0.021$)		
No violation was found	-	-
Three months ($n_2 = 720, \beta = 0.028$)		
No violation was found	-	-
Six months ($n_3 = 354, \beta = 0.045$)		
HAZ ₀ ≥ -1 & Male	0.038	26 (7.3)
Nine months ($n_4 = 226, \beta = 0.061$)		
WAZ ₄ $\in [-2, -1[$ & age ≥ 3	0.048	42 (18.6)
CD4 _{c0} $\in [100, 225[$ & CD4% ₀ < 15	0.000	12 (5.3)
CD4 _{c4} < 400 & CD4 _{c0} $\in [100, 225[$	0.000	12 (5.3)
CD4 _{c0} < 400 & WAZ ₄ $\in [-2, -1[$	0.034	26 (12.8)
CD4 _{c4} $\in [100, 225[$ & CD4% ₀ < 15	0.000	15 (6.6)
WAZ ₄ $\in [-2, -1[$ & CD4% ₀ $\in [10, 15[$	0.043	23 (10.2)
CD4 _{c4} $\in [100, 400[$ & HAZ ₀ $\in [-3, -2[$	0.056	18 (8.0)
$P(A_t = 0 A_{t-1} = 0, L_t, W, d_t(L_t) = 0)$		
One month ($n_1 = 1188, \beta = 0.020$)		
No violation was found	-	-
Three months ($n_2 = 832, \beta = 0.026$)		
No violation was found	-	-
Six months ($n_3 = 577, \beta = 0.033$)		
No violation was found	-	-
Nine months ($n_4 = 451, \beta = 0.039$)		
No violation was found	-	-

Abbreviations: β , Gruber bound; CD4c, CD4 count; HAZ, height for age z-score; n_t , sample size at time t ; n^* , subgroup size; Prob., probability of being treated; WAZ, weight for age z-score. Subscript indicates time-point. Smoothing over treatment history was considered.