QUANTIFIED CRAMÉR-WOLD CONTINUITY THEOREM FOR THE KANTOROVICH TRANSPORT DISTANCE

SERGEY G. BOBKOV^{1,3} AND FRIEDRICH GÖTZE^{2,3}

ABSTRACT. An upper bound for the Kantorovich transport distance between probability measures on multidimensional Euclidean spaces is given in terms of transport distances between one dimensional projections. This quantifies the Cramér-Wold continuity theorem for the weak convergence of probability measures.

1. Introduction

Given a sequence of random vectors $(X_n)_{n\geq 1}$ and a random vector X with values in \mathbb{R}^d , the Cramér-Wold continuity theorem indicates that $X_n \Rightarrow X$ weakly in distribution, if and only if this convergences holds true for all one dimensional projections, i.e. if and only if

 $\langle X_n, \theta \rangle \Rightarrow \langle X, \theta \rangle$ as $n \to \infty$

on the real line for any $\theta \in \mathbb{R}^d$ (cf. [4], [1]). Of large interest is the problem of how one can quantify this characterization by means of various distances responsible for the weak convergence. Indeed, this could potentially reduce a number of high dimensional questions to dimension one, perhaps under proper moment assumptions. Here we consider the problem with respect to the Kantorovich transport distance.

Let X and Y be random vectors in \mathbb{R}^d with distributions μ and ν having finite first absolute moments. The Kantorovich transport distance, also called the minimal distance between μ and ν , is defined with respect to the Euclidean metric on \mathbb{R}^d by

$$W(X,Y) = W(\mu,\nu) = \inf \mathbb{E} |X' - Y'| = \inf \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |x - y| \, d\pi(x,y).$$
(1.1)

Here the first infimum is taken over all pairs of random vectors X', Y' with distributions μ, ν , and the second one is running over all Borel probability measures π on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals μ and ν . By a well-known characterization of the convergence in W, this metric metrizes the topology of weak convergence in the space of all Borel probability measures μ on \mathbb{R}^d with bounded *p*-th absolute moments for any fixed p > 1 (cf. [10], Theorem 7.12).

²⁰¹⁰ Mathematics Subject Classification. Primary 60E, 60F.

Key words and phrases. Cramér-Wold continuity theorem, transport inequalities.

¹⁾ School of Mathematics, University of Minnesota, Minneapolis, MN, USA, bobkov@math.umn.edu.

²⁾ Faculty of Mathematics, Bielefeld University, Germany, goetze@math-uni.bielefeld.de.

³⁾ Research supported by the NSF grant DMS-2154001, the GRF – SFB 1283/2 2021 - 317210226, and the Hausdorff Research Institute for Mathematics.

According to the Kantorovich duality theorem (cf. [6]),

$$W(X,Y) = \sup_{\|u\|_{\text{Lip}} \le 1} |\mathbb{E}u(X) - \mathbb{E}u(Y)|$$

$$= \sup_{\|u\|_{\text{Lip}} \le 1} \left| \int_{\mathbb{R}^d} u \, d\mu - \int_{\mathbb{R}^d} u \, d\nu \right|, \qquad (1.2)$$

where the supremum runs over all functions $u : \mathbb{R}^d \to \mathbb{R}$ with Lipschitz semi-norm $||u||_{\text{Lip}} \leq 1$. Since the functions of the form $u(x) = v(\langle x, \theta \rangle)$ with $|\theta| = 1$ and $v : \mathbb{R} \to \mathbb{R}$ such that $||v||_{\text{Lip}} \leq 1$ participate in the supremum (1.2), we have

$$W(X,Y) \ge \sup_{|\theta|=1} W(X_{\theta}, Y_{\theta})$$
(1.3)

for the linear functionals

$$X_{\theta} = \langle X, \theta \rangle, \ Y_{\theta} = \langle Y, \theta \rangle.$$

Recall that in dimension one, in big contrast to the multidimensional situation, the Kantorovich distance has a simple description

$$W(X_{\theta}, Y_{\theta}) = \int_{-\infty}^{\infty} |F_{\theta}(x) - G_{\theta}(x)| \, dx$$

in terms of the distribution functions $F_{\theta}(x) = \mathbb{P}\{X_{\theta} \leq x\}, G_{\theta}(x) = \mathbb{P}\{Y_{\theta} \leq x\}.$

Here we reverse the inequality (1.3) in a somewhat similar form under a p-th moment assumption.

Theorem 1.1. Suppose that $(\mathbb{E}|X|^p)^{1/p} \leq b$ and $(\mathbb{E}|Y|^p)^{1/p} \leq b$ for some p > 1 and $b \geq 0$. Then

$$W(X,Y) \le 12 b^{1-\alpha} \sup_{|\theta|=1} W(X_{\theta}, Y_{\theta})^{\alpha}, \qquad (1.4)$$

where

$$\alpha = \frac{2}{dp^* + 2}, \quad p^* = \frac{p}{p-1}.$$
(1.5)

Note that the distance W is homogeneous with respect to (X, Y), and so is (1.4) when this inequality is written with an optimal value of b.

Letting $p \to \infty$ and assuming that $|X| \le 1$ and $|Y| \le 1$ a.s., we obtain a simpler relation

$$W(X,Y) \le 12 \sup_{|\theta|=1} W(X_{\theta}, Y_{\theta})^{\frac{2}{d+2}}.$$
 (1.6)

As another interesting case, suppose that $\mathbb{E} |X|^2 \leq d$ and $\mathbb{E} |Y|^2 \leq d$ (which holds for isotropic distributions). Then (1.4) yields

$$W(X,Y) \le 12\sqrt{d} \sup_{|\theta|=1} W(X_{\theta},Y_{\theta})^{\frac{1}{d+1}}.$$

2. Transport Distances and Convergence of Empirical Measures

It is not clear whether or not the exponent $\alpha = \alpha(p, d)$ defined in (1.5) is optimal in the inequality (1.4), even if one can add an additional (p, d)-dependent multiplicative factor. In order to illustrate the strength of the inequality we consider the following example involving empirical measures.

Let X_1, \ldots, X_n be a sample of size n drawn from μ , that is, independent random vectors in \mathbb{R}^d with distribution μ . They may be treated as independent copies of a random vector X. The associated empirical measures are defined by

$$\mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{X_k},$$
(2.1)

where δ_x denotes a delta-measure at the point x. Correspondingly, their linear projections represent one dimensional empirical measures

$$\mu_{n,\theta} = \frac{1}{n} \sum_{k=1}^{n} \delta_{\langle X_k, \theta \rangle}.$$
(2.2)

For simplicity, let us restrict ourselves to compactly supported distributions μ supported on the unit ball B_1 in \mathbb{R}^d . By concavity of $x \to x^{\alpha}$ in $x \ge 0$ for $\alpha \in (0, 1]$, it follows from (1.6) that

$$\mathbb{E} W(\mu_n, \mu) \le 12 \left[\mathbb{E} \sup_{|\theta|=1} W(\mu_{n,\theta}, \mu_{\theta}) \right]^{\alpha}$$
(2.3)

with $\alpha = \frac{2}{d+2}$, In the case $d \geq 3$, it is known (cf. e.g. [5], [3]) that $\mathbb{E} W(\mu_n, \mu)$ is of order at most $c_d n^{-1/d}$, and this rate cannot be improved as $n \to \infty$ for the uniform distribution. However, if d = 1 and μ is compactly supported, the rate is of the standard order $\mathbb{E} W(\mu_n, \mu) \sim \frac{1}{\sqrt{n}}$ up to μ -dependent factors (for a general two-sided bound we refer to [2], Theorem 3.5). Hence, we also have $\mathbb{E} W(\mu_{n,\theta}, \mu_{\theta}) \sim \frac{1}{\sqrt{n}}$ for every fixed θ . In order to determine the upper bound for the transport distance via the bound (2.3), let us prove the following result.

Theorem 2.1. If μ is supported on the unit ball B_1 in \mathbb{R}^d , then

$$\mathbb{E} \sup_{|\theta|=1} W(\mu_{n,\theta}, \mu_{\theta}) \le c\sqrt{d} \frac{1}{\sqrt{n}}$$
(2.4)

up to some absolute constant c > 0.

Applying this bound in (2.3) and using a lower bound with rate $c_d n^{-1/d}$ for the left-hand side with large n, we may conclude that necessarily $\alpha \geq 2/d$. Thus, the exponent $\alpha = \frac{2}{d+2}$ is asymptotically optimal for growing dimension d.

3. Reduction to Compactly Supported Lipschitz Functions

We need some preparation for the proof of Theorem 1.1. The argument is based on truncation, smoothing, the Plancherel theorem, together with the Kantorovich duality theorem (1.2).

Let U_r (r > 0) denote the collection of all functions $u : \mathbb{R}^d \to \mathbb{R}$ with $||u||_{\text{Lip}} \leq 1$, u(0) = 0, which are supported on the Euclidean ball $B_r = \{x \in \mathbb{R}^d : |x| \leq r\}$. As a first step, we show that the supremum in (1.2) may be restricted to the set U_r at the expense of a small error for large values of the parameter r under a p-th moment assumption. Define

$$W^{(r)}(X,Y) = \sup_{u \in U_r} |\mathbb{E}u(X) - \mathbb{E}u(Y)| = \sup_{u \in U_r} \left| \int_{\mathbb{R}^d} u \, d\mu - \int_{\mathbb{R}^d} u \, d\nu \right|,$$

assuming that the random vectors X and Y have distributions μ and ν .

Sergey G. Bobkov and Friedrich Götze

Lemma 3.1. Let $(\mathbb{E}|X|^p)^{1/p} \leq b$ and $(\mathbb{E}|Y|^p)^{1/p} \leq b$ for some p > 1. For any r > 0,

$$W(X,Y) \le 3 W^{(r)}(X,Y) + 4b \left(\frac{2b}{r}\right)^{p-1}$$

Proof. Let u be a Lipschitz function participating in the supremum (1.2) with u(0) = 0 (whout loss of generality). The latter ensures that $|u(x)| \leq |x|$ for all $x \in \mathbb{R}^d$.

Define the function $u_r = u\psi_r$, where using the notation $a^+ = \max(a, 0)$,

$$\psi_r(x) = \left(1 - \frac{2}{r}\operatorname{dist}(B_{r/2}, x)\right)^+, \quad x \in \mathbb{R}^d.$$

Clearly, $0 \le \psi_r \le 1$ and $\|\psi_r\|_{\text{Lip}} \le \frac{2}{r}$, as the distance function

$$x \to \operatorname{dist}(B, x) = \inf\{|x - y| : y \in B\}$$

has a Lipschitz semi-norm at most one for any non-empty set B in \mathbb{R}^d .

By the definition, $|u_r(x)| \leq |u(x)|$ for all $x \in \mathbb{R}^d$, and

$$u_r(x) = u(x) \text{ for } |x| \le r/2, \qquad u_r(x) = 0 \text{ for } |x| \ge r.$$
 (3.1)

Writing

$$u_r(x) - u_r(y) = (u(x) - u(y))\psi_r(x) + u(y)(\psi_r(x) - \psi_r(y)),$$

it follows that, for all $x \in \mathbb{R}^d$ and $y \in B_r$,

$$|u_r(x) - u_r(y)| \le |u(x) - u(y)| + |y|\frac{2}{r}|x - y| \le 3|x - y|.$$

A similar final inequality holds true for $x \in B_r$ and $y \in \mathbb{R}^d$. In addition, $u_r(x) - u_r(y) = 0$ in the case $x, y \notin B_r$. Therefore, $||u_r||_{\text{Lip}} \leq 3$.

Next, by (3.1),

$$\begin{aligned} \left| \int_{\mathbb{R}^d} u \, d\mu - \int_{\mathbb{R}^d} u_r \, d\mu \right| &\leq \int_{|x| > r/2} |u(x)| \, d\mu(x) + \int_{|x| > r/2} |u_r(x)| \, d\mu(x) \\ &\leq 2 \int_{|x| > r/2} |x| \, d\mu(x) \\ &= 2 \mathbb{E} \left| X \right| \mathbf{1}_{\{|X| > r/2\}} \leq 2 \frac{\mathbb{E} \left| X \right|^p}{(r/2)^{p-1}} \leq 2 b^p \left(\frac{2}{r} \right)^{p-1}. \end{aligned}$$

With a similar inequality for the measure ν , we obtain that

$$\left|\int_{\mathbb{R}^d} u \, d(\mu - \nu) - \int_{\mathbb{R}^d} u_r \, d(\mu - \nu)\right| \leq 4b^p \left(\frac{2}{r}\right)^{p-1}$$

implying

$$\left|\int_{\mathbb{R}^d} u \, d(\mu - \nu)\right| \le \left|\int_{\mathbb{R}^d} u_r \, d(\mu - \nu)\right| + 4b^p \left(\frac{2}{r}\right)^{p-1}.$$

Since $||u_r||_{\text{Lip}} \leq 3$, we have $\frac{1}{3}u_r \in U_r$, so that the last integral does not exceed $3W^{(r)}(X,Y)$ in absolute value. Thus,

$$\left| \int_{\mathbb{R}^d} u \, d(\mu - \nu) \right| \le 3 \, W^{(r)}(X, Y) + 4b^p \left(\frac{2}{r}\right)^{p-1}.$$

It remains to take the supremum on the left-hand side over all functions $u : \mathbb{R}^d \to \mathbb{R}$ such that $||u||_{\text{Lip}} \leq 1$ and u(0) = 0.

Cramér-Wold theorem

4. Fourier Transforms

Any (measurable) compactly supported function u on \mathbb{R}^d has a well-defined Fourier transform

$$\widehat{u}(t) = \int_{\mathbb{R}^d} e^{i\langle t, x \rangle} u(x) \, dx, \quad t \in \mathbb{R}^d, \tag{4.1}$$

which represents a C^{∞} -smooth function. Towards the proof of Theorem 1.1 let us state now the following integrability property.

Lemma 4.1. For any function $u : \mathbb{R}^d \to \mathbb{R}$ which is supported on the ball B_r and has a Lipschitz semi-norm $||u||_{\text{Lip}} \leq 1$,

$$\int_{\mathbb{R}^d} |\widehat{u}(t)|^2 \, |t|^2 \, dt \, \le \, \omega_d \, (2\pi r)^d. \tag{4.2}$$

In particular, this inequality holds true for any function u in U_r , Here and elsewhere ω_d stands for the *d*-dimensional volume of the unit ball B_1 .

Proof. First assume that $\hat{u}(t) = O(1/|t|^p)$ as $|t| \to \infty$ with p > 0 sufficiently large. Then (4.1) may be inverted in the form of the Fourier transform

$$u(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-i\langle t, x \rangle} \,\widehat{u}(t) \, dt.$$

In particular, u is smooth on \mathbb{R}^n . Moreover, this equality may be differentiated along every coordinate x_k to represent the corresponding partial derivatives as

$$\partial_{x_k} u(x) = -\frac{i}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-i\langle t, x \rangle} t_k \,\widehat{u}(t) \, dt, \quad k = 1, \dots, d.$$

Hence, by the Plancherel theorem,

$$\int_{\mathbb{R}^d} t_k^2 \, |\widehat{u}(t)|^2 \, dt = (2\pi)^d \int_{\mathbb{R}^d} (\partial_{x_k} u(x))^2 \, dx.$$

Summing over all $k \leq d$ and using $|\nabla u(x)| \leq 1$ for $x \in B_r$ and $\nabla u(x) = 0$ for |x| > r, we get

$$\int_{\mathbb{R}^d} |t|^2 \, |\widehat{u}(t)|^2 \, dt = (2\pi)^d \int_{\mathbb{R}^d} |\nabla u(x)|^2 \, dx \le (2\pi)^d \cdot \omega_d r^d,$$

which is the desired inequality (4.2).

In the general case, a smoothing argument can be used. Take a probability density won \mathbb{R}^d which is supported on the unit ball B_1 and has characteristic function $\widehat{w}(t)$ satisfying $\widehat{w}(t) = O(1/|t|^p)$ as $|t| \to \infty$. Given $\varepsilon > 0$, the probability density $w_{\varepsilon}(x) = \varepsilon^{-d}w(x/\varepsilon)$ is supported on the ball B_{ε} and has characteristic function $\widehat{w}_{\varepsilon}(t) = \widehat{w}(\varepsilon t)$. Consider the convolution

$$u_{\varepsilon}(x) = (u * w_{\varepsilon})(x) = \int_{\mathbb{R}^d} u(x - y) w_{\varepsilon}(y) dy, \quad x \in \mathbb{R}^d.$$

This function is supported on $B_{r+\varepsilon}$ and has a Lipschitz semi-norm $||u_{\varepsilon}||_{\text{Lip}} \leq ||u||_{\text{Lip}} \leq 1$. By the Lipschitz property, $|u(x)| \leq |u(0)| + r$ for any $x \in B_r$, implying that

$$\sup_{t\in\mathbb{R}^d} |\widehat{u}(t)| \le \int_{B_r} |u(x)| \, dx \le (|u(0)|+r) \, \omega_d r^d < \infty.$$

Hence, the Fourier transform of u_{ε} satisfies $\hat{u}_{\varepsilon}(t) = \hat{u}(t)\hat{w}(\varepsilon t) = O(1/|t|^p)$ as $|t| \to \infty$. Thus, one may apply the previous step to the function u_{ε} which gives

$$\int_{\mathbb{R}^d} |\widehat{u}(t)|^2 \, |\widehat{w}(\varepsilon t)|^2 \, |t|^2 \, dt \, \leq \, \omega_d \, (2\pi \, (r+\varepsilon))^d.$$

It remains to send $\varepsilon \to 0$ in this inequality and apply Fatou's lemma together with $\widehat{w}(\varepsilon t) \to 1$ as $\varepsilon \to 0$.

Next, let us connect the Kantorovich distance with the multivariate characteristic functions

$$f(t) = \mathbb{E} e^{i\langle t, X \rangle}, \quad g(t) = \mathbb{E} e^{i\langle t, Y \rangle} \qquad (t \in \mathbb{R}^d).$$

Lemma 4.2. Given random vectors X, Y in \mathbb{R}^d with characteristic functions f, g and finite first absolute moments, we have, for any $t \in \mathbb{R}^d$,

$$|f(t) - g(t)| \le |t| \sup_{|\theta|=1} W(X_{\theta}, Y_{\theta}).$$

Proof. In dimension one, using the property that the function $u_t(x) = \frac{1}{t} e^{itx}$ with parameter $t \neq 0$, has a Lipschitz semi-norm at most 1, it follows from (1.2) that

$$|f(t) - g(t)| \le |t| W(X, Y).$$

In dimension d, just note that, for any $\theta \in \mathbb{R}^d$, the functions $r \to f(r\theta)$ and $r \to g(r\theta)$ represent the characteristic functions of X_θ of Y_θ .

5. Proof of Theorem 1.1

With $b = \max(||X||_p, ||Y||_p)$, where

$$||X||_p = (\mathbb{E} |X|^p)^{1/p}, \quad ||Y||_p = (\mathbb{E} |Y|^p)^{1/p},$$

the inequality (1.4) is homogeneous with respect to (X, Y), so one may assume that b = 1. As a consequence,

$$W(X,Y) \le \mathbb{E}|X| + \mathbb{E}|Y| \le 2.$$

Let η be a random vector with uniform distribution in the ball B_1 , that is, with density $w(x) = \frac{1}{\omega_d} \mathbb{1}_{B_1}(x)$, and let h(t) denote its characteristic function. Consider the random vectors

$$X(\varepsilon) = X + \varepsilon \eta, \quad Y(\varepsilon) = Y + \varepsilon \eta \qquad (\varepsilon > 0)$$

assuming that η is independent of X and Y. Then, by the definition (1.1),

$$W(X,Y) \le W(X(\varepsilon),Y(\varepsilon)) + \varepsilon.$$

On the other hand, using

$$|X(\varepsilon)||_p \le 1 + \varepsilon, \quad ||Y(\varepsilon)||_p \le 1 + \varepsilon,$$

one may apply Lemma 3.1, which gives that, for any r > 0,

$$W(X(\varepsilon), Y(\varepsilon)) \le 3W^{(r)}(X(\varepsilon), Y(\varepsilon)) + 4(1+\varepsilon)\left(\frac{2(1+\varepsilon)}{r}\right)^{p-1}.$$

Therefore,

$$W(X,Y) \le 3W^{(r)}(X(\varepsilon),Y(\varepsilon)) + 4(1+\varepsilon)\left(\frac{2(1+\varepsilon)}{r}\right)^{p-1} + \varepsilon.$$
(5.1)

In order to estimate the first term on the right-hand side, note that $X(\varepsilon)$ and $Y(\varepsilon)$ have respective characteristic functions

$$f_{\varepsilon}(t) = f(t)h(\varepsilon t), \quad g_{\varepsilon}(t) = g(t)h(\varepsilon t) \qquad (t \in \mathbb{R}^d),$$

where f and g denote the characteristic functions of X and Y. Since h(t) is square integrable, while $|f(t)| \leq 1$ and $|g(t)| \leq 1$, these random vectors have square integrable densities which we denote by p_{ε} and q_{ε} respectively. Hence, given a function u in U_r , one may write

$$\mathbb{E} u(X(\varepsilon)) - \mathbb{E} u(Y(\varepsilon)) = \int_{\mathbb{R}^d} u(x) \left(p_{\varepsilon}(x) - q_{\varepsilon}(x) \right) dx$$

Now we are in position to apply the Plancherel theorem and rewrite the last integral as

$$\frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \widehat{u}(t) \left(\bar{f}(t) - \bar{g}(t) \right) h(\varepsilon t) \, dt,$$

and then thanks to Lemma 4.2 we have

$$\left|\mathbb{E}\,u(X(\varepsilon)) - \mathbb{E}\,u(Y(\varepsilon))\right| \le \frac{M}{(2\pi)^d} \int_{\mathbb{R}^d} \left|\widehat{u}(t)\right| \left|t\right| \left|h(\varepsilon t)\right| dt,\tag{5.2}$$

where

$$M = \sup_{|\theta|=1} W(X_{\theta}, Y_{\theta}).$$

Moreover, using

$$\begin{split} \int_{\mathbb{R}^d} |h(\varepsilon t)|^2 \, dt &= \varepsilon^{-d} \int_{\mathbb{R}^d} |h(t)|^2 \, dt \\ &= (2\pi)^d \, \varepsilon^{-d} \int_{\mathbb{R}^d} w(x)^2 \, dx \, = \, (2\pi)^d \, \varepsilon^{-d} \omega_d^{-1} \end{split}$$

and applying Cauchy's inequality in (5.2) together with Lemma 4.1, we obtain that

$$|\mathbb{E} u(X(\varepsilon)) - \mathbb{E} u(Y(\varepsilon))| \le M\left(\frac{r}{\varepsilon}\right)^{\frac{d}{2}}.$$

Taking the supremum over all $u \in U(r)$ on the left-hand side leads to the similar bound for $W^{(r)}(X(\varepsilon), Y(\varepsilon))$, and using this in (5.1) we are led to

$$W(X,Y) \leq 3M\left(\frac{r}{\varepsilon}\right)^{\frac{d}{2}} + 4(1+\varepsilon)\left(\frac{2(1+\varepsilon)}{r}\right)^{p-1} + \varepsilon.$$

To simplify optimization over the free parameters r and ε , let us assume that ε is bounded, so that $2(1 + \varepsilon) \le c$ for some constant c (to be chosen later on). Then we have

$$W(X,Y) \leq 3M\left(\frac{r}{\varepsilon}\right)^{\frac{d}{2}} + 2c\left(\frac{c}{r}\right)^{p-1} + \varepsilon.$$

Let us then replace r with cs and ε with c δ in the above inequality to get

$$W(X,Y) \leq 3M\left(\frac{s}{\delta}\right)^{\frac{d}{2}} + 2c\left(\frac{1}{s}\right)^{p-1} + c\delta.$$

Here, equalizing the terms $M\left(\frac{s}{\delta}\right)^{\frac{d}{2}}$ and $s^{-(p-1)}$, we find the unique value of s for which the above yields

$$W(X,Y) \le (3+2c) A\delta^{-\beta} + c\delta, \qquad (5.3)$$

where

$$\beta = \frac{(p-1)\frac{d}{2}}{p-1+\frac{d}{2}}, \quad A = M^{\frac{p-1}{p-1+\frac{d}{2}}}$$

The choice $\delta = A^{\frac{1}{\beta+1}}$ in (5.3) leads to

$$W(X,Y) \le (3+3c)A^{\frac{1}{\beta+1}},$$
 (5.4)

provided that $2(1 + c\delta) \leq c$. If we require that $\delta \leq \frac{1}{6}$, the latter condition is satisfied for c = 3, and we obtain from (5.4) that

$$W(X,Y) \le 12 A^{\frac{1}{\beta+1}},$$
 (5.5)

provided that $A^{\frac{1}{\beta+1}} \leq \frac{1}{6}$. In the other case, the right-hand side in (5.5) is greater than or equal to 2, so this inequality is fulfilled automatically due to the property $W(X,Y) \leq 2$.

It remains to note that

$$A^{\frac{1}{\beta+1}} = M^{\frac{2}{dp^*+2}}, \quad p^* = \frac{p}{p-1}.$$

6. Proof of Theorem 2.1

Let U denote for the space of all functions $u : [-1,1] \to \mathbb{R}$ with $||u||_{\text{Lip}} \leq 1$, such that u(0) = 0. We equip U with the uniform distance

$$||u - v||_{\infty} = \max\{|u(x) - v(x)| : |x|, |y| \le 1\},\$$

which turns this set into a compact space.

The empirical measures μ_n and $\mu_{n,\theta}$ defined in (2.1)-(2.2) are random with mean μ so that

$$\int_{\mathbb{R}^d} u \, d\mu_{n,\theta} = \frac{1}{n} \sum_{k=1}^n u(\langle X_k, \theta \rangle), \quad \int_{\mathbb{R}^d} u \, d\mu_{\theta} = \mathbb{E} \, u(\langle X, \theta \rangle).$$

According to the Kantorovich duality theorem (1.2),

$$\sup_{|\theta|=1} W(\mu_{n,\theta},\mu_{\theta}) = \frac{1}{n} \sup_{\theta \in S^{d-1}} \sup_{u \in U} \left| \sum_{k=1}^{n} \left(u(\langle X_k,\theta \rangle) - \mathbb{E} u(\langle X_k,\theta \rangle) \right) \right|,$$
(6.1)

where $S^{d-1} = \{\theta \in \mathbb{R}^d : |\theta| = 1\}$ denotes the unit sphere in \mathbb{R}^d .

In order to bound the expectation of the right-hand side in (6.1), one may use chaining arguments, in particular, a well-known theorem by Dudley which says the following (for a proof, let us refer to [8], [9]). Given a random variable ξ , define its Orlicz ψ_2 -norm

$$\|\xi\|_{\psi_2} = \inf \{\lambda > 0 : \mathbb{E} e^{\xi^2/\lambda^2} \le 2\}.$$

Suppose that $\xi(t)$ is a mean zero random process defined on some compact metric space (T, ρ) , which satisfies the Lipschitz property

$$\|\xi(t) - \xi(s)\|_{\psi_2} \le \Lambda \rho(t, s), \quad t, s \in T,$$
(6.2)

with some $\Lambda > 0$. Then with some absolute constant K we have

$$\mathbb{E} \sup_{t \in T} \xi(t) \le K\Lambda \int_0^D \sqrt{\log N(\varepsilon)} \, d\varepsilon, \tag{6.3}$$

8

where $D = \max\{\rho(t,s) : t, s \in T\}$ is the diameter and $N(\varepsilon) = N(T, \rho, \varepsilon)$ is the minimal number of closed balls in T of radius ε needed to cover the space (recall that $\log N(\varepsilon)$ is called the ε -entropy of (T, ρ)).

In view of (6.1), it is natural to consider the random process

$$\xi(t) = \xi(u,\theta) = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} \left(u(\langle X_k, \theta \rangle) - \mathbb{E} u(\langle X_k, \theta \rangle) \right), \quad t = (u,\theta) \in T = U \times S^{d-1}.$$

We equip T with the metric

$$\rho(t,s) = ||u - v||_{\infty} + |\theta - \theta'|, \quad t = (u,\theta), s = (v,\theta') \in T$$

Endowed with this metric T will be a compact space of diameter D = 4.

Now, given two points $t = (u, \theta)$, $s = (v, \theta')$ in T, one may write

$$\xi(t) - \xi(s) = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} (\eta_k - \mathbb{E}\eta_k),$$

where

$$\eta_k = u(\langle X_k, \theta \rangle) - v(\langle X_k, \theta' \rangle).$$

These random variables are independent and bounded. Indeed, writing

$$\eta_k = \left(u(\langle X_k, \theta \rangle) - v(\langle X_k, \theta \rangle) \right) + \left(v(\langle X_k, \theta \rangle) - v(\langle X_k, \theta' \rangle) \right)$$

and using the Lipschitz property of v together with the assumption $|X_k| \leq 1$ a.s., we have

$$|\eta_k| \leq \rho(t,s)$$
 a.s

Note that, for any random variable η such that $|\eta| \leq r$ a.s.,

$$\mathbb{E} e^{z(\eta - \mathbb{E}\eta)} \le e^{r^2 z^2/2} \quad \text{for all } z \in \mathbb{R}.$$

Hence, this holds for all $\eta = \eta_k$ with $r = \rho(t, s)$, and thus

$$\mathbb{E} e^{z(\xi(t) - \xi(s))} < e^{\rho^2(t,s)z^2/2}$$

Integrating this inequality with respect to z over the Gaussian measure with mean zero and variance σ^2 ($0 < \sigma < 1/\rho(t,s)$), we get $\mathbb{E} e^{\sigma^2(\xi(t)-\xi(s))^2/2} \le 1/\sqrt{\frac{1}{(\sigma\rho(t,s))^2}-1}$ which implies

$$\|\xi(t) - \xi(s)\|_{\psi_2} \le \rho(t, s)\sqrt{5/2}$$

Thus, the Lipschitz condition (6.2) is fulfilled with an absolute constant Λ , and we may apply the Dudley's bound (6.3). By the definition of the metric ρ in T, any closed ball of radius 2ε in this space contains the product of a closed ball in U and a closed ball in S^{d-1} , both of radius ε . Hence, the corresponding ε -entropies are connected by the relation

$$N(T,\rho,2\varepsilon) \le N(U,\varepsilon) N(S^{d-1},\varepsilon).$$
(6.4)

It is well-known (cf. [7]) that

$$\log N(U,\varepsilon) \le \frac{c}{\varepsilon}, \quad N(S^{d-1},\varepsilon) \le \left(\frac{c}{\varepsilon}\right)^d, \quad 0 < \varepsilon \le 2,$$

with some absolute constant c > 0. Using these bounds in (6.4) and then in (6.3), we conclude that the expectation of both sides in (6.1) does not exceed a multiple of $\sqrt{d/n}$. This gives the desired relation (2.4).

Sergey G. Bobkov and Friedrich Götze

References

- [1] Billingsley, P. Probability and Measure. Third ed., John Wiley & Sons. 1995.
- [2] Bobkov, S. G.; Ledoux, M. One-dimensional empirical measures, order statistics and Kantorovich transport distances. Memoirs of the AMS, 261 (2019), no. 1259, 126 pp.
- [3] Bobkov, S. G.; Ledoux, M. A simple Fourier analytic proof of the AKT optimal matching theorem. Ann. Appl. Probab. 31 (2021), no. 6, 2567–2584.
- [4] Cramér, H; Wold, H. Some Theorems on Distribution Functions. J. London Math. Soc. 11 (4) (1936), 290–294.
- [5] Dudley, R. M. The speed of mean Glivenko–Cantelli convergence. Ann. Math. Stat. 40 (1968), 40–50.
- [6] Dudley, R. M. Real analysis and probability. Cambridge Stud. Adv. Math., 74 Cambridge University Press, Cambridge, 2002, x+555 pp.
- [7] Kolmogorov, A. N.; Tikhomirov, V. M. ε-entropy and ε-capacity of sets in function spaces. (Russian) Uspekhi Matem. Nauk 14 (1959), no. 2 (86), 3–86.
- [8] Talagrand, M. Majorizing measures: the generic chaining. Ann. Probab. 24 (1996), no. 3, 1049– 1103.
- [9] Talagrand, M. Upper and lower bounds for stochastic processes. Modern methods and classical problems. 60. Springer, Heidelberg, 2014. xvi+626 pp.
- [10] Villani, C. Topics in optimal transportation. Grad. Stud. Math., 58 American Mathematical Society, Providence, RI, 2003, xvi+370 pp.