

Fractonic superfluids. III. Hybridizing higher moments

Han-Xie Wang^{1,*}, Shuai A. Chen^{2,*} and Peng Ye^{1,†}

¹Guangdong Provincial Key Laboratory of Magnetolectric Physics and Devices,
State Key Laboratory of Optoelectronic Materials and Technologies,
and School of Physics, Sun Yat-sen University, Guangzhou, 510275, China

²Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Straße 38, Dresden 01187, Germany
(Dated: Monday 16th December, 2024)

Fractonic many-body systems are characterized by mobility restrictions on single-particle hopping due to the conservation of higher moments (e.g., dipoles, angular momenta, and quadrupoles). This conservation allows the definition of new symmetries and their physical consequences, with spontaneous symmetry breaking (SSB) being especially intriguing in bosonic systems. Building on previous work [Yuan *et al.*, *Phys. Rev. Res.* **2**, 023267 (2020) (arXiv:1911.02876)] and [Chen *et al.*, *Phys. Rev. Res.* **3**, 013226 (2021) (arXiv:2010.03261)], we propose a new class of phases called hybrid fractonic superfluids (HFS), where multiple boson species interact and moment hybridization is conserved. We focus on two model series: Model Series A, where moments of the same order from different species are conserved, and Model Series B, where moments of different orders are conserved. In Model Series A, we analyze dipole moment hybridization and extend the analysis to higher-order moments. We investigate ground states, Goldstone modes, correlation functions, and the spontaneous breaking of charge symmetry, showing that at zero temperature, charge symmetry is partially broken when the spatial dimension is at least $N + 1$. This results in off-diagonal long-range order (ODLRO) driven by particle-hole bound states formed by bosons of different species. In Model Series B, we explore systems conserving mixed dipole-quadrupole moments. Additionally, we construct Bose-Hubbard-type lattice models where the small Hubbard U limit realizes HFS for both series. We analyze mean-field phase diagrams for Model Series A and discuss potential realizations in strongly tilted optical lattices. Finally, we suggest directions for future research, including generalized Mermin-Wagner theorems and large-scale numerical studies for Bose-Hubbard models. This article, together with previous work, forms a trilogy of fractonic superfluids, from which we attempt to uncover the exotic interplay of SSB physics and fracton physics through most of representative models and key properties of fractonic superfluids.

I. INTRODUCTION

In recent years, fracton topological order has been gaining increasing attention. Exactly solvable lattice models of fracton topological order are typically described by fracton codes, such as the three-dimensional X-cube model [1–20]. Topological excitations in fracton codes are commonly referred to as fractonic excitations, including fractons, subdimensional particles (e.g., lineons and planeons), and more intricate spatially extended excitations that are subject to mobility restriction [12, 14, 16]. In contrast to conventional topological orders, such as the toric code model, where topological excitations can be freely moved to any location via string operators, fractonic excitations in fracton topological order exhibit mobility restrictions. For instance, a single fracton cannot move freely along any direction, while a single lineon can only move along a straight line. Due to these mobility restrictions, fracton codes are believed to provide robust fault-tolerant quantum memory [2–4, 19]. The exploration of fracton topological order has also stimulated studies on subsystem symmetry-protected topological order (SPT) [21–32], substantially enlarging the territory of SPT phases.

Another active research direction involves treating fractons as constituent particles of many-body systems, where mobility restrictions arise from the conservation of higher-order

moments [33–88]. Such systems are referred to as *fractonic many-body systems*. It is well known that conventional total charge conservation is associated with a global $U(1)$ symmetry. Similarly, these new conserved quantities formed by higher moments are associated with $U(1)$ symmetry groups. Upon gauging, the resulting gauge fields are tensor gauge fields, and this new symmetry is alternatively referred to as “higher-rank symmetry”. To sum up, this line of research has evolved into a platform for exploring new / generalized symmetry, stimulating interests and ideas from condensed matter physics, gravity, effective field theory, and quantum information science.

On the other hand, spontaneous symmetry breaking (SSB) is a fundamental concept spanning a broad range of research areas in theoretical physics, from the Standard Model of particle physics to the Ginzburg-Landau theory of second-order phase transitions. In condensed matter physics, superfluids serve as canonical examples of systems with SSB, where the full charge $U(1)$ symmetry group is broken, and off-diagonal long-range order (ODLRO) is established. A celebrated phenomenon in this context is the Kosterlitz-Thouless topological phase transition, which occurs at finite temperature in two dimensions (2D). The study of SSB, particularly in the context of continuous global symmetries, has long been associated with important topics such as Goldstone mode counting, the Mermin-Wagner theorem concerning the stability of ODLRO at finite temperatures, and the infrared divergences induced by quantum fluctuations at zero temperature. In recent years, growing interest in generalized symmetries has led to significant advances across various subfields of theoretical physics.

* These authors contributed equally to this work.

† yepeng5@mail.sysu.edu.cn

Within the framework of generalized symmetries, long-range entangled states (e.g., topological order as pioneered by X.-G. Wen), which do not fall under the conventional classification of SSB phases, are now interpreted as SSB phases of higher-(form) symmetries. As the fracton physics is evolving rapidly, it is timely and urgent to investigate SSB physics in fracton systems. More specifically, for fractonic many-body quantum systems, with their exotic conserved quantities and corresponding new symmetries, key questions arise: *What are the physical consequences of the SSB of such new symmetries? What are the resulting phases of matter?*

Motivated by these questions, a series of exotic SSB phases of matter, termed *fractonic superfluids*, were proposed in Ref. [44] and have been systematically explored [44–49]. In these phases, the constituent bosons that are condensed are either completely immobile [44] or movable only along fixed directions [45–48]. Due to the similar restriction of mobility, these particles are also referred to as fractons and lineons respectively by borrowing the terminology in fractonic topological order and fracton codes. For simplicity, dipole moments, which are first-order moments, are often used in concrete model studies. The mobility restriction on fractons can be realized by imposing conservation of dipole moments, $\hat{Q}^{(i)} = \int d^d \mathbf{x} \hat{\rho} x_i$ [44]. Additionally, the conservation of angular charge moments (abbreviated as “angular moments”), $\hat{Q}_{a,b} = \int d^d \mathbf{x} (\hat{\rho}_a x_b - \hat{\rho}_b x_a)$, was explored in Refs. [45–48] to capture the partial restriction of mobility. Here, $\hat{\rho}$ represents the density operator, and a, b label the species of bosons. To make sense of higher-moment conservation laws while preserving the translation invariance of the Hamiltonians, total particle number conservation (i.e., “charge” conservation) is implicitly required in all cases.

In Refs. [44, 45], minimal models of fractonic superfluids are constructed and analyzed across various aspects, including symmetry transformations, Noether currents, ground-state wavefunctions, Gross-Pitaevskii equations, hydrodynamics, and vortex configurations. Through low-energy effective field theories, it has been shown that conservation of higher moments tends to suppress charge symmetry breaking, making the establishment of ODLRO more challenging under higher-moment conservation. Specifically, in systems with charge and dipole moment conservation, charge symmetry breaking occurs only in spatial dimensions $d \geq 3$ at zero temperature. Recently, Ref. [58] demonstrated more generally that charge symmetry breaking is forbidden at finite temperature even in four dimensions as long as dipole conservation is imposed. In Ref. [46], continuity equations and Navier-Stokes-like equations were derived from the Euler-Lagrange equations of fractonic superfluids. Critical velocity fields and density currents were obtained, leading to a Landau-like criterion. Ref. [47] examined the emergence of higher-rank symmetries and angular moment conservation at low energies through renormalization group analysis, providing insights into their experimental realization. Meanwhile, Ref. [48] investigated finite-temperature phase transitions in fractonic superfluids, showing that the topological vortices (symmetry defects) proposed in Ref. [45] form a hierarchical structure of bound states, resulting in hierarchical confinement-deconfinement transitions.

Bose-Hubbard-type lattice models incorporating higher-moment conservation are also of significant interest, as they extend the conventional Bose-Hubbard model, which realizes the superfluid phase in the weak interaction regime. Dipolar Bose-Hubbard lattice models have been constructed and studied extensively in various spatial dimensions, yielding profound physical insights (e.g., Refs. [59–64]). In the small- U limit, these lattice models at long wavelengths reduce to the many-boson model of fractonic superfluids discussed in Ref. [44].

In the fractonic superfluid systems and associated Bose-Hubbard lattice models discussed above, the formulation of conserved quantities plays a crucial role in shaping the low-energy physics. Different conserved quantities can lead to entirely distinct outcomes in the interplay between SSB and fracton physics, influencing phenomena such as vortex configurations, the Mermin-Wagner theorem, finite-temperature phase diagrams, and hydrodynamics. Therefore, it is worthwhile to explore novel conserved quantities that substantially extend beyond straightforward generalizations of previously studied cases.

In this paper, we introduce the concept of *hybrid fractonic superfluids* (HFS) by imposing the conservation of hybridized moments. HFS represent a class of higher-moment symmetry-breaking phases characterized by hybrid moments, which are constructed from the moments of bosons belonging to different species. Therefore, HFS models inherently involve at least two distinct species of bosons. We systematically investigate HFS and their realizations in Bose-Hubbard lattice models. Specifically, we focus on two representative series of HFS (referred to as *Model Series A* and *Model Series B*), distinguished by the mechanisms of moment hybridization. Notably, previous studies [45–48] on angular moments can also be regarded as a subclass of fractonic superfluids with hybrid moments and could be categorized as Model Series C if needed. Given the profound physical implications uncovered in these prior studies—which likely represent only the tip of the iceberg of hybridization—it is imperative to systematically explore the broader impacts of hybridization, a key motivation for the present work. This article, together with Refs. [44, 45], forms a trilogy of fractonic superfluids. From the trilogy as well as the extended studies in Refs. [46–49], we attempt to uncover the exotic interplay of SSB physics and fracton physics through most of representative models and key properties of fractonic superfluids.

More concretely, in Model Series A, we consider the hybridization of the *same* order moments of bosons of *different* species. Arbitrary models in Model Series A can be represented by a tuple $[d, N, m, m']$, where d is the spatial dimension, N is the order of the moments participating in the hybridization, m is the total number of species of bosons, and m' denotes the number of species of bosons involved in the hybridization. The conserved hybrid moments are $\hat{Q}_{\text{mix}}^{(i)} = \int d^d \mathbf{x} \left(\sum_{a=1}^{m'} \hat{\rho}_a \right) \mathcal{M}_{N,d}^{(i)}$. The symbol $\mathcal{M}_{N,d}^{(i)}$ represents a monomial obtained by multiplying coordinates. Here, the index i labels the i -th specific monomial in lexicographic ordering [89]. To preserve spatial translation symmetry as before, moments of orders lower than N must also be conserved

in addition to the conservation of hybrid moments. We specifically study the $[d, 1, 2, 2]$ model of Model Series A in which total dipole moments are conserved. In this model, we examine the ground states, Goldstone modes, Noether currents, and correlation functions. We find that one of the dispersion relations is linear. In two spatial dimensions, the correlation functions $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \rangle$ and $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decay as a power law, where $\hat{\Phi}_1$ and $\hat{\Phi}_2$ are respectively annihilation bosonic operators of different species. However, the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{0}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant at long distances. As a result, the system undergoes the spontaneous breaking of *relative* charge symmetry that is associated with the particle number difference of different species of bosons, thus supporting true off-diagonal long-range order (ODLRO). The spontaneous breaking of relative charge symmetry in $[d, 1, 2, 2]$ model is characterized by the order parameter $\langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \rangle$. This series of results are also extended to arbitrary $[d, N, m, m']$ models, and partial spontaneous breaking of charge symmetry occurs in spatial dimensions $d = N + 1$.

In Model Series B, we consider the hybridization of *different* order moments of bosons of *different* species. The hybrid moments are $\hat{Q}_{\text{mix}}^{(i_1 \dots i_m)} = \int d^d \mathbf{x} \sum_a^m \hat{\rho}_a \mathcal{M}_{N_a, d}^{(i_a)} \ell^{N_m - N_a}$. N_a denotes the orders of the moments of the species a which participate in the hybridization, and a constant ℓ with the dimension of length is added in order to compensate dimension imbalance between moments of different orders. Except for the order requirement, the other settings are the same as those in Model Series A. The moments whose orders are lower than N_a are also required to be conserved. We specifically study an example with conservation of hybridization of dipole and quadrupole moments, and also find one of the dispersion relations is linear. In two spatial dimensions, the correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decays in a power law, and the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant at long distances. Therefore, the spontaneous breaking of charge symmetry of bosons of species 1 occurs, and the system has a true ODLRO. This spontaneous breaking in Model Series B is characterized by the order parameter $\langle \hat{\Phi}_1(\mathbf{x}) \rangle$. A summary of comparison between different model realization with two species of bosons is given in Table I.

In this paper, we also construct Bose-Hubbard-type lattice models which correspond to Model Series A and Model Series B, and perform mean-field approximation for $[d, 1, 2, 2]$ lattice model. The mean-field results indicate that the introduction of hybridization causes new phases to appear in the system. The coupling coefficients of the two species of bosons and the hopping coefficients of the same species of bosons play similar roles in the phase transition. We also discuss the realization of $[d, 1, 2, 2]$ lattice model in strongly tilted optical lattices. In this method, Hubbard interaction terms between bosons of different species are required.

The remaining part of this paper is organized as follows. Sec. II introduces two typical hybrid moments. Based on the two hybrid moments, we provide Model Series A and Model Series B, and also present their minimal Hamiltonians. Effective field theories of the two models are performed in Sec. III

and IV. In Sec. V, the Bose-Hubbard-type lattice models are constructed, and relevant discussions are given. Sec. VI provides a summary of the principal findings and offers additional perspectives for future research.

II. CONSTRUCTION OF HAMILTONIAN OF HYBRID MOMENTS

In this section, we start with the conserved unhybrid moments and corresponding fractonic superfluid models with no hybridization. In a fractonic superfluid model [44] with no hybridization, there is only one species of bosons. By introducing two or more species of bosons in the system, we subsequently develop two models with typical hybridizations. With determining the hybrid moments, we construct the minimal Hamiltonians which satisfy the conservation laws.

A. Review of fractonic superfluids

Initially, the fractonic superfluid is put forward in the single-component boson system which preserves the multipole moments. We set $\hat{\Phi}^\dagger(\mathbf{x})$ and $\hat{\Phi}(\mathbf{x})$ to be bosonic creation and annihilation operators and $\hat{\rho}(\mathbf{x}) = \hat{\Phi}^\dagger(\mathbf{x}) \hat{\Phi}(\mathbf{x})$ is a density operator. The creation and annihilation operators satisfy the standard commutation relations $[\hat{\Phi}(\mathbf{x}), \hat{\Phi}^\dagger(\mathbf{y})] = \delta^d(\mathbf{x} - \mathbf{y})$, where $\mathbf{x} = (x_1, \dots, x_d)$ is a spatial coordinate and we neglect it in the following for symbol convenience. We formulate the charges related to the multipole moments systematically as

$$\hat{Q}_{n,d}^{(i)} = \int d^d \mathbf{x} \hat{\rho} \mathcal{M}_{n,d}^{(i)}, \quad (1)$$

where $\mathcal{M}_{n,d}^{(i)}$ denote the monomials obtained by multiplying the spatial rectangular coordinates with the coefficients equal to 1. The notation n stands for the number of multiplied spatial rectangular coordinates in $\mathcal{M}_{n,d}^{(i)}$, and d stands for spatial dimensions. Each $\hat{Q}_{n,d}^{(i)}$ denotes one component of the n -order moment in d spatial dimensions. The notation i denotes a index of the monomial for n th-order moments in the lexicographic ordering [89], and the index i is listed from 1 to $C_{d-1+n}^n = \frac{(d-1+n)!}{(d-1)!n!}$. For instance, in two spatial dimensions, we can sort the monomials of dipole moments as $\mathcal{M}_{2,2}^{(1)} = x_1^2$, $\mathcal{M}_{2,2}^{(2)} = x_1 x_2$, and $\mathcal{M}_{2,2}^{(3)} = x_2^2$. In addition, the conservation of charge $\hat{Q} = \int d^d \mathbf{x} \hat{\rho}$ is also required. Given the maximal order of the monomials as N , we can construct a symmetry group \mathcal{G} with an element

$$U = \exp \left[-i \left(\lambda \hat{Q} + \sum_{n=1}^N \sum_{i=1}^{C_{d-1+n}^n} \lambda_{n,d}^{(i)} \hat{Q}_{n,d}^{(i)} \right) \right] \quad (2)$$

which leads to a transformation

$$\hat{\Phi}' = U \hat{\Phi} U^\dagger = \hat{\Phi} e^{i(\lambda + \sum_n \sum_i \lambda_{n,d}^{(i)} \mathcal{M}_{n,d}^{(i)})}, \quad (3)$$

Table I. Comparison of patterns of charge symmetry breaking in four different model realizations with two species of bosons in two spatial dimensions at $T = 0$. The specific analysis of $[d, 1, 2, 2]$ is given in Sec. III, and the specific analysis of the hybrid dipole-quadrupole moment conserving model is given in Sec. III.

Superfluid models with two species of bosons	Two decoupled conventional superfluids	Two decoupled fractonic superfluids [44] with separate dipole conservations	$[d, 1, 2, 2]$ model (Eq. (20)) of Model Series A	Hybrid dipole-quadrupole conserving model (Eq. (35)) of Model Series B
Higher moment conservation	N/A	Dipole moments of two species	Hybrid dipole moments	Hybrid dipole-quadrupole moments
Broken charge symmetry, ODLRO, and Order parameter	$U(1)_{1,C} \otimes U(1)_{2,C}$ $\checkmark, \langle \hat{\Phi}_a(\mathbf{x}) \rangle$	N/A $\times, \text{N/A}$	$U(1)_{-C}$ $\checkmark, \langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \rangle$	$U(1)_{1,C}$ $\checkmark, \langle \hat{\Phi}_1(\mathbf{x}) \rangle$

where $\lambda, \lambda_{n,d}^{(i)} \in \mathbb{R}$ are group parameters. Hereafter, this symmetry will be alternatively called a higher-rank symmetry since the symmetry leads to higher-rank tensor gauge fields upon gauging. The kinetic term \mathcal{H}_0 is no longer Gaussian, e.g. the Hamiltonian constructed in the Appendix of Ref. [44]:

$$\mathcal{H}_0 = \sum_{i_1, \dots, i_{N+1}} d^d \mathbf{x} K^{i_1 \dots i_{N+1}} \left(\hat{\Phi}^\dagger \right)^{N+1} \left(\nabla_{i_1 \dots i_{N+1}} \log \hat{\Phi}^\dagger \right) \hat{\Phi}^{N+1} \left(\nabla_{i_1 \dots i_{N+1}} \log \hat{\Phi} \right), \quad (4)$$

where $\nabla_{i_1 \dots i_n} = \partial_{i_1} \dots \partial_{i_n}$ and symmetric coefficients $K^{i_1 \dots i_{N+1}}$. The operator $\log \hat{\Phi}$ is a notation to simplify the Hamiltonian, see Appendix of Ref. [44]. For convenience, we set the coefficients $K^{i_1 \dots i_{N+1}} > 0$. We can apply the potential term $V(\hat{\Phi}^\dagger, \hat{\Phi}) = -\mu \hat{\Phi}^\dagger \hat{\Phi} + \frac{g}{2} \hat{\Phi}^\dagger \hat{\Phi}^\dagger \hat{\Phi} \hat{\Phi}$, where μ is the chemical potential and $g > 0$ describes onsite repulsive interaction. A fractonic superfluid phase can be reached for a positive chemical potential.

To get an understanding on the fractonic superfluid phase, we can look at the following two examples. When the dipole moments are conserved, the kinetic term takes the form as

$$\mathcal{H}_0 = \sum_{i,j} d^d \mathbf{x} K^{ij} \left(\hat{\Phi}^\dagger \partial_i \partial_j \hat{\Phi}^\dagger - \partial_i \hat{\Phi}^\dagger \partial_j \hat{\Phi}^\dagger \right) \left(\hat{\Phi} \partial_i \partial_j \hat{\Phi} - \partial_i \hat{\Phi} \partial_j \hat{\Phi} \right). \quad (5)$$

The Hamiltonian is more complicated when the quadrupole moments are conserved:

$$\begin{aligned} \mathcal{H}_0 = & \sum_{i,j,k} d^d \mathbf{x} K^{ijk} \left(\hat{\Phi}^{\dagger 2} \partial_i \partial_j \partial_k \hat{\Phi}^\dagger - \hat{\Phi}^\dagger \partial_i \hat{\Phi}^\dagger \partial_j \partial_k \hat{\Phi}^\dagger \right. \\ & - \hat{\Phi}^\dagger \partial_j \hat{\Phi}^\dagger \partial_i \partial_k \hat{\Phi}^\dagger - \hat{\Phi}^\dagger \partial_k \hat{\Phi}^\dagger \partial_i \partial_j \hat{\Phi}^\dagger \\ & + 2 \partial_i \hat{\Phi}^\dagger \partial_j \hat{\Phi}^\dagger \partial_k \hat{\Phi}^\dagger \left. \right) \left(\hat{\Phi}^2 \partial_i \partial_j \partial_k \hat{\Phi} - \hat{\Phi} \partial_i \hat{\Phi} \partial_j \partial_k \hat{\Phi} \right. \\ & - \hat{\Phi} \partial_j \hat{\Phi} \partial_i \partial_k \hat{\Phi} - \hat{\Phi} \partial_k \hat{\Phi} \partial_i \partial_j \hat{\Phi} + 2 \partial_i \hat{\Phi} \partial_j \hat{\Phi} \partial_k \hat{\Phi} \left. \right). \quad (6) \end{aligned}$$

For the two examples, the dynamics of a single particle are strongly restricted while a bound state becomes free when the cooperative motions are required to conserve multipole moments: dipole moments for Hamiltonian in Eq. (5) and quadrupole moments for Eq. (6). The corresponding fractonic superfluid phase then has one Goldstone mode with a

high-order dispersion. For a general N as the largest order of the monomials, the effective theory describes the Goldstone mode with the dispersion relation $\omega \propto |\mathbf{k}|^{N+1}$. Therefore, this system breaks its charge symmetry in spatial dimensions $d \geq N + 2$.

B. Construction of Model Series A

Next, we begin to construct models with hybridization of moments. We first focus on Model Series A in which hybrid moments are formed by moments of the same order. Consider a system with m species of bosons, the operators $\hat{\Phi}_a^\dagger(\mathbf{x})$ and $\hat{\Phi}_a(\mathbf{x})$ create and annihilate a boson of species a respectively, and satisfy the commutation relations $[\hat{\Phi}_a(\mathbf{x}), \hat{\Phi}_b^\dagger(\mathbf{y})] = \delta_{ab} \delta^d(\mathbf{x} - \mathbf{y})$. In such a system, we consider the following conserved quantities in Eqs. (7, 8, 9, 10):

- The charges of bosons of species a are conserved. The range of a is $1 \leq a \leq m$.

$$\hat{Q}_a = \int d^d \mathbf{x} \hat{\rho}_a; \quad (7)$$

- The hybrid moments defined below are conserved. N is the order of the moments which participate in the hybridization. The range of the label i is $1 \leq i \leq C_{d-1+N}^N$.

$$\hat{Q}_{\text{mix}}^{(i)} = \int d^d \mathbf{x} \mathcal{M}_{N,d}^{(i)} \sum_{a=1}^{m'} \hat{\rho}_a; \quad (8)$$

- All the moments whose orders are lower than N are conserved. The ranges of a , n , and i are $1 \leq a \leq m$, $1 \leq n \leq N - 1$, and $1 \leq i \leq C_{d-1+n}^n$ respectively.

$$\hat{Q}_{a,n,d}^{(i)} = \int d^d \mathbf{x} \hat{\rho}_a \mathcal{M}_{n,d}^{(i)}; \quad (9)$$

- The moments of bosons of species $m' + 1 \leq a \leq m$ do not participate in the hybridization. Their N th-order moments are conserved. The ranges of a and i are $m' + 1 \leq a \leq m$, $1 \leq i \leq C_{d-1+N}^N$.

$$\hat{Q}_{a,N,d}^{(i)} = \int d^d \mathbf{x} \hat{\rho}_a \mathcal{M}_{N,d}^{(i)}. \quad (10)$$

Compared with the separate conservation of the N th-order moments for each species of bosons, the number of conserved quantities in the present hybridization case decreases. That means freeing up some mobility of bosons. We can see this better in the corresponding symmetry. We denote a group \mathcal{G} generated by the conserved quantities [Eqs. (7, 8, 9, 10)], and an element of the group

$$U = \exp \left[-i \sum_{a=1}^m \left(\lambda_a \hat{Q}_a + \sum_n^{N-1} \sum_i^{C_{d-1+n}^n} \lambda_{a,n}^{(i)} \hat{Q}_{a,n,d}^{(i)} \right) - i \sum_i^{C_{d-1+N}^N} \lambda_{\text{mix}}^{(i)} \hat{Q}_{\text{mix}}^{(i)} - i \sum_{a=m'+1}^m \sum_i^{C_{d-1+N}^N} \lambda_{a,N}^{(i)} \hat{Q}_{a,N,d}^{(i)} \right] \quad (11)$$

leads to the following transformation of field operators:

$$\begin{aligned} \hat{\Phi}'_a &= U \hat{\Phi}_a U^\dagger \\ &= \hat{\Phi}_a \exp \left\{ i \left[\lambda_a + \sum_n^{N-1} \sum_i^{C_{d-1+n}^n} \lambda_{a,n}^{(i)} \mathcal{M}_{n,d}^{(i)} + \sum_{b=1}^{m'} \delta_{ab} \sum_i^{C_{d-1+N}^N} \lambda_{\text{mix}}^{(i)} \mathcal{M}_{N,d}^{(i)} + \sum_{b=m'+1}^m \delta_{ab} \sum_i^{C_{d-1+N}^N} \lambda_{a,N}^{(i)} \mathcal{M}_{N,d}^{(i)} \right] \right\}, \end{aligned} \quad (12)$$

where $\lambda_a, \lambda_{a,n}, \lambda_{\text{mix}}^{(i)} \in \mathbb{R}$ are group parameters. The group \mathcal{G} can be written as the direct product of $U(1)$ groups,

$$\begin{aligned} \mathcal{G} &= \left(\prod_a^m \otimes U(1)_{a,C} \right) \left(\prod_a^m \prod_{n,i} \otimes U(1)_{a,n,i} \right) \\ &\left(\prod_i \otimes U(1)_{\text{mix-A},i} \right) \left(\prod_{a=m'+1}^m \prod_i \otimes U(1)_{a,N,i} \right), \end{aligned} \quad (13)$$

where group elements of $U(1)_{a,C}$, $U(1)_{a,n,i}$, and $U(1)_{\text{mix-A},i}$ are respectively labeled by λ_a , $\lambda_{a,n}^{(i)}$, and $\lambda_{\text{mix}}^{(i)}$.

A Hamiltonian satisfying this conservation law needs to commute with conserved quantities [Eqs. (7, 8, 9, 10)]. The minimal Hamiltonian can be written as:

$$\begin{aligned} \mathcal{H}_0 &= \sum_a^m \sum_{i_1, \dots, i_{N+1}}^d K_a^{i_1 \dots i_{N+1}} \left(\hat{\Phi}_a^\dagger \right)^{N+1} \left(\nabla_{i_1, \dots, i_{N+1}} \log \hat{\Phi}_a^\dagger \right) \\ &\hat{\Phi}_a^{N+1} \left(\nabla_{i_1, \dots, i_{N+1}} \log \hat{\Phi}_a \right) + \sum_{a \neq b}^{m'} \sum_{i_1, \dots, i_N}^d \Gamma_{a,b}^{i_1 \dots i_N} \left(\hat{\Phi}_a^\dagger \right)^N \\ &\left(\hat{\Phi}_b^\dagger \right)^N \left[\left(\nabla_{i_1, \dots, i_N} \log \hat{\Phi}_a^\dagger \right) - \left(\nabla_{i_1, \dots, i_N} \log \hat{\Phi}_b^\dagger \right) \right] \\ &\hat{\Phi}_a^N \hat{\Phi}_b^N \left[\left(\nabla_{i_1, \dots, i_N} \log \hat{\Phi}_a \right) - \left(\nabla_{i_1, \dots, i_N} \log \hat{\Phi}_b \right) \right]. \end{aligned} \quad (14)$$

For convenience, we also set $K_a^{i_1 \dots i_{N+1}} > 0$, $\Gamma_{a,b}^{i_1 \dots i_N} > 0$. In addition, $\Gamma_{a,b}^{i_1 \dots i_N} = \Gamma_{b,a}^{i_1 \dots i_N}$ is needed.

In the construction above, the spatial dimensions d , the order N , the number of boson species involved in hybridization m' , and the total number of boson species m are tunable for Model Series A. A tuple $[d, N, m, m']$ can denote one example of Model Series A.

C. $[d, 1, 2, 2]$ model of Model Series A

In this subsection, we discuss the simplest example of Model Series A, $[d, 1, 2, 2]$ model. $[d, 1, 2, 2]$ model can be regarded as a system in which the total dipole moments are conserved along with the conservation of the charges of each species,

$$\hat{Q}_a = \int d^d \mathbf{x} \hat{\rho}_a \quad (a = 1, 2), \quad (15)$$

$$\hat{Q}_{\text{mix}}^{(i)} = \int d^d \mathbf{x} (\hat{\rho}_1 x_i + \hat{\rho}_2 x_i) \quad (i = 1, \dots, d). \quad (16)$$

The total symmetry group \mathcal{G} can be written as

$$\mathcal{G} = \left(U(1)_{1,C} \otimes U(1)_{2,C} \right) \left(\prod_i \otimes U(1)_{\text{mix-A},i} \right). \quad (17)$$

An element of the group \mathcal{G} is given by

$$U = \exp \left[-i (\lambda_1 \hat{Q}_1 + \lambda_2 \hat{Q}_2 + \sum_i^d \lambda_{\text{mix}}^{(i)} \hat{Q}_{\text{mix}}^{(i)}) \right]. \quad (18)$$

And the transformation on field operators can be written as

$$\hat{\Phi}'_a = U \hat{\Phi}_a U^\dagger = \hat{\Phi}_a \exp \left[i (\lambda_a + \sum_i^d \lambda_{\text{mix}}^{(i)} x_i) \right]. \quad (19)$$

To satisfy this conservation law, the minimal Hamiltonian of $[d, 1, 2, 2]$ model with the potential terms is

$$\begin{aligned} \mathcal{H} &= \sum_a^2 \sum_{i,j}^d K_a^{ij} (\hat{\Phi}_a^\dagger \partial_i \partial_j \hat{\Phi}_a^\dagger - \partial_i \hat{\Phi}_a^\dagger \partial_j \hat{\Phi}_a^\dagger) \\ &\cdot (\hat{\Phi}_a \partial_i \partial_j \hat{\Phi}_a - \partial_i \hat{\Phi}_a \partial_j \hat{\Phi}_a) \\ &+ \sum_{a \neq b}^2 \sum_i^d \Gamma_{a,b}^i (\hat{\Phi}_a^\dagger \partial_i \hat{\Phi}_b^\dagger - \hat{\Phi}_b^\dagger \partial_i \hat{\Phi}_a^\dagger) \\ &\cdot (\hat{\Phi}_a \partial_i \hat{\Phi}_b - \hat{\Phi}_b \partial_i \hat{\Phi}_a) + V(\hat{\Phi}^\dagger, \hat{\Phi}), \end{aligned} \quad (20)$$

where $V(\hat{\Phi}^\dagger, \hat{\Phi}) = \sum_a^2 -\mu_a \hat{\Phi}_a^\dagger \hat{\Phi}_a + \frac{g}{2} \hat{\Phi}_a^\dagger \hat{\Phi}_a^\dagger \hat{\Phi}_a \hat{\Phi}_a$, and the coefficients satisfy $K_a^{ij} > 0$, $\Gamma_{a,b}^i > 0$, $K_a^{ij} = K_a^{ji}$, $\Gamma_{a,b}^i = \Gamma_{b,a}^i$, and $g > 0$.

We can also give the corresponding conserved quantities and Noether currents. For the conserved charge $\hat{Q}_a = \int d^d \mathbf{x} \hat{\Phi}_a^\dagger \hat{\Phi}_a = \int d^d \mathbf{x} \hat{\rho}_a$, we have the charge current densities,

$$\begin{aligned} \hat{J}_i^a &= i \sum_j^d K_a^{ij} \partial_j [\hat{\Phi}_a^{\dagger 2} (\hat{\Phi}_a \partial_i \partial_j \hat{\Phi}_a - \partial_i \hat{\Phi}_a \partial_j \hat{\Phi}_a) - \text{h.c.}] \\ &+ 2i \sum_{b \neq a}^2 \Gamma_{a,b}^i [(\hat{\rho}_a \hat{\Phi}_b \partial_i \hat{\Phi}_b^\dagger - \hat{\rho}_b \hat{\Phi}_a \partial_i \hat{\Phi}_a^\dagger) - \text{h.c.}]. \end{aligned} \quad (21)$$

For the hybrid moments $\hat{Q}_{\text{mix}}^{(j)} = \int d^d \mathbf{x} (\hat{\rho}_1 x_j + \hat{\rho}_2 x_j) = \int d^d \mathbf{x} \hat{\rho}_{\text{mix}}^{(j)}$, we have the current densities,

$$\begin{aligned} \hat{\mathcal{D}}_i^{(j)} &= x_j \hat{J}_i^1 + x_j \hat{J}_i^2 - i \sum_a^2 K_a^{ij} \\ &[\hat{\Phi}_a^{\dagger 2} (\hat{\Phi}_a \partial_i \partial_j \hat{\Phi}_a - \partial_i \hat{\Phi}_a \partial_j \hat{\Phi}_a) - \text{h.c.}]. \end{aligned} \quad (22)$$

The $\Gamma_{a,b}^i$ term is the emerging term from the hybridization of moments. This Hamiltonian density (Eq. (20)) is the simplest example of HFS in the Model A. The effective field theory of this Hamiltonian will be studied in Sec. III.

D. Construction of Model Series B

In the construction of Model Series B, in a system with m species of bosons, we consider the hybridization of moments of different orders. We can give conserved quantities of Model Series B:

- The charges of bosons of species a are conserved. The range of a is $1 \leq a \leq m$.

$$\hat{Q}_a = \int d^d \mathbf{x} \hat{\rho}_a, \quad (23)$$

- The hybrid moments defined below are conserved.

$$\hat{Q}_{\text{mix}}^{(i_1) \cdots (i_m)} = \int d^d \mathbf{x} \sum_a^m \hat{\rho}_a \mathcal{M}_{N_a, d}^{(i_a)} \ell^{N_m - N_a}, \quad (24)$$

Here, we use N_a to mark the orders of the moments of the species a which participate in the hybridization. For brevity, we let N_a increase as a increase, i.e. $N_1 < N_2 \cdots < N_m$. ℓ with a dimension of length denotes the coefficient by which moments of different orders are converted to each other. The introduction of ℓ ensures that the quantities in $\hat{Q}_{\text{mix}}^{(i_1) \cdots (i_m)}$ have the same dimensions. For each $\hat{Q}_{\text{mix}}^{(i_1) \cdots (i_m)}$, the index i_a can arbitrary taken in the range $1 \leq i_a \leq C_{d-1+N_a}^{N_a}$, but cannot be repeated with other $\hat{Q}_{\text{mix}}^{(i_1) \cdots (i_m)}$. The number of the conserved quantities $\hat{Q}_{\text{mix}}^{(i_1) \cdots (i_m)}$ is $C_{d-1+N_1}^{N_1}$.

- All the moments of species a whose orders are lower than N_a are conserved. The ranges of a , n , and i are $1 \leq a \leq m$, $1 \leq n \leq N_a - 1$, and $1 \leq i \leq C_{d-1+n}^n$ respectively.

$$\hat{Q}_{a,n,d}^{(i)} = \int d^d \mathbf{x} \hat{\rho}_a \mathcal{M}_{n,d}^{(i)}, \quad (25)$$

- The components of N_a -th-order moments which do not participate in the hybridization are conserved.

$$\hat{Q}_{a,N_a,d}^{(i_a)} = \int d^d \mathbf{x} \hat{\rho}_a \mathcal{M}_{N_a,d}^{(i_a)}. \quad (26)$$

Due to the fact that the numbers of components of different order moments are not equal in two spatial dimensions and above, some components of the N_a -th-order moments do not participate in the hybridization, and they are required to be conserved in Model Series B. The index i_a takes all $1 \leq i_a \leq C_{d-1+N_a}^{N_a}$ that are not taken in Eq. (24). The range of a is $1 \leq a \leq m$.

The setting of the conserved quantities above is similar to Model Series A. Since the hybridization of Model Series B is more complex, we require that all species of bosons are involved in the hybridization, i.e. $m' = m$. Since it is difficult to give a concrete expression of general Model Series B with the conservation of quantities in Eqs. (23, 24, 25, 26), we need to fix the coefficient m and N_a to simplify this model. Then, we can give the simplest Model Series B by setting $m = 2$, $N_1 = 1$, and $N_2 = 2$. In a system with two species of bosons, we mix dipole moments of bosons of species 1 and quadrupole moments of bosons of species 2. For convenience, the conserved quantities are given by:

$$\hat{Q}_a = \int d^d \mathbf{x} \hat{\rho}_a \quad (a = 1, 2), \quad (27)$$

$$\hat{Q}_2^{(i)} = \int d^d \mathbf{x} \hat{\rho}_2 x_i \quad (i = 1, \dots, d), \quad (28)$$

$$\hat{Q}_{\text{mix}}^{i,ii} = \int d^d \mathbf{x} (\hat{\rho}_1 x_i \ell + \hat{\rho}_2 x_i^2) \quad (i = 1, \dots, d), \quad (29)$$

$$\hat{Q}_2^{ij} = \int d^d \mathbf{x} \hat{\rho}_2 x_i x_j \quad (i, j = 1, \dots, d, \quad i \neq j). \quad (30)$$

To facilitate the visualization of the correspondence between components of different order moments, distinct symbols are employed in this example for the hybrid conserved quantities. We denote a group \mathcal{G} generated by the conserved quantities [Eqs. (27, 28, 29, 30)], and an element of the group

$$\begin{aligned} U &= \exp \left[-i (\lambda_1 \hat{Q}_1 + \lambda_2 \hat{Q}_2 + \sum_i^d \lambda_2^{(i)} \hat{Q}_2^i \right. \\ &\left. + \sum_i^d \lambda_{\text{mix}}^{(i)} \hat{Q}_{\text{mix}}^{i,ii} + \sum_{i \neq j}^d \lambda_2^{(ij)} \hat{Q}_2^{ij} \right] \end{aligned} \quad (31)$$

leads to the following transformations of field operators:

$$\hat{\Phi}'_1 = U \hat{\Phi}_1 U^\dagger = \hat{\Phi}_1 \exp \left[i (\lambda_1 + \sum_i^d \lambda_{\text{mix}}^{(i)} x_i \ell) \right], \quad (32)$$

$$\begin{aligned} \hat{\Phi}'_2 &= U \hat{\Phi}_2 U^\dagger = \hat{\Phi}_2 \exp \left[i (\lambda_2 + \sum_i^d \lambda_2^{(i)} x_i \right. \\ &\left. + \sum_i^d \lambda_{\text{mix}}^{(i)} x_i^2 + \sum_{i \neq j}^d \lambda_2^{(ij)} x_i x_j) \right], \end{aligned} \quad (33)$$

where $\lambda_1, \lambda_2, \lambda_2^{(i)}, \lambda_{\text{mix}}^{(i)}, \lambda_2^{(ij)} \in \mathbb{R}$ are group parameters. The

group \mathcal{G} can be written as the direct product of $U(1)$ groups,

$$\mathcal{G} = \left(\prod_a^2 \otimes U(1)_{a,C} \right) \left(\prod_i \otimes U(1)_{2,i} \right) \left(\prod_i \otimes U(1)_{\text{mix-B},i} \right) \left(\prod_{i \neq j} \otimes U(1)_{2,\{i,j\}} \right), \quad (34)$$

where group elements of $U(1)_{a,C}$, $U(1)_{2,i}$, $U(1)_{\text{mix-B},i}$, and $U(1)_{2,\{i,j\}}$ are respectively labeled by λ_a , $\lambda_2^{(i)}$, $\lambda_{\text{mix}}^{(i)}$, $\lambda_2^{(ij)}$.

A Hamiltonian satisfying this conservation law needs to commute with conserved quantities (Eqs. (27, 28, 29, 30)). We can write down the minimal Hamiltonian of Model Series B with the potential terms,

$$\begin{aligned} \mathcal{H} = & \sum_{i,j}^d K_1^{ij} \left(\hat{\Phi}_1^\dagger \partial_i \partial_j \hat{\Phi}_1^\dagger - \partial_i \hat{\Phi}_1^\dagger \partial_j \hat{\Phi}_1^\dagger \right) \left(\hat{\Phi}_1 \partial_i \partial_j \hat{\Phi}_1 - \partial_i \hat{\Phi}_1 \partial_j \hat{\Phi}_1 \right) \\ & + \sum_{i,j,k}^d K_2^{ijk} \left(\hat{\Phi}_2^{\dagger 2} \partial_i \partial_j \partial_k \hat{\Phi}_2^\dagger - \hat{\Phi}_2^\dagger \partial_i \hat{\Phi}_2^\dagger \partial_j \partial_k \hat{\Phi}_2^\dagger - \hat{\Phi}_2^\dagger \partial_j \hat{\Phi}_2^\dagger \partial_i \partial_k \hat{\Phi}_2^\dagger \right. \\ & \left. - \hat{\Phi}_2^\dagger \partial_k \hat{\Phi}_2^\dagger \partial_i \partial_j \hat{\Phi}_2^\dagger + 2 \partial_i \hat{\Phi}_2^\dagger \partial_j \hat{\Phi}_2^\dagger \partial_k \hat{\Phi}_2^\dagger \right) \\ & \cdot \left(\hat{\Phi}_2^2 \partial_i \partial_j \partial_k \hat{\Phi}_2 - \hat{\Phi}_2 \partial_i \hat{\Phi}_2 \partial_j \partial_k \hat{\Phi}_2 - \hat{\Phi}_2 \partial_j \hat{\Phi}_2 \partial_i \partial_k \hat{\Phi}_2 \right. \\ & \left. - \hat{\Phi}_2 \partial_k \hat{\Phi}_2 \partial_i \partial_j \hat{\Phi}_2 + 2 \partial_i \hat{\Phi}_2 \partial_j \hat{\Phi}_2 \partial_k \hat{\Phi}_2 \right) \\ & + \sum_i^d \Gamma_{1,2}^i \left(\ell \hat{\Phi}_1^\dagger \hat{\Phi}_2^\dagger \partial_i^2 \hat{\Phi}_2^\dagger - \ell \hat{\Phi}_1^\dagger \partial_i \hat{\Phi}_2^\dagger \partial_i \hat{\Phi}_2^\dagger - 2 \partial_i \hat{\Phi}_1^\dagger \hat{\Phi}_2^{\dagger 2} \right) \\ & \cdot \left(\ell \hat{\Phi}_1 \hat{\Phi}_2 \partial_i^2 \hat{\Phi}_2 - \ell \hat{\Phi}_1 \partial_i \hat{\Phi}_2 \partial_i \hat{\Phi}_2 - 2 \partial_i \hat{\Phi}_1 \hat{\Phi}_2^2 \right) + V(\hat{\Phi}_1^\dagger, \hat{\Phi}_1). \end{aligned} \quad (35)$$

For convenience, the coefficients need to satisfy: $K_1^{ij} = K_1^{ji} > 0$, $K_2^{ijk} = K_2^{ikj} = K_2^{jki} = K_2^{kji} = K_2^{kij} = K_2^{kji} > 0$, $\Gamma_{1,2}^i > 0$, and $g > 0$. The $\Gamma_{1,2}^i$ terms are the emerging terms from the hybridization of moments. In the same way as Model Series A, we can also obtain Noether currents of Model Series B. The $\Gamma_{1,2}^i$ terms liberate part of the boson mobility restriction. The effective field theory of this Hamiltonian is performed in Sec. IV.

III. HYBRID FRACTONIC SUPERFLUIDS (MODEL SERIES A)

In the preceding text, we have constructed the minimal Hamiltonians with conservation of hybrid moments. In the present and next sections, we subsequently study physical properties of Model Series A and Model Series B. In this section, we study physical properties of Model Series A. We start our analysis with the $[d, 1, 2, 2]$ model and then directly generalize to arbitrary $[d, N, m, m']$ models.

A. Classical ground state wavefunctions

Classically, the energy density \mathcal{E} of Model Series A has the form as:

$$\begin{aligned} \mathcal{E} = & \sum_a^2 \sum_{i,j}^d K_a^{ij} |\phi_a \partial_i \partial_j \phi_a - \partial_i \phi_a \partial_j \phi_a|^2 \\ & + \sum_{a \neq b}^2 \sum_i^d \Gamma_{a,b}^i |\phi_a \partial_i \phi_b - \phi_b \partial_i \phi_a|^2 \\ & + \sum_a^2 -\mu_a |\phi_a|^2 + \frac{g}{2} |\phi_a|^4. \end{aligned} \quad (36)$$

Considering the chemical potential $\mu_1 > 0$, $\mu_2 > 0$, at $|\phi_a| = \sqrt{\rho_{a0}} = \sqrt{\frac{\mu_a}{g}}$, the potential reaches the minimal value. Then, the energy density is

$$\begin{aligned} \mathcal{E} = & \sum_a^2 \sum_{i,j}^d K_a^{ij} \rho_{a0}^2 (\partial_i \partial_j \theta_a)^2 \\ & + \sum_{a \neq b}^2 \sum_i^d \Gamma_{a,b}^i \rho_{10} \rho_{20} (\partial_i \theta_a - \partial_i \theta_b)^2 - \sum_a^2 \frac{\mu_a^2}{2g}. \end{aligned} \quad (37)$$

As the coefficients $K_a^{ij} > 0$ and $\Gamma_{a,b}^i > 0$, to make these terms reach the minimal value, we need $\partial_i \partial_j \theta_a = 0$ and $\partial_i \theta_a = \partial_i \theta_b$ for any a and b . Then we can give the ground state $|\text{GS}_\beta^{\{\theta_{a0}\}}\rangle = \bigotimes_{\mathbf{x}} |\text{GS}_\beta^{\{\theta_{a0}\}}\rangle_{\mathbf{x}}$, where

$$|\text{GS}_\beta^{\{\theta_{a0}\}}\rangle_{\mathbf{x}} = \frac{1}{C} \prod_a^2 \exp \left[\sqrt{\rho_{a0}} e^{i(\theta_{a0} + \sum_i^d \beta_i x_i)} \hat{\Phi}_a^\dagger(\mathbf{x}) \right] |0\rangle. \quad (38)$$

C is the normalization factor, and $|0\rangle$ is the quantum vacuum state. $\beta = \{\beta_1, \dots, \beta_d\}$ has the dimension of momentum. A significant feature of this ground state is the formation of ODLRO. The correlation functions are given by: $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b(\mathbf{0}) \rangle = \sqrt{\rho_{a0} \rho_{b0}} e^{i(\theta_{b0} - \theta_{a0} - \sum_i \beta_i x_i)}$, $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b^\dagger(\mathbf{0}) \rangle = \rho_{a0} \rho_{b0}$, and $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b(\mathbf{0}) \rangle = \rho_{a0} \rho_{b0} e^{-2i \sum_i \beta_i x_i}$. For the hybrid fractonic superfluid phase in Model Series A, the order parameters are defined as the expectation values on the ground state: $\langle \hat{\Phi}_1(\mathbf{x}) \rangle = \sqrt{\rho_{10}} e^{i(\theta_{10} + \sum_i \beta_i x_i)}$ and $\langle \hat{\Phi}_2(\mathbf{x}) \rangle = \sqrt{\rho_{20}} e^{i(\theta_{20} + \sum_i \beta_i x_i)}$. After condensation, the Noether currents (Eq. (21, 22)) become

$$\begin{aligned} \hat{\rho}_a &= \rho_{a0}, \\ \hat{J}_i^a &= -2K_a^{ij} \rho_{a0}^2 \sum_j^d \partial_i \partial_j^2 \hat{\theta}_a + 4\Gamma_{1,2}^i \rho_{10} \rho_{20} \left(\partial_i \hat{\theta}_1 + \partial_i \hat{\theta}_2 \right), \end{aligned} \quad (39)$$

and

$$\begin{aligned}\hat{\rho}_{\text{mix}}^{(j)} &= (\rho_{10} + \rho_{20}) x_j, \\ \hat{D}_i^{(j)} &= x_j \hat{J}_i^1 + x_j \hat{J}_i^2 + 2 \sum_a K_a^{ij} \rho_{a0}^2 \partial_i \partial_j \hat{\theta}_a.\end{aligned}\quad (40)$$

B. Goldstone modes and quantum fluctuations

Quantum fluctuations tend to destroy true ODLRO. In this subsection, we introduce the effect of quantum fluctuations.

Considering quantum fluctuations, we have $\hat{\Phi}_a(\mathbf{x}) = \sqrt{\rho_{a0} + \hat{f}_a(\mathbf{x})} e^{i\hat{\theta}_a(\mathbf{x})}$, where $\hat{f}_a(\mathbf{x}, t) \ll \rho_{a0}$ are the quantum fluctuations of the density fields, and $\hat{\theta}_a$ is the quantum fluctuations of the phase fields. The commutation relation between $\hat{\theta}_a$ and \hat{f}_a is given by $[\hat{\theta}_a(\mathbf{x}), \hat{f}_b(\mathbf{y})] = -i\delta^{(d)}(\mathbf{x} - \mathbf{y}) \delta_{ab}$. Therefore, the conjugate momentum operator of $\hat{\theta}_a$ is $\hat{\pi}_a = -\hat{f}_a$, and the operators $\hat{\theta}_a$ and $\hat{\pi}_a$ satisfy the commutation relations, $[\hat{\theta}_a(\mathbf{x}), \hat{\pi}_b(\mathbf{y})] = i\delta_{ab}\delta^{(d)}(\mathbf{x} - \mathbf{y})$. To the second order, we can derive the effective Hamiltonian.

$$\begin{aligned}\mathcal{H} &= \sum_a \left\{ \frac{g}{2} \hat{\pi}_a^2 + \sum_{i,j} K_a^{ij} \left[\rho_{a0}^2 (\partial_i \partial_j \hat{\theta}_a)^2 + \frac{(\partial_i \partial_j \hat{\pi}_a)^2}{4} \right] \right. \\ &\quad + \sum_{b \neq a} \sum_i \Gamma_{a,b}^i \left[\frac{(\rho_{a0} \partial_i \hat{\pi}_b - \rho_{b0} \partial_i \hat{\pi}_a)^2}{4\rho_{a0}\rho_{b0}} \right. \\ &\quad \left. \left. + \rho_{a0}\rho_{b0} (\partial_i \hat{\theta}_b - \partial_i \hat{\theta}_a)^2 \right] \right\}.\end{aligned}\quad (41)$$

The time evolutions of $\hat{\theta}_a$ and $\hat{\pi}_a$ are given by

$$\begin{aligned}\hat{\theta}_a &= -i \left[\hat{\theta}_a, \int d^d \mathbf{x} \mathcal{H} \right] = g\hat{\pi}_a + \frac{1}{2} \sum_{i,j} K_a^{ij} \partial_i^2 \partial_j^2 \hat{\pi}_a \\ &\quad - \sum_{b \neq a} \sum_i \Gamma_{a,b}^i \left(\frac{\rho_{b0}}{\rho_{a0}} \partial_i^2 \hat{\pi}_a - \partial_i^2 \hat{\pi}_b \right),\end{aligned}\quad (42)$$

$$\begin{aligned}\hat{\pi}_a &= -i \left[\hat{\pi}_a, \int d^d \mathbf{x} \mathcal{H} \right] = -2 \sum_{i,j} K_a^{ij} \rho_{a0}^2 \partial_i^2 \partial_j^2 \hat{\theta}_a \\ &\quad + 2 \sum_{b \neq a} \sum_i \Gamma_{a,b}^i \rho_{a0} \rho_{b0} (\partial_i^2 \hat{\theta}_a - \partial_i^2 \hat{\theta}_b).\end{aligned}\quad (43)$$

Considering the isotropic case $K_a^{ij} = \frac{\kappa}{2}$, $\Gamma_{a,b}^i = \frac{\Gamma}{2}$, we define the coherence length

$$\xi_c = \pi \sqrt{\frac{2\Gamma}{g}}.\quad (44)$$

In the long-wave length limit, i.e., $|\mathbf{k}| \ll 2\pi\xi_c^{-1}$, we have $g\hat{\pi}_a \gg -\sum_{b \neq a} \sum_i \Gamma_{a,b}^i \partial_i^2 \hat{\pi}_a$, and thus, with finite ρ_{b0}/ρ_{a0} ,

the last term in Eq. (42) is negligible. In addition, the term $\frac{1}{2} \sum_{i,j} K_a^{ij} \partial_i^2 \partial_j^2 \hat{\pi}_a$ with higher-order gradient is also negligible. Thus, we can safely make the approximation: $\hat{\pi}_a = \hat{\theta}_a/g$. Plugging this approximation into Eq. (41), and excluding the higher derivative terms of $\hat{\theta}_a$ and $\hat{\pi}_a$, we can obtain the following effective Hamiltonian

$$\begin{aligned}\mathcal{H} &= \sum_a \left[\frac{g}{2} \hat{\pi}_a^2 + \frac{\kappa \rho_{a0}^2}{2} \sum_{i,j} (\partial_i \partial_j \hat{\theta}_a)^2 \right. \\ &\quad \left. + \frac{\Gamma \rho_{a0} \rho_{b0}}{2} \sum_{b \neq a} \sum_i (\partial_i \hat{\theta}_a - \partial_i \hat{\theta}_b)^2 \right].\end{aligned}\quad (45)$$

By performing a Fourier transform, we obtain the Hamiltonian in the momentum space,

$$H = \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) M(\mathbf{k}) \psi(\mathbf{k}),\quad (46)$$

where we set $\psi(\mathbf{k}) = [\hat{\pi}_1(\mathbf{k}) \ \hat{\pi}_2(\mathbf{k}) \ \hat{\theta}_1(\mathbf{k}) \ \hat{\theta}_2(\mathbf{k})]^T$, $\psi^\dagger(\mathbf{k}) = [\hat{\pi}_1(-\mathbf{k}) \ \hat{\pi}_2(-\mathbf{k}) \ \hat{\theta}_1(-\mathbf{k}) \ \hat{\theta}_2(-\mathbf{k})]$, $M(\mathbf{k}) = \text{diag}(M_1(\mathbf{k}), M_2(\mathbf{k}))$. The matrices $M_1(\mathbf{k})$, $M_2(\mathbf{k})$ are given by $M_1(\mathbf{k}) = \text{diag}(g, g)$ and

$$M_2(\mathbf{k}) = \begin{bmatrix} \kappa \rho_{10}^2 |\mathbf{k}|^4 + 2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 & -2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 \\ -2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 & \kappa \rho_{20}^2 |\mathbf{k}|^4 + 2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 \end{bmatrix}.\quad (47)$$

In the momentum space, the operators $\hat{\theta}_a(\mathbf{k})$ and the conjugate momentum densities $\hat{\pi}_b(\mathbf{k}')$ satisfy the commutation relations: $[\hat{\theta}_a(\mathbf{k}), \hat{\pi}_b(\mathbf{k}')] = i(2\pi)^d \delta_{ab} \delta^{(d)}(\mathbf{k} + \mathbf{k}')$. The derivation of this Hamiltonian (Eq. (46)) is in Appendix A.1. From this Hamiltonian, we can see that two species of bosons couple to each other. By performing the Bogoliubov transformation, we can decouple the coupling bosons in a new basis. For this purpose, we introduce a transformation matrix $T(\mathbf{k})$ which changes the basis from $\psi(\mathbf{k})$ into $\phi(\mathbf{k})$,

$$\phi(\mathbf{k}) = [\hat{\alpha}(\mathbf{k}) \ \hat{\beta}(\mathbf{k}) \ \hat{\alpha}^\dagger(-\mathbf{k}) \ \hat{\beta}^\dagger(-\mathbf{k})]^T = T(\mathbf{k}) \psi(\mathbf{k}),\quad (48)$$

where $\hat{\alpha}(\mathbf{k})$ and $\hat{\beta}(\mathbf{k})$ are annihilation operators of two independent modes, and they satisfy the commutation relations,

$$\begin{aligned}[\hat{\alpha}(\mathbf{k}), \hat{\alpha}^\dagger(\mathbf{k}')] &= [\hat{\beta}(\mathbf{k}), \hat{\beta}^\dagger(\mathbf{k}')] = (2\pi)^d \delta^{(d)}(\mathbf{k} - \mathbf{k}'), \\ [\hat{\alpha}(\mathbf{k}), \hat{\alpha}(\mathbf{k}')] &= [\hat{\beta}(\mathbf{k}), \hat{\beta}(\mathbf{k}')] = [\hat{\alpha}(\mathbf{k}), \hat{\beta}(\mathbf{k}')] \\ &= [\hat{\alpha}(\mathbf{k}), \hat{\beta}^\dagger(\mathbf{k}')] = [\hat{\beta}(\mathbf{k}), \hat{\alpha}^\dagger(\mathbf{k}')] = 0.\end{aligned}\quad (49)$$

The transformation matrix $T(\mathbf{k})$ is given in Appendix A.2. The matrix $T(\mathbf{k})$ diagonalizes the Hamiltonian (Eq. (46)), and the diagonalized Hamiltonian becomes

$$H = \frac{1}{2(2\pi)^d} \phi^\dagger(\mathbf{k}) (T^{-1}(\mathbf{k}))^\dagger M(\mathbf{k}) T^{-1}(\mathbf{k}) \phi(\mathbf{k}),\quad (50)$$

where the matrix $(T^{-1}(\mathbf{k}))^\dagger M(\mathbf{k}) T^{-1}(\mathbf{k}) = \text{diag}(D_1(\mathbf{k}), D_2(\mathbf{k}), D_3(\mathbf{k}), D_4(\mathbf{k}))$ is a diagonal matrix with the diagonal elements:

$$D_1(\mathbf{k}) = D_3(\mathbf{k}) = \left\{ \frac{g}{2} \left[\kappa (\rho_{10}^2 + \rho_{20}^2) |\mathbf{k}|^4 + 4\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 - \sqrt{\kappa^2 (\rho_{10}^2 - \rho_{20}^2) |\mathbf{k}|^8 + 16\Gamma^2 \rho_{10}^2 \rho_{20}^2 |\mathbf{k}|^4} \right] \right\}^{\frac{1}{2}}, \quad (51)$$

$$D_2(\mathbf{k}) = D_4(\mathbf{k}) = \left\{ \frac{g}{2} \left[\kappa (\rho_{10}^2 + \rho_{20}^2) |\mathbf{k}|^4 + 4\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 + \sqrt{\kappa^2 (\rho_{10}^2 - \rho_{20}^2) |\mathbf{k}|^8 + 16\Gamma^2 \rho_{10}^2 \rho_{20}^2 |\mathbf{k}|^4} \right] \right\}^{\frac{1}{2}}. \quad (52)$$

Then, we can obtain the following Hamiltonian formed by the creation and annihilation operators of Goldstone modes,

$$H = \frac{1}{(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}), \quad (53)$$

where the normal ordering is applied. The derivation of this Hamiltonian is in Appendix A 3. The dispersion relations are

$$\omega_1 = D_1(\mathbf{k}) \approx \sqrt{\frac{g}{2} \kappa (\rho_{10}^2 + \rho_{20}^2) |\mathbf{k}|^2}, \quad (54)$$

$$\omega_2 = D_2(\mathbf{k}) \approx \sqrt{4g\Gamma \rho_{10} \rho_{20} |\mathbf{k}|}. \quad (55)$$

C. Correlation functions

To check if the system spontaneously breaks charge symmetry, we need to calculate the correlation functions. For this system with two species of bosons, the two-operator correlation functions can be written as

$$\begin{aligned} \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle &= \rho_{10} e^{-\frac{1}{2} \langle [\hat{\theta}_1(\mathbf{x}) - \hat{\theta}_1(\mathbf{0})]^2 \rangle} \\ &= \rho_{10} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle}, \end{aligned} \quad (56)$$

$$\begin{aligned} \langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle &= \rho_{20} e^{-\frac{1}{2} \langle [\hat{\theta}_2(\mathbf{x}) - \hat{\theta}_2(\mathbf{0})]^2 \rangle} \\ &= \rho_{20} e^{\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle}, \end{aligned} \quad (57)$$

$$\begin{aligned} \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle &= \sqrt{\rho_{10} \rho_{20}} e^{-\frac{1}{2} \langle [\hat{\theta}_1(\mathbf{x}) - \hat{\theta}_2(\mathbf{0})]^2 \rangle} \\ &= \sqrt{\rho_{10} \rho_{20}} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \frac{1}{2} \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \frac{1}{2} \langle \hat{\theta}_2^2(\mathbf{0}) \rangle}. \end{aligned} \quad (58)$$

The four-operator correlation functions can be written as

$$\begin{aligned} &\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left[\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right. \\ &\quad \left. + 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle \right. \\ &\quad \left. - 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right], \end{aligned} \quad (59)$$

$$\begin{aligned} &\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left[\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right. \\ &\quad \left. - 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle \right. \\ &\quad \left. + 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right]. \end{aligned} \quad (60)$$

Using the equations above, we can calculate two-operator and four-operator correlation functions in different spatial dimensions. The calculation results of correlation functions are shown in Table II. The calculation details are in Appendix B.

In these correlation functions, we set $c_1 = \frac{2\sqrt{g\kappa(\rho_{10}^2 + \rho_{20}^2)}}{g}$, $c_2 = \frac{8\sqrt{g\Gamma\rho_{10}\rho_{20}}}{g}$. The coherence length $\xi_c = \pi\sqrt{\frac{2\Gamma}{g}}$ is given by Eq. (44). During the computation, we require that $|\mathbf{x}|$ is much larger than the coherence length ξ_c but much smaller than the system size L .

For the later convenience, we express the total charge symmetry in two equivalent ways:

$$\mathcal{G}_C = U(1)_{1,C} \otimes U(1)_{2,C} = U(1)_{+,C} \otimes U(1)_{-,C}, \quad (61)$$

where $U(1)_{+,C}$ corresponds to the conserved total charges $\hat{Q}_+ = \int d^d \mathbf{x} (\hat{\rho}_1 + \hat{\rho}_2)$ of the whole system of two species of bosons, and $U(1)_{-,C}$ corresponds to the conserved relative charges (i.e., charge difference) $\hat{Q}_- = \int d^d \mathbf{x} (\hat{\rho}_1 - \hat{\rho}_2)$.

In these results, we focus on the correlation functions $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$, $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{0}) \hat{\Phi}_1(\mathbf{0}) \rangle$, and $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle$, which correspond to the order parameters $\langle \hat{\Phi}_1(\mathbf{x}) \rangle$, $\langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \rangle$, and $\langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \rangle$ respectively. We can see that in one spatial dimension, the correlation functions $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ and $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decay exponentially. The correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{0}) \hat{\Phi}_1(\mathbf{0}) \rangle$ decays in a power law. This shows that there is no spontaneous breaking of charge symmetry in one spatial dimension.

In two spatial dimensions, the correlation functions $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ and $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decay in a power law. The correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{0}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant. According to the commutation relations

$$\begin{aligned} \left[\hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}), \hat{Q}_+ \right] &= 0, \\ \left[\hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}), \hat{Q}_- \right] &= 2\hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}), \end{aligned} \quad (62)$$

Table II. Correlation functions of $[d, 1, 2, 2]$ model at zero temperature with quantum fluctuations. This table illustrates the broken charge symmetry of the $[d, 1, 2, 2]$ model and the corresponding order parameters at $d = 1, 2, 3$. The analysis is detailed in Sec. III C. In Sec. III D, we generalize these results to arbitrary $[d, N, m, m']$ models

d	$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$	$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$	$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{0}) \hat{\Phi}_1(\mathbf{0}) \rangle$	$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle$	Broken charge symmetry	Order parameter
1	$\rho_{10} e^{-\frac{ \mathbf{x} }{2c_1} \left(\frac{ \mathbf{x} }{\xi_c}\right)^{-\frac{1}{\pi c_2}}}$	0	$\rho_{10} \rho_{20} \left(\frac{ \mathbf{x} }{\xi_c}\right)^{-\frac{1}{\pi c_2}}$	$\rho_{10} \rho_{20} e^{-\frac{2}{c_1} \mathbf{x} }$	N/A	N/A
2	$\rho_{10} \left(\frac{ \mathbf{x} }{\xi_c}\right)^{-\frac{1}{2\pi c_1}} e^{-\frac{1}{2\pi c_2 \xi_c}}$	$\sqrt{\rho_{10} \rho_{20}} \left(\frac{ \mathbf{x} }{\xi_c}\right)^{-\frac{1}{2\pi c_1}} e^{-\frac{1}{2\pi c_2 \xi_c}}$	$\rho_{10} \rho_{20} e^{-\frac{2}{\pi c_2 \xi_c}}$	$\rho_{10} \rho_{20} \left(\frac{ \mathbf{x} }{\xi_c}\right)^{-\frac{2}{\pi c_1}}$	$U(1)_{-,C}$	$\langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \rangle$
3	$\rho_{10} e^{-\frac{1}{4\pi c_1 \xi_c} e^{-\frac{1}{c_2 \xi_c^2}}}$	$\sqrt{\rho_{10} \rho_{20}} e^{-\frac{1}{4\pi c_1 \xi_c} e^{-\frac{1}{c_2 \xi_c^2}}}$	$\rho_{10} \rho_{20} e^{-\frac{4}{c_2 \xi_c^2}}$	$\rho_{10} \rho_{20} e^{-\frac{1}{\pi c_1 \xi_c}}$	$U(1)_{1,C} \otimes U(1)_{2,C}$	$\langle \hat{\Phi}_1(\mathbf{x}) \rangle, \langle \hat{\Phi}_2(\mathbf{x}) \rangle$

the system breaks the relative charge symmetry $U(1)_{-,C}$, and has a true ODLRO.

In three spatial dimensions and above, all the correlation functions mentioned above are constants at long distances. This means that the charge symmetry is completely broken in three spatial dimensions.

As a summary, the charge symmetry is not broken in one spatial dimension, and is completely broken in three spatial dimension and above. In two spatial dimensions, a partial breaking of the charge symmetry occurs.

D. Arbitrary $[d, N, m, m']$ models of Model Series A

In this subsection, we generalize the above analysis to $[d, N, m, m']$ models of Model Series A with the conserved quantities (Eq. (7, 8, 9, 10)). The corresponding Hamiltonian density is Eq. (14). In this case, we focus on an isotropic coupling constant $K_a^{i_1 \dots i_{N+1}} = \frac{\kappa}{2}, \Gamma_{a,b}^{i_1 \dots i_N} = \frac{\Gamma}{2}$. For simplicity, we take the chemical potential $\mu_a = \mu > 0$, and then we have degenerate vacuum with finite uniform density distribution $\rho_a = \rho_0$. We can also derive an effective theory for the quantum fluctuation field $\hat{\theta}_a$ after condensation,

$$\mathcal{H} = \sum_a^m \left[\frac{g}{2} \hat{\pi}_a^2 + \frac{\kappa \rho_0^{N+1}}{2} \left(\nabla^{N+1} \hat{\theta}_a \right)^2 \right] + \frac{\Gamma \rho_0^{2N}}{2} \sum_{a \neq b}^{m'} \sum_{i_1, \dots, i_N}^d \left(\partial_{i_1} \dots \partial_{i_N} \hat{\theta}_a - \partial_{i_1} \dots \partial_{i_N} \hat{\theta}_b \right)^2. \quad (63)$$

Through the same processes, we can obtain the Hamiltonian in the momentum space

$$H = \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) M(\mathbf{k}) \psi(\mathbf{k}), \quad (64)$$

where

$$\psi(\mathbf{k}) = [\hat{\pi}_1(\mathbf{k}) \dots \hat{\pi}_m(\mathbf{k}) \hat{\theta}_1(\mathbf{k}) \dots \hat{\theta}_m(\mathbf{k})]^T, \quad (65)$$

$$M(\mathbf{k}) = \text{diag}(M_1(\mathbf{k}), M_2(\mathbf{k}), M_3(\mathbf{k})). \quad (66)$$

The matrix $M_1(\mathbf{k})$ is a $m \times m$ diagonal matrix whose diagonal elements are g . The diagonal elements of the $m' \times m'$ matrix $M_2(\mathbf{k})$ are $[\kappa \rho_0^{N+1} |\mathbf{k}|^{2N+2} + 2(m' - 1) \Gamma \rho_0^{2N} |\mathbf{k}|^{2N}]$, and

all the off-diagonal elements are $[-2\Gamma \rho_0^{2N} |\mathbf{k}|^{2N}]$. $M_3(\mathbf{k})$ is a diagonal $(m - m') \times (m - m')$ matrix whose diagonal elements are $[\kappa \rho_0^{N+1} |\mathbf{k}|^{2N+2}]$. By diagonalizing this Hamiltonian, we end up with the dispersion relations:

$$\omega_a = \sqrt{\kappa g \rho_0^{N+1} |\mathbf{k}|^{N+1}} \quad (1 \leq a \leq m - m' + 1), \quad (67)$$

$$\omega_a = \sqrt{\kappa g \rho_0^{N+1} |\mathbf{k}|^{2N+2} + 2m' g \Gamma \rho_0^{2N} |\mathbf{k}|^{2N}} \approx \sqrt{2m' g \Gamma \rho_0^{2N} |\mathbf{k}|^N} \quad (m - m' + 2 \leq a \leq m). \quad (68)$$

The two-operator correlation function can be written as

$$\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \rangle = \rho_0 e^{-\frac{1}{2} \langle [\hat{\theta}_a(\mathbf{x}) - \hat{\theta}_a(\mathbf{0})]^2 \rangle} = \rho_0 e^{\langle \hat{\theta}_a(\mathbf{x}) \hat{\theta}_a(\mathbf{0}) \rangle - \langle \hat{\theta}_a^2(\mathbf{0}) \rangle}. \quad (69)$$

The four-operator correlation functions can be written as

$$\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b^\dagger(\mathbf{0}) \rangle = \rho_0^2 \exp \left[\langle \hat{\theta}_a(\mathbf{x}) \hat{\theta}_a(\mathbf{0}) \rangle + \langle \hat{\theta}_b(\mathbf{x}) \hat{\theta}_b(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_a(\mathbf{0}) \hat{\theta}_b(\mathbf{0}) \rangle - \langle \hat{\theta}_a^2(\mathbf{0}) \rangle - \langle \hat{\theta}_b^2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_a(\mathbf{x}) \hat{\theta}_b(\mathbf{0}) \rangle \right], \quad (70)$$

$$\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b(\mathbf{0}) \rangle = \rho_0^2 \exp \left[\langle \hat{\theta}_a(\mathbf{x}) \hat{\theta}_a(\mathbf{0}) \rangle + \langle \hat{\theta}_b(\mathbf{x}) \hat{\theta}_b(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_a(\mathbf{0}) \hat{\theta}_b(\mathbf{0}) \rangle - \langle \hat{\theta}_a^2(\mathbf{0}) \rangle - \langle \hat{\theta}_b^2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_a(\mathbf{x}) \hat{\theta}_b(\mathbf{0}) \rangle \right]. \quad (71)$$

If the species of bosons satisfy the conditions $1 \leq a, b \leq m'$ and $a \neq b$, the exponent of e in Eq. (70) is a integral of a function which is proportional to $(1/|\mathbf{k}|^N)$. The correlation function $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b^\dagger(\mathbf{0}) \rangle$ saturates to a constant in spatial dimensions $d \geq N + 1$. For the correlation functions $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \rangle$ and $\langle \hat{\Phi}_a^\dagger(\mathbf{x}) \hat{\Phi}_b^\dagger(\mathbf{x}) \hat{\Phi}_a(\mathbf{0}) \hat{\Phi}_b(\mathbf{0}) \rangle$, in the exponent of e , the integrand is the sum of a function proportional to $(1/|\mathbf{k}|^N)$ and another function proportional to $(1/|\mathbf{k}|^{N+1})$. The two correlation functions saturate to a constant only in dimensions $d \geq N + 2$. This implies a partial spontaneous breaking of charge symmetry in spatial dimensions $d = N + 1$. The system breaks the relative charge symmetry of the bosons involved in the hybridization.

IV. HYBRID FRACTONIC SUPERFLUIDS (MODEL SERIES B)

In this section, we focus on the details in Model Series B. Based on the Hamiltonian (Eq. (35)) of a hybrid dipole-quadrupole conserving model, we present the ground state along with its Goldstone modes and calculate correlation functions to determine at what minimum spatial dimension this model will break charge symmetry and have a true ODLRO.

A. Classical ground state wavefunctions

Classically, the energy density \mathcal{E} of the Hamiltonian (Eq. (35)) has the form as:

$$\begin{aligned} \mathcal{E} = & \sum_{i,j}^d K_1^{ij} |\phi_1 \partial_i \partial_j \phi_1 - \partial_i \phi_1 \partial_j \phi_2|^2 \\ & + \sum_{i,j,k}^d K_2^{ijk} \left| \phi_2^2 \partial_i \partial_j \partial_k \phi_2 - \phi_2 \partial_i \phi_2 \partial_j \partial_k \phi_2 \right. \\ & \left. - \phi_2 \partial_j \phi_2 \partial_i \partial_k \phi_2 - \phi_2 \partial_k \phi_2 \partial_i \partial_j \phi_2 + 2 \partial_i \phi_2 \partial_j \phi_2 \partial_k \phi_2 \right|^2 \\ & + \sum_i^d \Gamma_{1,2}^i |\ell \phi_1 \phi_2 \partial_i^2 \phi_2 - \ell \phi_1 \partial_i \phi_2 \partial_i \phi_2 - 2 \partial_i \phi_1 \phi_2^2|^2 \\ & - \mu_1 |\phi_1|^2 - \mu_2 |\phi_2|^2 + \frac{g}{2} |\phi_1|^4 + \frac{g}{2} |\phi_2|^4. \end{aligned} \quad (72)$$

If $\mu_1, \mu_2 > 0$, at $|\phi_1| = \sqrt{\rho_{10}} = \sqrt{\frac{\mu_1}{g}}$, $|\phi_2| = \sqrt{\rho_{20}} = \sqrt{\frac{\mu_2}{g}}$, the potential reaches the minimal value,

$$\begin{aligned} \mathcal{E} = & \sum_{i,j}^d \rho_{10}^2 (\partial_i \partial_j \theta_1)^2 + \sum_{i,j,k}^d \rho_{20}^3 (\partial_i \partial_j \partial_k \theta_2)^2 \\ & + \sum_i^d \rho_{10} \rho_{20}^2 (2 \partial_i \theta_1 - \ell \partial_i^2 \theta_2)^2 - \frac{\mu_1^2}{2g} - \frac{\mu_2^2}{2g}. \end{aligned} \quad (73)$$

To minimize the kinetic terms, we need to satisfy 3 equations: $\partial_i \partial_j \theta_1 = 0$, $\partial_i \partial_j \partial_k \theta_2 = 0$, and $2 \partial_i \theta_1 - \ell \partial_i^2 \theta_2 = 0$ for any i , j , and k . For the first two equations, we can give the form of θ_a as $\theta_1 = \theta_{10} + \sum_i^d \beta_i^{(1)} x_i$, and $\theta_2 = \theta_{20} + \sum_{i,j}^d \beta_{ij}^{(2)} x_i x_j$. And the last condition gives another limit of $\beta_i^{(1)}$ and $\beta_{ij}^{(2)}$: $2\beta_i^{(1)} - \ell \beta_{ii}^{(2)} = 0$. Thus, we can give the ground state,

$$\begin{aligned} & |\text{GS}_{\beta^{(1)}, \beta^{(2)}}^{\{\theta_{a0}\}} \rangle \\ = & \bigotimes_{\mathbf{x}} \frac{1}{C} \exp \left[\sqrt{\rho_{10}} e^{i(\theta_{10} + \sum_i^d \beta_i^{(1)} x_i)} \hat{\Phi}_1^\dagger(\mathbf{x}) \right] \\ & \exp \left[\sqrt{\rho_{20}} e^{i(\theta_{20} + \sum_{i,j}^d \beta_{ij}^{(2)} x_i x_j)} \hat{\Phi}_2^\dagger(\mathbf{x}) \right] |0\rangle. \end{aligned} \quad (74)$$

This ground state also has the formation of ODLRO. We can calculate correlation functions in the classical level,

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle = \rho_{10} e^{-i \sum_i^d \beta_i^{(1)} x_i}, \quad (75)$$

$$\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \rho_{20} e^{-i \sum_{i,j}^d \beta_{ij}^{(2)} x_i x_j}. \quad (76)$$

The order parameter can be determined with finite expectation value on the ground state,

$$\langle \hat{\Phi}_1(\mathbf{x}) \rangle = \sqrt{\rho_{10}} e^{i(\theta_{10} + \sum_i^d \beta_i^{(1)} x_i)}, \quad (77)$$

$$\langle \hat{\Phi}_2(\mathbf{x}) \rangle = \sqrt{\rho_{20}} e^{i(\theta_{20} + \sum_{i,j}^d \beta_{ij}^{(2)} x_i x_j)}. \quad (78)$$

B. Goldstone modes and quantum fluctuations

Considering quantum fluctuations, we have $\hat{\Phi}_a(\mathbf{x}) = \sqrt{\rho_{a0} + \hat{f}_a(\mathbf{x})} e^{i\hat{\theta}_a(\mathbf{x})}$, where $\hat{f}_a(\mathbf{x}) \ll \rho_{a0}$. Through the same processes in Model Series A, we can obtain the effective Hamiltonian under the long-wave approximation,

$$\begin{aligned} \mathcal{H} = & \frac{g}{2} \hat{\pi}_1^2 + \frac{g}{2} \hat{\pi}_2^2 + \frac{\kappa_1 \rho_{10}^2}{2} \sum_{i,j}^d \hat{\theta}_1 \partial_i^2 \partial_j^2 \hat{\theta}_1 \\ & - \frac{\kappa_2 \rho_{20}^3}{2} \sum_{i,j,k}^d \hat{\theta}_2 \partial_i^2 \partial_j^2 \partial_k^2 \hat{\theta}_2 + \frac{\Gamma \rho_{10} \rho_{20}^2}{2} \sum_i^d \left(\ell^2 \hat{\theta}_2 \partial_i^4 \hat{\theta}_2 \right. \\ & \left. - 4 \hat{\theta}_1 \partial_i^2 \hat{\theta}_1 + 2 \ell \hat{\theta}_1 \partial_i^3 \hat{\theta}_2 - 2 \ell \hat{\theta}_2 \partial_i^3 \hat{\theta}_1 \right), \end{aligned} \quad (79)$$

where we set $K_1^{ij} = \frac{\kappa_1}{2}$, $K_2^{ijk} = \frac{\kappa_2}{2}$, and $\Gamma_{1,2}^i = \frac{\Gamma}{2}$. $\hat{\pi}_a = -\hat{f}_a$ are the conjugate momentum densities of $\hat{\theta}_a$. Continuing the derivation process, we can finally obtain the regularized quantized and diagonalized Hamiltonian,

$$H = \frac{1}{(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}), \quad (80)$$

where the normal ordering is applied. The operators in Model Series B above still satisfies the same commutation relations as in Model Series A. The dispersion relations are

$$\omega_1 = D_1(\mathbf{k}), \quad \omega_2 = D_2(\mathbf{k}). \quad (81)$$

The dispersion relations of Model Series B are complex and anisotropic and are given in Appendix C18. We can only approximate the dispersion relations for given spatial dimensions. One of the dispersion relations is always linear,

$$\omega_1 \approx \sqrt{4g\Gamma\rho_{10}\rho_{20}^2} |\mathbf{k}|. \quad (82)$$

The other one of the dispersion relations is anisotropic. In one spatial dimension, we have

$$\omega_2 \approx \sqrt{\kappa_2 g \rho_{20}^3 + \frac{\kappa_1 g \rho_{10}^2 \ell^2}{4}} |k|^3. \quad (83)$$

In two or more spatial dimensions, except for some special directions, we have

$$\omega_2 \propto |\mathbf{k}|^2. \quad (84)$$

Since the derivation process is similar to that of Model Series A, we put the detailed derivation of Model Series B in Appendix C. The derivation of the dispersion relations in different spatial dimensions is given in Appendix D.

C. Correlation functions

To check the effect of hybridization in Model Series B on ODLRO, we calculate correlation functions. Unlike Model Series A, we only need two-operator correlation functions to characterize the spontaneous breaking of charge symmetry in Model Series B instead of four-operator correlation functions. The two-operator correlation functions can be written as

$$\begin{aligned} \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle &= \rho_{10} e^{-\frac{1}{2} \langle [\hat{\theta}_1(\mathbf{x}) - \hat{\theta}_1(\mathbf{0})]^2 \rangle} \\ &= \rho_{10} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle}, \end{aligned} \quad (85)$$

$$\begin{aligned} \langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle &= \rho_{20} e^{-\frac{1}{2} \langle [\hat{\theta}_2(\mathbf{x}) - \hat{\theta}_2(\mathbf{0})]^2 \rangle} \\ &= \rho_{20} e^{\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle}. \end{aligned} \quad (86)$$

Then, we can calculate the correlation function of the two species of bosons. Due to the anisotropy, we can just give a general trend of the correlation functions in two or more spatial dimensions. The calculation results of correlation functions are shown in Table III. The details of the analytical calculations and numerical calculations are in Appendix D. In these correlation functions, we set $c_{11} = \frac{4\Gamma\rho_{10}\rho_{20}}{\sqrt{g\Gamma\rho_{10}}}$, $c_{21} = \frac{4\sqrt{\kappa_1\rho_{10}^2\ell^2 + 4\kappa_2\rho_{20}^3}}{\sqrt{g\ell^2}}$ and the coherence length $\xi_c = 2\pi\sqrt{\frac{\Gamma\rho_{20}^2}{g\rho_{10}}}$.

Table III. Correlation functions of the hybrid dipole-quadrupole moment conserving model (Eq. (35)) at zero temperature with quantum fluctuations. This table illustrates the broken charge symmetry of this model and the corresponding order parameters at $d = 1, 2, 3$. The analysis is detailed in Sec. IV C.

d	$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$	$\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$	Broken charge symmetry	Order parameter
1	$\rho_{10} \left(\frac{ x }{\xi_c} \right)^{-\frac{c_{11}+c_{21}}{\pi c_{11}c_{21}}}$	0	N/A	N/A
2	$\rho_{10} e^{-\frac{c_{11}+c_{21}}{2\pi c_{11}c_{21}\xi_c}}$	power-law decay	$U(1)_{1,C}$	$\langle \hat{\Phi}_1(\mathbf{x}) \rangle$
3	constant	constant	$U(1)_{1,C} \otimes U(1)_{2,C}$	$\langle \hat{\Phi}_1(\mathbf{x}) \rangle, \langle \hat{\Phi}_2(\mathbf{x}) \rangle$

Through the calculations, we confirm that the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ decays in a power law in one spatial dimension and saturates a constant in two or more spatial dimensions. The correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decays exponentially in one spatial dimension, decays in a power law in two spatial dimensions, and saturates a constant in three or more spatial dimensions. The constant behaviors of these two correlation functions represent the spontaneous breaking of $U(1)_{1,C}$ symmetry and $U(1)_{2,C}$ symmetry respectively.

Therefore, this system has a true ODLRO in two spatial dimensions and above. Compared with dipole-conserving and quadrupole-conserving fractonic superfluid models with no hybridization, the spontaneous breaking of the charge symmetry $U(1)_{1,C}$ and $U(1)_{2,C}$ occurs in lower spatial dimensions. This decrease in dimension comes from the hybridization of conserved quantities as well as the coupling of the two species of bosons.

V. BOSE-HUBBARD-TYPE LATTICE MODELS

A lattice model whose lower energy physics is described by fractonic superfluid model with no hybridization [44] is constructed in the article [59]. The Hamiltonians in the two articles [44, 59] can be connected by the difference method. In the previous sections, we discuss the low energy physics of HSF (Eq. (20, 35)) by using effective field theory. In this section, we will use the difference method to obtain lattice models that satisfies the aforementioned conservation laws.

A. Construction of Lattice Model Series A

We specifically study the lattice model corresponding to the $[d, 1, 2, 2]$ model. From the Hamiltonian (Eq. (20)), one of the kinetic terms is given by

$$\left(\hat{\Phi}_a^\dagger \partial_i \partial_j \hat{\Phi}_a^\dagger - \partial_i \hat{\Phi}_a^\dagger \partial_j \hat{\Phi}_a^\dagger \right) \left(\hat{\Phi}_a \partial_i \partial_j \hat{\Phi}_a - \partial_i \hat{\Phi}_a \partial_j \hat{\Phi}_a \right). \quad (87)$$

We can set

$$\begin{aligned} \partial_i \hat{\Phi}_a(\mathbf{x}) &\rightarrow \frac{1}{\eta} \left(\hat{b}_{a, \mathbf{i} + \hat{x}_i} - \hat{b}_{a, \mathbf{i}} \right), \\ \partial_i \partial_j \hat{\Phi}_a(\mathbf{x}) &\rightarrow \frac{1}{\eta^2} \left(\hat{b}_{a, \mathbf{i} + \hat{x}_i + \hat{x}_j} - \hat{b}_{a, \mathbf{i} + \hat{x}_i} - \hat{b}_{a, \mathbf{i} + \hat{x}_j} + \hat{b}_{a, \mathbf{i}} \right), \end{aligned} \quad (88)$$

where in the lattice model we use \hat{x}_i to mark the spatial unit vectors. In the lattice model, $\mathbf{i} = \frac{1}{\eta}(x_1, \dots, x_d)$ represent the coordinates of lattice sites. The quantity η represents the lattice constant of the lattice model. Then, we can obtain the hopping terms in Lattice Model Series A,

$$\begin{aligned} &\left(\hat{\Phi}_a^\dagger \partial_i \partial_j \hat{\Phi}_a^\dagger - \partial_i \hat{\Phi}_a^\dagger \partial_j \hat{\Phi}_a^\dagger \right) \left(\hat{\Phi}_a \partial_i \partial_j \hat{\Phi}_a - \partial_i \hat{\Phi}_a \partial_j \hat{\Phi}_a \right) \\ &\rightarrow -\frac{1}{\eta^4} \left(\hat{b}_{a, \mathbf{i}}^\dagger \hat{b}_{a, \mathbf{i} + \hat{x}_i + \hat{x}_j}^\dagger \hat{b}_{a, \mathbf{i} + \hat{x}_i} \hat{b}_{a, \mathbf{i} + \hat{x}_j} + \text{h.c.} \right) + \text{onsite terms.} \end{aligned} \quad (89)$$

For the kinetic terms

$$\left(\hat{\Phi}_a^\dagger \partial_i \hat{\Phi}_b^\dagger - \hat{\Phi}_b^\dagger \partial_i \hat{\Phi}_a^\dagger \right) \left(\hat{\Phi}_a \partial_i \hat{\Phi}_b - \hat{\Phi}_b \partial_i \hat{\Phi}_a \right), \quad (90)$$

by the difference method (Eq. (88)), this kinetic terms become

$$\begin{aligned} &\left(\hat{\Phi}_a^\dagger \partial_i \hat{\Phi}_b^\dagger - \hat{\Phi}_b^\dagger \partial_i \hat{\Phi}_a^\dagger \right) \left(\hat{\Phi}_a \partial_i \hat{\Phi}_b - \hat{\Phi}_b \partial_i \hat{\Phi}_a \right) \\ &\rightarrow -\frac{1}{\eta^2} \left(\hat{b}_{b, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{a, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{a, \mathbf{i}}^\dagger \hat{b}_{b, \mathbf{i}} + \text{h.c.} \right) + \text{onsite terms.} \end{aligned} \quad (91)$$

Thus, we can give the lattice model which conserves both charges of bosons of two species and the hybrid moments (Eq. (16)),

$$H = H_{\text{hop}} + H_{\text{onsite}}, \quad (92)$$

where

$$H_{\text{hop}} = \sum_a^2 \left[\sum_{\mathbf{i}, \mathbf{i}, \mathbf{j}} -t_a^{(ij)} \hat{b}_{a, \mathbf{i} + \hat{x}_i + \hat{x}_j}^\dagger \hat{b}_{a, \mathbf{i}}^\dagger \hat{b}_{a, \mathbf{i} + \hat{x}_i} \hat{b}_{a, \mathbf{i} + \hat{x}_j} \right. \\ \left. + \sum_{b \neq a}^2 \sum_{\mathbf{i}, \mathbf{i}} -t_{ab}^{(i)} \hat{b}_{b, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{a, \mathbf{i} + \hat{x}_i} \hat{b}_{a, \mathbf{i}}^\dagger \hat{b}_{b, \mathbf{i}} \right] + \text{h.c.} \quad (93)$$

$$H_{\text{onsite}} = \sum_a^2 \left[-\mu_a \sum_{\mathbf{i}} \hat{n}_{a, \mathbf{i}} + \frac{U}{2} \sum_{\mathbf{i}} \hat{n}_{a, \mathbf{i}} (\hat{n}_{a, \mathbf{i}} - 1) \right]. \quad (94)$$

The $t_a^{(ij)}$ terms and the $t_{ab}^{(i)}$ terms in Lattice Model Series A correspond to the K_a^{ij} terms and the $\Gamma_{a,b}^i$ terms in Model Series A respectively. And, we have $t_a^{(ij)} = t_a^{(ji)} > 0$, $t_{ab}^{(i)} = t_{ba}^{(i)} > 0$. The diagonal and off-diagonal $t_a^{(ij)}$ terms are corresponding to the t and t' terms in Ref. [59] respectively. We can give the illustration (Fig. 1(a), 1(b), and 1(c)) of the hopping terms of Lattice Model Series A.

The mean-field theory of this lattice model is performed, which is in Appendix E. The mean-field results imply the existence of phases with condensates of bound states $\hat{b}_{1, \mathbf{i}} \hat{b}_{2, \mathbf{i}}^\dagger$. To derive the global phase diagram beyond meanfield theory, it will be meaningful to employ more advanced treatments, e.g., large-scale Monte-Carlo simulations.

B. Construction of Lattice Model Series B

In this subsection, we give a lattice model of the Hamiltonian (Eq. (35)). The hopping terms corresponding to the K_1^{ij} term are similar to the hopping term $t_a^{(ij)}$ in lattice Model Series A. In addition, the hopping terms corresponding to the K_2^{ijk} include more than 6 operators, so we won't discuss them. We will just discuss the hopping terms which are corresponding to the coupling terms $\Gamma_{a,b}^i$. For the kinetic terms

$$\left(\hat{\ell} \hat{\Phi}_1^\dagger \hat{\Phi}_2^\dagger \partial_i^2 \hat{\Phi}_2^\dagger - \hat{\ell} \hat{\Phi}_1^\dagger \partial_i \hat{\Phi}_2^\dagger \partial_i \hat{\Phi}_2^\dagger - 2 \partial_i \hat{\Phi}_1^\dagger \hat{\Phi}_2^{\dagger 2} \right) \\ \left(\hat{\ell} \hat{\Phi}_1 \hat{\Phi}_2 \partial_i^2 \hat{\Phi}_2 - \hat{\ell} \hat{\Phi}_1 \partial_i \hat{\Phi}_2 \partial_i \hat{\Phi}_2 - 2 \partial_i \hat{\Phi}_1 \hat{\Phi}_2^2 \right), \quad (95)$$

we set

$$\ell \rightarrow 2\eta, \quad \partial_i \hat{\Phi}_1(\mathbf{x}) \rightarrow \frac{1}{\eta} (\hat{b}_{1, \mathbf{i} + \hat{x}_i} - \hat{b}_{1, \mathbf{i}}), \\ \partial_i \hat{\Phi}_2(\mathbf{x}) \rightarrow \frac{1}{\eta} (\hat{b}_{2, \mathbf{i} + \hat{x}_i} - \hat{b}_{2, \mathbf{i}}) \text{ or } \frac{1}{\eta} (\hat{b}_{2, \mathbf{i}} - \hat{b}_{2, \mathbf{i} - \hat{x}_i}), \\ \partial_i^2 \hat{\Phi}_2(\mathbf{x}) \rightarrow \frac{1}{\eta^2} (\hat{b}_{2, \mathbf{i} + \hat{x}_i} - 2\hat{b}_{2, \mathbf{i}} + \hat{b}_{2, \mathbf{i} - \hat{x}_i}). \quad (96)$$




Fig. 1. Illustration of the hopping terms (Eq. (93, 100)) on the square lattice. In Fig. 1(a), 1(b), and 1(c), the balls of the same color represent bosons of the same species, and the balls of different colors represent bosons of different species. The hopping shown in Fig. 1(a) corresponds to diagonal $t_a^{(ij)}$ terms of Eq. (93). The hopping shown in Fig. 1(b) corresponds to off-diagonal $t_a^{(ij)}$ terms of Eq. (93). The hopping shown in Fig. 1(c) corresponds to $t_{ab}^{(i)}$ terms of Eq. (93). In Fig. 1(d), the blue ball represents one boson of species 1, and the red balls represent bosons of species 2. The hopping shown in Fig. 1(d) corresponds to the $t_{12}^{(i)}$ terms of Eq. (100)

Then, we can get the correspondence from the continuous field theory to the lattice model,

$$\hat{\ell} \hat{\Phi}_1 \hat{\Phi}_2 \partial_i^2 \hat{\Phi}_2 - \hat{\ell} \hat{\Phi}_1 \partial_i \hat{\Phi}_2 \partial_i \hat{\Phi}_2 - 2 \partial_i \hat{\Phi}_1 \hat{\Phi}_2^2 \\ \rightarrow \frac{2}{\eta} \left(\hat{b}_{1, \mathbf{i}} \hat{b}_{2, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i} - \hat{x}_i} - \hat{b}_{1, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i}}^2 \right). \quad (97)$$

The lattice hopping terms become

$$\left(\hat{b}_{1, \mathbf{i}}^\dagger \hat{b}_{2, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{2, \mathbf{i} - \hat{x}_i}^\dagger - \hat{b}_{1, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{2, \mathbf{i}}^{\dagger 2} \right) \\ \left(\hat{b}_{1, \mathbf{i}} \hat{b}_{2, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i} - \hat{x}_i} - \hat{b}_{1, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i}}^2 \right) \\ = -\hat{b}_{1, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{1, \mathbf{i}} \hat{b}_{2, \mathbf{i}}^{\dagger 2} \hat{b}_{2, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i} - \hat{x}_i} + \text{h.c.} + \text{onsite terms.} \quad (98)$$

This lattice model which conserves charges of two species, the hybrid moment, and other lower moments (Eq. (27, 28, 29, 30)) is given by

$$H = H_{\text{hop}} + H_{\text{onsite}}, \quad (99)$$

where

$$H_{\text{hop}} = \left[\sum_{\mathbf{i}, \mathbf{i}, \mathbf{j}} -t_1^{(ij)} \hat{b}_{1, \mathbf{i} + \hat{x}_i + \hat{x}_j}^\dagger \hat{b}_{1, \mathbf{i}}^\dagger \hat{b}_{1, \mathbf{i} + \hat{x}_i} \hat{b}_{1, \mathbf{i} + \hat{x}_j} \right. \\ \left. + \sum_{\mathbf{i}, \mathbf{i}} -t_{12}^{(i)} \hat{b}_{1, \mathbf{i} + \hat{x}_i}^\dagger \hat{b}_{1, \mathbf{i}} \hat{b}_{2, \mathbf{i}}^{\dagger 2} \hat{b}_{2, \mathbf{i} + \hat{x}_i} \hat{b}_{2, \mathbf{i} - \hat{x}_i} \right] + \text{h.c.} \quad (100)$$

$$H_{\text{onsite}} = \sum_a^2 \left[-\mu_a \sum_i \hat{n}_{a,i} + \frac{U}{2} \sum_i \hat{n}_{a,i} (\hat{n}_{a,i} - 1) \right]. \quad (101)$$

The $t_1^{(ij)}$ terms and the $t_{12}^{(i)}$ terms correspond to the K_1^{ij} terms and the $\Gamma_{1,2}^i$ terms in Model Series B (Eq. (35)). The K_2^{ijk} terms of Model Series B are ignored in Lattice Model Series B because there are too many operators. We can give the illustration (Fig. 1(d)) of the $t_{12}^{(i)}$ terms (Eq. (100)). Under a specific choice of coefficients, there would be bound states condensate formed by one boson of species 1, one boson of species 2, and one hole of species 1.

C. Realizations of Lattice Model Series A in strongly tilted optical lattice

We can give a pathway to realize the Hamiltonian (Eq. (92)) of Lattice Model Series A in strongly tilted optical lattices. We start with a microscopic Hamiltonian with two species of bosons in a tilted optical lattice chain. Since there is only one direction, we can simplify the notation $\hat{x}_1 = 1$ and write the Hamiltonian as:

$$H_{\text{ilted}} = -t \sum_{a,i} \left(\hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} + \text{h.c.} \right) - \mu \sum_{a,i} \hat{n}_{a,i} + \frac{U}{2} \sum_{a,i} \hat{n}_{a,i} (\hat{n}_{a,i} - 1) + U' \sum_i \hat{n}_{1,i} \hat{n}_{2,i} + V \sum_{a,i} i \hat{n}_{a,i}, \quad (102)$$

where t , μ , and U terms are just terms in Bose-Hubbard model, and the U' term denotes the interaction between bosons of different species. V is the strength of the tilt potential. In the limit $V \gg t, U$, we perform a third-order perturbation theory. The effective Hamiltonian becomes

$$H_{\text{eff}} = \sum_{a,i} \left(-\frac{t^2 U}{V^2} \hat{b}_{a,i} \hat{b}_{a,i+1}^\dagger \hat{b}_{a,i+2} + \text{h.c.} \right) + \sum_{a \neq b, i} \left(\frac{t^2 U'}{V^2} \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} \hat{b}_{b,i+1} \hat{b}_{b,i+2}^\dagger - \frac{t^2 U'}{V^2} \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} \hat{b}_{b,i} \hat{b}_{b,i+1}^\dagger + \text{h.c.} \right) + \sum_{a,i} \left(-\mu - \frac{U}{2} + V i \right) \hat{n}_{a,i} + \sum_{a,i} \left(\frac{U}{2} - \frac{2t_0^2 U}{V^2} \right) \hat{n}_{a,i}^2 + \frac{4t_0^2 U}{V^2} \hat{n}_{a,i} \hat{n}_{a,i+1} + \sum_{a \neq b, i} \left[\left(\frac{U'}{2} - \frac{2t_0^2 U'}{V^2} \right) \hat{n}_{a,i} \hat{n}_{b,i} + \frac{2t_0^2 U'}{V^2} \hat{n}_{a,i} \hat{n}_{b,i+1} \right]. \quad (103)$$

The detailed derivation of Eq. (103) is given in Appendix F. The two terms under the second summation of Eq. (103) commute with total dipole moments, but do not commute with dipole moments of bosons of separate series. From this, we find a way to realize Lattice Model Series A in a tilted lattice.

VI. SUMMARY AND OUTLOOK

In conclusion, this article serves as the third piece of the trilogy of fractonic superfluids in order to uncover the exotic interplay of SSB physics and fracton physics through most of representative models and key properties of fractonic superfluids. Specifically, in this paper, we study HFS where hybrid moments are conserved. We select two typical types of hybridization, i.e. hybridization of the same order moments and hybridization of different order moments. In continuous space, we construct two series of Hamiltonians which correspond to these two typical hybridizations, and denote them by Model Series A and Model Series B. Arbitrary models of Model Series A can be represented by a tuple $[d, N, m, m']$. The simplest example of Model Series A is a system conserves hybrid dipole-dipole moments, i.e., $[d, 1, 2, 2]$ model (Eq. (20)), and the simplest example of Model Series B is a system conserves hybrid dipole-quadrupole moments (Eq. (35)). We study the two specific examples and obtain the ground states and Goldstone modes of these two models through calculation. The conservation of hybrid moments affects the dispersion relations of the Goldstone modes. In two spatial dimensions, there are Goldstone modes with linear dispersion in both the two models. According to the calculation results of the correlation functions at zero temperature (Table II, III), we determine that in both models a partial spontaneous breaking of charge symmetry occurs in two spatial dimensions. This means that the two models have true ODLRO at zero temperature in two spatial dimensions. In Model Series A, the relative charge symmetry $U(1)_{1,C}$ is spontaneously broken, and in Model Series B, the charge symmetry $U(1)_{-C}$ of bosons of species 1 is spontaneously broken (Table I). We also extend the related results of $[d, 1, 2, 2]$ model to arbitrary $[d, N, m, m']$ models, and partial spontaneous breaking of charge symmetry occurs in spatial dimensions $d = N + 1$.

We construct Bose-Hubbard-type lattice models which correspond to Model Series A and Model Series B by using the difference method. We perform the mean-field theory of the Hamiltonian (Eq. (92)) of Lattice Model Series A, and give a method to realize it in a strongly tilted optical lattice. The mean-field results imply the existence of phases with condensates of bound states $\hat{b}_{1,i} \hat{b}_{2,i}^\dagger$, and we find that the coupling coefficient t_2 plays a similar role as the dipolar hopping coefficient t_1 .

This paper extends the field of research on fractonic superfluids and simultaneously connects with the study of the corresponding lattice models. There are many interesting directions for future investigation. By giving algebraic structures [40] with conserved hybrid moments, a rich variety of effective field theories can be constructed. By adding the concept of conserved hybridized moments, diffusion in spin models [43] can also be discussed. It is attractive to obtain and numerically solve the hydrodynamic equations [46] of systems with conservation of hybrid moments. We can also use renormalization groups analysis to determine whether systems with conserved hybrid moments can emerge at low energies [47]. The finite-temperature phase transitions [48] in these

hybrid-moment-conserving models are also worth investigating. The results of the partial spontaneous breaking of charge symmetry will generalize the Mermin-Wagner theorems under higher-rank symmetries [55, 58]. As mobility restriction leads to unconventional types of toric codes [90], it is interesting to study more generalization of discrete gauge theory and toric codes through more possibilities of mobility constraints due to conservations such hybridization discussed in the present work. Detecting such states in the context of stabilizer codes and cellular automata, e.g., in Refs. [32, 91, 92] will be an intriguing direction. This series of studies has the potential to be realized through cold atom experiments such as strongly tilted optical lattices. The associated gen-

eralized Bose-Hubbard models in various dimensions can be simulated in large-scale numerical simulations e.g., quantum Monte Carlo, DMRG, and tensor network.

ACKNOWLEDGMENTS

We thank Long Zhang for helpful discussions. This work was in part supported by National Natural Science Foundation of China (NSFC) Grant No. 12074438 and No. 12474149. The calculations reported were performed on resources provided by the Guangdong Provincial Key Laboratory of Magnetoelectric Physics and Devices, No. 2022B1212010008.

-
- [1] C. Chamon, Quantum glassiness in strongly correlated clean systems: An example of topological overprotection, *Phys. Rev. Lett.* **94**, 040402 (2005).
 - [2] S. Vijay, J. Haah, and L. Fu, A new kind of topological quantum order: A dimensional hierarchy of quasiparticles built from stationary excitations, *Phys. Rev. B* **92**, 235136 (2015).
 - [3] S. Vijay, J. Haah, and L. Fu, Fracton topological order, generalized lattice gauge theory, and duality, *Phys. Rev. B* **94**, 235157 (2016).
 - [4] W. Shirley, K. Slagle, Z. Wang, and X. Chen, Fracton models on general three-dimensional manifolds, *Phys. Rev. X* **8**, 031051 (2018).
 - [5] H. Ma, E. Lake, X. Chen, and M. Hermele, Fracton topological order via coupled layers, *Phys. Rev. B* **95**, 245126 (2017).
 - [6] W. Shirley, K. Slagle, and X. Chen, Foliated fracton order from gauging subsystem symmetries, *SciPost Phys.* **6**, 041 (2019).
 - [7] A. Prem, J. Haah, and R. Nandkishore, Glassy quantum dynamics in translation invariant fracton models, *Phys. Rev. B* **95**, 155133 (2017).
 - [8] A. Dua, I. H. Kim, M. Cheng, and D. J. Williamson, Sorting topological stabilizer models in three dimensions, *Phys. Rev. B* **100**, 155137 (2019).
 - [9] R. M. Nandkishore and M. Hermele, Fractons, *Annual Review of Condensed Matter Physics* **10**, 295 (2019).
 - [10] D. Bulmash and M. Barkeshli, Gauging fractons: Immobile non-abelian quasiparticles, fractals, and position-dependent degeneracies, *Phys. Rev. B* **100**, 155146 (2019).
 - [11] A. Prem, S.-J. Huang, H. Song, and M. Hermele, Cage-net fracton models, *Phys. Rev. X* **9**, 021010 (2019).
 - [12] M.-Y. Li and P. Ye, Fracton physics of spatially extended excitations, *Phys. Rev. B* **101**, 245134 (2020).
 - [13] K. Slagle, Foliated quantum field theory of fracton order, *Phys. Rev. Lett.* **126**, 101603 (2021).
 - [14] M.-Y. Li and P. Ye, Fracton physics of spatially extended excitations. ii. polynomial ground state degeneracy of exactly solvable models, *Phys. Rev. B* **104**, 235127 (2021).
 - [15] C. Zhou, M.-Y. Li, Z. Yan, P. Ye, and Z. Y. Meng, Evolution of dynamical signature in the x-cube fracton topological order, *Phys. Rev. Res.* **4**, 033111 (2022).
 - [16] M.-Y. Li and P. Ye, Hierarchy of entanglement renormalization and long-range entangled states, *Phys. Rev. B* **107**, 115169 (2023).
 - [17] G.-Y. Zhu, J.-Y. Chen, P. Ye, and S. Trebst, Topological fracton quantum phase transitions by tuning exact tensor network states, *Phys. Rev. Lett.* **130**, 216704 (2023).
 - [18] G. Canossa, L. Pollet, M. A. Martin-Delgado, H. Song, and K. Liu, Exotic symmetry breaking properties of self-dual fracton spin models, *Phys. Rev. Res.* **6**, 013304 (2024).
 - [19] H. Song, J. Schönmeier-Kromer, K. Liu, O. Viyuela, L. Pollet, and M. A. Martin-Delgado, Optimal thresholds for fracton codes and random spin models with subsystem symmetry, *Phys. Rev. Lett.* **129**, 230502 (2022).
 - [20] B.-X. Li, Y. Zhou, and P. Ye, Three-dimensional fracton topological orders with boundary toeplitz braiding, *Phys. Rev. B* **110**, 205108 (2024).
 - [21] T. Devakul, S. A. Parameswaran, and S. L. Sondhi, Correlation function diagnostics for type-i fracton phases, *Phys. Rev. B* **97**, 041110 (2018).
 - [22] T. Devakul and D. J. Williamson, Universal quantum computation using fractal symmetry-protected cluster phases, *Phys. Rev. A* **98**, 022332 (2018).
 - [23] Y. You, T. Devakul, F. J. Burnell, and S. L. Sondhi, Subsystem symmetry protected topological order, *Phys. Rev. B* **98**, 035112 (2018).
 - [24] T. Devakul, Y. You, F. J. Burnell, and S. L. Sondhi, Fractal Symmetric Phases of Matter, *SciPost Phys.* **6**, 007 (2019).
 - [25] T. Devakul, Classifying local fractal subsystem symmetry-protected topological phases, *Phys. Rev. B* **99**, 235131 (2019).
 - [26] D. T. Stephen, H. P. Nautrup, J. Bermejo-Vega, J. Eisert, and R. Raussendorf, Subsystem symmetries, quantum cellular automata, and computational phases of quantum matter, *Quantum* **3**, 142 (2019).
 - [27] A. T. Schmitz, S.-J. Huang, and A. Prem, Entanglement spectra of stabilizer codes: A window into gapped quantum phases of matter, *Phys. Rev. B* **99**, 205109 (2019).
 - [28] Y. You, T. Devakul, S. L. Sondhi, and F. J. Burnell, Fractonic chern-simons and bf theories, *Phys. Rev. Res.* **2**, 023249 (2020).
 - [29] J. F. San Miguel, A. Dua, and D. J. Williamson, Bifurcating subsystem symmetric entanglement renormalization in two dimensions, *Phys. Rev. B* **103**, 035148 (2021).
 - [30] F. J. Burnell, T. Devakul, P. Gorantla, H. T. Lam, and S.-H. Shao, Anomaly inflow for subsystem symmetries, *Phys. Rev. B* **106**, 085113 (2022).
 - [31] C. Zhou, M.-Y. Li, Z. Yan, P. Ye, and Z. Y. Meng, Detecting subsystem symmetry protected topological order through strange correlators, *Phys. Rev. B* **106**, 214428 (2022).
 - [32] J.-Y. Zhang, M.-Y. Li, and P. Ye, Higher-order cellular automata generated symmetry-protected topological phases and detection through multi point strange correlators, *PRX Quantum* **5**, 030342 (2024).

- [33] M. Pretko, Generalized electromagnetism of subdimensional particles: A spin liquid story, *Phys. Rev. B* **96**, 035119 (2017).
- [34] M. Pretko, Subdimensional particle structure of higher rank $u(1)$ spin liquids, *Phys. Rev. B* **95**, 115139 (2017).
- [35] M. Pretko, The fracton gauge principle, *Phys. Rev. B* **98**, 115134 (2018).
- [36] S. Pai and M. Pretko, Dynamical scar states in driven fracton systems, *Phys. Rev. Lett.* **123**, 136401 (2019).
- [37] H. Ma and M. Pretko, Higher-rank deconfined quantum criticality at the lifshitz transition and the exciton bose condensate, *Phys. Rev. B* **98**, 125105 (2018).
- [38] S. Pai and M. Pretko, Fractonic line excitations: An inroad from three-dimensional elasticity theory, *Phys. Rev. B* **97**, 235102 (2018).
- [39] S. Pai, M. Pretko, and R. M. Nandkishore, Localization in fractonic random circuits, *Phys. Rev. X* **9**, 021003 (2019).
- [40] A. Gromov, Towards classification of fracton phases: The multipole algebra, *Phys. Rev. X* **9**, 031035 (2019).
- [41] J. Wang and S.-T. Yau, Non-abelian gauged fracton matter field theory: Sigma models, superfluids, and vortices, *Phys. Rev. Res.* **2**, 043219 (2020).
- [42] A. Gromov, A. Lucas, and R. M. Nandkishore, Fracton hydrodynamics, *Phys. Rev. Res.* **2**, 033124 (2020).
- [43] J. Feldmeier, P. Sala, G. De Tomasi, F. Pollmann, and M. Knap, Anomalous diffusion in dipole- and higher-moment-conserving systems, *Phys. Rev. Lett.* **125**, 245303 (2020).
- [44] J.-K. Yuan, S. A. Chen, and P. Ye, Fractonic superfluids, *Phys. Rev. Res.* **2**, 023267 (2020).
- [45] S. A. Chen, J.-K. Yuan, and P. Ye, Fractonic superfluids. ii. condensing subdimensional particles, *Phys. Rev. Res.* **3**, 013226 (2021).
- [46] J.-K. Yuan, S. A. Chen, and P. Ye, Quantum hydrodynamics of fractonic superfluids with lineon condensate: From navier-stokes-like equations to landau-like criterion, *Chin. Phys. Lett.* **39**, 057101 (2022).
- [47] H. Li and P. Ye, Renormalization group analysis on emergence of higher rank symmetry and higher moment conservation, *Phys. Rev. Res.* **3**, 043176 (2021).
- [48] J.-K. Yuan, S. A. Chen, and P. Ye, Hierarchical proliferation of higher-rank symmetry defects in fractonic superfluids, *Phys. Rev. B* **107**, 205134 (2023).
- [49] S. A. Chen and P. Ye, Many-body physics of spontaneously broken higher-rank symmetry: from fractonic superfluids to dipolar hubbard model (2023), [arXiv:2305.00941 \[cond-mat.str-el\]](https://arxiv.org/abs/2305.00941).
- [50] D. Doshi and A. Gromov, Vortices as fractons, *Communications Physics* **4**, 44 (2021).
- [51] K. T. Grosvenor, C. Hoyos, F. Peña Benitez, and P. Surówka, Hydrodynamics of ideal fracton fluids, *Phys. Rev. Res.* **3**, 043186 (2021).
- [52] J. Wang, K. Xu, and S.-T. Yau, Higher-rank tensor non-abelian field theory: Higher-moment or subdimensional polynomial global symmetry, algebraic variety, noether's theorem, and gauging, *Phys. Rev. Res.* **3**, 013185 (2021).
- [53] R. Argurio, C. Hoyos, D. Musso, and D. Naegels, Fractons in effective field theories for spontaneously broken translations, *Phys. Rev. D* **104**, 105001 (2021).
- [54] J. Iaconis, A. Lucas, and R. Nandkishore, Multipole conservation laws and subdiffusion in any dimension, *Phys. Rev. E* **103**, 022142 (2021).
- [55] C. Stahl, E. Lake, and R. Nandkishore, Spontaneous breaking of multipole symmetries, *Phys. Rev. B* **105**, 155107 (2022).
- [56] L. Radzihovsky, Lifshitz gauge duality, *Phys. Rev. B* **106**, 224510 (2022).
- [57] A. Osborne and A. Lucas, Infinite families of fracton fluids with momentum conservation, *Phys. Rev. B* **105**, 024311 (2022).
- [58] A. Kapustin and L. Spodyneiko, Hohenberg-mermin-wagner-type theorems and dipole symmetry, *Phys. Rev. B* **106**, 245125 (2022).
- [59] E. Lake, M. Hermele, and T. Senthil, Dipolar bose-hubbard model, *Phys. Rev. B* **106**, 064511 (2022).
- [60] E. Lake, H.-Y. Lee, J. H. Han, and T. Senthil, Dipole condensates in tilted bose-hubbard chains, *Phys. Rev. B* **107**, 195132 (2023).
- [61] K. Giergiel, R. Lier, P. Surówka, and A. Kosior, Bose-hubbard realization of fracton defects, *Phys. Rev. Res.* **4**, 023151 (2022).
- [62] P. Zechmann, E. Altman, M. Knap, and J. Feldmeier, Fractonic luttinger liquids and supersolids in a constrained bose-hubbard model, *Phys. Rev. B* **107**, 195131 (2023).
- [63] J. Boesl, P. Zechmann, J. Feldmeier, and M. Knap, Deconfinement dynamics of fractons in tilted bose-hubbard chains, *Phys. Rev. Lett.* **132**, 143401 (2024).
- [64] W. Xu, C. Lv, and Q. Zhou, Multipolar condensates and multipolar Josephson effects, *Nature Communications* **15**, 4786 (2024).
- [65] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, Global dipole symmetry, compact lifshitz theory, tensor gauge theory, and fractons, *Phys. Rev. B* **106**, 045112 (2022).
- [66] K. T. Grosvenor, C. Hoyos, F. Peña-Benítez, and P. Surówka, Space-dependent symmetries and fractons, *Frontiers in Physics* **9**, 10.3389/fphy.2021.792621 (2022).
- [67] E. Lake and T. Senthil, Non-fermi liquids from kinetic constraints in tilted optical lattices, *Phys. Rev. Lett.* **131**, 043403 (2023).
- [68] J. Molina-Vilaplana, A post-Gaussian approach to dipole symmetries and interacting fractons, *Journal of High Energy Physics* **2023**, 65 (2023).
- [69] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, (2+1)-dimensional compact lifshitz theory, tensor gauge theory, and fractons, *Phys. Rev. B* **108**, 075106 (2023).
- [70] A. Anakru and Z. Bi, Non-fermi liquids from dipolar symmetry breaking, *Phys. Rev. B* **108**, 165112 (2023).
- [71] P. Glorioso, X. Huang, J. Guo, J. F. Rodriguez-Nieva, and A. Lucas, Goldstone bosons and fluctuating hydrodynamics with dipole and momentum conservation, *Journal of High Energy Physics* **2023**, 22 (2023).
- [72] A. Morningstar, N. O'Dea, and J. Richter, Hydrodynamics in long-range interacting systems with center-of-mass conservation, *Phys. Rev. B* **108**, L020304 (2023).
- [73] X. Huang, A Chern-Simons theory for dipole symmetry, *SciPost Phys.* **15**, 153 (2023).
- [74] E. Afxonidis, A. Caddeo, C. Hoyos, and D. Musso, Dipole symmetry breaking and fractonic Nambu-Goldstone mode, *SciPost Phys. Core* **6**, 082 (2023).
- [75] C. Stahl, M. Qi, P. Glorioso, A. Lucas, and R. Nandkishore, Fracton superfluid hydrodynamics, *Phys. Rev. B* **108**, 144509 (2023).
- [76] H. T. Lam, Classification of dipolar symmetry-protected topological phases: Matrix product states, stabilizer hamiltonians, and finite tensor gauge theories, *Phys. Rev. B* **109**, 115142 (2024).
- [77] J. H. Han, E. Lake, and S. Ro, Scaling and localization in multipole-conserving diffusion, *Phys. Rev. Lett.* **132**, 137102 (2024).
- [78] F. J. Burnell, S. Moudgalya, and A. Prem, Filling constraints on translation invariant dipole conserving systems, *Phys. Rev. B* **110**, L121113 (2024).

- [79] A. Chavda, D. Naegels, and J. Staunton, Fractonic Coset Construction for Spontaneously Broken Translations, [arXiv e-prints](#) , [arXiv:2409.15193 \(2024\)](#), [arXiv:2409.15193 \[hep-th\]](#).
- [80] J. Armas and E. Have, Ideal fracton superfluids, [SciPost Phys.](#) **16**, 039 (2024).
- [81] P. Zechmann, J. Boesl, J. Feldmeier, and M. Knap, Dynamical spectral response of fractonic quantum matter, [Phys. Rev. B](#) **109**, 125137 (2024).
- [82] A. Głódkowski, F. Peña-Benítez, and P. Surówka, Dissipative fracton superfluids, [Journal of High Energy Physics](#) **2024**, 285 (2024).
- [83] S. Angus, M. Kim, and J.-H. Park, Fractons, non-riemannian geometry, and double field theory, [Phys. Rev. Res.](#) **4**, 033186 (2022).
- [84] F. Peña Benítez, Fractons, symmetric gauge fields and geometry, [Phys. Rev. Res.](#) **5**, 013101 (2023).
- [85] E. Bertolini, N. Maggiore, and G. Palumbo, Covariant fracton gauge theory with boundary, [Phys. Rev. D](#) **108**, 025009 (2023).
- [86] E. Bertolini and H. Kim, Covariant Interacting Fractons, [arXiv e-prints](#) , [arXiv:2410.01397 \(2024\)](#), [arXiv:2410.01397 \[hep-th\]](#).
- [87] E. Bertolini and H. Kim, Strings as Hyper-Fractons, [arXiv e-prints](#) , [arXiv:2410.11678 \(2024\)](#), [arXiv:2410.11678 \[hep-th\]](#).
- [88] Y.-M. Hu and B. Lian, The Bosonic Quantum Breakdown Hubbard Model, [arXiv e-prints](#) , [arXiv:2401.04309 \(2024\)](#), [arXiv:2401.04309 \[cond-mat.str-el\]](#).
- [89] D. A. Cox, J. Little, and D. O’Shea, Gröbner bases, in *Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra* (Springer International Publishing, Cham, 2015) pp. 49–119.
- [90] Y.-T. Oh, J. Kim, E.-G. Moon, and J. H. Han, Rank-2 toric code in two dimensions, [Phys. Rev. B](#) **105**, 045128 (2022).
- [91] Z. Liang, Y. Xu, J. T. Iosue, and Y.-A. Chen, Extracting topological orders of generalized pauli stabilizer codes in two dimensions, [PRX Quantum](#) **5**, 030328 (2024).
- [92] Z. Liang, B. Yang, J. T. Iosue, and Y.-A. Chen, Operator algebra and algorithmic construction of boundaries and defects in (2+1)D topological Pauli stabilizer codes, [arXiv e-prints](#) , [arXiv:2410.11942 \(2024\)](#), [arXiv:2410.11942 \[quant-ph\]](#).

Appendix A: Derivations of Goldstone modes and dispersion relations for Model Series A

1. Derivation of the Hamiltonian in the momentum space

Based on the Hamiltonian (Eq. (45)), we can derive the Hamiltonian of the momentum space (Eq. (46)) in the following steps. In the momentum space, we can perform a Fourier transform,

$$\begin{aligned}
 H = & \frac{1}{(2\pi)^{2d}} \int d^d \mathbf{x} d^d \mathbf{k} d^d \mathbf{k}' e^{i\mathbf{k}\cdot\mathbf{x} + i\mathbf{k}'\cdot\mathbf{x}} \sum_a^2 \left[\frac{g}{2} \hat{\pi}_a(\mathbf{k}) \hat{\pi}_a(\mathbf{k}') + \frac{\kappa \rho_{a0}^2 |\mathbf{k}|^4}{2} \hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(\mathbf{k}') \right. \\
 & \left. + \frac{\Gamma \rho_{a0} \rho_{b0} |\mathbf{k}|^2}{2} \sum_{b \neq a}^2 (\hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(\mathbf{k}') + \hat{\theta}_b(\mathbf{k}) \hat{\theta}_b(\mathbf{k}') - \hat{\theta}_a(\mathbf{k}) \hat{\theta}_b(\mathbf{k}') - \hat{\theta}_b(\mathbf{k}) \hat{\theta}_a(\mathbf{k}')) \right]. \quad (\text{A1})
 \end{aligned}$$

Integrating over the real space, we can obtain a δ function so that only the integral of the momentum space is left in the Hamiltonian,

$$\begin{aligned}
 H = & \frac{1}{(2\pi)^d} \int d^d \mathbf{k} d^d \mathbf{k}' \delta(\mathbf{k} + \mathbf{k}') \sum_a^2 \left[\frac{g}{2} \hat{\pi}_a(\mathbf{k}) \hat{\pi}_a(\mathbf{k}') + \frac{\kappa \rho_{a0}^2 |\mathbf{k}|^4}{2} \hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(\mathbf{k}') \right. \\
 & \left. + \frac{\Gamma \rho_{a0} \rho_{b0} |\mathbf{k}|^2}{2} \sum_{b \neq a}^2 (\hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(\mathbf{k}') + \hat{\theta}_b(\mathbf{k}) \hat{\theta}_b(\mathbf{k}') - \hat{\theta}_a(\mathbf{k}) \hat{\theta}_b(\mathbf{k}') - \hat{\theta}_b(\mathbf{k}) \hat{\theta}_a(\mathbf{k}')) \right] \\
 = & \frac{1}{(2\pi)^d} \int d^d \mathbf{k} \sum_a^2 \left[\frac{g}{2} \hat{\pi}_a(\mathbf{k}) \hat{\pi}_a(-\mathbf{k}) + \frac{\kappa \rho_{a0}^2 |\mathbf{k}|^4}{2} \hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(-\mathbf{k}) \right. \\
 & \left. + \frac{\Gamma \rho_{a0} \rho_{b0} |\mathbf{k}|^2}{2} \sum_{b \neq a}^2 (\hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(-\mathbf{k}) + \hat{\theta}_b(\mathbf{k}) \hat{\theta}_b(-\mathbf{k}) - \hat{\theta}_a(\mathbf{k}) \hat{\theta}_b(-\mathbf{k}) - \hat{\theta}_b(\mathbf{k}) \hat{\theta}_a(-\mathbf{k})) \right]. \quad (\text{A2})
 \end{aligned}$$

The Hamiltonian (Eq. (46)) can be obtained by writing it in the matrix form and using the notation in the main text,

$$\begin{aligned}
H &= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \sum_a^2 \left[g \hat{\pi}_a(\mathbf{k}) \hat{\pi}_a(-\mathbf{k}) + \kappa \rho_{a0}^2 |\mathbf{k}|^4 \hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(-\mathbf{k}) \right. \\
&\quad \left. + \Gamma \rho_{a0} \rho_{b0} |\mathbf{k}|^2 \sum_{b \neq a}^2 (\hat{\theta}_a(\mathbf{k}) \hat{\theta}_a(-\mathbf{k}) + \hat{\theta}_b(\mathbf{k}) \hat{\theta}_b(-\mathbf{k}) - \hat{\theta}_a(\mathbf{k}) \hat{\theta}_b(-\mathbf{k}) - \hat{\theta}_b(\mathbf{k}) \hat{\theta}_a(-\mathbf{k})) \right] \\
&= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) \begin{bmatrix} g & 0 & 0 & 0 \\ 0 & g & 0 & 0 \\ 0 & 0 & \kappa \rho_{10}^2 |\mathbf{k}|^4 + 2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 & -2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 \\ 0 & 0 & -2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 & \kappa \rho_{20}^2 |\mathbf{k}|^4 + 2\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 \end{bmatrix} \psi(\mathbf{k}) \\
&= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) M(\mathbf{k}) \psi(\mathbf{k}). \tag{A3}
\end{aligned}$$

2. Matrix elements of the transformation matrix $T(\mathbf{k})$

In the main text, we give the commutation relations that the operators need to satisfy. To keep the commutation relations and diagonalize the matrix $M(\mathbf{k})$, the transformation matrix $T(\mathbf{k})$ needs to obey the equation below,

$$T(\mathbf{k}) \Lambda_1 T^\dagger(\mathbf{k}) = \Lambda_2 \tag{A4}$$

where

$$\Lambda_1 = \begin{bmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{bmatrix}, \quad \Lambda_2 = \text{diag}(1, 1, -1, -1). \tag{A5}$$

The transformation matrix $T(\mathbf{k})$ is given by

$$T(\mathbf{k}) = \begin{bmatrix} T_{11}(\mathbf{k}) & T_{12}(\mathbf{k}) & T_{13}(\mathbf{k}) & T_{14}(\mathbf{k}) \\ T_{21}(\mathbf{k}) & T_{22}(\mathbf{k}) & T_{23}(\mathbf{k}) & T_{24}(\mathbf{k}) \\ T_{31}(\mathbf{k}) & T_{32}(\mathbf{k}) & T_{33}(\mathbf{k}) & T_{34}(\mathbf{k}) \\ T_{41}(\mathbf{k}) & T_{42}(\mathbf{k}) & T_{43}(\mathbf{k}) & T_{44}(\mathbf{k}) \end{bmatrix}, \tag{A6}$$

where the matrix elements are

$$\begin{aligned}
T_{11}(\mathbf{k}) = -T_{31}(\mathbf{k}) &= \left(ig \left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 k^4 \rho_{10}^2 \rho_{20}^2} \right) / \right. \\
&\quad \left\{ \sqrt[4]{2} \left[g \left(16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2 + \left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 k^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)^2 \right) \right. \right. \\
&\quad \left. \left. \sqrt{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} - \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)} \right]^{\frac{1}{2}} \right\}, \tag{A7}
\end{aligned}$$

$$\begin{aligned}
T_{12}(\mathbf{k}) = -T_{32}(\mathbf{k}) &= \left(4ig\Gamma \rho_{10} \rho_{20} |\mathbf{k}|^2 \right) / \\
&\quad \left\{ \sqrt[4]{2} \left[g \left(16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2 + \left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 k^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)^2 \right) \right. \right. \\
&\quad \left. \left. \sqrt{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} - \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)} \right]^{\frac{1}{2}} \right\}, \tag{A8}
\end{aligned}$$

$$T_{13}(\mathbf{k}) = T_{33}(\mathbf{k}) = \left[\left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right) \right. \\ \left. \sqrt[4]{ \frac{\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} - \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2}}{g} } \right] / \\ \left(2^{3/4} \sqrt{16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} + \left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)^2 \right), \quad (\text{A9})$$

$$T_{14}(\mathbf{k}) = T_{34}(\mathbf{k}) \\ = \left(2\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} \sqrt[4]{ \frac{2\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 8\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} - 2\sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2}}{g} } \right) / \\ \left(\sqrt{16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} + \left(\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)^2 \right) \quad (\text{A10})$$

$$T_{21}(\mathbf{k}) = -T_{41}(\mathbf{k}) = - \left[ig \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 - \rho_{20}^2) + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right) \right] / \\ \left\{ 2^{3/4} |\mathbf{k}|^2 \left[g \left(16\Gamma^2 \rho_{10}^2 \rho_{20}^2 + \kappa^2 |\mathbf{k}|^4 (\rho_{10}^2 - \rho_{20}^2)^2 + \kappa (\rho_{10}^2 - \rho_{20}^2) \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right) \right. \right. \\ \left. \left. \sqrt{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)} \right]^{\frac{1}{2}} \right\}, \quad (\text{A11})$$

$$T_{22}(\mathbf{k}) = -T_{42}(\mathbf{k}) = \left(2i \sqrt[4]{2} \Gamma g \rho_{10} \rho_{20} \right) / \\ \left[\sqrt[4]{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)} \right. \\ \left. \sqrt{g \left(16\Gamma^2 \rho_{10}^2 \rho_{20}^2 + \kappa^2 |\mathbf{k}|^4 (\rho_{10}^2 - \rho_{20}^2)^2 + \kappa (\rho_{10}^2 - \rho_{20}^2) \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)} \right], \quad (\text{A12})$$

$$T_{23}(\mathbf{k}) = T_{43}(\mathbf{k}) \\ = \frac{\kappa |\mathbf{k}|^4 (\rho_{20}^2 - \rho_{10}^2) - \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2}}{2\sqrt[4]{2} |\mathbf{k}|^2 \sqrt{ \frac{g \left(16\Gamma^2 \rho_{10}^2 \rho_{20}^2 + \kappa^2 |\mathbf{k}|^4 (\rho_{10}^2 - \rho_{20}^2)^2 + \kappa \rho_{10}^2 \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} - \kappa \rho_{20}^2 \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)}{\sqrt{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)}} } }, \quad (\text{A13})$$

$$T_{24}(\mathbf{k}) = T_{44}(\mathbf{k}) \\ = \frac{2^{3/4} \Gamma \rho_{10} \rho_{20}}{\sqrt{ \frac{g \left(16\Gamma^2 \rho_{10}^2 \rho_{20}^2 + \kappa^2 |\mathbf{k}|^4 (\rho_{10}^2 - \rho_{20}^2)^2 + \kappa \rho_{10}^2 \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} - \kappa \rho_{20}^2 \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)}{\sqrt{g \left(\kappa |\mathbf{k}|^4 (\rho_{10}^2 + \rho_{20}^2) + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20} + \sqrt{\kappa^2 |\mathbf{k}|^8 (\rho_{10}^2 - \rho_{20}^2)^2 + 16\Gamma^2 |\mathbf{k}|^4 \rho_{10}^2 \rho_{20}^2} \right)}} } }. \quad (\text{A14})$$

3. Derivation of the Hamiltonian represented by the creation and annihilation operators

After diagonalized, the Hamiltonian (Eq. (50)) becomes

$$\begin{aligned} H &= \frac{1}{2(2\pi)^d} \phi^\dagger(\mathbf{k}) (T^{-1}(\mathbf{k}))^\dagger M(\mathbf{k}) T^{-1}(\mathbf{k}) \phi(\mathbf{k}) \\ &= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) (\hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + \hat{\alpha}(-\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k})) + D_2(\mathbf{k}) (\hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \hat{\beta}(-\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k})). \end{aligned} \quad (\text{A15})$$

Because of $D_1(\mathbf{k}) = D_1(-\mathbf{k})$, $D_2(\mathbf{k}) = D_2(-\mathbf{k})$, we rewrite the Hamiltonian as

$$\begin{aligned} H &= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) (\hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + \hat{\alpha}(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k})) + D_2(\mathbf{k}) (\hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \hat{\beta}(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k})) \\ &= \frac{1}{(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \int d^d \mathbf{k} \frac{D_1(\mathbf{k}) + D_2(\mathbf{k})}{2} \delta^{(d)}(\mathbf{0}) \\ &= \frac{1}{(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \frac{\tilde{V}}{(2\pi)^d} \int d^d \mathbf{k} \frac{D_1(\mathbf{k}) + D_2(\mathbf{k})}{2}, \end{aligned} \quad (\text{A16})$$

where \tilde{V} is the volume of the d -dimensional space, the last term is the vacuum zero point energy. Then, we obtain the dispersion relations of Goldstone mode (Eq. (55)).

Appendix B: The calculation details of correlation function of Model Series A

In this appendix, we will give the calculation details of correlation functions of Model Series A. From the main text and Appendix A 2, we have the Bogoliubov transformation from $\hat{\theta}$ and $\hat{\pi}$ to the independent modes $\hat{\alpha}$ and $\hat{\beta}$,

$$\begin{aligned} \psi(\mathbf{k}) &= [\hat{\pi}_1(\mathbf{k}) \ \hat{\pi}_2(\mathbf{k}) \ \hat{\theta}_1(\mathbf{k}) \ \hat{\theta}_2(\mathbf{k})]^T = T^{-1}(\mathbf{k}) \phi(\mathbf{k}) = \Lambda_1 T^\dagger(\mathbf{k}) \Lambda_2 \phi(\mathbf{k}) \\ &= \begin{bmatrix} -iT_{13}^*(\mathbf{k}) & -iT_{23}^*(\mathbf{k}) & iT_{33}^*(\mathbf{k}) & iT_{43}^*(\mathbf{k}) \\ -iT_{14}^*(\mathbf{k}) & -iT_{24}^*(\mathbf{k}) & iT_{34}^*(\mathbf{k}) & iT_{44}^*(\mathbf{k}) \\ iT_{11}^*(\mathbf{k}) & iT_{21}^*(\mathbf{k}) & -iT_{31}^*(\mathbf{k}) & -iT_{41}^*(\mathbf{k}) \\ iT_{12}^*(\mathbf{k}) & iT_{22}^*(\mathbf{k}) & -iT_{32}^*(\mathbf{k}) & -iT_{42}^*(\mathbf{k}) \end{bmatrix} \begin{bmatrix} \hat{\alpha}(\mathbf{k}) \\ \hat{\beta}(\mathbf{k}) \\ \hat{\alpha}^\dagger(-\mathbf{k}) \\ \hat{\beta}^\dagger(-\mathbf{k}) \end{bmatrix}. \end{aligned} \quad (\text{B1})$$

In the main text, we express two-operator and four-operator correlation functions of the boson operator $\hat{\Phi}_a$ in terms of two-operator correlation functions of the phase field operator $\hat{\theta}_a$. Two-operator correlation functions of the phase field operator $\hat{\theta}_a$ can be given in terms of the matrix elements of the matrix $T(\mathbf{k})$ in the following procedure,

$$\begin{aligned} \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle &= \langle \int \frac{d^d \mathbf{k} d^d \mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') \rangle = \langle \int \frac{d^d \mathbf{k} d^d \mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1^\dagger(-\mathbf{k}') \rangle \\ &= \int \frac{d^d \mathbf{k} d^d \mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \langle [iT_{11}^*(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + iT_{21}^*(\mathbf{k}) \hat{\beta}(\mathbf{k}) - iT_{31}^*(\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}) - iT_{41}^*(\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k})] \\ &\quad [-iT_{11}(-\mathbf{k}') \hat{\alpha}^\dagger(-\mathbf{k}') - iT_{21}(-\mathbf{k}') \hat{\beta}^\dagger(-\mathbf{k}') + iT_{31}(-\mathbf{k}') \hat{\alpha}(\mathbf{k}') + iT_{41}(-\mathbf{k}') \hat{\beta}(\mathbf{k}')] \rangle. \end{aligned} \quad (\text{B2})$$

For the expectation value of the creation and annihilation operators in the ground state, we have

$$\langle \hat{\alpha}(\mathbf{k}') \hat{\alpha}(\mathbf{k}) \rangle = \langle \hat{\beta}(\mathbf{k}') \hat{\beta}(\mathbf{k}) \rangle = \langle \hat{\alpha}^\dagger(-\mathbf{k}') \hat{\alpha}(\mathbf{k}) \rangle = \langle \hat{\beta}^\dagger(-\mathbf{k}') \hat{\beta}(\mathbf{k}) \rangle = 0, \quad (\text{B3})$$

$$\langle \hat{\alpha}(\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}') \rangle = \langle \hat{\beta}(\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k}') \rangle = (2\pi)^d \delta^{(d)}(\mathbf{k} + \mathbf{k}'). \quad (\text{B4})$$

Then we can obtain the correlation function

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle = \int \frac{d^d \mathbf{k}}{(2\pi)^d} e^{i\mathbf{k}\cdot\mathbf{x}} (|T_{11}(\mathbf{k})|^2 + |T_{21}(\mathbf{k})|^2). \quad (\text{B5})$$

Similarly, we can get

$$\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \int \frac{d^d \mathbf{k}}{(2\pi)^d} e^{i\mathbf{k} \cdot \mathbf{x}} \left(|T_{12}(\mathbf{k})|^2 + |T_{22}(\mathbf{k})|^2 \right). \quad (\text{B6})$$

The correlation function of different species of phase field operators is given by

$$\begin{aligned} \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle &= \left\langle \int \frac{d^d \mathbf{k} d^d \mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k} \cdot \mathbf{x}} \hat{\theta}_1(\mathbf{k}) \hat{\theta}_2^\dagger(-\mathbf{k}') \right\rangle \\ &= \int \frac{d^d \mathbf{k} d^d \mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k} \cdot \mathbf{x}} \left\langle \left[iT_{11}^*(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + iT_{21}^*(\mathbf{k}) \hat{\beta}(\mathbf{k}) - iT_{31}^*(\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}) - iT_{41}^*(\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k}) \right] \right. \\ &\quad \left. \left[-iT_{12}(-\mathbf{k}') \hat{\alpha}^\dagger(-\mathbf{k}') - iT_{22}(-\mathbf{k}') \hat{\beta}^\dagger(-\mathbf{k}') + iT_{32}(-\mathbf{k}') \hat{\alpha}(\mathbf{k}') + iT_{42}(-\mathbf{k}') \hat{\beta}(\mathbf{k}') \right] \right\rangle \\ &= \int \frac{d^d \mathbf{k}}{(2\pi)^d} e^{i\mathbf{k} \cdot \mathbf{x}} [T_{11}^*(\mathbf{k}) T_{12}(\mathbf{k}) + T_{21}^*(\mathbf{k}) T_{22}(\mathbf{k})]. \end{aligned} \quad (\text{B7})$$

For the $T(\mathbf{k})$ matrix of Model Series A in Appendix A 2, we can give some approximations,

$$\begin{aligned} |T_{11}(\mathbf{k})|^2 &\approx \frac{g}{2\sqrt{2g\kappa}(\rho_{10}^2 + \rho_{20}^2)|\mathbf{k}|^2} = \frac{1}{c_1 |\mathbf{k}|^2}, \\ |T_{12}(\mathbf{k})|^2 &\approx \frac{g}{2\sqrt{2g\kappa}(\rho_{10}^2 + \rho_{20}^2)|\mathbf{k}|^2} = \frac{1}{c_1 |\mathbf{k}|^2}, \\ |T_{21}(\mathbf{k})|^2 &\approx \frac{g}{8\sqrt{g\Gamma\rho_{10}\rho_{20}}|\mathbf{k}|} = \frac{1}{c_2 |\mathbf{k}|}, \\ |T_{22}(\mathbf{k})|^2 &\approx \frac{g}{8\sqrt{g\Gamma\rho_{10}\rho_{20}}|\mathbf{k}|} = \frac{1}{c_2 |\mathbf{k}|}, \\ T_{11}^*(\mathbf{k}) T_{12}(\mathbf{k}) &\approx \frac{g}{2\sqrt{2g\kappa}(\rho_{10}^2 + \rho_{20}^2)|\mathbf{k}|^2} = \frac{1}{c_1 |\mathbf{k}|^2}, \\ T_{21}^*(\mathbf{k}) T_{22}(\mathbf{k}) &\approx -\frac{g}{8\sqrt{g\Gamma\rho_{10}\rho_{20}}|\mathbf{k}|} = -\frac{1}{c_2 |\mathbf{k}|}. \end{aligned} \quad (\text{B8})$$

Based on the above derivation, we can calculate correlation functions in different spatial dimensions.

1. One spatial dimension

In one spatial dimension, we can plug in the results of the above approximation. The correlation functions of the operator $\hat{\theta}$ are

$$\begin{aligned} \langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle &= \langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle = \int \frac{dk}{2\pi} e^{ikx} \left(\frac{1}{c_1 |k|^2} + \frac{1}{c_2 |k|} \right) \\ &= \frac{1}{2\pi c_1} \int dk \frac{e^{ikx}}{|k|^2} + \frac{1}{2\pi c_2} \int dk \frac{e^{ikx}}{|k|} \\ &= \frac{1}{2\pi c_1} \left(\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{k^2} + \int_{\frac{2\pi}{L}}^{\infty} dk \frac{e^{ikx}}{k^2} \right) + \frac{1}{2\pi c_2} \left(\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{-k} + \int_{\frac{2\pi}{L}}^{\infty} \frac{e^{ikx}}{k} dk \right). \end{aligned} \quad (\text{B9})$$

We need to compute these integrals separately. Considering $L > 0$, $x > 0$, we have the first two integrals

$$\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{k^2} + \int_{\frac{2\pi}{L}}^{\infty} dk \frac{e^{ikx}}{k^2} = \frac{1}{2\pi} \left[e^{-2i\pi x L} + e^{2i\pi x L} + 2i\pi x \Gamma \left(0, -\frac{2i\pi x}{L} \right) - 2i\pi x \Gamma \left(0, -\frac{2i\pi x}{L} \right) \right] \stackrel{L \rightarrow \infty}{=} \frac{L}{\pi} - \pi x, \quad (\text{B10})$$

where $\Gamma(a, z) = \int_z^\infty t^{a-1} e^{-t} dt$ is the incomplete gamma function. The last two integrals are

$$\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{-k} + \int_{\frac{2\pi}{L}}^{\infty} \frac{e^{ikx}}{k} = \Gamma \left(0, -\frac{2i\pi x}{L} \right) + \Gamma \left(0, \frac{2i\pi x}{L} \right) \stackrel{L \rightarrow \infty}{=} -2 \left(\gamma + \ln \left(\frac{2\pi x}{L} \right) \right), \quad (\text{B11})$$

where γ is the Euler-Mascheroni constant. Then, we can get the correlation functions.

$$\langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle = \langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle \stackrel{L \rightarrow \infty}{=} \frac{1}{2\pi c_1} \left(\frac{L}{\pi} - \pi |x| \right) + \frac{1}{\pi c_2} \left(-\gamma + \ln \left(\frac{L}{2\pi |x|} \right) \right) \quad (\text{B12})$$

Similarly, we have

$$\langle \hat{\theta}_1(x) \hat{\theta}_2(0) \rangle \stackrel{L \rightarrow \infty}{=} \frac{1}{2\pi c_1} \left(\frac{L}{\pi} - \pi |x| \right) - \frac{1}{\pi c_2} \left(-\gamma + \ln \left(\frac{L}{2\pi |x|} \right) \right). \quad (\text{B13})$$

To get $\langle \hat{\theta}_a^2(0) \rangle = \lim_{x \rightarrow 0} \langle \theta_a(x) \theta_a(0) \rangle$, we just need to replace $|x|$ by ξ_c . ξ_c is the coherence length,

$$\langle \hat{\theta}_1^2(0) \rangle = \langle \hat{\theta}_2^2(0) \rangle = \frac{1}{2\pi c_1} \left(\frac{L}{\pi} - \pi \xi_c \right) + \frac{1}{\pi c_2} \left(-\gamma + \ln \left(\frac{L}{2\pi \xi_c} \right) \right), \quad (\text{B14})$$

$$\langle \hat{\theta}_1(0) \hat{\theta}_2(0) \rangle = \frac{1}{2\pi c_1} \left(\frac{L}{\pi} - \pi \xi_c \right) - \frac{1}{\pi c_2} \left(-\gamma + \ln \left(\frac{L}{2\pi \xi_c} \right) \right). \quad (\text{B15})$$

Then, we can calculate correlation functions of $\hat{\Phi}$. The two-operator correlation functions are

$$\langle \hat{\Phi}_1^\dagger(x) \hat{\Phi}_1(0) \rangle = \rho_{10} e^{\langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle - \langle \hat{\theta}_1^2(0) \rangle} = \rho_{10} e^{-\frac{|x|}{2c_1} + \frac{\xi_c}{2c_1}} \left(\frac{|x|}{\xi_c} \right)^{-\frac{1}{\pi c_2}} \approx \rho_{10} e^{-\frac{|x|}{2c_1}} \left(\frac{|x|}{\xi_c} \right)^{-\frac{1}{\pi c_2}}, \quad (\text{B16})$$

$$\langle \hat{\Phi}_2^\dagger(x) \hat{\Phi}_2(0) \rangle = \rho_{20} e^{\langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle - \langle \hat{\theta}_2^2(0) \rangle} = \rho_{20} e^{-\frac{|x|}{2c_1} + \frac{\xi_c}{2c_1}} \left(\frac{|x|}{\xi_c} \right)^{-\frac{1}{\pi c_2}} \approx \rho_{20} e^{-\frac{|x|}{2c_1}} \left(\frac{|x|}{\xi_c} \right)^{-\frac{1}{\pi c_2}}, \quad (\text{B17})$$

$$\langle \hat{\Phi}_1^\dagger(x) \hat{\Phi}_2(0) \rangle = \sqrt{\rho_{10} \rho_{20}} e^{\langle \hat{\theta}_1(x) \hat{\theta}_2(0) \rangle - \frac{1}{2} \langle \hat{\theta}_1^2(0) \rangle - \frac{1}{2} \langle \hat{\theta}_2^2(0) \rangle} = \sqrt{\rho_{10} \rho_{20}} e^{-\frac{|x|}{2c_1} + \frac{\xi_c}{2c_1} + \frac{2\gamma}{\pi c_2}} \left(\frac{L^2}{4\pi^2 \xi_c |x|} \right)^{-\frac{1}{\pi c_2}} \stackrel{L \rightarrow \infty}{=} 0. \quad (\text{B18})$$

The four-operator correlation functions are

$$\begin{aligned} & \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \\ &= \rho_{10} \rho_{20} \left(\frac{|x|}{\xi_c} \right)^{-\frac{4}{\pi c_2}}, \end{aligned} \quad (\text{B19})$$

$$\begin{aligned} & \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \\ &= \rho_{10} \rho_{20} e^{\frac{2\xi_c}{c_1}} e^{-\frac{2}{c_1} |x|} \approx \rho_{10} \rho_{20} e^{-\frac{2}{c_1} |x|}. \end{aligned} \quad (\text{B20})$$

Since all these correlation functions do not saturate to constants, Model Series A does not break the charge symmetry in one spatial dimension. Since the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle$ is power-law decay, the system is algebraically ordered in one spatial dimension.

2. Two spatial dimensions

In two spatial dimensions, we can plug in the results of the above approximation. The correlation functions of the operator $\hat{\theta}$ are

$$\begin{aligned} \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle &= \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \int \frac{d^2 \mathbf{k}}{(2\pi)^2} e^{i\mathbf{k} \cdot \mathbf{x}} \left(\frac{1}{c_1 |\mathbf{k}|^2} + \frac{1}{c_2 |\mathbf{k}|} \right) \\ &= \frac{1}{4\pi^2 c_1} \int dk d\theta \frac{e^{ik|\mathbf{x}| \cos \theta}}{k} + \frac{1}{4\pi^2 c_2} \int dk d\theta e^{ik|\mathbf{x}| \cos \theta} \\ &= \frac{1}{4\pi^2 c_1} \int_0^{2\pi} d\theta \int_{\frac{2\pi}{L}}^\infty dk \frac{e^{ik|\mathbf{x}| \cos \theta}}{k} + \frac{1}{4\pi^2 c_2} \int_0^{2\pi} d\theta \int_0^\infty dk e^{ik|\mathbf{x}| \cos \theta}. \end{aligned} \quad (\text{B21})$$

We need to compute these integrals separately,

$$\int_0^{2\pi} d\theta \int_{\frac{2\pi}{L}}^{\infty} dk \frac{e^{ik|\mathbf{x}|\cos\theta}}{k} \stackrel{L \rightarrow \infty}{=} -2\pi \left(\gamma + \ln \left(\frac{\pi |\mathbf{x}|}{L} \right) \right), \quad (\text{B22})$$

$$\int_0^{2\pi} d\theta \int_0^{\infty} dk e^{ik|\mathbf{x}|\cos\theta} = 2\pi \int_0^{\infty} dk J_0(k|\mathbf{x}|) = \frac{2\pi}{|\mathbf{x}|}, \quad (\text{B23})$$

where $J_n(z)$ is the Bessel function of the first kind. Then, we can get the correlation functions.

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle = \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \stackrel{L \rightarrow \infty}{=} \frac{1}{2\pi c_1} \left(-\gamma + \ln \left(\frac{L}{\pi |\mathbf{x}|} \right) \right) + \frac{1}{2\pi c_2 |\mathbf{x}|} \quad (\text{B24})$$

Similarly, we have

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \stackrel{L \rightarrow \infty}{=} \frac{1}{2\pi c_1} \left(-\gamma + \ln \left(\frac{L}{\pi |\mathbf{x}|} \right) \right) - \frac{1}{2\pi c_2 |\mathbf{x}|}. \quad (\text{B25})$$

To get $\langle \hat{\theta}_a^2(0) \rangle = \lim_{x \rightarrow 0} \langle \theta_a(x) \theta_a(0) \rangle$, we just need to replace $|x|$ by ξ_c ,

$$\langle \hat{\theta}_1^2(\mathbf{0}) \rangle = \langle \hat{\theta}_2^2(\mathbf{0}) \rangle = \frac{1}{2\pi c_1} \left(-\gamma + \ln \left(\frac{L}{\pi \xi_c} \right) \right) + \frac{1}{2\pi c_2 \xi_c}, \quad (\text{B26})$$

$$\langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle = \frac{1}{2\pi c_1} \left(-\gamma + \ln \left(\frac{L}{\pi \xi_c} \right) \right) - \frac{1}{2\pi c_2 \xi_c}. \quad (\text{B27})$$

Then, we can calculate correlation functions of $\hat{\Phi}$. The two-operator correlation functions are

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle = \rho_{10} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle} = \rho_{10} \left(\frac{|\mathbf{x}|}{\xi_c} \right)^{-\frac{1}{2\pi c_1}} e^{\frac{1}{2\pi c_2 |\mathbf{x}|} - \frac{1}{2\pi c_2 \xi_c}}, \quad (\text{B28})$$

$$\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \rho_{20} e^{\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle} = \rho_{20} \left(\frac{|\mathbf{x}|}{\xi_c} \right)^{-\frac{1}{2\pi c_1}} e^{\frac{1}{2\pi c_2 |\mathbf{x}|} - \frac{1}{2\pi c_2 \xi_c}}, \quad (\text{B29})$$

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \sqrt{\rho_{10} \rho_{20}} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \frac{1}{2} \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \frac{1}{2} \langle \hat{\theta}_2^2(\mathbf{0}) \rangle} = \sqrt{\rho_{10} \rho_{20}} \left(\frac{|\mathbf{x}|}{\xi_c} \right)^{-\frac{1}{2\pi c_1}} e^{-\frac{1}{2\pi c_2 |\mathbf{x}|} - \frac{1}{2\pi c_2 \xi_c}}. \quad (\text{B30})$$

The four-operator correlation functions are

$$\begin{aligned} & \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \\ &= \rho_{10} \rho_{20} e^{\frac{2}{\pi c_1 |\mathbf{x}|} - \frac{2}{\pi c_1 \xi_c}}, \end{aligned} \quad (\text{B31})$$

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle \quad (\text{B32})$$

$$= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \quad (\text{B33})$$

$$= \rho_{10} \rho_{20} \left(\frac{|\mathbf{x}|}{\xi_c} \right)^{-\frac{2}{\pi c_1}}. \quad (\text{B34})$$

Since the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle$ saturates to a constant when $|\mathbf{x}| \rightarrow \infty$, Model Series A spontaneously breaks the relative charge symmetry $U(1)_{-C}$ and has a true ODLRO in two spatial dimensions.

3. Three spatial dimensions

In three spatial dimensions, we can plug in the results of the above approximation. The correlation functions of the operator $\hat{\theta}$ are

$$\begin{aligned} \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle &= \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \int \frac{d^3\mathbf{k}}{(2\pi)^3} e^{i\mathbf{k}\cdot\mathbf{x}} \left(\frac{1}{c_1 |\mathbf{k}|^2} + \frac{1}{c_2 |\mathbf{k}|} \right) \\ &= \frac{1}{8\pi^3 c_1} \int_0^\pi d\psi \int_0^{2\pi} d\theta \int_0^\infty dk \left(\sin\psi e^{ik|\mathbf{x}| \cos\theta \sin\psi} \right) \\ &\quad + \frac{1}{8\pi^3 c_2} \int_0^\pi d\psi \int_0^{2\pi} d\theta \int_0^{\frac{2\pi}{\xi_c}} dk \left(k \sin\psi e^{ik|\mathbf{x}| \cos\theta \sin\psi} \right). \end{aligned} \quad (\text{B35})$$

We need to compute these integrals separately,

$$\int_0^\pi d\psi \int_0^{2\pi} d\theta \int_0^\infty dk \left(\sin\psi e^{ik|\mathbf{x}| \cos\theta \sin\psi} \right) = \int_0^\infty dk \frac{4\pi \sin(k|\mathbf{x}|)}{k|\mathbf{x}|} = \frac{2\pi^2}{|\mathbf{x}|}, \quad (\text{B36})$$

$$\int_0^\pi d\psi \int_0^{2\pi} d\theta \int_0^{\frac{2\pi}{\xi_c}} dk \left(k \sin\psi e^{ik|\mathbf{x}| \cos\theta \sin\psi} \right) = \int_0^{\frac{2\pi}{\xi_c}} dk \frac{4\pi k \sin(k|\mathbf{x}|)}{k|\mathbf{x}|} = \frac{8\pi \sin^2\left(\frac{\pi|\mathbf{x}|}{\xi_c}\right)}{|\mathbf{x}|^2}. \quad (\text{B37})$$

Then, we can get the correlation functions,

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle = \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \frac{1}{4\pi c_1 |\mathbf{x}|} + \frac{\sin^2\left(\frac{\pi|\mathbf{x}|}{\xi_c}\right)}{\pi^2 c_2 |\mathbf{x}|^2}. \quad (\text{B38})$$

Similarly, we have

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \frac{1}{4\pi c_1 |\mathbf{x}|} - \frac{\sin^2\left(\frac{\pi|\mathbf{x}|}{\xi_c}\right)}{\pi^2 c_2 |\mathbf{x}|^2}. \quad (\text{B39})$$

Since the coherence length is introduced by the previous integration, we can't just set $|\mathbf{x}| = \xi_c$ to get $\langle \hat{\theta}_a^2(\mathbf{0}) \rangle$. We need to calculate the integral,

$$\int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{1}{c_2 |\mathbf{k}|} = \frac{1}{8\pi^3 c_2} \int_0^{\frac{2\pi}{\xi_c}} dk \left(4\pi k^2 \frac{1}{k} \right) = \frac{1}{2\pi^2 c_2} \int_0^{\frac{2\pi}{\xi_c}} k dk = \frac{1}{c_2 \xi_c^2}. \quad (\text{B40})$$

Then, we can get

$$\langle \hat{\theta}_1^2(\mathbf{0}) \rangle = \langle \hat{\theta}_2^2(\mathbf{0}) \rangle = \frac{1}{4\pi c_1 \xi_c} + \frac{1}{c_2 \xi_c^2}, \quad (\text{B41})$$

$$\langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle = \frac{1}{4\pi c_1 \xi_c} - \frac{1}{c_2 \xi_c^2}. \quad (\text{B42})$$

Then, we can calculate correlation functions of $\hat{\Phi}$. The two-operator correlation functions are

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle = \rho_{10} e^{(\hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle) |\mathbf{x}| \xrightarrow{\infty}} \rho_{10} e^{-\frac{1}{4\pi c_1 \xi_c} - \frac{1}{c_2 \xi_c^2}}, \quad (\text{B43})$$

$$\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \rho_{20} e^{(\hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle) |\mathbf{x}| \xrightarrow{\infty}} \rho_{20} e^{-\frac{1}{4\pi c_1 \xi_c} - \frac{1}{c_2 \xi_c^2}}, \quad (\text{B44})$$

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \sqrt{\rho_{10} \rho_{20}} e^{(\hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) - \frac{1}{2} \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \frac{1}{2} \langle \hat{\theta}_2^2(\mathbf{0}) \rangle) |\mathbf{x}| \xrightarrow{\infty}} \sqrt{\rho_{10} \rho_{20}} e^{-\frac{1}{4\pi c_1 \xi_c} - \frac{1}{c_2 \xi_c^2}}. \quad (\text{B45})$$

The four-operator correlation functions are

$$\begin{aligned} & \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2^\dagger(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \\ & \stackrel{|\mathbf{x}| \rightarrow \infty}{=} \rho_{10} \rho_{20} e^{-\frac{4}{c_2 \xi_c^2}}, \end{aligned} \quad (\text{B46})$$

$$\begin{aligned} & \langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \hat{\Phi}_2(\mathbf{0}) \rangle \\ &= \rho_{10} \rho_{20} \exp \left(\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle + \langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - 2 \langle \hat{\theta}_1(\mathbf{0}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle + 2 \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \right) \\ & \stackrel{|\mathbf{x}| \rightarrow \infty}{=} \rho_{10} \rho_{20} e^{-\frac{1}{\pi c_1 \xi_c}}. \end{aligned} \quad (\text{B47})$$

Since all the correlation functions above saturate to constants when $|\mathbf{x}| \rightarrow \infty$, Model Series A breaks the charge symmetry and has a true ODLRO in three spatial dimensions.

Appendix C: Derivations of Goldstone modes and dispersion relations for Model Series B

In this appendix, we give the detailed derivation of Goldstone mode and dispersion relation for Model Series B in Sec. IV B.

1. Derivation of the effective Hamiltonian density

Considering quantum fluctuations, we have $\hat{\Phi}_a(\mathbf{x}) = \sqrt{\rho_{a0} + \hat{f}_a(\mathbf{x})} e^{i\hat{\theta}_a(\mathbf{x})}$, where $\hat{f}_a(\mathbf{x}) \ll \rho_{a0}$. The commutation relation between $\hat{\theta}_a$ and \hat{f}_a is given by $[\hat{\theta}_a(\mathbf{x}), \hat{f}_b(\mathbf{y})] = -i\delta^{(d)}(\mathbf{x} - \mathbf{y}) \delta_{ab}$. Therefore, the conjugate momentum operator of $\hat{\theta}_a$ is $\hat{\pi}_a = -\hat{f}_a$, and the operators $\hat{\theta}_a$ and $\hat{\pi}_a$ satisfy the commutation relations, $[\hat{\theta}_a(\mathbf{x}), \hat{\pi}_b(\mathbf{y})] = i\delta_{ab}\delta^d(\mathbf{x} - \mathbf{y})$. We can derive the effective Hamiltonian,

$$\begin{aligned} \mathcal{H} &= \frac{g}{2} \hat{\pi}_1^2 + \frac{g}{2} \hat{\pi}_2^2 + \sum_{i,j}^d K_1^{ij} \left[\rho_{10}^2 \left(\partial_i \partial_j \hat{\theta}_1 \right)^2 + \frac{(\partial_i \partial_j \hat{\pi}_1)^2}{4} \right] \\ &+ \sum_{i,j,k}^d K_2^{ijk} \left[\rho_{20}^3 \left(\partial_i \partial_j \partial_k \hat{\theta}_2 \right)^2 + \frac{1}{4} \rho_{20} \left(\partial_i \partial_j \partial_k \hat{\pi}_2 \right)^2 + \frac{1}{2} \left(\partial_i \hat{\pi}_2 \partial_j \partial_k \hat{\pi}_2 + \partial_j \hat{\pi}_2 \partial_i \partial_k \hat{\pi}_2 + \partial_k \hat{\pi}_2 \partial_i \partial_j \hat{\pi}_2 \right) \partial_i \partial_j \partial_k \hat{\pi}_2 \right] \\ &+ \sum_i^d \Gamma_{1,2}^i \left[4\rho_{10}\rho_{20}^2 \left(\partial_i \hat{\theta}_1 \right)^2 - 4\ell\rho_{10}\rho_{20}^2 \partial_i \hat{\theta}_1 \partial_i^2 \hat{\theta}_2 + \ell^2 \rho_{10}\rho_{20}^2 \left(\partial_i^2 \hat{\theta}_2 \right)^2 + \frac{\rho_{20}^2}{\rho_{10}} \left(\partial_i \hat{\rho}_1 \right)^2 - \ell \partial_i \hat{\pi}_1 \left(\partial_i \hat{\rho}_2 \right)^2 \right. \\ &\left. - \ell \rho_{20} \partial_i \hat{\pi}_1 \partial_i^2 \hat{\pi}_2 + \frac{\ell^2 \rho_{10}}{4\rho_{20}^2} \left(\partial_i \hat{\pi}_2 \right)^4 + \frac{\ell^2 \rho_{10}}{2\rho_{20}} \left(\partial_i \hat{\pi}_2 \right)^2 \partial_i^2 \hat{\pi}_2 + \frac{1}{4} \ell^2 \rho_{10} \left(\partial_i^2 \hat{\pi}_2 \right)^2 \right]. \end{aligned} \quad (\text{C1})$$

We can obtain the time evolution of $\hat{\theta}_a$,

$$\hat{\theta}_1 = g\hat{\pi}_1 + \sum_{i,j}^d \frac{1}{2} K_1^{ij} \partial_i^2 \partial_j^2 \hat{\pi}_1 + \sum_i^d -2\Gamma_{1,2}^i \frac{\rho_{20}^2}{\rho_{10}} \partial_i^2 \hat{\pi}_1 + 2\Gamma_{1,2}^i \ell \partial_i \hat{\pi}_2 \partial_i^2 \hat{\pi}_2 + \Gamma_{1,2}^i \ell \rho_{20} \partial_i^3 \hat{\pi}_2 \quad (\text{C2})$$

$$\begin{aligned} \hat{\theta}_2 &= g\hat{\pi}_2 + \sum_{i,j,k}^d \frac{1}{2} K_2^{ijk} \left[-\rho_{20} \partial_i^2 \partial_j^2 \partial_k^2 \hat{\pi}_2 - 3 \left(\partial_i \partial_j \partial_k \hat{\pi}_2 \right)^2 - 2\partial_i \partial_j \hat{\pi}_2 \partial_i \partial_j \partial_k^2 \hat{\pi}_2 - 2\partial_i \partial_k^2 \hat{\pi}_2 \partial_i \partial_j^2 \hat{\pi}_2 - 2\partial_i \partial_k \hat{\pi}_2 \partial_i \partial_j^2 \partial_k \hat{\pi}_2 \right. \\ &\left. - 2\partial_j^2 \partial_k \hat{\pi}_2 \partial_i^2 \partial_k \hat{\pi}_2 - 2\partial_j \partial_k^2 \hat{\pi}_2 \partial_i^2 \partial_j \hat{\pi}_2 - 2\partial_j \partial_k \hat{\pi}_2 \partial_i^2 \partial_j \partial_k \hat{\pi}_2 - \partial_i^2 \hat{\pi}_2 \partial_j^2 \partial_k^2 \hat{\pi}_2 - \partial_j^2 \hat{\pi}_2 \partial_i^2 \partial_k^2 \hat{\pi}_2 - \partial_k^2 \hat{\pi}_2 \partial_i^2 \partial_j^2 \hat{\pi}_2 \right] \\ &+ \sum_i^d \Gamma_{1,2}^i \left[2\ell \partial_i^2 \hat{\pi}_1 \partial_i \hat{\pi}_2 + 2\ell \partial_i \hat{\pi}_1 \partial_i^2 \hat{\pi}_2 - \ell \rho_{20} \partial_i^3 \hat{\pi}_1 - \frac{3\ell^2 \rho_{10}}{\rho_{20}^2} \left(\partial_i \hat{\pi}_2 \right)^2 \partial_i^2 \hat{\pi}_2 - \frac{1}{2} \ell^2 \rho_{10} \partial_i^4 \hat{\pi}_2 \right]. \end{aligned} \quad (\text{C3})$$

In the long-wave length limit, we have

$$g\hat{\pi}_1 \gg \sum_{i,j}^d \frac{1}{2} K_1^{ij} \partial_i^2 \partial_j^2 \hat{\pi}_1 + \sum_i^d -2\Gamma_{1,2}^i \frac{\rho_{20}^2}{\rho_{10}} \partial_i^2 \hat{\pi}_1 + 2\Gamma_{1,2}^i \ell \partial_i \hat{\pi}_2 \partial_i^2 \hat{\pi}_2 + \Gamma_{1,2}^i \ell \rho_{20} \partial_i^3 \hat{\pi}_2 \quad (\text{C4})$$

$$g\hat{\pi}_2 \gg \sum_{i,j,k}^d \frac{1}{2} K_2^{ijk} \left[-\rho_{20} \partial_i^2 \partial_j^2 \partial_k^2 \hat{\pi}_2 - 3(\partial_i \partial_j \partial_k \hat{\pi}_2)^2 - 2\partial_i \partial_j \hat{\pi}_2 \partial_i \partial_j \partial_k^2 \hat{\pi}_2 - 2\partial_i \partial_k^2 \hat{\pi}_2 \partial_i \partial_j^2 \hat{\pi}_2 - 2\partial_i \partial_k \hat{\pi}_2 \partial_i \partial_j^2 \partial_k \hat{\pi}_2 \right. \\ \left. - 2\partial_j^2 \partial_k \hat{\pi}_2 \partial_i^2 \partial_k \hat{\pi}_2 - 2\partial_j \partial_k^2 \hat{\pi}_2 \partial_i^2 \partial_j \hat{\pi}_2 - 2\partial_j \partial_k \hat{\pi}_2 \partial_i^2 \partial_j \partial_k \hat{\pi}_2 - \partial_i^2 \hat{\pi}_2 \partial_j^2 \partial_k^2 \hat{\pi}_2 - \partial_j^2 \hat{\pi}_2 \partial_i^2 \partial_k^2 \hat{\pi}_2 - \partial_k^2 \hat{\pi}_2 \partial_i^2 \partial_j^2 \hat{\pi}_2 \right] \\ + \sum_i^d \Gamma_{1,2}^i \left[2\ell \partial_i^2 \hat{\pi}_1 \partial_i \hat{\pi}_2 + 2\ell \partial_i \hat{\pi}_1 \partial_i^2 \hat{\pi}_2 - \ell \rho_{20} \partial_i^3 \hat{\pi}_1 - \frac{3\ell^2 \rho_{10}}{\rho_{20}^2} (\partial_i \hat{\pi}_2)^2 \partial_i^2 \hat{\pi}_2 - \frac{1}{2} \ell^2 \rho_{10} \partial_i^4 \hat{\pi}_2 \right]. \quad (\text{C5})$$

Here we just need to consider the lowest order and get

$$g \gg \sum_i^d 2\Gamma_{1,2}^i \frac{\rho_{20}^2}{\rho_{10}} k_i^2. \quad (\text{C6})$$

If we set an isotropic $K_1^{ij} = \frac{\kappa_1}{2}$, $K_2^{ijk} = \frac{\kappa_2}{2}$, and $\Gamma_{1,2}^i = \frac{\Gamma}{2}$, we can get the coherence length $\xi_c = 2\pi \sqrt{\frac{\Gamma \rho_{20}^2}{g \rho_{10}}}$. Then, we can drop the higher-order terms in Eq. (C1), and get the effective Hamiltonian,

$$\mathcal{H} = \frac{g}{2} \hat{\pi}_1^2 + \frac{g}{2} \hat{\pi}_2^2 + \frac{\kappa_1 \rho_{10}^2}{2} (\partial_i \partial_j \hat{\theta}_1)^2 + \frac{\kappa_2 \rho_{20}^3}{2} (\partial_i \partial_j \partial_k \hat{\theta}_2)^2 + \frac{\Gamma \rho_{10} \rho_{20}^2}{2} (\ell \partial_i^2 \hat{\theta}_2 - 2\partial_i \hat{\theta}_1)^2. \quad (\text{C7})$$

2. Derivation of the Hamiltonian in the momentum space

In the momentum space, we can perform a Fourier transform,

$$H = \frac{1}{(2\pi)^{2d}} \int d^d \mathbf{x} d^d \mathbf{k} d^d \mathbf{k}' e^{i\mathbf{k} \cdot \mathbf{x} + i\mathbf{k}' \cdot \mathbf{x}} \left[\frac{g}{2} \hat{\pi}_1(\mathbf{k}) \hat{\pi}_1(\mathbf{k}') + \frac{g}{2} \hat{\pi}_2(\mathbf{k}) \hat{\pi}_2(\mathbf{k}') + \frac{\kappa_1 \rho_{10}^2}{2} \sum_{i,j}^d k_i^2 k_j^2 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') \right. \\ \left. + \frac{\kappa_2 \rho_{20}^3}{2} \sum_{i,j,k}^d k_i^2 k_j^2 k_k^2 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') + \frac{\Gamma \rho_{10} \rho_{20}^2}{2} \sum_i^d \left[\ell^2 k_i^4 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') + 4k_i^2 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') + 2i\ell k_i^3 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') \right. \right. \\ \left. \left. - 2i\ell k_i^3 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') \right] \right]. \quad (\text{C8})$$

Integrating over the real space, we can obtain a δ function so that only the integral of the momentum space is left in the Hamiltonian,

$$H = \frac{1}{(2\pi)^d} \int d^d \mathbf{k} d^d \mathbf{k}' \delta(\mathbf{k} + \mathbf{k}') \left[\frac{g}{2} \hat{\pi}_1(\mathbf{k}) \hat{\pi}_1(\mathbf{k}') + \frac{g}{2} \hat{\pi}_2(\mathbf{k}) \hat{\pi}_2(\mathbf{k}') + \frac{\kappa_1 \rho_{10}^2}{2} |\mathbf{k}|^4 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') + \frac{\kappa_2 \rho_{20}^3}{2} |\mathbf{k}|^6 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') \right. \\ \left. + \frac{\Gamma \rho_{10} \rho_{20}^2}{2} \sum_i^d \left[\ell^2 k_i^4 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') + 4k_i^2 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') + 2i\ell k_i^3 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_2(\mathbf{k}') - 2i\ell k_i^3 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') \right] \right] \\ = \frac{1}{(2\pi)^d} \int d^d \mathbf{k} \left[\frac{g}{2} \hat{\pi}_1(\mathbf{k}) \hat{\pi}_1(-\mathbf{k}) + \frac{g}{2} \hat{\pi}_2(\mathbf{k}) \hat{\pi}_2(-\mathbf{k}) + \frac{\kappa_1 \rho_{10}^2}{2} |\mathbf{k}|^4 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(-\mathbf{k}) + \frac{\kappa_2 \rho_{20}^3}{2} |\mathbf{k}|^6 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(-\mathbf{k}) \right. \\ \left. + \frac{\Gamma \rho_{10} \rho_{20}^2}{2} \sum_i^d \left[\ell^2 k_i^4 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_2(-\mathbf{k}) + 4k_i^2 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(-\mathbf{k}) + 2i\ell k_i^3 \hat{\theta}_1(\mathbf{k}) \hat{\theta}_2(-\mathbf{k}) - 2i\ell k_i^3 \hat{\theta}_2(\mathbf{k}) \hat{\theta}_1(-\mathbf{k}) \right] \right]. \quad (\text{C9})$$

Using the same method we used to compute Model 1, we can write the Hamiltonian in matrix form,

$$\begin{aligned}
H &= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \left[g \hat{\pi}_1(\mathbf{k}) \hat{\pi}_1(-\mathbf{k}) + g \hat{\pi}_2(\mathbf{k}) \hat{\pi}_2(-\mathbf{k}) \right. \\
&\quad \left. + [\hat{\theta}_1(-\mathbf{k}) \quad \hat{\theta}_2(-\mathbf{k})] \begin{bmatrix} \kappa_1 \rho_{10}^2 |\mathbf{k}|^4 + 4\Gamma \rho_{10} \rho_{20}^2 |\mathbf{k}|^2 & -2i\Gamma \rho_{10} \rho_{20}^2 \ell \sum_i^d k_i^3 \\ 2i\Gamma \rho_{10} \rho_{20}^2 \ell \sum_i^d k_i^3 & \kappa_2 \rho_{20}^3 |\mathbf{k}|^6 + \Gamma \rho_{10} \rho_{20}^2 \ell^2 \sum_i^d k_i^4 \end{bmatrix} \begin{bmatrix} \hat{\theta}_1(\mathbf{k}) \\ \hat{\theta}_2(\mathbf{k}) \end{bmatrix} \right] \\
&= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) \begin{bmatrix} g & 0 & 0 & 0 \\ 0 & g & 0 & 0 \\ 0 & 0 & \kappa_1 \rho_{10}^2 |\mathbf{k}|^4 + 4\Gamma \rho_{10} \rho_{20}^2 |\mathbf{k}| & -2i\Gamma \rho_{10} \rho_{20}^2 \ell \sum_i^d k_i^3 \\ 0 & 0 & 2i\Gamma \rho_{10} \rho_{20}^2 \ell \sum_i^d k_i^3 & \kappa_2 \rho_{20}^3 |\mathbf{k}|^6 + \Gamma \rho_{10} \rho_{20}^2 \ell^2 \sum_i^d k_i^4 \end{bmatrix} \psi(\mathbf{k}) \\
&= \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} \psi^\dagger(\mathbf{k}) M(\mathbf{k}) \psi(\mathbf{k}). \tag{C10}
\end{aligned}$$

In the momentum space, the operators $\hat{\theta}_a(\mathbf{k})$ and the conjugate momentum densities $\hat{\pi}_b(\mathbf{k}')$ satisfy the commutation relations, $[\hat{\theta}_a(\mathbf{k}), \hat{\pi}_b(\mathbf{k}')] = i(2\pi)^d \delta_{ab} \delta^{(d)}(\mathbf{k} + \mathbf{k}')$.

3. Bogoliubov transformation and the transformation matrix $T(\mathbf{k})$

In this Hamiltonian, two species of bosons couple to each other. By the Bogoliubov transformation, we can convert the coupling bosons into two independent modes. We introduce a transformation matrix $T(\mathbf{k})$ which changes the basis from the basis $\psi(\mathbf{k})$ into the basis $\phi(\mathbf{k})$,

$$\phi(\mathbf{k}) = [\hat{\alpha}(\mathbf{k}) \quad \hat{\beta}(\mathbf{k}) \quad \hat{\alpha}^\dagger(-\mathbf{k}) \quad \hat{\beta}^\dagger(-\mathbf{k})]^T = T(\mathbf{k}) \psi(\mathbf{k}), \tag{C11}$$

where $\hat{\alpha}(\mathbf{k})$ and $\hat{\beta}(\mathbf{k})$ are annihilation operators of two independent modes, and they satisfy the commutation relations.

$$[\hat{\alpha}(\mathbf{k}), \hat{\alpha}^\dagger(\mathbf{k}')] = [\hat{\beta}(\mathbf{k}), \hat{\beta}^\dagger(\mathbf{k}')] = (2\pi)^d \delta^{(d)}(\mathbf{k} - \mathbf{k}') \tag{C12}$$

$$[\hat{\alpha}(\mathbf{k}), \hat{\alpha}(\mathbf{k}')] = [\hat{\beta}(\mathbf{k}), \hat{\beta}(\mathbf{k}')] = [\hat{\alpha}(\mathbf{k}), \hat{\beta}(\mathbf{k}')] = [\hat{\alpha}(\mathbf{k}), \hat{\beta}^\dagger(\mathbf{k}')] = [\hat{\beta}(\mathbf{k}), \hat{\alpha}^\dagger(\mathbf{k}')] = 0 \tag{C13}$$

To keep the commutation relations and diagonalize the matrix $M(\mathbf{k})$, the transformation matrix $T(\mathbf{k})$ needs to obey the equation below,

$$T(\mathbf{k}) \Lambda_1 T^\dagger(\mathbf{k}) = \Lambda_2 \tag{C14}$$

$$\tag{C15}$$

where

$$\Lambda_1 = \begin{bmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{bmatrix}, \quad \Lambda_2 = \text{diag}(1, 1, -1, -1). \tag{C16}$$

The diagonalized Hamiltonian becomes

$$H = \frac{1}{2(2\pi)^d} \phi^\dagger(\mathbf{k}) (T^{-1}(\mathbf{k}))^\dagger M(\mathbf{k}) T^{-1}(\mathbf{k}) \phi(\mathbf{k}), \tag{C17}$$

where the matrix $(T^{-1}(\mathbf{k}))^\dagger M(\mathbf{k}) T^{-1}(\mathbf{k}) = \text{diag}(D_1(\mathbf{k}), D_2(\mathbf{k}), D_3(\mathbf{k}), D_4(\mathbf{k}))$ is a diagonal matrix with the diagonal elements:

$$\begin{aligned}
D_1(\mathbf{k}) = D_3(\mathbf{k}) &= \left\{ \frac{g}{2} \left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + \kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
&+ \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\ell^2 \sum_i^d k_i^4 - 4|\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right\}^{\frac{1}{2}} \\
D_2(\mathbf{k}) = D_4(\mathbf{k}) &= \left\{ \frac{g}{2} \left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + \kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
&- \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\ell^2 \sum_i^d k_i^4 - 4|\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right\}^{\frac{1}{2}} \quad (C18)
\end{aligned}$$

The transformation matrix $T(\mathbf{k})$ is given by

$$T(\mathbf{k}) = \begin{bmatrix} T_{11}(\mathbf{k}) & T_{12}(\mathbf{k}) & T_{13}(\mathbf{k}) & T_{14}(\mathbf{k}) \\ T_{21}(\mathbf{k}) & T_{22}(\mathbf{k}) & T_{23}(\mathbf{k}) & T_{24}(\mathbf{k}) \\ T_{31}(\mathbf{k}) & T_{32}(\mathbf{k}) & T_{33}(\mathbf{k}) & T_{34}(\mathbf{k}) \\ T_{41}(\mathbf{k}) & T_{42}(\mathbf{k}) & T_{43}(\mathbf{k}) & T_{44}(\mathbf{k}) \end{bmatrix}, \quad (C19)$$

where the matrix elements are

$$\begin{aligned}
T_{11}(\mathbf{k}) = -T_{31}(\mathbf{k}) &= - \left\{ g \left[-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
&+ \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\ell^2 \sum_i^d k_i^4 - 4|\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right] \right\} / \\
&\left\{ 2\sqrt[4]{2}\Gamma \ell \rho_{10} \rho_{20}^2 \sum_i^d k_i^3 \left[g \left(4 + \left(\left(-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \right. \right. \\
&+ \left. \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\ell^2 \sum_i^d k_i^4 - 4|\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right)^2 \right] \right. \\
&\left. \left. \left(4\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \right) \right) \left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \right. \\
&+ \left. \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\ell^2 \sum_i^d k_i^4 - 4|\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right) \right]^{\frac{1}{2}} \right\}^{\frac{1}{2}}, \quad (C20)
\end{aligned}$$

$$\begin{aligned}
T_{12}(\mathbf{k}) = -T_{32}(\mathbf{k}) &= \left(i2^{3/4} g \right) / \left\{ \left[g \left(4 + \left(\left(-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \right. \right. \right. \\
&+ \left. \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right)^2 \right] \right. \\
&/ 4\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \left[g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
&+ \left. \left. \left. \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \Gamma \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right] \right]^{\frac{1}{2}} \right]^{\frac{1}{2}}, \quad (C21)
\end{aligned}$$

$$\begin{aligned}
T_{13}(\mathbf{k}) = T_{33}(\mathbf{k}) = & \left\{ i \left[-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \\
& \left. \left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \left. \left. \right) \right]^{\frac{1}{4}} \left. \right\} / \\
& \left\{ 2^{3/4} \Gamma \ell \rho_{10} \rho_{20}^2 \sum_i^d k_i^3 \left[g \left(\left(-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \left. \left. \right) \right]^2 / \right. \\
& \left. \left(4\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 + 4 \right) \right]^{\frac{1}{2}} \left. \right\}, \tag{C22}
\end{aligned}$$

$$\begin{aligned}
T_{14}(\mathbf{k}) = T_{34}(\mathbf{k}) = & \left\{ \sqrt[4]{2} \left[g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma k^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \left. \left. \right) \right]^{\frac{1}{4}} \left. \right\} / \\
& \left\{ \left[g \left(\left(-\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 - \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \left. \left. \right) \right]^2 / \right. \\
& \left. \left(4\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 + 4 \right) \right]^{\frac{1}{2}} \left. \right\}, \tag{C23}
\end{aligned}$$

$$\begin{aligned}
T_{21}(\mathbf{k}) = -T_{41}(\mathbf{k}) = & 2^{-\frac{3}{4}} \text{sgn} \left(\sum_i^d k_i^3 \right) \left\{ \left[g \left(\left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& + \sqrt{\left(\kappa_1 k^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \\
& \left. \left(\rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 \right) + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \right) \right] / \\
& \left[\left(\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \right) \right. \\
& \left. \left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& \left. \left. \left. - \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right] \right\}^{\frac{1}{2}}, \tag{C24}
\end{aligned}$$

$$\begin{aligned}
T_{22}(\mathbf{k}) = -T_{42}(\mathbf{k}) = & \left(2i \sqrt[4]{2} \Gamma \ell \rho_{10} \rho_{20}^2 \left| \sum_i^d k_i^3 \right| \right) / \\
& \left\{ \left[\left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& \left. \left. \left. - \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right] \right\}^{\frac{1}{2}} \\
& \left[\left(16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 - \left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right) \right. \\
& \left. \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
& \left. \left. + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right] / \\
& \left(\left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \right. \\
& \left. \left. \left. - \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right) \right] \right\}^{\frac{1}{2}}, \tag{C25}
\end{aligned}$$

$$\begin{aligned}
T_{23}(\mathbf{k}) = T_{43}(\mathbf{k}) = & - \left\{ i \operatorname{sgn} \left(\sum_i^d k_i^3 \right) \left[\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right. \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \left. \right] \Bigg\} / \\
& \left\{ 2\sqrt[4]{2} \left[- \left(g \left(\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right) \right) \right. \right. \\
& \left. \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right) \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \\
& \left. \left. - 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \right) \right] / \left(\left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right) \right. \right. \right. \\
& \left. \left. \left. - \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right) \right] \Bigg\}^{\frac{1}{2}}, \tag{C26}
\end{aligned}$$

$$\begin{aligned}
T_{24}(\mathbf{k}) = T_{44}(\mathbf{k}) = & \left(2^{3/4} \Gamma \ell \rho_{10} \rho_{20}^2 \left| \sum_i^d k_i^3 \right| \right) / \\
& \left\{ \left[- \left(g \left(\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right) \right) \right. \right. \\
& \left. \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 - \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right) \right. \\
& + \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \\
& \left. \left. - 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2 \right) \right] / \left(\left(g \left(\kappa_2 |\mathbf{k}|^6 \rho_{20}^3 + \kappa_1 |\mathbf{k}|^4 \rho_{10}^2 + 4\Gamma |\mathbf{k}|^2 \rho_{10} \rho_{20}^2 + \Gamma \ell^2 \rho_{10} \rho_{20}^2 \sum_i^d k_i^4 \right) \right. \right. \right. \\
& \left. \left. \left. - \sqrt{\left(\kappa_1 |\mathbf{k}|^4 \rho_{10}^2 - \rho_{20}^2 \left(\kappa_2 |\mathbf{k}|^6 \rho_{20} + \rho_{10} \left(\Gamma \ell^2 \sum_i^d k_i^4 - 4\Gamma |\mathbf{k}|^2 \right) \right) \right) \right)^2 + 16\Gamma^2 \ell^2 \rho_{10}^2 \rho_{20}^4 \left(\sum_i^d k_i^3 \right)^2} \right) \right) \right] \Bigg\}^{\frac{1}{2}}. \tag{C27}
\end{aligned}$$

4. Derivation of the Hamiltonian represented by the creation and annihilation operators

After diagonalized, the Hamiltonian becomes

$$H = \frac{1}{2(2\pi)^d} \int d^d \mathbf{k} D_1(\mathbf{k}) \left(\hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + \hat{\alpha}(-\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}) \right) + D_2(\mathbf{k}) \left(\hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \hat{\beta}(-\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k}) \right). \tag{C28}$$

Because of $D_1(\mathbf{k}) = D_1(-\mathbf{k})$, $D_2(\mathbf{k}) = D_2(-\mathbf{k})$, we rewrite the Hamiltonian as

$$\begin{aligned}
H &= \frac{1}{2(2\pi)^d} \int d^d\mathbf{k} D_1(\mathbf{k}) (\hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + \hat{\alpha}(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k})) + D_2(\mathbf{k}) (\hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \hat{\beta}(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k})) \\
&= \frac{1}{(2\pi)^d} \int d^d\mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \int d^d\mathbf{k} \frac{D_1(\mathbf{k}) + D_2(\mathbf{k})}{2} \delta^{(d)}(\mathbf{0}) \\
&= \frac{1}{(2\pi)^d} \int d^d\mathbf{k} D_1(\mathbf{k}) \hat{\alpha}^\dagger(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + D_2(\mathbf{k}) \hat{\beta}^\dagger(\mathbf{k}) \hat{\beta}(\mathbf{k}) + \frac{\tilde{V}}{(2\pi)^d} \int d^d\mathbf{k} \frac{D_1(\mathbf{k}) + D_2(\mathbf{k})}{2}, \tag{C29}
\end{aligned}$$

where \tilde{V} is the volume of the d -dimensional space, the last term is the vacuum zero point energy. The dispersion relations are

$$\omega_1 = D_1(\mathbf{k}), \quad \omega_2 = D_2(\mathbf{k}). \tag{C30}$$

Appendix D: The calculation details of correlation functions of Model Series B

From the main text, we have the Bogoliubov transformation from $\hat{\theta}$ and $\hat{\pi}$ to the independent modes $\hat{\alpha}$ and $\hat{\beta}$,

$$\begin{aligned}
\psi(\mathbf{k}) &= [\hat{\pi}_1(\mathbf{k}) \quad \hat{\pi}_2(\mathbf{k}) \quad \hat{\theta}_1(\mathbf{k}) \quad \hat{\theta}_2(\mathbf{k})]^T = T^{-1}(\mathbf{k}) \phi(\mathbf{k}) = \Lambda_1 T^\dagger(\mathbf{k}) \Lambda_2 \phi(\mathbf{k}) \\
&= \begin{bmatrix} -iT_{13}^*(\mathbf{k}) & -iT_{23}^*(\mathbf{k}) & iT_{33}^*(\mathbf{k}) & iT_{43}^*(\mathbf{k}) \\ -iT_{14}^*(\mathbf{k}) & -iT_{24}^*(\mathbf{k}) & iT_{34}^*(\mathbf{k}) & iT_{44}^*(\mathbf{k}) \\ iT_{11}^*(\mathbf{k}) & iT_{21}^*(\mathbf{k}) & -iT_{31}^*(\mathbf{k}) & -iT_{41}^*(\mathbf{k}) \\ iT_{12}^*(\mathbf{k}) & iT_{22}^*(\mathbf{k}) & -iT_{32}^*(\mathbf{k}) & -iT_{42}^*(\mathbf{k}) \end{bmatrix} \begin{bmatrix} \hat{\alpha}(\mathbf{k}) \\ \hat{\beta}(\mathbf{k}) \\ \hat{\alpha}^\dagger(-\mathbf{k}) \\ \hat{\beta}^\dagger(-\mathbf{k}) \end{bmatrix}. \tag{D1}
\end{aligned}$$

The two-operator correlation function can be written as

$$\langle \hat{\Phi}_1(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle = \rho_{10} e^{-\frac{1}{2} \langle [\hat{\theta}_1(\mathbf{x}) - \hat{\theta}_1(\mathbf{0})]^2 \rangle} = \rho_{10} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle}, \tag{D2}$$

where

$$\begin{aligned}
\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle &= \langle \int \frac{d^d\mathbf{k} d^d\mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1(\mathbf{k}') \rangle = \langle \int \frac{d^d\mathbf{k} d^d\mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \hat{\theta}_1(\mathbf{k}) \hat{\theta}_1^\dagger(-\mathbf{k}') \rangle \\
&= \int \frac{d^d\mathbf{k} d^d\mathbf{k}'}{(2\pi)^{2d}} e^{i\mathbf{k}\cdot\mathbf{x}} \langle [iT_{11}^*(\mathbf{k}) \hat{\alpha}(\mathbf{k}) + iT_{21}^*(\mathbf{k}) \hat{\beta}(\mathbf{k}) - iT_{31}^*(\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}) - iT_{41}^*(\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k})] \\
&\quad [-iT_{11}(-\mathbf{k}') \hat{\alpha}^\dagger(-\mathbf{k}') - iT_{21}(-\mathbf{k}') \hat{\beta}^\dagger(-\mathbf{k}') + iT_{31}(-\mathbf{k}') \hat{\alpha}(\mathbf{k}') + iT_{41}(-\mathbf{k}') \hat{\beta}(\mathbf{k}')] \rangle. \tag{D3}
\end{aligned}$$

For the expectation value of the creation and annihilation operators in the ground state, we have

$$\langle \hat{\alpha}(\mathbf{k}') \hat{\alpha}(\mathbf{k}) \rangle = \langle \hat{\beta}(\mathbf{k}') \hat{\beta}(\mathbf{k}) \rangle = \langle \hat{\alpha}^\dagger(-\mathbf{k}') \hat{\alpha}(\mathbf{k}) \rangle = \langle \hat{\beta}^\dagger(-\mathbf{k}') \hat{\beta}(\mathbf{k}) \rangle = 0, \tag{D4}$$

$$\langle \hat{\alpha}(\mathbf{k}) \hat{\alpha}^\dagger(-\mathbf{k}') \rangle = \langle \hat{\beta}(\mathbf{k}) \hat{\beta}^\dagger(-\mathbf{k}') \rangle = (2\pi)^d \delta^{(d)}(\mathbf{k} + \mathbf{k}'). \tag{D5}$$

Then we can obtain the correlation function,

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle = \int \frac{d^d\mathbf{k}}{(2\pi)^d} e^{i\mathbf{k}\cdot\mathbf{x}} (|T_{11}(\mathbf{k})|^2 + |T_{21}(\mathbf{k})|^2). \tag{D6}$$

Similarly, we can get

$$\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \int \frac{d^d\mathbf{k}}{(2\pi)^d} e^{i\mathbf{k}\cdot\mathbf{x}} (|T_{12}(\mathbf{k})|^2 + |T_{22}(\mathbf{k})|^2). \tag{D7}$$

Therefore, to calculate correlation functions, we need to calculate $|T_{11}(\mathbf{k})|^2$, $|T_{21}(\mathbf{k})|^2$, $|T_{12}(\mathbf{k})|^2$, and $|T_{22}(\mathbf{k})|^2$ in different spatial dimensions.

1. One spatial dimension

In one spatial dimension, the dispersion relations are

$$\omega_1 \approx \sqrt{4g\Gamma\rho_{10}\rho_{20}^2} |k|, \quad \omega_2 \approx \sqrt{\kappa_2 g \rho_{20}^3 + \frac{\kappa_1 g \rho_{10}^2 \ell^2}{4}} |k|^3 \quad (\text{D8})$$

We can approximate these four terms,

$$\begin{aligned} |T_{11}(\mathbf{k})|^2 &\approx \frac{\sqrt{g\Gamma\rho_{10}}}{4\Gamma\rho_{10}\rho_{20}} \frac{1}{|k|} = \frac{1}{c_{11}|k|}, & |T_{12}(\mathbf{k})|^2 &\approx \frac{\sqrt{g\Gamma\rho_{10}\ell^2}}{16\Gamma\rho_{10}\rho_{20}} |k| = \frac{|k|}{c_{12}}, \\ |T_{21}(\mathbf{k})|^2 &\approx \frac{\sqrt{g}\ell^2}{4\sqrt{\kappa_1\rho_{10}^2\ell^2 + 4\kappa_2\rho_{20}^3}} \frac{1}{|k|} = \frac{1}{c_{21}|k|}, & |T_{22}(\mathbf{k})|^2 &\approx \frac{\sqrt{g}}{\sqrt{\kappa_1\rho_{10}^2\ell^2 + 4\kappa_2\rho_{20}^3}} \frac{1}{|k|^3} = \frac{1}{c_{22}|k|^3}. \end{aligned} \quad (\text{D9})$$

The correlation function of $\hat{\theta}_1$ is

$$\begin{aligned} \langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle &= \int \frac{dk}{2\pi} e^{ikx} \left(\frac{1}{c_{11}|k|} + \frac{1}{c_{21}|k|} \right) = \frac{1}{2\pi} \left(\frac{1}{c_{11}} + \frac{1}{c_{21}} \right) \int dk \frac{e^{ikx}}{|k|} \\ &= \frac{1}{2\pi} \left(\frac{1}{c_{11}} + \frac{1}{c_{21}} \right) \left(\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{-k} + \int_{\frac{2\pi}{L}}^{\infty} dk \frac{e^{ikx}}{k} \right). \end{aligned} \quad (\text{D10})$$

Considering $L > 0, x > 0$, we have the two integrals

$$\int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{-k} + \int_{\frac{2\pi}{L}}^{\infty} \frac{e^{ikx}}{k} = \Gamma\left(0, -\frac{2i\pi x}{L}\right) + \Gamma\left(0, \frac{2i\pi x}{L}\right) \stackrel{L \rightarrow \infty}{=} -2\left(\gamma + \ln\left(\frac{2\pi x}{L}\right)\right), \quad (\text{D11})$$

where γ is the Euler-Mascheroni constant. Then, we can get the correlation functions,

$$\langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle \stackrel{L \rightarrow \infty}{=} \frac{1}{2\pi} \left(\frac{1}{c_{11}} + \frac{1}{c_{21}} \right) \left(-2\left(\gamma + \ln\left(\frac{2\pi|x|}{L}\right)\right) \right) = \frac{c_{11} + c_{21}}{\pi c_{11} c_{21}} \left(-\gamma + \ln \frac{L}{2\pi|x|} \right). \quad (\text{D12})$$

To get $\langle \hat{\theta}_1^2(0) \rangle = \lim_{x \rightarrow 0} \langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle$, we just need to replace $|x|$ by ξ_c . ξ_c is the coherence length,

$$\langle \hat{\theta}_1^2(0) \rangle = \langle \hat{\theta}_1(\xi_c) \hat{\theta}_1(0) \rangle = \frac{c_{11} + c_{21}}{\pi c_{11} c_{21}} \left(-\gamma + \ln \frac{L}{2\pi\xi_c} \right). \quad (\text{D13})$$

Then, we obtain the correlation function $\langle \hat{\Phi}_1^\dagger(x) \hat{\Phi}_1(0) \rangle$,

$$\langle \hat{\Phi}_1^\dagger(x) \hat{\Phi}_1(0) \rangle = \rho_{10} e^{\langle \hat{\theta}_1(x) \hat{\theta}_1(0) \rangle - \langle \hat{\theta}_1^2(0) \rangle} = \rho_{10} \left(\frac{|x|}{\xi_c} \right)^{-\frac{c_{11} + c_{21}}{\pi c_{11} c_{21}}}. \quad (\text{D14})$$

The correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ is a power-law decay. The correlation function of $\hat{\theta}_2$ is

$$\langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle = \int \frac{dk}{2\pi} e^{ikx} \left(\frac{|k|}{c_{12}} + \frac{1}{c_{22}|k|^3} \right) = \frac{1}{2\pi c_{12}} \int dk e^{ikx} |k| + \frac{1}{2\pi c_{22}} \int dk \frac{e^{ikx}}{|k|^3}. \quad (\text{D15})$$

Here we calculate the these integrals separately,

$$\int_{-\frac{2\pi}{\xi_c}}^{\frac{2\pi}{\xi_c}} dk e^{ikx} |k| = \frac{4\pi|x| \sin\left(\frac{2\pi|x|}{\xi_c}\right) - 4\xi_c \sin^2\left(\frac{\pi|x|}{\xi_c}\right)}{|x|^2 \xi_c}, \quad (\text{D16})$$

$$\int_{\frac{2\pi}{L}}^{\infty} dk \frac{e^{ikx}}{|k|^3} + \int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{e^{ikx}}{|k|^3} \stackrel{L \rightarrow \infty}{=} \frac{L^2}{4\pi^2} + \frac{|x|^2}{2} \left(-3 + 2\gamma + 2 \ln\left(\frac{2\pi|x|}{L}\right) \right). \quad (\text{D17})$$

Then, we obtain the correlation functions $\langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle$ and $\langle \hat{\theta}_2^2(0) \rangle$,

$$\begin{aligned} \langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle &= \int \frac{dk}{2\pi} e^{ikx} \left(\frac{|k|}{c_{12}} + \frac{1}{c_{22} |k|^3} \right) = \frac{1}{2\pi c_{12}} \int dk e^{ikx} |k| + \frac{1}{2\pi c_{22}} \int dk \frac{e^{ikx}}{|k|^3} \\ &= \frac{1}{2\pi c_{12}} \left(\frac{4\pi |x| \sin\left(\frac{2\pi|x|}{\xi_c}\right) - 4\xi_c \sin^2\left(\frac{\pi|x|}{\xi_c}\right)}{|x|^2 \xi_c} \right) + \frac{1}{2\pi c_{22}} \left(\frac{L^2}{4\pi^2} + \frac{|x|^2}{2} \left(-3 + 2\gamma + 2 \ln\left(\frac{2\pi|x|}{L}\right) \right) \right), \end{aligned} \quad (\text{D18})$$

$$\langle \hat{\theta}_2^2(0) \rangle = \int \frac{dk}{2\pi} \left(\frac{|k|}{c_{12}} + \frac{1}{c_{22} |k|^3} \right) = \int_{-\frac{2\pi}{\xi_c}}^{\frac{2\pi}{\xi_c}} dk |k| + \int_{\frac{2\pi}{L}}^{\infty} dk \frac{1}{|k|^3} + \int_{-\infty}^{-\frac{2\pi}{L}} dk \frac{1}{|k|^3} = \frac{1}{2\pi c_{12}} \frac{4\pi^2}{\xi_c^2} + \frac{1}{2\pi c_{22}} \frac{L^2}{4\pi^2}. \quad (\text{D19})$$

Then, we obtain the correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$,

$$\langle \hat{\Phi}_2^\dagger(x) \hat{\Phi}_2(0) \rangle = \rho_{20} e^{\langle \hat{\theta}_2(x) \hat{\theta}_2(0) \rangle - \langle \hat{\theta}_2^2(0) \rangle} \rightarrow 0. \quad (\text{D20})$$

The correlation function $\langle \hat{\Phi}_2^\dagger(x) \hat{\Phi}_2(0) \rangle$ decays exponentially with $|x|^2$. The correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ is a power-law decay. Model Series B does not break the charge symmetry and is algebraically ordered in one spatial dimension.

2. Two spatial dimensions

In two spatial dimensions, we set $k_1 = |\mathbf{k}| \cos \theta$, and $k_2 = |\mathbf{k}| \sin \theta$, for $|\mathbf{k}| \geq 0$, $0 \leq \theta < 2\pi$. Then, we can get the dispersion relations,

$$\omega_1 \approx \sqrt{4g\Gamma\rho_{10}\rho_{20}^2} |\mathbf{k}|, \quad (\text{D21})$$

$$\omega_2 \approx \begin{cases} \frac{1}{2} \rho_{20} \ell |\sin 2\theta| \sqrt{g\rho_{10}\Gamma(1 - \sin 2\theta)} |\mathbf{k}|^2 & \left(\theta \neq 0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi, \frac{5\pi}{4}, \frac{3\pi}{2}, \frac{7\pi}{4} \right) \\ \sqrt{\kappa_2 g \rho_{20}^3 + \frac{\kappa_1 g \rho_{10}^2 \ell^2}{4}} |\mathbf{k}|^3 & \left(\theta = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2} \right) \\ \sqrt{\kappa_2 g \rho_{20}^3 + \frac{\kappa_1 g \rho_{10}^2 \ell^2}{8}} |\mathbf{k}|^3 & \left(\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4} \right) \end{cases} \quad (\text{D22})$$

In two spatial dimensions, the dispersion relations are anisotropic. The same is true for the two terms $|T_{21}(\mathbf{k})|^2$ and $|T_{22}(\mathbf{k})|^2$,

$$|T_{11}(\mathbf{k})|^2 \approx \frac{\sqrt{g\Gamma\rho_{10}}}{4\Gamma\rho_{10}\rho_{20}} \frac{1}{|\mathbf{k}|}, \quad |T_{12}(\mathbf{k})|^2 \approx \frac{\sqrt{g\Gamma\rho_{10}} \ell^2 (\cos^3 \theta + \sin^3 \theta)^2}{16\Gamma\rho_{10}\rho_{20}} |\mathbf{k}|, \quad (\text{D23})$$

$$|T_{21}(\mathbf{k})|^2 \begin{cases} \approx \frac{\sqrt{g\Gamma\rho_{10}} \ell (3 \cos \theta + \cos 3\theta + 4 \sin^3 \theta)^2}{64\Gamma\rho_{10}\rho_{20} |\sin 2\theta| \sqrt{1 - \sin 2\theta}} & \left(\theta \neq 0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi, \frac{5\pi}{4}, \frac{3\pi}{2}, \frac{7\pi}{4} \right) \\ \approx \frac{\sqrt{g} \ell^2}{4\sqrt{\kappa_1 \rho_{10}^2 \ell^2 + 4\kappa_2 \rho_{20}^3}} \frac{1}{|\mathbf{k}|} & \left(\theta = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2} \right) \\ \approx \frac{\sqrt{g} \ell^2}{4\sqrt{2} \sqrt{\kappa_1 \rho_{10}^2 \ell^2 + 8\kappa_2 \rho_{20}^3}} \frac{1}{|\mathbf{k}|} & \left(\theta = \frac{\pi}{4}, \frac{5\pi}{4} \right) \\ = 0 & \left(\theta = \frac{3\pi}{4}, \frac{7\pi}{4} \right) \end{cases}, \quad (\text{D24})$$

$$|T_{22}(\mathbf{k})|^2 \begin{cases} \approx \frac{\sqrt{g\Gamma\rho_{10}}}{\Gamma\ell\rho_{10}\rho_{20}\sqrt{1-\sin 2\theta}|\sin 2\theta||\mathbf{k}|^2} & \left(\theta \neq 0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi, \frac{5\pi}{4}, \frac{3\pi}{2}, \frac{7\pi}{4}\right) \\ \approx \frac{\sqrt{g}}{\sqrt{\kappa_1\rho_{10}^2\ell^2 + 4\kappa_2\rho_{20}^3}} \frac{1}{|\mathbf{k}|^3} & \left(\theta = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}\right) \\ \approx \frac{\sqrt{2g}}{\sqrt{\kappa_1\rho_{10}^2\ell^2 + 8\kappa_2\rho_{20}^3}} \frac{1}{|\mathbf{k}|^3} & \left(\theta = \frac{\pi}{4}, \frac{5\pi}{4}\right) \\ = 0 & \left(\theta = \frac{3\pi}{4}, \frac{7\pi}{4}\right) \end{cases}. \quad (\text{D25})$$

In the long-wave length limit, to simplify the calculation, we can always choose the lowest order to give a rough approximation,

$$|T_{11}(\mathbf{k})|^2 \approx \frac{1}{c_{11}|\mathbf{k}|}, \quad |T_{12}(\mathbf{k})|^2 \lesssim \frac{|\mathbf{k}|}{c_{12}}, \quad |T_{21}(\mathbf{k})|^2 \lesssim \frac{1}{c_{21}|\mathbf{k}|}, \quad |T_{22}(\mathbf{k})|^2 \lesssim \frac{1}{c_{22}|\mathbf{k}|^3}, \quad (\text{D26})$$

where $c_{11} = \frac{4\Gamma\rho_{10}\rho_{20}}{\sqrt{g\Gamma\rho_{10}}}$, $c_{12} = \frac{16\Gamma\rho_{10}\rho_{20}}{\sqrt{g\Gamma\rho_{10}\ell^2}}$, $c_{21} = \frac{4\sqrt{\kappa_1\rho_{10}^2\ell^2 + 4\kappa_2\rho_{20}^3}}{\sqrt{g\ell^2}}$, $c_{22} = \frac{\sqrt{\kappa_1\rho_{10}^2\ell^2 + 8\kappa_2\rho_{20}^3}}{\sqrt{2g}}$.

Following this rough approximation, if the calculation results show that there is a spontaneous breaking of charge symmetry, there must be, but not the opposite. With the above approximation, we can compute correlation functions,

$$\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle = \int \frac{kd\mathbf{k}d\theta}{(2\pi)^2} e^{ik|\mathbf{x}|\cos\theta} \left(\frac{1}{c_{11}k} + \frac{1}{c_{21}k} \right) = \frac{c_{11} + c_{21}}{4\pi^2 c_{11} c_{21}} \int dk d\theta e^{ik|\mathbf{x}|\cos\theta} = \frac{c_{11} + c_{21}}{2\pi c_{11} c_{21} |\mathbf{x}|}. \quad (\text{D27})$$

To get $\langle \hat{\theta}_1^2(\mathbf{0}) \rangle = \lim_{|\mathbf{x}| \rightarrow 0} \langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle$, we just need to replace $|\mathbf{x}|$ by ξ_c . ξ_c is the coherence length,

$$\langle \hat{\theta}_1^2(\mathbf{0}) \rangle = \langle \hat{\theta}_1(\xi_c) \hat{\theta}_1(\mathbf{0}) \rangle = \frac{c_{11} + c_{21}}{2\pi c_{11} c_{21} \xi_c}. \quad (\text{D28})$$

Then, we can calculate the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$,

$$\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle = \rho_{10} e^{\langle \hat{\theta}_1(\mathbf{x}) \hat{\theta}_1(\mathbf{0}) \rangle - \langle \hat{\theta}_1^2(\mathbf{0}) \rangle} \stackrel{|\mathbf{x}| \rightarrow \infty}{=} \rho_{10} e^{-\frac{c_{11} + c_{21}}{2\pi c_{11} c_{21} \xi_c}}. \quad (\text{D29})$$

The correlation function of $\hat{\theta}_2$ is

$$\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle = \int \frac{kd\mathbf{k}d\theta}{(2\pi)^2} e^{ik|\mathbf{x}|\cos\theta} \left(\frac{k}{c_{12}} + \frac{1}{c_{22}k^3} \right) = \frac{1}{4\pi^2 c_{12}} \int dk d\theta e^{ik|\mathbf{x}|\cos\theta} k^2 + \frac{1}{4\pi^2 c_{22}} \int dk d\theta e^{ik|\mathbf{x}|\cos\theta} \frac{1}{k^2}. \quad (\text{D30})$$

Here we compute the two integrals separately.

$$\begin{aligned} \frac{1}{4\pi^2 c_{12}} \int dk d\theta e^{ik|\mathbf{x}|\cos\theta} k^2 &= \frac{1}{4\pi^2 c_{12}} \int_0^{\frac{2\pi}{\xi_c}} 2\pi J_0(k|\mathbf{x}|) k^2 \\ &= \frac{1}{4\pi^2 c_{12}} \frac{2\pi^3 \left[J_1\left(\frac{2\pi|\mathbf{x}|}{\xi_c}\right) \left(4|\mathbf{x}| - \xi_c H_0\left(\frac{2\pi|\mathbf{x}|}{\xi_c}\right)\right) + \xi_c J_0\left(\frac{2\pi|\mathbf{x}|}{\xi_c}\right) H_1\left(\frac{2\pi|\mathbf{x}|}{\xi_c}\right) \right]}{|\mathbf{x}|^2 \xi_c^2} \stackrel{|\mathbf{x}| \rightarrow \infty}{=} 0 \end{aligned} \quad (\text{D31})$$

where $J_a(z)$ is the Bessel function of the first kind, and $H_n(z)$ is the Struve function.

$$\frac{1}{4\pi^2 c_{22}} \int dk d\theta e^{ik|\mathbf{x}|\cos\theta} \frac{1}{k^2} = \frac{1}{4\pi^2 c_{22}} \int_{\frac{2\pi}{L}}^{\infty} \frac{2\pi J_0(k|\mathbf{k}|)}{k^2} \stackrel{L \rightarrow \infty}{=} \frac{1}{4\pi^2 c_{22}} (L - 2\pi|\mathbf{x}|) \quad (\text{D32})$$

Then, we can get

$$\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle \stackrel{L \rightarrow \infty, |\mathbf{x}| \rightarrow \infty}{=} \frac{1}{4\pi^2 c_{22}} (L - 2\pi|\mathbf{x}|). \quad (\text{D33})$$

We can also calculate $\langle \hat{\theta}_2^2(\mathbf{0}) \rangle$,

$$\langle \hat{\theta}_2^2(\mathbf{0}) \rangle = \frac{1}{4\pi^2 c_{12}} \int_0^{\frac{2\pi}{\xi_c}} dk \int_0^{2\pi} d\theta k^2 + \frac{1}{4\pi^2 c_{22}} \int_{\frac{2\pi}{L}}^{\infty} dk \int_0^{2\pi} d\theta \frac{1}{k^2} = \frac{4\pi^2}{3c_{12}\xi_c^3} + \frac{L}{4\pi^2 c_{22}}. \quad (\text{D34})$$

Then, we can calculate the correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$,

$$\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle = \rho_{20} e^{\langle \hat{\theta}_2(\mathbf{x}) \hat{\theta}_2(\mathbf{0}) \rangle - \langle \hat{\theta}_2^2(\mathbf{0}) \rangle} \Big|_{|\mathbf{x}| \rightarrow \infty} \rho_{20} e^{-\frac{4\pi^2}{3c_{12}\epsilon_c^3} - \frac{|\mathbf{x}|}{2\pi c_{22}}} \Big|_{|\mathbf{x}| \rightarrow \infty} 0. \quad (\text{D35})$$

According to the results of the above analytical calculation, the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant and $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ is a power-law decay when $|\mathbf{x}| \rightarrow \infty$. We can test these results numerically. By setting $\kappa_1 = 1$, $\kappa_2 = 1$, $\Gamma = 1$, $g = 1$, $\rho_{10} = 1$, $\rho_{20} = 1$, $\ell = 1$, $\xi_c = 2\pi$, $\frac{2\pi}{\xi_c} = 1$, $|\mathbf{x}| \sim 10^{10}$, $L \sim 10^{20}$, we can calculate the correlation function by using the numerical methods. Firstly, we can calculate $\langle \hat{\Phi}_1^\dagger(10^{10}, 0) \hat{\Phi}_1(\mathbf{0}) \rangle$ numerically. For the integrals, we have

$$\int_0^1 dk \int_0^{2\pi} d\theta |T_{11}(\mathbf{k})|^2 k e^{i10^{10}k \cos \theta} - \int_0^1 dk \int_0^{2\pi} d\theta |T_{11}(\mathbf{k})|^2 k \approx (5.90 \times 10^{-13} + 8.41 \times 10^{-16}i) - 1.42 \approx -1.42, \quad (\text{D36})$$

$$\int_0^1 dk \int_0^{2\pi} d\theta |T_{21}(\mathbf{k})|^2 k e^{i10^{10}k \cos \theta} - \int_0^1 dk \int_0^{2\pi} d\theta |T_{21}(\mathbf{k})|^2 k \approx (-5.23 \times 10^{-4} - 6.08 \times 10^{-4}i) - 0.351 \approx -0.351. \quad (\text{D37})$$

Then, we obtain the result,

$$\langle \hat{\Phi}_1^\dagger(10^{10}, 0) \hat{\Phi}_1(\mathbf{0}) \rangle \approx e^{\frac{1}{(2\pi)^2}(-1.42-0.351)} \approx 0.956. \quad (\text{D38})$$

From this we can believe that $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant when $|\mathbf{x}| \rightarrow \infty$. We can also calculate the correlation function $\langle \hat{\Phi}_2^\dagger(10^{10}, 0) \hat{\Phi}_2(\mathbf{0}) \rangle$. For the integrals, we have

$$\int_0^1 dk \int_0^{2\pi} d\theta |T_{12}(\mathbf{k})|^2 k e^{i10^{10}k \cos \theta} - \int_0^1 dk \int_0^{2\pi} d\theta |T_{12}(\mathbf{k})|^2 k \approx (2.34 \times 10^{-4} + 3.56 \times 10^{-4}i) - 0.0623 \approx -0.0623, \quad (\text{D39})$$

$$\int_{10^{-20}}^1 dk \int_0^{2\pi} d\theta |T_{22}(\mathbf{k})|^2 k \left(e^{i10^{10}k \cos \theta} - 1 \right) \approx -874 - 10.9i \approx -874, \quad (\text{D40})$$

$$\langle \hat{\Phi}_2^\dagger(10^{10}, 0) \hat{\Phi}_2(\mathbf{0}) \rangle \approx e^{\frac{1}{(2\pi)^2}(-874)} \approx 2.43 \times 10^{-10}. \quad (\text{D41})$$

From this we can know that $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ do not saturate to a constant when $|\mathbf{x}| \rightarrow \infty$. The correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ is shown in Fig. 2(a) below. The figure shows that the correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ decays power-law. Since the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ decays in a power law when Γ is 0, the value of Γ affects the value of the correlation function. We can also give the dependence of the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ on Γ , see the blue line of Fig. 2(b). The constant behavior of the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ shows the spontaneous breaking of the charge symmetry $U(1)_{1,C}$ of bosons of species 1. Model Series B has a true ODLRO in two spatial dimensions.

3. Three spatial dimensions

In three spatial dimensions, we set $k_1 = |\mathbf{k}| \cos \theta \sin \psi$, $k_2 = |\mathbf{k}| \sin \theta \sin \psi$, $k_3 = |\mathbf{k}| \cos \psi$. Due to the large error in the analytical approximation of the 3D results, we directly use numerical methods to calculate the correlation functions,

$$\int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{11}(\mathbf{k})|^2 k^2 \sin \psi e^{i10^{10}k \cos \theta \sin \psi} - \int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{11}(\mathbf{k})|^2 k^2 \sin \psi \approx (-0.0239 + 0.0113i) - 1.39 \approx -1.41, \quad (\text{D42})$$

$$\int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{21}(\mathbf{k})|^2 k^2 \sin \psi e^{i10^{10}k \cos \theta \sin \psi} - \int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{21}(\mathbf{k})|^2 k^2 \sin \psi \approx (-9.34 \times 10^{-4} - 9.71 \times 10^{-4}i) - 0.258 \approx -0.258. \quad (\text{D43})$$

From the above results, we can obtain the correlation function $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$,

$$\langle \hat{\Phi}_1^\dagger(10^{10}, 0, 0) \hat{\Phi}_1(\mathbf{0}) \rangle \approx e^{\frac{1}{(2\pi)^3}(-1.41-0.258)} \approx 0.993. \quad (\text{D44})$$

From this we can believe that $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ saturates to a constant when $|\mathbf{x}| \rightarrow \infty$.

$$\int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{12}(\mathbf{k})|^2 k^2 \sin \psi e^{i10^{10}k \cos \theta \sin \psi} - \int_0^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{12}(\mathbf{k})|^2 k^2 \sin \psi \\ \approx (1.05 \times 10^{-3} + 6.24 \times 10^{-5}i) - 0.0654 \approx -0.064, \quad (\text{D45})$$

$$\int_{10^{-20}}^1 dk \int_0^{2\pi} d\theta \int_0^\pi d\psi |T_{22}(\mathbf{k})|^2 k^2 \sin \psi \left(e^{i10^{10}k \cos \theta \sin \psi} - 1 \right) \approx -11.6, \quad (\text{D46})$$

$$\langle \hat{\Phi}_2^\dagger(10^{10}, 0, 0) \hat{\Phi}_2(\mathbf{0}) \rangle \approx e^{\frac{1}{(2\pi)^3}(-0.064-11.6)} \approx 0.954. \quad (\text{D47})$$

From this we can believe that $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ saturates to a constant when $|\mathbf{x}| \rightarrow \infty$. We can also give the dependence of the correlation function $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ on Γ , see the orange line of Fig. 2(b). The constant behaviors of the correlation functions $\langle \hat{\Phi}_1^\dagger(\mathbf{x}) \hat{\Phi}_1(\mathbf{0}) \rangle$ and $\langle \hat{\Phi}_2^\dagger(\mathbf{x}) \hat{\Phi}_2(\mathbf{0}) \rangle$ shows the spontaneous breaking of charge symmetry $U(1)_{1,C}$ and $U(1)_{2,C}$. This system has a true ODLRO in three spatial dimensions.




Fig. 2. Results obtained by numerical calculations. Fig. 2(a) shows the graph of the correlation function $\langle \hat{\Phi}_2^\dagger(x, 0) \hat{\Phi}_2(\mathbf{0}) \rangle$ decaying with increasing x in two spatial dimensions. In Fig. 2(b), the blue line shows the variation of correlation function $\langle \hat{\Phi}_1^\dagger(10^{10}, 0) \hat{\Phi}_1(\mathbf{0}) \rangle$ with increasing Γ in two spatial dimensions, and the orange line shows the variation of correlation function $\langle \hat{\Phi}_2^\dagger(10^{10}, 0, 0) \hat{\Phi}_2(\mathbf{0}) \rangle$ with increasing Γ in three spatial dimensions.

Appendix E: Mean field theory of Lattice Model Series A

According to Ref. [59], we can also give the mean field theory of Lattice Model Series A. We define the operators of dipoles,

$$\hat{d}_{a,j}^i = \hat{b}_{a,j+\hat{x}_i}^\dagger \hat{b}_{a,j}. \quad (\text{E1})$$

Then, the hopping terms can be written as:

$$H_{\text{hop}} = \sum_a \left[\sum_{\mathbf{i}, \mathbf{j}} -t_a^{(ij)} \left(\hat{d}_{a,\mathbf{i}}^i \right)^\dagger \hat{d}_{a,\mathbf{i}+\hat{x}_j}^i + \sum_{b \neq a} \sum_{\mathbf{i}, \mathbf{i}} -t_{ab}^{(i)} \left(\hat{d}_{a,\mathbf{i}}^i \right)^\dagger \hat{d}_{b,\mathbf{i}}^i \right] + \text{h.c.} \\ = - \sum_{a,b} \sum_{\mathbf{i}, \mathbf{j}, \mathbf{i}} \left(\hat{d}_{a,\mathbf{i}}^i \right)^\dagger \mathcal{A}_{\mathbf{ij}}^{ab,i} \hat{d}_{b,\mathbf{j}}^i, \quad (\text{E2})$$

where

$$[\mathcal{A}^{ab,i}]_{\mathbf{ij}} = \sum_j (\delta_{\mathbf{i},\mathbf{j}+\hat{x}_j} + \delta_{\mathbf{i},\mathbf{j}-\hat{x}_j}) t_a^{(ij)} \delta_{ab} + \delta_{\mathbf{i},\mathbf{j}} t_{ab}^{(i)} (1 - \delta_{a,b}). \quad (\text{E3})$$

We introduce a parameter $\Psi_{a,i}$ as

$$\Psi_{a,i} = \sum_b \sum_j \mathcal{A}_{\mathbf{ij}}^{ab,i} \langle \hat{d}_{b,\mathbf{j}}^i \rangle = \sum_j \langle \hat{d}_{a,\mathbf{i}-\hat{x}_j}^i + \hat{d}_{a,\mathbf{i}+\hat{x}_j}^i \rangle t_a^{(ij)} + \sum_{b \neq a} \langle \hat{d}_{b,\mathbf{i}}^i \rangle t_{ab}^{(i)}. \quad (\text{E4})$$

By mean-field approximation, we can obtain

$$H_{\text{hop}} \approx - \sum_a \sum_{\mathbf{i},i} \left[\Psi_{a,i} (\hat{d}_{a,\mathbf{i}}^i)^\dagger + \Psi_{a,i}^* \hat{d}_{a,\mathbf{i}}^i + \frac{|\Psi_{a,i}|^2}{2 \sum_j t_a^{(ij)} + \sum_{b \neq a} t_{ab}^{(i)}} \right]. \quad (\text{E5})$$

We consider the co-ordination number in the square lattices, and set $t_a^{(ij)} = t_1$, $t_{ab}^{(i)} = t_2$, which mean the coefficients of the hopping terms for the same species and different species respectively. The hopping terms become

$$H_{\text{hop}} = - \sum_a \sum_{\mathbf{i},i} \left[\Psi_{a,i} (\hat{d}_{a,\mathbf{i}}^i)^\dagger + \Psi_{a,i}^* \hat{d}_{a,\mathbf{i}}^i + \frac{|\Psi_{a,i}|^2}{2dt_1 + t_2} \right]. \quad (\text{E6})$$

After the mean-field approach, the Hamiltonian becomes

$$H = \sum_{\mathbf{i}} \left[h_{\mathbf{i}}^{(0)} + \sum_a \sum_i \left(\Psi_{a,i} (\hat{d}_{a,\mathbf{i}}^i)^\dagger + \Psi_{a,i}^* \hat{d}_{a,\mathbf{i}}^i \right) \right], \quad (\text{E7})$$

where

$$h_{\mathbf{i}}^{(0)} = \sum_a \left[-\mu_a \hat{n}_{a,\mathbf{i}} + \frac{U}{2} \hat{n}_{a,\mathbf{i}} (\hat{n}_{a,\mathbf{i}} - 1) + \sum_i \frac{|\Psi_{a,i}|^2}{2dt_1 + t_2} \right]. \quad (\text{E8})$$

For simplicity, we set $\mu_a = \mu$. Considering the energy of site i and one of its nearest neighbour and the filling numbers of bosons of species 1 and 2 are n ,

$$E_{a,(n,n)}^{(0)} = \langle n, n | \left(h_{a,\mathbf{i}}^{(0)} + h_{a,\mathbf{i}+\hat{x}_i}^{(0)} \right) | n, n \rangle = 2 \left[\mu n + \frac{U}{2} n(n-1) + \sum_i \frac{|\Psi_{a,i}|^2}{2dt_1 + t_2} \right]. \quad (\text{E9})$$

The expression for the second-order energy is given by

$$E_{a,(n,n)}^{(2)} = \sum_x |\Psi_{a,i}|^2 \sum_{(m,m') \neq (n,n)} \frac{\left| \langle n, n | \left(\hat{d}_{a,\mathbf{i}}^i \right)^\dagger + \hat{d}_{a,\mathbf{i}}^i | m, m' \rangle \right|^2}{E_{a,(m,m')}^{(0)} - E_{a,(n,n)}^{(0)}}. \quad (\text{E10})$$

According to the operator $\hat{d}_{a,\mathbf{i}}^i = \hat{b}_{a,\mathbf{i}} \hat{b}_{a,\mathbf{i}+\hat{x}_i}^\dagger$, we just need $(m, m') = (n-1, n+1)$ or $(n+1, n-1)$. We can calculate $E_{a,(n-1,n+1)}^{(0)}$ and $E_{a,(n+1,n-1)}^{(0)}$.

$$E_{a,(n-1,n+1)}^{(0)} = E_{a,(n+1,n-1)}^{(0)} = -\mu(n+1) + \frac{U}{2}(n+1)n - \mu(n-1) + \frac{U}{2}(n-1)(n-2) = E_{a,(n,n)}^{(0)} + U \quad (\text{E11})$$

Then, we can obtain the second-order energy,

$$\begin{aligned} E_{a,(n,n)}^{(2)} &= \sum_i |\Psi_{a,i}|^2 \left[\frac{\left| \langle n, n | \hat{b}_{a,\mathbf{i}}^\dagger \hat{b}_{a,\mathbf{i}+\hat{x}_i} + \hat{b}_{a,\mathbf{i}} \hat{b}_{a,\mathbf{i}+\hat{x}_i}^\dagger | n-1, n+1 \rangle \right|^2}{E_{a,(n-1,n+1)}^{(0)} - E_{a,(n,n)}^{(0)}} + \frac{\left| \langle n, n | \hat{b}_{a,\mathbf{i}}^\dagger \hat{b}_{a,\mathbf{i}+\hat{x}_i} + \hat{b}_{a,\mathbf{i}} \hat{b}_{a,\mathbf{i}+\hat{x}_i}^\dagger | n+1, n-1 \rangle \right|^2}{E_{a,(n+1,n-1)}^{(0)} - E_{a,(n,n)}^{(0)}} \right] \\ &= \sum_x |\Psi_{a,i}|^2 \frac{2n(n+1)}{U}. \end{aligned} \quad (\text{E12})$$

As the Landau formalism, we can give $E = \sum_a^2 \sum_i \left[\text{constant} + R |\Psi_{a,i}|^2 + W |\Psi_{a,i}|^4 + \dots \right]$. From $\frac{E_{a,(n,n)}^{(0)}}{2}$ and $E_{a,(n,n)}^{(2)}$, the coefficient R is

$$R = \frac{1}{2dt_1 + t_2} - \frac{2n(n+1)}{U}. \quad (\text{E13})$$

$R = 0$ is the critical point. At $R > 0$, this system is under the Mott insulator phase, while at $R < 0$, the condensation occurs. In the coefficient R , the coupling coefficient t_2 plays a similar role as t_1 . It is worth mentioning that if we set $t_1 = 0$, $t_2 > 0$ and set $\hat{d}_j^i = \hat{b}_{a,j}^\dagger \hat{b}_{b,j}$ where $b \neq a$, we can still obtain similar results with mean-field theory. This implies condensates with a variety of particle-hole bound states in this system as long as $t_2 > 0$. This phase is called multi-dipole condensate phase.

As t_1, t_2 increases, the bosons firstly form particle-hole bound states, and these bound states condense firstly. Here, a hole of a species of bosons means that the number of the species of bosons in a lattice is lower than the average number. As $(2dt_1 + t_2)$ tends to infinity, single bosons condense, that is, the HFS phase.

Appendix F: Derivation of Lattice Model A in a tilted lattice

In a microscopic Hamiltonian with two species of bosons in a tilted optical lattice chain, a U' term Hubbard term between different species is included:

$$\begin{aligned} H_{\text{tilted}} &= -t \sum_{a,i} \left(\hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} + \text{h.c.} \right) - \mu \sum_{a,i} \hat{n}_{a,i} + \frac{U}{2} \sum_{a,i} \hat{n}_{a,i} (\hat{n}_{a,i} - 1) + U' \sum_i \hat{n}_{1,i} \hat{n}_{2,i} + V \sum_{a,i} \mathbf{i} \hat{n}_{a,i} \\ &\equiv H_t + H_U + H_V. \end{aligned} \quad (\text{F1})$$

In the limit $V \gg t, U, U'$, we use a Nakajima transformation $e^S H e^{-S} = H + [S, H] + \frac{1}{2}[S, [S, H]] + \dots$, where S is anti-Hermitian. We take $S = \sum_{n=1}^{\infty} S_n$, where S_n is order n in $t/V, U/V$, and U'/V . In this derivation, we need a effective Hamiltonian to third order, and thus we can set S_n to 0 for $n \geq 4$. Then, the effective Hamiltonian becomes

$$\begin{aligned} H_{\text{eff}} &= H_V + H_t + H_U + [S_1, H_V] + [S_1, H_U] + [S_1, H_t] + [S_2, H_V] + [S_2, H_U] + [S_2, H_t] + [S_3, H_V] \\ &+ \frac{1}{2}[S_1, [S_1, H_V]] + \frac{1}{2}[S_1, [S_1, H_U]] + \frac{1}{2}[S_1, [S_1, H_t]] + \frac{1}{2}[S_1, [S_2, H_V]] + \frac{1}{2}[S_2, [S_1, H_V]] + \frac{1}{6}[S_1, [S_1, [S_1, H_V]]] \end{aligned} \quad (\text{F2})$$

Firstly, we need $[S_1, H_V] + H_t = 0$, and we can get: $S_1 = \frac{t}{V} \sum_{a,i} \left(\hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} - \text{h.c.} \right)$, and we have

$$[S_1, H_V] = -H_t, \quad [S_1, H_t] = 0 \quad (\text{F3})$$

$$[S_1, H_U] = \frac{t_0 U}{2V} \sum_{a,i} \{ \hat{n}_{a,i}, (\hat{b}_{a,i-1}^\dagger \hat{b}_{a,i} - \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} + \text{h.c.}) \} + \frac{t_0 U'}{V} \sum_{a \neq b, i} (\hat{b}_{a,i-1}^\dagger \hat{b}_{a,i} - \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} + \text{h.c.}) \hat{n}_{b,i}. \quad (\text{F4})$$

Plugging Eq. (F3) into the effective Hamiltonian, we obtain

$$H_{\text{eff}} = H_V + H_U + [S_1, H_U] + [S_2, H_V] + [S_2, H_U] + \frac{1}{2}[S_2, H_t] + [S_3, H_V] + \frac{1}{2}[S_1, [S_1, H_U]] + \frac{1}{2}[S_1, [S_2, H_V]]. \quad (\text{F5})$$

Then we require S_2 to satisfy $[S_2, H_V] + [S_1, H_U] = 0$. S_2 is given by:

$$S_2 = -\frac{t_0 U}{2V^2} \sum_{a,i} \{ \hat{n}_{a,i}, (\hat{b}_{a,i-1}^\dagger \hat{b}_{a,i} - \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} - \text{h.c.}) \} - \frac{t_0 U'}{V^2} \sum_{a \neq b, i} (\hat{b}_{a,i-1}^\dagger \hat{b}_{a,i} - \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} - \text{h.c.}) \hat{n}_{b,i}. \quad (\text{F6})$$

And thus, the effective Hamiltonian becomes $H_{\text{eff}} = H_V + H_U + [S_2, H_U] + \frac{1}{2}[S_2, H_t] + [S_3, H_V]$. To third order, we need $[S_3, H_V]$ to cancel the terms in $[S_2, H_U] + \frac{1}{2}[S_2, H_t]$ which do not conserve the total dipole moment. Here we define a projection operator P which projects onto total dipole-conserving terms, and we have

$$\begin{aligned} [S_2, H_U] + \frac{1}{2}[S_2, H_t] + [S_3, H_V] &= P([S_2, H_U] + \frac{1}{2}[S_2, H_t])P = \frac{1}{2}P[S_2, H_t]P \\ &= \sum_{a,i} \left(-\frac{2t_0^2 U}{V^2} \hat{n}_{a,i}^2 + \frac{4t_0^2 U}{V^2} \hat{n}_{a,i} \hat{n}_{a,i+1} \right) + \left(-\frac{t^2 U}{V^2} \hat{b}_{a,i} (\hat{b}_{a,i+1}^\dagger)^2 \hat{b}_{a,i+2} + \text{h.c.} \right) \\ &+ \sum_{a \neq b, i} \left(-\frac{2t_0^2 U'}{V^2} \hat{n}_{a,i} \hat{n}_{b,i} + \frac{2t_0^2 U'}{V^2} \hat{n}_{a,i} \hat{n}_{b,i+1} \right) + \left(\frac{t^2 U'}{V^2} \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} \hat{b}_{b,i+1} \hat{b}_{b,i+2}^\dagger - \frac{t^2 U'}{V^2} \hat{b}_{a,i}^\dagger \hat{b}_{a,i+1} \hat{b}_{b,i} \hat{b}_{b,i+1}^\dagger + \text{h.c.} \right) \end{aligned} \quad (\text{F7})$$

At last, the effective Hamiltonian is

$$\begin{aligned}
H_{\text{eff}} = & \sum_{a,\mathbf{i}} \left(-\frac{t^2 U}{V^2} \hat{b}_{a,\mathbf{i}} (\hat{b}_{a,\mathbf{i}+1}^\dagger)^2 \hat{b}_{a,\mathbf{i}+2} + \text{h.c.} \right) + \sum_{a \neq b, \mathbf{i}} \left(\frac{t^2 U'}{V^2} \hat{b}_{a,\mathbf{i}}^\dagger \hat{b}_{a,\mathbf{i}+1} \hat{b}_{b,\mathbf{i}+1} \hat{b}_{b,\mathbf{i}+2}^\dagger - \frac{t^2 U'}{V^2} \hat{b}_{a,\mathbf{i}}^\dagger \hat{b}_{a,\mathbf{i}+1} \hat{b}_{b,\mathbf{i}} \hat{b}_{b,\mathbf{i}+1}^\dagger + \text{h.c.} \right) \\
& + \sum_{a,\mathbf{i}} \left(-\mu - \frac{U}{2} + V\mathbf{i} \right) \hat{n}_{a,\mathbf{i}} + \sum_{a,\mathbf{i}} \left(\frac{U}{2} - \frac{2t_0^2 U}{V^2} \right) \hat{n}_{a,\mathbf{i}}^2 + \frac{4t_0^2 U}{V^2} \hat{n}_{a,\mathbf{i}} \hat{n}_{a,\mathbf{i}+1} + \sum_{a \neq b, \mathbf{i}} \left(\left(\frac{U'}{2} - \frac{2t_0^2 U'}{V^2} \right) \hat{n}_{a,\mathbf{i}} \hat{n}_{b,\mathbf{i}} + \frac{2t_0^2 U'}{V^2} \hat{n}_{a,\mathbf{i}} \hat{n}_{b,\mathbf{i}+1} \right)
\end{aligned} \tag{F8}$$

After the above derivation, we obtain effective hopping terms with the conservation of the total dipole moments on the tilted optical lattice.