

**$k$ -LOOSE ELEMENTS AND  $k$ -PAVING MATROIDS**

JAGDEEP SINGH

ABSTRACT. For a matroid of rank  $r$  and a non-negative integer  $k$ , an element is called  $k$ -loose if all circuits containing it have size greater than  $r - k$ . Zaslavsky and the author previously characterized all binary matroids with a 1-loose element. In this paper, we establish a linear bound on the size of a binary matroid, in terms of its rank, that contains a  $k$ -loose element. A matroid is  $k$ -paving if all its elements are  $k$ -loose. Rajpal showed that for a prime power  $q$ , the rank of a  $GF(q)$  matroid that is  $k$ -paving is bounded. We provide a bound on the rank of  $GF(q)$ -matroids that are cosimple and have two  $k$ -loose elements. Consequently, we deduce a bound on the rank of  $GF(q)$  matroids that are  $k$ -paving. Additionally, we provide a bound on the size of binary matroids that are  $k$ -paving.

## 1. INTRODUCTION

The notation and terminology follow [2] unless stated otherwise. All graphs and matroids considered here are finite and simple. For a matroid of rank  $r$  and a non-negative integer  $k$ , an element  $t$  of  $M$  is called  **$k$ -loose** if every circuit of  $M$  that contains  $t$  has size greater than  $r - k$ . A matroid  $M$  is called  **$k$ -paving** if every element of  $M$  is  $k$ -loose. Note that a matroid  $M$  is paving if and only if  $M$  is 1-paving. Acketa [1] provided a characterization of all 1-paving matroids that are binary, while Oxley [3] determined all ternary 1-paving matroids. Zaslavsky and the author previously characterized all binary matroids with a 1-loose element [6]. In Section 2, we show that the size of a binary matroid that contains a  $k$ -loose element is linear in terms of its rank. In particular, we prove the following.

**Theorem 1.1.** *For  $r \geq 5$  and  $0 \leq k \leq \frac{r-2}{3}$ , let  $M$  be a simple binary matroid of rank  $r$  with no coloops and a  $k$ -loose element. Then  $|E(M)| \leq (2^k - 1)(r + 1 - k) + (r + 1)$ .*

We say a matroid  $M$  is a  $GF(q)$ -matroid if  $M$  is representable over the field of  $q$  elements. Section 3 considers  $GF(q)$ -matroids that have two  $k$ -loose elements. We show that the rank of such a matroid is bounded unless the two  $k$ -loose elements form a cocircuit of size two. Following is the precise statement.

---

*Date:* December 16, 2024.

*2020 Mathematics Subject Classification.* 05B35.

*Key words and phrases.* binary matroid, paving matroid, loose elements.

**Theorem 1.2.** *Let  $M$  be a simple  $GF(q)$ -matroid with no coloops that has two  $k$ -loose elements  $e$  and  $f$ . Then  $r(M) \leq (q + 1)(k - 1) + 2q$ , or  $\{e, f\}$  is a cocircuit of  $M$ .*

An immediate consequence of Theorem 1.2 is the following.

**Corollary 1.3.** *Let  $M$  be a simple  $GF(q)$ -matroid with no coloops that is  $k$ -paving. Then  $M$  is a circuit or  $r(M) \leq (q + 1)(k - 1) + 2q$ .*

Rajpal [5, Proposition 8] demonstrated the existence of a rank bound for paving matroids representable over  $GF(q)$ . Corollary 1.3 provides such a bound. Observe that paving matroids are precisely 1-paving matroids. Therefore by Corollary 1.3, a paving matroid representable over  $GF(q)$  has rank at most  $2q$ , unless it is a circuit.

It is immediate from Corollary 1.3 that a binary matroid that is  $k$ -paving has rank at most  $3k + 1$ . In Section 4, we provide a bound on the size of binary  $k$ -paving matroids. In particular, we prove the following.

**Theorem 1.4.** *Let  $M$  be a simple binary  $k$ -paving matroid of rank  $r$ , where  $M$  is not a circuit and  $r \geq k + 4$ , and let  $t = \frac{3k+1-r}{2}$ . Then*

$$|E(M)| \leq (r + 1) + \sum_{i=0}^t 2 \binom{k}{i}.$$

Note that every simple binary matroid that has rank at most  $k + 2$  is  $k$ -paving, and every affine binary matroid of rank  $k + 3$  is  $k$ -paving. We apply Theorem 1.4 to 3-paving binary matroids and obtain the following.

**Corollary 1.5.** *Let  $M$  be a simple binary 3-paving matroid of rank at least seven. Then  $r(M) \leq 10$  and  $|E(M)| \leq 17$ . In particular, the size of  $M$  is at most 16, 17, 12, and 13 when the rank of  $M$  is 7, 8, 9, and 10 respectively.*

## 2. $k$ -LOOSE ELEMENTS IN BINARY MATROIDS

Theorem 1.1 gives a linear bound on the size of a binary matroid containing a  $k$ -loose element, in terms of its rank. Specifically, if a binary matroid  $M$  has a 1-loose element, then the size of  $M$  is at most  $2r + 1$ . If  $M$  has a 2-loose element, then the size of  $M$  does not exceed  $4r - 2$ .

*Proof of Theorem 1.1.* Let  $e$  be a  $k$ -loose element of  $M$ . Note that if  $k = 0$ , then by [4, Lemma 3.1],  $M$  is a circuit and the result follows. Therefore  $k \geq 1$ . Suppose that  $e$  is not  $(k - 1)$ -loose. It follows that there is a basis  $B = \{e_1, \dots, e_r\}$  of  $M$  such that, in the standard binary representation  $P = [I_r | Q]$  of  $M$  with respect to  $B$ , the column  $e^P$  labeled by  $e$  has its top  $k$  entries equal to zero and all other entries non-zero. We call the set of indices of the non-zero entries of a column  $f^P$  in  $P$  the *support*,  $\text{supp}(f^P)$ , of  $f^P$ . The entries of  $f^P$  excluding the top  $k$  entries are called the *root entries* of  $f^P$  and the set of indices of the non-zero root entries of  $f^P$  is the *root support* of  $f^P$ . Let  $Q - e^P$  denote the columns in  $Q$  excluding  $e^P$ .

**2.0.1.** *Let  $f^P$  be a column in  $Q - e^P$ . Then the sum of the number of zeroes in the top  $k$  entries of  $f^P$  and the number of non-zero root entries of  $f^P$  is at most  $k + 1$ .*

Suppose otherwise. Then  $e^P + f^P$  has at least  $k + 2$  zeroes, so at most  $r - (k + 2)$  columns from  $I_r$  can be added to  $e^P + f^P$  to obtain a column with all entries equal to zero. It follows that  $e$  is in a circuit of size at most  $r - k$ , a contradiction. Therefore 2.0.1 holds.

Let  $F$  be a set of all columns in  $Q - e^P$  such that, for any pair  $f^P, g^P$  in  $F$ , the sum  $f^P + g^P$  has its top  $k$  entries equal to zero. It is clear that the top  $k$  entries of the columns in  $F$  are identical.

**2.0.2.** *Let  $f^P, g^P$  be columns in  $F$ . Then, the symmetric difference of their supports,  $\text{supp}(f^P) \Delta \text{supp}(g^P)$ , contains at most two elements.*

Suppose not. Then  $f^P + g^P$  has its top  $k$  entries equal to zero and contains at least three non-zero entries. It follows that  $e^P + f^P + g^P$  has at least  $k + 3$  zeroes so it can be combined linearly over  $\text{GF}(2)$  with at most  $r - (k + 3)$  columns in  $I_r$  to obtain the column that has all the entries zero. Therefore  $e$  is in a circuit of size at most  $r - k$ , a contradiction. Thus 2.0.2 holds.

**2.0.3.**  $|F| \leq (r + 1 - k)$ .

Choose a column  $f^P$  in  $F$  such that the root support of  $f^P$ , say  $S$ , has the largest possible size. Suppose  $|S| = s$ . It follows by 2.0.1 that  $s \leq k + 1$ . In view of 2.0.2, a column  $g^P$  in  $F - f^P$  can fall into one of the following three categories, namely,  $\text{supp}(g^P) \subseteq \text{supp}(f^P)$  with  $|\text{supp}(f^P) - \text{supp}(g^P)| = 1$ ,  $\text{supp}(g^P) \subseteq \text{supp}(f^P)$  with  $|\text{supp}(f^P) - \text{supp}(g^P)| = 2$ , or  $\text{supp}(g^P) \not\subseteq \text{supp}(f^P)$ . We call these columns type  $A$ , type  $B$ , and type  $C$  respectively. By 2.0.2, it follows that the number of type  $B$  columns is at most  $\max\{3, s - 1\}$ . Note that if  $s \leq 1$ , there is no type  $B$  column. If  $s = 2$ , there is at most one type  $B$  column. It follows that the number of type  $B$  columns does not exceed  $s$ .

First, suppose that  $F$  contains a type  $C$  column. By 2.0.2, a type  $C$  column has precisely one non-zero root entry outside  $S$  and exactly one zero entry in  $S$ . If  $F$  contains two type  $C$  columns such that their entries in  $S$  differ, then by 2.0.2, all their entries outside  $S$  are identical. In this case, the number of type  $C$  columns is at most  $s$ . On the other hand, if all type  $C$  columns in  $F$  have identical entries in  $S$ , the number of type  $C$  columns is at most  $r - (s + k)$ . Therefore, the number of columns of type  $C$  in  $F$  is at most  $\max\{s, r - (s + k)\}$ . Given that  $s \leq k + 1$  and  $k \leq \frac{r-2}{3}$ , it follows that the number of type  $C$  columns in  $F$  is at most  $r - (s + k)$ . Since  $F$  contains a type  $C$  column, it is immediate from 2.0.2 that there is at most one type  $A$  column in  $F$ . Furthermore, if a type  $A$  column exists, then we have at most  $s - 1$  type  $B$  columns. It follows that the sum of the number of type  $A$  columns and the number of type  $B$  columns does not exceed  $s$ . Therefore the size of  $F$  is at most  $r - (s + k) + s + 1 = r + 1 - k$ .

Now, suppose that no type  $C$  column exists. By 2.0.2, it follows that if a type  $B$  column exists, then the number of type  $A$  columns is at most 2 so  $F$  has size at most  $s + 3$ . Since  $s \leq k + 1, r \geq 5$ , and  $k \leq \frac{r-2}{3}$ , it follows that  $s + 3 \leq r + 1 - k$ . Finally note that the number of type  $A$  columns is at most  $s$  so 2.0.3 holds.

Observe that if there is a column  $h^P$  in  $Q - e^P$  that has its top  $k$  entries equal to zero, then by 2.0.1, the column  $h^P$  has at most one non-zero root entry so  $h^P$  has at most one non-zero entry. Since  $M$  is simple, this is a contradiction. It follows that there are at most  $2^k - 1$  sets  $F$  in  $Q - e^P$ . Therefore  $|E(M)| \leq (2^k - 1)(r + 1 - k) + (r + 1)$ .

If  $e$  is  $(k - 1)$ -loose, we choose the smallest non-negative integer  $t$  such that  $e$  is  $t$ -loose. By the same argument we obtain  $|E(M)| \leq (2^t - 1)(r + 1 - t) + (r + 1)$ . Since  $t \leq k - 1, r \geq 5$ , and  $k \leq \frac{r-2}{3}$ , it follows that  $(2^t - 1)(r + 1 - t) + (r + 1) \leq (2^k - 1)(r + 1 - k) + (r + 1)$ .  $\square$

### 3. $k$ -LOOSE ELEMENTS IN $GF(q)$ -MATROIDS

Zaslavsky and the author in [6] showed that a ternary matroid with no coloops that contains a 1-loose element has linear size relative to its rank. In particular they proved the following.

**Theorem 3.1.** *For  $r \geq 5$ , let  $M$  be a simple ternary matroid of rank  $r$  with no coloops and a 1-loose element. Then  $|E(M)| \leq \lfloor \frac{41r-101}{2} \rfloor$  if  $r > 10$ . If  $r \leq 10$ , then  $|E(M)| \leq \lfloor \frac{35r-35}{2} \rfloor$ .*

Furthermore, they proved that if a ternary matroid with no coloops and no cocircuits of size two contains two 1-loose elements, then the rank of the matroid is bounded. This result was shown to hold for all  $GF(q)$ -matroids.

**Theorem 3.2.** *Let  $M$  be a simple  $GF(q)$ -matroid with no coloops that has two 1-loose elements  $e$  and  $f$ . Then  $r(M) \leq 2q$ , or  $\{e, f\}$  is a cocircuit of  $M$ .*

We extend Theorem 3.2 to  $k$ -loose elements and prove Theorem 1.2. Following is the restatement of Theorem 1.2.

**Theorem.** *Let  $M$  be a simple  $GF(q)$ -matroid with no coloops that has two  $k$ -loose elements  $e$  and  $f$ . Then  $r(M) \leq (q + 1)(k - 1) + 2q$ , or  $\{e, f\}$  is a cocircuit of  $M$ .*

*Proof.* We may assume that  $\{e, f\}$  is not a cocircuit of  $M$  else the result follows directly. Since  $\{e, f\}$  is not a cocircuit, there is basis  $B$  of  $M$  that is disjoint from  $\{e, f\}$ . Note that, in the standard  $GF(q)$ -representation  $P = [I_r | Q]$  of  $M$  with respect to  $B$ , the column  $e^P$  corresponding to  $e$  and the column  $f^P$  corresponding to  $f$  have at most  $k$  zeroes. By row and column scaling, we may assume that  $e^P$  consists entirely of ones except possibly at the top  $k$  entries.

Let  $f^{P'}$  denote the column obtained by removing the top  $k$  entries of  $f^P$ . Suppose  $f^{P'}$  has  $k + 2$  non-zero entries that are all equal. Then  $f^P, e^P$ , and

at most  $r - (k + 2)$  columns from  $I_r$  can be combined linearly over  $GF(q)$  to obtain a column that has all its entries equal to zero. This would create a circuit of  $M$  that contains  $\{e, f\}$  and has size at most  $r - k$ , a contradiction. Therefore,  $f^{P'}$  has at most  $k + 1$  non-zero entries that are equal. Since the field  $GF(q)$  has  $q - 1$  non-zero elements, it follows that  $f^{P'}$  has at most  $(k + 1)(q - 1)$  non-zero entries. Additionally,  $f^{P'}$  has at most  $k$  zeroes so it has at most  $kq + q - 1$  entries. Therefore,  $f^P$  has at most  $kq + q + k - 1$  entries so the rank of  $M$  is at most  $(q + 1)(k - 1) + 2q$ .  $\square$

Following is an immediate consequence.

**Corollary 3.3.** *Let  $M$  be a simple and a cosimple  $GF(q)$ -matroid that has two  $k$ -loose elements. Then  $r(M) \leq (q + 1)(k - 1) + 2q$ .*

The following corollary of Theorem 1.2 determines the bound whose existence was established by Rajpal [5, Proposition 8].

**Corollary 3.4.** *Let  $M$  be a simple  $GF(q)$ -matroid with no coloops that is  $k$ -paving. Then  $M$  is a circuit, or  $r(M) \leq (q + 1)(k - 1) + 2q$ .*

*Proof.* Note that every element of  $M$  is  $k$ -loose. If  $r(M) > (q + 1)(k - 1) + 2q$ , it follows from Theorem 1.2 that every pair of elements of  $M$  is in a cocircuit of size two. It follows that  $M$  is a circuit.  $\square$

Following special cases of Corollary 3.4 are worth noting. The binary case was shown by Rajpal [5, Proposition 6].

**Corollary 3.5.** *Let  $M$  be a simple binary matroid with no coloops that is  $k$ -paving. Then  $M$  is a circuit, or  $r(M) \leq 3k + 1$ .*

**Corollary 3.6.** *Let  $M$  be a simple ternary matroid with no coloops that is  $k$ -paving. Then  $M$  is a circuit, or  $r(M) \leq 4k + 2$ .*

#### 4. $k$ -PAVING BINARY MATROIDS

It follows by Theorem 1.1 and Corollary 3.5 that the size of a binary  $k$ -paving matroid, excluding circuits, is at most  $(2^k - 1)(2k + 2) + (3k + 2)$ . However the bound obtained this way is crude as it represents the maximum size of a binary matroid with a  $k$ -loose element and rank at most  $3k + 1$ . Theorem 1.4 obtains a sharper bound. Following is its restatement.

**Theorem.** *Let  $M$  be a simple binary  $k$ -paving matroid of rank  $r$ , where  $M$  is not a circuit and  $r \geq k + 4$ , and let  $t = \frac{3k+1-r}{2}$ . Then*

$$|E(M)| \leq (r + 1) + \sum_{i=0}^t 2 \binom{k}{i}.$$

*Proof.* Suppose that  $M$  has a circuit  $C$  of size  $r - k + 1$ . For an element  $e$  of  $C$ , let  $B$  be a basis of  $M$  containing  $C - e$ . Consider the standard binary representation  $P = [I_r | Q]$  of  $M$  with respect to the basis  $B$ .

**4.0.1.** *Let  $h$  be the sum of  $m$  columns in  $Q$ . Then  $h$  has at most  $k + (m - 1)$  zeroes.*

Assume that  $h$  has at least  $k + m$  entries that are zero. Then, at most  $r - (k + m)$  columns in  $I_r$  can be added to  $h$  to obtain a column with all entries equal to zero. This would result in a circuit of size at most  $r - k$  in  $M$ , a contradiction. Therefore 4.0.1 holds.

Note that the column  $e^P$  in  $Q$  corresponding to the element  $e$  of  $M$  has exactly  $k$  zeroes. By row and column scaling, we may assume that the top  $k$  entries of  $e^P$  are zero. We denote the set of indices of the zero entries of a column  $f^P$  in  $P$  by  $\text{zeroes}(f^P)$ . The entries of  $f^P$  excluding the top  $k$  entries are called the *root entries* of  $f^P$ . Let  $g^P$  be a column in  $Q - e^P$  such that  $|\text{zeroes}(e^P) \cap \text{zeroes}(g^P)| = l$  is maximum.

**4.0.2.**  $r \leq 3k + 1 - 2l$ .

By 4.0.1,  $g^P$  has at most  $k - l$  root entries that are zero and at most  $k + 1 - l$  root entries that are one so  $g^P$  has at most  $k + (k - l) + (k + 1 - l) = 3k + 1 - 2l$  entries. Therefore 4.0.2 holds.

Let  $F$  be a set of all columns in  $Q - e^P$  such that, for any pair  $f^P, g^P$  in  $F$ , the sum  $f^P + g^P$  has its top  $k$  entries equal to zero. Evidently, the top  $k$  entries of the columns in  $F$  are identical.

**4.0.3.**  $|F| \leq 2$ .

Suppose not, and let  $a, b$ , and  $c$  be three columns in  $F$ . Note that, by 4.0.1,  $a$  and  $b$  have at most one root entry that is equal. Since  $r \geq k + 4$ , we have at least three root entries of  $a$ , say at indices  $j, k$ , and  $l$ , such that  $b$  has different entries than  $a$  at each of  $j, k$ , and  $l$ . Note that the  $j^{\text{th}}, k^{\text{th}}$  and  $l^{\text{th}}$  entries of  $c$  are either the same as those of  $a$  or  $b$ . This means that at least two root entries of  $c$  are the same as those of one of  $\{a, b\}$ . This is a contradiction to 4.0.1 so 4.0.3 holds.

**4.0.4.**  $|E(M)| \leq (r + 1) + \sum_{i=0}^l 2^{\binom{k}{i}}$ .

Observe that each column in  $Q - e^P$  has at most  $l$  zero entries among the top  $k$  entries. It follows that there are at most  $\sum_{i=0}^l \binom{k}{i}$  sets  $F$  in  $Q - e^P$ . Therefore, by 4.0.3,  $|E(M)| \leq (r + 1) + \sum_{i=0}^l 2^{\binom{k}{i}}$  so 4.0.4 holds.

By 4.0.2,  $l \leq \frac{3k+1-r}{2} = t$  so by 4.0.4, the result follows.

If  $M$  has no circuit of size  $r - k + 1$ , let  $w$  denote the smallest number such that  $M$  has a circuit of size  $r - w + 1$ . Since all circuits of  $M$  have size greater than  $r - k$ , it is clear that  $w < k$ . By the same argument, it follows that  $|E(M)| \leq (r + 1) + \sum_{i=0}^s 2^{\binom{w}{i}}$ , where  $s = \frac{3w+1-r}{2}$ . Since  $w < k$ , it follows that  $s \leq t$  and  $|E(M)| \leq (r + 1) + \sum_{i=0}^s 2^{\binom{w}{i}} \leq (r + 1) + \sum_{i=0}^t 2^{\binom{k}{i}}$ .  $\square$

We apply Theorem 1.4 to 3-paving binary matroids and prove Corollary 1.5.

*Proof of Corollary 1.5.* It is clear from Corollary 3.5 that  $r(M) \leq 10$ . Note that if  $r(M)$  is 7, then  $t$  equals 1 so by Theorem 1.4,  $|E(M)|$  is at most 16. Similarly, if  $r(M) = 8, 9$ , and 10, then  $|E(M)|$  is at most 17, 12, and 13 respectively.  $\square$

## REFERENCES

- [1] D.M. Acketa, On Binary paving matroids, *Discrete Math.* **70** (1988), 109–110.
- [2] J. Oxley *Matroid Theory*, Second edition, Oxford University Press, New York, 1992.
- [3] J. Oxley, Ternary paving matroids, *Discrete Math.* **91** (1991), 77–86.
- [4] J. Oxley and J. Singh, The smallest class of binary and ternary matroids closed under direct sums and complements, *SIAM J. Discrete Math.* **36** (2022), 2051–2072.
- [5] S. Rajpal, On binary  $k$ -paving matroids and Reed-Muller codes, *Discrete Math.* **190** (1998), 191–200.
- [6] J. Singh and T. Zaslavsky, Loose elements in binary and ternary matroids, in preparation.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MISSISSIPPI STATE UNIVERSITY,  
STARKVILLE, MISSISSIPPI

*Email address:* `singhjagdeep070@gmail.com`