PARTIAL REGULARITY AND HIGHER INTEGRABILITY FOR *A*-QUASICONVEX VARIATIONAL PROBLEMS

ZHOULIN LI AND BOGDAN RAIȚĂ

Dedicated to Jan Kristensen on the occasion of his 60th birthday

ABSTRACT. We prove that minimizers of variational problems

ı

n

ninimize
$$\mathcal{E}(v) = \int_{\Omega} f(x, v(x)) \, \mathrm{d}x$$
 for $\mathscr{A}v = 0$,

are partially continuous provided that the integrands f are strongly \mathscr{A} -quasiconvex in a suitable sense. We consider p-growth problems with 1 , linear pde operators $<math>\mathscr{A}$ of constant rank, and Dirichlet boundary conditions, in the sense that admissible fields are of the form $v = v_0 + \varphi$, with \mathscr{A} -free $\varphi \in C_c^{\infty}(\Omega)$. Our analysis also covers the "potentials case"

inimize
$$\mathcal{F}(u) = \int_{\Omega} f(x, \mathscr{B}u(x)) \, \mathrm{d}x$$
 for $u \in C_c^{\infty}(\Omega)$,

where \mathscr{B} is a different linear pde operator of constant rank. We also prove appropriate higher integrability of minimizers for both types of problems.

CONTENTS

1.	Introduction	1
2.	Preliminaries	6
3.	Coercivity and lower semicontinuity	10
4.	Existence of generalized minimizers	12
5.	Local linear estimates	15
6.	Higher integrability	19
7.	Partial regularity	23
References		31

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be an open bounded set and $1 \leq p < \infty$. Consider the problem to

$$(P\mathscr{A}_{\Omega}) \qquad \text{minimize} \quad \mathcal{E}(v,\Omega) = \int_{\Omega} f(x,v(x)) \, \mathrm{d}x \quad \text{for } v \in v_0 + C^{\infty}_{c,\mathscr{A}}(\Omega)$$

where $v_0 \in L^p(\Omega, V)$ is \mathscr{A} -free and $C^{\infty}_{c,\mathscr{A}}(\Omega) = \{\varphi \in C^{\infty}_c(\Omega, V) \colon \mathscr{A}\varphi = 0\}.$

Here \mathscr{A} is a homogeneous vectorial differential operator with constant coefficients, defined on \mathbb{R}^n from V to W, which are both finite dimensional inner product spaces. We denote its characteristic polynomial by $\mathscr{A}(\xi) \in \operatorname{Lin}(V, W)$ for $\xi \in \mathbb{R}^n$. For the precise definitions, see Section 2.1

Functionals of this type have been studied extensively over the last several decades, first in the prototypical case $\mathscr{A} = \operatorname{curl}$ (which leads to functionals defined on Sobolev spaces) and later in the case of general differential constraints, emanating from the study of

²⁰²⁰ Mathematics Subject Classification. Primary: 49N60, Secondary: 35B65, 49J45, 28B05, 35E20.

Key words and phrases. Regularity theory, *A*-Quasiconvexity, Strong quasiconvexity, Constant rank operators, Partial continuity, Partial regularity, Systems of linear pde, Degenerate variational problems.

Z. LI AND B. RAIȚĂ

compensated compactness [Tar78; Mur81]. The lower semicontinuity of these functionals with respect to weak convergence is well understood both in the curl-free case [Mor52; Dac82; AF84; FM93; AD92; KP94; Kri99; FMP98; KR10] as well as in the general \mathscr{A} free case [Mur81; Dac06; FM99; FLM04; BDS15; ADR20; Arr21; GR22; KR22]. The overarching theme in these works is that the lower semicontinuity of functionals in $(P\mathscr{B}_{\Omega})$ is equivalent to the so-called \mathscr{A} -quasiconvexity condition introduced in [FM99].

All these results are subject to the so called **constant rank** condition on \mathscr{A} , i.e.

(CR)
$$\operatorname{rank} \mathscr{A}(\xi) = \operatorname{const.} \text{ for } \xi \in \mathbb{R}^n \setminus \{0\}$$

Except for notable examples [Mül99; LMM11; SW12], the lower semicontinuity problem is widely open in the absence of the constant rank condition. We will make the same assumption, together with the assumption that the wave cone of \mathscr{A} [Tar78; Mur81] is spanning, i.e.

(SC)
$$\operatorname{span} \bigcup_{\xi \in \mathbb{R}^n \setminus \{0\}} \ker \mathscr{A}(\xi) = V.$$

This second assumption is mild (see Section 4.1 in [KR22]).

Under these assumptions, the regularity of minimizers is far from fully understood. Our aim is to reach the high benchmark established in the case $\mathscr{A} = \text{curl}$ by works such as [Eva86; AF87; AF89; GM86; Mar89; CFM98; GK19b]. These works use stronger quasiconvexity conditions to infer partial regularity results, which is typical in the vectorial calculus of variations even for strictly convex problems. Strong quasiconvexity also leads to higher integrability of minimizers above the growth parameter, see [GG82] for a classic reference. For more details see the comprehensive exposition in [Min06].

On the other hand, there are only a handful of contributions to regularity for the general Problem $(P\mathscr{A}_{\Omega})$ [Fra19; GK19a; Gme20; Gme21; CG22; BK22; Sch25], all of which impose stringent restrictions on the setup.

Our main result establishes partial continuity and higher integrability of minimizers of $(P\mathscr{A}_{\Omega})$ under a suitable assumption of strong quasiconvexity on the integrand f:

Theorem A. Let \mathscr{A} be a differential operator satisfying the (CR) and (SC) conditions and let $1 . Suppose that <math>f: \Omega \times V \to \mathbb{R}$ satisfies the growth, smoothness, and strong \mathscr{A} -quasiconvexity assumptions (H1), (H2), and (H3) below. Then for any generalized minimizer v of $(P\mathscr{A}_{\Omega})$, there exists an open subset $\Omega' \subset \Omega$ and a constant $\sigma_0 > 0$ such that $\mathscr{L}^n(\Omega \setminus \Omega') = 0$ and $v \in C^{0,\alpha}(\Omega') \cap L^{p+\sigma}_{loc}(\Omega)$ for any $\alpha \in (0,1)$ and $\sigma \in (0,\sigma_0)$.

Here \mathscr{A} is assumed to have homogeneous entries in the derivatives for simplicity of exposition, but the same result holds for differential operators with homogeneous rows (see [Rai24] and Remark 2.3). The case p = 1 can be addressed by a non-trivial upgrade of our method, which will make the object of our future work.

Most of the earlier results concerning general differential operators that we mentioned pertain to the related problem on potential operator level to

$$(P\mathscr{B}_{\Omega}) \qquad \text{minimize} \quad \mathcal{F}(u,\Omega) = \int_{\Omega} f(x,\mathscr{B}u(x)) \,\mathrm{d}x \quad \text{for } u \in u_0 + C_c^{\infty}(\Omega,U),$$

where $u_0 \in W^{k-1}(\Omega, U)$ is such that $\mathscr{B}u \in L^p(\Omega, V)$; here k is the order of \mathscr{B} .

We will show that establishing partial continuity of minimizers of (\mathcal{PA}_{Ω}) can be reduced to proving partial continuity of minimizers $\mathscr{B}u$ of (\mathcal{PB}_{Ω}) (see Remark 4.4). Here \mathscr{A} and \mathscr{B} satisfy a homological relation described in Section 2.1 (see also [Rai19b; Rai24]). The reduction of (\mathcal{PA}_{Ω}) to (\mathcal{PB}_{Ω}) in the context of regularity theory is likely new.

Earlier results pertaining to partial regularity of minimizers of (\mathcal{PB}_{Ω}) in [CG22; Fra19] concern *elliptic* operators \mathcal{B} of first order. Ellipticity is a restrictive assumption which allows the use of strong harmonic analysis estimates. We prove partial continuity and higher integrability of minimizers $\mathcal{B}u$ of (\mathcal{PB}_{Ω}) for arbitrary constant rank operators:

Theorem B. Let \mathscr{B} be a differential operator satisfying (CR) and let 1 . Suppose $that <math>f: \Omega \times V \to \mathbb{R}$ satisfies the growth, smoothness, and strong quasiconvexity assumptions (H1), (H2), and (H3)' below. Then for any generalized minimizer $\mathscr{B}u$ of $(\mathcal{P}\mathscr{B}_{\Omega})$, there exists an open subset $\Omega' \subset \Omega$ satisfying $\mathscr{L}^n(\Omega \setminus \Omega') = 0$ and a constant $\sigma_0 > 0$ such that $\mathscr{B}u \in C^{0,\alpha}(\Omega') \cap L^{p+\sigma}_{\text{loc}}(\Omega)$ for any $\alpha \in (0,1), \sigma \in (0,\sigma_0)$. Moreover, for each $\omega \in \Omega$, there is a representative $\tilde{u} \in W^{k,p}(\Omega)$ such that $\mathscr{B}u = \mathscr{B}\tilde{u}$ in ω and $\tilde{u} \in C^{k,\alpha}(\omega \cap \Omega') \cap W^{k,p+\sigma}(\omega)$.

Unlike for elliptic operators \mathscr{B} , the potential u associated with a minimizer $\mathscr{B}u$ can be wildly irregular since non-elliptic operators \mathscr{B} have large kernels. In fact, it was shown in [CG22] that ellipticity is necessary for minimizers u of $(P\mathscr{B}_{\Omega})$ to be partially $C^{k,\alpha}$ regular. This degeneracy also makes the proof of existence of generalized minimizers of Problem $(P\mathscr{B}_{\Omega})$ more involved, see the details in Section 4.2.

The natural coercivity given by the appropriate strong quasiconvexity assumption becomes insufficient to choose a good representative (Remark 3.2). This is also reflected in the Caccioppolli inequality in Proposition 6.1 which we found insufficient for performing an excess-decay estimate approach. Our main new idea is to look at a potential operator \mathscr{C} of \mathscr{B} (Theorem 2.1) and to attempt to impose the condition $\mathscr{C}^* u = 0$ in $(P\mathscr{B}_{\Omega})$. It is unclear if this is possible in general open sets, so we only worked with local estimates, see Section 5. One difficulty in proving these comes from the fact that the order of \mathscr{C} is in general larger than the order of \mathscr{B} , so it was unclear what kind of Calderón–Zygmund estimates to expect. The nonlinear estimate in (5.2) lead us to the improved Caccioppolli inequality in Proposition 6.2, which is strong enough to enable us to prove both the higher integrability in Section 6 and the partial regularity in Section 7.

Both our main results above also apply to local minimizers in the sense of [CG22; Fra19], i.e. vector fields $v \in L^p(\Omega)$ such that $\mathscr{A}v = 0$ which satisfy

$$\int_{\Omega} f(x, v(x)) \, \mathrm{d}x \leqslant \int_{\Omega} f(x, v(x) + \varphi(x)) \, \mathrm{d}x \quad \varphi \in C_c^{\infty}(\Omega) \text{ with } \mathscr{A}\varphi = 0.$$

For the \mathscr{B} -gradients setting we consider $v = \mathscr{B}u$ and $\varphi = \mathscr{B}\phi$ with $\phi \in C_c^{\infty}(\Omega)$.

We remark that in contrast to Theorem A, in Theorem B, we require no spanning cone condition. This is so since in the \mathscr{B} -gradient framework, this condition is satisfied automatically, see Lemma 2.4. We also clarify that in both Theorem A and B the notion of generalized minimality is considered with respect to the natural sequential weakly L^p -closure of the admissible classes. The details are provided in Section 4.

We now list our assumptions on integrands $f: \Omega \times V \to \mathbb{R}$:

- (H1) $|f(x,z)| \leq L(1+|z|^p)$ for any $(x,z) \in \Omega \times V$, where L > 0;
- (H2) For any $x \in \Omega$ the function $f(x, \cdot)$ is C^2 , and for any $z \in V$ the functions $f(\cdot, z)$, $\partial_z f(\cdot, z)$ and $\partial_z^2 f(\cdot, z)$ are all continuous with

$$|\partial_z f(x,z) - \partial_z f(y,z)| \leq L|x-y|(1+|z|^2)^{\frac{p-1}{2}}$$

for any $z \in V$ and any $x, y \in \Omega$;

(H3) The function f is strongly \mathscr{A} -quasiconvex in the following sense: there exists $\ell > 0$ such that for any $z \in V$, $\varphi \in C_c^{\infty}(B_1)$ with $\mathscr{A}\varphi = 0$ and a.e. $x_0 \in \Omega$, we have

$$\int_{B_1} f(x_0, z + \varphi) \, \mathrm{d}x \ge f(x_0, z) + \ell \int_{B_1} (1 + |z|^2 + |\varphi|^2)^{\frac{p-2}{2}} |\varphi|^2 \, \mathrm{d}x.$$

(H3)' The function f is strongly quasiconvex with respect to \mathscr{B} -gradients in the following sense: there exists $\ell > 0$ such that for any $z \in V$, $\phi \in C_c^{\infty}(B_1)$ and a.e. $x_0 \in \Omega$, we have

$$\int_{B_1} f(x_0, z + \mathscr{B}\phi) \,\mathrm{d}x \ge f(x_0, z) + \ell \int_{B_1} (1 + |z|^2 + |\mathscr{B}\phi|^2)^{\frac{p-2}{2}} |\mathscr{B}\phi|^2 \,\mathrm{d}x.$$

Z. LI AND B. RAIȚĂ

Notice that (H1) and (H3) together imply, by Lemma 2.3 in [KK16],

(1.1)
$$|\partial_z f(x,z)| \leq CL(1+|z|^{p-1}) \text{ for any } (x,z) \in \Omega \times V,$$

an estimate which will be used frequently in the sequel.

It was observed in [CK17] that strong quasiconvexity at a point is sufficient for proving coercivity for autonomous integrands. This interesting observation led us to realize that our strong quasiconvexity assumptions ensure existence in the case of autonomous integrands:

Theorem C. Let $f: V \to \mathbb{R}$. Under the assumptions of Theorem A (resp. B), generalized minimizers of Problem $(P\mathscr{A}_{\Omega})$ (resp. $(P\mathscr{B}_{\Omega})$) exist.

The technical description of our notion of minimality can be found in Definition 4.1. When applied to the case of exterior derivatives, this result is more general than Theorem 5.1 in [BDS15] as far as both the coercivity condition and the regularity of the boundary data are concerned. In fact, to the best of our knowledge, our results are the first partial regularity results for minimizers of integral functionals defined on exact differential forms [BDS15; BS16a; BS16b; Sil19]:

Theorem D. Let $1 and <math>u_0 \in L^p(\Omega)$ with $du_0 \in L^p(\Omega)$. Consider the problem to

(1.2)
$$\qquad \text{minimize} \quad \int_{\Omega} f(du(x)) \, \mathrm{d}x \quad \text{for } u \in u_0 + C_c^{\infty}(\Omega, \Lambda^{k-1} \mathbb{R}^n),$$

where $f: \Lambda^k \mathbb{R}^n \to \mathbb{R}$ satisfies the growth, smoothness, and strong quasiconvexity assumptions (H1), (H2), and (H3)' with $\mathscr{B} = d$. Then generalized minimizers du of (1.2) exist. For any such minimizer, there exists an open subset $\Omega' \subset \Omega$ satisfying $\mathscr{L}^n(\Omega \setminus \Omega') = 0$ and a constant $\sigma_0 > 0$ such that $du \in C^{0,\alpha}(\Omega') \cap L^{p+\sigma}_{loc}(\Omega)$ for any $\alpha \in (0,1)$ and $\sigma \in (0,\sigma_0)$. Moreover, for each $\omega \Subset \Omega$, there is a representative $\tilde{u} \in W^{1,p}(\Omega)$ such that $du = d\tilde{u}$ in ω and $\tilde{u} \in C^{1,\alpha}(\omega \cap \Omega') \cap W^{1,p+\sigma}(\omega)$.

Despite precise Hodge decompositions on domains in this case [ISS99; Sch06], we cannot obtain the existence of a representative \tilde{u} such that $d\tilde{u} = du$ in Ω due to the low boundary regularity of Ω (open set). This would be possible if Ω is assumed Lipschitz for instance.

Our last main result concerns the partial continuity and higher integrability for functionals defined on closed differential forms:

Theorem E. Let $1 and <math>v_0 \in L^p(\Omega)$ satisfy $dv_0 = 0$. Suppose that $f: \Lambda^k \mathbb{R}^n \to \mathbb{R}$ satisfies the growth, smoothness, and strong d-quasiconvexity assumptions (H1), (H2), and (H3) for $\mathscr{A} = d$. Then there exist generalized minimizers v of the problem to

minimize
$$\int_{\Omega} f(v(x)) \, \mathrm{d}x$$
 for $v \in v_0 + C^{\infty}_{c,d}(\Omega)$

For any such minimizer v, there exists an open set $\Omega' \subset \Omega$ and a constant $\sigma_0 > 0$ such that $\mathscr{L}^n(\Omega \setminus \Omega') = 0$ and $v \in C^{0,\alpha}(\Omega') \cap L^{p+\sigma}_{\text{loc}}(\Omega)$ for any $\alpha \in (0,1)$ and $\sigma \in (0,\sigma_0)$.

This result is an immediate application of Theorem A. Both Theorems D and E extend to the case of vector valued differential forms. While there has been work on regularity for functionals and nonlinear elliptic systems defined on exact or closed differential forms [Uhl77; Ham92; BS13; LOP24], we believe that Theorems D and E are new. In fact, it may well be that Theorem A is new for $\mathscr{A} = \operatorname{div}, \mathscr{A} : \mathscr{D}'(\mathbb{R}^n, \mathbb{R}^{N \times n}) \to \mathscr{D}'(\mathbb{R}^n, \mathbb{R}^n).$

Example 1.1. In the following, we list a few more examples of pde constraints \mathscr{A} of constant rank appearing in variational models for various applications. To the best of our knowledge, the regularity results established in Theorem A are new for all the following examples:

(a) Divergence on symmetric matrices: $\mathscr{A}: \mathscr{D}'(\mathbb{R}^n, \mathbb{R}^{n \times n}_{sym}) \to \mathscr{D}'(\mathbb{R}^n, \mathbb{R}^n)$ is the rowwise divergence operator. Symmetric div-quasiconvex hulls were studied in connection with elasticity theory in [CMO18; CMO20; BGS22]. (b) The linearly relaxed Euler system, $\mathscr{A}: \mathscr{D}'(\mathbb{R}^{1+d}, \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}_0^d \times \mathbb{R}) \to \mathscr{D}'(\mathbb{R}^{1+d}, \mathbb{R}^{1+d}),$

$$\mathscr{A}_E(\rho, m, M, q) = (\partial_t m + \operatorname{div} M + Dq, \, \partial_t \rho + \operatorname{div} m),$$

which was used for selecting measure-valued solution of the isentropic Euler system [Gal21; GW21; Gal23; GW23; GWW23], see also [DS09; SW12].

- (c) Generalized Saint-Venant compatibility operators: The higher order operators defined in Section 2.2, Equation (14) of [Rai+24] are used in inverse problems and tensor tomography [Sha12; PSU14; PSU23].
- (d) Higher order divergence: The operator $\mathscr{A} = \operatorname{div}^k = (D^k)^*$, i.e. the adjoint of the k-th order gradient gradient, $\mathscr{A} : \mathscr{D}'(\mathbb{R}^n, \operatorname{Slin}^k(\mathbb{R}^n, \mathbb{R}^N)) \to \mathscr{D}'(\mathbb{R}^n, \mathbb{R}^N)$. This is an ubiquitous example since any operator of order k can be written as $\operatorname{div}^k \circ T$ for some linear map T. This operator has several invariance properties which were used to prove very general statements, e.g. [Van08; Van13; Van22].

Our work is just an intersection node between two large networks of results in the calculus of variations: regularity theory and variational problems under linear pde constraints. While no amount of citations would do justice to either field, we will nevertheless endeavor to point the reader in a few fascinating directions that are related to our theme. We begin by mentioning regularity results for the widely studied curl-free case, which is Problem $(P\mathscr{A}_{\Omega})$ with $\mathscr{A} = \text{curl or Problem } (P\mathscr{B}_{\Omega})$ for $\mathscr{B} = D$, with standard growth conditions including linear growth [CN03; KT03; KM16; Li22; Li23] and (p,q)-growth conditions [Mar89; FM97; ELM99; ELM04; BS20; Sch09; Sch14; CKN21; De 22; Irv23; DS23]. Problems of the type $(P \mathscr{A}_{\Omega})$ with the same Dirichlet boundary conditions were considered recently in [RRT23; Rai+24], see also [GN04; PP04; CC15; DPR18; ST23; RT22]. Contributions related to concentration phenomena in *A*-free sequences include [FK10; KK16; DR16; DR17; Arr+19; GRS22; KR22; DT23; Arr+24; GRS24]. Symmetric div-quasiconvex hulls of sets, with an outlook to geometrically linear elasticity, were studied in [CMO18; CMO20; BGS22]. Interesting contributions to the study of the Euler equations in the \mathscr{A} -free frameworks were made in [Chi+17; GW21; Gal23; GW23; GWW23]. Homogenization results for \mathscr{A} -quasiconvex integrals can be found in [BFL00; MMS15; DF16a; DF16b; DF18; FFV21; DKP24]. Applications to image processing can be found in [Pag+22; DFL23]. Results on lower semicontinuity and regularity of minimizers of integrals defined on spaces of mixed smoothness can be found in [KSW17; Pro19; Pro23]. Other interesting contributions that do not fit in any of the above categories can be found in [DF02; San04; Baí+13; Krä+17; Pro18; KV21; SW21; LSS23; Sch24].

This paper is organized as follows: In Section 2 we collect preliminaries on differential operators and function space inequalities, in Section 3 we prove coercivity for autonomous integrands and lower semicontinuity for non-autonomous integrands, paving the way to the existence proofs in Section 4, where we also prove Theorem C. In Section 5 we prove local estimates for linear systems which are crucial for our main results. In Section 6 we prove two Caccioppolli inequalities and show how they lead to higher integrability. Finally, in Section 7 we prove the excess decay estimates that lead to partial regularity and complete the proofs of Theorems A and B.

Acknowledgement. The authors thank Tatiana Toro for insightful discussions. B.R. thanks SLMath and Tatiana Toro for the hospitality and financial support during a visit when significant parts of the current research were conducted. Part of this paper is based upon work supported by the National Science Foundation under Grant No. DMS-1928930 and by the Alfred P. Sloan Foundation under grant G-2021-16778, while Z.L. was in residence at the Simons Laufer Mathematical Sciences Institute (formerly MSRI) in Berkeley, California, during the Spring 2024 semester.

Z. LI AND B. RAIȚĂ

2. Preliminaries

2.1. Differential operators. We will work with differential operators

$$\mathscr{A} = \sum_{|\alpha|=h} A_{\alpha} \partial^{\alpha}, \quad \mathscr{B} = \sum_{|\beta|=k} B_{\beta} \partial^{\beta}, \quad \mathscr{C} = \sum_{|\gamma|=l} C_{\gamma} \partial^{\gamma},$$

where A_{α} , B_{β} , C_{γ} are matrix coefficients. The three operators will be assumed to satisfy certain exact relations in Fourier variable that we now explain. To this end, we will use the notation

$$\mathscr{A} = \sum_{|\alpha|=h} A_{\alpha} \xi^{\alpha}, \quad \mathscr{B} = \sum_{|\beta|=k} B_{\beta} \xi^{\beta}, \quad \mathscr{C} = \sum_{|\gamma|=l} C_{\gamma} \xi^{\gamma},$$

for the characteristic (matrix) polynomials of respectively \mathscr{A}, \mathscr{B} , and \mathscr{C} .

Recall that an operator \mathscr{B} is said to be of **constant rank** if rank $\mathscr{B}(\xi) = \text{const.}$ for all $\xi \in \mathbb{R}^n \setminus \{0\}$. We will need the following result from [Rai19b]:

Theorem 2.1. Let \mathscr{B} be an operator as above which is assumed to have constant rank. Then there exists an operator \mathscr{C} as above, also of constant rank, such that

(2.1)
$$\operatorname{im} \mathscr{C}(\xi) = \ker \mathscr{B}(\xi) \quad \text{for all } \xi \in \mathbb{R}^n \setminus \{0\}$$

Moreover, if $B_{\beta} \in \text{Lin}(U, V)$, we can choose \mathscr{C} such that $C_{\gamma} \in \text{Lin}(U, U)$ and l > k.

The reason why we can define \mathscr{C} on U as well is by construction and we can choose l > k because we can simply replace \mathscr{C} with $\Delta^m \mathscr{C}$ for large enough m. By duality and iteration we can obtain the following:

Corollary 2.2. Let \mathscr{A} be an operator as above which is assumed to have constant rank. Then there exist operators \mathscr{B} , \mathscr{C} as above, also of constant rank, such that

(2.2)
$$\operatorname{im} \mathscr{B}(\xi) = \ker \mathscr{A}(\xi), \quad \operatorname{im} \mathscr{C}(\xi) = \ker \mathscr{B}(\xi) \quad \text{for all } \xi \in \mathbb{R}^n \setminus \{0\}$$

Moreover, if $A_{\alpha} \in \operatorname{Lin}(V, W)$, we can choose \mathscr{B}, \mathscr{C} such that $B_{\beta}, C_{\gamma} \in \operatorname{Lin}(V, V)$ and l > k > h.

Remark 2.3. It was observed in [Rai24] that \mathscr{A} can have homogeneous rows (i.e. if we write $\mathscr{A}(\xi) = (\mathscr{A}_{ij}(\xi))$, then \mathscr{A}_{ij} is h_i -homogeneous) and the conclusion of Corollary 2.2 remains identical. It can be showed that in these circumstances the conclusion of the main result Theorem A remains the same. We do not include the details in the present work.

We also clarify that U, V, W above are finite dimensional inner product spaces. In the case of Corollary 2.2, we have that U = V. With this assumption, we have that in both Theorem 2.1 and Corollary 2.2, we have $\mathscr{A}(\xi) \in \operatorname{Lin}(V,W), \ \mathscr{B}(\xi) \in \operatorname{Lin}(U,V),$ $\mathscr{C}(\xi) \in \operatorname{Lin}(U,U)$ for $\xi \in \mathbb{R}^n$.

We will also use the notation $\mathscr{B}u = T(D^k u)$, for a linear map $T \in \operatorname{Lin}(\operatorname{SLin}^k(\mathbb{R}^n, U), V)$, which only amounts to collecting all the coefficients of \mathscr{B} in a tensor. We also record that $\mathscr{B}(\xi)u_0 = T(u_0 \otimes \xi^{\otimes k})$. With this notation we have the following observation:

Lemma 2.4. For any homogeneous linear differential operator \mathscr{B} with constant coefficients we have that

$$\operatorname{span} \bigcup_{\xi \in \mathbb{R}^n \setminus \{0\}} \operatorname{im} \mathscr{B}(\xi) = \{ \mathscr{B}u(x) \colon u \in C_c^{\infty}(\mathbb{R}^n), x \in \mathbb{R}^n \} = \operatorname{im} T.$$

Proof. First consider the case when $\mathscr{B} = D^k$, so $V = \mathrm{SLin}^k(\mathbb{R}^n, U)$ and T = 1. We then have

$$\operatorname{span} \bigcup_{\xi \in \mathbb{R}^n \setminus \{0\}} U \otimes \xi^{\otimes k} = \{ D^k u(x) \colon u \in C_c^{\infty}(\mathbb{R}^n), x \in \mathbb{R}^n \} = \operatorname{SLin}^k(\mathbb{R}^n, U),$$

which settles this case. To prove the general case, apply T to the last equality to get

$$\operatorname{span} \bigcup_{\xi \in \mathbb{R}^n \setminus \{0\}} T(U \otimes \xi^{\otimes k}) = \{ T(D^k u(x)) \colon u \in C_c^{\infty}(\mathbb{R}^n), x \in \mathbb{R}^n \} = T(\operatorname{SLin}^k(\mathbb{R}^n, U)),$$

which concludes the proof.

Remark 2.5. The relevance of Lemma 2.4 is that in the \mathscr{B} -gradient setting of Theorem B, we can simply assume that $f: \Omega \times \operatorname{im} T \to \mathbb{R}$. The spanning cone condition is satisfied automatically. In fact, not even for the \mathscr{A} -free setting, the spanning cone condition (SC) is not at all restrictive, as can be seen from Lemma 4.2 in [KR22].

2.2. Function spaces. Let $\Omega \subset \mathbb{R}^n$ be an open set. We define $L^p_{0,\mathscr{A}}(\Omega)$ as the strong L^p -closure of the set $C^{\infty}_{c,\mathscr{A}}(\Omega) = \{\varphi \in C^{\infty}_c(\Omega, V) : \mathscr{A}\varphi = 0\}$. This is a Banach space, as it is a closed subspace of $L^p(\Omega, V)$. We also define $X_0^{\mathscr{B},p}(\Omega)$ as the L^p -closure of the set $\mathscr{B}C^{\infty}_c(\Omega, U) = \{\mathscr{B}\phi : \phi \in C^{\infty}_c(\Omega, U)\}$, which is also a Banach space.

Let $m \ge 0$ be an integer. We define $\dot{W}^{m,p}(\Omega)$ as the closure of $C_c^{\infty}(\Omega)$ in the (semi-)norm $u \mapsto \|D^k u\|_{L^p(\Omega)}$. Its dual space is $\dot{W}^{-m,p}(\Omega) = \dot{W}^{m,p}(\Omega)^*$. If Ω is bounded, the full Sobolev norm is an equivalent norm on $\dot{W}^{m,p}(\Omega)$ by Poincaré's inequality and therefore $\dot{W}^{-m,p}(\Omega)$ coincides with the classic negative Sobolev space $W^{-m,p}(\Omega)$. If $\Omega = \mathbb{R}^n$, we have the simple Fourier description

$$||f||_{\dot{W}^{m,p}(\mathbb{R}^n)} = ||D^m f||_{L^p(\mathbb{R}^n)} = ||\mathcal{F}^{-1}(|\xi|^m \dot{f})||_{L^p(\mathbb{R}^n)}$$

which holds on $C_c^{\infty}(\mathbb{R}^n)$ and is extended by density to $\dot{W}^{m,p}(\mathbb{R}^n)$. On the dual side we have

$$\|f\|_{\dot{W}^{-m,p}(\mathbb{R}^n)} \sim_{n,p,m} \|\mathcal{F}^{-1}(|\xi|^{-m}\hat{f})\|_{L^p(\mathbb{R}^n)}$$

which can be made precise, e.g. with the techniques from §5.2 in [Tri83]:

Theorem 2.6. Given any non-negative integer $m \in \mathbb{N}$ and any 1 , we have the following estimate for <math>(-m)-homogeneous multipliers:

 $\|\mathcal{F}^{-1}|\xi|^{-m}\hat{\varphi}(\xi)\|_{L^p(\mathbb{R}^n)}\sim_{n,p,m}\|\varphi\|_{\dot{W}^{-m,p}(\mathbb{R}^n)},\quad for \ any \ \varphi\in\dot{W}^{-m,p}(\mathbb{R}^n).$

Proof. Denote by $F_{p,q}^s$ ($\dot{F}_{p,q}^s$) the (homogeneous) Triebel-Lizorkin space with parameters s and p, q. See §5.2 in [Tri83] for the precise definition. For any $\sigma \in \mathbb{R}$, define the lifting operator \dot{I}^{σ} by

$$\dot{I}^{\sigma}\varphi := \mathcal{F}^{-1}(|\cdot|^{\sigma}\hat{\varphi}(\cdot)), \quad \varphi \in \mathcal{S}'(\mathbb{R}^n),$$

where $\mathcal{S}'(\mathbb{R}^n)$ is the space of Schwartz distributions.

By Theorem 1 in $\S5.2.3$ in [Tri83], the map

$$\dot{I}^{-m} \colon \dot{F}^0_{p',2}(\mathbb{R}^n) \to \dot{F}^m_{p',2}(\mathbb{R}^n)$$

is an isomorphism. By the duality of homogeneous Triebel-Lizorkin spaces (see $\S2.11.2$ and $\S5.2.5$ in [Tri83]), we know that the following map

$$I^{-m} = (I^{-m})^* \colon \dot{F}_{p,2}^{-m}(\mathbb{R}^n) = (\dot{F}_{p',2}^m(\mathbb{R}^n))^* \to (\dot{F}_{p',2}^0(\mathbb{R}^n))^* = \dot{F}_{p,2}^0(\mathbb{R}^n)$$

is also an isomorphism.

Theorem 1 in $\S5.2.3$ in [Tri83] also tells

$$\dot{F}_{p,2}^{0}(\mathbb{R}^{n}) = L^{p}(\mathbb{R}^{n}), \quad \dot{F}_{p',2}^{m}(\mathbb{R}^{n}) = \dot{W}^{m,p'}(\mathbb{R}^{n}),$$

where p' = p/(p-1) is the conjugate exponent of p.

Since $\dot{W}^{-m,p}(\mathbb{R}^n)$ is the dual space of $\dot{W}^{m,p'}(\mathbb{R}^n) \simeq \dot{F}^m_{p',2}(\mathbb{R}^n)$, we have that the negative Sobolev space is isomorphic to $\dot{F}^{-m}_{p,2}(\mathbb{R}^n)$. It follows from the above that the map

$$\dot{I}^{-m} \colon \dot{W}^{-m,p}(\mathbb{R}^n) \to L^p(\mathbb{R}^n)$$

is an isomorphism, which implies the desired inequality.

The following result was essentially proved in [Rai19b; GR22]. We sketch a proof for the convenience of the reader.

Theorem 2.7 (Full space Hodge decomposition). Let 1 .

(a) Let $\mathscr{A}, \mathscr{B}, \mathscr{C}$ be operators which satisfy the assumptions of Corollary 2.2. Then, each $v \in L^p(\mathbb{R}^n, V)$ can be decomposed as

$$v = \mathscr{B}u + \mathscr{A}^* w$$

where $u \in \dot{W}^{k,p}(\mathbb{R}^n, U)$, $w \in \dot{W}^{h,p}(\mathbb{R}^n, W)$ are such that $\mathscr{C}^*u = 0$ and

$$||D^{k}u||_{L^{p}(\mathbb{R}^{n})} + ||D^{h}w||_{L^{p}(\mathbb{R}^{n})} \leq C||v||_{L^{p}(\mathbb{R}^{n})}.$$

(b) Let \mathscr{B} and \mathscr{C} be operators that satisfy the assumptions of Theorem 2.1. We have that each $\varphi \in C_c^{\infty}(\mathbb{R}^n, U)$ can be written as $\varphi = \tilde{\varphi} + \psi$, where $\mathscr{C}^* \tilde{\varphi} = 0$, $\mathscr{B} \tilde{\varphi} = \mathscr{B} \varphi$, and $\tilde{\varphi} \in \dot{W}^{k,p}(\mathbb{R}^n, U)$ with

$$\|D^k \tilde{\varphi}\|_{L^p(\mathbb{R}^n)} \leqslant C \|\mathscr{B}\varphi\|_{L^p(\mathbb{R}^n)}$$

Sketch. We know that for constant rank operators \mathscr{A} we can write

$$\operatorname{Proj}_{\operatorname{im}\mathscr{A}(\xi)} = \mathscr{A}(\xi)\mathscr{A}^{\dagger}(\xi),$$

where $\mathscr{A}^{\dagger} \in C^{\infty}(\mathbb{R}^n \setminus \{0\})$ is (-h)-homogeneous, see [FM99; Rai19b]. Using this fact, elementary linear algebra, and the exact relations (2.2) we can write

$$\hat{v} = \mathscr{B}(\xi)\mathscr{B}^{\dagger}(\xi)\hat{v} + \mathscr{A}^{*}(\xi)\mathscr{A}^{*\dagger}(\xi)\hat{v},$$

so we can define

$$\hat{u} = \mathscr{B}^{\dagger}(\xi)\hat{v}, \quad \hat{w} = \mathscr{A}^{*}(\xi)\mathscr{A}^{*\dagger}(\xi)\hat{v}.$$

The estimates follow from

$$\widehat{D^k u} = \mathscr{B}\left(\frac{\xi}{|\xi|}\right) \hat{v} \otimes \left(\frac{\xi}{|\xi|}\right)^{\otimes k}$$

and the analogous identity for w and the Hörmander–Mikhlin multiplier theorem. Finally, since im $\mathscr{B}^{\dagger}(\xi) = \operatorname{im} \mathscr{B}^{*}(\xi)$, we also have that $\mathscr{C}^{*}(\xi)\hat{u} = 0$, so $\mathscr{C}^{*}u = 0$, proving part (a). Part (b) follows from part (a) by replacing \mathscr{A} with \mathscr{C}^* and \mathscr{B} with \mathscr{B}^* .

2.3. Auxiliary results. We first recall some properties of the V-function defined as $V_p: X \to X$ by $V_p(z) := (1+|z|^2)^{\frac{p-2}{4}} z$, where X is any finite dimensional inner-product space. We may use V(z) for simplicity when the exponent p is clear from the context.

Lemma 2.8. For any $z, w \in X$, we have the following:

- (a) $|V_p(z)|^2 \sim_p (|z|^2 + |z|^p) \sim_p \max\{|z|^2, |z|^p\}$ if $p \ge 2$, and $|V_p(z)|^2 \sim_p \min\{|z|^2, |z|^p\}$ if 1 ;
- (b) $|V_p(z+w)| \leq_p |V_p(z) + V_p(w)|;$
- (c) $|V_p(tz)| \leq \max\{t, t^{\frac{p}{2}}\}|V_p(z)|$ for any t > 0;
- (d) $|V_n(\cdot)|^2$ satisfies Jensen's inequality up to a constant:

$$\left| V_p \left(\oint_{\omega} g \, \mathrm{d}x \right) \right|^2 \leqslant C(p) \oint_{\omega} |V_p(g)|^2 \, \mathrm{d}x$$

for any bounded domain $\omega \subset \mathbb{R}^n$ and any $g \in L^p(\omega)$ with some constant $C(p) \ge 1$.

Proof. Properties (a)-(c) can be easily proved in the case $p \ge 2$, and see Lemma 2.1 in [CFM98] for 1 .

The inequality in (d) holds true when $p \ge 2$ since $|V_p(\cdot)|^2 \sim_p |\cdot|^2 + |\cdot|^p$ and the latter is convex. In the case $1 , we take <math>E_p(\cdot) := (1 + |\cdot|^2)^{\frac{p}{2}} - 1$ as the competitor, which is convex as well. \square

For any $z, w \in V$ and any $x_0 \in \Omega$, define the shifted integrand by

$$f_w(x_0, z) := f(x_0, z + w) - f(x_0, w) - \partial_z f(x_0, w) \cdot z,$$

where the function $f: \Omega \times V \to \mathbb{R}$ satisfies (H1)-(H2).

Lemma 2.9. Suppose that the integrand $f: \Omega \times V \to \mathbb{R}$ satisfies (H1)-(H2). Given any $w \in V$ with $|w| \leq M$ for some m > 0, the following estimates hold true:

(2.3)
$$|f_w(x_0, z)| \leq C(n, p, \dim V, M, f)|V_p(z)|^2$$

$$(2.4) |f_w(x_0,z) - f_w(x_0,y)| \leq C(n,p,\dim V,M,f)(1+|z|^2+|y|^2)^{\frac{p-2}{2}}|z+y||z-y|,$$

(2.5)
$$\int_{B_r} f_w(x_0, z + \varphi) \, \mathrm{d}x \ge C(p, \ell, M) \int_{B_r} |V_p(\varphi)|^2 \, \mathrm{d}x,$$

for any $\varphi \in C_c^{\infty}(B_r)$ with $\mathscr{A}\varphi = 0$. Moreover, if we consider the \mathscr{B} -derivative of a map, the inequality (2.5) holds true for any $\varphi = \mathscr{B}\phi$ with $\phi \in W_0^{k,p}(B_r)$.

Proof. Inequality (2.3) can be proved by considering the two cases $|z| \ge 1$ and |z| < 1 separately. In the first case, the Lipchitz continuity (1.1) of $f(x_0, \cdot)$ implies

$$|f_w(x_0, z)| = \left| \int_0^1 \partial_z f(x_0, w + tz) \cdot z \, \mathrm{d}t - \partial_z f(x_0, w) \cdot z \right|$$

$$\leqslant \int_0^1 C(2 + |w + tz|^{p-1} + |w|^{p-1}) \, \mathrm{d}t |z| \leqslant C |V_p(z)|^2.$$

In the second one, by Taylor's theorem we have

$$|f_w(x_0, z)| = \left| \int_0^1 (1-t) \partial_z^2 f(x_0, w + tz)[z, z] \, \mathrm{d}t \right| \le C |z|^2 \le C |V_p(z)|^2.$$

To show (2.4), we need the estimate

$$|\partial_z f_w(x_0, z)| \leq C(1 + |z|^2)^{\frac{p-2}{2}} |z|$$

The latter can be proved as well by considering $|z| \ge 1$ and |z| < 1 separately. Then (2.4) follows from this estimate and Lemma 2.11.

The strong \mathscr{A} -quasiconvexity (**H3**) of f implies (2.5). Then use the Lipschitz continuity of $f(x_0, \cdot)$ to extend it to $\varphi = \mathscr{B}\phi$ with $\phi \in W_0^{k,p}(B_r)$.

We conclude this section with two auxiliary results that will be used later. The first one is an iteration lemma used in the proof of the Caccioppolli-type inequality and is adapted from Lemma 6.1 in [Giu03].

Lemma 2.10. Suppose that the function $\Phi: (0, R] \to \mathbb{R}_+$ is bounded, and $\Psi_i: (0, R] \to \mathbb{R}_+, i = 1, \ldots, k$, are decreasing with $\Psi_i(\sigma t) \leq \sigma^{-\beta i} \Psi(t), \beta > 0$, for any $t \in (0, R]$ and any $\sigma \in (0, 1)$. If the following inequality

(2.6)
$$\Phi(r) \leqslant \theta \Phi(s) + \sum_{i=1}^{k} \Psi_i(s-r) + B$$

holds true for any $r, s \in (0, R]$ with r < s and some given $\theta \in (0, 1)$, B > 0, then there exists $C = C(\theta, k, \beta, \tau) > 0$ for any $\tau \in (0, 1)$ such that $\Phi(\tau R)$ is controlled as follows:

$$\Phi(\tau R) \leqslant C\left(\sum_{i=1}^{k} \Psi_i(R) + B\right).$$

Proof. Take $\lambda \in (0, 1)$ to be determined later, and set

$$r_0 = \tau R,$$
 $r_{j+1} - r_j = (1 - \lambda)\lambda^j (1 - \tau)R$ for $j \in \mathbb{N}.$

Then we apply (2.6) to $r_j, r_j + 1$ iteratively with $\tau' := 1 - \tau$ and obtain

$$\Phi(r_0) \leqslant \theta \Phi(r_1) + \sum_{i=1}^k \Psi_i((1-\lambda)\tau'R) + B$$

$$\leqslant \theta(\theta \Phi(r_2) + \sum_{i=1}^k \Psi_i((1-\lambda)\lambda\tau'R) + B) + \sum_{i=1}^k \Psi_i((1-\lambda)\tau'R) + B$$

$$\leqslant \dots \leqslant \theta^{L+1} \Phi(r_{L+1}) + \sum_{j=0}^L \theta^j \left(\sum_{i=1}^k \Psi_i((1-\lambda)\lambda^j\tau'R) + B\right).$$

Since $\Psi_i((1-\lambda)\lambda^j\tau'R) \leq (\tau')^{\beta i}(1-\lambda)^{-\beta i}\lambda^{-\beta ij}\Psi_i(R)$, under the condition $\theta < \lambda^{\beta i}, i = 1, \ldots, k$, the above inequality becomes

$$\Phi(r_0) \leqslant \theta^{L+1} \Phi(r_{L+1}) + \Psi(R) \sum_{i=1}^k \frac{(\tau')^{\beta i}}{(1-\lambda)^{\beta i}} \left(\sum_{j=0}^L \frac{\theta^j}{\lambda^{\beta i j}} \right) + \frac{1-\theta^{L+1}}{1-\theta} B$$

$$\leqslant \theta^{L+1} \Phi(r_{L+1}) + \Psi(R) \sum_{i=1}^k \frac{(\tau')^{\beta i}}{(1-\lambda)^{\beta i}} \frac{1-(\theta\lambda^{-\beta i})^{L+1}}{1-\theta\lambda^{-\beta i}} + \frac{1-\theta^{L+1}}{1-\theta} B.$$

The desired result can be obtained by taking $L \to \infty$.

The second one is Lemma 8.3 in [Giu03], and we include it here for completeness:

Lemma 2.11. Suppose that X is a finite-dimensional vector space and $z, w \in X$ with $z \neq w$. Then for any $-1 < q < \infty$ and $s, \gamma \ge 0$, there exists $c = c(q, s, \gamma) \ge 1$ such that

$$(2.7) \quad \frac{1}{c_1} (\gamma^2 + |z|^2 + |w|^2)^{\frac{q}{2}} \leqslant \int_0^1 (1-t)^s (\gamma^2 + |tz + (1-t)w|^2)^{\frac{q}{2}} dt \\ \leqslant c_1 (\gamma^2 + |z|^2 + |w|^2)^{\frac{q}{2}}.$$

3. Coercivity and lower semicontinuity

We next prove that strong quasiconvexity implies coercivity for autonomous integrands, under standard growth conditions.

Theorem 3.1. Suppose that function $f: V \to \mathbb{R}$ satisfies (H1) and (H3). Then there exist constants $a_i \in \mathbb{R}$, i = 1, 2, 3, with $a_1 = a_1(p, \ell) > 0$ and $a_2 = a_2(n, \dim V, p, f(0), L, \ell)$ and $a_3 = a_3(n, \dim V, p, L, \ell) > 0$ such that, for any bounded open set $\Omega \subset \mathbb{R}^n$ and \mathscr{A} -free map $v_0 \in L^p(\Omega, V)$, the following holds:

(3.1)
$$\int_{\Omega} f(v_0 + \varphi) \,\mathrm{d}x \ge a_1 \int_{\Omega} |\varphi|^p \,\mathrm{d}x + a_2 - a_3 \int_{\Omega} |v_0|^p \,\mathrm{d}x$$

for any $\varphi \in C^{\infty}_{c,\mathscr{A}}(\Omega, V)$.

Proof. First, by (H3), the strong \mathscr{A} -quasiconvexity of f, we have

(3.2)
$$\int_{\Omega} f(v_0 + \varphi) \, \mathrm{d}x = \int_{\Omega} f(\varphi) \, \mathrm{d}x + \int_{\Omega} (f(v_0 + \varphi) - f(\varphi)) \, \mathrm{d}x$$
$$\geqslant \int_{\Omega} (\ell |V_p(\varphi)|^2 + f(0)) \, \mathrm{d}x + \int_{\Omega} \int_0^1 f'(\varphi + tv_0) \, \mathrm{d}t \cdot v_0 \, \mathrm{d}x$$
$$=: I + II.$$

The two terms I and II are estimated separately in the following. Since we know that $|z|^p \leq_p |V_p(z)|^2 + 1$ by (a), the first term can be bounded from below by

(3.3)
$$I \ge C_1 \int_{\Omega} |\varphi|^p \,\mathrm{d}x + (f(0) - C_1') \mathscr{L}^n(\Omega),$$

where $C_1, C_2 > 0$ depending on p, ℓ .

To control the second term, the Lipschitz continuity of f in V (see (1.1)) is needed. With this bound of f' and Young's inequality, we have

(3.4)

$$|II| \leq CL \int_{\Omega} \int_{0}^{1} (1 + |\varphi + tv_{0}|^{p-1}) |v_{0}| \, \mathrm{d}t \, \mathrm{d}x$$

$$\leq C \int_{\Omega} (1 + |\varphi|^{p-1} |v_{0}| + |v_{0}|^{p}) \, \mathrm{d}x$$

$$\leq C_{2} \int_{\Omega} (1 + \varepsilon |\varphi|^{p} + C(\varepsilon, p) |v_{0}|^{p}) \, \mathrm{d}x,$$

where $C_2 = C_2(n, \dim V, p, L) > 0$. Take ε to be small enough such that $\varepsilon C_2 < \frac{C_1}{2}$, then it follows from (3.2), (3.3) and (3.4) that

$$\begin{aligned} \oint_{\Omega} f(v_0 + \varphi) \, \mathrm{d}x &\geq C_1 \oint_{\Omega} |\varphi|^p \, \mathrm{d}x + (f(0) - C_1') \mathscr{L}^n(\Omega) \\ &- C_2 \left(\varepsilon \int_{\Omega} |\varphi|^p \, \mathrm{d}x + C(\varepsilon, p) \oint_{\Omega} |v_0|^p \, \mathrm{d}x + \mathscr{L}^n(\Omega) \right) \\ &\geq \frac{C_1}{2} \oint_{\Omega} |\varphi|^p \, \mathrm{d}x + C_3 - C_2 C(\varepsilon, p) \oint_{\Omega} |v_0|^p \, \mathrm{d}x. \end{aligned}$$

The desired inequality (3.1) then follows with $a_1 = \frac{C_1}{2}$, $a_2 = C_3$ and $a_3 = C_2 C(\varepsilon, p)$.

Remark 3.2. Notice that for any $g \in \mathscr{D}'(\Omega, U)$ with $\mathscr{B}g \in L^p(\Omega, V)$ and $\phi \in C_c^{\infty}(\Omega, U)$, the following coercivity inequality also holds true by the same argument:

$$\int_{\Omega} f(\mathscr{B}(g+\phi)) \,\mathrm{d}x \ge a_1 \int_{\Omega} |\mathscr{B}\phi|^p \,\mathrm{d}x + a_2 - a_3 \int_{\Omega} |\mathscr{B}g|^p \,\mathrm{d}x.$$

Lemma 3.3. Suppose that $f: \Omega \times V \to \mathbb{R}$ is \mathscr{A} -quasiconvex ((**H3**) with $\ell = 0$) and satisfies (**H1**) for $1 . Then for any <math>v_0 \in L^p(\Omega, V)$ and any $C^{\infty}_{c,\mathscr{A}}(\Omega) \ni v_j \to v$ in $L^p(\Omega)$ we have that

$$\liminf_{j \to \infty} \int_{\Omega} f(x, v_0 + v_j) \, \mathrm{d}x \ge \int_{\Omega} f(x, v_0 + v) \, \mathrm{d}x.$$

The reason why we isolate this result as a separate lemma is to highlight that it does not follow from the main results of [FM99], which does not cover signed integrands explicitly. The lemma also does not follow by an immediate application of the lower semicontinuity results in [GR22] either, as those do not cover effects of concentration at the boundary. The simple argument below shows that this is not an issue in the context of Dirichlet boundary conditions.

Proof. Consider an open bounded set $\Omega' \supseteq \Omega$. Extend $v_0, \{v_j\}$ and v by 0 to $\Omega' \setminus \Omega$ and denote the extension by $\bar{v}_0, \{\bar{v}_j\}$ and \bar{v} , respectively. Also let $\bar{f}(x, z) = \chi_{\Omega}(x)f(x, z)$. It is easy to see that $\bar{v}_j \rightarrow \bar{v}$ in $L^p(\Omega')$. Note that $\mathscr{A}(\bar{v}_0 + \bar{v}_j) = \mathscr{A}\bar{v}_0 \in W^{-h,p}_{\text{loc}}(\Omega')$, so Theorem 5.6 in [KR20] applies in Ω' , giving

$$\liminf_{j \to \infty} \int_{\Omega'} \bar{f}(x, \bar{v}_0 + \bar{v}_j) \, \mathrm{d}x \ge \int_{\Omega'} \bar{f}(x, \bar{v}_0 + \bar{v}) \, \mathrm{d}x$$

The claim is then immediate.

Z. LI AND B. RAIȚĂ

4. EXISTENCE OF GENERALIZED MINIMIZERS

We begin by defining our notion of generalized minimizers, which requires $L_{0,\mathscr{A}}^{p}(\Omega)$ and $X_{0}^{\mathscr{B},p}(\Omega)$, the two weakly closed L^{p} -based classes defined in Section 2.2.

Definition 4.1. Let $v_0 \in L^p(\Omega, V)$ be such that $\mathscr{A}v_0 = 0$ (resp. $u_0 \in \mathscr{D}'(\Omega, U)$ be such that $v_0 = \mathscr{B}u_0 \in L^p(\Omega, V)$). Let $f \colon \Omega \times V \to \mathbb{R}$ be measurable. We say that $v \in v_0 + L^p_{0,\mathscr{A}}(\Omega)$ (resp. $v \in \mathscr{B}u_0 + X^{\mathscr{B},p}_0(\Omega)$) is a generalized minimizer of Problem $(P\mathscr{A}_\Omega)$ (resp. $(P\mathscr{B}_\Omega)$) if

$$\int_{\Omega} f(x, v(x)) \, \mathrm{d}x \leqslant \int_{\Omega} f(x, v_0(x) + \varphi(x)) \, \mathrm{d}x \quad \text{for all } \varphi \in L^p_{0,\mathscr{A}}(\Omega) \ (\text{resp. } X^{\mathscr{B}, p}_0(\Omega)).$$

4.1. \mathscr{A} -free setting. We begin by proving that \mathscr{A} -quasiconvexity and coercivity imply existence of minimizers of Problem $(P\mathscr{A}_{\Omega})$ by the direct method.

Proposition 4.2. Suppose that $f: \Omega \times V \to \mathbb{R}$ is \mathscr{A} -quasiconvex ((**H3**) with $\ell = 0$) and satisfies (**H1**) for 1 . Assume in addition that there exists <math>C > 0 such that for any $v_0 \in L^p(\Omega, V)$, $\varphi \in C_c^{\infty}(\Omega, V)$ with $\mathscr{A}v_0 = 0 = \mathscr{A}\varphi$, we have that

(4.1)
$$\int_{\Omega} f(x, v_0 + \varphi) \, \mathrm{d}x \ge C \left(\int_{\Omega} |\varphi|^p - 1 - \int_{\Omega} |v_0|^p \right).$$

Then Problem $(P\mathscr{A}_{\Omega})$ admits a generalized minimizer in the sense of Definition 4.1.

Proof. Let $v_0 \in L^p(\Omega, V)$ be \mathscr{A} -free. We claim that

$$\inf\left\{\int_{\Omega} f(x, v_0 + v) \colon v \in C^{\infty}_{c,\mathscr{A}}(\Omega)\right\} = \min\left\{\int_{\Omega} f(x, v_0 + v) \colon v \in L^p_{0,\mathscr{A}}(\Omega)\right\}.$$

First, it is obvious that the inequality " \geq " holds. Second, we show that the minimization problem on the right has a solution in $L^p_{0,\mathscr{A}}(\Omega)$. To see this note that by (4.1), we can find a minimizing sequence $\{v_j\} \subset L^p_{0,\mathscr{A}}(\Omega)$ which is bounded in $L^p(\Omega)$ by the assumed coercivity (4.1). It follows by the \mathscr{A} -quasiconvexity of f and Lemma 3.3 that

$$\liminf_{j \to \infty} \int_{\Omega} f(x, v_0 + v_j) \, \mathrm{d}x \ge \int_{\Omega} f(x, v_0 + \tilde{v}) \, \mathrm{d}x,$$

where \tilde{v} is a weak limit point of $\{v_j\}$ and thus $\tilde{v} \in L^p_{0,\mathscr{A}}(\Omega)$ by Mazur's Lemma. The map \tilde{v} is thus a minimizer of the problem on the right.

Let now $\phi_j \in C^{\infty}_{c,\mathscr{A}}(\Omega)$ be such that $\phi_j \to \tilde{v}$ in $L^p(\Omega, V)$. The Lipschitz continuity of f in (1.1) implies

$$\int_{\Omega} f(x, v_0 + \phi_j) \, \mathrm{d}x \to \int_{\Omega} f(x, v_0 + \tilde{v}) \, \mathrm{d}x.$$

which shows that $\{\phi_j\}$ is a minimizing sequence for the problem on the left, so the claimed equality holds true.

Proof of Theorem C, A-free setting. Follows immediately from Proposition 4.2, Lemma 3.3, and Theorem 3.1.

We proceed to show that any minimizer \tilde{v} has a far better structure. In particular, the following will be deduced in Proposition 4.3: as far as problem $(P\mathscr{A}_{\Omega})$ is concerned, we can assume without loss of generality that Ω is smooth and that the competitors have the form $v = \phi + \mathscr{B}u$, where $\phi \in C^{\infty}(\bar{\Omega})$ is \mathscr{A} -free and $u \in C^{\infty}_{c}(\Omega, V)$.

Proposition 4.3. Let \tilde{v} be the minimizer described above and let $\omega \in \Omega$. We have that

$$\inf\{\mathcal{E}(v,\omega)\colon v\in v_0+\tilde{v}+C^{\infty}_{c,\mathscr{A}}(\omega,V)\}=\mathcal{E}(v_0+\tilde{v},\omega)$$

i.e. that $v_0 + \tilde{v}$ is a local minimizer of $(P\mathscr{A}_{\Omega})$. Moreover, it is possible to write $v_0 + \tilde{v} = \mathscr{B}\tilde{u} + \phi$ in ω , where

(a) $\tilde{u} \in W^{k,p}(\mathbb{R}^n, U)$ satisfies $\mathscr{C}^* \tilde{u} = 0$ in \mathbb{R}^n ,

(b) $\phi \in L^p(\mathbb{R}^n, V)$ satisfies $\mathscr{B}^*\phi = 0$ in \mathbb{R}^n and $\mathscr{A}\phi = 0$ in $\bar{\omega}$.

In particular, $\phi \in C^{\infty}(\bar{\omega}, V)$.

Proof. Take $\rho \in C_c^{\infty}(\Omega)$ be equal to 1 in an open neighborhood of $\bar{\omega}$ and perform the Hodge decomposition of $\rho(v_0 + \tilde{v})$ as in Theorem 2.7(a):

$$\rho(v_0 + \tilde{v}) = \mathscr{B}\tilde{u} + \phi_i$$

where $\tilde{u} \in \dot{W}^{k,p}(\mathbb{R}^n)$ with $\mathscr{C}^* u = 0$ in \mathbb{R}^n and $\phi = \mathscr{A}^* \psi$ for some $\psi \in \dot{W}^{h,p}(\mathbb{R}^n)$. Therefore $\mathscr{B}^* \phi = \mathscr{B}^* \mathscr{A}^* \psi = 0$ in \mathbb{R}^n . Since

$$\mathscr{A}(v_0 + \tilde{v}) = 0 \text{ in } \Omega \quad \text{and} \quad \mathscr{AB}\tilde{u} = 0 \text{ in } \mathbb{R}^n,$$

we obtain $\mathscr{A}\phi = 0$ in $\bar{\omega}$.

The assertion concerning the smoothness (in fact, analyticity) of ϕ follows from the fact that the system

$$\mathscr{A}\phi = 0, \quad \mathscr{B}^*\phi = 0$$

is elliptic. We verify the ellipticity explicitly here: let $\xi \neq 0$, so that ker $\mathscr{A}(\xi) \cap \ker \mathscr{B}^*(\xi) =$ im $\mathscr{B}(\xi) \cap (\operatorname{im} \mathscr{B}(\xi))^{\perp} = \{0\}.$

Remark 4.4. In particular, we obtain that $\tilde{u} \in W^{k,p}(\omega)$ is \mathscr{C}^* -free and that

$$\inf\left\{\int_{\omega} f(x,\mathscr{B}u+\phi)\colon u\in\tilde{u}+C_c^{\infty}(\omega)\right\}=\int_{\omega} f(x,\mathscr{B}\tilde{u}+\phi),$$

where $\phi \in C^{\infty}(\bar{\omega}, V)$ is \mathscr{A} -free and \mathscr{B}^* -free in $\bar{\omega}$. This shows that the regularity claim of Theorem A reduces to the regularity claim of Theorem B.

4.2. \mathscr{B} -gradient setting. In this subsection, we first show the existence of a generalized minimizer for $(P\mathscr{B}_{\Omega})$ under quasiconvexity and coercivity assumptions.

Proposition 4.5. Suppose that $f: \Omega \times V \to \mathbb{R}$ is quasiconvex for \mathscr{B} -gradients $((\mathbf{H3})'$ with $\ell = 0)$ and satisfies $(\mathbf{H1})$ for $1 . Assume in addition that for any <math>u_0 \in \mathscr{D}'(\Omega, U)$ such that $\mathscr{B}u_0 \in L^p(\Omega, V)$ and $\phi \in C_c^{\infty}(\Omega, U)$, we have that

$$f_{\Omega} f(x, \mathscr{B}(u_0 + \phi)) \ge C \left(f_{\Omega} |\mathscr{B}\phi|^p - 1 - f_{\Omega} |\mathscr{B}u_0|^p \right).$$

Then Problem $(P\mathscr{B}_{\Omega})$ admits a generalized minimizer in the sense of Definition 4.1.

The proof is analogous to that of Proposition 4.2, establishing that

$$\inf\left\{\int_{\Omega} f(x,\mathscr{B}(u_0+u))\colon u\in C_c^{\infty}(\Omega,U)\right\} = \min\left\{\int_{\Omega} f(x,\mathscr{B}u_0+v)\colon v\in X_0^{\mathscr{B},p}(\Omega)\right\}.$$

We do not repeat the argument here.

Proof of Theorem C, *B*-gradient setting. Follows immediately from Proposition 4.5, Lemma 3.3, and Remark 3.2.

We will show in addition that any minimizer $\mathscr{B}u_0 + \tilde{v}$ has a better structure. In Proposition 4.8 we will deduce that, as far as problem $(P\mathscr{B}_{\Omega})$ is concerned, we can assume without loss of generality that Ω is smooth and $u_0 \in W^{k,p}(\Omega, U)$, reducing our setup of Theorem B to the one in [BDS15].

We begin by choosing a good representative \tilde{u} for $\tilde{v} = \mathscr{B}\tilde{u}$:

Proposition 4.6. Let 1 and <math>f satisfy the assumptions of Proposition 4.5. For any $u_0 \in \mathscr{D}'(\Omega, U)$ with $\mathscr{B}u_0 \in L^p(\Omega, V)$, there exists $\tilde{u}_0 \in \mathscr{D}'(\mathbb{R}^n, U)$ such that

- (a) $\mathscr{B}\tilde{u}_0$ is a generalized minimizer of $(P\mathscr{B}_{\Omega})$,
- (b) $\tilde{u}_0 u_0 \in W^{k,p}(\Omega, U),$
- (c) $\mathscr{C}^*(\tilde{u}_0 u_0) = 0.$

Proof. Let $u_0 + \varphi_j$ with $\varphi_j \in C_c^{\infty}(\Omega, U)$ be a minimizing sequence. By the coercivity assumption we have the L^p -boundedness of $\{\mathscr{B}\varphi_j\}$. Next, project φ_j to the kernel of \mathscr{C}^* as in Theorem 2.7(b) to get

$$\varphi_j = \tilde{\varphi}_j + \psi_j,$$

where $\mathscr{C}^* \tilde{\varphi}_j = 0$, $\mathscr{B} \tilde{\varphi}_j = \mathscr{B} \varphi_j$ in \mathbb{R}^n . The Sobolev estimate for $\tilde{\varphi}_j$ implies the boundedness of $\{\tilde{\varphi}_j\}$ in $W^{k,p}(\Omega, U)$. Then there exists $\pi \in W^{k,p}(\Omega, U)$ with $\mathscr{C}^* \pi = 0$ such that, up to a subsequence,

$$\tilde{\varphi}_j \rightharpoonup \pi \text{ in } W^{k,p}(\Omega, U).$$

Write $\tilde{u}_0 = u_0 + \pi$, and it is easy to see that $\mathscr{B}(u_0 + \varphi_j) = \mathscr{B}(u_0 + \bar{\varphi}_j) \rightharpoonup \mathscr{B}\tilde{u}_0$ in $L^p(\Omega, V)$. By Lemma 3.3

$$\liminf_{j \to \infty} \int_{\Omega} f(x, \mathscr{B}(u_0 + \tilde{\varphi}_j)) \ge \int_{\Omega} f(x, \mathscr{B}\tilde{u}_0).$$

Notice that the left-hand side converges to the infimum of $(P\mathscr{B}_{\Omega})$ since $\mathscr{B}\tilde{\varphi}_j = \mathscr{B}\varphi_j$, and $\mathscr{B}\tilde{u}_0 \in \mathscr{B}u_0 + X_0^{\mathscr{B},p}(\Omega)$, which indicates

$$\int_{\Omega} f(x, \mathscr{B}\tilde{u}_0) \, \mathrm{d}x \ge \liminf_{j \to \infty} \int_{\Omega} f(x, \mathscr{B}(u_0 + \tilde{\varphi}_j)) \, \mathrm{d}x,$$

and thus the equality holds. The claim follows.

Lemma 4.7. We follow the notation in Proposition 4.6. Let $\omega \in \Omega$, and then there holds $\inf \{ \mathcal{F}(u, \omega) \colon u \in \tilde{u}_0 + C_c^{\infty}(\omega, U) \} = \mathcal{F}(\tilde{u}_0, \omega).$

In other words, \tilde{u}_0 is a local minimizer of $(P\mathscr{B}_{\Omega})$.

In particular, to prove Theorem B it suffices to consider functionals defined on smooth sets.

Proof of Lemma 4.7. Fix an arbitrary $\varphi \in C_c^{\infty}(\omega, U)$. Since $u_0 + \pi$ is a generalized minimizer, and $\mathscr{B}(u_0 + \pi + \varphi) \in \mathscr{B}u_0 + X_0^{\mathscr{B},p}(\Omega)$, we have

$$\mathcal{F}(u_0 + \pi, \Omega) \leqslant \mathcal{F}(u_0 + \pi + \varphi, \Omega),$$

which implies

$$\int_{\omega} f(x, \mathscr{B}(u_0 + \pi)) \, \mathrm{d}x \leqslant \int_{\omega} f(x, \mathscr{B}(u_0 + \pi + \varphi)) \, \mathrm{d}x$$

as desired.

Finally, we show that on any subdomain $\omega \in \Omega$, we can replace the minimizer with a minimizer in the right Sobolev class, provided that we assume integrability of the lower order derivatives of u_0 .

Proposition 4.8. We use the notation of Proposition 4.6 and denote $u_0 + \pi$ by \tilde{u}_0 . Assume that $u_0 \in W^{k-1,p}(\Omega, U)$. For any $\omega \Subset \Omega$, there exists $u_\omega \in W^{k,p}(\Omega, U)$ such that

(a)
$$\mathscr{B}u_{\omega} = \mathscr{B}\tilde{u}_0$$
 in ω

(b)
$$\mathscr{C}^* u_{\omega} = 0$$
 in Ω .

- (c) $\|u_{\omega}\|_{W^{k,p}(\Omega)} \leq c \|\mathscr{B}u_0\|_{L^p(\Omega)} + \|u_0\|_{W^{k-1,p}(\Omega)},$
- (d) u_{ω} is a minimizer of the problem

$$\inf\{\mathcal{F}(u,\omega)\colon u\in u_{\omega}+C_c^{\infty}(\omega)\}=\mathcal{F}(u_{\omega},\omega).$$

Proof. Consider a cut-off $\rho \in C_c^{\infty}(\Omega)$ such that $\rho = 1$ in ω . We now apply Theorem 2.7(b) to decompose $\rho \tilde{u}_0$ as follows:

$$\rho \tilde{u}_0 = u_\omega + \psi,$$

where $u_{\omega} \in \dot{W}^{k,p}(\mathbb{R}^n, U)$ such that $\mathscr{C}^* u_{\omega} = 0$ and $\mathscr{B} u_{\omega} = \mathscr{B}(\rho \tilde{u}_0) \in L^p(\mathbb{R}^n)$. The third assertion follows from the estimate of Theorem 2.7(b) and the product rule. The last claim follows from $\mathscr{B} u_{\omega} = \mathscr{B} \tilde{u}_0$ in ω and Lemma 4.7.

In particular, the conclusion of this subsection is that we can assume that Ω is smooth, $u_0 \in W^{k,p}(\Omega, U)$, and that minimizers are \mathscr{C}^* -free in the statement of Theorem B.

5. Local linear estimates

5.1. Local Korn-type inequality. We will establish some linear estimates for the minimizers of Problem $(P\mathscr{B}_{\Omega})$ which are a generalization of Lemma 2.10 in [Rai19a]. Below, \mathscr{B} and \mathscr{C} are as in Theorem 2.1 so k, l are the orders of \mathscr{B}, \mathscr{C} respectively.

Proposition 5.1. Suppose that $\Omega \subset \mathbb{R}^n$ is a bounded open set, and $v \in \mathscr{D}'(\Omega)$ satisfies

$$\mathscr{B}v \in L^p_{\mathrm{loc}}(\Omega) \quad and \quad \mathscr{C}^*v \in \dot{W}^{-q,p}_{\mathrm{loc}}(\Omega)$$

for some 1 ; let <math>q = l - k. Then we have $v \in W^{k,p}_{\text{loc}}(\Omega)$, and for any open subsets $\omega \in \Omega' \in \Omega$, there holds

(5.1)
$$\|D^{k}v\|_{L^{p}(\omega)} \leq C(\|\mathscr{B}v\|_{L^{p}(\Omega')} + \|\mathscr{C}^{*}v\|_{\dot{W}^{-q,p}(\Omega')} + \|v\|_{W^{k-1,p}(\Omega')})$$

with $C = C(n, p, \mathscr{B}, \mathscr{C}, \omega, \Omega') > 0$. In particular, if $\mathscr{C}^* v = 0$ in Ω , for any ball $B_R = B(x_0, R) \in \Omega$ and any $\tau \in (0, 1)$, we have

(5.2)
$$\int_{B_{\tau R}} |D^k v|^p \, \mathrm{d}x \leqslant C \left(\int_{B_R} |\mathscr{B}v|^p \, \mathrm{d}x + \sum_{i=0}^{k-1} \int_{B_R} \frac{|D^i v|^p}{R^{p(k-i)}} \, \mathrm{d}x \right)$$
$$\int_{B_{\tau R}} |V_p (D^k v)|^2 \, \mathrm{d}x \leqslant C \left(\int_{B_R} |V_p (\mathscr{B}v)|^2 \, \mathrm{d}x + \sum_{i=0}^{k-1} \int_{B_R} \left| V_p \left(\frac{D^i v}{R^{k-i}} \right) \right|^2 \, \mathrm{d}x \right)$$

with $C = C(n, p, \mathscr{B}, \tau) > 0$.

Proof. Take any subsets ω and Ω' as in the assumption, and a cut-off function $\rho \in C_c^{\infty}(\Omega')$ with

$$0 \leq \rho \leq 1, \quad \rho \equiv 1 \text{ in } \omega.$$

Define $w := \rho v$ and an operator $\mathscr{P} := (D^q \mathscr{B}, \mathscr{C}^*)$ as in Subsection 5.2.

Given any multi-index α with $|\alpha| = k + q - 1 = l - 1$, we have

$$\begin{split} \widehat{\partial^{\alpha}w}(\xi) &= (-i)^{k+q-1}\xi^{\alpha}\widehat{w}(\xi) \\ &= (-i)^{k+q-1}\xi^{\alpha}\mathscr{P}^{\dagger}(\xi)\mathscr{P}(\xi)\widehat{w}(\xi) \\ &= i\xi^{\alpha}|\xi|^{-(k+q)}\mathscr{P}^{\dagger}\left(\frac{\xi}{|\xi|}\right)\widehat{\mathscr{P}w}(\xi) \\ &=: m(\xi)\widehat{\mathscr{P}w}(\xi), \end{split}$$

where $\mathscr{P}^{\dagger}(\xi)$ is the left-inverse of $\mathscr{P}(\xi)$, and the multiplier $m(\xi)$ is smooth away from the origin and (-1)-homogeneous. Let $K := \mathscr{F}^{-1}m$, which is (1-n)-homogeneous and integrable. Then the partial derivative $\partial^{\alpha}(\rho w)$ can be expressed as follows:

$$\partial^{\alpha}w = \partial^{\alpha}(\rho v) = K * (\mathscr{P}(\rho v)) = K * (\rho \mathscr{P}v) + \sum_{j=1}^{k+q} K * (b_j[D^j\rho, D^{k+q-j}v]),$$

where $\{b_j[\cdot, \cdot]\}_{j=1}^{k+q}$ are bilinear forms with constant coefficients. Notice that $D^j \rho \equiv 0$ on ω , and thus

sing spt
$$(b_j[D^j\rho, D^{k+q-j}v]) \subset \mathbb{R}^n \setminus \omega$$
,

and we have that the terms in the sum are smooth on ω . To control the first term, notice that for any $\varphi \in C_c^{\infty}(\omega)$ there holds

$$\langle K * (\rho \mathscr{P}v), \varphi \rangle = \langle \mathscr{P}v, \rho(K' * \varphi) \rangle$$

= $(-1)^q \langle \mathscr{B}v, (\operatorname{div})^q (\rho(K' * \varphi)) \rangle + \langle \mathscr{C}^*v, \rho(K' * \varphi) \rangle,$

where $K'(\cdot) = K(-\cdot)$ is the dual convolution kernel of K. The second term can be easily controlled as follows

$$\left\langle \mathscr{C}^* v, \rho(K' * \varphi) \right\rangle \leqslant \|\mathscr{C}^* v\|_{\dot{W}^{-q,p}(\Omega')} \|\rho(K' * \varphi)\|_{\dot{W}^{q,p'}(\Omega')} \\ \leqslant C \|\mathscr{C}^* v\|_{\dot{W}^{-q,p}(\Omega')} \|K' * \varphi\|_{W^{q,p'}(\Omega')}.$$

By Lemma 7.12 in [GT01], we have

$$\|K' * \varphi\|_{L^{p'}(\Omega')} \leqslant C(n) \mathscr{L}^n(\Omega')^{\frac{1}{n}} \|\varphi\|_{L^{p'}(\Omega')},$$

and the estimate of the higher-order terms requires Theorem 2.6:

$$\sum_{i=1}^{q} \|K' * \varphi\|_{\dot{W}^{i,p'}(\Omega')} \leqslant \sum_{i=1}^{q} \|K' * \varphi\|_{\dot{W}^{i,p'}(\mathbb{R}^n)} \leqslant C \sum_{i=1}^{q} \|\varphi\|_{\dot{W}^{i-1,p'}(\Omega')}$$

This implies

$$\|K' * \varphi\|_{W^{q,p'}(\Omega')} \leq C \|\varphi\|_{W^{q-1,p'}(\Omega')}$$

and to obtain $\partial^{\alpha} v \in W^{-(q-1),p}_{\text{loc}}(\Omega)$, we need to estimate the $L^{p'}$ -norm of $(\text{div})^q(\rho(K' * \varphi))$. This term expands as

$$(\operatorname{div})^{q}(\rho(K' * \varphi)) = \sum_{i=0}^{q} c_{i}[D^{i}\rho, D^{q-i}(K' * \varphi)],$$

where $\{c_i[\cdot, \cdot]\}_{i=0}^q$ are bilinear forms with constant coefficients. The terms with i < q can be estimated as follows

$$\begin{split} \left\| \sum_{i=0}^{q-1} c_i [D^i \rho, D^{q-i}(K' * \varphi)] \right\|_{L^{p'}(\Omega')} &\leq C \sum_{i=0}^{q-1} \sup |D^i \rho| \|K' * \varphi\|_{\dot{W}^{q-i,p'}(\mathbb{R}^n)} \\ &\leq C \sum_{i=0}^{q-1} \|\varphi\|_{\dot{W}^{q-i-1,p'}(\mathbb{R}^n)} = C \|\varphi\|_{W^{q-1,p'}(\omega)} \end{split}$$

where the fourth line follows from Theorem 2.6. When i = q, we apply Lemma 7.12 in [GT01] to estimate the corresponding term

$$\langle \mathscr{B}v, (D^{q}\rho)K' * \varphi \rangle \leqslant \|\mathscr{B}v\|_{L^{p}(\Omega')} \| (D^{q}\rho)K' * \varphi \|_{L^{p'}(\Omega')} \leqslant \|\mathscr{B}v\|_{L^{p}(\Omega')} \sup |D^{q}\rho| \|K' * \varphi \|_{L^{p'}(\Omega')} \leqslant C(n)\mathscr{L}^{n}(\Omega')^{\frac{1}{n}} \|\mathscr{B}v\|_{L^{p}(\Omega')} \|\varphi\|_{L^{p'}(\omega)}.$$

From the above, we know that $D^{k+q-1}v \in W_{\text{loc}}^{-(q-1),p}(\Omega)$. Then with a procedure similar to the one above and \mathscr{P} replaced by D^{k+q-i} , we can show inductively that $D^{k+q-i-1}v \in W_{\text{loc}}^{-(q-i-1),p}(\Omega)$ for $i = 1, \ldots, q-1$. Furthermore, with \mathscr{P} replaced by D^{i+1} , we can show $D^iv \in W_{\text{loc}}^{i,p}(\Omega)$ for $i = k-1, \ldots, 0$ with only applying Lemma 7.12 in [GT01] in the last step.

Having established that $v \in W_{\text{loc}}^{k,p}(\Omega)$, we are now in the position to show the inequality (5.1). Take ρ as above and a multi-index α with $|\alpha| = k$, and define again $w = \rho v$. Then we have

$$\widehat{\partial^{\alpha} w}(\xi) = m_{\alpha}(\xi) \widehat{\mathscr{P} w}(\xi),$$

where m_{α} is a (-q)-homogeneous multiplier. Then by Theorem 2.6 we have the following global estimate on \mathbb{R}^n :

(5.3)
$$\begin{aligned} \|\partial^{\alpha}w\|_{L^{p}(\mathbb{R}^{n})} \leqslant C\|I^{-q}\mathscr{P}w\|_{L^{p}(\mathbb{R}^{n})} \leqslant C\|\mathscr{P}w\|_{\dot{W}^{-q,p}(\mathbb{R}^{n})}\\ \leqslant C(\|\mathscr{B}w\|_{L^{p}(\mathbb{R}^{n})} + \|\mathscr{C}^{*}w\|_{\dot{W}^{-q,p}(\mathbb{R}^{n})}).\end{aligned}$$

Since $\|\partial^{\alpha}v\|_{L^{p}(\omega)} \leq \|\partial^{\alpha}w\|_{L^{p}(\mathbb{R}^{n})}$, it is sufficient to estimate the right-hand side of the above inequality.

It is easy to see that

$$|\mathscr{B}w||_{L^{p}(\mathbb{R}^{n})} = ||\mathscr{B}(\rho v)||_{L^{p}(\mathbb{R}^{n})} \leq ||\mathscr{B}v||_{L^{p}(\Omega')} + C||v||_{W^{k-1,p}(\Omega')}$$

The other term can be controlled by

$$\|\mathscr{C}^*w\|_{\dot{W}^{-q,p}(\mathbb{R}^n)} \leqslant \|\rho\mathscr{C}^*v\|_{\dot{W}^{-q,p}(\Omega')} + \sum_{i=1}^{l} \|d_i[D^i\rho, D^{l-i}v]\|_{\dot{W}^{-q,p}(\Omega')},$$

where $\{d_i[\cdot,\cdot]\}_{i=1}^l$ are bilinear forms with constant coefficients. We test these terms with any $\varphi \in C_c^{\infty}(\Omega')$ separately:

$$\begin{aligned} \langle \rho \mathscr{C}^* v, \varphi \rangle &= \langle \mathscr{C}^* v, \rho \varphi \rangle \leqslant \| \mathscr{C}^* v \|_{\dot{W}^{-q,p}(\Omega')} \| \rho \varphi \|_{\dot{W}^{q,p'}(\Omega')} \\ &\leqslant C \| \mathscr{C}^* v \|_{\dot{W}^{-q,p}(\Omega')} \| \varphi \|_{W^{q,p'}(\Omega')} \leqslant C \| \mathscr{C}^* v \|_{\dot{W}^{-q,p}(\Omega')} \| \varphi \|_{\dot{W}^{q,p'}(\Omega')}, \end{aligned}$$

where the last inequality follows from Poincaré's inequality;

$$\begin{split} \left\langle c_i[D^i\rho, D^{l-i}v], \varphi \right\rangle &= \left\langle D^{l-i}v, c_i'[D^i\rho, \varphi] \right\rangle \\ &= (-1)^{l-i-t_i} \left\langle D^{t_i}v, (\operatorname{div})^{l-i-t_i}c_i'[D^i\rho, \varphi] \right\rangle \\ &\leqslant C \|D^{t_i}v\|_{L^p(\Omega')} \|\varphi\|_{W^{q,p'}(\Omega')} \\ &\leqslant C \|D^{t_i}v\|_{L^p(\Omega')} \|\varphi\|_{\dot{W}^{q,p'}(\Omega')}, \end{split}$$

where $\{c'_i[\cdot, \cdot]\}_{i=1}^l$ are also bilinear forms with constant coefficients and $t_i := \min\{l-i, k-1\}$. This proves (5.1), which immediately implies the L^p -inequality in (5.2).

To prove the second estimate in (5.2), we argue as follows. It suffices to prove that

$$\int_{\omega} |V_p(D^k v)|^2 \,\mathrm{d}x \leqslant C\left(\int_{\Omega'} |V_p(\mathscr{B}v)|^2 \,\mathrm{d}x + \sum_{i=0}^{k-1} \int_{\Omega'} |V_p(D^i v)|^2 \,\mathrm{d}x\right)$$

provided that $\mathscr{C}^* v = 0$, and the inequality on the concentric balls $B_{\tau R}$ and B_R can be done via scaling from B_{τ} and B_1 . We note that $t \mapsto (1+t^2)^{(p-2)/2}t^2$ is a Young function which satisfies both the Δ_2 and ∇_2 conditions [RR02], so we can use the extrapolation inequality from Lemma 2.2 in [CG22]: it suffices to verify this inequality in weighted Lebesgue spaces with respect to Muckenhoupt weights $\mu \in A_p$ for a certain $p \in (1, \infty)$:

$$\int_{\omega} |D^{k}v|^{p} \mu \,\mathrm{d}x \leqslant C \left(\int_{\Omega'} |\mathscr{B}v|^{p} \mu \,\mathrm{d}x + \sum_{i=0}^{k-1} \int_{\Omega'} |D^{i}v|^{p} \mu \,\mathrm{d}x \right)$$

This inequality follows along the same lines as the proof of (5.1) (when $\mu = 1$): the boundedness in (5.3) follows since Calderón–Zygmund operators are bounded on $L^p(d\mu)$; the duality argument rests on the following Fourier characterization of $W^{a,p}(d\mu)$:

$$\|u\|_{(\dot{W}^{a,p}(\mathrm{d}\mu))^*} \sim \|\mathcal{F}^{-1}(|\xi|^a \hat{u})\|_{L^{p'}(\mathrm{d}\mu)},$$

with p' = p/(p-1). This is clear for a = 0 (duality of Lebesgue space with respect to arbitrary measures), follows by boundedness of Calderón–Zygmund operators for a > 0, and by duality for a < 0. The proof is complete.

5.2. Estimates for linear systems. We investigate the regularity of the linear system

$$\mathscr{B}^*(A\mathscr{B}h) = 0, \, \mathscr{C}^*h = 0.$$

The following estimate is in a similar spirit to those for linear elliptic systems (see, for example, [Gia83] Chapter III) while also requiring the Korn-type inequality (5.2) since we

are dealing with constant rank operators instead of elliptic. We will use this in Section 7, where the main partial regularity claim is proved.

Theorem 5.2. Suppose that \mathscr{B} and \mathscr{C} are as above, and $\Omega \subset \mathbb{R}^n$ is an open set. If $h \in \mathscr{D}'(\Omega)$ satisfies the following

(5.4)
$$\mathscr{B}^*(A\mathscr{B}h) = 0, \quad \mathscr{C}^*h = 0$$

in Ω , where $A: V \times V \to \mathbb{R}$ is a symmetric linear operator with constant coefficients and satisfies

$$A[z, z] \ge \lambda |z|^2$$
, $|A[z, w]| \le \Lambda |z||w|$, for any $z, w \in V$

with $0 < \lambda \leq \Lambda < \infty$. Then we have $h \in C^{\infty}(\Omega)$ and the following estimate

(5.5)
$$r \sup_{B_r} |D^{k+1}h| \leq C \oint_{B_{2r}} |D^k h - b| \,\mathrm{d}x$$

for any constant tensor field b and any ball B_r with $B_{2r} \subseteq \Omega$ and 0 < r < 1, where $C = C(n, \dim V, \frac{\Lambda}{\lambda}) > 0$.

Proof. With out loss of generality, we can assume A = I, since the general case can be done by replacing \mathscr{B} with $A^{\frac{1}{2}}\mathscr{B}$. We consider the operator $\mathscr{P} := (D^q \mathscr{B}, \mathscr{C}^*)$. It is easy to see that \mathscr{P} is an elliptic operator of order l, and from (5.4) we know that h satisfies

(5.6)
$$\mathscr{P}^*\mathscr{P}h = 0$$
 in Ω

in the sense of distribution. The operator $\mathscr{P}^*\mathscr{P}$ is elliptic, and thus we have the smoothness of h in Ω .

Take a ball B_r with $B_{2r} \in \Omega$, and a k-homogeneous polynomial P such that

$$(D^{i}P)_{2r} = (D^{i}h)_{2r}, i = 0, 1, \dots, k-1, \text{ and } D^{k}P = b.$$

where b is any vector as in (5.5). The function h - P satisfies the first system in (5.4) as well, and we test it with $-\partial^s(\rho^{2k}\partial^s(h-P))$, where $s \in \{1, \ldots, n\}$ and $\rho \in C_c^{\infty}(B_{2r})$ satisfies

$$0 \leq \rho \leq 1$$
, $\rho \equiv 1$ in B_r , and $|D^i \rho| \leq Cr^{-i}$, $i = 1, \dots, k$.

This test function gives

$$\int_{\Omega} [\rho^k \partial^s \mathscr{B}(h-P), \rho^k \partial^s \mathscr{B}(h-P)] \, \mathrm{d}x \leqslant \int_{\Omega} [\rho^k \partial^s \mathscr{B}(h-P), \sum_{i=1}^k R_i] \, \mathrm{d}x,$$

where R_i is a linear combination of products of the *i*-th order derivatives of ρ^k and the (k-i)-th order derivatives of $\partial^s(h-P)$. Thus, by the Cauchy-Schwarz inequality we have

$$\int_{B_{2r}} [\rho^k \partial^s \mathscr{B}(h-P), \rho^k \partial^s \mathscr{B}(h-P)] \,\mathrm{d}x$$

$$\leqslant \frac{1}{2} \int_{B_{2r}} [\rho^k \partial^s \mathscr{B}(h-P), \rho^k \partial^s \mathscr{B}(h-P)] \,\mathrm{d}x + C \sum_{i=1}^k \frac{1}{r^{2i}} \int_{B_{2r}} |\partial^s D^{k-i}(h-P)|^2 \,\mathrm{d}x$$

which furthermore implies, with Poincaré's inequality and the choice of P,

$$\int_{B_r} |\partial^s \mathscr{B}h|^2 \,\mathrm{d}x \leqslant \int_{B_{2r}} [\rho^k \partial^s \mathscr{B}(h-P), \rho^k \partial^s \mathscr{B}(h-P)] \,\mathrm{d}x \leqslant \frac{C}{r^2} \int_{B_{2r}} |D^k(h-P)|^2 \,\mathrm{d}x.$$

Then we can conclude that

$$\int_{B_r} |D\mathscr{B}h|^2 \,\mathrm{d}x \leqslant \frac{C}{r^2} \int_{B_{2r}} |D^k h - b|^2 \,\mathrm{d}x$$

It is actually possible to estimate $D^{k+1}h$: Apply Proposition 5.1 to $\partial^s(h-P), s = 1, \ldots, n$, and then we obtain

$$\begin{split} \int_{B_{\frac{r}{2}}} |D^{k+1}h|^2 \, \mathrm{d}x &\leq C \left(\int_{B_r} |D\mathscr{B}h|^2 \, \mathrm{d}x + \sum_{i=0}^{k-1} \int_{B_r} \frac{|D^i D(h-P)|^2}{r^{2(k-i)}} \, \mathrm{d}x \right) \\ &\leq \frac{C}{r^2} \int_{B_{2r}} |D^k h - b|^2 \, \mathrm{d}x, \end{split}$$

where the second line is obtained with Poincaré's inequality.

Notice that $\partial^s h, s = 1, \dots, n$, also solves the first system in (5.4), and with an argument similar to the one above we obtain

$$\int_{B_{2\sigma^2 r}} |D^{k+2}h|^2 \, \mathrm{d}x \leq \frac{C(\sigma)}{r^4} \int_{B_{2r}} |D^k h - b|^2 \, \mathrm{d}x$$

for any $\sigma \in (0,1)$. Take $d \in \mathbb{N}$ such that 2d > n, and then $\sigma \in (0,1)$ with $\sigma^{d+1} \ge \frac{1}{2}$. Morrey's inequality then implies the following estimate

$$r^{n} \sup_{B_{r}} |D^{k+1}h|^{2} \leq C \sum_{i=0}^{d} \int_{B_{r}} r^{2i} |D^{k+i+1}h|^{2} \, \mathrm{d}x \leq \frac{C}{r^{2}} \int_{B_{2r}} |D^{k}h - b|^{2} \, \mathrm{d}x.$$

The desired estimate (5.5) can be further concluded as in Corollary 7.1 and Theorem 7.3 in [Giu03]. $\hfill \Box$

6. Higher integrability

The purpose of this section is to establish higher integrability for generalized minimizers of $(\mathcal{P}\mathscr{A}_{\Omega})$ and $(\mathcal{P}\mathscr{B}_{\Omega})$. We follow the commonly used strategy for proving higher integrability ([Gia83], Chapter V), which involves first showing a Caccioppolli-type inequality and then the application of Poincaré's inequality and Gehring's lemma. However, the Korntype inequality (5.1) is required since the constant operator \mathscr{B} is here in the place of the gradient operator.

6.1. Caccioppolli-type inequalities. We prove a Caccioppolli-type inequality, which applies to non-autonomous integrands and arbitrary constant rank operators.

Proposition 6.1. Suppose that the integrand f satisfies assumptions(**H1**)-(**H3**) and that $u \in W^{k-1,p}(\Omega, U)$ such that $\mathcal{B}u \in L^p(\Omega, V)$ is a generalized minimizer of $(P\mathcal{B}_{\Omega})$, and the map $a: \Omega \to V$ is such that $\mathcal{B}a$ is constant and $|\mathcal{B}a| \leq M$ for some M > 0. Then for any $\tau \in (0,1)$ there exists a constant $C = C(n, \dim V, \mathcal{B}, M, p, L, \ell, \tau) > 0$ such that the following estimate holds true for any ball $B(x_0, R)(=: B_R) \in \Omega$:

$$(6.1) \quad \int_{B_{\tau R}} |V_p(\mathscr{B}(u-a))|^2 \,\mathrm{d}x \leqslant C \sum_{i=0}^{k-1} \int_{B_R} \left| V_p\left(\frac{D^i(u-a)}{R^{k-i}}\right) \right|^2 \,\mathrm{d}x \\ + CR \sum_{i=0}^{k-1} \int_{B_R} \left\langle \frac{D^i(u-a)}{R^{k-i}} \right\rangle^{p-1} \frac{|D^i(u-a)|}{R^{k-i}} \,\mathrm{d}x + CR \int_{B_R} \left\langle \mathscr{B}(u-a) \right\rangle^{p-1} |\mathscr{B}(u-a)| \,\mathrm{d}x.$$

Proof. Fix a ball $B_R \Subset \Omega$ and r, s > 0 such that $\frac{R}{2} < r < s < R$. Take a cut off function $\rho \in C_c^{\infty}(B_s)$ with

 $0 \leq \rho \leq 1$, $\rho \equiv 1$ in B_r , $|D^i \rho| \leq c_i (s-r)^{-i}, i = 1, \dots, k$.

Suppose that the map $a: \Omega \to V$ is as in the assumption.

Z. LI AND B. RAIȚĂ

Then define $\tilde{f}(z) := f(x_0, \mathscr{B}a + z) - f(x_0, \mathscr{B}a) - f_z(x_0, \mathscr{B}a) \cdot z$, $\tilde{u} := u - a$, $\varphi := \rho \tilde{u}$, and $\psi := (1 - \rho)\tilde{u} = \tilde{u} - \varphi$. From the strong quasiconvexity of f and the estimate (2.5), we have

$$\begin{split} \int_{B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x &\leqslant \int_{B_s} |V_p(\mathscr{B}\varphi)|^2 \, \mathrm{d}x \\ &\lesssim_{M,p,\ell} \int_{B_s} (f(x_0,\mathscr{B}(\varphi+a)) - f(x_0,\mathscr{B}a) - f_z(x_0,\mathscr{B}a) \cdot \mathscr{B}\varphi) \, \mathrm{d}x \\ &= \int_{B_s} \tilde{f}(\mathscr{B}\varphi) \, \mathrm{d}x = \int_{B_s} \tilde{f}(\mathscr{B}\tilde{u} - \mathscr{B}\psi) \, \mathrm{d}x \\ &= \int_{B_s} \tilde{f}(\mathscr{B}\tilde{u}) \, \mathrm{d}x + \int_{B_s} (\tilde{f}(\mathscr{B}\tilde{u} - \mathscr{B}\psi) - \tilde{f}(\mathscr{B}\tilde{u})) \, \mathrm{d}x. \end{split}$$

The definition of \tilde{f} and the minimality of u then imply

$$\begin{split} \int_{B_s} \tilde{f}(\mathscr{B}\tilde{u}) \, \mathrm{d}x &= \int_{B_s} f(x, \mathscr{B}u) \, \mathrm{d}x + \int_{B_s} (\tilde{f}(\mathscr{B}\tilde{u}) - f(x, \mathscr{B}u)) \, \mathrm{d}x \\ &\leqslant \int_{B_s} f(x, \mathscr{B}u - \mathscr{B}\varphi) \, \mathrm{d}x + \int_{B_s} (\tilde{f}(\mathscr{B}\tilde{u}) - f(x, \mathscr{B}u)) \, \mathrm{d}x. \end{split}$$

The first term in the second line is further controlled as follows:

$$\begin{split} \int_{B_s} f(x, \mathscr{B}u - \mathscr{B}\varphi) \, \mathrm{d}x &= \int_{B_s} f(x, \mathscr{B}\psi + \mathscr{B}a) \, \mathrm{d}x \\ &= \int_{B_s} f(x_0, \mathscr{B}\psi + \mathscr{B}a) \, \mathrm{d}x + \int_{B_s} (f(x, \mathscr{B}\psi + \mathscr{B}a) - f(x_0, \mathscr{B}\psi + \mathscr{B}a)) \, \mathrm{d}x \\ &= \int_{B_s} (\tilde{f}(\mathscr{B}\psi) + f(x_0, \mathscr{B}a) + f_z(x_0, \mathscr{B}a) \cdot \mathscr{B}\psi) \, \mathrm{d}x \\ &+ \int_{\mathscr{B}_s} (f(x, \mathscr{B}\psi + \mathscr{B}a) - f(x_0, \mathscr{B}\psi + \mathscr{B}a)) \, \mathrm{d}x. \end{split}$$

Combining the three estimates above, we obtain

(6.2)

$$\int_{B_{r}} |V_{p}(\mathscr{B}\tilde{u})|^{2} dx \leq C \int_{B_{s}} \tilde{f}(\mathscr{B}\varphi) dx$$

$$\leq C \left(\int_{B_{s}} \tilde{f}(\mathscr{B}\psi) dx + \int_{B_{s}} (f(x, \mathscr{B}\psi + \mathscr{B}a) - f(x_{0}, \mathscr{B}\psi + \mathscr{B}a)) dx$$

$$+ \int_{B_{s}} (f(x_{0}, \mathscr{B}u) - f(x, \mathscr{B}u)) dx + \int_{B_{s}} (\tilde{f}(\mathscr{B}\tilde{u} - \mathscr{B}\psi) - \tilde{f}(\mathscr{B}\tilde{u})) dx$$

$$=: I + II + III + IV.$$

By (2.3) we know

(6.3)

$$I \leq C \int_{B_s \setminus B_r} |V_p(\mathscr{B}\psi)|^2 \,\mathrm{d}x$$

$$\leq C \int_{B_s \setminus B_r} |V_p(\mathscr{B}\tilde{u}|^2 \,\mathrm{d}x + C \sum_{i=1}^k \int_{B_s} \left| V_p\left(\frac{D^{k-i}\tilde{u}}{(s-r)^i}\right) \right|^2 \,\mathrm{d}x.$$

The terms II and III can be controlled with the continuity of $\partial_z f$ in x as assumed in (H2):

$$II + III = -\int_{B_s} \int_0^1 (\partial_z f(x, \mathscr{B}u - t\mathscr{B}\varphi) - \partial_z f(x_0, \mathscr{B}u - t\mathscr{B}\varphi)) \cdot \mathscr{B}\varphi \,\mathrm{d}t$$

$$(6.4) \qquad \leqslant C \int_{B_s} \int_0^1 |x - x_0| (1 + |\mathscr{B}u - t\mathscr{B}\varphi|^2)^{\frac{p-1}{2}} |\mathscr{B}\varphi| \,\mathrm{d}x \\ \leqslant Cs \int_{B_s} (1 + |\mathscr{B}\tilde{u}|^2 + |\mathscr{B}\varphi|^2)^{\frac{p-1}{2}} |\mathscr{B}\varphi| \,\mathrm{d}x \\ \leqslant Cs \int_{B_s} \langle \mathscr{B}\tilde{u} \rangle^{p-1} \, |\mathscr{B}\tilde{u}| \,\mathrm{d}x + Cs \sum_{i=0}^{k-1} \int_{B_s} \left\langle \frac{D^i \tilde{u}}{(s-r)^{k-i}} \right\rangle^{p-1} \frac{|D^i \tilde{u}|}{(s-r)^{k-i}} \,\mathrm{d}x.$$

Notice that the integrand in IV vanishes on B_r , and by (2.4) we have

(6.5)
$$IV \leqslant C \int_{B_s \setminus B_r} (1 + |\mathscr{B}\psi|^2 + |\mathscr{B}\tilde{u}|^2)^{\frac{p-2}{2}} (|\mathscr{B}\psi| + |\mathscr{B}\tilde{u}|) |\mathscr{B}\psi| \, \mathrm{d}x$$
$$\leqslant C \int_{B_s \setminus B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x + C \sum_{i=1}^k \int_{B_s} \left| V_p\left(\frac{D^{k-i}\tilde{u}}{(s-r)^i}\right) \right|^2 \, \mathrm{d}x.$$

Then (6.2)-(6.5) together imply

$$\begin{split} \int_{B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x &\leq C \int_{B_s \setminus B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x + C \sum_{i=1}^k \int_{B_s} \left| V_p\left(\frac{D^{k-i}\tilde{u}}{(s-r)^i}\right) \right|^2 \, \mathrm{d}x \\ &+ Cs\left(\int_{B_s} \left\langle \mathscr{B}\tilde{u} \right\rangle^{p-1} |\mathscr{B}\tilde{u}| \, \mathrm{d}x + \sum_{i=0}^{k-1} \int_{B_s} \left\langle \frac{D^i\tilde{u}}{(s-r)^{k-i}} \right\rangle^{p-1} \frac{|D^i\tilde{u}|}{(s-r)^{k-i}} \, \mathrm{d}x \right), \end{split}$$

and we add $C \int_{B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x$ to both sides to get

$$\begin{split} \int_{B_r} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x &\leqslant \frac{C}{C+1} \int_{B_s} |V_p(\mathscr{B}\tilde{u})|^2 \, \mathrm{d}x + \frac{C}{C+1} \sum_{i=1}^k \int_{B_s} \left| V_p\left(\frac{D^{k-i}\tilde{u}}{(s-r)^i}\right) \right|^2 \, \mathrm{d}x \\ &+ \frac{C}{C+1} s \left(\int_{B_s} \langle \mathscr{B}\tilde{u} \rangle^{p-1} \left| \mathscr{B}\tilde{u} \right| \, \mathrm{d}x + \sum_{i=0}^{k-1} \int_{B_s} \left\langle \frac{D^i\tilde{u}}{(s-r)^{k-i}} \right\rangle^{p-1} \frac{|D^i\tilde{u}|}{(s-r)^{k-i}} \, \mathrm{d}x \right). \end{split}$$

The desired inequality (6.1) follows from Lemma 2.10 with

$$\begin{split} \Phi(r) &= \int_{B_r} |V_p(\mathscr{B}\tilde{u})|^2 \,\mathrm{d}x, \\ \Psi_i(t) &= \int_{B_R} \left| V_p\left(\frac{D^{k-i}\tilde{u}}{(s-r)^i}\right) \right|^2 \mathrm{d}x + R \int_{B_s} \left\langle \frac{D^i\tilde{u}}{(s-r)^{k-i}} \right\rangle^{p-1} \frac{|D^i\tilde{u}|}{(s-r)^{k-i}} \,\mathrm{d}x, \\ B &= CR \int_{B_R} \left\langle \mathscr{B}\tilde{u} \right\rangle^{p-1} |\mathscr{B}\tilde{u}| \,\mathrm{d}x, \end{split}$$

which concludes the proof.

We have the following improved Caccioppolli-type inequality for good minimizers u of $(P\mathscr{B}_{\Omega})$, which follows from Proposition 6.1 and inequality (5.2).

Proposition 6.2. Let an integrand f satisfy (H1)-(H3) and $u \in W^{\mathscr{B},p}(\Omega,U)$ be such that $\mathscr{B}u$ is a generalized minimizer of $(P\mathscr{B}_{\Omega})$ satisfying $\mathscr{C}^*u = 0$. Take a map $a \colon \Omega \to V$ with $\mathscr{B}a$ being constant and $|\mathscr{B}a| \leq M$ for some M > 0. Then for any $\tau \in (0,1)$ there exists a constant $c_{M,\tau} = c(n, \dim V, \mathscr{B}, M, p, L, \ell, \tau) > 0$ such that the following estimate holds true for any ball $B(x_0, R)(=: B_R) \subseteq \Omega$:

(6.6)
$$\int_{B_{\tau R}} |V_p(D^k(u-a))|^2 \, \mathrm{d}x \leqslant c_{M,\tau} \left(\sum_{i=0}^{k-1} \int_{B_R} \left| V_p\left(\frac{D^i(u-a)}{R^{k-i}}\right) \right|^2 \, \mathrm{d}x \right)$$

Z. LI AND B. RAIȚĂ

$$+R\sum_{i=0}^{k-1}\int_{B_R}\left\langle\frac{D^i(u-a)}{R^{k-i}}\right\rangle^{p-1}\frac{|D^i(u-a)|}{R^{k-i}}\,\mathrm{d}x+R\int_{B_R}\left\langle\mathscr{B}(u-a)\right\rangle^{p-1}|\mathscr{B}(u-a)|\,\mathrm{d}x\right).$$

Remark 6.3. Notice that Proposition 6.1 and 6.2 also hold true for (good) minimizers of $(P\mathscr{A}_{\Omega})$, which follows from Remark 4.4. The extra function ϕ can be considered as part of the x-dependence.

6.2. Higher integrability for minimizers. The Caccioppolli-type inequality (6.6) implies higher integrability of minimizers of our variational problems $(P\mathscr{B}_{\Omega})$ and $(P\mathscr{A}_{\Omega})$. The proof in the case 1 requires the following Poincaré inequality with respect to V. It will also be combined with the Caccioppolli inequality (6.6) in the excess decay estimate in Subsection 7.2 and can be shown by modifying Theorem 2.4 in [CFM98].

Theorem 6.4. If $1 , there exist <math>\alpha \in (\frac{2}{p}, 2)$ and $\sigma > 0$ such that any map $u \in W^{1,p}(B_{3R}, V)$ satisfies

$$\sum_{i=0}^{k-1} \left(\oint_{B_R} \left| V\left(\frac{D^i u - (D^i u)_R}{R^{k-i}} \right) \right|^{2(1+\sigma)} \mathrm{d}x \right)^{1/(2(1+\sigma))} \leqslant C \left(\oint_{B_{3R}} |V(D^k u)|^\alpha \,\mathrm{d}x \right)^{1/\alpha} \mathrm{d}x$$

with C = C(n, p, k, N) > 0.

The higher integrability result states as follows:

Proposition 6.5. Let an integrand f satisfy (H1)-(H3) and that $u \in W^{k,p}(\Omega, U)$ be a generalized minimizer of $(P\mathscr{B}_{\Omega})$ satisfying $\mathscr{C}^*u = 0$. Then there exists $\sigma_0 > 0$ depending on n, p and $c_{0,1/2}$ in (6.6) such that $D^k u \in L^{p+\sigma}_{loc}(\Omega)$ for any $\sigma \in (0, \sigma_0)$.

Proof. It is sufficient to derive a weak reverse Hölder inequality, and then the desired integrability result follows from the generalized Gehring lemma (see, for example, §V.1 in [Gia83]).

In the case where $p \ge 2$, we have

$$\langle z \rangle^{p-1} |z| \lesssim |z| + |z|^p \lesssim 1 + |z|^p \lesssim 1 + |z|^p$$

for any vector z. In addition, the estimate (a) in Lemma 2.8 implies that

 $|z|^p \lesssim |V_p(z)|^2 \lesssim 1 + |z|^p.$

Fix an arbitrary ball $B_R \in \Omega$ with $R \leq 1$ and take a (k-1)-polynomial a such that

$$(D^i a)_{B_R} = (D^i u)_{B_R}, \quad i = 0, \dots, k-1.$$

Then from (6.6) with $\tau = \frac{1}{2}$ we obtain

$$\begin{split} \oint_{B_{R/2}} |D^k u|^p \, \mathrm{d}x &\leq C \left(\sum_{i=0}^{k-1} \oint_{B_R} \frac{|D^i (u-a)|^p}{R^{p(k-i)}} \, \mathrm{d}x + R \oint_{B_R} |\mathscr{B}(u-a)|^p \, \mathrm{d}x + 1 \right) \\ &\leq C \left(\left(\oint_{B_R} |D^k u|^{p_*} \, \mathrm{d}x \right)^{\frac{p}{p_*}} + R \oint_{B_R} |\mathscr{B}(u-a)|^p \, \mathrm{d}x + 1 \right), \end{split}$$

where the second line follows from the Sobolev-Poincaré inequality and $p_* = \frac{np}{n+p}$. Take $R \ll 1$ such that CR < 1. Then this weak reverse Hölder inequality together with the generalized Gehring lemma ([Gia83], Chapter V, Proposition 1.1) implies the local higher integrability of $D^k u$.

When $1 , we show a weak reverse Hölder inequality with respect to <math>V_p(z)$ instead. Notice that

$$\langle z \rangle^{p-1} |z| \lesssim |z| + |z|^p \lesssim 1 + |V_p(z)|^2.$$

Fix an arbitrary ball B_R with $B_{3R} \in \Omega$ and $R \leq 1$, take a (k-1)-polynomial a as above and let $\tau = \frac{1}{2}$. Then (6.6) implies

$$\begin{split} \int_{B_{R/2}} &|V_p(D^k u)|^2 \,\mathrm{d}x \leqslant C \left(\sum_{i=0}^{k-1} \oint_{B_R} \left| V_p \left(\frac{D^i(u-a)}{R^{k-i}} \right) \right|^2 \,\mathrm{d}x \\ &+ \sum_{i=0}^{k-1} \oint_{B_R} \left(1 + \left| V_p \left(\frac{D^i(u-a)}{R^{k-i}} \right) \right|^2 \right) \,\mathrm{d}x + R \oint_{B_R} (1 + |V_p(\mathscr{B}(u-a))|^2) \,\mathrm{d}x \right) \\ &\leqslant C \left(\left(\oint_{B_{3R}} |V_p(D^k u)|^\beta \,\mathrm{d}x \right)^{\frac{2}{\beta}} + R \oint_{B_R} |V_p(D^k u)|^2 \,\mathrm{d}x + 1 \right), \end{split}$$

where the third line follows from Theorem 6.4 and $\beta \in (\frac{2}{p}, 2)$. Taking $R \ll 1$ such that CR < 1 and applying the generalized Gehring lemma again, we obtain the local higher integrability of $V_p(D^k u)$, and thus of $D^k u$.

7. PARTIAL REGULARITY

To prove Theorem B, it is enough to prove the corresponding excess decay estimate (Propositions 7.1 and 7.2). These imply the claimed partial Hölder regularity using a routine argument, which can be found in [Gia83], Chapter IV. We consider the superquadratic case $p \ge 2$ and sub-quadratic one p < 2 separately, and base our proofs on [Eva86] and [CFM98].

Proof of Theorem B. Follows from Propositions 7.1 or 7.2 (depending on p), Proposition 6.5 (for higher integrability), and Proposition 4.8 (to verify that minimizers can be assumed \mathscr{C}^* -free).

Proof of Theorem A assuming Theorem B. Follows from Remarks 4.4 and 6.3.

To finish the paper, it remains to prove Propositions 7.1 and 7.2.

7.1. Super-quadratic case. Let $\alpha \in (0, 1)$. We consider the following excess functional:

(7.1)
$$\mathscr{E}(u, x_0, R) := R^{2\alpha} + \int_{B_R(x_0)} |V_p(D^k u - (D^k u)_{x_0, R})|^2 \, \mathrm{d}x.$$

Proposition 7.1. Suppose that the integrand f satisfies (H1)-(H3) with $p \ge 2$, and that $u \in W^{k,p}(\Omega)$ is such that $\mathscr{B}u$ a generalized minimizer of $(P\mathscr{B}_{\Omega})$ with $\mathscr{C}^*u = 0$. For any given M > 0, $\alpha \in (0,1)$ and $\tau \in (0,\frac{1}{4})$, there exists $\varepsilon > 0$ such that the following holds true: for any ball $B(x_0, R)(=: B_R) \Subset \Omega$, the conditions

$$|(D^{\kappa}u)_R| < M, \quad \mathscr{E}(u, x_0, R) < \varepsilon$$

implies

$$\mathscr{E}(\tau R) \leqslant C_0 \tau^{2\alpha} \mathscr{E}(R)$$

for some $C_0 = C_0(n, p, \mathscr{B}, \dim V, L, \ell, M) > 0$, where \mathscr{E} is defined as in (7.1).

Proof. The proposition is proved by contradiction and the proof is divided into four steps. **Step 1. Contradiction assumption.** Suppose that the proposition is not true for a certain $\tau \in (0, \frac{1}{4})$. Then there exists a sequence of balls $\{B(x_m, r_m)\}_{m \in \mathbb{N}}$ with $B(x_m, r_m) \subseteq \Omega$ such that

- (7.2) $|(D^k u)_{x_m, r_m}| < M, \quad \mathscr{E}(u, x_m, r_m) = \lambda_m^2 \xrightarrow{m \to \infty} 0,$
- (7.3) $\mathscr{E}(u, x_m, \tau r_m) > C_0 \tau^{2\alpha} \mathscr{E}(u, x_m, r_m),$

where the constant C_0 is to be determined. Without loss of generality, we can assume $r_m \to 0$.

Define the following map on $B_1 = B(0, 1)$ for each $m \in \mathbb{N}$:

$$v_m(y) := \frac{1}{\lambda_m r_m^k} (u(x_m + r_m y) - r_m^k a_m(y)),$$

where the map a_m on B_1 is a k-polynomial such that

$$\int_{B(x_m, r_m)} D^i(u - a_m) \, \mathrm{d}x = 0, \quad 0 \leqslant i \leqslant k - 1, \quad D^k a_m = (D^k u)_{x_m, r_m} := A_m.$$

Notice that $\mathscr{C}^* u = 0$ implies $\mathscr{C}^* a_m = 0$. Then we have, by direct calculation and (7.3),

(7.4)
$$D^k v_m = \frac{1}{\lambda_m} (D^k u (x_m + r_m y) - A_m), \quad \mathscr{C}^* v_m = 0,$$

 $(D^i v_m)_1 = 0, \quad 0 \le i \le k,$

(7.5)
$$r_m^{2\alpha} + \int_{B_1} (1 + \lambda_m^2 |D^k v_m|^2)^{\frac{p-2}{2}} \lambda_m^2 |D^k v_m|^2 \, \mathrm{d}y = \mathscr{E}(u, x_m, r_m) = \lambda_m^2,$$

(7.6)
$$(\tau r_m)^{2\alpha} + \int_{B_\tau} |V(\lambda_m (D^k v_m - (D^k v_m)_\tau))|^2 \,\mathrm{d}x > C_0 \lambda_m^2 \tau^{2\alpha}.$$

In addition, (7.5) implies $\int_{B_1} |D^k v_m|^2 dy \leq 1$. Using the choice of a_m and Poincaré's inequality, we know

$$||v_m||_{W^{k,2}(B_1)} \leqslant C ||D^k v_m||_{L^2(B_1)} \leqslant C.$$

Step 2. Blow up. Now we know that $\{v_m\}$ is uniformly bounded in $W^{k,2}(B_1)$ with $\mathscr{C}^*v_m = 0$, then up to a subsequence there exists a map $v \in W^{k,2}(B_1)$ such that

$$\begin{cases} v_m \rightharpoonup v & \text{in } W^{k,2}(B_1), \\ \mathscr{C}^* v = 0 & \text{in } \mathscr{D}'(B_1). \end{cases}$$

In particular, we have the strong convergence $v_m \to v$ in $W^{k-1,2}(B_1)$. Notice that $\{A_m\}$ is a bounded sequence, so without loss of generality, we can assume that $A_m \to A_0$ for some $A_0 \in \mathbb{R}^{N^k \times n}$ with $|A_0| \leq M$. Then it is easy to see

$$|(\mathscr{B}v_m)_1| = |(T(D^k v_m))_1| = |T((D^k v_m)_1)| \leq C(\mathscr{B}, M),$$

and $|(\mathscr{B}v)_1| = \lim_{m \to \infty} |(\mathscr{B}v_m)_1| \leq C(\mathscr{B}, M),$

where $T: \mathbb{R}^{N^k \times n} \to V$ is a linear operator with $T(D^k u) = \mathscr{B}u$.

Step 3. The harmonicity of v. This step is to show that v solves the following system on B_1 in a weak sense:

$$\mathscr{B}^*(\partial_z^2 f(x_0, T(A_0))\mathscr{B}v) = 0.$$

Given any suitable test map $\psi \in C_c^{\infty}(B_1)$, we test the Euler-Lagrange system satisfied by u on $B(x_m, r_m)$ with $\psi_m(\cdot) := \psi((\cdot - x_m)/r_m)$:

$$0 = \int_{B(x_m, r_m)} \partial_z f(x, \mathscr{B}u) \cdot \mathscr{B}\psi_m \, \mathrm{d}x = \int_{B_1} \partial_z f(x_m + r_m y, \lambda_m \mathscr{B}v_m + T(A_m)) \cdot \mathscr{B}\psi \, \mathrm{d}y$$

(7.7)
$$= \int_{B_1} (\partial_z f(x_m + r_m y, \lambda_m \mathscr{B}v_m + T(A_m)) - \partial_z f(x_m + r_m y, T(A_m))) \cdot \mathscr{B}\psi \, \mathrm{d}y$$
$$+ \int_{B_1} (\partial_z f(x_m + r_m y, T(A_m)) - \partial_z f(x_m, T(A_m))) \cdot \mathscr{B}\psi \, \mathrm{d}y := \int_{B_1} I \, \mathrm{d}y + \int_{B_1} II \, \mathrm{d}y.$$

Notice that

$$\int_{B_1} II \, \mathrm{d}y \leqslant r_m L (1 + |T(A_m)|^2)^{\frac{p-1}{2}} \leqslant C(p, L, M) r_m$$

by the Lipchitz continuity of $\partial_z f(\cdot, z)$ as in (**H2**). Since $r_m \leq \lambda_m^{\frac{1}{\alpha}}$, we have

(7.8)
$$\lim_{m \to \infty} \frac{1}{\lambda_m} \oint_{B_1} II \, \mathrm{d}y = 0$$

To analyze the other term, we split B_1 into

$$E_m^+ := \{ y \in B_1 \colon \lambda_m | \mathscr{B}v_m | \ge 1 \},\$$

$$E_m^- := \{ y \in B_1 \colon \lambda_m | \mathscr{B}v_m | < 1 \}.$$

The measure of E_m^+ is controlled with (7.5):

$$|E_m^+| \leqslant \int_{E_m^+} \lambda_m^2 |\mathscr{B}v_m|^2 \, \mathrm{d}y \leqslant C \int_{E_m^+} \lambda_m^2 |D^k v_m|^2 \, \mathrm{d}y \leqslant C \lambda_m^2.$$

It also indicates that $|E_m^+| \to 0$ as $m \to \infty$. Thus, with the growth rate of $\partial_z f$ we have

(7.9)
$$\frac{1}{\lambda_m} \int_{E_m^+} I \, \mathrm{d}y \leqslant \frac{C}{\lambda_m} \int_{E_m^+} (1 + |\lambda_m \mathscr{B} v_m|^{p-1}) \, \mathrm{d}y$$
$$\leqslant \frac{C}{\lambda_m} \left(|E_m^+| + \left(\int_{E_m^+} \lambda_m^p |D^k v_m|^p \, \mathrm{d}y \right)^{\frac{p-1}{p}} |E_m^+|^{\frac{1}{p}} \right)$$
$$\overset{(7.5)}{\leqslant} C\lambda_m \xrightarrow{m \to \infty} 0.$$

Notice that $\lambda_m \mathscr{B} v_m \to 0$ a.e. in B_1 up to a subsequence and $\mathbb{1}_{E_m^-} \to 1$ in $L^2(B_1)$. Then considering the boundedness of the arguments and the regularity of f in (**H2**), we use the dominated convergence theorem to obtain

$$(7.10) \quad \frac{1}{\lambda_m} \int_{E_m^-} I \, \mathrm{d}y = \int_{E_m^-} \int_0^1 \partial_z^2 f(x_m + r_m y, T(A_m) + t\lambda_m \mathscr{B}v_m) \, \mathrm{d}t[\mathscr{B}v_m, \mathscr{B}\psi] \, \mathrm{d}y$$
$$= \int_{E_m^-} \int_0^1 (\partial_z^2 f(x_m + r_m y, T(A_m) + t\lambda_m \mathscr{B}v_m) - \partial_z^2 f(x_0, T(A_0))) \, \mathrm{d}t[\mathscr{B}v_m, \mathscr{B}\psi] \, \mathrm{d}y$$
$$+ \int_{E_m^-} \partial_z^2 f(x_0, T(A_0))[\mathscr{B}v_m, \mathscr{B}\psi] \, \mathrm{d}y \xrightarrow{m \to \infty} \int_{B_1} \partial_z^2 f(x_0, T(A_0))[\mathscr{B}v, \mathscr{B}\psi] \, \mathrm{d}y$$

With (7.7)-(7.10) we can conclude the following equality

$$\int_{B_1} \partial_z^2 f(x_0, T(A_0))[\mathscr{B}v, \mathscr{B}\psi] \,\mathrm{d}x = 0,$$

which indicates that v solves the following system

(7.11)
$$\begin{cases} \mathscr{B}^*(\partial_z^2 f(x_0, T(A_0))\mathscr{B}v) = 0\\ \mathscr{C}^*v = 0 \end{cases}$$

on the ball B_1 .

Step 4. Excess decay estimate. From Theorem 5.2, we know that $v \in C^{\infty}(B_1)$ and satisfies the following inequality

(7.12)
$$\sup_{B_{\frac{1}{2}}} |D^{k+1}v| \leq C \left(\oint_{B_1} |D^k v - b|^2 \, \mathrm{d}x \right)^{\frac{1}{2}}$$

with some constant C > 0 and any constant vector b. For the given $\tau \in (0, \frac{1}{4})$, take k-polynomials $\{b_m\}$ such that

(7.13)
$$\int_{B_{2\tau}} D^i (v_m - b_m) \, \mathrm{d}x = 0, \quad 0 \le i \le k - 1,$$

(7.14) and
$$D^k b_m = (D^k v_m)_{2\tau}$$

By the convergence $v_m \rightharpoonup v$ in $W^{k,2}(B_1)$, we have b_m converges to some b pointwise with

(7.15)
$$\int_{B_{2\tau}} D^i(v-b) \, \mathrm{d}x = 0, \quad 0 \leqslant i \leqslant k-1,$$

(7.16) and
$$D^k b = (D^k v)_{2\tau}$$

With (7.12), we can control the mean oscillation of $D^k v$ on $B_{2\tau}$ as follows:

(7.17)
$$\begin{aligned} \int_{B_{2\tau}} |D^{k}v - (D^{k}v)_{2\tau}|^{2} \, \mathrm{d}x &\leq 16\tau^{2} \sup_{B_{2\tau}} |D^{k+1}v|^{2} \\ &\leq C\tau^{2} \int_{B_{1}} |D^{k}v - (D^{k}v)_{1}|^{2} \, \mathrm{d}x \\ &\leq C\tau^{2} \liminf_{m \to \infty} \int_{B_{1}} |D^{k}v_{m} - (D^{k}v_{m})_{1}|^{2} \, \mathrm{d}x \leq C\tau^{2}, \end{aligned}$$

where the third line is by the lower semicontinuity of the integral $\int |\cdot|^2$ and the weak convergence $D^k v_m \rightharpoonup D^k v$ on B_1 .

On the other hand, the strong convergence $D^i v_m \to D^i v$ in $L^2(B_1)$, $0 \leq i \leq k-1$, implies

(7.18)
$$\lim_{m \to \infty} \int_{B_{2\tau}} \frac{|D^i(v_m - b_m)|^2}{\tau^{2(k-i)}} \, \mathrm{d}x = \int_{B_{2\tau}} \frac{|D^i(v - b)|^2}{\tau^{2(k-i)}} \, \mathrm{d}x$$
$$\leqslant \dots \leqslant C \int_{B_{2\tau}} |D^k v - (D^k v)_{2\tau}|^2 \, \mathrm{d}x \leqslant C\tau^2,$$

where the third line follows from Poincaré's inequality and (7.17).

To obtain a contradiction in the case p > 2, we need to show

(7.19)
$$\lim_{m \to \infty} \lambda_m^{p-2} \int_{B_{2\tau}} |D^i(v_m - b_m)|^p \, \mathrm{d}x = 0, \quad 0 \le i \le k-1.$$

It is not necessary when p = 2. Take p^* as follows

$$p^* := \begin{cases} \frac{np}{n-p}, & 2 p, & p \ge n, \end{cases}$$

with a fixed r > n, and then $\theta \in (0, 1)$ such that

$$\frac{1}{p} = \frac{\theta}{p^*} + \frac{1-\theta}{2}.$$

For any $i \in \{0, 1, \dots, k-1\}$, by Hölder's inequality and the Sobolev-Poincaré inequality we have

$$\begin{split} \lambda_{m}^{p-2} \oint_{B_{2\tau}} |D^{i}(v_{m} - b_{m})|^{p} \, \mathrm{d}x \\ & \leq \lambda_{m}^{p-2} \left(\int_{B_{2\tau}} |D^{i}(v_{m} - b_{m})|^{2} \, \mathrm{d}x \right)^{\frac{p(1-\theta)}{2}} \left(\int_{B_{2\tau}} |D^{i}(v_{m} - b_{m})|^{p^{*}} \, \mathrm{d}x \right)^{\frac{p\theta}{p^{*}}} \\ & \leq C(\tau) \lambda_{m}^{p-2} \left(\int_{B_{2\tau}} |D^{i}(v_{m} - b_{m})|^{2} \, \mathrm{d}x \right)^{\frac{p(1-\theta)}{2}} \left(\int_{B_{2\tau}} |D^{k}(v_{m} - b_{m})|^{p} \, \mathrm{d}x \right)^{\theta}. \end{split}$$

Notice that the first term $(\int_{B_{2\tau}} |D^i(v_m - b_m)|^2 dx)^{p(1-\theta)/2}$ is dominated by $C(\tau) + o(1)$ as $m \to \infty$. The second one can be controlled by (7.5) as follows

$$\int_{B_{2\tau}} |D^k(v_m - b_m)|^p \, \mathrm{d}x \leqslant C(\tau) \int_{B_1} |D^k v_m - (D^k v_m)_1|^p \, \mathrm{d}x \leqslant C(\tau) \lambda_m^{2-p}.$$

Thus, we have

$$\lambda_m^{p-2} \oint_{B_{2\tau}} |D^i(v_m - b_m)|^p \,\mathrm{d}x \leqslant C(\tau) \lambda_m^{(p-2)(1-\theta)} \to 0 \quad \text{as } m \to \infty.$$

The choice of b_m and the fact $(\mathscr{B}v_m)_1 = 0$ imply

$$\begin{split} \lambda_m |D^k b_m| &= \lambda_m |(D^k v_m)_{2\tau}| = \lambda_m |(D^k v_m)_{2\tau} - (D^k v_m)_1| \\ &\leqslant \frac{\lambda_m}{(2\tau)^n} \left(\int_{B_1} |D^k v_m - (D^k v_m)_1|^2 \,\mathrm{d}x \right)^{\frac{1}{2}} \leqslant \frac{\sqrt{\varepsilon}}{(2\tau)^n} < 1 \end{split}$$

when $\varepsilon < (2\tau)^{2n}$. It is then easy to see that

$$\lambda_m(\mathscr{B}v_m(y) - \mathscr{B}b_m(y)) = \mathscr{B}u(x_m + r_m y) - T(A_m) - \lambda_m(\mathscr{B}v_m)_{2\tau}$$

with $|T(A_m) + \lambda_m(\mathscr{B}v_m)_{2\tau}| \leq C(\mathscr{B}, M)$. Then (b) and (d) in Lemma 2.8, and Proposition 6.2 imply

$$\begin{split} \int_{B_{\tau}} |V(\lambda_m (D^k v_m - (D^k v_m)_{\tau}))|^2 \, \mathrm{d}x &\leq C \int_{B_{\tau}} |V(\lambda_m (D^k (v_m - b_m)))|^2 \, \mathrm{d}x \\ &\leq C \sum_{i=0}^{k-1} \int_{B_{2\tau}} \left| V\left(\frac{\lambda_m D^i (v_m - b_m)}{(2\tau)^{k-i}}\right) \right|^2 \, \mathrm{d}x \\ &+ C \tau r_m \sum_{i=0}^{k-1} \int_{B_{2\tau}} \left\langle \frac{\lambda_m D^i (v_m - b_m)}{(2\tau)^{k-i}} \right\rangle^{p-1} \left| \frac{\lambda_m D^i (v_m - b_m)}{(2\tau)^{k-i}} \right| \, \mathrm{d}x \\ &+ C \tau r_m \int_{2\tau} \left\langle \lambda_m \mathscr{B}(v_m - b_m) \right\rangle^{p-1} \left| \lambda_m \mathscr{B}(v_m - b_m) \right| \, \mathrm{d}x =: I + II + III. \end{split}$$

With (a) in Lemma 2.8, (7.18) and (7.19) we know

$$\begin{split} I &\leqslant C \oint_{B_{2\tau}} \left(\lambda_m^2 \frac{|D^i(v_m - b_m)|^2}{(2\tau)^{2(k-i)}} + \lambda_m^p \frac{|D^i(v_m - b_m)|^p}{(2\tau)^{p(k-i)}} \right) \mathrm{d}x \\ &\leqslant C \lambda_m^2 \tau^2 + \lambda_m^2 o(1), \quad \text{as } m \to \infty. \end{split}$$

The estimates of II is similar:

$$II \leqslant C\tau r_m \sum_{i=0}^{k-1} \oint_{B_{2\tau}} \left(\lambda_m \frac{|D^i(v_m - b_m)|}{(2\tau)^{(k-i)}} + \lambda_m^p \frac{|D^i(v_m - b_m)|^p}{(2\tau)^{p(k-i)}} \right) \mathrm{d}x$$
$$\leqslant C\tau r_m \sum_{i=0}^{k-1} \left(\lambda_m \left(\oint_{B_{2\tau}} \frac{|D^i(v_m - b_m)|^2}{(2\tau)^{2(k-i)}} \,\mathrm{d}x \right)^{\frac{1}{2}} + \lambda_m^2 o(1) \right)$$
$$\leqslant C\tau r_m (\lambda_m \tau + \lambda_m^2 o(1)), \quad \text{as } m \to \infty.$$

Finally, control III with (7.5) similarly:

$$III \leqslant C\tau r_m (\lambda_m (2\tau)^{-\frac{n}{2}} + \lambda_m^2 (2\tau)^{-n}).$$

Notice that $r_m \leqslant \lambda_m^{1/\alpha}$ by (7.2), then we have

$$\begin{aligned} &\int_{B_{\tau}} |V(\lambda_m (D^k v_m - (D^k v_m)_{\tau}))|^2 \,\mathrm{d}x \\ &\leqslant C \lambda_m^2 (\tau^2 + o(1) + \tau^2 \lambda_m^{\frac{1}{\alpha} - 1} + \tau \lambda_m^{\frac{1}{\alpha}} o(1) + \tau \lambda_m^{\frac{1}{\alpha} - 1} (2\tau)^{-\frac{n}{2}} + \tau \lambda_m^{\frac{1}{\alpha}} (2\tau)^n) \leqslant C_1 \lambda_m^2 \tau^{2\alpha} \end{aligned}$$

if $\varepsilon(>\lambda_m^2)$ is taken to be small enough and *m* large enough. The above inequality contradicts (7.6) if $C_0 > C_1 + 1$. The proof of the proposition is now complete. 7.2. Sub-quadratic case. The proof in the sub-quadratic set-up $(1 is similar to that in the super-quadratic one. As mentioned above, it is sufficient to show the excess decay estimate as in Proposition 7.1, where the excess <math>\mathscr{E}$ is defined as in (7.1). We omit some details and only mention the difference here.

Proposition 7.2. Suppose that the integrand f satisfies (H1)-(H3) with $p \in (1, 2)$, and that $u \in W^{k,p}(\Omega)$ is such that $\mathscr{B}u$ is a generalized minimizer of $(P\mathscr{B}_{\Omega})$ with $\mathscr{C}^*u = 0$. For any given M > 0, $\alpha \in (0, 1)$ and $\tau \in (0, \frac{1}{12})$, there exists $\varepsilon > 0$ such that the following holds true: for any ball $B(x_0, R)(=: B_R) \subseteq \Omega$, the conditions

(7.20)
$$|(D^k u)_R| < M, \quad \mathscr{E}(u, x_0, R) < \varepsilon$$

implies

(7.21)
$$\mathscr{E}(\tau R) \leqslant C_0 \tau^2 \mathscr{E}(R)$$

for some $C_0 = C_0(n, p, \mathscr{B}, \dim V, L, \ell, M) > 0.$

Proof. The proof is done by contradiction as well. We assume that the proposition does not hold for a certain $\tau \in (0, \frac{1}{12})$ and take a sequence of balls $\{B(x_m, r_m)\}_{m \in \mathbb{N}}$ with $B(x_m, r_m) \subseteq \Omega$ such that

(7.22)
$$|(D^k u)_{x_m, r_m}| < M, \quad \mathscr{E}(u, x_m, r_m) = \lambda_m^2 \stackrel{m \to \infty}{\longrightarrow} 0,$$

(7.23)
$$\mathscr{E}(u, x_m, \tau r_m) > C_0 \tau^{2\alpha} \mathscr{E}(u, x_m, r_m)$$

where the constant C_0 is to be determined. Without loss of generality, we can assume $r_m \to 0$. Then define $\{a_m\}$ and $\{v_m\}$ as in Proposition 7.1, and the counterparts of (7.4)-(7.6) hold true:

(7.24)
$$D^{k}v_{m} = \frac{1}{\lambda_{m}}(D^{k}u(x_{m}+r_{m}y)-A_{m}), \quad \mathscr{C}^{*}v_{m} = 0,$$
$$(D^{i}v_{m})_{1} = 0, \quad 0 \leq i \leq k,$$
$$(7.25) \quad r_{m}^{2\alpha} + \int_{B_{1}}|V(\lambda_{m}D^{k}v_{m})|^{2} \,\mathrm{d}y = \mathscr{E}(u,x_{m},r_{m}) = \lambda_{m}^{2},$$

(7.26)
$$(\tau r_m)^{2\alpha} + \int_{B_\tau} |V(\lambda_m (D^k v_m - (D^k v_m)_\tau))|^2 \,\mathrm{d}x > C_0 \lambda_m^2 \tau^{2\alpha}.$$

Moreover, from (7.25) and (a) in Lemma 2.8 we have $f_{B_1}|D^k v_m|^p dy \leq C(p)$, and then Poincaré's inequality implies the boundedness of $\{v_m\}$ in $W^{k,p}(B_1)$.

Similarly, there exists a subsequence (not relabeled) such that

$$\begin{cases} v_m \rightharpoonup v & \text{in } W^{k,p}(B_1), \\ v_m \rightarrow v & \text{in } W^{k-1,p}(B_1), \\ \mathscr{C}^* v = 0 & \text{in } \mathscr{D}'(B_1), \end{cases}$$

and $A_m \to A_0$ for some $A_0 \in \mathbb{R}^{N^k \times n}$ with $|A_0| \leq M$. With the same argument, we can deduce that v solves the following system

(7.27)
$$\begin{cases} \mathscr{B}^*(\partial_z^2 f(x_0, T(A_0))\mathscr{B}v) = 0\\ \mathscr{C}^*v = 0 \end{cases}$$

on B_1 , which implies

(7.28)
$$\sup_{B_{\frac{1}{2}}} |D^{k+1}v| \leq C \left(\int_{B_1} |D^k v - b|^p \, \mathrm{d}x \right)^{\frac{1}{p}}$$

with some constant C > 0 and any constant vector b. Take $\{b_m\}$ as in the proof of Proposition 7.1, and the sequence converges to some k-polynomial b that satisfies (7.15) and (7.16).

The L^2 -oscillation of $D^k v$ on $B_{2\tau}$ can be then controlled as follows:

$$\begin{aligned} \int_{B_{2\tau}} |D^{k}v - (D^{k}v)_{2\tau}|^{2} \, \mathrm{d}x &\leq 16\tau^{2} \sup_{B_{2\tau}} |D^{k+1}v|^{2} \\ &\leq C\tau^{2} \left(\int_{B_{1}} |D^{k}v - (D^{k}v)_{1}|^{p} \, \mathrm{d}x \right)^{\frac{2}{p}} \\ &\leq C\tau^{2} \left(\liminf_{m \to \infty} \int_{B_{1}} |D^{k}v_{m} - (D^{k}v_{m})_{1}|^{p} \, \mathrm{d}x \right)^{\frac{2}{p}} \leq C\tau^{2}. \end{aligned}$$

Now we estimate the integral term in $\mathscr{E}(u, x_m, r_m)$ with the Caccioppolli-type inequality (6.6). Notice that

$$\lambda_m(\mathscr{B}v_m(y) - \mathscr{B}b_m(y)) = \mathscr{B}u(x_m + r_m y) - T(A_m) - \lambda_m(\mathscr{B}v_m)_{2\tau},$$

with $|T(A_m) + \lambda_m(\mathscr{B}v_m)_{2\tau}| \leq C(\mathscr{B}, M),$

as in Proposition 7.1. Then apply this inequality to u on $B_{2\tau r_m}$ and rescale it to $B_{2\tau}$ to get

$$\begin{split} \int_{B_{\tau}} |V(\lambda_m D^k v_m - \lambda_m (D^k v_m)_{\tau})|^2 \, \mathrm{d}x &\leq C \int_{B_{\tau}} |V(\lambda_m D^k (v_m - b_m))|^2 \, \mathrm{d}x \\ &\leq C \left(\sum_{i=0}^{k-1} \int_{B_{2\tau}} \left| V \left(\frac{\lambda_m D^i (v_m - b_m)}{(2\tau)^{k-i}} \right) \right|^2 \, \mathrm{d}x \\ &+ 2\tau r_m \sum_{i=0}^{k-1} \int_{B_{2\tau}} \left\langle \frac{\lambda_m D^i (v_m - b_m)}{(2\tau)^{k-i}} \right\rangle^{p-1} \frac{\lambda_m |D^i (v_m - b_m)|}{(2\tau)^{k-i}} \, \mathrm{d}x \\ &+ 2\tau r_m \int_{B_{2\tau}} \left\langle \lambda_m \mathscr{B}(v_m - b_m) \right\rangle^{p-1} \lambda_m |\mathscr{B}(v_m - b_m)| \, \mathrm{d}x \right) =: I + II + III. \end{split}$$

Since 1 , the term II can be estimated as follows:

$$II \leqslant C\tau\lambda_{m}r_{m}\sum_{i=0}^{k-1} \oint_{B_{2\tau}} \left(\frac{|D^{i}(v_{m}-b_{m})|}{(2\tau)^{k-i}} + \frac{|D^{i}(v_{m}-b_{m})|^{p}}{(2\tau)^{p(k-i)}}\right) dx$$

$$(7.30) \qquad \leqslant C\tau\lambda_{m}r_{m} \oint_{B_{2\tau}} (|D^{k}(v_{m}-b_{m})| + |D^{k}(v_{m}-b_{m})|^{p}) dx$$

$$= C\tau\lambda_{m}r_{m} \oint_{B_{2\tau}} (|D^{k}v_{m}-(D^{k}v_{m})_{2\tau}| + |D^{k}v_{m}-(D^{k}v_{m})_{2\tau}|^{p}) dx$$

$$\leqslant C\tau^{1-2n}\lambda_{m}r_{m} \left(\left(\int_{B_{1}} |D^{k}v_{m}|^{p} dx\right)^{\frac{1}{p}} + \int_{B_{1}} |D^{k}v_{m}|^{p} dx\right) \leqslant C\tau^{1-2n}\lambda_{m}r_{m},$$

where the second line follows from Poincaré's inequality and the last estimate from the uniform boundedness of $\{v_m\}$ in $W^{k,p}(B_1)$. It is also easy to see that

(7.31)
$$III \leqslant C\tau^{1-2n}\lambda_m r_m \left(\left(\oint_{B_1} |D^k v_m|^p \,\mathrm{d}x \right)^{\frac{1}{p}} + \oint_{B_1} |D^k v_m|^p \,\mathrm{d}x \right) \leqslant C\tau^{1-2n}\lambda_m r_m.$$

To estimate I, we control each term (i = 0, ..., k - 1) with (b) in Lemma 2.8

$$\oint_{B_{2\tau}} \left| V\left(\frac{\lambda_m D^i \tilde{v}_m}{(2\tau)^{k-i}}\right) \right|^2 \mathrm{d}x \leqslant C\left(\oint_{B_{2\tau}} \left| V\left(\frac{\lambda_m D^i (\tilde{v}_m - \tilde{v})}{(2\tau)^{k-i}}\right) \right|^2 \mathrm{d}x$$

Z. LI AND B. RAIȚĂ

$$+ \oint_{B_{2\tau}} \left| V\left(\frac{\lambda_m D^i \tilde{v}}{(2\tau)^{k-i}}\right) \right|^2 \mathrm{d}x \right) := I_1 + I_2,$$

where $\tilde{v}_m = v_m - b_m$ and $\tilde{v} = v - b$. It is easy to see $\tilde{v}_m \rightharpoonup \tilde{v}$ in $W^{k,p}(B_1)$. The estimate (7.29), and the counterparts of (7.15) and (7.16), together with (a) in Lemma 2.8 imply

(7.32)
$$I_2 \leqslant C\lambda_m^2 \oint_{B_{2\tau}} \frac{|D^i \tilde{v}|^2}{(2\tau)^{2(k-i)}} \,\mathrm{d}x \leqslant C\lambda_m^2 \oint_{B_{2\tau}} |V(D^k v - (D^k v)_{2\tau})|^2 \,\mathrm{d}x \leqslant C\lambda_m^2 \tau^2.$$
Take $\theta \in (0, 1)$ such that

Take $\theta \in (0, 1)$ such that

$$\frac{1}{2} = \theta + \frac{1-\theta}{2(1+\sigma)}$$

where σ is as in Theorem 6.4. Then with Hölder's inequality we have

$$\begin{split} I_1 &\leqslant C \left(\oint_{B_{2\tau}} \left| V \left(\frac{\lambda_m D^i(\tilde{v}_m - \tilde{v})}{(2\tau)^{k-i}} \right) \right| \mathrm{d}x \right)^{2\theta} \left(\oint_{B_{2\tau}} \left| V \left(\frac{\lambda_m D^i(\tilde{v}_m - \tilde{v})}{(2\tau)^{k-i}} \right) \right|^{2(1+\sigma)} \mathrm{d}x \right)^{\frac{1-\theta}{1+\sigma}} \\ &=: E_1^{2\theta} \cdot E_2^{1-\theta}. \end{split}$$

Apply (a) in Lemma 2.8 again to obtain

$$E_1 \leqslant C\lambda_m \oint_{B_{2\tau}} \frac{|D^i(\tilde{v}_m - \tilde{v})|}{(2\tau)^{k-i}} \,\mathrm{d}x = \lambda_m \tau^{-k} \cdot o(1)$$

as $m \to \infty$. The term E_2 can be controlled with Theorem 6.4:

$$\begin{split} E_{2} &\leqslant C \int_{B_{6\tau}} |V(\lambda_{m}D^{k}(\tilde{v}_{m} - \tilde{v})|^{2} \,\mathrm{d}x \\ &\leqslant C \int_{B_{6\tau}} (|V(\lambda_{m}D^{k}v_{m}|^{2} + |V(\lambda_{m}D^{k}(b_{m} - b))|^{2} + |V(\lambda_{m}D^{k}v|^{2}) \,\mathrm{d}x \\ &\leqslant C \left(\tau^{-n} \int_{B_{1}} |V(\lambda_{m}D^{k}v_{m})|^{2} \,\mathrm{d}x + \lambda_{m}^{2} \int_{B_{6\tau}} |D^{k}(b_{m} - b)|^{2} \,\mathrm{d}x + \lambda_{m}^{2} \int_{B_{6\tau}} |D^{k}v|^{2} \,\mathrm{d}x \right) \\ &\leqslant C \lambda_{m}^{2} (\tau^{-n} + o(1)) + C \lambda_{m}^{2} \int_{B_{6\tau}} |D^{k}v|^{2} \,\mathrm{d}x. \end{split}$$

With (7.28) it is easy to obtain

$$\begin{split} & \int_{B_{6\tau}} |D^k v|^2 \, \mathrm{d}x \leqslant 2 \int_{B_{6\tau}} |D^k v - (D^k v)_{\frac{1}{2}}|^2 \, \mathrm{d}x + 2|(D^k v)_{\frac{1}{2}}|^2 \\ & \leqslant C \left(\int_{B_1} |D^k v|^p \, \mathrm{d}x \right)^{\frac{2}{p}} \leqslant C, \end{split}$$

which helps us conclude

(7.33)
$$I_1 \leqslant \lambda_m^2 \tau^{-k} (\tau^{-n} + 1) \cdot o(1)$$

as $m \to \infty$. Combining (7.30)-(7.33) and $r_m < \lambda_m^{\frac{1}{\alpha}}$, we have

(7.34)
$$\int_{B_{\tau}} |V(\lambda_m D^k v_m - (D^k v_m)_{\tau})|^2 dx \leq C \lambda_m^2 (\tau^{1-2n} \lambda_m^{\frac{1}{\alpha}-1} + \tau^{-k} (\tau^{-n} + 1) \cdot o(1)) \leq C_1 \lambda_m^2 \tau^{2\alpha}$$

when $\varepsilon \geq \lambda_m^2$ is taken to be small enough and m is large enough. This contradicts (7.26) if we take $C_0 > C_1 + 1$. The proof of the proposition is now complete.

References

- [AD92] L. Ambrosio and G. Dal Maso. "On the relaxation in $BV(\Omega; \mathbb{R}^m)$ of quasi-convex integrals". In: J. Funct. Anal. 109.1 (1992), pp. 76–97. DOI: 10.1016/0022-1236(92)90012-8.
- [ADR20] Adolfo Arroyo-Rabasa, Guido De Philippis, and Filip Rindler. "Lower semicontinuity and relaxation of linear-growth integral functionals under PDE constraints". In: Advances in calculus of variations 13.3 (2020), pp. 219–255.
- [AF84] E. Acerbi and N. Fusco. "Semicontinuity problems in the calculus of variations". In: Arch. Rational Mech. Anal. 86.2 (1984), pp. 125–145. DOI: 10.1007/BF00275731.
- [AF87] E. Acerbi and N. Fusco. "A regularity theorem for minimizers of quasiconvex integrals". In: Arch. Rational Mech. Anal. 99.3 (1987), pp. 261–281. DOI: 10.1007/BF00284509.
- [Arr+19] Adolfo Arroyo-Rabasa, Guido De Philippis, Jonas Hirsch, and Filip Rindler. "Dimensional estimates and rectifiability for measures satisfying linear PDE constraints". In: *Geometric and Functional Analysis* 29 (2019), pp. 639–658.
- [Arr+24] Adolfo Arroyo-Rabasa, Guido De Philippis, Jonas Hirsch, Filip Rindler, and Anna Skorobogatova. "Higher integrability for measures satisfying a PDE constraint". In: Transactions of the American Mathematical Society 377 (2024), pp. 6195–6224.
- [Arr21] Adolfo Arroyo-Rabasa. "Characterization of Generalized Young Measures Generated by A-free Measures". In: Archive for rational mechanics and analysis 242 (2021), pp. 235–325.
- [Baí+13] Margarida Baía, Milena Chermisi, José Matias, and Pedro M Santos. "Lower semicontinuity and relaxation of signed functionals with linear growth in the context of A-quasiconvexity". In: Calc. Var. Partial Differential Equations 47.3-4 (2013), pp. 465–498.
- [BDS15] Saugata Bandyopadhyay, Bernard Dacorogna, and Swarnendu Sil. "Calculus of variations with differential forms." In: Journal of the European Mathematical Society (EMS Publishing) 17.4 (2015).
- [BFL00] Andrea Braides, Irene Fonseca, and Giovanni Leoni. "A-quasiconvexity: relaxation and homogenization". In: ESAIM: Control, Optimisation and Calculus of Variations 5 (2000), pp. 539– 577.
- [BGS22] Linus Behn, Franz Gmeineder, and Stefan Schiffer. "On symmetric div-quasiconvex hulls and divsym-free L[∞]-truncations". In: Annales de l'Institut Henri Poincaré C 40.6 (2022), pp. 1267– 1317.
- [BK22] Matthias Bärlin and Konrad Keßler. "Partial Regularity for A-quasiconvex Functionals". In: arXiv preprint arXiv:2203.00153 (2022).
- [BS13] Lisa Beck and Bianca Stroffolini. "Regularity results for differential forms solving degenerate elliptic systems". In: *Calculus of Variations and Partial Differential Equations* 46.3 (2013), pp. 769–808.
- [BS16a] Saugata Bandyopadhyay and Swarnendu Sil. "Exterior convexity and classical calculus of variations". In: ESAIM: Control, Optimisation and Calculus of Variations 22.2 (2016), pp. 338– 354.
- [BS16b] Saugata Bandyopadhyay and Swarnendu Sil. "Notions of affinity in calculus of variations with differential forms". In: *Advances in Calculus of Variations* 9.3 (2016), pp. 293–304.
- [BS20] Peter Bella and Mathias Schäffner. "On the regularity of minimizers for scalar integral functionals with (p, q)-growth". In: Analysis & PDE 13.7 (2020), pp. 2241–2257.
- [CC15] Allan Chan and Sergio Conti. "Energy scaling and branched microstructures in a model for shape-memory alloys with SO (2) invariance". In: *Mathematical Models and Methods in Applied Sciences* 25.06 (2015), pp. 1091–1124.
- [CFM98] M. Carozza, N. Fusco, and G. Mingione. "Partial regularity of minimizers of quasiconvex integrals with subquadratic growth". In: Ann. Mat. Pura Appl. (4) 175 (1998), pp. 141–164. DOI: 10.1007/BF01783679.
- [CG22] Sergio Conti and Franz Gmeineder. "A-quasiconvexity and partial regularity". In: Calculus of Variations and Partial Differential Equations 61.6 (2022), p. 215.
- [Chi+17] Elisabetta Chiodaroli, Eduard Feireisl, Ondřej Kreml, and Emil Wiedemann. "A-free rigidity and applications to the compressible Euler system". In: Annali di Matematica Pura ed Applicata (1923-) 196.4 (2017), pp. 1557–1572.
- [CK17] C.-Y. Chen and J. Kristensen. "On coercive variational integrals". In: Nonlinear analysis 153 (2017), pp. 213–229. DOI: 10.1016/j.na.2016.09.011.
- [CKN21] Menita Carozza, Jan Kristensen, and Antonia Passarelli di Napoli. "A trace preserving operator and applications". In: Journal of Mathematical Analysis and Applications 501.1 (2021), p. 124170.

- [CMO18] Sergio Conti, Stefan Müller, and Michael Ortiz. "Data-driven problems in elasticity". In: Archive for Rational Mechanics and Analysis 229 (2018), pp. 79–123.
- [CMO20] Sergio Conti, Stefan Müller, and Michael Ortiz. "Symmetric div-quasiconvexity and the relaxation of static problems". In: Archive for Rational Mechanics and Analysis 235.2 (2020), pp. 841–880.
- [CN03] Menita Carozza and Antonia Passarelli di Napoli. "Partial regularity of local minimizers of quasiconvex integrals with sub-quadratic growth". In: Proceedings of the Royal Society of Edinburgh Section A: Mathematics 133.6 (2003), pp. 1249–1262.
- [Dac06] Bernard Dacorogna. Weak continuity and weak lower semicontinuity of non-linear functionals. Vol. 922. Springer, 2006.
- [Dac82] Bernard Dacorogna. "Quasiconvexity and relaxation of nonconvex problems in the calculus of variations". In: *Journal of functional analysis* 46.1 (1982), pp. 102–118.
- [De 22] Cristiana De Filippis. "Quasiconvexity and partial regularity via nonlinear potentials". In: Journal de Mathématiques Pures et Appliquées 163 (2022), pp. 11–82.
- [DF02] Bernard Dacorogna and Irene Fonseca. "AB quasiconvexity and implicit partial differential equations". In: Calculus of Variations and Partial Differential Equations 14.2 (2002), pp. 115– 149.
- [DF16a] Elisa Davoli and Irene Fonseca. "Homogenization of integral energies under periodically oscillating differential constraints". In: Calculus of Variations and Partial Differential Equations 55 (2016), pp. 1–60.
- [DF16b] Elisa Davoli and Irene Fonseca. "Periodic homogenization of integral energies under spacedependent differential constraints". In: *Portugaliae Mathematica* 73.4 (2016), pp. 279–317.
- [DF18] Elisa Davoli and Irene Fonseca. "Relaxation of p-growth integral functionals under spacedependent differential constraints". In: Trends in Applications of Mathematics to Mechanics (2018), pp. 1–21.
- [DFL23] Elisa Davoli, Irene Fonseca, and Pan Liu. "Adaptive image processing: first order PDE constraint regularizers and a bilevel training scheme". In: *Journal of Nonlinear Science* 33.3 (2023), p. 41.
- [DKP24] Elisa Davoli, Martin Kružík, and Valerio Pagliari. "Homogenization of high-contrast composites under differential constraints". In: Advances in Calculus of Variations 17.2 (2024), pp. 277–318.
- [DPR18] Guido De Philippis, Luca Palmieri, and Filip Rindler. "On the two-state problem for general differential operators". In: Nonlinear Analysis 177 (2018), pp. 387–396.
- [DR16] Guido De Philippis and Filip Rindler. "On the structure of *A*-free measures and applications". In: Annals of Mathematics (2016), pp. 1017–1039.
- [DR17] Guido De Philippis and Filip Rindler. "Characterization of generalized Young measures generated by symmetric gradients". In: Archive for Rational Mechanics and Analysis 224 (2017), pp. 1087–1125.
- [DS09] Camillo De Lellis and László Székelyhidi Jr. "The Euler equations as a differential inclusion". In: Annals of mathematics (2009), pp. 1417–1436.
- [DS23] Cristiana De Filippis and Bianca Stroffolini. "Singular multiple integrals and nonlinear potentials". In: *Journal of Functional Analysis* 285.2 (2023), p. 109952.
- [DT23] Luigi De Rosa and Riccardo Tione. "Fine properties of symmetric and positive matrix fields with bounded divergence". In: *Advances in Mathematics* 427 (2023), p. 109130.
- [ELM04] Luca Esposito, Francesco Leonetti, and Giuseppe Mingione. "Sharp regularity for functionals with (p, q) growth". In: *Journal of Differential Equations* 204.1 (2004), pp. 5–55.
- [ELM99] Luca Esposito, Francesco Leonetti, and Giuseppe Mingione. "Higher integrability for minimizers of integral functionals with (p, q) growth". In: *journal of differential equations* 157.2 (1999), pp. 414–438.
- [Eva86] L. C. Evans. "Quasiconvexity and partial regularity in the calculus of variations". In: Arch. Rational Mech. Anal. 95.3 (1986), pp. 227–252. DOI: 10.1007/BF00251360.
- [FFV21] Rita Ferreira, Irene Fonseca, and Raghavendra Venkatraman. "Homogenization of quasi-crystalline functionals via two-scale-cut-and-project convergence". In: SIAM Journal on Mathematical Analysis 53.2 (2021), pp. 1785–1817.
- [FK10] Irene Fonseca and Martin Kružík. "Oscillations and concentrations generated by A-free mappings and weak lower semicontinuity of integral functionals". In: ESAIM: Control, Optimisation and Calculus of Variations 16.2 (2010), pp. 472–502.
- [FLM04] Irene Fonseca, Giovanni Leoni, and Stefan Müller. "A-quasiconvexity: weak-star convergence and the gap". In: Annales de l'Institut Henri Poincaré C 21.2 (2004), pp. 209–236.
- [FM93] I. Fonseca and S. Müller. "Relaxation of quasiconvex functionals in $BV(\Omega, \mathbb{R}^p)$ for integrands $f(x, u, \nabla u)$ ". In: Arch. Rational Mech. Anal. 123.1 (1993), pp. 1–49. DOI: 10.1007/BF00386367.

- [FM97] Irene Fonseca and Jan Maly. "Relaxation of multiple integrals below the growth exponent". In: Annales de l'Institut Henri Poincaré C, Analyse non linéaire. Vol. 14. 3. Elsevier. 1997, pp. 309–338.
- [FM99] I. Fonseca and S. Müller. "A-quasiconvexity, lower semicontinuity, and Young measures". In: SIAM J. Math. Anal. 30.6 (1999), pp. 1355–1390. DOI: 10.1137/S0036141098339885.
- [FMP98] Irene Fonseca, Stefan Müller, and Pablo Pedregal. "Analysis of concentration and oscillation effects generated by gradients". In: SIAM journal on mathematical analysis 29.3 (1998), pp. 736–756.
- [Fra19] F. Franceschini. "Partial regularity for BV² local minimizers". MA thesis. Università degli Studi di Pisa, 2019. URL: https://etd.adm.unipi.it/theses/available/etd-06242019-103617/.
- [Gal21] Dennis Gallenmüller. "Müller-Zhang truncation for general linear constraints with first or second order potential". In: Calculus of Variations and Partial Differential Equations 60.3 (2021), p. 118.
- [Gal23] Dennis Gallenmüller. "Measure-valued low Mach number limits of ideal fluids". In: SIAM Journal on Mathematical Analysis 55.2 (2023), pp. 1145–1169.

[GG82] M. Giaquinta and E. Giusti. "On the regularity of the minima of variational integrals". In: Acta Math. 148 (1982), pp. 31–46.

- [Gia83] M. Giaquinta. Multiple integrals in the calculus of variations and nonlinear elliptic systems. Vol. 105. Annals of Mathematics Studies. Princeton, NJ: Princeton University Press, 1983.
- [Giu03] E. Giusti. Direct methods in the calculus of variations. River Edge, NJ: World Scientific, 2003. DOI: 10.1142/5002.
- [GK19a] F. Gmeineder and J. Kristensen. "Sobolev regularity for convex functionals on BD". In: Calc. Var. Partial Differential Equations 58.56 (2019). DOI: 10.1007/s00526-019-1491-6.
- [GK19b] Franz Gmeineder and Jan Kristensen. "Partial regularity for BV minimizers". In: Archive for Rational Mechanics and Analysis 232.3 (2019), pp. 1429–1473.
- [GM86] M. Giaquinta and G. Modica. "Partial regularity of minimizers of quasiconvex integrals". In: Ann. Inst. H. Poincaré Anal. Non Linéaire 3.3 (1986), pp. 185–208.
- [Gme20] F. Gmeineder. "The regularity of minima for the Dirichlet problem on BD". In: Arch. Rational Mech. Anal. 237 (2020), pp. 1099–1171. DOI: 10.1007/s00205-020-01507-5.
- [Gme21] F. Gmeineder. "Partial regularity for symmetric quasiconvex functionals on BD". In: J. Math. Pures Appl. 145 (2021), pp. 83–129. DOI: 10.1016/j.matpur.2020.09.005.
- [GN04] Adriana Garroni and Vincenzo Nesi. "Rigidity and lack of rigidity for solenoidal matrix fields". In: Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences 460.2046 (2004), pp. 1789–1806.
- [GR22] A. Guerra and B. Raiță. "Quasiconvexity, null Lagrangians, and Hardy space integrability under constant rank constraints". In: Arch. Ration. Mech. Anal. 245.1 (2022), pp. 279–320. DOI: 10.1007/s00205-022-01775-3.
- [GRS22] André Guerra, Bogdan Raiță, and Matthew RI Schrecker. "Compensated compactness: continuity in optimal weak topologies". In: *Journal of Functional Analysis* 283.7 (2022), p. 109596.
- [GRS24] André Guerra, Bogdan Raiță, and Matthew Schrecker. "Compensation phenomena for concentration effects via nonlinear elliptic estimates". In: Ars Inveniendi Analytica (2024).
- [GT01] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Reprint of the 1998 edition. Springer-Verlag, Berlin, 2001.
- [GW21] Dennis Gallenmüller and Emil Wiedemann. "On the selection of measure-valued solutions for the isentropic Euler system". In: Journal of Differential Equations 271 (2021), pp. 979–1006.
- [GW23] Dennis Gallenmüller and Emil Wiedemann. "Which measure-valued solutions of the monoatomic gas equations are generated by weak solutions?" In: Archive for Rational Mechanics and Analysis 247.4 (2023), p. 61.
- [GWW23] Dennis Gallenmüller, Raphael Wagner, and Emil Wiedemann. "Probabilistic descriptions of fluid flow: a survey". In: Journal of Mathematical Fluid Mechanics 25.3 (2023), p. 52.
- [Ham92] C Hamburger. "Regularity of differential forms minimizing degenerate elliptic functionals*". In: j. reine angew. Math 431 (1992), pp. 7–64.
- $[Irv23] \qquad \text{Christopher Irving. "BMO ε-regularity results for solutions to Legendre-Hadamard elliptic systems". In: Calculus of Variations and Partial Differential Equations 62.5 (2023), p. 166.$
- [ISS99] Tadeusz Iwaniec, Chad Scott, and Bianca Stroffolini. "Nonlinear Hodge theory on manifolds with boundary". In: Annali di Matematica pura ed applicata 177.1 (1999), pp. 37–115.
- [KK16] Bernd Kirchheim and Jan Kristensen. "On rank one convex functions that are homogeneous of degree one". In: Archive for rational mechanics and analysis 221 (2016), pp. 527–558.
- [KM16] Tuomo Kuusi and Giuseppe Mingione. "Partial regularity and potentials". In: Journal de l'École polytechnique—Mathématiques 3 (2016), pp. 309–363.

- [KP94] David Kinderlehrer and Pablo Pedregal. "Gradient Young measures generated by sequences in Sobolev spaces". In: The Journal of Geometric Analysis 4.1 (1994), pp. 59–90.
- [KR10] J. Kristensen and F. Rindler. "Relaxation of signed integral functionals in BV". In: Calc. Var. Partial Differential Equations 37.1-2 (2010), pp. 29–62. DOI: 10.1007/s00526-009-0250-5.
- [KR20] J. Kristensen and Bogdan Raiță. An introduction to generalized Young measures. Lecture note 45/2020, Max-Planck-Institut für Mathematik in den Naturwissenschaften Leipzig. 2020.
- [KR22] Jan Kristensen and Bogdan Raiță. "Oscillation and concentration in sequences of PDE constrained measures". In: Archive for Rational Mechanics and Analysis 246.2 (2022), pp. 823– 875.
- [Krä+17] Jan Krämer, Stefan Krömer, Martin Kružík, and Gabriel Pathó. "A-quasiconvexity at the boundary and weak lower semicontinuity of integral functionals". In: Advances in Calculus of Variations 10.1 (2017), pp. 49–67.
- [Kri99] Jan Kristensen. "Lower semicontinuity in spaces of weakly differentiable functions". In: *Mathematische Annalen* 4.313 (1999), pp. 653–710.
- [KSW17] Krystian Kazaniecki, Dmitriy M Stolyarov, and Michał Wojciechowski. "Anisotropic ornstein noninequalities". In: Analysis & PDE 10.2 (2017), pp. 351–366.
- [KT03] Jan Kristensen and Ali Taheri. "Partial regularity of strong local minimizers in the multidimensional calculus of variations". In: Archive for rational mechanics and analysis 170.1 (2003), pp. 63–89.
- [KV21] Konstantinos Koumatos and Andreas P Vikelis. "A-quasiconvexity, gårding inequalities, and applications in pde constrained problems in dynamics and statics". In: SIAM Journal on Mathematical Analysis 53.4 (2021), pp. 4178–4211.
- [Li22] Zhuolin Li. "Partial regularity for ω -minimizers of quasiconvex functionals". In: Calculus of Variations and Partial Differential Equations 61.5 (2022), p. 178.
- [Li23] Zhuolin Li. "Regularity for almost minimizers in variational problems". PhD thesis. University of Oxford, 2023.
- [LMM11] Jihoon Lee, Paul FX Müller, and Stefan Müller. "Compensated compactness, separately convex functions and interpolatory estimates between Riesz transforms and Haar projections". In: *Communications in Partial Differential Equations* 36.4 (2011), pp. 547–601.
- [LOP24] Mikyoung Lee, Jihoon Ok, and Juncheol Pyo. "Calderón-Zygmund estimates for nonlinear equations of differential forms with BMO coefficients". In: Journal of Differential Equations 397 (2024), pp. 262–288.
- [LSS23] Christina Lienstromberg, Stefan Schiffer, and Richard Schubert. "A Data-Driven Approach to Viscous Fluid Mechanics: The Stationary Case". In: Archive for Rational Mechanics and Analysis 247.2 (2023), p. 30.
- [Mar89] Paolo Marcellini. "Regularity of minimizers of integrals of the calculus of variations with non standard growth conditions". In: Archive for Rational Mechanics and Analysis 105 (1989), pp. 267–284.
- [Min06] G. Mingione. "Regularity of minima: an invitation to the dark side of the calculus of variations". In: *Appl. Math.* 51.4 (2006), pp. 355–426. DOI: 10.1007/s10778-006-0110-3.
- [MMS15] José Matias, Marco Morandotti, and Pedro M Santos. "Homogenization of functionals with linear growth in the context of A-quasiconvexity". In: Applied Mathematics & Optimization 72.3 (2015), pp. 523–547.
- [Mor52] Jr. Morrey C. B. "Quasi-convexity and the lower semicontinuity of multiple integrals". In: Pacific J. Math 2 (1952), pp. 25-53. DOI: 10.2140/pjm.1952.2.25.
- [Mül99] Stefan Müller. "Rank-one convexity implies quasiconvexity on diagonal matrices". In: International Mathematics Research Notices 1999.20 (1999), pp. 1087–1095.
- [Mur81] F. Murat. "Compacité par compensation: condition nécessaire et suffisante de continuité faible sous une hypothèse de rang constant". French. In: Ann. Scuola Norm. Sup. Pisa CI. Sci. (4) 8.1 (1981), pp. 69–102.
- [Pag+22] Valerio Pagliari, Kostas Papafitsoros, Bogdan Raibţă, and Andreas Vikelis. "Bilevel Training Schemes in Imaging for Total Variation–Type Functionals with Convex Integrands". In: SIAM Journal on Imaging Sciences 15.4 (2022), pp. 1690–1728.
- [PP04] Mariapia Palombaro and Marcello Ponsiglione. "The three divergence free matrix fields problem". In: Asymptotic Analysis 40.1 (2004), pp. 37–49.
- [Pro18] Adam Prosinski. "Closed A-p Quasiconvexity and Variational Problems with Extended Real-Valued Integrands". In: ESAIM: Control, Optimisation and Calculus of Variations 24.4 (2018), pp. 1605–1624.
- [Pro19] Adam Prosinski. "Calculus of variations in the mixed smoothness setting". PhD thesis. University of Oxford, 2019.

[Pro23]	Adam Prosinski. "Existence of minimisers of variational problems posed in spaces of mixed
	smoothness". In: Calculus of Variations and Partial Differential Equations 62.1 (2023), p. 6.
[PSU14]	Gabriel P Paternain, Mikko Salo, and Gunther Uhlmann. "Tensor tomography: progress and challenges" In: Chinese Annals of Mathematica, Series B 35.3 (2014), pp. 300–428
[PSU23]	Cabriel P Paternain Mikko Salo and Cunther Uhlmann, <i>Geometric inverse problems</i> Vol 201
[1 5025]	Cambridge University Press 2023
[Rai+24]	Bogdan Raită, Angkana Rüland, Camillo Tissot, and Antonio Tribuzio. "On Scaling Prop-
ι · j	erties for a Class of Two-Well Problems for Higher Order Homogeneous Linear Differential
	Operators". In: SIAM Journal on Mathematical Analysis 56.3 (2024), pp. 3720-3758.
[Rai19a]	B. Raiță. "Critical L^p -differentiability of $BV^{\mathbb{A}}$ -maps and canceling operators". In: Trans. Amer.
	Math. Soc. 372.10 (2019).
[Rai19b]	B. Raiță. "Potentials for <i>A</i> -quasiconvexity". In: <i>Calc. Var. PDE</i> 58.105 (2019). DOI: 10.1007/s00526-019-1544-x.
[Rai24]	Bogdan Raiță. "A simple construction of potential operators for compensated compactness".
[BB 09]	In: The Quarterly Journal of Mathematics (2024). Malemonti Madhucudana Rao and Zhong Dao Ron. Annligations of Onling engage CBC Press.
[11102]	2002
[RRT23]	Bogdan Raită, Angkana Rüland, and Camillo Tissot. "On Scaling Properties for Two-State
[]	Problems and for a Singularly Perturbed T 3 Structure". In: Acta Applicandae Mathematicae
	184.1 (2023), p. 5.
[RT22]	Angkana Rüland and Antonio Tribuzio. "On the energy scaling behaviour of a singularly
	perturbed Tartar square". In: Archive for Rational Mechanics and Analysis 243.1 (2022),
[C 04]	pp. 401-431.
[San04]	Society of Edinburgh Section A: Mathematics 134.6 (2004) pp. 1210–1227
[Sch06]	Günter Schwarz Hodge Decomposition-A method for solving boundary value problems. Springer
[Benoo]	2006.
[Sch09]	Thomas Schmidt. "Regularity of relaxed minimizers of quasiconvex variational integrals with
	(p, q)-growth". In: Archive for rational mechanics and analysis 193.2 (2009), pp. 311–337.
[Sch14]	T. Schmidt. "Partial regularity for degenerate variational problems and image restoration mod-
	els in BV". In: Indiana Univ. Math. J. 63.1 (2014), pp. 213–279. DOI: 10.1512/iumj.2014.63.5174.
[Sch24]	Stefan Schiffer. "A sufficient and necessary condition for A-quasiaffinity". In: Rendiconti del
[Sch25]	Stefan Schiffer "A free truncation and higher integrability of minimisers" In: Calculus of
[50125]	Variations and Partial Differential Equations 64.1 (2025), pp. 1–17.
[Sha12]	Vladimir Altafovich Sharafutdinov. Integral geometry of tensor fields. Vol. 1. Walter de Gruyter,
	2012.
[Sil19]	Swarnendu Sil. "Calculus of variations: a differential form approach". In: Advances in Calculus
	of Variations 12.1 (2019), pp. 57–84.
[ST23]	Massimo Sorella and Riccardo Tione. "The four-state problem and convex integration for
[GW10]	linear differential operators". In: Journal of Functional Analysis 284.4 (2023), p. 109785.
[5 W 12]	fuid flows" In: Archive for Patienal Mechanics and Analysis 206 (2012) pp. 323–366
[SW21]	Jack Skipper and Emil Wiedemann. "Lower semi-continuity for A-quasiconvex functionals
[2 = 1]	under convex restrictions". In: ESAIM: Control, Optimisation and Calculus of Variations 27
	(2021), p. 107.
[Tar78]	L. Tartar. "Une nouvelle méthode de résolution d'équations aux dérivées partielles non linéaires".
	French. In: Journées d'Analyse non linéaire. Lecture Notes in Math. 665. Berlin: Springer,
	1978, pp. 228–241.
[Tri83]	H. Triebel. Theory of functions spaces. Vol. 78. Monographs in Mathematics. Basel: Birkhauser
[Uh177]	Veriag, 1965. DOI: 10.1007/976-5-0540-0416-1. Karen Uhlenbeck "Regularity for a class of non-linear elliptic systems" In: Acta Mathematica
lomul	138 (1977), pp. 219–240.
[Van08]	Jean Van Schaftingen. "Estimates for L^1 1-vector fields under higher-order differential condi-
	tions". In: Journal of the European Mathematical Society 10.4 (2008), pp. 867–882.
[Van13]	Jean Van Schaftingen. "Limiting Sobolev inequalities for vector fields and canceling linear
	differential operators." In: Journal of the European Mathematical Society (EMS Publishing)
[van22]	Jean van Schaltingen. "Injective Ellipticity, Cancelling Operators, and Endpoint". In: Geomet-
	но ини Аниции Авреиз ој Ринсионии китишонии Еттистриез. Сентиго, ПШУ (2022), р. 239.

Max Planck Institute for Mathematics in the Sciences, Inselstrasse 22, Leipzig, 04103, Germany *E-mail address*, Z. Li: **zhuolin.li@mis.mpg.de**

Department of Mathematics and Statistics, Georgetown University,

 $3700\ \mathrm{Reservoir}\ \mathrm{Road}\ \mathrm{NW},\ \mathrm{Washington},\ \mathrm{D.C.},\ 20007$, USA

 $\mathit{E}\text{-mail}\ address,\ B.\ Raiță: br607@georgetown.edu$