# Adaptive Nonparametric Perturbations of Parametric Bayesian Models

Bohan Wu<sup>\*,1</sup>, Eli N. Weinstein<sup>\*,2</sup>, Sohrab Salehi<sup>3,4</sup>, Yixin Wang<sup>5</sup>, David M. Blei<sup>1,2,6</sup>

December 18, 2024

#### Abstract

Parametric Bayesian modeling offers a powerful and flexible toolbox for scientific data analysis. Yet the model, however detailed, may still be wrong, and this can make inferences untrustworthy. In this paper we study nonparametrically perturbed parametric (NPP) Bayesian models, in which a parametric Bayesian model is relaxed via a distortion of its likelihood. We analyze the properties of NPP models when the target of inference is the true data distribution or some functional of it, such as in causal inference. We show that NPP models can offer the robustness of nonparametric models while retaining the data efficiency of parametric models, achieving fast convergence when the parametric model is close to true. To efficiently analyze data with an NPP model, we develop a generalized Bayes procedure to approximate its posterior. We demonstrate our method by estimating causal effects of gene expression from single cell

<sup>\*</sup>Equal contribution

<sup>&</sup>lt;sup>1</sup>Department of Statistics, Columbia University, New York, NY

<sup>&</sup>lt;sup>2</sup>Data Science Institute, Columbia University, New York, NY

<sup>&</sup>lt;sup>3</sup>Computational Oncology, Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY

<sup>&</sup>lt;sup>4</sup>Irving Institute for Cancer Dynamics, Columbia University, New York, NY

<sup>&</sup>lt;sup>5</sup>Department of Statistics, University of Michigan, Ann Arbor, MI

<sup>&</sup>lt;sup>6</sup>Department of Computer Science, Columbia University, New York, NY

contact: bw2766@columbia.edu, ew2760@columbia.edu, david.blei@columbia.edu

RNA sequencing data. NPP modeling offers an efficient approach to robust Bayesian inference and can be used to robustify any parametric Bayesian model.

# **1** Introduction

Parametric Bayesian modeling offers a flexible toolbox for incorporating scientific knowledge and hypotheses into data analysis. Doing so has many benefits, such as increased data efficiency and interpretability. Yet it can also present risks, as the model remains only an approximation of reality. How can we ensure inferences are trustworthy even when the model might be wrong?

In this article, we introduce a method for robustifying a parametric Bayesian model, to obtain reliable inferences even when it might be wrong. We are specifically concerned with the possibility that the model likelihood is misspecified. Concretely, assume the data is i.i.d. from some true underlying distribution,  $x_{1:n} \stackrel{iid}{\sim} p_0$ . A parametric Bayesian model of the data is specified by a prior over an unknown parameter  $\pi(\theta)$ , together with a likelihood  $p_{\theta}$ ,

$$\begin{aligned} \theta &\sim \pi(\theta), \\ x_{1:n} &\stackrel{iid}{\sim} \mathbf{p}_{\theta}. \end{aligned}$$
 (1)

The problem is that the model may not be able to describe  $p_0$ . The induced prior over the distribution of the data,  $\pi(p_\theta)$ , only has support on a subset of all possible distributions,  $\mathscr{M}_{pm} = \{p_\theta : \theta \in \Theta\} \subset \mathscr{P}$ . If the likelihood is misspecified, such that  $p_0 \notin \mathscr{M}_{pm}$ , inferences drawn from the posterior may be unreliable, as no amount of data can overwhelm such an overconfident prior.

In practice, the statistician who uses a particular parametric likelihood  $p_{\theta}$  to analyze their data usually does not do so because they actually believe that their chosen likelihood perfectly describes the truth. They do so because the likelihood offers a convenient simplification of reality, often one grounded in specific domain knowledge. That is to say, they believe the likelihood is "probably roughly true."

The problem is how to encode such a mature opinion into a Bayesian model. In this paper,

we study a general approach. Rather than insist the distribution of the data matches the likelihood exactly, we allow for perturbations, so the data is generated as  $x \sim p_{\theta} + \delta$  instead of as  $x \sim p_{\theta}$ . We place a prior on  $\delta$  that is centered at 0 but has support everywhere, so  $p_{\theta} + \delta$  can be any distribution in  $\mathscr{P}$ . The resulting nonparametrically perturbed parametric (NPP) Bayesian model encodes the belief that the likelihood is "probably roughly true."

We show that NPP models offer a robust way of drawing inferences, while preserving the data efficiency of the original parametric model. We then develop a practical way of approximating the NPP posterior, which maintains its nice properties. The approximation does not require any change to how the parametric model's posterior is computed, instead only needing access to samples from the predictive distribution. We demonstrate the method in simulation and with an application to gene expression in cancer.

#### **1.1 Related Work**

Nonparametric perturbations of parametric Bayesian models have been proposed previously. At a high level, our contribution to this literature is to formalize NPP models under general assumptions, study their theoretical properties, and develop an efficient computational approximation.

Berger and Guglielmi (2001) consider nonparametric perturbations with a Polya tree, but in practice their method is limited to low-dimensional Euclidean data. Lyddon et al. (2018) consider perturbations using a Dirichlet process. We prove their approach sacrifices data efficiency, and propose an alternative that does not. Amin et al. (2021) develop nonparametric perturbations that retain data efficiency, but their method is restricted in scope to discrete sequence data. Miller (2019), in unpublished work, consider perturbations based on Dirichlet process mixture models, but they do not provide theoretical analysis.

We estimate functionals of the true data distribution  $p_0$ , following previous work on robust Bayesian modeling such as Lyddon et al. (2019); Pompe (2021); Dellaporta et al. (2022); Yiu et al. (2023). Other robust Bayes approaches consider a different problem, in which the target of inference is not identified from  $p_0$  because  $p_0$  has been corrupted by a population of outliers (de Finetti, 1961; Wang et al., 2017; Wang and Blei, 2018; Miller and Dunson, 2018; Bhatia et al., 2024).

Our theoretical results relate to semiparametric Bayesian causal inference methods that employ ideas like influence functions, doubly robust estimators, and Neyman orthogonality (Antonelli et al., 2020; Walker, 2024; Yiu et al., 2023). These methods offer provably robust and efficient causal inferences. We achieve similar theoretical guarantees using a very different methodological approach. Though the goals are similar, our approach is not in conflict with these influence function-based methods. In Section 4 we explain how the two methods can be combined.

Our posterior approximation builds on generalized Bayesian posteriors, which replace the model likelihood with an alternative loss (Jiang and Tanner, 2008; Zhang, 2006; Bissiri et al., 2016; Jewson et al., 2018; Shao et al., 2018; Miller, 2021; Knoblauch et al., 2022). Following Weinstein and Miller (2023), we construct a generalized Bayes factor between a parametric and nonparametric model by using a statistical divergence. Unlike in Weinstein and Miller (2023), we only need samples from the parametric model's posterior to estimate this generalized Bayes factor.

# 2 Method

#### 2.1 Target of inference

Suppose we observe  $x_{1:n} \stackrel{iid}{\sim} p_0$ . Our target of inference is a functional of the data generating distribution,  $\psi(p_0)$ , where  $\psi : \mathscr{P} \to \mathbb{R}^s$ . Here are some examples:

- Summary statistics. The target of inference could be the mean, in which case the functional is  $\psi(p_0) = \int x p_0(x) dx$ . Likewise we could infer the variance, etc.
- Loss minimizers. The target of inference could be the minimizer of a loss or utility,  $\psi(\mathbf{p}_0) = \operatorname{argmin}_{\alpha} \int \ell(x; \alpha) \mathbf{p}_0(x) dx$ . The loss can even be specified in terms of the parametric model  $\mathbf{p}_{\theta}$ , e.g. if  $\ell(x; \alpha) = \log \mathbf{p}_{\alpha}(x)$  then the target of inference is value of the model parameter closest to the truth in KL,  $\psi(\mathbf{p}_0) = \operatorname{argmin}_{\theta} \operatorname{KL}(\mathbf{p}_0 || \mathbf{p}_{\theta})$ .
- Causal effects. The target of inference could be a functional of  $p_0$  describing the effect of

an intervention (Kennedy, 2023). For example, consider data x = (a, y, w) consisting of a treatment  $a \in \{0, 1\}$ , outcome *y*, and confounder *w*. The average treatment effect is,

$$\psi(\mathbf{p}_0) = \mathbb{E}_{\mathbf{p}_0}[Y; \operatorname{do}(a=1)] - \mathbb{E}_{\mathbf{p}_0}[Y; \operatorname{do}(a=0)]$$
  
=  $\int \int y[\mathbf{p}_0(y \mid a=1, w) - \mathbf{p}_0(y \mid a=0, w)] dy \mathbf{p}_0(w) dw.$ 

The inference challenge is that if the parametric model is misspecified, we cannot learn the true data distribution, and hence cannot learn the true functional,  $\psi(p_0)$ .

#### 2.2 NPP Models

We define an NPP Bayesian model as a mixture of a parametric Bayesian model and a perturbed version of the same model,

$$\theta \sim \pi(\theta), \quad b \sim \text{Bernoulli}(\eta),$$

$$\begin{cases} p = p_{\theta} & \text{if } b = 1 \\ p \sim \pi_{\text{pert}}(p \mid p_{\theta}) & \text{if } b = 0 \end{cases}, \quad (2)$$

$$x_{1:n} \stackrel{iid}{\sim} p.$$

There are several ways to perturb the likelihood of a Bayesian model. For example,  $\pi_{pert}(p \mid p_{\theta})$  could be a Dirichlet process mixture (Antoniak, 1974; Miller, 2019),

$$\sum_{k=1}^{\infty} w_k \delta_{\phi_k} \sim \mathrm{DP}(\mathbf{p}_{\theta}, \alpha), \quad \mathbf{p} = \sum_{k=1}^{\infty} w_k \mathrm{Normal}(\phi_k, \tau).$$
(3)

Other possible perturbation models are Polya trees (Berger and Guglielmi, 2001) and Gaussian process density models (Adams et al., 2009). We assume the perturbation model is nonparametric, placing its support on any distribution over *x*, including  $p_{\theta}$ .

We want to use an NPP model instead of a conventional nonparametric model because parametric models are efficient for small data. We can think of NPP models as nonparametric models with a special spike-and-slab prior. The prior has a "spike" component that puts its mass just on the

parametric model (b = 1), and a "slab" component extending over all distributions (b = 0). This prior is nonparametric but regularizes inferences towards the parametric model. This choice of an adaptive prior enables data efficient learning.

#### **2.3 The NPP Posterior**

We perform inference with an NPP model by computing the posterior over the target functional,  $\Pi(\psi(p) \mid x_{1:n})$ . The posterior is a mixture between a parametric and nonparametric posterior,

$$\Pi(\Psi(\mathbf{p}) | x_{1:n}) = \Pi_{\mathrm{pm}}(\Psi(\mathbf{p}_{\theta}) | x_{1:n}) \eta_n + \Pi_{\mathrm{pert}}(\Psi(\mathbf{p}) | x_{1:n}) (1 - \eta_n),$$
(4)

where  $\Pi_{pm}(\psi(p_{\theta}) | x_{1:n})$  is the posterior from the parametric model,  $\Pi_{pert}(\psi(p) | x_{1:n})$  is the posterior from the nonparametric perturbation model, and  $\eta_n := \Pi(b = 1 | x_{1:n})$ . The weight  $\eta_n$  adaptively trades off between the parametric and nonparametric posterior, based on the data.

NPP models provide a *robust* and *efficient* way of learning about  $\psi(p_0)$ . To make these notions precise, we compare NPP models to two alternatives: (1) the original parametric model, given by  $\theta \sim \pi(\theta), x_{1:n} \sim p_{\theta}$ , and (2) a nonparametric model with a generic prior,  $p \sim \pi_{np}(p), x_{1:n} \sim p$ , where  $\pi_{np}(p)$  has support over all distributions on  $\mathscr{X}$ . We use  $\Pi_{pm}(\cdot | x_{1:n})$  to denote the parametric model posterior,  $\Pi_{np}(\cdot | x_{1:n})$  to denote the nonparametric model posterior, and  $\Pi(\cdot | x_{1:n})$  to denote the NPP model posterior. We will show that the parametric model is efficient but not robust, while the nonparametric model is robust but not efficient. Formal statements and proofs of the following results are in Section 4.

**Misspecified case.** We want inferences about  $\psi(p_0)$  to be *robust* to model misspecification, such that they always converge to the truth with enough data. The NPP model is robust: Theorem 1 shows that  $\Pi(\psi(p) | x_{1:n}) \rightarrow \delta_{\psi(p_0)}$  even when  $p_0 \notin \mathscr{M}_{pm}$ . To see why, recall from Eq. (4) that the NPP posterior is a mixture of a parametric and nonparametric posterior. The parametric posterior will not converge to the truth,  $\Pi_{pm}(\psi(p_{\theta}) | x_{1:n}) \not\rightarrow \delta_{\psi(p_0)}$ , but the nonparametric component will,  $\Pi_{pert}(\psi(p) | x_{1:n}) \rightarrow \delta_{\psi(p_0)}$ . The overall NPP posterior converges to the truth because the mixing

weight  $\eta_n$  asymptotically places all weight on the nonparametric component: if  $p_0 \notin \mathcal{M}_{pm}$  then  $\eta_n \to 0$  a.s. as  $n \to \infty$  (Proposition 1).

To see why intuitively, write  $\eta_n$  in terms of a Bayes factor comparing the marginal likelihood of the data under the parametric model to the marginal likelihood under the nonparametric model,

$$\frac{\eta_n}{1-\eta_n} = \frac{\prod_{\text{pm}}(x_{1:n})\eta}{\prod_{\text{pert}}(x_{1:n})(1-\eta)} =: BF_n,$$
(5)

where

$$\Pi_{\rm pm}(x_{1:n}) = \int \prod_{i=1}^n p_{\theta}(x_i) \pi(\theta) d\theta,$$

and

$$\Pi_{\text{pert}}(x_{1:n}) = \int \int \prod_{i=1}^{n} \mathbf{p}(x_i) \pi_{\text{pert}}(\mathbf{p}|\mathbf{p}_{\theta}) \pi(\theta) d\mathbf{p} d\theta$$

When the parametric model is misspecified, the Bayes factor prefers the more expressive nonparametric model, since it can match the data distribution, while the parametric model cannot. Hence,  $BF_n \rightarrow 0$ , so the mixing weight  $\eta_n \rightarrow 0$  a.s. as  $n \rightarrow \infty$  (Proposition 1).

In summary, the NPP model is robust:  $\Pi(\psi(p) | x_{1:n}) \to \delta_{\psi(p_0)}$  even when when  $p_0 \notin \mathscr{M}_{pm}$ . In this way, the NPP model behaves like a generic nonparametric model, for which we also expect  $\Pi_{np}(\psi(p) | x_{1:n}) \to \delta_{\psi(p_0)}$ . A parametric model, by contrast, is not robust: in general,  $\Pi_{pm}(\psi(p_\theta) | x_{1:n}) \neq \delta_{\psi(p_0)}$  when  $p_0 \notin \mathscr{M}_{pm}$ .

Well-specified case. We next consider the case where the parametric model is well-specified,  $p_0 \in \mathscr{M}_{pm}$ . In this case, the NPP model converges to the truth at an efficient rate: the posterior  $\Pi(\psi(p) \mid x_{1:n})$  contracts at a rate  $1/\sqrt{n}$  around  $\psi(p_0)$  when  $p_0 \in \mathscr{M}_{pm}$ . To see this, return to the decomposition in Eq. (4). Both the parametric posterior  $\Pi_{pm}(\psi(p_\theta) \mid x_{1:n})$  and the nonparametric posterior  $\Pi_{pert}(\psi(p) \mid x_{1:n})$  will converge to the truth, but the parametric posterior will converge faster, at a rate  $1/\sqrt{n}$ , while the nonparametric component will in general converge more slowly. The NPP posterior inherits the rate of the parametric posterior because the mixing weight asymptotically places all weight on the parametric component: if  $p_0 \in \mathcal{M}_{pm}$  then  $\eta_n \to 1$  a.s. as  $n \to \infty$  (Proposition 1).

To see why  $\eta_n \to 1$  intuitively, consider again the Bayes factor, Eq. (5). Both the parametric and the nonparametric model describe the data distribution, but the parametric model uses fewer parameters than the nonparametric model: a finite number, rather than infinite. Moreover, the parametric model is nested inside the nonparametric model, since it corresponds to sampling  $p = p_{\theta}$ from  $\pi_{pert}(p | p_{\theta})$ . By Bayesian Occam's razor, the Bayes factor prefers the simpler parametric model,  $BF_n \to \infty$  (Dawid, 2011; Hong and Preston, 2012; MacKay, 2003, Chap. 28). Hence  $\eta_n \to 1$ .

In summary, when the parametric model is correctly specified,  $p_0 \in \mathcal{M}_{pm}$ , the NPP posterior  $\Pi(\psi(p) | x_{1:n})$  converges to the truth  $\psi(p_0)$  at an efficient parametric rate of  $1/\sqrt{n}$ . So here, the NPP posterior behaves like a parametric Bayes posterior, for which we also expect a convergence rate of  $1/\sqrt{n}$  by the Berstein-von Mises theorem. In a nonparametric model, however, the convergence rate is generally slower, e.g.  $\Pi_{np}(\psi(p) | x_{1:n})$  may converge at a rate of  $1/\sqrt[3]{n}$  or worse (Ghosal et al., 2000).

**Finite samples.** When the parametric model is misspecified, the NPP model is only guaranteed to converge at a nonparametric rate. Heuristically, however, we expect the NPP posterior to converge towards the truth about as quickly as the parametric posterior, up until the parametric model stops providing a good description of the data. Specifically, the Bartlett-Lindley effect says the Bayes factor will place more weight on the simpler model up until there is enough data to determine that model is wrong (Miller and Dunson, 2018).

#### 2.4 Generalized Bayes Approximation

We have seen that NPP models offer both robustness and efficiency, but this comes at a computational cost. Computing the NPP posterior (Eq. (4)) requires computing not only the parametric model posterior, but also (1) the mixing weight  $\eta_n$ , and (2) the nonparametric posterior,  $\Pi_{pert} (\psi(p) | x_{1:n})$ . In this section we introduce approximations to both terms that are efficient to compute, practical

to implement and preserve robustness and efficiency.

**Mixing weight.** The mixing weight  $\eta_n$  depends on the Bayes factor BF<sub>n</sub> (Eq. (5)), which compares the marginal likelihood of the parametric model to the marginal likelihood of a nonparametric alternative. Computing marginal likelihoods is often challenging, especially for nonparametric models. We propose to replace the Bayes factor with a generalized Bayes factor, which uses divergences instead of marginal likelihoods to evaluate the parametric model (Miller, 2021; Weinstein and Miller, 2023).

Let  $D(p, p_0)$  denote a divergence between probability distributions, which satisfies  $D(p, p_0) = 0$ when  $p = p_0$  and  $D(p, p_0) > 0$  when  $p \neq p_0$ . Let  $D_n(p, p_0)$  denote an estimate of  $D(p, p_0)$  based on the data. Concretely, the divergences we consider are the Wasserstein distance, the maximum mean discrepancy and the kernelized Stein discrepancy, though other choices are also possible (Gretton et al., 2012; Liu et al., 2016). We use the divergence to construct a generalized Bayes factor,

$$gBF_n := \Xi \left( \frac{\mathbb{E}[D_n(p_\theta, p_0)]}{\mathbb{E}[D_n(p_\theta, p_0) \mid x_{1:n}]} (n+1)^{-r} \right) \frac{\eta}{1-\eta}.$$
(6)

Here,  $\mathbb{E}[D_n(p_{\theta}, p_0)]$  is the expected divergence between the parametric likelihood and the true data distribution under the prior  $\theta \sim \pi(\theta)$ , while  $\mathbb{E}[D_n(p_{\theta}, p_0)]$  is the expected value under the posterior  $\theta \sim \Pi_{pm}(\theta \mid x_{1:n})$ . The rate r > 0 is a hyperparameter, and the function  $\Xi(x) = \exp(1 - \frac{1}{x})x$  is monotonic and satisfies  $\Xi(x) \to 0$  as  $x \to 0$  and  $\Xi(x) \to \infty$  as  $x \to \infty$ . Although Eq. (6) does not contain the standard Bayes factor as a special case, it is related: like the standard marginal likelihood, it averages over an estimate model-data mismatch. Now, to approximate the NPP posterior, we propose to use  $\hat{\eta}_n = 1/(1 + gBF_n^{-1})$ .

The aim of the approximation  $\hat{\eta}_n$  is not to directly estimate the mixing weight  $\eta_n$ , but rather to match its statistical behavior. When the parametric model is misspecified,  $\hat{\eta}_n \rightarrow 0$ , just like  $\eta_n$ . The reason is that  $D(p_{\theta}, p_0) > 0$  for all  $\theta$  when  $p_0 \notin \mathscr{M}_{pm}$ , so  $\mathbb{E}[D_n(p_{\theta}, p_0) | x_{1:n}]$  will converge to a positive value. On the other hand, when the parametric model is well-specified,  $\hat{\eta}_n \rightarrow 1$ , just like  $\eta_n$ . The reason is that the posterior will concentrate at  $p_0$ , so  $\mathbb{E}[D_n(p_\theta, p_0) | x_{1:n}] \to 0$ . So long as the rate hyperparameter *r* is chosen small enough that  $\mathbb{E}[D_n(p_\theta, p_0) | x_{1:n}](n+1)^r \to 0$  also, we will have  $\hat{\eta}_n \to 1$ . So, the generalized Bayes factor is asymptotically *consistent* in detecting misspecification.

In addition to achieving the same asymptotic limits as the original Bayes factor, the generalized Bayes factor also approaches those limits at a similar rate, thanks to the choice of function  $\Xi(x)$  (Appendix A). Before we observe data, the generalized Bayes factor coincides with the prior,  $g_{BF_0} = BF_0 = \eta/(1-\eta)$ .

But the key advantage of the generalized Bayes factor is computation: it requires only the ability to draw samples from the prior and posterior of the parametric model, to approximate  $\mathbb{E}[D_n(p_{\theta}, p_0)]$  and  $\mathbb{E}[D_n(p_{\theta}, p_0) | x_{1:n}]$ . The marginal likelihood of the parametric model does not appear. The nonparametric model does not appear at all.

The generalized Bayes factor can also be understood as a form of predictive check on the parametric model (Gelman et al., 1996; Moran et al., 2024; Li and Huggins, 2024). Specifically, the term  $\mathbb{E}[D_n(p_{\theta}, p_0) | x_{1:n}]$  compares the posterior over the parametric model's predictive distribution,  $\Pi_{pm}(p_{\theta} | x_{1:n})$ , to data. It uses  $D_n$  as measure of model-data discrepancy.

**Choice of divergence and rate.** We consider several types of divergence, which prioritize different kinds of mismatch between the model and data.

Wasserstein distance. The Wasserstein measures the difference between two distributions as the cost of transforming one into the other by transporting probability mass. The *p*-Wasserstein distance is defined as W<sup>p</sup><sub>p</sub>(p,q) := inf<sub>π∈Π(p,q)</sub> E<sub>π</sub>[||X − Y||<sup>p</sup><sub>2</sub>], where Π(p,q) represents the set of all couplings between p and q, i.e. the set of all joint distributions with p and q as marginals, and || · ||<sub>2</sub> denotes the Euclidean metric on ℝ<sup>κ</sup>. We can estimate the distance between the model and the data using *m* samples drawn from the model. We draw the same number of samples as data points, *m* = *n*, by default.

• Maximum mean discrepancy (MMD). The maximum mean discrepancy focuses on the worst case difference in expected value that two distributions assign to the same function in a reproducing kernel Hilbert space (RKHS). It is given by

$$MMD^{2}(p,q) := \sup_{\|f\|_{\mathscr{H}_{k}} \le 1} |\mathbb{E}_{p}[f(X)] - \mathbb{E}_{q}[f(X)]|^{2}$$
(7)

$$= \mathbb{E}_{X,X'\sim p}\left[k(X,X')\right] - 2\mathbb{E}_{X\sim p,Y\sim q}\left[k(X,Y)\right] + \mathbb{E}_{Y,Y'\sim q}\left[k(Y,Y')\right].$$
(8)

where  $\mathscr{H}_k$  is an RKHS and  $||f||_{\mathscr{H}_k}$  is the norm of a function in that RKHS. We can estimate the MMD by drawing *m* samples from the model and comparing them to the data using the RKHS's kernel *k*. We again set m = n by default.

Kernelized Stein discrepancy (KSD). The kernelized Stein discrepancy focuses on differences in the (Stein) score function between two distributions, i.e. differences in the gradient of their log densities. It is given by KSD(p,q) = E<sub>X,X'~p</sub> [Δ<sub>q,p</sub>(X)<sup>T</sup>k(X,X')Δ<sub>q,p</sub>(X')] where Δ<sub>q,p</sub>(x) := ∇<sub>x</sub> log p(x) - ∇<sub>x</sub> log q(x). To estimate the KSD we do not need samples from the model.

Both the MMD and KSD rely on a kernel, which allows user control over the aspects of a distribution to prioritize, though it also introduces the need for some degree of hyperparameter selection (which is absent in the Wasserstein distance). Although the MMD and the KSD are sensitive to a rescaling of the kernel, the generalized Bayes factor is scale-invariant, since it normalizes by the prior discrepancy. For the MMD and KSD we have a consistent generalized Bayes factor so long as we choose a rate 0 < r < 1/2, i.e. slower than the parametric rate (Theorem 3). For Wasserstein, we need to choose a slower rate when the dimension  $\kappa$  of the data space is greater than four, namely  $r \in (0, n^{-2/(\kappa \lor 4)})$  for  $\mathscr{X} \subseteq \mathbb{R}^{\kappa}$  (Theorem S2 in Appendix D.3.1).

**Nonparametric posterior.** Next we consider approximating the nonparametric term  $\Pi_{pert}(\psi(p) | x_{1:n})$  in the NPP posterior. This term can be challenging: the robustness and efficiency of the NPP

model depends on the nonparametric perturbation  $\pi_{pert}(p \mid p_{\theta})$  covering all distributions on  $\mathscr{X}$ , so computing the posterior requires integrating over a large space. But now the mixing weight,  $\hat{\eta}_n$ , does not depend on the nonparametric perturbation, and so we no longer need so much flexibility. We propose instead to use any semiparametric model that can consistently estimate  $\psi(p_0)$ . That is, we consider a model  $p \sim \hat{\pi}_{pert}(p), x_{1:n} \sim p$  such that  $\hat{\Pi}_{pert}(\psi(p) \mid x_{1:n}) \rightarrow \delta_{\psi(p_0)}$ . We propose to approximate the term  $\Pi_{pert}(\psi(p) \mid x_{1:n})$  in the NPP posterior with  $\hat{\Pi}_{pert}(\psi(p) \mid x_{1:n})$ .

To see how this approximation can simplify computation, consider data consisting of a treatment and a response,  $x_i = (y_i, a_i)$ , and a target functional that is the expected response to a given treatment,  $\psi(p) = \mathbb{E}[Y \mid a_*] = \int y p(y \mid a_*) dy$ . An NPP model would require us to place a prior over all distributions on  $\mathscr{A} \times \mathscr{Y}$ . But to estimate the functional consistently, we can instead use a semiparametric model  $y_i \sim \text{Normal}(f(a_i), 1), f \sim \pi_f(f)$ , and place a nonparametric prior on functions  $f : \mathscr{A} \to \mathscr{Y}$ . For example,  $\pi_f$  could be a Gaussian process prior or a Bayesian additive regression tree. These semiparametric models cannot describe all distributions over  $\mathscr{A} \times \mathscr{Y}$ , e.g. they cannot describe a distribution where the conditional  $p(y \mid a_*)$  is not Gaussian. But they can still estimate any conditional expectation  $\mathbb{E}[Y \mid a_*]$ . Moreover, the posterior is convenient to calculate, as we can rely on existing software for Gaussian processes and Bayesian additive regression trees.

Summary. In summary, our approximation to the NPP is

$$\Pi(\Psi(\mathbf{p}) \mid x_{1:n}) \approx \hat{\Pi}(\Psi(\mathbf{p}) \mid x_{1:n}) := \Pi_{\mathrm{pm}}(\Psi(\mathbf{p}_{\theta}) \mid x_{1:n}) \,\hat{\eta}_n + \hat{\Pi}_{\mathrm{pert}}(\Psi(\mathbf{p}) \mid x_{1:n}) \,(1 - \hat{\eta}_n), \quad (9)$$

where  $\hat{\eta}_n$  comes from the generalized Bayes factor and  $\hat{\Pi}_{pert}$  comes from a semiparametric model. We refer to this approximation as the *generalized NPP* (*gNPP*). In Section 4 we prove formally that the gNPP preserves the robustness and efficiency of the NPP: for any  $p_0$ ,  $\hat{\Pi}(\psi(p) | x_{1:n}) \rightarrow \delta_{\psi(p_0)}$ , and for  $p_0 \in \mathscr{M}_{pm}$ ,  $\hat{\Pi}(\psi(p) | x_{1:n})$  converges at a rate  $1/\sqrt{n}$ .

The key to the gNPP is replacing the Bayes factor between parametric and nonparametric models with a generalized Bayes factor based on a divergence. In doing so, we avoid the need to

compute the marginal likelihood of the parametric model or of the nonparametric perturbation. We also reduce the requirements and constraints on the perturbation model we can use: it no longer needs to be fully nonparametric, and the parametric model does not need to be nested inside it.

# **3** Synthetic Data Illustration

We illustrate the behavior of NPP models and the gNPP approximation in a synthetic data setting. We follow the basic setup of Lyddon et al. (2018). Consider a Gaussian parametric model,

$$\boldsymbol{\theta} \sim \mathcal{N}(0,1), \quad x_{1:n} \stackrel{iid}{\sim} \mathcal{N}(\boldsymbol{\theta},1).$$
 (10)

We study inference in the well-specified setting where the true distribution is  $\mathcal{N}(0,1)$ , and in the misspecified setting, where the true distribution is a *skew* normal with the same mean and variance but skew parameter 10. Details on the simulations and models are in Appendix **B**, and code for these experiments is at https://github.com/bohanwu424/npp.

#### 3.1 Polya Tree NPP

We first study an NPP model that perturbs the parametric model with a Polya tree. The advantage of Polya trees is that they admit closed form marginal likelihoods, making accurate approximation the Bayes factor in the NPP posterior possible. This provides a careful check of our theoretical results. The Polya tree NPP model is,

$$\theta \sim \mathcal{N}(0,1) \quad z \sim \text{Bernoulli}(\eta = 0.1),$$

$$\begin{cases} p = \mathcal{N}(\theta,1) & \text{if } b = 1 \\ p \sim \text{PolyaTree}(\mathcal{N}(\theta,1),h) & h \sim \text{Exponential}(1) & \text{if } b = 0 \end{cases},$$

$$x_{1:n} \stackrel{iid}{\sim} p.$$

$$(11)$$



Figure 1: a. **Synthetic data results.** The log Bayes factor (black) and log generalized Bayes factor (colors) comparing the parametric model to a nonparametric alternative; positive values indicate the parametric model is favored. Solid lines describe the well-specified case and dashed lines the misspecified case. b,c. The KL divergence between the true data density and the posterior predictive of each model, in the misspecified case (b) and the well-specified case (c). d,e. The absolute difference between the posterior mean estimate of the median and the true median, for each model, in the misspecified case (d) and the well-specified case (e). Here we use the MMD gNPP approximation. f. Calibration of the MMD gNPP approximation. We plot how often, across independent simulations, the posterior credible interval covers the true median. The nominal coverage is 90% (dashed).

Here, we parameterize the Polya tree as in Berger and Guglielmi (2001); it is a distribution over distributions, with mean  $\mathcal{N}(\theta, 1)$  and scale parameter *h*. We draw the scale parameter *h* from an exponential prior. We compute  $\Pi_{\text{pert}}(x_{1:n})$  using the procedure in Berger and Guglielmi (2001).

We first examine convergence of the posterior distribution to the truth, measuring the KL

divergence between the true distribution and the posterior predictive of the NPP model, averaged over 100 independent datasets. We compare to the performance of a parametric model (Eq. (11) given b = 1) and to the performance of a nonparametric model (Eq. (11) given b = 0).

The NPP model is efficient: in the well-specified case, it matches the faster convergence of the parametric model (Figure 1b). The NPP model is also robust: in the misspecified case, it converges to the true distribution, even though the parametric model does not (Figure 1b). Note that at very small amounts of data (n = 5), the parametric model provides better estimates than the nonparametric model despite its misspecification, as it offers a useful approximation of the underlying distribution. The NPP inherits some of these benefits, outperforming the nonparametric model at n = 5.

The NPP achieves its adaptive behavior due to the Bayes factor (Figure 1a, black), which prefers the parametric model when that model is well-specified (solid) and the nonparametric model otherwise (dashed). With small amounts of data (n = 5), the Bayes factor prefers the parametric model even if it is misspecified, and even though the prior  $\eta = 0.1$  prefers the nonparametric model. This is an example of the Bartlett-Lindley effect: the parametric model is preferred even when it is misspecified since it provides a better approximation given the amount of available data.

#### **3.2** Generalized Bayes Approximation to the NPP

We next study the generalized Bayes approximation to the posterior (gNPP). To design our approximation, we must first choose a functional of interest. Following Lyddon et al. (2018), we focus on the median,  $\psi(p) = \operatorname{argmin}_{\alpha} \mathbb{E}_p[|X - \alpha|]$ . The parametric model assumes the mean and median of the distribution are both  $\theta$ , but in the misspecified case, when the true data distribution  $p_0$  is skew normal, the median is about -0.2. For the nonparametric component, we use a Bayesian bootstrap, a Dirichlet process with concentration 0 (Rubin, 1981). For the generalized Bayes factor, we consider the 2-Wasserstein divergence, the MMD with an inverse multiquadric (IMQ) kernel, and the KSD with the same kernel. We compare to the parametric model and to a nonparametric model, the Bayesian bootstrap on its own.

We quantify the performance of models' point estimates in terms of the absolute difference between the posterior mean of the median and the true median,  $|\mathbb{E}[\psi(p) | x_{1:n}] - \psi(p_0)|$ . The gNPP approximation is efficient: in the well-specified case, it matches the fast convergence of the parametric model. Figure 1e shows the MMD version; the others are similar, see Figure S1. The gNPP approximation is also robust: in the misspecified case, it converges to the true median, even though the parametric model does not, see Figure 1e and Figure S1. In the misspecified case, it also slightly outperforms the nonparametric model at small amounts of data, when using the MMD and KSD.

The gNPP achieves adaptation thanks to the generalized Bayes factor (Figure 1a), which matches the asymptotic behavior of the Bayes factor. At small *n* we see a Bartlett-Lindley effect in the MMD and KSD gNPPs, as they prefer the misspecified parametric model at n = 5 despite the prior  $\eta = 0.1$ . This explains their performance improvement over the nonparametric model. Note the effect is not theoretically guaranteed to always occur. In this case it is absent from the Wasserstein gNPP.

We also examined uncertainty quantification. We checked whether posterior credible sets are calibrated and achieve frequentist coverage. We computed how often, across independent simulations, the true median fell in the posterior 95% credible interval; with correct calibration, it should fall in the interval 95% of time. In the well-specified case, the gNPP is well calibrated, as is the parametric and nonparametric model (Figure S2). In the misspecified case, the parametric model (Figure 1f, Figure S2).

A key hyperparameter in the gNPP is the convergence rate r. We set r = 0.49 for the MMD and KSD versions, based on the intuition that r should be as large as possible to enable efficient convergence in the misspecified case, while still below 0.5 to preserve convergence in the wellspecified case. While this default worked well for the MMD and KSD versions, in the Wasserstein version it led to a gBF that failed to favor the nonparametric model even with n = 500 datapoints, and even though the gBF slowly trended towards larger value (Figure S3). Instead, setting r = 0.1 led to improved performance by effectively discriminating between the well-specified and misspecified cases at lower values of *n* (Figure 1a, blue). So although r = 0.49 may be a reasonable default, problem-specific tuning can improve performance.

## 4 Theory

We establish the frequentist properties of NPP models and the gNPP approximation. In particular, we prove they are (a) robust, in the sense that the posterior converges to the true data distribution, and (b) efficient, in the sense that the posterior converges at a parametric rate when the underlying parametric model is well-specified. Throughout we assume a population distribution  $p_0$  over  $\mathscr{X} \subseteq \mathbb{R}^{\kappa}$  that produces observations  $x_{1:n}$ . We also assume a unique parameter  $\theta_0$  in an open set  $\Theta \subseteq \mathbb{R}^d$  that minimizes the KL divergence to the population distribution,  $D_{\text{KL}}(p_0 || p_{\theta})$ . Detailed statements and proofs of each result are in Appendix D.

#### 4.1 NPP models

Model selection consistency. We first analyze the mixing weight  $\eta_n$  in the NPP. Proposition 1 establishes general conditions for its model selection consistency:  $\eta_n$  asymptotically selects the parametric model when that model is correct, and the nonparametric model otherwise. The assumptions (Assumptions S5 to S7) follow the "prior mass and testing" framework for Bayesian nonparametrics (Ghosal and Van der Vaart, 2001). Informally, we need the NPP's prior to put sufficient mass in a reverse KL ball around p<sub>0</sub> (Assumption S6), and, when the parametric model is well-specified, the nonparametric prior should put much less mass around the truth than the parametric model (Assumptions S5 and S7). The result then follows directly from Ghosal et al. (2008, Corollary 3.1).

**Proposition 1** ( $\eta_n$  is consistent for model selection). Assume the marginal density  $p_{pm}(x_{1:n}) := \int_{\Theta} p_{\theta}(x_{1:n}) d\Pi_{pm}(\theta)$  is well-defined. Under Assumptions S5 to S7 given in Appendix C.2,

*1.*  $\eta_n \to 1$  *a.s.*  $[\mathbb{P}_0^{\infty}]$ , if  $p_0 \in \mathscr{M}_{pm}$ .

2.  $\eta_n \to 0 \text{ a.s. } [\mathbb{P}_0^{\infty}], \text{ if } p_0 \notin \mathscr{M}_{pm}.$ 

We check if the assumptions of Proposition 1 are satisfied for non-parametric perturbations based on a Dirichlet process and on a Dirichlet process mixture model.

**Example 1** (Dirichlet process perturbations are not consistent). Consider an NPP model (Eq. (2)) with a Dirichlet process perturbation  $p \sim DP(p_{\theta}, \alpha)$ . For this choice of perturbation, the mixing weight fails to satisfy model selection consistency. The technical assumption that fails is Assumption S6 which requires the DP prior to put sufficient mass around  $\{p : D_{KL}(p_0 || p) < \varepsilon^2\}$  for small  $\varepsilon$ . But, since the DP prior is only supported on discrete measures, it puts zero mass on this reverse KL ball for any non-discrete  $p_0$ .

One implication of this result is that the robust Bayes approach studied by Lyddon et al. (2018), in which a parametric model is perturbed by a Dirichlet process, is robust but not necessarily efficient.

**Example 2** (Dirichlet process normal mixture perturbations are consistent). Consider a nonparametric perturbation based on a Dirichlet process normal mixture, as described in Eqs. (2) and (3). Unlike the Dirichlet process, this perturbation describes continuous densities over  $\mathscr{X}$ . When  $p_0$  is smooth, Proposition S3 justifies the assumptions of Proposition 1, implying  $\eta_n$  is model selection consistent.

Semiparametric BvM. We now establish robustness and efficiency for the NPP posterior. Recall the target of inference is defined via a functional  $\psi : \mathscr{P}(\mathscr{X}) \mapsto \mathbb{R}^s, s < \infty$ . To establish efficiency, we prove a semiparametric Bernstein-von Mises (BvM) theorem: when  $p_0 \in \mathscr{M}_{pm}$ , the posterior over the functional  $\psi(p)$  is asymptotically normal, with a standard deviation proportional to  $1/\sqrt{n}$ . To establish robustness, we show that even when  $p_0 \notin \mathscr{M}_{pm}$ , the posterior concentrates at  $\psi(p_0)$ .

The NPP inherits its asymptotic normality from the underlying parametric model. The standard Bernstein-von Mises theorem says the parametric posterior  $\Pi_{pm}(d\theta \mid x_{1:n})$  is asymptotically normal, so under smoothness conditions, the posterior over  $\psi(p_{\theta})$  will be asymptotically normal as well.

**Assumption 1** (The parametric model is asymptotically normal). As  $n \to \infty$ ,

- (a)  $\sqrt{n}(\hat{\theta}_{MLE} \theta_0) \xrightarrow{w} \mathcal{N}(0, V_{\theta_0}^{-1})$ , where  $\xrightarrow{w}$  denotes weak convergence and  $\hat{\theta}_{MLE}$  is the maximum likelihood estimate of  $\theta$ , and
- (b)  $d_{TV}\left(\sqrt{n}\left(\theta \hat{\theta}_{MLE}\right), \mathcal{N}\left(0, V_{\theta_0}^{-1}\right)\right) \stackrel{\mathbb{P}_0}{\to} 0$  for  $\theta \sim \prod_{\text{pm}} \left(\cdot \mid x_{1:n}\right)$ , with convergence in first and second moments in  $[\mathbb{P}_0]$ -probability.

Conditions (a) and (b) assume the parametric MLE and posterior are asymptotically normal at the  $\sqrt{n}$ -rate. Condition (b) is the Bernstein-von Mises theorem for regular parametric models, and guarantees the frequentist coverage of posterior credible regions will asymptotically match its nominal level. The convergence in first two moments follows, for example, from convergence of the posterior to a normal distribution in the Wasserstein metric.

**Assumption 2** (The target functional is smooth). The function  $\chi(\theta) := \psi(\mathbf{p}_{\theta}) - \psi(\mathbf{p}_{\hat{\theta}_{MLE}})$  is twice differentiable, with  $\dot{\chi}_{\theta_0}^{\top} V_{\theta_0}^{-1} \dot{\chi}_{\theta_0} \succ 0$ ,  $\ddot{\chi}_{\theta}$  being continuous at  $\theta_0$  and  $\|\ddot{\chi}_{\theta_0}\|_2 < \infty$ .

We now establish our main result. To describe convergence of the NPP posterior to a normal distribution, we use the bounded Lipschitz distance  $d_{BL}$ , which metrizes the topology of weak convergence (Definition S4 and Theorem S1) (Castillo and Rousseau, 2015).

**Theorem 1** (NPP models are efficient and robust). Let Proposition 1 and Assumptions 1 and 2 hold. Let  $\Pi(\tilde{\psi}_n(\mathbf{p}) | x_{1:n})$  denote the pushforward measure of the NPP posterior through  $\tilde{\psi}_n(\mathbf{p}) := \sqrt{n}(\psi(\mathbf{p}) - \psi(\mathbf{p}_{\hat{\theta}_{MLE}}))$ . If  $\mathbf{p}_0 \in \mathcal{M}_{pm}$ , then

$$d_{BL}\left(\Pi\left(\tilde{\psi}_{n}(\mathbf{p}) \mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\boldsymbol{\chi}}_{\theta_{0}}^{\top} \boldsymbol{V}_{\theta_{0}}^{-1} \dot{\boldsymbol{\chi}}_{\theta_{0}}\right)\right) \xrightarrow{\mathbb{P}_{0}} 0 \quad (Efficiency).$$
(12)

Further assume the nonparametric perturbation  $\Pi_{pert}(\psi(p) \mid x_{1:n})$  is consistent at  $\psi(p_0)$ . Then if  $p_0 \notin \mathscr{M}_{pm}$ ,

$$d_{BL}\left(\Pi\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_0)}\right) \xrightarrow{\mathbb{P}_0} 0 \quad (Robustness).$$
(13)

*Hence the NPP posterior*  $\Pi(\psi(\mathbf{p}) \mid x_{1:n})$  *is also consistent at*  $\psi(\mathbf{p}_0)$ *.* 

When the parametric model is correctly specified, Theorem 1 tell us that the NPP posterior of the target estimand achieves the  $\sqrt{n}$ -rate and the optimal variance in the asymptotic minimax sense (Van der Vaart, 2000). When the parametric model is misspecified, the NPP posterior is still robust as it converges to the true  $\psi(p_0)$ . Sufficient conditions for the consistency of  $\Pi_{pert}(\psi(p) | x_{1:n})$  are (1) consistency of the nonparametric posterior for all  $p_{\theta}$ , e.g. Dirichlet process normal mixture models, and (2) continuity of the functional  $\psi$  with respect to the weak topology.

The efficiency guarantee could be extended to the misspecified setting if the nonparametric posterior  $\Pi_{pert}(\psi(p) \mid x_{1:n})$  achieves semiparametric efficiency for the target estimand  $\psi$ . This is satisfied, for example, by the one-step corrected posterior (Yiu et al., 2023).

#### 4.2 gNPP approximation

We establish the efficiency and robustness of the gNPP approximation, proving an analogue of Theorem 1 for our generalized Bayes inference approach (Theorem 4). To do so, we first show that the generalized Bayes factor satisfies model selection consistency (Theorem 2). Our results apply to a broad class of divergences, but we verify the conditions and provide explicit contraction rates for the Wasserstein, MMD and KSD (Theorems S2 to S4). The contraction rates justify our guidance for setting the rate hyperparameter r.

Model selection consistency. We first analyze the model selection consistency of the mixing weight  $\hat{\eta}_n$  in the gNPP. Recall  $\hat{\eta}_n := 1/(1 + gBF_n^{-1})$ , where  $gBF_n$  depends on a posterior expected empirical divergence  $\mathbb{E}_{pm} [D_{m,n}(p_{\theta}, p_0) | x_{1:n}]$  (Eq. (6)). We are interested in the asymptotic behavior of this posterior divergence, showing it correctly detects model misspecification.

To begin, we study general divergences D that are semimetrics or depend on one, in the sense that  $D(p,q) = \rho^k(p,q)$  for  $k \in \mathbb{N}$  and  $\rho$  a semimetric. The Wasserstein and MMD fit this form; the KSD is asymmetric and hence does not, so we later extend our arguments. We also assume in the general theory that the estimated divergence  $D_{m,n}(p,q)$  takes the form of a *plug-in* estimator  $D_{m,n}(p,q) = D(p^m,q^n)$ , where  $p^m$  is the *m*-sample empirical distribution of p, and  $q^n$  is the *n*-sample empirical distribution of q. The empirical *p*-Wasserstein distance is directly given by this plug-in estimator, while the plug-in form of the MMD is its V-statistic. We later extend our arguments to the MMD U-statistic, which matches the plug-in form with a vanishing error,  $O(n^{-1} + m^{-1})$ .

We make regularity assumptions on the divergence.

**Assumption 3** (The divergence is well-behaved). *Assume*  $D(p,q) = \rho^k(p,q)$  *where*  $k \in \mathbb{N}$  *and*  $\rho$  *is a semimetric, and that* 

- (a)  $\rho$  is continuous in the weak topology and  $\sup_{\theta \in \Theta} \rho(p_{\theta}, p_0) < \infty$ .
- (b) There exists an  $M_n n^{-1/2}$  neighborhood of  $\theta_0$  where  $M_n \to \infty$  such that the mapping  $\theta \mapsto \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0})$  is twice differentiable and  $\|\nabla^2_{\theta} \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0})\|_2$  is uniformly bounded.

Assumption 3(a) requires the divergence to be uniformly bounded. For example, with the Wasserstein distance, the condition is satisfied if  $\mathscr{X}$  is compact. For MMD, the condition is satisfied if the kernel is uniformly bounded. Assumption 3(b) requires local Lipschitz smoothness of the function  $\theta \mapsto \rho(p_{\theta}, p_{\theta_0})$  within an asymptotically vanishing ball around  $\theta_0$ , where the radius is at least  $n^{-\alpha/2}$ , with  $\alpha < 1$ . This condition is typically satisfied by smooth parametric models (Barp et al., 2019).

We also assume that the empirical divergence converges to the true divergence, at least when the model likelihood is in a neighborhood of the truth.

Assumption 4 (The empirical divergence converges). There exists a sequence  $r_{m,n} \rightarrow 0$  such that

- (a) For any sequence  $\theta_n \to \theta_0$ , as  $m, n \to \infty$  with  $m/(m+n) \to c \in (0, 1)$ , we have  $\mathbb{E}[D_{m,n}(p_{\theta_n}, p_0)] = D(p_{\theta_0}, p_0) + O(r_{m,n})$ . The expectation is over m samples from  $p_{\theta_n}$  and n samples from  $p_0$ .
- (b) There exists a function  $\mathscr{V}(\theta)$  that is finite at  $\theta_0$  and continuous at  $\theta_0$  such that for all (m,n) large enough,  $Var(D_{m,n}(p_{\theta},p_0)) \leq r_{m,n}^2 \mathscr{V}(\theta)$ , where Var denotes the variance.

Assumption 4 requires an asymptotic bound on the expectation of  $D_{m,n}(p_{\theta}, p_0)$  and a finite-sample variance bound around  $\theta_0$ , both converging at a rate  $r_{m,n}$ . These assumptions are local, applying

only to a neighborhood of  $\theta_0$ . They are also weaker than assuming uniform convergence of the mean or the variance of  $D_{m,n}(p_{\theta}, p_0)$  in a neighborhood around  $\theta_0$ .

When we specialize Assumption 4 to specific divergences, we can verify the assumption more closely. For example, for the MMD, the plug-in V-statistic matches the U-statistic up to an ignorable error, and for the U-statistic, Assumption 4(a) is satisfied, since the statistic is unbiased, and Assumption 4(b) follows from the variance formula of two-sample U-statistics and kernel regularity conditions.

We now study the posterior expected value of the empirical divergence, under the parametric model. To make the analysis more tractable theoretically, we assume the samples used to approximate  $p_0$  in  $D_{m,n}(p_{\theta}, p_0)$  are independent of those used to compute the posterior. This can be achieved by splitting data and cross-fitting (Chernozhukov et al., 2018). However, we expect the model selection consistency of  $\hat{\eta}_n$  (Theorem 3) to remain valid even without sample splitting, and in the empirical studies we do not apply it.

**Theorem 2** (The posterior expected empirical divergence converges at a rate  $r_{m,n} \vee n^{-1}$ ). Let Assumptions 1, 3 and 4 be satisfied. As  $m, n \to \infty$  with  $\frac{n}{n+m} \to c \in (0,1)$ ,  $\mathbb{E}_{pm}[D_{m,n}(p_{\theta},p_0) | x_{1:n}]$ converges in  $[\mathbb{P}_0^{\infty}]$ -probability to  $D(p_{\theta_0},p_0)$  at the rate of  $r_{m,n} \vee n^{-1}$ :

$$\mathbb{E}_{pm}[D_{m,n}(p_{\theta}, p_0) \mid x_{1:n}] = D(p_{\theta_0}, p_0) + O_{\mathbb{P}_0}(r_{m,n} \vee n^{-1}).$$
(14)

So the posterior empirical divergence converges to the true minimal divergence between the model class  $\mathcal{M}_{pm}$  and  $p_0$ , at a rate that is the slower of (a)  $r_{m,n}$ , the convergence rate of  $D_{m,n}$ , and (b)  $n^{-1}$ .

Theorem 2 applies to general divergences. For the Wasserstein, MMD, and KSD in particular, we derive the following contraction rates (detailed theorems and conditions are in Appendix D.3).

1. *p*-Wasserstein Distance. The empirical Wasserstein divergence  $\mathbb{E}_{pm} \left[ W_p^p(\mathbf{p}_{\theta}^m, \mathbf{p}_0^n) \mid x_{1:n} \right]$  converges to  $W_p^p(\mathbf{p}_{\theta_0}, \mathbf{p}_0)$  at a rate of  $O_{\mathbb{P}_0} \left( n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)} \right)$ , where  $\kappa$  is the dimension of  $\mathscr{X}$  (Appendix D.3.1).

- 2. **MMD.** The empirical MMD  $\mathbb{E}_{pm} \left[ \text{MMD}_U^2 \left( p_{\theta}^m, p_0^n \right) | x_{1:n} \right]$ , based on the U-statistic for the MMD, converges to MMD<sup>2</sup>  $\left( p_{\theta_0}, p_0 \right)$  at a rate of  $O_{\mathbb{P}_0} \left( n^{-1/2} + m^{-1/2} \right)$  (Appendix D.3.2).
- 3. **KSD.** The empirical KSD  $\mathbb{E}_{pm} [KSD_U(p_0^n, p_\theta) | x_{1:n}]$ , based on the U-statistic for the KSD, converges to  $KSD(p_0, p_{\theta_0})$  at a rate of  $O_{\mathbb{P}_0}(n^{-1/2})$  (Appendix D.3.3).

For simplicity, in our implementation, we set m = n for the two-sample empirical divergences (Wasserstein and MMD), so the rate  $r_{m,n}$  simplifies to  $r_n := r_{n,n}$ . In this case, for the Wasserstein, we can take  $r_n = n^{-2/(\kappa \vee 4)}$ , and for the MMD and KSD, we can take  $r_n = n^{-1/2}$ .

We have shown that the posterior expected empirical divergence converges to the true divergence at a known rate,  $r_n$ . We can now set the rate hyperparameter r in the generalized Bayes factor to be slower than this known rate, i.e.  $r \in (0, \frac{2}{\kappa \lor 4})$  for the Wasserstein and  $r \in (0, 1/2)$  for MMD and KSD. Then, the generalized Bayes factor is model selection consistent.

**Theorem 3** ( $\hat{\eta}_n$  is consistent for model selection). Let  $\mathbb{E}_{pm}[D_n(p_\theta, p_0)]$  be bounded in  $[\mathbb{P}_0^{\infty}]$ -probability,  $\eta > 0$  and assume  $\mathbb{E}_{pm}[D_n(p_\theta, p_0) | x_{1:n}] = D(p_{\theta_0}, p_0) + r_n$ , where  $D(\cdot, \cdot)$  is a statistical divergence. Choose r > 0 such that  $r_n(n+1)^r = o(1)$ . Then, as  $n \to \infty$ :

- 1. *if*  $\mathbf{p}_0 \in \mathscr{M}_{pm}$ , then  $\hat{\boldsymbol{\eta}}_n \to 1$  a.s.  $[\mathbb{P}_0^{\infty}]$ ,
- 2. *if*  $\mathbf{p}_0 \notin \mathscr{M}_{pm}$ , then  $\hat{\boldsymbol{\eta}}_n \to 0$  a.s.  $[\mathbb{P}_0^{\infty}]$ .

Note  $\mathbb{E}_{pm}[D_n(p_{\theta}, p_0)]$  is bounded in probability when, for example,  $D_n(p_{\theta}, p_0)$  is uniformly bounded or satisfies a uniform law of large numbers.

**Semiparametric BvM.** The gNPP approximation uses the generalized Bayes factor to adaptively trade off between parametric and nonparametric models. We have shown the generalized Bayes factor is model selection consistent. As a result, the gNPP approximation converges efficiently when the parametric model is well-specified, but still converges to the truth when it is misspecified.

**Theorem 4** (gNPP approximations are efficient and robust). Let Assumptions 1 to 4 hold for m = nand  $r_n := r_{n,n}$ . Assume also that  $\mathbb{E}_{pm}[D_n(p_{\theta}, p_0)]$  is bounded in  $[\mathbb{P}_0]$ -probability. Let r < 1 be a positive constant such that  $r_n(n+1)^r = o(1)$ . Let  $\hat{\Pi}(\tilde{\psi}_n(\mathbf{p}) | x_{1:n})$  denote the pushforward measure of  $\tilde{\psi}_n(\mathbf{p}) := \sqrt{n}(\psi(\mathbf{p}) - \psi(\mathbf{p}_{\hat{\theta}_{MLE}}))$  onto the gNPP posterior. If  $\mathbf{p}_0 \in \mathcal{M}_{pm}$ , then

$$d_{BL}\left(\hat{\Pi}\left(\tilde{\Psi}_{n}(\mathbf{p}) \mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\chi}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\chi}_{\theta_{0}}\right)\right) \xrightarrow{\mathbb{P}_{0}} 0 \quad (Efficiency).$$

$$(15)$$

Furthermore, assume that the nonparametric posterior  $\hat{\Pi}_{pert}(\psi(p) \mid x_{1:n})$  is consistent at  $\psi(p_0)$ . Then if  $p_0 \notin \mathcal{M}_{pm}$ ,

$$d_{BL}\left(\hat{\Pi}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_0)}\right) \stackrel{\mathbb{P}_0}{\to} 0 \quad (Robustness), \tag{16}$$

hence the gNPP posterior  $\Pi(\psi(\mathbf{p}) | x_{1:n})$  is also consistent at  $\psi(\mathbf{p}_0)$ .

We can replace Assumptions 3 and 4 with assumptions specific to the Wasserstein, MMD and KSD, as detailed in Appendix D.3.

## 5 Application: Estimating the Effects of Gene Expression

A cell's behavior depends crucially on its genes' *expression levels*, or the number of RNA transcripts of each gene in the cell. An important question in cell biology is how the expression level of one gene affects another. Answering such questions can help scientists understand cells' behavior, and may help develop drugs which modify gene expression levels or activity.

We develop a robust method to estimate the causal effect of one gene's expression level on another's. To do so, we leverage observational data about the expression level of genes in individual cells, measured with single cell RNA sequencing (Kolodziejczyk et al., 2015). We focus on analyzing primary patient data, which is often hard to obtain and hence limited in scale. Bayesian parametric models offer data-efficient inference, but robustness is crucial given the importance of the estimates to human health.

We specify the causal effect as a functional of the data distribution. We start from a parametric Bayesian model that incorporates prior information from an auxiliary dataset, a *single cell atlas*. We robustify this parametric model by creating a gNPP approximation to the NPP posterior, with



Figure 2: **Causal model.** To analyze the effects of gene expression, we assume this causal graphical model. a. The initial model where all variables are observed. b. The model after intervening on the treatment variable *A*. The goal is to estimate the effect of the treatment gene on the outcome gene, as highlighted in blue.

a BART-based nonparametric perturbation and an MMD-based generalized Bayes factor.

We apply the gNPP model to conduct an exploratory analysis of data from an ovarian cancer patient. We estimate the causal effects of genes that are potential therapeutic targets. Our results reveal that interventions on some genes may have unexpected effects in this patient.

Additional details are in Appendix E. Code and data can be found at https://github. com/bohanwu424/npp.

#### 5.1 Setup

**Estimand.** We are interested in the effect of the expression level of a treatment gene on the expression level of an outcome gene. The challenge for causal inference is that there may be confounders, factors which affect the expression of both the treatment and the outcome genes. An important source of confounding is *transcription factors*, proteins which directly modify the expression level of many other genes. To correct for this confounding, we adjust for the expression level of the transcription factor genes in each cell.

Formally, we consider the causal graph in Figure 2, where a is the treatment gene expression level, y is the outcome gene expression level, and w is a vector of expression levels for confounding

genes, i.e. transcription factors. We are interested in the effect of an intervention which sets the expression level of *a* to  $a_{\star}$ . From Figure 2, the distribution of the outcome after the intervention is given by the backdoor adjustment,  $p(y | do(a_{\star})) = \int p(y | a_{\star}, w)p(w)dw$  (Pearl, 2009).

We compare the average outcome when the treatment gene is highly expressed versus when it is unexpressed. To define "highly expressed" we take the 98th quantile of the empirical distribution of treatment expression, denoted  $q_{98}(a)$ . We define the *average treatment effect*,

ATE (p) = 
$$\mathbb{E}_{p}[Y; \operatorname{do}(a_{\star} = q_{98}(a))] - \mathbb{E}_{p}[Y; \operatorname{do}(a_{\star} = 0)]$$
  
=  $\int \int yp(y \mid q_{98}(a), w)p(w)dydw - \int \int yp(y \mid 0, w)p(w)dydw.$  (17)

Our target functional is  $\psi(p) = ATE(p)$ , where p(a, y, w) is the joint distribution over treatment, outcome and confounders.

**Parametric model** We first introduce a parametric Bayesian model to estimate the causal effect  $\psi(p_0)$ . This parametric model incorporates information from a single cell atlas dataset, a large collection of scRNAseq data from donors without cancer.

Rather than using the full vector of confounding gene expression w, the model uses a lowdimensional representation z(w) derived from the atlas. Specifically, we set  $z(\cdot)$  to be the projection of w onto the first ten principal components of the atlas. Such low-dimensional representations of gene expression – whether derived from PCA or other approaches – have been found to provide an effective summary of *cell type* and *cell state* (Lopez et al., 2018). By using representations learned from a large atlas, biologists aim to better estimate the cell type and state underlying the noisy expression levels in a small dataset (Hao et al., 2021). We apply this idea to obtain a lowdimensional representation of confounding.

Our parametric Bayesian model depends linearly on the treatment and on the low-dimensional

representation of the confounders,

$$\mathbf{p}_{\boldsymbol{\theta}}(\boldsymbol{y} \mid \boldsymbol{a}, \boldsymbol{w}) = \mathcal{N}(\boldsymbol{c} + \boldsymbol{\tau} \boldsymbol{a} + \boldsymbol{\gamma}^{\mathsf{T}} \boldsymbol{z}(\boldsymbol{w}), \boldsymbol{\sigma}^{2}), \tag{18}$$

where  $\theta := [c, \tau, \gamma]$  and  $\mathcal{N}(\mu, \sigma^2)$  denotes a normal distribution with mean  $\mu$  and variance  $\sigma^2$ . The parameter *c* is an intercept,  $\tau$  is the coefficient on the treatment and  $\gamma$  is a vector of coefficients on the confounder representation. We use an improper prior  $\pi(\theta) \propto 1$  and specify  $\sigma \sim \text{HalfNormal}(0, 1)$ . To compute the empirical divergence of the prior, we approximate the improper prior by a uniform distribution over the hypercube with slice [-100, 100]. We place a Dirichlet process prior on p(a, w) with concentration approaching zero, such that the posterior over the ATE is obtained by the Bayesian bootstrap.

We expect the parametric model to learn efficiently even from limited numbers of cells, since the cell representation z(w) is low-dimensional and hence so is the model. However, the parametric model may be misspecified, for instance if the ovarian cancer patient has a cells in a pathological state that is not present in the healthy atlas. In this case, the representation z(w) may project out important variation in w, ignoring a source of confounding and leading to unreliable inferences.

**gNPP model** We introduce a gNPP model for robust inference. For the nonparametric model, we use a Bayesian additive regression tree (BART) with a fixed propensity score correction, an established method for nonparametric Bayesian causal inference (Hill, 2011; Hahn et al., 2020). Since single cell RNA expression data is highly non-normal, we adjust the BART model with a Yeo-Johnson transformation (Yeo and Johnson, 2000). We place a Dirichlet process prior on p(a, w) with concentration zero, such that the posterior over the ATE is obtained by a Bayesian bootstrap. The model does not incorporate prior information from the atlas. For the generalized Bayes factor, we use the MMD with rate r = 0.49.

**Data and preprocessing** We study an scRNAseq dataset consisting of 544 T cells from a patient with ovarian cancer (Vázquez-García et al., 2022). The atlas contains 261,611 T cells collected

across 17 sites/tissues from 12 organ donors without cancer (Domínguez Conde et al., 2022). We log transform and standardize the expression data following standard practice in the field (Heumos et al., 2023). As potential confounders, we adjust for 157 transcription factors with highly variable expression. To find highly variable genes, we use the method of Stuart et al. (2019) as implemented in scanpy (Wolf et al., 2018). Briefly, the read counts are z-score normalized (using a regularized standard deviation), then genes are ranked by their empirical variance.

### 5.2 Results

A route to treating cancer is to intervene on a patient's immune system, such that it can better attack the cancer. One potential strategy is to modify regulatory T cells, which suppress immune attack, to be more like cytotoxic T cells, which conduct immune attack. FOXP3 is a transcription factor that makes T cells regulatory, so it may be a good drug target for achieving this goal (Revenko et al., 2022). Will decreasing the expression of FOXP3 make this ovarian cancer patient's T cells more cytotoxic? To address this question, we estimate the causal effect of FOXP3 on the expression of key genes that T cells use to clear tumors. We also consider interventions on another gene in Appendix E.2.

**GZMH** We first consider the effect of FOXP3 on granzyme H and K (GZMH and GZMK), proteases which digests tumor proteins. For granzyme H (GZMH), under the parametric model, the posterior probability that the ATE is positive is low, 0.12 (Figure 3a). This suggests the ATE is likely negative, so an intervention that decreases FOXP3 expression will increase GZMH expression. In other words, the parametric model infers that intervening on FOXP3 will make these T cells more capable of attacking tumors. However, there are reasons to question this conclusion. The parametric model's  $R^2$  is low, at 0.12, suggesting it explains only a small portion of the variance in the data. Moreover, its residual histogram and QQ plot suggest strong deviations from normality (Figure S4). So the parametric model may be poorly specified.

The gNPP draws inferences that are robust to misspecification. We find a small generalized



Figure 3: Effect of FOXP3 on GZMH. a. Posterior probability of the ATE being positive under the parametric, nonparametric, and gNPP models. *n* denotes the size of the (subsampled) dataset. Values are the estimated median from 10 independent data subsamples and model samples. b. Generalized mixing weights,  $\hat{\eta}_n$ . The estimated confidence interval (CI) is across independent data subsamples and model samples.

mixing weights of 0.18, placing most of the posterior weight on the nonparametric model (Figure 3b). However, the posterior probability of a positive ATE under the gNPP model remains nearly unchanged, at 0.12. So, the conclusion that decreasing FOXP3 expression will increase GZMH expression is robust to possible model misspecification.

**GZMK** We next consider the effect of FOXP3 on granzyme K (GZMK). The posterior probability that the ATE is positive, under the parametric model, is 0.43 (Figure 4a). The fact that this value is close to 0.5 suggests there is substantial uncertainty in the effect of FOXP3 on GZMK, and there may be no effect at all. The gNPP, however, revises this estimate downward to 0.33, providing more confidence that decreasing FOXP3 expression will increase GZMK expression. In this case, the gNPP still places considerable weight on the parametric model, with a generalized Bayes factor of 0.69 (Figure 4b).



Figure 4: Effect of FOXP3 on GZMK. a. Posterior probability of the ATE being positive under the parametric, nonparametric, and gNPP models. *n* denotes the size of the (subsampled) dataset. b. Generalized mixing weights,  $\hat{\eta}_n$ . CI: confidence interval across independent data subsamples and model samples.

We subsampled the data down to smaller numbers of cells and reran the analysis, averaging over ten independent subsamples. We found that with smaller amounts of data the gNPP model relies even more heavily on the parametric model, producing nearly the same posterior probability of a positive ATE (Figure 4a). The gNPP's estimates only begin to diverge from the parametric model at around n = 200 cells.

**NKG7** We next consider the effect of FOXP3 on the granule protein NKG7, another key component of T cells' attack. The parametric model suggests that decreasing FOXP3 expression will actually decrease NKG7, with the posterior probability of a positive ATE of 0.99 (Figure 5a). Biologically this is unexpected, since regulatory T cells do not express much NKG7 in general (Szabo et al., 2019). Correspondingly, the gNPP increases our uncertainty in the finding, revising the posterior probability down to 0.77 (Figure 5a). Here, the gNPP has strong confidence that the parametric model is wrong, with a generalized mixing weight of 0.1.



Figure 5: Effect of FOXP3 on NKG7. a. Posterior probability of the ATE being positive under the parametric, nonparametric, and gNPP models. *n* denotes the size of the (subsampled) dataset. b. Generalized mixing weights,  $\hat{\eta}_n$ . CI: confidence interval across independent data subsamples and model samples.

The gNPP adaptively smooths between the parametric and nonparametric posteriors, so its posterior probability of a positive ATE is quite stable here even if we sub-sample the data to below 100 datapoints (Figure 5a). The nonparametric model, by contrast, produces more variable estimates: at n = 100, it produces a posterior probability of 0.35, compared to the full dataset value of 0.76.

# 6 Discussion

We proposed NPP models to robustify parametric Bayesian models, guarding against misspecification without sacrificing data efficiency. We also developed a generalized Bayes approximation to NPP model posteriors that enjoys more scalable computation and implementation. The basic technique is: (1) use a backup nonparametric or semiparametric model that is consistent for the target of inference and (2) mix the parametric posterior with the nonparametric posterior based on how well the parametric model describes the data.

The robustness and efficiency of NPP models bear analogy to the properties of doubly robust estimators used for causal inference. Specifically, consider a doubly robust estimator that uses a parametric outcome model and a nonparametric propensity model. When the outcome model is well-specified, the estimator will converge at a parametric rate. When the outcome model is misspecified, the estimator will still converge to the true effect, but at a slower rate (Bang and Robins, 2005; Antonelli et al., 2020). An NPP model with a parametric outcome model perturbed by a nonparametric outcome model offers the same guarantees, even without a propensity model. But the NPP framework is not restricted to causal functionals.

Overall, NPP models offer a practical approach to combining scientific domain knowledge with flexible machine learning models, by regularizing inferences from a nonparametric model towards a parametric model.

## Acknowledgments

We wish to thank Jeffrey Miller for useful discussions, including introducing us to the idea of nonparametrically perturbing parametric Bayesian models.

## References

- Adams, R. P., Murray, I., and MacKay, D. J. C. (2009). Nonparametric Bayesian density modeling with Gaussian processes.
- Amin, A. N., Weinstein, E. N., and Marks, D. S. (2021). A generative nonparametric Bayesian model for whole genomes. In *Advances in Neural Information Processing Systems*.
- Antonelli, J., Papadogeorgou, G., and Dominici, F. (2020). Causal inference in high dimensions: A marriage between Bayesian modeling and good frequentist properties. *Biometrics*.
- Antoniak, C. E. (1974). Mixtures of Dirichlet processes with applications to Bayesian nonparametric problems. *Annals of Statistics*, pages 1152–1174.

- Bang, H. and Robins, J. M. (2005). Doubly robust estimation in missing data and causal inference models. *Biometrics*, 61:962–973.
- Barp, A., Briol, F. X., Duncan, A. B., Girolami, M., and Mackey, L. (2019). Minimum Stein discrepancy estimators. Advances in Neural Information Processing Systems, 32.
- Berger, J. and Guglielmi, A. (2001). Bayesian and conditional frequentist testing of a parametric model versus nonparametric alternatives. *Journal of the American Statistical Association*, 96:174–184.
- Bhatia, K., Ma, Y.-A., Dragan, A. D., Bartlett, P. L., and Jordan, M. I. (2024). Bayesian robustness: A nonasymptotic viewpoint. *Journal of the American Statistical Association*, 119(546):1112– 1123.
- Bissiri, P. G., Holmes, C. C., and Walker, S. G. (2016). A general framework for updating belief distributions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 78:1103–1130.
- Blackwell, D. and MacQueen, J. B. (1973). Ferguson distributions via Pólya urn schemes. *Annals of Statistics*, 1(2):353–355.
- Briol, F.-X., Barp, A., Duncan, A. B., and Girolami, M. (2019). Statistical inference for generative models with maximum mean discrepancy.
- Castillo, I. and Rousseau, J. (2015). A Bernstein–von Mises theorem for smooth functionals in semiparametric models. *Annals of Statistics*, 43:2353–2383.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *Econometrics Journal*, 21:C1–C68.
- Chipman, H. A., George, E. I., and McCulloch, R. E. (2010). BART: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4:266–298.
- Dawid, A. P. (2011). Posterior model probabilities. In Bandyopadhyay, P. S. and Forster, M. R., editors, *Philosophy of Statistics*, volume 7, pages 607–630. North-Holland, Amsterdam.

- de Finetti, B. (1961). The Bayesian approach to the rejection of outliers. *Proc. Fourth Berkeley Symp. on Math. Statist. and Prob*, 1.
- Dellaporta, C., Knoblauch, J., Damoulas, T., and Briol, F.-X. (2022). Robust Bayesian inference for simulator-based models via the MMD posterior bootstrap. *International Conference on Artificial Intelligence and Statistics*, pages 943–970.
- Domínguez Conde, C., Xu, C., Jarvis, L., Rainbow, D., Wells, S., Gomes, T., Howlett, S., Suchanek, O., Polanski, K., King, H., et al. (2022). Cross-tissue immune cell analysis reveals tissue-specific features in humans. *Science*, 376(6594):eabl5197.
- Gelman, A., Meng, X. L., and Stern, H. (1996). Posterior predictive assessment of model fitness via realized discrepancies. *Statistica Sinica*, 6:733–807.
- Ghosal, S., Ghosh, J. K., and Van der Vaart, A. W. (2000). Convergence rates of posterior distributions. *Annals of Statistics*, 28(2):500–531.
- Ghosal, S., Lember, J., and Van der Vaart, A. W. (2008). Nonparametric Bayesian model selection and averaging. *Electronic Journal of Statistics*, 2:63–89.
- Ghosal, S. and Van der Vaart, A. W. (2001). Entropies and rates of convergence for maximum likelihood and bayes estimation for mixtures of normal densities. *Annals of Statistics*, 29(5):1233–1263.
- Ghosal, S. and Van der Vaart, A. W. (2017). *Fundamentals of nonparametric Bayesian inference*. Cambridge University Press.
- Gorham, J. and Mackey, L. (2017). Measuring sample quality with kernels. In *International Conference on Machine Learning (ICML)*, pages 1292–1301.
- Gretton, A., Borgwardt, K. M., Rasch, M. J., Schölkopf, B., and Smola, A. (2012). A kernel two-sample test. *Journal of Machine Learning Research*, 13.
- Guntuboyina, A. and Sen, B. (2012).  $L_1$  covering numbers for uniformly bounded convex functions. *Journal of Machine Learning Research*, 23.

- Hahn, P. R., Murray, J. S., and Carvalho, C. M. (2020). Bayesian regression tree models for causal inference: Regularization, confounding, and heterogeneous effects (with discussion). *Bayesian Analysis*, 15:965–1056.
- Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk,
  A. J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. *Cell*, 184(13):3573–3587.
- Heumos, L., Schaar, A. C., Lance, C., Litinetskaya, A., Drost, F., Zappia, L., Lücken, M. D., Strobl, D. C., Henao, J., Curion, F., et al. (2023). Best practices for single-cell analysis across modalities. *Nature Reviews Genetics*, 24(8):550–572.
- Hill, J. (2011). Bayesian nonparametric modeling for causal inference. *Journal of Computational and Graphical Statistics*, 20:217–240.
- Hong, H. and Preston, B. (2005). Nonnested model selection criteria. *unpublished*.
- Hong, H. and Preston, B. (2012). Bayesian averaging, prediction and nonnested model selection. *Journal of Econometrics*, 167(2):358–369.
- Jewson, J., Smith, J. Q., and Holmes, C. (2018). Principles of Bayesian inference using general divergence criteria. *Entropy*, 20(6):442.
- Jiang, W. and Tanner, M. A. (2008). Gibbs posterior for variable selection in high-dimensional classification and data mining. *Annals of Statistics*, 36(5):2207–2231.
- Kennedy, E. H. (2023). Semiparametric doubly robust targeted double machine learning: a review.
- Knoblauch, J., Jewson, J., and Damoulas, T. (2022). An optimization-centric view on Bayes'rule: Reviewing and generalizing variational inference. *Journal of Machine Learning Research*, 23(132):1–109.
- Kolodziejczyk, A. A., Kim, J. K., Svensson, V., Marioni, J. C., and Teichmann, S. A. (2015). The technology and biology of single-cell RNA sequencing. *Molecular Cell*, 58(4):610–620.

- Kumari, S., Arora, M., Singh, J., Chauhan, S. S., Kumar, S., and Chopra, A. (2021). Lselectin expression is associated with inflammatory microenvironment and favourable prognosis in breast cancer. *3 Biotech*, 11(2):38.
- Li, J. and Huggins, J. H. (2024). Calibrated model criticism using split predictive checks.
- Liu, Q., Lee, J. D., and Jordan, M. (2016). A kernelized Stein discrepancy for goodness-of-fit tests. *33rd International Conference on Machine Learning*, 1.
- Lopez, R., Regier, J., Cole, M. B., Jordan, M. I., and Yosef, N. (2018). Deep generative modeling for single-cell transcriptomics. *Nature Methods*, 15(12):1053–1058.
- Luxburg, U. V. and Bousquet, O. (2004). Distance-based classification with Lipschitz functions. *Journal of Machine Learning Research*.
- Lyddon, S., Walker, S., and Holmes, C. (2018). Nonparametric learning from Bayesian models with randomized objective functions. *Advances in Neural Information Processing Systems*.
- Lyddon, S. P., Holmes, C. C., and Walker, S. G. (2019). General Bayesian updating and the losslikelihood bootstrap. *Biometrika*, 106:465–478.
- MacKay, D. J. (2003). *Information Theory, Inference and Learning Algorithms*. Cambridge university press.
- Manole, T. and Niles-Weed, J. (2024). Sharp convergence rates for empirical optimal transport with smooth costs. *Annals of Applied Probability*, 34.
- Miller, J. (2019). Flexible perturbation models for robustness to misspecification. Unpublished.
- Miller, J. W. (2021). Asymptotic normality, concentration, and coverage of generalized posteriors. *Journal of Machine Learning Research*, 22:1–53.
- Miller, J. W. and Dunson, D. B. (2018). Robust Bayesian inference via coarsening. *Journal of the American Statistical Association*.

Moran, G. E., Blei, D. M., and Ranganath, R. (2024). Holdout predictive checks for Bayesian model criticism. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 86.
Pearl, J. (2009). *Causality*. Cambridge University Press.
- Pompe, E. (2021). Introducing prior information in weighted likelihood bootstrap with applications to model misspecification.
- Quiroga, M., Garay, P. G., Alonso, J. M., Loyola, J. M., and Martin, O. A. (2023). Bayesian additive regression trees for probabilistic programming.
- Revenko, A., Carnevalli, L. S., Sinclair, C., Johnson, B., Peter, A., Taylor, M., Hettrick, L., Chapman, M., Klein, S., Solanki, A., Gattis, D., Watt, A., Hughes, A. M., Magiera, L., Kar, G., Ireland, L., Mele, D. A., Sah, V., Singh, M., Walton, J., Mairesse, M., King, M., Edbrooke, M., Lyne, P., Barry, S. T., Fawell, S., Goldberg, F. W., and MacLeod, A. R. (2022). Direct targeting of FOXP3 in tregs with AZD8701, a novel antisense oligonucleotide to relieve immunosuppression in cancer. *Journal for ImmunoTherapy of Cancer*, 10(4):e003892.
- Rubin, D. B. (1981). The Bayesian bootstrap. Annals of Statistics, pages 130–134.
- Serfling, R. J. (2009). Approximation Theorems of Mathematical Statistics. John Wiley & Sons.
- Shao, S., Jacob, P. E., Ding, J., and Tarokh, V. (2018). Bayesian model comparison with the Hyvärinen score: Computation and consistency. *J. Am. Stat. Assoc.*, pages 1–24.
- Shen, W., Tokdar, S., and Ghosal, S. (2013). Adaptive Bayesian multivariate density estimation with Dirichlet mixtures. *Biometrika*, 100:623–640.
- Sriperumbudur, B. K., Fukumizu, K., and Lanckriet, G. R. (2011). Universality, characteristic kernels and RKHS embedding of measures. *Journal of Machine Learning Research*, 12.
- Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive integration of single-cell data. *Cell*, 177(7):1888–1902.
- Szabo, P. A., Levitin, H. M., Miron, M., Snyder, M. E., Senda, T., Yuan, J., Cheng, Y. L., Bush, E. C., Dogra, P., Thapa, P., Farber, D. L., and Sims, P. A. (2019). Single-cell transcriptomics of human t cells reveals tissue and activation signatures in health and disease. *Nature Communications*, 10.
- Tsybakov, A. B. (2008). Introduction to Nonparametric Estimation. Springer New York.

Van der Vaart, A. W. (2000). Asymptotic Statistics, volume 3. Cambridge university press.

Van der Vaart, A. W. and Wellner, J. A. (2023). *Weak Convergence and Empirical Processes*. Springer International Publishing.

Villani, C. (2003). Topics in Optimal Transportation. American Mathematical Society.

- Vázquez-García, I., Uhlitz, F., Ceglia, N., Lim, J. L., Wu, M., Mohibullah, N., Niyazov, J., Ruiz, A. E. B., Boehm, K. M., Bojilova, V., Fong, C. J., Funnell, T., Grewal, D., Havasov, E., Leung, S., Pasha, A., Patel, D. M., Pourmaleki, M., Rusk, N., Shi, H., Vanguri, R., Williams, M. J., Zhang, A. W., Broach, V., Chi, D. S., Paula, A. D. C., Gardner, G. J., Kim, S. H., Lennon, M., Roche, K. L., Sonoda, Y., Zivanovic, O., Kundra, R., Viale, A., Derakhshan, F. N., Geneslaw, L., Bhaloo, S. I., Maroldi, A., Nunez, R., Pareja, F., Stylianou, A., Vahdatinia, M., Bykov, Y., Grisham, R. N., Liu, Y. L., Lakhman, Y., Nikolovski, I., Kelly, D., Gao, J., Schietinger, A., Hollmann, T. J., Bakhoum, S. F., Soslow, R. A., Ellenson, L. H., Abu-Rustum, N. R., Aghajanian, C., Friedman, C. F., McPherson, A., Weigelt, B., Zamarin, D., and Shah, S. P. (2022). Ovarian cancer mutational processes drive site-specific immune evasion. *Nature*, 612.
- Walker, C. D. (2024). Parametrization, prior independence, and the Semiparametric Bernstein-von Mises theorem for the partially linear model.
- Wang, C. and Blei, D. M. (2018). A general method for robust Bayesian modeling. *Bayesian Analysis*, 13.
- Wang, Y., Kucukelbir, A., and Blei, D. M. (2017). Robust probabilistic modeling with Bayesian data reweighting. In 34th International Conference on Machine Learning, ICML 2017, volume 7.
- Weinstein, E. N. and Miller, J. W. (2023). Bayesian data selection. *Journal of Machine Learning Research*, 24:1–72.
- Wolf, F. A., Angerer, P., and Theis, F. J. (2018). Scanpy: large-scale single-cell gene expression data analysis. *Genome Biology*, 19:1–5.

- Yeo, I. K. and Johnson, R. A. (2001). A uniform strong law of large numbers for U -statistics with application to transforming to near symmetry. *Statistics and Probability Letters*, 51.
- Yeo, I. N. and Johnson, R. A. (2000). A new family of power transformations to improve normality or symmetry. *Biometrika*, 87.
- Yiu, A., Fong, E., Holmes, C., and Rousseau, J. (2023). Semiparametric posterior corrections.
- Zhang, T. (2006). Information-theoretic upper and lower bounds for statistical estimation. *IEEE Transactions on Information Theory*, 52(4):1307–1321.

## **Supplementary Material**

## A Scaling of the Generalized Bayes Factor

In Section 2, we define the generalized Bayes factor using the transformation  $\Xi$ . Here, we show that the transformed variable aligns with the typical scaling behavior of the Bayes factor (Hong and Preston, 2005; Dawid, 2011). First note,

$$-\log gBF_n = \frac{\mathbb{E}\left[D_n\left(p_{\theta}, p_0\right) \mid x_{1:n}\right]}{\mathbb{E}\left[D_n\left(p_{\theta}, p_0\right)\right]} \left(n+1\right)^r - 1 + \log\left(\frac{\mathbb{E}\left[D_n\left(p_{\theta}, p_0\right) \mid x_{1:n}\right]}{\mathbb{E}\left[D_n\left(p_{\theta}, p_0\right)\right]} \left(n+1\right)^r\right).$$

Suppose the assumptions of Theorem 4 hold. Then the empirical divergence satisfies

$$\mathbb{E}\left[\mathsf{D}_{n}(\mathsf{p}_{\theta},\mathsf{p}_{0}) \mid x_{1:n}\right] = \mathsf{D}(\mathsf{p}_{\theta_{0}},\mathsf{p}_{0}) + r_{n},$$

for some rate  $r_n \to 0$  such that  $r_n(n+1)^r = o(1)$ . In practice, for the divergences we study,  $r_n = n^{-\tilde{r}}$ , and we choose  $0 < r < \tilde{r}$ . This yields the following scaling behavior: if  $p_0 \in \mathscr{M}_{pm}$ , then  $\log gBF_n = O((\tilde{r} - r)\log n)$ ; if  $p_0 \notin \mathscr{M}_{pm}$ , then  $-\log gBF_n = O(n^r)$ .

This aligns with the standard Bayes factor, comparing a high-dimensional model to a nested low-dimensional model, where, if  $p_0$  is in the low-dimensional model class, then  $\log BF_n = O(\log n)$ ; otherwise  $-\log BF_n = O(n)$  (Hong and Preston, 2005; Dawid, 2011).

The scaling behavior of the generalized Bayes factor also reveals the tradeoffs involved in setting the hyperparameter r. Choosing smaller values of r leads to faster convergence in the well-specified case, but slower convergence in the misspecified case.

## **B** Details on Synthetic Experiments

In the NPP, we estimate the KL divergence between  $p_0$  and the posterior predictive using Monte Carlo, with 1000 heldout samples from the true distribution. We use m = n to estimate the divergence in the gNPP, for the MMD version and Wasserstein version. We use the IMQ kernel  $k(x,y) = (c^2 + ||x - y||_2^2)^{-0.5}$  with the bandwidth *c* set to the median distance between datapoints (Gorham and Mackey, 2017).



Figure S1: **Point estimation error of the gNPP.** Absolute difference between the posterior mean of the median and the true median, using the Wasserstein (a,b) and KSD (c, d) divergences, in the well-specified (a,c) and mis-specified cases (b,d).



Figure S2: **Calibration of the gNPP.** We calculate how often the credible interval of the gNPP posterior includes the true median.



Figure S3: **Dependence of the generalized Bayes factor on the rate** r. We plot the log generalized Bayes factor with the Wasserstein distance, under different choices of rate hyperparameter r (colors), in the well-specified case (solid) and misspecified case (dashed). Values above zero indicate the parametric model is favored.

# **C** Review

We review key background for our theoretical results.

### C.1 Distances and Divergences

This section collect some definitions and results about statistical divergences.

**Definition S1** (KL divergence). *The Kullback-Leibler (KL) divergence between two probability distributions* p *and* q *is defined as* 

$$D_{\mathrm{KL}}(\mathbf{p} \parallel \mathbf{q}) = \int \log\left(\frac{\mathbf{p}(t)}{\mathbf{q}(t)}\right) \mathbf{p}(t) dt.$$

**Definition S2** (Hellinger distance). *The Hellinger distance between two probability distributions* p *and* q *is defined as* 

$$d_H(\mathbf{p},\mathbf{q}) = \sqrt{\frac{1}{2} \int \left(\sqrt{\mathbf{p}(x)} - \sqrt{\mathbf{q}(x)}\right)^2} dx$$

**Definition S3** (Total variation distance). *The total variation distance between two probability distributions* p *and* q *is defined as* 

$$d_{TV}(\mathbf{p},\mathbf{q}) = \frac{1}{2} \int |\mathbf{p}(x) - \mathbf{q}(x)| dx.$$

**Definition S4** (Bounded Lipschitz distance). *For a real-valued function*  $f : \mathbb{R}^d \to \mathbb{R}$ *, the bounded Lipschitz norm is defined as* 

$$||f||_{BL} = ||f||_{\infty} + ||f||_{L}, \tag{19}$$

where  $||f||_{\infty} = \sup_{x} |f(x)|$  and  $||f||_{L} = \sup_{x \neq y} \frac{|f(x) - f(y)|}{||x - y||_{2}}$ .

For two distributions p,q on  $\mathbb{R}^d$ , the bounded Lipschitz metric is defined as

$$d_{BL}(\mathbf{p},\mathbf{q}) = \sup_{\|f\|_{BL} \le 1} \left| \int f(x)(\mathbf{p}(x) - \mathbf{q}(x)) dx \right|.$$
(20)

**Theorem S1.** (*Van der Vaart and Wellner*, 2023, *Theorem 1.12.4*) For a sequence of distributions  $p_n$  and a distribution  $p_0$  on  $\mathbb{R}^d$ , the following are equivalent:

- (a)  $p_n$  converges weakly to  $p_0$ .
- (b)  $d_{BL}(\mathbf{p}_n,\mathbf{p}_0) \rightarrow 0.$

### C.2 Bayesian Nonparametrics

In this section we provide background on the frequentist analysis of Bayesian nonparametric models.

**Notation.** Let  $\mathscr{X} \subseteq \mathbb{R}^d$ . We denote the Hellinger distance, KL divergence, and  $L_2$  norm between two densities  $p, q \in \mathscr{P}(\mathscr{X})$  as  $d_H(p,q)$ ,  $D_{KL}(p || q)$ , and  $||p - q||_2$ , respectively, as reviewed in Appendix C.1. We also consider a generic metric *d* on the space of probability densities, such as the Hellinger distance,  $L_1$  distance, or  $L_2$  distance for uniformly bounded densities.

The  $\varepsilon$ -covering numbers and  $\varepsilon$ -packing numbers of a metric space  $(\mathscr{P}, \rho)$  denoted by  $N(\varepsilon, \mathscr{P}, \rho)$ and  $D(\varepsilon, \mathscr{P}, \rho)$  are defined as the minimal number of balls of radius  $\varepsilon$  needed to cover  $\mathscr{P}$ , and the maximal number of  $\varepsilon$ -separated points, respectively.

For any  $\varepsilon > 0$ , we define the neighborhood of  $p_0$  as follows:

$$B_{0}(\mathbf{p}_{0}, \boldsymbol{\varepsilon}, \mathscr{P}) = \left\{ \mathbf{p} \in \mathscr{P} : d_{H}(\mathbf{p}_{0}, \mathbf{p}) < \boldsymbol{\varepsilon}^{2} \right\},$$
  

$$B_{k}(\mathbf{p}_{0}, \boldsymbol{\varepsilon}, \mathscr{P}) = \left\{ \mathbf{p} \in \mathscr{P} : D_{\mathrm{KL}}(\mathbf{p}_{0} \parallel \mathbf{p}) < \boldsymbol{\varepsilon}^{2}, \mathbb{E}_{\mathbf{p}_{0}} \left[ \left| \log(\mathbf{p}_{0}/\mathbf{p}) - D_{\mathrm{KL}}(\mathbf{p}_{0} \parallel \mathbf{p}) \right|^{k} \right] < \boldsymbol{\varepsilon}^{k} \right\}, \quad \text{for } k > 0$$
(21)

The  $B_0$  neighborhood defines an  $\varepsilon$ -open ball around  $p_0$  under the Hellinger distance. The  $B_k$  neighborhoods (for k > 0) consider the KL divergence and higher-order moments of the loglikelihood ratio. These definitions help us reasons about the posterior contraction rates. We keep the metric *d* purposefully vague, but a common choice is the Hellinger divergence.

We define the nonparametric prior  $\Pi_{np}(p)$  and  $\mathcal{M}_{np}$  as the support of  $\Pi_{np}$ .

When we say the parametric model is regular, we mean that the model is finite-dimensional and satisfies sufficient conditions for the Bernstein von-Mises theorem. Sufficient conditions can be found in Ch10 of Van der Vaart (2000), Ch 8 of Ghosal and Van der Vaart (2017) or Section 2 of Miller (2021).

**Prior mass and entropy assumptions.** Let  $\varepsilon_{n,pm}$ ,  $\varepsilon_{n,np}$  be two sequences of positive numbers tending to zero, such that  $\varepsilon_{n,pm} < \varepsilon_{n,np}$ .

Assumption S5 (Model Entropy). There exists constant  $C_{pm}$ ,  $C_{np}$  such that

$$\sup_{\varepsilon \ge \varepsilon_{n,pm}} \log N\left(\frac{\varepsilon}{3}, B_0\left(p_0, 2\varepsilon, \mathcal{M}_{pm}\right), d_H\right) \le C_{pm} n \varepsilon_{n,pm}^2, \quad and$$

$$\sup_{\varepsilon \ge \varepsilon_{n,np}} \log N\left(\frac{\varepsilon}{3}, B_0\left(p_0, 2\varepsilon, \mathcal{M}_{np}\right), d_H\right) \le C_{np} n \varepsilon_{n,np}^2.$$
(22)

Assumption S6 (Prior Mass). For the KL neighborhood in Eq. (21), the priors satisfy

$$\Pi_{pm} \left( B_2 \left( p_0, \varepsilon_{n,pm}, \mathscr{M}_{pm} \right) \right) \ge \exp \left( -n \varepsilon_{n,pm}^2 \right), \quad if \ p_0 \in \mathscr{M}_{pm},$$

$$\Pi_{np} \left( B_2 \left( p_0, \varepsilon_{n,np}, \mathscr{M}_{np} \right) \right) \ge \exp \left( -n \varepsilon_{n,np}^2 \right), \quad if \ p_0 \notin \mathscr{M}_{pm}.$$
(23)

Assumptions S5 and S6 establish the standard prior mass and entropy conditions necessary for posterior contraction rates (Ghosal et al., 2000). Assumption S5 defines the parametric and nonparametric rates. Assumption S6 requires the parametric prior to put sufficient mass around the truth when the parametric model is correctly specified. Similarly, the nonparametric prior is required to put sufficient mass around the truth when the parametric model is incorrectly specified.

It is useful to think of  $\varepsilon_{n,pm}$ ,  $\varepsilon_{n,np}$  as the posterior contraction rates of the parametric and nonparametric models, respectively. When the parametric model is regular, a choice of  $\varepsilon_{n,pm}$  is  $n^{-1/2}d^{1/2}\log n$  by the Bernstein von-Mises theorem (Van der Vaart, 2000). The nonparametric contraction rate is slower by at least a logarithmic factor. For example, the posterior of a Dirichlet process mixture of normal priors contracts a rate of  $n^{-1/2}(\log n)^{(d+1+1/r_0)/2}$  where *d* is the dimension and  $r_0$  is some measure of smoothness of the true mixing distribution (Shen et al., 2013).

Assumption S7 (Rate Difference). We assume that for every M > 0,

$$\Pi_{\rm np}\left(B_0(p_0, M\varepsilon_{n,{\rm np}}, \mathcal{M}_{\rm np})\right) = o\left(\exp\left(-3n\varepsilon_{n,{\rm pm}}^2\right)\right). \tag{24}$$

Assumption S7 establishes an upper bound for the nonparametric prior mass within a ball of radius  $M\varepsilon_{n,np}$ , which matches the prior mass lower bound in Assumption S6. As we approach the truth at a rate of  $M\varepsilon_{n,np}$ , the prior mass decreases at a rate exceeding  $\exp\left(-3n\varepsilon_{n,pm}^2\right)$  within this neighborhood. When the parametric model contracts at a rate of  $\varepsilon_{n,pm}^2 = \frac{d \log n}{n}$  under the regularity conditions, this requirement implies  $\Pi_{np} \left(B_0(\mathbb{P}_0, M\varepsilon_{n,np}, \mathcal{M}_{np})\right) = o\left(n^{-3d}\right)$ . This is typically met in most nonparametric models, where the decay rate of the left side of Eq. (24) is on the order of  $\exp\left(-M'n\varepsilon_{n,pp}^2\right)$ .

# **D Proof of Section 4**

### **D.1 Proofs of Section 4.1**

We prove that for Dirichlet process perturbation, model selection consistency does not hold (Example 1).

**Proposition S2.** *Example 1 fails to satisfy model selection consistency as in Proposition 1.* 

*Proof of Proposition* S2. Consider the Polya urn construction of the Dirichlet process,

$$x_1 \sim p_{\theta}, \quad x_k \sim \frac{1}{1+h} p_{\theta} + \frac{h}{1+h} \hat{p}_{x_{1:k-1}}, \quad \text{for } k > 1.$$
 (25)

where  $p_{x_{1:k-1}}$  is the empirical distribution of  $x_{1:k-1}$ , and  $h = 1/\alpha$  (Blackwell and MacQueen, 1973). For distinct  $x_1, \dots, x_n$ , we have

$$\eta_{n} = \frac{\eta \int p_{\theta}(x_{1:n}) d\Pi_{pm}(\theta)}{(1-\eta) \int \tilde{p}(x_{1:n}) d\Pi_{pert}(\tilde{p})} = \frac{\eta}{1-\eta} \frac{\int p_{\theta}(x_{1:n}) d\Pi_{pm}(\theta)}{\left(\frac{1}{1+h}\right)^{n-1} \int p_{\theta}(x_{1:n}) d\Pi_{pm}(\theta)} = \frac{\eta}{1-\eta} \left(\frac{1}{1+h}\right)^{1-n}.$$
(26)

The resulting  $\eta_n$  does not depend on whether the parametric model is correct, thus it does not satisfy model selection consistency.

We next establish that for Dirichlet process mixture model perturbation, model selection consistency holds if  $p_0$  is sufficiently smooth (Example 2). Loosely, a distribution is said to be *supersmooth* when it is a Gaussian mixture over a thin-tailed mixing distribution, and  $\beta$ -smooth when the density is thin-tailed and has sufficiently regular derivatives.

**Definition S5.** A density  $p_0$  on  $\mathbb{R}^d$  is said to be

- supersmooth if there exists  $(F_0, \Sigma_0)$  such that  $p_0 = F_0 \otimes \mathcal{N}(0, \Sigma_0)$  and  $1 F_0([-z, z]^d) \lesssim \exp(-c_0 z^{r_0})$  for every z > 0, with  $c_0 > 0$  and  $r_0 \ge 2$ .
- *β*-smooth if the following holds:

1. The mixed partial derivative  $D^k p_0$  of order up to  $k_+ \leq \lfloor \beta \rfloor$  satisfying

$$\left| (D^{k} \mathbf{p}_{0})(x+y) - D^{k} \mathbf{p}_{0}(x) \right| \leq L(x) \exp\left(c_{1} \|y\|^{2}\right) \|y\|^{\beta - \lfloor \beta \rfloor}, k_{+} = \lfloor \beta \rfloor, x, y \in \mathbb{R}^{d},$$

$$P_{0} \left(\frac{L + |D^{k} \mathbf{p}_{0}|}{\mathbf{p}_{0}}\right)^{(2\beta + \varepsilon)/\beta} < \infty,$$
(27)

for some function  $L : \mathbb{R}^d \mapsto [0, \infty)$  and  $c_1, \varepsilon > 0$ .

2. For every ||x|| > a,  $p_0 \le c \exp(-b||x||^r)$ , for some  $a, b, c, \tau > 0$ .

**Proposition S3.** Assume that the true density  $p_0$  is supersmooth or  $\beta$ -smooth. Assume that with positive  $[\Pi_{pm}]$ -probability, the following holds

- 1.  $p_{\theta}$  has continuous and positive Lebesgue density over  $\mathbb{R}^{d}$ .
- 2. there exists positive constants  $z, b_1, C_1$  such that  $P_{\theta}\left([-z, z]^d\right) \ge 1 b_1 \exp\left(-C_1 z^{r_0}\right)$  for some  $r_0 \ge 2$ .

*Then Assumptions S5 and S6 hold for the Dirichlet process normal mixture perturbation (Eqs.* (2) *and* (3)) *and* 

1. 
$$\varepsilon_{n,np} = n^{-1/2} (\log n)^{(d-1+1/r_0)/2}$$
 if  $p_0 = F_0 * \mathcal{N}(0, \sigma_0^2 I_d)$  is supersmooth with  $\sigma_0^2 > 0$ ,

2. 
$$\varepsilon_{n,np} = n^{-\beta/(2\beta+d^*)} (\log n)^{t_0}$$
 if  $p_0$  is  $\beta$ -smooth, where  $d^* = d \wedge 2$  and  $t_0(\beta d^* + \beta d^*/\tau + d^* + \beta)/(2\beta + d^*)$ .

When the true density  $p_0$  is  $\beta$ -Hölder smooth, Shen et al. (2013) shows that Dirichlet process normal mixtures converge to  $p_0$  at a rate of  $\varepsilon_{n,np} = n^{-\beta/(2\beta+d^*)} (\log n)^{t_0}$  for some constant  $t_0 >$ 0. Under the same rate, Proposition S3 explicitly verifies Assumptions S5 and S6. To satisfy Proposition 1, Assumption S7 must also hold; for this it suffices for  $\varepsilon_{n,np}$  to lower-bound other posterior contraction rates (Ghosal and Van der Vaart, 2017, Theorem 8.35). This is reasonable to expect, as  $\varepsilon_{n,np}$  matches the minimax rate for estimating  $\beta$ -Hölder smooth densities up to a polylogarithmic factor (Tsybakov, 2008). *Proof of Proposition* S3. Let  $\Pi_{pert}$  be the prior for p. First, we verify Assumption S6.

$$\Pi_{\text{pert}}\left(B_{2}\left(\mathbf{p}_{0},\boldsymbol{\varepsilon}_{n,\text{np}},\mathcal{M}_{\text{np}}\right)\mid\boldsymbol{\theta},h\right)\geq\exp\left(-n\boldsymbol{\varepsilon}_{n,\text{np}}^{2}\right).$$
(28)

Let *E* be the event that

$$\left\{\boldsymbol{\theta}: 1 - \mathbf{P}_{\boldsymbol{\theta}}\left(\left[-z, z\right]^{d}\right) \le b_{1} \exp\left(-C_{1} z^{r_{0}}\right), \mathbf{p}_{\boldsymbol{\theta}}(x) > 0 \text{ and is continuous } \forall x \in \mathbb{R}^{d}\right\}.$$
 (29)

Under the assumption, we have  $\Pi_{pm}(E) > 0$ .

Conditioning on the event E,  $\Pi(p | \theta)$  is a Dirichlet process mixture that satisfies the sufficient conditions for Proportions 9.14 of Ghosal and Van der Vaart (2017). Then by the same proposition, Assumption S6 holds with choices of  $\varepsilon_{n,np}$  given in the main statement. In particular, there exist positive constants A, C such that

$$\Pi_{\text{pert}}\left(B_{2}\left(p_{0},A\varepsilon_{n,\text{np}},\mathscr{M}_{\text{np}}\right)\mid\theta\right)\geq\exp\left(-Cn\varepsilon_{n,\text{np}}^{2}\right).$$
(30)

Since  $\Pi_{pm}(E) > 0$ , after marginalizing out  $\theta$ , we have

$$\Pi_{\text{pert}}\left(B_2\left(\mathbf{p}_0, A\varepsilon_{n,\text{np}}, \mathcal{M}_{\text{np}}\right)\right) \ge \exp\left(-C' n\varepsilon_{n,\text{np}}^2\right),\tag{31}$$

for some other constant C' > 0.

Moreover, condition on E, Assumption S5(a) holds using the argument in Example 9.4 of Ghosal and Van der Vaart (2017). Recall that Assumption S5 states that

$$\sup_{\varepsilon \ge \varepsilon_{n,\mathrm{np}}} \log N\left(\frac{\varepsilon}{3}, B_0\left(\mathrm{p}_0, 2\varepsilon, \mathcal{M}_{\mathrm{np}}\right), d_H\right) \le C_{\mathrm{np}} n \varepsilon_{n,\mathrm{np}}^2.$$
(32)

Since Eq. (32) does not depend on  $\theta$ , it holds true if we remove the conditioning on E.

*Proof of Theorem 1*. First note the bounded Lipschitz metric is convex in the first argument. For  $\eta \in [0,1]$  and probability measures  $\mathbb{P}_0, \mathbb{P}_1, \mathbb{Q}$ ,

$$d_{BL}(\eta \mathbb{P}_{0} + (1 - \eta) \mathbb{P}_{1}, \mathbb{Q}) = \sup_{\|f\|_{BL} \leq 1} \left| \int f(d(\eta \mathbb{P}_{0} + (1 - \eta) \mathbb{P}_{1}) - d\mathbb{Q}) \right|$$

$$\leq \eta \sup_{\|f\|_{BL} \leq 1} |f(d\mathbb{P}_{0} - d\mathbb{Q})| + (1 - \eta) \sup_{\|f\|_{BL} \leq 1} |f(d\mathbb{P}_{1} - d\mathbb{Q})|$$

$$= \eta d_{BL}(\mathbb{P}_{0}, \mathbb{Q}) + (1 - \eta) d_{BL}(\mathbb{P}_{1}, \mathbb{Q}).$$
(33)

Assumption 1(b) implies the total-variation convergence

$$d_{TV}\left(\Pi_{\text{pm}}\left(d\theta \mid x_{1:n}\right), \mathcal{N}\left(\hat{\theta}_{\text{MLE}}, \left(nV_{\theta_{0}}\right)^{-1}\right)\right) = O_{\mathbb{P}_{0}}(n^{-1/2}).$$
(34)

If we treat  $\theta \mapsto \sqrt{n}\chi(\theta)$  as a pushforward map, then by the triangle inequality,

$$d_{BL}\left(\Pi_{pm}\left(\tilde{\psi}_{n}(\mathbf{p}_{\theta}) \mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\chi}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\chi}_{\theta_{0}}\right)\right)$$

$$\leq \underbrace{d_{BL}\left(\Pi_{pm}\left(\tilde{\psi}_{n}(\mathbf{p}_{\theta}) \mid x_{1:n}\right), \left(\sqrt{n}\chi\right) \# \mathcal{N}\left(\hat{\theta}_{MLE}, \left(nV_{\theta_{0}}\right)^{-1}\right)\right)}_{A_{n}}$$

$$+ \underbrace{d_{BL}\left(\left(\sqrt{n}\chi\right) \# \mathcal{N}\left(\hat{\theta}_{MLE}, \left(nV_{\theta_{0}}\right)^{-1}\right), \mathcal{N}\left(0, \dot{\chi}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\chi}_{\theta_{0}}\right)\right)}_{B_{n}}.$$
(35)

Let  $\tilde{\Pi}_n$  be the law of  $\sqrt{n}(\theta - \hat{\theta}_{MLE})$  where  $\theta \sim \Pi_{pm}(\cdot | x_{1:n})$ . We upper bound  $A_n$  with the total variation bound in Assumption 1(b).

$$A_{n} \leq d_{TV} \left( \left( \sqrt{n} \chi \right)_{\#} \Pi_{pm} \left( \cdot \mid x_{1:n} \right), \left( \sqrt{n} \chi \right)_{\#} \mathscr{N} \left( \hat{\theta}_{MLE}, \left( nV_{\theta_{0}} \right)^{-1} \right) \right)$$

$$= 2 \sup_{A} \left| \Pi_{pm} \left( n^{-1/2} \chi^{-1}(A) \mid x_{1:n} \right) - \mathscr{N} \left( \chi^{-1}(A); \hat{\theta}_{MLE}, V_{\theta_{0}}^{-1} \right) \right|$$

$$\leq 2 \sup_{B} \left| \tilde{\Pi}_{n}(B) - \mathscr{N} \left( B; 0, V_{\theta_{0}}^{-1} \right) \right| = d_{TV} \left( \tilde{\Pi}_{n}, \mathscr{N} \left( 0, V_{\theta_{0}}^{-1} \right) \right) = o_{\mathbb{P}_{0}}(1).$$
(36)

The third line follows from the definition of TV distance.

Now we bound  $B_n$ . For  $\theta \sim \mathcal{N}\left(\hat{\theta}_{MLE}, (nV_{\theta_0})^{-1}\right)$ , we can reparametrize  $\theta = \hat{\theta}_{MLE} + (nV_{\theta_0})^{-1/2}Z$ for  $Z \sim \mathcal{N}(0, I_d)$ . By a Taylor expansion, there exists  $\tilde{\theta}_n$  in an  $O_{\mathbb{P}_0}(n^{-1/2})$ -neighborhood of  $\hat{\theta}_{MLE}$ such that

$$\boldsymbol{\chi}(\boldsymbol{\theta}) = \boldsymbol{\chi}(\hat{\boldsymbol{\theta}}_{\text{MLE}}) + \dot{\boldsymbol{\chi}}_{\hat{\boldsymbol{\theta}}_{\text{MLE}}}^{T} \left( n V_{\boldsymbol{\theta}_{0}} \right)^{-1/2} Z + n^{-1} \| V_{\boldsymbol{\theta}_{0}} \|_{2} Z^{T} \dot{\boldsymbol{\chi}}_{\tilde{\boldsymbol{\theta}}_{n}} Z.$$
(37)

We note that  $\chi(\hat{\theta}_{MLE}) = 0$ . Since  $\hat{\theta}_{MLE} \xrightarrow{\mathbb{P}_0} \theta_0$ ,  $\tilde{\theta}_n \xrightarrow{\mathbb{P}_0} \theta_0$ . Thus,

$$\sqrt{n}\boldsymbol{\chi}(\boldsymbol{\theta}) = \dot{\boldsymbol{\chi}}_{\hat{\boldsymbol{\theta}}_{\mathrm{MLE}}}^{T} V_{\boldsymbol{\theta}_{0}}^{-1/2} Z + O_{\mathbb{P}_{0}}(n^{-1/2}) \xrightarrow{\mathrm{w}} \mathcal{N}\left(0, \dot{\boldsymbol{\chi}}_{\boldsymbol{\theta}_{0}}^{\top} V_{\boldsymbol{\theta}_{0}}^{-1} \dot{\boldsymbol{\chi}}_{\boldsymbol{\theta}_{0}}\right).$$
(38)

Finally, we use the equivalence between  $d_{BL}$  and weak convergence to obtain the bound  $B_n = o_{\mathbb{P}_0}(1)$ .

For p drawn from the NPP model, we have

$$d_{BL}\left(\Pi\left(\tilde{\psi}_{n}(\mathbf{p})\mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\chi}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\chi}_{\theta_{0}}\right)\right) \leq \eta_{n} d_{BL}\left(\Pi_{pm}\left(\tilde{\psi}_{n}(\mathbf{p}_{\theta})\mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\chi}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\chi}_{\theta_{0}}\right)\right) + (1 - \eta_{n})$$

$$(39)$$
When  $\mathbf{p}_{0} \in \mathcal{M}_{pm}, \eta_{n} \to 1$  by Proposition 1, and  $d_{BL}\left(\Pi\left(\tilde{\psi}_{n}(\mathbf{p})\mid x_{1:n}\right), \mathcal{N}\left(0, V_{\theta_{0}}^{-1}\right)\right) \xrightarrow{\mathbb{P}_{0}} 0.$ 

When  $p_0 \notin \mathscr{M}_{pm}$ ,  $\eta_n \to 0$  by Proposition 1. Then

$$d_{BL}\left(\Pi\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right)$$

$$\leq \eta_{n} d_{BL}\left(\Pi_{\mathrm{pm}}\left(\psi(\mathbf{p}_{\theta}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) + (1 - \eta_{n}) d_{BL}\left(\Pi_{\mathrm{pert}}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) \qquad (40)$$

$$\leq \eta_{n} + (1 - \eta_{n}) d_{BL}\left(\Pi_{\mathrm{pert}}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) \stackrel{\mathbb{P}_{0}}{\to} 0.$$

### D.2 Proofs of Section 4.2

The first result shows the expected divergence  $\mathbb{E}_{pm} \left[ \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0}) \mid x_{1:n} \right]$  vanishes at a rate of  $n^{-1}$ .

**Lemma S1.** Let Assumptions 1 and 3 be satisfied. Then  $\mathbb{E}_{pm}\left[\rho(\mathbf{p}_{\theta},\mathbf{p}_{\theta_0}) \mid x_{1:n}\right] = O_{\mathbb{P}_0}(n^{-1}).$ 

*Proof.* Let  $B_{M_n n^{-1/2}}(\theta_0)$  be the neighborhood defined in Assumption 3(b). Let  $E_n$  be the event that  $\hat{\theta}_{MLE} \in B_{M_n n^{-1/2}}(\theta_0)$ . By Assumption 1(a),  $\mathbb{P}_0(E_n) \to 1$  as  $n \to \infty$ . For the rest of the proof, we establish the result conditioning on  $E_n$ .

Let  $S_{\theta} := \nabla_{\theta} \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0}), H_{\theta} := \nabla_{\theta}^2 \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0})$ . By Taylor expansion, there exists a vector  $\tilde{\theta} \in B\left(\theta_0, \|\hat{\theta}_{MLE} - \theta_0\|_2\right)$  such that

$$\rho(\mathbf{p}_{\hat{\theta}_{MLE}}, \mathbf{p}_{\theta_0}) \le S_{\theta_0} \left( \hat{\theta}_{MLE} - \theta_0 \right) + \frac{1}{2} \left\| H_{\tilde{\theta}}^{1/2} \left( \hat{\theta}_{MLE} - \theta_0 \right) \right\|_2^2.$$
(41)

We note that  $\nabla_{\theta} \rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0})_{\theta=\theta_0} = 0$  since  $\theta_0$  minimizes  $\rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_0})$  on the open set  $\Theta$ . By the bounded Hessian assumption (Assumption 3(b)), there exists a constant  $C_{\theta_0}$  s.t.

$$\rho(\mathbf{p}_{\hat{\boldsymbol{\theta}}_{\mathrm{MLE}}}, \mathbf{p}_{\boldsymbol{\theta}_0}) \le C_{\boldsymbol{\theta}_0} \left\| \hat{\boldsymbol{\theta}}_{\mathrm{MLE}} - \boldsymbol{\theta}_0 \right\|_2^2 = O_{\mathbb{P}_0}(n^{-1}).$$
(42)

Similarly, there exists  $\tilde{\theta} \in B\left(\theta_0, \|\hat{\theta}_{\text{MLE}} - \theta_0\|_2\right)$  such that

$$\left\|\nabla_{\theta}\rho(\mathbf{p}_{\theta},\mathbf{p}_{\theta_{0}})_{\theta=\hat{\theta}_{\mathrm{MLE}}}\right\|_{2} = \left\|H_{\tilde{\theta}'}(\hat{\theta}_{\mathrm{MLE}}-\theta_{0})\right\|_{2} \le C_{\theta_{0}}\|\hat{\theta}_{\mathrm{MLE}}-\theta_{0}\|_{2}.$$
(43)

By Taylor expansion, we have

$$\mathbb{E}_{pm}\left[\rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_{0}}) \mid x_{1:n}\right] = \rho(\mathbf{p}_{\hat{\theta}_{MLE}}, \mathbf{p}_{\theta_{0}}) + \nabla_{\theta}\rho(\mathbf{p}_{\theta}, \mathbf{p}_{\theta_{0}})_{\theta=\hat{\theta}_{MLE}} \mathbb{E}_{pm}\left[\theta - \hat{\theta}_{MLE} \mid x_{1:n}\right] \\ + \mathbb{E}_{pm}\left[\frac{1}{2} \left\| H_{\hat{\theta}_{MLE}}^{1/2} \left(\hat{\theta}_{MLE} - \theta_{0}\right) \right\|_{2}^{2} + o_{\mathbb{P}_{0}} \left( \left\| \theta - \hat{\theta}_{MLE} \right\|_{2}^{2} \right) \mid x_{1:n} \right].$$

$$(44)$$

Given the BvM theorem for  $\Pi_{pm}(\cdot | x_{1:n})$  (Assumption 1), we have

$$\mathbb{E}_{\mathrm{pm}}\left[\left\|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\right\|_{2}^{2} | \boldsymbol{x}_{1:n}\right] = O_{\mathbb{P}_{0}}\left(\frac{d}{n}\right), \quad \text{and} \quad \left\|\mathbb{E}_{\mathrm{pm}}\left[\boldsymbol{\theta} | \boldsymbol{x}_{1:n}\right] - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\right\|_{2} = O_{\mathbb{P}_{0}}\left(\sqrt{\frac{d}{n}}\right). \tag{45}$$

Finally, we combine the above inequalities to conclude that

$$\mathbb{E}_{pm}\left[\rho(\mathbf{p}_{\theta},\mathbf{p}_{\theta_{0}}) \mid x_{1:n}\right] = \rho(\mathbf{p}_{\hat{\theta}_{MLE}},\mathbf{p}_{\theta_{0}}) + \frac{1}{2}\mathbb{E}_{pm}\left[\left\|H_{\hat{\theta}_{MLE}}^{1/2}\left(\hat{\theta}_{MLE}-\theta_{0}\right)\right\|_{2}^{2} \mid x_{1:n}\right] + O_{\mathbb{P}_{0}}\left(\frac{d}{n}\right)$$

$$\leq \rho(\mathbf{p}_{\hat{\theta}_{MLE}},\mathbf{p}_{\theta_{0}}) + \frac{C_{\theta_{0}}}{2}\mathbb{E}_{pm}\left[\left\|\theta-\hat{\theta}_{MLE}\right\|_{2}^{2} \mid x_{1:n}\right] + O_{\mathbb{P}_{0}}\left(\frac{d}{n}\right) = O_{\mathbb{P}_{0}}\left(\frac{d}{n}\right). \tag{46}$$

*Proof of Theorem 2.* By the triangle inequality, we have

$$\mathbb{E}_{pm} [D_{m,n}(p_{\theta}, p_{0}) | x_{1:n}] - D(p_{0}, p_{\theta_{0}})$$

$$\leq \mathbb{E}_{pm} [D_{m,n}(p_{\theta}, p_{0}) | x_{1:n}] - \mathbb{E}_{pm} [D(p_{\theta}, p_{0}) | x_{1:n}]$$

$$+ \mathbb{E}_{pm} [D(p_{\theta}, p_{0}) | x_{1:n}] - D(p_{\theta_{0}}, p_{0}).$$
(47)

If  $d(\cdot, \cdot)$  is a semimetric, then we can directly bound the second term with Lemma S1.

$$\left|\mathbb{E}_{pm}\left[D\left(p_{\theta}, p_{0}\right) \mid x_{1:n}\right] - D\left(p_{\theta_{0}}, p_{0}\right)\right| \leq \mathbb{E}_{pm}\left[D\left(p_{\theta}, p_{\theta_{0}}\right) \mid x_{1:n}\right] = O_{\mathbb{P}_{0}}\left(n^{-1}\right).$$
(48)

If  $d(\cdot, \cdot)^{1/k}$  is a semimetric, then we apply the binomial expansion  $x^k - y^k = (x - y) \sum_{j=0}^{k-1} {k \choose j} (x - y)^{k-1-j} y^j$  to obtain

$$\begin{aligned} \left| \mathbb{E}_{pm} \left[ D(p_{\theta}, p_{0}) \mid x_{1:n} \right] - D(p_{\theta_{0}}, p_{0}) \right| \\ &= \left| \mathbb{E}_{pm} \left[ \left( \rho(p_{\theta}, p_{0}) - \rho(p_{\theta_{0}}, p_{0}) \right) \sum_{j=0}^{k-1} {k \choose j} \left( \rho(p_{\theta}, p_{0}) - \rho(p_{\theta_{0}}, p_{0}) \right)^{k-1-j} \rho(p_{\theta_{0}}, p_{0})^{j} \mid x_{1:n} \right] \right| \\ &\leq C \mathbb{E}_{pm} \left[ \left| \rho(p_{\theta}, p_{0}) - \rho(p_{\theta_{0}}, p_{0}) \right| \mid x_{1:n} \right] \\ &\leq C \mathbb{E}_{pm} \left[ \rho(p_{\theta}, p_{\theta_{0}}) \mid x_{1:n} \right] = O_{\mathbb{P}_{0}} \left( n^{-1} \right). \end{aligned}$$

$$(49)$$

where the inequality uses the uniform boundedness of  $ho(\cdot,\cdot).$ 

Let  $M_n$  be a sequence such that  $M_n \to \infty$  and  $M_n = o(n^{1/2})$ . To control the fluctuation of

 $\mathbb{E}_{pm}[D_{m,n}(p_{\theta},p_0) | x_{1:n}]$  around  $d(p_{\theta},p_0)$ , we decompose it into two terms in and outside of a ball of radius  $M_n n^{-1/2}$ :

$$\mathbb{E}_{pm} \left[ \mathbb{D}_{m,n} \left( \mathbf{p}_{\theta}, \mathbf{p}_{0} \right) - d(\mathbf{p}_{\theta}, \mathbf{p}_{0}) \mid x_{1:n} \right] = \underbrace{\mathbb{E}_{pm} \left[ \left( \mathbb{D}_{m,n} \left( \mathbf{p}_{\theta}, \mathbf{p}_{0} \right) - d(\mathbf{p}_{\theta}, \mathbf{p}_{0}) \right) I_{B_{M_{n}n^{-1/2}}^{c}(\theta_{0})} \mid x_{1:n} \right]}_{A_{m,n}}$$

$$+ \underbrace{\mathbb{E}_{pm} \left[ \left( \mathbb{D}_{m,n} \left( \mathbf{p}_{\theta}, \mathbf{p}_{0} \right) - d(\mathbf{p}_{\theta}, \mathbf{p}_{0}) \right) I_{B_{M_{n}n^{-1/2}}(\theta_{0})} \mid x_{1:n} \right]}_{B_{m,n}}.$$
(50)

Since we can write  $D_{m,n}(p_{\theta}, p_0) = D(p_{\theta}^m, p_0^n), |D_{m,n}(p_{\theta}, p_0) - d(p_{\theta}, p_0)|$  is uniformly bounded. Then applying Assumption 1 yields

$$\begin{aligned} A_{m,n} &\leq C \Pi_{pm} \left( \| \theta - \theta_0 \|_2 > M_n n^{-1/2} | x_{1:n} \right) \\ &\leq C \Pi_{pm} \left( \sqrt{n} \| \theta - \hat{\theta}_{MLE} \|_2 > M_n - \sqrt{n} \| \theta_0 - \hat{\theta}_{MLE} \|_2 | x_{1:n} \right) \\ &\leq C \inf_{\lambda > 0} \left\{ \exp \left( -\lambda (M_n - \sqrt{n} \| \theta_0 - \hat{\theta}_{MLE} \|_2) \right) \mathbb{E}_{pm} \left[ \exp \left( \lambda \sqrt{n} \| \theta - \hat{\theta}_{MLE} \|_2 \right) | x_{1:n} \right] \right\} \\ &\leq C \inf_{\lambda > 0} \left\{ \exp \left( -\lambda M_n \right) \exp \left( \lambda \sqrt{n} \| \theta_0 - \hat{\theta}_{MLE} \|_2 \right) \mathbb{E}_{pm} \left[ \exp \left( \lambda \sqrt{n} \| \theta - \hat{\theta}_{MLE} \|_2 \right) | x_{1:n} \right] \right\} \\ &= O_{\mathbb{P}_0} \left( \exp \left( -C' M_n \right) \right) O_{\mathbb{P}_0} (1) O_{\mathbb{P}_0} (1) = O_{\mathbb{P}_0} \left( \exp \left( -C' M_n \right) \right), \end{aligned}$$
(51)

for some constants *C* and *C'*. The third line applies the Chernoff bound. The last line uses the fact that weak convergence implies the convergence of characteristic functions (to a sub-Gaussian limit). Moreover, asymptotic normality of MLE implies  $\exp(\lambda \sqrt{n} || \theta_0 - \hat{\theta}_{MLE} ||_2) = O_{\mathbb{P}_0} \left( \exp(-C'M_n) \right)$ . If we take  $M_n = n^{1/3}$ , the bound above implies  $A_{m,n} = O_{\mathbb{P}_0} \left( \exp\left(-C'n^{1/3}\right) \right) = O_{\mathbb{P}_0} \left( n^{-1} \right)$ 

To bound  $B_{m,n}$ , we take the standard approach by analyzing its mean and variance with respect

to the approximating samples. For the mean, we apply Fubini's theorem

$$\mathbb{E}[B_{m,n}] = \mathbb{E}_{pm} \left[ \mathbb{E}\left[ D_{m,n}(\mathbf{p}_{\theta}, \mathbf{p}_{0}) - d(\mathbf{p}_{\theta}, \mathbf{p}_{0}) \right] I_{B_{n^{-\alpha/2}}(\theta_{0})} \mid x_{1:n} \right]$$

$$\leq \sup_{\theta \in B_{n^{-\alpha/2}}(\theta_{0})} \mathbb{E}\left[ D_{m,n}(\mathbf{p}_{\theta}, \mathbf{p}_{0}) - d(\mathbf{p}_{\theta}, \mathbf{p}_{0}) \right].$$
(52)

Take  $\theta_n \in \operatorname{argmax}_{\theta \in \bar{B}_{n-\alpha/2}(\theta_0)} \mathbb{E} [D_{m,n} (p_{\theta}, p_0) - d(p_{\theta}, p_0)]$ . As  $n \to \infty$ , the sequence  $\theta_n \to \theta_0$  by the fact that  $\|\theta_n - \theta_0\|_2 \le n^{-\alpha/2}$ . Then we apply Assumption 4(a) to conclude that  $\mathbb{E} [D_{m,n} (p_{\theta_n}, p_0) - d(p_{\theta_n}, p_0)] = O(r_{m,n})$ .

Now we control the variance. First, we upper bound the variance using Jensen's inequality.

$$\operatorname{Var}(B_{m,n}) \leq \mathbb{E}_{\operatorname{pm}}\left[\operatorname{Var}\left(\operatorname{D}_{m,n}\left(\operatorname{p}_{\theta},\operatorname{p}_{0}\right)\right)I_{B_{n^{-\alpha/2}}(\theta_{0})} \mid x_{1:n}\right].$$
(53)

By Assumption 4, we refine the upper bound as follows

$$\operatorname{Var}(B_{m,n}) \le r_{m,n}^2 \sup_{\theta \in \bar{B}_{n^{-\alpha/2}}(\theta_0)} \mathscr{V}(\theta).$$
(54)

As  $n \to \infty$ ,  $\sup_{\theta \in \overline{B}_{n^{-\alpha/2}}(\theta_0)} \mathscr{V}(\theta) \to \mathscr{V}(\theta_0)$  by the continuity of  $\mathscr{V}$  at  $\theta_0$ , thus  $\operatorname{Var}(B_{m,n}) = O(r_{m,n}^2)$ . By applying Chebyshev's inequality, we conclude that  $B_{m,n} = O_{\mathbb{P}_0}(r_{m,n})$ .

Putting the bounds together, we have

$$\mathbb{E}_{pm}[D_{m,n}(p_{\theta}, p_0) \mid x_{1:n}] - D(p_{\theta_0}, p_0) = O_{\mathbb{P}_0}(r_{m,n} \vee n^{-1}).$$
(55)

Proof of Theorem 3. Recall that we defined the generalized Bayes factor as

$$gBF_n := \Xi \left( \frac{\mathbb{E}[D_n(\mathbf{p}_{\theta}, \mathbf{p}_0)]}{\mathbb{E}[D_n(\mathbf{p}_{\theta}, \mathbf{p}_0) \mid x_{1:n}]} (n+1)^{-r} \right) \frac{\eta}{1-\eta}.$$
(56)

The assumptions imply  $r_n(n+1)^r = o(1)$  and  $\mathbb{E}_{pm}[D_n(p_\theta, p_0)] = O_{\mathbb{P}_0}(1)$ . By the continuous mapping theorem, gBF<sub>n</sub> scales like

$$gBF_n = \Xi\left(\frac{O_{\mathbb{P}_0}(1)}{D(p_{\theta_0}, p_0)(n+1)^r + o(1)}\right)\frac{\eta}{1-\eta}.$$
(57)

If  $p_0 \in \mathscr{M}_{pm}$ , then  $D(p_{\theta_0}, p_0) = 0$  and  $gBF_n \xrightarrow{\mathbb{P}_0} \infty$  by the continuous mapping theorem. Analogously, if  $p_0 \notin \mathscr{M}_{pm}$ , then  $D(p_{\theta_0}, p_0) > 0$  and  $gBF_n \xrightarrow{\mathbb{P}_0} 0$ .

We conclude with the desired result by applying the limit of  $gBF_n$  to  $\hat{\eta}_n$ .

*Proof of Theorem 4.* By Assumptions 1, 3 and 4, we have  $\mathbb{E}_{pm}[D_n(p_\theta, p_0) | x_{1:n}] = D(p_{\theta_0}, p_0) + r_n$ .

By the decomposition provided in Eq. (4) and Eq. (6), we have

$$d_{BL}\left(\hat{\Pi}\left(\tilde{\psi}_{n}(\mathbf{p}) \mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\boldsymbol{\chi}}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\boldsymbol{\chi}}_{\theta_{0}}\right)\right) \leq \hat{\eta}_{n} \underbrace{d_{BL}\left(\Pi_{\mathrm{pm}}\left(\tilde{\psi}_{n}(\mathbf{p}_{\theta}) \mid x_{1:n}\right), \mathcal{N}\left(0, \dot{\boldsymbol{\chi}}_{\theta_{0}}^{\top} V_{\theta_{0}}^{-1} \dot{\boldsymbol{\chi}}_{\theta_{0}}\right)\right)}_{\mathscr{A}_{n}} + (1 - \hat{\eta}_{n}) \underbrace{(58)}$$

In the proof of Theorem 1, we showed that the convergence  $\mathscr{A}_n = o_{\mathbb{P}_0}(1)$  under Assumptions 1 and 2. When  $p_0 \in \mathscr{M}_{pm}$ ,  $\hat{\eta}_n \stackrel{\mathbb{P}_0}{\to} 1$  by Theorem 3, and  $d_{BL}\left(\hat{\Pi}\left(\tilde{\psi}_n(\mathbf{p}) \mid x_{1:n}\right), \mathscr{N}(0, V_{\theta_0}^{-1})\right) \stackrel{\mathbb{P}_0}{\to} 0$ . When  $p_0 \notin \mathscr{M}_{pm}$ ,  $\hat{\eta}_n \stackrel{\mathbb{P}_0}{\to} 0$  by Theorem 3. Then

$$d_{BL}\left(\hat{\Pi}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right)$$

$$\leq \hat{\eta}_{n} d_{BL}\left(\Pi_{\mathrm{pm}}\left(\psi(\mathbf{p}_{\theta}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) + (1 - \hat{\eta}_{n}) d_{BL}\left(\hat{\Pi}_{\mathrm{pert}}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) \qquad (59)$$

$$\leq \hat{\eta}_{n} + (1 - \hat{\eta}_{n}) d_{BL}\left(\hat{\Pi}_{\mathrm{pert}}\left(\psi(\mathbf{p}) \mid x_{1:n}\right), \delta_{\psi(\mathbf{p}_{0})}\right) \stackrel{\mathbb{P}_{0}}{\to} 0.$$

### **D.3** Empirical Divergences

#### D.3.1 Wasserstein

We establish the convergence rate of  $\mathbb{E}_{pm} \left[ W_p^p(\mathbf{p}_{\theta}^m, \mathbf{p}_0^n) \mid x_{1:n} \right]$  over a sample space  $\mathscr{X} \subseteq \mathbb{R}^{\kappa}$ . This implies that Theorem 4 holds for the Wasserstein divergence, by replacing Theorem 2 with Theorem S2 in the proof of Theorem 4.

**Theorem S2** (Posterior expected Wasserstein convergence rate). Suppose  $p \ge 1$ . Let Assumptions 1, 3 and S8 be satisfied. Then

$$\mathbb{E}_{pm}\left[\mathbf{w}_{p}^{p}(\mathbf{p}_{\theta}^{m},\mathbf{p}_{0}^{n}) \mid x_{1:n}\right] - \mathbf{w}_{p}^{p}\left(\mathbf{p}_{\theta_{0}},\mathbf{p}_{0}\right) = O_{\mathbb{P}_{0}}\left(n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)}\right).$$
(60)

In Assumption S8, we assume that both the parametric family and the true distribution have compact support and finite moments up to order 2p.

**Assumption S8.** The distributions  $\{p_{\theta}\}_{\theta \in \Theta}$  and  $p_0$  satisfy

- (Support) The distributions are supported on a compact subset X ⊆ ℝ<sup>κ</sup> and have positive densities in the interior of their respective supports.
- (Moments) The distributions have finite moments up to order 2p. Additionally, the mapping  $\theta \mapsto \mathbb{E}_{p_{\theta}} \left[ \|X\|_2^{2p} \right]$  is continuous at  $\theta_0$ .

#### Proofs

**Lemma S2.** Let  $X_1, \dots, X_n \stackrel{iid}{\sim} p, Y_1, \dots, Y_m \stackrel{iid}{\sim} q$  be two independent samples with finite moments up to order 2p, specifically  $\mathbb{E}_p\left[\|X\|_2^j\right] + \mathbb{E}_q\left[\|Y\|_2^j\right] =: M_j^j < \infty$  for  $j \in [2p]$ . For any n, m > 0,

$$Var\left(\mathbf{W}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m})\right) \leq 2^{p-1}M_{2p}^{2p}\left(\frac{1}{n}+\frac{1}{m}\right)^{2},$$
 (61)

where  $p^n, q^m$  are empirical measures formed by  $X_i$ 's and  $Y_i$ 's.

*Proof of Lemma S2.* By the definition of  $W_p$ , we have

$$\mathbf{w}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m}) = \min_{\sum_{i=1}^{n} w_{ij} = \frac{1}{m}, \sum_{j=1}^{m} w_{ij} = \frac{1}{n}} \sum_{i=1}^{n} \sum_{j=1}^{m} w_{ij} \|X_{i} - Y_{j}\|_{2}^{p} \le \frac{\sum_{i=1}^{n} \sum_{j=1}^{m} \|X_{i} - Y_{j}\|_{2}^{p}}{nm}.$$
 (62)

Define  $Z_{ij} := ||X_i - Y_j||_2^p$ . Since each  $(X_i, Y_j)$  pairs are independent, the set of variables  $\{Z_{ij}\}$  are i.i.d. Under the bounded moment condition, the variance satisfies

$$\operatorname{Var}\left(Z_{ij}\right) \le 2^p M_{2p}^{2p} < \infty.$$
(63)

Aggregating the variance shows

$$\operatorname{Var}\left(\mathbf{W}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m})\right) \leq \frac{nm2^{p}M_{2p}^{2p}}{n^{2}m^{2}} = \frac{2^{p}M_{2p}^{2p}}{nm} \leq 2^{p-1}M_{2p}^{2p}\left(\frac{1}{n} + \frac{1}{m}\right)^{2}.$$
(64)

where the last inequality uses  $(nm)^{-1/2} \le \frac{1}{2}(n^{-1} + m^{-1})$ .

**Lemma S3.** If p,q are probability densities on a convex compact set  $\mathscr{X} \subseteq \mathbb{R}^{\kappa}$  with nonempty interior, then for  $p \ge 1$ , we have

$$\sup_{\mathbf{p},\mathbf{q}\in\mathscr{P}(\mathscr{X})}\mathbb{E}\left[\left|\mathbf{w}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m})-\mathbf{w}_{p}^{p}(\mathbf{p},\mathbf{q})\right|\right]=O\left(n^{-2/(\kappa\vee4)}+m^{-2/(\kappa\vee4)}\right).$$
(65)

for  $X_1, \cdots, X_n \stackrel{iid}{\sim} p$  and  $Y_1, \cdots, Y_m \stackrel{iid}{\sim} q$ .

*Proof of Lemma S3.* Recall  $p^n := \frac{1}{n} \sum_{i=1}^n \delta_{X_i}, q^m := \frac{1}{m} \sum_{j=1}^m \delta_{Y_j}$ . We note that the function  $x \mapsto ||x||_2^p$  is convex by an application of Holder's inequality. Then, following Villani (2003, Remark 1.13), there exists Kantorovich potentials  $\phi_n, \psi_m : \mathscr{X} \to \mathbb{R}$  such that  $|\phi_n|, |\psi_m| \le 1$  and

$$W_p^p(\mathbf{p}^n, \mathbf{q}^m) = \mathbb{E}_{\mathbf{p}^n} \left[ \phi_n(X) \right] + \mathbb{E}_{\mathbf{q}^m} \left[ \psi_m(Y) \right].$$
(66)

Let  $\Phi(\mathbf{p},\mathbf{q})$  be the set of pairs  $(\phi,\psi) \in L^1(\mathbf{p}) \times L^1(\mathbf{q})$  such that  $\phi(x) + \psi(y) \le ||x-y||_2^p$  for all

 $x, y \in \mathscr{X}$ . Since p,q are compactly supported, we have  $(\phi_n, \psi_n) \in \Phi(p,q)$ . By Kantorovich duality, we have

$$W_{p}^{p}(\mathbf{p},\mathbf{q}) \geq \mathbb{E}_{\mathbf{p}}[\phi_{n}(X)] + \mathbb{E}_{\mathbf{q}}[\psi_{m}(Y)] \\ = W_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m}) + \int \phi_{n}(x)(\mathbf{p}(x) - \mathbf{p}^{n}(x))dx + \int \psi_{m}(x)(\mathbf{q}(x) - \mathbf{q}^{m}(x))dx.$$
(67)

and

$$W_{p}^{p}(p^{n},q^{m}) \leq W_{p}^{p}(p,q) + \int \phi_{n}(x)(p(x)-p^{n}(x))dx + \int \psi_{m}(x)(q(x)-q^{m}(x))dx.$$
(68)

Define  $\tilde{\phi}_n(x) = \phi_n(x) - \frac{L}{4} ||x||_2^2$  and  $\tilde{\psi}_m(y) = \psi_m(y) - \frac{L}{4} ||y||_2^2$ . By Manole and Niles-Weed (2024, Lemma 5),  $\tilde{\phi}_n, \tilde{\psi}_m$  are concave, *L*-Lipschitz, and uniformly bounded by *L*.

Let  $\mathscr{F}_{L,U}(K)$  be the set of *L*-Lipschitz convex functions over a convex set  $K \subseteq \mathbb{R}^{\kappa}$ , where  $\|f(x)\|_{\infty} \leq U$ . By Manole and Niles-Weed (2024, Lemma 5), the functions  $\tilde{\phi}_n, \tilde{\psi}_m$  are concave, *L*-Lipschitz, and uniformly bounded by *L*, thus  $-\tilde{\phi}_n/L, -\tilde{\psi}_m/L$  belong to  $\mathscr{F}_{1,1}(\mathscr{X})$ .

Since  $\mathscr{X}$  is convex and compact, we define

$$\Delta_{n,m} := \sup_{f \in \mathscr{F}_{1,1}(\mathscr{X})} \int f(x)(\mathbf{p}(x) - \mathbf{p}^n(x))dx + \sup_{g \in \mathscr{F}_{1,1}(\mathscr{X})} \int g(x)(\mathbf{q}(x) - \mathbf{q}^m(x))dx \tag{69}$$

Using the fact that  $-\tilde{\phi}_n/L, -\tilde{\psi}_m/L \in \mathscr{F}_{1,1}(\mathscr{X})$ , we have

$$W_{p}^{p}(p^{n},q^{m}) - W_{p}^{p}(p,q) \le L\Delta_{n,m} + \frac{L}{4} \int ||x||_{2}^{2}((p_{n}(x) - p(x)) + (q_{m}(x) - q(x)))dx.$$
(70)

For the lower bound, there exists a pair of Kantorovich potentials  $(\phi_p, \psi_q) \in \Phi(p,q)$  such that  $\|\phi_p\|_{\infty} \vee \|\psi_q\|_{\infty} \leq 1$  and  $W_p^p(p,q) = \mathbb{E}_p[\phi_p(X)] + \mathbb{E}_q[\psi_q(Y)]$ . Thus,

$$W_{p}^{p}(p^{n},q^{m}) - W_{p}^{p}(p,q) \ge \int \phi_{p}(x)(p(x) - p^{n}(x))dx + \int \psi_{q}(x)(q(x) - q^{m}(x))dx.$$
(71)

Combining the above displays, we have

$$\mathbb{E}\left[\left|\mathbf{w}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m})-\mathbf{w}_{p}^{p}(\mathbf{p},\mathbf{q})\right|\right]$$

$$\leq L\mathbb{E}_{\mathbf{p}\otimes\mathbf{q}}\left[\Delta_{n,m}\right]+\frac{L}{4}\int \|x\|_{2}^{2}((\mathbf{p}_{n}(x)-\mathbf{p}(x))+(\mathbf{q}_{m}(x)-\mathbf{q}(x)))dx$$

$$+\mathbb{E}_{\mathbf{p}}\left[\left|\int \phi_{\mathbf{p}}(x)(\mathbf{p}(x)-\mathbf{p}^{n}(x))dx\right|\right]+\mathbb{E}\left[\left|\int \psi_{\mathbf{q}}(x)(\mathbf{q}(x)-\mathbf{q}^{m}(x))dx\right|\right].$$
(72)

Since  $\|\phi_p\|_{\infty} \vee \|\psi_q\|_{\infty} \leq 1$ , the functional  $p \mapsto \mathbb{E}_p[\phi_p(X)]$  is a bounded linear functional. By Chebyshev's inequality,  $\mathbb{P}(|\mathbb{E}_{p^n}[\phi_p(X)] - \mathbb{E}_p[\phi_p(X)]| \geq t) \leq \frac{\operatorname{Var}_p(\phi_p(X))}{nt^2} < \frac{4}{nt^2}$ . The variance bound is uniform around all distributions p over  $\mathscr{X}$ , thus we have

$$\sup_{\mathbf{p}} \mathbb{E}_{\mathbf{p}} \left[ \left| \int \phi_{\mathbf{p}}(x) (\mathbf{p}(x) - \mathbf{p}^{n}(x)) dx \right| \right] = O\left(n^{-1/2}\right).$$
(73)

Similarly, we have

$$\sup_{\mathbf{q}} \mathbb{E}_{\mathbf{q}}\left[\left|\int \psi_{\mathbf{q}}(x)(\mathbf{q}(x) - \mathbf{q}^{m}(x))dx\right|\right] = O\left(m^{-1/2}\right),\tag{74}$$

and

$$\sup_{\mathbf{p},\mathbf{q}} \int \|x\|_2^2 ((\mathbf{p}_n(x) - \mathbf{p}(x)) + (\mathbf{q}_m(x) - \mathbf{q}(x))) dx = O\left(n^{-1/2} + m^{-1/2}\right).$$
(75)

As a result, we have

$$\sup_{\mathbf{p},\mathbf{q}} \mathbb{E}_{\mathbf{p}\otimes\mathbf{q}} \left[ \left| \mathbf{W}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m}) - \mathbf{W}_{p}^{p}(\mathbf{p},\mathbf{q}) \right| \right] = L \sup_{\mathbf{p},\mathbf{q}} \mathbb{E}_{\mathbf{p}\otimes\mathbf{q}} \left[ \Delta_{n,m} \right] + O\left( n^{-1/2} + m^{-1/2} \right).$$
(76)

To upper bound  $\mathbb{E}_{p\otimes q}[\Delta_{n,m}]$ , we note that it is a sum of expectation suprema of empirical processes indexed by convex Lipschitz functions. Thus, we can bound the expectation of suprema by applying Dudley's chaining technique in terms of metric entropy of the class  $\mathscr{F}_{1,1}(\mathbf{X})$ . One such results is Luxburg and Bousquet (2004, Theorem 16) which states

$$\mathbb{E}_{p}\left[\sup_{f\in\mathscr{F}(\mathscr{X})}\int f(x)(p(x)-p^{n}(x))dx\right] \leq 2\tau + \frac{4\sqrt{2}}{\sqrt{n}}\int_{\tau/4}^{\sup_{f,f'\in\mathscr{F}}\|f-f'\|_{L^{2}(p_{n})}}\sqrt{\log N(\varepsilon,\mathscr{F},L^{2}(p_{n}))}d\varepsilon.$$
(77)

Since  $L^2$ -distance is strictly small than the  $L^{\infty}$ -distance,

$$N\left(\varepsilon,\mathscr{F}_{1,1}(\mathscr{X}),L^{2}(\mathbf{p}_{n})\right) \leq N\left(\varepsilon,\mathscr{F}_{1,1}(\mathscr{X}),L^{\infty}\right) \leq N\left(\varepsilon,\mathscr{F}_{1,1}([-1,1]^{\kappa}),L^{\infty}\right),\tag{78}$$

where the last inequality uses the compactness of  $\mathscr{X}$ .

By Guntuboyina and Sen (2012, Theorem 1), there exists  $\varepsilon_0 > 0$  such that when  $\varepsilon \le \varepsilon_0(B+2)$ ,

$$\log N(\varepsilon, \mathscr{F}_{1,1}([-1,1]^{\kappa}), L^{\infty}) \le C\left(\frac{B+2}{\varepsilon}\right)^{\kappa/2},\tag{79}$$

where C is a universal constant.

Take  $B = 0 \vee (\frac{2}{\varepsilon_0} - 2)$ . Combining the above three displays, we have

$$\mathbb{E}_{p\otimes q}\left[\Delta_{n,m}\right] \leq 2\tau_{n} + \frac{4\sqrt{2}}{\sqrt{n}} \int_{\tau_{n}/4}^{2} C\left(\frac{2}{\varepsilon_{0}}\right)^{\kappa/4} \varepsilon^{-\kappa/4} d\varepsilon + 2\tau_{m} + \frac{4\sqrt{2}}{\sqrt{m}} \int_{\tau_{m}/4}^{2} C\left(\frac{2}{\varepsilon_{0}}\right)^{\kappa/4} \varepsilon^{-\kappa/4} d\varepsilon = 2\tau_{n} + \frac{C'}{\sqrt{n}} \left(\frac{2}{\varepsilon_{0}}\right)^{\kappa/4} \left|2^{1-\kappa/4} - (\tau_{n}/4)^{1-\kappa/4}\right| + 2\tau_{m} + \frac{C'}{\sqrt{m}} \left(\frac{2}{\varepsilon_{0}}\right)^{\kappa/4} \left|2^{1-\kappa/4} - (\tau_{m}/4)^{1-\kappa/4}\right| \leq C'' \varepsilon_{0}^{-\kappa/4} \left(n^{-1/2} + m^{-1/2}\right) + 2\tau_{n} + 2\tau_{m} + \varepsilon_{0}^{-\kappa/4} \left(\frac{\tau_{n}^{1-\kappa/4}}{\sqrt{n}} + \frac{\tau_{m}^{1-\kappa/4}}{\sqrt{m}}\right).$$
(80)

Choose  $\tau_n = \varepsilon_0 n^{-2/\kappa}$ . Then for constants  $C_{\varepsilon_0,\kappa}, C'_{\varepsilon_0,\kappa}$ ,

$$\mathbb{E}_{p\otimes q}\left[\Delta_{n,m}\right] \le C_{\varepsilon_0,\kappa}\left(n^{-1/2} + m^{-1/2}\right) + C'_{\varepsilon_0,\kappa}\left(n^{-2/\kappa} + m^{-2/\kappa}\right).$$
(81)

The constants only depend on  $\varepsilon_0$  and  $\kappa$ , so the bound is preserved after we take the supremum over p and q. Combining the bound on  $\sup_{p,q} \mathbb{E}_{p \otimes q}[\Delta_{n,m}]$  with Eq. (76), we get:

$$\sup_{\mathbf{p},\mathbf{q}} \mathbb{E}_{\mathbf{p}\otimes\mathbf{q}} \left[ \left| \mathbf{W}_{p}^{p}(\mathbf{p}^{n},\mathbf{q}^{m}) - \mathbf{W}_{p}^{p}(\mathbf{p},\mathbf{q}) \right| \right] = O\left( n^{-1/2} + m^{-1/2} \right) + O\left( n^{-2/\kappa} + m^{-2/\kappa} \right).$$
(82)

The final statement comes from the observation that the  $O\left(n^{-1/2} + m^{-1/2}\right)$  term dominates when  $\kappa < 4$  and the  $O\left(n^{-2/\kappa} + m^{-2/\kappa}\right)$  terms dominates when  $\kappa \ge 4$ .

**Lemma S4.** Assume that  $diam(\mathscr{X}) < \infty$ . Then  $\sup_{p,q \in \mathbb{P}(\mathscr{X})} W_p^p(p,q) < \infty$ .

*Proof of Lemma S4.* Under the definition of  $W_p$ , we have  $\sup_{p,q \in \mathbb{P}(\mathscr{X})} W_p^p(p,q) \leq \operatorname{diam}(\mathscr{X})^p < \infty$ . □

We also establish the convergence rate for the sample approximation to the  $W_p$  distance under the degenerate and non-degenerate cases.

**Lemma S5.** Assume that the distributions  $p_{\theta}$  and  $p_0$  are supported on a compact subset  $\mathscr{X} \subseteq \mathbb{R}^{\kappa}$  have positive densities in the interior of their supports. Additionally, assume that the mapping  $\theta \mapsto \mathbb{E}_{p_{\theta}} \left[ \|X\|_{2}^{2p} \right]$  is continuous at  $\theta_0$ . Suppose that  $\lim_{m,n\to\infty} \frac{m}{m+n} = c \in (0,1)$ . Then, Assumption 4 is satisfied for  $D_{m,n}(p,q) = W_p^p(p^m,q^n)$  and  $D(p,q) = W_p^p(p,q)$  with  $r_{m,n} = n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)}$ .

*Proof of Lemma S5.* By Lemma S3, Assumption 4(a) is satisfied with  $r_{m,n} = n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)}$ . Take  $\mathscr{V}(\theta) = 2^p \mathbb{E}_{p_{\theta}} \left[ \|X\|_2^p \right] + 2^p \mathbb{E}_{p_0} \left[ \|Y\|_2^p \right]$ . From Lemma S2, for any (m, n), we have

$$\operatorname{Var}\left(\operatorname{W}_{p}^{p}(\operatorname{p}^{n},\operatorname{q}^{m})\right) \leq \left(\frac{1}{m}+\frac{1}{n}\right)^{2} \mathscr{V}(\boldsymbol{\theta}).$$
(83)

Since  $\mathscr{X}$  is compact,  $\mathscr{V}(\theta_0)$  is finite and  $\mathscr{V}(\theta)$  is continuous at  $\theta_0$  by assumption. Hence, Assumption 4(b) is verified with  $r_{m,n} = m^{-1} + n^{-1}$ . Taking  $r_{m,n} = n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)}$  provides an upper bound for both conditions.

*Proof of Theorem S2.* When  $p \ge 1$ , the  $W_p$  metric is a metric and continuous under the weak topology. Lemma S4 implies the uniform boundedness in Assumption 3. Lemma S5 verify

Assumption 4 a rate of  $r_{m,n} = n^{-2/(\kappa \vee 4)} + m^{-2/(\kappa \vee 4)}$ . This allows us to conclude that Theorem 2 that

$$\mathbb{E}_{pm}\left[\mathbb{W}_{p}^{p}(\mathbb{p}_{\theta}^{m},\mathbb{p}_{0}^{n})\mid x_{1:n}\right]-\mathbb{W}_{p}^{p}\left(\mathbb{p}_{\theta_{0}},\mathbb{p}_{0}\right)=O_{\mathbb{P}_{0}}\left(n^{-2/(\kappa\vee4)}+m^{-2/(\kappa\vee4)}\right).$$
(84)

#### **D.3.2 MMD**

We establish the convergence rate for  $\mathbb{E}_{pm} \left[ MMD_U^2 \left( p_{\theta}^m, p_0^n \right) \mid x_{1:n} \right]$ , based on the following U-statistic approximation to MMD (Gretton et al., 2012):

$$MMD_{U}^{2}(\mathbf{p}^{m},\mathbf{q}^{n}) := \frac{\sum_{i=1}^{m} \sum_{j \neq i} k(x_{i},x_{j})}{m(m-1)} + \frac{\sum_{i=1}^{n} \sum_{j \neq i} k(y_{i},y_{j})}{n(n-1)} - 2\frac{\sum_{i=1}^{m} \sum_{j=1}^{n} k(x_{i},y_{j})}{mn}, \quad (85)$$

with i.i.d samples  $x_{1:m}$  drawn from p and  $y_{1:n}$  drawn from q. We then use this to prove Theorem 4 holds for the MMD, by replacing Theorem 2 with Theorem S3 in the proof of Theorem 4. Note for large-scale applications, MMD is often preferred over Wasserstein because the computation is quadratic rather than cubic in *n*, and the choice of kernel offers additional flexibility and control.

**Theorem S3** (Posterior expected MMD convergence rate). Let Assumptions 1, 3, S9 and S10 be satisfied. Then

$$\mathbb{E}_{pm}\left[MMD_{U}^{2}\left(p_{\theta}^{m}, p_{0}^{n}\right) \mid x_{1:n}\right] - MMD^{2}\left(p_{\theta_{0}}, p_{0}\right) = O_{\mathbb{P}_{0}}\left(n^{-1/2} + m^{-1/2}\right).$$
(86)

Assumptions S9 and S10 impose standard regularity conditions on the RKHS and the parametric model (Gretton et al., 2012).

Assumption S9. The RKHS kernel  $k : \mathscr{X} \times \mathscr{X} \mapsto \mathbb{R}$  is a symmetric, positive semi-definite, characteristic kernel such that  $\sup_{\theta \in \Theta} \mathbb{E}_{X,X' \sim p_{\theta}} \left[ k^{2}(X,X') \right] < \infty$  and  $\mathbb{E}_{Y,Y' \sim p_{\theta}} \left[ k^{2}(Y,Y') \right] < \infty$ .

We require a characteristic kernel to make the MMD a valid statistical divergence (Sriperumbudur et al., 2011). The bounded second moments are sometimes referred to as the *Hilbert-Schmidt* 

*condition*, which is a sufficient assumption for the eigendecomposition of the kernel operator. Examples of kernels that satisfy Assumption S9 include the Gaussian kernel  $k(x,y) = \exp\left(-\|x-y\|_2^2/\gamma^2\right)$ and the Laplace kernel  $k(x,y) = \exp\left(-\|x-y\|_2/\gamma\right)$ .

**Assumption S10.** The mapping  $\theta \mapsto \mathbb{E}_{X,X' \sim p_{\theta}} \left[ k^2(X,X') \right]$  is continuous at  $\theta_0$ .

This assumption requires the parametric model behaves smoothly around  $\theta_0$ , with respect to the RKHS geometry.

#### Proofs

**Lemma S6** (RKHS-Cauchy-Schwarz). *For any*  $x, y \in \mathcal{X}$ ,  $k^2(x,y) \leq k(x,x)k(y,y)$ .

Proof. Under the RKHS formalism,

$$k^{2}(x,y) = \langle k(x,\cdot), k(y,\cdot) \rangle^{2} \leq \langle k(x,\cdot), k(x,\cdot) \rangle^{2} \langle k(y,\cdot), k(y,\cdot) \rangle^{2} = k(x,x)k(y,y).$$
(87)

The following lemma establishes the convergence rate of the empirical MMD, applying the results of Gretton et al. (2012).

**Lemma S7.** Let Assumptions S9 and S10 be satisfied. Then Assumption 4 is satisfied for  $D_{m,n}(p,q) = MMD_U^2(p^m,q^n)$  and  $D(p,q) = MMD^2(p,q)$  with  $r_{m,n} = m^{-1/2} + n^{-1/2}$ .

*Proof of Lemma S7.* Since  $D_{m,n}(p,q)$  is an unbiased estimate of D(p,q), Assumption 4(a) is satisfied for any positive sequence  $r_{m,n} \rightarrow 0$ . For Assumption 4(b), we need to bound the variance of the empirical MMD. First, by Eq. (85), we have

$$\operatorname{Var}\left(\operatorname{MMD}_{U}^{2}(\operatorname{p}^{m},\operatorname{q}^{n})\right) \leq 2\operatorname{Var}_{X_{i}^{iid}}\left(\frac{\sum_{i=1}^{m}\sum_{j\neq i}k(X_{i},X_{j})}{m(m-1)}\right) + 2\operatorname{Var}_{Y_{i}^{iid}}\left(\frac{\sum_{i=1}^{n}\sum_{j\neq i}k(Y_{i},Y_{j})}{n(n-1)}\right) + 8\operatorname{Var}_{X_{i}^{iid}}_{Y_{i}^{iid}}\left(\frac{\sum_{i=1}^{m}\sum_{j=1}^{n}k(X_{i},Y_{j})}{mn}\right).$$

$$(88)$$

The last term is simply the variance of a sum of i.i.d. variables, thus

$$\operatorname{Var}_{X_{i} \stackrel{iid}{\sim} p, Y_{i} \stackrel{iid}{\sim} q} \left( \frac{\sum_{i=1}^{m} \sum_{j=1}^{n} k(X_{i}, Y_{j})}{mn} \right) \leq \frac{\mathbb{E}_{X \sim p, Y \sim q} \left[ k^{2}(X, Y) \right]}{mn}.$$
(89)

By Lemma S6,

$$\operatorname{Var}_{X_{i}^{iid} p, Y_{i}^{iid} q}\left(\frac{\sum_{i=1}^{m} \sum_{j=1}^{n} k(X_{i}, Y_{j})}{mn}\right) \leq \frac{\mathbb{E}_{X \sim p}\left[k(X, X)\right] \mathbb{E}_{Y \sim q}\left[k(Y, Y)\right]}{mn}.$$
(90)

The first two terms in Eq. (88) are variances of one-sample U-statistics. By Serfling (2009, Section 5.2.1, Lemma A), we have the following bound:

$$\operatorname{Var}_{X_{i} \stackrel{iid}{\sim} p}\left(\frac{\sum_{i=1}^{m} \sum_{j \neq i} k(X_{i}, X_{j})}{m(m-1)}\right) \leq \frac{2\operatorname{Var}_{X, X' \sim p}\left(k(X, X')\right)}{m} \leq \frac{2\mathbb{E}_{X, X' \sim p}\left[k^{2}(X, X')\right]}{m}.$$
 (91)

Analogously, we have

$$\operatorname{Var}_{Y_{i} \stackrel{iid}{\sim} q}\left(\frac{\sum_{i=1}^{n} \sum_{j \neq i} k(Y_{i}, Y_{j})}{n(n-1)}\right) \leq 2 \frac{\mathbb{E}_{Y, Y' \sim q}\left[k^{2}(Y, Y')\right]}{n}.$$
(92)

Putting the bounds together and applying Jensen's inequality with some algebra, we obtain a simple bound without the cross term,

$$\operatorname{Var}\left(\operatorname{MMD}_{U}^{2}(\operatorname{p}^{m},\operatorname{q}^{n})\right) \leq \frac{8\mathbb{E}_{X,X'\sim \operatorname{p}}\left[k^{2}(X,X')\right]}{m} + \frac{8\mathbb{E}_{Y,Y'\sim \operatorname{q}}\left[k^{2}(Y,Y')\right]}{n}.$$
(93)

Define

$$\mathscr{V}(\boldsymbol{\theta}) := 8 \left( \mathbb{E}_{X, X' \sim \mathbf{p}_{\boldsymbol{\theta}}} \left[ k^2(X, X') \right] + \mathbb{E}_{Y, Y' \sim \mathbf{p}_0} \left[ k^2(Y, Y') \right] \right).$$
(94)

The function satisfies  $\mathscr{V}(\theta_0) < \infty$  by Assumption S9 and  $\mathscr{V}(\theta)$  is continuous at  $\theta_0$  by Eq. (96).

Then Eq. (93) implies the bound,

$$\operatorname{Var}\left(\operatorname{MMD}_{U}^{2}(\operatorname{p}_{\theta}^{m},\operatorname{p}_{0}^{n})\right) \leq \left(\frac{1}{m} + \frac{1}{n}\right) \mathscr{V}(\theta) \leq r_{m,n}^{2} \mathscr{V}(\theta).$$
(95)

for 
$$r_{m,n} = m^{-1/2} + n^{-1/2}$$
.

**Lemma S8.** Let Assumption S9 be satisfied. Then  $\sup_{\theta \in \Theta} MMD_U^2(p_{\theta}^m, p_0^n) < \infty$ .

*Proof.* This follows directly from the following representation for the MMD (Gretton et al., 2012):

$$\mathrm{MMD}^{2}(\mathbf{p},\mathbf{q}) = \mathbb{E}_{X,X'\sim\mathbf{p}}\left[k(X,X')\right] - 2\mathbb{E}_{X\sim\mathbf{p},Y\sim\mathbf{q}}\left[k(X,Y)\right] + \mathbb{E}_{Y,Y'\sim\mathbf{q}}\left[k(Y,Y')\right].$$
(96)

By Lemma S6, we have

$$\sup_{\theta \in \Theta} \operatorname{MMD}_{U}^{2}(\mathbf{p}_{\theta}^{m}, \mathbf{p}_{0}^{n}) \leq 2 \sup_{\theta \in \Theta} \mathbb{E}_{X, X' \sim \mathbf{p}_{\theta}} \left[ k(X, X') \right] + 2 \mathbb{E}_{Y, Y' \sim \mathbf{p}_{0}} \left[ k(Y, Y') \right] < \infty.$$
(97)

Since the expectation of  $k^2(x, x')$  is uniformly bounded in each case, MMD<sup>2</sup> ( $\mathscr{H}_k, p_{\theta}, p_0$ ) is uniformly bounded by Jensen's inequality.

*Proof of Theorem S3*. The divergence  $MMD_U^2(p_{\theta}^m, p_0^n)$  is slightly different from the MMD between empirical distributions,  $MMD^2(p_{\theta}^m, p_0^n)$ . Only the latter applies in the setting of Theorem 2. But the difference between them is negligible. By Lemma 2 of Briol et al. (2019), however, this difference is bounded by a factor of  $m^{-1} + n^{-1}$ ,

$$\sup_{\theta \in \Theta} \left| \text{MMD}_{U}^{2} \left( \mathbf{p}_{\theta}^{m}, \mathbf{p}_{0}^{n} \right) - \text{MMD} \left( \mathbf{p}_{\theta}^{m}, \mathbf{p}_{0}^{n} \right) \right| \leq 2(m^{-1} + n^{-1}) \sup_{x, x' \in \mathscr{X}} k(x, x') = O(m^{-1} + n^{-1}).$$
(98)

Using this fact, we have

$$\mathbb{E}_{pm}\left[\mathsf{MMD}_{U}^{2}\left(\mathsf{p}_{\theta}^{m},\mathsf{p}_{0}^{n}\right) \mid x_{1:n}\right] - \mathbb{E}_{pm}\left[\mathsf{MMD}^{2}\left(\mathsf{p}_{\theta}^{m},\mathsf{p}_{0}^{n}\right) \mid x_{1:n}\right]$$

$$\leq \sup_{\theta \in \Theta}\left|\mathsf{MMD}_{U}^{2}\left(\mathsf{p}_{\theta}^{m},\mathsf{p}_{0}^{n}\right) - \mathsf{MMD}^{2}\left(\mathsf{p}_{\theta}^{m},\mathsf{p}_{0}^{n}\right)\right| = O(m^{-1} + n^{-1}).$$
(99)

MMD is a pseudometric and continuous under the weak topology. Lemma S8 implies the uniform boundedness in Assumption 3. Lemma S5 verifies Assumption 4 with a rate of  $r_{m,n} = n^{-1/2} + m^{-1/2}$ . This allows us to conclude that

$$\mathbb{E}_{pm}\left[MMD_{U}^{2}\left(\mathbf{p}_{\theta}^{m},\mathbf{p}_{0}^{n}\right) \mid x_{1:n}\right] - MMD^{2}\left(\mathbf{p}_{\theta_{0}},\mathbf{p}_{0}\right) = O_{\mathbb{P}_{0}}\left(n^{-1/2} + m^{-1/2}\right).$$
(100)

Combining the two displays above yields the desired convergence rate.

### D.3.3 KSD

We establish the convergence rate for  $\mathbb{E}_{pm} [KSD_U(p_0^n, p_\theta) | x_{1:n}]$ , where  $KSD_U$  is a one-sample U-statistic (Liu et al., 2016):

$$KSD_U(p^n, q) := \frac{1}{n(n-1)} \sum_{1 \le i \ne j \le n} u_{q,k}(x_i, x_j),$$
(101)

where  $x_{1:n} \stackrel{iid}{\sim} p$  and  $u_{q,k}(x,x') := \nabla \log q(x)k(x,x')\nabla \log q(x') + 2\nabla \log q(x)^{\top} \nabla_{x'}k(x,x') + tr(\nabla_{x,x'}k(x,x'))$ . This implies that Theorem 4 holds for the KSD, by replacing Theorem 2 with Theorem S4 in the proof of Theorem 4.

**Theorem S4** (Posterior expected KSD convergence rate). Let Assumptions 1 and S11 to S13 be satisfied. As  $n \to \infty$ ,  $\mathbb{E}_{pm} \left[ \text{KSD}_U(p_0^n, p_\theta) \mid x_{1:n} \right]$  converges in  $[\mathbb{P}_0^{\infty}]$ -probability to  $\text{KSD}(p_0, p_{\theta_0})$  at the rate of

$$\mathbb{E}_{pm}[\mathrm{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta}) \mid x_{1:n}] = \mathrm{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta_0}) + O_{\mathbb{P}_0}(n^{-1/2}).$$
(102)

Assumptions S11 to S13 involve regularity conditions on the kernel and the parametric model. Recall that  $\Delta_{q,p}(x) := \nabla_x \log p(x) - \nabla_x \log q(x)$ 

**Assumption S11.** The kernel k(x,x') is symmetric, integrally positive definite, and belongs to the Stein class of all continuous densities  $p_0$  and  $p_{\theta}$ . Additionally,  $\sup_{\theta \in \Theta} \|p_0(x)\Delta_{p_{\theta},p_0}(x)\|_2^2 < \infty$ .

Assumption S11 is satisfied when the tail of  $p_0$  decays exponentially and k is positive definite and characteristic. The Stein class requirement ensures that KSD operates as a valid statistical divergence (Liu et al., 2016). For example, the squared-exponential kernel  $k(x,y) = \exp(-||x-y||_2^2/\gamma^2)$ belongs to the Stein class for smooth densities on  $\mathbb{R}^d$ .

We also need assumptions on the regularity of the KSD.

# Assumption S12. There exists an $M_n n^{-1/2}$ neighborhood of $\theta_0$ for some $M_n \to \infty$ such that

- (a) the mapping  $\theta \mapsto \mathbb{E}_{X,X' \sim p_0} \left[ u_{p_{\theta},k}(X,X') \right]$  is twice differentiable with bounded Hessian (in the  $L_2$  sense).
- (b) the mapping  $\theta \mapsto Var_{X,X' \sim p_0} \left[ u_{p_{\theta},k}(X,X') \right]$  is bounded and continuous at  $\theta_0$ .

When  $p_0$  and  $p_{\theta}$  are smooth densities, Assumption S12(a) is equivalent to stating that  $\theta \mapsto KSD(p_0, p_{\theta})$  is twice differentiable with a bounded Hessian in a shrinking neighborhood of  $\theta_0$ . Assumption S12(b) is satisfied when the parametric model is regular at  $\theta_0$ .

To ensure a notion of boundedness for the empirical KSD, we require an additional assumption:

**Assumption S13.** We assume that  $\sup_{x,x' \in \mathbb{R}} k(x,x') < \infty$  and  $\operatorname{KSD}_U(p_0^n, p_\theta) \xrightarrow{a.s.} \operatorname{KSD}(p_0, p_\theta)$  uniformly in  $\theta$ .

The first part of Assumption S13 requires *k* to be uniformly bounded, which holds for e.g. Gaussian or Laplace kernels. The second part of Assumption S13 requires more effort to verify; it essentially involves establishing a uniform law of large numbers (ULLN) for  $KSD_U(p_0^n, p_\theta)$  across all  $\theta \in \Theta$ , as done in Barp et al. (2019).

Sufficient conditions for Assumption S13 includes (1) the domination of  $u_{p_{\theta},k}(x,x')$  by an integrable and symmetric kernel g(x,x'), and (2) the existence of a sequence of sets where the mappings  $\theta \mapsto \mathbb{E}_{X' \sim p_0} \left[ u_{p_{\theta},k}(x,X') \right]$  and  $\theta \mapsto u_{p_{\theta},k}(x,x')$  are equicontinuous for all  $x \in \mathscr{X}$  and all  $(x,x') \in \mathscr{X} \times \mathscr{X}$ , respectively (Yeo and Johnson, 2001). Alternatively, it suffices that the function class  $\{u_{\mathbb{P}_{\theta},k}\}_{\theta \in \Theta}$  is  $[\mathbb{P}_0 \times \mathbb{P}_0]$ -Glivenko-Cantelli.

#### Proofs

**Lemma S9.** Let Assumptions 1, S11 and S12 be satisfied. Then  $\mathbb{E}_{pm}[KSD(p_0, p_{\theta}) | x_{1:n}] = KSD(p_0, p_{\theta_0}) + O_{\mathbb{P}_0}(n^{-1}).$ 

*Proof of Lemma S9.* Let  $B_{M_n n^{-1/2}}(\theta_0)$  be the neighborhood defined in Assumption S12. Let  $E_n$  be the event that  $\hat{\theta}_{MLE} \in B_{M_n n^{-1/2}}(\theta_0)$ . By Assumption 1(a),  $\mathbb{P}_0(E_n) \to 1$  as  $n \to \infty$ . For the rest of the proof, we condition on  $E_n$ .

Let  $S_{\theta} := \nabla_{\theta} KSD(p_{\theta}, p_{\theta_0}), H_{\theta} := \nabla_{\theta}^2 KSD(p_{\theta}, p_{\theta_0})$ . By Assumption S12, there exists L > 0 such that

$$\sup_{\theta \in B_{M_n n^{-1/2}}(\theta_0)} \|H_{\theta}\|_2 \le L.$$
(103)

By Taylor expanding  $KSD(p_0, p_{\theta})$  around the MLE, we have

$$KSD(p_0, p_{\theta}) = KSD(p_0, p_{\hat{\theta}_{MLE}}) + S_{\hat{\theta}_{MLE}}^T (\theta - \hat{\theta}_{MLE}) + (\theta - \hat{\theta}_{MLE})^T H_{\hat{\theta}_{MLE}} (\theta - \hat{\theta}_{MLE}) + o(\|\theta - \hat{\theta}_{MLE}\|_2^2)$$
(104)

By the bonded Hessian condition, we have a sandwich inequality,

$$\left| \mathsf{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta}) - \mathsf{KSD}(\mathbf{p}_0, \mathbf{p}_{\hat{\theta}_{\mathsf{MLE}}}) \right| \le L \|\theta - \hat{\theta}_{\mathsf{MLE}}\|_2^2 + o(\|\theta - \hat{\theta}_{\mathsf{MLE}}\|_2^2).$$
(105)

Applying Assumption 1(b) to the posterior expectation of the above display, we have

$$\mathbb{E}_{pm}[KSD(p_0, p_{\theta}) | x_{1:n}] = KSD(p_0, p_{\hat{\theta}_{MLE}}) + O_{\mathbb{P}_0}(n^{-1}).$$
(106)

For KSD $(p_0, p_{\hat{\theta}_{MLE}})$ , a Taylor expansion around  $\theta_0$  yields,

$$\begin{split} \text{KSD}(\mathbf{p}_{0},\mathbf{p}_{\hat{\theta}_{\text{MLE}}}) &= \text{KSD}(\mathbf{p}_{0},\mathbf{p}_{\theta_{0}}) + S_{\tilde{\theta}}^{T}(\hat{\theta}_{\text{MLE}} - \theta_{0}) \\ &= \text{KSD}(\mathbf{p}_{0},\mathbf{p}_{\theta_{0}}) + LO_{\mathbb{P}_{0}}(\|\hat{\theta}_{\text{MLE}} - \theta_{0}\|_{2}^{2}) = \text{KSD}(\mathbf{p}_{0}\mathbf{p}_{\theta_{0}}) + O_{\mathbb{P}_{0}}(n^{-1}). \end{split}$$
(107)

**Lemma S10.** Let Assumption S11 be satisfied. Assume that  $\sup_{x,x' \in \mathscr{X}} k(x,x') < \infty$ . Then  $\sup_{\theta \in \Theta} \text{KSD}(p_0, p_\theta) < \infty$ .

Proof of Lemma S10. From its definition, we can upper bound the KSD by Cauchy-Schwarz,

$$KSD(\mathbf{p}_{0},\mathbf{p}_{\theta}) = \mathbb{E}_{X,X'\sim\mathbf{p}_{0}}\left[\Delta_{\mathbf{p}_{0},\mathbf{p}_{\theta}}(X)^{T}k(X,X')\Delta_{\mathbf{p}_{0},\mathbf{p}_{\theta}}(X')\right] \leq \sup_{x,x'\in\mathscr{X}}\left\|k(x,x')\right\|_{2}\|\mathbf{p}_{0}(x)\Delta_{\mathbf{p}_{\theta},\mathbf{p}_{0}}(x)\|_{2}^{2}.$$
(108)

Since k(a,b) < C for all  $a, b \in \mathbb{R}$ , for  $x, x' \in \mathscr{X}$ ,  $||k(x,x')||_2 \le \operatorname{tr}(k(x,x')) \le \kappa C$  which uses the assumption that k(x,x') is positive definite. Since  $||p_0(x)\Delta_{p_\theta,p_0}(x)||_2^2$  is uniformly bounded by Assumption S11,  $\operatorname{KSD}(p_0,p_\theta)$  is uniformly bounded.

**Lemma S11.** Let Assumptions S11 and S12 be satisfied. Then Assumption 4 is satisfied for  $D_n(p,q) = KSD_U(p^n,q)$  and D(p,q) = KSD(p,q) with  $r_n = n^{-1/2}$ .

*Proof of Lemma S11*. (Liu et al., 2016, Theorem 3.6) shows that under Assumption *S11*,  $KSD_U(p_0^n, p_\theta)$  is a valid U-statistic for  $KSD(p_0, p_\theta)$ ,

$$\mathbb{E}\left[\mathrm{KSD}_{U}\left(\mathbf{p}_{0}^{n},\mathbf{p}_{\theta}\right)\right] = \mathrm{KSD}(\mathbf{p}_{0},\mathbf{p}_{\theta}). \tag{109}$$

Thus Assumption 4(a) holds true for any  $r_n$ .

To verify Assumption 4(b), we apply the variance bound for one-sample U-statistics (Serfling, 2009, Section 5.2.1, Lemma A) to obtain:

$$\operatorname{Var}\left(\operatorname{KSD}_{U}\left(\operatorname{p}_{0}^{n},\operatorname{p}_{\theta}\right)\right) \leq \frac{2\operatorname{Var}_{X,X'\sim\operatorname{p}_{0}}\left(u_{\operatorname{p}_{\theta},k}(X,X')\right)}{n} \leq \frac{2\mathbb{E}_{X,X'\sim\operatorname{p}_{0}}\left[u_{\operatorname{p}_{\theta},k}^{2}(X,X')\right]}{n}.$$
(110)

Set  $\mathscr{V}(\theta) = \mathbb{E}_{X,X' \sim p_0} \left[ u_{p_{\theta},k}^2(X,X') \right]$  and  $r_n = n^{-1/2}$  provides the desired result.  $\Box$ 

*Proof of Theorem S4.* By the triangle inequality, we have

$$\mathbb{E}_{pm} \left[ \text{KSD}_{U}(p_{0}^{n}, p_{\theta}) \mid x_{1:n} \right] - \text{KSD} \left( p_{0}, p_{\theta_{0}} \right) \\ \leq \underbrace{\mathbb{E}_{pm} \left[ \text{KSD}_{U} \left( p_{0}^{n}, p_{\theta} \right) - \text{KSD} \left( p_{0}, p_{\theta} \right) \mid x_{1:n} \right]}_{A_{n}} + \underbrace{\mathbb{E}_{pm} \left[ \text{KSD} \left( p_{0}, p_{\theta} \right) \mid x_{1:n} \right] - \text{KSD} \left( p_{0}, p_{\theta_{0}} \right)}_{R_{n}}.$$
(111)

Lemma S9 proved that  $R_n = O_{\mathbb{P}_0}(n^{-1})$ . To bound  $A_n$ , we bound its expectation and variance with respect to the randomness in  $KSD_U(\mathbf{p}_0^n, \mathbf{p}_\theta)$ 

By Fubini's theorem, we exchange the integrals and apply the unbiasedness of U-statistics:

$$\mathbb{E}[A_n] = \mathbb{E}_{pm}\left[\mathbb{E}\left[\mathrm{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta}) - \mathrm{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta})\right] \mid x_{1:n}\right] = 0.$$
(112)

For the variance, we have

$$\operatorname{Var}\left[A_{n}\right] = \mathbb{E}_{\operatorname{pm}}\left[\operatorname{Var}\left[\operatorname{KSD}_{U}\left(\operatorname{p}_{0}^{n}, \operatorname{p}_{\theta}\right)\right] \mid x_{1:n}\right] \le 2n^{-1} \mathbb{E}_{\operatorname{pm}}\left[\operatorname{Var}_{X, X' \sim \operatorname{p}_{0}}\left[u_{\operatorname{p}_{\theta}, k}(X, X')\right] \mid x_{1:n}\right].$$
(113)

By Assumption S12, the posterior expectation  $\mathbb{E}_{pm} \left[ \operatorname{Var}_{X,X' \sim p_0} \left[ u_{p_{\theta},k}(X,X') \right] | x_{1:n} \right]$  is bounded in probability. Thus, by Chebyshev's bound, we conclude that  $A_n = O_{\mathbb{P}_0}(n^{-1/2})$ .

Let  $M_n$  be a sequence such that  $M_n \to \infty$  and  $M_n = o(n^{1/2})$ . To control the fluctuation of  $\mathbb{E}_{pm} [KSD_U(p_0^n, p_\theta) | x_{1:n}]$  around  $KSD(p_0, p_\theta)$ , we decompose it into two terms inside and outside of a ball of radius  $M_n n^{-1/2}$ :

$$\mathbb{E}_{pm} \left[ \text{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta}) - \text{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta}) \mid x_{1:n} \right] = \underbrace{\mathbb{E}_{pm} \left[ \left( \text{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta}) - \text{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta}) \right) I_{B_{M_n n^{-1/2}(\theta_0)}} \mid x_{1:n} \right]}_{A_n} + \underbrace{\mathbb{E}_{pm} \left[ \left( \text{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta}) - \text{KSD}(\mathbf{p}_0, \mathbf{p}_{\theta}) \right) I_{B_{M_n n^{-1/2}(\theta_0)}} \mid x_{1:n} \right]}_{B_n}.$$

$$(114)$$

To bound  $A_n$ , we use the ULLN for  $KSD_U(p_0^n, p_\theta)$ . By Holder's inequality, we have

$$A_{n} \leq \sup_{\theta \in \Theta} |\mathrm{KSD}_{U}(\mathbf{p}_{0}^{n},\mathbf{p}_{\theta}) - \mathrm{KSD}(\mathbf{p}_{0},\mathbf{p}_{\theta})| \Pi_{\mathrm{pm}}\left( \|\theta - \theta_{0}\|_{2} > M_{n}n^{-1/2} | x_{1:n} \right)$$
  
=  $o_{\mathbb{P}_{0}}(1)\Pi_{\mathrm{pm}}\left( \|\theta - \theta_{0}\|_{2} > M_{n}n^{-1/2} | x_{1:n} \right).$  (115)

Then we bound the posterior probability using Assumption 1.

$$\Pi_{\mathrm{pm}} \left( \|\boldsymbol{\theta} - \boldsymbol{\theta}_{0}\|_{2} > M_{n} n^{-1/2} | x_{1:n} \right)$$

$$\leq \Pi_{\mathrm{pm}} \left( \sqrt{n} \|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\|_{2} > M_{n} - \sqrt{n} \|\boldsymbol{\theta}_{0} - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\|_{2} | x_{1:n} \right)$$

$$\leq \inf_{\eta} \exp \left( -\eta \left( M_{n} - \sqrt{n} \|\boldsymbol{\theta}_{0} - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\|_{2} \right) \right) \mathbb{E}_{\mathrm{pm}} \left[ \exp \left( \eta \sqrt{n} \|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}_{\mathrm{MLE}}\|_{2} \right) | x_{1:n} \right]$$

$$= O_{\mathbb{P}_{0}} \left( \exp \left( -C'M_{n} \right) \right),$$
(116)

for some constants *C* and *C'*. The third line applies Hoeffding's inequality. The fourth line uses the fact that weak convergence implies the convergence of characteristic functions (to a sub-Gaussian limit). Thus, if we take  $M_n = n^{1/3}$ , we get  $A_n = O_{\mathbb{P}_0}(n^{-1})$ .

To bound  $B_n$ , we analyze its mean and variance with respect to the randomness in KSD<sub>U</sub> ( $p^n$ , q). To control the mean, we apply Fubini's theorem

$$\mathbb{E}\left[B_{n}\right] = \mathbb{E}_{pm}\left[\mathbb{E}\left[\mathrm{KSD}_{U}\left(\mathbf{p}_{0}^{n},\mathbf{p}_{\theta}\right) - \mathrm{KSD}(\mathbf{p}_{0},\mathbf{p}_{\theta})\right]I_{B_{n}-\alpha/2}(\theta_{0}) \mid x_{1:n}\right] = 0.$$
(117)

Now we control the variance. Some calculation shows

$$\operatorname{Var}(B_n) = \mathbb{E}_{\operatorname{pm}}\left[\operatorname{Var}\left(\operatorname{KSD}_U(\mathbf{p}_0^n, \mathbf{p}_{\theta})\right) I_{B_{n^{-\alpha/2}}(\theta_0)} \mid x_{1:n}\right].$$
(118)

Let  $\mathscr{V}(\theta) = \operatorname{Var}_{X,X' \sim p_0} \left[ u_{p_{\theta},k}(X,X') \right]$ . By Eq. (110), we refine the upper bound as follows

$$\operatorname{Var}(B_n) \le 2n^{-1} \sup_{\boldsymbol{\theta} \in \bar{B}_{n^{-\alpha/2}}(\boldsymbol{\theta}_0)} \mathscr{V}(\boldsymbol{\theta}).$$
(119)
As  $n \to \infty$ ,  $\sup_{\theta \in \bar{B}_{n^{-\alpha/2}}(\theta_0)} \mathscr{V}(\theta) \to \mathscr{V}(\theta_0)$  by the continuity of  $\mathscr{V}$  at  $\theta_0$ , thus  $B_n = O(n^{-1/2})$  by Chebyshev's inequality.

Combining the bounds, we obtain

$$\mathbb{E}_{pm}[\text{KSD}_U(p_0^n, p_\theta) \mid x_{1:n}] = \text{KSD}(p_0, p_{\theta_0}) + O_{\mathbb{P}_0}(n^{-1/2}).$$
(120)

## **E** Details on Empirical Studies

## E.1 gNPP

**Semiparametric model.** The semiparametric model in the gNPP follows the model proposed by Hahn et al. (2020) for causal inference, with an additional transformation step to account for non-normality. The model is,

$$y_i = T_{\lambda}^{-1}(\mu(w_i, \hat{a}(w_i)) + \tau a_i + \varepsilon_i), \quad \varepsilon_i \sim \mathcal{N}(0, \sigma^2)$$
(121)

where:

- The function  $T_{\lambda}(\cdot)$  is the Yeo-Johnson transformation (Yeo and Johnson, 2000). It corrects for non-normality in the outcome expression level.
- The function μ(·,·) is a sum of piecewise constant regression trees. We place a BART prior on μ, following previous work in Bayesian causal inference (Hill, 2011).
- The function â(w<sub>i</sub>) is a propensity model, obtained by training a five-layer neural network to predict a<sub>i</sub> from w<sub>i</sub> under a mean-squared loss. Hahn et al. (2020) show that including â(·) in Bayesian causal inference reduces estimator bias.
- The coefficient  $\tau$  determines the effect of the treatment. We place an improper flat prior on

 $\tau$ . For simplicity we assume the (transformed) outcome depends linearly on  $a_i$  even though it may depend nonlinearly on  $w_i$ .

• We place a half-Normal prior on the variance  $\sigma$ .

The Yeo-Johnson transformation,  $T_{\lambda}$ , is a monotonic function used to reduce skewness and approximate normality (Yeo and Johnson, 2000). The parameter  $\lambda$  is fit via maximum likelihood. The transformation is defined as:

$$T_{\lambda}(y) = \begin{cases} \frac{((y+1)^{\lambda}-1)}{\lambda}, & \text{if } y \ge 0, \lambda \ne 0, \\ \log(y+1), & \text{if } y \ge 0, \lambda = 0, \\ \frac{-((-y+1)^{2-\lambda}-1)}{2-\lambda}, & \text{if } y < 0, \lambda \ne 2, \\ -\log(-y+1), & \text{if } y < 0, \lambda = 2. \end{cases}$$

In the model, we assign a BART prior to  $\mu$ . The BART function is represented as a sum of piecewise constant binary regression trees. Each tree  $T_l$  partitions the covariate space  $\mathscr{A} \times \mathscr{X}$ , with each partition element  $A_b$  assigned a parameter  $m_{lb}$ . The function  $g_l(x)$  takes the value  $m_{lb}$  if  $x \in A_b$  and 0 otherwise. The overall function is then  $\mu(x) = \sum_{l=1}^{L} g_l(x)$ .

We use 50 trees, each constrained by a prior that favors small trees and leaf parameters near zero, making them "weak learners." The prior specification follows that of Chipman et al. (2010), where the probability of a node splitting at depth *h* is  $\eta (1 + h)^{-\beta}$  with  $\eta \in (0, 1)$  and  $\beta \in [0, \infty)$ . The splitting variable and cut-point are chosen uniformly at random. Large trees have low prior probability, with typical values  $\eta = 0.95$  and  $\beta = 2$ . Leaf parameters  $m_{lb}$  follow independent priors  $\mathcal{N}(0, \sigma_m^2)$ , where  $\sigma_m = \sigma_0/\sqrt{L}$ . The induced marginal prior for  $\mu(a, w)$  is centered at zero, with 95% of the prior mass within  $\pm 2\sigma_0$ .

We sample from the BART posterior via MCMC, using the PyMC-BART package (Quiroga et al., 2023). In general we found the chains were well-mixed, after a burn-in period of 1000 steps (Figure S5). Our reported results pool samples from four separately initialized chains.

To account for posterior uncertainty in  $w_{1:n}$ , we introduce a Bayesian bootstrap model for

the distribution of  $w_{1:n}$  (Rubin, 1981). The posterior of ATE(p) is obtained as the pushforward distribution under the product posterior of the conditional distribution p and the covariate distribution  $p_w$ :

$$\mathbf{p} \sim \Pi_{\text{BART}}\left(\mathbf{p} \mid x_{1:n}\right), \quad \mathbf{p}_{w} \sim \text{BB}\left(\frac{1}{n}\sum_{i=1}^{n}\delta_{w_{i}}\right),$$
 (122)

where p implicitly contains the parameters for the conditional distribution, and  $p_w$  is the distribution of *w*. We sample from the posterior of the ATE by simulating from the posterior predictive of Eq. (121).

Let  $\Pi_{BART}(ATE(p) | x_{1:n})$  be the posterior distribution of the ATE under the BART model. The gNPP posterior is then given by

$$\widehat{\Pi}\left(\operatorname{ATE}\left(\mathbf{p}\right) \mid x_{1:n}\right) = \Pi_{pm}\left(\operatorname{ATE}\left(\mathbf{p}_{\theta}\right) \mid x_{1:n}\right)\widehat{\eta}_{n} + \Pi_{BART}\left(\operatorname{ATE}\left(\mathbf{p}\right) \mid x_{1:n}\right)\left(1 - \widehat{\eta}_{n}\right)$$

where  $\hat{\eta}_n$  is the generalized mixing weight based on the MMD.



Figure S4: Diagnostic plots of the causal linear model for the effect of FOXP3 (treatment) on GZMH (outcome).



Figure S5: Traceplots of the BART model MCMC inference for the effect of FOXP3 (treatment) on GZMH (outcome), corresponding to the parameter  $\tau$ . Different lines correspond to separate chains (4 total).

**Parametric model.** The parametric model assumes a linear relationship between the target gene expression (y), the treatment gene (a), and the cell-type/state representation (z). The parametric model is specified as:

$$\mathbf{p}_{\theta}(y \mid a, w) = \mathcal{N}\left(c + \tau a + \gamma^{T} z, \sigma^{2}\right), \tag{123}$$

where *c* is the intercept  $\tau$  is the coefficient of the treatment gene *a*,  $\gamma$  is a vector of coefficients for the confounding genes in *w*, and  $\sigma^2$  is the noise variance.

Let  $\theta \equiv [c, \tau, \gamma]$ . Assuming a flat prior  $p(\theta) \propto 1$  and that the variables  $c, \tau, \gamma$  are apriori independent from *a* and *x*, the posterior distribution is given by:

$$\mathbf{p}(\boldsymbol{\theta} \mid \boldsymbol{x}_{1:n}) \propto \mathbf{p}_{\boldsymbol{\theta}}(\boldsymbol{y}_{1:n} \mid \boldsymbol{a}_{1:n}, \boldsymbol{z}_{1:n}) = \mathcal{N}\left(\hat{\boldsymbol{\theta}}, \hat{\boldsymbol{V}}\boldsymbol{\sigma}^{2}\right),$$
(124)

where

$$\begin{bmatrix} \hat{c} \\ \hat{\tau} \\ \hat{\gamma} \end{bmatrix} = \hat{\theta} \equiv (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T y, \quad \hat{V} \equiv (\tilde{X}^T \tilde{X})^{-1}, \quad \text{for } \tilde{X} \equiv \begin{bmatrix} 1 & a_1 & z_1^T \\ 1 & a_2 & z_2^T \\ \vdots & \vdots & \vdots \\ 1 & a_n & z_n^T \end{bmatrix}.$$
(125)

Under the parametric model (123), the CATE is constant across x so the posterior of ATE(p)

simplifies to

ATE 
$$(p_{\theta}) = \tau(q_{98}(a) - q_0(a)), \quad \tau \sim \Pi(\tau \mid x_{1:n}).$$

Eq. (124) shows that the joint posterior of  $(c, \tau, \gamma)$  follows a multivariate normal distribution. After marginalization, we obtain the posterior for ATE, denoted  $\Pi_{pm}$  (ATE  $(p_{\theta}) | x_{1:n}$ ).

$$\Pi_{\rm pm}\left({\rm ATE}\left({\rm p}_{\theta}\right) \mid x_{1:n}\right) = \mathcal{N}\left(\hat{\tau}(q_{98}(a) - q_0(a)), \hat{V}_{22}(q_{98}(a) - q_0(a))^2\right),\tag{126}$$

where  $\hat{V}_{22}$  is the second diagonal entry of  $\hat{V}$  defined in Eq. (125).

## E.2 Effect of TCF7 on SELL

Besides considering interventions on FOXP3, we also considered the effect of interventions on *TCF7* (Transcription Factor 7) on *SELL* (L-selectin). Elevated levels of SELL are associated with favorable survival outcomes in breast cancer (Kumari et al., 2021). The gNPP posterior suggests that increasing the expression of TCF7 is likely to increase the expression of SELL (Figure S6a). In this case, the generalized mixing weight places strong weight on the parametric model (Figure S6b).



Figure S6: Effect of TCF7 on SELL. a. Posterior probability of the ATE being positive under the parametric, nonparametric, and gNPP models. *n* denotes the size of the (subsampled) dataset. Values are the median across 10 independent data subsamples and model samples. b. Generalized mixing weights,  $\hat{\eta}_n$ . The estimated confidence interval (CI) is across independent data subsamples and model samples.