# Probing Decaying Dark Matter Using the Post-EoR HI21-cm signal

Mohit Yadav<sup>\*</sup> and Tapomoy Guha Sarkar<sup>†</sup>

Department of Physics, Birla Institute of Technology and Science - Pilani, Rajasthan, India

We propose the HI 21-cm power spectrum from the post-reionization epoch as a probe of a cosmological model with decaying dark matter particles. The unstable particles are assumed to undergo a 2-body decay into a massless and massive daughter. We assume, that a fraction f of the total dark matter budget to be, unstable and quantify the decay using the life-time  $\Gamma^{-1}$  and the relative mass splitting  $\epsilon$  between the parent and the massive daughter. The redshift space anisotropic power spectrum of the post-reionization 21-cm signal brightness temperature, as a tracer of the dark matter clustering, imprints the decaying dark matter model through its effect on background evolution and the suppression of power on small scales. We find that with an idealized futuristic intensity mapping experiment with a SKA-I Mid like radio-array,  $\epsilon$  and  $\Gamma$  can be measured at 3.1% and 4.64% around their fiducial values of  $\epsilon = 0.01$  and  $\Gamma = 0.074 \text{Gyr}^{-1}$  respectively.

#### INTRODUCTION

Dark matter (DM) comprises roughly  $\sim 25\%$  of the total energy density of the universe and plays the most crucial role in cosmological structure formation. Its existence is supported by a diverse range of observations, including galactic rotation curves [1–3], large-scale structure surveys [4–6], strong and weak gravitational lensing [7-25], and anisotropies in the cosmic microwave background (CMB)[26–33]. In the standard cosmological framework, the ACDM model proposes that dark matter is cold, collisionless, and stable, offering a robust framework for understanding the expansion history of the universe and structure formation. Despite its remarkable success, the ACDM model faces observational challenges [34, 35]. Notable amongst these issues is the Hubble tension [36, 37], indicating a discrepancy between the Hubble constant  $(H_0)$  measured from distance ladder estimates using Cepheid variables [38–43] and CMBR observations [44–48]. There is also the consistent mild tension  $\sim 2 - 3\sigma$  in the clustering amplitude quantified through  $S_8 = \sigma_8 (\Omega_m/0.3)^{0.5}$  measured CMB and weak lensing surveys, with the latter consistently reporting lower values of  $S_8$  [49–56]. These cosmological tensions seem to have no resolution within the standard  $\Lambda CDM$ paradigm, and attempts have been made to modify both the dark energy [34, 57-73] and dark matter sectors [74-91]. Any modification of the pre-recombination physics aims to decrease the sound horizon at recombination to alleviate the Hubble tension. However, this has the possibility of spoiling the bounds on other parameters and may even aggravate the  $S_8$  tension. To resolve the  $S_8$ tension, it is typically required to reduce the amplitude of matter fluctuations on small scales. Such suppression can be achieved through modifications of the dark matter properties at low redshifts [92–101]. Unstable dark matter with a massless relativistic daughter can potentially cause such a suppression owing to the erasure of power on small scales [102-108]

There is no strong argument from fundamental physics [109–113] or observations [114–117] to believe that dark

matter is entirely stable. Decaying Dark Matter (DDM) models, which permit dark matter to decay into lighter particles such as dark radiation and warm dark matter, represent a natural extension of the standard model. Such decays alter the evolving energy density of the universe, consequently affecting its expansion history and inhibiting structure formation on small scales. These effects make DDM particularly important for addressing the persistent cosmological tensions that challenge the  $\Lambda CDM$  model. Through the redistribution of energy between radiation and matter, DDM possesses the capacity to mitigate these tensions while adhering to current cosmological data. Moreover, DDM scenarios may also resolve issues with small-scale structures, including "cuspy core" [118] and "missing satellite" problem [119], by reducing the clustering on small scales.

The dynamics of DDM is defined by two principal parameters: the decay rate ( $\Gamma$ ), which determines the lifetime of the particle, and the fraction of rest mass energy ( $\epsilon$ ) allocated to the decay products[120–126]. These parameters imprint on the background evolution, and development of cosmological perturbations, thereby influencing observables like the matter power spectrum. Given our interest in late time cosmology, we are interested in the tomographic study of the post-reionization universe using the redshifted HI 21-cm signal.

The post-reionization IGM for z < 6 is predominantly ionized. While this is true for the diffuse low density gas, a small fraction of neutral hydrogen survives in confined over-dense regions and is self-shielded from the background ionizing radiation. These clumped, dense selfshielded, damped Lyman- $\alpha$  systems (DLAs) [127] are located in regions of highly non-linear matter over density peaks [128–130], and are believed to store bulk (~ 80%) of the HI at z < 4 [131] with HI column density greater than  $2 \times 10^{20}$  atoms/cm<sup>2</sup> [132–134]. These DLA clouds, source the 21-cm radiation emission in the post reionization epoch. The post-reionization HI 21-cm signal has been extensively studied [135–143] and is a powerful cosmological probe [143–147].

The first detection of intensity mapping of 21-cm signal

in the redshift range  $z \sim 0.53 - 1.12$  was achieved using radio observations with the Green Bank Telescope (GBT) [148] indicating that the 21-cm intensity field traces the distribution of galaxies at  $z \approx 1$  [149]. The 21-cm signal was further detected in cross-correlation between the 21-cm signal and galaxies in the WiggleZ Dark Energy Survey [150] at  $z \approx 0.8$ . An observational upper limit on the 21 cm auto-power spectrum has also been reported [151]. The 21-cm power spectrum in auto-correlation has yet to be discovered. This is a primary science goal of several presently operational as well as new or upgraded radio interferometric arrays. Some of these telescopes which aim to detect the cosmological 21cm signal are: the Giant Metrewave Radio Telescope (GMRT)[152], the Ooty Wide Field Array (OWFA)[153], The Canadian Hydrogen Intensity Mapping Experiment (CHIME)[154], the Meer-Karoo Array Telescope (MeerKAT)[155], the Square Kilometer Array (SKA)[156].

The parameters of a model involving decaying dark matter shall affect the cosmological background evolution and matter clustering. In this paper we study the post-reionization HI 21-cm anisotropic power spectrum in redshift space in a cosmology with decaying dark matter and make error projections of model parameters using a futuristic intensity mapping experiment with a SKA 1-Mid like radio interferometer.

### COSMOLOGY WITH DECAYING DARK MATTER (DDM)

In the simplest case, an unstable dark matter particle decays into a single relativistic component known as dark radiation. Such models are very well constrained from CMBR observations [157–159]. We shall consider a 2-body decay scenario [105, 158, 160–163] where a fraction f of the dark matter component consists of unstable particles (denoted by the subscript "0") decays into a massless dark photon (denoted by the subscript "1") [164, 165] and massive daughter particles (as is denoted by the subscript "2") with mass  $m_2$ . The parent unstable particle of mass  $m_0$  assumed to be at rest, decays exponentially with a lifetime  $\tau = 1/\Gamma$ . If  $\epsilon$  denote the fraction of energy of the parent particle transferred to the massless daughter, then from conservation of energy and momentum,

$$\epsilon = \frac{1}{2} \left( 1 - \frac{m_0^2}{m_2^2} \right) \qquad \& \qquad \beta_2 = \frac{v_2}{c} = \frac{\epsilon}{1 - \epsilon} \qquad (1)$$

where,  $v_2$  is the magnitude of the "kick" velocity of the massive daughter. In the non-relativistic limit,  $v_2 = \epsilon c$ . Thus, the dark-matter sector now comprises four species of particles: Stable cold dark matter (CDM), parent unstable decaying dark matter (DDM), massive daughter, and massless dark photon. The evolution of the densities ( $\rho_{\rm cdm}, \rho_0, \rho_1, \rho_2$ ) of these particles in an expanding universe is governed by [122-126, 166]

$$\frac{d\rho_{\rm cdm}}{dt} + 3\frac{\dot{a}}{a}\rho_{\rm cdm} = 0 \tag{2}$$

$$\frac{d\rho_0}{dt} + 3\frac{\dot{a}}{a}\rho_0 = -\Gamma\rho_0 \tag{3}$$

$$\frac{d\rho_1}{dt} + 4\frac{\dot{a}}{a}\rho_1 = \epsilon\Gamma\rho_0 \tag{4}$$

$$\frac{d\rho_2}{dt} + 3(1+w_2(a))\frac{\dot{a}}{a}\rho_2 = (1-\epsilon)\Gamma\rho_0 \qquad (5)$$

where,  $w_2(a)$  is the equation of state for the massive daughter particle. The first two equations are not coupled and can be solved with suitable initial conditions, as

$$\rho_{\rm cdm}(a) = (1-f) \frac{\Omega_{mo}}{a^3} \& \rho_0(a) = \mathcal{A} \frac{e^{-\Gamma t(a)}}{a^3}, \quad (6)$$

where  $\mathcal{A}$  is a normalization set by the initial condition on  $\rho_0(a)$ . Assuming that dark matter does not decay prior to recombination ( $z_{\rm rec} \sim 1090$ ), this normalization is obtained by the condition  $\rho(a_{\rm rec}) = f\rho_{\rm crit}\Omega_{mo}(1+z_{\rm rec})^3$ , where  $\rho_{\rm crit} = 3H_0^2/8\pi G$  and,  $\Omega_{mo}$  is the total dark matter density parameter at the present epoch.



Figure 1. A schematic representation of the 4-momentum  $p^{\mu} = (E, \mathbf{p})$  in a two-body decay with a massive parent particle decaying into a massless and massive daughter

Using the solution  $\rho_0(a)$  from Eq. 6 we have

$$d\left(\rho_{1}a^{4}\right) = \epsilon \mathcal{A}\Gamma e^{-\Gamma t} a dt \tag{7}$$

On integration this gives us

$$\rho_1(a) = \frac{\epsilon \mathcal{A}}{a^4} \left( \int_{a_{\rm rec}}^a e^{-\Gamma t(a')da'} - ae^{-\Gamma t(a)} + a_{\rm rec}e^{-\Gamma t(a_{\rm rec})} \right)$$
(8)

The evolution of the massive daughter is complicated since its equation of state  $w_2(a)$  is dynamic. There is a possibility for this particle to be relativistic initially, and slow down later. If this transition from a relativistic to non-relativistic regime occurs at  $a_D$  for a given particle, then one has to consider the sum of contributions to  $\rho_2(a)$  from different particles. Thus we have [166]

$$\rho_2(a) = \frac{\mathcal{A}\Gamma\sqrt{1-2\epsilon}}{a^3} \int_{a_*}^a \mathcal{K}\left(a, a_D\right) da_D \tag{9}$$

where

$$\mathcal{K}(a, a_D) \equiv \frac{e^{-\Gamma t(a_D)}}{a_D H_D} \sqrt{\frac{\beta_2^2}{1 - \beta_2^2} \left(\frac{a_D}{a}\right)^2 + 1} \qquad (10)$$

We note that to evaluate  $\rho_0, \rho_1, \rho_2$ , we require t(a) and H(a) which are interdependent on each other, while H(a) itself depends on  $\rho_0, \rho_1, \rho_2$ . Thus, these equations are solved iteratively to obtain H(a).



Figure 2. The Hubble parameter H(z) for different ( $\bar{\tau} = \frac{\Gamma^{-1}}{1 \text{Gyr}}, \epsilon$ ) shows the expected match with  $\Lambda \text{CDM}$  model at high redshifts and departure at low redshifts. The inset shows the evolution of the parent DDM and daughter particles. Here, we have assumed f = 1. The observational results at z = 0.38, 0.51, and 0.61 are from galaxy clustering data of BOSS SDSS III [167]. We have also shown some high redshift data from Lyman- $\alpha$  forest at z = 2.3 [168] and z = 2.4 [169].

$$H^{2}(a) = \frac{H_{0}^{2}}{\rho_{\text{crit}}} \sum \rho_{i}(a)$$
(11)

where we have CDM, baryonic matter, radiation and  $\Lambda$  contributing to the sum other than the contributions from DDM and its daughter particles.

Fig. 2, shows the expansion history for different DDM parameters. The decay of DDM starts at z = 1090. Thus, for sufficiently small  $\epsilon$  and large  $\tau$ , the Hubble parameter follows the  $\Lambda$ CDM behaviour at high redshifts. Only at low redshifts and much after the epoch of recombination does the DDM density falls, while the daughter particles are produced as seen in the inset of Fig. 2. The lifetime  $\Gamma^{-1}$ , thus controls the redshift when the expansion history deviates from  $\Lambda$ CDM whereas  $\epsilon$  determines the fraction of DDM density that goes into the daughters. At low redshifts, the Hubble rate falls below  $\Lambda$ CDM, due to the excess of radiation. It must be noted that we have kept the dark energy sector fixed, *i.e.*  $\Omega_{\Lambda} = 0.69$  is held fixed. However, if we impose an additional constraint at very high redshift that the angular diameter distance to the epoch of recombination is strongly constrained from observations, then it would require  $\Omega_{\Lambda}$  to be strongly correlated with  $\epsilon$  and  $\Gamma$ , whereby the necessary increase in  $\Omega_{\Lambda}$  will actually make H(z) larger than the  $\Lambda$ CDM expectation in the dark energy dominated era.

A crucial cosmological impact of two-body decaying dark-matter scenarios is that small-scale structures are wiped out. This leads to a suppression of the linear matter power spectrum at scales smaller than a 'cut off scale'  $k > k_f$ . The two parameters of the DDM model  $(\epsilon, \Gamma)$ has two distinct roles to play in this suppression. The parameter  $\epsilon$  which measures the fraction of the mass of the DDM particle that transformed into kinetic energy, fixes the value of  $k_f$ . The higher the energy, the smaller the cutoff  $k_f$  since  $k_f \propto \frac{1}{\sqrt{v}}$ . If the lifetime of DDM  $\tau = \Gamma^{-1}$ is small, the abundance of the free-streaming decay products, shall be large. Thus,  $\Gamma$  determines the amount of small-scale power suppression. In addition, the effects of decay become more pronounced at low redshifts for a given  $\Gamma$ , due to the greater abundance of the daughter particles at low redshifts. Thus, the power suppression is more at lower redshifts. All these effects are incorporated into the transfer function of the matter power spectrum at z < 2.35 [170, 171].

#### THE POST-REIONIZATION 21-CM SIGNAL

The IGM is mostly ionized at redshifts  $z \leq 6$ . In this epoch, neutral hydrogen predominantly resides in the self-shielded DLAs, which source the 21-cm radiation. The signal from the individual DLA clouds is extremely weak (< 10µJy). In an intensity mapping experiment, one is typically interested in the low-resolution large-field image. Thus, the discrete nature of DLA sources is not considered. The corresponding Poisson noise owing to this discrete sampling is neglected, assuming that the number density of the DLA emitters is very large [142]. The diffuse collective signal is expected to be detected in radio-interferometric observations. The modeling of the post-reionization HI 21-cm signal is simplified by several assumptions, which are based on observations or derived from numerical simulations.

At low redshifts, the Wouthuysen field effect enhances the population of the triplet state of HI, causing the ratio of the spin temperature and the CMB temperature  $T_s/T_{\gamma}$ to be much larger than unity. As a result, the 21-cm radiation appears as an emission signal against the background of the CMB [4, 172, 173]. Extensive literature on Lyman- $\alpha$  absorption lines in quasar spectra reveals that, in the post-reionization epoch the mean neutral fraction  $\bar{x}_{\rm HI} = \Omega_{gas}/\Omega_b \sim 2.45 \times 10^{-2}$ , which does not evolve in the entire redshift range  $z \leq 6$  [131].

Galaxies are located at the peaks of the dark matter density field and are known to be biased tracers of the dark matter distribution [174–176]. The dark matter distribution is also believed to be generated by a Gaussian random process, and non-Gaussianity, if any, is believed to be small (ignored in this work). The statistical properties of such a random field are thus quantified using the 2-point function/ power spectrum. The HI in the postreionization epoch is housed predominantly (~ 80%) in DLAs which are believed to be galaxies. It is therefore expected that HI distribution shall trace the underlying dark matter density field with a bias  $b_T(k,z) = \left[\frac{P_{\rm HI}(k,z)}{P(k,z)}\right]^{1/2}$  where  $P_{\rm HI}(k,z)$  and P(k,z) denote the HI and dark matter power spectra, respectively.

On large scales, the bias is known to be scaleindependent, but on small scales it is k-dependent. The post-reionization HI bias has been extensively studied using N-body simulations [177–180]. It is seen that the 4

large scale linear bias grows monotonically with redshift for 1 < z < 4 [181]. This feature is shared by galaxy bias as well [175, 182, 183]. There is a steep rise in the 21-cm bias on small scales. This is due to the absence of small mass halos as is expected from the CDM power spectrum and, consequently the HI being distributed only in larger halos. We have adopted the fitting formula for the bias  $b_T(k, z)$  as a function of both redshift z and scale k obtained from numerical simulations [179] of the postreionization signal.

$$b_T(k,z) = \sum_{m=0}^{4} \sum_{n=0}^{2} c(m,n) k^m z^n$$
 (12)

Adopting all the assumptions discussed above, the power spectrum of post-reionization HI 21-cm brightness temperature fluctuations from redshift z is given by [146, 173]

$$P_{\rm HI}(k,z,\mu) = \frac{1}{\alpha_{\parallel}\alpha_{\perp}^{2}} \mathcal{C}^{2}(z) \left[1 + f_{g}\tilde{\mu}^{2}/b_{T}\right]^{2} G_{F_{oG}}(k_{\parallel}) P\left(\tilde{k},z\right); \text{ where } \frac{\mathcal{C}(z)}{4.0\text{mK}} = b_{T}(k,z)\bar{x}_{\rm HI}(1+z)^{2} \left(\frac{\Omega_{b0}h^{2}}{0.02}\right) \left(\frac{0.7}{h}\right) \frac{H_{0}}{H(z)}$$

$$\alpha_{\parallel} = \frac{H^{f}}{H^{r}}, \ \alpha_{\perp} = \frac{D_{A}^{r}(z)}{D_{A}^{f}(z)}, \ \mu = \frac{k_{\parallel}}{k}, \ \tilde{\mu}^{2} = \frac{\mu^{2}}{F^{2} + \mu^{2}(1-F^{2})}, \ \tilde{k} = \frac{k}{\alpha_{\perp}}\sqrt{1 + \mu^{2}(F^{-2}-1)} \text{ and } F = \frac{\alpha_{\parallel}}{\alpha_{\perp}}$$

$$(13)$$

where  $\mu = \hat{\mathbf{k}} \cdot \hat{\mathbf{n}}$ ,  $\beta_T(k, z) = f_g(k, z)/b_T(k, z)$ . The term  $f_g(z, k)\mu^2$  has its origin in the HI peculiar velocities [137, 173] which, as we mentioned, is sourced by the dark matter fluctuations. On the large cosmological scales of interest, HI peculiar velocities are assumed to be determined by the dark matter distribution. Thus, peculiar velocity manifests itself as a redshift space distortion anisotropy (RSD) in the 21-cm power spectrum [184, 185]). Other than this usual anisotropy of the power spectrum due to redshift space distortion, we have also incorporated the nonlinear effect of peculiar velocity dispersion - Finger of God effect [186] and the Alcock-Paczynski (AP) effect [187, 188]. The Finger of God anisotropy is taken to be of the form  $G_{FoG}\left(k_{\parallel},\sigma_{p}\right) = \left(1 + \frac{1}{2}k_{\parallel}^{2}\sigma_{p}^{2}\right)^{-2}$ , where  $\sigma_{p}$  is the pair velocity dispersion. The AP effect manifests as geometric distortion of cosmological structures and arises due to departures of cosmological distances as predicted in the fiducial cosmology from their actual values along the line of sight and in the transverse direction.

Fig.3(a) shows the dimensionless anisotropic 21-cm power spectrum  $k^3 P_{\rm HI}/2\pi^2$  at the fiducial redshift z = 2 in the  $(k_{\parallel}, k_{\perp})$  space. We have taken the fiducial values of the parameters  $(\Omega_m, \epsilon, \Gamma, H_0, f) =$  $(0.3, 0.01, 0.074 \text{Gyr}^{-1}, 67.76 \text{Km/s/Mpc}, 0.5)$ . We find that owing to the RSD and AP effect, the power spectrum has significant departure from spherical symmetry on small scales  $\approx k \geq 1 \text{Mpc}^{-1}$ .

## **RESULTS AND DISCUSSIONS**

The noise estimates for the 21-cm auto power spectrum is obtained by considering a radio-interferometric observation. We adopt the visibility-visibility correlation formalism [189] and assume that the visibility data is suitably binned such that the correlation between baselines owing to the primary beam of the telescope can be ignored and the covariance matrix is diagonal. Thus we have for redshift z corresponding to an observational frequency  $\nu = 1420/(1+z)$ MHz or wavelength  $\lambda = 0.21(1+z)$ m

$$\delta P_{\rm HI}(k,\mu,z) = \frac{P_{\rm HI}(k,\mu,z) + N_T(k,\mu,z)}{\sqrt{N_c}}$$
(14)

where the first term corresponds to the cosmic variance and the instrumental noise is given by

$$N_T = \frac{\lambda^2 T_{sys}^2 r^2 dr/d\nu}{A_e t_{\mathbf{k}}} \tag{15}$$

Here r denotes the comoving distance to the source,  $A_e$  is the effective area of the individual antenna dish,  $T_{sys}$  is the system temperature assumed to be dominated by the sky temperature and

$$t_{\mathbf{k}} = T_o N_{ant} \left( N_{ant} - 1 \right) A_e \rho / 2\lambda^2 \tag{16}$$

is the fraction of the total observation time  $T_o$  for a given pointing spent on the given mode. We have considered a radio-array with  $N_{ant}$  antennae spread out in



Figure 3. Figure (a) shows the dimensionless anisotropic 21-cm power spectrum  $k^3 P_{\rm HI}/2\pi^2$  for a DDM cosmology at the fiducial redhsift z = 2. The departure from spherical symmetry is seen on small scales. Figure (b) shows the signal to noise ratio  $P_{\rm HI}(k_{\parallel}, k_{\perp}, z)/\delta P_{\rm HI}(k_{\parallel}, k_{\perp}, z)$ .

a plane, such that the total number of visibility pairs  $N_{ant}(N_{ant}-1)/2$  are distributed over different baselines according to a normalized baseline distribution function  $\rho(k_{\perp},\nu)$ . The noise is suppressed by a factor  $\sqrt{N_c}$  where  $N_c$  is the number of modes in a given survey volume. Thus we have

$$N_c = 2\pi k^2 \Delta k \Delta \mu r^2 (dr/d\nu) B \lambda^2 / A_e (2\pi)^3 \tag{17}$$

The noise estimates are based on a futuristic SKA1-Mid like intensity mapping experiment. We consider an interferometer with 250 dish antennae each of diameter 15m and antenna efficiency of 0.7. The baseline distribution function is obtained by assuming that the antennae are distributed over  $1km^2$  with a density falling off as  $1/r^2$  with radial distance from the centre. This increases the noise on small scales due to poor visibility sampling. We consider about a 32MHz bandwidth observation at the observing frequency of 470MHz corresponding to the probing redshift of z = 2. We also consider a k- binning with  $\Delta k = k/N_{bin} \ln(k_{max}/k_{min})$ , with  $N_{bin} = 40$  and where  $k_{max}$  and  $k_{min}$  are set by the largest baseline and the dish size, respectively. We consider 400 hrs observations per pointing with multiple pointings for a full sky survey. For the given observa-

Table I. Telescope parameters used in our analysis.

$N_{\rm ant}$	Antennae efficiency	$D_{\rm dish}$	$T_{\rm o}$	$T_{\rm sys}$	В
250	0.7	$15 \mathrm{m}$	$400~{\rm hrs}$	60K	32MHz

tional parameters, a possible range of scales to be probed is  $0.021 \text{Mpc}^{-1} \leq k < 1.5 \text{Mpc}^{-1}$  at a fiducial redshift z = 2, which corresponds to an observational frequency of 473MHz. Fig.3 (b) shows the signal to noise ratio for the 21-cm power spectrum. We find that a SNR > 7 can be reached in the k-range  $0.1 \text{Mpc}^{-1} \leq k \leq 1.1 \text{Mpc}^{-1}$ .



Figure 4. Error projections at the 68% and 95% and 99% confidence levels for the parameters  $(H_o, \Gamma, \epsilon, f)$ 

While poor visibility sampling is the cause of the cosmic variance noise on large scales, it is the instrument noise that dominates on small scales.

The sensitivity of the 21-cm power spectrum to the DDM parameters occurs due to the suppression of power on small scales and also due to the change in the background expansion rate H(z) and the growth rate of perturbations that affect the normalization C(z) and the anisotropy of the power spectrum. We choose  $q_m = (\epsilon, \Gamma, H_o, f)$  as the parameters to be constrained from a 21-cm intensity mapping experiment described above. The Fisher matrix

$$F_{mn} = \sum_{k,\mu} \frac{1}{\delta P_{\rm HI}(k,\mu,z)^2} \frac{\partial P_{\rm HI}(k,\mu,z)}{\partial q_m} \frac{\partial P_{\rm HI}(k,\mu,z)}{\partial q_n}$$
(18)

allows us to put errors from the Cramer-Rao bound  $\delta q_m = \sqrt{F_{mm}^{-1}}$ , assuming a Gaussian likelihood. If we assume the fiducial cosmology to be the  $\Lambda CDM$  model, then bounds on DDM model parameters can be obtained using the Fisher matrix. We find that for the intensity mapping experiment described above, the  $1 - \sigma$  bounds on the parameters are  $\epsilon \leq 0.07$  and  $\Gamma \leq 0.072 \text{Gyr}^{-1}$ . These bounds are consistent with the literature [190]. We choose fiducial parameter values which are motivated by the bounds obtained on these parameters from our finding and studies of other probes (like the BOSS Lyman-

alpha studies [190]). Fig. 4 shows the 68% and 95% and 99% error contours. The  $1-\sigma$  errors are given in table 2. We have marginalized over the well-constrained parameter  $\Omega_{mo}$ . The correlation between the parameters are on expected lines with increase in  $\epsilon$  and  $\Gamma$  imply greater suppression of power and over greater range of scales respectively. We find that for the given fiducial cosmology and a futuristic SKA like observation we can measure  $\epsilon$  and  $\Gamma$  at 3.1% and 4.64% respectively.

We note here that the dark energy sector has been kept entirely untouched in our analysis and we have assumed a constant dark energy equation of state w = -1. Any dynamical dark energy equation of state would affect both the background evolution and the structure formation. The amplitude of the power spectrum C(z) is sensitive to the IGM astrophysics and astrophysical parameters like the neutral fraction, or the large-scale HIbias have strong degeneracy with  $H_o$  and  $\sigma_8$ . We, therefore focus only on the parameters of the DDM model.

Parameters	$\epsilon$	$\Gamma (\mathrm{GYr}^{-1})$	$H_0 \ (\rm km/s/Mpc)$	f
$1 - \sigma$ error	$0.010 \pm 3.07 \times 10^{-4}$	$0.074 \pm 3.426 \times 10^{-3}$	$67.76\pm0.036$	$0.50\pm0.01$

Table II. Fiducial values and  $1\sigma$  uncertainties for parameters.

While, 21-cm intensity mapping is a promising probe of cosmology beyond the  $\Lambda CDM$  model, such experiments are plagued with the problem of foregrounds originating from sources like galactic Synchrotron radiation, free-free emission, or extra-galactic radio point sources [191–193]. These foregrounds are  $\approx 5$  orders of magnitude larger [189, 194] than the signal and present a formidable challenge towards detecting the 21-cm signal. While the issue of foregrouds is less challenging for the post-reionization signal as compared to the epoch of reionization, significant foreground cleaning is required before the cosmological signal can be explored. The difference of the spectral properties of the foregrounds from that of the 21-cm signal is often exploited to remove the foregrounds from observed visibility data or from the multifrequency angular power spectrum [193, 195–201]. Foreground from sources in and around the field of view can be removed by avoiding the modes in the "foreground wedge" while estimating the 21-cm power spectrum [202–206]. Apart from foregrounds, RFI's pose a critical problem towards detecting the signal. Further, bandpass calibration errors introduce spurious spectral structure into the foreground signal. This difficulty has led to many proposals for precise bandpass calibration [207–215]. We conclude by noting that these major observational issues would degrade the signal, and a more complete realistic analysis invoking some of these issues is necessary.

- \* E-mail: p20210462@pilani.bits-pilani.ac.in
  † E-mail: tapomoy1@gmail.com
- J. R. Brownstein and J. W. Moffat, The Astrophysical Journal 636, 721 (2006), arXiv:astro-ph/0506370 [astroph].
- [2] V. C. Rubin and J. Ford, W. Kent, The Astrophysical Journal 159, 379 (1970).
- [3] A. Botteon, R. Cassano, and e. Eckert, Astronomy and Astrophysics 630, A77 (2019), arXiv:1908.07527 [astroph.CO].
- [4] A. Loeb and M. Zaldarriaga, Physical Review Letters 92, 211301 (2004), arXiv:astro-ph/0312134.
- [5] W. J. Percival, S. Cole, D. J. Eisenstein, R. C. Nichol, J. A. Peacock, A. C. Pope, and A. S. Szalay, Monthly Notices of the Royal Astronomical Society 381, 1053–1066 (2007).
- [6] H. Okada, T. Totani, and S. Tsujikawa, Physical Review D 87, 10.1103/physrevd.87.103002 (2013).
- [7] B. Jain and U. Seljak, The Astrophysical Journal 484, 560–573 (1997).
- [8] L. Hui and N. Y. Gnedin, Monthly Notices of the Royal Astronomical Society 292, 27–42 (1997).
- [9] U. Seljak and M. Zaldarriaga, Physical Review Letters 82, 2636–2639 (1999).
- [10] M. Bartelmann and P. Schneider, Physics Reports 340, 291–472 (2001).
- [11] D. Huterer and M. White, The Astrophysical Journal 578, L95–L98 (2002).

- [12] W. Hu and T. Okamoto, The Astrophysical Journal 574, 566–574 (2002).
- [13] L. V. Waerbeke and Y. Mellier, Gravitational lensing by large scale structures: A review (2003), arXiv:astroph/0305089 [astro-ph].
- [14] A. Heavens, Monthly Notices of the Royal Astronomical Society 343, 1327–1334 (2003).
- [15] M. Takada and B. Jain, Monthly Notices of the Royal Astronomical Society 344, 857–886 (2003).
- [16] L. Knox, A. Albrecht, and Y. S. Song, Weak lensing and supernovae: Complementary probes of dark energy (2004), arXiv:astro-ph/0408141 [astro-ph].
- [17] D. Huterer, M. Takada, G. Bernstein, and B. Jain, Monthly Notices of the Royal Astronomical Society 366, 101–114 (2006).
- [18] S. Miyazaki, T. Hamana, R. S. Ellis, N. Kashikawa, R. J. Massey, J. Taylor, and A. Refregier, The Astrophysical Journal 669, 714–728 (2007).
- [19] D. Munshi, P. Valageas, L. Van Waerbeke, and A. Heavens, Physics Reports 462, 67–121 (2008).
- [20] H. Hoekstra and B. Jain, Annual Review of Nuclear and Particle Science 58, 99–123 (2008).
- [21] A. Vallinotto, S. Das, D. N. Spergel, and M. Viel, Physical Review Letters 103, 10.1103/physrevlett.103.091304 (2009).
- [22] M. Takada and B. Jain, Monthly Notices of the Royal Astronomical Society 395, 2065–2086 (2009).
- [23] K. M. Smith, A. Cooray, S. Das, O. Dore, D. Hanson, C. Hirata, M. Kaplinghat, B. Keating, M. LoVerde, N. Miller, and et al., AIP Conference Proceedings 10.1063/1.3160886 (2009).
- [24] T. G. Sarkar, Journal of Cosmology and Astroparticle Physics 2010 (02), 002–002.
- [25] M. Kamionkowski and E. D. Kovetz, Annual Review of Astronomy and Astrophysics 54, 227–269 (2016).
- [26] R. K. Sachs and A. M. Wolfe, Astrophysical Journal 147, 73 (1967).
- [27] E. Komatsu, J. Dunkley, M. R. Nolta, C. L. Bennett, B. Gold, G. Hinshaw, N. Jarosik, D. Larson, M. Limon, L. Page, and et al., The Astrophysical Journal Supplement Series 180, 330–376 (2009).
- [28] N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Bartolo, and et al., Astronomy and Astrophysics 641, 6 (2020).
- [29] K. Subramanian, The physics of cmbr anisotropies (2004), arXiv:astro-ph/0411049 [astro-ph].
- [30] W. Hu, Lecture notes on cmb theory: From nucleosynthesis to recombination (2008), arXiv:0802.3688 [astroph].
- [31] S. Tanaka, S. Yoshiura, K. Kubota, K. Takahashi, A. J. Nishizawa, and N. Sugiyama, Detectability of cmb weak lensing and hi cross correlation and constraints on cosmological parameters (2020), arXiv:1904.10363 [astroph.CO].
- [32] W. Hu, The Astrophysical Journal 557, L79–L83 (2001).
- [33] Planck Collaboration, Ade, P. A. R., Aghanim, N., Arnaud, M., and A. et;al, Astronomy and Astrophysics 594, A14 (2016).
- [34] L. Perivolaropoulos and F. Skara, New Astronomy Reviews 95, 101659 (2022).
- [35] J., T. Padmanabhan, and J. V. Narlikar, arXiv preprint astro-ph/9511102 (1995).

- [36] J.-P. Hu and F.-Y. Wang, Universe 9, 94 (2023).
- [37] E. Abdalla and e. Abellán, Journal of High Energy Astrophysics 34, 49 (2022), arXiv:2203.06142 [astroph.CO].
- [38] W. L. Freedman, The Astrophysical Journal **919**, 16 (2021), arXiv:2106.15656 [astro-ph.CO].
- [39] A. J. Lee, W. L. Freedman, B. F. Madore, K. A. Owens, A. J. Monson, and T. J. Hoyt, The Astrophysical Journal 907, 112 (2021), arXiv:2012.04536 [astro-ph.CO].
- [40] W. L. Freedman and B. F. Madore, Journal of Cosmology and Astroparticle Physics **2023** (11), 050.
- [41] A. Sandage, G. Tammann, A. Saha, B. Reindl, F. Macchetto, and N. Panagia, The Astrophysical Journal 653, 843 (2006).
- [42] R. L. Beaton, W. L. Freedman, B. F. Madore, G. Bono, E. K. Carlson, G. Clementini, M. J. Durbin, A. Garofalo, D. Hatt, I. S. Jang, *et al.*, The Astrophysical Journal 832, 210 (2016).
- [43] A. G. Riess, L. M. Macri, S. L. Hoffmann, D. Scolnic, S. Casertano, A. V. Filippenko, B. E. Tucker, M. J. Reid, D. O. Jones, J. M. Silverman, and et al., The Astrophysical Journal 826, 56 (2016).
- [44] L. Balkenhol and D. Dutcher *et.al* (SPT-3G Collaboration), Phys. Rev. D **104**, 083509 (2021).
- [45] B. Jain and A. Taylor, Physical Review Letters 91, 141302 (2003).
- [46] D. Huterer, Physical Review D 65, 063001 (2002).
- [47] N. Martinet, J. Harnois-Déraps, E. Jullo, and P. Schneider, Astronomy & Astrophysics 646, A62 (2021).
- [48] L. Amendola, M. Kunz, and D. Sapone, Journal of Cosmology and Astroparticle Physics 2008 (04), 013.
- [49] Planck Collaboration, N. Aghanim, Y. Akrami, and e. Ashdown, Astronomy and Astrophysics 641, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [50] K. Kuijken, C. Heymans, A. Dvornik, H. Hildebrandt, J. T. A. de Jong, A. H. Wright, T. Erben, M. Bilicki, B. Giblin, H. Y. Shan, F. Getman, A. Grado, H. Hoekstra, L. Miller, N. Napolitano, M. Paolilo, M. Radovich, P. Schneider, W. Sutherland, M. Tewes, C. Tortora, E. A. Valentijn, and G. A. Verdoes Kleijn, Astronomy and Astrophysics **625**, A2 (2019), arXiv:1902.11265 [astro-ph.GA].
- [51] B. Giblin, C. Heymans, M. Asgari, H. Hildebrandt, H. Hoekstra, B. Joachimi, A. Kannawadi, K. Kuijken, C.-A. Lin, L. Miller, T. Tröster, J. L. van den Busch, A. H. Wright, M. Bilicki, C. Blake, J. de Jong, A. Dvornik, T. Erben, F. Getman, N. R. Napolitano, P. Schneider, H. Shan, and E. Valentijn, Astronomy and Astrophysics 645, A105 (2021), arXiv:2007.01845 [astro-ph.CO].
- [52] H. Hildebrandt, J. L. van den Busch, A. H. Wright, C. Blake, B. Joachimi, K. Kuijken, T. Tröster, M. Asgari, M. Bilicki, J. T. A. de Jong, A. Dvornik, T. Erben, F. Getman, B. Giblin, C. Heymans, A. Kannawadi, C. A. Lin, and H. Y. Shan, Astronomy and Astrophysics 647, A124 (2021), arXiv:2007.15635 [astro-ph.CO].
- [53] H. Aihara and e. Arimoto, Publications of the Astronomical Society of Japan 70, 10.1093/pasj/psx066 (2017).
- [54] T. Hamana, M. Shirasaki, S. Miyazaki, C. Hikage, M. Oguri, S. More, R. Armstrong, A. Leauthaud, R. Mandelbaum, H. Miyatake, A. J. Nishizawa, M. Simet, M. Takada, H. Aihara, J. Bosch, Y. Komiyama, R. Lupton, H. Murayama, M. A. Strauss,

and M. Tanaka, Publications of the Astronomical Society of Japan **72**, 16 (2020), arXiv:1906.06041 [astro-ph.CO].

- [55] The Dark Energy Survey Collaboration, arXiv e-prints , astro-ph/0510346 (2005), arXiv:astro-ph/0510346 [astro-ph].
- [56] A. Amon, D. Gruen, e. Troxel, and DES Collaboration, Physical Review D 105, 023514 (2022), arXiv:2105.13543 [astro-ph.CO].
- [57] K. Dutta, A. Roy, A. A. Ruchika, Sen, and M. M. Sheikh-Jabbari, General Relativity and Gravitation 52, 15 (2020).
- [58] L. Visinelli, S. Vagnozzi, and U. Danielsson, Symmetry 11, 1035 (2019), arXiv:1907.07953 [astro-ph.CO].
- [59] R.-Y. Guo, J.-F. Zhang, and X. Zhang, Journal of Cosmology and Astroparticle Physics **2019** (2), 054, arXiv:1809.02340 [astro-ph.CO].
- [60] S. Kumar, R. C. Nunes, and S. K. Yadav, European Physical Journal C 79, 576 (2019), arXiv:1903.04865 [astro-ph.CO].
- [61] V. Poulin, T. L. Smith, T. Karwal, and M. Kamionkowski, Physical Review Letters 122, 221301 (2019), arXiv:1811.04083 [astro-ph.CO].
- [62] F. Niedermann and M. S. Sloth, arXiv e-prints , arXiv:1910.10739 (2019), arXiv:1910.10739 [astroph.CO].
- [63] J. Sakstein and M. Trodden, Physical Review Letters 124, 161301 (2020), arXiv:1911.11760 [astro-ph.CO].
- [64] E. Di Valentino, E. V. Linder, and A. Melchiorri, Physical Review D 97, 043528 (2018), arXiv:1710.02153 [astro-ph.CO].
- [65] A. Banihashemi, N. Khosravi, and A. H. Shirazi, Physical Review D 99, 083509 (2019), arXiv:1810.11007 [astro-ph.CO].
- [66] A. Banihashemi, N. Khosravi, and A. H. Shirazi, Physical Review D 101, 123521 (2020), arXiv:1808.02472 [astro-ph.CO].
- [67] E. Di Valentino, A. Melchiorri, O. Mena, and S. Vagnozzi, Physics of the Dark Universe 30, 100666 (2020), arXiv:1908.04281 [astro-ph.CO].
- [68] W. Yang, S. Pan, L. Xu, and D. F. Mota, Monthly Notices of the Royal Astronomical Society 482, 1858 (2019), arXiv:1804.08455 [astro-ph.CO].
- [69] W. Yang, S. Pan, and A. Paliathanasis, Monthly Notices of the Royal Astronomical Society 482, 1007 (2019), arXiv:1804.08558 [gr-qc].
- [70] W. Yang, S. Pan, E. Di Valentino, R. C. Nunes, S. Vagnozzi, and D. F. Mota, Journal of Cosmology and Astroparticle Physics **2018** (9), 019, arXiv:1805.08252 [astro-ph.CO].
- [71] E. Mörtsell and S. Dhawan, Journal of Cosmology and Astroparticle Physics 2018 (9), 025, arXiv:1801.07260 [astro-ph.CO].
- [72] G. Alestas, L. Kazantzidis, and L. Perivolaropoulos, Physical Review D 101, 123516 (2020), arXiv:2004.08363 [astro-ph.CO].
- [73] G. Choi, M. Suzuki, and T. T. Yanagida, Physics Letters B 805, 135408 (2020), arXiv:1910.00459 [hep-ph].
- [74] M. Raveri, W. Hu, T. Hoffman, and L.-T. Wang, Physical Review D 96, 103501 (2017), arXiv:1709.04877 [astro-ph.CO].
- [75] P. Ko, N. Nagata, and Y. Tang, Physics Letters B 773, 513 (2017), arXiv:1706.05605 [hep-ph].
- [76] W. J. C. da Silva, H. S. Gimenes, and R. Silva, Astropar-

ticle Physics **105**, 37 (2019), arXiv:1809.07797 [astro-ph.CO].

- [77] M. A. Buen-Abad, R. Emami, and M. Schmaltz, Physical Review D 98, 083517 (2018), arXiv:1803.08062 [hepph].
- [78] J. Alcaniz, N. Bernal, A. Masiero, and F. S. Queiroz, Physics Letters B 812, 136008 (2021).
- [79] F. D'Eramo, R. Z. Ferreira, A. Notari, and J. L. Bernal, Journal of Cosmology and Astroparticle Physics 2018 (11), 014, arXiv:1808.07430 [hep-ph].
- [80] K. Vattis, S. M. Koushiappas, and A. Loeb, Physical Review D 99, 121302 (2019), arXiv:1903.06220 [astroph.CO].
- [81] L. A. Anchordoqui, V. Barger, H. Goldberg, X. Huang, D. Marfatia, L. H. M. da Silva, and T. J. Weiler, Physical Review D 92, 061301 (2015), arXiv:1506.08788 [hepph].
- [82] J. Buch, P. Ralegankar, and V. Rentala, Journal of Cosmology and Astroparticle Physics **2017** (10), 028, arXiv:1609.04821 [hep-ph].
- [83] T. Bringmann, F. Kahlhoefer, K. Schmidt-Hoberg, and P. Walia, Physical Review D 98, 023543 (2018), arXiv:1803.03644 [astro-ph.CO].
- [84] K. L. Pandey, T. Karwal, and S. Das, Journal of Cosmology and Astroparticle Physics 2020 (7), 026, arXiv:1902.10636 [astro-ph.CO].
- [85] A. G. Doroshkevich, A. A. Klypin, and M. Y. Khlopov, Soviet Astronomy 32, 127 (1988).
- [86] D. Hooper, F. S. Queiroz, and N. Y. Gnedin, Physical Review D 85, 063513 (2012), arXiv:1111.6599 [astroph.CO].
- [87] A. El-Zant, W. El Hanafy, and S. Elgammal, The Astrophysical Journal 871, 210 (2019), arXiv:1809.09390 [gr-qc].
- [88] N. Khosravi, S. Baghram, N. Afshordi, and N. Altamirano, Physical Review D 99, 103526 (2019), arXiv:1710.09366 [astro-ph.CO].
- [89] J. Renk, M. Zumalacárregui, F. Montanari, and A. Barreira, Journal of Cosmology and Astroparticle Physics 2017 (10), 020, arXiv:1707.02263 [astro-ph.CO].
- [90] M. Ballardini, M. Braglia, F. Finelli, D. Paoletti, A. A. Starobinsky, and C. Umiltà, Journal of Cosmology and Astroparticle Physics **2020** (10), 044, arXiv:2004.14349 [astro-ph.CO].
- [91] M. Braglia, M. Ballardini, W. T. Emond, F. Finelli, A. E. GümrükçüoVglu, K. Koyama, and D. Paoletti, Physical Review D 102, 023529 (2020), arXiv:2004.11161 [astro-ph.CO].
- [92] A. Schneider, A. Refregier, S. Grandis, D. Eckert, N. Stoira, T. Kacprzak, M. Knabenhans, J. Stadel, and R. Teyssier, Journal of Cosmology and Astroparticle Physics **2020** (4), 020, arXiv:1911.08494 [astro-ph.CO].
- [93] G. Parimbelli, G. Scelfo, S. Giri, A. Schneider, M. Archidiacono, S. Camera, and M. Viel, Journal of Cosmology and Astroparticle Physics **2021** (12), 044.
- [94] D. Pappadopulo, J. T. Ruderman, and G. Trevisan, Physical Review D 94, 035005 (2016), arXiv:1602.04219 [hep-ph].
- [95] S. Heimersheim, N. Schöneberg, D. C. Hooper, and J. Lesgourgues, Journal of Cosmology and Astroparticle Physics 2020, 016.
- [96] F.-Y. Cyr-Racine, K. Sigurdson, J. Zavala, T. Bringmann, M. Vogelsberger, and C. Pfrommer, Physical Review D 93, 123527 (2016), arXiv:1512.05344 [astro-

ph.CO].

- [97] M. Vogelsberger, J. Zavala, F.-Y. Cyr-Racine, C. Pfrommer, T. Bringmann, and K. Sigurdson, Monthly Notices of the Royal Astronomical Society 460, 1399 (2016), arXiv:1512.05349 [astro-ph.CO].
- [98] H. Rubira, A. Mazoun, and M. Garny, JCAP 01, 034, arXiv:2209.03974 [astro-ph.CO].
- [99] M. Joseph, D. Aloni, M. Schmaltz, E. N. Sivarajan, and N. Weiner, Physical Review D 108, 023520 (2023), arXiv:2207.03500 [astro-ph.CO].
- [100] A. Pourtsidou, C. Skordis, and E. J. Copeland, Physical Review D 88, 083505 (2013).
- [101] V. Poulin, J. L. Bernal, E. D. Kovetz, and M. Kamionkowski, Physical Review D 107, 123538 (2023), arXiv:2209.06217 [astro-ph.CO].
- [102] K. Enqvist, S. Nadathur, T. Sekiguchi, and T. Takahashi, Journal of Cosmology and Astroparticle Physics 2015 (9), 067, arXiv:1505.05511 [astro-ph.CO].
- [103] K. Enqvist, S. Nadathur, T. Sekiguchi, and T. Takahashi, Journal of Cosmology and Astroparticle Physics 2020 (4), 015, arXiv:1906.09112 [astro-ph.CO].
- [104] R. Murgia, A. Merle, M. Viel, M. Totzauer, and A. Schneider, Journal of Cosmology and Astroparticle Physics **2017** (11), 046, arXiv:1704.07838 [astroph.CO].
- [105] G. F. Abellán, R. Murgia, and V. Poulin, Physical Review D 104, 123533 (2021), arXiv:2102.12498 [astroph.CO].
- [106] A. Chen, D. Huterer, S. Lee, e. Ferté, and DES Collaboration, Physical Review D 103, 123528 (2021), arXiv:2011.04606 [astro-ph.CO].
- [107] G. Choi and T. T. Yanagida, Physics Letters B 827, 136954 (2022), arXiv:2104.02958 [hep-ph].
- [108] J. Bucko, S. K. Giri, and A. Schneider, Astronomy and Astrophysics 672, A157 (2023), arXiv:2211.14334 [astro-ph.CO].
- [109] N. E. Bomark and L. Roszkowski, Physical Review D 90, 011701 (2014), arXiv:1403.6503 [hep-ph].
- [110] A. Ibarra, A. Ringwald, and C. Weniger, Journal of Cosmology and Astroparticle Physics **2009** (1), 003, arXiv:0809.3196 [hep-ph].
- [111] T. Asaka, K. Ishiwata, and T. Moroi, Physical Review D 75, 065001 (2007), arXiv:hep-ph/0612211 [hep-ph].
- [112] A. Ibarra, D. Tran, and C. Weniger, International Journal of Modern Physics A 28, 1330040 (2013), arXiv:1307.6434 [hep-ph].
- [113] D. Hooper and S. Profumo, Physics Reports 453, 29 (2007), arXiv:hep-ph/0701197 [hep-ph].
- [114] M. G. Aartsen, M. Ackermann, e. Adams, and IceCube Collaboration, Physical Review Letters **113**, 101101 (2014), arXiv:1405.5303 [astro-ph.HE].
- [115] L. Accardo, M. Aguilar, D. Aisa, e. Alvino, and AMS Collaboration, Physical Review Letters 113, 121101 (2014).
- [116] M. Aguilar, D. Aisa, A. Alvino, and e. A. Ambrosi, Physical Review Letters 113, 121102 (2014).
- [117] L. A. Anchordoqui, V. Barger, I. Cholis, H. Goldberg, D. Hooper, A. Kusenko, J. G. Learned, D. Marfatia, S. Pakvasa, T. C. Paul, and T. J. Weiler, Journal of High Energy Astrophysics 1, 1 (2014), arXiv:1312.6587 [astro-ph.HE].
- [118] W. De Blok, Advances in Astronomy **2010** (2010).
- [119] J. S. Bullock, arXiv e-prints, arXiv:1009.4505 (2010), arXiv:1009.4505 [astro-ph.CO].

- [120] A. Ibarra, D. Tran, and C. Weniger, Int. J. Mod. Phys. A 28, 1330040 (2013), arXiv:1307.6434 [hep-ph].
- [121] G. Franco Abellán, R. Murgia, and V. Poulin, Phys. Rev. D 104, 123533 (2021), arXiv:2102.12498 [astroph.CO].
- [122] S. J. Clark, K. Vattis, and S. M. Koushiappas, Phys. Rev. D 103, 043014 (2021).
- [123] G. Blackadder and S. M. Koushiappas, Physical Review D 93, 023510 (2016), arXiv:1510.06026 [astro-ph.CO].
- [124] K. Vattis, S. M. Koushiappas, and A. Loeb, Phys. Rev. D 99, 121302 (2019), arXiv:1903.06220 [astro-ph.CO].
- [125] S. J. Clark, K. Vattis, and S. M. Koushiappas, Phys. Rev. D 103, 043014 (2021), arXiv:2006.03678 [astroph.CO].
- [126] J. Bucko, S. K. Giri, F. H. Peters, and A. Schneider, Astronomy and Astrophysics 683, A152 (2024), arXiv:2307.03222 [astro-ph.CO].
- [127] A. M. Wolfe, E. Gawiser, and J. X. Prochaska, ARAA 43, 861 (2005), arXiv:astro-ph/0509481.
- [128] J. Cooke, A. M. Wolfe, E. Gawiser, and J. X. Prochaska, Astrophysical Journal Letters 636, L9 (2006), arXiv:astro-ph/0511509.
- [129] M. A. Zwaan, J. M. van der Hulst, F. H. Briggs, M. A. W. Verheijen, and E. V. Ryan-Weber, MNRAS 364, 1467 (2005), arXiv:astro-ph/0510127.
- [130] K. Nagamine, A. M. Wolfe, L. Hernquist, and V. Springel, Astrophysical Journal 660, 945 (2007), arXiv:astro-ph/0510729.
- [131] J. X. Prochaska, S. Herbert-Fort, and A. M. Wolfe, ApJ 635, 123 (2005), arXiv:astro-ph/0508361.
- [132] K. M. Lanzetta, A. M. Wolfe, and D. A. Turnshek, The Astrophysical Journal 440, 435 (1995).
- [133] L. J. Storrie-Lombardi, R. G. McMahon, and M. J. Irwin, MNRAS 283, L79 (1996), arXiv:astro-ph/9608147.
- [134] C. Peroux, R. G. McMahon, L. J. Storrie-Lombardi, and M. J. Irwin, MNRAS 346, 1103 (2003), arXiv:astroph/0107045.
- [135] J. S. B. Wyithe and A. Loeb, Monthly Notices of the Royal Astronomical Society **397**, 1926 (2009), arXiv:0808.2323 [astro-ph].
- [136] S. Bharadwaj and S. K. Sethi, Journal of Astrophysics and Astronomy 22, 293 (2001), arXiv:astroph/0203269.
- [137] S. Bharadwaj, B. B. Nath, and S. K. Sethi, Journal of Astrophysics and Astronomy 22, 21 (2001), arXiv:astroph/0003200.
- [138] S. Wyithe and A. Loeb, ArXiv e-prints (2007), arXiv:0708.3392.
- [139] A. Loeb and J. S. B. Wyithe, Physical Review Letters 100, 161301 (2008), arXiv:0801.1677.
- [140] S. Wyithe and A. Loeb, ArXiv e-prints (2008), arXiv:0808.2323.
- [141] E. Visbal, A. Loeb, and S. Wyithe, Journal of Cosmology and Astro-Particle Physics 10, 30 (2009), arXiv:0812.0419.
- [142] S. Bharadwaj and P. S. Srikant, Journal of Astrophysics and Astronomy 25, 67 (2004), arXiv:astro-ph/0402262.
- [143] T. Chang, U. Pen, J. B. Peterson, and P. Mc-Donald, Physical Review Letters **100**, 091303 (2008), arXiv:0709.3672.
- [144] S. Wyithe, A. Loeb, and P. Geil, ArXiv e-prints (2007), arXiv:0709.2955.
- [145] P. Bull, P. G. Ferreira, P. Patel, and M. G. Santos, The Astrophysical Journal 803, 21 (2015).

- [146] S. Bharadwaj, S. K. Sethi, and T. D. Saini, Physical Rev D 79, 083538 (2009), arXiv:0809.0363.
- [147] Y. Mao, M. Tegmark, M. McQuinn, M. Zaldarriaga, and O. Zahn, Physical Rev D 78, 023529 (2008), arXiv:0802.1710.
- [148] T.-C. Chang, U.-L. Pen, K. Bandura, and J. B. Peterson, arXiv e-prints, arXiv:1007.3709 (2010), arXiv:1007.3709 [astro-ph.CO].
- [149] M. Davis, J. A. Newman, and S. M. F. A. C. Phillips, in *Deep Fields: Proceedings of the ESO Workshop Held* at Garching, Germany, 9-12 October 2000 (Springer, 2003) pp. 241–246.
- [150] K. W. Masui, E. R. Switzer, N. Banavar, K. Bandura, C. Blake, L. M. Calin, T. C. Chang, X. Chen, Y. C. Li, Y. W. Liao, A. Natarajan, U. L. Pen, J. B. Peterson, J. R. Shaw, and T. C. Voytek, The Astrophysical Journal Letters **763**, L20 (2013), arXiv:1208.0331 [astro-ph.CO].
- [151] E. R. Switzer *et al.*, Mon. Not. Roy. Astron. Soc. **434**, L46 (2013), arXiv:1304.3712 [astro-ph.CO].
- [152] Https://www.gmrt.ncra.tifr.res.in.
- [153] Http://rac.ncra.tifr.res.in/ort/owfa.html.
- [154] Https://chime-experiment.ca/en.
- [155] Https://www.sarao.ac.za/science/meerkat/aboutmeerkat/.
- [156] Https://www.skao.int/en.
- [157] J. Hubert, A. Schneider, D. Potter, J. Stadel, and S. K. Giri, Journal of Cosmology and Astroparticle Physics 2021 (10), 040, arXiv:2104.07675 [astro-ph.CO].
- [158] T. Simon, G. F. Abellán, P. Du, V. Poulin, and Y. Tsai, Physical Review D 106, 023516 (2022), arXiv:2203.07440 [astro-ph.CO].
- [159] J. Bucko, S. K. Giri, and A. Schneider, Astronomy and Astrophysics 672, A157 (2023), arXiv:2211.14334 [astro-ph.CO].
- [160] M.-Y. Wang and A. R. Zentner, Physical Review D 85, 043514 (2012), arXiv:1201.2426 [astro-ph.CO].
- [161] A. H. G. Peter and A. J. Benson, Physical Review D 82, 123521 (2010), arXiv:1009.1912 [astro-ph.GA].
- [162] D. Cheng, M. C. Chu, and J. Tang, Journal of Cosmology and Astroparticle Physics 2015 (7), 009, arXiv:1503.05682 [astro-ph.CO].
- [163] S. Mau, E. O. Nadler, e. Wechsler, and DES Collaboration, The Astrophysical Journal **932**, 128 (2022), arXiv:2201.11740 [astro-ph.CO].
- [164] J. L. Feng, A. Rajaraman, and F. Takayama, Physical Review Letters **91**, 011302 (2003), arXiv:hepph/0302215 [hep-ph].
- [165] M. Kopp, C. Skordis, D. B. Thomas, and S. Ilić, Physical Review Letters **120**, 221102 (2018), arXiv:1802.09541 [astro-ph.CO].
- [166] G. Blackadder and S. M. Koushiappas, Phys. Rev. D 90, 103527 (2014).
- [167] S. Satpathy, S. Alam, S. Ho, and e. White, Martin, Monthly Notices of the Royal Astronomical Society 469, 1369 (2017), arXiv:1607.03148 [astro-ph.CO].
- [168] N. G. Busca, T. Delubac, J. Rich, S. Bailey, and A. Font-Ribera, Astronomy and Astrophysics 552, A96 (2013), arXiv:1211.2616 [astro-ph.CO].
- [169] T. Delubac, J. E. Bautista, N. G. Busca, J. Rich, D. Kirkby, S. Bailey, and e. Font-Ribera, Andreu, Astronomy and Astrophysics 574, A59 (2015), arXiv:1404.1801 [astro-ph.CO].
- [170] D. Potter, J. Stadel, and R. Teyssier, Compu-

tational Astrophysics and Cosmology 4, 2 (2017), arXiv:1609.08621 [astro-ph.IM].

- [171] J. Bucko, S. K. Giri, F. H. Peters, and A. Schneider, Astronomy and Astrophysics 683, A152 (2024), arXiv:2307.03222 [astro-ph.CO].
- [172] P. Madau, A. Meiksin, and M. J. Rees, Astrophysical Journal 475, 429 (1997), arXiv:astro-ph/9608010.
- [173] S. Bharadwaj and S. S. Ali, MNRAS 352, 142 (2004), arXiv:astro-ph/0401206.
- [174] A. Dekel and O. Lahav, Astrophysical Journal 520, 24 (1999), arXiv:astro-ph/9806193.
- [175] H. J. Mo, Y. P. Jing, and S. D. M. White, MNRAS 282, 1096 (1996), arXiv:astro-ph/9602052.
- [176] K. Yoshikawa, A. Taruya, Y. P. Jing, and Y. Suto, Astrophysical Journal 558, 520 (2001), arXiv:astroph/0104361.
- [177] J. S. Bagla, N. Khandai, and K. K. Datta, Monthly Notices of the Royal Astronomical Society 407, 567–580 (2010).
- [178] T. Guha Sarkar, S. Mitra, S. Majumdar, and T. R. Choudhury, Monthly Notices of the Royal Astronomical Society 421, 3570–3578 (2012).
- [179] D. Sarkar, S. Bharadwaj, and S. Anathpindika, Monthly Notices of the Royal Astronomical Society 460, 4310–4319 (2016).
- [180] I. P. Carucci, F. Villaescusa-Navarro, and M. Viel, Journal of Cosmology and Astroparticle Physics 2017 (04), 001–001.
- [181] F. A. Marín, N. Y. Gnedin, H.-J. Seo, and A. Vallinotto, The Astrophysical Journal 718, 972–980 (2010).
- [182] J. N. Fry, Astrophysical Journal Letters 461, L65 (1996).
- [183] H. J. Mo, S. Mao, and S. D. M. White, MNRAS 304, 175 (1999), arXiv:astro-ph/9807341.
- [184] N. Kaiser, Monthly Notices of the Royal Astronomical Society 227, 1 (1987).
- [185] A. Hamilton, in *The evolving universe* (Springer, 1998) pp. 185–275.
- [186] J. Jackson, Monthly Notices of the Royal Astronomical Society 156, 1P (1972).
- [187] C. Alcock and B. Paczynski, Nature 281, 358 (1979).
- [188] M. Lopez-Corredoira, The Astrophysical Journal 781, 96 (2014).
- [189] M. McQuinn, O. Zahn, M. Zaldarriaga, L. Hernquist, and S. R. Furlanetto, The Astrophysical Journal 653, 815 (2006), arXiv:astro-ph/0512263 [astro-ph].
- [190] L. Fuß and M. Garny, Journal of Cosmology and Astroparticle Physics **2023** (10), 020, arXiv:2210.06117 [astro-ph.CO].
- [191] T. Di Matteo, R. Perna, T. Abel, and M. J. Rees, Astrophys. J. 564, 576 (2002), arXiv:astro-ph/0109241.
- [192] P. A. Shaver, R. A. Windhorst, P. Madau, and A. G. de Bruyn, Astronomy and Astrophysics 345, 380 (1999), arXiv:astro-ph/9901320 [astro-ph].
- [193] A. Ghosh, S. Bharadwaj, S. S. Ali, and J. N. Chengalur, MNRAS **418**, 2584 (2011), arXiv:1108.3707 [astroph.CO].
- [194] M. G. Santos, A. Cooray, and L. Knox, The Astrophysical Journal 625, 575 (2005), arXiv:astro-ph/0408515 [astro-ph].
- [195] G. Paciga, T.-C. Chang, Y. Gupta, R. Nityanada, J. Odegova, U.-L. Pen, J. B. Peterson, J. Roy, and K. Sigurdson, Monthly Notices of the Royal Astronomical Society **413**, 1174 (2011), arXiv:1006.1351 [astro-

ph.CO].

- [196] A. Datta, J. D. Bowman, and C. L. Carilli, The Astrophysical Journal **724**, 526 (2010), arXiv:1005.4071 [astro-ph.CO].
- [197] E. Chapman, F. B. Abdalla, G. Harker, V. Jelić, P. Labropoulos, S. Zaroubi, M. A. Brentjens, A. G. de Bruyn, and L. V. E. Koopmans, Monthly Notices of the Royal Astronomical Society **423**, 2518 (2012), arXiv:1201.2190 [astro-ph.CO].
- [198] F. G. Mertens, A. Ghosh, and L. V. E. Koopmans, Monthly Notices of the Royal Astronomical Society 478, 3640 (2018), arXiv:1711.10834 [astro-ph.CO].
- [199] C. M. Trott, R. Mondal, G. Mellema, S. G. Murray, B. Greig, J. L. B. Line, N. Barry, and M. F. Morales, Astron. Astrophys. 666, A106 (2022), arXiv:2208.06082 [astro-ph.CO].
- [200] K. K. Datta, T. R. Choudhury, and S. Bharadwaj, Monthly Notices of the Royal Astronomical Society 378, 119 (2007), arXiv:astro-ph/0605546 [astro-ph].
- [201] K. M. A. Elahi, S. Bharadwaj, S. Pal, A. Ghosh, S. S. Ali, S. Choudhuri, A. Chakraborty, A. Datta, N. Roy, M. Choudhury, and P. Dutta, mnras 525, 3439 (2023), arXiv:2308.08284 [astro-ph.CO].
- [202] J. C. Pober, A. R. Parsons, J. E. Aguirre, Z. Ali, R. F. Bradley, C. L. Carilli, D. DeBoer, M. Dexter, N. E. Gugliucci, D. C. Jacobs, P. J. Klima, D. MacMahon, J. Manley, D. F. Moore, I. I. Stefan, and W. P. Walbrugh, Astrophysical Journal Letters **768**, L36 (2013), arXiv:1301.7099 [astro-ph.CO].
- [203] J. C. Pober, A. Liu, J. S. Dillon, J. E. Aguirre, J. D. Bowman, R. F. Bradley, C. L. Carilli, D. R. DeBoer, J. N. Hewitt, D. C. Jacobs, M. McQuinn, M. F. Morales, A. R. Parsons, M. Tegmark, and D. J. Werthimer, The Astrophysical Journal **782**, 66 (2014), arXiv:1310.7031

[astro-ph.CO].

- [204] A. Liu, A. R. Parsons, and C. M. Trott, Physical Review D 90, 023018 (2014), arXiv:1404.2596 [astro-ph.CO].
- [205] J. S. Dillon and et. al., Physical Review D 91, 123011 (2015), arXiv:1506.01026 [astro-ph.CO].
- [206] S. Pal, S. Bharadwaj, A. Ghosh, and S. Choudhuri, Monthly Notices of the Royal Astronomical Society 501, 3378 (2021), arXiv:2012.04998 [astro-ph.CO].
- [207] J. M. Goodman, Radio Science 44, RS0A01 (2009), arXiv:0807.1912 [astro-ph].
- [208] S. Kazemi, S. Yatawatta, S. Zaroubi, P. Lampropoulos, A. G. de Bruyn, L. V. E. Koopmans, and J. Noordam, Monthly Notices of the Royal Astronomical Society 414, 1656 (2011), arXiv:1012.1722 [astro-ph.IM].
- [209] I. S. Sullivan, M. F. Morales, and et. al., The Astrophysical Journal 759, 17 (2012), arXiv:1209.1653 [astroph.IM].
- [210] S. Kazemi and S. Yatawatta, Monthly Notices of the Royal Astronomical Society 435, 597 (2013), arXiv:1307.5040 [astro-ph.IM].
- [211] J. S. Dillon and et. al., Monthly Notices of the Royal Astronomical Society 499, 5840 (2020), arXiv:2003.08399 [astro-ph.IM].
- [212] N. S. Kern, Dillon, and et. al., The Astrophysical Journal 890, 122 (2020), arXiv:1910.12943 [astro-ph.IM].
- [213] R. Byrne, M. F. Morales, B. J. Hazelton, and M. Wilensky, Monthly Notices of the Royal Astronomical Society 503, 2457 (2021), arXiv:2004.08463 [astro-ph.IM].
- [214] P. H. Sims, J. C. Pober, and J. L. Sievers, Monthly Notices of the Royal Astronomical Society 517, 935 (2022), arXiv:2206.13553 [astro-ph.IM].
- [215] A. Ewall-Wice, J. S. Dillon, B. Gehlot, A. Parsons, T. Cox, and D. C. Jacobs, The Astrophysical Journal 938, 151 (2022), arXiv:2110.11994 [astro-ph.CO].