The symbiotic star V694 Mon has transitioned from the accreting-only state to the steady burning phase

ULISSE MUNARI¹

¹INAF Astronomical Observatory of Padova, 36012 Asiago (VI), Italy

ABSTRACT

I propose that the drastic photometric and spectroscopic changes affecting the symbiotic star V694 Mon since 2018, are due to its transition from the accreting-only state to steady hydrogen-burning on the surface of the white dwarf, closely mimicking the pattern followed by V4368 Sgr. The phase of peak optical brightness and weakest emission lines has probably been reached in early 2024. The high-velocity absorptions powered by jet-ejection and the wild flickering, which dominated the century-long quiescence, should not reappear as long has nuclear burning will hold (time-scale of decades). The 3,500 L_{\odot} burning luminosity suggests a 0.60 M_{\odot} mass for the WD.

Keywords: symbiotic stars — novae — nuclear burning — accretion

1. EARLY HISTORY OF V694 MON

V694 Mon (= MWC 560 = SS73 5) is a symbiotic star (SySt), in which an M4III cool giant is orbited by a white dwarf (WD) surrounded by a massive accretion disk (Lucy et al. 2020). It was first noted by Merrill & Burwell (1943) and Sanduleak & Stephenson (1973) as showing Balmer emission lines flaked by wide and deep blue-shifted absorptions. Bond et al. (1984) reported about terminal velocities up to -3000 km/s and noted the persistent presence of photometric flickering, with an amplitude of ~0.2 mag and time-scales of a few minutes. According to the historic lightcurve compiled by Luthardt (1991), V694 Mon was highly variable around a mean B=12.0 mag brightness during 1925-1990. Suddenly in 1990, V694 Mon underwent a $\Delta B \sim 2$ mag surge in brightness, never returned to prior fainter levels, and continued the incessant flickering and the wild and chaotic variability around some persisting periodicities (Munari et al. 2016).

The 1990 surge in brightness coincided with an exacerbation of the velocity and complexity of the blue-shifted absorptions, that reached a record -6500 km/s (Szkody et al. 1990). Tomov et al. (1990) attributed the wildly variable absorptions to discrete jet-like ejections with a relatively high degree of collimation and with the direction of the ejection near the line of sight, with the binary system being seen close to face-on conditions. V694 Mon then remained self-similar from 1990 to 2018, with typical optical spectra and line profiles for 2016 being shown in the top and right panels of Fig. 1: a hot continuum from the accretion disk dominates in the blue, the molecular bands of the M giant emerge at reddest wavelengths, and strong FeII and Balmer emission lines are flanked by high-velocity and blue-shifted absorptions, while large and wild flickering has been persistently affecting photometric observations (Tomov et al. 1996; Sokoloski et al. 2001).

2. THE DRASTIC CHANGE WHICH STARTED IN 2018

The appearance radically changed in 2018, when V694 Mon entered a steep rise in brightness accompanied by the disappearance of both flickering and high-velocity absorptions (Goranskij et al. 2018). Flickering and high-velocity absorptions never reappeared afterward (Zamanov et al. 2019; Munari & Dallaporta 2021), while the once-strong emission lines progressively weakened toward near-disappearance.

The brightness of V694 Mon has been increasing since 2018, leveling off to a maximum reached during early 2024, as illustrated by the *B*-band lightcurve from ANS Collaboration presented in the lower-left panel of Fig. 1. Contrary to



Figure 1. Comparing the accreting (2016) and burning (2024) states of V694 Mon. Top: Asiago 1.22m+B&C low-res spectra averaged over 2016 and early 2024. The latter is compared with that of the A7 Ib supergiant HD 148743. Lower right: high resolution H β profiles of V694 Mon for 2016 and 2024 obtained with the Asiago 1.82m+Echelle telescope. Lower left: ANS Collaboration B-band lightcurve of V694 Mon, with the orange line highlighting the smooth rise to burning peak brightness.

1925-2018, the post-2018 lightcurve is much sharper, confirming the complete disappearance flickering. The average spectrum and line profiles of V694 Mon for early 2024 are shown in top and right panels of Fig. 1: the overall appearance is rather similar to that of an A7 Ib supergiant, with sharp lines, no trace of the high-velocity absorption components, and emission limited to a weak H α .

3. NUCLEAR BURNING

I propose that the drastic changes experienced in 2018 are due to V694 Mon transitioning from the accreting-only state (dominating since object discovery) to non-explosive, non-degenerate, thermal-equilibrium and stable hydrogen burning on the surface of its WD.

The median brightness of V694 Mon for early 2024 is V=8.235. At $E_{\rm B-V}=0.13$ reddening and Gaia DR3 distance of 2.35 kpc, the absolute magnitude of the burning WD is $M_V=-4.02$. Assuming that the bolometric correction is the same as for normal A7 lb stars (B.C.=-0.07), the luminosity of the burning WD is $L=3,500 L_{\odot}$ which corresponds to a white dwarf mass of $M_{\rm WD}\sim0.60 M_{\odot}$ following Wolf et al. (2013).

According to the evolutionary scenario for SySts proposed by Munari (2019), the longest intervals are spent accreting onto the WD (at rates which change with time). In the (great) majority of SySts the accreted envelope is non-degenerate, and once the conditions for nuclear burning are reached, the burning ignites non-explosively and continues in thermal equilibrium as long as enough hydrogen fuel is present in the shell, a phase lasting decades to centuries,

with the rise to optical maximum taking a few years to complete (Iben 1982; Fujimoto 1982a,b). The shell is *not* ejected and remains in hydrostatic equilibrium, progressively contracting in radius and rising in surface temperature as the hydrogen fuel is consumed. Once the burning shell shrinks to the radius and temperature of the nuclei of planetary nebulae, accretion-induced ingestion by the shell of limited amounts of mass may trigger short-lived ZAND outbursts. The exhaustion of the hydrogen fuel in the shell eventually stops the burning, and the SySt resumes the long-lasting accretion phase in preparation for a new cycle. The alternative of the shell being instead accreted by the WD in *degenerate* conditions, leads to the *explosive* burning of a nova outburst, like those repeatedly experienced by the symbiotic recurrent novae T CrB, RS Oph, V3890 Sgr, or V745 Sco.

Comparing with the closely similar spectroscopic and photometric evolution presented by other SySts undergoing similar non-explosive, non-degenerate, thermal-equilibrium hydrogen burning (eg. V4368 Sgr), the maximum optical brightness and weakest emission lines may have been reached by V694 Mon in early 2024. If this is truly the case, V694 Mon will now enter a (very) slow decline, characterized by a progressive increase in the photospheric temperature of the burning shell and rise in intensity and excitation of the emission lines and nebular continuum.

Flickering should not return for long, at least as long as the nuclear burning will continue on the surface of the WD. As emphasized by Sokoloski (2003), flickering plays a relevant diagnostic role in SySts: while ubiquitous among accreting-only objects, it is completely absent among burning SySts.

REFERENCES

- Bond, H. E., Pier, J., Pilachowski, C., Slovak, M., & Szkody, P. 1984, in Bulletin of the American Astronomical Society, Vol. 16, 516
- Fujimoto, M. Y. 1982a, ApJ, 257, 752, doi: 10.1086/160029
- —. 1982b, ApJ, 257, 767, doi: 10.1086/160030
- Goranskij, V. P., Zharova, A. V., Barsukova, E. A., & Burenkov, A. N. 2018, The Astronomer's Telegram, 12227, 1
- Iben, I., J. 1982, ApJ, 259, 244, doi: 10.1086/160164
- Lucy, A. B., Sokoloski, J. L., Munari, U., et al. 2020, MNRAS, 492, 3107, doi: 10.1093/mnras/stz3595
- Luthardt, R. 1991, Information Bulletin on Variable Stars, 3563, 1
- Merrill, P. W., & Burwell, C. G. 1943, ApJ, 98, 153, doi: 10.1086/144557
- Munari, U. 2019, in The Impact of Binary Stars on Stellar Evolution, Cambridge Univ. Press, ed. G. Beccari & H. Boffin, arXiv:1909.01389,
 - doi: 10.48550/arXiv.1909.01389

- Munari, U., & Dallaporta, S. 2021, The Astronomer's Telegram, 15066, 1
- Munari, U., Dallaporta, S., Castellani, F., et al. 2016, NewA, 49, 43, doi: 10.1016/j.newast.2016.06.004
- Sanduleak, N., & Stephenson, C. B. 1973, ApJ, 185, 899, doi: 10.1086/152464
- Sokoloski, J. L. 2003, in Astronomical Society of the Pacific Conference Series, Vol. 303, Symbiotic Stars Probing Stellar Evolution, ed. R. L. M. Corradi, J. Mikolajewska, & T. J. Mahoney, 202,

doi: 10.48550/arXiv.astro-ph/0209101

- Sokoloski, J. L., Bildsten, L., & Ho, W. C. G. 2001, MNRAS, 326, 553, doi: 10.1046/j.1365-8711.2001.04582.x
- Szkody, P., Mateo, M., & Schmeer, P. 1990, IAUC, 4987, 1
- Tomov, T., Kolev, D., Georgiev, L., et al. 1990, Nature, 346, 637, doi: 10.1038/346637a0
- Tomov, T., Kolev, D., Ivanov, M., et al. 1996, A&AS, 116, 1
- Wolf, W. M., Bildsten, L., Brooks, J., & Paxton, B. 2013, ApJ, 777, 136, doi: 10.1088/0004-637X/777/2/136
- Zamanov, R., Stoyanov, K., Nikolov, G., et al. 2019, The Astronomer's Telegram, 13236, 1