Detecting the Coupling of Axion Dark Matter to Neutron Spins at Spallation Sources via Rabi Oscillation

Peter Fierlinger, ^{1,*} Jie Sheng, ^{2,3,†} Yevgeny V. Stadnik, ^{4,‡} and Chuan-Yang Xing^{5,§}

¹ TUM School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany

² Tsung-Dao Lee Institute & School of Physics and Astronomy, Shanghai Jiao Tong University, China

³ Key Laboratory for Particle Astrophysics and Cosmology (MOE)

& Shanghai Key Laboratory for Particle Physics and Cosmology,

Shanghai Jiao Tong University, Shanghai 200240, China

⁴ School of Physics, The University of Sydney, Sydney, New South Wales 2006, Australia

⁵ College of Science, China University of Petroleum (East China), Qingdao 266580, China

We propose a novel detection method for axion dark matter using the Rabi oscillation of neutron spins in beam-based measurements. If axions couple to neutron spins, a background oscillating axion dark matter field would drive transitions between spin-up and spin-down neutron states in a magnetic field when the axion particle energy matches the energy gap between the spin states. The transition can be detected in a double-Stern-Gerlach-type apparatus, with the first splitter producing a pure spin-polarized neutron beam and the second splitter selecting spin-flipped signals. Our approach offers enhanced detection capability for axions within the $10^{-12} - 10^{-10} \,\mathrm{eV}$ mass window with the capability to surpass the sensitivity of current laboratory experiments.

Introduction – Dark matter (DM), comprising approximately 80% of the Universe's matter content [1], is supported by numerous astronomical and cosmological observations [2–5]. However, our understanding of the nature of DM remains limited. Current models about DM span a wide mass spectrum, ranging from ultralight wave-like DM [6] to particle-like DM [7] to macroscopic objects like primordial black holes [8, 9]. Among them, the axion has been a well-motivated DM candidate for a long time [6, 10–13]. The axion was originally proposed to solve the strong CP problem [14–17]. String theory also predicts the existence of axion-like particles [18].

Depending on theoretical models, axion DM can interact with various Standard Model particles, including photons, gluons, electrons, protons, and neutrons. Currently, most ground-based direct-detection experiments focus on detecting the axion-photon coupling. Helioscope experiments [19–21] search for axions produced in the Sun and are sensitive to a wide range of axion masses below the keV scale. Haloscope experiments [19, 22–25], on the other hand, concentrate on detecting axion DM in the Galactic halo and exhibit exceptional sensitivity to axions with Compton wavelengths of $\sim 1\,\mathrm{m}$.

In contrast, terrestrial experiments targeting axion couplings to fermions, particularly the pure coupling to neutrons, are relatively scarce. The most sensitive current searches for axion-nucleon couplings utilize magnetometry-based schemes to measure the anomalous magnetic-field-like effects of the axion DM field on nuclear or neutron spins [26–37]. However, these schemes mainly apply to axion DM with lighter masses $\lesssim 10^{-12} \, \text{eV}$. For higher masses, different detection approaches are needed. One such potentially promising direction is the proposed CASPEr experiment [38, 39].

In this paper, we propose a novel detection method for axion DM using the Rabi oscillation of neutron spins in

beam-based measurements. In a uniform magnetic field, neutrons form a two-level system involving their spin-up and spin-down states. Due to the axion-neutron coupling, the axion DM background field acts as an oscillating perturbation on the neutron spins with the oscillation frequency set by the energy of the axion DM particles, which owing to the non-relativistic nature of the Galactic DM is approximately given by the axion rest mass energy. As a result, neutron spins undergo transitions between the spin-up and spin-down states (spin-flips), with the maximum effect occurring when the axion mass-energy matches the energy level difference in the two-level system. This phenomenon is known as Rabi oscillation. We propose an experimental setup employing a double-Stern-Gerlach-type apparatus to precisely measure axion-DMinduced spin-flips in high-intensity neutron beams. New major neutron sources, such as the European Spallation Source (ESS) [40], the Spallation Neutron Source (SNS) in the U.S. [41], and the China Spallation Neutron Source (CSNS) [42] can provide strong neutron beams and facilitate the axion DM search. Ref. [43] recently proposed the use of high-intensity neutron beams to probe the axionneutron coupling using the Ramsey method; however, that method is mainly limited to rather small axion DM masses $\lesssim 10^{-14} \, \text{eV}$. In contrast, the Rabi oscillation approach we propose in the present work enables searches for axion DM with masses as high as 10^{-10} eV.

Rabi Oscillation of Neutron Spins in Axion Background – Rabi oscillation refers to the transition between two states of a two-level system under an oscillatory perturbation. The two-level system can be constructed by applying a uniform magnetic field \mathbf{B}_0 to neutron beams, as neutron spins couple to the magnetic field \mathbf{B}_0 via the Hamiltonian,

$$H_0 = -g_n \frac{e}{2m_p} \mathbf{S} \cdot \mathbf{B}_0 \,. \tag{1}$$

Here, $g_n = -3.826$ is the g-factor of the neutron, m_p is the proton mass, and ${\bf S}$ is defined in terms of the vector of Pauli spin matrix operators ${\boldsymbol \sigma}$ according to ${\bf S} = {\boldsymbol \sigma}/2$. The energy eigenstates of this two-level system are the spin-up $|\uparrow\rangle$ and spin-down $|\downarrow\rangle$ states of neutrons along the direction of the magnetic field, with eigenenergies ω_{\uparrow} and ω_{\downarrow} , respectively. The energy level difference ω_0 is $\omega_0 \equiv |\omega_{\uparrow} - \omega_{\downarrow}| = |g_n|eB_0/2m_p$ with B_0 the magnitude of ${\bf B}_0$.

An axion DM background field can play the role of the oscillatory perturbation field in the Rabi oscillation of neutron spins. Its interaction with nucleon fields N=(p,n) is described by

$$\mathcal{L}_{\rm int} = -\frac{C_N}{2f_a} \partial_\mu a \bar{N} \gamma^\mu \gamma^5 N \,, \tag{2}$$

where C_N is a model- and species-dependent dimensionless constant, f_a is the axion decay constant, and \overline{N} is the Dirac adjoint of the nucleon field. To be cold DM, ultralight axions can be produced non-thermally via misalignment [10-12] in the early Universe, and then form an oscillating field, $a = a_0 \cos(m_a t - \mathbf{p}_a \cdot \mathbf{r})$ with a mass m_a and momentum $\mathbf{p}_a = m_a \mathbf{v}_a$. The velocity \mathbf{v}_a is due to the relative movement between our solar system and the Galactic halo, as well as the random velocities of the axion DM particles themselves, with a typical magnitude $v_a \equiv |\mathbf{v}_a| \simeq 10^{-3}$ in the laboratory frame. The amplitude a_0 is directly related to the local DM energy density $\rho_a \simeq 0.4 \,\mathrm{GeV/cm^3}$ [44] via $\rho_a = m_a^2 a_0^2/2$, neglecting small axion kinetic energy terms and assuming that axions make up the entirety of the DM. The oscillations of the axion DM field are coherent on timescales up to $\tau_{\rm coh} \sim 2\pi/(m_a v_a^2) \sim 2\pi \times 10^6/m_a$. In the non-relativistic limit, the spatial components of the interaction in Eq. (2) can be simplified as [45],

$$H_{\text{int}} = \boldsymbol{\sigma} \cdot \mathbf{B}_a \sin(m_a t), \text{ with } \mathbf{B}_a \equiv \frac{C_N}{2f_a} a_0 \mathbf{p}_a.$$
 (3)

Consider a beam of neutrons propagating in a uniform magnetic field with an axion DM background field acting as a perturbation. The general form of a neutron state at time t is a superposition of spin-up and spin-down states,

$$|\Psi(t)\rangle = c_{\uparrow}(t)e^{-i\omega_{\uparrow}t}\left|\uparrow\right\rangle + c_{\downarrow}(t)e^{-i\omega_{\downarrow}t}\left|\downarrow\right\rangle\,, \tag{4}$$

with two time-dependent coefficients $c_{\uparrow,\downarrow}$. Assuming the neutrons are prepared to be spin-up initially with $c_{\uparrow}(0) = 1$ and $c_{\downarrow}(0) = 0$, these coefficients can be determined by solving the Schrödinger equation with the axion DM perturbation using time-dependent perturbation theory [46].

In principle, the angle between the direction of the applied magnetic field \mathbf{B}_0 and the effective axion magnetic field \mathbf{B}_a can be arbitrary, while only the component of \mathbf{B}_a perpendicular to \mathbf{B}_0 contributes to the spin-flip of neutrons. To maximize the spin-flip effect, the uniform

magnetic field \mathbf{B}_0 can be applied perpendicular to the direction of axion momentum, or equivalently \mathbf{B}_a , which on average is directed either along or opposite to the orbital motion of the Solar System about the Galactic Centre. In this case, the differential equations for the coefficients $c_{\uparrow,\downarrow}$ can be analytically solved by applying the rotatingwave approximation and the probability for the neutron to be in the spin-down state at time t is [46],

$$P_{\downarrow}(\delta;t) = |c_{\downarrow}(t)|^2 \simeq \frac{|\mathbf{B}_a|^2 \sin^2\left(\frac{\sqrt{|\mathbf{B}_a|^2 + \delta^2}}{2}t\right)}{|\mathbf{B}_a|^2 + \delta^2}, \quad (5)$$

where $\delta \equiv m_a - \omega_0$ is the difference between the axion mass-energy and the two-level energy difference. As a typical feature of Rabi oscillation, this probability peaks at resonance when $m_a = \omega_0$,

$$P_{\downarrow}^{\max}(t) = \sin^2\left(\frac{1}{2}|\mathbf{B}_a|t\right) \simeq \left(\frac{C_n a_0|\mathbf{p}_a|}{4f_a}t\right)^2$$
, (6)

where the second equality holds for a tiny axion coupling C_n/f_a or a sufficiently short time t. In deriving Eqs. (5) and (6), we have treated oscillations of the axion DM field as monochromatic, which is a valid approximation for axion masses as high as $10^{-8}-10^{-7}\,\mathrm{eV}$ for the beam parameters we consider in our present paper (see below for more details). Near the peak at $\delta=0$, the probability in Eq. (5) vanishes when $\sqrt{|\mathbf{B}_a|^2+\delta^2}\,t/2=\pm\pi$. Thus, the minimum width of the resonance peak $\Delta\delta$ can be estimated as $\Delta\delta\simeq 4\times 10^{-14}\,\mathrm{eV}\times (0.1\,\mathrm{s}/t)^1$, which is negligibly small for $m_a\gtrsim 10^{-12}\,\mathrm{eV}$ and the interaction times of interest in the present work. This narrow resonance width indicates that the Rabi oscillation approach not only aids in detecting the axion, but also enables precise measurement of its mass.

Proposed Experimental Setup – The experimental setup we propose for detecting an axion DM background field via Rabi oscillation of neutron beams is illustrated in Fig. 1. Modern advanced spallation neutron sources, such as the ESS [40], the SNS [41], and the CSNS [42], can provide extremely intense neutron beams of up to about $10^{12} \, n/\text{s}$. Initially, the neutron spins are randomly oriented. Firstly, using a mirror polarizer, the neutron beam can be pre-polarized to a polarization state of about 99.9% spin-down, losing only about 20% of the neutron number in the process. Then, a Stern-Gerlach apparatus, which generates an inhomogeneous magnetic field with a vertically directed gradient, is employed to eliminate the remaining unwanted spin-up neutrons. As

 $^{^1}$ The interaction time $t\simeq 0.1\,\mathrm{s}$ here corresponds to the neutron flight time τ within $\mathbf{B}_0,$ discussed in more detail in the following sections.




FIG. 1. Schematic experimental setup for detecting neutron spin oscillation. The incoming neutron beam is pre-polarized and then split into two beams with opposing spin states by a Stern-Gerlach apparatus. The spin-down neutrons traverse a region of uniform magnetic field \mathbf{B}_0 , where the axion DM background field induces spin-flips. The spin-flipped neutrons are focused by a magnetic lens, then separated by a second Stern-Gerlach apparatus and subsequently detected.

neutrons traverse this magnetic-field gradient, their magnetic dipoles experience net spin-dependent forces, resulting in the splitting of spin-up and spin-down neutrons into separate beams.

The spin-down neutron beam subsequently propagates through a magnetic field region of length l containing a uniform constant magnetic field directed vertically upwards. The coupling between neutron spins and the magnetic field creates a two-level system, with all the neutrons initially occupying the spin-down ground state.

As discussed in the previous section, if the axion DM mass approaches the energy gap between the two spin states, the oscillating axion background field will induce the Rabi oscillation of neutrons in the beam due to the axion-neutron coupling. Consequently, a small portion of the neutrons will transition to the higher-energy spin-up state.

After a long-distance flight, the neutron beam will spread out in the transverse directions due to velocity variations. We can add a magnetic focusing lens [47] to focus the beam into a smaller size. Finally, a second Stern-Gerlach apparatus, with the same vertically varying magnetic field, is positioned after the magnetic focusing lens to separate neutrons that have undergone a spin-flip. The number of spin-flipped neutrons is then counted using a neutron detector.

The main backgrounds in our experimental setup are controllable. Firstly, during the flight time of $\sim 0.1\,\mathrm{s}$, approximately 10^{-4} of the neutrons decay. The trajectories of charged decay products $(p^+$ and $e^-)$ will be strongly deflected by the Lorentz force due to the magnetic field in the second Stern-Gerlach device, while any remaining decay products can be easily distinguished in the neutron detector by virtue of their relatively low energies (sub-MeV versus a few MeV for the products of the reaction $n + {}^6\mathrm{Li} \to {}^3\mathrm{H} + {}^4\mathrm{He}$). Secondly, the homogeneity of the magnetic field along the flight path is

at the 10^{-3} level. Neutrons may undergo spin-flips due to the magnetic-field gradient, i.e., non-adiabatic Majorana transitions. However, such transitions are positively correlated with the neutron velocity, as faster neutrons experience more rapid changes in the local magnetic field, whereas DM-induced spin-flips are negatively correlated with the velocity, since the DM-induced effect grows with time as shown in Eq. (6). This anti-correlation can help distinguish DM-induced signals from Majorana transition backgrounds. Finally, mHz background-rate neutron detectors based on standard techniques without spatial discrimination have been used previously, e.g., for ultracold neutron detection. Neutron detectors with lower background rates of 10^{-5} Hz (and below) should be feasible to construct by employing particle-physics tracking methods. The reason for this high level of discrimination is that background events emerge mainly from radioactivity in the detector materials, which can be discriminated by proper spatial resolution and fiducial cuts in the volume of interest. As a result, our experiment can be carried out with measurements lasting for one day at a given magnetic-field strength B_0 corresponding to a single specific axion DM mass. The strength of the applied magnetic field can then be incrementally varied to scan over a range of axion DM particle masses.

Projected Sensitivity – Experimental sensitivity is enhanced with increased event rates. Neutron spallation sources can provide strong pulsed neutron beams. For example, at the ESS, around 10^{11} neutrons can be emitted per pulse within $\mathcal{O}(100)$ ms [48]. Considering the 20% neutron number loss in the pre-polarization stage and a factor of 4 loss in intensity due to the divergence of the neutron beam in the transverse directions, $N_n \approx 2 \times 10^{10}$ of these are usable. After running the experiment for one day, $N_{\text{pulse}} \simeq 10^6$ pulses can be emitted and the statistical data can reach $N_n N_{\text{pulse}} \simeq 2 \times 10^{16}$. Typically, neutrons emitted from spallation neutron sources




FIG. 2. Projected sensitivity to the axion-nucleon coupling (assuming that $C_N = C_n = C_p$) assuming one day of running time at a fixed magnetic field strength B_0 . Different axion masses within the red shaded region can be probed by varying the applied magnetic field strength in the range $2.5 \times 10^{-6} - 10^{-3}$ T. The total measurement time required to scan the entire mass range is ≈ 3 years. Constraints from ChangE [35, 36] (light blue), NASDUCK [31, 33] (light yellow), fifth-force experiment [49] (light green), torsion balance experiment [50] (light pink), SNO [51] (light cyan), as well as neutron star cooling [52] (gray), are also shown.

exhibit high velocities with a broad velocity distribution. However, after undergoing a series of scattering and absorption processes in a decelerator, the neutrons can be slowed down to speeds around $v_n \sim \mathcal{O}(1000) \,\mathrm{m/s}$, leading to a longer interaction time with the axion DM field. Taking the length of the magnetic field region to be $l = 40 \,\mathrm{m}$ as an example, the interacting time of the Rabi oscillation is $\tau \equiv l/v_n \sim 0.1 \,\mathrm{s}$. Assuming that background neutron spin-state transitions are sufficiently under control, one can estimate the projected sensitivity to the axion coupling parameters C_n/f_a from the relationship that the transition probability in Eq. (6) is less than $P_{\downarrow}(\tau) \lesssim 1/(N_n N_{\text{pulse}})$. The sensitivity to the axion decay constant is optimized when the axion massenergy matches the energy difference between different spin states in the applied magnetic field, i.e., $\delta = 0$, ac-

$$\frac{f_a}{C_n} \gtrsim \sqrt{\frac{\rho_a v_a^2 \tau^2 N_n N_{\text{pulse}}}{8}}$$

$$= 1.3 \times 10^7 \,\text{GeV} \left(\frac{\tau}{0.1 \,\text{s}}\right) \left(\frac{N_n}{2 \times 10^{10}}\right)^{\frac{1}{2}} \left(\frac{N_{\text{pulse}}}{10^6}\right)^{\frac{1}{2}}.$$
(7)

The sensitivity improves for a smaller neutron velocity and larger neutron number. A smaller velocity results in a longer interaction time with the axion DM field, while a larger neutron number increases the overall probability of observing a spin-flip transition.

For axion mass-energies much larger than the resonance peak width $\Delta \delta$, $m_a \gtrsim \Delta \delta$, a specific value

of B_0 can only measure the corresponding axion mass $m_a = |g_n|eB_0/2m_p = 1.2 \times 10^{-10}B_0/(10^{-3} \,\mathrm{T})\,\mathrm{eV}$. Since the width $\Delta\delta$ depends only on the single-shot time-offlight τ and not on the axion mass m_a , lower axion masses can be scanned over much more quickly than higher axion masses. The magnetic field intensity that can be established stably and uniformly is in the range of $\sim 10^{-9} - 10^{-3} \,\mathrm{T}$. Electric dipole moment (EDM) searches using stored ultracold neutrons, such as the one in [53], require volumes with magnetic-field homogeneity at the sub- 10^{-3} level for spin coherence. In the case of our proposed neutron beam measurements, we employ a similar 0.5 m diameter and similar (albeit shorter) coil geometry. By incrementally varying the magnetic-field strength over the range $2.5 \times 10^{-6} - 10^{-3}$ T (when the resonance is narrow and well-defined), one can achieve the projected sensitivity to axion DM in the mass range $3 \times 10^{-13} - 10^{-10} \,\mathrm{eV}$ shown as the red shaded region in Fig. 2. For our estimates, we have assumed $l = 40 \,\mathrm{m}$ and an integration time of 1 day at each chosen value of B_0 (each corresponding to a particular narrow range of axion masses). In this case, the entire indicated mass range can be scanned in ≈ 3 years of total measurement time.

The projected sensitivity to the axion-neutron coupling in the mass range $10^{-12}-10^{-10}\,\mathrm{eV}$ with our proposed setup can be stronger than the constraints from indirect terrestrial detection methods, such as searches for solar axions in the Sudbury Neutrino Observatory (SNO) experiment [51] (light cyan), the fifth-force search experiment [49] (light green) and the torsion balance experiment [50] (light pink), as well as atomic co-magnetometry experiments, such as NASDUCK [31, 33] (light yellow) and ChangE [35, 36] (light blue). Besides, the axion-neutron coupling also receives strong astrophysical constraints from neutron star cooling [52], which is shown as the gray region in Fig. 2. However, this constraint contains some model dependencies and theoretical uncertainties.

Conclusion – In this paper, we propose utilizing the Rabi oscillation of neutron spins in high-intensity neutron beams at spallation sources, such as the ESS, SNS and CSNS, to detect the axion DM background, and present a novel experimental design to realize it. In our setup, firstly, a Stern-Gerlach apparatus fully polarizes a 99.9% pre-polarized neutron beam. Subsequently, the polarized neutrons enter a region of uniform magnetic field for free propagation. Due to the presence of axion dark matter, the axion-neutron coupling introduces an oscillatory perturbation to the neutron, causing the neutron spin to flip. Finally, a second Stern-Gerlach apparatus allows us to extract the signal of the spin-flip. After background and experimental sensitivity estimation and assuming the experiment runs for a day at a given applied magnetic field strength, our proposal can achieve a sensitivity up to two orders of magnitude stronger than the current best ground-based experimental constraints on the axionneutron coupling over the mass range $10^{-12} - 10^{-10} \, \text{eV}$, reaching the level $f_a/C_n \sim 10^7 \, \text{GeV}$.

By increasing the running time at a fixed magnetic field strength and the intensity of the neutron-beam flux, the sensitivity can be further enhanced. Once a signal is observed, our proposal also has the capability to determine the mass of the axion DM particle.

The axion DM background field can induce an oscillating neutron EDM at the one-loop level via the axiongluon coupling [54]. The EDM oscillates at a frequency corresponding to the axion mass-energy along the neutron spin direction. If a uniform electric field E is applied perpendicular to the homogeneous magnetic field \mathbf{B}_0 inside the magnetically controlled region, the timevarying perturbation due to the coupling between the oscillating neutron EDM and the electric field will induce Rabi oscillations of the neutron spins. In order to minimise undesired systematic effects associated with the additional magnetic field ${\bf B}_{\rm rel}={\bf E}\times{\bf v}/c^2$ seen by moving neutrons with velocity ${\bf v}$, this electric field should be applied along the propagation direction of the neutron beam itself, which severely limits the maximum strength of the applied electric field. On the other hand, for the Ramsey approach discussed in [43] for much lower axion masses, a strong electric field can be applied along \mathbf{B}_0 (which is directed transverse to the neutron beam), with undesired effects of the induced magnetic field $\mathbf{B}_{\mathrm{rel}}$ being strongly suppressed, since the effects of the small component of \mathbf{B}_{rel} directed along \mathbf{B}_0 on the neutrons' Larmor precession frequency are minimised in that case.

ACKNOWLEDGEMENTS

P. F. is supported by the DFG Cluster of Excellence "Origins" (EXC 2094). J. S. is supported by the National Natural Science Foundation of China (Nos. 12375101, 12425506, 12090060, 12090064) and the SJTU Double First Class start-up fund WF220442604. The work of Y. V. S. was supported by the Australian Research Council under the Discovery Early Career Researcher Award No. DE210101593. C.-Y. X. is supported by the Fundamental Research Funds for the Central Universities (No. 24CX06048A). Y. V. S. gratefully acknowledges the hospitality of Tsung-Dao Lee Institute during the writing of this paper.

- * peter.fierlinger@tum.de
- † shengjie04@sjtu.edu.cn
- [‡] yevgenystadnik@gmail.com
- § cyxing@upc.edu.cn
- [1] Planck Collaboration, N. Aghanim et al., "Planck 2018 results. VI. Cosmological parameters," Astron. Astro-

- phys. 641 (2020) A6, [arXiv:1807.06209 [astro-ph.CO]].[Erratum: Astron.Astrophys. 652, C4 (2021)].
- [2] G. Bertone, D. Hooper, and J. Silk, "Particle dark matter: Evidence, candidates and constraints," Phys. Rept. 405 (2005) 279–390, [arXiv:hep-ph/0404175].
- [3] B.-L. Young, "A survey of dark matter and related topics in cosmology," Front. Phys. (Beijing) 12 no. 2, (2017) 121201. [Erratum: Front.Phys.(Beijing) 12, 121202 (2017)].
- [4] A. Arbey and F. Mahmoudi, "Dark matter and the early Universe: a review," Prog. Part. Nucl. Phys. 119 (2021) 103865, [arXiv:2104.11488 [hep-ph]].
- [5] J. Billard et al., "Direct detection of dark matter—APPEC committee report*," Rept. Prog. Phys. 85 no. 5, (2022) 056201, [arXiv:2104.07634 [hep-ex]].
- [6] L. Hui, "Wave Dark Matter," Ann. Rev. Astron. Astrophys. 59 (2021) 247-289, [arXiv:2101.11735 [astroph.CO]].
- [7] S. Profumo, L. Giani, and O. F. Piattella, "An Introduction to Particle Dark Matter," Universe 5 no. 10, (2019) 213, [arXiv:1910.05610 [hep-ph]].
- [8] B. Carr and F. Kuhnel, "Primordial Black Holes as Dark Matter: Recent Developments," Ann. Rev. Nucl. Part. Sci. 70 (2020) 355–394, [arXiv:2006.02838 [astro-ph.CO]].
- [9] B. Carr and F. Kuhnel, "Primordial black holes as dark matter candidates," SciPost Phys. Lect. Notes 48 (2022)
 1, [arXiv:2110.02821 [astro-ph.CO]].
- [10] J. Preskill, M. B. Wise, and F. Wilczek, "Cosmology of the Invisible Axion," Phys. Lett. B 120 (1983) 127–132.
- [11] L. F. Abbott and P. Sikivie, "A Cosmological Bound on the Invisible Axion," Phys. Lett. B 120 (1983) 133–136.
- [12] M. Dine and W. Fischler, "The Not So Harmless Axion," Phys. Lett. B 120 (1983) 137–141.
- [13] C. B. Adams et al., "Axion Dark Matter," in Snowmass 2021. 3, 2022. [arXiv:2203.14923 [hep-ex]].
- [14] R. D. Peccei and H. R. Quinn, "CP Conservation in the Presence of Instantons," Phys. Rev. Lett. 38 (1977) 1440–1443.
- [15] S. Weinberg, "A New Light Boson?," Phys. Rev. Lett. 40 (1978) 223–226.
- [16] F. Wilczek, "Problem of Strong P and T Invariance in the Presence of Instantons," Phys. Rev. Lett. 40 (1978) 279–282.
- [17] R. D. Peccei, "The Strong CP problem and axions," Lect. Notes Phys. 741 (2008) 3-17, [arXiv:hep-ph/0607268].
- [18] P. Svrcek and E. Witten, "Axions In String Theory," JHEP 06 (2006) 051, [arXiv:hep-th/0605206].
- [19] P. Sikivie, "Experimental Tests of the Invisible Axion," Phys. Rev. Lett. 51 (1983) 1415–1417. [Erratum: Phys.Rev.Lett. 52, 695 (1984)].
- [20] CAST Collaboration, K. Altenmüller et al., "A new upper limit on the axion-photon coupling with an extended CAST run with a Xe-based Micromegas detector," [arXiv:2406.16840 [hep-ex]].
- [21] IAXO Collaboration, E. Armengaud et al., "Physics potential of the International Axion Observatory (IAXO)," JCAP 06 (2019) 047, [arXiv:1904.09155 [hep-ph]].
- [22] CAPP Collaboration, S. Ahn et al., "Extensive Search for Axion Dark Matter over 1 GHz with CAPP'S Main Axion Experiment," Phys. Rev. X 14 no. 3, (2024) 031023, [arXiv:2402.12892 [hep-ex]].
- [23] QUAX Collaboration, A. Rettaroli et al., "Search for axion dark matter with the QUAX-LNF tunable

- haloscope," Phys. Rev. D 110 no. 2, (2024) 022008, [arXiv:2402.19063 [physics.ins-det]].
- [24] A. P. Quiskamp, G. Flower, S. Samuels, B. T. McAllister, P. Altin, E. N. Ivanov, M. Goryachev, and M. E. Tobar, "Near-quantum limited axion dark matter search with the ORGAN experiment around 26 μeV," [arXiv:2407.18586 [hep-ex]].
- [25] ADMX Collaboration, C. Bartram et al., "Axion Dark Matter eXperiment around 3.3 μeV with Dine-Fischler-Srednicki-Zhitnitsky Discovery Ability," [arXiv:2408.15227 [hep-ex]].
- [26] C. Abel et al., "Search for Axionlike Dark Matter through Nuclear Spin Precession in Electric and Magnetic Fields," Phys. Rev. X 7 no. 4, (2017) 041034, [arXiv:1708.06367 [hep-ph]].
- [27] T. Wu et al., "Search for Axionlike Dark Matter with a Liquid-State Nuclear Spin Comagnetometer," Phys. Rev. Lett. 122 no. 19, (2019) 191302, [arXiv:1901.10843 [hep-ex]].
- [28] A. Garcon et al., "Constraints on bosonic dark matter from ultralow-field nuclear magnetic resonance," Sci. Adv. 5 no. 10, (2019) eaax4539, [arXiv:1902.04644 [hepex]].
- [29] I. M. Bloch, Y. Hochberg, E. Kuflik, and T. Volansky, "Axion-like Relics: New Constraints from Old Comagnetometer Data," JHEP 01 (2020) 167, [arXiv:1907.03767 [hep-ph]].
- [30] M. Jiang, H. Su, A. Garcon, X. Peng, and D. Budker, "Search for axion-like dark matter with spin-based amplifiers," Nature Phys. 17 no. 12, (2021) 1402–1407, [arXiv:2102.01448 [hep-ph]].
- [31] NASDUCK Collaboration, I. M. Bloch, G. Ronen, R. Shaham, O. Katz, T. Volansky, and O. Katz, "New constraints on axion-like dark matter using a Floquet quantum detector," Sci. Adv. 8 no. 5, (2022) abl8919, [arXiv:2105.04603 [hep-ph]].
- [32] J. Lee, M. Lisanti, W. A. Terrano, and M. Romalis, "Laboratory Constraints on the Neutron-Spin Coupling of feV-Scale Axions," Phys. Rev. X 13 no. 1, (2023) 011050, [arXiv:2209.03289 [hep-ph]].
- [33] NASDUCK Collaboration, I. M. Bloch, R. Shaham, Y. Hochberg, E. Kuflik, T. Volansky, and O. Katz, "Constraints on axion-like dark matter from a SERF comagnetometer," Nature Commun. 14 no. 1, (2023) 5784, [arXiv:2209.13588 [hep-ph]].
- [34] C. Abel et al., "Search for ultralight axion dark matter in a side-band analysis of a ¹⁹⁹ Hg free-spin precession signal," SciPost Phys. 15 no. 2, (2023) 058, [arXiv:2212.02403 [nucl-ex]].
- [35] K. Wei et al., "Dark matter search with a strongly-coupled hybrid spin system," [arXiv:2306.08039 [hep-ph]].
- [36] Z. Xu et al., "Constraining ultralight dark matter through an accelerated resonant search," Commun. Phys. 7 no. 1, (2024) 226, [arXiv:2309.16600 [hep-ph]].
- [37] D. Gavilan-Martin et al., "Searching for dark matter with a 1000 km baseline interferometer," [arXiv:2408.02668 [hep-ph]].
- [38] D. Budker, P. W. Graham, M. Ledbetter, S. Rajendran, and A. Sushkov, "Proposal for a Cosmic Axion Spin Precession Experiment (CASPEr)," Phys. Rev. X 4 no. 2, (2014) 021030, [arXiv:1306.6089 [hep-ph]].
- [39] D. F. Jackson Kimball et al., "Overview of the Cosmic Axion Spin Precession Experiment (CASPEr)," Springer Proc. Phys. 245 (2020) 105–121, [arXiv:1711.08999

- [physics.ins-det]].
- [40] R. Garoby et al., "The European Spallation Source Design," Phys. Scripta 93 no. 1, (2018) 014001.
- [41] T. Mason, D. Abernathy, I. Anderson, J. Ankner, T. Egami, G. Ehlers, A. Ekkebus, G. Granroth, M. Hagen, K. Herwig, J. Hodges, C. Hoffmann, C. Horak, L. Horton, F. Klose, J. Larese, A. Mesecar, D. Myles, J. Neuefeind, M. Ohl, C. Tulk, X.-L. Wang, and J. Zhao, "The spallation neutron source in oak ridge: A powerful tool for materials research," Physica B: Condensed Matter 385-386 (2006) 955-960. https://www.sciencedirect.com/science/article/pii/S092145260601177X.
- [42] W. Sheng, F. Shou-Xian, F. Shi-Nian, L. Wei-Bin, O. Hua-Fu, Q. Qing, T. Jing-Yu, and W. Jie, "Introduction to the overall physics design of csns accelerators," Chinese Physics C 33 no. S2, (Jun, 2009) 1. https://dx.doi.org/10.1088/1674-1137/33/S2/001.
- [43] P. Fierlinger, M. Holl, D. Milstead, V. Santoro, W. M. Snow, and Y. V. Stadnik, "Proposal for a Ramsey Neutron-Beam Experiment to Search for Ultralight Axion Dark Matter at the European Spallation Source," Phys. Rev. Lett. 133 no. 18, (2024) 181001, [arXiv:2404.15521 [hep-ph]].
- [44] Particle Data Group Collaboration, S. Navas et al., "Review of particle physics," Phys. Rev. D 110 no. 3, (2024) 030001.
- [45] Y. V. Stadnik and V. V. Flambaum, "Axion-induced effects in atoms, molecules, and nuclei: Parity nonconservation, anapole moments, electric dipole moments, and spin-gravity and spin-axion momentum couplings," Phys. Rev. D 89 no. 4, (2014) 043522, [arXiv:1312.6667 [hep-ph]].
- [46] J. J. Sakurai and J. Napolitano, Modern Quantum Mechanics. Quantum physics, quantum information and quantum computation. Cambridge University Press, 10, 2020.
- [47] H. Shimizu, T. Oku, H. Sato, C. Otani, H. Kato, Y. Suda, H. Nakagawa, T. Kamiyama, Y. Kiyanagi, and T. Wakabayashi, "A magnetic neutron lens," Physica B: Condensed Matter 276-278 (2000) 63-64. https://www.sciencedirect.com/science/article/ pii/S0921452699013265.
- [48] V. Santoro et al., "The HIBEAM program: search for neutron oscillations at the ESS," [arXiv:2311.08326 [physics.ins-det]].
- [49] G. Vasilakis, J. M. Brown, T. W. Kornack, and M. V. Romalis, "Limits on new long range nuclear spin-dependent forces set with a K He-3 co-magnetometer," Phys. Rev. Lett. 103 (2009) 261801, [arXiv:0809.4700 [physics.atom-ph]].
- [50] E. G. Adelberger, B. R. Heckel, S. A. Hoedl, C. D. Hoyle, D. J. Kapner, and A. Upadhye, "Particle Physics Implications of a Recent Test of the Gravitational Inverse Square Law," Phys. Rev. Lett. 98 (2007) 131104, [arXiv:hep-ph/0611223].
- [51] A. Bhusal, N. Houston, and T. Li, "Searching for Solar Axions Using Data from the Sudbury Neutrino Observatory," Phys. Rev. Lett. 126 no. 9, (2021) 091601, [arXiv:2004.02733 [hep-ph]].
- [52] M. Buschmann, C. Dessert, J. W. Foster, A. J. Long, and B. R. Safdi, "Upper Limit on the QCD Axion Mass from Isolated Neutron Star Cooling," Phys. Rev. Lett. 128 no. 9, (2022) 091102, [arXiv:2111.09892 [hep-ph]].

- [53] C. A. Baker et al., "An Improved experimental limit on the electric dipole moment of the neutron," Phys. Rev. Lett. 97 (2006) 131801, [arXiv:hep-ex/0602020].
- Lett. 97 (2006) 131801, [arXiv:hep-ex/0602020].
 [54] M. Pospelov and A. Ritz, "Theta vacua, QCD sum rules, and the neutron electric dipole moment," Nucl. Phys. B

573 (2000) 177-200, [arXiv:hep-ph/9908508].