

MACHINE PROOFS FOR ADAMS DIFFERENTIALS AND EXTENSION PROBLEMS AMONG CW SPECTRA

WEINAN LIN, GUOZHEN WANG AND ZHOU LI XU

ABSTRACT. In this document, we describe the process of obtaining numerous Adams differentials and extensions using computational methods, as well as how to interpret the dataset uploaded to Zenodo [7]. Detailed proofs of the machine-generated results are also provided. The dataset includes information on 210 CW spectra, 624 maps, and 98 cofiber sequences. Leveraging these results, and with the addition of some ad hoc arguments derived through human insight, we successfully resolved the Last Kervaire Invariant Problem in dimension 126 [8].

1. INTRODUCTION

In this project, we first compute the Adams E_2 -pages and maps for many CW spectra using the computer program `./Adams`, developed by the first author [5]. All relevant spectra and maps have been uploaded to the Zenodo repository [7] and are described in detail in Section 2. Additionally, another program, `./ss`, also developed by the first author [5], computes Adams differentials and extensions. The resulting data are included in the Zenodo repository and described in Section 3.

The program `./ss` also generates proofs for all computed results, which are organized into a comprehensive resource we refer to as the Table of Proofs. This program partially employs the Generalized Leibniz Rule and the Generalized Mahowald Trick—two theorems established in the paper [8] addressing the Last Kervaire Invariant Problem. These theorems enable the program to derive new differentials and extensions from previously computed ones. Detailed instructions on how to interpret and navigate the Table of Proofs are provided in Section 4.

In Section 5, we present several charts and tables of machine-generated Adams differentials that are particularly relevant to the solution of the Last Kervaire Invariant Problem.

2. ADAMS E_2 DATA

2.1. Adams E_2 -pages. Tables 1–2 provide a list of the CW spectra considered in this work, along with the maximum internal degrees computed for the Adams E_2 -pages of the corresponding spectra.

For a CW spectrum X , the Adams E_2 -page data for X is available in the database file `X_AdamsSS.db` within `kervaire_database.zip`. Alternatively, the data can be accessed as three plain-text table files:

- `X_AdamsE2_generators.csv` (for generators),
- `X_AdamsE2_relations.csv` (for relations),
- `X_AdamsE2_basis.csv` (for \mathbb{F}_2 -basis).

CW spectrum	max(t)	CW spectrum	max(t)
S0	261	C2sigma	200
tmf	261	Cetasigma	200
C2	200	Ctheta5sq	200
Ceta	200	C4	200
Cnu	200	C8	200
Csigma	200	Cetasq	200
CW_2_eta	200	Cetacube	200
CW_eta_2	200	Ceta4	200
CW_eta_nu	200	Ceta5	200
CW_nu_eta	200	Ckappa	200
CW_sigma_nu	200	Cg	200
CW_nu_sigma	200	CW_2_V_eta	200
CW_2_eta_nu	200	CW_eta_V_nu	200
CW_nu_eta_2	200	CW_nu_V_sigma	200
CW_sigma_nu_eta	200	CW_2_V_nu	180
CW_eta_nu_sigma	200	CW_2_V_sigma	180
CW_sigma_nu_eta_2	200	CW_eta_V_sigma	179
CW_2_eta_nu_sigma	200	CW_2_A_eta	180
Csigmasq	200	CW_eta_A_nu	180
C2h4	200	CW_nu_A_sigma	180
DC2h4	198	CW_2_A_nu	180
CW_sigmasq_2_sigmasq	200	CW_2_A_sigma	180
Ctheta4	200	CW_eta_A_sigma	180
C2h5	200	CW_2_eta_V_eta	171
DC2h5	200	CW_eta_2_A_eta	171
CW_theta4_2_theta4	196	CW_eta_nu_V_nu	170
Ctheta5	200	CW_nu_eta_A_nu	170
C2h6	200	CW_nu_sigma_V_sigma	170
DC2h6	200	CW_sigma_nu_A_sigma	171
CW_theta5_2_theta5	200	Joker	200
CW_2_theta5_2_Eq_eta_theta5	200	CW_eta_2_eta_Eq_2_nu	200
C2_C2	180	CW_eta_2_eta_Eq_nu_2	200
Ceta_Ceta	160	CW_eta_nu_eta_Eq_nu_nu	200
Cnu_Cnu	180	CW_nu_sigma_nu_Eq_sigma_sigma	200
Csigma_Csigma	180	C2_Ceta	200
C2_Ceta_Cnu	180	Ceta_Cnu	198
CW_2_eta_theta5sq	180	Cnu_Csigma	200
CW_2sigma_sigma	180	C2_Cnu	180
CW_etasigma_2_eta_nu	180	C2_Csigma	180
CW_etasigma_2_eta_nu_sigma	180	Ceta_Csigma	180
CW_nu_eta_2_etasigma	180	RP3_6	199
CW_sigma_2sigma	180	Fphi4	180
CW_sigma_nu_eta_2_etasigma	180	Fphi6	180
CW_theta5sq_eta_2	180	Fphi8	177
C2nu	200	Fphi10	177

TABLE 1. CW spectra

CW spectrum	max(t)	CW spectrum	max(t)	CW spectrum	max(t)
Fphi12	177	RP1_86	140	RP59_256	140
Fphi14	176	RP1_88	140	RP61_256	140
Fphi16	175	RP1_90	140	RP63_256	140
Fphi	200	RP1_92	140	RP65_256	140
RP1_4	200	RP1_94	140	RP67_256	140
RP1_6	183	RP1_96	140	RP69_256	140
RP1_8	180	RP1_98	140	RP71_256	140
RP1_10	180	RP1_100	140	RP73_256	140
RP1_12	178	RP1_102	140	RP75_256	140
RP1_14	160	RP1_104	140	RP77_256	140
RP1_16	160	RP1_106	140	RP79_256	140
RP1_18	140	RP1_108	140	RP81_256	140
RP1_20	140	RP1_110	140	RP83_256	140
RP1_22	140	RP1_112	140	RP85_256	140
RP1_24	140	RP1_114	140	RP87_256	140
RP1_26	140	RP1_116	140	RP89_256	140
RP1_28	140	RP1_256	200	RP91_256	140
RP1_30	140	RP3_256	160	RP93_256	140
RP1_32	140	RP5_256	140	RP95_256	140
RP1_34	140	RP7_256	140	RP97_256	140
RP1_36	140	RP9_256	140	RP99_256	140
RP1_38	140	RP11_256	140	RP101_256	140
RP1_40	140	RP13_256	140	RP103_256	140
RP1_42	140	RP15_256	140	RP105_256	140
RP1_44	140	RP17_256	140	RP107_256	140
RP1_46	140	RP19_256	140	RP109_256	140
RP1_48	140	RP21_256	140	RP111_256	140
RP1_50	140	RP23_256	140	RP113_256	140
RP1_52	140	RP25_256	140	RP115_256	140
RP1_54	140	RP27_256	140	RP117_256	140
RP1_56	140	RP29_256	140	CP1_4	180
RP1_58	140	RP31_256	140	CP1_128	160
RP1_60	140	RP33_256	140	tmf_C2	261
RP1_62	140	RP35_256	140	tmf_Ceta	261
RP1_64	140	RP37_256	140	tmf_Cnu	261
RP1_66	140	RP39_256	140	tmf_C2_Ceta	200
RP1_68	140	RP41_256	140	tmf_C2_Cnu	200
RP1_70	140	RP43_256	140	tmf_Ceta_Cnu	200
RP1_72	140	RP45_256	140	tmf_C2_Ceta_Cnu	200
RP1_74	140	RP47_256	140	tmf_RP1_256	200
RP1_76	140	RP49_256	140		
RP1_78	140	RP51_256	140		
RP1_80	140	RP53_256	140		
RP1_82	140	RP55_256	140		
RP1_84	140	RP57_256	140		

TABLE 2. CW spectra

id	name	stem	s
0	[0]	0	0
1	(h_1[1])	2	1
2	(h_2^2[1])	7	2
3	(h_0^3 h_3[1])	8	4
4	(c_0[1])	9	3
5	((Ph_1)[1])	10	5

TABLE 3. C2_AdamsE2_generators.csv

rel	stem	s
0,1,0	0	1
0,1,1;1,2,0	2	2
2,1,1	5	2
0,1,2	7	3
1,1,2;4,1,0	8	3

TABLE 4. C2_AdamsE2_relations.csv

index	mon	stem	s	d2
0	0	0	0	
0	1,1,0	1	1	
0	1	2	1	
0	1,2,0	2	2	
0	2,1,0	3	1	

TABLE 5. C2_AdamsE2_basis.csv

These files are included in `kervaire_csv.rar`, available in the Zenodo repository. We illustrate these tables with examples. Tables 3–5 present the first few rows of the corresponding files for the spectrum C2 (the naming convention will be explained later).

Notation 2.1. In the table of generators, a row with $id = i$ is the i 'th generator of the E_2 -page of X . In this section we will denote this generator by x_i if X is a ring spectrum or v_i if X is a module over some ring spectrum (in our dataset there are only two ring spectra `S0` and `tmf`). The columns `stem`, `s` are the degrees of the generators. The column `name` is the latex name we give to the i 'th generator. However, currently not all generators have latex names. The column `name` might be sometimes vacant (filled with `[NULL]`).

Notation 2.2. In the table of relations, consider the column `rel` for relations. If X is a ring spectrum, then `0,1;1,2,5,4` denotes the relation

$$x_0^1 + x_1^2 x_5^4 = 0.$$

If X is a module over the ring spectra R , then `0,1,3;1,2,5,4,7` denotes the relation

$$x_0^1 v_3 + x_1^2 x_5^4 v_7 = 0$$

where v_i is the i 'th generator of \mathbf{X} and x_j is the j 'th generator of the ring spectrum \mathbf{R} . In summary, the semicolon can be parsed as the plus sign and the monomials are sequences of sub- and upper-scripts whose lengths are even if \mathbf{X} is a ring or odd if \mathbf{X} is a module. The columns `stem,s` are the degrees of the relations which can be determined by the degrees of the generators.

Notation 2.3. In the table of basis, the column `mon` is parsed in the same way as we parse `rel` for relations. The only difference is that each `mon` is a monomial so there should be no semicolon in the column `mon`. The columns `stem,s` are the degrees of the monomials. The column `index` is **important**. When there are multiple monomial basis elements in the same bidegree, they will be indexed by $0, 1, \dots, k$. We write

$$[i] \in \text{Ext}^{\mathbf{s}, \mathbf{stem+s}}(\mathbf{X})$$

to denote the monomial basis element with index i in $\text{Ext}^{\mathbf{s}, \mathbf{stem+s}}(\mathbf{X})$. In the Table of Proofs (see Section 4), the element will be written as

$$\boxed{\mathbf{X} (\mathbf{stem, s}) [i]}$$

in the `info` column. We can also write $[i_1, i_2, \dots]$ to denote the linear combination of monomials with indices i_1, i_2, \dots and $[]$ to denote the zero element. The `d2` column represents the d_2 differentials of the basis elements in the format $[i_1, i_2, \dots]$ we just described but in degree $(\mathbf{stem-1, s+2})$. If it is empty, it means that the d_2 differential is trivial. If it is filled with `[NULL]`, it means that the d_2 differential is not calculated yet by the program `./Adams` using the secondary Steenrod operations.

Next we explain our naming scheme for the above CW spectra. We chose to use plain text names for these objects because we want to use consistent names for computer codes, command-line arguments for `./Adams` and `./ss`, the website links of plots and the plain text machine proofs. Table 6 is a list of keywords we use in the name and the corresponding elements in $\pi_*(S^0)$ the stable homotopy groups of spheres.

Notation 2.4. Let $\mathbf{a}, \mathbf{b}, \mathbf{c}, \dots$ be keywords, a, b, c, \dots be the corresponding elements in $\pi_*(S^0)$ and $|a|, |b|, |c|$ be the topological degrees of a, b, c, \dots . The following examples demonstrate our naming for the CW spectra.

- S^0 is the sphere spectrum S^0 . \mathbf{tmf} is the topological modular form tmf .
- \mathbf{Ca} is the cofiber of a .
- $\mathbf{CW_a_b}$ is a three-cell complex such that the following two cofiber sequences exist.

$$\begin{aligned} \mathbf{Ca} &\rightarrow \mathbf{CW_a_b} \rightarrow \Sigma^{|a|+|b|+2} S^0 \\ S^0 &\rightarrow \mathbf{CW_a_b} \rightarrow \Sigma^{|a|+1} \mathbf{Cb} \end{aligned}$$

The existence of such complex corresponds to $a \cdot b = 0$ in $\pi_*(S^0)$.

- $\mathbf{CW_a_b_c}$ is a four-cell complex such that the following two cofiber sequences exist.

$$\begin{aligned} \mathbf{CW_a_b} &\rightarrow \mathbf{CW_a_b_c} \rightarrow \Sigma^{|a|+|b|+|c|+3} S^0 \\ \mathbf{Ca} &\rightarrow \mathbf{CW_a_b_c} \rightarrow \Sigma^{|a|+|b|+2} \mathbf{Cc} \end{aligned}$$

The existence of such complex corresponds to $0 \in \langle a, b, c \rangle$ in $\pi_*(S^0)$.

- Inductively we can define $\mathbf{CW_a_..._z}$ as long as it exists.

keyword	$\pi_*(S^0)$
eta	η
nu	ν
sigma	σ
2nu	2ν
2sigma	2σ
etasigma	$\eta\sigma$
sigmasq	σ^2
theta4	θ_4
theta5	θ_5
theta5sq	θ_5^2
etasq	η^2
etacube	η^3
eta4	η_4
eta5	η_5
kappa	κ
g	$\bar{\kappa}$

TABLE 6. Keywords in the name of CW spectra

- Ca_Cb is the smash product of Ca and Cb . tmf_X is the smash product of tmf and X .
- CW_a_V_b is a three-cell complex such that the following two cofiber sequences exist.

$$\begin{aligned} \text{Ca} &\rightarrow \text{CW_a_V_b} \xrightarrow{\text{top cell } \#1} \Sigma^{|b|+1} S^0 \\ \text{Cb} &\rightarrow \text{CW_a_V_b} \xrightarrow{\text{top cell } \#2} \Sigma^{|a|+1} S^0 \end{aligned}$$

- DX is a k -fold suspension of the Spanier–Whitehead dual of X such that the lowest dimension of the cells is 0.
- CW_a_A_b is equivalent to DCW_a_V_b .
- C2h4 is the same as CW_2_sigmasq . C2h5 is the same as CW_2_theta4 . C2h6 is the same as CW_2_theta5 .
- CW_a_b_V_c is a four-cell complex such that the following two cofiber sequences exist.

$$\begin{aligned} \text{CW_a_b} &\rightarrow \text{CW_a_b_V_c} \rightarrow \Sigma^{|c|+1} S^0 \\ \text{Cc} &\rightarrow \text{CW_a_b_V_c} \rightarrow \Sigma^{|a|+1} \text{Cb} \end{aligned}$$

- CW_a_b_c_Eq_d_e is a five-cell complex such that the following two cofiber sequences exist.

$$\text{CW_a_b_V_d} \rightarrow \text{CW_a_b_c_Eq_d_e} \rightarrow \Sigma^{|a|+|b|+|c|+3} S^0$$

$$\text{Ca} \rightarrow \text{CW_a_b_c_Eq_d_e} \rightarrow \Sigma^{\min\{|a|+|b|+2, |d|+1\}} \text{CW_c_A_e}$$

The existence of such complex corresponds to $d \cdot e \in \langle a, b, c \rangle$ in $\pi_*(S^0)$.

- Joker is equivalent to $\text{CW_2_eta_2_Eq_eta_eta}$.
- RPm_n is equivalent to the stunted real projective space $\mathbb{R}P_m^n$ for positive integers m, n .

- $\text{CP}_{\mathbf{m}}_{\mathbf{n}}$ is equivalent to the stunted complex projective space $\mathbb{C}P_m^n$ for positive integers \mathbf{m}, \mathbf{n} .
- $F\phi$ is the suspension of $F\phi$ the fiber of the Kahn-Priddy map $\phi : \mathbb{R}P_1^\infty \rightarrow S^0$ (or the cofiber of ϕ).
- $F\phi_{\mathbf{k}}$ is the suspension of the k -skeleton of $F\phi$ for an integers \mathbf{k} .

Remark 2.5. All the CW spectra we considered are connective spectra. The current programs only work well with connective spectra.

Remark 2.6. If there are two or more CW spectra with different homotopy types that satisfy the same naming condition, then the name represents all choices and we do not assume which one we pick. The current important input of the program is the existence of these cofiber sequences and the ones derived from these using the properties of triangulated categories. We will make assumptions on homotopy classes in the future when a new feature of the program is complete and it will enable the `./ss` program to rigorously determine if there exists at least one homotopy class that satisfies multiple constraints.

Most of the CW spectra we consider are known to exist in the literature. For example, `CW_theta5sq_eta_2` is a subcomplex of the quadratic construction of `CW_theta5_2 = DC2h6` constructed in [1] [3]. The only exception is

```
CW_sigma_nu_eta_2_etasigma
```

together with its dual and subquotients. This CW complex is a six-cell complex with one cell in each dimension $d \in \{0, 8, 12, 14, 15, 24\}$ and in cohomology the 8-cell supports an Sq^{16} -operation to the top cell. The following is the proof of its existence.

We start with its 15-skeleton

```
CW_sigma_nu_eta_2.
```

In fact, the 4-fold Toda bracket $\langle \sigma, \nu, \eta, 2 \rangle \subset \pi_{13} = 0$ is strictly defined and has zero indeterminacy – it is simply zero.

First for degree reason if

$$h_4[8] \in E_2^{1,15+1}(\text{CW_sigma_nu_eta_2})$$

is not a permanent cycle it must support a d_7 -differential hitting $Pd_0[0]$. However, `CW_sigma_nu_eta_2` is a skeleton of `tmf` and the image of $Pd_0[0]$ in `tmf` is $w_1 d_0$ which is a nontrivial permanent cycle. Hence $Pd_0[0]$ should not be hit by d_7 and $h_4[8]$ must be permanent. Consider the extension spectral sequence of the following cofiber sequence

$$\text{CW_sigma_nu_eta} \rightarrow \text{CW_sigma_nu_eta_2} \rightarrow \Sigma^{15} S^0.$$

For degree reason $h_1 h_3 \in E_2^{1,8+1}(S^0)$ must be hit by an extension from $h_4[8]$. Hence we conclude that the top two cells of the cofiber of $h_4[8]$ is either $C(\eta\sigma)$ or $C(\eta\sigma + \epsilon)$. To rule out $C(\eta\sigma + \epsilon)$ it is sufficient to prove that in sphere

$$\kappa = \langle \nu, \eta, 2, \epsilon \rangle$$

while

$$0 = \langle \nu, \eta, 2, \eta\sigma \rangle.$$

Both 4-fold Toda brackets have zero indeterminacy.

The first one is well-known and can also be read off directly from the module structure of the E_2 -page of `CW_nu_eta` because it comes from a Massey product

$$\langle h_2, h_1, h_0, c_0 \rangle = d_0.$$

For the second one, we only need to rule out κ for filtration reasons. We have

$$\eta\langle\nu, \eta, 2, \eta\sigma\rangle \subset \langle\langle\eta, \nu, \eta\rangle, 2, \eta\sigma\rangle = \langle\nu^2, 2, \eta\sigma\rangle \supset \langle\nu^2, 2, \eta\rangle\sigma = 0.$$

It then follows from the fact that the middle 3-fold Toda bracket has zero indeterminacy.

Remark 2.7. Alternatively, given that both 4-fold Toda brackets

$$\langle\sigma, \nu, \eta, 2\rangle, \langle\nu, \eta, 2, \eta\sigma\rangle$$

are strictly zero, the existence of `CW_sigma_nu_eta_2_etasigma` is equivalent to the fact that the 5-fold Toda bracket

$$\langle\sigma, \nu, \eta, 2, \eta\sigma\rangle$$

in π_{22} contains zero. We check that this is the case.

Observe that $\pi_{22} \cong \mathbb{Z}/2 \otimes \mathbb{Z}/2$, generated by $[h_2c_1]$ and $[Pd_0]$. We first note that the element h_2c_1 detects a homotopy class that lies in the indeterminacy of this 5-fold Toda bracket.

In fact, we have a Massey product on the E_2 -page

$$h_2c_1 = \langle h_3, h_2, h_1, h_2^3 \rangle,$$

which follows from $c_0 = \langle h_2, h_1, h_0, h_1^2 \rangle$ by applying Sq^0 and multiplying by h_2 . This implies that h_2c_1 detects a homotopy class in the 4-fold Toda bracket $\langle\sigma, \nu, \eta, \nu^3\rangle$, which lies in the indeterminacy.

It remains to check that $[Pd_0]$ is not contained in this 5-fold Toda bracket. Since $[Pd_0]$ is detected by `tmf`, we map this 5-fold Toda bracket to `tmf`. We know $\sigma = 0$ in `tmf`, therefore this 5-fold Toda bracket becomes $\langle 0, \nu, \eta, 2, 0 \rangle$ in `tmf`, which is contained in

$$\pi_{11} \cdot \pi_{11} + \pi_{13} \cdot \pi_9 + \langle \pi_{11}, \eta, \pi_9 \rangle,$$

and can be checked to be zero.

2.2. Maps between Adams E_2 -pages. Tables 7–17 give the list of maps between CW spectra we have considered and the maximum computed internal degrees of these maps. The maximum computed internal degree of a map $f : X \rightarrow \Sigma^k Y$ refers to the maximum T such that such that the induced map

$$f : E_2^{s,t}(X) \rightarrow E_2^{s+\text{AF}(f), t-k+\text{AF}(f)}(Y)$$

is computed in the range $t \leq T$.

For a map `X__Y` (we will explain the naming later), the data can be found in the database file `map_AdamsSS_X_to_Y.db` in `kervaire_database.zip` or in the plain-text table file `map_X_to_Y.csv`. (A few filenames created by a newer version of `./Adams` remove `to` from the names.) Table 18 lists some rows from `map_C2_to_C2h4.csv` to illustrate the following notation.

Notation 2.8. In the table of maps, there are only two columns `id`, `map`. The `map` column gives us the image of the `id`'th generator from $E_2^{*,*}(X)$ to $E_2^{*,*}(Y)$, which

map	max(t)
S0_tmf	261
C2_S0	200
Ceta_S0	200
Cnu_S0	200
Csigma_S0	200
C2_CW_2_eta	180
Ceta_CW_eta_2	180
Ceta_CW_eta_nu	180
Cnu_CW_nu_eta	180
Cnu_CW_nu_sigma	200
CW_nu_sigma_Csigma	199
CW_sigma_nu_Cnu	199
CW_2_eta_Ceta	200
CW_eta_2_C2	180
CW_eta_nu_Cnu	200
CW_nu_eta_Ceta	200
C2_Q_CW_nu_eta_2	194
Cnu_Q_CW_2_eta_nu	197
CW_2_eta_nu_CW_eta_nu	190
CW_2_eta_CW_2_eta_nu	200
CW_eta_2_Q_CW_nu_eta_2	200
CW_eta_nu_Q_CW_2_eta_nu	200
CW_nu_eta_2_CW_eta_2	200
CW_nu_eta_2_CW_nu_eta_2	187
C2_CW_sigma_nu_by_eta	179
Ceta_Csigma_by_nu	188
Csigma_Ceta_by_nu	194
Csigma_CW_2_eta_by_nu	193
CW_2_eta_nu_sigma_CW_eta_nu_sigma	200
CW_2_eta_nu_CW_2_eta_nu_sigma	200
CW_eta_2_Csigma_by_nu	188
CW_eta_nu_sigma_CW_nu_sigma	200
CW_eta_nu_sigma_S0_by_2	200
CW_eta_nu_CW_eta_nu_sigma	200
CW_nu_eta_2_S0_by_sigma	200
CW_nu_eta_S0_by_sigma	197
CW_nu_sigma_C2_by_eta	197
CW_nu_sigma_S0_by_eta	199
CW_sigma_nu_eta_2_CW_nu_eta_2	200
CW_sigma_nu_eta_CW_nu_eta	200
CW_sigma_nu_eta_CW_sigma_nu_eta_2	200
Ceta_Q_CW_2_eta	200
Ceta_Q_CW_nu_eta	200
Cnu_Q_CW_eta_nu	200
Cnu_Q_CW_sigma_nu	200

TABLE 7. Maps between CW spectra

map	max(t)
Csigma__CW_sigma_nu	190
Csigma__Q_CW_nu_sigma	200
CW_sigma_nu__CW_sigma_nu_eta	200
Csigma__Csigmasq	193
Csigmasq__S0	200
C2h4__Csigmasq	200
Csigmasq__DC2h4	196
Csigmasq__Csigma	200
Csigmasq__C2_by_sigmasq	184
Csigmasq__Q_C2h4	200
Csigmasq__S0_by_sigmasq	200
C2__C2h4	200
DC2h4__C2	193
C2h4__S0	200
C2__Q_DC2h4	200
CW_sigmasq_2_sigmasq__C2h4	188
DC2h4__CW_sigmasq_2_sigmasq	198
C2h4__S0_by_sigmasq	199
C2__C2h5	200
C2h5__Ctheta4	200
Ctheta4__DC2h5	200
DC2h5__C2	200
C2h5__S0	200
C2__Q_DC2h5	200
CW_theta4_2_theta4__C2h5	196
DC2h5__CW_theta4_2_theta4	196
C2h5__S0_by_theta4	200
Ctheta4__Q_C2h5	200
Ctheta4__S0	200
Ctheta4__C2_by_theta4	168
Ctheta4__S0_by_theta4	200
C2__C2h6	200
C2h6__Ctheta5	200
Ctheta5__DC2h6	200
DC2h6__C2	200
C2h6__S0	200
C2__Q_DC2h6	198
CW_theta5_2_theta5__C2h6	188
DC2h6__CW_theta5_2_theta5	188
C2h6__S0_by_theta5	198
Ctheta5__Q_C2h6	200
Ctheta5__S0	200
CW_2_V_eta__CW_2_theta5_2_Eq_eta_theta5	192
CW_2_theta5_2_Eq_eta_theta5__Ctheta5	170
Ceta__CW_2_theta5_2_Eq_eta_theta5	200

TABLE 8. Maps between CW spectra

map	max(t)
C2__Q_CW_2_theta5_2_Eq_eta_theta5	136
CW_2_theta5_2_Eq_eta_theta5_DC2h6	173
C2h6__CW_2_theta5_2_Eq_eta_theta5	198
Ctheta5__Q_CW_2_theta5_2_Eq_eta_theta5	198
DC2h6__Q_CW_2_theta5_2_Eq_eta_theta5	191
C2_C2__C2	180
C2_Ceta_Cnu__C2_Ceta	180
C2_Ceta_Cnu__C2_Cnu	180
C2_Ceta_Cnu__Ceta_Cnu	180
C2_Ceta_Cnu__tmf_C2_Ceta_Cnu	180
C2_Ceta__C2_Ceta_Cnu	180
C2_Ceta__tmf_C2_Ceta	180
C2_Cnu__C2_Ceta_Cnu	160
C2_Cnu__tmf_C2_Cnu	180
C2__C2_C2	180
CW_2_eta__CW_2_eta_theta5sq	160
CW_2_eta__Q_RP1_6	180
CW_sigma_nu_eta_2_etasigma__CW_nu_eta_2_etasigma	180
CW_theta5sq_eta_2__CW_eta_2	180
Ceta_Ceta__Ceta	157
Ceta_Cnu__C2_Ceta_Cnu	180
Ceta_Cnu__tmf_Ceta_Cnu	180
Ceta__Ceta_Ceta	160
Cnu_Cnu__Cnu	180
Cnu__Cnu_Cnu	180
Csigma_Csigma__CW_sigma_2sigma	180
Csigma_Csigma__Csigma	180
Csigma__CW_sigma_2sigma	180
Csigma__Csigma_Csigma	180
tmf_C2_Ceta_Cnu__tmf_C2_Ceta	200
tmf_C2_Ceta_Cnu__tmf_C2_Cnu	200
tmf_C2_Ceta_Cnu__tmf_Ceta_Cnu	200
tmf_C2_Ceta__tmf_C2_Ceta_Cnu	200
tmf_C2_Cnu__tmf_C2_Ceta_Cnu	200
tmf_Ceta_Cnu__tmf_C2_Ceta_Cnu	200
Ctheta5sq__C2_by_eta	197
C2nu__S0	200
C2sigma__S0	200
Cetasigma__S0	200
Ctheta5sq__S0	200
C4__S0	200
C8__S0	200
Cetacube__S0	200
Cetasq__S0	200
Ceta4__S0	200

TABLE 9. Maps between CW spectra

map	max(t)
Ceta5__S0	200
Cg__S0	200
Ckappa__S0	200
C2nu__CP1_4	178
C2sigma__CW_2sigma_sigma	160
CW_2_eta_theta5sq__Ctheta5sq	180
CW_nu_eta_2_etasigma_Cetasigma	180
CW_sigma_2sigma_C2sigma	180
Cetasigma__CW_etasigma_2_eta_nu	160
Ctheta5sq__CW_theta5sq_eta_2	180
C2_C2__CW_2_V_eta	160
C2_Cnu__C2_Ceta_Cnu	180
C2_Q_CW_2_A_eta	139
CW_2_A_eta__C2_C2	160
CW_eta_A_nu__Ceta_Ceta	160
CW_nu_A_sigma__Cnu_Cnu	160
Ceta_Ceta__CW_eta_V_nu	160
Ceta__Q_CW_2_A_eta	160
Ceta__Q_CW_eta_A_nu	160
Cnu_Cnu__CW_nu_V_sigma	160
Cnu__Q_CW_eta_A_nu	160
Cnu__Q_CW_nu_A_sigma	160
Csigma__Q_CW_nu_A_sigma	160
CW_2_V_eta__S1	160
CW_2_V_eta__S2	160
CW_eta_V_nu__S2	136
CW_eta_V_nu__S4	160
CW_nu_V_sigma__S4	160
CW_nu_V_sigma__S8	160
Ceta__Joker	193
Ceta__Q_Joker	198
CW_2_eta__Joker	200
CW_eta_2__Q_Joker	199
Joker__Ceta	200
Joker__CW_eta_2	200
C2_Q_Joker	197
CW_2_eta_V_eta__Joker	154
CW_eta_2_A_eta__Q_Joker	153
Joker__CW_eta_2_A_eta	154
CW_2_V_eta__Joker	192
Joker__CW_2_A_eta	182
CW_2_A_eta__Q_Joker	180
CW_sigma_nu_eta_2__tmf	200
C2_tmf_C2	200
Ceta__tmf_Ceta	200

TABLE 10. Maps between CW spectra

	max(t)
map	
Cnu__tmf_Cnu	200
tmf_C2__tmf	260
tmf_Ceta__tmf	259
tmf_Cnu__tmf	257
RP1_256__tmf_RP1_256	150
CW_nu_sigma_nu_Eq_sigma_sigma__CW_sigma_nu	190
C2_Ceta__C2	200
C2_Ceta__Ceta	200
Ceta_Cnu__Ceta	198
Ceta_Cnu__Cnu	198
Cnu_Csigma__Cnu	200
Cnu_Csigma__Csigma	200
Cnu__Cnu_Csigma	180
Csigma__Cnu_Csigma	180
C2__C2_Ceta	180
Ceta__C2_Ceta	180
Ceta__Ceta_Cnu	180
Cnu__Ceta_Cnu	180
Cnu__CW_eta_nu_eta_Eq_nu_nu	200
Cnu__Q_CW_eta_nu_eta_Eq_nu_nu	196
Csigma__CW_nu_sigma_nu_Eq_sigma_sigma	190
Csigma__Q_CW_nu_sigma_nu_Eq_sigma_sigma	192
CW_eta_nu_eta_Eq_nu_nu__Cnu	200
CW_eta_nu_eta_Eq_nu_nu__CW_nu_eta	199
CW_eta_nu__CW_eta_nu_eta_Eq_nu_nu	200
CW_nu_sigma_nu_Eq_sigma_sigma__Csigma	200
CW_nu_eta__Q_CW_eta_nu_eta_Eq_nu_nu	195
CW_nu_sigma__CW_nu_sigma_nu_Eq_sigma_sigma	200
CW_sigma_nu__Q_CW_nu_sigma_nu_Eq_sigma_sigma	194
C2__CW_eta_2_eta_Eq_2_nu	193
C2__Q_CW_eta_2_eta_Eq_nu_2	196
Cnu__Q_CW_eta_2_eta_Eq_2_nu	199
CW_2_eta__Q_CW_eta_2_eta_Eq_2_nu	198
CW_2_eta__Q_CW_eta_2_eta_Eq_nu_2	198
CW_eta_2_eta_Eq_2_nu__Cnu	193
CW_eta_2_eta_Eq_2_nu__CW_2_eta	193
CW_eta_2_eta_Eq_nu_2__C2	200
CW_eta_2_eta_Eq_nu_2__CW_2_eta	200
CW_eta_2__CW_eta_2_eta_Eq_2_nu	193
CW_eta_2__CW_eta_2_eta_Eq_nu_2	200
Cnu__CW_eta_2_eta_Eq_nu_2	200
C2__CW_2_V_eta	177
CW_2_V_eta__C2_Ceta	200
CW_2_V_eta__CW_eta_2_eta_Eq_2_nu	200
CW_eta_V_nu__CW_eta_2_eta_Eq_nu_2	200

TABLE 11. Maps between CW spectra

map	max(t)
CW_eta_V_nu__CW_eta_nu_eta_Eq_nu_nu	184
CW_eta_V_nu__Ceta_Cnu	186
CW_nu_V_sigma__CW_nu_sigma_nu_Eq_sigma_sigma	190
CW_nu_V_sigma__Cnu_Csigma	200
Ceta__CW_2_V_eta	200
Ceta__CW_eta_V_nu	186
Cnu__CW_eta_V_nu	200
Cnu__CW_nu_V_sigma	200
Csigma__CW_nu_V_sigma	200
C2__CW_2_V_nu	180
C2__CW_2_V_sigma	180
Ceta__CW_eta_V_sigma	179
Cnu__CW_2_V_nu	180
Csigma__CW_2_V_sigma	180
Csigma__CW_eta_V_sigma	179
C2_Cnu__CW_2_A_nu	180
C2_Csigma__CW_2_A_sigma	180
CW_2_A_nu__C2	180
CW_2_A_nu__Cnu	180
CW_2_A_nu__Q_C2_Cnu	180
CW_2_A_sigma__C2	180
CW_2_A_sigma__Csigma	180
CW_2_V_eta__CW_2_eta_V_eta	154
CW_2_eta__CW_2_eta_V_eta	154
CW_eta_2_A_eta__CW_2_A_eta	153
CW_eta_2_A_eta__CW_eta_2	153
CW_eta_V_nu__CW_eta_nu_V_nu	151
CW_eta_nu_V_nu__CW_eta_nu_eta_Eq_nu_nu	129
CW_eta_nu__CW_eta_nu_V_nu	151
CW_eta_nu_eta_Eq_nu_nu__CW_nu_eta_A_nu	153
CW_nu_V_sigma__CW_nu_sigma_V_sigma	156
CW_nu_eta_A_nu__CW_eta_A_nu	151
CW_nu_eta_A_nu__CW_nu_eta	151
CW_nu_eta_A_nu__Q_CW_eta_nu_eta_Eq_nu_nu	151
CW_nu_sigma_V_sigma__CW_nu_sigma_nu_Eq_sigma_sigma	156
CW_nu_sigma__CW_nu_sigma_V_sigma	156
CW_nu_sigma_nu_Eq_sigma_sigma__CW_sigma_nu_A_sigma	157
CW_sigma_nu_A_sigma__CW_nu_A_sigma	153
CW_sigma_nu_A_sigma__CW_sigma_nu	153
CW_sigma_nu_A_sigma__Q_CW_nu_sigma_nu_Eq_sigma_sigma	153
Ceta_Cnu__CW_eta_A_nu	179
Ceta_Csigma__CW_eta_A_sigma	176
Cnu_Csigma__CW_nu_A_sigma	182
CW_2_A_sigma__Q_C2_Csigma	180
CW_2_V_nu__C2_Cnu	180

TABLE 12. Maps between CW spectra

map	max(t)
CW_2_V_sigma__C2_Csigma	180
CW_eta_A_sigma__Ceta	180
CW_eta_A_sigma__Csigma	163
CW_eta_A_sigma__Q_Ceta_Csigma	158
CW_eta_V_sigma__Ceta_Csigma	158
C2_Ceta__CW_2_A_eta	181
CW_2_A_eta__C2	180
CW_2_A_eta__Ceta	180
CW_2_A_eta__Q_C2_Ceta	180
CW_2_A_eta__Q_CW_eta_2_eta_Eq_nu_2	180
CW_eta_2_eta_Eq_nu_2__CW_2_A_eta	183
Ceta__Q_CW_eta_2_eta_Eq_2_nu	197
Ceta__Q_CW_eta_2_eta_Eq_nu_2	197
Ceta__Q_CW_eta_nu_eta_Eq_nu_nu	194
Cnu__Q_CW_nu_sigma_nu_Eq_sigma_sigma	188
CW_eta_2_eta_Eq_2_nu__CW_eta_A_nu	181
CW_eta_A_nu__Ceta	180
CW_eta_A_nu__Cnu	180
CW_eta_A_nu__Q_CW_eta_2_eta_Eq_2_nu	180
CW_eta_A_nu__Q_CW_eta_nu_eta_Eq_nu_nu	180
CW_eta_A_nu__Q_Ceta_Cnu	180
CW_eta_nu_eta_Eq_nu_nu__CW_eta_A_nu	184
CW_nu_A_sigma__Cnu	180
CW_nu_A_sigma__Csigma	180
CW_nu_A_sigma__Q_CW_nu_sigma_nu_Eq_sigma_sigma	180
CW_nu_A_sigma__Q_Cnu_Csigma	180
CW_nu_sigma_nu_Eq_sigma_sigma__CW_nu_A_sigma	188
CW_eta_2__Fphi	199
CW_eta_2__Fphi4	180
Fphi4__Fphi6	180
Fphi6__Fphi8	155
Fphi8__Fphi10	177
Fphi10__Fphi12	175
Fphi12__Fphi14	174
Fphi14__Fphi16	172
Fphi4__RP1_4	180
Fphi6__RP1_6	180
Fphi8__RP1_8	177
Fphi10__RP1_10	177
Fphi12__RP1_12	175
Fphi14__RP1_14	161
Fphi16__RP1_16	161
RP5_8__RP3_4	173
CW_eta_2__RP1_6	180
RP1_6__CW_2_eta	180

TABLE 13. Maps between CW spectra

map	max(t)	map	max(t)
Fphi__RP1_256	196	RP1_54__RP1_56	140
RP1_256__S0	199	RP1_56__RP1_58	140
CP1_128__S0	160	RP1_58__RP1_60	140
RP1_256__CP1_128	160	RP1_60__RP1_62	140
CP1_2__CP1_4	160	RP1_62__RP1_64	140
CP1_4__CP3_4	160	RP1_64__RP1_66	140
CP1_4__CP1_128	160	RP1_66__RP1_68	140
RP1_4__CP1_2	200	RP1_68__RP1_70	140
RP1_8__CP1_4	180	RP1_70__RP1_72	140
RP1_256__RPm7_0	172	RP1_72__RP1_74	140
RP1_10__RP7_10	180	RP1_74__RP1_76	140
RP1_16__RP9_16	157	RP1_76__RP1_78	140
RP9_16__RP1_8	156	RP1_78__RP1_80	140
RP3_4__RP1_2	198	RP1_80__RP1_82	140
RP5_6__RP3_4	198	RP1_82__RP1_84	140
RP1_4__RP2_4	160	RP1_84__RP1_86	140
RP3_5__RP3_6	160	RP1_86__RP1_88	140
RP3_4__RP3_6	188	RP1_88__RP1_90	140
RP1_2__RP1_4	199	RP1_90__RP1_92	140
RP1_4__RP1_6	183	RP1_92__RP1_94	140
RP1_6__RP1_8	175	RP1_94__RP1_96	140
RP1_8__RP1_10	174	RP1_96__RP1_98	140
RP1_10__RP1_12	178	RP1_98__RP1_100	140
RP1_12__RP1_256	178	RP1_100__RP1_102	140
RP1_12__RP1_14	140	RP1_102__RP1_104	140
RP1_14__RP1_16	160	RP1_104__RP1_106	140
RP1_16__RP1_18	140	RP1_106__RP1_108	140
RP1_18__RP1_20	140	RP1_108__RP1_110	140
RP1_20__RP1_22	140	RP1_110__RP1_112	140
RP1_22__RP1_24	140	RP1_112__RP1_114	140
RP1_24__RP1_26	140	RP1_114__RP1_116	140
RP1_26__RP1_28	140	RP1_116__RP1_256	140
RP1_28__RP1_30	140	RP1_6__RP3_6	179
RP1_30__RP1_32	140	RP3_6__RP5_6	191
RP1_32__RP1_34	140	RP1_4__RP3_4	200
RP1_34__RP1_36	140	RP1_6__RP5_6	162
RP1_36__RP1_38	140	RP1_8__RP5_8	176
RP1_38__RP1_40	140	RP1_12__RP9_12	178
RP1_40__RP1_42	140	RP1_14__RP11_14	140
RP1_42__RP1_44	140	RP1_16__RP13_16	140
RP1_44__RP1_46	140	RP1_18__RP15_18	140
RP1_46__RP1_48	140	RP1_20__RP17_20	140
RP1_48__RP1_50	140	RP1_22__RP19_22	140
RP1_50__RP1_52	140	RP1_24__RP21_24	140
RP1_52__RP1_54	140	RP1_26__RP23_26	140

TABLE 14. Maps between CW spectra

map	max(t)	map	max(t)
RP1_28__RP25_28	140	RP1_256__RP3_256	160
RP1_30__RP27_30	140	RP3_256__RP5_256	140
RP1_32__RP29_32	140	RP5_256__RP7_256	140
RP1_34__RP31_34	140	RP7_256__RP9_256	140
RP1_36__RP33_36	140	RP9_256__RP11_256	140
RP1_38__RP35_38	140	RP11_256__RP13_256	140
RP1_40__RP37_40	140	RP13_256__RP15_256	140
RP1_42__RP39_42	140	RP15_256__RP17_256	140
RP1_44__RP41_44	140	RP17_256__RP19_256	140
RP1_46__RP43_46	140	RP19_256__RP21_256	140
RP1_48__RP45_48	140	RP21_256__RP23_256	140
RP1_50__RP47_50	140	RP23_256__RP25_256	140
RP1_52__RP49_52	140	RP25_256__RP27_256	140
RP1_54__RP51_54	140	RP27_256__RP29_256	140
RP1_56__RP53_56	140	RP29_256__RP31_256	140
RP1_58__RP55_58	140	RP31_256__RP33_256	140
RP1_60__RP57_60	140	RP33_256__RP35_256	140
RP1_62__RP59_62	140	RP35_256__RP37_256	140
RP1_64__RP61_64	140	RP37_256__RP39_256	140
RP1_66__RP63_66	136	RP39_256__RP41_256	140
RP1_68__RP65_68	136	RP41_256__RP43_256	140
RP1_70__RP67_70	136	RP43_256__RP45_256	140
RP1_72__RP69_72	137	RP45_256__RP47_256	140
RP1_74__RP71_74	137	RP47_256__RP49_256	140
RP1_76__RP73_76	137	RP49_256__RP51_256	140
RP1_78__RP75_78	137	RP51_256__RP53_256	140
RP1_80__RP77_80	137	RP53_256__RP55_256	140
RP1_82__RP79_82	140	RP55_256__RP57_256	140
RP1_84__RP81_84	140	RP57_256__RP59_256	140
RP1_86__RP83_86	140	RP59_256__RP61_256	140
RP1_88__RP85_88	140	RP61_256__RP63_256	140
RP1_90__RP87_90	140	RP63_256__RP65_256	140
RP1_92__RP89_92	140	RP65_256__RP67_256	140
RP1_94__RP91_94	140	RP67_256__RP69_256	140
RP1_96__RP93_96	140	RP69_256__RP71_256	140
RP1_98__RP95_98	140	RP71_256__RP73_256	140
RP1_100__RP97_100	140	RP73_256__RP75_256	140
RP1_102__RP99_102	140	RP75_256__RP77_256	140
RP1_104__RP101_104	140	RP77_256__RP79_256	140
RP1_106__RP103_106	140	RP79_256__RP81_256	140
RP1_108__RP105_108	140	RP81_256__RP83_256	140
RP1_110__RP107_110	140	RP83_256__RP85_256	140
RP1_112__RP109_112	140	RP85_256__RP87_256	140
RP1_114__RP111_114	140	RP87_256__RP89_256	140
RP1_116__RP113_116	140	RP89_256__RP91_256	140

TABLE 15. Maps between CW spectra

map	max(t)	map	max(t)
RP91_256__RP93_256	140	RP57_60__RP57_256	84
RP93_256__RP95_256	140	RP59_62__RP59_256	84
RP95_256__RP97_256	140	RP5_8__RP5_256	136
RP97_256__RP99_256	140	RP61_64__RP61_256	80
RP99_256__RP101_256	140	RP63_66__RP63_256	80
RP101_256__RP103_256	140	RP65_68__RP65_256	76
RP103_256__RP105_256	140	RP67_70__RP67_256	76
RP105_256__RP107_256	140	RP69_72__RP69_256	72
RP107_256__RP109_256	140	RP71_74__RP71_256	72
RP109_256__RP111_256	140	RP73_76__RP73_256	68
RP111_256__RP113_256	140	RP75_78__RP75_256	68
RP113_256__RP115_256	140	RP77_80__RP77_256	64
RP115_256__RP117_256	140	RP79_82__RP79_256	64
RP101_104__RP101_256	40	RP7_10__RP7_256	136
RP103_106__RP103_256	40	RP81_84__RP81_256	60
RP105_108__RP105_256	36	RP83_86__RP83_256	60
RP107_110__RP107_256	36	RP85_88__RP85_256	56
RP109_112__RP109_256	32	RP87_90__RP87_256	56
RP111_114__RP111_256	32	RP89_92__RP89_256	52
RP113_116__RP113_256	28	RP91_94__RP91_256	52
RP115_118__RP115_256	28	RP93_96__RP93_256	48
RP117_120__RP117_256	24	RP95_98__RP95_256	48
RP11_14__RP11_256	132	RP97_100__RP97_256	44
RP13_16__RP13_256	128	RP99_102__RP99_256	44
RP15_18__RP15_256	128	RP9_12__RP9_256	132
RP17_20__RP17_256	124	RP3_6__RP3_256	140
RP19_22__RP19_256	124	RP3_256__RP1_2	160
RP21_24__RP21_256	120	RP5_256__RP1_4	140
RP23_26__RP23_256	120	RP7_256__RP1_6	140
RP25_28__RP25_256	116	RP9_256__RP1_8	140
RP27_30__RP27_256	116	RP11_256__RP1_10	140
RP29_32__RP29_256	112	RP13_256__RP1_12	140
RP31_34__RP31_256	112	RP15_256__RP1_14	140
RP33_36__RP33_256	108	RP17_256__RP1_16	140
RP35_38__RP35_256	108	RP19_256__RP1_18	140
RP37_40__RP37_256	104	RP21_256__RP1_20	140
RP39_42__RP39_256	104	RP23_256__RP1_22	140
RP41_44__RP41_256	100	RP25_256__RP1_24	140
RP43_46__RP43_256	100	RP27_256__RP1_26	140
RP45_48__RP45_256	96	RP29_256__RP1_28	140
RP47_50__RP47_256	96	RP31_256__RP1_30	140
RP49_52__RP49_256	92	RP33_256__RP1_32	140
RP51_54__RP51_256	92	RP35_256__RP1_34	140
RP53_56__RP53_256	88	RP37_256__RP1_36	140
RP55_58__RP55_256	88	RP39_256__RP1_38	140

TABLE 16. Maps between CW spectra

map	max(t)
RP41_256__RP1_40	140
RP43_256__RP1_42	140
RP45_256__RP1_44	140
RP47_256__RP1_46	140
RP49_256__RP1_48	140
RP51_256__RP1_50	140
RP53_256__RP1_52	140
RP55_256__RP1_54	140
RP57_256__RP1_56	140
RP59_256__RP1_58	140
RP61_256__RP1_60	140
RP63_256__RP1_62	140
RP65_256__RP1_64	140
RP67_256__RP1_66	140
RP69_256__RP1_68	140
RP71_256__RP1_70	140
RP73_256__RP1_72	140
RP75_256__RP1_74	140
RP77_256__RP1_76	140
RP79_256__RP1_78	140
RP81_256__RP1_80	140
RP83_256__RP1_82	140
RP85_256__RP1_84	140
RP87_256__RP1_86	140
RP89_256__RP1_88	140
RP91_256__RP1_90	140
RP93_256__RP1_92	140
RP95_256__RP1_94	140
RP97_256__RP1_96	140
RP99_256__RP1_98	140
RP101_256__RP1_100	140
RP103_256__RP1_102	140
RP105_256__RP1_104	140
RP107_256__RP1_106	140
RP109_256__RP1_108	140
RP111_256__RP1_110	140
RP113_256__RP1_112	140
RP115_256__RP1_114	140
RP117_256__RP1_116	140

TABLE 17. Maps between CW spectra

is parsed in the same way as we parse `rel` in the table of relations. For example, we can read from Table 18 that the image of $v_{36} \in E_2^{*,*}(\text{C2})$ is

$$v_{63} + x_1 x_{48} v_0 \in E_2^{*,*}(\text{C2h4}).$$

id	map
34	61
35	62
36	63;1,1,48,1,0
37	66;2,1,53
38	67

TABLE 18. `map_C2_to_C2h4.csv`

We explain our naming scheme for these maps.

Notation 2.9. Let X, Y, Z be names of CW spectra. The following examples demonstrate our naming for the maps between CW spectra.

- If X has strictly fewer cells than Y and there exists a unique k and a unique inclusion such that X is a subcomplex of $\Sigma^k Y$, then X_Y denotes this subcomplex map $X \hookrightarrow \Sigma^k Y$.
- If X has strictly more cells than Y and there exists a unique k and a unique quotient such that $\Sigma^k Y$ is a quotient complex of X , then X_Y denotes this quotient map $X \rightarrow \Sigma^k Y$.
- If we have a cofiber sequence

$$X \rightarrow \Sigma^m Y \rightarrow \Sigma^n Z \rightarrow \Sigma X$$

whose first map is a subcomplex inclusion denoted by X_Y and whose second map is a quotient map denoted by Y_Z , then the third boundary map is denoted by Z_Q_Y .

- For key words a, b, c, d, e , the map $CW_d_e_CW_a_b_by_c$ is equivalent to

$$CW_d_e_Q_CW_a_b_c_d_e.$$

Similarly, if we have a map with name $X_Y_by_c$ it means that X has a unique bottom cell and Y has a unique top cell and the map attaches the bottom cell of X to the top cell of Y by c . In other words, these two cells form a subquotient of the cofiber of the map which is equivalent to Cc .

- For numbers n, l, p, q where $p < q = n - 1 < n < l$, the map $Rp_n_l_Rp_p_q$ is equivalent to $Rp_n_l_Q_Rp_p_l$. In other words, the following three maps form a cofiber sequence.

$$(Rp_q_Rp_l, Rp_l_Rp_n, Rp_n_Rp_p)$$

- In plain-text names, we use the letter m to denote the negative sign. For example, $RP1_256_RPm7_0$ corresponds to the map

$$RP_1^{256} \rightarrow \Sigma RP_{-7}^0.$$

- Note that $RP_0^n \equiv S^0 \vee RP_1^n$. We use $RP1_256_S0$ to denote the truncated Kahn-Priddy map, which is a composition of the following maps.

$$RP_1^{256} \rightarrow RP_0^{256} \rightarrow \Sigma RP_{-1}^{-1} = S^0.$$

Similarly, We use $CP1_128_S0$ to denote the composition of the following maps.

$$CP_1^{128} \rightarrow CP_0^{128} \rightarrow \Sigma CP_{-1}^{-1} = S^{-1}.$$

- $S0_tmf$ and X_tmf_X are the Hurewicz maps.

X	$\max(t)$
S0	176
C2	161
Ceta	159
Cnu	157
Csigma	152
C2h6	167
DC2h4	134
DC2h5	138
DC2h6	145
CW_2_eta	159
CW_2_eta_nu	149
CW_2_eta_nu_sigma	149
CW_2_eta_theta5sq	156
CW_eta_2	144
CW_eta_nu	149
CW_nu_eta	148
CW_nu_eta_2	149
Joker	135
RP1_4	135
RP1_6	128
RP1_8	125

TABLE 19. Computed d_2 differentials

All the maps except the Hurewicz maps $S0_tmf$ and X_tmf_X we considered are maps in the cofiber sequences generated by inclusions $X \subset Y$ of CW complexes, i.e., the homology maps they induce are injective, surjective or trivial.

2.3. Adams d_2 differentials. Table 19 demonstrates the range of d_2 differentials we have computed and used in the computer programs. The maximum internal degree refers to the maximum T such that all the d_2 differentials of $x \in E_2^{s,t \leq T}$ are known for the corresponding CW spectra.

We refer the reader to [4] and [9] for the algorithm that computes d_2 differentials using the secondary Steenrod operations encoded in a differential graded algebra. The first author also implemented this algorithm in the program `./Adams`. The d_2 data we computed is already contained the table of basis (see Notation 2.3).

3. ADAMS SPECTRAL SEQUENCES AND EXTENSION SPECTRAL SEQUENCES

In this section we describe how the computer program `./ss` organizes the data of the Adams spectral sequence and the extension spectral sequence and how to read the differentials and extensions generated by the computer program.

The program `./ss` organizes the Adams spectral sequence by the following.

$$0 = B_1^{s,t} \subseteq B_2^{s,t} \subseteq B_3^{s,t} \subseteq \cdots \subseteq Z_3^{s,t} \subseteq Z_2^{s,t} \subseteq E_2^{s,t}$$

	$h_0^2 x_{123,9}$	d_2^{-1}	$x_{124,9}$
	$h_5 x_{92,10}$	d_5^{-1}	$x_{124,6}$
11	$x_{123,11,2} + x_{123,11} + h_0 h_6 [B_4]$	d_7	$h_1 x_{121,17}$
	$x_{123,11} + h_0 h_6 [B_4]$	d_3	$h_0 x_{122,13}$
	$h_0 h_6 [B_4]$	d_2	$h_0^2 h_6 M d_0$
	$h_0 x_{123,9}$	d_2^{-1}	$x_{124,8}$
10	$x_{123,10} + h_6 [B_4]$	d_3^{-1}	$x_{124,7}$
	$h_6 [B_4]$	d_2	$h_0 h_6 M d_0$
9	$x_{123,9} + h_0 x_{123,8}$	d_{12}	?
	$h_0 x_{123,8}$	d_3	$h_0 x_{122,11} + h_0 h_6 M d_0$
8	$x_{123,8}$	d_3	$x_{122,11} + h_6 M d_0$

TABLE 20. $E_2^{*,*}(S^0)$ stem=123

where B_r is the subgroup of the image of d_r differentials and Z_r is the subgroup of d_r -cycles. We can interpret Adams differentials as isomorphisms

$$(3.1) \quad d_r : Z_{r-1}^{s,t}/Z_r^{s,t} \xrightarrow{\cong} B_r^{s+r,t+r-1}/B_{r-1}^{s+r,t+r-1}$$

and therefore we can also consider the inverse

$$(3.2) \quad d_r^{-1} : B_r^{s,t}/B_{r-1}^{s,t} \xrightarrow{\cong} Z_{r-1}^{s-r,t-r+1}/Z_r^{s-r,t-r+1}.$$

Table 20 is an illustration of how we store the spectral sequence in memory. In Table 20 we can directly see that $B_2^{11,123+11}(S^0) = B_3 = B_4$ is generated by $h_0^2 x_{123,9}$, $B_5^{11,123+11}(S^0) = B_6$ is generated by $h_0^2 x_{123,9}$ and $h_5 x_{92,10}$, and $Z_6^{11,123+11}(S^0) = Z_5 = Z_4 = Z_3$ is generated by $h_0^2 x_{123,9}$, $h_5 x_{92,10}$ and $x_{123,11,2} + x_{123,11} + h_0 h_6 [B_4]$, etc.

We write differentials in terms of elements in $E_2^{*,*}$. The value of $d_r^{\pm 1}$ should be viewed as a coset. By (3.1) and (3.2) we see that

- if we get $d_r(x) = y_1$ and $d_r(x) = y_2$ for two different reasons, then we have $y_1 - y_2 \in B_{r-1}$;
- if we get $d_r^{-1}(x) = z_1$ and $d_r^{-1}(x) = z_2$ for two different reasons, then we have $z_1 - z_2 \in Z_r$.

Consider a cofiber sequence

$$X \xrightarrow{f} Y \xrightarrow{g} Z.$$

The program `./ss` organizes the extension spectral sequence (see [8]) by the following.

$$B_\infty^{s,t} \subseteq {}^f B_0^{s,t} \subseteq {}^f B_1^{s,t} \subseteq \dots \subseteq {}^g Z_1^{s,t} \subseteq {}^g Z_0^{s,t} \subseteq Z_\infty^{s,t}.$$

where B_r is the subgroup of the image of d_r differentials and Z_r is the subgroup of d_r -cycles. We can interpret extensions as isomorphisms

$$(3.3) \quad d_r^f : {}^f Z_{r-1}^{s,t}(X)/{}^f Z_r^{s,t}(X) \xrightarrow{\cong} {}^f B_r^{s+r,t+r}(Y)/{}^f B_{r-1}^{s+r,t+r}(Y)$$

when $r > 0$ and

$$(3.4) \quad d_0^f : Z_\infty^{s,t}(X)/{}^f Z_0^{s,t}(X) \xrightarrow{\cong} {}^f B_0^{s+r,t+r}(Y)/B_\infty^{s+r,t+r}(Y)$$

when $r = 0$.

We write differentials in terms of elements in $E_2^{*,*}$. The value of $(d_r^f)^{\pm 1}$ should be viewed as a coset. By (3.3) and (3.4) we see that

stem	s	base	diff	level
15	1	0	0	9998
0	17	0	[NULL]	9000
11	6	0	[NULL]	9000
14	3	0	0	2
15	2	0	0	9997

TABLE 21. `S0_AdamsE2_ss.csv`

iC	stem	s	base	diff	level
1	87	42	0	1	1
1	127	3	0	[NULL]	9996
1	126	7	1	[NULL]	9998
1	118	19	0	1,3	10000
1	134	4	0	[NULL]	9996
1	118	21	0	0	10000

TABLE 22. `cofseq_S0_C2_S0.csv`

- if $r > 0$ and we get $d_r^f(x) = y_1$ and $d_r^f(x) = y_2$ for two different reasons, then we have $y_1 - y_2 \in {}^f B_{r-1}(Y)$;
- if $r = 0$ and we get $d_0^f(x) = y_1$ and $d_0^f(x) = y_2$ for two different reasons, then we have $y_1 - y_2 \in B_\infty(Y)$.
- we leave it to the reader to think about what happens to $(d_r^f)^{-1}$ in similar situations.

The program `./ss` can compute a lot differentials and extensions based on the data structure we describe above and various structures such as the Leibniz rule and naturality. The proofs are given in the Table of Proofs (Section 4). There are three manually added differentials: $d_5 h_0^{24} h_6 = h_0^2 P^6 d_0$ and $d_6 h_0^{55} h_7 = h_0^2 x_{126,60}$ in the Adams spectral sequence of S^0 (from the image of J), and $d_3 v_2^{16} = \beta^5 g$ in the Adams spectral sequence of tmf , derived from power operations (Bruner–Rognes [2]).

The Adams differentials for a spectrum X produced by the machine are available in the database `X_AdamsSS.db` within `kervaire_database.zip`. Alternatively, the data can be accessed in the plain-text table file `X_AdamsE2_ss.csv` within `kervaire_csv.rar`.

The extensions in a cofiber sequence `X__Y__Z` are available in the database `cofseq_X__Y__Z.db` within `kervaire_database.zip`. Alternatively, the data can be accessed in the plain-text table file `cofseq_X__Y__Z.csv` within `kervaire_csv.rar`.

For illustration, Table 21 is a table of some rows from the table of Adams spectral sequence `S0_AdamsE2_ss.csv` while Table 22 is a table of some rows from the table of extension spectral sequence `cofseq_S0_C2_S0.csv`.

Notation 3.5. In the table of Adams spectral sequence, columns `stem`, `s` refers to the degrees of the elements in the `base` column. The columns `base`, `diff` should be parsed as linear combinations of the monomial basis in the corresponding bidegree. If `level=r<5000`, the `base` column is hit by a d_r -differential from `diff`. If `level=10000-r>5000`, the `base` column supports a d_r -differential hitting `diff`. If

`level=9000`, it means that `base` is a permanent cycle. If `diff` is filled with `[NULL]`, it means that the value of `diff` is undetermined. The reader can see that we assume that all visible nontrivial differentials in the data are of length shorter than 1000.

Notation 3.6. In the table of extension spectral sequence corresponding to a cofiber sequence

$$X \xrightarrow{f} Y \xrightarrow{g} Z,$$

the column `iC=0,1,2` refers to `X,Y,Z` where `base` lives respectively. The other columns are similar to those in Notation 3.5. If `iC=1` and `level=r<5000`, then `base` is hit by an f -extension which jumps filtration by r . If `iC=1` and `level=10000-r>5000`, then `base` supports an g -extension which jumps filtration by r .

Remark 3.7. All the Adams differentials and extensions can be also visually accessed from the website [6].

4. PROOFS

In this section we explain the structure and notations in our machine generated proofs. We call it the Table of Proofs. The Table of Proofs is provided in two different forms in the Zenodo repo: a single sqlite3 database file `proofs.db` and 22 plain-text tables `proofs-part1.csv`–`proofs-part22.csv`. The plain-text table is divided into 22 table files because there are more than 21 million rows in the table and some table viewers like Microsoft Excel has a row limit of one million. We encourage the reader to use the open source software DB Browser for SQLite to access the table in `proofs.db`. This tool can help you quickly search for what you want in big tables. You can also execute advanced SQL queries in the software.

The following are two typical rows in the Table of Proofs.

id	depth	reason	name	stem	s	t	r	x	dx	info
450592	0	D	Ceta	107	16	123	3	0,2	0,1	NULL
4378830	0	D	Cnu_CW_nu_eta_2_C2:1	15	2	17	2	0	0	NULL

TABLE 23. Two rows from the Table of Proofs

The first column `id` refers to the line number in the table. The columns `depth` and `reason` refer to the structure of the proof which we will explain in more details later. The column `name` refers to the name of a CW spectra or a map in a cofiber sequence.

- If `name` is the name of a CW spectrum like the first row, then the columns `stem,s,r,x,dx` refer to an Adams differential of the spectrum. In our example, this is

$$d_3([0, 2]) = [0, 1]$$

for

$$[0, 2] \in E_2^{16, 107+16}(\text{Ceta})$$

where $[0, 2]$ denotes the linear combination of the 0th and 2nd basis elements in $E_2^{16, 107+16}$ in the basis table of `Ceta` and $[0, 1]$ denotes the linear combination in the bidegree that this differential hits.

- If `name` is the name of a map in a cofiber sequence, it is of the form `X__Y__Z:n` where

$$X \xrightarrow{f} \Sigma^? Y \xrightarrow{g} \Sigma^? Z \xrightarrow{h} \Sigma X$$

is a cofiber sequence and $n = 0, 1, 2$ correspond to f, g, h respectively. The columns `stem, s, r, x, dx` refer to an extension of the map. In our example (second row of Table 23), consider the cofiber sequence

$$Cnu \xrightarrow{f} CW_nu_eta_2 \xrightarrow{g} \Sigma^6 C2 \xrightarrow{h} \Sigma Cnu.$$

The second row of the table indicates that there is a g -extension from the permanent cycle

$$[0] \in E_2^{2,15+2}(CW_nu_eta_2)$$

to the permanent cycle

$$[0] \in E_2^{4,15+4-6}(C2)$$

which jumps filtration by $r = 2$.

Next we explain the structure of the Table of Proofs. A typical proof of an Adams differential has the following form.

Proof Structure 4.1. Assume that the value of an Adams differential $d_r(x)$ has four candidates y_1, y_2, y_3, y_4 . (Usually the number of candidates is a power of two.)

- If $d_r(x) = y_1$, propagate (by the Leibniz rule, naturality, ...). Get a contradiction.
- If $d_r(x) = y_2$, propagate. Get a contradiction.
- If $d_r(x) = y_3$, propagate. Get a contradiction.

Then we can conclude that $d_r(x) = y_4$.

The following 4 rows from the Table of Proofs is a concrete form of Proof Structure 4.1.

id	depth	reason	name	stem	s	t	r	x	dx	info
504950	1	T	Csigma	116	13	129	4	1	3	proof1
504951	1	T	Csigma	116	13	129	4	1	0	proof2
504952	1	T	Csigma	116	13	129	4	1	0,3	proof3
504953	0	D	Csigma	116	13	129	4	1		NULL

Here the `depth` column can be thought as the nesting level of the latex `itemize` environment. Rows with `reason='T'` correspond to assumptions ('T' is a symbol for 'Try'). Each assumption row has a multi-line proof in the `info` column which explains why the assumption leads to a contradiction. For example, the actual content of `proof1` is the following.

```
Get Csigma (116,13) d_4[1]=[3]. Apply the Leibniz rule with
S0 (20,4) d_4[0]=[] and get Csigma (136,17) d_4[]=[3].
However, Csigma (135,21) [3] is not in B_3.
```

In `proof1` the element `Csigma (135,21) [3]` refers to the element

$$[3] \in E_2^{21,135+21}(Csigma).$$

Rows with **reason**=‘D’ correspond to the conclusions after we rule out enough candidates (‘T’ rows) of the value of Adams differentials or extensions (‘D’ is a symbol for ‘Deduction’).

Microsoft Excel tip: Since the text in the **info** column usually contains multiple lines, users who view the **csv** files with Excel should adjust the height of the edit box indicated by the following screenshot in order to view the whole content of the text.

A	B	C	D	E	F	G	H	I	J	K	L	M
477829	477828	1 T	Cnu	165	24	189	2	3 2,4,7	Get 'Cnu			
477830	477829	1 T	Cnu	165	24	189	2	3 2,5,6,7	Get 'Cnu			
477831	477830	1 T	Cnu	165	24	189	2	3 2,4,5,7	Get 'Cnu			
477832	477831	1 T	Cnu	165	24	189	2	3 2 3 7	Get 'Cnu			

Next we give examples of more complex proofs of Adams differentials.

Proof Structure 4.2. Assume that the value of an Adams differential $d_r(x)$ has four candidates $0, y_1, y_2, y_3$.

- If $d_r(x) = 0$, propagate. Assume that $d_{r+1}(x)$ has four possible values z_1, z_2, z_3, z_4 .
 - Assume $d_{r+1}(x) = z_1$. Propagate. Get a contradiction.
 - Assume $d_{r+1}(x) = z_2$. Propagate. Get a contradiction.
 - Assume $d_{r+1}(x) = z_3$. Propagate. Get a contradiction.
 - Assume $d_{r+1}(x) = z_4$. Propagate. Get a contradiction.
- If $d_r(x) = y_1$, propagate. Get a contradiction.
- If $d_r(x) = y_2$, propagate. Get a contradiction.

Then we can conclude that $d_r(x) = y_3$.

The following 8 rows from the Table of Proofs is an example of Proof Structure 4.2.

id	depth	reason	name	stem	s	t	r	x	dx	info
11406044	1	T	C2_Cnu	94	7	101	3	0		proof1
11406045	2	T	C2_Cnu	94	7	101	4	0		proof2
11406046	2	T	C2_Cnu	94	7	101	4	0	0	proof3
11406047	2	T	C2_Cnu	94	7	101	4	0	1	proof4
11406048	2	T	C2_Cnu	94	7	101	4	0	0,1	proof5
11406049	1	T	C2_Cnu	94	7	101	3	0	1	proof6
11406050	1	T	C2_Cnu	94	7	101	3	0	0,1	proof7
11406051	0	D	C2_Cnu	94	7	101	3	0	0	NULL

The reader can imagine that there could be proofs with even more complex structure. Consider the following.

Proof Structure 4.3. Assume that the value of an Adams differential $d_r(x)$ has four candidates $0, y_1, y_2, y_3$.

- If $d_r(x) = 0$, propagate. Assume that $d_{r+1}(x)$ has two possible values $z, 0$.
 - Assume $d_{r+1}(x) = z$. Propagate. Get a contradiction.

We get $d_{r+1}(x) = 0$. Assume that $d_{r+2}(x)$ has two possible values w_1, w_2 .

- Assume $d_{r+2}(x) = w_1$. Propagate. Get a contradiction.

- Assume $d_{r+2}(x) = w_2$. Propagate. Get a contradiction.
- If $d_r(x) = y_1$, propagate. Get a contradiction.
- If $d_r(x) = y_2$, propagate. Get a contradiction.

Then we can conclude that $d_r(x) = y_3$.

A proof with the above structure would be 8 rows from the Table of Proofs with (depth, reason)=(1,T), (2,T), (1,D), (2,T), (2,T), (1,T), (1,T), (0,D). If at some point in the proof we got a permanent cycle, we can even make assumptions about the value of extensions. The reader can find such examples in the Table of Proofs.

Next we describe other types of rows that can be derived from former differentials directly for different reasons.

- Rows with **reason**=‘d2’. This means that the d_2 differential is calculated by ./Adams using secondary Steenrod operation algorithm.
- Rows with **reason**=‘N’. This means that we get a differential from a previous one by naturality.
- Rows with **reason**=‘G’. This means that we get a trivial differential for degree reason.
- Rows with **reason**=‘XX’. This corresponds to the case that if we have $d_{r-1}(x) = 0$ then we automatically get $d_r(x^2) = 0$.
- Rows with **reason**=‘XY’. This corresponds to the case that if we have $d_{r-1}(x) = 0$ then no matter what $d_r(x)$ is (when there are two candidates or more), we always get the same value for $d_r(xy)$ by Leibniz rule or $d_r(f(x))$ by naturality.
- Rows with **reason**=‘ToCs’. This means that in the extension spectral sequence, we have tried all possible values of an extension (linear combinations of elements in the E_2 -page) and the outcome is that certain element in the Adams spectral sequence should survive to the E_∞ -page and it becomes a summand of the value of the extension.
- Rows with **reason**=‘OutCsI’. This means that in the extensions spectral sequence, two permanent cycles should be the same on the E_∞ -page. Hence the difference dx should be hit by an Adams differential. In those rows the columns **stem**, **s**, **t** refer to the degree of dx in stead of **x**.
- Rows with **reason**=‘CsCm’. These are extensions obtained from some simple homotopy relations. For example, if $x \in E_\infty(S^0)$ is the target of an η -extension, then it should detect a ν -torsion ($d_n^\nu(\eta) = 0$ for all n) because we know that $\eta \cdot \nu = 0 \in \pi_*(S^0)$. The reader can find more property of extensions in [8].
- Rows with **reason**=‘Syn’. This means that the differential is obtained by the Generalized Leibniz Rule from the former row. The map used in the Generalized Leibniz Rule and the conditions that the machine checked is written in the **info** column.
- Rows with **reason**=‘SynCs’. This means that the row is an extension and is obtained by the Generalized Mahowald Trick. The cofiber sequence used in the Generalized Mahowald Trick and the conditions that the machine checked is written in the **info** column.
- Rows with **reason**=‘SynIn’. This means that by the Generalized Mahowald Trick we determine that an element in the Adams spectral sequence should be a permanent cycle and become a summand of an extension.

- Rows with `reason='TI'`, `'DI'`, `'GI'`. These are essentially the same as rows with `reason='T'`, `'D'`, `'G'` except that the columns `stem`, `s`, `t` refer to the degree of dx in stead of x . For consecutive rows with `reason='TI'` the machine are enumerating possible sources of some differential (when we know dx is hit by some d_r differential beforehand) instead of the possible targets and the conclusion row is one with `reason='DI'`.

Remark 4.4. If the reader wants to search for a proof of certain differential, then usually you need to search for the bidegree of the source of the differential with `reason='D'` or `reason='T'`. Occasionally the machine knows the target should be hit by a certain differential first and in this case you need to search for the bidegree of the target of the differential with `reason='DI'` or `reason='TI'`.

5. CHARTS AND TABLES

In this section we list the Adams differentials of S^0 in the range $122 \leq stem \leq 127$, $0 \leq s \leq 25$ and Adams differentials of $C\nu$ in the range $stem = 126$, $9 \leq s \leq 14$. These differentials would play an important role in our solution of the Last Kervaire Invariant Problem [8].

s	Elements	d_r	value
25	$e_0 g^3 \Delta h_1 g$	d_2^{-1}	$g^2 \Delta^2 m$
	$d_0^3 g [B_4]$	d_4	$d_0^4 x_{65,13}$
24	$d_0 g \Delta^2 g^2$	d_2	$d_0 e_0 g^3 \Delta h_2^2$
23	$P h_1 x_{113,18,2}$	d_3^{-1}	$x_{123,20}$
	$h_0^2 d_0 x_{108,17}$	d_3	$d_0^4 M g$
22	$e_0 g^2 M g$	d_3^{-1}	$h_0^2 h_3 x_{116,16}$
	$x_{122,22}$	d_3	$h_0^2 d_0 x_{107,19}$
	$h_0 d_0 x_{108,17}$	d_2	$h_0^2 d_0 x_{107,18}$
21	$h_0^2 d_0 e_0 x_{91,11}$		Permanent
	$d_0 x_{108,17}$	d_2	$d_0 e_0 \Delta h_2^2 [B_4] + h_0 d_0 x_{107,18}$
20	$h_0^4 x_{122,16}$	d_2^{-1}	$h_0 h_3 x_{116,16}$
	$g^3 (C_0 + h_0^6 h_5^2)$	d_3	$g^3 \Delta h_2^2 n$
	$h_0 d_0 e_0 x_{91,11}$	d_2	$h_0^6 x_{121,16}$
19	$h_0^3 x_{122,16}$	d_2^{-1}	$h_3 x_{116,16}$
	$d_0 e_0 x_{91,11}$	d_2	$h_0 d_0^2 x_{93,12}$
18	$h_0^2 x_{122,16}$	d_3^{-1}	$h_0^2 x_{123,13,2}$
	$h_1 x_{121,17}$	d_7^{-1}	$x_{123,11,2} + x_{123,11} + h_0 h_6 [B_4]$
	$d_0 x_{108,14}$	d_3	$h_0 d_0^2 x_{93,12} + h_0^5 x_{121,16}$
17	$h_0 x_{122,16}$	d_3^{-1}	$h_0 x_{123,13,2}$
	$g^3 [H_1]$	d_3^{-1}	$\Delta h_2^2 x_{93,8}$
16	$x_{122,16} + h_0 x_{122,15,2}$	d_3^{-1}	$x_{123,13,2}$
	$\Delta h_2^2 x_{92,10}$	d_3^{-1}	$x_{123,13}$
	$h_0 x_{122,15,2}$		Permanent
15	$x_{122,15}$	d_3	$g^3 A$
	$x_{122,15,2}$	d_2	$h_0^2 h_4 x_{106,14}$
14	$h_0 x_{122,13}$	d_3^{-1}	$x_{123,11,2}$
13	$h_0^2 h_6 M d_0$	d_2^{-1}	$x_{123,11}$
	$h_1^2 x_{120,11}$		Permanent
	$x_{122,13}$	d_3	$h_0 h_4 x_{106,14}$
12	$h_0 h_6 M d_0$	d_2^{-1}	$x_{123,10}$
	$h_0 x_{122,11}$	d_3^{-1}	$h_0 x_{123,8}$
	$h_5 x_{91,11}$		Permanent
11	$x_{122,11} + h_6 M d_0$	d_3^{-1}	$x_{123,8}$
	$h_6 M d_0$		Permanent
9-10			
8	$h_1 x_{121,7}$	d_6	?
0-7			

TABLE 24. The classical Adams spectral sequence of S^0 for $s \leq 25$ in stem 122

s	Elements	d_r	value
25			
24	$h_1 Ph_1 x_{113,18,2}$	d_2^{-1}	$x_{124,22}$
	$d_0^2 \Delta h_2^2 Mg$	d_2^{-1}	$d_0 x_{110,18}$
	$d_0 M P x_{56,10}$	d_4	$M P x_{69,18}$
23	$g^2 \Delta^2 m$	d_2	$e_0 g^3 \Delta h_1 g$
21-22			
20	$x_{123,20}$	d_3	$Ph_1 x_{113,18,2}$
19	$e_0 g^2 x_{66,7} + h_0^6 x_{123,13,2}$	d_2^{-1}	$x_{124,17}$
	$h_0^6 x_{123,13,2}$	d_2^{-1}	$h_0^3 x_{124,14,2}$
	$h_0 e_0 x_{106,14} + h_0^2 h_3 x_{116,16}$	d_3^{-1}	$e_0 x_{107,12}$
	$h_0^2 h_3 x_{116,16}$	d_3	$e_0 g^2 M g$
18	$h_0^5 x_{123,13,2}$	d_2^{-1}	$h_0^2 x_{124,14,2}$
	$e_0 x_{106,14} + h_0 h_3 x_{116,16}$		Permanent
	$h_0 h_3 x_{116,16}$	d_2	$h_0^4 x_{122,16}$
17	$h_0^2 x_{123,15} + h_0^4 x_{123,13,2}$	d_2^{-1}	$h_0 x_{124,14}$
	$h_0^4 x_{123,13,2}$	d_2^{-1}	$h_0 x_{124,14,2} + h_0 x_{124,14}$
	$d_0 x_{109,13}$		Permanent
	$h_3 x_{116,16}$	d_2	$h_0^3 x_{122,16}$
16	$h_0 x_{123,15} + h_0^3 x_{123,13,2}$	d_2^{-1}	$x_{124,14}$
	$h_0^3 x_{123,13,2}$	d_2^{-1}	$x_{124,14,2} + x_{124,14}$
	$h_1 x_{122,15,2}$	d_3^{-1}	$h_4 x_{109,12}$
15	$x_{123,15}$	d_4^{-1}	$x_{124,11,2} + x_{124,11}$
	$h_4 x_{108,14}$	d_5^{-1}	$h_0 x_{124,9}$
	$h_0^2 x_{123,13,2}$	d_3	$h_0^2 x_{122,16}$
14	$h_0^3 x_{123,11}$	d_2^{-1}	$h_0 x_{124,11}$
	$h_0 x_{123,13,2}$	d_3	$h_0 x_{122,16}$
	$\Delta h_2^2 x_{93,8}$	d_3	$g^3 [H_1]$
13	$h_0^2 x_{123,11}$	d_2^{-1}	$x_{124,11}$
	$x_{123,13,2}$	d_3	$x_{122,16} + h_0 x_{122,15,2}$
	$x_{123,13}$	d_3	$\Delta h_2^2 x_{92,10}$
12	$h_0 x_{123,11} + h_0^2 h_6 [B_4]$	d_2^{-1}	$x_{124,9,2} + h_0 x_{124,8}$
	$x_{123,12}$	d_3^{-1}	$x_{124,9} + h_0 x_{124,8}$
	$h_0^2 h_6 [B_4]$	d_5^{-1}	$h_6 A$
11	$h_0^2 x_{123,9}$	d_2^{-1}	$x_{124,9}$
	$h_5 x_{92,10}$	d_5^{-1}	$x_{124,6}$
	$x_{123,11,2} + x_{123,11} + h_0 h_6 [B_4]$	d_7	$h_1 x_{121,17}$
	$x_{123,11} + h_0 h_6 [B_4]$	d_3	$h_0 x_{122,13}$
	$h_0 h_6 [B_4]$	d_2	$h_0^2 h_6 M d_0$
10	$h_0 x_{123,9}$	d_2^{-1}	$x_{124,8}$
	$x_{123,10} + h_6 [B_4]$	d_3^{-1}	$x_{124,7}$
	$h_6 [B_4]$	d_2	$h_0 h_6 M d_0$
9	$x_{123,9} + h_0 x_{123,8}$	d_{12}	?
	$h_0 x_{123,8}$	d_3	$h_0 x_{122,11} + h_0 h_6 M d_0$
8	$x_{123,8}$	d_3	$x_{122,11} + h_6 M d_0$
0-7			

TABLE 25. The classical Adams spectral sequence of S^0 for $s \leq 25$ in stem 123

s	Elements	d_r	value
25	$h_0^{11}x_{124,14,2}$	d_2^{-1}	$h_0^9x_{125,14}$
	$ix_{101,18}$	d_2	$d_0^2\Delta h_2^2x_{65,13} + h_0Pd_0x_{101,18}$
	$d_0e_0\Delta^3h_1g$	d_2	$d_0^2g^3m$
24	$h_0^{10}x_{124,14,2}$	d_2^{-1}	$h_0^8x_{125,14}$
	$h_0^2d_0x_{110,18}$	d_2^{-1}	$h_0e_0x_{108,17}$
23	$h_0^9x_{124,14,2}$	d_2^{-1}	$h_0^7x_{125,14}$
	$d_0g\Delta h_2^2[B_4] + h_0d_0x_{110,18}$	d_2^{-1}	$e_0x_{108,17}$
	$h_0d_0x_{110,18}$	d_3^{-1}	$x_{125,20}$
	$d_0x_{110,19}$		Permanent
22	$h_0^8x_{124,14,2}$	d_2^{-1}	$h_0^6x_{125,14}$
	$g^2\Delta^2t$	d_3^{-1}	$gx_{105,15}$
	$x_{124,22}$	d_2	$h_1Ph_1x_{113,18,2}$
	$d_0x_{110,18}$	d_2	$d_0^2\Delta h_2^2Mg$
21	$h_0^7x_{124,14,2}$	d_2^{-1}	$h_0^5x_{125,14}$
	$h_0d_0^2[\Delta\Delta_1g]$	d_2^{-1}	$d_0gx_{91,11}$
20	$h_0^6x_{124,14,2}$	d_2^{-1}	$h_0^4x_{125,14}$
	$d_0^2[\Delta\Delta_1g]$	d_5^{-1}	$h_1x_{124,14}$
19	$h_0^5x_{124,14,2}$	d_2^{-1}	$h_0^3x_{125,14}$
	$d_0x_{110,15}$		Permanent
18	$h_0x_{124,17} + h_0^4x_{124,14,2}$	d_2^{-1}	$x_{125,16}$
	$h_0^4x_{124,14,2}$	d_2^{-1}	$h_0^2x_{125,14}$
17	$h_0^3x_{124,14}$	d_2^{-1}	$x_{125,15}$
	$h_0^2x_{124,15}$	d_2^{-1}	$h_0x_{125,14}$
	$x_{124,17}$	d_2	$e_0g^2x_{66,7} + h_0^6x_{123,13,2}$
	$h_0^3x_{124,14,2}$	d_2	$h_0^6x_{123,13,2}$
16	$h_0x_{124,15}$	d_2^{-1}	$x_{125,14}$
	$h_1x_{123,15}$	d_3^{-1}	$h_3x_{118,12}$
	$h_0^2x_{124,14}$		Permanent
	$e_0x_{107,12}$	d_3	$e_0g^2x_{66,7} + h_0e_0x_{106,14} + h_0^2h_3x_{116,16}$
	$h_0^2x_{124,14,2}$	d_2	$h_0^5x_{123,13,2}$
15	$x_{124,15}$	d_4^{-1}	$h_6x_{62,10}$
	$h_3^2x_{110,13} + h_0x_{124,14}$		Permanent
	$h_0x_{124,14}$	d_2	$h_0^2x_{123,15} + h_0^4x_{123,13,2}$
	$h_0x_{124,14,2}$	d_2	$h_0^2x_{123,15}$
14	$h_1x_{123,13}$	d_2^{-1}	$x_{125,12}$
	$h_1x_{123,13,2}$	d_2^{-1}	$x_{125,12,2}$
	$\Delta h_2^2x_{94,8}$		Permanent
	$x_{124,14}$	d_2	$h_0x_{123,15} + h_0^3x_{123,13,2}$
	$x_{124,14,2}$	d_2	$h_0x_{123,15}$
13	$h_0^5x_{124,8}$	d_2^{-1}	$h_0^3x_{125,8}$
	$[H_1](\Delta e_1 + C_0 + h_0^6h_5^2)$	d_3^{-1}	$x_{125,10,2}$
	$e_0\Delta h_6g$		Permanent
	$h_4x_{109,12}$	d_3	$h_1x_{122,15,2}$

TABLE 26. The classical Adams spectral sequence of S^0 for $13 \leq s \leq 25$ in stem 124

	$h_0x_{124,11,2} + h_0x_{124,11}$	d_2^{-1}	$x_{125,10}$
12	$h_0^2x_{124,10,2} + h_0^4x_{124,8}$	d_2^{-1}	$h_0x_{125,9,2}$
	$h_0^4x_{124,8}$	d_2^{-1}	$h_0^2x_{125,8}$
	$h_1x_{123,11,2}$	d_3^{-1}	$x_{125,9}$
	$h_0x_{124,11}$	d_2	$h_0^3x_{123,11}$
	$h_0x_{124,10,2} + h_0^3x_{124,8}$	d_2^{-1}	$x_{125,9,2}$
11	$h_0^3x_{124,8}$	d_2^{-1}	$h_0x_{125,8}$
	$x_{124,11,3}$	d_3^{-1}	$x_{125,8,2}$
	$x_{124,11,2} + x_{124,11}$	d_4	$x_{123,15}$
	$x_{124,11}$	d_2	$h_0^2x_{123,11}$
10	$h_1x_{123,9} + h_0^2x_{124,8}$	d_2^{-1}	$x_{125,8}$
	$x_{124,10,2} + h_0x_{124,9}$	d_4^{-1}	$h_6[H_1]$
	$x_{124,10} + h_0^2x_{124,8}$	d_4^{-1}	$h_6[H_1] + h_0x_{125,5}$
	$h_0^2x_{124,8}$		Permanent
	$h_0x_{124,9}$	d_5	$h_4x_{108,14}$
9	$x_{124,9,2} + h_0x_{124,8}$	d_3	$h_0x_{123,11} + h_0^2h_6[B_4]$
	$x_{124,9} + h_0x_{124,8}$	d_3	$x_{123,12}$
	$h_0x_{124,8}$	d_2	$h_0^2x_{123,9}$
8	$x_{124,8}$	d_2	$h_0x_{123,9}$
7	$h_0x_{124,6}$	d_2^{-1}	$x_{125,5}$
	h_6A	d_5	$h_0^2h_6[B_4]$
	$x_{124,7}$	d_3	$x_{123,10} + h_6[B_4]$
6	$x_{124,6}$	d_5	$h_5x_{92,10}$
0-5			

TABLE 27. The classical Adams spectral sequence of S^0 for $s \leq 12$ in stem 124

s	Elements	d_r	value
25	$d_0^2e_0g[B_4]$	d_4^{-1}	$h_1x_{125,20}$
	$x_{125,25,2} + x_{125,25} + g^4\Delta h_1g$	d_4^{-1}	$x_{126,21} + \text{possibly } h_1x_{125,20}$
	$g^4\Delta h_1g$		Permanent
	$x_{125,25}$	d_2	$h_0x_{124,26}$
24	$e_0g\Delta^2g^2$	d_2	$d_0g^4\Delta h_2^2$
23	$h_0^2e_0x_{108,17}$	d_3	$d_0^3e_0Mg$
	$h_0^9x_{125,14}$	d_2	$h_0^{11}x_{124,14,2}$
22	g^3Mg	d_4^{-1}	$x_{126,18}$
	$ix_{102,15} + h_0^8x_{125,14}$	d_4^{-1}	$gx_{106,14} + e_0x_{109,14,2}$
	$h_0^8x_{125,14}$	d_2	$h_0^{10}x_{124,14,2}$
	$h_0e_0x_{108,17}$	d_2	$h_0^2d_0x_{110,18}$
21	$x_{125,21}$	d_4^{-1}	$d_0x_{112,13}$
	$h_0^7x_{125,14}$	d_2	$h_0^9x_{124,14,2}$
	$e_0x_{108,17}$	d_2	$d_0g\Delta h_2^2[B_4] + h_0d_0x_{110,18}$
20	$h_0d_0gx_{91,11}$	d_2^{-1}	$x_{126,18,2}$
	$x_{125,20}$	d_3	$d_0g\Delta h_2^2[B_4]$
	$h_0^6x_{125,14}$	d_2	$h_0^8x_{124,14,2}$

TABLE 28. The classical Adams spectral sequence of S^0 for $20 \leq s \leq 25$ in stem 125

	$gx_{105,15}$	d_3	$g^2\Delta^2t$
19	$h_0^3x_{125,14}$	d_2	$h_0'x_{124,14,2}$
	$d_0gx_{91,11}$	d_2	$h_0d_0^2[\Delta\Delta_1g]$
18	$d_0^2x_{97,10}$	d_5^{-1}	$h_1x_{125,12,2}$
	$h_0^4x_{125,14}$	d_2	$h_0^6x_{124,14,2}$
17	$h_0^2Q_2x_{68,8}$	d_2^{-1}	$h_0D_2x_{68,8}$
	$h_0^3x_{125,14}$	d_2	$h_0^5x_{124,14,2}$
16	$h_0Q_2x_{68,8}$	d_2^{-1}	$D_2x_{68,8}$
	$h_1x_{124,15}$	d_4^{-1}	$h_0^2x_{126,10}$
	$x_{125,16}$	d_2	$h_0x_{124,17} + h_0^4x_{124,14,2}$
	$h_0^2x_{125,14}$	d_2	$h_0^4x_{124,14,2}$
15	$h_0^3x_{125,12}$	d_2^{-1}	$h_0h_3x_{119,11}$
	$Q_2x_{68,8}$	d_4^{-1}	$h_1x_{125,10}$
	$h_1x_{124,14}$	d_5	$d_0^2[\Delta\Delta_1g]$
	$x_{125,15}$	d_2	$h_0^3x_{124,14}$
	$h_0x_{125,14}$	d_2	$h_0^2x_{124,15}$
14	$h_0^2x_{125,12}$	d_2^{-1}	$h_3x_{119,11}$
	$h_1h_4x_{109,12}$		Permanent
	$x_{125,14}$	d_2	$h_0x_{124,15}$
13	$h_0^4x_{125,9,2}$	d_2^{-1}	$x_{126,11}$
	$h_0^3x_{125,8}$	d_2^{-1}	$h_0x_{126,10}$
	$nx_{94,8}$	d_4^{-1}	$h_1x_{125,8}$
	$h_0x_{125,12}$	d_4^{-1}	$h_1x_{125,8,2}$
	$h_3x_{118,12}$	d_3	$h_1x_{123,15}$
12	$h_0h_6x_{62,10}$	d_2^{-1}	$x_{126,10}$
	$h_0^3x_{125,9,2} + h_0^4x_{125,8}$	d_3^{-1}	$x_{126,9}$
	$h_0^4x_{125,8}$	d_3^{-1}	$x_{126,9} + h_0x_{126,8,3}$
	$x_{125,12}$	d_2	$h_1x_{123,13}$
	$x_{125,12,2}$	d_2	$h_1x_{123,13,2}$
11	$h_1x_{124,10,2}$	d_3^{-1}	$x_{126,8}$
	$h_1x_{124,10}$	d_3^{-1}	$x_{126,8,2}$
	$h_0^2x_{125,9,2}$	d_5	?
	$h_6x_{62,10}$	d_4	$x_{124,15}$
	$h_0^3x_{125,8}$	d_2	$h_0^5x_{124,8}$
10	$h_0x_{125,9}$	d_2^{-1}	$x_{126,8,3}$
	$x_{125,10,2}$	d_3	$[H_1](\Delta e_1 + C_0 + h_0^6h_5^2)$
	$x_{125,10}$	d_2	$h_0x_{124,11,2} + h_0x_{124,11}$
	$h_0x_{125,9,2}$	d_2	$h_0^2x_{124,10,2} + h_0^4x_{124,8}$
	$h_0^2x_{125,8}$	d_2	$h_0^4x_{124,8}$
9	$h_6(\Delta e_1 + C_0 + h_0^6h_5^2)$		Permanent
	$h_5x_{94,8}$	d_7	?
	$x_{125,9}$	d_3	$h_1x_{123,11,2}$
	$x_{125,9,2}$	d_2	$h_0x_{124,10,2} + h_0^3x_{124,8}$
	$h_0x_{125,8}$	d_2	$h_0^3x_{124,8}$
8	$x_{125,8,2}$	d_3	$x_{124,11,3}$
	$x_{125,8}$	d_2	$h_1x_{123,9} + h_0^2x_{124,8}$
7	$h_0^2x_{125,5}$	d_3^{-1}	$x_{126,4}$
6	$h_6[H_1]$	d_4	$x_{124,10,2} + h_0x_{124,9}$
	$h_0x_{125,5}$	d_4	$x_{124,10,2} + x_{124,10} + h_0x_{124,9} + h_0^2x_{124,8}$
5	$x_{125,5}$	d_2	$h_0x_{124,6}$
0-4			

TABLE 29. The classical Adams spectral sequence of S^0 for $s \leq 19$ in stem 125

s	Elements	d_r	value
25	$h_0^7 x_{126,18}$	d_3^{-1}	$h_0^{21} h_7$
24	$d_0 e_0 \Delta h_2^2 M g$	d_2^{-1}	$d_0 x_{113,18}$
	$h_0^6 x_{126,18}$	d_3^{-1}	$h_0^{20} h_7$
	$g^4 \Delta h_2 c_1$	d_3^{-1}	$g^3 C''$
	$d_0 P d_0 M^2$	d_4^{-1}	$d_0 e_0 [\Delta \Delta_1 g]$
23	$h_0^5 x_{126,18}$	d_3^{-1}	$h_0^{19} h_7$
	$x_{126,23}$	d_4^{-1}	$e_0 x_{110,15}$
22	$h_0 x_{126,21} + h_0^4 x_{126,18}$	d_3^{-1}	$h_1 x_{126,18,2}$
	$h_0^4 x_{126,18}$	d_3^{-1}	$h_0^{18} h_7$
21	$h_0^3 x_{126,18}$	d_3^{-1}	$h_0^{17} h_7$
	$h_1 x_{125,20}$	d_4	$d_0^2 e_0 g [B_4]$
	$x_{126,21}$	d_4	$x_{125,25,2} + x_{125,25} + g^4 \Delta h_1 g + \text{possibly } d_0^2 e_0 g B_4$
20	$h_0^2 x_{126,18}$	d_3^{-1}	$h_0^{16} h_7$
	$d_0 x_{112,16}$	d_5^{-1}	$x_{127,15}$
19	$h_0 x_{126,18}$	d_3^{-1}	$h_0^{15} h_7$
	$g^3 x_{66,7}$	d_3^{-1}	$g x_{107,12}$
18	$x_{126,18} + e_0 x_{109,14,2}$	d_7	?
	$e_0 x_{109,14,2}$	d_4	$g^3 M g$
	$g x_{106,14}$	d_4	$i x_{102,15} + g^3 M g + h_0^8 x_{125,14}$
	$x_{126,18,2}$	d_2	$h_0 d_0 g x_{91,11}$
17	$h_0^{15} h_6^2$	d_2^{-1}	$h_0^{14} h_7$
	$h_1^2 x_{124,15}$	d_2^{-1}	$h_6 x_{64,14}$
	$x_{126,17}$	d_8	?
	$d_0 x_{112,13}$	d_4	$x_{125,21}$
16	$h_0^{14} h_6^2$	d_2^{-1}	$h_0^{13} h_7$
	$h_0^2 D_2 x_{68,8}$	d_3^{-1}	$x_{127,13}$
	$h_1^2 x_{124,14}$	d_6^{-1}	$h_2 x_{124,9} + h_0^2 x_{127,8}$
15	$h_0^{13} h_6^2$	d_2^{-1}	$h_0^{12} h_7$
	$h_0 D_2 x_{68,8}$	d_2	$h_0^2 Q_2 x_{68,8}$
14	$h_0^{12} h_6^2$	d_2^{-1}	$h_0^{11} h_7$
	$x_{126,14}$	d_4^{-1}	$h_2 x_{124,9}$
	$h_1 h_3 x_{118,12}$	d_5^{-1}	$h_1 x_{126,8,2}$
	$D_2 x_{68,8}$	d_2	$h_0 Q_2 x_{68,8}$
13	$h_0^{11} h_6^2$	d_2^{-1}	$h_0^{10} h_7$
	$h_1 x_{125,12,2}$	d_5	$d_0^2 x_{97,10}$
	$h_0 h_3 x_{119,11}$	d_2	$h_0^3 x_{125,12}$
12	$d_1 x_{94,8}$	d_2^{-1}	$x_{127,10}$
	$h_0 x_{126,11}$	d_2^{-1}	$h_3 x_{120,9}$
	$h_0^{10} h_6^2$	d_2^{-1}	$h_0^9 h_7$
	$h_0^2 x_{126,10}$	d_4	$h_1 x_{124,15}$
	$h_3 x_{119,11}$	d_2	$h_0^2 x_{125,12}$
11	$h_0^2 x_{126,9}$	d_2^{-1}	$h_0 x_{127,8}$
	$h_0^9 h_6^2$	d_2^{-1}	$h_0^8 h_7$
	$h_1 x_{125,10,2} + h_1 x_{125,10}$		Permanent
	$h_1 x_{125,10}$	d_4	$Q_2 x_{68,8}$
	$x_{126,11}$	d_2	$h_0^4 x_{125,9,2}$
	$h_0 x_{126,10}$	d_2	$h_0^5 x_{125,8}$

TABLE 30. The classical Adams spectral sequence of S^0 for $11 \leq s \leq 25$ in stem 126

10	$h_0x_{126,9}$	d_2^{-1}	$x_{127,8}$
	$h_0^2x_{126,8}$	d_2^{-1}	$h_0x_{127,7}$
	$h_0^8h_6^2$	d_2^{-1}	$h_0^7h_7$
	$h_0^2x_{126,8,3}$	d_3^{-1}	$x_{127,7,2} + x_{127,7}$
	$x_{126,10}$	d_2	$h_0h_6x_{62,10}$
9	$h_0x_{126,8}$	d_2^{-1}	$x_{127,7}$
	$h_0^7h_6^2$	d_2^{-1}	$h_0^6h_7$
	$h_1x_{125,8}$	d_4	$nx_{94,8}$
	$h_1x_{125,8,2}$	d_4	$h_0x_{125,12}$
	$x_{126,9}$	d_3	$h_0^3x_{125,9,2} + h_0^4x_{125,8}$
	$h_0x_{126,8,3}$	d_3	$h_0^3x_{125,9,2}$
8	$h_0^6h_6^2$	d_2^{-1}	$h_0^5h_7$
	$h_6(C' + X_2)$	d_{17}	?
	$x_{126,8,4} + x_{126,8}$	d_6	?
	$x_{126,8}$	d_3	$h_1x_{124,10,2}$
	$x_{126,8,2}$	d_3	$h_1x_{124,10}$
	$x_{126,8,3}$	d_2	$h_0x_{125,9}$
7	$h_0^5h_6^2$	d_2^{-1}	$h_0^4h_7$
	$h_1h_6[H_1]$	d_{18}	?
6	$h_0^4h_6^2$	d_2^{-1}	$h_0^3h_7$
	$x_{126,6}$	d_3	$h_5x_{94,8} + \text{possibly } h_6(\Delta e_1 + C_0 + h_0^6h_5^2)$
5	$h_0^3h_6^2$	d_2^{-1}	$h_0^2h_7$
	$x_{126,4}$	d_3	$h_0^2x_{125,5}$
4	$h_0^2h_6^2$	d_2^{-1}	h_0h_7
	$x_{126,4}$	d_3	$h_0^2x_{125,5}$
3	$h_0h_6^2$	d_2^{-1}	h_7
	h_6^2	d_7	?
0-1			

TABLE 31. The classical Adams spectral sequence of S^0 for $s \leq 10$ in stem 126

s	Elements	d_r	value
25	$h_0^{24}h_7$	d_3	$h_0^{10}x_{126,18}$
	$ix_{104,18}$	d_2	$d_0^3x_{84,15,2} + h_0d_0x_{112,22}$
	$d_0g\Delta^3h_1g$	d_2	$d_0e_0g^3m$
24	$h_0^2d_0x_{113,18}$	d_2^{-1}	$h_0gx_{108,17}$
	$h_1x_{126,23}$	d_3^{-1}	$x_{128,21}$
	$h_0^{23}h_7$	d_3	$h_0^9x_{126,18}$
23	$e_0g\Delta h_2^2[B_4] + h_0d_0x_{113,18}$	d_2^{-1}	$gx_{108,17}$
	$h_0d_0x_{113,18}$	d_4	$h_0d_0x_{112,22}$
	$d_0Pd_0x_{91,11}$	d_3	$h_1x_{125,25}$
	$h_0^{22}h_7$	d_3	$h_0^8x_{126,18}$
22	$d_0x_{113,18,2}$	d_4^{-1}	$d_0e_0x_{97,10}$
	$h_0^{21}h_7$	d_3	$h_0^7x_{126,18}$
	$d_0x_{113,18}$	d_2	$d_0e_0\Delta h_2^2Mg$
21	$h_0^6x_{127,15}$	d_2^{-1}	$h_0^5x_{128,14}$
	$x_{127,21} + g^3C''$		Permanent
	$h_0^{20}h_7$	d_3	$h_0^6x_{126,18}$
	g^3C''	d_3	$g^4\Delta h_2c_1$

TABLE 32. The classical Adams spectral sequence of S^0 for $21 \leq s \leq 25$ in stem 127

20	$h_0^5 x_{127,15}$	d_2^{-1}	$h_0^4 x_{128,14}$
	$d_0 e_0 [\Delta \Delta_1 g]$	d_4	$d_0 P d_0 M^2$
	$h_0^{19} h_7$	d_3	$h_0^5 x_{126,18}$
19	$h_0^4 x_{127,15}$	d_2^{-1}	$h_0^3 x_{128,14}$
	$h_1 x_{126,18}$		Permanent
	$e_0 x_{110,15}$	d_4	$x_{126,23}$
	$h_1 x_{126,18}, 2$	d_3	$h_0 x_{126,21} + h_0^4 x_{126,18}$
	$h_0^{18} h_7$	d_3	$h_0^4 x_{126,18}$
18	$h_0^3 x_{127,15}$	d_2^{-1}	$h_0^2 x_{128,14}$
	$h_0^3 h_6 x_{64,14}$	d_2^{-1}	$h_0^2 h_6 x_{65,13}$
	$g^2 \Delta h_1 H_1$	d_3^{-1}	$g x_{108,11}$
	$h_1 x_{126,17}$		Permanent
	$h_0^{17} h_7$	d_3	$h_0^3 x_{126,18}$
17	$h_0^2 h_2 x_{124,14}$	d_2^{-1}	$x_{128,15}$
	$h_0^2 x_{127,15}$	d_2^{-1}	$x_{128,15} + h_0 x_{128,14}$
	$h_0^2 h_6 x_{64,14}$	d_2^{-1}	$h_0 h_6 x_{65,13}$
	$g x_{107,13}$	d_4^{-1}	$h_0 h_3 x_{121,11}$
	$h_0^{16} h_7$	d_3	$h_0^2 x_{126,18}$
16	$h_0 x_{127,15} + h_0 h_2 x_{124,14}$	d_2^{-1}	$x_{128,14}$
	$h_0 h_6 x_{64,14}$	d_2^{-1}	$h_6 x_{65,13}$
	$h_0 h_2 x_{124,14}$		Permanent
	$x_{127,16}$		Permanent
	$h_0^{15} h_7$	d_3	$h_0 x_{126,18}$
	$g x_{107,12}$	d_3	$g^3 x_{66,7}$
15	$h_1 x_{126,14}$	d_2^{-1}	$x_{128,13,2}$
	$h_2 x_{124,14}$		Permanent
	$x_{127,15}$	d_5	$d_0 x_{112,16}$
	$h_0^{14} h_7$	d_2	$h_0^{15} h_6^2$
	$h_6 x_{64,14}$	d_2	$h_1^2 x_{124,15}$
14	$h_0 g \Delta h_6 g$	d_2^{-1}	$x_{128,12,2}$
	$h_0 h_3 x_{120,12}$	d_2^{-1}	$h_3 x_{121,11}$
	$h_0^{13} h_7$	d_2	$h_0^{14} h_6^2$
13	$h_0^3 x_{127,10}$	d_2^{-1}	$h_0 x_{128,10}$
	$g \Delta h_6 g$	d_3^{-1}	$x_{128,10,2}$
	$h_3 x_{120,12}$	d_4^{-1}	$h_2 x_{125,8,2}$
	$x_{127,13}$	d_3	$h_0^2 D_2 x_{68,8}$
	$h_0^{12} h_7$	d_2	$h_0^{13} h_6^2$
12	$h_0^2 x_{127,10}$	d_2^{-1}	$x_{128,10}$
	$h_1 x_{126,11}$	d_3^{-1}	$h_1 x_{127,8}$
	$h_0^{11} h_7$	d_2	$h_0^{12} h_6^2$
11	$h_0 h_3 x_{120,9}$	d_3^{-1}	$h_3 D_2 h_6$
	$h_0 h_2 x_{124,9}$		Permanent
	$h_0 x_{127,10}$		Permanent
	$h_0^{10} h_7$	d_2	$h_0^{11} h_6^2$
10	$h_1^2 x_{125,8}$		Permanent
	$h_2 x_{124,9} + h_0^2 x_{127,8}$	d_6	$h_1^2 x_{124,14}$
	$h_0^2 x_{127,8}$	d_4	$x_{126,14}$
	$x_{127,10}$	d_2	$d_1 x_{94,8}$
	$h_3 x_{120,9}$	d_2	$h_0 x_{126,11}$
	$h_0^9 h_7$	d_2	$h_0^{10} h_6^2$

TABLE 33. The classical Adams spectral sequence of S^0 for $10 \leq s \leq 20$ in stem 127

	$h_0^2 x_{127,7}$	d_2^{-1}	$h_0 x_{128,6}$
	$h_1 x_{126,8}$		Permanent
9	$h_1 x_{126,8,2}$	d_5	$h_1 h_3 x_{118,12}$
	$h_0 x_{127,8}$	d_2	$h_0^5 x_{126,9}$
	$h_0^8 h_7$	d_2	$h_0^9 h_6^2$
	$h_0 x_{127,7,2} + h_0 x_{127,7} + h_0^2 x_{127,6}$	d_2^{-1}	$x_{128,6}$
8	$h_0^2 x_{127,6}$	d_2^{-1}	$h_0 x_{128,5}$
	$h_2 h_6 A$		Permanent
	$h_2 x_{124,7}$	d_9	?
	$x_{127,8}$	d_2	$h_0 x_{126,9}$
	$h_0 x_{127,7}$	d_2	$h_0^7 x_{126,8}$
	$h_0^7 h_7$	d_2	$h_0^8 h_6^2$
7	$h_0 x_{127,6}$	d_2^{-1}	$x_{128,5}$
	$h_1 x_{126,6}$	d_{10}	?
	$x_{127,7,2} + x_{127,7}$	d_3	$h_0^2 x_{126,8,3}$
	$x_{127,7}$	d_2	$h_0 x_{126,8}$
	$h_0^6 h_7$	d_2	$h_0^6 h_6^2$
6	$x_{127,6}$	d_5	?
	$h_0^5 h_7$	d_2	$h_0^6 h_6^2$
5	$h_0^4 h_7$	d_2	$h_0^5 h_6^2$
4	$h_0^3 h_7$	d_2	$h_0^4 h_6^2$
3	$h_1 h_6^2$	d_8	?
	$h_0^2 h_7$	d_2	$h_0^3 h_6^2$
2	$h_0 h_7$	d_2	$h_0^5 h_6^2$
1	h_7	d_2	$h_0 h_6^2$
0			

TABLE 34. The classical Adams spectral sequence of S^0 for $s \leq 9$ in stem 127

s	Elements	d_r	value
14	$h_0^{12} h_6^2 [0]$	d_2^{-1}	$h_0^{11} h_7 [0]$
	$x_{126,14} [0]$	d_3^{-1}	$((x_{123,11,2}) + (x_{123,11}) + h_0 h_6 [B_4]) [4]$
	$Q_2 D_2 (h_3 [4]) + D_2 x_{68,8} [0]$	d_3^{-1}	$((x_{123,11,2}) + h_5 (x_{92,10})) [4]$
	$D_2 x_{68,8} [0]$	d_2	$h_0 Q_2 x_{68,8} [0]$
13	$h_0^{11} h_6^2 [0]$	d_2^{-1}	$h_0^{10} h_7 [0]$
	$h_1 x_{120,11} (h_1 [4])$	d_{12}	?
	$h_1 x_{125,12,2} [0]$	d_5	$d_5^2 x_{97,10} [0]$
	$h_6 x_{56,10} (h_0 h_2 [4])$	d_3	$h_1 x_{124,15} [0]$
	$((x_{122,13}) + h_1^2 (x_{120,11}) + h_0^2 h_6 (Md_0)) [4]$	d_2	$x_{125,15} [0] + h_0^3 x_{125,12} [0]$
12	$h_0 h_3 x_{119,11} [0]$	d_2	$h_0^3 x_{125,12} [0]$
	$d_1 x_{94,8} [0]$	d_2^{-1}	$x_{127,10} [0]$
	$h_0^{10} h_6^2 [0]$	d_2^{-1}	$h_0^9 h_7 [0]$
	$((h_5 (x_{91,11}) + h_0 (x_{122,11})) [4])$	d_3	$Q_2 x_{68,8} [0]$
11	$h_3 x_{119,11} [0]$	d_2	$h_0^2 x_{125,12} [0]$
	$h_0^9 h_6^2 [0]$	d_2^{-1}	$h_0^8 h_7 [0]$
	$x_{126,11} [0]$	d_3^{-1}	$x_{127,8} [0]$
	$h_1 x_{125,10,2} [0] + h_1 x_{125,10} [0]$		Permanent
10	$h_1 x_{125,10} [0]$	d_{14}	?
	$h_0^2 x_{126,8} [0]$	d_2^{-1}	$h_0 x_{127,7} [0]$
	$h_0^8 h_6^2 [0]$	d_2^{-1}	$h_0^7 h_7 [0]$
9	$x_{126,10} [0]$	d_3	$nx_{94,8} [0]$
	$h_0 x_{126,8} [0]$	d_2^{-1}	$x_{127,7} [0]$
	$h_0^6 h_6^2 [0]$	d_2^{-1}	$h_0^6 h_7 [0]$
	$h_1 x_{125,8} [0]$	d_{16}	?
	$h_0 x_{126,8,3} [0]$	d_4	$h_0 x_{125,12} [0]$
	$x_{126,9} [0]$	d_3	$h_0^4 x_{125,8} [0]$

TABLE 35. The classical Adams spectral sequence of S^0/ν for $9 \leq s \leq 14$ in stem 126

REFERENCES

- [1] M. G. Barratt, J. D. S. Jones, and M. E. Mahowald. The Kervaire invariant problem. In *Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982)*, volume 19 of *Contemp. Math.*, pages 9–22. Amer. Math. Soc., Providence, RI, 1983.
- [2] Robert R. Bruner and John Rognes. *The Adams spectral sequence for topological modular forms*, volume 253 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, [2021] ©2021.
- [3] Robert Burklund and Zhouli Xu. The Adams differentials on the classes h_j^3 . *To appear in Inventiones Mathematicae. arXiv.2302.11869.*, 2023.
- [4] Dexter Chua. The E_3 page of the Adams spectral sequence. *arXiv preprint, arXiv.2105.07628*, 2021.
- [5] Weinan Lin. Github repo. <https://github.com/WayneLin92/SSeqCpp> Github Webpage 2024.
- [6] Weinan Lin. Interactive plot of the Adams spectral sequence. <https://waynelin92.github.io/ss/kervaire.html> Github Webpage 2024.
- [7] Weinan Lin, Guozhen Wang, and Zhouli Xu. Machine proofs for Adams differentials and extension problems among CW spectra. Zenodo. <https://doi.org/10.5281/zenodo.14272279> December 2024.
- [8] Weinan Lin, Guozhen Wang, and Zhouli Xu. On the Last Kervaire Invariant Problem. *Preprint, 2024.*
- [9] Christian Nassau. On the secondary Steenrod algebra. *New York J. Math.*, 18:679–705, 2012.