Nonlinear optical spectroscopy of open quantum systems

Haoran Sun,¹ Upendra Harbola,² Shaul Mukamel,³ and Michael Galperin^{4, *}

¹Department of Chemistry & Biochemistry, University of California San Diego, La Jolla, CA 92093, USA

²Department of Inorganic and Physical Chemistry,

Indian Institute of Science, Bangalore 560012, India

³Department of Chemistry, University of California Irvine, Irvine, CA 92697, USA

⁴School of Chemistry, Tel Aviv University, Tel Aviv 69978, Israel

Development of experimental techniques at nanoscale resulted in ability to perform spectroscopic measurements on single-molecule current carrying junctions. These experiments are natural meeting point for research fields of optical spectroscopy and molecular electronics. We present a pedagogical comparison between perturbation theory expansion of standard nonlinear optical spectroscopy and (non-self-consistent) perturbative diagrammatic formulation of the nonequilibrium Green's functions method (NEGF is widely used in molecular electronics) indicating their similarities and differences. Comparing the two approaches we argue that optical spectroscopy of open quantum systems has to be analyzed within the more general Green's function.

I. INTRODUCTION

Interaction of matter with light is on the forefront of research due to its its ability to answer fundamental questions and its applicational promises. In particular, spectroscopy is an invaluable characterization tool in studies of properties of molecules and materials [1-7]. Also, lightmatter interaction allows to control electron and energy transfer, nuclear dynamics, and chemistry in molecular systems as was shown, for example, in recent measurements in systems with strong light-matter interaction. The latter opens directions for practical utilization of light as a tool for quantum technologies [8–10], transport [11–16] and energy transfer [17, 18] enhancement, light engineering [19, 20], and as a catalyst for chemical reactions [21–23]. Another novel direction is due to recent experimental developments in X-ray spectroscopies. X-ray light is capable to provide spatially localized and element specific information extending capabilities of established nonlinear spectroscopy techniques [24, 25]. Finally, quantum properties of light (e.g. photon entanglement) recently attracted attention of researchers as a tool which allows to circumvent many experimental shortcomings of spectroscopies relying on classical properties of light [26, 27].

Originally, spectroscopy experiments where done on molecules in gas phase and/or on surfaces. Theoretically, the former situation corresponds to isolated quantum system (molecule) whose dynamics is described within the time-dependent Schrödinger equation. The latter is formally an open system although the only effect of the bath (surface) on molecule was related to dissipation. Dynamics of such systems traditionally is described within the quantum master equation in its simplest, Redfield/Lindblad, form and light-matter interaction is taken into account employing bare perturbation theory (PT) expansions. This level of treatment is at the heart of nonlinear optical spectroscopy theory which is widely and successfully used for description of experiments and perturbation theory expansions define the very classification of optical processes [28–30].

Recent developments in experimental techniques made it possible to perform spectroscopic measurements in current carrying single-molecule junctions. For example, such are bias-induced electroluminescence [31–34] and Raman scattering [35–39]. Molecular cavity spectroscopy is also moving in this direction [40, 41]. Open character of these systems plays important role in its responses. Thus, considerations employing bare perturbation theory expansions in light-matter interaction and Markov (weak system-bath coupling) descriptions of system evolution are not always adequate [42, 43]. Similarly, widely employed in cavity spectroscopy non-Hermitian quantum mechanics considerations (i.e. theory taking into account dissipation and ignoring fluctuations) may lead to qualitative failures [44, 45].

Nonequilibrium Green's functions (NEGF) is the method adequately describing evolution of an open quantum mechanical system. It is the method of choice in molecular electronics studies which until recently were mostly focused on description of electron transport in junctions. Theoretical descriptions of spectroscopic measurements in single-molecule junctions resulted in application of methods from nonlinear optical spectroscopy and quantum transport communities, and naturally raised question of comparison between them [46]. Note that while NEGF studies of optical signals obtained employing the bare perturbation theory are available on the literature [47], consistent NEGF treatment for optics and transport is new.

In our recent work we showed that shortcomings of the bare perturbation theory expansion can be corrected within the NEGF consideration [48]. The correction deals with imposing conservation laws following celebrated works by Kadanoff and Baym [49, 50]. The approach implies self-consistent procedure in treating lightmatter interaction which reflects physics of mutual influence of light and matter and thus requires significant

^{*} mgalperin@tauex.tau.ac.il

changes in classification of optical processes accepted in nonlinear spectroscopy studies. At the same time, it was noted in a number of publications that lowest (second order) bare diagrammatic expansion does satisfy conservation laws in open systems [51–53]. Bare diagrammatic expansion while not equivalent nevertheless is close to philosophy of bare perturbation theory consideration accepted in nonlinear optical spectroscopy.

Here, we discuss general (any order) approach to formulating conserving bare diagrammatic expansion within NEGF and compare it with the bare perturbation theory formulation standard in spectroscopy studies. The comparison allows to extend traditional classification of optical processes to the case of open quantum systems. Structure of the paper is the following. In Section II we introduce model of an open quantum system and discuss treatment of its responses within the bare perturbation theory and within the NEGF. Numerical results illustrating conserving character of the bare diagrammatic approach are presented in Section III. Conclusions are drawn in Section IV.

II. NONLINEAR OPTICAL SPECTROSCOPY

A. Model

We consider open quantum system S driven by external radiation field rad (see Fig. 1). Hamiltonian of the model is

$$\hat{H} = \hat{H}_S + \hat{H}_{rad} + \hat{V}_{S,rad} \tag{1}$$

where \hat{H}_S and \hat{H}_{rad} are Hamiltnians of the system and radiation field, respectively. $\hat{V}_{S,rad}$ couples between the two subsystems. The system is a junction which consists of molecule M coupled set of contacts K. Radiation field is modeled as a set of modes α . Explicit expressions are

FIG. 1. Optical spectroscopy of an open quantum system.

(here and below $\hbar = k_B = 1$)

$$\hat{H}_{S} = \sum_{m \in M} \varepsilon_{m} \hat{d}_{m}^{\dagger} \hat{d}_{m} + \sum_{K} \sum_{k \in K} \varepsilon_{k} \hat{c}_{k}^{\dagger} \hat{c}_{k} + \sum_{m \in M} \sum_{K} \sum_{k \in K} \left(V_{mk} \hat{d}_{m}^{\dagger} \hat{c}_{k} + \text{H.c.} \right) \hat{H}_{rad} = \sum_{\alpha \in rad} \omega_{\alpha} \hat{a}_{\alpha}^{\dagger} \hat{a}_{\alpha} \hat{V}_{S,rad} = -\hat{P} \sum_{\alpha \in rad} \left(E_{\alpha} \hat{a}_{\alpha} + \text{H.c.} \right)$$
(2)

Here, \hat{d}_m^{\dagger} (\hat{d}_m) and \hat{c}_k^{\dagger} (\hat{c}_k) creates (annihilates) electron in molecular orbital m and contact state k, respectively.

$$\hat{P} \equiv \sum_{m_1, m_2 \in M} \mu_{m_1 m_2} \hat{d}^{\dagger}_{m_1} \hat{d}_{m_2}$$
(3)

is the molecular polarization operator with $\mu_{m_1m_2}$ being the matrix element of molecular dipole. $\hat{a}^{\dagger}_{\alpha}$ (\hat{a}_{α}) creates (annihilates) excitation in mode α of the field, and $E_{\alpha} = i (2\pi \hbar \omega_{\alpha}/V)^{1/2}$ is the field amplitude.

We note in passing that non-interacting (molecular orbitals) form of the Hamiltonian \hat{H}_M is chosen for simplicity. Generalization to include intra-molecular interactions and/or couplings to Bose degrees of freedom other than photons (phonons, electron-hole excitations, thermal baths, etc.) would require either consideration of additional diagrams or implementation of many-body flavors of NEGF [54].

Below we will be interested in simulation of electron (e) and photon (p) particle (I) and energy (J) fluxes

$$I_{e}^{K}(t) \equiv -\sum_{k \in K} \frac{d}{dt} \langle \hat{c}_{k}^{\dagger}(t) \hat{c}_{k}(t) \rangle$$

$$J_{e}^{K}(t) \equiv -\sum_{k \in K} \varepsilon_{k} \frac{d}{dt} \langle \hat{c}_{k}^{\dagger}(t) \hat{c}_{k}(t) \rangle$$

$$I_{p}(t) \equiv -\sum_{\alpha \in rad} \frac{d}{dt} \langle \hat{a}_{\alpha}^{\dagger}(t) \hat{a}_{\alpha}(t) \rangle$$

$$J_{p}(t) \equiv -\sum_{\alpha \in rad} \omega_{\alpha} \frac{d}{dt} \langle \hat{a}_{\alpha}^{\dagger}(t) \hat{a}_{\alpha}(t) \rangle$$
(4)

Here, $\langle \ldots \rangle$ indicates quantum mechanical and statistical average and operators are in the Heisenberg picture.

B. The PT formulation

Markov evolution and bare perturbative expansion are the two main assumptions of the traditional theory of nonlinear optical spectroscopy. The former is caused by similarity between Markov form of the quantum master equation (QME),

$$\frac{d}{dt}|\rho(t)\rangle = -i\mathcal{L}(t)|\rho(t)\rangle, \qquad (5)$$



and the Schrödinger equation which was employed at earlier stages of spectroscopy theory. Here,

$$\mathcal{L}(t) = \mathcal{L}_M + \sum_K (\mathcal{L}_K + \mathcal{L}_{MK}) + \mathcal{L}_{rad} + \mathcal{L}_{S,rad}$$

$$\equiv \mathcal{L}_0 + \mathcal{L}_{S,rad}$$
(6)

is Liouvillian which describes evolution of molecular system (free isolated molecule dynamics \mathcal{L}_M) and radiation field (free radiation field dynamics \mathcal{L}_{rad}) in presence of contacts (dissipators \mathcal{L}_K). Coupling between molecule and the field is described by $\mathcal{L}_{S,rad}$.

Markov QME (5) yields simple solution which in interaction picture is

$$|\rho^{I}(t)\rangle = T \exp\left[-i \int_{t_{0}}^{t} dt' \mathcal{L}^{I}_{S,rad}(t')\right] |\rho^{I}(t_{0})\rangle$$
(7)

Expansion of evolution operator in Taylor series yields expression for system density which accounts for lightmatter interaction in order-by-order (bare perturbation theory) form

$$|\rho^{I}(t)\rangle = \sum_{n=0}^{\infty} (-i)^{n} \int_{t_{0}}^{t} dt_{1} \int_{t_{0}}^{t_{1}} dt_{2} \dots \int_{t_{0}}^{t_{n-1}} dt_{n}$$
$$\mathcal{L}^{I}_{S,rad}(t_{1}) \mathcal{L}^{I}_{S,rad}(t_{2}) \dots \mathcal{L}^{I}_{S,rad}(t_{n}) \hat{\rho}^{I}(t_{0}) \quad (8)$$
$$\equiv \sum_{n=0}^{\infty} |\rho^{I(n)}(t)\rangle$$

Using this form in expressions (3)-(4) yields bare pertur-

bation theory expansions for system properties

$$I_{e}^{K}(t) = \langle\!\langle N_{e} | i\mathcal{L}_{K} | \rho(t) \rangle\!\rangle \equiv \sum_{n=0}^{\infty} I_{e}^{K(n)}(t)$$

$$J_{e}^{K}(t) = \langle\!\langle H_{e} | i\mathcal{L}_{K} | \rho(t) \rangle\!\rangle \equiv \sum_{n=0}^{\infty} J_{e}^{K(n)}(t)$$

$$I_{p}(t) = \langle\!\langle N_{p} | i\mathcal{L}_{S,rad} | \rho(t) \rangle\!\rangle \equiv \sum_{n=0}^{\infty} I_{p}^{(n)}(t)$$

$$J_{p}(t) = \langle\!\langle H_{p} | i\mathcal{L}_{S,rad} | \rho(t) \rangle\!\rangle \equiv \sum_{n=0}^{\infty} J_{p}^{(n)}(t)$$
(9)

Here, $\hat{N}_e = \sum_{m \in M} \hat{d}_m^{\dagger} \hat{d}_m$ and $\hat{N}_p = \sum_{\alpha \in rad} \hat{a}_{\alpha}^{\dagger} \hat{a}_{\alpha}$ is the electron and photon number operator, respectively. $\hat{H}_e = \sum_{m \in M} \varepsilon_m \hat{d}_m^{\dagger} \hat{d}_m$ and $\hat{H}_p \equiv \hat{H}_{rad} = \sum_{\alpha \in rad} \omega_\alpha \hat{a}_{\alpha}^{\dagger} \hat{a}_{\alpha}$. Note opposite sign as compared to traditional definition of optical flux in spectroscopy studies. Expansion of $I_p(t)$, Eq.(9), is the basis of description and classification of optical processes in traditional nonlinear optical spectroscopy.

In many situations radiation field is treated classically. In this case (5) is equation for molecular density operator only (this is contrary to total - light and matter - density operator of quantum consideration), contribution $\mathcal{L}_{S,rad}$ becomes time-dependent, and term \mathcal{L}_{rad} drops.

C. The NEGF formulations

Central object in NEGF formulation is single-particle Green's function of the open quantum system

$$G_{m_1m_2}(\tau_1, \tau_2) = -\langle T_c \, \hat{d}_{m_1}(\tau_1) \, \hat{d}_{m_2}^{\dagger}(\tau_2) \rangle \tag{10}$$

where T_c is the Keldysh contour ordering operator and $\tau_{1,2}$ are the contour variables. Its dynamics is described by the Dyson equation which can be written in two equivalent forms [55]

$$G_{m_1m_2}(\tau_1,\tau_2) = G_{m_1m_2}^{(0)}(\tau_1,\tau_2) + \sum_{m_3,m_4} \int_c d\tau_3 \int_c d\tau_4 \, G_{m_1m_3}^{(0)}(\tau_1,\tau_3) \, \Sigma_{m_3m_4}^{ep}(\tau_3,\tau_4) \, G_{m_4m_2}(\tau_4,\tau_2) \tag{11}$$

$$=G_{m_1m_2}^{(0)}(\tau_1,\tau_2) + \sum_{m_3,m_4} \int_c d\tau_3 \int_c d\tau_4 \, G_{m_1m_3}^{(0)}(\tau_1,\tau_3) \, \Sigma_{m_3m_4}^{ep\,(r)}(\tau_3,\tau_4) \, G_{m_4m_2}^{(0)}(\tau_4,\tau_2) \tag{12}$$

Here, $G^{(0)}$ is Green's function of the junction in absence of radiation field, Σ^{ep} and $\Sigma^{ep(r)}$ are proper and reducible forms of the self-energy which describes electron-photon interaction. Due to many-body character of the interaction, exact expression for the self-energy is not available. Diagrammatic technique is a way to obtain approximate (up to a particular order in the interaction) solutions. For the approximation to be conserving the self-energy should be Φ -derivable [55]

$$\Sigma_{m_1 m_2}^{ep}(\tau_1, \tau_2) = -\frac{\delta \Phi^{ep}}{\delta G_{m_2 m_1}(\tau_2, \tau_1)}$$
(13)

$$\Sigma_{m_1m_2}^{ep(r)}(\tau_1, \tau_2) = -\frac{\delta \Phi^{ep(r)}}{\delta G_{m_2m_1}^{(0)}(\tau_2, \tau_1)}$$
(14)

Here, Φ^{ep} is the sum of all connected and topologically inequivalent skeleton diagrams, while $\Phi^{ep(r)}$ is the sum of all connected and topologically inequivalent vacuum diagrams (no dressing). Diagrams up to fourth order in



FIG. 2. Φ functional for electron-photon interaction. Shown are (a)-(b) second and (c)-(d) fourth order contributions to Φ^{ep} . Panel (e) shows additional fourth order contribution to $\Phi^{ep(r)}$. Directed solid line represents electron propagator G or $G^{(0)}$. Wavy line represents photon propagator $F^{(0)}$, both directions are possible for each wavy line.

electron-photon interaction are shown in Fig. 2. Explicit expressions for the self-energies are given in Appendix A and corresponding diagrams are shown in Fig. 7. Note that Φ functional presented in Fig. 2b does not contribute to self-energy because zero-frequency photons do not exist. However, the functional does contribute to kernel expression (see explanation below); the contribution is presented in Fig. 8b. To calculate energy fluxes, Eqs. (4), one needs to know also two-particle Green's function

$$G_{m_1m_2,m_3m_4}(\tau_1,\tau_2;\tau_3,\tau_4) \equiv -\langle T_c \, \hat{d}_{m_1}(\tau_1) \, \hat{d}_{m_2}(\tau_2) \, \hat{d}^{\dagger}_{m_4}(\tau_4) \, \hat{d}^{\dagger}_{m_3}(\tau_3) \rangle$$
(15)

Its dynamics is described by (we use shorthand notation with 1 standing for m_1 and τ_1 , etc.)

$$G(1,2;3,4) = G(1,3)G(2,4) - G(1,4)G(2,3)$$

$$+ \int d1' \int d2' \int d3' \int d4' G(1,1')G(3',3)K^{ep}(1',2';3',4') [G(2,2')G(4',4) - G(2,4';4,2')]$$

$$= G(1,3)G(2,4) - G(1,4)G(2,3)$$

$$+ \int d1' \int d2' \int d3' \int d4' G^{(0)}(1,1')G^{(0)}(3',3)K^{ep}(r)(1',2';3',4')G^{(0)}(2,2')G^{(0)}(4',4)$$

$$(16)$$

$$(17)$$

Here, K^{ep} and $K^{ep(r)}$ are irreducible and reducible (in two-particle sense) forms of the kernel.

Similar to self-energy, for the approximation to be conserving the kernel should be Φ -derivable (second derivative) [55]

$$K^{ep}(1,2;3,4) = \frac{\delta^2 \Phi^{ep}}{\delta G(4,2) \,\delta G(3,1)} \equiv -\frac{\delta \Sigma^{ep}(1,3)}{\delta G(4,2)} \quad (18)$$

Reducible kernel, $K^{ep(r)}$, is related to the irreducible one, K^{ep} , via the Bethe-Salpeter equation

$$K^{ep(r)}(1,2;3,4) = K^{ep}(1,2;3,4)$$

- $\int d1' \int d2' \int d3' \int d4' K^{ep}(1,2';3,4')$ (19)
 $\times G(4',1') G(3',2') K^{ep(r)}(1',2;3',4)$

Diagrams up to fourth order in electron-photon interaction and explicit expressions for the irreducible kernel are given in Appendix B and corresponding diagrams are shown in Fig. 8.

The knowledge of Green's functions (11) and (16) or (12) and (17) allows us to calculate molecular polarization, and particle and energy fluxes

$$\begin{split} I_{e}^{K}(t) &= 2 \operatorname{Re} \sum_{m,m' \in M} \int_{t_{0}}^{t} dt' \left(\Sigma_{mm'}^{K<}(t,t') \, G_{m'm}^{>}(t',t) - \Sigma_{mm'}^{K>}(t,t') \, G_{m'm}^{<}(t',t) \right) \\ J_{e}^{K}(t) &= 2 \operatorname{Re} \sum_{k \in K} \varepsilon_{k} \sum_{m,n \in M} \int_{t_{0}}^{t} dt' \, V_{mk} V_{km'} \left(g_{k}^{<}(t,t') \, G_{m'm}^{>}(t',t) - g_{k}^{>}(t,t') \, G_{m'm}^{<}(t',t) \right) \\ I_{p}(t) &= -2 \operatorname{Im} \sum_{m,n,m',n' \in M} \int_{t_{0}}^{t} dt' \left(\Pi_{mn,m'n'}^{<}(t,t') \, G_{mn',nm'}^{-+++}(t,t';t,t') - \Pi_{mn,m'n'}^{>}(t,t') \, G_{mn',nm'}^{+++-}(t,t',t,t') \right) \end{split}$$
(20)

$$&= -2 \operatorname{Re} \sum_{\alpha \in rad} \int_{t_{0}}^{t} dt' \left| E_{\alpha} \right|^{2} \left(F_{\alpha}^{(0)}(t,t') \, G_{PP}^{>}(t't) - F_{\alpha}^{(0)}(t,t') \, G_{PP}^{<}(t't) \right) \\ J_{p}(t) &= -2 \operatorname{Im} \sum_{\alpha \in rad} \omega_{\alpha} \sum_{m,n,m',n' \in M} \int_{t_{0}}^{t} dt' \, \mu_{mn}^{*} \mu_{m'n'} \left| E_{\alpha} \right|^{2} \left(F_{\alpha}^{(0)}(t,t') \, G_{mn',nm'}^{++++}(t,t';t,t') - F_{\alpha}^{(0)}(t,t') \, G_{mn',nm'}^{++++}(t,t';t,t') - F_{\alpha}^{(0)}(t,t') \, G_{mn',nm'}^{++++}(t,t';t,t') \right) \end{split}$$

Here, $G^{>/<}$ is the lesser/greater projection of the Green's function (11) or (12), Σ^K and g_k are, respectively, the self-energy due to coupling to contact K and Green's function of free electron in state k in the contact

$$g_k(\tau_1, \tau_2) \equiv -i \langle T_c \, \hat{c}_k(\tau_1) \, \hat{c}_k^{\dagger}(\tau_2) \rangle_0$$

$$\Sigma_{m_1 m_2}^K(\tau_1, \tau_2) \equiv \sum_{k \in K} V_{m_1 k} g_k(\tau_1, \tau_2) V_{k m_2}$$
(21)

 $G_{mn',m'n}$ are the - + +- and + - -+ projections $(-/+indicates forward/backward branch of the Keldysh contour) of the two-particle Green's function and <math>\Pi$ and $F_{\alpha}^{(0)}$ are the photon-induced interaction and Green's function of free photon defined in Eqs. (A5) and (A6), respectively. G_{PP} is correlation function of the molecular polarization operator

$$G_{PP}(\tau_1, \tau_2) \equiv -i \langle T_c \, \hat{P}(\tau) \, \hat{P}^{\dagger}(\tau') \rangle \tag{22}$$

The following observations are notable:

- 1. Because electron-photon is the only many-body interaction of the model (1)-(2) we discuss only its contributions to approximate expressions for self-energy and kernel. In presence of additional many-body interactions (e.g., electron-electron or electron-phonon) additional contributions to these expressions should be taken into account. Moreover, interactions resulting in time-nonlocal correlations in electron subsystem may lead to contributions mixed with those of the radiation field. This would affect discussion of optical processes presented below.
- 2. Formulation (11) and (16) is self-consistent procedure. For example, single-particle Green's function (11) depends on self-energy, and self-energy, Eqs. (A1) and (A3), depends on Green's function. This self-consistency accounts for mutual influence of electrons and radiation field in determining state of the system. The formulation is standard in studies of quantum transport and we used it in our previous work to discuss nonlinear spectroscopy in open systems [48].

- 3. Formulation (12) and (17) treats interaction with radiation field order-by-order. Thus, results for molecular polarization and fluxes are obtained as sum of contributions of different orders in the lightmatter interaction. In this sense, (12) and (17) is closer in spirit to scattering type consideration of traditional nonlinear spectroscopy.
- 4. For the non-interacting molecular model represented by Hamiltonian \dot{H}_S , Eq.(2), consideration of the two-particle Green's function (15) and higher correlations is only relevant in the self-consistent formulation (11) and (16). Single-particle Green's function (10) is the only correlation function required in the formulation (12) and (17), because all higher order correlation functions can be expressed in terms of single-particle Green's function employing the Wick's theorem.
- 5. As we discussed in Refs. [42, 48], employing bare perturbation expansion, which is standard in nonlinear spectroscopy studies, in the formulation (11) and (16) leads to violation of conservation laws. Below we use formulation (12) and (17) to compare NEGF with the bare perturbation theory treatment of traditional nonlinear optical spectroscopy.

D. Comparison between PT and NEGF

Here, we follow classification of optical processes introduced in Chapter 9 of Ref. [28]. We note in passing that classification of second order process as absorption and fourth order as spontaneous emission used in Ref. [28] is based on assumptions of occupied ground and unoccupied excited molecular states. For nonequilibrium with bias induced occupation of excited state and/or depletion of ground state the classification does not hold anymore. Nevertheless, below we use traditional names for the second and fourth order optical processes.

1. Absorption of a quantum field

In the PT formulation, linear absorption is given by n = 1 contribution to photon flux in Eq.(9)

$$I_{p}^{abs}(t) = \langle\!\langle N_{p} | \int_{t_{0}}^{t} dt_{1} \mathcal{L}_{S,rad}^{I}(t) \mathcal{L}_{S,rad}^{I}(t_{1}) | \rho(t_{0}) \rangle\!\rangle$$
(23)
= $\int_{t_{0}}^{t} dt_{1} \operatorname{Tr} \{ [[\hat{N}_{p}, \hat{H}_{S,rad}^{I}(t)], \hat{H}_{S,rad}^{I}(t_{1})] \hat{\rho}(t_{0}) \}$

Here, $\hat{\rho}$ is the total density matrix of the system. Note that initially molecule and radiation field are assumed to be decoupled

$$\hat{\rho}(t_0) = \hat{\rho}_M(t_0) \otimes \hat{\rho}_{rad}(t_0) \tag{24}$$

Note also that usually absorption is considered driven by a particular mode α of radiation field (only one mode of radiation field is included in the Liouvillian/Hamiltonian).

Within the (12) and (17) NEGF formulation, absorption is given by the second order (in light-matter coupling) contribution to photon flux. It is obtained from (20) by substituting

$$G_{PP}^{(0)}(\tau,\tau') \equiv -i \sum_{m_1,m_2,m_3,m_4 \in M} \mu_{m_1m_2} \mu^*_{m_3m_4} \qquad (25)$$
$$\times G_{m_3m_1}^{(0)}(\tau',\tau) G_{m_2m_4}^{(0)}(\tau,\tau')$$

in place of G_{PP} in the expression. This leads to

$$I_{p}^{(2)}(t) = -2 \operatorname{Re} \sum_{\alpha \in rad} \int_{t_{0}}^{t} dt' |E_{\alpha}|^{2}$$

$$\times \left(F_{\alpha}^{(0)} < (t,t') \, G_{PP}^{(0)}(t',t) - F_{\alpha}^{(0)} < (t,t') \, G_{PP}^{(0)}(t',t) \right)$$
(26)

It is straightforward to show that after evaluating commutators in the PT result, Eq.(23), and separating molecular and radiation field degrees of freedom one gets exactly structure of the NEGF expression, Eq.(26). The only difference is lack of hybridization information in the PT correlation function of molecular polarization (often this information is added to the correlation function in an approximate *ad hoc* manner), while the NEGF expression accounts for the hybridization exactly.

2. Spontaneous light emission spectroscopy

In the PT formulation, SLE is given by n = 3 contribution to photon flux in Eq.(9)

$$I_{p}^{SLE}(t) = \langle\!\langle N_{p} | \int_{t_{0}}^{t} dt_{1} \int_{t_{0}}^{t_{1}} dt_{2} \int_{t_{0}}^{t_{2}} dt_{3}$$

$$\mathcal{L}_{S,rad}^{I}(t) \mathcal{L}_{S,rad}^{I}(t_{1}) \mathcal{L}_{S,rad}^{I}(t_{2}) \mathcal{L}_{S,rad}^{I}(t_{3}) | \rho(t_{0}) \rangle$$

$$= \int_{t_{0}}^{t} dt_{1} \int_{t_{0}}^{t_{1}} dt_{2} \int_{t_{0}}^{t_{2}} dt_{3} \operatorname{Tr} \{ [[[[\hat{N}_{p}, \hat{H}_{S,rad}^{I}(t_{3})] \rho(t_{0})], \\ \hat{H}_{S,rad}^{I}(t_{1})], \hat{H}_{S,rad}^{I}(t_{2})], \hat{H}_{S,rad}^{I}(t_{3})] \hat{\rho}(t_{0}) \}$$

Within the (12) and (17) NEGF formulation, SLE is given by the fourth order (in light-matter coupling) contribution to photon flux. It is obtained from (20) by substituting

$$G_{PP}^{(2)}(\tau,\tau') \equiv -i \sum_{m_1,m_2,m_3,m_4 \in M} \mu_{m_1m_2} \mu_{m_3m_4}^* \left(G_{m_3m_1}^{(0)}(\tau',\tau) G_{m_2m_4}^{(2)}(\tau,\tau') + G_{m_3m_1}^{(2)}(\tau',\tau) G_{m_2m_4}^{(0)}(\tau,\tau') - \sum_{n_1,n_2,n_3,n_4} \int_c d\tau_1 \int_c d\tau_2 \int_c d\tau_3 \int_c d\tau_4 \qquad (28) \\ G_{m_2n_1}^{(0)}(\tau,\tau_1) G_{n_3m_1}^{(0)}(\tau_3,\tau) K_{n_1n_2;n_3n_4}^{ep(r)(2)}(\tau_1,\tau_2;\tau_3,\tau_4) \\ G_{m_3n_2}^{(0)}(\tau',\tau_2) G_{n_4m_4}^{(0)}(\tau_4,\tau') \right)$$

in place of G_{PP} in the expression. Here,

$$G_{m_1m_2}^{(2)}(\tau_1,\tau_2) = \sum_{m_3,m_4} \int_c d\tau_3 \int_c d\tau_4$$
(29)
$$G_{m_1m_3}^{(0)}(\tau_1,\tau_3) \Sigma_{m_3m_4}^{ep(r)(2)}(\tau_3,\tau_4) G_{m_4m_2}^{(0)}(\tau_4,\tau_2)$$

and expression for $K^{ep(r)(2)}$ is given in Eq.(B1). This leads to

$$I_{p}^{(4)}(t) = -2 \operatorname{Re} \sum_{\alpha \in rad} \int_{t_{0}}^{t} dt' |E_{\alpha}|^{2}$$

$$\times \left(F_{\alpha}^{(0)} < (t,t') \, G_{PP}^{(2)}(t',t) - F_{\alpha}^{(0)}(t,t') \, G_{PP}^{(2)}(t',t) \right)$$
(30)

Direct comparison between the two approaches is less straightforward in this case because multi-time electron correlation function is evaluated within different schemes: the quantum regression theorem (QRT) in the QME case vs. the Bethe-Salpeter equation in the NEGF formulation. Nevertheless, the following conclusions are possible

- 1. Standard nonlinear spectroscopy as well as (12) and (17) NEGF formulation are similar because both are based on PT expansion of multi-time correlation function of molecular polarization. Their main difference is in the way to evaluate the multi-time correlation function.
- 2. By the very construction, QRT evaluates multitime correlation function as a sequence of blocks between adjacent times when molecule interacts with radiation field. Electronic correlations due to coupling to Fermi baths for pair of times in different blocks are missing in such scheme. They are preserved in the NEGF formulation.
- 3. In particular, inability by QRT to describe correlation across the blocks implies that radiation field induced interactions (such as, e.g., presented by diagram in Fig. 7c) cannot be accounted properly in the standard nonlinear spectroscopy formulation. We note in passing that limitations of the QRT when applied to description of open quantum systems were extensively discussed in the literature [56–59].



FIG. 3. (Color online) Model of an open donor-bridgeacceptor system with light assisted tunneling.

- 4. Another point of critical difference between the PT and NEGF is related to accounting for all possible dicouplings within PT, while only connected diagrams contribute within the NEGF. Thus, among other processes the PT accounts for virtual excitation in the radiation field, which neither affect the system, nor related to its physics.
- 5. Finally, we note the difference in language of standard nonlinear spectroscopy PT formulation and that of diagrammatic NEGF expansion. Famous double sided Feynman diagrams of the PT formulation are related to particular arrangements of times when molecule interacts with radiation field on the Keldysh contour. Within the NEGF, such arrangements are called projections while word diagrams is used to indicate different physical processes (see Fig. 7).

III. NUMERICAL RESULTS

In Ref. [42] we demonstrated violation of conservation laws in an open system when bare perturbation theory is employed in (11) and (16) formulation of the NEGF. Here, we illustrate that perturbative diagrammatic expansion (12) and (17) is conserving.

We employ the same donor-bridge-acceptor molecular model as in Ref. [42] (see Fig. 3). Parameters of the simulation are temperature ($k_BT = 0.25$), positions of molecular levels ($\varepsilon_1 = -5$, $\varepsilon_2 = 5$, $\varepsilon_3 = -2$). Electron escape rates into metallic contacts are taken within the wide-band approximation ($\Gamma_1^L = \Gamma_3^R = 1$, $\Gamma_2^L = \Gamma_2^R = 0.1$). Energy escape rate into radiation bath is modeled as

$$\gamma(\omega) = \gamma_0 \left(\frac{\omega}{\omega_0}\right)^2 e^{(\omega_0^2 - \omega^2)/\omega_C^2} \tag{31}$$

with $\gamma_0 = 0.01$ and $\omega_C = 10$. Molecule is pumped by a monochromatic laser characterized by its frequency tuned to resonance with molecular transition between levels 2 and 3, $\omega_0 = \varepsilon_2 - \varepsilon_3 = 7$, intensity $N_0 = 1$, and bandwidth $\delta = 0.1$. Fermi energy is taken as origin, $E_F = 0$, and bias is assumed to be applied symmetrically, $\mu_{L,R} = E_F \pm |e|V_{sd}/2$. Simulations are performed on energy grid spanning region from -100 to 100 with step 0.01. Here, energy is given in terms of an arbitrary unit of energy E_0 . Results for particle and energy fluxes are presented in terms of flux units $I_0 = 1/t_0$ and $J_0 = E_0/t_0$, respectively, where $t_0 = \hbar/E_0$ is unit of time.

Figure 4 shows photon flux vs. applied bias. As mentioned earlier second order optical process (panel a) is pure absorption and fourth order optical process (panel b) is pure emission only at and near equilibrium. Significant changes in the total photon flux (panel c) occur when chemical potentials cross molecular resonances.

Figure 5 illustrates charge conservation. At steadystate amount of charge entering the system at left interface should be compensated by amount of charge leaving the system at right interface. That is, $I_e^L = -I_e^R$. As expected, currents at the two interfaces (solid and dashed lines) sum to zero (dotted line). Note that conserving character of the approximation holds at every order of perturbation theory (see panels a, b, and c).

Figure 6 illustrates energy conservation. At steadystate amount of energy entering the system should be compensated by amount of charge leaving the system. Energy enters (leaves) the system from (to) left and right contacts and from (to) radiation field modes. That is, $J_e^L + J_e^R + J_p = 0$. As expected, energy currents at the two interfaces (solid and dashed lines) plus energy current from radiation modes (dash-dotted line) sum to zero (dotted line). Again, conservation holds separately for contributions of every order in light-matter interaction.



FIG. 4. (Color online) Photon flux in DBA junction of Fig. 3 vs applied bias. Shown are (a) second and (b) fourth contributions. Panel (c) shows total photon flux.



FIG. 5. (Color online) Electron (particle) currents in DBA junction of Fig. 3 vs applied bias. Shown are currents at left I_e^L (solid line, blue) and right I_e^R (dashed line, red) interfaces for (a) zero, (b) second, and (c) fourth orders in light-matter interaction. Panel (d) shows total electron currents. In each panel, sum of the left and right currents is indicated by dotted line (grey).

IV. CONCLUSIONS

Development of experimental techniques at nanoscale resulted in combination of research fields of optical spectroscopy and molecular electronics. Today, theoretical studies of spectroscopy in open quantum systems use methods traditionally coming from nonlinear optical spectroscopy and quantum transport. The former employ Markov QME description of molecular dynamics and are based on bare perturbation theory expansion of light-matter interaction. Molecular electronics, which historically developed from quantum transport measurements, employs diagrammatic expansion of the NEGF method. Usual NEGF formulation is based on consideration of dressed diagrams which corresponds to accounting for mutual influence of light and matter degrees of freedom on their dynamics. In our previous studies we noted that employing bare perturbation expansion in such formulation leads to violation of conservation laws.

Here, we consider an alternative - bare diagrams for-



FIG. 6. (Color online) Energy currents in DBA junction of Fig. 3 vs applied bias. Shown are electron energy currents at left J_e^L (solid line, blue) and right J_e^R (dashed line, red) interfaces, and photon energy current J_p (dash-dotted line, green) for (a) zero, (b) second, and (c) fourth orders in light-matter interaction. Panel (d) shows corresponding total energy currents. In each panel, sum of the energy currents is indicated by dotted line (grey).

mulation of the NEGF. This description is close in spirit to the bare PT of traditional nonlinear spectroscopy and thus is advantageous in attempt to explore connections between theoretical methods of the two communities. Comparing the two methods we find close similarity in overall structure of the approaches: both are based on bare perturbation theory expansion in light-matter interaction, in each order of the expansion both require evaluation of a multi-time correlation function of molecular polarization operators. Main difference between the methods is in the way the latter evaluation is performed: Markov QME description of traditional nonlinear optical spectroscopy relies on the quantum regression theorem (QRT), while NEGF employs Wick's theorem and diagrammatic methods.

We note that limitation of the QRT in description of dynamics in open quantum systems was widely discussed in the literature. In particular, we argue that QRT describes dynamics as sequence of propagations between times when molecule interacts with the radiation field and as such is not adequate in description of coherences in molecular system across the times. For example, it cannot properly describe two-time correlation function with one time before and the other after the time of interaction. Note that such correlation functions naturally appear in non-Markov NEGF description. Note also that in case of quantum light the limitation affects description of radiation field induced intra-molecular interactions (see, e.g., diagram in Fig. 7c).

On formal level we note difference in terminology between the PT and NEGF treatments. In particular, traditional nonlinear optical spectroscopy employs double sided Feynman diagrams for description and classification of optical processes. These diagrams are particular arrangements of light-matter interaction times on the Keldysh contour. Within the NEGF, such arrangements are called projections, while word diagram is reserved for description of different physical processes in light-matter interaction.

We conclude that traditional way of processes classification accepted in nonlinear optical spectroscopy is preserved when describing open quantum systems. However, it is important to employ non-Markov description of dynamics in such systems. Technically, this results in using diagrammatic methods of the NEGF in evaluation of multi-time correlation functions instead of the QRT as is employed in traditional spectroscopy studies.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation under Grant No. CHE-2154323.

Appendix A: Self-energies due to electron-photon interaction



FIG. 7. Self-energy due to electron-photon interaction. Shown are second (panel a) and fourth (panels b and c) order contributions to Σ^{ep} . Panels d and e show additional fourth order contributions to $\Sigma^{ep(r)}$.

Second order expressions are

$$\Sigma_{m_1m_2}^{ep(2)}(\tau_1,\tau_2) = i \sum_{n_1,n_2 \in M} G_{n_1n_2}(\tau_1,\tau_2) \left[\Pi_{n_1m_1,n_2m_2}(\tau_1,\tau_2) + \Pi_{m_2n_2,m_1n_1}(\tau_2,\tau_1) \right]$$
(A1)

$$\Sigma_{m_1m_2}^{ep(r)(2)}(\tau_1,\tau_2) = i \sum_{n_1,n_2 \in M} G_{n_1n_2}^{(0)}(\tau_1,\tau_2) \left[\Pi_{n_1m_1,n_2m_2}(\tau_1,\tau_2) + \Pi_{m_2n_2,m_1n_1}(\tau_2,\tau_1) \right]$$
(A2)

Fourth order expressions

$$\Sigma_{m_{1}m_{2}}^{ep(4)}(\tau_{1},\tau_{2}) = \sum_{n_{1},n_{2},n_{3},n_{4},m_{3},m_{4}} \int_{c} d\tau_{3} \int_{c} d\tau_{4}$$
(A3)

$$\begin{pmatrix} G_{n_{1}n_{2}}(\tau_{1},\tau_{2}) G_{m_{3}m_{4}}(\tau_{3},\tau_{4}) G_{n_{4}n_{3}}(\tau_{4},\tau_{3}) \\ \times [\Pi_{n_{1}m_{1},n_{3}m_{3}}(\tau_{1},\tau_{3}) + \Pi_{m_{3}n_{3},m_{1}n_{1}}(\tau_{3},\tau_{1})] [\Pi_{m_{2}n_{2},m_{4}n_{4}}(\tau_{2},\tau_{4}) + \Pi_{n_{4}m_{4},n_{2}m_{2}}(\tau_{4},\tau_{2})] \\ - G_{n_{1}m_{3}}(\tau_{1},\tau_{3}) G_{n_{3}n_{4}}(\tau_{3},\tau_{4}) G_{m_{4}n_{2}}(\tau_{4},\tau_{2}) \\ \times [\Pi_{n_{1}m_{1},n_{4}m_{4}}(\tau_{1},\tau_{4}) + \Pi_{m_{4}n_{4},m_{1}n_{1}}(\tau_{4},\tau_{1})] [\Pi_{n_{3}m_{3},n_{2}m_{2}}(\tau_{3},\tau_{2}) + \Pi_{m_{2}n_{2},m_{3}n_{3}}(\tau_{2},\tau_{3})] \end{pmatrix}$$

$$\Sigma_{m_{1}m_{2}}^{ep(r)(4)}(\tau_{1},\tau_{2}) = \sum_{n_{1},n_{2},n_{3},n_{4},m_{3},m_{4}} \int_{c} d\tau_{3} \int_{c} d\tau_{4}$$
(A4)

$$\begin{pmatrix} G_{n_{1}n_{2}}^{(0)}(\tau_{1},\tau_{2}) G_{m_{3}m_{4}}^{(0)}(\tau_{3},\tau_{4}) G_{n_{4}n_{3}}^{(0)}(\tau_{4},\tau_{3}) \\ \times [\Pi_{n_{1}m_{1},n_{3}m_{3}}(\tau_{1},\tau_{3}) + \Pi_{m_{3}n_{3},m_{1}n_{1}}(\tau_{3},\tau_{1})] [\Pi_{m_{2}n_{2},m_{4}n_{4}}(\tau_{2},\tau_{4}) + \Pi_{n_{4}m_{4},n_{2}m_{2}}(\tau_{4},\tau_{2})] \\ - G_{n_{1}m_{3}}^{(0)}(\tau_{1},\tau_{3}) G_{n_{3}n_{4}}^{(0)}(\tau_{3},\tau_{4}) G_{m_{4}n_{2}}^{(0)}(\tau_{4},\tau_{2}) \\ \times \left\{ [\Pi_{n_{1}m_{1},n_{4}m_{4}}(\tau_{1},\tau_{4}) + \Pi_{m_{4}n_{4},m_{1}n_{1}}(\tau_{4},\tau_{1})] [\Pi_{n_{3}m_{3},n_{2}m_{2}}(\tau_{3},\tau_{2}) + \Pi_{m_{2}n_{2},m_{3}n_{3}}(\tau_{2},\tau_{3})] \\ + [\Pi_{n_{1}m_{1},n_{2}m_{2}}(\tau_{1},\tau_{2}) + \Pi_{m_{3}n_{3},m_{1}n_{1}}(\tau_{3},\tau_{1})] [\Pi_{n_{3}m_{3},n_{4}m_{4}}(\tau_{3},\tau_{4}) + \Pi_{m_{4}n_{4},m_{3}n_{3}}(\tau_{4},\tau_{3})] \right\} \end{pmatrix}$$

Here,

$$\Pi_{m_1 n_1, m_2 n_2}(\tau_1, \tau_2) = \mu_{m_1 n_1}^* \mu_{m_2 n_2} \sum_{\alpha \in rad} |E_\alpha|^2 F_\alpha^{(0)}(\tau_1, \tau_2)$$
(A5)

is photon-induced interaction between electronic transitions m_1n_1 and m_2n_2 , and

$$F_{\alpha}^{(0)}(\tau_1, \tau_2) = -i \langle T_c \, \hat{a}_{\alpha}(\tau_1) \, \hat{a}_{\alpha}^{\dagger}(\tau_2) \rangle_0 \tag{A6}$$

is Green's function of free photon of mode α .

Appendix B: Irreducible kernel due to electron-photon interaction



FIG. 8. Irreducible kernel due to electron-photon interaction. Shown are second (panels a-b) and fourth (panels b-h) order contributions to K^{ep} .

Second order expression is

$$K_{m_{1}m_{2};m_{3}m_{4}}^{ep\,(2)}(\tau_{1},\tau_{2};\tau_{3},\tau_{4}) = -i\delta(\tau_{1},\tau_{4})\delta(\tau_{2},\tau_{3})\left[\Pi_{m_{4}m_{1},m_{2}m_{3}}(\tau_{1},\tau_{2}) + \Pi_{m_{3}m_{2},m_{1}m_{4}}(\tau_{2},\tau_{1})\right] + i\delta(\tau_{1},\tau_{3})\delta(\tau_{2},\tau_{4})\left[\Pi_{m_{3}m_{1},m_{2}m_{4}}(\tau_{1},\tau_{2}) + \Pi_{m_{4}m_{2},m_{1}m_{3}}(\tau_{2},\tau_{1})\right]$$
(B1)

Fourth order expression is

$$K_{m_{1}m_{2},m_{3}m_{4}}^{ep(4)}(\tau_{1},\tau_{2};\tau_{3},\tau_{4}) =$$
(B2)

$$-\delta(\tau_{1},\tau_{4})\delta(\tau_{2},\tau_{3})\sum_{n_{5},n_{6},m_{5},m_{6}}\int_{c}d\tau_{5}\int_{c}d\tau_{6}G_{n_{5}n_{6}}(\tau_{5},\tau_{6})G_{m_{6}m_{5}}(\tau_{6},\tau_{5}) \times [\Pi_{m_{4}m_{1},m_{5}n_{5}}(\tau_{1},\tau_{5}) + \Pi_{n_{5}m_{5},m_{1}m_{4}}(\tau_{5},\tau_{1})][\Pi_{m_{3}m_{2},n_{6}m_{6}}(\tau_{2},\tau_{6}) + \Pi_{m_{6}n_{6},m_{2}m_{3}}(\tau_{6},\tau_{2})] - \sum_{n_{1},n_{2},n_{3},n_{4}}G_{n_{1}n_{3}}(\tau_{1},\tau_{3})G_{n_{2}n_{4}}(\tau_{2},\tau_{4}) \times \left([\Pi_{n_{1}m_{1},n_{4}m_{4}}(\tau_{1},\tau_{4}) + \Pi_{m_{4}n_{4},m_{1}n_{1}}(\tau_{4},\tau_{1})][\Pi_{m_{3}n_{3},m_{2}n_{2}}(\tau_{3},\tau_{2}) + \Pi_{n_{2}m_{2},n_{3}m_{3}}(\tau_{2},\tau_{3})] + [\Pi_{n_{1}m_{1},n_{2}m_{2}}(\tau_{1},\tau_{2}) + \Pi_{n_{2}m_{2},m_{1}n_{1}}(\tau_{2},\tau_{1})][\Pi_{m_{3}n_{3},n_{4}m_{4}}(\tau_{3},\tau_{4}) + \Pi_{m_{4}n_{4},n_{3}m_{3}}(\tau_{4},\tau_{3})] \right) + \delta(\tau_{2},\tau_{3})\sum_{m,n,n_{1},n_{4}}\int_{c}d\tau G_{n_{1}m_{1}}(\tau_{1},\tau_{1}) + \Pi_{nm,m_{1}m_{4}}(\tau,\tau_{1})][\Pi_{n_{2}m_{2},n_{3}m_{3}}(\tau_{2},\tau_{3}) + \Pi_{m_{3}n_{2},m_{2}n_{2}}(\tau_{3},\tau_{2})] + \delta(\tau_{2},\tau_{3})\sum_{m,n,n_{1},n_{4}}\int_{c}d\tau G_{n_{1}n_{1}}(\tau_{1},\tau_{1}) + \Pi_{m_{4}n_{4},m_{1}n_{1}}(\tau_{4},\tau_{1})][\Pi_{m_{n}m_{2}m_{3}}(\tau,\tau_{3}) + \Pi_{m_{3}m_{3},n_{4}m_{4}}(\tau_{3},\tau_{4})] + \sum_{n_{1},n_{2},n_{3},n_{4}}G_{n_{1}n_{4}}(\tau_{1},\tau_{4}) + \Pi_{n_{4}n_{4},m_{1}n_{1}}(\tau_{2},\tau_{1})][\Pi_{m_{4}n_{4},n_{3}m_{3}}(\tau_{4},\tau_{3}) + \Pi_{m_{3}n_{3},n_{4}m_{4}}(\tau_{3},\tau_{4})]$$

- [1] T. W. Ebbesen, Acc. Chem. Res. 49, 2403 (2016).
- [2] J.-F. Li, C.-Y. Li, and R. F. Aroca, Chem. Soc. Rev. 46, 3962 (2017).
- [3] F. Neubrech, C. Huck, K. Weber, A. Pucci, and H. Giessen, Chem. Rev. 117, 5110 (2017).
- [4] A. B. Zrimsek, N. Chiang, M. Mattei, S. Zaleski, M. O. McAnally, C. T. Chapman, A.-I. Henry, G. C. Schatz, and R. P. Van Duyne, Chem. Rev. **117**, 7583 (2017).
- [5] C. Zhan, X.-J. Chen, J. Yi, J.-F. Li, D.-Y. Wu, and Z.-Q. Tian, Nat. Rev. Chem. 2, 216 (2018).
- [6] N. Picqué and T. W. Hänsch, Nat. Photonics 13, 146 (2019).
- [7] C. R. Baiz, B. Blasiak, J. Bredenbeck, M. Cho, J.-H. Choi, S. A. Corcelli, A. G. Dijkstra, C.-J. Feng, S. Garrett-Roe, N.-H. Ge, M. W. D. Hanson-Heine, J. D. Hirst, T. L. C. Jansen, K. Kwac, K. J. Kubarych, C. H. Londergan, H. Maekawa, M. Reppert, S. Saito, S. Roy, J. L. Skinner, G. Stock, J. E. Straub, M. C. Thielges, K. Tominaga, A. Tokmakoff, H. Torii, L. Wang, L. J. Webb, and M. T. Zanni, Chem. Rev. **120** (2020), 10.1021/acs.chemrev.9b00813.
- [8] H. J. Kimble, Nature **453**, 1023 (2008).
- [9] S. Ritter, C. Nölleke, C. Hahn, A. Reiserer, A. Neuzner, M. Uphoff, M. Mücke, E. Figueroa, J. Bochmann, and G. Rempe, Nature 484, 195 (2012).
- [10] Z. Yang and W. Xiong, Adv. Quantum Technol. 5, 2100163 (2022).
- [11] E. Orgiu, J. George, J. A. Hutchison, E. Devaux, J. F. Dayen, B. Doudin, F. Stellacci, C. Genet, J. Schachen-

mayer, C. Genes, G. Pupillo, P. Samorì, and T. W. Ebbesen, Nature Mater. 14, 1123 (2015).

- [12] G. Lerario, D. Ballarini, A. Fieramosca, A. Cannavale, A. Genco, F. Mangione, S. Gambino, L. Dominici, M. De Giorgi, G. Gigli, and D. Sanvitto, Light: Science & Applications 6, e16212 (2017).
- [13] G. G. Rozenman, K. Akulov, A. Golombek, and T. Schwartz, ACS Photonics 5, 105 (2018).
- [14] S. Hou, M. Khatoniar, K. Ding, Y. Qu, A. Napolov, V. M. Menon, and S. R. Forrest, Adv. Mater. **32**, 2002127 (2020).
- [15] N. Krainova, A. J. Grede, D. Tsokkou, N. Banerji, and N. C. Giebink, Phys. Rev. Lett. **124**, 177401 (2020).
- [16] M. Balasubrahmaniyam, A. Simkhovich, A. Golombek, G. Sandik, G. Ankonina, and T. Schwartz, Nature Mater. 22, 338 (2023).
- [17] K. Akulov, D. Bochman, A. Golombek, and T. Schwartz, J. Phys. Chem. C **122**, 15853 (2018).
- [18] B. Xiang, R. F. Ribeiro, M. Du, L. Chen, Z. Yang, J. Wang, J. Yuen-Zhou, and W. Xiong, Science **368**, 665 (2020).
- [19] A. Yariv, Y. Xu, R. K. Lee, and A. Scherer, Opt. Lett. 24, 711 (1999).
- [20] Y.-C. Liu, B.-B. Li, and Y.-F. Xiao, Nanophotonics 6, 789 (2017).
- [21] J. A. Hutchison, T. Schwartz, C. Genet, E. Devaux, and T. W. Ebbesen, Angew. Chem. Int. Ed. **51**, 1592 (2012).
- [22] X. Shi, K. Ueno, T. Oshikiri, Q. Sun, K. Sasaki, and H. Misawa, Nature Nanotechnol. 13, 953 (2018).

- [23] B. Munkhbat, M. Wersäll, D. G. Baranov, T. J. Antosiewicz, and T. Shegai, Sci. Adv. 4, eaas9552 (2018).
- [24] P. M. Kraus, M. Zürch, S. K. Cushing, D. M. Neumark, and S. R. Leone, Nat. Rev. Chem. 2, 82 (2018).
- [25] J. R. Rouxel, Y. Zhang, and S. Mukamel, Chem. Sci. 10, 898 (2019).
- [26] K. E. Dorfman, F. Schlawin, and S. Mukamel, Rev. Mod. Phys. 88, 045008 (2016).
- [27] B. Gu and S. Mukamel, ACS Photonics 9, 938 (2022).
- [28] S. Mukamel, Principles of Nonlinear Optical Spectroscopy, Vol. 6 (Oxford University Press, 1995).
- [29] U. Harbola, B. K. Agarwalla, and S. Mukamel, J. Chem. Phys. 141, 074107 (2014).
- [30] B. K. Agarwalla, U. Harbola, W. Hua, Y. Zhang, and S. Mukamel, The Journal of Chemical Physics 142, 212445 (2015).
- [31] H. Imada, K. Miwa, M. Imai-Imada, S. Kawahara, K. Kimura, and Y. Kim, Nature 538, 364 (2016).
- [32] H. Imada, K. Miwa, M. Imai-Imada, S. Kawahara, K. Kimura, and Y. Kim, Phys. Rev. Lett. **119**, 013901 (2017).
- [33] K. Miwa, H. Imada, M. Imai-Imada, K. Kimura, M. Galperin, and Y. Kim, Nano Lett. 19, 2803 (2019).
- [34] R. Chikkaraddy, R. Arul, L. A. Jakob, and J. J. Baumberg, Nat. Photonics 17, 865 (2023).
- [35] Z. Ioffe, T. Shamai, A. Ophir, G. Noy, I. Yutsis, K. Kfir, O. Cheshnovsky, and Y. Selzer, Nature Nanotechnol. 3, 727 (2008).
- [36] D. R. Ward, N. J. Halas, J. W. Ciszek, J. M. Tour, Y. Wu, P. Nordlander, and D. Natelson, Nano Lett. 8, 919 (2008).
- [37] T. Shamai and Y. Selzer, Chem. Soc. Rev. 40, 2293 (2011).
- [38] D. R. Ward, D. A. Corley, J. M. Tour, and D. Natelson, Nature Nanotechnol. 6, 33 (2011).
- [39] R. B. Jaculbia, H. Imada, K. Miwa, T. Iwasa, M. Takenaka, B. Yang, E. Kazuma, N. Hayazawa, T. Taketsugu, and Y. Kim, Nature Nanotechnol. 15, 105 (2020).
- [40] R. Chikkaraddy, B. de Nijs, F. Benz, S. J. Barrow, O. A. Scherman, E. Rosta, A. Demetriadou, P. Fox, O. Hess,

and J. J. Baumberg, Nature 535, 127 (2016).

- [41] F. Benz, M. K. Schmidt, A. Dreismann, R. Chikkaraddy, Y. Zhang, A. Demetriadou, C. Carnegie, H. Ohadi, B. de Nijs, R. Esteban, J. Aizpurua, and J. J. Baumberg, Science **354**, 726 (2016).
- [42] Y. Gao and M. Galperin, J. Chem. Phys. 144, 174113 (2016).
- [43] Y. Gao and M. Galperin, J. Chem. Phys. 144, 244106 (2016).
- [44] S. Mukamel, A. Li, and M. Galperin, J. Chem. Phys. 158, 154106 (2023).
- [45] N. Seshadri, A. Li, and M. Galperin, J. Chem. Phys. 160, 044116 (2024).
- [46] M. Galperin, Chem. Soc. Rev. 46, 4000 (2017).
- [47] H. K. Yadalam, S. Mitra, and U. Harbola, J. Phys. Chem. A 123, 10594 (2019).
- [48] S. Mukamel and M. Galperin, J. Phys. Chem. C 123, 29015 (2019).
- [49] G. Baym and L. P. Kadanoff, Phys. Rev. 124, 287 (1961).
- [50] G. Baym, Phys. Rev. **127**, 1391 (1962).
- [51] J. K. Viljas, J. C. Cuevas, F. Pauly, and M. Häfner, Phys. Rev. B 72, 245415 (2005).
- [52] L. de la Vega, A. Martïn-Rodero, N. Agraït, and A. L. Yeyati, Phys. Rev. B 73, 075428 (2006).
- [53] H. K. Yadalam, S. Mukamel, and U. Harbola, J. Phys. Chem. Lett. 11, 1762 (2020).
- [54] G. Cohen and M. Galperin, J. Chem. Phys. 152, 090901 (2020).
- [55] G. Stefanucci and R. van Leeuwen, Nonequilibrium Many-Body Theory of Quantum Systems. A Modern Introduction. (Cambridge University Press, 2013).
- [56] P. Talkner, Annals of Physics 167, 390 (1986).
- [57] G. Guarnieri, A. Smirne, and B. Vacchini, Phys. Rev. A 90, 022110 (2014).
- [58] J. Jin, C. Karlewski, and M. Marthaler, New J. Phys. 18, 083038 (2016).
- [59] M. Cosacchi, T. Seidelmann, M. Cygorek, A. Vagov, D. Reiter, and V. Axt, Phys. Rev. Lett. **127**, 100402 (2021).