

Multiple Imputation for Nonresponse in Complex Surveys Using Design Weights and Auxiliary Margins

Kewei Xu*

Jerome P. Reiter†

Abstract

Survey data typically have missing values due to unit and item nonresponse. Sometimes, survey organizations know the marginal distributions of certain categorical variables in the survey. As shown in previous work, survey organizations can leverage these distributions in multiple imputation for nonignorable unit nonresponse, generating imputations that result in plausible completed-data estimates for the variables with known margins. However, this prior work does not use the design weights for unit nonrespondents; rather, it relies on a set of fabricated weights for these units. We extend this previous work to utilize the design weights for all sampled units. We illustrate the approach using simulation studies.

Key Words: Item; Missing; Nonignorable; Unit

1 Introduction

Survey data usually suffer from both unit and item nonresponse. As a result, survey organizations have to make strong assumptions about the reasons for missingness, for example, the

*Department of Statistical Science, Box 90251, Duke University, Durham, NC 27708-0251

†Department of Statistical Science, 202 Old Chemistry Building, Duke University, Durham, NC 27708-0251

data are missing at random. One way to lessen reliance on assumptions is to utilize information in auxiliary data sources. For example, and pertinent to the setting of our work, survey organizations may know the population percentages or totals of some categorical variables in the survey. These could be available from recent censuses, administrative databases, or other high quality surveys (Sadinle and Reiter, 2017). Indeed, such information is frequently used in methods for handling survey nonresponse, such as calibration and raking.

In this short note, we consider settings where analysts seek to integrate auxiliary information on marginal distributions in multiple imputation for nonresponse (Rubin, 1987) in complex surveys. We build on the missing data with auxiliary margins, or MD-AM, framework introduced by (Akande et al., 2021) and extended to complex surveys by Akande and Reiter (2022), (Tang et al., 2024), and (Yang and Reiter, 2025). The latter three works impute missing values in ways that ensure completed-data, survey-weighted inferences are plausible given the known margins. However, these works make a simplifying condition on the unit nonrespondents, namely that the analyst does not use their design weights and replaces them with a convenient constant. However, this creates difficulties for design-based analyses, as the resulting weights for the full sample do not correspond to the design. Further, treating all unit nonrespondents exchangeably discards information that potentially could inform imputations.

We propose MD-AM models that allow use of the design-based weights for all sampled units. The basic idea is as follows. We first impute all missing values due to item nonresponse using multiple imputation by chained equations (Van Buuren, 2018; Raghunathan et al., 2001). In each partially completed dataset, we then impute the unit nonrespondents' values of the categorical variables that have auxiliary margins. To do so, we specify initial probabilities for imputations based on logistic regressions estimated using the completed data. We adjust these probabilities so that, in expectation, imputations result in plausible Horvitz and Thompson (1952) estimates in the completed data given the known margins, using the design weights for all sampled units. Finally, for the variables without auxiliary

margins, we use random hot deck imputation proposed by Yang and Reiter (2025).

The remainder of this article is organized as follows. Section 2 provides a brief review of the hybrid missing MD-AM model (Tang et al., 2024; Yang and Reiter, 2025), which we extend to incorporate design weights for unit nonrespondents. Section 3 introduces the methodology. Section 4 illustrates the methods using simulation studies.

2 Hybrid Missingness MD-AM Modeling

In reviewing the hybrid missingness MD-AM model, we closely follow the notation in Yang and Reiter (2025). Let \mathcal{P} be a finite population comprising N units. For $i = 1, \dots, N$, let $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})$ denote the p survey variables for unit i ; let $I_i = 1$ if unit i is selected for inclusion in the survey and $I_i = 0$ otherwise; let $\pi_i = \Pr(I_i = 1)$ represent the probability that unit i is selected into the survey; and, let $w_i^d = 1/\pi_i$ be its design weight. Let $n = \sum_{i=1}^N I_i$ be the intended sample size of the survey. We refer to the sampled units as \mathcal{S} .

Let U be the unit nonresponse indicator so that, for each unit $i \in \mathcal{S}$, $U_i = 1$ if the unit provides no responses to any survey questions and $U_i = 0$ otherwise. Let $\mathbf{R} = (R_1, \dots, R_p)$ be item nonresponse indicators corresponding to (X_1, \dots, X_p) so that, for any unit i with $U_i = 0$ and any variable X_j , $R_{ij} = 1$ if unit i does not respond to question X_j and $R_{ij} = 0$ otherwise. When $U_i = 1$, (R_{i1}, \dots, R_{ij}) are undefined.

The survey organization has auxiliary information \mathcal{A} on the marginal distributions of a subset of the p survey variables. For $j = 1, \dots, p$, let \mathcal{A}_j comprise the known marginal in \mathcal{P} for X_j , when it exists. We write $X_j \in \mathcal{A}$ whenever X_j has a known margin in \mathcal{A} . For convenience, we presume each \mathcal{A}_j comprises totals; incorporating percentages is a trivial modification. For any categorical X_j taking on levels $c = \{1, \dots, m_j\}$ and $X_j \in \mathcal{A}$, let $T_{jc} = \sum_{i=1}^N I(x_{ij} = c)$ be the total number of units in \mathcal{P} at level c . Here, $I(\cdot) = 1$ when the condition inside the parenthesis is true and $I(\cdot) = 0$ otherwise.

The hybrid missingness MD-AM model specifies a joint distribution of $(\mathbf{X}, \mathbf{R}, U)$. It

uses a pattern mixture model for unit nonresponse coupled with selection models for item nonresponse. Specifically, the model presumes $U \sim \text{Bernoulli}(\pi_u)$, where $\pi_u = \Pr(U = 1)$ is the marginal probability of unit nonresponse. For any $j = 1, \dots, p$, let $g_j(-)$ represent the model for X_j given X_1, \dots, X_{j-1} (and possibly the weights W), and let Ω_j and θ_j be the corresponding model parameters. For example, $g_j(-)$ could be a logistic regression of X_j on some function of X_1, \dots, X_{j-1} , possibly including interactions, as well as a main effect for U when $X_j \in \mathcal{A}$. Thus, for $\mathbf{X}|U$, the hybrid missingness MD-AM model uses

$$X_1 | U \sim g_1(\Omega_1, \theta_1 UI(X_1 \in \mathcal{A})) \quad (2.1)$$

$$X_j | X_1, \dots, X_{j-1}, U \sim g_j(X_1, \dots, X_{j-1}, \Omega_j, \theta_j UI(X_j \in \mathcal{A})), \text{ for } j = 2, \dots, p. \quad (2.2)$$

Because the distribution for any $X_j \in \mathcal{A}$ differs for unit nonrespondents and respondents when $\theta_j \neq 0$, the model encodes potentially missing not at random mechanisms for unit nonresponse. The key identifying assumption is that the predictor function for any $X_j \in \mathcal{A}$ not include interactions between U and elements of (X_1, \dots, X_{j-1}) . This assumption is a version of the additive nonignorable missingness mechanism (Hirano et al., 2001; Sadinle and Reiter, 2019).

For the model for each R_j , let $h_j(-)$ be some predictor function that excludes the corresponding X_j but may include main and interaction effects involving other variables in \mathbf{X} . For $j = 1, \dots, p$, we have

$$R_j | X_1, \dots, X_p, U = 0 \sim \text{Bernoulli}(\pi_{R_j}), \quad \text{logit}(\pi_{R_j}) = h_j(X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_p, \Phi_j). \quad (2.3)$$

The models in (2.3) encode itemwise conditionally independent nonresponse (ICIN) mechanisms (Sadinle and Reiter, 2017), which are known to have identifiable parameters.

The estimation algorithm in Tang et al. (2024) does not use the design weights for unit nonrespondents. Instead, they replace the weights for unit nonrespondents with a constant

that makes $\sum_{i=1}^N w_i = N$, so that they base imputations and inferences on

$$w_i = \begin{cases} w_i^d & \text{if } U_i = 0 \\ \frac{N - \sum_{j \in \mathcal{S}} w_j^d}{\sum_{j \in \mathcal{S}} U_j} & \text{if } U_i = 1. \end{cases} \quad (2.4)$$

For any unit $i \in \mathcal{S}$ and for all j , let $x_{ij}^* = x_{ij}$ when $(R_{ij} = 0, U_i = 0)$, and let x_{ij}^* be an imputed value when $R_{ij} = 1$ or $U_i = 1$. For any $X_j \in \mathcal{A}$, Akande and Reiter (2022), Tang et al. (2024), and Yang and Reiter (2025) suppose the imputations to adhere to (2.1) – (2.3), with the additional constraint that

$$\hat{T}_{jc}^* = \sum_{i \in \mathcal{S}} w_i I(x_{ij}^* = c) \sim N(T_{jc}, V_{jc}), \quad (2.5)$$

Here, V_{jc} is set by the analyst and determines how closely \hat{T}_{jc}^* matches T_{jc} in any completed dataset. As we also assume (2.5), we discuss setting V_{jc} in Section 3.

3 Methodology

We now present our modifications of the hybrid missingness MD-AM model to use the design weights for all sampled individuals, including unit nonrespondents. For convenience, we presume that $X_j \in \mathcal{A}$ for $j = 1, \dots, k < p$, and the remaining $p - k$ variables do not have auxiliary margins.

3.1 Variables with Item Nonresponse

We first create multiple imputations for all values of x_{ij} missing due to item nonresponse for units with $U_i = 0$. Following Yang and Reiter (2025), we use a multiple imputation by chained equations (MICE) algorithm (Van Buuren, 2018; Raghunathan et al., 2001) estimated using the units with $U_i = 0$. The specification of the MICE algorithm does not require any extraordinary considerations. In our simulations, we use the “mice” package in

R, following its default settings including the ordering of variables. Each conditional model is a function of all other survey variables except X_j , including possibly interaction terms. The conditional models also could include subsets of $(R_1, \dots, R_{j-1}, R_{j+1}, \dots, R_p)$ as predictors; these are fully observed for units with $U_i = 0$. We run the MICE procedure to create L completed datasets for the units with $U_i = 0$.

3.2 Variables with Margins for Unit Nonrespondents

After imputing values for item nonresponse, in each completed dataset we impute the unit nonrespondents' values for all $X_j \in \mathcal{A}$. The analyst orders these variables based on ease of modeling. For notational convenience, we suppose that X_1 is imputed first, X_2 is imputed second, and so on sequentially until we impute all $X_j \in \mathcal{A}$.

To begin, for each unit i with $U_i = 1$ and in each of the L completed datasets, the analyst sets an initial probability distribution for imputing x_{i1} . For example, in each completed dataset, the analyst estimates a (multinomial) logistic regression of X_1 on some function of W , such as a spline (Chen et al., 2010), and uses the predicted probabilities derived from the estimated model. Alternatively, the analyst could regress X_1 on an intercept only or compute marginal probabilities of X_1 using survey-weighted ratio estimators. For these two alternatives, the initial probabilities are identical for all unit nonrespondents. This approach may be especially appropriate when the analyst does not have strong reason to expect associations between the weights and the missing values of X_1 for unit nonrespondents. Regardless of how it is derived, we refer to the initial distribution for imputing x_{i1} as the working distribution, notated as $\{p_{i1c} : c = 1, \dots, m_1; U_i = 1; i \in \mathcal{S}\}$. Because of imputation for item nonresponse, the working distributions for each completed dataset may differ; however, for convenience, we forego notation designating the completed dataset. The analysts performs the computations of this section for each of the L completed datasets.

Imputing each missing x_{i1} using its working distribution does not utilize the information in \mathcal{A}_1 . Thus, the resulting imputations could generate completed-data Horvitz and Thompson

(1952) estimates that are far from the known totals for some levels of X_1 . We therefore modify $\{p_{i1c}\}$ so that the resulting imputations result in reasonable completed-data point estimates of these totals.

To do so, we first simulate plausible values of \hat{T}_{1c} for each level c by appealing to large-sample central limit theorems. For $c = 1, \dots, (m_1 - 1)$, we sample \hat{T}_{1c} from $N(T_{1c}, V_{1c})$ and set $\hat{T}_{1m_1} = N - \sum_{c=1}^{m_1-1} \hat{T}_{1c}$. We then find modified probabilities $\{\tilde{p}_{i1c} : c = 1, \dots, m_1; U_i = 1; i \in \mathcal{S}\}$ so that, for each $c = 1, \dots, m_1$, the expectation of each completed-data estimator \hat{T}_{1c}^* from (2.5) over imputations for unit nonresponse approximately equals \hat{T}_{1c} . Put another way, recognizing that

$$\begin{aligned} E\left(\sum_{i \in \mathcal{S}} w_i I(x_{ij}^* = c) I(U_i = 1)\right) &= \sum_{i \in \mathcal{S}} w_i I(x_{ij}^* = c) I(U_i = 0) + E\left(\sum_{i \in \mathcal{S}} w_i I(x_{ij}^* = c) I(U_i = 1)\right) \\ &= \sum_{i \in \mathcal{S}} w_i I(x_{ij}^* = c) I(U_i = 0) + \sum_{i \in \mathcal{S}} w_i \tilde{p}_{i1c} I(U_i = 1), \end{aligned} \quad (3.1)$$

we want

$$\sum_{i \in \mathcal{S}} w_i \tilde{p}_{i1c} I(U_i = 1) = \hat{T}_{1c} - \sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = c) I(U_i = 0). \quad (3.2)$$

To keep the imputation probabilities tied to the working distribution, for $c = 1, \dots, m_1 - 1$, we set $\tilde{p}_{i1c} = f_{1c} p_{i1c}$, where each constant $f_{1c} > 0$ is given by

$$f_{1c} = \frac{\hat{T}_{1c} - \sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = c, U_i = 0)}{\sum_{i \in \mathcal{S}} p_{i1c} w_i I(U_i = 1)}. \quad (3.3)$$

We impute x_{i1}^* for each unit with $U_i = 1$ using a random draw from the adjusted probability mass function, $\{\tilde{p}_{i1c} : c = 1, \dots, m_1\}$, where $\tilde{p}_{i1m_1} = 1 - \sum_{c=1}^{m_1-1} f_{1c} p_{i1c}$. If $\sum_{c=1}^{m_1-1} f_{1c} p_{i1c} > 1$, we compute (3.3) for $c = 1, \dots, m_1$ and set $\tilde{p}_{i1c} = f_{1c} p_{i1c} / \sum_{c=1}^{m_1} f_{1c} p_{i1c}$.

When each p_{i1c} equals some common value p_{1c} for all units with $U_i = 1$, each $\tilde{p}_{i1c} =$

$f_{1c}p_{1c} = \tilde{p}_{1c}$ is given conveniently by

$$\tilde{p}_{1c} = \frac{\hat{T}_{1c} - \sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = c, U_i = 0)}{\sum_{i \in \mathcal{S}} w_i I(U_i = 1)}. \quad (3.4)$$

We now turn to imputing $X_2 \in \mathcal{A}$ for unit nonrespondents. We follow a similar strategy: in each completed dataset, set each unit nonrespondent's working probabilities for imputation of x_{i2}^* given x_{i1}^* , then adjust them to make $(\hat{T}_{21}^*, \dots, \hat{T}_{2m_2}^*)$ approximately match a sampled value of $(\hat{T}_{21}, \dots, \hat{T}_{2m_2})$.

Let $p_{i2c} = \Pr(X_{i2} = c | x_{i1}^* = d, w_i)$ be the working probability of $X_2 = c$ for unit i , given its imputed $x_{i1}^* = d$ and w_i . Following the MD-AM model in (2.2), we determine these from a (multinomial) logistic regression of X_2 on X_1 and possibly some function of W , with coefficients estimated from the completed data for units with $U_i = 0$. If we disregard W , $p_{i2c} = \Pr(X_2 = c | X_1 = d)$ for $c = 1, \dots, m_2$ and $d = 1, \dots, m_1$; that is, the conditional probability is the same for all units with a common value of $x_{i1}^* = d$.

For $c = 1, \dots, m_2 - 1$, we sample a plausible value of \hat{T}_{2c} by drawing from a $N(T_{2c}, V_{2c})$, setting the estimated total for the final level as $\hat{T}_{2m_2} = N - \sum_{c=1}^{m_2-1} \hat{T}_{2c}$. Akin to (3.3), we then find suitable constants $f_{2c} > 0$ to adjust each p_{i2c} . We have

$$f_{2c} = \frac{\hat{T}_{2c} - \sum_{i \in \mathcal{S}} w_i I(x_{i2}^* = c, U_i = 0)}{\sum_{i \in \mathcal{S}} p_{i2c} w_i I(U_i = 1)}. \quad (3.5)$$

We impute each x_{i2}^* given x_{i1}^* for units with $U_i = 1$ using a random draw from the adjusted probability mass function $\{\tilde{p}_{i2c} : c = 1, \dots, m_2\}$, where $\tilde{p}_{i2c} = f_{2c}p_{i2c}$ for $c = 1, \dots, m_2 - 1$ and $\tilde{p}_{i2m_2} = 1 - \sum_{c=1}^{m_2-1} f_{2c}p_{i2c}$. If $\sum_{c=1}^{m_2-1} f_{2c}p_{i2c} > 1$, we compute (3.5) for $c = 1, \dots, m_2$ and set $\tilde{p}_{i2c} = f_{2c}p_{i2c} / \sum_{c=1}^{m_2} f_{2c}p_{i2c}$. This process ensures the completed-data Horvitz and Thompson (1952) estimators approximately match the sampled $(\hat{T}_{21}, \dots, \hat{T}_{2m_2})$ in expectation, while also incorporating relationships between X_2 and (X_1, W) implied in the working probabilities. We independently repeat the estimation and imputation process in each completed dataset.

When $\{p_{i2c}\}$ derives from a logistic regression of X_2 on X_1 (and possibly W), using

$\{\tilde{p}_{i2c}\}$ is equivalent to adjusting the intercept in that estimated regression, leaving other coefficients alone. Essentially, this implies a model where the log-odds for X_2 are the same for unit respondents and nonrespondents, per the hybrid missingness MD-AM assumption in (2.2) of no interaction between U and X_2 .

When disregarding W in the logistic regression for X_2 on X_1 —so that within any completed dataset, $p_{i2c} = \Pr(X_2 = c|x_1 = d, U = 1)$ for all unit nonrespondents with $x_{i1}^* = d$ in that dataset—we can find $\{\tilde{p}_{2c}\}$ by solving a system of equations implied by (2.2). Specifically, we encode the constant log-odds assumption as a set of $(m_1 - 1)(m_2 - 1)$ equations. For $c = 2, \dots, m_2 - 1$ and $d = 2, \dots, m_1 - 1$, we have

$$\begin{aligned} & \log\left(\frac{\Pr(X_2=c|X_1=d,U=1)}{\Pr(X_2=1|X_1=d,U=1)}\right) - \log\left(\frac{\Pr(X_2=c|X_1=d,U=1)}{\Pr(X_2=c|X_1=1,U=1)}\right) \\ &= \log\left(\frac{\Pr(X_2=c|X_1=d,U=0)}{\Pr(X_2=1|X_1=d,U=0)}\right) - \log\left(\frac{\Pr(X_2=c|X_1=d,U=0)}{\Pr(X_2=c|X_1=1,U=0)}\right) \end{aligned} \quad (3.6)$$

We can estimate the conditional probabilities for units with $U_i = 0$ via survey-weighted ratio estimators using the completed data; for example,

$$\Pr(X_2 = c|X_1 = d, U = 0) = \frac{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = d, x_{i2}^* = c, U_i = 0)}{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = d, U_i = 0)}. \quad (3.7)$$

We also encode the condition (2.5) on $(\hat{T}_{21}^*, \dots, \hat{T}_{2m_2}^*)$ as a set of m_2 equations. We have

$$\sum_{d=1}^{m_1} \sum_{i \in \mathcal{S}} \Pr(X_2 = c|X_1 = d, U = 1) w_i I(x_{i2}^* = c, x_{i1}^* = d, U_i = 1) = \hat{T}_{2c} - \sum_{i \in \mathcal{S}} w_i I(x_{i2}^* = c, U_i = 0). \quad (3.8)$$

In total, we have $m_1(m_2 - 1)$ equations to solve for the $m_1(m_2 - 1)$ conditional probabilities, $\{\Pr(X_2 = c|X_1 = d, U = 1) : c = 2, \dots, m_2; d = 1, \dots, m_1\}$, which we use for imputation.

We now describe the computations for (3.6) and (3.8) for binary X_2 and X_1 . We first

compute

$$\Pr(X_2 = 1|X_1 = 1, U = 0) = \frac{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 1, x_{i2}^* = 1, U_i = 0)}{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 1, U_i = 0)} \quad (3.9)$$

$$\Pr(X_2 = 1|X_1 = 0, U = 0) = \frac{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 0, x_{i2}^* = 1, U_i = 0)}{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 0, U_i = 0)}. \quad (3.10)$$

To ease notation, define

$$\hat{\beta}_{1,obs} = \text{logit}(\Pr(X_2 = 1|X_1 = 1, U = 0)) - \text{logit}(\Pr(X_2 = 1|X_1 = 0, U = 0)) \quad (3.11)$$

$$\hat{\beta}_{0,obs} = \text{logit}(\Pr(X_2 = 1|X_1 = 1, U = 0)) - \hat{\beta}_{1,obs} \quad (3.12)$$

Define the unobserved quantities $\hat{\beta}_{1,miss}$ and $\hat{\beta}_{0,miss}$ for the unit nonrespondents analogously, replacing $U_i = 0$ with $U_i = 1$ in (3.11) and (3.12).

We set the log odds ratios equal for the unit respondents and nonrespondents, that is, we set $\hat{\beta}_{1,miss} = \hat{\beta}_{1,obs}$. To determine $\hat{\beta}_{0,miss}$, we numerically solve the system of nonlinear equations using the modified Powell method implemented in the package described by Hasselman (2018). We have

$$\frac{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 1, U_i = 1)}{1 + \exp(-(\hat{\beta}_{0,miss} + \hat{\beta}_{1,miss}))} + \frac{\sum_{i \in \mathcal{S}} w_i I(x_{i1}^* = 0, U_i = 1)}{1 + \exp(-\hat{\beta}_{0,miss})} = \hat{T}_2 - \sum_{i \in \mathcal{S}} w_i I(x_{i2}^* = 1, U_i = 0).$$

The imputation probabilities are then

$$\Pr(X_2 = 1|x_{i1}^* = 1, U_i = 1) = \frac{1}{1 + \exp[-(\hat{\beta}_{0,miss} + \hat{\beta}_{1,miss})]} \quad (3.13)$$

$$\Pr(X_2 = 1|x_{i1}^* = 0, U_i = 0) = \frac{1}{1 + \exp(-\hat{\beta}_{0,miss})}. \quad (3.14)$$

When $k > 2$, we can apply the process used in (3.5) to each $X_j \in \mathcal{A}$. For example, if $X_3 \in \mathcal{A}$, the analyst specifies a set of working probabilities, $p_{i3c} = \Pr(X_{i3} = c|x_{i1}^*, x_{i2}^*, w_i)$ for $c = 1, \dots, m_3$ via a (multinomial) logistic regression of X_3 on (X_1, X_2) and possibly W . The analyst samples values of $(\hat{T}_{31}, \dots, \hat{T}_{3m_3})$ and finds the values of f_{3c} using expressions

analogous to (3.5), replacing quantities based on X_2 with those based on X_3 . When the working probabilities are identical across nonresponding units, analysts can solve a system of equations akin to those in (3.6) and (3.8).

3.3 Variables Without Margins for Unit Nonrespondents

After creating L partially completed datasets using Section 3.1 and 3.2, we impute values x_{ij}^* for unit nonrespondents for $X_j \notin \mathcal{A}$. We use the random hot deck imputation procedure developed by Yang and Reiter (2025). Here we briefly summarize the procedure, closely following their presentation.

For any unit $i \in \mathcal{S}$ with $U_i = 0$, in any completed dataset, let $\mathbf{x}_i^{\mathcal{A}} = \{x_{ij} : X_j \in \mathcal{A}; j = 1, \dots, p; U_i = 0\}$ be the values of the (possibly imputed) survey variables in the completed data for those $X_j \in \mathcal{A}$. Similarly, for any unit $i' \in \mathcal{S}$ with $U_{i'} = 1$, let $\mathbf{x}_{i'}^{\mathcal{A}*} = \{x_{i'j}^* : X_j \in \mathcal{A}; j = 1, \dots, p; U_{i'} = 1\}$. For each unit i' with $U_{i'} = 1$, in each completed dataset we construct its donor set, $\mathcal{D}_{i'} = \{(x_{i1}, \dots, x_{ip}) : \mathbf{x}_i^{\mathcal{A}} = \mathbf{x}_{i'}^{\mathcal{A}*}, U_i = 0, i \in \mathcal{S}\}$. We randomly sample one record i from $\mathcal{D}_{i'}$ and append its values of $\{(x_{i1}, \dots, x_{ik}) : X_j \notin \mathcal{A}\}$ to $\mathbf{x}_{i'}^{\mathcal{A}*}$ to make the full imputation $\mathbf{x}_{i'}^*$ for unit i' . We repeat this process for all units in \mathcal{S} with $U_{i'} = 1$, resulting in a completed dataset with values for all n units in \mathcal{S} . We repeat this process in each of the L datasets.

4 Simulation Studies

To demonstrate the methodology, we mimic the simulation study used by Yang and Reiter (2025). They found that, by leveraging auxiliary margins, multiple imputation via the hybrid missingness MD-AM model can offer more accurate inferences than multiple imputation that does not utilize \mathcal{A} . However, they presume design weights are unused or unavailable for unit nonrespondents, whereas we use them in the imputation procedure.

We generate a population \mathcal{P} comprising $N = 3,373,378$ units. This is the number of

individuals in the 2022 American Community Survey data used by Yang and Reiter (2025). We make one of the variables in this file a size variable $Z = (z_1, \dots, z_N)$; this is observed for all N individuals. To generate a \mathcal{S} , we sample each record from \mathcal{P} independently with $\pi_i = 1/10z_i$, where $i = 1, \dots, N$. This Poisson sampling design results in approximately $n = 6000$ units in any \mathcal{S} , although the precise number varies. Let $W = 10Z$ be the survey weights.

Following Yang and Reiter (2025), for each unit $i = 1, \dots, N$, we generate four binary variables $(x_{i1}, x_{i2}, x_{i3}, x_{i4})$ and two continuous variables (x_{i5}, x_{i6}) . We make X_1 depend on W to ensure associations between the weights and the survey variables. The data generation model is displayed in (4.1)–(4.5).

$$U \sim \text{Bernoulli}(\pi_U) \quad \text{logit}(\pi_U) = \nu_0 \quad (4.1)$$

$$X_1|U \sim \text{Bernoulli}(\pi_{x_1}) \quad \text{logit}(\pi_{x_1}) = \omega_{10} + \omega_{11}W + \theta_1U \quad (4.2)$$

$$X_2|X_1, U \sim \text{Bernoulli}(\pi_{x_2}) \quad \text{logit}(\pi_{x_2}) = \omega_{20} + \omega_{21}X_1 + \theta_2U \quad (4.3)$$

$$X_j|X_1, \dots, X_{j-1}, U \sim \text{Bernoulli}(\pi_{x_j}) \quad \text{logit}(\pi_{x_j}) = \omega_{j0} + \sum_{k=1}^{j-1} \omega_{jk}X_k \quad \text{for } j = 3, 4 \quad (4.4)$$

$$X_j|X_1, \dots, X_{j-1}, U \sim \text{Normal}(\omega_{j0} + \sum_{k=1}^{j-1} \omega_{jk}X_k, \sigma_j^2) \quad \text{for } j = 5, 6. \quad (4.5)$$

We set $\nu_0 = -1.2$ for a unit nonresponse rate around 23%. We consider $\theta_1 \in \{-2, -0.5\}$. The marginal distribution of X_1 for unit respondents and nonrespondents is substantially different when $\theta_1 = -2$ and modestly different when $\theta_1 = -0.5$. We use $\theta_2 = -2$, which ensures the unit respondents and nonrespondents have different marginal distributions of X_2 . The remaining parameter values are available in Appendix A.

We independently sample 500 datasets using Poisson sampling. For each sample \mathcal{S} , we generate the unit nonrespondents by sampling Bernoulli random draws of U_i , where $i \in \mathcal{S}$, from the distribution of $U|X_1, X_2$ implied by (4.1)–(4.3). We make \mathbf{x}_i completely missing for any unit in \mathcal{S} where the drawn $U_i = 1$. For item nonresponse, we use the ICIN mechanisms

defined in (4.6) for X_j where $j \in \{2, 3, 4, 6\}$, and we let X_1 and X_5 be fully observed.

$$R_j | X_1, X_2, X_3, X_4, X_5, X_6, U = 0 \sim \text{Bernoulli}(\pi_{R_j}), \quad \text{logit}(\pi_{R_j}) = \phi_{j0} + \sum_{k \neq j} \phi_{jk} X_k. \quad (4.6)$$

We blank every x_{ij} for which the sampled $R_{ij} = 1$. We set the parameters in (4.6) so that approximately 25% of each eligible survey variable is missing for units with $U_i = 0$. Specific parameter values are available in Appendix A.

We presume that $T_1 = \sum_{i=1}^N x_{i1}$ and $T_2 = \sum_{i=1}^N x_{i2}$ are known and use them as \mathcal{A} . We use a bespoke implementation of the “mice” package in R (Van Buuren, 2018) to generate $L = 10$ completed data sets for item nonresponse. As values for V_1 and V_2 , we use MICE to make one completed dataset for all of \mathcal{S} and compute the expressions for the usual design-based estimators of the variances of \hat{T}_1 and \hat{T}_2 under Poisson sampling. We create donor pools for the hot deck per Section 3.3 by matching on (X_1, X_2) .

After generating $L = 10$ completed datasets, we use the inferential methods in Rubin (1987) for the estimands in Yang and Reiter (2025), including marginal totals for (X_1, \dots, X_6) and several probabilities involving multiple categorical variables. In each completed dataset, we compute Horvitz and Thompson (1952) point and variance estimators for a Poisson sampling design based on the design weights for all sampled records. The results will be forthcoming in a revised version of this manuscript.

A Simulation Study Parameters

This section presents the parameter values used in the simulation studies of Section 4, which are identical to those in Yang and Reiter (2025). We set $\nu_u = -1.2$; $(\omega_{10}, \omega_{11}) = (0.06, -0.0002)$ and $\theta_1 \in (-2, -0.5)$; $(\omega_{20}, \omega_{21}) = (0.2, 0.4)$ and $\theta_2 = -2$; $(\omega_{30}, \omega_{31}, \omega_{32}) = (0.2, 0.3, 0.1)$; $(\omega_{40}, \dots, \omega_{43}) = (0.2, 0.4, 0.4, 0.1)$; $(\omega_{50}, \dots, \omega_{54}) = (0.4, 1.2, -0.9, 0.1, 0.2)$ and $\sigma_5 = 0.5$; $(\omega_{60}, \dots, \omega_{65}) = (0.4, 1.2, -0.9, 0.1, -0.1, 0.1)$ and $\sigma_6 = 0.5$; $(\phi_{j0}, \dots, \phi_{j5}) = (-1.4, 0.1, 0.1, 0.1, 0.1, 0)$ for $j \in \{2, 3, 4\}$; and, $(\phi_{60}, \dots, \phi_{65}) = (-1.4, 0.1, 0.1, 0.1, 0.1, 0.1)$.

References

- Akande, O., G. Madson, D. S. Hillygus, and J. P. Reiter (2021). Leveraging auxiliary information on marginal distributions in nonignorable models for item and unit nonresponse. *Journal of the Royal Statistical Society Series A* 184, 643–662.
- Akande, O. and J. P. Reiter (2022). Multiple imputations for nonignorable item nonresponse in complex surveys using auxiliary margins. In *Statistics in the Public Interest: In Memory of Stephen E. Fienberg*, pp. 289–306. Cham: Springer International Publishing.
- Chen, Q., M. R. Elliott, and R. Little (2010). Bayesian penalized spline model-based inference for finite population proportion in unequal probability sampling. *Survey Methodology* 36, 23–34.
- Hasselmann, B. (2018). nleqslv: solve systems of nonlinear equations. R package version 3.2.
- Hirano, K., G. W. Imbens, G. Ridder, and D. B. Rubin (2001). Combining panel data sets with attrition and refreshment samples. *Econometrica* 69(6), 1645–1659.
- Horvitz, D. G. and D. J. Thompson (1952). A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association* 47, 663–685.
- Raghunathan, T. E., J. M. Lepkowski, J. Van Hoewyk, P. Solenberger, et al. (2001). A multivariate technique for multiply imputing missing values using a sequence of regression models. *Survey Methodology* 27, 85–96.
- Rubin, D. B. (1987). *Multiple Imputation for Nonresponse in Surveys*. New York: John Wiley.
- Sadinle, M. and J. P. Reiter (2017). Itemwise conditionally independent nonresponse modelling for incomplete multivariate data. *Biometrika* 104, 207–220.
- Sadinle, M. and J. P. Reiter (2019). Sequentially additive nonignorable missing data modelling using auxiliary marginal information. *Biometrika* 106, 889–911.

Tang, J., D. S. Hillygus, and J. P. Reiter (2024). Using auxiliary marginal distributions in imputations for nonresponse while accounting for survey weights, with application to estimating voter turnout. *Journal of Survey Statistics and Methodology* 12, 155–182.

Van Buuren, S. (2018). *Flexible Imputation of Missing Data*. CRC Press.

Yang, Y. and J. P. Reiter (2025). Imputation of nonignorable missing data in surveys using auxiliary margins via hot deck and sequential imputation. *Survey Methodology*, forthcoming.

This figure "theta2.png" is available in "png" format from:

<http://arxiv.org/ps/2412.10988v1>

This figure "theta0.5.png" is available in "png" format from:

<http://arxiv.org/ps/2412.10988v1>