

Symmetric instability in a Boussinesq fluid on a rotating planet

Yaoxuan Zeng¹ and Malte F. Jansen¹

¹Department of the Geophysical Sciences, The University of Chicago, Chicago, IL 60637, USA

Key Points:

- Symmetric instability with arbitrary background shear and stratification is studied with linear instability analysis
- Three types of instability criteria are explored with theoretical analysis and numerical simulations
- In icy moon oceans with low Rossby number flows slantwise convection occurs if and only if stratification is unstable along axis of rotation

arXiv:2412.11027v2 [physics.flu-dyn] 28 Mar 2025

Corresponding author: Yaoxuan Zeng, yzeng@uchicago.edu

Abstract

Symmetric instability has broad applications in geophysical and planetary fluid dynamics. It plays a crucial role in the formation of mesoscale rainbands at mid-latitudes on Earth, instability in the ocean’s mixed layer, and slantwise convection on gas giants and icy moon oceans. Here, we apply linear instability analysis to an arbitrary zonally symmetric Boussinesq flow on a rotating spherical planet, with applicability to icy moon oceans. We characterize the instabilities into three types: (1) gravitational instability, occurring when stratification is unstable along angular momentum surfaces, (2) inertial instability, occurring when angular momentum shear is unstable along buoyancy surfaces, and (3) a mixed symmetric instability, occurring when neither of the previous conditions are fulfilled, but the potential vorticity has the opposite sign as planetary rotation. We note that $N^2 < 0$ where N is the Brunt–Väisälä frequency—a typical criterion used to trigger convective adjustment in global ocean models—is neither necessary nor sufficient for instability. Instead, $b_z \sin \theta_0 < 0$, where b_z is the stratification along the planetary rotation axis and θ_0 is the local latitude, is always sufficient for instability and also necessary in the low Rossby number limit. In this limit, relevant for convection in icy moon oceans, the most unstable mode is slantwise convection parallel to the planetary rotation axis. This slantwise convection differs from the parameterized convection in existing general circulation models, whose convective schemes parameterize convection in the direction of gravity. Our results suggest that convection schemes in global ocean models must be revised before being applied to icy moon oceans.

Plain Language Summary

Flows on rotating planets can become unstable because of the combined effects of rotation and density stratification, a phenomenon known as symmetric instability. This instability shapes the flow patterns seen in planetary atmospheres and oceans. In our study, we use theoretical analysis and numerical simulations to study the instability criteria and the most unstable modes for axisymmetric flows on rotating planets. For flows with low Rossby numbers—typical of icy moon oceans—instability occurs if and only if the stratification is unstable along the rotation axis, leading to slantwise convection that aligns with the planetary rotation axis. This result is consistent with the rotation-aligned structures in global numerical simulations for the icy moon oceans. Additionally, it suggests that the traditional convection parameterization in ocean models, which only considers unstable stratification and heat transport in the direction of gravity, is not applicable for the icy moon oceans.

1 Introduction

Symmetric instability describes the instability of axisymmetric flow. In an axisymmetric rotating fluid with background density stratification and angular momentum gradient, if a fluid parcel is perturbed from its origin, two restoring forces come into play: the buoyancy force and the inertial acceleration (Coriolis and centrifugal forces). These forces individually may result in gravitational instability or inertial instability. Moreover, even when the fluid is both gravitationally and inertially stable, the combined effects of the two force anomalies may result in symmetric instability (Hoskins, 1974; Haine & Marshall, 1998).

Symmetric instability has broad applications in geophysical fluid dynamics. On Earth, it is relevant to the formation of mesoscale rainbands in the midlatitude atmosphere (e.g., Emanuel, 1983, 1985) and instability in the ocean’s mixed layer (e.g., Straneo et al., 2002; Callies & Ferrari, 2018). Beyond Earth, symmetric instability is closely tied to slantwise convection in the atmospheres of gas giants (e.g., Stone, 1967; Busse, 1970; Stone, 1971; Walton, 1975; O’Neill & Kaspi, 2016) and in the oceans of their icy satellites (e.g., Soder-

lund, 2019; Ashkenazy & Tziperman, 2021; Kang et al., 2022; Bire et al., 2022; Zeng & Jansen, 2024a), where convection is tilted along angular momentum surfaces.

Symmetric instability is often studied under certain assumptions about the background field. In most literature, it is examined in inertially stable ($\eta/f > 0$) fluids with stable gravitational stratification ($N^2 > 0$), where $N^2 = -(g/\rho)(\partial\rho/\partial R)$, $\eta = f - (1/R)(\partial u/\partial\theta)$ is the absolute vorticity, g is gravity, ρ is density, u is the zonal component of the velocity, $f = 2\Omega \sin\theta$ is the component of planetary rotation parallel to gravity, $\theta \in (-\pi/2, \pi/2]$ is the latitude, R is the planetary radius, and $\partial/\partial R$ denotes the derivative along the gravitational direction. Under these assumptions, instability occurs when the planetary vorticity has the opposite sign to the potential vorticity. In the Northern Hemisphere, symmetric instability arises when the potential vorticity, defined as $q = \nabla\mathbf{b} \cdot (2\boldsymbol{\Omega} + \nabla \times \mathbf{v})$, is negative, where $\boldsymbol{\Omega}$ represents planetary rotation, \mathbf{v} is the three-dimensional (3-D) relative velocity in the rotating frame, and b is the buoyancy (Eliassen, 1951; Ooyama, 1966; Hoskins, 1974; Stevens, 1983). Although assuming $N^2 > 0$ is reasonable for most regions of Earth’s atmosphere and ocean, this condition may not hold on other planetary bodies such as icy moon oceans and gas giant atmospheres. Studies on rotating convective instability have examined scenarios with $N^2 < 0$, and have found that rotating systems can remain stable even when $N^2 < 0$ (Flasar & Gierasch, 1978; Hathaway et al., 1979). In Flasar & Gierasch (1978) and Hathaway et al. (1979), the background zonal shear is assumed to be purely vertical. However, meridional shears resulting from eddy angular momentum transport (e.g., Busse, 1970; Aurnou et al., 2007; Soderlund, 2019; Zeng & Jansen, 2021) are likely significant in the oceans of icy moons and the atmospheres of gas giants, thereby influencing the instability criteria.

Beyond the background field, various approximations have been employed in studies of symmetric instability. In early studies, the traditional approximation (Gerkema et al., 2008) has been commonly applied, wherein the problem is examined on an f -plane that considers only the component of planetary rotation parallel to gravity. In this framework, the instability criteria for gravitationally and inertially stable fluids can be expressed as $\text{Ri} < \eta/f$, the same as negative PV in the Northern Hemisphere (c.f. Hoskins, 1974; Haine & Marshall, 1998), where $\text{Ri} = N^2/(\partial u/\partial R)^2$ is the Richardson number. However, the traditional approximation is not valid for planets with deep fluid layers, such as the atmospheres of gas giants and the oceans of icy moons, for flows with strong vertical motion, or for near-equatorial flows where $f \rightarrow 0$ (Gerkema et al., 2008). Later studies have considered a “tilted f -plane” that takes into account both vertical and horizontal components of the planetary rotation, $\mathbf{f} = f\mathbf{e}_R + \tilde{f}\mathbf{e}_\theta$ where $\tilde{f} = 2\Omega \cos\theta$, \mathbf{e}_R denotes the upwards direction parallel to gravity, and \mathbf{e}_θ denotes the latitudinal direction (Sun, 1995; Straneo et al., 2002; Fruman & Shepherd, 2008; Itano & Maruyama, 2009; Jeffery & Wingate, 2009). Sun (1995) concluded that as latitude decreases, i.e., the angle between planetary rotation and gravity becomes larger, the effect of \tilde{f} in modulating the maximum growth rate of symmetric instability increases. Itano & Maruyama (2009) found that the parameter regime for the occurrence of symmetric instability is less sensitive to \tilde{f} when $\text{Ri} > 0.25$ and $\eta/f > 1$, but is considerably influenced in other regions.

The instabilities discussed above are in general not well resolved in large-scale circulation models, requiring parameterization. However, most parameterizations are designed for Earth-like flows, and whether they are applicable to other planets remains unclear. Here, we discuss symmetric instability conditions and unstable mode properties over a wide planetary parameter regime with slowly varying, but otherwise arbitrary zonal velocity shear and density stratification. We apply a local linear instability analysis to the adiabatic, inviscid, non-hydrostatic Boussinesq equations formulated in a cylindrical coordinate system, which is the most natural choice for analyzing fast-rotating planets, such as icy moon oceans, where motions are largely aligned with the rotational axis (Ashkenazy & Tziperman, 2021; Bire et al., 2022). We retain all Coriolis and metric terms

and linearize around a zonally symmetric background state in hydrostatic and gradient wind balance, without imposing additional assumptions on this state. In particular, we do not require the background stratification to be stable in the gravitational direction ($N^2 > 0$), which, as we show, is in general neither a necessary nor sufficient condition for stability. We consider zonally-symmetric perturbations, which allow for gravitational, inertial, and mixed symmetric instabilities, as discussed in this paper, while excluding baroclinic instabilities associated with nonzero zonal wavenumbers (Stone, 1966). In Section 2, the theoretical analysis for linear instability criteria, the most unstable mode, and the maximum growth rate are discussed. Section 3 presents the numerical simulation results for comparison with the theoretical analysis. Section 4 provides a discussion and concluding remarks.

2 Linear instability analysis

2.1 Instability criteria

The adiabatic, inviscid, nonhydrostatic equations for a Boussinesq fluid in cylindrical coordinate are:

$$\left(\frac{\partial}{\partial t} + \frac{u}{r} \frac{\partial}{\partial \phi} + v \frac{\partial}{\partial r} + w \frac{\partial}{\partial z} \right) u + 2\Omega v + \frac{uv}{r} = -\frac{1}{r} \frac{\partial \Phi}{\partial \phi}, \quad (1)$$

$$\left(\frac{\partial}{\partial t} + \frac{u}{r} \frac{\partial}{\partial \phi} + v \frac{\partial}{\partial r} + w \frac{\partial}{\partial z} \right) v - 2\Omega u - \frac{u^2}{r} = -\frac{\partial \Phi}{\partial r} + b \cos \theta, \quad (2)$$

$$\left(\frac{\partial}{\partial t} + \frac{u}{r} \frac{\partial}{\partial \phi} + v \frac{\partial}{\partial r} + w \frac{\partial}{\partial z} \right) w = -\frac{\partial \Phi}{\partial z} + b \sin \theta, \quad (3)$$

$$\left(\frac{\partial}{\partial t} + \frac{u}{r} \frac{\partial}{\partial \phi} + v \frac{\partial}{\partial r} + w \frac{\partial}{\partial z} \right) b = 0, \quad (4)$$

$$\frac{1}{r} \frac{\partial u}{\partial \phi} + \frac{\partial v}{\partial r} + \frac{\partial w}{\partial z} + \frac{v}{r} = 0, \quad (5)$$

where r is the radial direction, increasing outward from the rotational axis of the planet, ϕ is the longitude, denoting the zonal (azimuthal) direction, and z is the vertical (rotational) direction (Figure 1). u , v , w are velocity components in the \mathbf{e}_ϕ , \mathbf{e}_r , and \mathbf{e}_z directions, respectively. $\Phi = p'/\rho_0$, where p' is the pressure anomaly and ρ_0 is the reference density. $b = -g\rho'/\rho_0$ is buoyancy, where ρ' is the density anomaly (c.f. Vallis, 2017).

We consider an axisymmetric state around the planetary rotation axis where all variables are invariant in the zonal direction ($\partial/\partial\phi = 0$, i.e. zonally symmetric), and assume a background state with an arbitrary background zonal flow $\bar{u}(r, z)$ and buoyancy field $\bar{b}(r, z)$ that are invariant in time: $b = \bar{b}(r, z) + b'(r, z, t)$, $\Phi = \bar{\Phi}(r, z) + \Phi'(r, z, t)$, $u = \bar{u}(r, z) + u'(r, z, t)$, $v = v'(r, z, t)$, $w = w'(r, z, t)$. Assuming the perturbations are small, the zeroth-order balance reveals the gradient wind balance and hydrostatic balance for the background state:

$$-2\Omega\bar{u} - \frac{\bar{u}^2}{r} = -\frac{\partial\bar{\Phi}}{\partial r} + \bar{b} \cos \theta, \quad (6)$$

$$0 = -\frac{\partial\bar{\Phi}}{\partial z} + \bar{b} \sin \theta. \quad (7)$$




Figure 1. Sketch of the coordinate system. Black coordinates show the cylindrical coordinate system applied in this paper (\mathbf{e}_r , \mathbf{e}_ϕ , \mathbf{e}_z), where \mathbf{e}_z is the planetary rotation axis and the origin locates at the center of the planet. Grey coordinates show the spherical coordinate system (\mathbf{e}_ϕ , \mathbf{e}_θ , \mathbf{e}_R), where \mathbf{e}_R is opposite to gravity. Note that the zonal direction \mathbf{e}_ϕ is the same in both cylindrical and spherical coordinates.

With

$$\bar{f} = 2\Omega + 2\bar{\omega}, \quad (8)$$

being a modified Coriolis parameter where $\bar{\omega} \equiv \bar{u}/r$ is the angular velocity of the background zonal flow, Equations 6 & 7 yield

$$\bar{f} \frac{\partial \bar{u}}{\partial z} = \frac{\partial \bar{b}}{\partial r} \sin \theta - \frac{\partial \bar{b}}{\partial z} \cos \theta. \quad (9)$$

Equation 9 is similar to the thermal wind balance in a rapidly rotating (geostrophic) fluid (Kaspi et al., 2009; Bire et al., 2022), but here \bar{f} is modified by the background flow, with the effects of centrifugal force incorporated, thus giving the shear of a flow in gradient-wind balance rather than geostrophic balance.

In the first-order balance, we obtain linearized perturbation equations (Equations A21-A25). In this paper, we focus on local instability analysis, with the length scales of the perturbations small compared to the planetary radius. We therefore look for plane-wave solutions, where all variables are proportional to $\exp(ik_r r + ik_z z - i\xi t)$, where ξ is the angular frequency, and k_r and k_z are wavenumbers in the radial and vertical directions, respectively. We assume that variations in the background field are small across the perturbation length scale (Equation A13), allowing us to evaluate them locally at $(r = r_0, z = z_0)$, with k_r and k_z treated as constants. Under these assumptions, we have the dispersion relation (see Appendix A for detailed derivation)

$$\xi^2 = \frac{b_z \sin \theta_0 k_r^2 + (f_0 M_r + b_r \cos \theta_0) k_z^2 - 2b_r \sin \theta_0 k_r k_z}{k_r^2 + k_z^2}, \quad (10)$$

where $b_z = \partial \bar{b} / \partial z$ and $b_r = \partial \bar{b} / \partial r$ are stratification, $f_0 = 2\Omega + 2\bar{\omega}$ is the modified Coriolis parameter, and $M_r = f_0 + r_0 (\partial \bar{\omega} / \partial r)$ describes the radial gradient of the background angular momentum $\bar{m} = (\Omega + \bar{\omega}) r^2$, all evaluated at (r_0, z_0) . $\theta_0 = \arctan(z_0 / r_0)$ is the local latitude.

Stability requires that the frequency ξ has no imaginary part, which means the right-hand-side of Equation 10 is positive definite. Specifically, this requires the quadratic function $b_z \sin \theta_0 k_r^2 + (f_0 M_r + b_r \cos \theta_0) k_z^2 - 2b_r \sin \theta_0 k_r k_z > 0$ for all k_r and k_z . Consequently, the stability matrix

$$\begin{pmatrix} b_z \sin \theta_0 & -b_r \sin \theta_0 \\ -b_r \sin \theta_0 & f_0 M_r + b_r \cos \theta_0 \end{pmatrix}$$

must be positive definite. This stability matrix is essentially the same as the one derived in previous studies on symmetric instability on a tilted f -plane (e.g., [Itano & Maruyama, 2009](#)) although here it is expressed in cylindrical coordinates instead of spherical coordinates to emphasize the symmetry in flows that are strongly rotationally constrained, and the Coriolis parameter f_0 is modified by the angular velocity of the background flow $\bar{\omega}_0$. The necessary and sufficient conditions for instability are that either the trace or determinant of this matrix be negative, i.e., either

$$b_z \sin \theta_0 + f_0 M_r + b_r \cos \theta_0 < 0, \quad (11)$$

or

$$b_z \sin \theta_0 (f_0 M_r + b_r \cos \theta_0) - b_r^2 \sin^2 \theta_0 < 0. \quad (12)$$

Equation 12 can also be expressed in terms of a condition for the background potential vorticity. The background potential vorticity is

$$q_0 = \nabla \bar{b} \cdot (2\mathbf{\Omega} + \nabla \times \bar{\mathbf{v}}) = M_r b_z - M_z b_r = M_r b_z - \frac{b_r}{f_0} (b_r \sin \theta_0 - b_z \cos \theta_0), \quad (13)$$

where $M_z = r_0 (\partial \bar{\omega} / \partial z)$ is the vertical gradient of the background angular momentum, and the background gradient wind shear (Equation 9) is applied. We can hence rewrite Equation 12 as

$$q_0 f_0 \sin \theta_0 < 0, \quad (14)$$

i.e. instability occurs if the potential vorticity has the opposite sign as the generalized background planetary vorticity. This instability criterion is commonly referred to as the symmetric instability criterion (e.g., [Hoskins, 1974](#); [Haine & Marshall, 1998](#)), although it should be noted that in general, Equation 14 by itself is a sufficient but not necessary condition for instability.

2.2 Most unstable mode and growth rate

When ξ is imaginary, the perturbation fields will grow exponentially. Note that ξ^2 is a real number (Equation 10); therefore, ξ is either purely real (corresponding to oscillatory solutions) or purely imaginary (corresponding to exponentially growing or decaying solutions). By substituting $\xi = i\sigma$ in the dispersion relation (Equation 10), the e-folding growth rate σ of the unstable mode can be expressed as

$$\sigma = \left(-\frac{b_z \sin \theta_0 \tan^2 \delta + 2b_r \sin \theta_0 \tan \delta + f_0 M_r + b_r \cos \theta_0}{1 + \tan^2 \delta} \right)^{1/2}, \quad (15)$$

where $\delta \in (-\pi/2, \pi/2]$ is the angle between the unstable mode and \mathbf{e}_r so that $\tan \delta = -k_r/k_z$. Equation 15 indicates that for a given background field and latitude, the growth rate is only a function of $\tan \delta$, i.e. the direction of the unstable mode, but not the magnitude of the wavenumber, i.e. the size of the mode. The result that the growth rate depends only on the direction of the mode is consistent with previous studies of symmetric instability on an f -plane (e.g., Ooyama, 1966) and a tilted f -plane (e.g., Sun, 1995). By calculating the derivatives of the function $\sigma(\tan \delta)$ and performing some algebra, we find that the maximum growth rate is obtained when

$$\tan \delta_m = \frac{b_z \sin \theta_0 - f_0 M_r - b_r \cos \theta_0 - [(f_0 M_r + b_r \cos \theta_0 - b_z \sin \theta_0)^2 + 4b_r^2 \sin^2 \theta_0]^{1/2}}{2b_r \sin \theta_0}. \quad (16)$$

Substituting Equation 16 into Equation 15, we obtain the maximum growth rate as

$$\sigma_m = \left(-\frac{f_0 M_r + b_r \cos \theta_0 + b_z \sin \theta_0 - \sqrt{(f_0 M_r + b_r \cos \theta_0 - b_z \sin \theta_0)^2 + 4b_r^2 \sin^2 \theta_0}}{2} \right)^{1/2}. \quad (17)$$

2.3 Instability diagram

In this section, we discuss the instability criteria assuming $b_r \sin \theta_0 \neq 0$. The special case where $b_r \sin \theta_0 = 0$ is discussed in Appendix B, where it is shown that the most unstable mode in this case always aligns with the radial (r) or vertical (z) directions. The instability criteria for fluid parcels perturbed purely along the \mathbf{e}_r and \mathbf{e}_z directions are $f_0 M_r + b_r \cos \theta_0 < 0$ and $b_z \sin \theta_0 < 0$, respectively, which are derived by solving $\xi^2 < 0$ (Equation 10) when $\delta = 0$ and $\delta = \pi/2$, respectively. It is convenient to define a 2-D nondimensionalized phase space based on the stability in these two orthogonal directions:

$$x = \frac{b_z \sin \theta_0}{|b_r \sin \theta_0|}, \quad y = \frac{f_0 M_r + b_r \cos \theta_0}{|b_r \sin \theta_0|}. \quad (18)$$

The stability matrix then becomes

$$\begin{pmatrix} x & -1 \\ -1 & y \end{pmatrix},$$

and positive definiteness requires that

$$x + y > 0, \text{ and } xy > 1, \quad (19)$$

which means only the region above the hyperbolic $xy = 1$ in the first quadrant ($x > 0, y > 0$) is stable (Figure 2a).

2.4 Gravitational instability, inertial instability, and mixed symmetric instability

In a rotating fluid with background stratification and shear, two restoring forces, associated with gravity and rotation, act when a fluid parcel is displaced: buoyancy force




Figure 2. Growth rate and orientation of the most unstable mode with positive b_r here shown for latitude $\theta_0=60^\circ$. (a): growth rate of the most unstable mode, nondimensionalized with $|b_r|^{1/2}$: $\sigma_m/|b_r|^{1/2} = ((-x - y + ((x - y)^2 + 4)^{1/2})|\sin \theta_0|/2)^{1/2}$. (b): sketch for the definition of α and β . (c) and (d): the angle (in degrees) between the most unstable mode and the buoyancy surface (α , c) and the angular momentum surface (β , d). In (a), (c), and (d), the solid lines separate the plane into regions where gravitational instability, inertial instability, mixed symmetric instability, or a combination of them occur (see Figure 3 and discussion in Section 2.4).

and inertial acceleration. If a fluid parcel is displaced along a constant angular momentum surface, the inertial acceleration anomaly is zero and the only restoring force is the buoyancy force anomaly. As a result, pure gravitational instability can occur when the stratification is unstable along constant angular momentum surfaces. Similarly, if the displacement of the fluid parcel is along a constant buoyancy surface, the only restoring force is the inertial acceleration anomaly, hence pure inertial instability can occur when the angular momentum shear is unstable along constant buoyancy surfaces. Here, unstable stratification indicates buoyancy decreases in the opposite direction to gravity, and unstable angular momentum shear indicates the angular momentum gradient is opposite to the background planetary angular momentum gradient, $f_0 \mathbf{e}_r$ (Figure 3a & b). Therefore, gravitational instability can occur when

$$(\nabla \bar{b} \cdot \mathbf{e}_M)(-\mathbf{e}_R \cdot \mathbf{e}_M) = (M_r b_z - M_z b_r)(M_r \sin \theta_0 - M_z \cos \theta_0) < 0, \quad (20)$$

where $\mathbf{e}_M = -M_z \mathbf{e}_r + M_r \mathbf{e}_z$ is the direction along a constant angular momentum surface. Inertial instability can occur when

$$(\nabla \bar{m} \cdot \mathbf{e}_b)(f_0 \mathbf{e}_r \cdot \mathbf{e}_b) = f_0 b_z r_0^2 (M_r b_z - M_z b_r) < 0, \quad (21)$$




Figure 3. Different types of instabilities and the instability diagram. (a) and (b): sketches for gravitational instability (a) and inertial instability (b), respectively (see text for detailed explanation). (c) and (d): instability diagram for $b_r > 0$ (c) and $b_r < 0$ (d), respectively. Solid lines indicate gravitational, inertial, and mixed symmetric instability criteria. The thicker solid lines indicate the hyperbolic curve $xy = 1$ (Equation 24) and the line $x = 0$ (Equation 23). The thinner solid lines indicate the line $y + \cot^2 \theta_0 x - 2|\cot \theta_0| \text{sgn}(b_r) = 0$ (Equation 22), which is always tangential to the hyperbolic curve $xy = 1$. The dashed lines indicate $N^2 = 0$, with regions to the left corresponding to $N^2 < 0$. The black lines are for $\theta_0 = 60^\circ$ and red lines are for $\theta_0 = 30^\circ$ for comparison. The subscripts indicate different types of instability, where G indicates gravitational instability (Equation 20), I indicates inertial instability (Equation 21), and M indicates mixed symmetric instability (Equation 14). The difference between S_{GI_a} and S_{GI_b} is the sign of the background potential vorticity, where in S_{GI_a} , $q_0 f_0 \sin \theta_0 < 0$; in S_{GI_b} , $q_0 f_0 \sin \theta_0 > 0$. For pure gravitational instability (S_G) or pure inertial instability (S_I), we also always have $q_0 f_0 \sin \theta_0 < 0$. The black dots are simulation parameters for s_G to s_S (Table 1). Note that for the low Rossby number cases s_G and s_S , the y -axis value is 1.15×10^4 , outside the y -values shown in the plot.

where $\mathbf{e}_b = -b_z \mathbf{e}_r + b_r \mathbf{e}_z$ is the direction along a constant buoyancy surface. Using the gradient wind shear (Equation 9), and the definition of x and y (Equation 18), the gravitational instability criterion (Equation 20) becomes

$$[y + \cot^2 \theta_0 x - 2|\cot \theta_0 \text{sgn}(b_r)](xy - 1) < 0, \quad (22)$$

where sgn denotes the sign function, and the inertial instability criterion (Equation 21) becomes

$$x(xy - 1) < 0. \quad (23)$$

Comparison with the results derived in Section 2.1 shows that neither the gravitational or inertial instability criteria are necessary for instability. Instability can still occur as a result of gravitational and inertial force anomalies when $q_0 f_0 \sin \theta_0 < 0$, or equivalently,

$$xy - 1 < 0. \quad (24)$$

We therefore refer this instability (Equation 24) when the flow is both gravitationally stable and inertially stable as a mixed symmetric instability.

In the instability diagram, the phase plane is divided into several regions where pure gravitational instability (Equation 22), pure inertial instability (Equation 23), or mixed symmetric instability (Equation 24) occur. In all regions except for the stable region S_S located above the hyperbolic curve $xy = 1$ in the first quadrant, at least one type of instability is present (see Figure 3c & d).

The alignment of the most unstable mode with the buoyancy and angular momentum surfaces is indicative of different types of instabilities. To illustrate how the most unstable mode aligns with the buoyancy and angular momentum surfaces, we define $\alpha \in (-\pi/2, \pi/2]$ as the angle between the direction of the most unstable mode and the buoyancy surface, and $\beta \in (-\pi/2, \pi/2]$ as the angle between the direction of the most unstable mode and the angular momentum surface, with the most unstable mode counterclockwise of the buoyancy/angular momentum surface defined as positive (Figure 2b). By comparing Figure 3c and Figure 2c & d, regions where the most unstable mode aligns with the buoyancy surface ($\alpha = 0$) can be identified as inertially unstable, and regions where the most unstable mode aligns with the angular momentum surface ($\beta = 0$) can be identified as gravitationally unstable. However, the opposite is not always true, and in many parts of the parameter regime, the most unstable mode does not align with either surface. The special case where the Rossby number is small and the most unstable mode always aligns with the angular momentum surface will be discussed in Section 2.5.

Most previous studies on symmetric instability have neglected regimes where $N^2 < 0$ or $\eta/f < 0$, as they typically assume stable stratification in the gravitational direction and stable zonal velocity shear in the latitudinal direction. However, we find that $N^2 < 0$ is neither sufficient nor necessary for instability. In Figure 3c & d, the dashed lines indicate $N^2 = 0$, with regions to the left of these lines corresponding to $N^2 < 0$. Therefore, when $b_r < 0$, the system can remain stable even if $N^2 < 0$. Meanwhile, we also find that $\eta/f < 0$ is a sufficient (but not necessary) condition for instability. When $\eta/f > 0$, the sufficient and necessary condition for instability becomes $q_0 f_0 \sin \theta_0 < 0$ (see Figure C1 and detailed derivation in Appendix C). This result is consistent with Flasar & Gierasch (1978) and Hathaway et al. (1979), who assumed that the background flow is in thermal wind balance with no meridional shear, such that $\eta/f = 1 > 0$. Under these assumptions, they found that symmetric instability would occur if and only if $q_0 f_0 \sin \theta_0 <$

0 (equivalent to Equation 38 in [Hathaway et al. \(1979\)](#) after substituting in the background thermal wind shear), consistent with our findings.

2.5 Low Rossby number limit

In the low Rossby number regime, $\text{Ro} = U/(f_0 L) \ll 1$ where U is the velocity scale and L is the length scale of motion, the rotational effect dominates over nonlinear advection (c.f. [Vallis, 2017](#)). Many planetary flows in geophysical fluid dynamics likely fall into this regime, such as the atmosphere outside the equatorial region on gas giants like Jupiter ([Kaspi et al., 2018](#)) and the ocean flows on icy moons ([Jansen et al., 2023](#)). In the low Rossby number limit, we have $\bar{w}, r\partial\bar{w}/\partial r, r\partial\bar{w}/\partial z \ll f_0$, which implies $M_r \approx f_0 \approx 2\Omega$ and $M_z \ll f_0$. In gradient wind balance, we moreover have the scaling that $b_r, b_z \sim f_0 M_z \ll f_0^2$. Consequently, the criterion for inertial instability, Equation 21, reduces to

$$f_0^2 b_z^2 < 0, \quad (25)$$

which cannot be satisfied, because small Rossby number flow is always inertially stable. Instability then occurs if, and only if,

$$b_z \sin \theta_0 < 0, \quad (26)$$

i.e. for an unstable stratification in the direction parallel to planetary rotation, which coincides with the angular momentum surface in the small Ro limit. The most unstable mode (Equation 16) is obtained when

$$|\tan \delta_m| \approx \left| \frac{-f_0^2 - [(f_0^2)^2]^{1/2}}{2b_r \sin \theta_0} \right| = \left| \frac{f_0^2}{b_r \sin \theta_0} \right| \gg 1, \quad \text{i.e. } \delta_m \approx \frac{\pi}{2}, \quad (27)$$

and the corresponding growth rate is

$$\sigma_m \approx (-b_z \sin \theta_0)^{1/2}. \quad (28)$$

Therefore, the most unstable mode is associated with slantwise convection aligned with the planetary angular momentum surface. This mode, parallel to the rotational axis, is likely to be important in the oceans of icy moons, which are thought to be characterized by low Rossby numbers. Such slantwise convection aligned with the rotational axis has been identified in numerical simulations of icy moon oceans (e.g., [Ashkenazy & Tziperman, 2021](#); [Kang et al., 2022](#); [Bire et al., 2022](#); [Zeng & Jansen, 2024a](#)).

3 Numerical simulations

3.1 Simulation set up

To verify the theoretical results, we numerically integrate Equations 1–5 for a zonally symmetric flow ($\partial/\partial\phi = 0$) using Dedalus, which can solve initial-value partial differential equations using spectral methods ([Burns et al., 2020](#)). We nondimensionalize the equations using the rotational time scale ($T = f_0^{-1}$) and the domain length scale. We neglect variations of the background field and the metric term in the continuity equation, allowing the use of a local Cartesian coordinate system and double-periodic boundary conditions (c.f. [Appendix D](#)). We prescribe a background state which satisfies the gradient wind balance (Equations 6 & 7), and solve for the evolution of the nondimensionalized perturbation fields ($\hat{u}, \hat{v}, \hat{w}, \hat{b}, \hat{\Phi}$). We apply a resolution with 256 grid points in both r and z directions. Simulations without viscosity and diffusivity develop unphysical grid-scale noise, once nonlinear effects become significant (Figure D1). In addition

Simulation	\hat{b}_r	\hat{b}_z	\hat{M}_r	\hat{q}_0	\hat{N}^2	Instability	$\hat{\sigma}_m$
s_G	1e-4	-3.33e-5	1	-3.33e-5	2.11e-5	Grav	5.37e-3
s_{GI_a}	1	0.15	2.5	-0.416	0.630	Grav, Iner	0.333
s_{GI_b}	1	-0.5	-4	0.884	6.70e-2	Grav, Iner	1.93
s_I	1	1	0	-0.366	1.37	Iner	0.450
s_M	1	-0.333	0	-1.03	0.211	Mixed	0.920
s_S	-1e-4	3.33e-5	1	3.33e-5	-2.11e-5	Stable	/

Table 1. Numerical simulation setups. The latitudes for all simulations are $\theta_0 = 60^\circ$. The simulation parameters are nondimensionalized with the rotational time scale (f_0^{-1}) and the domain length scale, and hats are used to indicate nondimensionalized quantities (See [Appendix A](#)). The types of instability in each simulation are indicated in the simulation name and the “Instability” column, where G and Grav indicate gravitational instability, I and Iner indicate inertial instability, M and Mixed indicate mixed symmetric instability, and S and Stable indicate the system is stable. Simulations s_G and s_S represent the low Rossby number limit.




Figure 4. Background states in the simulations, with $\hat{b} = \hat{b}_r \hat{r} + \hat{b}_z \hat{z}$ and $\hat{M} = \hat{M}_r \hat{r} + \hat{M}_z \hat{z}$, where the domain-averaged values are subtracted. The black dashed lines indicate the direction parallel to gravity (\mathbf{e}_R), and the magenta dashed lines indicate the direction perpendicular to gravity (\mathbf{e}_θ). Notice that the much weaker buoyancy gradients in s_G and s_S place these simulations in the small Rossby number regime, such that the angular momentum surfaces are approximately aligned with the rotation axes (i.e. the vertical axis in this coordinate system).

to inviscid simulations, we therefore perform simulations with a Leith sub-grid parameterization ([Leith, 1996](#)) to represent the effects of sub-grid-scale eddy mixing, and compare their results with the simulations without viscosity and diffusivity. It takes some time for the most unstable mode to grow and become dominant, before which nonlinear effects may already induce turbulence in the system. To address this issue, we initialize the simulations with their most unstable modes (as inferred from an integration of the linearized equations). A detailed description of the simulation setup can be found in [Appendix D](#).

We carry out six simulations with the background field representing each region in the parameter space, where different types of instability (or no instability) occur ([Figure 3](#)). The simulation parameters are summarized in [Table 1](#) and the prescribed background fields are shown in [Figure 4](#).




Figure 5. Time series for domain-averaged eddy kinetic energy (EKE), $(\hat{u}^2 + \hat{v}^2 + \hat{w}^2)/2$. The blue and red solid lines show the simulation results without diffusivity and viscosity, and the blue and red dashed lines show the simulation results with eddy diffusivity and viscosity. The black dashed line shows the predicted maximum growth rate of EKE ($2\hat{\sigma}_m$ due to the square, Equation 17). The time is normalized by the maximum growth rate in each simulation (for s_S , the normalization uses the maximum growth rate for simulation s_G).

3.2 Simulation results

For all simulations without diffusivity and viscosity, the eddy kinetic energy (EKE) grows exponentially, consistent with the maximum growth rate predicted by Equation 17. When the Leith closure is applied, the EKE growth rate decreases slightly due to viscous and diffusive dissipation (Figure 5). During the exponential growth phase of simulations with an unstable background state, the orientation of the perturbation fields aligns closely with the most unstable mode predicted by Equation 16 (Figure 6). Eventually, the non-linear advection effects become dominant, the flow becomes turbulent, and the growth rates of EKE decrease significantly. Notably, the EKE growth eventually converges for simulations with and without eddy viscosity and diffusivity (Figure 5).

The simulation with stable background state (s_S) does not show exponential growth but an oscillation behavior. The initial condition used in simulation s_S is the most unstable mode in simulation s_G , which has $\tan \delta = \pi/2$. In this case, we can estimate the oscillation period P according to the dispersion relation (Equation 10) as $P = 584.79$, or $3.14/\hat{\sigma}_m$ with $\hat{\sigma}_m$ being the maximum growth rate of s_G , consistent with the oscillation period in the simulation (Figure 5a).

The numerical simulations also show that $N^2 < 0$ is neither a sufficient nor necessary condition for instability. In simulation s_S , $N^2 < 0$, but the system is stable. In simulations s_G and s_{GI_b} , $N^2 > 0$, but the systems are unstable, with either $q_0 f_0 \sin \theta_0 > 0$ (s_{GI_b}) or $q_0 f_0 \sin \theta_0 < 0$ (s_G).

In simulations with $Ro \ll 1$ (s_G and s_S), the instability criteria reduce to $b_z \sin \theta_0 < 0$. In the unstable case (simulation s_G), the most unstable mode is then parallel to the rotational axis, i.e. slantwise convection parallel to planetary rotation.

4 Discussions and conclusions

We conduct local linear instability analysis with slowly-varying background shear and stratification, and find that the necessary and sufficient conditions for instability of zonally symmetric Boussinesq flow on a rotating planet are the background field is (1) gravitationally unstable, or (2) inertially unstable, or (3) the background potential vorticity has a different sign from the planetary vorticity, which is consistent with previous studies on symmetric instability (e.g., Hoskins, 1974). In our framework, we define the gravitational instability criterion as unstable buoyancy stratification along angular mo-




Figure 6. Snapshots of the normalized buoyancy anomaly fields from simulations with eddy diffusivity and viscosity. The buoyancy anomaly is normalized by the maximum value of the background buoyancy field ($\hat{b}/\max(|\hat{b}|)$) in each simulation. The time of each snapshot is indicated above the respective columns and are normalized by the maximum growth rate $\hat{\sigma}_m$ in each simulation. The black solid lines indicate the orientation of the most unstable mode predicted by Equation 16.

mentum surfaces (Equation 20 & Figure 3a) and the inertial instability criterion as unstable angular momentum shear along buoyancy surfaces (Equation 21 & Figure 3b), which ensures both to be sufficient conditions for instability. Mathematically, the criterion for instability can most compactly be expressed as either $b_z \sin \theta_0 < 0$ (i.e. unstable stratification along the planetary rotation axis) or $q_0 f_0 \sin \theta_0 < 0$ where q_0 is the background potential vorticity. When instability occurs, the growth rate is not sensitive to the magnitude of the radial and vertical wavenumbers, k_r and k_z , but is only a function of their ratio, k_r/k_z , i.e., the tilting direction of the mode.

In the low Rossby number limit, instability occurs if and only if $b_z \sin \theta_0 < 0$, and the most unstable mode is slantwise convection parallel to the planetary rotation axis. The low Rossby number limit proves to be particularly valuable for understanding slantwise convection in icy moon oceans, as this phenomenon is neither properly resolved nor parameterized in global ocean simulations (c.f. Zeng & Jansen, 2024a).

Negative stratification in the gravitational direction, $N^2 < 0$, is neither sufficient nor necessary for instability. This suggests that the traditional convective adjustment schemes widely applied in ocean General Circulation Models (e.g., Marotzke, 1991), which parameterizes upright convection when $N^2 < 0$, may not be suitable for slantwise convection in icy moon oceans. Given that $b_z \sin \theta_0 < 0$ is a sufficient condition for symmetric instability, and becomes a sufficient and necessary condition under the low Rossby number limit, we propose that stratification along the rotational axis should be treated as the criterion for static stability and used to trigger convection parameterizations in planetary ocean models.

Although we consider Boussinesq fluids in this manuscript, the results are also relevant for anelastic fluids, such as atmospheric convection on gas giants (e.g., O’Neill & Kaspi, 2016), as long as the vertical scale of the unstable modes is much smaller than the scale height of the atmosphere (i.e., the vertical density variation is small). Flows on gas giants also satisfy $Ro \ll 1$, indicating that rotationally aligned modes may be the most unstable, as suggested by our results. Observations of the gravitational fields of Jupiter and Saturn suggest that the zonal jets on these planets are aligned with the rotational axis (Kaspi et al., 2018; Galanti et al., 2019; Kaspi et al., 2023), indicating that the prevailing modes are parallel to planetary rotation. The in-situ temperature profile measurement from the Galileo probe indicates a neutral stratification along constant angular momentum surfaces, which are parallel to the rotational axis outside the high-shear region in the upper weather layer (O’Neill et al., 2017). Numerical simulations of gas giant atmospheres also reveal slantwise convection aligned with planetary rotation (Christensen, 2002; Kaspi et al., 2009; Heimpel et al., 2016).

Our study focuses exclusively on zonally symmetric instabilities. However, asymmetric modes could also play a significant role in fluid motions on a rotating planet. For instance, equatorial convective rolls driven by heating from the bottom of the fluid are believed to be important in driving the equatorial jets on gas giants (e.g., Busse & Hood, 1982; Busse & Or, 1986). Therefore, it is important to characterize the growth rate of different symmetric and asymmetric unstable modes to determine under which regime the symmetric mode dominates. Stone (1966) studied the growth rate of symmetric instability, Kelvin-Helmholtz instability, and baroclinic instability on an f -plane with hydrostatic balance. He concludes that on an f -plane, symmetric instability prevails when $0.25 < Ri < 0.95$; Kelvin-Helmholtz instability dominates for $Ri < 0.25$; and baroclinic instability is dominant when $Ri > 0.95$. When considering the horizontal component of planetary rotation, Jeffery & Wingate (2009) suggest that the transition point between symmetric and baroclinic instability dominance can exceed $Ri = 0.95$. Understanding how the symmetric mode, such as slantwise convection in the low Rossby number limit, interacts with baroclinic instability under arbitrary background stratification and shear, and conditions that separate symmetric and asymmetric mode dominance, are important future research directions. We applied a linear instability analysis method,

which is valuable to understand stability criteria for small-amplitude perturbations but is not applicable to large-amplitude perturbations or non-linear instability problems. [Bowman & Shepherd \(1995\)](#) applied the energy-Casimir stability method ([Holm et al., 1985](#); [Shepherd, 1990](#); [Cho et al., 1993](#)) to study nonlinear symmetric instability where the hydrostatic approximation is made and the rotation is aligned with gravity. However, extending this approach to a more generalized setup (e.g., [Fruman & Shepherd, 2008](#)) presents challenges that remain to be addressed in future work. In our current study, solutions are sought on an infinitely large domain (local plane-wave solutions). Future work should also explore how boundary conditions might modify the characteristics of the most unstable modes.

Despite these limitations, our work provides criteria and physical insight into instabilities within a generalized framework applicable across diverse planetary parameter regimes. These results enhance our understanding of fluid motions on various planetary bodies and can improve parameterizations of sub-grid-scale transports in large-scale models, such as slantwise convection in icy moon oceans.

Appendix A Derivation of the dispersion relation

After subtracting the gradient wind balance and the hydrostatic balance for the background flow (Equations 6 & 7) from the original Equations 1-5, we have the perturbation equations:

$$\left(\frac{\partial}{\partial t} + v' \frac{\partial}{\partial r} + w' \frac{\partial}{\partial z}\right) u' + \left(2\Omega + \frac{\partial \bar{u}}{\partial r} + \frac{\bar{u}}{r}\right) v' + \frac{\partial \bar{u}}{\partial z} w' = 0, \quad (\text{A1})$$

$$\left(\frac{\partial}{\partial t} + v' \frac{\partial}{\partial r} + w' \frac{\partial}{\partial z}\right) v' - \left(2\Omega + \frac{2\bar{u}}{r}\right) u' - \cos \theta b' + \frac{\partial \Phi'}{\partial r} = 0, \quad (\text{A2})$$

$$\left(\frac{\partial}{\partial t} + v' \frac{\partial}{\partial r} + w' \frac{\partial}{\partial z}\right) w' - \sin \theta b' + \frac{\partial \Phi'}{\partial z} = 0, \quad (\text{A3})$$

$$\left(\frac{\partial}{\partial t} + v' \frac{\partial}{\partial r} + w' \frac{\partial}{\partial z}\right) b' + \frac{\partial \bar{b}}{\partial r} v' + \frac{\partial \bar{b}}{\partial z} w' = 0, \quad (\text{A4})$$

$$\frac{1}{r} \frac{\partial(v'r)}{\partial r} + \frac{\partial w'}{\partial z} = 0. \quad (\text{A5})$$

We assume that the background fields vary slowly over the characteristic length scale of the perturbations, L , so that they can be treated as constants. This criterion can be formally obtained by performing the Taylor expansion of all background fields locally around ($r = r_0, z = z_0$):

$$\begin{aligned} 2\Omega + \frac{2\bar{u}}{r} &= 2\Omega + 2\bar{\omega} \\ &= 2\Omega + 2\bar{\omega}(r_0, z_0) + 2 \left. \frac{\partial \bar{\omega}}{\partial r} \right|_{(r_0, z_0)} r' + 2 \left. \frac{\partial \bar{\omega}}{\partial z} \right|_{(r_0, z_0)} z' + \dots, \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} 2\Omega + \frac{\partial \bar{u}}{\partial r} + \frac{\bar{u}}{r} &= 2\Omega + 2\bar{\omega} + r \frac{\partial \bar{\omega}}{\partial r} \\ &= 2\Omega + 2\bar{\omega}(r_0, z_0) + r_0 \left. \frac{\partial \bar{\omega}}{\partial r} \right|_{(r_0, z_0)} \\ &\quad + \left(3 \left. \frac{\partial \bar{\omega}}{\partial r} \right|_{(r_0, z_0)} + r_0 \left. \frac{\partial^2 \bar{\omega}}{\partial r^2} \right|_{(r_0, z_0)}\right) r' + \left(2 \left. \frac{\partial \bar{\omega}}{\partial z} \right|_{(r_0, z_0)} + r_0 \left. \frac{\partial^2 \bar{\omega}}{\partial r \partial z} \right|_{(r_0, z_0)}\right) z' + \dots, \end{aligned} \quad (\text{A7})$$

$$\frac{\partial \bar{u}}{\partial z} = r \frac{\partial \bar{\omega}}{\partial z} = r_0 \left. \frac{\partial \bar{\omega}}{\partial z} \right|_{(r_0, z_0)} + \left(r_0 \frac{\partial^2 \bar{\omega}}{\partial r \partial z} + \frac{\partial \bar{\omega}}{\partial z} \right)_{(r_0, z_0)} r' + r_0 \left. \frac{\partial^2 \bar{\omega}}{\partial z^2} \right|_{(r_0, z_0)} z' + \dots, \quad (\text{A8})$$

$$\frac{\partial \bar{b}}{\partial r} = \left. \frac{\partial \bar{b}}{\partial r} \right|_{(r_0, z_0)} + \left. \frac{\partial^2 \bar{b}}{\partial r^2} \right|_{(r_0, z_0)} r' + \left. \frac{\partial^2 \bar{b}}{\partial r \partial z} \right|_{(r_0, z_0)} z' + \dots, \quad (\text{A9})$$

$$\frac{\partial \bar{b}}{\partial z} = \left. \frac{\partial \bar{b}}{\partial z} \right|_{(r_0, z_0)} + \left. \frac{\partial^2 \bar{b}}{\partial r \partial z} \right|_{(r_0, z_0)} r' + \left. \frac{\partial^2 \bar{b}}{\partial z^2} \right|_{(r_0, z_0)} z' + \dots, \quad (\text{A10})$$

$$\begin{aligned} \sin \theta &= \sin \theta_0 + \left. \frac{\partial \sin \theta}{\partial r} \right|_{(r_0, z_0)} r' + \left. \frac{\partial \sin \theta}{\partial z} \right|_{(r_0, z_0)} z' + \dots \\ &= \sin \theta_0 - \sin \theta_0 \cos \theta_0 \frac{r'}{\sqrt{r_0^2 + z_0^2}} + \cos^2 \theta_0 \frac{z'}{\sqrt{r_0^2 + z_0^2}} + \dots, \end{aligned} \quad (\text{A11})$$

$$\begin{aligned} \cos \theta &= \cos \theta_0 + \left. \frac{\partial \cos \theta}{\partial r} \right|_{(r_0, z_0)} r' + \left. \frac{\partial \cos \theta}{\partial z} \right|_{(r_0, z_0)} z' + \dots \\ &= \cos \theta_0 + \sin^2 \theta_0 \frac{r'}{\sqrt{r_0^2 + z_0^2}} - \sin \theta_0 \cos \theta_0 \frac{z'}{\sqrt{r_0^2 + z_0^2}} + \dots, \end{aligned} \quad (\text{A12})$$

where $\theta_0 = \arctan(z_0/r_0)$ is the local latitude, and $|r'| \sim |z'| \sim L$. We study the local instability of fluid in the outer region of the planet, where we assume $L \ll r_0$. The left-hand-sides of Equations A6–A12 remain approximately constant over the scale L if

$$\begin{aligned} \left(\frac{|\bar{\omega}|}{|d\bar{\omega}|} \right)_{(r_0, z_0)} &\gg L, & \left(\frac{|d\bar{\omega}|}{|d^2\bar{\omega}|} \right)_{(r_0, z_0)} &\gg L, \\ \left(\frac{|d\bar{b}|}{|d^2\bar{b}|} \right)_{(r_0, z_0)} &\gg L, & \frac{|\cos \theta_0|}{|\sin^2 \theta_0|}, \frac{|\sin \theta_0|}{|\cos^2 \theta_0|} &\gg \frac{L}{r_0}, \end{aligned} \quad (\text{A13})$$

where d and d^2 represent the first and second spatial derivatives, respectively. Equation A13 indicates that the approximation holds when \bar{b} and $\bar{\omega}$ have a large radius of curvature relative to the characteristic perturbation length scale L , the variation in $\bar{\omega}$ is small compared to its average, and the region we consider is not too close to either the equator or the pole. Under this limit, we define

$$\begin{aligned} f_0 &= 2\Omega + 2\bar{\omega}(r_0, z_0), & M_r &= f_0 + r_0 \left. \frac{\partial \bar{\omega}}{\partial r} \right|_{(r_0, z_0)}, & M_z &= r_0 \left. \frac{\partial \bar{\omega}}{\partial z} \right|_{(r_0, z_0)}, \\ b_r &= \left. \frac{\partial \bar{b}}{\partial r} \right|_{(r_0, z_0)}, & b_z &= \left. \frac{\partial \bar{b}}{\partial z} \right|_{(r_0, z_0)}. \end{aligned} \quad (\text{A14})$$

We nondimensionalize Equations A1–A5 using the rotational time scale (f_0^{-1}) and the perturbation length scale (L), such that

$$\begin{aligned} \hat{t} &= t f_0, & (\hat{u}, \hat{v}, \hat{w}) &= (u', v', w')/(f_0 L), & \hat{b} &= b'/(f_0^2 L), & \hat{\Phi} &= \Phi'/(f_0^2 L^2), \\ (\hat{r}, \hat{z}) &= (r', z')/L, & (\hat{M}_r, \hat{M}_z) &= (M_r, M_z)/f_0, & (\hat{b}_r, \hat{b}_z) &= (b_r, b_z)/f_0^2. \end{aligned} \quad (\text{A15})$$

The nondimensionalized equations become

$$\left(\frac{\partial}{\partial \hat{t}} + \hat{v} \frac{\partial}{\partial \hat{r}} + \hat{w} \frac{\partial}{\partial \hat{z}}\right) \hat{u} + \hat{M}_r \hat{v} + \hat{M}_z \hat{w} = 0, \quad (\text{A16})$$

$$\left(\frac{\partial}{\partial \hat{t}} + \hat{v} \frac{\partial}{\partial \hat{r}} + \hat{w} \frac{\partial}{\partial \hat{z}}\right) \hat{v} - \hat{u} - \cos \theta_0 \hat{b} + \frac{\partial \hat{\Phi}}{\partial \hat{r}} = 0, \quad (\text{A17})$$

$$\left(\frac{\partial}{\partial \hat{t}} + \hat{v} \frac{\partial}{\partial \hat{r}} + \hat{w} \frac{\partial}{\partial \hat{z}}\right) \hat{w} - \sin \theta_0 \hat{b} + \frac{\partial \hat{\Phi}}{\partial \hat{z}} = 0, \quad (\text{A18})$$

$$\left(\frac{\partial}{\partial \hat{t}} + \hat{v} \frac{\partial}{\partial \hat{r}} + \hat{w} \frac{\partial}{\partial \hat{z}}\right) \hat{b} + \hat{b}_r \hat{v} + \hat{b}_z \hat{w} = 0, \quad (\text{A19})$$

$$\frac{\partial \hat{v}}{\partial \hat{r}} + \frac{\partial \hat{w}}{\partial \hat{z}} + \frac{\hat{v}}{r_0/L + \hat{r}} = 0. \quad (\text{A20})$$

We neglect the last term in Equation A20 because $r_0/L \gg 1$ for the local instability problem. In the linearized instability problem, we assume that the perturbation fields are much smaller than the background fields, and obtain the first-order linear perturbation equations:

$$\frac{\partial \hat{u}}{\partial \hat{t}} + \hat{M}_r \hat{v} + \hat{M}_z \hat{w} = 0, \quad (\text{A21})$$

$$\frac{\partial \hat{v}}{\partial \hat{t}} - \hat{u} - \cos \theta_0 \hat{b} + \frac{\partial \hat{\Phi}}{\partial \hat{r}} = 0, \quad (\text{A22})$$

$$\frac{\partial \hat{w}}{\partial \hat{t}} - \sin \theta_0 \hat{b} + \frac{\partial \hat{\Phi}}{\partial \hat{z}} = 0, \quad (\text{A23})$$

$$\frac{\partial \hat{b}}{\partial \hat{t}} + \hat{b}_r \hat{v} + \hat{b}_z \hat{w} = 0, \quad (\text{A24})$$

$$\frac{\partial \hat{v}}{\partial \hat{r}} + \frac{\partial \hat{w}}{\partial \hat{z}} = 0. \quad (\text{A25})$$

We look for plane-wave solutions of the linearized perturbation equations A21–A25:

$$(\hat{u}, \hat{v}, \hat{w}, \hat{b}, \hat{\Phi}) = (A_u, A_v, A_w, A_b, A_\Phi) \exp(i\hat{k}_r \hat{r} + i\hat{k}_z \hat{z} - i\hat{\xi} \hat{t}), \quad (\text{A26})$$

where $A_u, A_v, A_w, A_b, A_\Phi$ are constant amplitudes, and $(\hat{k}_r, \hat{k}_z) = (k_r, k_z)L$.

Under these conditions and substituting Equation A26 into Equations A16–A20, we get a linearized equation system. The linearized equations must have zero determinant to have non-trivial solutions, which requires

$$\begin{vmatrix} -i\hat{\xi} & \hat{M}_r & \hat{M}_z & 0 & 0 \\ -1 & -i\hat{\xi} & 0 & -\cos \theta_0 & i\hat{k}_r \\ 0 & 0 & -i\hat{\xi} & -\sin \theta_0 & i\hat{k}_z \\ 0 & \hat{b}_r & \hat{b}_z & -i\hat{\xi} & 0 \\ 0 & i\hat{k}_r & i\hat{k}_z & 0 & 0 \end{vmatrix} = 0. \quad (\text{A27})$$

This gives

$$\hat{\xi}[(\hat{k}_r^2 + \hat{k}_z^2)\hat{\xi}^2 - \hat{b}_z \sin \theta_0 \hat{k}_r^2 - (\hat{M}_r + \hat{b}_r \cos \theta_0)\hat{k}_z^2 + 2\hat{b}_r \sin \theta_0 \hat{k}_r \hat{k}_z] = 0. \quad (\text{A28})$$

Neglecting the trivial solutions $\hat{\xi} = 0$ and $\hat{k}_r = \hat{k}_z = 0$, we have the dispersion relation:

$$\hat{\xi}^2 = \frac{\hat{b}_z \sin \theta_0 \hat{k}_r^2 + (\hat{M}_r + \hat{b}_r \cos \theta_0)\hat{k}_z^2 - 2\hat{b}_r \sin \theta_0 \hat{k}_r \hat{k}_z}{\hat{k}_r^2 + \hat{k}_z^2}. \quad (\text{A29})$$

Restoring Equation A29 back to dimensionalized form, we have the dispersion relation:

$$\xi^2 = \frac{b_z \sin \theta_0 k_r^2 + (f_0 M_r + b_r \cos \theta_0)k_z^2 - 2b_r \sin \theta_0 k_r k_z}{k_r^2 + k_z^2}. \quad (\text{A30})$$

Appendix B Instability analysis for $b_r \sin \theta_0 = 0$

In this section, we discuss the special case when $b_r \sin \theta_0 = 0$, which essentially suggests $b_r = 0$ since $\sin \theta_0 = 0$ violates Equation A13. In this case, the dispersion relation (Equation 10) becomes

$$\xi^2 = \frac{b_z \sin \theta_0 k_r^2 + f_0 M_r k_z^2}{k_r^2 + k_z^2}. \quad (\text{B1})$$

Therefore, the instability criteria reduce to $b_z \sin \theta_0 < 0$ or $f_0 M_r < 0$, i.e. the instability for perturbations in the vertical or radial directions (where the vertical is here defined as parallel to the rotational axis, while the radial is orthogonal to the rotation axis). When the system is unstable, the most unstable mode can be inferred by taking the limit $b_r \rightarrow 0$ of Equation 16, which gives

$$\lim_{b_r \rightarrow 0} \tan \delta_m \rightarrow \begin{cases} -\frac{b_r \sin \theta_0}{b_z \sin \theta_0 - f_0 M_r - b_r \cos \theta_0} \rightarrow 0 & \text{if } b_z \sin \theta_0 - f_0 M_r > 0, \\ \frac{b_z \sin \theta_0 - f_0 M_r - b_r \cos \theta_0}{b_r \sin \theta_0} \rightarrow \pm\infty & \text{if } b_z \sin \theta_0 - f_0 M_r < 0, \end{cases} \quad (\text{B2})$$

with the maximum growth rate being

$$\lim_{b_r \rightarrow 0} \sigma_m \rightarrow \begin{cases} (-f_0 M_r)^{1/2} & \text{if } b_z \sin \theta - f_0 M_r > 0, \\ (-b_z \sin \theta_0)^{1/2} & \text{if } b_z \sin \theta - f_0 M_r < 0. \end{cases} \quad (\text{B3})$$

This indicates that the most unstable mode always aligns with either the radial ($\delta_m = 0$) or vertical ($\delta_m = \pi/2$) directions, depending on the relative magnitude of the growth rate in these two directions, $(-f_0 M_r)^{1/2}$ and $(-b_z \sin \theta_0)^{1/2}$.

Appendix C Instability analysis for $N^2 < 0$ and $\eta/f < 0$

In this section, we show that $N^2 < 0$ is not a sufficient condition for instability in a zonally symmetric flow, while $\eta/f < 0$ is a sufficient condition for instability.

N^2 can be expressed as




Figure C1. Instability diagram for $\theta_0 = 60^\circ$ (other latitudes have similar results). Blue lines indicate $N^2 = 0$, and red lines indicate $\eta/f = 0$, with solid lines indicating the results for $b_r > 0$ and dashed lines indicating the results for $b_r < 0$. The black lines indicate $xy = 1$. The shading indicates the stable region. The black arrow indicates the parameters for simulation s_S . Note that for s_S , the y -axis value is 1.15×10^4 , outside the y -values shown in the plot.

$$\begin{aligned}
 N^2 &= \frac{\partial \bar{b}}{\partial R} \\
 &= b_z \sin \theta_0 + b_r \cos \theta_0 \\
 &= |b_r \sin \theta_0| \left[\frac{b_z \sin \theta_0}{|b_r \sin \theta_0|} + \frac{b_r \cos \theta_0}{|b_r| |\sin \theta_0|} \right] \\
 &= |b_r \sin \theta_0| [x + |\cot \theta_0| \text{sgn}(b_r)],
 \end{aligned} \tag{C1}$$

where $x = b_z \sin \theta_0 / |b_r \sin \theta_0|$ is defined in Equation 18, and we used the definition that $\theta_0 \in (-\pi/2, \pi/2]$ so that $\cos \theta_0 > 0$. Therefore, $N^2 < 0$ is equivalent to

$$x < -|\cot \theta_0| \text{sgn}(b_r). \tag{C2}$$

The blue lines mark Equation C2 in Figure C1 when $\theta_0 = 60^\circ$. As indicated in the plot, when $b_r < 0$, $N^2 < 0$ is not a sufficient condition for instability as the system can remain stable even when $N^2 < 0$, which is consistent with results shown in the numerical simulation s_S .

We next consider the case $\eta/f < 0$, where, for easier comparison with previous work (e.g., [Itano & Maruyama, 2009](#)), we neglect the geometric terms so that $f_0 = 2\Omega$ and $M_r = f_0 + \partial \bar{u} / \partial r$. Note that

$$\frac{1}{R} \frac{\partial \bar{u}}{\partial \theta} = \frac{\partial \bar{u}}{\partial z} \cos \theta_0 - \frac{\partial \bar{u}}{\partial r} \sin \theta_0 = M_z \cos \theta_0 - (M_r - f_0) \sin \theta_0. \tag{C3}$$

As a result,

$$\begin{aligned}
 \frac{\eta}{f} &= \frac{f_0 \sin \theta_0 - \frac{1}{R} \frac{\partial \bar{u}}{\partial \theta_0}}{f_0 \sin \theta_0} \\
 &= \frac{M_r \sin \theta_0 - M_z \cos \theta_0}{f_0 \sin \theta_0} \\
 &= \frac{M_r \sin \theta_0 - \frac{b_r \sin \theta_0 - b_z \cos \theta_0}{f_0} \cos \theta_0}{f_0 \sin \theta_0} \\
 &= \frac{f_0 M_r - b_r \cos \theta_0 + b_z \sin \theta_0 \cot^2 \theta_0}{f_0^2} \\
 &= \frac{|b_r \sin \theta_0|}{f_0^2} \left[\frac{f_0 M_r + b_r \cos \theta_0}{|b_r \sin \theta_0|} + \frac{b_z \sin \theta_0 \cot^2 \theta_0}{|b_r \sin \theta_0|} - \frac{2b_r \cos \theta_0}{|b_r \sin \theta_0|} \right] \\
 &= \frac{|b_r \sin \theta_0|}{f_0^2} [y + \cot^2 \theta_0 x - 2|\cot \theta_0| \text{sgn}(b_r)]. \tag{C4}
 \end{aligned}$$

where $y = (f_0 M_r + b_r \cos \theta_0)/|b_r \sin \theta_0|$ is defined in Equation 18. Therefore, $\eta/f < 0$ is equivalent to

$$y < -\cot^2 \theta_0 x + 2|\cot \theta_0| \text{sgn}(b_r). \tag{C5}$$

The separating line described by Equation C5 is always tangential to the hyperbolic curve, $xy = 1$ (see the red lines in Figure C1 for the case with $\theta_0 = 60^\circ$). $\eta/f < 0$ is therefore always a sufficient (although not necessary) condition for instability.

Appendix D Details of the simulation setup

The perturbation equations A16–A20 form the system that we numerically integrate in time with Dedalus (Burns et al., 2020), where we choose the domain scale of the numerical simulations as the length scale L . We neglect the last term in Equation A20, consistent with the scaling analysis in Appendix A. To search for plane-wave solutions in an infinitely large domain, we apply a Cartesian coordinate system with double-periodic boundary conditions. The equations are solved on a grid with 256×256 grid points.

The nonlinear effects may induce turbulence in the system before the most unstable mode becomes dominant when simulations are initialized with white noise. To address this issue and emphasize the initial exponential growth phase, we integrate the linearized equations (Equations A16–A20 without the advection terms, $\hat{v}\partial/\partial\hat{r}$ and $\hat{w}\partial/\partial\hat{z}$) with white noise in the \hat{b} field as the initial condition for at least $10/\hat{\sigma}_m$, where $\hat{\sigma}_m$ is the expected maximum growth rate, until the most unstable mode dominates. Subsequently, we use these modes, albeit with reduced amplitude, as the initial conditions for nonlinear simulations in each scenario (first column in Figure 6). Only the results of these nonlinear simulations are presented in this paper. In s_S , we adopt the same initial conditions as in s_G for the nonlinear simulation, representing both stable and unstable scenarios with $\text{Ro} \ll 1$.

To address the problem of grid-scale noise in the non-linear simulations without viscosity and diffusivity (Figure D1), we employ sub-grid parameterizations to represent the effects of sub-grid-scale eddy mixing via a flow-dependent eddy diffusivity (κ_e) and viscosity (ν_e), computed following the parameterization of Leith (1996). The Leith sub-grid parameterization is based on the theory of 2-D turbulence, where enstrophy cascades toward smaller scales and is ultimately dissipated at the grid scale. As a result, the eddy diffusivity and viscosity are parameterized as follows:

$$\nu_e = \kappa_e = \left(\frac{K_{Leith}}{\pi}\right)^3 L_{grid}^3 \sqrt{\left[\frac{\partial}{\partial \hat{r}} \left(\frac{\partial \hat{v}}{\partial \hat{z}} - \frac{\partial \hat{w}}{\partial \hat{r}}\right)\right]^2 + \left[\frac{\partial}{\partial \hat{z}} \left(\frac{\partial \hat{v}}{\partial \hat{z}} - \frac{\partial \hat{w}}{\partial \hat{r}}\right)\right]^2}, \quad (\text{D1})$$

where L_{grid} is the grid scale and K_{Leith} is a proportionality parameter, chosen as $K_{Leith} = 1$ in our simulations. In addition to simulations with Leith viscosity and diffusivity, we also perform simulations without any explicit viscosity and diffusivity, and compare their results (see Figures 5, 6, and D1). During the initial, approximately linear, growth stage, the two simulation sets behave similar albeit with a somewhat reduced growth rate in the simulations with eddy viscosity and diffusivity. Once the simulations become strongly nonlinear, the simulations without Leith viscosity and diffusivity develop excessive noise at small scales, although the large-scale results remain qualitatively similar over the time-scale considered in our simulations.




Figure D1. Same as Figure 6, but for simulations with no diffusivity and viscosity.

Open Research Section

Dedalus (Burns et al., 2020) code was used to perform numerical simulations used in this study. The data that support the findings of this study are openly available on Zenodo at Zeng & Jansen (2024b).

Acknowledgments

We thank Noboru Nakamura, Wanying Kang, Hao Fu, and Yixiao Zhang for helpful discussions and comments.

References

- Ashkenazy, Y., & Tziperman, E. (2021). Dynamic Europa ocean shows transient Taylor columns and convection driven by ice melting and salinity. *Nature communications*, *12*(1), 6376.
- Aurnou, J., Heimpel, M., & Wicht, J. (2007). The effects of vigorous mixing in a convective model of zonal flow on the ice giants. *Icarus*, *190*(1), 110–126.
- Bire, S., Kang, W., Ramadhan, A., Campin, J.-M., & Marshall, J. (2022). Exploring ocean circulation on icy moons heated from below. *Journal of Geophysical Research: Planets*, *127*(3), e2021JE007025.
- Bowman, J. C., & Shepherd, T. G. (1995). Nonlinear symmetric stability of planetary atmospheres. *Journal of Fluid Mechanics*, *296*, 391–407.
- Burns, K. J., Vasil, G. M., Oishi, J. S., Lecoanet, D., & Brown, B. P. (2020). Dedalus: A flexible framework for numerical simulations with spectral methods. *Physical Review Research*, *2*(2), 023068.
- Busse, F. H. (1970, November). Thermal instabilities in rapidly rotating systems. *Journal of Fluid Mechanics*, *44*(3), 441–460. Retrieved 2024-01-22, from https://www.cambridge.org/core/product/identifier/S0022112070001921/type/journal_article doi: 10.1017/S0022112070001921
- Busse, F. H., & Hood, L. (1982). Differential rotation driven by convection in a rapidly rotating annulus. *Geophysical & Astrophysical Fluid Dynamics*, *21*(1-2), 59–74.
- Busse, F. H., & Or, A. (1986). Convection in a rotating cylindrical annulus: thermal Rossby waves. *Journal of Fluid Mechanics*, *166*, 173–187.
- Callies, J., & Ferrari, R. (2018, January). Baroclinic Instability in the Presence of Convection. *Journal of Physical Oceanography*, *48*(1), 45–60. Retrieved 2024-01-22, from <https://journals.ametsoc.org/view/journals/phoc/48/1/jpo-d-17-0028.1.xml> doi: 10.1175/JPO-D-17-0028.1
- Cho, H., Shepherd, T., & Vladimirov, V. (1993). Application of the direct Liapunov method to the problem of symmetric stability in the atmosphere. *Journal of the Atmospheric Sciences*, *50*(6), 822–836.
- Christensen, U. R. (2002). Zonal flow driven by strongly supercritical convection in rotating spherical shells. *Journal of Fluid Mechanics*, *470*, 115–133.
- Eliassen, A. (1951). Slow thermally or frictionally controlled meridional circulation in a circular vortex. *Astrophysica Norvegica*, *v. 5*, p. 19, 5, 19.
- Emanuel, K. A. (1983). The Lagrangian parcel dynamics of moist symmetric instability. *Journal of Atmospheric Sciences*, *40*(10), 2368–2376.
- Emanuel, K. A. (1985). Convective adjustment in baroclinic atmospheres. *The Journal of Atmospheres*, *2441*, 163–171.
- Flasar, F. M., & Gierasch, P. J. (1978, January). Turbulent convection within rapidly rotating superadiabatic fluids with horizontal temperature gradients. *Geophysical & Astrophysical Fluid Dynamics*, *10*(1), 175–212. Retrieved 2024-05-06, from <http://www.tandfonline.com/doi/abs/10.1080/03091927808242636> doi: 10.1080/03091927808242636

- Fruman, M. D., & Shepherd, T. G. (2008, June). Symmetric Stability of Compressible Zonal Flows on a Generalized Equatorial β Plane. *Journal of the Atmospheric Sciences*, *65*(6), 1927–1940. Retrieved 2024-05-07, from <https://journals.ametsoc.org/doi/10.1175/2007JAS2582.1> doi: 10.1175/2007JAS2582.1
- Galanti, E., Kaspi, Y., Miguel, Y., Guillot, T., Durante, D., Racioppa, P., & Iess, L. (2019). Saturn’s deep atmospheric flows revealed by the cassini grand finale gravity measurements. *Geophysical Research Letters*, *46*(2), 616–624.
- Gerkema, T., Zimmerman, J., Maas, L., & Van Haren, H. (2008). Geophysical and astrophysical fluid dynamics beyond the traditional approximation. *Reviews of Geophysics*, *46*(2).
- Haine, T. W. N., & Marshall, J. (1998, April). Gravitational, Symmetric, and Baroclinic Instability of the Ocean Mixed Layer. *Journal of Physical Oceanography*, *28*(4), 634–658. Retrieved 2024-02-05, from [http://journals.ametsoc.org/doi/10.1175/1520-0485\(1998\)028<0634:GSABIO>2.0.CO;2](http://journals.ametsoc.org/doi/10.1175/1520-0485(1998)028<0634:GSABIO>2.0.CO;2) doi: 10.1175/1520-0485(1998)028<0634:GSABIO>2.0.CO;2
- Hathaway, D. H., Gilman, P. A., & Toomre, J. (1979, January). Convective instability when the temperature gradient and rotation vector are oblique to gravity. I. Fluids without diffusion. *Geophysical & Astrophysical Fluid Dynamics*, *13*(1), 289–316. Retrieved 2024-05-06, from <http://www.tandfonline.com/doi/abs/10.1080/03091927908243778> doi: 10.1080/03091927908243778
- Heimpel, M., Gastine, T., & Wicht, J. (2016). Simulation of deep-seated zonal jets and shallow vortices in gas giant atmospheres. *Nature Geoscience*, *9*(1), 19–23.
- Holm, D. D., Marsden, J. E., Ratiu, T., & Weinstein, A. (1985). Nonlinear stability of fluid and plasma equilibria. *Physics reports*, *123*(1-2), 1–116.
- Hoskins, B. J. (1974, July). The role of potential vorticity in symmetric stability and instability. *Quarterly Journal of the Royal Meteorological Society*, *100*(425), 480–482. Retrieved 2024-01-22, from <https://rmets.onlinelibrary.wiley.com/doi/10.1002/qj.49710042520> doi: 10.1002/qj.49710042520
- Itano, T., & Maruyama, K. (2009). Symmetric Stability of Zonal Flow under Full-Component Coriolis Force -Effect of the Horizontal Component of the Planetary Vorticity-. *Journal of the Meteorological Society of Japan. Ser. II*, *87*(4), 747–753. Retrieved 2024-05-07, from http://www.jstage.jst.go.jp/article/jmsj/87/4/87_4_747/_article doi: 10.2151/jmsj.87.747
- Jansen, M. F., Kang, W., Kite, E. S., & Zeng, Y. (2023). Energetic constraints on ocean circulations of icy ocean worlds. *The Planetary Science Journal*, *4*(6), 117.
- Jeffery, N., & Wingate, B. (2009, December). The Effect of Tilted Rotation on Shear Instabilities at Low Stratifications. *Journal of Physical Oceanography*, *39*(12), 3147–3161. Retrieved 2024-05-06, from <http://journals.ametsoc.org/doi/10.1175/2009JPO4138.1> doi: 10.1175/2009JPO4138.1
- Kang, W., Mittal, T., Bire, S., Campin, J.-M., & Marshall, J. (2022). How does salinity shape ocean circulation and ice geometry on enceladus and other icy satellites? *Science advances*, *8*(29), eabm4665.
- Kaspi, Y., Flierl, G. R., & Showman, A. P. (2009). The deep wind structure of the giant planets: Results from an anelastic general circulation model. *Icarus*, *202*(2), 525–542.
- Kaspi, Y., Galanti, E., Hubbard, W. B., Stevenson, D., Bolton, S., Iess, L., . . . others (2018). Jupiter’s atmospheric jet streams extend thousands of kilometres deep. *Nature*, *555*(7695), 223–226.
- Kaspi, Y., Galanti, E., Park, R., Duer, K., Gavriel, N., Durante, D., . . . others (2023). Observational evidence for cylindrically oriented zonal flows on jupiter. *Nature Astronomy*, *7*(12), 1463–1472.
- Leith, C. (1996). Stochastic models of chaotic systems. *Physica D: Nonlinear Phenomena*, *98*(2-4), 481–491.

- Marotzke, J. (1991). Influence of convective adjustment on the stability of the thermohaline circulation. *Journal of Physical Oceanography*, *21*(6), 903–907.
- O’Neill, M. E., & Kaspi, Y. (2016, October). Slantwise convection on fluid planets. *Geophysical Research Letters*, *43*(20). Retrieved 2024-01-22, from <https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2016GL071188> doi: 10.1002/2016GL071188
- O’Neill, M. E., Kaspi, Y., & Fletcher, L. N. (2017). Galileo probe interpretation indicating a neutrally stable layer in the jovian troposphere. *Geophysical Research Letters*, *44*(9), 4008–4017.
- Ooyama, K. (1966). On the stability of the baroclinic circular vortex: A sufficient criterion for instability. *Journal of Atmospheric Sciences*, *23*(1), 43–53.
- Shepherd, T. G. (1990). Symmetries, conservation laws, and hamiltonian structure in geophysical fluid dynamics. In *Advances in geophysics* (Vol. 32, pp. 287–338). Elsevier.
- Soderlund, K. M. (2019). Ocean dynamics of outer solar system satellites. *Geophysical Research Letters*, *46*(15), 8700–8710.
- Stevens, D. E. (1983). On symmetric stability and instability of zonal mean flows near the equator. *Journal of the Atmospheric Sciences*, *40*(4), 882–893.
- Stone, P. H. (1966). On non-geostrophic baroclinic stability. *Journal of the Atmospheric Sciences*, *23*(4), 390–400.
- Stone, P. H. (1967). An application of baroclinic stability theory to the dynamics of the jovian atmosphere. *Journal of Atmospheric Sciences*, *24*(6), 642–652.
- Stone, P. H. (1971, January). The symmetric baroclinic instability of an equatorial current. *Geophysical Fluid Dynamics*, *2*(1), 147–164. Retrieved 2024-05-06, from <https://www.tandfonline.com/doi/full/10.1080/03091927108236055> doi: 10.1080/03091927108236055
- Straneo, F., Kawase, M., & Riser, S. C. (2002, February). Idealized Models of Slantwise Convection in a Baroclinic Flow. *Journal of Physical Oceanography*, *32*(2), 558–572. Retrieved 2024-01-22, from [http://journals.ametsoc.org/doi/10.1175/1520-0485\(2002\)032<0558:IMOSCI>2.0.CO;2](http://journals.ametsoc.org/doi/10.1175/1520-0485(2002)032<0558:IMOSCI>2.0.CO;2) doi: 10.1175/1520-0485(2002)032<0558:IMOSCI>2.0.CO;2
- Sun, W. (1995, January). Unsymmetrical symmetric instability. *Quarterly Journal of the Royal Meteorological Society*, *121*(522), 419–431. Retrieved 2024-05-07, from <https://rmets.onlinelibrary.wiley.com/doi/10.1002/qj.49712152209> doi: 10.1002/qj.49712152209
- Vallis, G. K. (2017). *Atmospheric and oceanic fluid dynamics*. Cambridge University Press.
- Walton, I. C. (1975, April). The viscous nonlinear symmetric baroclinic instability of a zonal shear flow. *Journal of Fluid Mechanics*, *68*(4), 757–768. Retrieved 2024-01-22, from https://www.cambridge.org/core/product/identifier/S0022112075001218/type/journal_article doi: 10.1017/S0022112075001218
- Zeng, Y., & Jansen, M. (2024b, November). *Simulation data for "symmetric instability in a boussinesq fluid on a rotating planet"*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.15086582> doi: 10.5281/zenodo.15086582
- Zeng, Y., & Jansen, M. F. (2021). Ocean circulation on enceladus with a high-versus low-salinity ocean. *The Planetary Science Journal*, *2*(4), 151.
- Zeng, Y., & Jansen, M. F. (2024a). The effect of salinity on ocean circulation and ice–ocean interaction on enceladus. *The Planetary Science Journal*, *5*(1), 13.