# Distributed Facility Location Games with Candidate Locations

Feiyue Sun

School of Mathematical Sciences, Ocean University of China, Qingdao, China 1127327435@qq.com

Abstract. We study the distributed facility location games with candidate locations, where agents on a line are partitioned into groups. Both desirable and obnoxious facility location settings are discussed. In distributed location problems, distortion can serve as a standard for quantifying performance, measuring the degree of difference between the actual location plan and the ideal location plan. For the desirable setting, under the max of sum cost objective, we give a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\sqrt{2} + 1$ . Under the sum of max cost objective, we give a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$ . Under the max of max cost, we get a strategyproof distributed mechanism with 3-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$ . For the obnoxious setting, under three social objectives, we present that there is no strategyproof mechanism with bounded distortion in the case of discrete candidate locations, and no group strategyproof mechanism with bounded distortion in the case of continuous candidate locations.

Keywords: Distributed facility location· Mechanism design· Distortion· Strategyproofness.

# 1 Introduction

Facility location problems are one of the most important in social choice, and involve selecting the optimal location for a facility to meet the target requirements of different groups, given the known positions of the participants. Procaccia and Tennenholtz [\[6\]](#page-11-0) used facility location to present the problem of approximate mechanism design without monetary payments. Due to incomplete information, strategic agents may misreport their private information (such as location) to reduce their costs, then a mechanism refers to a function that maps the reported position information from agents to the facility location. Therefore, it is necessary to design incentive-compatible mechanisms to encourage agents to truthfully report their positions, ensuring that some social objective is as good as possible. If a mechanism can guarantee that no benefit of the agents can be obtained by misreporting their locations, it is said to be strategyproof. Facility

location often leads to outcomes that do not accurately select the optimal location owing to multiple reasons.

The distributed facility location problem involves grouping agents into several regions. The mechanism in distributed facilities is carried out in two steps: first, selecting a point within each group that represents the reported positions of the agents in that group, and then choosing the location of the facility from among these points. Filos-Ratsikas et al. [\[10\]](#page-11-1) conducted the first study on distributed distortion from the perspective of voting. The decision is made by the following mechanism: the mechanism first selects a representative voter for each region, and then chooses one of the representatives as the winner. The distortion is the worst-case ratio of the social objective function value achieved by the mechanism to the optimal social objective function value, and used as a systematic way of comparing different mechanisms. Procaccia and Rosenschein [\[1\]](#page-11-2) originally proposed the notion of distortion to measure the loss in social welfare due to mechanisms by the preference rankings of the agents over the possible outputs, rather than their true cardinal costs.

In the classic models, all points in the metric space are feasible for building facilities. However, this is often impractical in many applications. For example, due to land use restrictions, the facilities can only be built in some feasible regions. Therefore, the facility location game with candidate locations is studied to find one or more locations to establish facilities in order to optimize objective functions (such as minimizing costs, maximizing profits, etc.) within a given set of candidate locations. It is considered both in the discrete setting, where the set of candidates (alternatives) is some finite subset of  $\mathbb{R}$ , as well as in the continuous setting, where the set of alternatives is the whole  $\mathbb R$  or some interval. For the single-facility case, Feldman et al. [\[16\]](#page-12-0) considered the candidate voting problem, which can be converted into a single-facility location problem, and where the facility can only be placed at candidate locations. They proposed a deterministic group strategy-proof mechanism with approximation ratio of 3 for minimizing social cost.

In the desirable facility location game, each agent wants to stay as close to the facility as possible. And obnoxious facility location game is considered where each agent wants to keep as far away from the located facility as possible. Objective functions for obnoxious facility location game are often referred to as social utility functions. The obnoxious facility location game was first introduced by Cheng et al. [\[13\]](#page-11-3) . For deterministic mechanisms on a line, a 3-approximation group strategy-proof mechanism was given and the lower bound was proved to be 2.

Above all, we study the distributed facility location games with candidate locations, where agents on a line are partitioned into groups. Both desirable and obnoxious facility location games are discussed.

#### 1.1 Our work

We study in distributed facility location with candidate locations, where a set of agents are positioned on the line and are partitioned into groups. Both desirable and obnoxious facilities are discussed.

In the desirable facility location problem, three objective functions, which are the maximum cost of any agent within any group, the maximum of the sum of costs of the agents in each group, and the sum of the maximum costs of the agents in each group, are discussed. And we proposed three strategyproof mechanisms under discrete candidate locations as opposed to continuous cases by [\[12\]](#page-11-4). We remark that our problem with discrete locations is more complex than that continuous cases, because the latter problem corresponds to a version in which every point in the line is a candidate location.

Under the max of sum cost objective, we give a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\sqrt{2}+1$ . Under the sum of max cost objective, we give a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$ . We get a strategyproof distributed mechanism with 3-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$  under the max of max cost. A summary of our results is illustrated in Table [1.](#page-2-0)

<span id="page-2-0"></span>Table 1. Upper and lower bounds of distortions for distributed facility location with discrete candidate locations.

	Upper bound	Lower bound
Max-Max		
$Sum-Max$		
Max-Sum		$\sqrt{2} + 1$

Furthermore, we discuss the distributed obnoxious facility problem with candidate locations, based on three objective functions, which are the minimum utility of any agent within any group, the minimum of the sum of utilities of the agents in each group, and the sum of the minimum utilities of the agents in each group, respectively. We present that there is no strategyproof mechanism with bounded distortion in the case of discrete candidate locations, and no group strategyproof mechanism with bounded distortion in the case of continuous candidate locations.

### 1.2 Related work

As one of the most important in social choice theory, facility location problems have been widely studied. Anshelevich and Postl [\[3\]](#page-11-5) examined randomized social

choice functions that require only the ordinal information and select an alternative that is good in expectation with respect to the costs from the metric. Boutilier et al. [\[4\]](#page-11-6) adopted a utilitarian perspective on social choice, assuming that agents have (possibly latent) utility functions over some space of alternatives. They proposed optimal social choice functions under three different models, and underscored the important role played by scoring functions.

Strategyproof facility location problems were studied by Procaccia and Tennenholtz [\[6\]](#page-11-0) as part of their approximate mechanism design. They argued that approximation can be leveraged to obtain truthfulness without resorting to payments, and established tight upper and lower bounds for the approximation ratio given by strategyproof mechanisms without payments, with respect to both deterministic and randomized mechanisms. Subsequently, the literature for strategyproof is also rather extensive. Moulin [\[5\]](#page-11-7) proposed a complete characterization of strategy-proof mechanisms when participants are located on a line. Fotakis and Tzamos [\[7\]](#page-11-8) were interested in mechanisms without payments that are (group) strategyproof for given cost function, and achieve a good approximation ratio for the social cost and/or the maximum cost of the agents. Serafino and Ventre [\[8\]](#page-11-9) proposed the heterogeneous facility location problem under two different objective functions, social cost (i.e., sum of all agents' costs) and maximum cost, and studied the approximation ratio of both deterministic and randomized truthful algorithms under the simplifying assumption that the underlying network topology would be a line. Lu et al. [\[9\]](#page-11-10) designed strategy-proof mechanisms without payment that have a small approximation ratio for social cost. Tang et al. [\[17\]](#page-12-1) provided a deterministic strategy-proof mechanism with an approximation ratio of 3 for minimizing the maximum cost. Additionally, for the two-facility case, they introduced an anonymous group strategy-proof mechanism with approximation ratio of  $(2n-3)$  for minimizing social cost, and approximation ratio of 3 for minimizing the maximum cost.

The distortion of mechanisms has been studied for several social choice problems. Based on minimizing social costs, Filos-Ratsikas and Voudouris [\[11\]](#page-11-11) provided distortions for continuous and discrete cases respectively. Anshelevich et al. [\[2\]](#page-11-12) studied the distortion of distributed mechanisms for variations of well-known objectives, such as the (average) total cost and the maximum cost, and chose a single alternative aiming to optimize various objectives that were functions of the distances between agents and alternatives in the metric space, under the constraint that this choice must be made in a distributed way; Filos-Ratsikas et al. [\[12\]](#page-11-4) improved the distortion result of the unrestricted mechanism to 2 when minimizing social costs for continuous cases, and also provided the distortions of the unrestricted mechanism and the strategyproof mechanism under the objective functions of the maximum cost of any agent within any group, the maximum of the sum of costs of the agents in each group, and the sum of the maximum costs of the agents in each group. Zou and Li [\[14\]](#page-11-13) investigated the facility location problem with double preferences, where participants may lie about both their own locations and their preferences for the facility. When both the locations and preferences of participants are private information, they gave a group

strategy-proof deterministic mechanism with approximation ratio of  $1/3$ . Ibara and Nagamochi [\[15\]](#page-12-2) showed that there is no bounded distortion strategy-proof mechanism in the line metric if the number of candidates is at least three, which implies the lower bound 3 for maxSum objective and 5 for maxSOS objective for obnoxious facility location game.

# 2 Preliminaries

A tuple  $I = (N, \mathbf{x}, D, Z)$  (abbreviated as I) is considered as an instance, where

- N is a set of n agents.
- $-\mathbf{x} = (x_1, \dots, x_i, \dots, x_n)$  is a position profile consisting of position  $x_i \in \mathbb{R}$ with agent  $i \in N$ .
- $-D = \{d_1, \dots, d_k\}$  is a set of  $k \ge 1$  groups into which all agents are divided. Each group d contains a set  $N_d \subseteq N$  of agents such that  $N_{d_i} \cap N_{d_j} = \emptyset$  for  $i \neq j$  and  $\bigcup_{d \in D} N_d = N$ . And  $\lambda_i$  is the number of  $N_{d_i}$  agents in group  $d_i$ .
- $Z$  is a set of candidate locations, including both continuous and discrete cases.

A distributed mechanism M is designed to output the facility location from Z. In detail, M includes two steps: firstly, select a point (called representative) from Z within each group d based on the reported positions of the agents in d, denoted by  $z_d$ , and then choose the location  $\omega$  of the facility from among all representatives  $z_d$  as the output. For  $z \in Z$ , the distance between  $x_i$  and z is denoted by  $\delta(x_i, z) = |x_i - z|$ .

#### 2.1 Social objectives

In desirable facility location, we consider minimizing the following three social objectives:

– The Sum of Max cost  $(SoM)$  of z is the sum of the maximum cost of all agents in each group:

$$
SoM(z|I) := \sum_{d \in D} \max_{i \in N_d} \delta(x_i, z). \tag{1}
$$

– The Max of Sum cost  $(MoS)$  of location z is the maximum sum of costs of all agents in each group:

$$
MoS(z|I) := \max_{d \in D} \sum_{i \in N_d} \delta(x_i, z). \tag{2}
$$

– The Max of Max cost  $(M \circ M)$  of location z is the maximum cost among all the agents:

$$
MoM(z|I) := \max_{i \in N} \delta(x_i, z) = \max_{d \in D} \max_{i \in N_d} \delta(x_i, z).
$$
 (3)

In obnoxious facility location, we consider maximizing the following three social objectives:

– The Sum of Min utility (*Som*) of location z is the sum of the minimum utility of all agents in each group:

$$
Som(z|I) := \sum_{d \in D} \min_{i \in N_d} \delta(x_i, z). \tag{4}
$$

– The Min of Sum utility  $(m \circ S)$  of location z is the minimum of the sum of all agents in each group:

$$
moS(z|I) := \min_{d \in D} \sum_{i \in N_d} \delta(x_i, z). \tag{5}
$$

– The Min of Min utility (mom) of location z is the minimum utility of all agents:

$$
mom(z|I) := \min_{i \in N} \delta(x_i, z) = \min_{d \in D} \min_{i \in N_d} \delta(x_i, z). \tag{6}
$$

We denote social objectives as  $SO(z|I)$  to facilitate a unified definition of distortion.

### 2.2 Strategyproofness and distortion

A significant characterization of designing mechanisms is resilient to strategic manipulation, i.e., they prevent agents to affect the output in their favor (reaching smaller cost in desirable facility location or increasing her utility in obnoxious setting) by misreporting positions.

Generally there are two important properties for both desirable and obnoxious mechanism: strategyproofness (SP) and group strategyproofness (GSP). SP implies that the mechanism does not allow the agents to gain any benefit by reporting false positions, that is, the truth-telling strategy is dominant for all agents. Formally, a mechanism M is strategyproof: for any agent  $i \in N$ , if agent i reports a false position  $x_i'$ , it holds that

$$
\delta(x_i, M(N, \mathbf{x}, D, Z)) \le \delta(x_i, M(N, (\mathbf{x}_{-i}, x'_i), D, Z))
$$
\n(7)

where  $\mathbf{x}_{-i} = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$  is the location profile without  $x_i$  for any agent i. Correspondingly, in the obnoxious case, the strategygproof mechanism should be satisfied by

$$
\delta(x_i, M(N, \mathbf{x}, D, Z)) \ge \delta(x_i, M(N, (\mathbf{x}_{-i}, x_i'), D, Z))).
$$
\n(8)

A stronger property is group strategyproofness (GSP), that is, no any set of agents can misreport the positions simultaneously such that each member of the set gains benefit. Formally speaking, given any location profile  $\mathbf{x} = (\mathbf{x}_{-S}, \mathbf{x}_{S})$ with any no-empty subset  $S \in N$ , there exists at least one agent  $i \in S$  such that

$$
\delta(x_i, M(N, \mathbf{x}, D, Z)) \le \delta(x_i, M(N, (\mathbf{x}_{-S}, \mathbf{x}'_S), D, Z))
$$
\n(9)

where  $\mathbf{x}'_S$  is the misreported location profile of all agents in S. Correspondingly, in the obnoxious case, GSP holds that

$$
\delta(x_i, M(N, \mathbf{x}, D, Z)) \ge \delta(x_i, M(N, (\mathbf{x}_{-S}, \mathbf{x}'_S), D, Z)).
$$
\n(10)

The distortion of a distributed mechanism M with respect to some social objective is the worst-case ratio between the social objective of the location chosen by the mechanism and the minimum social objective. Set  $opt(I) :=$  $\min_{z\in Z} SO(z|I),$ 

$$
dist(M|I) = \sup_{I} \frac{SO(M|I)}{opt(I)}.
$$
\n(11)

Similarly, in the the obnoxious case, the distortion of a mechanism M is defined by the social utility. Set  $opt(I) := \max_{z \in Z} SO(z|I)$ 

$$
dist(M|I) = \sup_{I} \frac{opt(I)}{SO(M|I)}.\t(12)
$$

# 3 Desirable facility location games with candidate locations

In this section, we discuss the distributed desirable facility location games with candidate locations under three social objectives, i.e. the Sum of Max cost, the Max of Sum cost and the Max of Max cost. Consider that  $n$  agents located at  $x_i \in \mathbb{R}$  are divided into k groups. The facility location would be selected from m discrete candidate locations, denoted by  $Z = \{z_j | z_j \in \mathbb{R}, j = 1, \dots, m\}$ . Under the Sum of Max cost, we give a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$ . Under the Max of Sum cost, we propose a strategyproof distributed mechanism with 5-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\sqrt{2}+1$ . We get a strategyproof distributed mechanism with 3-distortion, and prove that no strategyproof mechanism can have a distortion better than  $\frac{\sqrt{5}+1}{2}$  under the Max of Max cost.

### 3.1 Sum of Max

In this subsection, we focus on the sum-of-max objective,

$$
SoM(z|I) = \sum_{d \in D} \max_{i \in N_d} \delta(x_i, z). \tag{13}
$$

Theorem 1. For Sum-of-Max, the distortion of any strategyproof mechanism is at least  $\frac{1+\sqrt{5}}{2}$ .

Proof. The proof process is detailed in Appendix 1.

**Mechanism 1** (Median-of-Rightmost): For each group  $d \in D$ **do**  $z_d := \arg\min_{z \in Z} \delta(r_d, z);$ **return**  $\omega := \frac{1}{2}k$ -th leftmost<sup>1</sup> representative;

Theorem 2. Mechanism 1 is strategyproof.

*Proof.* Consider any instance I, and let  $\omega$  be the facility location chosen by the mechanism. For each group d,  $x_i$  denotes the location of agent  $i \in N_d$  and  $r_d$ denotes the rightmost agent in d. Mechanism 1 chooses the rightmost agent in d, then the misreported location  $x'_i \leq r_d$  implies no decrease in the cost of agent *i*. We only discuss  $x'_i > r_d$ .

- Case 1:  $\omega < z_d$ . If  $z'_d := \arg\min_{z \in Z}$ z∈Z  $\delta(x'_i, z) \in [\omega, z_d)$ , due to the definition of  $z_d$ , i.e. no candidate in  $(r_d - \delta(r_d, z_d), r_d + \delta(r_d, z_d))$ ,  $z'_d$  only locates in  $[\omega, r_d - \delta(r_d, z_d)]$ .  $\delta(x'_i, z_d) < \delta(x'_i, z'_d)$  could be derived by  $z'_d < z_d$  and  $x'_i > r_d$ , and there is a contradiction with the definition of  $z'_d$ . Therefore,  $z'_d$ must satisfy  $z_d' \ge z_d > \omega$ .
- Case 2:  $z_d \leq \omega$ . We have  $r_d < \omega$  with the definition of  $z_d$ . To choose the different alternative  $z_d'$ , agent i only misreports her location in  $x_i' > r_d +$  $\delta(r_d, z_d)$ , then  $z'_d \geq z_d$ .

Above all, for any agent  $i \in N$ , the social cost could not decrease by misreporting the location. Therefore, Mechanism 1 is strategyproof.

Theorem 3. For Sum-of-Max, the distortion of Mechanism 1 is at most 5.

Proof. The proof process is detailed in Appendix 2.

#### 3.2 Max of Sum

In this subsection, we focus on the max-of-sum objective,

$$
MoS(z|I) = \max_{d \in D} \sum_{i \in N_d} \delta(x_i, z). \tag{14}
$$

Theorem 4. For Max-of-Sum, the distortion of any strategyproof mechanism is at least  $1 + \sqrt{2}$ .

Proof. The proof process is detailed in Appendix 3.

**Mechanism 2** (Rightmost-of-Median): For each group  $d_i \in D$ **do**  $x_{\frac{\lambda_i}{2}} := \frac{1}{2}\lambda_i$ -th leftmost<sup>3</sup> agent;  $z_d := \arg\min_{z \in Z}$  $\delta(x_{\frac{\lambda_i}{2}},z);$ return  $\omega := \text{rightmost representative};$ 

<sup>&</sup>lt;sup>1</sup> Mechanism 1 chooses the  $\lceil \frac{k}{2} \rceil$ -th leftmost representative as the output. To simplify our notation, we will use  $\frac{1}{2}\overline{k}$ -th to denote the ceiling.

<sup>&</sup>lt;sup>3</sup> Mechanism chooses the  $\lceil \frac{1}{2} \lambda_i \rceil$ -th leftmost agent. For simplicity, we will use  $\frac{1}{2} \lambda_i$ -th to denote the ceiling.

### Theorem 5. Mechanism 2 is strategyproof.

*Proof.* Consider any instance I, and let  $\omega$  be the facility location chosen by the mechanism,  $D_{\omega}$  be a set of groups with  $z_d = \omega$ . Without loss of generality, we assume that  $x_i < \omega$ . Mechanism 2 chooses the rightmost representative, then for any  $i \in d \notin D_\omega$ , the misreported location  $x_i'$  implies no effect on the output  $\omega$ . We only discuss  $d_i \in D_\omega$ .

- Case 1:  $x_i < x_{\lambda_j}$ . If agent *i* misreports her position to change the output, he/she would choose that  $x'_i > x_{\frac{\lambda_j}{2}}$ . Notice that  $\delta(x'_i, \omega) < \delta(x_{\frac{\lambda_j}{2}}, \omega)$ , and the output could be still.
- Case 2:  $x_i > x_{\frac{\lambda_j}{2}}$ . If agent i misreports her position to change the output, he/she would choose that  $x'_i < x_{\frac{\lambda_j}{2}}$ . If Mechanism 2 reaches a new output  $\omega'$ by  $x'_i$ , then  $\omega' < x_{\frac{\lambda_j}{2}} < \omega$ . Combining with  $\delta(x_{\frac{\lambda_j}{2}}, \omega') > \delta(x_{\frac{\lambda_j}{2}}, \omega)$ , we have

$$
\delta(x_i, \omega') > \delta(x_{\frac{\lambda_j}{2}}, \omega') > \delta(x_{\frac{\lambda_j}{2}}, \omega) > \delta(x_i, \omega),\tag{15}
$$

i.e.  $\delta(x_i, \omega) < \delta(x_i, \omega')$ . The social cost could not decrease by misreporting the location.

Besides, if  $x_i = x_{\frac{\lambda_j}{2}}$ , there is no need to misreport for agent *i*. In conclusion, Mechanism 2 is strategyproof.

Theorem 6. For Max-of-Sum, the distortion of Mechanism 2 is at most 5.

Proof. The proof process is detailed in Appendix 4.

### 3.3 Max of Max

In this subsection, we focus on the max-of-max objective,

$$
MoM(z|I) = \max_{d \in D} \max_{i \in N_d} \delta(x_i, z).
$$
\n(16)

Theorem 7. For Max-of-Max, the distortion of any strategyproof mechanism is at least  $\frac{1+\sqrt{5}}{2}$ .

Proof. We suppose towards a contradiction that there is a strategyproof mechanism  $M_3$  with distortion strictly smaller than  $\frac{1+\sqrt{5}}{2}$ , i.e. dist $(M_3|I) < \frac{1+\sqrt{5}}{2}$ , where  $\omega$  and  $\sigma$  be the location chosen by the mechanism and the optimal one. respectively.

Consider an instance  $I_7$  with a group with one agent at 0 and the other agent at 1, and there are three alternatives at  $z_1 = \frac{3-\sqrt{5}}{2}$ ,  $z_2 = 1$ ,  $z_3 = \frac{1+\sqrt{5}}{2}$ . We have that  $M \circ M(z_3 | I_7) = \frac{\sqrt{5}+1}{2} > M \circ M(z_2 | I_7) = 1 > M \circ M(z_1 | I_7) =$ 

 $\frac{\sqrt{5}-1}{2}$ , and  $opt(I_7) = SoM(z_1|I_7)$ . We claim that  $M_3$  must choose  $z_1 = 1$  as the representative as otherwise the distortion would be  $\frac{1+\sqrt{5}}{2}$ .

Consider another instance  $I_8$  with a group with one agent at 1 and the other agent at 2, and there are three alternatives at  $z_1 = \frac{3-\sqrt{5}}{2}$ ,  $z_2 = 1$ ,  $z_3 = \frac{1+\sqrt{5}}{2}$ . As a result,  $M_1$  must choose  $z_3 = z_3 = \frac{1+\sqrt{5}}{2}$  as the representative as otherwise the distortion would be at least  $\frac{1+\sqrt{5}}{2}$ . This follows by arguments similar to those for instance  $I_1$ .

To reach a contradiction, we consider the following instance  $I_9$  with two groups and  $Z = \{z_1 = \frac{3-\sqrt{5}}{2}, z_2 = 1, z_3 = \frac{1+\sqrt{5}}{2}\}$ :

- In  $d_1$ , there are two agents at 0 and 1, respectively.
- In  $d_2$ , there are two agents at 1 and 2, respectively.

By  $I_7$  and  $I_8$ , the representatives of two groups must be  $z_1$  and  $z_3$ , respectively. Consequently, the final winner would be  $z_1$  or  $z_3$ . However,  $MoM(z_1|I_9)$  =  $SolI(z_3|I_9) = \frac{1+\sqrt{5}}{2}$ , while  $opt(I_9) = SoM(z_2|I_9) = 1$ , reaching a distortion of  $\frac{1+\sqrt{5}}{2}$ . Hence, we prove this theorem as desired.

**Mechanism 3** (Median-of-Median): For each group  $d_i \in D$ 

**do** 
$$
x_{\frac{\lambda_i}{2}} := \frac{1}{2}\lambda_i
$$
-th leftmost agent;  
\n $z_d := \arg\min_{z \in Z} \delta(x_{\frac{\lambda_i}{2}}, z);$   
\n**return**  $\omega := \frac{1}{2}k$ -th leftmost representative;

Theorem 8. Mechanism 3 is strategyproof.

The proof is the same as the mechanism DM in [\[11\]](#page-11-11).

Theorem 9. For Max-of-Max, the distortion of Mechanism 3 is at most 3.

The proof can be referred in [\[2\]](#page-11-12).

# 4 Obnoxious facility location games with candidate locations

In this section, we discuss the distributed single obnoxious facility location problem under several minimum utility objectives. Consider a set of n agents located along [0, 1]. One facility would be selected from alternatives set  $Z = [0, 1]$  (continuous candidate locations) and  $Z = \{z_j | z_j \in [0,1], j = 1, \cdots, m\}$  (discrete candidate locations), respectively. The agent utility is her distance to the output by the mechanism.

### 4.1 Sum of Min

<span id="page-9-0"></span>We focus on the sum-of-min objective,

$$
Som(z|I) = \sum_{d \in D} \min_{i \in N_d} \delta(x_i, z). \tag{17}
$$

Theorem 10. For Sum-of-Min, no group strategyproof mechanism for obnoxious facility location in  $Z = [0, 1]$  can have a bounded distortion.

<span id="page-10-0"></span>Proof. The proof process is detailed in Appendix 5.

Theorem 11. For Sum-of-Min, no strategyproof mechanism for obnoxious facility location in  $Z = \{z_i \in [0, 1], i = 1, \dots, m\}$  can have a bounded distortion.

Proof. We suppose that there is a strategyproof mechanism M with bounded distortion of B, i.e.  $\frac{Som(o)}{Som(\omega)} \leq B$ . Consider two candidate locations  $z_1 = 0, z_2 = 1$ , and two agents in one group, where  $x_1 = \epsilon^2$  and  $x_2 = 1 - \epsilon$  with  $0 < \epsilon <$  $\min\{\frac{\sqrt{5}-1}{2},\frac{1}{B}\}\.$  We have that  $x_1 < x_2$ , and  $opt(I) = Som(1) = \epsilon$ , By distortion B, mechanism M outputs point 1. Otherwises,  $\frac{Som(1)}{Som(0)} = \frac{1}{\epsilon} > B$ . And so, social utility of agent at  $x_2$  is  $\epsilon$ .

If agent at  $x_2$  misreports her location as  $1 - \epsilon^3$ , the optimal solution of new profile is point 0. By similar arguments above, the finial output must return point 0. Therefore, agent at  $x_2$  can change the output of M and increase her utility by misreporting. There is a contradiction to the strategyproofness.

Theorem [10](#page-9-0) and [11](#page-10-0) can be also applicable to the objective function of Min-of-Min  $mom(\omega|I) = mom(\omega) = \min_{d \in D} \min_{i \in N_d} \delta(x_i, \omega).$ 

### 4.2 Min of Sum

We focus on the social welfare of the min-of-sum objective,

$$
moS(z|I) = moS(z) = \min_{d \in D} \sum_{i \in N_d} \delta(x_i, z).
$$

Theorem 12. For Min-of-Sum, no group strategyproof mechanism for obnoxious facility location in  $Z = [0, 1]$  can have a bounded distortion.

Proof. The proof process is detailed in Appendix 6.

Theorem 13. For Min-of-Sum, no strategyproof mechanism for obnoxious facility location in  $Z = \{z_i \in [0,1], i = 1, \cdots, m\}$  can have a bounded distortion.

Proof. We suppose that there is a strategyproof mechanism M with bounded distortion of B, i.e.  $\frac{m\delta S(o)}{m\delta S(\omega)} \leq B$ . Consider two candidate locations  $z_1 = 0, z_2 = 1$ , and there are two groups and two agents with one agent in each group, where  $d_1 = \{i = 1, x_1 = \epsilon^2\}$  and  $d_2 = \{i = 2, x_2 = \epsilon^2\}$  with  $0 < \epsilon < \min\{\frac{\sqrt{5}-1}{2}, \frac{1}{B}\}.$ We have that  $x_1 < x_2$ , and  $opt(I) = moS(1) = \epsilon$ , By distortion B, mechanism M outputs point 1. Otherwises,  $\frac{moS(1)}{moS(0)} = \frac{1}{\epsilon} > B$ . And so, social utility of agent at  $x_2$  is  $\epsilon^2$ .

If agent at  $x_2$  misreports her location as  $1 - \epsilon^3$ , the optimal solution of new profile is point 0. By similar arguments above, the finial output must return point 0. Therefore, agent at  $x_2$  can change the output of M and increase her utility by misreporting. There is a contradiction to the strategyproofness.

## 5 Conclusions

In this paper, we purposed bounds on the distortion of single-winner distributed mechanisms for many different objectives, and for obnoxious single facility games, we showed no bounded strategyproof mechanism for discrete candidate locations and no bounded group strategyproof mechanism for continuous candidate locations under several minimum utility objectives, some of which are novel. Still, there are several challenging open questions, as well as new directions for future research. Starting with our results, it would be interesting to close the gaps between the lower and upper bounds presented in Table 1 for the various scenarios we considered. For obnoxious cases, we would consider to design 2-candidate mechanism under some maximum objectives.

# <span id="page-11-2"></span>References

- 1. Procaccia A, Rosenschein J: The distortion of cardinal preferences in voting. In: International Workshop on Cooperative Information Agents (CIA), pp. 317-–331. Springer(2006)
- <span id="page-11-12"></span>2. Anshelevich E, Filos-Ratsikas A, Voudouris A A: The distortion of distributed metric social choice. Artificial intelligence 308, 103713 (2022)
- <span id="page-11-5"></span>3. Anshelevich E, Postl J: Randomized social choice functions under metric preferences. Journal of Artificial Intelligence Research 58, 797-–827 (2017)
- <span id="page-11-6"></span>4. Boutilier C, Caragiannis I, Haber S, et al: Optimal social choice functions: A utilitarian view. Artificial Intelligence 227, 190—213 (2015)
- <span id="page-11-7"></span>5. Moulin H: On strategy-proofness and single peakedness. Public Choice 35(4), 437– 455 (1980)
- <span id="page-11-0"></span>6. Procaccia A, Tennenholtz M, Approximate Mechanism Design without Money. ACM Transactions on Economics and Computation 1 4(18), 1–26 (2013)
- <span id="page-11-8"></span>7. Fotakis D, Tzamos C: Strategyproof facility location for concave cost functions. Algorithmica 76 1, 143–167 (2016)
- <span id="page-11-9"></span>8. Serafino P, Ventre C: Heterogeneous facility location without money. Theoretical Computer Science 636, 27–46 (2016)
- <span id="page-11-10"></span>9. Lu P, Sun X, Wang Y, Zhu Z: Asymptotically optimal strategy-proof mechanisms for two-facility games. In: Proceedings of the 11th ACM Conference on Electronic Commerce (EC), pp. 315–324. ACM, New York (2010)
- <span id="page-11-1"></span>10. Filos-Ratsikas A, Micha E, Voudouris A: The distortion of distributed voting. Artificial Intelligence, 286: 103343 (2020)
- <span id="page-11-11"></span>11. Filos-Ratsikas A, Voudouris A: Approximate mechanism design for distributed facility location. In: Algorithmic Game Theory: 14th International Symposium, pp. 49–63. Springer International Publishing, Aarhus (2021)
- <span id="page-11-4"></span>12. Filos-Ratsikas A, Kanellopoulos P, Voudouris A, et al: Settling the distortion of distributed facility location. arxiv preprint arxiv, 2301.01604 (2023)
- <span id="page-11-3"></span>13. Cheng Y, Yu W, Zhang G: Strategy-proof approximation mechanisms for an obnoxious facility game on networks. Theoretical Computer Science 497, 154–163 (2013)
- <span id="page-11-13"></span>14. Zou S, Li M: Facility location games with dual preference. In: Proceedings of the 2015 international conference on autonomous agents and multiagent systems, pp. 615–623. Istanbul (2015)
- <span id="page-12-2"></span>15. Ibara K, Nagamochi H: Characterizing mechanisms in obnoxious facility game. In: International conference on combinatorial optimization and applications. pp. 301– 311. Springer Berlin Heidelberg, Heidelberg (2012)
- <span id="page-12-0"></span>16. Feldman M, Fiat A, Golomb I: On voting and facility location. In: Proceedings of the 17th ACM Conference on Economics and Computation (ACM-EC), pp. 269–286. (2016)
- <span id="page-12-1"></span>17. Tang Z, Wang C, Zhang M, Zhao Y: Mechanism design for facility location games with candidate locations. In: Proceedings of the 14th International Conference on Combinatorial Optimization and Applications (COCOA), pp. 440–452. (2020)

# 6 Appendix

# 6.1 Appendix 1

Proof. We suppose towards a contradiction that there is a strategyproof mechanism  $M_1$  with distortion strictly smaller than  $\frac{1+\sqrt{5}}{2}$ , i.e. dist $(M_1|I) < \frac{1+\sqrt{5}}{2}$ .

Consider an instance  $I_1$  with one agent at  $\frac{3-\sqrt{5}}{2}$  in a group and three alternatives at  $z_1 = 0, z_2 = 1, z_3 = 2$ . We have that  $S \circ M(z_3|I_1) = \frac{\sqrt{5}+1}{2}$  $SolN(z_2|I_1) = \frac{\sqrt{5}-1}{2} > SoM(z_1|I_1) = \frac{3-\sqrt{5}}{2}$ , and  $opt(I_1) = SoM(z_1|I_1) = \frac{3-\sqrt{5}}{2}$ . We claim that  $M_1$  must choose  $z_1 = 0$  as the representative, otherwise the distortion would be  $\frac{1+\sqrt{5}}{2}$ .

Consider another instance  $I_2$  with one agent at  $\frac{1+\sqrt{5}}{2}$  in a group and three alternatives at  $z_1 = 0$ ,  $z_2 = 1$ ,  $z_3 = 2$ . As a result,  $M_1$  must choose  $z_3 = 2$  as the representative, otherwise the distortion would be at least  $\frac{1+\sqrt{5}}{2}$ . This follows by arguments similar to those for instance  $I_1$ .

To reach a contradiction, we consider the following instance  $I_3$  with two groups:

- In  $d_1$ , there are one agent at  $\frac{3-\sqrt{5}}{2}$  and three alternatives at  $z_1 = 0$ ,  $z_2 = 1$ ,
- $z_3 = 2$ .<br>– In  $d_2$ , there are one agent at  $\frac{1+\sqrt{5}}{2}$  and three alternatives at  $z_1 = 0$ ,  $z_2 = 1$ ,  $z_3 = 2.$

By  $I_1$  and  $I_2$ , the representatives of two groups must be  $z_1 = 0$  and  $z_3 =$ 2, respectively. Consequently, the final winner would be  $z_1$  or  $z_3$ . However,  $S \circ M(z_1|I_3) = S \circ M(z_3|I_3) = \frac{3-\sqrt{5}}{2} + \frac{1+\sqrt{5}}{2} = 2$ , while  $opt(I_3) = S \circ M(z_2|I_3) =$  $\sqrt{5} - 1$ , reaching a distortion of  $\frac{2}{\sqrt{5}}$ .  $\frac{2}{5-1} = \frac{1+\sqrt{5}}{2}.$ 

# 6.2 Appendix 2

*Proof.* Let  $\omega$  and  $\delta$  be the location chosen by the mechanism and the optimal one, respectively.  $i_d$  denotes the agent in the group d, whose distance from w is the maximum, and  $i_{d^*}$  is the agent with the farthest distance from  $o$  in each

group  $d$ . We have that

$$
S o M(\omega) = \sum_{D} \delta(x_{i_d}, \omega) \le \sum_{D} \delta(x_{i_d}, o) + \sum_{D} \delta(o, \omega)
$$
  

$$
\le \sum_{D} \delta(x_{i_{d^*}}, o) + k \delta(o, \omega) \le S o M(o) + k \delta(o, \omega).
$$
 (18)

– Case 1:  $o < \omega$ . Let  $L = \{d \in D, z_d < \omega\}$ ,  $R = \{d \in D, z_d \geq \omega\}$ , and it is easily derived that  $|R| \geq \frac{k}{2}$ , which  $|\cdot|$  represents the number of groups. If  $\delta(r_d, o) < \delta(r_d, \omega)$  for any  $d \in R$ , then  $z_d$  could be o, and  $z_d < \omega$ . Thus, there is the contraction with  $d \in R$ . We have that  $\delta(r_d, o) \geq \delta(r_d, \omega)$ ,  $d \in R$ . By the triangle inequality, we obtain

<span id="page-13-0"></span>
$$
\delta(o,\omega) \le \delta(o,r_d) + \delta(r_d,\omega) \le 2\delta(o,r_d), d \in R. \tag{19}
$$

Hence,

$$
SoM(o) = \sum_{D} \delta(o, x_{i_{d^*}}) \ge \sum_{R} \delta(o, x_{i_{d^*}}) \ge \sum_{R} \delta(o, r_d) \ge \frac{k}{4} \delta(o, \omega), \quad (20)
$$

with inequality [\(18\)](#page-13-0), we have  $S \circ M(\omega) \leq 5 \cdot S \circ M(\omega)$  as desired.

– Case 2:  $\omega < \omega$ . Let  $L' = \{d \in D, z_d \le \omega\}$ ,  $R' = \{d \in D, z_d > \omega\}$ . Then, we get that  $|L'| \geq \frac{k}{2}$ . We have that

$$
\delta(o,\omega) \le \delta(o,r_d) + \delta(r_d,\omega) \le 2\delta(o,r_d), d \in L'.
$$
\n(21)

It holds that

$$
SoM(o) = \sum_{D} \delta(o, x_{i_{d^*}}) \ge \sum_{L'} \delta(o, x_{i_{d^*}}) \ge \sum_{L'} \delta(o, r_d) \ge \frac{k}{4} \delta(o, \omega) \tag{22}
$$

Therefore, we obtain that  $Sol(\omega) \leq 5 \cdot SoM(o)$  with inequality [\(18\)](#page-13-0).

#### 6.3 Appendix 3

Proof. We suppose towards a contradiction that there is a strategyproof mechanism  $M_2$  with distortion strictly smaller than  $1 + \sqrt{2}$ , i.e. dist $(M_2|I) < 1 + \sqrt{2}$ .

Consider an instance  $I_4$  with a group with  $(1+\sqrt{2})p$  agents at 0 and p agents at 1, where  $p$  is an arbitrarily large integer<sup>2</sup>. And three alternatives locate at  $z_1 = 0, z_2 = 1, z_3 = 2$ . We have that  $MoS(z_3|I_4) = (2\sqrt{2} + 3)p > MoS(z_2|I_4) =$  $(\sqrt{2} + 1)p > Mos(z_1|I_4) = p$ , and  $opt(I_4) = Mos(z_1|I_4)$ . We claim that  $M_2$ must choose  $z_1 = 0$  as the representative as otherwise the distortion would be  $\frac{1 + \sqrt{2}}{1 + \sqrt{2}}$ 

Consider another instance  $I_5$  with a group with p agents at 1 and  $(1 + \sqrt{2})p$ agents at 2, and three alternatives are  $z_1 = 0$ ,  $z_2 = 1$  and  $z_3 = 2$ . Therefore,  $M_2$ must choose  $z_3 = 2$  as the representative. This follows by arguments similar to those for instance  $I_4$ .

At last, we consider the following instance  $I_6$  with two groups and  $Z = \{z_1 =$  $0, z_2 = 1, z_3 = 2$ :

<sup>2</sup>  $(1 + \sqrt{2})p$  should be an integer  $[(1 + \sqrt{2})p]$ . For simplicity, we drop the ceiling.

- In  $d_1$ , there are  $(1 + \sqrt{2})p$  agents at 0 and p agents at 1.
- In  $d_2$ , there are p agents at 1 and  $(1 + \sqrt{2})p$  agents at 2.

By  $I_3$  and  $I_4$ , the representatives of two groups must be  $z_1 = 0$  and  $z_3 =$ 2, respectively. Consequently, the final winner would be  $z_1$  or  $z_3$ . However,  $M_0S(z_1|I_6) = M_0S(z_3|I_6) = (3 + 2\sqrt{2})p$ , while  $opt(I_6) = S_0M(z_2|I_6) = (1 +$  $\sqrt{2}$ )p, reaching a distortion of  $(1+\sqrt{2})$ . There is a contradiction with dist $(M_2|I)$  <  $\sqrt{2}$ .

### 6.4 Appendix 4

*Proof.* Let  $\omega$  and  $\delta$  be the location chosen by the mechanism and the optimal one, respectively.  $d^*$  denotes the group where  $\sum_{i \in N_d} \delta(x_i, \omega)$  is the maximum. We have that

<span id="page-14-1"></span><span id="page-14-0"></span>
$$
MoS(\omega) = \sum_{i \in N_{d^*}} \delta(x_i, \omega) \le \sum_{i \in N_{d^*}} \delta(x_i, o) + \sum_{i \in N_{d^*}} \delta(o, \omega)
$$
  

$$
\le MoS(o) + \lambda_j \delta(o, \omega).
$$
 (23)

 $- \text{ Case 1: } o < \omega. \text{ If } x_{\frac{\lambda_j}{2}} \leq \omega \text{ for any } d \in D_\omega, \text{ we have that } \delta(x_{\frac{\lambda_j}{2}}, o) \geq \delta(x_{\frac{\lambda_j}{2}}, \omega).$ Let  $S_d = \{i \in N_d, x_i \geq x_{\frac{\lambda_j}{2}}\}$ , and  $|S_d| \geq \frac{\lambda_j}{2}$ . By the triangle inequality, we obtain that  $\delta(o, \omega) = \delta(o, x_{\frac{\lambda_j}{2}}) + \delta(x_{\frac{\lambda_j}{2}}, \omega) \leq 2\delta(o, x_{\frac{\lambda_j}{2}})$ , and

$$
\lambda_j \delta(o, \omega) \le 2\lambda_j \delta(o, x_{\frac{\lambda_j}{2}}) \le 4 \sum_{i \in S_d} \delta(o, x_{\frac{\lambda_j}{2}}) \le 4 \sum_{i \in N_d} \delta(o, x_i) \le 4 \cdot MoS(o).
$$
\n(24)

If there exists a group d such that  $x_{\frac{\lambda_j}{2}} > \omega$ . Let  $S'_d = \{i \in N_d, x_i \geq x_{\frac{\lambda_j}{2}}\},\$ and  $|S'_a$  $\vert d'_{d} \vert \geq \frac{\lambda_{j}}{2}$ . We have that

$$
MoS(o) \geq \sum_{i \in N_d} \delta(x_i, o) \geq \sum_{i \in S'_d} \delta(x_i, o) \geq \sum_{i \in S'_d} \delta(o, \omega) \geq \frac{\lambda_j}{2} \delta(o, \omega), \quad (25)
$$

i.e.  $\lambda_j \delta(o, \omega) \leq 2 \cdot MoS(o)$ . With inequality [\(23\)](#page-14-0) and [\(24\)](#page-14-1),  $MoS(\omega) \leq 5 \cdot$  $MoS(o).$ 

- Case 2:  $o > \omega$ . Let  $T_d = \{i \in N_d, x_i \leq x_{\frac{\lambda_i}{2}}\}$ , and  $|T_d| \geq \frac{\lambda_j}{2}$ , and we have that  $\delta(x_{\frac{\lambda_j}{2}}, o) \geq \delta(x_{\frac{\lambda_j}{2}}, \omega)$ . If  $x_{\frac{\lambda_j}{2}} \leq \omega$ , we obtain that

<span id="page-14-2"></span>
$$
\lambda_j \delta(o, \omega) \le \lambda_j \delta(o, x_{\frac{\lambda_j}{2}}) \le 2 \sum_{T_d} \delta(x_i, o) \le 2 \sum_{N_d} \delta(x_i, o) \le 2 \cdot MoS(o). \tag{26}
$$

If  $\omega < x_{\lambda_j} < \delta$ , by the triangle inequality, we draw the conclusions that  $\delta(o, \omega) = \overline{\delta(o, x_{\frac{\lambda_j}{2}})} + \delta(x_{\frac{\lambda_j}{2}}, \omega) \leq 2\delta(o, x_{\frac{\lambda_j}{2}}),$  and

$$
\lambda_j \delta(o, \omega) \le 2\lambda_j \delta(o, x_{\frac{\lambda_j}{2}}) \le 4 \sum_{T_d} \delta(o, x_{\frac{\lambda_j}{2}}) \le 4 \sum_{N_d} \delta(o, x_i) \le 4 \cdot MoS(o)
$$
\n(27)

with inequality [\(23\)](#page-14-0) and [\(26\)](#page-14-2),  $MoS(\omega) \leq 5 \cdot MoS(o)$ . Above all, the conclusion can be proved.

### 6.5 Appendix 5

*Proof.* We suppose that there is a group strategyproof mechanism  $M$  with bounded distortion of B, i.e.  $\frac{Som(o)}{Som(\omega)} \leq B$ . Consider the instance with  $N = N_1 +$  $N_2$  agents in one group, where  $N_1 = \left[\frac{1-\epsilon-\epsilon^2}{\epsilon^3}\right]$  $\frac{\epsilon-\epsilon^2}{\epsilon^3}$  + 2 and  $N_2 = \left[\frac{1}{\epsilon^2}\right] + 1 = \left[\frac{\epsilon-\epsilon^3}{\epsilon^3}\right]$  $\frac{-\epsilon^3}{\epsilon^3}]+2$ with  $0 < \epsilon < \min\{\frac{\sqrt{5}-1}{2}, \frac{1}{B}\}\.$  Let the leftmost agent  $x_l = \epsilon^2$  and the rightmost one  $x_r = 1 - \epsilon$ . In  $N_1$  agents, two agents locate at  $x_l$  and  $x_r$ , respectively, and others of  $N_1$  are uniformly distributed in the interval  $(\epsilon^2, 1 - \epsilon)$ . This ensures the distance between two adjacent points in  $N_1$  that

$$
\frac{x_r - x_l}{N_1 - 1} \le \frac{1 - \epsilon - \epsilon^2}{\frac{1 - \epsilon - \epsilon^2}{\epsilon^3}} < \epsilon^2. \tag{28}
$$

Set  $N_2$  agents all at  $x_r$ , and so, the optimal solution is point 1,  $opt(I)$  $Som(1) = \epsilon$ . By the approximate radio B, we have  $Som(\omega) > \epsilon^2$ . Hence, M must choose the facility in  $(1 - \epsilon + \epsilon^2, 1]$ .

If all  $N_2$  agents at  $x_r$  misreport their locations as uniformly distributed in the interval  $[1 - \epsilon, 1 - \epsilon^3]$  with two of these agents at both endpoints. Therefore, the distance between two adjacent points in the new profile of  $N_2$  agents is also smaller than  $\epsilon^2$ :

$$
\frac{1 - \epsilon^3 - 1 + \epsilon}{N_2 - 1} \le \frac{\epsilon - \epsilon^3}{\frac{\epsilon - \epsilon^3}{\epsilon^3}} < \epsilon^2. \tag{29}
$$

And we have the distance between two adjacent points of  $N$  agents becomes smaller than  $\epsilon^2$ , denote the new profile as  $I'$ ,  $opt(I') = Som(0|I') = \epsilon^2$ . Combing with B, we derive  $Som(\omega) > \epsilon^3$ , and M must return the interval  $[0, \epsilon^2 - \epsilon^3)$ . Consequently, by misreporting, the group of  $N_2$  agents can change the output and all increase their utilities, a contradiction to the group strategyproofness of M.

### 6.6 Appendix 6

*Proof.* We suppose that there is a group strategyproof mechanism  $M$  with bounded distortion of B, i.e.  $\frac{m\delta(s)}{m\delta(s)} \leq B$ . Consider the instance with  $N =$  $N_1+N_2$  agents in N groups, i.e. one agent in each group, where  $N_1=[\frac{1-\epsilon-\epsilon^2}{\epsilon^3}]$  $\frac{\epsilon-\epsilon^2}{\epsilon^3}]+2$ and  $N_2 = \left[\frac{1}{\epsilon^2}\right]+1 = \left[\frac{\epsilon - \epsilon^3}{\epsilon^3}\right]$  $\frac{-\epsilon^3}{\epsilon^3}$  + 2 with  $0 < \epsilon < \min\{\frac{\sqrt{5}-1}{2}, \frac{1}{B}\}\$ . Let the leftmost agent  $x_l = \epsilon^2$  and the rightmost one  $x_r = 1 - \epsilon$ . In  $N_1$  agents (or groups), two agents locate at  $x_l$  and  $x_r$ , respectively, and others of  $N_1$  are uniformly distributed in the interval  $(\epsilon^2, 1 - \epsilon)$ . This ensures the distance between two adjacent points in  $N_1$  that

$$
\frac{x_r - x_l}{N_1 - 1} \le \frac{1 - \epsilon - \epsilon^2}{\frac{1 - \epsilon - \epsilon^2}{\epsilon^3}} < \epsilon^2. \tag{30}
$$

Set  $N_2$  agents all at  $x_r$ , and so, the optimal solution is point 1,  $opt(I)$  $m\delta(1) = \epsilon$ . By the approximate radio B, we have  $m\delta(\omega) > \epsilon^2$ . Hence, M must choose the facility in  $(1 - \epsilon + \epsilon^2, 1]$ .

If all  $N_2$  agents at  $x_r$  misreport their locations as uniformly distributed in the interval  $[1 - \epsilon, 1 - \epsilon^3]$  with two of these agents at both endpoints. Therefore, the distance between two adjacent points in the new profile of  $N_2$  agents is also smaller than  $\epsilon^2$ :

$$
\frac{1 - \epsilon^3 - 1 + \epsilon}{N_2 - 1} \le \frac{\epsilon - \epsilon^3}{\frac{\epsilon - \epsilon^3}{\epsilon^3}} < \epsilon^2. \tag{31}
$$

And we have the distance between two adjacent points of  $N$  agents becomes smaller than  $\epsilon^2$ , denote the new profile as  $I'$ ,  $opt(I') = mos(0|I') = \epsilon^2$ . Combing with B, we derive  $moS(\omega) > \epsilon^3$ , and M must return the interval  $[0, \epsilon^2 - \epsilon^3)$ . Consequently, by misreporting, the group of  $N_2$  agents can change the output and all increase their utilities, a contradiction to the group strategyproofness of  $\cal M.$ 

