GAUSS MAPS OF MÖBIUS SURFACES IN THE *n*-DIMENSIONAL SPHERE

DAVID BRANDER, SHIMPEI KOBAYASHI, AND PENG WANG

ABSTRACT. In this note we discuss Gauss maps for Möbius surfaces in the *n*-sphere, and their applications in the study of Willmore surfaces. One such "Gauss map", naturally associated to a Willmore surface that has a dual Willmore surface, is the Lorentzian 2-plane bundle given by a lift of the suface and its dual. More generally, we define the concept of a Lorentzian 2-plane lift for an arbitrary Möbius surface, and show that the conformal harmonicity of this lift is equivalent to the Willmore condition for the surface. This clarifies some previous work of F. Hélein, Q. Xia-Y Shen, X. Ma and others, and, for instance, allows for the treatment of the Björling problem for Willmore surfaces in the presence of umbilics.

INTRODUCTION

Gauss maps of submanifolds have been extensively used in the study of special submanifolds, including minimal submanifolds. To study surfaces in an *n*-dimensional Möbius space, the *conformal Gauss map*, which represents the central sphere congruence, proves particularly useful [2]. It is known that a surface is Willmore if and only if its conformal Gauss map is conformally harmonic [5], [10].

The conformal Gauss map can be regarded as a map into the Grassmannian symmetric space of 4-dimensional Lorentzian subspaces in the (n + 2)-dimensional Minkowski space. This formulation enables an application of integrable systems methods. From the harmonicity of the conformal Gauss map, one can derive a family of flat connections, leading to a family of moving frames, i.e., a map into the loop group of the indefinite orthogonal group SO⁺(1, n+1)as in (3.2). This perspective was initiated in [11] and further developed in [8].

While the conformal Gauss map uniquely characterizes a Möbius surface and its harmonicity defines a Willmore surface, it has some limitations. For example, defining the conformal Gauss map requires information about a 4-dimensional Lorentzian subspace determined by second-order contact. Additionally, at umbilic points, the conformal Gauss map degenerates, complicating the analysis. To address these issues, a more geometrically intuitive Gauss map has been proposed. This alternative map targets the Grassmannian of 2-dimensional Lorentzian planes in the 4-dimensional Lorentzian subspace, [11], [14].

Date: December 17, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary 53A31; 53C43, Secondary 58E20; 53C35.

Key words and phrases. Willmore surfaces; Gauss maps; harmonic maps; flat connections .

The first named author is partially supported by Independent Research Fund Denmark, grant 9040-00196B. The second named author is partially supported by JSPS KAKENHI Grant Number JP18K03265, JP22K03304. The third named author is supported by the Project 12371052 of NSFC.

This Gauss map can be easily constructed for special Willmore surfaces (the so-called S-Willmore surfaces, [10] and Section 3.2 below). It is defined by Lorentzian planes and denoted by $Z = [Y \wedge \hat{Y}]$, where Y is a lift of a Willmore surface, and \hat{Y} is its dual Willmore surface, and Z is a conformal harmonic map. S. Ma extended this approach to a broader class of Willmore surfaces, including S-Willmore surfaces, and introduced the concept of the "adjoint surface", where \hat{Y} is another Willmore surface, not necessarily dual to Y ([14]). Moreover, this idea was thoroughly explored in [11], where a family of such Gauss maps was considered. F. Héléin characterized Willmore surfaces under the condition of "roughly harmonicity" and applied integrable systems methods.

In this note, we define a generalization of this Gauss map for all surfaces in Möbius space, referred to as an \mathbb{L} -lift (Definition 2). While the choice of an \mathbb{L} -lift is not unique, this flexibility allows for a special selection such that an \mathbb{L} -lift is conformal. Note that the conformality is characterized by the Riccati equation; thus, such an \mathbb{L} -lift exists locally but not globally, in general. (However, there exist many globally defined conformal \mathbb{L} -lifts, see Remark 2.1). Then, using a conformal \mathbb{L} -lift, we establish the equivalence between the harmonicity of the \mathbb{L} -lift and the Willmore property of the surface. We finally provide a characterization of Willmore surfaces through integrable systems methods (Theorem 3.1).

In Section 3, we focus on S-Willmore surfaces and the adjoint transformation of a Willmore surface, analyzing them via an \mathbb{L} -lift with a Riccati equation. Furthermore, we investigate the properties of the moving frame associated with these structures (Proposition 3.5).

The characterization of Willmore surfaces via the \mathbb{L} -lift presented in this short paper forms the basis for addressing the Björling problem. (The problem involves reconstructing surfaces from boundary curves and the Gauss map data along those curves). Our approach will be further explored in a forthcoming paper [3], focusing on minimal surfaces in \mathbb{S}^n and \mathbb{H}^n .

Acknowledgments: The third named author is thankful to Prof. Xiang Ma and Prof. Zhenxiao Xie for valuable discussions.

1. Preliminaries

1.1. Möbius surfaces in \mathbb{S}^n and the conformal Gauss maps. The following Möbius equivalence is known [2]: $\mathbb{S}^n \cong \mathbb{P}(\mathcal{C}^{n+1}_+)$ by $x \leftrightarrow [(1, x)]$, where

$$\mathcal{C}_{+}^{n+1} = \left\{ v \in \mathbb{R}^{n+1,1} \mid \langle v, v \rangle = -v_0^2 + \sum_{j=1}^{n+1} v_j^2 = 0, \, v_0 > 0 \right\} \subset \mathbb{R}^{n+1,1}$$

is the forward light cone in the n + 2-dimensional Minkowski space $\mathbb{R}^{n+1,1}$. We define a *Möbius surface* to be an immersion $y: M \to \mathbb{S}^n \cong \mathbb{P}(\mathcal{C}^{n+1}_+)$, where the Riemann surface M has the conformal structure induced by any Möbius equivalent choice of Riemannian metric on \mathbb{S}^n .

Let M be a Riemann surface with conformal coordinate z and $y: M \to \mathbb{S}^n$ be a conformal immersion. Let $Y: \mathbb{D} \subset M \to \mathcal{C}^{n+1}_+$ be a local lift of y, that is, [1, y] = [Y] holds. It is known that the conformality of y is equivalent to $\langle Y_z, Y_z \rangle = 0$ and $\langle Y_z, Y_{\bar{z}} \rangle > 0$, where the subscript z (resp. \bar{z}) denotes the partial derivative with respect to z (resp. \bar{z}) and \langle , \rangle denotes the complex bilinear extension of the inner product of $\mathbb{R}^{n+1,1}$. Let us decompose the trivial bundle $M \times \mathbb{R}^{n+1,1}$ as $V \oplus V^{\perp}$ with

(1.1)
$$V = \operatorname{span} \left\{ Y, \operatorname{Re} Y_z, \operatorname{Im} Y_z, Y_{z\bar{z}} \right\},$$

and V^{\perp} denotes the orthogonal complement of V in $\mathbb{R}^{n+1,1}$. Clearly V is independent of the choice of a lift Y and the coordinate z, thus V is a Lorentzian rank four sub-bundle. Let us denote $V_{\mathbb{C}}$ (resp. $V_{\mathbb{C}}^{\perp}$) as the complexification of V (resp. V^{\perp}). Thus we reach the following definition:

Definition 1. For a Möbius surface $y : M \to \mathbb{S}^n$, let V be the sub-bundle in (1.1) and denote an evaluation at $p \in M$ by V_p . Then the map

$$V: M \to \operatorname{Gr}_{3,1} = \operatorname{SO}^+(1, n+1) / (\operatorname{SO}^+(1, 3) \times \operatorname{SO}(n-2)),$$

defined by $V(p) = [V_p]$ will be called the *conformal Gauss map* of y. Here SO(n-2) and $SO^+(1, n+1)$ denote degree n orthogonal group and the identity component of degree n+2 indefinite orthogonal group, respectively.

Remark 1.1. It is well known that the conformal Gauss map V is obtained by the mean curvature 2-spheres of a Möbius surface in \mathbb{S}^n , the central sphere congruence, through the identification $\mathbb{S}^n \cong P(\mathcal{C}^{n+1}_+)$, and it is conformal for any Möbius surface.

1.2. The canonical lift, the Willmore energy and Willmore surfaces. The canonical lift with respect to z is defined by a lift Y such that

(1.2)
$$|dY|^2 = |dz|^2$$

holds, that is, $\langle Y_z, Y_{\bar{z}} \rangle = 1/2$ holds. For the canonical lift Y, it is easy to see that Y_{zz} is orthogonal to $Y_z, Y_{\bar{z}}$ and $Y_{z\bar{z}}$, and there exists a complex function $s : \mathbb{D} \to \mathbb{C}$ and a section $\kappa \in \Gamma(V_{\mathbb{C}}^{\perp})$ such that

(1.3)
$$Y_{zz} + \frac{1}{2}sY = \kappa$$

holds, that is, Y satisfies the *inhomogeneous Hill's equation*. The function s defined above will be called the *Schwarzian derivative* of the immersion y with respect to the coordinate z and it will be denoted by $S_z(y) := s$, and the section κ will be called the *Hopf differential* of the immersion y. The reasons why we respectively call s and κ the Schwarzian derivative and the Hopf differential will be given below. (See Formula (23) and (24) of [6]).

Proposition 1.2 ([6]). The Schwarzian derivative s and the Hopf differential κ of an immersion y are invariant under Möbius transformations. Moreover, the quadratic differential $\kappa \frac{dz^2}{|dz|}$ is globally defined on M with values in $\mathcal{L}V^{\perp} \otimes \mathbb{C}$ and $\mathcal{L} = (K\bar{K})^{-1/2}$, where K is the canonical bundle ¹ of M and $\kappa \frac{dz^2}{|dz|} = \frac{\mathbb{I}^{(2,0)}}{|dy|}$ holds, where $\mathbb{I}^{(2,0)}$ denotes the (2,0)-part of the normal bundle-valued second fundamental form of y.

Remark 1.3. For another complex coordinate w, the canonical lift \tilde{Y} with respect to w is given by $\tilde{Y} = Y |w_z|$ in terms of the canonical lift Y with respect to coordinate z because

¹The real line bundle $\mathcal{L} = (K\bar{K})^{-1/2}$ has been called the 1-density bundle, see [6].

of (1.2). Then by the invariance of $\kappa \frac{dz^2}{|dz|}$, the coefficient function $\tilde{\kappa}$ with respect to w can be represented as

(1.4)
$$\widetilde{\kappa} = \kappa \frac{|w_z|}{w_z^2}.$$

Moreover, the Schwarzian derivative $S_z(y)$ satisfies the transformation rule:

(1.5)
$$S_w(y)dw^2 = (S_z(y) - S_z(w))dz^2,$$

where $S_z(w)$ is the classical Schwarzian derivative of w with respect to z. More explicitly, $s = S_z(y)$ and $\tilde{s} = S_w(y)$ satisfy the following transformation rule,

(1.6)
$$\tilde{s} = \left(\frac{1}{w'}\right)^2 \left(s - \left(\frac{w''}{w'}\right)' + \frac{1}{2} \left(\frac{w''}{w'}\right)^2\right).$$

Here we use the abbreviation $w' = w_z$.

Theorem 1.4 ([5], [10]). The Möbius transformation invariant energy

$$W(y) = 2\sqrt{-1} \int_M \langle \kappa, \,\overline{\kappa} \rangle \, dz \wedge d\bar{z}$$

will be called the Willmore energy of a conformal immersion $y : M \to \mathbb{S}^n$. Moreover, a surface which attains a critical point of the Willmore energy will be called a Willmore surface. Then a conformal immersion $y : \mathbb{D} \to \mathbb{S}^n$ is Willmore if and only if

(Willmore)
$$D_{\bar{z}}D_{\bar{z}}\kappa + \frac{1}{2}\bar{s}\kappa = 0$$

holds. Here κ and s are the Hopf differential and the Schwarzian derivative of the surface y, respectively. Moreover, $y : \mathbb{D} \to \mathbb{S}^n$ is Willmore if and only if the conformal Gauss map $V : \mathbb{D} \to \operatorname{Gr}_{3,1}$ is conformal harmonic.

2. LORENTZIAN 2-PLANE LIFTS (L-LIFTS) OF A MÖBIUS SURFACE

2.1. Lorentzian 2-plane lifts (L-lifts). The conformal Gauss map V becomes degenerate at umbilic points. Hélein [11] defined an alternative map which does not have that problem, essentially as follows.

For a Willmore surface y with a dual Willmore surface \hat{y} , there is (provided $Y \wedge \hat{Y} \neq 0$) a naturally defined bundle of Lorentzian 2-planes defined by $[Y \wedge \hat{Y}] : M \to \operatorname{Gr}_{1,1} = \operatorname{SO}^+(1, n + 1)/(\operatorname{SO}^+(1, 1) \times \operatorname{SO}(n))$. We generalize this first to an arbitrary Möbius surface:

Definition 2. Let $y : M \to \mathbb{S}^n$ be a Möbius surface, and let V be as defined above. A Lorentzian 2-plane lift or \mathbb{L} -lift is a map $Z : M \to \operatorname{Gr}_{1,1}$ such that, for any lift Y of y, we have $Y(p) \subset Z(p) \subset V_p$ for all $p \in M$.

Remark 2.1. We will be interested in conformal \mathbb{L} -lifts. These exist on a neighbourhood of a point (see below), but global existence would induce a bundle decomposition $V = Z \oplus Z^{\perp}$ and it may be only true under a topological assumption. However, it is known that global conformal \mathbb{L} -lifts exist for special classes of Willmore surfaces:

(1) Every minimal surface in the space forms \mathbb{S}^n , \mathbb{R}^n and \mathbb{H}^n has a natural conformal \mathbb{L} -lift associated to its dual surface [11, 12, 15].

- (2) Under some conditions, in the sense of Bryant [5] or Babich-Bobenko [1], a minimal surface in \mathbb{R}^n or \mathbb{H}^n , can extend smoothly across infinity by conformally embedding \mathbb{R}^n or \mathbb{H}^n into \mathbb{S}^n . But the L-lifts associated to the dual surface cannot extend to the possible ends or infinite boundary since these points are the unibilic points of the surfaces.
- (3) The homogeneous Willmore torus constructed by Eiiri [10] is not S-Willmore and it admits two different global adjoint surfaces, thus two different global conformal L-lifts.

If Y is a canonical lift, then we may construct a Lorentzian 2-plane lift by considering a section $\widehat{Y} \in \Gamma(V)$ satisfying

(2.1)
$$\langle \hat{Y}, \hat{Y} \rangle = 0, \quad \langle \hat{Y}, Y \rangle = -1.$$

Then a 2-plane bundle given by $Z = [Y \wedge \widehat{Y}]$ is well defined: the choice of \widehat{Y} does not depend on the conformal coordinate: if w = w(z) then the canonical lift Y changes to $\tilde{Y} = |w_z|Y$ and $\hat{\tilde{Y}}$ defined by $|w_z|^{-1}\hat{Y}$ also satisfies (2.1), and thus $Y \wedge \hat{Y} = \tilde{Y} \wedge \hat{\tilde{Y}}$ is well defined.

Lemma 2.2. Let $Z: M \to \operatorname{Gr}_{1,1}$ be a Lorentzian 2-plane subbundle of $M \times \mathbb{R}^{n+1,1}$, spanned by null vector fields Y and \hat{Y} satisfying $\langle Y, \hat{Y} \rangle = -1$. Then Z is conformal if and only if the function θ defined by:

$$\theta := 2\langle Y \wedge Y_z, \widehat{Y} \wedge \widehat{Y}_z \rangle$$

is zero.

Proof. A computation using $\langle Y, Y \rangle = \langle \hat{Y}, \hat{Y} \rangle = 0$ and $\langle Y, \hat{Y} \rangle = -1$.

2.2. Existence of conformal L-lifts and the Riccati equation. We now consider the problem of finding \widehat{Y} such that $Z = [Y \wedge \widehat{Y}]$ is conformal. To get an explicit condition on \widehat{Y} , at least locally, we choose a canonical frame for V, and write down the equations that the coefficients of \widehat{Y} in this basis must satisfy.

For the bundle $V = \text{span} \{Y, \text{Re} Y_z, \text{Im} Y_z, Y_{z\bar{z}}\}$, we define an alternative basis

$$\{Y, \operatorname{Re} Y_z, \operatorname{Im} Y_z, N\},\$$

where N is the unique section of $\Gamma(V)$ such that

(2.2)
$$\langle N, N \rangle = 0, \quad \langle N, Y \rangle = -1, \quad \langle N, Y_z \rangle = \langle N, Y_{\overline{z}} \rangle = 0.$$

Since V is a rank four sub-bundle, the above equations determine the unique section N. However, the conditions on N are not invariant under coordinate changes.

Given Y and \hat{Y} as in the previous section, with

(2.3)
$$\langle \hat{Y}, \hat{Y} \rangle = 0, \quad \langle \hat{Y}, Y \rangle = -1,$$

in $\mathbb{R}^{n+1,1}$, let us define a complex function $\mu : \mathbb{D} \to \mathbb{C}$ as follows:

(2.4)
$$\widehat{Y} = N + \overline{\mu}Y_z + \mu Y_{\overline{z}} + \frac{1}{2}|\mu|^2 Y.$$

Э

Under the change of coordinate w = w(z) the canonical lift Y changes as $|w_z|Y$ and \widehat{Y} changes $|w_z|^{-1}\widetilde{Y}$. Thus the complex function μ has the following transformation rule:

(2.5)
$$\mu = \tilde{\mu}w_z + \frac{w_{zz}}{w_z}.$$

We now consider the derivative of \widehat{Y} from (2.4), and by using (1.3) we obtain

$$\widehat{Y}_z = \frac{\mu}{2}\widehat{Y} + \theta\left(Y_{\overline{z}} + \frac{\overline{\mu}}{2}Y\right) + \rho\left(Y_z + \frac{\mu}{2}Y\right) + L,$$

where we define

(2.6)
$$\theta = \mu_z - \frac{1}{2}\mu^2 - s \in \mathbb{C},$$

(2.7)
$$\rho = \overline{\mu}_z - 2\langle \kappa, \overline{\kappa} \rangle \in \mathbb{C},$$

(2.7)
$$\rho = \bar{\mu}_z - 2\langle \kappa, \bar{\kappa} \rangle \in \mathbb{C},$$

(2.8)
$$L = 2D_{\bar{z}}\kappa + \bar{\mu}\kappa \in \Gamma(V_{\mathbb{C}}^{\perp}),$$

and D denotes the normal connection. Note that the function s and the section κ are respectively the Schwarzian derivative and the Hopf differential. A straightforward computation shows that θ in (2.6) and ρ in (2.7) can be computed as

$$\theta = 2\langle Y \wedge Y_z, \widehat{Y} \wedge \widehat{Y}_z \rangle, \quad \rho = 2\langle Y \wedge Y_{\overline{z}}, \widehat{Y} \wedge \widehat{Y}_z \rangle,$$

and thus θdz^2 and $\rho dz d\bar{z}$ are well-defined (2,0)- and (1,1)-forms, respectively.

It is known that $\theta = 0$ can be locally solved for a complex function μ :

Lemma 2.3 (Riccati equation). For a given function s, there exist local solutions μ to the following differential equation

(2.9)
$$\mu_z - \frac{1}{2}\mu^2 - s = 0$$

The equation (2.9) will be commonly called the Riccati equation. Moreover, if μ is a solution of the Riccati equation (2.9), then $\tilde{\mu}$ in (2.5) is a solution of $\tilde{\mu}_w - \frac{1}{2}\tilde{\mu}^2 - \tilde{s} = 0$, where \tilde{s} is given in (1.6).

Proof. The proof of existence can be found in [11, Lemma 3], and using the transformation rule of the Schwarzian derivative for \tilde{s} in (1.6) and $\tilde{\mu}$ in (2.5) implies that $\tilde{\mu}$ is again a solution of the Riccati equation. \square

Remark 2.4. Any solution μ of the Riccati equation (2.9) is only locally defined in general. But see also Remark 2.1.

A straightforward computation shows that for $Z = [Y \land \widehat{Y}]$

(2.10)
$$\langle dZ, dZ \rangle = \theta dz^2 + \frac{1}{2}(\rho + \bar{\rho})(dzd\bar{z} + d\bar{z}dz) + \bar{\theta}d\bar{z}^2$$

holds, where θ , ρ is given in (2.6) and (2.7). Combining Lemma 2.3, we obtain:

Lemma 2.5. Let $Z = [Y \land \widehat{Y}]$ be an \mathbb{L} -lift of a Möbius surface y. Moreover, let μ be the complex function defined in (2.4). Then the following statements are equivalent:

- (1) Z is conformal.
- (2) μ is a solution of the Riccati equation (2.9).

(3) Y and \widehat{Y} satisfies $\widehat{Y}_z \in \operatorname{span}_{\mathbb{C}}\{\widehat{Y}, Y, Y_z\}$ modulo $V_{\mathbb{C}}^{\perp}$, where $V_{\mathbb{C}}^{\perp}$ denotes the complement of the complexification of the 4-dimensional Lorentzian subspace V.

Note that from (2.10), if Z is conformal, the conformal factor is $2 \operatorname{Re} \rho$.

2.3. The moving frame and the structure equations. From now on we only consider a solution μ of the Riccati equation for a Mobius surface in \mathbb{S}^n , that is, we choose a section $\widehat{Y} \in \Gamma(V)$ such that $\langle \widehat{Y}, \widehat{Y} \rangle = 0$ and $\langle \widehat{Y}, Y \rangle = -1$ hold, and $Z = [Y \wedge \widehat{Y}]$ is conformal. Then \widehat{Y} is well-defined, $\mu dz = 2\langle \widehat{Y}, Y_z \rangle dz$ becomes a complex connection 1-form on a surface yand \widehat{Y} defines another surface $\widehat{y} = [\widehat{Y}]$. Note that the surface \widehat{y} can be degenerate in general, and moreover \widehat{y} could coincide with y at some points.

Let $\{\psi_1, \ldots, \psi_{n-2}\}$ be an orthonormal frame of the normal bundle V^{\perp} and define complex functions γ_j, k_j and $d_{ij} (1 \le i, j \le n-2)$ by

(2.11)
$$L := \sum_{j=1}^{n-2} \frac{1}{2} \gamma_j \psi_j, \quad \kappa := \sum_{j=1}^{n-2} \frac{1}{2} k_j \psi_j, \quad D_z \psi_i := \sum_{j=1}^{n-2} \frac{1}{2} d_{ji} \psi_j.$$

Note that $L = 2D_{\bar{z}}\kappa + \bar{\mu}\kappa$, and since $\{\psi_1, \ldots, \psi_{n-2}\}$ is an orthonormal basis, $d_{ij} + d_{ji} = 0$ $(i, j = 1, \ldots, n-2)$ hold.² Then the fundamental system of a Möbius surface in \mathbb{S}^n can be summarized as follows:

Proposition 2.6. Let Y be the canonical lift of a Möbius surface y and $Z = [Y \land \widehat{Y}]$ a conformal \mathbb{L} -lift. Moreover, define four sections of V by

(2.12)
$$X^{\pm} = \frac{1}{\sqrt{2}} (\pm Y + \hat{Y}), \quad P^{+} = \operatorname{Re}(2Y_{\bar{z}} + \bar{\mu}Y), \quad P^{-} = \operatorname{Im}(2Y_{\bar{z}} + \bar{\mu}Y),$$

and denote by $\{\psi_1, \ldots, \psi_{n-2}\}$ the orthonormal frame of the normal bundle V^{\perp} in (2.11). Set

(2.13)
$$F = (X^+, X^-, P^+, P^-, \psi_1, \dots, \psi_{n-2}) : \mathbb{D} \to \mathrm{SO}^+(1, n+1).$$

and take $\alpha = F^{-1}dF$ the Maurer-Cartan form of F and decompose it by (1,0)- and (0,1)-parts as $\alpha = \alpha' + \alpha''$ ($\alpha'' = \overline{\alpha'}$) and set

(2.14)
$$\alpha' = \begin{pmatrix} A & B^{\sharp} \\ B & C \end{pmatrix} dz, \quad B^{\sharp} = \operatorname{diag}(1, -1)B^{T},$$

where A, B and C respectively take values in $\mathfrak{so}(1,1,\mathbb{C})$, $M(n,2,\mathbb{C})$ and $\mathfrak{so}(n,\mathbb{C})$. Then the matrix valued functions A, B and C can be explicitly computed by the complex functions ρ ,

²The section L is equal to 2ζ of [4, p. 419]. We have chosen factors, respectively, 1/2 in (2.11) comparing to [4, Proposition 7.4], since the Maurer-Cartan form in the following can be written in a simple manner.

$$\mu, \gamma_j, k_j (j = 1, \dots, n-2) \text{ and } d_{ij} (i, j = 1, \dots, n-2) \text{ in } (2.7), (2.9), (2.11):$$

$$(2.15) \quad A = \frac{1}{2} \begin{pmatrix} 0 & \mu \\ \mu & 0 \end{pmatrix},$$

$$(2.16) \quad B = \frac{1}{2\sqrt{2}} \begin{pmatrix} \rho^+ & \rho^- \\ \gamma & \gamma \end{pmatrix}, \quad \rho^{\pm} = (\rho \pm 1) \begin{pmatrix} 1 \\ -\sqrt{-1} \end{pmatrix}, \quad \gamma = \begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_{n-2} \end{pmatrix},$$

$$(2.17) \quad C = \frac{1}{2} \begin{pmatrix} 0 & \sqrt{-1} \mu & -k^T \\ -\sqrt{-1} \mu & 0 \\ k & \sqrt{-1} k & d \end{pmatrix}, k = \begin{pmatrix} k_1 \\ \vdots \\ k_{n-2} \end{pmatrix}, d = (d_{ij}) \in \mathfrak{so}(n-2, \mathbb{C}).$$

Moreover, the compatibility condition $F_{z\bar{z}} = F_{\bar{z}z}$ is equivalent to the conformal Gauss, Codazzi and Ricci equations for a Möbius surface $y : \mathbb{D} \to \mathbb{S}^n$:

- (Gauss) $\frac{1}{2}s_{\bar{z}} = 3\langle \kappa, D_z\overline{\kappa} \rangle + \langle D_z\kappa, \overline{\kappa} \rangle,$
- (Codazzi) $\operatorname{Im}(D_{\bar{z}}D_{\bar{z}}\kappa + \frac{1}{2}\bar{s}\kappa) = 0,$

(Ricci)
$$D_{\bar{z}}D_{z}\psi_{j} - D_{z}D_{\bar{z}}\psi_{j} = 2\langle\psi_{j},\kappa\rangle\overline{\kappa} - 2\langle\psi_{j},\overline{\kappa}\rangle\kappa, \quad (j = 1,\ldots,n-2).$$

Note that $R_{\bar{z}z}^D \psi_j = D_{\bar{z}} D_z \psi_j - D_z D_{\bar{z}} \psi_j$ holds and s and κ are the Schwarzian derivative and the Hopf differential, respectively. The orthonormal frame F in (2.13) will be called the adapted frame.

Remark 2.7. Note that the sections P^{\pm} in (2.12) can also be represented as

(2.18)
$$P^+ = \operatorname{Re}(2Y_z + \mu Y), \quad P^- = -\operatorname{Im}(2Y_z + \mu Y).$$

Proof. For the canonical lift Y, the unique section N in (2.2) and any section $L \in \Gamma(V_{\mathbb{C}}^{\perp})$, the following equations hold:

(2.19)
$$\begin{cases} Y_{zz} = -\frac{s}{2}Y + \kappa, \\ Y_{z\overline{z}} = -\langle \kappa, \overline{\kappa} \rangle Y + \frac{1}{2}N, \\ N_z = -2\langle \kappa, \overline{\kappa} \rangle Y_z - sY_{\overline{z}} + 2D_{\overline{z}}\kappa, \\ L_z = D_z L + 2\langle L, D_{\overline{z}}\kappa \rangle Y - 2\langle L, \kappa \rangle Y_{\overline{z}}, \end{cases}$$

where s and κ are respectively the Schwarzian derivative and the Hopf differential. Note that the first equation is nothing but the inhomogeneous Hill's equation in (1.3), and the other three equations follow from a straightforward computation. Consider the new framing $\{Y, \hat{Y}, P^{\pm}\}$ of V, by (2.12). We reformulate (2.19) as (Here we set $P := P^+ - \sqrt{-1}P^-$)

$$\begin{cases} Y_z = -\frac{\mu}{2}Y + \frac{1}{2}P, \\ \widehat{Y}_z = \frac{\mu}{2}\widehat{Y} + \frac{\rho}{2}P + L, \\ P_z = \frac{\mu}{2}P + 2\kappa, \\ P_{\overline{z}} = -\frac{\overline{\mu}}{2}P + \overline{\rho}Y + \widehat{Y}, \\ L_z = D_zL + \langle L, L \rangle Y - \langle L, \kappa \rangle \overline{P} \\ 8 \end{cases}$$

So we obtain

$$\begin{cases} X_z^{\pm} &= \frac{\mu}{2} X^{\mp} + \frac{\pm 1+\rho}{2\sqrt{2}} (P^+ - \sqrt{-1}P^-) + \frac{1}{\sqrt{2}} L, \\ P_z^+ &= \frac{-\sqrt{-1}\mu}{2} P^- + \frac{1+\rho}{2\sqrt{2}} X^+ - \frac{-1+\rho}{2\sqrt{2}} X^- + \kappa, \\ P_z^- &= \frac{\sqrt{-1}\mu}{2} P^+ - \frac{\sqrt{-1}(1+\rho)}{2\sqrt{2}} X^+ + \frac{\sqrt{-1}(-1+\rho)}{2\sqrt{2}} X^- + \sqrt{-1}\kappa, \\ L_z &= D_z L + \langle L, L \rangle Y - \langle L, \kappa \rangle (P^+ + \sqrt{-1}P^-). \end{cases}$$

Then the equations (2.15), (2.16) and (2.17) follow from this, i.e., the Gauss equation, Codazzi equation and Ricci equations have been derived in [6].

Remark 2.8. The adapted frame F depends on the choice of a solution μ of the Riccati equation, and it is locally defined. The compatibility $F_{z\bar{z}} = F_{\bar{z}z}$ is equivalent to $d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0$, where $\alpha = F^{-1}dF$ is the left Maurer-Cartan form of F. Moreover, it can be interpreted as a flatness of the connection $d + \alpha$.

Corollary 2.9. The identity $B^T B = \frac{1}{2} \langle L, L \rangle \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ holds, where L is the section in (2.8). Moreover, by setting $B = (b_1 b_2)$, $b_1 - b_2 \neq 0$ holds.

3. WILLMORE SURFACES AND CONFORMAL HARMONIC L-LIFTS

3.1. Willmore surfaces, \mathbb{L} -lifts and Ruh-Vilms type theorem. As we have seen that the connection 1-form $d + \alpha$ with $\alpha = F^{-1}dF$ is a fundamental object for a Möbius surface y, and we now introduce a family of connections 1-forms $d + \alpha^{\lambda}$ parameterized by $\lambda \in S^1$ as follows:

(3.1)
$$\alpha^{\lambda} = \lambda^{-1} \alpha'_{\mathfrak{p}} + \alpha_{\mathfrak{k}} + \lambda \alpha''_{\mathfrak{p}},$$

where $\mathfrak{g} = \operatorname{Lie}(\operatorname{SO}^+(1, n+1))$ will be decomposed as $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ with $\mathfrak{k} = \operatorname{Lie}(\operatorname{SO}^+(1, 1) \times \operatorname{SO}(n))$, and $\alpha_{\mathfrak{k}}$ and $\alpha_{\mathfrak{p}}$ are the \mathfrak{k} - and the \mathfrak{p} -valued 1-forms. Moreover \prime and $\prime\prime$ denote the (1, 0)- and the (0, 1)-parts, respectively. Note that $\alpha^{\lambda}|_{\lambda=1} = \alpha$, and $d + \alpha^{\lambda}$ is flat for $\lambda = 1$. Then the flatness of a whole family $d + \alpha^{\lambda}$ parameterized by $\lambda \in S^1$ gives an additional constraint for a Möbius surface y. We now characterize Willmore surfaces in terms of a conformal \mathbb{L} -lift and the family of connections $d + \alpha^{\lambda}$.

Theorem 3.1 (Ruh-Vilms type theorem). Let Y be the canonical lift of a conformal immersion $y : \mathbb{D} \to \mathbb{S}^n$ and $Z = [Y \land \widehat{Y}]$ a conformal \mathbb{L} -lift. Then the following statements are equivalent:

- (1) The surface is a Willmore surface.
- (2) The section L in (2.8) satisfies $2D_{\bar{z}}L \bar{\mu}L = 0$.
- (3) The local \mathbb{L} -lift Z is conformal harmonic.
- (4) The family of connections $d + \alpha^{\lambda}$ in (3.1) is flat for all $\lambda \in S^1$.

Proof. The equivalence of (3) and (4) is a standard formulation of a harmonic map from a Riemann surface into a symmetric space, since the decomposition of $\mathfrak{g} = \text{Lie}(\text{SO}^+(1, n+1))$, by $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ with $\mathfrak{k} = \text{Lie}(\text{SO}^+(1, 1) \times \text{SO}(n))$, is associated to the symmetric space $\text{Gr}_{1,1}$, see for example [7].

(1) \Leftrightarrow (2): By definition and the Riccati equation of μ , we have

$$2D_{\bar{z}}L - \bar{\mu}L = 2(2D_{\bar{z}}D_{\bar{z}}\kappa + \bar{\mu}_{\bar{z}}\kappa + \bar{\mu}D_{\bar{z}}\kappa) - \bar{\mu}(2D_{\bar{z}}\kappa + \bar{\mu}\kappa) = 2(2D_{\bar{z}}D_{\bar{z}}\kappa + \bar{s}\kappa)$$

By the Willmore equation, we obtain the conclusion.

(2) \Leftrightarrow (4): We compute the flatness conditions $d\alpha^{\lambda} + \frac{1}{2}[\alpha^{\lambda} \wedge \alpha^{\lambda}] = 0$. It is equivalent to the Gauss, Codazzi and Ricci equations in Proposition 2.6, together with the extra condition $B_{\bar{z}} + \bar{C}B - B\bar{A} = 0$, which is equivalent to

$$\rho_{\bar{z}} - \bar{\mu}\rho - 2\langle L, \bar{\kappa} \rangle = 0, \quad \gamma_{i\bar{z}} + \frac{1}{2} \sum_{j} \gamma_{j} \overline{d_{ij}} - \frac{\bar{\mu}}{2} \gamma_{i} = 0, \quad (1 \le i \le n-2).$$

The second equation is equivalent to $2D_{\bar{z}}L - \bar{\mu}L = 0$. Since $\rho_{\bar{z}} - \bar{\mu}\rho - 2\langle L, \bar{\kappa} \rangle = 0$ holds for all Willmore surfaces ((4.10) of [14]), we complete the proof.

From (4) in Theorem 3.1, there exists a family of frames F^{λ} from \mathbb{D} to the following twisted loop group

(3.2)
$$\Lambda \mathrm{SO}^+(1, n+1)_{\sigma} = \left\{ g : S^1 \to \mathrm{SO}^+(1, n+1) \mid g(-\lambda) = \sigma g(\lambda) \right\},$$

where σ is the involution of the symmetric space $Gr_{1,1}$ given by

(3.3)
$$\sigma g = \operatorname{Ad}\operatorname{diag}(-1, -1, 1, \dots, 1)g, \quad g \in \operatorname{SO}^+(1, n+1).$$

Definition 3. The family of moving frames $F^{\lambda} : \mathbb{D} \to \Lambda \mathrm{SO}^+(1, n+1)_{\sigma}$ given by $\alpha^{\lambda} = (F^{\lambda})^{-1} dF^{\lambda}$ of (4) in Theorem 3.1 will be called the *extended frame* of a Willmore surface y.

Note that the extended frame F^{λ} evaluated $\lambda = 1$ is the adapted frame of y up to a left multiplication of a constant factor in SO⁺(1, n + 1).

3.2. Adjoint transformation, S-Willmore surfaces and L-lifts. Let us recall the Willmore condition $D_{\bar{z}}D_{\bar{z}}\kappa + \frac{1}{2}\bar{s}\kappa = 0$. By the Codazzi equation, the Willmore condition is also equivalent to $\operatorname{Re}(D_{\bar{z}}D_{\bar{z}}\kappa + \frac{1}{2}\bar{s}\kappa) = 0$. Then we define the special class of Willmore surfaces, [10, Theorem 1.1].

Definition 4. A Willmore surface $y : M \to \mathbb{S}^n$ is called an *S*-Willmore surface if $D_{\bar{z}}\kappa \parallel \kappa$ holds, that is, there exists some function r such that $D_{\bar{z}}\kappa + r\kappa = 0$ holds.

As seen in Proposition 1.4, the Willmore condition for a Möbius surface with a Riccati equation $\mu_z - \frac{1}{2}\mu^2 - s = 0$ is characterized as

(3.4)
$$D_{\bar{z}}L - \frac{\bar{\mu}}{2}L = 0,$$

where $L = 2D_{\bar{z}}\kappa + \bar{\mu}\kappa$. Then this condition means that the covariant derivative of L is parallel to itself $D_{\bar{z}}L \parallel L$, and thus we could consider the following sufficient conditions: $\langle L, L \rangle = 0$ or L = 0. This observation gives the following results.

Lemma 3.2. Let y be a Möbius surface in \mathbb{S}^n with a conformal \mathbb{L} -lift $Z = [Y \land \widehat{Y}]$. Then the following statements hold:

(1) The surface y is S-Willmore with a dual surface $\hat{y} = [\hat{Y}]$ if and only if L = 0.

(2) If y is a Willmore surface and $\langle L,L\rangle = 0$ is satisfied, then the surface $\hat{y} = [\hat{Y}]$ with \hat{Y} in (2.4) is a Willmore surface, and it will be called the adjoint transformation of y.

Proof. The statement (1) is a straightforward computation by using Definition 4. The statement (2) has been proven in [14, Section 4.2]. \Box

Remark 3.3.

- (1) The adjoint transformation of a Willmore surface was defined in [14, Definition 4.2]. From Theorem 4.8 in [14], there always exist adjoint transformations of a Willmore surface y and then $\hat{y} = [\hat{Y}]$ is also a Willmore surface if it is not degenerate, and \hat{y} will be called an *adjoint surface*. Moreover adjoint transformations of a Willmore surface y are not unique, since solutions of $\theta = 0$ and $\langle L, L \rangle = 0$ are not unique, in general.
- (2) When n = 3, any Willmore surface is an S-Willmore surface, since L is a complex function and $\langle L, L \rangle = 0$ is equivalent to L = 0 and the existence of adjoint transformations as given in the above (2). For n > 4, there exist Willmore surfaces which are not S-Willmore, [9].

We now consider singularities of the \mathbb{L} -lift $Z = [Y \land \widehat{Y}]$ of a Willmore surface Y and its adjoint transformation \widehat{Y} . It may occur on umbilic points.

Proposition 3.4. [13, 14] Let y be a Willmore surface on M and $M_0 \subset M$ be the umbilic free subset of y on M. Then for any $z_0 \in M_0$, there exists an open set $U \subset M_0$ containing z_0 such that the canonical lifts of y and the adjoint transformation \hat{y} , which will be denoted by Y and \hat{Y} , define the conformal harmonic \mathbb{L} -lift $Z = [Y \land \hat{Y}]$ on U.

On the other hand, for an umbilic point $z_1 \in M$ and an open set $U_1 \subset M$ containing z_1 , the \mathbb{L} -lift $Z = [Y \land \hat{Y}]$ may not be well-defined on U_1 in some choice of μ . Therefore, we need to think about the original Willmore conditions (3.4) in general to have a well-defined conformal \mathbb{L} -lift.

Recall that the $n \times 2$ matrix B in the Maurer-Cartan form is given in (2.16) and its identity in Corollary 2.9. From Lemma 3.2 we obtain:

Proposition 3.5. (1) Let y be a Willmore surface and \hat{y} be an adjoint transformation of y such that $Z = [Y \land \hat{Y}]$ is well-defined. Then the $n \times 2$ matrix B in the Maurer-Cartan form of the adapted frame in (2.16) satisfies

$$B^T B = 0.$$

In this case, the conformal harmonic \mathbb{L} -lift Z will be called isotropic.

(2) Let y be an S-Willmore surface and \hat{y} a dual surface such that $Z = [Y \land \hat{Y}]$ is welldefined. Then the $n \times 2$ matrix B in the Maurer-Cartan form of the adapted frame in (2.16) satisfies

$$B^T B = 0, \quad \operatorname{rank} B = 1.$$

In this case, the conformal harmonic \mathbb{L} -lift Z will be called the S-isotropic.

The results are summarized as the following table:

Surface classes	Conformal L-lifts	$n \times 2$ matrix B
Willmore on M	Conformal harmonic	$B^T B = \frac{1}{2} \langle L, L \rangle \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$
Willmore with adjoint transformation on M_0	Isotropic conformal harmonic	$B^T B = O_2$
S-Willmore on M_0	S-isotropic conformal harmonic	$B^T B = O_2$ and rank $B = 1$

Since μ is not a unique solution of the Riccati equation, thus the conformal harmonic \mathbb{L} -lift Z of a surface (S-) Willmore is not necessarily (S-) isotropic, that is, μ may not be a solution of $\langle L, L \rangle = 0$ (or L = 0) but a solution of $2D_{\bar{z}}L - \bar{\mu}L = 0$. In fact, a conformal harmonic (S-) isotropic \mathbb{L} -lift $Z = [Y \wedge \hat{Y}]$ may not be well-defined around an umbilic point by Proposition 3.4.

Open questions: We would like to ask some open problems that originally come from Hélein's work [11, 12].

- (1) For a closed, oriented, Willmore surface y in S^n , when will it admit a global isotropic conformal harmonic \mathbb{L} -lift?
- (2) We might split this problem into two sub-questions:
 - (a) If y is S-Willmore, does it always admit a global isotropic conformal harmonic \mathbb{L} -lift?
 - (b) If y is not S-Willmore, does it always admit a global isotropic conformal harmonic \mathbb{L} -lift?

References

- M Babich and A Bobenko. Willmore tori with umbilic lines and minimal surfaces in hyperbolic space. Duke Math. J., 72:151–185, 1993.
- [2] W Blaschke. Vorlesungen über Differentialgeometrie, III. Springer, Berlin. Springer, 1929.
- [3] D Brander, S-P Kobayashi, and P Wang. The björling problem for minimal surfaces in \mathbb{H}^n and \mathbb{S}^n . In preparation, 2024.
- [4] D Brander and P Wang. On the Björling problem for Willmore surfaces. J. Diff. Geom., 108:411–457, 2018.
- [5] R Bryant. A duality theorem for Willmore surfaces. J. Diff. Geom, 20:23–53, 1984.
- [6] F Burstall, F Pedit, and U Pinkall. Schwarzian derivatives and flows of surfaces. Contemporary Mathematics, 308:39–61, 2002.
- [7] J Dorfmeister, F Pedit, and H Wu. Weierstrass type representation of harmonic maps into symmetric spaces. Comm. Anal. Geom., 6:633–668, 1998.
- [8] J Dorfmeister and P Wang. Willmore surfaces in spheres via loop groups I:generic cases and some examples. arXiv:1301.2756v4, 2013.
- [9] N Ejiri. A counter example for weiner's open question. Indiana Univ. Math J., 31:209–211, 1982.
- [10] N Ejiri. Willmore surfaces with a duality in $S^n(1)$. Proc. London Math. Soc. (3), 57:383–416, 1988.
- [11] F Hélein. Willmore immersions and loop groups. J. Diff. Geom., 50:331-385, 1998.
- [12] F Hélein. A Weierstrass representation for Willmore surfaces. In Harmonic morphisms, harmonic maps, and related topics (Brest 1997), Chapman & Hall/CRC Res. Notes Math., pages 287–302, 2000.
- [13] X Ma. Willmore surfaces in S^n : transforms and vanishing theorems. PhD thesis, Technischen Universität Berlin, 2005.

- [14] X Ma. Adjoint transforms of Willmore surfaces in S^n . Manuscripta Math., 120:163–179, 2006.
- [15] QL Xia and YB Shen. Weierstrass type representation of Willmore surfaces in S^n . Acta Math. Sinica, 20:1029–1046, 2004.

Department of Applied Mathematics and Computer Science Technical University of Den-Mark 2800 Lyngby (Copenhagen) Denmark

Email address: dbra@dtu.dk

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810, JAPAN

Email address: shimpei@math.sci.hokudai.ac.jp

SCHOOL OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF ANALYTICAL MATHEMATICS AND APPLICATIONS (MINISTRY OF EDUCATION), FUJIAN KEY LABORATORY OF ANALYTICAL MATHEMATICS AND APPLICATIONS (FJKLAMA), FUJIAN NORMAL UNIVERSITY, FUZHOU 350117, P. R. CHINA

Email address: netwangpeng@163.com, pengwang@fjnu.edu.cn