Treatment Evaluation at the Intensive and Extensive Margins[∗]

Phillip Heiler† Asbjørn Kaufmann‡ Bezirgen Veliyev§

This paper provides a solution to the evaluation of treatment effects in selective samples when neither instruments nor parametric assumptions are available. We provide sharp bounds for average treatment effects under a conditional monotonicity assumption for all principal strata, i.e. units characterizing the complete intensive and extensive margins. Most importantly, we allow for a large share of units whose selection is indifferent to treatment, e.g. due to non-compliance. The existence of such a population is crucially tied to the regularity of sharp population bounds and thus conventional asymptotic inference for methods such as Lee bounds can be misleading. It can be solved using smoothed outer identification regions for inference. We provide semiparametrically efficient debiased machine learning estimators for both regular and smooth bounds that can accommodate high-dimensional covariates and flexible functional forms. Our study of active labor market policy reveals the empirical prevalence of the aforementioned indifference population and supports results from previous impact analysis under much weaker assumptions.

Keywords: Double/debiased machine learning; Lee bounds; Partial identification; Prin-

cipal strata; Sample selection

JEL classification: C13, C14, C21

[∗]We would like to thank Isaiah Andrews, Andres Aradillas-Lopez, Kevin Chen, Juan Carlos Escanciano, Patrik Guggenberger, Mark Henry, Keisuke Hirano, Edward Kennedy, Giovanni Mellace, Peter Schochet, Vira Semenova, Neil Shephard, Mikkel Sølvsten, Elie Tamer, Davide Viviano, Chris Walker, as well as the participants of the Aarhus, Harvard, and PennState Econometrics Seminars for their comments and suggestions that helped to improve the work. All remaining errors are ours.

[†]Aarhus University. Department of Economics and Business Economics, TrygFonden's Centre for Child Research, Fuglesangs Allé 4, 8210 Aarhus V, Denmark, email: [pheiler@econ.au.dk.](mailto:pheiler@econ.au.dk) Part of this research was conducted during main affiliation with Harvard University, Department of Economics. 02138 Cambridge MA, United States.

[‡]Aarhus University. Department of Economics and Business Economics, Fuglesangs Allé 4, 8210 Aarhus V, Denmark.

[§]Aarhus University. Department of Economics and Business Economics, Fuglesangs Allé 4, 8210 Aarhus V, Denmark.

1. Introduction

This paper deals with the evaluation of causal effects of a binary treatment when the outcome is only selectively observed and no instruments are available. Such missing outcome data or sample selection is ubiquitous and a threat to internal validity [\(Heckman,](#page-44-0) [1974,](#page-44-0) [1979\)](#page-44-1). Typical examples include outcome data such as wages that are only observed if units are employed [\(Lee, 2009\)](#page-46-0), missing survey responses [\(Bernhardt et al., 2024\)](#page-44-2), or attrition [\(Zhang and Rubin, 2003\)](#page-47-0). In the context of impact evaluation, a treatment can often change the sample selection status for some units. For example, if the treatment is an active labor market policy such as job training, it likely affects the probability of employment for some participants, e.g. due to accumulation of human capital (positive) or opportunity costs to job search (negative). For some units, however, the policy might be irrelevant with regards to employment or sample selection, but still relevant in terms of their earnings or other outcomes. Hence, there are potentially heterogeneous units at the intensive and extensive margins. These units can be classified within principal strata defined by their potential selection status.

We consider set identification, estimation, and inference for all principal strata, encompassing the complete intensive and extensive margins, under a weak conditional monotonicity assumption. Weak monotonicity allows for the simultaneous presence of always-takers, compliers, defiers, and never-takers.[1](#page-1-0)

The main challenge is the presence of units whose selection probabilities are unaffected by the treatment. The existence of such units is most apparent when the treatment is subject to additional non-compliance after assignment and intent-to-treat effects are analyzed. In this case, units who have information or preferences that lead them to not participate after treatment assignment will likely also not change their selection behavior, e.g. selection into employment, as a consequence. Further, there are policies that do not boost or hinder employability for some units, e.g. due to a lack of signal for their potential employers in the short run, but increase human capital boosting productivity and earnings. Similarly, in field experiments where responses cannot be enforced, an intervention might only affect response probabilities to follow-up surveys for some units.

From a discrete choice perspective, our setup admits units who are indifferent to the

¹In contrast to the instrumental variables literature, principal strata here refer to potential selection statuses when treatment is exogenously set to zero and one respectively. See Section [3.](#page-8-0)

treatment at any conceivable level of their private information. From a model selection perspective, unaffected units are equivalent to a sparsity constraint in the selection equation. Modern model selection such as *L*1-regularization, subset selection, deep neural networks, boosted trees, and random forests are able to recover such potentially sparse structures in the selection equation.

Figure [1.1](#page-2-0) contains an example from the job training program analyzed in this paper (Job Corps). It provides the histogram of the estimated relative selection probabilities with and without assignment to training using boosted trees. A value of 1*.*00 indicates that assignment does not predict employment status for a given unit. In this case we have that 2616 out of 9415 units stacking up at an exact. Inspect one. This pattern is striking and persistent across evaluation periods and estimation methods, see Sections [3](#page-8-0) and [8.](#page-34-0) In this study, there is significant non-compliance. Thus a large presence of estimated unaffected units seems plausible.

Figure 1.1: Job Corps Relative Sample Selection Probabilities

Histogram of the estimated relative conditional employment selection probabilities under treatment and control. Time period $t = 180$, method XGBoost, $n = 9415$. There are 2616 units at exact numerical 1.00 $(P(X^0) = 28.61\%)$. For more details consider Section [8.](#page-34-0)

In general, sample selection with unaffected units yield *mixture* distributions for relative conditional selection probabilities obtained in a first stage with point mass at 1*.*00. While without harm from an identification perspective, it directly translates into a fundamental problem for estimation and inference. In particular, sharp population bounds are nonsmooth functionals for which no uniformly unbiased estimator exists, i.e. the problem

is irregular by construction [\(Hirano and Porter, 2012\)](#page-45-0). More concretely, we show that not having a mixture is in fact necessary and sufficient for the population bound to be pathwise differentiable in the sense of [Bickel et al.](#page-44-3) [\(1993\)](#page-44-3). This means that standard errors and statistical inference obtained from methods such as Lee bounds with "naive" asymptotics [\(Lee, 2009\)](#page-46-0) are invalid in the presence of unaffected units.

We propose a solution to the problem involving repeated smoothing steps that effectively act as covers for the sharp identified sets. The resulting smooth outer identification regions can be efficiently estimated at the parametric rate and always yield valid confidence regions for the true parameter of interest as the direction of any bias is known. Thus, the degree of smoothing can be chosen to maximize power in finite samples. Under regularity, the smooth bounds can converge to the sharp identified set. On a high level, our results reveal that there is an identification-precision trade-off: Aiming for an outer identification region with high precision over a sharp region with low precision can significantly improve inference in finite samples.

For both the regular and the smoothed irregular case, we provide efficient influence functions and corresponding estimators using debiased machine learning. Under simple high level conditions, the estimators are asymptotically normal and reach their respective semiparametric efficiency bounds. The influence function for the intensive margin is equal to the one implied by the moment estimator in [Semenova](#page-46-1) [\(2024\)](#page-46-1) under *unknown* propensity scores and no moment selection. To our knowledge, this is the first paper to provide efficient influence functions that can be used for semiparametric estimation of compliers, defiers, and the combined extensive margin bounds in the sample selection model without exclusion. In contrast to any regular bounds, the smooth counterparts require weaker convergence assumptions for the involved nuisance parameters and, even in the irregular case, no margin condition that controls the distribution of selection effects.

We also explore the role of the propensity score for efficiency, extending the results in [Hahn](#page-44-4) [\(1998\)](#page-44-4) to sample selection without exclusion. The efficiency bound for the principal strata treatment effect bounds do not depend on the knowledge of the propensity score. We quantify the efficiency gap when using the moment functions under knowledge of the propensity score as suggested by [Semenova](#page-46-1) [\(2024\)](#page-46-1) and [Heiler](#page-45-1) [\(2024\)](#page-45-1). These estimators are generally inefficient except in knife-edge cases. However, there is a trade-off from a practical perspective: Analogously to inverse probability weighting (IPW) estimation of average treatment effects, these estimators do not require conditional (truncated) outcome means as additional nuisance inputs. Thus, depending on the difficulty in estimating the latter, using these simpler moment functions might be preferable if propensity scores are known. All aforementioned properties regarding (ir)regularity and smoothing also directly apply to these inefficient moment functions and their smoothed counterparts.

Monte Carlo simulations suggest that, in irregular designs, inference using the smooth methods compares favorably to the alternatives from the literature that either ignore the irregularity of the problem or rely on trimming [\(Heiler, 2024\)](#page-45-1), or moment selection/switching methods [\(Andrews and Soares, 2010;](#page-43-0) [Semenova, 2024\)](#page-46-1).

We extend the theory to other quantile-trimmed bounds obtained under additional stochastic dominance assumptions [\(Zhang and Rubin, 2003;](#page-47-0) [Huber and Mellace, 2015\)](#page-45-2). The components of our influence functions can also be used to construct heterogeneous treatment effect bounds in the sense of [Heiler](#page-45-1) [\(2024\)](#page-45-1) for any principal strata.

The empirical study is a comprehensive re-evaluation of the National Job Corps Study [\(Burghardt et al., 1999;](#page-44-5) [Schochet et al., 2008\)](#page-46-2). We demonstrate the empirical relevance of flexible estimation via machine learning methods, in particular with regards to sparsity in the selection equation. Aggregate results of the new semiparametrically efficient smoothing bounds with optimized learners rule out moderate to large negative impact of Job Corps on hourly earnings at the end of the evaluation period. The results over various time periods are surprisingly close to the more restrictive impact analysis by [Lee](#page-46-0) [\(2009\)](#page-46-0) that relies on a stronger monotonicity assumption rejected by the data [\(Semenova,](#page-46-1) [2024\)](#page-46-1).

The paper is organized as follows. Section [2](#page-5-0) discusses the relevant literature. Section [3](#page-8-0) introduces the nonparametric sample selection model and provides an in-depth discussion of conditional monotonicity. Section [4](#page-14-0) presents the sharp and smooth outer identification regions for all principal strata. Section [5](#page-21-0) provides the results on pathwise differentiability. Section [6](#page-24-0) contains the assumptions for debiased machine learning estimation and asymptotic inference as well as the efficiency gap under known propensity scores. Section [7](#page-32-0) extends the methodology to stochastic dominance bounds and heterogeneous effect bounds. Section [8](#page-34-0) presents the empirical study. Section [9](#page-43-1) concludes.

2. Literature

There is a large literature on sample selection models without exclusion under restrictive parametric assumptions [\(Heckman, 1979;](#page-44-1) [Staub, 2014\)](#page-47-1). Bounds for causal effects under varying weaker assumptions also have a rich tradition, see [Molinari](#page-46-3) [\(2020\)](#page-46-3) for a comprehensive survey. We focus on research that does not rely on the use of additional exclusion restrictions or instruments and is connected to the monotonicity assumption. We follow a principal stratification approach that considers identification of causal effects for all latent subgroups characterized by their potential selection behavior as a function of a binary treatment [\(Frangakis and Rubin, 2002\)](#page-44-6). This circumvents comparing systematically different groups when conditioning on realized selection behavior.

Bounds on causal effects for the intensive margin or always-takers under monotonicity have first been introduced by [Zhang and Rubin](#page-47-0) [\(2003\)](#page-47-0), see also [Lee](#page-46-0) [\(2009\)](#page-46-0) for an extension to a form of conditional monotonicity that excludes units whose selection behavior is unaffected by the treatment. Such "Lee" or "Zhang-Rubin-Lee" bounds [\(Andersen](#page-43-2) [et al., 2023\)](#page-43-2) are defined by conditional expectations trimmed at conditional quantiles. [Imai](#page-45-3) [\(2008\)](#page-45-3) shows the sharpness of such trimming bounds. [Semenova](#page-46-1) [\(2024\)](#page-46-1) introduces orthogonal moments for estimation of intensive margin bounds that can be used in highdimensional setups, see also [Olma](#page-46-4) [\(2021\)](#page-46-4) for nonparametric estimation of truncated conditional expectations. [Huber and Mellace](#page-45-2) [\(2015\)](#page-45-2) derive bounds for compliers under strong monotonicity. We extend their ideas allowing for compliers and defiers simultaneously. [Honoré and Hu](#page-45-4) [\(2020,](#page-45-4) [2022\)](#page-45-5) also discuss bounds in sample selection models without exclusion restrictions under additional parametric assumptions. Our bounds do not require any parametric structure. [Bartalotti et al.](#page-44-7) [\(2023\)](#page-44-7) provide bounds using monotonicity and stochastic dominance within a marginal treatment effect (MTE) framework. They focus on strong monotonicity only and do not provide estimation or inference theory. [Heiler](#page-45-1) [\(2024\)](#page-45-1) considers heterogeneous intensive margin bounds and misspecification robust inference under a strong margin assumption effectively ruling out subgroups for which selection probabilities do not depend on the treatment. Our new moment functions can also be used in conjunction with the approach by [Heiler](#page-45-1) [\(2024\)](#page-45-1) to provide heterogeneous effect bounds for any margin. [Okamoto](#page-46-5) [\(2023\)](#page-46-5) discusses sensitivity analysis of intensive margin and MTE bounds under partial violation of the conventional monotonicity assumption. His stochastic monotonicity assumption is distinct from our

approach as it assumes strong monotonicity for a limited share of the population instead of a weaker form of monotonicity. The use of intensive margin bounds has also been advocated by [Chen and Roth](#page-44-8) [\(2024\)](#page-44-8) when encountering dependent variables with (many) zeroes. All these papers consider either strong monotonicity or the restrictive version of weak monotonicity [\(Lee, 2009\)](#page-46-0), with [Semenova](#page-46-1) [\(2024\)](#page-46-1) being a notable exception.

Monotonicity also plays a crucial role for instrumental variable (IV) based methods [\(Imbens and Angrist, 1994;](#page-45-6) [Angrist and Imbens, 1995;](#page-44-9) [Angrist et al., 2000;](#page-43-3) [Abadie, 2003;](#page-43-4) [Heckman and Vytlacil, 2007;](#page-45-7) [Słoczyński, 2020;](#page-47-2) [Heiler, 2022\)](#page-45-8). Similar to the sample selection setup, it is an assumption on latent subtypes in the population to react towards a change in the instrument weakly in a particular direction in terms of their treatment selection. It rationalizes the same choices as a structural model for treatment selection that is additive separable in observables and unobservables both in the population [\(Vyt](#page-47-3)[lacil, 2002\)](#page-47-3) as well as numerically if estimated nonparametrically [\(Kline and Walters,](#page-46-6) [2019\)](#page-46-6). This also applies to our setup. For nonparametric identification of local average treatment effects or other MTE-type parameters, the instrument has to be able to move certain units in terms of their treatment choice (first stage). Combining compliers and defiers for IV using a weak monotonicity assumption has been considered by [Kolesár](#page-46-7) [\(2013\)](#page-46-7) and [Słoczyński](#page-47-2) [\(2020\)](#page-47-2). The key distinction to the IV setup is that we are interested in the causal effect of the treatment accounting for endogenous selection. Thus, our treatment takes on the role of the instrument while the selection is fully endogenous as the treatment in the IV case. The weak monotonicity assumption posited in this paper allows for the presence of all latent subtypes in the population but limits their presence within ex-ante unknown partitions of the covariate space. This can also by motivated by a structural selection model with mixed indices within these partitions. The IV framework does not allow for nonparametric identification in any population without movers (no first stage). We explicitly want to allow for units whose selection is not affected by the treatment. This can then generates the mixture distribution for the "first stage" relative conditional selection probabilities with point mass at one.

Such mixtures imply that the (generalized) ZR-Lee bounds are no longer smooth in the underlying data distribution and thus there exist no estimator sequence that is locally asymptotically unbiased in the sense of [Hirano and Porter](#page-45-0) [\(2012\)](#page-45-0). We construct a particular cover for the identification region that circumvents the non-regularity of the underlying functional of interest. Our identified set has non-empty interior and a zero duality gap as discussed in [Kaido and Santos](#page-45-9) [\(2014\)](#page-45-9). Thus, its semiparametric efficiency bound can be characterized and \sqrt{n} -consistent regular estimators of the support function exist in a uniform sense. As a by-product of the strict convexity of the smooth identified set, confidence intervals for the effects (not bounds) can be made more precise using modified critical values [\(Imbens and Manski, 2004\)](#page-45-10). Under a strong margin assumption, our bounds converge converge to the standard sharp ZR-Lee bounds at a known rate.

Our estimands are ratios of a weighted conditional expectations normalized by their share. For the intensive margin, estimating the share is related to estimating an expected conditional outcome under unconfoundedness when setting the treatment status to the best conditional mean. For the latter in isolation, [Luedtke and van der Laan](#page-46-8) [\(2016\)](#page-46-8) show that for the estimand to be pathwise differentiable, it is necessary and sufficient that units which are indifferent between treatments have an *exceptional* law with zero conditional variance under both regimes. We extend their approach to a class of ratio estimands and show that, for the principal strata bounds, the point mass phenomenon is both necessary and sufficient for irregularity, i.e. there are no *exceptional* laws as the latter would imply a violation of the overlap assumption.

[Levis et al.](#page-46-9) [\(2023\)](#page-46-9) also suggest smoothing for obtaining regular estimators of [Balke](#page-44-10) [and Pearl](#page-44-10) [\(1997\)](#page-44-10) bounds in the presence of covariates. They propose confidence intervals with a worst-case bias correction arising from the particular approximation. Unlike their approach, our smoothing will always widen the identified set and thus one can navigate the identification-precision trade-off without the need for a worst-case bias component.

[Semenova](#page-46-1) [\(2024\)](#page-46-1) uses the same weak monotonicity assumption as this paper but focuses exclusively on always-takers and does not address regularity and efficiency. To address the problem of misclassifying units, she uses a weak margin assumption and a shrinkage/moment-switching approach. The latter selects between different moment functions based on the estimated selection probabilities with a vanishing sequence of shrinkage parameters that obey a rate condition. There is no guidance on how to choose this parameter in finite samples. Moreover, while asymptotically valid in handling potential misclassification, this moment selection effectively narrows the identified set leading to potential undercoverage in finite samples. Our approach guarantees at least nominal coverage and can be optimized with respect to power. For always-taker bounds, [Heiler](#page-45-1)

[\(2024\)](#page-45-1) suggests to round trimming threshold towards the closest value on a grid. While this approach similarly yields an outer identification region in the context of relative selection probabilities very close to one, it is unclear how to use in the case of a point mass for units whose selection is unaffected by the treatment, i.e. an exact one, as rounding could be done in two directions leading to different impact of misclassification errors.

In other contexts, [Lee and Weidner](#page-46-10) [\(2021\)](#page-46-10) and [Pakel and Weidner](#page-46-11) [\(2023\)](#page-46-11) also note that there can be a trade-off between identification and precision. They argue that confidence sets obtained from outer bounds can well be tighter compared to using sharp bounds. A similar intuition applies for our smooth outer identification region. For a data combination problem, [d'Haultfoeuille et al.](#page-44-11) [\(2024\)](#page-44-11) also consider outer bounds whose conservativeness arise from regularization.

We characterize the efficient influence functions and semiparametric efficiency bounds with and without knowledge of the propensity score. Our results generalize [Hahn](#page-44-4) [\(1998\)](#page-44-4) to sample selection under weak conditional monotonicity and either (i) regular sharp bounds or (ii) regular smooth bounds that cover the irregular sharp bounds. We also contribute and make use of the expanding literature on the use of debiased machine learning methods for estimation of causal effects in economics [\(Chernozhukov et al., 2018,](#page-44-12) [2022\)](#page-44-13) and its combination with partial identification [\(Heiler, 2024;](#page-45-1) [Semenova, 2023,](#page-46-12) [2024\)](#page-46-1). More specifically, we show that the particularities of the relative selection probabilities and the rate at which they can be learned from data crucially interact with the (ir)regularity of the parameters of interest.

3. Model and Monotonicity

Assume we observe iid data $W = (YS, S, D, X)'$ where $S \in \{0, 1\}$ is a selection variable indicating whether an outcome is observed or not. $Y \in \mathcal{Y}$ is the partially unobserved outcome of interest. $D \in \{0,1\}$ is a binary treatment of interest. $X \in \mathcal{X}$ is a vector of predetermined covariates. We are interested in the causal effect of the treatment on the outcome defined in terms of potential outcomes and potential selection indicators. The realized but partially unobserved outcome is defined as

$$
Y = DY(1) + (1 - D)Y(0).
$$
\n(3.1)

Selection is connected to potential selection indicators as

$$
S = DS(1) + (1 - D)S(0). \tag{3.2}
$$

Figure 3.1: Nonparametric Sample Selection Model

A graphical representation of a nonparametric sample selection model. Nodes denote variables and edges are structural relationships. A missing arrow from one node to another is an exclusion restriction. Unobserved independent components at each node are omitted. Unobserved variables and its edges are dashed. Gray nodes are partially unobserved.

Figure [3.1](#page-9-0) depicts a prototypical sample selection model with the relevant exclusion restrictions. Such a model implies the following conditional independence assumption for the treatment which we maintain throughout:

Assumption 3.1 (Conditional Independence)

$$
D \perp\!\!\!\perp Y(d), S(d) \mid X = x \text{ for all } x \in \mathcal{X} \text{ and } d = 0, 1,
$$

where \perp denotes statistical independence. Thus, treatment is assumed to be exogenous conditional on *X*. However, selection is left essentially unrestricted, i.e. it can be fully endogenous or exogenous. In particular, there are no exclusion restrictions available for *S*, i.e. we cannot use instruments or similar to account for endogenous selection.

The potential selection indicators define principal strata which represent how the treatment affects selection behavior. Table [3.1](#page-10-0) contains all strata using the nomenclature from the instrumental variables literature.

To proceed, we postulate the existence of an unknown but identified partitioning of the covariate space $\mathcal X$ on which a weak monotonicity assumption applies to these principal strata^{[2](#page-9-1)}.

²Note that this definition differs slightly from the conditional monotonicity assumption of [Semenova](#page-46-1) [\(2024\)](#page-46-1) which does not yield a unique partitioning.

	Lable 0.1. I Hillipal bulata		
Type		S(0)	S(1)
(AT)	Always-takers		
(C)	Compliers		
(D)	Defiers		
(NT)	Never-takers		

Table 3.1: Principal Strata

Assumption 3.2 (Weak/Conditional Monotonicity) *Assume there are subsets of the covariate space* $\tilde{\mathcal{X}}^+$ *,* $\tilde{\mathcal{X}}^- \subseteq \mathcal{X}$ *such that*

$$
S(1) \ge S(0) \text{ if } x \in \tilde{\mathcal{X}}^+,
$$

$$
S(1) \le S(0) \text{ if } x \in \tilde{\mathcal{X}}^-.
$$

Assumption [3.2](#page-9-2) yields a distinct partitioning of $\mathcal{X} = \mathcal{X}^+ \cup \mathcal{X}^- \cup \mathcal{X}^0$ where

$$
\mathcal{X}^0 = \tilde{\mathcal{X}}^+ \cap \tilde{\mathcal{X}}^-,
$$

$$
\mathcal{X}^+ = \tilde{\mathcal{X}}^+ \backslash \mathcal{X}^0,
$$

$$
\mathcal{X}^- = \tilde{\mathcal{X}}^- \backslash \mathcal{X}^0.
$$

The presence of a potentially non-empty \mathcal{X}^0 , i.e. $P(\mathcal{X}^0) > 0$, will be crucial in what follows. Substantially, weak monotonicity allows for the presence of all principal strata including defiers. However, it is a partial restriction of types within partitions. Table [3.2](#page-10-1) contains the mixture of types in the different partitions as well the population.

Table 3.2: Partitions, Observed Strata, and Admitted Latent Types

			$S \setminus D$	$\overline{0}$					
			$\boldsymbol{0}$	NT/C	NT/D				
			$\mathbf{1}$	AT/D	AT/C				
				(a) $x \in \mathcal{X}$					
$S \setminus D$	$\overline{0}$	$\overline{1}$	$S \setminus D$	$\overline{0}$	$\overline{1}$		$S \setminus D$	$\overline{0}$	-1
θ	NT/C	NT	$\boldsymbol{0}$	NT	NΤ		θ	NT	NT/D
	AT	AT/C	$\mathbf 1$	AT	AT		$\mathbf{1}$	AT/D	AT
(b) $x \in \mathcal{X}^+$			(c) $x \in \mathcal{X}^0$			(d) $x \in \mathcal{X}^-$			

For example, in partition \mathcal{X}^0 , the population consists only of always-takers and nevertakers while on \mathcal{X}^+ and \mathcal{X}^- only defiers or compliers are ruled out, respectively. This assumption seems most relevant if we have some discrete covariates and/or continuous variable that affect selection behavior discontinuously e.g. by threshold effects. Assumption [3.2](#page-9-2) is also consistent with a single-index structure in the selection equation that allows for sparsity with respect to the treatment on some parts of the covariate space.

Example (Single Index Model): Consider a simple semiparametric selection model with treatment and selection that are additive separable in observables and unobservables

$$
D = \mathbb{1}(f(X) + U_D \ge 0),
$$

\n
$$
Y = \mu(X, D) + U_Y,
$$

\n
$$
S = \mathbb{1}(g(D, X) + U_S \ge 0),
$$
\n(3.3)

where $U_D \perp\!\!\!\perp U_Y, U_S$ conditional on $X = x$ then implies conditional independence as in Assumption [3.1.](#page-9-3) Weak monotonicity as in Assumption [3.2](#page-9-2) here is implied by an unrestricted $g(D, X)$. This can generate a mixed index structure

$$
g(d,x) = \begin{cases} g_{+}(d,x) & \text{if } x \in \mathcal{X}^{+}, \\ g_{-}(d,x) & \text{if } x \in \mathcal{X}^{-}, \\ g_{0}(x) & \text{if } x \in \mathcal{X}^{0}. \end{cases}
$$
(3.4)

where $g_{+}(1, x) > g_{+}(0, x)$ and $g_{-}(0, x) > g_{-}(1, x)$. The fact that $g_{0}(x)$ does not depend on *d* can be seen as a sparsity assumption within a partition of the covariate space. For example, consider the case of a simple parametric index in the selection equation with $X = (X_1, X_2)'$ where X_1 is continuous and X_2 discrete with support points $\{-1, 0, 1\}$

$$
S = \mathbb{1}(\gamma_0 + \gamma_1 X_1 + \gamma_2 D \mathbb{1}(X_2 = -1) + \gamma_3 D \mathbb{1}(X_2 = 1) + U_S \ge 0).
$$
 (3.5)

Here X_2 can separate the monotonicity types. In particular, $\gamma_2 < 0 < \gamma_3$ or $\gamma_2 > 0 >$ *γ*₃ imply conditional monotonicity. If the signs of γ ₂ and γ ₃ are identical, then either $\mathcal{X}^+ = \{\emptyset\}$ or $\mathcal{X}^- = \{\emptyset\}$, implying the conventional *unconditional* or *strong* monotonicity $P(S(1) \geq S(0)) = 1.$

Whenever $P(X^0) > 0$, conditional monotonicity and independence generate a *mixture*

distribution for the potential or relative conditional selection probabilities

$$
p_0(x) = \frac{P(S=1|D=0, X=x)}{P(S=1|D=1, X=x)} = \frac{P(S(0)=1|X=x)}{P(S(1)=1|X=x)}.
$$
\n(3.6)

In particular, for \mathcal{X}^+ and \mathcal{X}^- we have that $p_0(x) < 1$ and $p_0(x) > 1$, respectively, while there is potentially a point mass for which $p_0(x) = 1$ corresponding to $P(\mathcal{X}^0) > 0$. Thus, conditional monotonicity allows for mixture distributions which are not necessarily smooth in $p_0(x)$. We seek to conduct inference on causal effects for all principal strata under such mixtures.

The figure contains histograms for the estimated $p_0(x)$ using five different estimation methods on the research sample at week $t = 208$. (a) Logit is a logistic regression with linear additive index. (b) Logit Interacted is a fully treatment interacted logistic model. (c) XGBoost are gradient boosted trees with cross-validated boosting steps using xgboost. (d) Neural Network is a batch-trained, fully connected artificial feed-forward neural network with three ReLu hidden layers, drop-out regularization, and a final sigmoid layer using keras. (e) Random Forest is an honest probability forest using grf with default parameters.

Modern model selection methods such as *L*1-regularization, subset selection, neural networks, boosting, and random forests are able to recover and produce potentially sparse structures in the selection equation. We present an example from relative employment probabilities and a job training treatment in Figure [3.2.](#page-12-0) It contains the predicted relative selection probabilities from Section [8](#page-34-0) using five different estimation methods: (a) logistic regression, (b) logistic regression with fully interacted treatment, (c) gradient boosted trees (XGBoost), (d) neural network, and (e) random forest, see Section [8](#page-34-0) and Appendix [B](#page-79-0) for more details.

The methods capable of sparsity yield a clear mixture pattern for the estimated distribution of $p_0(x)$. In particular, (c), (d), and (e) produce an exact share of numerical ones with up to 45.16% of the sample estimates. (a) imposes strong monotonicity and cannot produce exact ones in finite samples by construction. We suspect that (a) is most likely to be misspecified. (b) has relevant density around one, but also generates overly extreme values for the relative selection probabilities. The sparse methods agree on around 45% – 60% of the close or equal to one classifications. This qualitative pattern can be observed along a long sequence of post-treatment periods. Over time there is an average distribution shift suggesting more positive employment effects. The sparsity pattern, however, is pervasive. We take this as a signal that the mixture distributions are likely reflecting of or a good approximation to an underlying structure and are not only spurious by-products of particular estimation algorithms motivating Assumption [3.2](#page-9-2) with $P(X^0) > 0$.

Remark: On a cautionary note, empirical distributions of individual predictions such as in Figure [3.2](#page-12-0) should not be over-interpreted. On one hand, we can have clear theoretical reasons why true $p_0(x)$ ought to be one for many units, e.g. units that did not intend to join the program are likely to be unaffected by a non-binding assignment to treatment. On the other hand, these are just finite-sample estimates from models that are all likely misspecified to a smaller or larger extend. Extreme values that suggest economically unreasonable unit selection effects are likely driven by a high-variance model. We can take a more agnostic perspective with regards to the evidence provided by the empirical distribution of $\hat{p}_0(x)$. In particular, in high dimensions or with complicated functional forms, convergence to true relative selection probabilities is hard to achieve for any model. In finite samples, producing a sparse representation by removing the impact of the treatment in parts of the covariate space could just be a method's solution to optimize fit. Methods for predicting conditional selection probabilities are not necessarily designed to best predict the implied relative $p_0(x)$. However, if a learner performs best among conventional performance metrics for probabilities and does so by producing sparsity with respect the treatment, we treat it as a potentially better approximation and need to deal with the point mass regardless of whether this is a true or approximate model.

4. Identification of Sharp and Smooth Bounds

4.1. Sharp Bounds

4.1.1. The General Case

We denote $a = O(b)$ and $a = O_p(b)$ as $a \leq b$ and $a \leq_P b$, respectively. We first present identification of conditional causal effect bounds for any principal stratum, i.e. parameters of the form

$$
\beta(x, s_0, s_1) = E[Y(1) - Y(0)|S(0) = s_0, S(1) = s_1, X = x].
$$
\n(4.1)

This covers all relevant components for any of the principal strata and margin effects.^{[3](#page-14-1)} Let $q_d(u, x)$ be the *u*-quantile of *Y* conditional on $S = 1, D = d, X = x$. For $d \in \{0, 1\}$, we define

$$
\beta_{1,d}(x,u) = E[Y|S=1, D=d, X=x, Y \le q_d(u,x)], \qquad (4.2)
$$

$$
\beta_{0,d}(x,u) = E[Y|S=1, D=d, X=x, Y \ge q_d(u,x)], \qquad (4.3)
$$

which yields

$$
\beta_{1,d}(x,1) = \beta_{0,d}(x,0) = E[Y|S=1, D=d, X=x].
$$
\n(4.4)

In line with the literature on monotonicity bounds, we assume a continuous outcome.

Assumption 4.1 (Continuity) For all $x \in \mathcal{X}$ and $d \in \{0,1\}$, the conditional outcome *distribution* $P(Y \le y | S = 1, D = d, X = x)$ *is continuous.*

³Trivially, for a principal stratum effect to be well defined, we require that the corresponding population share is non-zero. We assume this throughout.

Moreover, there must be comparable units in terms of covariates in both selected treatment groups.

Assumption 4.2 (Multiple Overlap) Let $m(x) = P(D = 1|X = x)$ and $s(d, x) =$ $P(S = 1|D = d, X = x)$ *. For all* $x \in \mathcal{X}$ and any $d \in \{0, 1\}$ *, we have that* $0 < m(x) < 1$ *and* $0 < s(d, x) < 1$ *.*

Assumptions [3.1](#page-9-3) and [3.2](#page-9-2) imply the specific bounds in terms of truncated means of observed conditional distributions. Assumptions [4.1](#page-14-2) and [4.2](#page-14-3) ensure that the corresponding conditioning sets are non-empty and the relevant truncation quantiles are unique. In particular, we obtain the following identification result for the sharp upper and lower bounds for all principal strata.

Proposition 4.1 (Identification) Let $\mathcal{Y}_d(x)$ be the support of $Y(d)$ conditional on $X =$ *x with* $\overline{y}_d(x)$ *and* $\underline{y}_d(x)$ *its respective infimum and supremum for both* $d \in \{0, 1\}$ *. Under Assumptions [3.1,](#page-9-3) [3.2,](#page-9-2) [4.1,](#page-14-2) and [4.2,](#page-14-3) the sharp lower bounds for the principal strata are given by*

$$
\beta_L(x, s_0, s_1) = \begin{cases}\n\beta_{1,1}(x, \min\{p_0(x), 1\}) - \beta_{0,0}(x, 1 - \min\{1/p_0(x), 1\}) & \text{if } s_0 = 1, s_1 = 1 \\
\beta_{1,1}(x, 1 - \min\{p_0(x), 1\}) - \beta_{0,0}(x, \min\{1/p_0(x), 1\}) & \text{if } s_0 \neq s_1 \\
\underline{y}_1(x) - \overline{y}_0(x) & \text{if } s_0 = 0, s_1 = 0,\n\end{cases}
$$

and the sharp upper bounds by

$$
\beta_U(x, s_0, s_1) = \begin{cases}\n\beta_{0,1}(x, 1 - \min\{p_0(x), 1\}) - \beta_{1,0}(x, \min\{1/p_0(x), 1\}) & \text{if } s_0 = 1, s_1 = 1 \\
\beta_{0,1}(x, \min\{p_0(x), 1\}) - \beta_{1,0}(x, 1 - \min\{1/p_0(x), 1\}) & \text{if } s_0 \neq s_1 \\
\overline{y}_1(x) - \underline{y}_0(x) & \text{if } s_0 = 0, s_1 = 0.\n\end{cases}
$$

For never-takers, all lower and upper bounds are generally uninformative due to never observing them as part of a selected group. For defiers and compliers, the bounds are only informative when at least part of the conditional support is bounded, while for always-takers bounds are also informative without any support condition.

Unconditional bounds for any principal stratum are obtained by integrating the con-

ditional bounds with respect to stratum-specific covariate distributions

$$
\beta_B(s_0, s_1) = \int \beta_B(x, s_0, s_1) dP(x|S(0) = s_0, S(1) = s_1)
$$
\n(4.5)

for $B \in \{L, U\}$, where

$$
dP(x|S(0) = s_0, S(1) = s_1) = \begin{cases} \frac{\min\{s(0,x), s(1,x)\}}{E[\min\{s(0,X), s(1,X)\}]} dP(x) & \text{if } s_0 = 1, s_1 = 1\\ \frac{\max\{0, s(1,x) - s(0,x)\}}{E[\max\{0, s(1,X) - s(0,X)\}]} dP(x) & \text{if } s_0 = 0, s_1 = 1\\ \frac{\max\{0, s(0,x) - s(1,X)\}}{E[\max\{0, s(0,X) - s(1,X)\}]} dP(x) & \text{if } s_1 = 1, s_1 = 0\\ \frac{1 - \max\{s(0,x), s(1,x)\}}{E[1 - \max\{s(0,X), s(1,X)\}]} dP(x) & \text{if } s_1 = 0, s_1 = 0. \end{cases}
$$

4.1.2. Example I: Intensive Margin Bounds

The upper and lower conditional intensive margin or always-taker bounds correspond to the ones given by [Lee](#page-46-0) [\(2009\)](#page-46-0) and [Semenova](#page-46-1) [\(2024\)](#page-46-1). Namely, we have that

$$
\beta_L(x, 1, 1) = \beta_{L,1}(x, 1, 1) - \beta_{L,0}(x, 1, 1) \tag{4.6}
$$

where

$$
\beta_{L,1}(x,1,1) = \begin{cases} E[Y|S=1,D=1,X=x,Y \le q_1(p_0(x),x)] & \text{if } x \in \mathcal{X}^+, \\ E[Y|S=1,D=1,X=x] & \text{if } x \in \mathcal{X}^- \cup \mathcal{X}^0, \\ \beta_{L,0}(x,1,1) = \begin{cases} E[Y|S=1,D=0,X=x] & \text{if } x \in \mathcal{X}^+ \cup \mathcal{X}^0, \\ E[Y|S=1,D=0,X=x,Y \ge q_0(1-1/p_0(x),x)] & \text{if } x \in \mathcal{X}^-. \end{cases} \end{cases}
$$

Similarly, the conditional upper bounds are given by

$$
\beta_U(x, 1, 1) = \beta_{U,1}(x, 1, 1) - \beta_{U,0}(x, 1, 1),\tag{4.7}
$$

where

$$
\beta_{U,1}(x,1,1) = \begin{cases} E[Y|S=1, D=1, X=x, Y \ge q_1(1-p_0(x),x)] & \text{if } x \in \mathcal{X}^+ \\ E[Y|S=1, D=1, X=x] & \text{if } x \in \mathcal{X}^- \cup \mathcal{X}^0, \\ E[Y|S=1, D=0, X=x] & \text{if } x \in \mathcal{X}^+ \cup \mathcal{X}^0, \\ E[Y|S=1, D=0, X=x, Y \le q_0(1/p_0(x),x)] & \text{if } x \in \mathcal{X}^-. \end{cases}
$$

The unconditional bounds for the effect at the intensive margin can then be recovered as

$$
\beta_B(1,1) = \int \beta_B(x,1,1) dP(x|S(0) = S(1) = 1)
$$

=
$$
\frac{\int \beta_B(x,1,1) \min\{s(0,x),s(1,x)\} dP(x)}{\int \min\{s(0,x),s(1,x)\} dP(x)}.
$$
 (4.8)

4.1.3. Example II: Extensive Margin Bounds

The extensive margin is the combination of compliers and defiers where $s_0 \neq s_1$. Thus, for $B \in \{L, U\}$, the extensive margin effect bounds can be written as

$$
\beta_B(em) = \int \beta_B(x, em)dP(x|S(0) \neq S(1)). \tag{4.9}
$$

where

$$
dP(x|S(0) \neq S(1))
$$

=
$$
\frac{P(S(0) \neq S(1)|X = x)dP(x)}{P(S(0) \neq S(1))}
$$

=
$$
\frac{[P(S(0) = 0, S(1) = 1|X = x) + P(S(0) = 1, S(1) = 0|X = x)] dP(x)}{\int [P(S(0) = 0, S(1) = 1|X = x) + P(S(0) = 1, S(1) = 0|X = x)] dP(x)}
$$

=
$$
\frac{|s(1, x) - s(0, x)| dP(x)}{\int |s(1, x) - s(0, x)| dP(x)}.
$$

Combining compliers and defiers from Proposition [4.1](#page-15-0) then yields the conditional lower bound

$$
\beta_{L,1}(x, em) = \begin{cases} E[Y|S=1, D=1, X=x, Y \le q_1(1-p_0(x), x)] & \text{if } x \in \mathcal{X}^+, \\ \underline{y}_1(x) & \text{if } x \in \mathcal{X}^- \cup \mathcal{X}^0, \end{cases}
$$

and

$$
\beta_{L,0}(x, em) = \begin{cases} \overline{y}_0(x) & \text{if } x \in \mathcal{X}^+ \cup \mathcal{X}^0, \\ E[Y|S=1, D=0, X=x, Y \ge q_0(1/p_0(x), x)] & \text{if } x \in \mathcal{X}^-.\end{cases}
$$

The conditional upper bound $\beta_U(x, em)$ is found analogously from from Proposition [4.1.](#page-15-0) As a result, the unconditional bounds at the extensive margin are given by

$$
\beta_B(em) = \frac{\int \beta_B(x, em) |s(1, x) - s(0, x)| dP(x)}{\int |s(1, x) - s(0, x)| dP(x)} \tag{4.10}
$$

4.2. Smooth Bounds

The bounds in Proposition [4.1](#page-15-0) are sharp. However, they, as well as the strata densities, contain a variety of non-differentiable components. This poses a challenge to statistical inference. We demonstrate the specific regularity problem and its consequences for inference in Section [5.](#page-21-0) We now introduce the smooth bounds that provide an outer identification region to circumvent these problems. In particular, the identification region is constructed to eliminate the impact of misclassifying different monotonicity types while still remaining close to the sharp identified set. We focus on the intensive margin bounds for simplicity but the method works equivalently for all principal strata. More specifically, the form of the bounds in Proposition [4.1](#page-15-0) reveal two sources where classification of the partition will enter, (i) conditional effects bounds $\beta_B(x, 1, 1)$, and (ii) conditional alwaystaker share min ${s(0, x), s(1, x)}$. We suggest to apply repeated, asymmetric smoothing to both components. For (i), consider first the conditional bound $\beta_L(x, 1, 1)$. Note that, for any $x \in \mathcal{X}$, the following parameter is a valid lower bound for $\beta_L(x, 1, 1)$

$$
\beta_{L,h}(x,1,1) = E[Y|S=1, D=1, X=x, Y \le q_1(g_{1,h}(p_0(x)),x)]
$$

$$
- E[Y|S=1, D=0, X=x, Y \ge q_0(1-g_{1,h}(1/p_0(x)),x)]
$$

$$
= \beta_{1,1}(x, g_{1,h}(p_0(x))) - \beta_{0,0}(x, 1-g_{1,h}(1/p_0(x)))
$$
(4.11)

where $g_{1,h}(z)$ is a smooth monotonic function indexed by a smoothing parameter $h \in$ $\mathcal{H}_n \subset \mathbb{R}^+$ such that $0 \leq g_{1,h}(z) \leq \min\{z, 1\}$. This ensures that

$$
\beta_{L,h}(x,1,1) \le \beta_L(x,1,1). \tag{4.12}
$$

Next, consider that the smoothing of the conditional always-taker share is given by $\min\{s(0, x), s(1, x)\} = s(1, x) \min\{p_0(x), 1\}.$ We obtain lower and upper bounds using smooth functions $g_{1,h}(z)$ and $g_{3,h}(z)$ such that $g_{1,h}(z) \le \min\{z,1\} \le g_{3,h}(z)$. In which direction to smooth, depends on the sign of $\beta_{L,h}(x,1,1)$. We decompose this into the difference of two non-negative terms via $z = \max\{z, 0\} - \max\{-z, 0\}$. We approximate these two functions with a smoothing function $g_{4,h}(z) \leq \max\{z, 0\}$, and the earlier $g_{2,h}(-z) \ge \max\{-z,0\}$. The unconditional smooth lower effect bound is then given by

$$
\beta_{L,h}(1,1) = \frac{E\left[g_{4,h}(\beta_{L,h}(x,1,1))g_{1,h}(p_0(X))s(1,X)\right]}{E[g_{3,h}(p_0(X))s(1,X)]}
$$

$$
-\frac{E[g_{2,h}(-\beta_{L,h}(x,1,1))g_{3,h}(p_0(X))s(1,X)]}{E[g_{1,h}(p_0(X))s(1,X)]}.
$$
(4.13)

The aforementioned relations together yield

$$
\beta_{L,h}(1,1) \le \beta_L(1,1). \tag{4.14}
$$

Analogously, the conditional smooth upper bound is given by

$$
\beta_{U,h}(x,1,1) = \beta_{0,1}(x,1-g_{1,h}(p_0(x))) - \beta_{1,0}(x,g_{1,h}(1/p_0(x)))
$$
\n(4.15)

which obeys the inequality $\beta_{U,h}(x,1,1) \geq \beta_U(x,1,1)$, and the unconditional smooth upper bound is defined as

$$
\beta_{U,h}(1,1) = \frac{E[g_{2,h}(\beta_{U,h}(x,1,1))g_{3,h}(p_0(X))s(1,X)]}{E[g_{1,h}(p_0(X))s(1,X)]} - \frac{E[g_{4,h}(-\beta_{U,h}(x,1,1))g_{1,h}(p_0(X))s(1,X)]}{E[g_{3,h}(p_0(X))s(1,X)]},
$$
\n(4.16)

which analogously satisfies the reversed inequality

$$
\beta_{U,h}(1,1) \ge \beta_U(1,1). \tag{4.17}
$$

Next, we propose a class of functions that satisfy the imposed conditions and show that the distance between smooth and sharp unconditional bounds is negligible as $h \to 0$ with bounded support.

Assumption 4.3 (Outcome Regularity I) *For* $d \in \{0, 1\}$ *, the outcome distribution* $P(Y \leq y | S = 1, D = d, X = x)$ *has uniformly bounded support, i.e.* $\sup_{x \in \mathcal{X}} (|\underline{y}_d(x)| +$ $|\overline{y}_d(x)|$) < ∞ .

We also impose the following assumption about approximation functions.

Assumption 4.4 (Approximation Functions) *Approximation functions gi,h, i* = 1*, . . . ,* 4 *satisfy for all* $z \in \mathbb{R}$

$$
g_{1,h}(z) \le \min\{z, 1\} \le g_{3,h}(z)
$$
 and $g_{4,h}(z) \le \max\{z, 0\} \le g_{2,h}(z)$,

and for each $i = 1, \ldots, 4$

$$
\sup_{z \in \mathbb{R}} |g_{i,h}(z) - g_i(z)| \lesssim h,
$$

where $g_1(z) = g_3(z) = \min\{z, 1\}$ *and* $g_2(z) = g_4(z) = \max\{z, 0\}$ *. In addition,* $g_{1,h}(z) \ge 0$ *for all* $z \in \mathbb{R}$ *and* $h \in \mathcal{H}_n$.^{[4](#page-20-0)}

There exist many functions with these properties. Figure [4.1](#page-21-1) depicts an example using a LogSumExp function for smoothing via $q_{1,h}(z)$. We obtain the following Theorem.

Theorem 4.1 *Let* P *be the set of probability measures satisfying Assumptions [3.1,](#page-9-3) [3.2,](#page-9-2)* [4.1,](#page-14-2) [4.2,](#page-14-3) [4.3,](#page-20-1) and [4.4.](#page-20-2) Denote $\underline{g}_n = \inf_x g_{1,h}(p_0(x))$. Then, for $B \in \{L, U\}$, and $(s_0, s_1) \neq$ (0*,* 0)*, we have that*

$$
\sup_{P \in \mathcal{P}} |\beta_{B,h}(s_0, s_1) - \beta_B(s_0, s_1)| \lesssim \frac{h}{\underline{g}_n^2}.
$$

Theorem [4.1](#page-20-3) gives the approximation error of the smooth bounds. Under the strong overlap condition, g_n is bounded away from zero and thus the difference decays linearly

⁴All quantile trimming thresholds are well defined as long as $\inf_{x \in \mathcal{X}} g_{1,h}(p_0(x)) \geq 0$ and $\inf_{x \in \mathcal{X}} g_{1,h}(1/p_0(x)) \geq 0$. Given the strong multiple overlap assumption, one can always find an upper bound for *h* such that this applies. These large-*h* values are not relevant from a practical perspective and we consider *h* to be below such a ceiling in what follows.

Figure 4.1: Smooth Approximation: *g*1*,h*(*z*) Example

The figure contains min $\{p_0(x), 1\}$ as well as two smooth approximations using $g_{1,h}(z) = 1 - h \log(1 +$ $\exp(-(z-1)/h)$ for $h = 0.30, h = 0.15,$ and $h = 0.0015$ (Asy). The size of the *h*-cover at $p_0(x) = 1$ determines how close the bounds are to the point identified effect for $x \in \mathcal{X}_0$. The *h*-cover for units with $p_0(x) \neq 1$ introduces an additional distance compared to the sharp trimming bounds.

in *h*. Without strong overlap, there is an implicit restriction on the rate at which the smooth relative probabilities $g_{1,h}(p_0(x))$ can approach zero.

5. Regularity and Semiparametric Efficiency Bounds

In this section, we present our main results on regularity and semiparametric efficiency. Again, we focus on the lower bound for the intensive margin for simplicity. Table [6.1](#page-26-0) provides the influence function for the always-taker lower bound and its smoothed counterpart. All other influence functions are in Appendix [A.](#page-48-0) We obtain the following theorem.

Assumption 5.1 (Outcome Regularity II) *For all* $x \in \mathcal{X}$ *and* $d \in \{0, 1\}$ *, the conditional outcome distribution* $P(Y \leq y | S = 1, D = d, X = x)$ *has finite support on* $[\underline{y}_d, \overline{y}_d]$ and a continuous density $f(y|X=x, D=d, S=1)$ bounded from below and above.

Assumption 5.2 (Strong Multiple Overlap) *There exist constants* $m, s \in (0, 1/2)$ *such that*

$$
\underline{m} < \inf_{x \in \mathcal{X}} m(x) \le \sup_{x \in \mathcal{X}} m(x) < 1 - \underline{m},
$$
\n
$$
\underline{s} < \inf_{x \in \mathcal{X}, d \in \{0, 1\}} s(d, x) \le \sup_{x \in \mathcal{X}, d \in \{0, 1\}} s(d, x) < 1 - \underline{s}.
$$

Assumptions [5.1](#page-21-2) and [5.2](#page-21-3) are stronger versions of continuity and overlap in Assumptions [4.1](#page-14-2) and [4.2,](#page-14-3) respectively. Without strong overlap, there is an additional irregularity problem for the population bounds equivalently to irregular identification of average treatment effects under unconfoundedness with many extreme propensities scores [\(Khan](#page-46-13) [and Tamer, 2010;](#page-46-13) [Heiler and Kazak, 2021\)](#page-45-11). We abstract from such issues in this paper to focus on the irregularity obtained from $P(\mathcal{X}^0) > 0$.

Theorem 5.1 *Suppose that Assumptions* [3.1,](#page-9-3) [3.2,](#page-9-2) [5.1,](#page-21-2) and [5.2](#page-21-3) hold. For $B \in \{L, U\}$, $\beta_B(1,1)$, $\beta_B(0,1)$, $\beta_B(0,1)$, $\beta_B(em)$ are pathwise differentiable if and only if $P(X^0) = 0$.

Theorem [5.1](#page-22-0) characterizes that the necessary and sufficient condition for the unconditional lower bound introduced in [\(4.8\)](#page-17-0) to be pathwise differentiable is the absence of the point mass on the set \mathcal{X}^0 , i.e. $P(\mathcal{X}^0) = 0.5$ $P(\mathcal{X}^0) = 0.5$ We also present the semiparametric efficiency bound for $\beta_L(1,1)$. It does not depend on the knowledge of the propensity score.

Corollary 5.1 *Let* $1_{\mathcal{X}^+} = \mathbb{1}(X \in \mathcal{X}^+)$ *and* $1_{\mathcal{X}^-} = \mathbb{1}(X \in \mathcal{X}^-)$ *. Suppose that Assumptions* [3.1,](#page-9-3) [3.2,](#page-9-2) [5.1](#page-21-2) and [5.2](#page-21-3) hold, and $P(X^0) = 0$. The semiparametric efficiency bound

⁵When $P(X^0) > 0$, the result in Theorem [5.1](#page-22-0) that $\beta_L(1,1)$ is not pathwise differentiable is related but different to [Luedtke and van der Laan](#page-46-8) [\(2016\)](#page-46-8). Their target is essentially the denominator of $\beta_L(1,1)$ in [\(4.8\)](#page-17-0). However, divergence of the derivative of denominator and/or divergence of the numerator separately does not necessarily imply divergence of the ratio. Moreover, the numerator includes the conditional treatment effect bounds $\beta_L(1,1,x)$ and consequently conditional quantiles $q_d(u,x)$ which changes the analysis.

Var_a(β ^{*L*}(1*,* 1)) *for* β ^{*L*}(1*,* 1) *is defined by*

$$
E[\min(s(0, X), s(1, X)]^{2}Var_{a}(\beta_{L}(1, 1)) = E\left[\frac{s(1, X)\sigma_{1}^{2}(X)}{m(X)} + \frac{s(0, X)\sigma_{0}^{2}(X)}{1 - m(X)}\right]
$$

+
$$
E\left[1_{\mathcal{X}^{+}(\beta_{L}(X, 1, 1) - \beta_{L}(1, 1))^{2}}\frac{s(0, X)(1 - s(0, X)m(X))}{1 - m(X)}\right]
$$

+
$$
E\left[1_{\mathcal{X}^{-}(\beta_{L}(X, 1, 1) - \beta_{L}(1, 1))^{2}}\frac{s(1, X)(1 - s(1, X) + s(1, X)m(X))}{m(X)}\right]
$$

+
$$
E\left[1_{\mathcal{X}^{+}}\frac{s(1, X)q_{1}(p_{0}(X), X)^{2}p_{0}(X)(1 - p_{0}(X))}{m(X)}\right]
$$

+
$$
E\left[1_{\mathcal{X}^{-}}\frac{s(0, X)q_{0}(1 - 1/p_{0}(X), X)^{2}p_{0}(X)^{-1}(1 - p_{0}(X)^{-1})}{1 - m(X)}\right]
$$

+
$$
E\left[1_{\mathcal{X}^{+}}(q_{1}(p_{0}(X), X) - \beta_{1,1}(X, p_{0}(X)))^{2}\left(\frac{s(0, X)(1 - s(0, X))}{1 - m(X)} + \frac{p_{0}(X)^{2}s(1, X)(1 - s(1, X))}{m(X)}\right)\right]
$$

+
$$
E\left[1_{\mathcal{X}^{-}}(q_{0}(1 - 1/p_{0}(X), X) - \beta_{0,0}(X, 1 - 1/p_{0}(X)))^{2}\right]
$$

$$
\times \left(\frac{p_{0}(X)^{-2}s(0, X)(1 - s(0, X))}{1 - m(X)} + \frac{s(1, X)(1 - s(1, X))}{m(X)}\right)\right]
$$

-
$$
2E\left[1_{\mathcal{X}^{+}}\frac{q_{1}(p_{0}(X), X)\beta_{1,1}(X, p_{0}(X))s(1, X)p_{0}(X)(1 - p_{0}(X)))}{1 - m(X)}\right]
$$

-
$$
2E\left[1_{\mathcal{X}^{-}}\frac{q
$$

where

$$
\sigma_1^2(x) = Var[Y1_{\{Y \le q_1(\min\{p_0(x), 1\}, x)\}}|S=1, D=1, X=x],
$$

$$
\sigma_0^2(x) = Var[Y1_{\{Y \ge q_0(\max\{1-1/p_0(x), 0\}, x)\}}|S=1, D=0, X=x].
$$

The setting with full selection, $S = 1$ almost surely, renders any truncation redundant and thus $\beta_L(x, 1, 1) = \beta_U(x, 1, 1) = \beta(x, 1, 1)$ are equal to the conditional average treatment effect. Consequently, the efficiency bound in Corollary [5.1](#page-22-2) becomes

$$
E\left[\frac{\sigma_1^2(X)}{m(X)} + \frac{\sigma_0^2(X)}{1 - m(X)} + (\beta(X, 1, 1) - \beta(1, 1))^2\right],\tag{5.1}
$$

with

$$
\sigma_d^2(x) = Var[Y|S=1, D=d, X=x],
$$

which is exactly the efficiency bound for the ATE obtained in Theorem 2 of [Hahn](#page-44-4) [\(1998\)](#page-44-4), i.e. Corollary [5.1](#page-22-2) nests this as a special case. The smooth bounds [\(4.13\)](#page-19-0) circumvent the irregularity problem by relaxing the width of the identified set using smooth approximators, i.e. the *g*-functions. If the latter are sufficiently smooth, then the resulting outer

,

bounds are pathwise differentiable. In particular, we obtain the following Theorem:

Theorem [5.2](#page-21-3) *Suppose that Assumptions* [3.1,](#page-9-3) [3.2,](#page-9-2) [4.4,](#page-20-2) [5.1,](#page-21-2) 5.2 *hold.* If $\sup_{z,j} g'_{i,h}(z) \lesssim 1$ *for some* $h > 0$ *, then* $\beta_{L,h}(1,1)$ *is pathwise differentiable.*

The corresponding efficiency bound is in Appendix [5.2.](#page-23-0) The additional smoothness assumption applies to the LogSumExp example shown in Section [4.2.](#page-18-0)

6. Estimation and Inference

6.1. Estimators

We propose to construct estimators by using empirical analogues to the efficient influ-ence functions in Table [6.1](#page-26-0) with estimated nuisances. Let $E_n[X] = n^{-1} \sum_i^n X_i$. For given nuisance estimates $\hat{\eta}$, any estimator $\hat{\beta}_B = \hat{\beta}_B(s_0, s_1)$ solves

$$
E_n[\psi_{\beta_B}(W,\hat{\eta},\hat{\beta}_B)] = 0 \tag{6.1}
$$

Note that all principal strata and margin influence functions have the following linear structure

$$
\psi_{\beta_B}(W,\eta,\beta_B) = \psi_{\beta_B}^{[S]}(W,\eta) - \psi_{\beta_B}^{[B]}(W,\eta)\beta_B.
$$
\n(6.2)

Thus, the influence function-based estimators have a ratio structure of the form

$$
\hat{\beta}_B = \frac{E_n[\psi_{\beta_B}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_B}^{[S]}(W,\hat{\eta})]}.
$$
\n(6.3)

For the smooth bounds, parameters are given by a sum

$$
\beta_{B,h} = \beta_{B,+,h} + \beta_{B,-,h},\tag{6.4}
$$

whose components also have linear influence functions with structure

$$
\psi_{\beta_{B,+,h}}(W,\eta,\beta_B) = \psi_{\beta_{B,+,h}}^{[S]}(W,\eta) - \psi_{\beta_{B,+,h}}^{[B]}(W,\eta)\beta_{B,+,h},\tag{6.5}
$$

$$
\psi_{\beta_{B,-,h}}(W,\eta,\beta_{B,-,h}) = \psi_{\beta_{B,-,h}}^{[S]}(W,\eta) - \psi_{\beta_{B,-,h}}^{[B]}(W,\eta)\beta_{B,-,h},\tag{6.6}
$$

which combined with estimated nuisances yield moment estimators

$$
\hat{\beta}_{B,h} = \hat{\beta}_{B,+,h} + \hat{\beta}_{B,-,h}
$$
\n
$$
= \frac{E_n[\psi_{\beta_{B,+,h}}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_{B,+,h}}^{[S]}(W,\hat{\eta})]} + \frac{E_n[\psi_{\beta_{B,-,h}}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_{B,-,h}}^{[S]}(W,\hat{\eta})]}.
$$
\n(6.7)

For the intensive margin, we also consider the moment functions suggested by [Semenova](#page-46-1) [\(2024\)](#page-46-1) and [Heiler](#page-45-1) [\(2024\)](#page-45-1) under known propensity scores that have influence function

$$
\tilde{\psi}_{\beta_B}(W,\eta,\beta_B) = \psi_{\beta_B}(W,\eta,\beta_B) - \delta(D,X,\eta),\tag{6.8}
$$

with mean-zero term $\delta(D, X, \eta)$ characterized by

$$
E[\min\{s(0, X), s(1, X)\}]\delta(D, X, \eta)
$$

= $\mathbb{1}_{(X \in \mathcal{X}^+)}s(0, X)\left[\beta_{1,1}(X, p_0(X))\left(1 - \frac{D}{m(X)}\right)\right]$

$$
- \beta_{0,0}(X, 0)\left(1 - \frac{1 - D}{1 - m(X)}\right)\right]
$$

+ $\mathbb{1}_{(X \in \mathcal{X}^-)}s(1, X)\left[\beta_{1,1}(X, 1)\left(1 - \frac{D}{m(X)}\right)\right]$

$$
- \beta_{0,0}(X, 1 - 1/p_0(X))\left(1 - \frac{1 - D}{1 - m(X)}\right)\right].
$$
 (6.9)

Solving for the parameter at the sample average then yields the alternative estimator

$$
\tilde{\beta}_B = \frac{E_n[\tilde{\psi}_{\beta_B}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_B}^{[S]}(W,\hat{\eta})]}
$$
(6.10)

6.2. Definitions and Assumptions

In this section, we provide and discuss the assumptions and results for large sample inference for the regular and irregular case. Denote $||\cdot||_p$ as the L_p norm. Let $\eta \in T$ where *T* is a convex subset of some normed vector space. We assume all nuisance functions are cross-fitted as in [Chernozhukov et al.](#page-44-12) [\(2018\)](#page-44-12), Definition 3.1 and use the LogSumExp func-tion for smoothing, see Section [4.2.](#page-18-0) Denote nuisance realization set $\mathcal{T}_n = (\mathcal{S}_{0,n} \times \mathcal{S}_{1,n} \times$ $\mathcal{Q}_{0,n} \times \mathcal{Q}_{1,n} \times \mathcal{M}_n \times \mathcal{B}_{0,0,n} \times \mathcal{B}_{0,1,n} \times \mathcal{B}_{1,0,n} \times \mathcal{B}_{1,0,n}$ C T as the set that with high prob-

Partition	Component	Non-centered Influence Function		
\mathcal{X}^+	$\psi_{\beta}^{[L^+]}$	$\frac{SD}{m(X)}Y1\{Y \leq q_1(p_0(X), X)\} - \frac{S(1-D)}{1-m(X)}Y$ $-\frac{SD}{m(X)}q_1(p_0(X),X)[1\{Y \leq q_1(p_0(X),X)\}-p_0(X)]$ $+q_1(p_0(X), X) \left \frac{1-D}{1-m(X)}(S-s(0, X)) - p_0(X) \frac{D}{m(X)}(S-s(1, X)) \right $ $+ s(0, X) \left[\beta_{1,1}(X, p_0(X)) \left(1 - \frac{D}{m(X)} \right) - \beta_{0,0}(X, 0) \left(1 - \frac{1 - D}{1 - m(X)} \right) \right]$		
	$\psi_{\beta}^{[S^+]}$	$s(0, X) + \frac{(1-D)(S-s(0, X))}{1-m(X)}$		
\mathcal{X}^-	$\psi_{\beta}^{\lfloor L^{-} \rfloor}$	$\frac{SD}{m(X)}Y - \frac{S(1-D)}{1-m(X)}Y1\{Y \ge q_0(1-1/p_0(X),X)\}\$ $-\frac{S(1-D)}{1-m(X)}q_0(1-1/p_0(X),X)\left[1/p_0(X)-1\{Y\geq q_1(p_0(X),X)\}\right]$ $-q_1(p_0(X), X) \left \frac{D}{m(X)}(S-s(1, X)) - p_0(X)^{-1} \frac{1-D}{1-m(X)}(S-s(0, X)) \right $ $+ s(1, X) \left[\beta_{1,1}(X, 1) \left(1 - \frac{D}{m(X)} \right) - \beta_{0,0}(X, 1 - 1/p_0(X)) \left(1 - \frac{1 - D}{1 - m(X)} \right) \right]$		
	$\psi_{\beta}^{[S^-]}$	$s(1,X) + \frac{D(S-s(1,X))}{m(X)}$		
$\psi_{\beta} = [\mathbb{1}(X \in \mathcal{X}^+) \psi_{\beta}^{L^+} + \mathbb{1}(X \in \mathcal{X}^-) \psi_{\beta}^{L^-}] - [\mathbb{1}(X \in \mathcal{X}^+) \psi_{\beta}^{S^+} + \mathbb{1}(X \in \mathcal{X}^-) \psi_{\beta}^{S^-}] \beta$				
Nuisance		Definition		
m(x) s(d,x) $q_d(u,x)$ $\beta_{1,d}(x,u)$ $\beta_{0,d}(x,u)$		$P(D = 1 X = x)$ $P(S = 1 D = d, X = x)$ $\inf\{q \in \mathcal{Y} : u \leq P(Y \leq q S = 1, D = d, X = x)\}\$ $E[Y S = 1, D = d, X = x, Y \leq q_d(u, x)]$ $E[Y S=1, D=d, X=x, Y>q_d(u,x)]$		

Table 6.1: Influence Functions: Regular Bound $\beta = \beta_L(1, 1)$

ability contains estimators $\hat{\eta} = \{\hat{s}(0, X), \hat{s}(1, X), \hat{q}_0(u, X), \hat{q}_1(u, X), \hat{m}(u, X), \hat{\beta}_{0,0}(u, X),\}$ $\hat{\beta}_{0,1}(u, X), \hat{\beta}_{1,0}(u, X), \hat{\beta}_{1,1}(u, X)$ for nuisance quantities $\eta = \{s(0, X), s(1, X), q_0(u, X),$ $q_1(u, X), m(X), \beta_{0,0}(u, X), \beta_{0,1}(u, X), \beta_{1,0}(u, X), \beta_{1,1}(u, X)$ }. Let their corresponding L_p error rates be

$$
\lambda_{s,n,p} = \sup_{d \in \{0,1\}} \sup_{\hat{s}(d) \in \mathcal{S}_{d,n}} E[|\hat{s}(d, X) - s(d, X)|^p]^{1/p},
$$

\n
$$
\lambda_{q,n,p} = \sup_{d \in \{0,1\}} \sup_{u \in \tilde{U}} \sup_{\hat{q}_d(u) \in \mathcal{Q}_{d,n}} E[|\hat{q}_d(u, X) - q_d(u, X)|^p]^{1/p},
$$

\n
$$
\lambda_{m,n,p} = \sup_{\hat{m} \in \mathcal{M}_n} E[|\hat{m}(X) - m(X)|^p]^{1/p},
$$

\n
$$
\lambda_{b,n,p} = \sup_{j,d \in \{0,1\}} \sup_{u \in \tilde{U}} \sup_{\hat{\beta}_{j,d}(u) \in \mathcal{B}_{j,d,n}} E[|\hat{\beta}_{j,d}(u, X) - \beta_{j,d}(u, X)|^p]^{1/p},
$$

Function	Definition
$f^{[L]}(j,k)$	$\frac{SD}{m(X)} \frac{g'_{k,h}(\beta_{L,h}(X))g_{j,h}(p_0(X))}{g_{1,h}(p_0(X))} \left[Y1\right] (Y \leq q_1(g_{1,h}(p_0(X)), X) - g_{1,h}(p_0(X))\beta_{1,1}(x, g_{1,h}(p_0(x)))$ $-\frac{S(1-D)}{1-m(X)}\frac{g'_{k,h}(\beta_{L,h}(X))g_{j,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X))}$ $\times [Y1(Y \ge q_0(1 - g_{1,h}(1/p_0(X)), X)) - g_{1,h}(1/p_0(X))\beta_{0,0}(x, 1 - g_{1,h}(p_0(x)))]$ $+g_{k,h}(\beta_{L,h}(X)) g_{j,h}(p_0(X))s(1,X) + g'_{j,h}(p_0(X))\frac{1-D}{1-m(X)}(S-s(0,X)) $ $+g_{k,h}(\beta_{L,h}(X)) g_{j,h}(p_0(X))-p_0(X)g'_{j,h}(p_0(X))\frac{D}{m(X)}(S-s(1,X)) $ $-\frac{SD}{m(X)} \frac{g'_{k,h}(\beta_{L,h}(X))g_{j,h}(p_0(X))}{g_{1,h}(p_0(X))} q_1(g_{1,h}(p_0(X)),X) \left[\mathbb{1}(Y \leq q_1(g_{1,h}(p_0(X)),X) - g_{1,h}(p_0(X)) \right]$ $\qquad \qquad + \tfrac{S(1-D)}{1-m(X)} \tfrac{g'_{k,h}(\beta_{L,h}(X))g_{j,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X))}-q_0\big(1-g_{1,h}\big(1/p_0(X)\big),X\big)$ $\times \left[\mathbb{1}(Y \geq q_0(1-g_{1,h}(1/p_0(X)), X) - g_{1,h}(1/p_0(X)) \right]$ $+g'_{k,h}(\beta_{L,h}(X))\left \frac{1-D}{1-m(X)}(S-s(0,X))-p_0(X)\frac{D}{m(X)}(S-s(1,X))\right $ $\times \Big \frac{g'_{1,h}(p_0(X))g_{j,h}(p_0(X))}{g_{1,h}(p_0(X))}(q_1(g_{1,h}(p_0(X)),X)-\beta_{1,1}(x,g_{1,h}(p_0(x)))\Big $ $+\frac{g'_{1,h}(1/p_0(X))g_{j,h}(p_0(X))}{p_0^2(X)g_{1,h}(1/p_0(X))}(q_0(1-g_{1,h}(1/p_0(X)),X)-\beta_{0,0}(x,1-g_{1,h}(p_0(x))))\Big $
$f^{[S]}(j)$	$ g_{j,h}(p_0(X))s(1,X)+g'_{j,h}(p_0(X))\frac{1-D}{1-m(X)}(S-s(0,X)) $ $ g_{j,h}(p_0(X)) + p_0(X)g'_{j,h}(p_0(X))\frac{D}{m(X)}(S-s(1,X)) $
$\begin{array}{l} {\psi}_{\beta_{L,+},h}^{[L]} \\ {\psi}_{\beta_{L,+},h}^{[S]} \\ {\psi}_{\beta_{L,-},h}^{[L]} \\ {\psi}_{\beta_{L,-},h}^{[S]} \\ \end{array}$ $\psi_{\beta_{L,+,h}}$ $\psi_{\beta_{L,-,h}}$	$f^{[L]}(1,4)$ $f^{[S]}(3)$ $f^{[L]}(3,5)$ $f^{[S]}(1)$ $\begin{array}{l} \psi_{\beta_{L,+},h}^{[S]} - \psi_{\beta_{L,+},h}^{[L]} \beta_{L,+,h} \\ \psi_{\beta_{L,-},h}^{[S]} - \psi_{\beta_{L,-},h}^{[L]} \beta_{L,-,h} \end{array}$
Nuisance	Definition
m(x) s(d,x) $q_d(u,x)$ $\beta_{L,h}(x)$ $\beta_{1,d}(x,u)$ $\beta_{0,d}(x,u)$	$P(D = 1 X = x)$ $P(S = 1 D = d, X = x)$ $\inf\{q \in \mathcal{Y} : u \leq P(Y \leq q S = 1, D = d, X = x)\}\$ $\beta_{1,1}(x,g_{1,h}(p_0(x)) - \beta_{0,0}(x,1-g_{1,h}(p_0(x)))$ $E[Y S = 1, D = d, X = x, Y \leq q_d(u, x)]$ $E[Y S = 1, D = d, X = x, Y \ge q_d(u, x)]$
	g -functions $g_{1,h}(z) \leq \min\{z,1\} \leq g_{3,h}(z)$ $g_{4,h}(z) \le \max\{z,0\} \le g_{2,h}(z)$ $g_{5,h}(z) = -g_{2,h}(-z)$

Table 6.2: Influence Functions: Smooth Bound $\beta_{L,h}(1,1) = \beta_{L,+h}(1,1) + \beta_{L,-h}(1,1)$

where \tilde{U} is a compact subset of $(0,1)$ containing the relevant quantile trimming threshold support unions.^{[6](#page-27-0)}

 6 For the regular case, this will be $([\text{supp}(p_0(X)) \cup \text{supp}(1-p_0(X))] \cap \mathcal{X}^+) \cup ([\text{supp}(1/p_0(X)) \cup \text{supp}(1-p_0(Y))]$ $1/p_0(X)$] ∩ \mathcal{X}^-) while for the smoothed bounds, the subset will depend on h and is defined equivalently with all bounds replaced by their $g_{1,h}$ -smoothed counterparts.

Assumption 6.1 (Outcome Regularity III) *The outcome has a continuous conditional density* $f(y|X=x, D=d, S=s)$ *that is uniformly bounded from above and away from zero with bounded first derivative for any* $x \in \mathcal{X}$ and $s, d \in \{0, 1\}$.

Assumption [6.1](#page-26-1) here strengthens the smoothness and moment assumptions. This is required to bound (higher order) terms in some of the expansions and a central limit theorem for transformations of the outcome with bounded weights.

6.3. Regular Bounds

We first consider the regular case that sets $P(X^0) = 0$. For this, we require the density around of $s(1, x) - s(0, x)$ in a neighborhood around zero to be well-behaved. This neighborhood is allowed shrink at a rate proportional to the error over the realization set for the selection probabilities. In particular,

Assumption 6.2 (Margin) *There exist a finite constant* $C > 0$ *and* $\alpha > 0$ *such that*

$$
P(|s(1, x) - s(0, x)| \le \delta) \le C\delta^{\alpha} \quad \text{for any} \ \ 0 \le \delta \le c_n
$$

and $c_n \lesssim \lambda_{s,n,2}^{1/(2+\alpha)}$ $_{s,n,2}^{1/(2+\alpha)}$.

Note that this implies that $P(X^0) = 0$ which we know is necessary for the existence of a first-order uniformly unbiased estimator by Theorem [5.1.](#page-22-0) Any continuous bounded density for $s(1, x) - s(0, x)$ is sufficient yielding $\alpha = \infty$. However, a density may also diverge around zero at a controlled rate. We further require the following learning rates.

Assumption 6.3 (Machine Learning Bias I) Let $e_n = o(1)$. For all folds, the nui*sance parameters obtained via cross-fitting belong to a shrinking neighborhood* \mathcal{T}_n *around η with probability of at least* $1 - e_n$ *, such that*

$$
\lambda_{s,n,1} + \lambda_{s,n,2}^{\frac{\alpha}{2+\alpha}} + \lambda_{q,n,1} + \lambda_{q,n,2} + \lambda_{m,n,2} + \lambda_{b,n,2} = o(1)
$$

and

$$
\sqrt{n}(\lambda_{s,n,2}^{\frac{2\alpha}{2+\alpha}} + \lambda_{q,n,2}^2 + \lambda_{m,n,2}^2 + \lambda_{b,n,2}^2) = o(1).
$$

One can see that we require L_1 and L_2 consistency for quantiles and conditional selection probabilities. The *L*¹ requirement is due to the variance of the trimming indicator for the outcome. Moreover, to control the machine learning bias, the squared L_2 rates have to be sufficiently fast. For most nuisances, these are equivalent to the usual $o(n^{-1/4})$ root mean squared error (RMSE) requirement in the debiased machine learning literature [\(Chernozhukov et al., 2018\)](#page-44-12). The more demanding rate requirement for the selection probabilities is due to difficulties of classifying positive and negative monotonicity types around the boundary. Given these rates, Assumption [6.2](#page-28-0) is only really plausible for $\alpha > 2$ as, even in the parametric case where $\lambda_{s,n,2} \sim n^{-1/2}$, $\sqrt{n}\lambda_{s,n,2}^{2\alpha/(2+\alpha)} \nrightarrow 0$ when $\alpha \leq 2$. For more demanding selection probabilities in high dimensions or with complicated functional forms, the density needs to be increasingly well-behaved around the margin. As $\alpha \to \infty$, the requirement reduces to the typical $o(n^{-1/4})$ rate as in [Heiler](#page-45-1) [\(2024\)](#page-45-1). We obtain the following Theorem.

Theorem 6.1 *Under Assumptions [3.1,](#page-9-3) [3.2,](#page-9-2) [5.2,](#page-21-3) [6.1,](#page-26-1) [6.2,](#page-28-0) and [6.3,](#page-28-1) the regular estimator is asymptotically normal and semiparametrically efficient, i.e.*

$$
\sqrt{n}(\hat{\beta}_L(1,1) - \beta_L(1,1)) \stackrel{d}{\rightarrow} \mathcal{N}\big(0, Var_a(\beta_L(1,1))\big),
$$

where $Var_a(\beta_L(1,1))$ *is the semiparametric efficiency bound in Theorem [5.1.](#page-22-0)*

6.4. Smooth Bounds

We now consider the large-sample behavior of the smooth bounds. We impose the following learning requirements.

Assumption 6.4 (Machine Learning Bias II) *Let* $e_n = o(1)$ *. For all folds, the nuisance parameters obtained via cross-fitting belong to a shrinking neighborhood* \mathcal{T}_n *around η with probability of at least* $1 - e_n$ *, such that*

$$
\lambda_{s,n,1} + \lambda_{s,n,2} + \lambda_{q,n,1} + \lambda_{q,n,2} + \lambda_{m,n,2} + \lambda_{b,n,2} = o(1)
$$

and

$$
\sqrt{n}(\lambda_{s,n,2}^2 + \lambda_{q,n,2}^2 + \lambda_{m,n,2}^2 + \lambda_{b,n,2}^2) = o(1).
$$

We obtain the following Theorem.

Theorem 6.2 *Under Assumptions [3.1,](#page-9-3) [3.2,](#page-9-2) [5.2,](#page-21-3) [6.1,](#page-26-1) and [6.4,](#page-29-0) and h >* 0 *such that* $\sup_{z,j} g'_{j,h}(z) \lesssim 1$, the smooth bound estimator is asymptotically normal and semipara*metrically efficient, i.e.*

$$
\sqrt{n}(\hat{\beta}_{L,h}(1,1)-\beta_{L,h}(1,1)) \stackrel{d}{\rightarrow} \mathcal{N}\big(0,Var_{a}(\beta_{L,h}(1,1))\big),\,
$$

where $Var_a(\beta_{L,h})$ *is the semiparametric efficiency bound for* $\beta_{L,h}$ *.*

We can see that, without restricting the distribution of $s(1, x) - s(0, x)$, i.e. allowing for point mass $P(X^0) > 0$ or arbitrary density around zero, the smooth bounds are asymptotically normal under weaker assumptions than the non-smooth bounds with restricted distributions. In particular, they only require L_1 and L_2 consistency as well as root mean squared error rates for all nuisance quantities of order $o(n^{-1/4})$ as in standard debiased machine learning [\(Chernozhukov et al., 2018\)](#page-44-12). This is fundamentally due to the smoothing turning the target parameter into a pathwise differentiable object.

Our smooth estimator differs from [Semenova](#page-46-1) [\(2024\)](#page-46-1) in the following way: [Semenova](#page-46-1) [\(2024\)](#page-46-1) uses a moment shifting or shrinkage procedure based on the selection probabilities. These shifting methods are known to be quite sensitive to the choice of tuning/selection parameter [\(Andrews and Soares, 2010\)](#page-43-0). Moreover, even when $P(X^0) = 0$, the estimator by [Semenova](#page-46-1) [\(2024\)](#page-46-1) in finite samples should tend to narrow estimated identified sets as observations close to $p_0(x) = 1$ use a moment for the standard difference in conditional outcome means without truncation. Thus, while asymptotically valid due to correct classification in the limit, it will tend to be over-reject a correct null hypothesis in finite samples. The smoothed estimators, on the other hand, are constructed to be always wider than the estimators using no smoothing. They therefore estimate an outer identification region and the estimated identified will always be at least as wide as the naive generalized Lee bounds. This suggests better finite-sample size control at the expense of some power, see Appendix [C](#page-87-0) for Monte Carlo simulations.

6.5. The Efficiency Gap and Known Propensity Scores

If propensity scores are known, the moment functions from [Semenova](#page-46-1) [\(2024\)](#page-46-1) or [Heiler](#page-45-1) [\(2024\)](#page-45-1) without correction can be used as well. The corresponding estimator [\(6.10\)](#page-25-0) does not require estimation of $\beta_{j,d}(u, X)$ for any $j, d \in \{0, 1\}$ and thus all $\lambda_{b,n,p}$ terms that enter Assumption [6.3](#page-28-1) and [6.4](#page-29-0) can be omitted. However, there is an information loss from ignoring the conditional variation in the truncated means. In particular, we obtain the following Theorem.

Theorem [6.3](#page-28-1) *Under Assumptions 3.1, 3.2, 5.2, 6.1, 6.2, and 6.3 with* $\lambda_{b,n,p} = 0$ *known,* $\tilde{\beta}_L(1,1)$ *is asymptotically normal*

$$
\sqrt{n}(\tilde{\beta}_L(1,1)-\beta_L(1,1)) \stackrel{d}{\rightarrow} \mathcal{N}\big(0, Var_b(\beta_L(1,1))\big),\,
$$

with efficiency gap

$$
E[\min\{s(0, X), s(1, X)\}]^{2}(Var_{b}(\beta_{L}(1, 1)) - Var_{a}(\beta_{L}(1, 1)))
$$

=
$$
E\bigg[\mathbb{1}_{\{X \in \mathcal{X}^{+}\}}s(0, X)^{2}\bigg(\beta_{1,1}(X, p_{0}(X))\sqrt{\frac{1 - m(X)}{m(X)}} - \beta_{0,0}(X, 0)\sqrt{\frac{m(X)}{1 - m(X)}}\bigg)^{2}\bigg]
$$

+
$$
E\bigg[\mathbb{1}_{\{X \in \mathcal{X}^{-}\}}s(1, X)^{2}\bigg(\beta_{1,1}(X, 1)\sqrt{\frac{1 - m(X)}{m(X)}} - \beta_{0,0}(X, 1 - 1/p_{0}(X))\sqrt{\frac{m(X)}{1 - m(X)}}\bigg)^{2}\bigg].
$$

Theorem [6.3](#page-30-0) demonstrates that, under known propensity scores, there is a trade-off between efficiency and learning rates for the truncated conditional means $\beta_{j,d}(u, X)$. A lack of efficiency from not including information about the conditional truncated potential outcome means in the respective influence function arises as the variance of the truncated variables are bounded from below by their conditional analogues. This is equivalent to the comparison between the variance of a dependent variable and its residual variance in standard regression analysis. Theorem [6.3](#page-30-0) also nests the case without sample selection. In particular, when $S = 1$ almost surely, the ATE is point identified $\beta_L(1,1) = \beta_U(1,1)$ and all truncated means simplify to their untruncated counterparts $\beta_{d,1}(x,1) = \beta_{d,0}(x,0) = E[Y|D=d, X=x]$. In this case all correction terms vanish as well and the alternative estimator collapses to

$$
\tilde{\beta}_L(1,1) = E_n \left[\frac{YD}{m(X)} \right] - E_n \left[\frac{Y(1-D)}{1-m(X)} \right].
$$

This is the standard Horvitz-Thompson-type IPW estimator for the ATE using true propensities which is known to not reach the semiparametric efficiency bound [\(Hahn,](#page-44-4)

[1998\)](#page-44-4). More precisely, Theorem [6.3](#page-30-0) yields efficiency gap

$$
Var_b(\beta_L(1,1)) - Var_a(\beta_L(1,1)) = E\bigg[\bigg(\beta_{1,1}(X,1)\sqrt{\frac{1-m(X)}{m(X)}} - \beta_{0,0}(X,0)\sqrt{\frac{m(X)}{1-m(X)}}\bigg)^2\bigg].
$$

This comparison to the point-identified case suggests that, in the regular regime, one could potentially obtain an efficient estimator for the always-taker ATE bounds without modelling the conditional truncated quantiles analogously to [Hirano et al.](#page-45-12) [\(2003\)](#page-45-12) using nonparametrically estimated treatment propensities with sufficiently rich basis functions that asymptotically encode all information about the truncated conditional outcome means. We leave the development of such an alternative estimation approach for future research.

7. Extensions

7.1. Mean Dominance

The bounds in Proposition [4.1](#page-15-0) can further be refined using a mean dominance assumptions. Mean dominance to bound causal effects in sample selection models has been previously suggested by [Zhang and Rubin](#page-47-0) [\(2003\)](#page-47-0) for always-takers and by [Huber and](#page-45-2) [Mellace](#page-45-2) [\(2015\)](#page-45-2) for compliers under strong monotonicity without covariates.

Assumption 7.1 (Mean Dominance) *The conditional potential outcomes at the intensive margin are larger than at the extensive margin, i.e.*

$$
E[Y(d)|S(0) = 1, S(1) = 1, X = x] \ge E[Y(d)|S(0) \neq S(1), X = x]
$$
\n(7.1)

for any $d = 0, 1$.

This yields the following modified sharp bounds^{[7](#page-32-1)}.

Proposition 7.1 (Identification with Mean Dominance) *Let* Y*d*(*x*) *be the support of* $Y(d)$ *conditional on* $X = x$ *and* $\overline{y}_d(x)$ *and* $\underline{y}_d(x)$ *its respective infimum and supremum*

⁷If only either complier or defier bounds are of interest, Assumption [7.1](#page-32-2) can be relaxed to hold for the respective group only.

for $d \in \{0, 1\}$ *. Under Assumptions* [3.1,](#page-9-3) [3.2,](#page-9-2) [4.1,](#page-14-2) [4.2,](#page-14-3) and [7.1,](#page-32-2) the lower bounds for the *principal strata are given by*

$$
\beta_L(x, s_0, s_1) = \begin{cases}\n\beta_{1,1}(x, 1) - \beta_{0,0}(x, 1 - \min\{1/p_0(x), 1\}) & \text{if } s_0 = 1, s_1 = 1 \\
\beta_{1,1}(x, 1 - \min\{p_0(x), 1\}) - \beta_{0,0}(x, 0) & \text{if } s_0 \neq s_1 \\
\underline{y}_1(x) - \overline{y}_0(x) & \text{if } s_0 = 0, s_1 = 0,\n\end{cases}
$$

and the upper bounds by

$$
\beta_U(x, s_0, s_1) = \begin{cases} \beta_{0,1}(x, 1 - \min\{p_0(x), 1\}) - \beta_{1,0}(x, 1) & \text{if } s_0 = 1, s_1 = 1 \\ \beta_{0,1}(x, 0) - \beta_{1,0}(x, 1 - \min\{1/p_0(x), 1\}) & \text{if } s_0 \neq s_1 \\ \overline{y}_1(x) - \underline{y}_0(x) & \text{if } s_0 = 0, s_1 = 0. \end{cases}
$$

As the share of the principal strata are already identified via conditional monotonicity, aggregation to unconditional bounds uses the same conditional probability weights as in the case without mean dominance in [\(4.5\)](#page-16-0) and thus the same moment function for the denominator or density part in [\(6.3\)](#page-24-1). For the numerator, all truncated quantities can be estimated as in the pure monotonicity case with the components in the influence function corresponding to the now untruncated means replaced by their standard untruncated augmented IPW counterparts for the conditional outcome means [\(Robins et al., 1994\)](#page-46-14).

7.2. Heterogeneous Bounds

In many evaluation problems, objects of interest are heterogeneous effects, e.g. the effect at a particular margin for an observable subgroup. [Heiler](#page-45-1) [\(2024\)](#page-45-1) argues that, in the nonparametric sample selection model and more broadly, there are two types of heterogeneity, 1) heterogeneous effects and 2) heterogeneity in the severity of the identification problem, i.e. the width of the identified set. Exploiting these in combination can yield more precise inference and significant effect bounds even when unconditional aggregate bounds do not reject a null-effect. Our influence functions for any principal strata or margin can readily be used for the estimation of such heterogeneous effect bounds following the method of [Heiler](#page-45-1) [\(2024\)](#page-45-1). In particular, let $f: \mathcal{X} \to \mathcal{Z}$ and define $Z = f(X)$ a low-dimensional subgroup or mapping from the covariate space. Recall that all presented influence functions have the following linear structure

$$
\psi_{\beta_B}(W,\eta,\beta_B) = \psi_{\beta_B}^{[B]}(W,\eta) - \psi_{\beta_B}^{[S]}(W,\eta)\beta_B,
$$
\n(7.2)

where

$$
E[\psi_{\beta_B}^{[B]}(W,\eta)|X=x] = \beta_L(x,s_0,s_1)P(S(0) = s_0, S(1) = s_1|X=x),
$$

\n
$$
E[\psi_{\beta_B}^{[S]}(W,\eta)|X=x] = P(S(0) = s_0, S(1) = s_1|X=x).
$$
\n(7.3)

Thus, for any $z \in \mathcal{Z}$,

$$
E[Y(1) - Y(0)|S(0) = s_0, S(1) = s_1, Z = z]
$$

\n
$$
\geq E[\beta_L(X, s_0, s_1)|S(0) = s_0, S(1) = s_1, Z = z]
$$

\n
$$
= \frac{E[\beta_L(X, s_0, s_1)P(S(0) = s_0, S(1) = s_1|X)|Z = z]}{E[P(S(0) = s_0, S(1) = s_1|X)|Z = z]}
$$

\n
$$
= \frac{E[\psi_{\beta_B}^{[B]}(W, \eta)|Z = z]}{E[\psi_{\beta_B}^{[S]}(W, \eta)|Z = z]},
$$
\n(7.4)

by the law of iterated expectations. Hence we can obtain conditional effect bounds by aggregating the components of the presented influence functions within covariate partitions or by (nonparametric) projection onto (basis transformations of) *Z*. Regularity and/or the necessity for smoothing can be obtained under analogous conditions to the ones presented in this paper. For more details consider [Heiler](#page-45-1) [\(2024\)](#page-45-1).

8. Empirical Study: Labor Market Policy

8.1. Job Corps and Data

Job Corps (JC) is a US Department of Labor program aimed at empowering young individuals of economically or otherwise disadvantaged background with important labor market skills and qualifications. The intense program combines general education, vocational training, extracurricular activities, placement services, and more. Most participants are in residential slots in centers of varying size all over the US. The National Job Corps Study, executed by Mathematica Policy Research in the mid-90s, analyzed the effects of the Job Corps program on labor market prospects. The experiment implemented stratified randomized assignment of applicants, incorporating over 15400 units

between ages 16 to 24. Data on various outcomes such as income, job status, educational achievements, and criminal activities were gathered at different intervals. The outcomes of the original Job Corps study were mixed. Initially, the program boosted the participants' educational achievements and income. However, most aggregate effects waned over time. There is evidence for increased earnings in the older participant population [\(Schochet et al., 2008\)](#page-46-2). There is also well-document heterogeneity in terms of vocational and academic training returns as well a gender, see e.g. [Flores et al.](#page-44-14) [\(2012\)](#page-44-14) or [Heiler and](#page-45-13) [Knaus](#page-45-13) [\(2023\)](#page-45-13).

We employ the public use files which contain all survey data up to 208 weeks after the initial assignment [\(Burghardt et al., 1999;](#page-44-5) [Schochet et al., 2008\)](#page-46-2). Our main outcome measure is log hourly earnings as in [Lee](#page-46-0) [\(2009\)](#page-46-0). If workers are priced around their marginal productivity, this measure can be used as a proxy for increases in the latter caused by assignment to Job Corps. We extract an extended set of covariates containing detailed information regarding demographics, employment, criminal history, education, health, expectations, regional characteristics, and other JC related information.^{[8](#page-35-0)} All impact analysis is conditional on having a non-missing earnings entry, including zero, at weeks $t \in \{1, \ldots, 208\}$ identical to [Lee](#page-46-0) [\(2009\)](#page-46-0) leaving a sample of [9](#page-35-1).415 units.⁹

In the following, results for nuisance functions are presented for the research sample. For any of the impact analysis as well as general descriptive statistics, we use the nationally representative design weights as in [Schochet et al.](#page-46-2) [\(2008\)](#page-46-2). For the impact evaluation, we analyze the effect per eligible participant, i.e. the causal effect of assignment to Job Corps. We focus on the average treatment effect at the intensive margin as studied by [\(Lee, 2009;](#page-46-0) [Semenova, 2024\)](#page-46-1). As participation or outside alternatives after assignment are not controlled, all Job Corps effects here are relative to the control state of not being assigned, including partaking in any alternative program and/or working.

Figure [8.1](#page-36-0) contains the weekly average earnings and employment rate for the 208

⁸We collect all variables used in [Lee](#page-46-0) [\(2009\)](#page-46-0) and additional information on individual background and behavior similar to [Flores et al.](#page-44-14) [\(2012\)](#page-44-14) and [Heiler and Knaus](#page-45-13) [\(2023\)](#page-45-13), as well as all variables that were used for stratification in the original experiment. Some of the latter have previously been overlooked but are necessary for correct specification of the propensity score, see Appendix [B.1](#page-79-1) for more details.

 9 [Lee](#page-46-0) [\(2009\)](#page-46-0) suggests that non-response at this stage is a second order issue and initial treatment-control balance is preserved, see his Remark 2. We reassess this claim with the extended data using both formal tests for average differences as well as standardized differences and find little evidence for imbalance, see Appendix [B.1.](#page-79-1) The only exception is significantly larger worries to attend Job Corps in the treatment group. Our prior is that worries are more likely negatively related to any treatment effect. Thus, we expect any potential bias to be negative, i.e. earnings and employment effects are likely to be at least as large.

Weakly earnings, hours worked, and employment averages of control and treated units. The *n* = 9415 sample is restricted to units that report hours and earnings for all periods $t \in \{1, \ldots, 208\}$. All calculations use nationally representative design weights.

weeks after treatment assignment. There are systematic trends and differences between treatment and control groups in both earnings and employment over time. There is a clear upwards trend in earnings and employment over time. The control group tends to be ahead early after assignment but catches up later in time exceeding the trajectory of the control group for both measures. However, any differences between treatment and control earnings effectively consists of a mixture of units (the working) that could be made up of varying proportions of always-takers, compliers, and defiers, effectively leading to selected comparisons. Raw differences in employment rates are also uninformative about heterogeneity and composition of the intensive and extensive margins. We take this into account when constructing the bounds in what follows.

8.2. Models and Estimation

We give a brief overview over the estimation methods used. See Appendix [B.2](#page-80-0) for more details. Treatment propensities are known by the design of the stratified experiment, see Appendix [B.1.](#page-79-0) All other nuisance parameters are estimated using 5-fold cross-fitting with all available observations and pre-treatment predictors at each time period reported. No cross-time information was used.

For the conditional selection probabilities we compare five different main specifications: (a) Logistic regression, (b) logistic regression with interacted treatment, (c) gradient boosted trees (XGBoost), (d) artificial neural network, and (e) random forest. These methods differ with respect to their capabilities to (i) allow for weak monotonicity, (ii) produce sparsity, (iii) be robust to extreme observations, and (iv) fit and predict employment probabilities and status well. Table [8.1](#page-37-0) contains an aggregate comparison between the different methods with respect to these properties. By construction, all models except for the simple logit allow for weak monotonicity. XGBoost, neural network, and random forest can produce sparsity.^{[10](#page-37-1)} XGBoost is the model that has, by far, the best out-of-sample accuracy and loss as measured by the negative likelihood function outperforming *all* other methods throughout 202/206 out of 208 periods, see Appendix [B.2](#page-80-0) for the detailed results. This is in line with many studies that demonstrate boosted trees as or among the best performing methods for tabular data [\(Shwartz-Ziv and Armon, 2022\)](#page-46-0).

Method	Weak Monotonicity	Sparsity	Extreme Predictions	Accuracy (rank)	Likelihood (rank)
Logit					
Logit Interacted					Ð
XGBoost					
Neural Network					
Random Forest					

Table 8.1: Selection Model Estimates: Comparison

This table contains the different estimation methods, their properties as well as an overview over the resulting $p_0(x)$ estimates. Weak Monotonicity refers to Assumption [3.2.](#page-9-0) Logit has that either $P(\mathcal{X}^+)$ or $P(X^{-})$ equal zero. Sparsity is achieved whenever $P(X^{0})$ can be larger than zero. Extreme Predictions refers to an average of at least 10 selection estimates outside of the (0*.*25*,* 4) interval. Accuracy and Likelihood contain the rank in the respective performance metric based on the cross-fitting error averaged over all periods $t \in \{1, \ldots, 208\}.$

For conditional means, conditional quantiles, and truncated means required for the bound estimators, we use off-the-shelf honest regression and quantile forests across all specifications [\(Athey et al., 2019\)](#page-44-0). This was chosen to isolate the contribution of differ-ences in relative selection probabilities which is the main issue addressed in this paper.^{[11](#page-37-2)}

For the impact analysis, we consider the bounds that (i) remove observations with $p_0(x) = 1$ and use the corresponding influence function ("Trim"), (ii) the moment shifting or selection method by [Semenova](#page-46-1) [\(2024\)](#page-46-1) ("Shift"), (iii) smooth bounds ("Smooth"), and (iv) conventional Lee bounds using strong monotonicity. For the shifting method, we

¹⁰Both the interacted logit and neural network tend to produce many or some relatively extreme estimates for the relative selection probabilities, see also Figure [B.2](#page-83-0) in Appendix [B.3.](#page-83-1) It seems questionable whether e.g. extreme relative selection probabilities that suggest e.g. a quadrupling in relative employment chances due to Job Corps are credible or mostly a by-product of an overly high variance model.

¹¹Quantile forests are also computationally attractive in high dimensions and have been used for sample selection bounds in previous research [\(Heiler, 2024;](#page-45-0) [Andersen et al., 2023\)](#page-43-0).

(a) contains the share of units for which each method returns relative selection probabilities $p_0(x)$ estimated within a 0.01 neighborhood of one at each post-assignment period $t = 1, \ldots, 208$. (b) contains the results for XGBoost with relative selection probabilities $p_0(x)$ estimated exactly equal or within the 0.01 neighborhood of one respectively.

use the tuning parameter as suggested by [Semenova](#page-46-1) [\(2024\)](#page-46-1), $\rho_n = n^{-1/4}/\log n$. For both trimming and shifting, we report results with and without using the propensity score correction that affects efficiency. For the smooth bounds, we report the analysis on a grid of smoothing parameters.[12](#page-38-0) For inference on the effect we use [Imbens and Manski](#page-45-1) [\(2004\)](#page-45-1) confidence intervals for all methods.

8.3. Relative Selection Probabilities $p_0(x)$

In this section, we consider the predicted relative selection probabilities close or equal to one for the methods that can produce sparsity. Figure [8.2a](#page-38-1) depicts the share of probabilities for XGBoost, neural network, and random forest. The average amount of close-to-one units averaged over all time periods are 51*.*91% (XGBoost), 18*.*75% (Neural Net), and 42*.*36% (Random Forest). Thus, there is a significant amount of units that can affect the various bounds differently. In particular, for the XGBoost, almost all these probabilities are exact numerical ones, i.e. the underlying model is sparse in some parts of the covariate space with respect to the treatment. Figure [8.2b](#page-38-1) depicts the close-toone and exact-one unit shares over all time periods. One can see that the trajectory of exact-one closely follows the close-to-one shares with an average rate of 44.36%.

¹²On any fixed grid, inference is valid for any *h*, so one could pick the *h* that provides the smallest confidence intervals to optimize power.

8.4. Employment Effects

Figure [8.3](#page-39-0) contains the estimated average assignment effect on employment using the efficient estimator for all probability models. Estimates are almost identical. We replicate the finding that, in the short run, assignment reduces the likelihood of employment which only bounces back after about 1.5 years (the average duration of JC participation was around 8 months at the time). After about 2.25 years, positive employment effects tend to stabilize. In particular, most estimates are significantly negative for weeks 2 to 76 and significantly positive for weeks 116 to 208 ($p < 0.05$).

Figure 8.3: Job Corps: Average Employment Effect Estimates

Estimated average treatment effect of assignment to Job Corps on employment *E*[*S*(1) − *S*(0)] using the efficient influence function with different selection probability models for $t \in \{1, \ldots, 208\}$. All calculations use nationally representative design weights.

8.5. Causal Effects Bounds

We now present the impact results for assignment to JC on log hourly earnings. We first present the results for the main evaluation in $t = 208$ using the new methods and contrast them with results based on the literature and replications with comparable sample and design weights. We then present the results using the best performing nuisance specification for methods that allow for conditional monotonicity for periods $t \in \{45, 90, 135, 180, 208\}$ as in [Lee](#page-46-2) [\(2009\)](#page-46-2). Robustness checks and results for additional methods, time periods, and smoothing parameters can be found in Appendix [B.3.1.](#page-84-0)

8.5.1. Main Effects at *t* = 208

Table [8.3](#page-42-0) contains the impact analysis for various methods and specifications at period $t = 208$. All entries except for Lee (2009) are based on own calculations and with identical outcome and design weights. [Schochet et al.](#page-46-3) [\(2008\)](#page-46-3) is a simple treatment control difference for the sample of reported hourly log earnings. It is of similar magnitude when compared with the equivalent mean difference in the restricted Lee (2009) sample in line with the balancing analysis.

Switch, regular, and smooth bounds all admit flexible estimation of nuisance parameters. Using the best performing XGBoost model for selection, there are 4252 observations with exact $\hat{p}_0(x) = 1$. Thus, trimming bounds effectively have to use a much reduced sample. Nevertheless, estimates are in a similar range but with larger standard errors. The moment switching approach by [Semenova](#page-46-1) [\(2024\)](#page-46-1) with known propensity scores provides overly wide bounds. This seems to be due to a strong sensitivity when there are many estimates for $p_0(x)$ around one. With unknown scores, estimates are tighter, however with standard errors much larger compared to alternatives.

Our preferred specification is smooth bounds with $h = 1$. The impact estimate of [−0*.*005*,* 0*.*113] with 95% confidence interval of (−0*.*040*,* 0*.*148) rules out large negative earnings effects and is surprisingly close and of similar width compared to results obtained under strong monotonicity as in [Lee](#page-46-2) [\(2009\)](#page-46-2). As strong monotonicity is rejected by the data [\(Semenova, 2024\)](#page-46-1), this provides more evidence on the robustness of this final impact evaluation. We also include a larger smoothing specification for comparison; for more values see Appendix [B.3.1.](#page-84-0) As expected, the identified set becomes larger. In this case, the standard errors are still comparable such that overall less smoothing seems preferable.

8.5.2. Secondary Effects and Robustness

We now analyze the effect for multiple periods using all methods. We present the results for the two best nuisance models (XGBoost and neural network). For the switching and the trimming bounds, we only consider the unknown propensity score estimates, as they are more reliable across specifications. For smooth bounds we present two smoothing parameter results as above. Results for the other methods and parameters are robust with the exception of the interacted logistic model, that failed to converge in some folds.^{[13](#page-40-0)}

¹³See Appendix [B.3](#page-83-1) for some histograms and descriptive statistics.

Simple Monotonicity Bounds										
		$t=45$	$t=90$	$t=135$	$t=180$	$t=208$				
Lee (2009)		$(-0.097, 0.152)$			$[-0.074, 0.125]$ $[0.042, 0.043]$ $[-0.016, 0.076]$ $[-0.033, 0.087]$ $[-0.012, 0.089]$ $(-0.005, 0.092)$ $(-0.051, 0.099)$ $(-0.064, 0.108)$ $(-0.037, 0.112)$					
Results for XGBoost Selection Model										
		$t=45$	$t=90$	$t=135$	$t=180$	$t = 208$				
Smooth	$h=5$	$[-0.109, 0.145]$ $(-0.172, 0.199)$	$[-0.112, 0.212]$ $(-0.161, 0.256)$	$[-0.093, 0.213]$ $(-0.139, 0.250)$	$[-0.072, 0.215]$ $(-0.107, 0.251)$	$[-0.057, 0.275]$ $(-0.095, 0.312)$				
	$h=1$	$[-0.000, 0.161]$ $(-0.068, 0.209)$	[0.008, 0.096] $(-0.036, 0.135)$	$[-0.028, 0.090]$ $(-0.066, 0.125)$	$[-0.029, 0.093]$ $(-0.064, 0.128)$	$[-0.005, 0.113]$ $(-0.040, 0.148)$				
Trim		$[-0.257, 0.126]$ $(-0.403, 0.269)$	$[-0.175, 0.004]$ $(-0.745, 0.573)$	$[-0.021, 0.152]$ $(-0.197, 0.327)$	$[-0.180, -0.011]$ $(-0.328, 0.134)$	$[-0.036, 0.077]$ $(-0.201, 0.245)$				
Shift		$[-0.324, -0.076]$ $(-0.470, 0.067)$	$[-0.021, -0.011]$ $(-0.194, 0.162)$		$[0.022, 0.067]$ $[0.075, 0.122]$ $[0.138, 0.162]$ $(-0.116, 0.204)$ $(-0.055, 0.253)$ $(-0.003, 0.303)$					
$ \hat{p}_0(x) - 1 \leq \frac{n^{-1/4}}{\log(n)}$		2592	8680	6350	4871	5688				
Results for Neural Network Selection Model										
		$t=45$	$t=90$	$t=135$	$t=180$	$t=208$				
Smooth	$h=5$	$[-0.177, 0.089]$ $(-0.234, 0.137)$	$(-0.183, 0.255)$	$(-0.132, 0.220)$	$[-0.137, 0.214]$ $[-0.089, 0.187]$ $[-0.092, 0.182]$ $[-0.058, 0.228]$ $(-0.125, 0.215)$	$(-0.094, 0.263)$				
	$h=1$	$[-0.469, 0.777]$ $(-0.543, 0.855)$	$[-0.005, 0.105]$ $(-0.049, 0.147)$	$[-0.023, 0.107]$ $(-0.061, 0.144)$	$[-0.045, 0.112]$ $(-0.080, 0.149)$	$[-0.022, 0.127]$ $(-0.058, 0.165)$				

Table 8.2: Job Corps Impact Results, by week

Trim [−0*.*347*,* 0*.*192] [−0*.*031*,* 0*.*003] [−0*.*107*,* 0*.*046] [−0*.*141*,* 0*.*038] [−0*.*157*,* 0*.*072]

Shift [−0*.*424*,* −0*.*122] [−0*.*054*,* 0*.*011] [0*.*049*,* 0*.*108] [−0*.*062*,* 0*.*076] [0*.*046*,* 0*.*121]

(−0*.*481*,* 0*.*319) (−0*.*179*,* 0*.*149) (−0*.*223*,* 0*.*164) (−0*.*266*,* 0*.*159) (−0*.*284*,* 0*.*203)

using smoothed, trimmed, shifted, and regular (generalized) Lee bounds using *n* = 9415 for all weeks. Identified sets are in brackets and 95% [Imbens and Manski](#page-45-1) [\(2004\)](#page-45-1) confidence intervals in parenthesis below. The threshold $\frac{n^{-1/4}}{\log(n)}$ $\frac{n^{-1/4}}{\log(n)}$ is the switching threshold suggested by [Semenova](#page-46-1) [\(2024\)](#page-46-1) to choose between moments. Smoothing parameters are scaled by 100. All calculations use design weights.

Table 8.3: Impact Results at $t = 208$: Overview

Impact estimates using different methods and samples with key assumptions for validity of the impact estimate. All estimates and intervals except for Lee (2009) are based on own calculations using design weights. Confidence intervals for point-identified methods use standard critical values. Confidence intervals for bounds use the [Imbens](#page-45-2)and [Manski](#page-45-2) [\(2004\)](#page-45-2) refinement. All methods use known propensity scores of the stratified experiment if necessary. ¹Independence of potential selection from potential outcomes. ²28 covariates are used to generate a single predictive score. Bounds are calculated within 5 strata of this score. 3 Months employed is used as excluded variable in the selection model. ⁴Semiparametric model also includes transformations of the original variables, see Lee (2009). ⁵Best performing nuisance parameter specifications, see Appendix [B.2](#page-80-1) for details. ⁶Trim are the bounds that remove all units for which $\hat{p}_0(x) = 1$ with unknown/efficient and known/inefficient propensity score and related moment function. ⁷Smoothing parameters *^h* scaled by 100.

see Appendix [B.3.1.](#page-84-0)

The intervals for the smooth bounds at $h = 1$ are, again, surprisingly close to the original [Lee](#page-46-2) (2009) specification. Even a strictly positive identified set at $t = 90$ is recovered with XGBoost, albeit with larger width which seems more credible given the estimates of the surrounding periods. The Lee unconditional monotonicity can be rejected from the data [\(Semenova, 2024\)](#page-46-1). However, it seems that the general finding for the later periods, with log hourly earnings effects in the range of around −0*.*03 to 0*.*11, is robust to conditional monotonicity without parametric assumptions.

9. Conclusion

This paper demonstrates the importance of heterogeneity in selection behavior with respect to estimation and inference on causal effects at the intensive and extensive margins in selected samples. Allowing for different types of monotonicity crucially affects the (ir)regularity of sharp effect bounds and thus the ability to provide precise inference for the associated causal effects. This paper provides a solution in the form of outer identification regions that can be estimated efficiently from a semiparametric perspective. The approach can handle modern machine learning methods that are able to better approximate such important heterogeneity in sample selection. We discover an empirically relevant trade-off between identification strength versus precision that is likely to apply to a much larger set of models and parameters beyond the ones considered in this paper, as in [Lee and Weidner](#page-46-4) [\(2021\)](#page-46-4) and [Pakel and Weidner](#page-46-5) [\(2023\)](#page-46-5).

References

- Abadie, A. (2003). Semiparametric Instrumental Variable Estimation of Treatment Response Models. *Journal of Econometrics*, 113(2):231–263.
- Andersen, S. C., Beuchert, L., Heiler, P., and Nielsen, H. S. (2023). A guide to impact evaluation under sample selection and missing data: Teacher's aides and adolescent mental health. *arXiv preprint arXiv:2308.04963*.
- Andrews, D. W. and Soares, G. (2010). Inference for parameters defined by moment inequalities using generalized moment selection. *Econometrica*, 78(1):119–157.
- Angrist, J. D., Graddy, K., and Imbens, G. W. (2000). The interpretation of instrumental variables estimators in simultaneous equations models with an application to the demand for fish. *Review of Economic Studies*, 67(3):499–527.
- Angrist, J. D. and Imbens, G. W. (1995). Two-stage Least Squares Estimation of Average Causal Effects in Models with Variable Treatment Intensity. *Journal of the American Statistical Association*, 90(430):431–442.
- Athey, S., Tibshirani, J., and Wager, S. (2019). Generalized random forests. *Annals of Statistics*, 47(2):1148–1178.
- Balke, A. and Pearl, J. (1997). Bounds on treatment effects from studies with imperfect compliance. *Journal of the American Statistical Association*, 92(439):1171–1176.
- Bartalotti, O., Kédagni, D., and Possebom, V. (2023). Identifying marginal treatment effects in the presence of sample selection. *Journal of Econometrics*, 234(2):565–584.
- Bernhardt, R., Munro, D., and Wolcott, E. L. (2024). How does the dramatic rise of nonresponse in the current population survey impact labor market indicators? *Journal of Applied Econometrics*, 39(3):498–512.
- Bickel, P. J., Klaassen, C. A., Bickel, P. J., Ritov, Y., Klaassen, J., Wellner, J. A., and Ritov, Y. (1993). *Efficient and Adaptive Estimation for Semiparametric Models*. Johns Hopkins University Press Baltimore.
- Burghardt, J., McConnell, S., Meckstroth, A., Shochet, P., Johnson, T., and Homrighausen, J. (1999). National job corps study: Report on study implementation. *Mathematica Policy Research, Inc.*
- Chen, J. and Roth, J. (2024). Logs with zeros? some problems and solutions. *Quarterly Journal of Economics*, 139(2):891–936.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *Econometrics Journal*, 21(1):C1–C68.
- Chernozhukov, V., Escanciano, J. C., Ichimura, H., Newey, W. K., and Robins, J. M. (2022). Locally robust semiparametric estimation. *Econometrica*, 90(4):1501–1535.
- d'Haultfoeuille, X., Gaillac, C., and Maurel, A. (2024). Partially linear models under data combination. *The Review of Economic Studies*.
- Flores, C. A., Flores-Lagunes, A., Gonzalez, A., and Neumann, T. C. (2012). Estimating the effects of length of exposure to instruction in a training program: The case of job corps. *Review of Economics and Statistics*, 94(1):153–171.
- Frangakis, C. E. and Rubin, D. B. (2002). Principal stratification in causal inference. *Biometrics*, 58(1):21–29.
- Hahn, J. (1998). On the Role of the Propensity Score in Efficient Semiparametric Estimation of Average Treatment Effects. *Econometrica*, 66(2):315–331.
- Heckman, J. (1974). Shadow prices, market wages, and labor supply. *Econometrica*, 42(4):679–694.
- Heckman, J. J. (1979). Sample Selection Bias as a Specification Error. *Econometrica*, 47(1):153–161.
- Heckman, J. J. and Vytlacil, E. J. (2007). Econometric Evaluation of Social Programs, Part II: Using the Marginal Treatment Effect to Organize Alternative Econometric Estimators to Evaluate Social Programs, and to Forecast Their Effects in New Environments. *Handbook of Econometrics*, 6:4875–5143.
- Heiler, P. (2022). Efficient covariate balancing for the local average treatment effect. *Journal of Business & Economic Statistics*, 40(4):1569–1582.
- Heiler, P. (2024). Heterogeneous treatment effect bounds under sample selection with an application to the effects of social media on political polarization. *Journal of Econometrics*, 244(1).
- Heiler, P. and Kazak, E. (2021). Valid inference for treatment effect parameters under irregular identification and many extreme propensity scores. *Journal of Econometrics*, 222(2):1083–1108.
- Heiler, P. and Knaus, M. C. (2023). Effect or treatment heterogeneity? policy evaluation with aggregated and disaggregated treatments.
- Hines, O., Dukes, O., Diaz-Ordaz, K., and Vansteelandt, S. (2022). Demystifying statistical learning based on efficient influence functions. *American Statistician*, 76(3):292–304.
- Hirano, K., Imbens, G. W., and Ridder, G. (2003). Efficient Estimation of Average Treatment Effects Using the Estimated Propensity Score. *Econometrica*, 71(4):1161– 1189.
- Hirano, K. and Porter, J. R. (2012). Impossibility results for nondifferentiable functionals. *Econometrica*, 80(4):1769–1790.
- Honoré, B. E. and Hu, L. (2020). Selection without exclusion. *Econometrica*, 88(3):1007– 1029.
- Honoré, B. E. and Hu, L. (2022). Sample selection models without exclusion restrictions: Parameter heterogeneity and partial identification. *Journal of Econometrics*, page 105360.
- Horowitz, J. L. and Manski, C. F. (1995). Identification and robustness with contaminated and corrupted data. *Econometrica*, pages 281–302.
- Huber, M. and Mellace, G. (2015). Sharp bounds on causal effects under sample selection. *Oxford Bulletin of Economics and Statistics*, 77(1):129–151.
- Imai, K. (2008). Sharp bounds on the causal effects in randomized experiments with "truncation-by-death". *Statistics & Probability Letters*, 78(2):144–149.
- Imbens, G. W. and Angrist, J. D. (1994). Identification and Estimation of Local Average Treatment Effects. *Econometrica*, 62(2):467–475.
- Imbens, G. W. and Manski, C. F. (2004). Confidence intervals for partially identified parameters. *Econometrica*, 72(6):1845–1857.
- Kaido, H. and Santos, A. (2014). Asymptotically efficient estimation of models defined by convex moment inequalities. *Econometrica*, 82(1):387–413.
- Kennedy, E. H. (2023). Semiparametric doubly robust targeted double machine learning: a review.
- Khan, S. and Tamer, E. (2010). Irregular Identification, Support Conditions, and Inverse Weight Estimation. *Econometrica*, 78(6):2021–2042.
- Kline, P. and Walters, C. R. (2019). On Heckits, LATE, and Numerical Equivalence. *Econometrica*, 87(2):677–696.
- Kolesár, M. (2013). Estimation in an instrumental variables model with treatment effect heterogeneity. *Working paper*.
- Lee, D. S. (2009). Training, wages, and sample selection: Estimating sharp bounds on treatment effects. *Review of Economic Studies*, 76(3):1071–1102.
- Lee, S. and Weidner, M. (2021). Bounding treatment effects by pooling limited information across observations. *arXiv preprint arXiv:2111.05243*.
- Levis, A. W., Bonvini, M., Zeng, Z., Keele, L., and Kennedy, E. H. (2023). Covariateassisted bounds on causal effects with instrumental variables. *arXiv preprint arXiv:2301.12106*.
- Luedtke, A. R. and van der Laan, M. J. (2016). Statistical inference for the mean outcome under a possibly non-unique optimal treatment strategy. *Annals of Statistics*, 44(2):713 – 742.
- Molinari, F. (2020). Microeconometrics with partial identification. *Handbook of Econometrics*, 7:355–486.
- Okamoto, Y. (2023). Bounds on treatment effects under stochastic monotonicity assumption in sample selection models. *arXiv preprint arXiv:2311.00439*.
- Olma, T. (2021). Nonparametric estimation of truncated conditional expectation functions. *arXiv preprint arXiv:2109.06150*.
- Pakel, C. and Weidner, M. (2023). Bounds on average effects in discrete choice panel data models. *arXiv preprint arXiv:2309.09299*.
- Robins, J. M., Rotnitzky, A., and Zhao, L. P. (1994). Estimation of Regression Coefficients when Some Regressors are not Always Observed. *Journal of the American Statistical Association*, 89(427):846–866.
- Schochet, P. Z., Burghardt, J., and McConnell, S. (2008). Does job corps work? impact findings from the national job corps study. *American Economic Review*, 98(5):1864–86.
- Semenova, V. (2023). Debiased machine learning of set-identified linear models. *Journal of Econometrics*, 235(2):1725–1746.
- Semenova, V. (2024). Generalized Lee bounds. *Journal of Econometrics (forthcoming)*.
- Shwartz-Ziv, R. and Armon, A. (2022). Tabular data: Deep learning is not all you need. *Information Fusion*, 81:84–90.
- Słoczyński, T. (2020). When should we (not) interpret linear iv estimands as late? *arXiv preprint arXiv:2011.06695*.
- Staub, K. E. (2014). A causal interpretation of extensive and intensive margin in generalized tobit models. *Review of Economics and Statistics*, 96(2):371–375.
- Vytlacil, E. (2002). Independence, Monotonicity, and Latent Index Models: An Equivalence Result. *Econometrica*, 70(1):331–341.
- Zhang, J. L. and Rubin, D. B. (2003). Estimation of causal effects via principal stratification when some outcomes are truncated by "death". *Journal of Educational and Behavioral Statistics*, 28(4):353–368.

A. Proofs and Supplementary Material

A.1. Proof of Proposition [4.1](#page-15-0)

We present the derivations for the lower bound. First note that the only observed strata for which we have outcome information are the selected treated and control groups. Moreover, for any $x \in \mathcal{X}^+$

$$
E[Y|DS = 1, X = x]
$$

= $E[Y(1)|S(1) = 1, X = x]$
= $E[Y(1)|S(0) = 1, S(1) = 1, X = x]p_0(X) + E[Y(1)|S(0) = 0, S(1) = 1, X = x](1 - p_0(X))$

and analogously, for any $x \in \mathcal{X}^-$,

$$
E[Y|(1 - D)S = 1, X = x]
$$

= $E[Y(0)|S(0) = 1, X = x]$
= $E[Y(0)|S(0) = 1, S(1) = 1, X = x]1/p_0(X) + E[Y(0)|S(0) = 1, S(1) = 0, X = x](1 - 1/p_0(X)).$

Thus, by [Horowitz and Manski](#page-45-3) [\(1995\)](#page-45-3), Corollary 4.1, we obtain the following sharp lower bounds

$$
E[Y(1)|S(0) = 1, S(1) = 1, X = x]
$$

\n
$$
\ge E[Y|DS = 1, Y \le q_1(p_0(X), X), X = x] \text{ for any } x \in \mathcal{X}^+
$$

and

$$
-E[Y(0)|S(0) = 1, S(1) = 1, X = x]
$$

\n
$$
\ge -E[Y|(1 - D)S = 1, Y \le q_0(1 - 1/p_0(X), X), X = x] \text{ for any } x \in \mathcal{X}^-.
$$

Moreover, for the always-takers, the remaining components are point identified as

$$
E[Y(0)|S(0) = 1, S(1) = 1, X = x] = E[Y|(1 - D)S = 1, X = x]
$$
 for any $x \in \mathcal{X}^+$,

$$
E[Y(1)|S(0) = 1, S(1) = 1, X = x] = E[Y|DS = 1, X = x]
$$
 for any $x \in \mathcal{X}^-$.

On \mathcal{X}^0 , the same applies to $E[Y(d)|X=x]$, as $D \perp \!\!\!\perp S \mid X=x$ for all $x \in \mathcal{X}^0$. For compliers and defiers, we can use the same mixtures as above to obtain

$$
E[Y(1)|S(0) = 0, S(1) = 1, X = x]
$$

\n
$$
\geq E[Y|DS = 1, Y \leq q_1(1 - p_0(X), X), X = x] \text{ for any } x \in \mathcal{X}^+,
$$

\n
$$
-E[Y(0)|S(0) = 1, S(1) = 0, X = x]
$$

\n
$$
\geq -E[Y|(1 - D)S = 1, Y \leq q_0(1/p_0(X), X), X = x] \text{ for any } x \in \mathcal{X}^-.
$$

For never-takers, no observed stratum is informative and thus, the lower bound is only informed by the support

$$
E[Y(1)|S(0) = 0, S(1) = 0, X = x] \ge \underline{y}_1(x),
$$

-
$$
E[Y(0)|S(0) = 0, S(1) = 0, X = x] \ge -\overline{y}_0(x).
$$

This completes all lower bounds. For the upper bounds, the corresponding expression can be obtained by using the complementary trimming thresholds and replacing the upper (lower) support bounds by their respective counterparts.

A.2. Proof of Proposition [7.1](#page-32-0)

Proposition [7.1](#page-32-0) follows directly from the previous proof by plugging in Assumption [7.1](#page-32-1) into the corresponding expressions.

A.3. Example for Smooth Bounds

Let $g(z) = h \log(1+e^{z/h})$ be a log-sum-exponential function, where $h > 0$ is a smoothing parameter. For each $h > 0$ and $z \in \mathbb{R}$, we have

$$
\max\{z, 0\} < g(z) \le \max\{z, 0\} + h\log(2),\tag{A.1}
$$

where the equality is attained if and only if $z = 0$. Based on this LogSumExp function *g*, we introduce the following four smoothing functions.

$$
g_{1,h}(z) = 1 - g(1 - z),
$$

\n
$$
g_{2,h}(z) = g(z),
$$

\n
$$
g_{3,h}(z) = 1 - g(1 - z) + h \log(2),
$$

\n
$$
g_{4,h}(z) = g(z) - h \log(2).
$$

Relying on inequalities in [\(A.1\)](#page-49-0), it is straightforward to show that

$$
g_{1,h}(z) < \min\{z, 1\} \le g_{3,h}(z),
$$

 $g_{4,h}(z) \le \max\{z, 0\} < g_{2,h}(z).$

Further, for each $i \in \{1, \ldots, 4\}$, and $h > 0$, we also have that

$$
\sup_{z \in \mathbb{R}} |g_{i,h}(z) - g_i(z)| \le h \log(2),\tag{A.2}
$$

where $g_1(z) = g_3(z) = \min\{z, 1\}$ and $g_2(z) = g_4(z) = \max\{z, 0\}.$

A.4. Proof of Theorem [4.1](#page-20-0)

We only prove this result for $\beta_{L,h}(1,1)$ since other case are analogous. We introduce the following intermediate Lee lower bound.

$$
\tilde{\beta}_{L,h} = \frac{E[\beta_{L,h}(X)\min\{s(0,X), s(1,X)\}]}{E[\min\{s(0,X), s(1,X)\}]}.
$$
\n(A.3)

It suffices to prove that

$$
\sup_{P \in \mathcal{P}} |\tilde{\beta}_{L,h} - \beta_L(1,1)| \lesssim h,\tag{A.4}
$$

and

$$
\sup_{P \in \mathcal{P}} |\beta_{L,h}(1,1) - \tilde{\beta}_L^h| \lesssim h. \tag{A.5}
$$

First, we deal with the convergence in [\(A.4\)](#page-50-0) and write this term as

$$
\tilde{\beta}_L^h - \beta_L(1,1) = \frac{E\left[\left(\beta_{L,h}(X,1,1) - \beta_L(X,1,1) \right) \min\{ s(0,X), s(1,X) \} \right]}{E[\min\{ s(0,X), s(1,X) \}]}.
$$
\n(A.6)

Next, we note that

$$
\beta_{L,h}(X,1,1) - \beta_L(X,1,1) = R_1^h(x) - R_2^h(x),\tag{A.7}
$$

where

$$
R_1^h(x) = \frac{1}{g_{1,h}(p_0(x))} \int_{\underline{y}_1(x)}^{q_1(g_{1,h}(p_0(x)),x)} y f_1(y|x) dy - \frac{1}{\min\{p_0(x), 1\}} \int_{\underline{y}_1(x)}^{q_1(\min\{p_0(x), 1\},x)} y f_1(y|x) dy,
$$

\n
$$
R_2^h(x) = \frac{1}{g_{1,h}(1/p_0(x))} \int_{q_0(1-g_{1,h}(1/p_0(x)),x)}^{y_0(x)} y f_0(y|x) dy
$$

\n
$$
- \frac{1}{\min\{1/p_0(x), 1\}} \int_{q_0(1-\min\{1/p_0(x), 1\},x)}^{y_0(x)} y f_0(y|x) dy.
$$

We further decompose the $R_1^h(x)$ term as

$$
R_1^h(x) = \left(\frac{1}{g_{1,h}(p_0(x))} - \frac{1}{\min\{p_0(x), 1\}}\right) \int_{\underline{y}_1(x)}^{q_1(g_{1,h}(p_0(x)), x)} y f_1(y|x) dy
$$

$$
- \frac{1}{\min\{p_0(x), 1\}} \int_{q_1(g_{1,h}(p_0(x)), x)}^{q_1(\min\{p_0(x), 1\}, x)} y f_1(y|x) dy.
$$

Relying on Assumption [4.3](#page-20-1) and the inequality $g_{1,h}(z) \le \min\{z, 1\}$, we obtain that

$$
|R_1^h(x)| \lesssim \frac{1}{\underline{g}_n^2} \sup_{x \in \mathcal{X}} |g_{1,h}(p_0(x)) - \min\{p_0(x), 1\}|
$$
 (A.8)

Similarly, we obtain that

$$
|R_2^h(x)| \lesssim \frac{1}{\underline{g}_n^2} \sup_{x \in \mathcal{X}} |g_{1,h}(1/p_0(x)) - \min\{1/p_0(x), 1\}|. \tag{A.9}
$$

Now, using Assumption [4.4](#page-20-2) and recalling the expression in [\(A.6\)](#page-50-1), we deduce that

$$
\sup_{P \in \mathcal{P}} |\tilde{\beta}_{L,h} - \beta_L(1,1)| \lesssim \frac{h}{\underline{g}_n^2}.
$$
\n(A.10)

Next, we turn to the term $\beta_{L,h}(1,1) - \tilde{\beta}_{L,h}$ and observe that

$$
\beta_{L,h}(1,1) - \tilde{\beta}_{L,h} = \left(\frac{A_h}{C_h} + \frac{B_h}{D_h}\right) - \left(\frac{\tilde{A}_h + \tilde{B}_h}{D}\right),\,
$$

where

$$
A_h = E[g_{4,h}(\beta_{L,h}(X,1,1))g_{1,h}(p_0(X))s(1,X)], \quad C_h = E[g_{3,h}(p_0(X))s(1,X)],
$$

\n
$$
B_h = E[g_{4,h}(\beta_{L,h}(X,1,1))g_{3,h}(p_0(X))s(1,X)], \quad D_h = E[g_{1,h}(p_0(X))s(1,X)],
$$

\n
$$
\tilde{A}_h = E[\max{\beta_{L,h}(X,1,1),0}\min{\{p_0(X),1\}}s(1,X)], \quad C = E[\min{\{s(0,X),s(1,X)\}}].
$$

\n
$$
\tilde{B}_h = E[\min{\beta_{L,h}(X,1,1),0}\min{\{p_0(X),1\}}s(1,X)].
$$

In view of Assumptions [4.3](#page-20-1) and [4.3,](#page-20-1) we note that

$$
|A_h - \tilde{A}_h| + |C_h - C| + |B_h - \tilde{B}_h| + |D_h - C| \lesssim \sum_{i=1}^5 \sup_{z \in \mathbb{R}} (|g_{i,h}(z) - g_i(z)|).
$$
 (A.11)

Further, from $C > 0$ and the assumption $|g_{1,h}(z) - \min\{z, 1\}| \lesssim h$, we deduce that D_h is bounded away from zero for all small h , which implies the same for C_h . Consequently, we obtain that

$$
|\beta_{L,h}(1,1) - \tilde{\beta}_{L,h}| \lesssim |A_h - \tilde{A}_h| + |C_h - C| + |B_h - \tilde{B}_h| + |D_h - C| \lesssim h. \tag{A.12}
$$

We finish the proof by adding both bounds.

A.5. Proof of Theorem [5.1](#page-22-0)

We only prove this theorem for always-takers/intensive margin (lower and upper bounds), extensive margin (lower bound), and compliers (lower bound). The proofs of other cases are analogous.

A.5.1. Notations and Auxiliary Results

In this part, we introduce some notations and preliminary results necessary for the proof of Theorem [5.1.](#page-22-0)

We are interested in the estimands $\beta_L(1,1)$, $\beta_U(1,1)$, $\beta_L(0,1)$ and $\beta_L(em)$. We note that all these estimands depend on $f_d(y|x)$, $s_d(x)$ for $d \in \{0,1\}$ and $f(x)$, where $f_d(y|x)$ is the conditional density of *Y* given $S = 1, D = d, X = x$.

In semiparametric efficiency proofs, we consider a submodel. Letting $\varepsilon = \theta - \theta_0$, we consider a parametric submodel for $d \in \{0, 1\}$.

$$
f_d(y|x,\theta) = (1 + \varepsilon \tau_d(y|x,\theta_0)) f_d(y|x), \quad s(d,x,\theta) = s(d,x) + \varepsilon s'(d,x,\theta_0),
$$

$$
f(x,\theta) = (1 + \varepsilon t(x)) f(x),
$$

which are equal to $f_d(y|x)$, $s_d(x)$ and $f(x)$ when $\theta = \theta_0$, and where

$$
\tau_d(y|x,\theta) = \frac{\partial \log f_d(y|x,\theta)}{\partial \theta}, \quad s'(d,x,\theta) = \frac{\partial s(d,x,\theta)}{\partial \theta},
$$

$$
t(x,\theta) = \frac{\partial \log f(x,\theta)}{\partial \theta},
$$

such that for $d \in \{0, 1\}$

$$
\int \tau_d(y|x,\theta_0) f_d(y|x) dy = 0 \text{ for all } x, \quad |\tau_d(y|x,\theta_0)| < \infty \text{ and } |s'(d,x,\theta_0)| < \infty \text{ for all } x, y,
$$

$$
\int t(x,\theta_0) f(x) dx = 0, \quad |t(x,\theta_0)| < \infty \text{ for all } x.
$$

In the following result, we use the decompositions $\beta_L(x, 1, 1) = \beta_{L,1}(x, 1, 1) - \beta_{L,0}(x, 1, 1)$, $\beta_U(x, 1, 1) = \beta_{U,1}(x, 1, 1) - \beta_{U,0}(x, 1, 1), \ \beta_L(x, 0, 1) = \beta_{L,1}(x, 0, 1) - \beta_{L,0}(x, 0, 1),$ which are derived from the components in Proposition [4.1.](#page-15-0) In addition, we have from Example [4.1.3](#page-17-0) that $\beta_L(x, em) = \beta_{L,1}(x, 0, 1)$.

Lemma A.1 *Suppose that the assumptions of Theorem [5.1](#page-22-0) hold and that* $P(X^0) = 0$. Then, for each $x \in \mathcal{X}$, $\beta_L(x,1,1)$, $\beta_U(x,1,1)$, $\beta_L(x,0,1)$, and $\beta_L(x,em)$ are pathwise *differentiable. In particular, we have that*

$$
\frac{\partial \beta_{L,1}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{\min\{p_{0}(x),1\}} \int_{y_{1}}^{q_{1}(\min\{p_{0}(x),1),x)} y_{T1}(y|x,\theta_{0}) f_{1}(y|x)dy \n- 1(x \in \mathcal{X}^{+}) \frac{q_{1}(p_{0}(x),x)}{p_{0}(x)} \int_{y_{1}}^{q_{1}(p_{0}(x),x)} \tau_{1}(y|x,\theta_{0}) f_{1}(y|x)dy \n+ 1(x \in \mathcal{X}^{+}) \frac{(q_{1}(p_{0}(x),x) - \beta_{L,1}(x,1,1))}{p_{0}(x)} \frac{\partial \theta}{\partial \theta},
$$
\n
$$
\frac{\partial \beta_{L,0}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{\min\{1/p_{0}(x),1\}} \int_{q_{0}(1-\min\{1/p_{0}(x),1\},x)}^{q_{0}} y_{T0}(y|x,\theta_{0}) f_{0}(y|x)dy \n- 1(x \in \mathcal{X}^{-}) p_{0}(x) q_{0}(1 - 1/p_{0}(x),x) \int_{q_{0}(1-1/p_{0}(x),x)}^{q_{0}(y,x,\theta_{0})} \tau_{0}(y|x,\theta_{0}) f_{0}(y|x)dy \n- 1(x \in \mathcal{X}^{-}) \frac{(q_{0}(1 - 1/p_{0}(x),x) - \beta_{L,0}(x,1,1))}{p_{0}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta},
$$
\n
$$
\frac{\partial \beta_{U,1}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{\min\{p_{0}(x),1\}} \int_{q_{1}(1-\min\{p_{0}(x),1\},x)}^{q_{1}(y|x,\theta_{0}) f_{1}(y|x)dy} + 1(x \in \mathcal{X}^{+}) \frac{(q_{1}(1 - p_{0}(x),x) - \beta_{U,1}(x,1,1))}{p_{0}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta},
$$
\n
$$
\frac{\partial \beta_{U,0}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{\min\{1/p_{0}(x),1\}} \int_{q_{0}}^{q_{0}(\min\{1/p
$$

Proof. We observe that $\beta_L(x, 1, 1)$, $\beta_U(x, 1, 1)$ and $\beta_L(x, 0, 1)$ have components of form $\beta_{1,1}(x, h(x)), \ \beta_{0,0}(x, h(x)), \ \beta_{0,1}(x, h(x))$ and $\beta_{1,0}(x, h(x))$ for different choices of $h(x)$.

 $-1(x \in \mathcal{X}^-) \frac{(q_0(1/p_0(x), x) - \beta_{L,0}(x, 0, 1))}{(1-x)}$

 $p_0(x)(1 - p_0(x))$

 $\frac{\partial p_0(x,\theta_0)}{\partial \theta}$.

Below, we spell out the details for the $\beta_L(x, 1, 1)$ case which is based on $\beta_{1,1}(x, h(x))$ and $\beta_{0,0}(x, h(x))$ with $h(x) = \min\{p_0(x), 1\}$ and $h(x) = 1 - \min\{1/p_0(x), 1\}$, respectively. Let $\beta(x) = \beta_L(x, 1, 1)$ *.* In the decomposition $\beta(x, \theta) = \beta_1(x, \theta) - \beta_0(x, \theta)$, we note that

$$
\beta_1(x,\theta) = \frac{1}{\min\{p(x,\theta),1\}} \int_{\underline{y}_1}^{q_1(\min\{p(x,\theta),1\},x)} y f_1(y|x,\theta) dy.
$$

This term hints to the use of product and Leibniz integral rules for differentiation. Since it contains minimum function, we cannot take derivatives directly. Instead, we carefully mimic these rules via decompositions. Letting $m1(x, \theta) = \min\{p_0(x, \theta), 1\}$, we have that

$$
\begin{aligned} &\beta_1(x,\theta)=\beta_1(x)+\frac{\varepsilon}{m1(x)}\int_{\underline{y}_1}^{q_1(m1(x,\theta),x)}y\tau_1(y|x,\theta_0)f_1(y|x)dy\\ &+\left(\frac{1}{m1(x,\theta)}-\frac{1}{m1(x)}\right)\int_{\underline{y}_1}^{q_1(m1(x,\theta),x)}yf_1(y|x,\theta)dy+\frac{1}{m1(x)}\int_{q_1(m1(x),x)}^{q_1(m1(x,\theta),x)}yf_1(y|x))dy. \end{aligned}
$$

Regarding the last term, we further write this as

$$
\int_{q_1(m1(x,\theta),x)}^{q_1(m1(x,\theta),x)} y f_1(y|x) dy = q_1(m1(x),x) \int_{\underline{y}_1}^{q_1(m1(x,\theta),x)} (f_1(y|x) - f_1(y|x,\theta)) dy
$$

+ $q_1(m1(x),x) \int_{\underline{y}_1}^{q_1(m1(x,\theta),x)} f_1(y|x,\theta) dy$
- $q_1(m1(x),x) \int_{\underline{y}_1}^{q_1(m1(x),x)} f_1(y|x)) dy$
+ $\int_{q_1(m1(x),x)}^{q_1(m1(x),x)} (y - q_1(m1(x),x)) f_1(y|x)) dy$.

Denoting $\Delta(x, \theta) = s(1, x, \theta) - s(0, x, \theta)$, we also observe that

$$
m1(x, \theta) - m1(x) = (p_0(x, \theta) - p_0(x))1(\Delta(x) > 0)
$$

$$
+ (1 - p_0(x, \theta))(1(\Delta(x, \theta) < 0) - 1(\Delta(x) < 0)).
$$

Combining these identities, we obtain that $\beta_1(x,\theta) = \hat{\beta}_1(x,\theta) + \bar{\beta}_1(x,\theta)$, where

$$
\hat{\beta}_{1}(x,\theta) = \beta_{1}(x) + \frac{\varepsilon}{m_{1}(x)} \int_{\underline{y}_{1}}^{q_{1}(m_{1}(x,\theta),x)} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x))dy \n- \frac{\varepsilon q_{1}(m_{1}(x),x)}{m_{1}(x)} \int_{\underline{y}_{1}}^{q_{1}(m_{1}(x,\theta),x)} \tau_{1}(y|x,\theta_{0})f_{1}(y|x)dy \n+ \mathbb{1}(\Delta(x) > 0) \frac{q_{1}(m_{1}(x),x) - \beta_{1}(x,\theta)}{p_{0}(x)} [p_{0}(x,\theta) - p_{0}(x)], \n\tilde{\beta}_{1}(x,\theta) = (\mathbb{1}(\Delta(x,\theta) < 0) - \mathbb{1}(\Delta(x) < 0)) \frac{q_{1}(m_{1}(x),x) - \beta_{1}(x,\theta)}{m_{1}(x)} [1 - p_{0}(x,\theta)] \n+ \frac{1}{m_{1}(x)} \int_{q_{1}(m_{1}(x),x)}^{q_{1}(m_{1}(x),x)} (y - q_{1}(m_{1}(x),x))f_{1}(y|x))dy.
$$

Here $\hat{\beta}_1(x,\theta)$ is the main term and now it is easy to derive that $\varepsilon(\hat{\beta}_1(x,\theta) - \beta_1(x))$ converges to the claimed limiting term by noting that

$$
\int_{\underline{y}_1}^{q_1(m1(x),x)} \tau_1(y|x,\theta_0) f_1(y|x) dy = 0 \text{ if } x \in \mathcal{X}^-.
$$
\n(A.13)

Further, $\bar{\beta}_1(x,\theta)$ is the remainder term. If $x \in \mathcal{X}^-$, we have that $\varepsilon^{-1} \bar{\beta}_1(x,\theta) = 0$ for all small $|\varepsilon|$. The reason is that $\Delta(x, \theta) < 0$ and $p(x, \theta) > 1$.

If $x \in \mathcal{X}^+$, we have that $|\varepsilon^{-1}\overline{\beta}_1(x,)| \leq C|\varepsilon|$ for all small $|\varepsilon|$ and a constant $C > 0$ in view of Assumption [5.1.](#page-21-0) The reason is that

$$
|q_1(m1(x,\theta),x) - q_1(m1(x),x)| \leq c_1^{-1} |p_0(x,\theta) - p_0(x) \leq C_1 |\varepsilon|
$$

for positive constants c_1, c_2 .

On the other hand, we can do a similar decomposition for the term $\beta_0(x)$. To be precise, we have that $\beta_0(x,\theta) = \hat{\beta}_0(x,\theta) + \bar{\beta}_0(x,\theta)$, where $m0(x,\theta) = \min\{1/p_0(x,\theta), 1\}$ and

$$
\hat{\beta}_0(x,\theta) = \beta_0(x) + \frac{\varepsilon}{m0(x)} \int_{q_0(1-m0(x,\theta),x)}^{y_0(1-m0(x,\theta),x)} y\tau_0(y|x,\theta_0) f_0(y|x)) dy \n- \frac{\varepsilon q_0(1-m0(x),x)}{m0(x)} \int_{q_0(1-m0(x,\theta),x)}^{y_0} \tau_0(y|x,\theta_0) f_0(y|x) dy \n- \mathbb{1}(\Delta(x) < 0) \frac{q_0(1-m0(x),x) - \beta_0(x,\theta)}{p_0(x,\theta)} [p_0(x,\theta) - p_0(x)], \n\bar{\beta}_0(x,\theta) = (\mathbb{1}(\Delta(x,\theta) > 0) - \mathbb{1}(\Delta(x) > 0)) \frac{q_0(1-m0(x),x) - \beta_0(x,\theta)}{m1(x)} [1 - 1/p_0(x,\theta)] \n- \frac{1}{m0(x)} \int_{q_0(1-m0(x),x)}^{q_0(1-m0(x,\theta),x)} (y - q_0(1-m0(x),x)) f_0(y|x)) dy.
$$

 $\text{Similarly, we obtain that } \varepsilon^{-1}\hat{\beta}_0(x,\theta) \text{ converges to the claim limiting term and } \varepsilon^{-1}\bar{\beta}_0(x,\theta)$ turns out to be negligible.

The results for other cases are analogous and therefore omitted.

A.5.2. Proof of Theorem [5.1](#page-22-0) - pathwise differentiability part

The proof is divided into several steps to improve readibility.

Step 0: Preliminaries and Decomposition

Under the assumption $P(X^0) = 0$, we prove that $\beta_B(s_0, s_1)$ is pathwise differentiable. For brevity, we will only derive this for $\beta_L(1,1)$ *,* $\beta_U(1,1)$ *,* $\beta_L(0,1)$ *,* and $\beta_L(em)$ *,* since proofs of other cases are analogous.

We consider a parametric submodel described in Section [A.5.1.](#page-52-0) We can represent each

 $\beta \in {\beta_L(1,1), \beta_U(1,1), \beta_L(0,1), \beta_L(em)}$ as $\beta(\theta) = N(\theta)/D(\theta)$, where its numerator and denominator are parameterized as

$$
N(\theta) = \int_{\mathcal{X}} \beta(x,\theta) g_{\beta}(x,\theta) f(x,\theta) dx, \quad D(\theta) = \int_{\mathcal{X}} g_{\beta}(x,\theta) f(x,\theta) dx.
$$
 (A.14)

The functions $\beta(x, \theta)$ and $g_{\beta}(x, \theta)$ for each case are found in Proposition [4.1,](#page-15-0) [\(4.5\)](#page-16-0) and Example [4.1.3.](#page-17-0) We note that

$$
\frac{\beta(\theta) - \beta(\theta_0)}{\varepsilon} = \frac{1}{D(\theta)} \left[\frac{N(\theta) - N(\theta_0)}{\varepsilon} - \beta \frac{D(\theta) - D(\theta_0)}{\varepsilon} \right].
$$
 (A.15)

Since $\lim_{\varepsilon\to 0} D(\theta) = D(\theta_0)$, it suffices to prove the convergence of the term in the brackets.

Step 1: Proof of the Convergence in [\(A.15\)](#page-56-0)

First, we prove this convergence for the case $\beta = \beta_L(1,1)$ with $g_\beta(x) = \min\{s(0,x), s(1,x)\}.$ Denoting $\Delta(x) = s(1, x) - s(0, x)$ and $\Delta(x, \theta) = s(1, x, \theta) - s(0, x, \theta)$, we note that

$$
\min\{s(0, x, \theta), s(1, x, \theta)\} = s(\mathbb{1}(\Delta(x, \theta) < 0), x, \theta)
$$
\n
$$
= s(0, x, \theta) + \Delta(x, \theta)\mathbb{1}(\Delta(x, \theta) < 0)
$$
\n
$$
= s(\mathbb{1}(\Delta(x) < 0), x, \theta) + \Delta(x, \theta)(\mathbb{1}(\Delta(x, \theta) < 0) - \mathbb{1}(\Delta(x) < 0)).
$$

Recalling the decomposition from the proof of Lemma [A.1,](#page-52-1) we also obtain for $\beta(x, \theta)$ that

$$
\beta(x,\theta) = \hat{\beta}(x,\theta) + \bar{\beta}(x,\theta),
$$

where $\hat{\beta}(x,\theta) = \text{and } \bar{\beta}(x,\theta) = \bar{\beta}_1(x,\theta) - \bar{\beta}_0(x,\theta).$

Then, we have that

$$
N(\theta) - N(\theta_0) = \Big(E_{\theta}[\hat{\beta}(X,\theta)s(\mathbb{1}(\Delta(X) < 0), X, \theta)] - E[\beta(X)s(\mathbb{1}(\Delta(X) < 0), X)] \Big) + E_{\theta}[\beta(X,\theta)\Delta(X,\theta)(\mathbb{1}(\Delta(X,\theta) < 0) - \mathbb{1}(\Delta(X) < 0))] + E_{\theta}[\bar{\beta}(X,\theta)s(\mathbb{1}(\Delta(X) < 0), X, \theta)] = \hat{N}(\theta) + \tilde{N}(\theta) + \bar{N}(\theta),
$$

Similarly, we decompose $D(\theta) - D(\theta_0)$ into $\hat{D}(\theta) + \tilde{D}(\theta)$ by setting $\beta(x, \theta)$ and $\beta(x)$ to one. It turns out that $\hat{N}(\theta)$ and $\hat{D}(\theta)$ are main terms, which include the component $D = \mathbb{1}(\Delta(X) < 0)$ that does not depend on θ and hence is easier to handle. On the other hand, the terms $\tilde{N}(\theta) - \beta \tilde{D}(\theta)$ and $\bar{N}(\theta) - \beta \bar{D}(\theta)$ include non-smooth components that depend on θ and turn out to be negligible.

In particular, we show that

$$
\lim_{\varepsilon \to 0} \frac{\hat{N}(\theta) - \beta \hat{D}(\theta)}{\varepsilon} = \int_{\mathcal{X}} (\beta(x) - \beta) \left(s'(\mathbb{1}(\Delta(x) < 0), x) + s(\mathbb{1}(\Delta(x) < 0), x) t(x, \theta_0) \right) f(x) dx, \\
+ \int_{\mathcal{X}} \frac{\partial \beta(x, \theta_0)}{\partial \theta} s(\mathbb{1}(\Delta(x) < 0), x) f(x) dx, \\
\lim_{\varepsilon \to 0} \frac{\tilde{N}(\theta) - \beta \tilde{D}(\theta)}{\varepsilon} = 0, \quad \lim_{\varepsilon \to 0} \frac{\bar{N}(\theta) - \beta \bar{D}(\theta)}{\varepsilon} = 0.
$$

The limiting result on $(\hat{N} - \beta \hat{D})(\theta)$ quickly follows by writing

$$
(\hat{N} - \beta \hat{D})(\theta) = \int_{\mathcal{X}} (\hat{\beta}(x,\theta) - \beta) s(\mathbb{1}(\Delta(x) < 0), x, \theta) f(x, \theta) dx - \int_{\mathcal{X}} (\beta(x) - \beta) s(\mathbb{1}(\Delta(x) < 0), x) f(x) dx,
$$

and exploiting Lemma [A.1](#page-52-1) with

$$
\lim_{\varepsilon \to 0} \frac{\hat{\beta}(x,\theta) - \beta(x)}{\varepsilon} = \frac{\partial \beta(x,\theta_0)}{\partial \theta}, \quad \lim_{\varepsilon \to 0} \frac{f(x,\theta) - f(x)}{\varepsilon} = t(x,\theta_0) f(x),
$$

$$
\lim_{\varepsilon \to 0} \frac{s(1_{\{\Delta(x) < 0\}}, x, \theta) - s(1_{\{\Delta(x) < 0\}}, x)}{\varepsilon} = s'(1_{\{\Delta(x) < 0\}}, x, \theta_0),
$$

along with product rule on differentiation and the boundedness of $\beta(x)$, $s(d, x)$ and $f(x)$.

Next, we deal with the term decompose $\tilde{N}(\theta) - \beta \tilde{D}(\theta)$. From Section [A.5.1,](#page-52-0) we first recall that $s(d, x, \theta) = s(d, x) + \varepsilon s'(d, x, \theta_0)$, which yields

$$
\Delta(x,\theta) - \Delta(x) = \varepsilon \left[s'(1,x,\theta_0) - s'(0,x,\theta_0) \right],\tag{A.16}
$$

and hence $|\Delta(x,\theta) - \Delta(x)| \leq C_1 |\varepsilon|$ for some constant $C_1 > 0$ since $s'(d, x, \theta_0)$ is bounded. We also obtain that

$$
|\mathbb{1}(\Delta(x,\theta) < 0) - \mathbb{1}(\Delta(x) < 0)| \le \mathbb{1}(|\Delta(x)| \le |\Delta(x,\theta) - \Delta(x)|),\tag{A.17}
$$

which exploits that the left-hand side is zero when both terms have same signs.

Consequently, using the identity $\mathcal{X}^+ \cup \mathcal{X}^- = \{x \in \mathcal{X} : |\Delta(x)| > 0\}$, we deduce that

$$
|\tilde{N}(\theta) - \beta \tilde{D}(\theta)| \le E_{\theta} [|\beta(X, \theta) - \beta||\Delta(X, \theta)||1(|\Delta(X)| \le |\Delta(X, \theta) - \Delta(X)|)1(|\Delta(X)| > 0)]
$$

$$
\le 2C_1C_2|\varepsilon|E_{\theta}[1(|\Delta(X)| \le 2C_1|\varepsilon|)1(|\Delta(X)| > 0)]
$$

$$
\le 2C_1C_2C_3|\varepsilon|E[1(0 < |\Delta(X)| \le 2C_1|\varepsilon|)] = o(\varepsilon),
$$

where we used $|\beta(x,\theta) - \beta| \leq C_2$ and $|\Delta(x,\theta)| \leq |\Delta(x)| + |\Delta(x,\theta) - \Delta(x)| \leq 2C_1|\varepsilon|$ under the indicator function $\mathbb{1}(|\Delta(X)| \leq |\Delta(X, \theta) - \Delta(X)|$) and $|1 + \varepsilon t(x)| \leq C_3$, and $\lim_{\varepsilon \to 0} P(0 < |\Delta(X)| \leq 2C_1 |\varepsilon|) = 0.$

We can deal with the term $\bar{N}(\theta) - \beta \bar{D}(\theta)$ similarly. In particular, the terms involving

the difference of two indicator functions are treated exactly as above. The other terms involving conditional quantiles are dealt with exploiting Assumption [5.1](#page-21-0) and turn out to be of order $O(\varepsilon^2)$.

As a result, we have shown the convergence in [\(A.15\)](#page-56-0) for the cases $\beta = \beta_L(1,1)$ and the obtained limiting integral can be written as

$$
\frac{\partial \beta(\theta_0)}{\partial \theta} = D(\theta_0)^{-1} \int_{\mathcal{X}^+} \left[(\beta(x) - \beta)(s'(0, x, \theta_0) + s(0, x)t(x, \theta_0)) + \frac{\partial \beta(x, \theta_0)}{\partial \theta} s(0, x) \right] f(x) dx + D(\theta_0)^{-1} \int_{\mathcal{X}^-} \left[(\beta(x) - \beta) (s'(1, x, \theta_0) + s(1, x)t(x, \theta_0)) + \frac{\partial \beta(x, \theta_0)}{\partial \theta} s(1, x) \right] f(x) dx.
$$

The case $\beta = \beta_U(1,1)$ has the same $g_\beta(x,\theta) = \min\{s(0,x,\theta), s(1,x,\theta)\}\)$ but based on β ^{*U*} (1, 1, *x*). Hence, its proof is analogous.

For the cases $\beta \in {\beta_L(0,1), \beta_L(em)}$, we have $g_\beta(x, \theta) = \max\{s(1, x, \theta) - s(0, x, \theta), 0\}$ and $g_{\beta}(x, \theta) = |s(1, x, \theta) - s(0, x, \theta)|$, respectively. We observe that

$$
\max\{s(1, x, \theta) - s(0, x, \theta), 0\} = s(1, x, \theta) - \min\{s(0, x, \theta), s(1, x, \theta)\},\
$$

$$
|s(1, x, \theta) - s(0, x, \theta)| = s(1, x, \theta) + s(0, x, \theta) - 2\min\{s(0, x, \theta), s(1, x, \theta)\},\
$$

and hence the convergence in [\(A.15\)](#page-56-0) for these two cases is handled analogously to the earlier min $\{s(0, x, \theta), s(1, x, \theta)\}\$ function case.

Step 2: Calculating the Tangent Space and the Score The density of the observed data (*SY, S, D, X*) is given by

$$
L(y, s, d, x) = [f_1(y|x)^d f_0(y|x)^{1-d} s(1, x)^d s(0, x)^{1-d}]^s
$$

$$
\times \left[(1 - s(1, x))^d (1 - s(0, x))^{1-d} \right]^{1-s} m(x)^d (1 - m(x))^{1-d} f(x)
$$

Now, we consider a regular parametric submodel indexed by θ with density

$$
L(y, s, d, x | \theta) = [f_1(y | x, \theta)^d f_0(y | x, \theta)^{1-d} s(1, x, \theta)^d s(0, x | \theta)^{1-d}]^s
$$

$$
\times \left[(1 - s(1, x, \theta))^{d} (1 - s(0, x, \theta))^{1-d} \right]^{1-s} m(x, \theta)^d (1 - m(x, \theta))^{1-d} f(x, \theta),
$$

which equals $L(y, s, d, x)$ when $\theta = \theta_0$.

If $m(x, \theta) = m(x)$ is known (does not depend on θ), its score is given by

$$
Q_1(y, s, d, x | \theta) = s d\tau_1(y | x, \theta) + s(1 - d)\tau_0(y | x, \theta) + t(x, \theta)
$$

+
$$
d \frac{(s - s(1, x, \theta))s'(1, x, \theta)}{s(1, x, \theta)(1 - s(1, x, \theta))} + (1 - d) \frac{(s - s(0, x, \theta))s'(0, x, \theta)}{s(0, x, \theta)(1 - s(0, x, \theta))}.
$$

In view of this, the tangent space of this model is

$$
\mathcal{S}_1 = \{sd\tau_1(y,x) + s(1-d)\tau_0(y,x) + \tau_x(x) + a(x)d(s-s(1,x)) + b(x)(1-d)(s-s(0,x))\},\
$$

where $a(x)$, $b(x)$ are any bounded functions, and $\tau_d(x)$, $d = 0, 1, t(x)$ are bounded functions that satisfy

$$
\int \tau_d(y, x) f_d(y|x) dy = 0, \forall x, \text{ and } \int t(x) f(x) dx = 0.
$$

If $m(x, \theta)$ is unknown (may depend on θ), its score is given by

$$
Q_2(y, s, d, x | \theta) = Q_1(y, s, d, x | \theta) + \frac{d - m(x, \theta)}{m(x, \theta)(1 - m(x, \theta))} \frac{\partial m(x, \theta)}{\partial \theta}.
$$

Hence, the tangent space of this model is

$$
S_2 = \{sd\tau_1(y,x) + s(1-d)\tau_0(y,x) + \tau_x(x) + a(x)d(s-s(1,x))
$$

+ $b(x)(1-d)(s-s(0,x)) + c(x)(d-m(x))\},\$

where $c(x)$ is any bounded function, and other functions were introduced earlier.

Step 3: Detailed Calculation of the Pathwise Derivative

For each $\beta \in {\beta_L(1,1), \beta_U(1,1), \beta_L(0,1), \beta_L(em)}$, our remaining task is to come up with the efficient influence function $\psi_{\beta}(W)$ such that for $i \in \{1, 2\}$

$$
\frac{\partial \beta(\theta_0)}{\partial \theta} = E[\psi_{\beta}(W)Q_i(W|\theta_0)].
$$
\n(A.18)

For both sides of this equality, we proceed on a case-by-case basis. In this step, based on Lemma [A.1](#page-52-1) and Step 1, we decompose the left-hand side of this as

$$
\frac{\partial \beta(\theta_0)}{\partial \theta} = (E[g_{\beta}(X)])^{-1} (T_1 + T_2 + T_3 + T_4 + T_5).
$$
\n(A.19)

First, we let $\beta = \beta_L(1, 1)$, where $g_\beta(x) = \min\{s(0, x), s(1, x)\}\$. Then, we have that

$$
T_{1} = \int_{\mathcal{X}} \int_{\underline{y}}^{q_{1}(\min\{p_{0}(x),1\},x)} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
T_{2} = -\int_{\mathcal{X}} \int_{q_{0}(\max\{1-1/p_{0}(x),0\},x)}^{q_{j}} y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{3} = \int_{\mathcal{X}} (\beta(x)-\beta) \left(s'(1_{\{\Delta(x)<0\}},x)+s(1_{\{\Delta(x)<0\}},x)t(x,\theta_{0})\right) f(x)dx,
$$

\n
$$
T_{4} = -\int_{\mathcal{X}^{+}} \int_{\underline{y}}^{q_{1}(p_{0}(x),x)} q_{1}(p_{0}(x),x)\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \int_{q_{0}(1-1/p_{0}(x),x)}^{q_{j}} q_{0}(1-1/p_{0}(x),x)\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{5} = \int_{\mathcal{X}^{+}} [q_{1}(p_{0}(x),x)-\beta_{1}(x)] \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \frac{[q_{0}(1-1/p_{0}(x),x)-\beta_{0}(x)]}{p_{0}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx.
$$

Second, we let $\beta = \beta_U(1,1)$, where $g_\beta(x) = \min\{s(0,x), s(1,x)\}\)$. Then, we have that

$$
T_{1} = \int_{\mathcal{X}} \int_{q_{1}(\max\{1-p_{0}(x),0\},x)}^{g} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
T_{2} = -\int_{\mathcal{X}} \int_{\underline{y}}^{q_{0}(\min\{1/p_{0}(x),1\},x)} y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{3} = \int_{\mathcal{X}} (\beta(x)-\beta) \left(s'(1_{\{\Delta(x)<0\}},x)+s(1_{\{\Delta(x)<0\}},x)t(x,\theta_{0})\right)f(x)dx,
$$

\n
$$
T_{4} = -\int_{\mathcal{X}^{+}} \int_{q_{1}(1-p_{0}(x),x)}^{y} q_{1}(1-p_{0}(x),x)\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \int_{\underline{y}}^{q_{0}(1/p_{0}(x),x)} q_{0}(1/p_{0}(x),x)\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{5} = \int_{\mathcal{X}^{+}} [q_{1}(1-p_{0}(x),x)-\beta_{1}(x)] \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \frac{[q_{0}(1/p_{0}(x),x)-\beta_{0}(x)]}{p_{0}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx.
$$

Third, we let $\beta = \beta_L(0,1)$, where $g_\beta(x) = \max\{s(1,x) - s(0,x), 0\}$. Then, we obtain that

$$
T_{1} = \int_{\mathcal{X}^{+}} \int_{\underline{y}_{1}}^{q_{1}(1-p_{0}(x),x)} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
T_{2} = -\int_{\mathcal{X}^{-}} \int_{q_{0}(1/p_{0}(x),x)}^{q_{0}(x)} y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{3} = \int_{\mathcal{X}^{+}} (\beta(x)-\beta)(s'(1,x,\theta_{0}) - s'(0,x,\theta_{0}) + (s(1,x)-s(0,x))t(x,\theta_{0}))f(x)dx,
$$

\n
$$
T_{4} = -\int_{\mathcal{X}^{+}} \int_{\underline{y}_{1}}^{q_{1}(1-p_{0}(x),x)} q_{1}(1-p_{0}(x),x)\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \int_{q_{0}(1/p_{0}(x),x)}^{q} q_{0}(1/p_{0}(x),x)\tau_{0}(y|x,\theta_{0})f_{0}(y|x)s(0,x)f(x)dydx,
$$

\n
$$
T_{5} = -\int_{\mathcal{X}^{+}} (q_{1}(1-p_{0}(x),x)-\beta_{1}(x)) \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx
$$

\n
$$
+ \int_{\mathcal{X}^{-}} \frac{[q_{0}(1/p_{0}(x),x)-\beta_{0}(x)]}{p_{0}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta} s(1,x)f(x)dx.
$$

Finally, we let $\beta = \beta_L(em)$, where $g_{\beta}(x) = |s(1, x) - s(0, x)|$. Then, we obtain that the same T_1, T_2, T_4 and T_5 terms as in the previous case due to $\beta_L(x, em) = \beta_L(x, 0, 1)$. The only new term here is

$$
T_3 = \int_{\mathcal{X}} (\beta(x) - \beta) (s'(1, x, \theta_0) - s'(0, x, \theta_0) + (s(1, x) - s(0, x))t(x, \theta_0)) f(x) dx,
$$

+
$$
\int_{\mathcal{X}^-} (\beta(x) - \beta) (s'(0, x, \theta_0) - s'(1, x, \theta_0) + (s(0, x) - s(1, x))t(x, \theta_0)) f(x) dx.
$$

Step 4: Proposing the Efficient Influence Function

In this step, for each β , we propose $\psi_{\beta}(W)$ on the right-hand side of [\(A.18\)](#page-59-0) in the form:

$$
\psi_{\beta}(W) = (E[g_{\beta}(X)])^{-1}(M_1 + M_2 + M_3 + M_4 + M_5). \tag{A.20}
$$

We obtained influence functions by following the heuristic approach of [Hines et al.](#page-45-4) [\(2022\)](#page-45-4). However, not to make the proof even longer, we refrain from showing these details and instead simply write these in the style of [Hahn](#page-44-1) [\(1998\)](#page-44-1).

First, let $\beta = \beta_L(1, 1)$, where $g_\beta(x) = \min\{s(0, x), s(1, x)\}\)$. Then, we have that

$$
M_{1} = \frac{SD}{m(X)} \left[Y1\{Y \leq q_{1}(\min\{p_{0}(X), 1\}, X)\} - \min\{p_{0}(X), 1\} \beta_{1}(X) \right],
$$

\n
$$
M_{2} = -\frac{S(1 - D)}{1 - m(X)} \left[Y1\{Y \geq q_{0}(1 - \min\{1/p_{0}(X), 1\}, X)\} - \min\{1/p_{0}(X), 1\} \beta_{0}(X) \right],
$$

\n
$$
M_{3} = \mathbb{I}(X \in \mathcal{X}^{+})(\beta(X) - \beta) \left[s(0, X) + \frac{1 - D}{1 - m(X)} (S - s(0, X)) \right]
$$

\n
$$
+ \mathbb{I}(X \in \mathcal{X}^{-})(\beta(X) - \beta) \left[s(1, X) + \frac{D}{m(X)} (S - s(1, X)) \right],
$$

\n
$$
M_{4} = - \mathbb{I}(X \in \mathcal{X}^{+}) \frac{SD}{m(X)} q_{1}(p_{0}(X), X) \left[\mathbb{I}(Y \leq q_{1}(p_{0}(X), X)) - p_{0}(X) \right]
$$

\n
$$
+ \mathbb{I}(X \in \mathcal{X}^{-}) \frac{S(1 - D)}{1 - m(X)} q_{0}(1 - p_{0}(X)^{-1}, X) \left[\mathbb{I}(Y \geq q_{0}(1 - p_{0}(X)^{-1}, X)) - p_{0}(X)^{-1} \right],
$$

\n
$$
M_{5} = \mathbb{I}(X \in \mathcal{X}^{+}) \left[q_{1}(p_{0}(X), X) - \beta_{1}(X) \right] \left[\frac{1 - D}{1 - m(X)} (S - s(0, X)) - p_{0}(X) \frac{D}{m(X)} (S - s(1, X)) \right]
$$

\n
$$
+ \mathbb{I}(X \in \mathcal{X}^{-}) \left[q_{0}(1 - p_{0}(X)^{-1}, X) - \beta_{0}(X) \right] \left[p_{0}(X)^{-1} \frac{1 - D}{1 - m(X)} (S - s(0, X)) - \frac{D}{m(X)} (S - s(1, X)) \right].
$$

Second, let $\beta = \beta_U(1,1)$, where $g_\beta(x) = \min\{s(0,x), s(1,x)\}\$. Then, we have that

$$
M_{1} = \frac{SD}{m(X)} \left[Y \mathbb{1}(Y \geq q_{1}(1 - \min\{p_{0}(X), 1\}, X)) - \min\{p_{0}(X), 1\} \beta_{1}(X) \right],
$$

\n
$$
M_{2} = -\frac{S(1 - D)}{1 - m(X)} \left[Y \mathbb{1}(Y \leq q_{0}(\min\{1/p_{0}(X), 1\}, X)) - \min\{1/p_{0}(X), 1\} \beta_{0}(X) \right],
$$

\n
$$
M_{3} = \mathbb{1}(X \in \mathcal{X}^{+})(\beta(X) - \beta) \left[s(0, X) + \frac{1 - D}{1 - m(X)}(S - s(0, X)) \right]
$$

\n
$$
+ \mathbb{1}(X \in \mathcal{X}^{-})(\beta(X) - \beta) \left[s(1, X) + \frac{D}{m(X)}(S - s(1, X)) \right],
$$

\n
$$
M_{4} = - \mathbb{1}(X \in \mathcal{X}^{+}) \frac{SD}{m(X)} q_{1}(1 - p_{0}(X), X) \left[\mathbb{1}(Y \geq q_{1}(1 - p_{0}(X), X)) - p_{0}(X) \right]
$$

\n
$$
+ \mathbb{1}(X \in \mathcal{X}^{-}) \frac{S(1 - D)}{1 - m(X)} q_{0}(1/p_{0}(X), X) \left[\mathbb{1}(Y \leq q_{0}(p_{0}(X)^{-1}, X)) - p_{0}(X)^{-1} \right],
$$

\n
$$
M_{5} = \mathbb{1}(X \in \mathcal{X}^{+}) \left[q_{1}(1 - p_{0}(X), X) - \beta_{1}(X) \right] \left[\frac{1 - D}{1 - m(X)}(S - s(0, X)) - p_{0}(X) \frac{D}{m(X)}(S - s(1, X)) \right]
$$

\n
$$
+ \mathbb{1}(X \in \mathcal{X}^{-}) \left[q_{0}(p_{0}(X)^{-1}, X) - \beta_{0}(X) \right] \left[p_{0}(X)^{-1} \frac{1 - D}{1 - m(X)}(S - s(0, X)) - \frac{D}{m(X)}(S - s(1, X)) \right].
$$

Third, let $\beta = \beta_L(0,1)$, where $g_\beta(x) = \max\{s(1,x) - s(0,x), 0\}$. Then, we obtain that

$$
M_1 = 1(X \in \mathcal{X}^+) \frac{SD}{m(X)} \left[Y1(Y \le q_1(1 - p_0(X), X)) - (1 - p_0(X))\beta_1(X) \right],
$$

\n
$$
M_2 = -1(X \in \mathcal{X}^-) \frac{S(1 - D)}{1 - m(X)} \left[Y1(Y \ge q_0(1/p_0(X), X)) - (1 - 1/p_0(X))\beta_0(X) \right],
$$

\n
$$
M_3 = 1(X \in \mathcal{X}^+) (\beta(X) - \beta) \left[s(1, X) - s(0, X) + \frac{D}{m(X)}(S - s(1, X)) - \frac{1 - D}{1 - m(X)}(S - s(0, X)) \right]
$$

\n
$$
+ 1(X \in \mathcal{X}^-) (\beta(X) - \beta) \left[s(0, X) - s(1, X) + \frac{1 - D}{1 - m(X)}(S - s(0, X)) - \frac{D}{m(X)}(S - s(1, X)) \right],
$$

\n
$$
M_4 = -1(X \in \mathcal{X}^+) \frac{SD}{m(X)} q_1(1 - p_0(X), X) \left[1(Y \le q_1(1 - p_0(X), X)) - (1 - p_0(X)) \right]
$$

\n
$$
+ 1(X \in \mathcal{X}^-) \frac{S(1 - D)}{1 - m(X)} q_0(p_0(X)^{-1}, X) \left[1(Y \ge q_0(p_0(X)^{-1}, X)) - (1 - p_0(X)^{-1}) \right],
$$

\n
$$
M_5 = -1(X \in \mathcal{X}^+) \left[q_1(1 - p_0(X), X) - \beta_1(X) \right] \left[\frac{1 - D}{1 - m(X)}(S - s(0, X)) - p_0(X) \frac{D}{m(X)}(S - s(1, X)) \right]
$$

\n
$$
+ 1(X \in \mathcal{X}^-) \left[q_0(p_0(X)^{-1}, X) - \beta_0(X) \right] \left[p_0(X)^{-1} \frac{1 - D}{1 - m(X)}(S - s(0, X)) - \frac{D}{m(X)}(S - s(1, X)) \right]
$$

Finally, let $\beta = \beta_L(em)$, where $g_\beta(x) = |s(1, x) - s(0, x)|$. Again, we obtain the same M_1, M_2, M_4 and M_5 terms as in the previous case due to $\beta_L(x, em) = \beta_L(x, 0, 1)$. The only new term is

$$
M_3 = \mathbb{1}(X \in \mathcal{X}^+)(\beta(X) - \beta) \left[s(1, X) - s(0, X) + \frac{D}{m(X)}(S - s(1, X)) - \frac{1 - D}{1 - m(X)}(S - s(0, X)) \right].
$$

Step 5: Checking the Equality in [\(A.18\)](#page-59-0)

We note that $M \in \mathcal{S}_1$ and $M \in \mathcal{S}_2$. In addition, for each $i = 1, \ldots, 5$, we have that

$$
T_i = E[M_i Q_1(Y, S, D, X | \theta_0)],
$$

$$
T_i = E[M_i Q_2(Y, S, D, X | \theta_0)],
$$

which shows that β is pathwise differentiable irrespective of $m(x, \theta)$ is known or unknown.

A.5.3. Pathwise differentiability in proof of Theorem [5.1](#page-22-0)

In this section, we show that $\beta \in {\beta_L(1,1), \beta_L(em)}$ is not pathwise differentiable if $P(X^0) > 0$ holds. Other cases are similar and therefore skipped. Our counterexample proposes $\tau_d(y|x, \theta_0)$, $s'(d, x, \theta_0)$, $t(x, \theta_0)$ (see Section [A.5.1\)](#page-52-0) such that

$$
\lim_{\varepsilon \downarrow 0} \frac{\beta(\theta) - \beta(\theta_0)}{\varepsilon} \neq \lim_{\varepsilon \uparrow 0} \frac{\beta(\theta) - \beta(\theta_0)}{\varepsilon}.
$$
\n(A.21)

We borrow the proof idea from [Luedtke and van der Laan](#page-46-6) [\(2016\)](#page-46-6). However, our case is more delicate because the divergence of the derivative of denominator $D(\theta)$ and numerator $N(\theta)$ does not necessarily imply the divergence of the ratio β . Recalling [\(A.15\)](#page-56-0) and since

 $\lim_{\varepsilon \to 0} D(\theta) = D(\theta_0)$, it suffices to show that $L_1 \neq L_2$, where

$$
L_1 \equiv \lim_{\varepsilon \downarrow 0} \frac{N(\theta) - N(\theta_0) - \beta(D(\theta) - D(\theta_0))}{\varepsilon},
$$

$$
L_2 \equiv \lim_{\varepsilon \uparrow 0} \frac{N(\theta) - N(\theta_0) - \beta(D(\theta) - D(\theta_0))}{\varepsilon}.
$$

First, we let $\beta = \beta_L(1,1)$ *.* In our decomposition introduced after [\(A.15\)](#page-56-0), we now include extra \mathcal{X}^0 terms in

$$
\tilde{N}(\theta) - \beta \tilde{D}(\theta) = E_{\theta}[1_{\{X \in \mathcal{X}^{0}\}}(\beta(X, \theta) - \beta)E_{\theta}[S|D = 1_{\{\Delta(X) < 0\}}, X]]
$$
\n
$$
- E[1_{\{X \in \mathcal{X}^{0}\}}(\beta(X) - \beta)E[S|D = 1_{\{\Delta(X) < 0\}}, X]],
$$
\n
$$
\tilde{N}(\theta) - \beta \tilde{D}(\theta) = E_{\theta}[1_{\{X \in \mathcal{X}^{0}\}}(\beta(X, \theta) - \beta) \Delta(X, \theta)(1_{\{\Delta(X, \theta) < 0\}} - 1_{\{\Delta(X) < 0\}})],
$$

while retaining $\mathcal{X}^+ \cup \mathcal{X}^-$ terms in $\hat{N}(\theta) - \beta \hat{D}(\theta)$, $\tilde{N}(\theta) - \beta \tilde{D}(\theta)$ and $\bar{N}(\theta) - \beta \bar{D}(\theta)$.

We let $\tau_d(y|x, \theta_0) = 0$, $t(x, \theta_0) = 0$, $s'(0, x, \theta_0) = 0$ and $s'(1, x, \theta_0) = 1$ for all d, y and $x \in \mathcal{X}$. Then, our results obtained in Step 1 of Theorem [5.1](#page-22-0) imply

$$
\lim_{\varepsilon \to 0} \frac{\hat{N}(\theta) - \beta \hat{D}(\theta)}{\varepsilon} = \int_{\mathcal{X}^{-}} (\beta(x) - \beta) f(x) dx + \int_{\mathcal{X}^{+} \cup \mathcal{X}^{-}} \frac{\partial \beta(x, \theta_{0})}{\partial \theta} s(1_{\{\Delta(x) < 0\}}, x) f(x) dx,
$$
\n
$$
\lim_{\varepsilon \to 0} \frac{\tilde{N}(\theta) - \beta \tilde{D}(\theta)}{\varepsilon} = 0 = \lim_{\varepsilon \to 0} \frac{\bar{N}(\theta) - \beta \bar{D}(\theta)}{\varepsilon},
$$

where some simplifications are due to $t(x, \theta_0) = s'(0, x, \theta_0) = 0$ and $s'(1, x, \theta_0) = 1$.

Before we proceed to the other two terms, for $x \in \mathcal{X}^0$, we will show that $\beta(x, \theta)$ is not pathwise differentiable. In particular, we will prove that

$$
\lim_{\varepsilon \downarrow 0} \frac{\beta(x,\theta) - \beta(x,\theta_0)}{\varepsilon} = L(x) + (\beta_1(x) - \overline{y}_1) \frac{s'(1,x,\theta_0) - s'(0,x,\theta_0)}{s(1,x)},
$$

$$
\lim_{\varepsilon \uparrow 0} \frac{\beta(x,\theta) - \beta(x,\theta_0)}{\varepsilon} = L(x) + (\beta_0(x) - \underline{y}_0) \frac{s'(1,x,\theta_0) - s'(0,x,\theta_0)}{s(1,x)},
$$

where

$$
L(x) = \int y\tau_1(y|x,\theta_0)f_1(y|x)dy - \int y\tau_0(y|x,\theta_0)f_0(y|x)dy.
$$

We have $p_0(x) = 1$ for $x \in \mathcal{X}^0$. Since $s'(1, x, \theta) = 1 > 0 = s'(0, x, \theta_0)$, we have that $p_0(x, \theta) < 1$ for $\varepsilon > 0$, and $p_0(x, \theta) > 1$ for $\varepsilon < 0$. Then, it suffices to show that

$$
\lim_{\varepsilon \downarrow 0} \frac{\beta_{1,1}(x, p_0(x, \theta), \theta) - \beta_{1,1}(x, 1, \theta)}{\varepsilon} = (\beta_1(x) - \overline{y}_1) \frac{s'(1, x, \theta_0) - s'(0, x, \theta_0)}{s(1, x)},
$$

$$
\lim_{\varepsilon \uparrow 0} \frac{\beta_{0,0}(x, 1 - 1/p_0(x, \theta), \theta) - \beta_{0,0}(x, 0, \theta)}{\varepsilon} = -(\beta_0(x) - \underline{y}_0) \frac{s'(1, x, \theta_0) - s'(0, x, \theta_0)}{s(1, x)}.
$$

To prove the first convergence, we first write this as

$$
\frac{\beta_{1,1}(x, p_0(x,\theta), \theta) - \beta_{1,1}(x,1,\theta)}{\varepsilon} = \frac{1 - p_0(x,\theta)}{p_0(x,\theta)} \int_{\underline{y}_1}^{q_1(p_0(x,\theta),x)} y f_1(y|x,\theta) dy - \int_{q_1(p_0(x,\theta),x)}^{\overline{y}_1} y f_1(y|x,\theta) dy,
$$

and hence the result follows from $\lim_{\varepsilon \downarrow 0} \varepsilon^{-1}(1-p_0(x,\theta)) = (s'(1,x,\theta_0) - s'(0,x,\theta_0))/s(1,x)$, and due to ε^{-1} times the second integral converging to $\overline{y}_1(s'(1, x, \theta_0) - s'(0, x, \theta_0))/s(1, x)$.

The second convergence follows similarly by using the decomposition

$$
\beta_{0,0}(x, 1 - 1/p_0(x, \theta), \theta) - \beta_{0,0}(x, 0, \theta) = (p_0(x, \theta) - 1) \int_{q_0(1 - p_0^{-1}(x, \theta), x)}^{\overline{y}_0} y f_0(y|x, \theta) dy - \int_{\underline{y}_0}^{q_0(1 - p_0^{-1}(x, \theta), x)} y f_0(y|x, \theta) dy.
$$

Having established this result, for $x \in \mathcal{X}^0$, we deduce that

$$
\lim_{\varepsilon \downarrow 0} \frac{\check{N}(\theta) - \beta \check{D}(\theta)}{\varepsilon} = \int_{\mathcal{X}^0} (\beta_1(x) - \overline{y}_1) f(x) dx,
$$

$$
\lim_{\varepsilon \uparrow 0} \frac{\check{N}(\theta) - \beta \check{D}(\theta)}{\varepsilon} = \int_{\mathcal{X}^0} (\beta_0(x) - \underline{y}_0) f(x) dx.
$$

Finally, we deal with the term $\ddot{N}(\theta) - \beta \ddot{D}(\theta)$. For $x \in \mathcal{X}^0$ we note that $\Delta(x) = 0$ and $\Delta(x, \theta) > 0$ if $\varepsilon > 0$. This quickly implies that

$$
\lim_{\varepsilon \downarrow 0} \frac{\ddot{N}(\theta) - \beta \ddot{D}(\theta)}{\varepsilon} = 0.
$$

On the other hand, we have $\Delta(x, \theta) < 0$ if $\varepsilon < 0$, which leads to

$$
\lim_{\varepsilon \uparrow 0} \frac{\ddot{N}(\theta) - \beta \ddot{D}(\theta)}{\varepsilon} = \int_{\mathcal{X}^0} (\beta(x) - \beta) f(x) dx.
$$

where we used $\beta(x, \theta) \to \beta(x)$ and $f(x, \theta) = f(x)$ and $\Delta(x) = 0$ for $x \in \mathcal{X}^0$.

Combining all these limits and denoting we obtain that

$$
L_1 - L_2 = \int_{\mathcal{X}^0} (\beta - (\overline{y}_1 - \underline{y}_0) f(x) dx = P(\mathcal{X}^0)(\beta - (\overline{y}_1 - \underline{y}_0)) < 0,
$$

where we used that $P(X^0) > 0$ and $\beta < (\overline{y}_1 - \underline{y}_0)$. The last inequality follows from *f*_{*d}*(*y*|*x*) being continuous density with support on $[\underline{y}_d, \overline{y}_d]$ and hence $\beta_{1,1}(x, u) < \overline{y}_1$ and</sub> $\beta_{0,0}(x,u) > y_0.$

Next, we consider the extensive margin case by letting $\beta = \beta_L(em)$. We again set $\tau_d(y|x, \theta_0) = 0$, $t(x, \theta_0) = 0$, $s'(0, x, \theta_0) = 0$ and $s'(1, x, \theta_0) = 1$.

We will only look at the difference between the terms evaluated on the set \mathcal{X}^0 since the terms convergence on $\mathcal{X}^+ \cup \mathcal{X}^-$ due to the results obtained in Step 1 of Theorem [5.1.](#page-22-0) We note that $|s(1, x) - s(0, x)| = 0$ for $x \in \mathcal{X}^0$, which implies that

$$
N(\theta_0) - \beta D(\theta_0) = \int_{\mathcal{X}^0} (\beta(x) - \beta)|s(1, x) - s(0, x)|f(x)dx = 0.
$$

On the other hand, using $f(x, \theta) = f(x)$ due to $t(x, \theta_0) = 0$, we have that

$$
N(\theta) - \beta D(\theta) = \int_{\mathcal{X}^0} (\beta(x, \theta) - \beta) |s(1, x, \theta) - s(0, x, \theta)| f(x) dx.
$$

Since $|s(1, x, \theta) - s(0, x, \theta)| = |\varepsilon|$ for $x \in \mathcal{X}^0$, we obtain that

$$
\varepsilon^{-1}[N(\theta) - \beta D(\theta) - (N(\theta_0) - \beta D(\theta_0))] = \text{sign}(\varepsilon) \int_{\mathcal{X}^0} (\beta(x, \theta) - \beta) f(x) dx.
$$

Due to $\lim_{\varepsilon \to 0} \beta(x, \theta) = \beta(x) = \underline{y}_1 - \overline{y}_0$, we obtain that

$$
\lim_{\varepsilon \downarrow 0} \varepsilon^{-1} \left[N(\theta) - \beta D(\theta) - (N(\theta_0) - \beta D(\theta_0)) \right] = \int_{\mathcal{X}^0} (\beta(x) - \beta) f(x) dx < 0
$$
\n
$$
\lim_{\varepsilon \uparrow 0} \varepsilon^{-1} \left[N(\theta) - \beta D(\theta) - (N(\theta_0) - \beta D(\theta_0)) \right] = - \int_{\mathcal{X}^0} (\beta(x) - \beta) f(x) dx > 0,
$$

where we used that $\beta(x) = y_1 - \overline{y}_0$ for $x \in \mathcal{X}^0$ and $\beta > y_1 - \overline{y}_0$. The last inequality is the consequence of $f_d(y|x)$ being continuous density with support on $[\underline{y}_d, \overline{y}_d]$ and hence $\beta_{1,1}(x, u) > \underline{y}_1$ for $u > 0$ and $\beta_{0,0}(x, u) < \overline{y}_0$ for $u < 1$ and $P(\mathcal{X}^+ \cup \mathcal{X}^-) > 0$.

A.6. Proof of Corollary [5.1](#page-22-1)

The efficiency lower bound is equal to $E[M^2]$, where the contribution terms to the total variance are

$$
E[M_1^2] + E[M_2^2] = E\left[\frac{s(1, X)\sigma_1^2(X)}{m(X)} + \frac{s(0, X)\sigma_0^2(X)}{1 - m(X)}\right],
$$

\n
$$
E[M_3^2] = E\left[1_{\mathcal{X}} + (X)(\beta_L(X) - \beta)^2 \frac{s(0, X)(1 - s(0, X)m(X))}{1 - m(X)}\right]
$$

\n
$$
+ E\left[1_{\mathcal{X}} - (X)(\beta_L(X) - \beta)^2 \frac{s(1, X)(1 - s(1, X) + s(1, X)m(X))}{m(X)}\right]
$$

\n
$$
E[M_4^2] = E\left[1_{\mathcal{X}} + \frac{s(1, X)q_1(p_0(X), X)^2 p_0(X)(1 - p_0(X))}{m(X)}\right]
$$

\n
$$
+ E\left[1_{\mathcal{X}} - \frac{s(0, X)q_0(1 - 1/p_0(X), X)^2 p_0(X)^{-1}(1 - p_0(X)^{-1})}{1 - m(X)}\right]
$$

\n
$$
E[M_5^2] = E\left[1_{\mathcal{X}} + (q_1(p_0(X), X) - \beta_1(X))^2 \left(\frac{s(0, X)(1 - s(0, X))}{1 - m(X)} + \frac{p_0(X)^2 s(1, X)(1 - s(1, X))}{m(X)}\right)\right]
$$

\n
$$
+ E\left[1_{\mathcal{X}} - (q_0(1 - 1/p_0(X), X) - \beta_0(X))^2 \left(\frac{p_0(X)^{-2} s(0, X)(1 - s(0, X))}{1 - m(X)} + \frac{s(1, X)(1 - s(1, X))}{m(X)}\right)\right]
$$

\n
$$
E[M_1M_4] = - E\left[1_{\mathcal{X}} + \frac{q_1(p_0(X), X)\beta_1(X)s(1, X)p_0(X)(1 - p_0(X))}{m(X)}\right]
$$

\n
$$
E[M_2M_4] = - E\left[1_{\mathcal{X}} - \frac{q_0(1 - 1/p_0(X), X)\beta_0(X)s(0, X)p_0(X)^{-1}(1 - p_0(X)^{-1})}{1 - m(X)}\right]
$$

\n
$$
E[M_3M_5] = E\left[1_{\math
$$

A.7. Proof of Theorem [5.2](#page-23-0)

First, we provide an auxiliary result similar to Lemma [A.1.](#page-52-1) For $B \in \{L, U\}$, we use the decomposition $\beta_{B,h}(x, 1, 1) = \beta_{B,1,h}(x, 1, 1) - \beta_{B,0,h}(x, 1, 1)$.

Lemma A.2 *Suppose that the assumptions of Theorem [5.2](#page-23-0) hold. Then, for each* $x \in \mathcal{X}$, $\beta_{L,h}(x,1,1)$ *and* $\beta_{U,h}(x,1,1)$ *are pathwise differentiable. In particular, we have that*

$$
\frac{\partial \beta_{L,1,h}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{g_{1,h}(p_{0}(x))} \int_{\underline{y}_{1}}^{q_{1}(g_{1,h}(p_{0}(x)),x)} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)dy \n- \frac{q_{1}(g_{1,h}(p_{0}(x)),x)}{g_{1,h}(p_{0}(x))} \int_{\underline{y}_{1}}^{q_{1}(g_{1,h}(p_{0}(x)),x)} \tau_{1}(y|x,\theta_{0})f_{1}(y|x)dy \n+ (q_{1}(g_{1,h}(p_{0}(x)),x) - \beta_{L1,h}(x,1,1)) \frac{g'_{1,h}(p_{0}(x))}{g_{1,h}(p_{0}(x))} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta}, \n\frac{\partial \beta_{L,0,h}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{g_{1,h}(1/p_{0}(x))} \int_{q_{0}(1-g_{1,h}(1/p_{0}(x)),x)}^{y_{0}} y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)dy \n- \frac{q_{0}(1-g_{1,h}(1/p_{0}(x)),x)}{g_{1,h}(1/p_{0}(x))} \int_{q_{0}(1-g_{1,h}(1/p_{0}(x)),x)}^{y_{0}} \tau_{0}(y|x,\theta_{0})f_{0}(y|x)dy \n- (q_{0}(1-g_{1,h}(1/p_{0}(x)),x) - \beta_{L0,h}(x,1,1,\theta_{0})) \frac{g'_{1,h}(1/p_{0}(x))}{g_{1,h}(1/p_{0}(x))p_{0}^{2}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta},
$$

and

$$
\frac{\partial \beta_{U,1,h}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{g_{1,h}(p_{0}(x))} \int_{q_{1}(1-g_{1,h}(p_{0}(x)),x)}^{q_{1}} y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)dy \n- \frac{q_{1}(1-g_{1,h}(p_{0}(x)),x)}{g_{1,h}(p_{0}(x))} \int_{q_{1}(1-g_{1,h}(p_{0}(x)),x)}^{q_{1}} \tau_{1}(y|x,\theta_{0})f_{1}(y|x)dy \n+ (q_{1}(1-g_{1,h}(p_{0}(x)),x) - \beta_{U1,h}(x,1,1)) \frac{g'_{1,h}(p_{0}(x))}{g_{1,h}(p_{0}(x))} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta}, \n\frac{\partial \beta_{U,0,h}(x,1,1,\theta_{0})}{\partial \theta} = \frac{1}{g_{1,h}(1/p_{0}(x))} \int_{\underline{y}_{0}}^{q_{0}(g_{1,h}(1/p_{0}(x)),x)} y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)dy \n- \frac{q_{0}(g_{1,h}(1/p_{0}(x)),x)}{g_{1,h}(1/p_{0}(x))} \int_{\underline{y}_{0}}^{q_{0}(g_{1,h}(1/p_{0}(x)),x)} \tau_{0}(y|x,\theta_{0})f_{0}(y|x)dy \n- (q_{0}(g_{1,h}(1/p_{0}(x)),x) - \beta_{U0,h}(x,1,1)) \frac{g'_{1,h}(1/p_{0}(x))}{g_{1,h}(1/p_{0}(x))p_{0}^{2}(x)} \frac{\partial p_{0}(x,\theta_{0})}{\partial \theta}.
$$

Proof. Its proof is a simpler version of the proof of Lemma [A.1](#page-52-1) since the irregularity issues are not present here. Details are omitted for brevity.

The rest of the proof is analogous and we only present the most important parts. First, we deal with the lower bound case. We write $\beta_{L,h}$ instead of $\beta_{L,h}(1,1)$, and split it as

$$
\beta_{L,h} = \beta_{L,+,h} + \beta_{L,-,h} \tag{A.22}
$$

with

$$
\beta_{L,+,h} = \frac{E[g_{4,h}(\beta_{L,h}(X))g_{1,h}(p_0(X))s(1,X)]}{E[g_{3,h}(p_0(X))s(1,X)]}, \quad \beta_{L,-,h} = \frac{E[g_{5,h}(\beta_{L,h}(X))g_{3,h}(p_0(X))s(1,X)]}{E[g_{1,h}(p_0(X))s(1,X)]}.
$$

where we use the notation $g_{5,h}(z) = -g_{2,h}(-z)$. This leads to

$$
\frac{\partial \beta_{L,h}(\theta_0)}{\partial \theta} = (T_1^+ + T_2^+ + T_3^+ + T_4^+ + T_5^+)/E[g_{3,h}(p_0(X))s(1,X)]
$$

+ $(T_1^- + T_2^- + T_3^- + T_4^- + T_5^-)/E[g_{1,h}(p_0(X))s(1,X)],$

$$
\begin{split} T_{1}^{+}=&\int_{\mathcal{X}}\int_{\underline{y}_{1}}^{q_{1}(g_{1,h}(p_{0}(x)),x)}g_{4,h}^{'}(\beta_{L,h}(x))y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)s(1,x)f(x)dydx,\\ T_{1}^{-}=&\int_{\mathcal{X}}\int_{\underline{y}_{1}}^{q_{1}(g_{1,h}(p_{0}(x)),x)}g_{5,h}^{'}(\beta_{L,h}(x))y\tau_{1}(y|x,\theta_{0})f_{1}(y|x)\frac{g_{3,h}(p_{0}(x))}{g_{1,h}(p_{0}(x))}\nonumber\\ x_{2}^{+}=-\int_{\mathcal{X}}\int_{g_{0}(1-g_{1,h}(1/p_{0}(x)),x)}^{g_{0}}g_{1,h}^{'}(\beta_{L,h}(x))y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)g_{1,h}(p_{0}(x))s(1,x)f(x)dydx,\\ T_{2}^{-}=-\int_{\mathcal{X}}\int_{g_{0}(1-g_{1,h}(1/p_{0}(x)),x)}^{g_{1}}g_{1,h}(1/p_{0}(x))y\tau_{0}(y|x,\theta_{0})f_{0}(y|x)g_{3,h}(p_{0}(x))s(1,x)f(x)dydx,\\ T_{3}^{+}=&\int_{\mathcal{X}}g_{4,h}(\beta_{L,h}(x))((g_{1,h}(p_{0}(x))s(1,x))')'+g_{1,h}(p_{0}(x))s(1,x)t(x))f(x)dx\\ &-\int_{\mathcal{X}}\beta_{L,+h}((g_{1,h}(p_{0}(x))s(1,x))')'+g_{1,h}(p_{0}(x))s(1,x)t(x))f(x)dx,\\ T_{3}^{+}=&\int_{\mathcal{X}}g_{5,h}(\beta_{L,h}(x))((g_{3,h}(p_{0}(x))s(1,x))')'+g_{3,h}(p_{0}(x))s(1,x)t(x))f(x)dx\\ &-\int_{\mathcal{X}}\beta_{L,-h}((g_{3,h}(p_{0}(x))s(1,x))')'+g_{3,h}(p_{0}(x))s(1,x)t(x))f(x)dx,\\ T_{4}^{+}=-\int_{\mathcal{X}}\int_{\underline{y}_{1}}^{q_{1}(g_{1,h}(p_{0}(x))x)}q_{0}(1-g_{1,h}(1/p_{0}(x)),x)\frac{g_{4,h}(g_{L,h}(x))}{g_{1,h}(1/p_{0}(x))}\tau_{0
$$

Next, we find the corresponding influence functions for $\beta_{L,+,h}$ and $\beta_{L,-,h}$, separately. For $\beta_{L,+,h}$, we consider the following candidate influence function

$$
M_{L,+,h} = (M_1^+ + M_2^+ + M_3^+ + M_4^+ + M_5^+)/E[g_{3,h}(p_0(X))s(1,X)],
$$
\n(A.23)

$$
\begin{split} M_{1}^{+}=&\frac{SDg'_{4,h}(\beta_{L,h}(X))}{m(X)}\left[Y1(Y\leq q_{1}(g_{1,h}(p_{0}(X)),X))-g_{1,h}(p_{0}(X))\beta_{L,1,h}(X)\right],\\ M_{2}^{+}=&-\frac{S(1-D)}{1-m(X)}\frac{g'_{4,h}(\beta_{L,h}(X))g_{1,h}(p_{0}(X))}{p_{0}(X)g_{1,h}(1/p_{0}(X))}\\ &\times\left[Y1(Y\geq q_{0}(1-g_{1,h}(1/p_{0}(X)),X))-g_{1,h}(1/p_{0}(X))\beta_{L,0,h}(X)\right],\\ M_{3}^{+}=g_{4,h}(\beta_{L,h}(X))\left[g_{1,h}(p_{0}(X))s(1,X)+g'_{1,h}(p_{0}(X))\frac{1-D}{1-m(X)}(S-s(0,X))\right]\\ &+g_{4,h}(\beta_{L,h}(X))\left[(g_{1,h}(p_{0}(X))-p_{0}(X)g'_{1,h}(p_{0}(X)))\frac{1-D}{m(X)}(S-s(1,X))\right]\\ &-\beta_{L,+h}\left[g_{3,h}(p_{0}(X))s(1,X)+g'_{3,h}(p_{0}(X))\frac{1-D}{1-m(X)}(S-s(0,X))\right]\\ &-\beta_{L,+h}\left[(g_{3,h}(p_{0}(X))-p_{0}(X)g'_{3,h}(p_{0}(X)))\frac{D}{m(X)}(S-s(1,X))\right],\\ M_{4}^{+}=&-\frac{SDg'_{4,h}(S_{L,h}(X))}{m(X)}q_{1}(g_{1,h}(p_{0}(X)),X)\left[1(Y\leq q_{1}(g_{1,h}(p_{0}(X)),X))-g_{1,h}(p_{0}(X))\right]\\ &+\frac{S(1-D)}{1-m(X)}\frac{g'_{4,h}(S_{L,h}(X))g_{1,h}(1/p_{0}(X))}{p_{0}(X)g_{1,h}(1/p_{0}(X))}q_{0}(1-g_{1,h}(1/p_{0}(X)),X)\\ &\times\left[1(Y\geq q_{0}(1-g_{1,h}(1/p_{0}(X)),X)-\beta_{L,1,h}(X))g'_{1,h}(p_{0}(X))\right]\\ &+(q_{0}(1-g_{1,h}(1/p_{0}(X)),X)-\beta_{L,0,h}(X))\frac{g'_{1,h}(1/p_{0}(X))g_{1,h}(p_{0}(X))}{g_{1,h}(1/p_{0}(X))p_{0}^{2}(X)}
$$

Similarly, for $\beta_{L,-,h},$ we have the following candidate influence function

$$
M_{L,-,h} = (M_1^- + M_2^- + M_3^- + M_4^- + M_5^-)/E[g_{1,h}(p_0(X))s(1,X)],
$$
\n(A.24)

$$
\begin{split} M_1^- = & \frac{SDg^{\ell}_{5,h}(\beta_{L,h}(X))}{m(X)} \frac{g_{3,h}(p_0(X))}{g_{1,h}(p_0(X))} \left[Y1(Y \leq q_1(g_{1,h}(p_0(X)),X)) - g_{1,h}(p_0(X)) \beta_{L,1,h}(X) \right], \\ M_2^- = & - \frac{S(1-D)}{1-m(X)} \frac{g^{\ell}_{5,h}(\beta_{L,h}(X))g_{3,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X)),X)) - g_{1,h}(1/p_0(X)) \beta_{L,0,h}(X) \right], \\ M_3^- = & g_{5,h}(\beta_{L,h}(X)) \left[g_{3,h}(p_0(X))s(1,X) + g^{\ell}_{3,h}(p_0(X)) \frac{1-D}{1-m(X)}(S-s(0,X)) \right] \\ & + g_{5,h}(\beta_{L,h}(X)) \left[(g_{3,h}(p_0(X)) - p_0(X)g^{\ell}_{3,h}(p_0(X))) \frac{1-D}{m(X)}(S-s(1,X)) \right] \\ & - \beta_{L,-,h} \left[g_{1,h}(p_0(X))s(1,X) + g^{\ell}_{1,h}(p_0(X)) \frac{1-D}{1-m(X)}(S-s(0,X)) \right] \\ & - \beta_{L,-,h} \left[(g_{1,h}(p_0(X)) - p_0(X)g^{\ell}_{1,h}(p_0(X))) \frac{D}{m(X)}(S-s(1,X)) \right], \\ M_4^- = & - \frac{SDg^{\ell}_{5,h}(\beta_{L,h}(X))}{m(X)} \frac{g_{3,h}(p_0(X))}{g_{1,h}(p_0(X))} q_1(g_{1,h}(p_0(X)),X) \\ & \times \left[1(Y \leq q_1(g_{1,h}(p_0(X)),X)) - g_{1,h}(p_0(X)) \right] \\ & + \frac{S(1-D)}{1-m(X)} \frac{g^{\ell}_{5,h}(\beta_{L,h}(X))g_{3,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X))} q_0(1-g_{1,h}(1/p_0(X)),X) \\ & \times \left[1(Y \geq q_0(1-g_{1,h}(1/p_0(X)),X) - \beta_{L,1,h}(X)) \frac{g^{\ell}_{1,h}(p_0(X))}{g_{1,h}(p_0(X))} g_{3,h}(p_0(X)) \right. \\ & + (q_0(1-g_{1,h}(1
$$

For each $i \in \{1, \ldots, 5\}$, we have that

$$
T_i^+ = E[M_i^+ Q(Y, S, D, X | \theta_0)], \quad T_i^- = E[M_i^- Q(Y, S, D, X | \theta_0)].
$$

As a result, $\beta_{L,+,h}$ and $\beta_{L,-,h}$ are pathwise differentiable and their efficient influence functions are given by $M_{L,+h}$ and $M_{L,-h}$, respectively. Due to the linearity property of the influence function, $\beta_{L,h}$ is pathwise differentiable and its efficient influence function is $M_{L,+,h} + M_{L,-,h}$.

In addition, $M_{L,+h} \in S_1$ and $M_{L,-h} \in S_1$, where S_1 denotes the score of the observed data (SY, S, D, X) Hence, the efficiency bound for $\beta_{L,h}$ is equal to $E[(M^h_+ + M^h_-)^2]$.

Results for the smooth upper bound $\beta_{U,h}$ are similar and therefore omitted. Below, we only write its influence function. Again, we use the decompositions $\beta_{U,h} = \beta_{U,+h} + \beta_{U,-h}$ and $M_{U,h} = M_{U,+,h} + M_{U,-,h}$. First, we have that

$$
M_{U,+,h} = (M_1^+ + M_2^+ + M_3^+ + M_4^+ + M_5^+)/E[g_{1,h}(p_0(X))s(1,X)],
$$
\n(A.25)

$$
\begin{split} M_{1}^{+}=&\frac{SDg_{2,h}^{t}(\beta_{U,h}(X))}{m(X)}\frac{g_{3,h}(p_{0}(X))}{g_{1,h}(p_{0}(X))}[Y1(Y\geq q_{1}(1-g_{1,h}(p_{0}(X)),X))-g_{1,h}(p_{0}(X))\beta_{U,1,h}(X)]\,,\\ M_{2}^{+}=&-\frac{S(1-D)}{1-m(X)}\frac{g_{2,h}^{t}(\beta_{U,h}(X))g_{3,h}(p_{0}(X))}{p_{0}(X)g_{1,h}(1/p_{0}(X))}\\ &\times[Y1(Y\leq q_{0}(g_{1,h}(1/p_{0}(X)),X))-g_{1,h}(1/p_{0}(X))\beta_{U,0,h}(X)]\,,\\ M_{3}^{+}=g_{2,h}(\beta_{U,h}(X))\left[g_{3,h}(p_{0}(X));s(1,X)+g_{3,h}'(p_{0}(X))\frac{1-D}{1-m(X)}(S-s(0,X))\right]\\ &+g_{2,h}(\beta_{U,h}(X))\left[(g_{3,h}(p_{0}(X))-p_{0}(X)g_{3,h}'(p_{0}(X)))\frac{D}{m(X)}(S-s(1,X))\right]\\ &-\beta_{U,+h}\left[g_{1,h}(p_{0}(X));s(1,X)+g_{1,h}'(p_{0}(X))\frac{1-D}{1-m(X)}(S-s(0,X))\right]\\ &-\beta_{U,+h}\left[(g_{1,h}(p_{0}(X))-p_{0}(X)g_{1,h}'(p_{0}(X)))\frac{D}{m(X)}(S-s(1,X))\right],\\ M_{4}^{+}=&-\frac{SDg_{2,h}'(\beta_{U,h}(X))}{m(X)}\frac{g_{3,h}(p_{0}(X))}{g_{1,h}(p_{0}(X))}q_{1}(1-g_{1,h}(p_{0}(X)),X)\\ &\times\left[1(Y\geq q_{1}(1-g_{1,h}(p_{0}(X)),X))-g_{1,h}(1/p_{0}(X))\right]\\ &+\frac{S(1-D)}{1-m(X)}\frac{g_{2,h}'(\beta_{U,h}(X))g_{3,h}(p_{0}(X))}{p_{0}(X,g_{1,h}(1/p_{0}(X))}-g_{0}(g_{1,h}(1/p_{0}(X)),X)\\ &\times\left[1(Y\leq q_{0}(g_{1,h}(1/p_{0}(X)),X)-\beta_{U,0,h}(X))\frac{g_{1,h}(1/p_{0}(X))g_{3,h}(p_{0}(X))}{g_{1,h}(p_{0}(X))
$$

Further, we have that

$$
M_{U,-,h} = (M_1^- + M_2^- + M_3^- + M_4^- + M_5^-)/E[g_{3,h}(p_0(X))s(1,X)],
$$
\n(A.26)
where we use the notation $g_{6,h}(z) = -g_{4,h}(-z)$ and

$$
\begin{split} M_1^- = & \frac{SDg'_{6,h}(\beta_{U,h}(X))}{m(X)} \left[Y1(Y \geq q_1(1-g_{1,h}(p_0(X)),X)) - g_{1,h}(p_0(X)) \beta_{U,1,h}(X) \right], \\ M_2^- = & - \frac{S(1-D)}{1-m(X)} \frac{g'_{6,h}(\beta_{U,h}(X))g_{1,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X))} \\ & \qquad \times \left[Y1(Y \leq q_0(g_{1,h}(1/p_0(X)),X)) - g_{1,h}(1/p_0(X)) \beta_{U,0,h}(X) \right], \\ M_3^- = & g_{6,h}(\beta_{U,h}(X)) \left[g_{1,h}(p_0(X))s(1,X) + g'_{1,h}(p_0(X)) \frac{1-D}{1-m(X)}(S-s(0,X)) \right] \\ & + g_{6,h}(\beta_{U,h}(X)) \left[(g_{1,h}(p_0(X)) - p_0(X)g'_{1,h}(p_0(X))) \frac{D}{m(X)}(S-s(1,X)) \right] \\ & - \beta_{U,-h} \left[g_{3,h}(p_0(X))s(1,X) + g'_{3,h}(p_0(X)) \frac{1-D}{1-m(X)}(S-s(0,X)) \right] \\ & - \beta_{U,-h} \left[(g_{3,h}(p_0(X)) - p_0(X)g'_{3,h}(p_0(X))) \frac{D}{m(X)}(S-s(1,X)) \right], \\ M_4^- = & - \frac{SDg'_{6,h}(\beta_{U,h}(X))}{m(X)} q_1(1-g_{1,h}(p_0(X)),X) \left[1(Y \geq q_1(1-g_{1,h}(p_0(X)),X)) - g_{1,h}(p_0(X)) \right] \\ & + \frac{S(1-D)}{1-m(X)} \frac{g'_{6,h}(\bar{\beta}(X))g_{1,h}(p_0(X))}{p_0(X)g_{1,h}(1/p_0(X))} q_0(g_{1,h}(1/p_0(X)),X) \\ & \qquad \times \left[1(Y \leq q_0(g_{1,h}(1/p_0(X)),X) - \beta_{U,1,h}(X))g'_{1,h}(p_0(X)) \right] \\ & + (q_0(g_{1,h}(1/p_0(X)),X) - \beta_{U,0,h}(X)) \frac{g'_{1,h}(1/p_0(X))g_{1,h}(p_0(X))}{g_{1,h}(1/p_0(X))p_0^2(X)} \right] \\ &
$$

A.8. Proof of Theorem [6.1](#page-28-0)

Define $G_n[X] = n^{-1/2} \sum_i^n (X_i - E[X_i])$. We suppress the parameter $\beta_B = \beta_B(s_0, s_1)$ in what follows whenever it does not cause confusion. For any $B \in \{L, U\}$, denote $M(\eta) = \psi_{\beta_B}(W, \eta)$ as the influence function for β_B . Also denote the uncentered influence functions for the corresponding numerator and denominator as $\psi^{[B]}(W,\eta) = \psi^{[B]}(\eta)$ and $\psi^{[S]}(W,\eta) = \psi^{[S]}(\eta)$. Recall that the sample analogues using estimated nuisances $\hat{\eta}$ $\hat{\eta}(X)$ are then given by

$$
\hat{\beta}_B = \frac{E_n[\psi^{[B]}(W,\hat{\eta})]}{E_n[\psi^{[S]}(W,\hat{\eta})]}.
$$

Thus, we obtain the following linearization

$$
\sqrt{n}(\hat{\beta}_B - \beta_B) = \sqrt{n} E_n[M(\hat{\eta}, \beta_B)] \left(1 + O\left(\sup_{b \in \{B, S\}} |E_n[\psi^{[b]}(\hat{\eta})] - E[\psi^{[b]}(\eta)]| \right) \right)
$$

$$
= G_n[M(\hat{\eta}, \beta_B)] \left(1 + O\left(\sup_{b \in \{B, S\}} |E_n[\psi^{[b]}(\hat{\eta})] - E[\psi^{[b]}(\eta)]| \right) \right)
$$

by Assumption [6.3.](#page-28-1) Further decomposing the leading term yields

$$
G_n[M(\hat{\eta}, \beta_B)] = G_n[M(\eta, \beta_B)] + (G_n[M(\hat{\eta}, \beta_B)] - G_n[M(\eta, \beta_B)])
$$

$$
+ \sqrt{n}E[M(\hat{\eta}, \beta_B) - M(\eta, \beta_B)].
$$

For any values of the nuisances parameters, note that the moments can be decomposed as the (finite) sum over its (finite) product components. In particular, we can write

$$
M(W, \eta) = \sum_{r} m_r(W, \eta)
$$

=
$$
\prod_{j \in \mathcal{J}(r)} m_{r,j}(W, \eta),
$$

where $\mathcal{J}(r)$ is the set of factors for multiplication in the *r*-th summand of $M(\eta)$. We now further omit *W* in what follows in all components for convenience. Decomposing the difference in moments evaluated at the difference nuisance parameters then yields

$$
M(\hat{\eta}) - M(\eta) = \sum_{r} (m_r(\hat{\eta}) - m_r(\eta))
$$

= $\sum_{r} \Biggl(\prod_{j \in \mathcal{J}(r)} m_{r,j}(\hat{\eta}) - \prod_{j \in \mathcal{J}(r)} m_{r,j}(\hat{\eta}) \Biggr)$
= $\sum_{r} \Biggl(\sum_{j} (m_{r,j}(\hat{\eta}) - m_{r,j}(\eta)) \prod_{k \neq j} m_{r,k}(\eta)$
+ $\sum_{j,k > j} (m_{r,j}(\hat{\eta}) - m_{r,j}(\eta)) (m_{r,k}(\hat{\eta}) - m_{r,k}(\eta)) \prod_{l \neq k,j} m_{r,l}(\eta)$
:
+ $\prod_{j} (m_{r,j}(\hat{\eta}) - m_{r,j}(\eta)) \Biggr).$

This decomposition has the following properties: For any *j*, all $\prod_{k\neq j} m_{r,k}(\eta)$ are either bounded or have at least two moments by Assumption [5.2](#page-21-0) and [6.1.](#page-26-0) Moreover, by the pathwise differentiability as shown in Proof of Theorem [5.1,](#page-22-0) we have that

$$
\sup_{\tilde{\eta}\in\mathcal{E}_n}\sum_r\sum_jE[(m_{r,j}(\tilde{\eta})-m_{r,j}(\eta))\prod_{k\neq j}m_{r,k}(\eta)]=0.
$$

The only component that needs a different bound compared to [Semenova](#page-46-0) [\(2024\)](#page-46-0) is the $m_{r,j}$ corresponding to the classification errors for \mathcal{X}^+ and \mathcal{X}^- . In particular, we have to bound

$$
E[|\mathbb{1}(\hat{p}_0(X) < 1) - \mathbb{1}(p_0(X) < 1)|] = E[|\mathbb{1}(\hat{s}(1, X) - \hat{s}(0, X) > 0) - \mathbb{1}(s(1, X) - s(0, X) > 0)|]
$$
\n
$$
= E[|\mathbb{1}(\hat{\Delta}(X) > 0) - \mathbb{1}(\Delta(X) > 0)|].
$$

We use the same decomposition as in [\(A.17\)](#page-57-0) to obtain

$$
E[|\mathbb{1}(\hat{\Delta}(X) > 0) - \mathbb{1}(\Delta(X) > 0)|] \le P(|\Delta(X)| > |\hat{\Delta}(X) - \Delta(X)|)
$$
\n
$$
= P(|\Delta(X)| > |\hat{\Delta}(X) - \Delta(X)|, |\hat{\Delta}(X) - \Delta(X)| > \delta)
$$
\n
$$
+ P(|\Delta(X)| > |\hat{\Delta}(X) - \Delta(X)|, |\hat{\Delta}(X) - \Delta(X)| \le \delta)
$$
\n
$$
\le P(|\hat{\Delta}(X) - \Delta(X)| > \delta) + P(|\Delta(X)| \le \delta)
$$
\n
$$
\le \frac{E[|\hat{\Delta}(X) - \Delta(X)|^2]}{\delta^2} + P(|\Delta(X)| \le \delta)
$$
\n
$$
\le \lambda_{s,n,2}^2 \delta^{-2} + C\delta^{\alpha},
$$

for any arbitrary $\delta > 0$ by Assumption [6.2.](#page-28-2) Optimizing rates by setting $\delta^{2+\alpha} \sim \lambda_{s,n,2}^2$ then yields

$$
E[|\mathbb{1}(\hat{p}_0(X) < 1) - \mathbb{1}(p_0(X) < 1)|] \lesssim \lambda_{s,n,2}^{\frac{2\alpha}{2+\alpha}},
$$

where $2\alpha/(2 + \alpha) \in (1, 2)$ for any $\alpha > 2$.

We now use this in combination with Semenova (2024), Version 3, Lemma A.7, for the corresponding remaining terms given a correct classification to obtain

$$
\sqrt{n}\sup_{\tilde{\eta}\in\mathcal{E}_n}|E[M(\tilde{\eta},\beta_B)-M(\eta,\beta_B)]\lesssim \sqrt{n}(\lambda_{q,n,2}^2+\lambda_{s,n,2}^2+\lambda_{m,n,2}^2++\lambda_{b,n,2}^2+\lambda_{s,n,2}^{\frac{2\alpha}{2+\alpha}}).
$$

Moreover, as the nuisances are cross-fitted, we also obtain as in [Chernozhukov et al.](#page-44-0) [\(2018\)](#page-44-0) or [Kennedy](#page-46-1) [\(2023\)](#page-46-1), Lemma 1 that

$$
\sup_{\tilde{\eta}\in\mathcal{E}_n} E[(M(\tilde{\eta},\beta_B)-M(\eta,\beta_B))^2]^{1/2}\lesssim \lambda_{q,n,1}+\lambda_{q,n,2}+\lambda_{s,n,1}+\lambda_{s,n,2}+\lambda_{m,n,2}+\lambda_{b,n,2}+\lambda_{s,n,2}^{\frac{\alpha}{2+\alpha}}.
$$

Hence, by Assumption [6.3](#page-28-1) and Slutzky's Theorem, we obtain that

$$
\sqrt{n}(\hat{\beta}_B - \beta_B) \stackrel{d}{\rightarrow} \mathcal{N}(0, Var_a(\beta_B)),
$$

where $Var_a(\beta_B)$ is the semiparametric efficiency bound from Theorem [5.1.](#page-22-0)

A.9. Proof of Theorem [6.2](#page-29-0)

First, we provide some definitions and auxiliary convergence results and then apply an analogous decomposition as in the proof of Theorem [6.1](#page-28-0) with these new, modified terms. Let $M_h(\eta) = M_h(W, \eta, \beta_{B,h})$ be the influence curve for $\beta_{B,h}$ based on Table [6.2.](#page-27-0) Recall that

$$
\beta_{B,h} = \beta_{B,+,h} + \beta_{B,-,h}
$$

and sample analogue using estimated nuisances $\hat{\eta} = \hat{\eta}(X)$ given by

$$
\hat{\beta}_{B,h} = \hat{\beta}_{B,+,h} + \hat{\beta}_{B,-,h}
$$
\n
$$
= \frac{E_n[\psi_{\beta_{B,+,h}}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_{B,+,h}}^{[S]}(W,\hat{\eta})]} + \frac{E_n[\psi_{\beta_{B,-,h}}^{[B]}(W,\hat{\eta})]}{E_n[\psi_{\beta_{B,-,h}}^{[S]}(W,\hat{\eta})]}.
$$

Thus, we obtain the following linearization

$$
\sqrt{n}(\hat{\beta}_{B,h} - \beta_{B,h}) = G_n[M_h(\hat{\eta}, \beta_{B,h})] \bigg(1 + O\bigg(\sup_{b \in \{B,S\}, j \in \{+,-\},} |E_n[\psi_{\beta_{B,j,h}}^{[b]}(\hat{\eta})] - E[\psi_{\beta_{B,j,h}}^{[b]}(\eta)]|\bigg)\bigg),
$$

by Assumption [6.4.](#page-29-1) Further decomposing the leading term yields

$$
G_n[M(\hat{\eta}, \beta_{B,h})] = G_n[M(\eta, \beta_{B,h})] + (G_n[M(\hat{\eta}, \beta_{B,h})] - G_n[M(\eta, \beta_{B,h})])
$$

$$
+ \sqrt{n}E[M(\hat{\eta}, \beta_{B,h}) - M(\eta, \beta_{B,h})].
$$

We again omit W in what follows in all components for convenience. Decomposing the difference in moments evaluated at the difference nuisance parameters then yields

$$
M_h(\hat{\eta}) - M_h(\eta) = \sum_r (m_{h,r}(\hat{\eta}) - m_r(\eta))
$$

= $\sum_r \Bigg(\prod_{j \in \mathcal{J}(r)} m_{h,r,j}(\hat{\eta}) - \prod_{j \in \mathcal{J}(r)} m_{h,r,j}(\hat{\eta}) \Bigg)$
= $\sum_r \Bigg(\sum_j (m_{h,r,j}(\hat{\eta}) - m_{h,r,j}(\eta)) \prod_{k \neq j} m_{h,r,k}(\eta)$
+ $\sum_{j,k > j} (m_{h,r,j}(\hat{\eta}) - m_{h,r,j}(\eta)) (m_{r,k}(\hat{\eta}) - m_{h,r,k}(\eta)) \prod_{l \neq k,j} m_{h,r,l}(\eta)$
:
+ $\prod_j (m_{h,r,j}(\hat{\eta}) - m_{h,r,j}(\eta)) \Bigg).$

This decomposition has again the following properties. For any *j*, all $\prod_{k \neq j} m_{h,r,k}(\eta)$ are either bounded or have at least two moments by Assumption [5.2](#page-21-0) and [6.1.](#page-26-0) Moreover, by pathwise differentiability as in Theorem [5.2,](#page-23-0) we have that

$$
\sup_{\tilde{\eta}\in\mathcal{E}_n}\sum_r\sum_jE[(m_{h,r,j}(\tilde{\eta})-m_{h,r,j}(\eta))\prod_{k\neq j}m_{h,r,k}(\eta)]=0.
$$

h is fixed and bounded away from zero, hence h^{-1} is bounded from above. Note that, for any $j \in \{1, ..., 5\}$

$$
\label{eq:1.1} \begin{array}{c} g_{j,h}'(z)=1/(1+\exp(-z/h))\lesssim 1,\\ \\ g_{j,h}''(z)=g_{j,h}'(z)(1-g_{j,h}'(z))h^{-1}\lesssim h^{-1}\lesssim 1. \end{array}
$$

By definition of the components of the moment functions, they are either equivalent to the components of $M(W, \eta)$ and independent of h whenever no smoothing is involved or, if it is, product of functions that are either bounded or have at least two moments times the differences of $g_{j,h}$ -smoothed functions instead of their originals. By the fact that the latter have uniformly bounded derivatives, the mean value theorem implies that the differences between the $g_{j,h}$ functions evaluated at different nuisances are equal to an almost surely bounded function times the differences in nuisances. Thus, for all but one component, we can use the equivalent bounds to proof of Theorem [6.1.](#page-28-0)

For the classification error, however, we now can use a different rate as there are no indicators in the corresponding $m_{h,r,j}(\hat{\eta}) - m_{h,r,j}(\eta)$ components anymore. First note that, as *h* is bounded away from zero, $g_{1,h}(p_0(x))$ and $1 - g_{1,h}(1/p_0(x))$ are uniformly bounded away from one and zero respectively. The same holds true almost surely on the realization set using $\hat{p}_0(x)$ instead. Thus, the truncation thresholds in both β_h or $\hat{\beta_h}$ are contained in a subset of (0*,* 1), respectively, and hence we can bound

$$
E[(g_{4,h}(\hat{\beta}_h(X)) - g_{4,h}(\beta_h(X)))^2] \lesssim \sup_{\tilde{\beta}_h \in \mathcal{B}_n} E[(\hat{\beta}_h(X) - \beta_h(X))^2]
$$

$$
\lesssim \sup_{j,d \in \{0,1\}} \sup_{\tilde{\beta} \in \mathcal{B}_n} \sup_{u \in (0,1)} E[(\tilde{\beta}_{j,d}(X,u) - \beta_{j,d}(X,u))^2]
$$

$$
= \lambda_{b,n,2}^2.
$$

A.10. Proof of Theorem [6.3](#page-30-0)

We first show the proof under the positive monotonicity assumption. We suppress the nuisance functions, parameter, and principal strata in any moment function in what follows. First note that the estimator $\tilde{\beta}_L$ of [Semenova](#page-46-0) [\(2024\)](#page-46-0) is based on proposing separate moments for numerator and denominator. The corresponding variance bound is established in Theorem 1 of [Semenova](#page-46-0) [\(2024\)](#page-46-0) by an application of the delta method. First, we consider the moment function suggested for the known propensity score case, namely, denoting $W = (YS, S, D, X)$ the uncentered moments for numerator and denominator are given by

$$
\tilde{\psi}_{\beta}^{[L]}(W) = \frac{SD}{m(X)} Y1\{Y \le q_1(p_0(X), X)\} - \frac{S(1 - D)}{(1 - m(X)}Y - \frac{SD}{m(X)}q_1(p_0(X), X)[1\{Y \le q_1(p_0(X), X)\} - p_0(X)] + q_1(p_0(X), X)\left[\frac{1 - D}{1 - m(X)}(S - s(0, X)) - p_0(X)\frac{D}{m(X)}(S - s(1, X))\right]
$$

$$
\tilde{\psi}_{\beta}^{[S]} = s(0, X) + \frac{1 - D}{1 - m(X)}(S - s(0, X)).
$$

The centered versions of these moments are given by

$$
\tilde{\psi}_{\beta}^{[L]}(W) - E[\tilde{\psi}_{\beta}^{[L]}(W)]
$$
\n
$$
= \frac{SD}{m(X)} Y1\{Y \le q_1(p_0(X), X)\} - \frac{S(1 - D)}{(1 - m(X))} Y - \beta E[s(0, X)]
$$
\n
$$
- \frac{SD}{m(X)} q_1(p_0(X), X) [1\{Y \le q_1(p_0(X), X)\} - p_0(X)]
$$
\n
$$
+ q_1(p_0(X), X) \left[\frac{1 - D}{1 - m(X)} (S - s(0, X)) - p_0(X) \frac{D}{m(X)} (S - s(1, X)) \right],
$$
\n
$$
\tilde{\psi}_{\beta}^{[S]}(W) - E[\tilde{\psi}_{\beta}^{[S]}(W)] = s(0, X) - E[s(0, X)] + \frac{1 - D}{1 - m(X)} (S - s(0, X)).
$$

Define the centered ratio moment as

$$
\tilde{\psi}_{\beta}(W) = E[s(0, X)]^{-1} (\tilde{\psi}_{\beta}^{[L]}(W) - \beta_L \tilde{\psi}_{\beta}^{[S]}(W) - E[\tilde{\psi}_{\beta}^{[L]}(W) - \beta_L \tilde{\psi}_{\beta}^{[S]}(W)])
$$

Thus, we can write the difference between our influence function and the linearized ratio moment as

$$
\delta^{+}(D, X) = \psi_{\beta}^{[L]}(W) - \tilde{\psi}_{\beta}^{[L]}(W)
$$

= $E[s(0, X)]^{-1} s(0, X) \left[\beta_{1,1}(X, p_0(X)) \left(1 - \frac{D}{m(X)} \right) - \beta_{0,0}(X, 0) \left(1 - \frac{1 - D}{1 - m(X)} \right) \right]$

It turns out that this is the difference to the moment function under unknown propensity scores as suggested by [Semenova](#page-46-0) [\(2024\)](#page-46-0), Section 6.6, i.e. the linearized version of the latter is equivalent to $\psi_{\beta}^{[L]}$ $\phi_{\beta}^{[L]}(W)$. Now note that, as $\psi_{\beta}^{[L]}$ $\beta_{\beta}^{[L]}(W)$ is efficient when $P(\mathcal{X}_0) = 0$, we must have that $Cov(\psi_{\beta}^{[L]})$ $\tilde{\psi}_{\beta}^{[L]}(W), \tilde{\psi}_{\beta}^{[L]}$ $\mu_{\beta}^{[L]}(W))=E[\psi_{\beta}^{[L]}]$ $\int_{\beta}^{\lfloor L \rfloor} (W)^2$] and thus the variance gap in the regular case is given by

$$
E[\delta^+(D, X)^2]
$$

= $-E[s(0, X)]^{-2}E\left[s(0, X)^2(\beta_{1,1}(X, p_0(X))\sqrt{\frac{1 - m(X)}{m(X)}} - \beta_{0,0}(X, 0)\sqrt{\frac{m(X)}{1 - m(X)}})\right]^2$.

For conditional monotonicity, the proof is analogous within partitions where now

$$
E[\min\{s(0, X), s(1, X)\}]\delta(D, X)
$$

= $\mathbb{1}_{X \in \mathcal{X}^{+}} s(0, X) \left[\beta_{1,1}(X, p_{0}(X)) \left(1 - \frac{D}{m(X)}\right) - \beta_{0,0}(X, 0) \left(1 - \frac{1 - D}{1 - m(X)}\right)\right]$
+ $\mathbb{1}_{X \in \mathcal{X}^{-}} s(1, X) \left[\beta_{1,1}(X, 1) \left(1 - \frac{D}{m(X)}\right) - \beta_{0,0}(X, 1 - 1/p_{0}(X)) \left(1 - \frac{1 - D}{1 - m(X)}\right)\right],$

which yields variance difference

$$
E[\min\{s(0, X), s(1, X)\}]^{2} (E[\psi_{\beta}^{[L]}(W)^{2}] - E[\tilde{\psi}_{\beta}^{[L]}(W)^{2}])
$$

=
$$
-E\left[\mathbb{1}_{\{X \in \mathcal{X}^{+}\}}s(0, X)^{2}\left(\beta_{1,1}(X, p_{0}(X))\sqrt{\frac{1 - m(X)}{m(X)}} - \beta_{0,0}(X, 0)\sqrt{\frac{m(X)}{1 - m(X)}}\right)^{2}\right]
$$

$$
-E\left[\mathbb{1}_{\{X \in \mathcal{X}^{-}\}}s(1, X)^{2}\left(\beta_{1,1}(X, 1)\sqrt{\frac{1 - m(X)}{m(X)}} - \beta_{0,0}(X, 1 - 1/p_{0}(X))\sqrt{\frac{m(X)}{1 - m(X)}}\right)^{2}\right].
$$

B. Supplementary Material for Section [8](#page-34-0)

B.1. Data

We use an extended set of covariates containing detailed information regarding demographics, employment, criminal history, education, health, expectations, regional characteristics, and other JC related information similar to [Lee](#page-46-2) [\(2009\)](#page-46-2), [Flores et al.](#page-44-1) [\(2012\)](#page-44-1), and [Heiler and Knaus](#page-45-0) [\(2023\)](#page-45-0). In particular, they contain all the pre-treatment covariates from [Lee](#page-46-2) [\(2009\)](#page-46-2) and [Heiler and Knaus](#page-45-0) [\(2023\)](#page-45-0), including all randomization variables. We removed all observations with a missing earnings or working hours entry at any week $t \in \{1, \ldots, 208\}$, i.e. the analysis is conditional on that. All missing covariate entries were imputed by their mean in the final research sample as in [Lee](#page-46-2) [\(2009\)](#page-46-2). The missingness rates for covariates are relatively low for most variables other than earnings.

The Job Corps has varying treatment and control probabilities even conditional on the research sample based on gender (*F EMALE*), week of randomization (*RAND*_*WK*, area concentration of nonresidential female students (*IN*57), and prediction regarding the residential status (*NONRES*), see [Burghardt et al.](#page-44-2) [\(1999\)](#page-44-2). Based on this, we constructed treatment propensities for the research sample. We also constructed design weights to account for differences between research and national JC population for the impact evaluation. Table [B.1](#page-79-0) contains the research sample propensities using the variables *RAND*_*WK*, *F EMALE*, *IN*57, and *NONRES* from the public use files.

rable D.1. Research Sample Treatment I ropensities								
RAND WK	≤ 41							
<i>FEMALE</i>								
<i>IN</i> 57								
<i>NONRES</i>		0				0		
$P(D = 1 X = x)$	15.4 $8 + 15.4$	10.7 $8 + 10.7$	15.4 $5 + 15.4$	10.7 $5 + 10.7$	15.4 $8 + 15.4$	10.7 $8 + 10.7$	15.4 $8 + 15.4$	10.7 $8 + 10.7$
RAND WK	≥ 41							
<i>FEMALE</i>								
IN57								
<i>NONRES</i>		0						
$P(D = 1 X = x)$	17 $9 + 17$	11.1 $9 + 11.1$	17 $5 + 17$	11.1 $5 + 11.1$	17 $9 + 17$	11.1 $9 + 11.1$	17 $8 + 17$	11.1 $8 + 11.1$

Table B.1: Research Sample Treatment Propensities

We make use of 77 pre-treatment covariates on top of the treatment assignment vari-

Figure B.1: Relative Performance Metrics for all Selection Models

able. Table [B.2](#page-82-0) contains the average covariates conditional on observing the earnings and hours at all periods for treated and control groups. We also report the standardized mean differences as well as *p*-values for the two-sided hypothesis of equal means between treated and control. The sample is still well-balanced with the exception of worries to attend Job Corps (HADWORRY) that are higher in the treatment group.

B.2. Estimation

This section contains details on the estimation methods in Table [8.1.](#page-37-0) Logit is a logistic regression that contains all confounders. Logit interacted is a logistic regression that contains all confounders interacted with the treatment variable. XGBoost are gradient boosted trees using the binary logistic objective functions with maximum depth of 5 and learning rate $\eta = 0.1$ and cross-validated boosting steps. Neural Network is a 2 hidden layer fully connected feed forward artificial neural network with rectified linear activation functions and a final sigmoid layer. Every layer has 0.2 dropout regularization. It is trained using binary cross-entropy with batch size 128 and 50 epochs. The latter were selected for the best cross-validation error in period *t* = 208. Random Forest are honest probability forests using 1000 trees and default parameters in grf.

Table [B.1a](#page-80-0) and [B.1b](#page-80-0) contain the accuracy and entropy for all periods $t \in \{1, \ldots, 208\}$. Higher accuracy and lower entropy loss are better. One can see that XGBoost is the dominant model except for the very first period. The summary ranks over all periods are in Table [8.1.](#page-37-0)

For the conditional quantile models, we use honest quantile forests with 1000 trees and default parameters from grf. The model was obtained using the fully selected sample and including the treatment indicator as well as all confounders.

For the truncated conditional expectations note that

$$
E\left[\frac{Y1(Y \le q_1(p_0(X), X))}{p_0(X)}\middle| SD = 1, X = x\right] = E[Y|SD = 1, Y \le q_1(p_0(X), X)]
$$

Thus, the truncated conditional expectations can be obtained from an auxiliary regression that uses dependent variable $Y1 \ (Y \leq q_1(p_0(X), X))/p_0(X)$ and regressors X on the $S = D = 1$ sample. The same applies equivalently to the other three truncated expectations. The specific models were estimated with a fully tuned regression forest. The unknown relative selection probabilities and quantiles were first estimated on the same training set on which the auxiliary model is estimated. This assures that the models for the truncated conditional expectations respect the same independencies as conventional cross-fitting.

	Treated	Control	std.diff	p-marginal	p-multiple
FEMALE	0.4609	0.4662	-0.0095	0.6630	1.0000
AGE	19.2696	19.1506	0.0288	0.1958	1.0000
MARRIED	0.0202	0.0234	-0.0217	0.3185	1.0000
TOGETHER	0.0394	0.0410	-0.0081	0.7088	1.0000
SEPARATED	0.0247	0.0216	0.0201	0.3479	1.0000
HASCHLD	0.1932	0.1960	-0.0071	0.7408	1.0000
NCHLD HGC	0.2731 10.3163	0.2697 10.2935	0.0053 0.0091	0.8072 0.6806	1.0000 1.0000
HGC_MOTH	11.7133	11.6768	0.0113	0.6040	1.0000
HGC FATH	11.6424	11.7148	-0.0227	0.2987	1.0000
EVARRST	0.2452	0.2448	0.0010	0.9636	1.0000
HH INC2	0.2115	0.2124	-0.0028	0.8974	1.0000
HH INC3	0.1188	0.1165	0.0086	0.6877	1.0000
HH_INC4	0.2489	0.2490	-0.0001	0.9962	1.0000
HH INC5	0.1822	0.1838	-0.0050	0.8170	1.0000
PERS_INC2	0.1303	0.1334	-0.0094	0.6613	1.0000
PERS_INC3 PERS_INC4	0.0535	0.0473	0.0289	0.1748	1.0000
	0.0321	0.0346	-0.0141	0.5126	1.0000
CURRJOB.y	0.2015	0.1956	0.0143	0.5045	1.0000
MOSINJOB YR_WORK	3.6672 0.6484	3.5971 0.6392	0.0157 0.0180	0.4637 0.4036	1.0000 1.0000
EARN_YR	2962.0485	2868.9746	0.0168	0.4137	1.0000
HRSWK_JR	22.2540	21.3020	0.0433	0.0428	1.0000
WKEARNR	113.2187	104.8441	0.0307	0.1151	1.0000
RACE W	0.2717	0.2680	0.0080	0.7081	1.0000
RACE B	0.5029	0.5006	0.0044	0.8383	1.0000
RACE_H	0.1727	0.1761	-0.0087	0.6846	1.0000
RACE_O	0.0727	0.0738	-0.0041	0.8502	1.0000
$EDUC_GR$	1.4706	1.4720	-0.0016	0.9406	1.0000
LIVESPOU	0.0605	0.0645	-0.0164	0.4475	1.0000
EVERWORK	0.8079	0.7935	0.0322	0.1354	1.0000
$JOB0_3$	0.1912	0.1902	0.0026	0.9029	1.0000
$JOB3_9$	0.2817	0.2741	0.0168	0.4322	1.0000
$JOB9_12$	0.1756	0.1749	0.0017	0.9364	1.0000
MOSTWELF GOT_AFDC	0.2065 0.3199	0.2040 0.3200	0.0060 -0.0002	0.7785 0.9915	1.0000
GOT FS	0.4564	0.4685	-0.0234	0.2787	1.0000 1.0000
ED0 6	0.2782	0.2566	0.0474	0.0269	1.0000
ED6 12	0.3603	0.3866	-0.0528	0.0139	0.9599
PUBLICH	0.2137	0.2018	0.0294	0.1693	1.0000
BADHLTH	0.1315	0.1399	-0.0239	0.2670	1.0000
HARDUSE	0.0655	0.0625	0.0118	0.5811	1.0000
POTUSE	0.2513	0.2411	0.0233	0.2760	1.0000
PMSA	0.3282	0.3187	0.0201	0.3483	1.0000
MSA	0.4655	0.4595	0.0116	0.5879	1.0000
HS _D	0.2000	0.1984	0.0040	0.8540	1.0000
GED _{_D}	0.0486	0.0572	-0.0375	0.0835	1.0000
VOC_D	0.0218	0.0197	0.0145	0.4970	1.0000
ANY $ED1$ NTV_LANG	0.6821 1.2063	0.6815 1.2016	0.0012 0.0085	0.9540 0.6939	1.0000 1.0000
R HEAD	0.1156	0.1256	-0.0309	0.1530	1.0000
HHMEMB	3.0350	2.9214	0.0622	0.0039	0.2717
HEALTH	1.7064	1.7137	-0.0088	0.6838	1.0000
PY_CIG	0.5283	0.5072	0.0402	0.0601	1.0000
${\rm PY}$ ALCHL	0.5547	0.5314	0.0444	0.0382	1.0000
$\bar{P}Y_P$ OT	0.3075	0.2966	0.0231	0.2802	1.0000
HADWORRY	0.3726	0.3339	0.0768	0.0003	0.0239
HEAR JC	3.3126	3.2811	0.0179	0.4058	1.0000
KNEWCNTR	0.5184	0.5334	-0.0286	0.1833	1.0000
E_MATH	0.7051	0.6911	0.0282	0.1904	1.0000
E _{READ}	0.5454	0.5444	0.0020	0.9269	1.0000
E ALONG E CONTRL	0.6160	0.6111	0.0094	0.6616	1.0000
E ESTEEM	0.5887 0.5926	0.6040 0.5922	-0.0293 0.0008	0.1720 0.9706	1.0000 1.0000
E SPCJOB	0.9696	0.9743	-0.0167	0.4437	1.0000
E FRIEND	0.7220	0.7070	0.0305	0.1567	1.0000
KNEW_JC	0.6858	0.6884	-0.0053	0.8064	1.0000
NONRES	0.1514	0.1581	-0.0206	0.3451	1.0000
PRARRI	0.7502	0.7443	0.0263	0.2244	1.0000
RAND_WK	34.4925	34.9669	-0.0285	0.1859	1.0000
IN57	0.3239	0.3298	-0.0133	0.5311	1.0000

Table B.2: Job Corps Data: Covariates Balance

Column 1 and Column 2 contain the average covariates within treatment and control group respectively. Standardized differences are in Column 3. Unadjusted *p*-values for differences in means and adjusted for multiple comparisons are in Column 5 and 6 respectively. Sample $n = 9415$ obtained from non-missing weekly hours worked and weakly earnings over all weeks $t \in \{1, \ldots, 208\}$. Calculations use research sample and treatment propensity weights.

Figure B.2: Example Distributions for $\hat{p}_0(x)$

The figures contains histograms for the estimated $p_0(x)$ using five different estimation methods on the research sample at weeks $t \in \{45, 90, 135, 180, 208\}$. glm1 are logistic regressions with linear additive index. glm2 are fully treatment interacted logistic models. xgb are gradient boosted trees with crossvalidated boosting steps using xgboost. nnet are batch-trained, fully connected artificial feed-forward neural network with three ReLu hidden layers, drop-out regularization, and final sigmoid layers using keras. grf are honest probability forests using grf with default parameters. The spike at $p_0(x) = 1$ for glm2 at $t = 45$ is due to non-convergence within some folds.

B.3.1. Effect Bounds

					$\check{ }$	
		$t=45$	$t=90$	$t=135$	$t=180$	$t = 208$
Smooth	$h=5$	$[-0.182, 0.223]$ $(-0.247, 0.263)$	$[-0.133, 0.206]$ $(-0.182, 0.247)$	$[-0.086, 0.149]$ $(-0.117, 0.181)$	$[-0.096, 0.151]$ $(-0.123, 0.181)$	$[-0.085, 0.171]$ $(-0.113, 0.201)$
	$h=1$	$[-0.164, 0.214]$ $(-0.232, 0.257)$	[0.018, 0.081] $(-0.025, 0.123)$	$[-0.015, 0.106]$ $(-0.054, 0.144)$	$[-0.041, 0.111]$ $(-0.077, 0.148)$	$[-0.030, 0.131]$ $(-0.067, 0.168)$
	$h=0.5$	$[-3.601, 0.213]$ $(-3.660, 0.256)$	$[-3.210, 0.064]$ $(-3.267, 0.108)$	$[-0.043, 0.102]$ $(-0.089, 0.139)$	$[-0.064, 0.110]$ $(-0.108, 0.147)$	$[-0.062, 0.130]$ $(-0.106, 0.167)$
Trim	Known	$[-0.135, 0.195]$ $(-0.172, 0.237)$	[0.052, 0.101] $(-0.021, 0.188)$	$[-0.007, 0.121]$ $(-0.046, 0.155)$	$[-0.034, 0.119]$ $(-0.071, 0.152)$	$[-0.024, 0.142]$ $(-0.061, 0.176)$
	Unknown	$[-0.222, 0.112]$ $(-0.342, 0.227)$	$[-0.052, -0.004]$ $(-0.187, 0.140)$	$[-0.105, 0.041]$ $(-0.214, 0.149)$	$[-0.112, 0.076]$ $(-0.224, 0.188)$	$[-0.124, 0.068]$ $(-0.239, 0.183)$
Shift	Known	$[-0.582, -0.242]$ $(-0.828, -0.018)$	[0.237, 1.895] $(-0.008, 2.084)$	[0.464, 3.011] (0.267, 3.136)	[0.541, 3.136] (0.352, 3.255)	[0.566, 3.257] (0.380, 3.373)
	Unknown	$[-0.221, 0.113]$ $(-0.341, 0.229)$	$[-0.023, -0.023]$ $(-0.202, 0.155)$	$[-0.105, 0.042]$ $(-0.214, 0.150)$	$[-0.111, 0.077]$ $(-0.223, 0.189)$	$[-0.121, 0.070]$ $(-0.236, 0.185)$
	$ \hat{p}_0(x) - 1 \leq n^{-1/4} / \log(n)$	$\overline{4}$	5320	4	3	5

Table B.3: Job Corps Results: Bounds with Selection Model Logit

		$t=45$	$t=90$	$t = 135$	$t=180$	$t=208$
Smooth	$h=5$	$[-12.082, 93.473]$ $(-13.379, 99.367)$	$[-0.351, 0.204]$ $(-1.610, 1.484)$	$[-4.914, 42.535]$ $(-6.244, 44.915)$	$[-2.766, 1.120]$ $(-3.312, 1.581)$	$[-2.146, 0.529]$ $(-2.314, 0.634)$
	$h=1$	$[-13.300, 86.263]$ $(-14.801, 91.929)$	$[-0.394, 0.276]$ $(-1.651, 1.554)$	$[-6.885, 54.294]$ $(-8.329, 57.852)$	$[-2.963, 1.066]$ $(-3.513, 1.536)$	$[-2.105, 0.459]$ $(-2.273, 0.565)$
	$h=0.5$	$[-13.750, 100.822]$ $(-15.404, 107.023)$	$[-2.167, 1.668]$ $(-4.717, 4.178)$	$[-6.267, 56.598]$ $(-7.643, 60.231)$	$[-3.668, 3.270]$ $(-4.228, 3.732)$	$[-2.940, 0.449]$ $(-3.111, 0.556)$
Trim	Known	$[-0.222, 0.296]$ $(-0.271, 0.348)$	[0.006, 0.109] $(-0.035, 0.152)$	$[-0.211, 0.316]$ $(-0.254, 0.363)$	[0.025, 0.076] $(-0.011, 0.114)$	[0.000, 0.101] $(-0.036, 0.140)$
	Unknown	$[-0.380, 0.228]$ $(-0.498, 0.344)$	$[-0.084, 0.039]$ $(-0.190, 0.146)$	$[-0.370, 0.372]$ $(-0.483, 0.486)$	$[-0.045, 0.021]$ $(-0.150, 0.128)$	$[-0.092, 0.037]$ $(-0.200, 0.146)$
Shift	Known	$[-0.558, 0.416]$ $(-1.061, 0.720)$	[0.196, 1.677] $(-0.019, 1.850)$	[1.676, 2.059] (0.096, 2.748)	[0.475, 2.117] (0.299, 2.248)	[0.473, 2.293] (0.298, 2.423)
	Unknown	$[-0.400, 0.141]$ $(-0.524, 0.261)$	$[-0.115, -0.010]$ $(-0.225, 0.101)$	$[-0.177, 0.335]$ $(-0.285, 0.442)$	$[-0.050, 0.063]$ $(-0.155, 0.169)$	$[-0.057, 0.047]$ $(-0.165, 0.157)$
$ \hat{p}_0(x)-1 $	$\leq n^{-1/4}/\log(n)$	1255	422	2749	467	510

Table B.4: Job Corps Results: Bounds with Selection Model Logit Interacted

Table B.5: Job Corps Results: Bounds with Selection Model XGBoost

		$t=45$	$t=90$	$t = 135$	$t=180$	$t=208$
Smooth	$h=5$	$[-0.109, 0.145]$ $(-0.172, 0.199)$	$[-0.112, 0.212]$ $(-0.161, 0.256)$	$[-0.093, 0.213]$ $(-0.139, 0.250)$	$[-0.072, 0.215]$ $(-0.107, 0.251)$	$[-0.057, 0.275]$ $(-0.095, 0.312)$
	$h=1$	$[-0.000, 0.161]$ $(-0.068, 0.209)$	[0.008, 0.096] $(-0.036, 0.135)$	$[-0.028, 0.090]$ $(-0.066, 0.125)$	$[-0.029, 0.093]$ $(-0.064, 0.128)$	$[-0.005, 0.113]$ $(-0.040, 0.148)$
	$h=0.5$	$[-3.641, 3.045]$ $(-3.705, 3.101)$	$[-3.224, 0.099]$ $(-3.275, 0.141)$	$[-0.078, 0.058]$ $(-0.126, 0.096)$	$[-0.093, 0.069]$ $(-0.141, 0.105)$	$[-0.091, 0.064]$ $(-0.139, 0.100)$
Trim	Known	$[-0.138, 0.160]$ $(-0.194, 0.217)$	[0.005, 0.014] $(-0.051, 0.081)$	$[-0.016, 0.054]$ $(-0.072, 0.106)$	$[-0.022, 0.087]$ $(-0.074, 0.140)$	$[-0.008, 0.054]$ $(-0.063, 0.111)$
	Unknown	$[-0.257, 0.126]$ $(-0.403, 0.269)$	$[-0.175, 0.004]$ $(-0.745, 0.573)$	$[-0.021, 0.152]$ $(-0.197, 0.327)$	$[-0.180, -0.011]$ $(-0.328, 0.134)$	$[-0.036, 0.077]$ $(-0.201, 0.245)$
Shift	Known	$[-0.756, 0.091]$ $(-1.085, 0.303)$	[0.238, 0.392] $(-0.093, 0.542)$	[0.549, 1.266] (0.291, 1.396)	[0.559, 1.911] (0.368, 2.052)	[0.646, 1.514] (0.438, 1.650)
	Unknown	$[-0.324, -0.076]$ $(-0.470, 0.067)$	$[-0.021, -0.011]$ $(-0.194, 0.162)$	[0.022, 0.067] $(-0.116, 0.204)$	[0.075, 0.122] $(-0.055, 0.253)$	[0.138, 0.162] $(-0.003, 0.303)$
$ \hat{p}_0(x)-1 $	$\leq n^{-1/4}/\log(n)$	2592	8680	6350	4871	5688

		$t=45$	$t=90$	$t=135$	$t=180$	$t = 208$
Smooth	$h=5$	$[-0.177, 0.089]$ $(-0.234, 0.137)$	$[-0.137, 0.214]$ $(-0.183, 0.255)$	$[-0.089, 0.187]$ $(-0.132, 0.220)$	$[-0.092, 0.182]$ $(-0.125, 0.215)$	$[-0.058, 0.228]$ $(-0.094, 0.263)$
	$h=1$	$[-0.469, 0.777]$ $(-0.543, 0.855)$	$[-0.005, 0.105]$ $(-0.049, 0.147)$	$[-0.023, 0.107]$ $(-0.061, 0.144)$	$[-0.045, 0.112]$ $(-0.080, 0.149)$	$[-0.022, 0.127]$ $(-0.058, 0.165)$
	$h=0.5$	$[-4.014, 3.815]$ $(-4.089, 3.906)$	$[-3.201, 0.103]$ $(-3.256, 0.145)$	$[-0.186, 0.182]$ $(-0.250, 0.223)$	$[-0.213, 0.210]$ $(-0.303, 0.282)$	$[-0.091, 0.096]$ $(-0.140, 0.132)$
Trim	Known	$[-0.252, 0.287]$ $(-0.315, 0.351)$	[0.050, 0.082] $(-0.038, 0.169)$	$[-0.014, 0.132]$ $(-0.076, 0.192)$	$[-0.036, 0.133]$ $(-0.090, 0.191)$	$[-0.035, 0.187]$ $(-0.091, 0.258)$
	Unknown	$[-0.347, 0.192]$ $(-0.481, 0.319)$	$[-0.031, 0.003]$ $(-0.179, 0.149)$	$[-0.107, 0.046]$ $(-0.223, 0.164)$	$[-0.141, 0.038]$ $(-0.266, 0.159)$	$[-0.157, 0.072]$ $(-0.284, 0.203)$
Shift	Known	$[-0.689, -0.033]$ $(-0.958, 0.189)$	[0.262, 2.277] (0.023, 2.462)	[0.476, 2.819] (0.283, 2.948)	[0.569, 2.947] (0.385, 3.070)	[0.585, 3.080] (0.402, 3.200)
	Unknown	$[-0.424, -0.122]$ $(-0.571, 0.019)$	$[-0.054, 0.011]$ $(-0.205, 0.162)$	[0.049, 0.108] $(-0.078, 0.235)$	$[-0.062, 0.076]$ $(-0.187, 0.199)$	[0.046, 0.121] $(-0.084, 0.252)$
$ \hat{p}_0(x)-1 $	$\leq n^{-1/4}/\log(n)$	737	4842	3220	2180	3669

Table B.6: Job Corps Results: Bounds with Selection Model Neural Network

Table B.7: Job Corps Results: Bounds with Selection Model Random Forest

		$t=45$	$t=90$	$t = 135$	$t=180$	$t=208$
Smooth	$h=5$	$[-0.126, 0.112]$ $(-0.187, 0.160)$	$[-0.141, 0.214]$ $(-0.190, 0.256)$	$[-0.081, 0.198]$ $(-0.129, 0.234)$	$[-0.076, 0.222]$ $(-0.112, 0.259)$	$[-0.074, 0.264]$ $(-0.113, 0.303)$
	$h=1$	$[-0.086, 0.185]$ $(-0.156, 0.237)$	[0.030, 0.075] $(-0.015, 0.114)$	$[-0.022, 0.096]$ $(-0.062, 0.132)$	$[-0.011, 0.083]$ $(-0.047, 0.118)$	[0.016, 0.098] $(-0.019, 0.135)$
	$h=0.5$	$[-3.666, 3.805]$ $(-3.731, 3.866)$	$[-3.182, 3.047]$ $(-3.234, 3.091)$	$[-0.071, 0.076]$ $(-0.122, 0.116)$	$[-0.082, 0.050]$ $(-0.132, 0.086)$	$[-0.061, 0.064]$ $(-0.109, 0.101)$
Trim	Known	$[-0.162, 0.202]$ $(-0.233, 0.273)$	[0.043, 0.100] $(-0.037, 0.189)$	$[-0.024, 0.138]$ $(-0.105, 0.209)$	[0.013, 0.121] $(-0.055, 0.192)$	[0.036, 0.120] $(-0.033, 0.196)$
	Unknown	$[-0.248, 0.116]$ $(-0.383, 0.244)$	$[-0.038, 0.019]$ $(-0.173, 0.159)$	$[-0.114, 0.054]$ $(-0.246, 0.178)$	$[-0.074, 0.034]$ $(-0.202, 0.160)$	$[-0.051, 0.044]$ $(-0.181, 0.177)$
Shift	Known	$[-0.679, -0.047]$ $(-0.958, 0.177)$	[0.213, 1.646] $(-0.028, 1.829)$	[0.446, 2.178] (0.247, 2.325)	[0.553, 2.429] (0.367, 2.571)	[0.542, 2.401] (0.357, 2.540)
	Unknown	$[-0.374, -0.110]$ $(-0.524, 0.033)$	$[-0.029, -0.028]$ $(-0.207, 0.150)$	[0.088, 0.090] $(-0.066, 0.244)$	[0.139, 0.145] $(-0.012, 0.295)$	[0.156, 0.165] (0.005, 0.315)
$ \hat{p}_0(x)-1 $	$\leq n^{-1/4}/\log(n)$	1052	6848	6420	6211	6342

C. Monte Carlo Study

In this section, we analyze properties of methods for inference under conditional monotonicity in finite samples. In particular, across a set of principal strata shares we consider the bias, root mean-squared error, size, and power of trimming bounds, switching bounds, and smooth bounds. Further, we analyze properties of smooth bounds along different values for the smoothing parameter *h*. We consider the following data-generating process

$$
S(0) = \mathbb{1}\{X_2 \ge V\}, \quad S(1) = \mathbb{1}(X_1 + X_2 \ge V), \quad Y(0) = 0,
$$

\n
$$
Y(1) = [U\mathbb{1}(S(1) = 1, S(0) = 1) + (U + \gamma)\mathbb{1}(S(1) = 1, S(0) = 0)] \mathbb{1}(X_1 = 1),
$$

\n
$$
D \sim \text{Bernoulli}(0.5), \quad U \sim \text{Uniform}(0, 1), \quad V \sim \mathcal{N}(0, 1),
$$

\n
$$
X_1 \sim \text{Categorical}(P(\mathcal{X}^+), P(\mathcal{X}^0), P(\mathcal{X}^-)) \text{ with support } X_1 \in \{1, 0, -1\},
$$

\n
$$
X_2 \sim \text{truncated } \mathcal{N}_{[-4,4]}(0, 1).
$$

This design implies that the target parameter, i.e. the treatment effect on always-takers, equals the true lower bound of the identified set. This makes one-sided tests on the lower bound equivalent to testing inclusion of the target parameter in the identified set. The target equals the lower bound because the always-takers and compliers are separated by a positive constant γ . This implies that units satisfying positive monotonicity are ordered such that always-takers never exhibit larger outcomes under treatment than compliers. This homogeneous design is setup in favor of the switching method by [Semenova](#page-46-0) [\(2024\)](#page-46-0) as the small mistakes when switching moments around the margin based on selection probabilities do not introduce systematic bias. It also favors low bias over low variance smoothing as the conditional variance is constant around the truncation threshold. We employ true nuisance functions throughout. All smooth approximations are based on the LogSumExp function.

We consider samples sizes $N = 400$ and $N = 2000$ across different strata shares $(P(X^+), P(X^0), P(X^-))$. The first design $(1/2, 0, 1/2)$ is regular, the second design (1*/*3*,* 1*/*3*,* 1*/*3) is irregular, and the third design (5*/*100*,* 95*/*100*,* 0) is highly irregular. Table [\(C.1\)](#page-89-0) and Figure [C.1](#page-90-0) reports results based on 2000 replications.

C.1. Results

Under the regular design reported in Table [C.1a,](#page-89-0) all methods are unbiased. Bias and RMSE decrease with sample size. All methods are approximately correctly sized. Table [C.1a](#page-89-0) and Figure [C.1a](#page-90-0) shows that methods with unknown propensity score display lower RMSE higher power as predicted by Theorem [6.3.](#page-30-0)

Under the irregular design reported in Table [C.1b,](#page-89-0) all methods are unbiased except the trim method with known propensity score. Methods with unknown propensity score again display lower RMSE. The switch and the smooth $(h = 10^{-9})$ methods are approximately correctly sized. The trim method with unknown propensity score is incorrectly sized as expected. Figure [C.1b](#page-90-0) shows that the switch method with unknown propensity score and the smooth method display higher power than the switch method with known propensity score. The power of trim methods is not well behaved.

Under the highly irregular design reported in Table [C.1c,](#page-89-0) all methods are again unbiased except the trim method with known propensity score. RMSE is again smaller for methods with unknown propensity score. The switch and the smooth methods are approximately correctly size, and the trim method with unknown propensity score is incorrectly sized, as in the irregular design. Figure [C.1c](#page-90-0) shows again that the switch method with unknown propensity score and the smooth method display higher power than with unknown propensity score. Again, the power of trim methods is not well behaved.

In summary, in this data-generating process, the switch and smooth methods are unbiased and approximately correctly sized across designs. RMSE is lower and power is higher for methods with unknown propensity score. As expected, the trim methods are only well behaved in regular designs. Other results might obtain for alternative datagenerating processes, for example the switch methods are theoretically biased, while the smooth method remains unbiased, under alternative distributions of $p_0(x)$.

The bias decreases in *h* across designs. RMSE also generally decreases except in the irregular design from $h = 10^{-2}$ to $h = 10^{-9}$. The size tends to one and power tends to zero as *h* increases. Future designs can evaluate potential bias-variance trade-offs and size distortion of switching methods when effects are heterogeneous.

Table C.1: Simulation results

Method	Bias	RMSE	Size
Switch (Known)	$-0.001;0.000$	0.065; 0.029	0.056; 0.052
Switch (Unknown)	$-0.001;0.000$	0.060; 0.027	0.058; 0.055
Trim (Known)	$-0.002; -0.001$	0.065; 0.029	0.053; 0.051
Trim (Unknown)	$-0.002;0.000$	0.059; 0.027	0.052; 0.054
Smooth			
$h = 5 \times 10^{-2}$	$-0.053; -0.051$	0.078; 0.057	0.005; 0.000
$h = 10^{-2}$	$-0.010; -0.011$	0.061; 0.028	0.042; 0.017
$h = 10^{-9}$	$-0.002;0.000$	0.060; 0.027	0.052; 0.054

(a) $(P(X^+), P(X^0), P(X^-)) = (1/2, 0, 1/2)$

(b) $(P(X^+), P(X^0), P(X^-)) = (1/3, 1/3, 1/3)$

Method	Bias	RMSE	Size
Switch (Known)	0.002; 0.000	0.052; 0.022	0.052; 0.042
Switch (Unknown)	0.002; 0.000	0.049; 0.021	0.054; 0.048
Trim (Known)	0.137; 0.136	0.158; 0.140	0.547; 0.982
Trim (Unknown)	0.001; 0.000	0.048; 0.021	0.020; 0.020
Smooth			
$h = 5 \times 10^{-2}$	$-0.043; -0.042$	0.064; 0.047	0.006:0.000
$h = 10^{-2}$	$-0.008; -0.008$	0.047; 0.023	0.034; 0.021
$h = 10^{-9}$	0.001; 0.000	0.048; 0.022	0.054; 0.046

(c) $(P(X^+), P(X^0), P(X^-)) = (5/100, 95/100, 0)$

Result before semicolon is $N = 400$ and after is $N = 2000$.

Solid lines are $N = 400$ and dotted lines are $N = 2000$.